



Corrigendum to

“Modelling of mineral dust for interglacial and glacial climate conditions with a focus on Antarctica” published in Clim. Past, 11, 765–779, 2015

N. Sudarchikova¹, U. Mikolajewicz¹, C. Timmreck¹, D. O’Donnell², G. Schurges^{3,*}, D. Sein⁴, and K. Zhang⁵

¹Max Planck Institute for Meteorology, Environmental Modeling, Hamburg, Germany

²Finnish Meteorological Institute, Climate Change Research, Helsinki, Finland

³Lund University, Department of Physical Geography and Ecosystem Science, Lund, Sweden

⁴Alfred Wegener Institute, Climate Dynamics, Bremerhaven, Germany

⁵Pacific Northwest National Laboratory, Atmospheric Sciences & Global Change Division, Richland, WA, USA

* now at: University of Copenhagen, Department of Geosciences and Natural Resource Management, Copenhagen, Denmark

Correspondence to: N. Sudarchikova (natalia.sudarchikova@mpimet.mpg.de)

Published: 2 September 2016

Table 2. Global mass budget for different climate conditions.

	CTRL	6 kyr	115 kyr	126 kyr	LGM
Emissions, [Tg yr ⁻¹]					
Total	1540	1961	1898	2385	3106
NH	1478	1845	1834	2273	2946
SH	62	117	65	112	160
Australia	60	115	63	110	84
South America	1	1.1	1.2	1.2	75
Southern Africa	1.2	0.9	1.1	0.8	1.6
Deposition, [Tg yr ⁻¹]					
Total dep.	1539	1959	1896	2376	3119
Wet dep.	643	828	790	1123	1293
Dry dep.	214	296	239	363	456
Sedimentation	682	835	867	890	1370
Deposition in Antarctica, [Tg yr ⁻¹]					
Total dep.	0.053	0.150	0.061	0.141	0.540

The correct e-mail address of the main author is natalia.sudarchikova@mpimet.mpg.de.

There was a typo in Table 2 in the NH LGM emissions. The correct NH LGM emissions amount to 2946 Tg yr⁻¹.