

# On Integrable Backgrounds Self-dual under Fermionic T-duality

---

Ido Adam<sup>1</sup>, Amit Dekel<sup>2</sup> and Yaron Oz<sup>2</sup>

<sup>1</sup> *Max-Planck-Institut für Gravitationsphysik  
Albert Einstein Institut  
Am Mühlenberg 1, 14476 Golm, Germany  
and*

*Kavli Institute for Theoretical Physics  
University of California  
Santa Barbara, CA 93106-4030, USA*

<sup>2</sup> *Raymond and Beverly Sackler School of Physics and Astronomy  
Tel-Aviv University, Ramat-Aviv 69978, Israel  
E-mails: idoadam@aei.mpg.de, amitde@post.tau.ac.il,  
yaronoz@post.tau.ac.il*

**ABSTRACT:** We study the fermionic T-duality symmetry of integrable Green-Schwarz sigma-models on AdS backgrounds with Ramond-Ramond fluxes in various dimensions. We show that sigma-models based on supercosets of PSU supergroups, such as  $AdS_2 \times S^2$  and  $AdS_3 \times S^3$  are self-dual under fermionic T-duality, while supercosets of OSp supergroups such as non-critical  $AdS_2$  and  $AdS_4$  models, and the critical  $AdS_4 \times CP^3$  background are not. We present a general algebraic argument to when a supercoset is expected to have a fermionic T-duality symmetry, and when it will fail to have one.

**KEYWORDS:** Duality in Gauge Field Theories, String Duality.

---

## Contents

<b>1. Introduction and summary</b>	<b>1</b>
<b>2. <math>AdS_p \times S^p</math> target-spaces</b>	<b>2</b>
2.1 The $AdS_2 \times S^2$ target-space	3
2.2 The $AdS_3 \times S^3$ target-space	5
<b>3. Models based on the ortho-symplectic supergroup</b>	<b>8</b>
3.1 The Green-Schwarz sigma-model on $AdS_2$ with four supersymmetries	8
3.2 The $AdS_4$ target-space	10
3.3 The $AdS_4 \times CP^3$ target-space	11
3.3.1 Current expansion in fermions	13
3.4 The $AdS_2$ target-space with eight supersymmetries	15
<b>4. A general analysis</b>	<b>16</b>
<b>A. The <math>osp(2 2)</math> algebra</b>	<b>18</b>
<b>B. The <math>osp(2 4)</math> algebra</b>	<b>19</b>
<b>C. The <math>osp(6 4)</math> algebra in <math>so(1,2) \oplus u(3)</math> basis</b>	<b>20</b>
<b>D. The <math>psu(1,1 2)</math> algebra</b>	<b>22</b>
<b>E. The <math>psu(1,1 2) \oplus psu(1,1 2)</math> algebra</b>	<b>23</b>

---

## 1. Introduction and summary

Recently, Alday and Maldacena proposed [1] that planar  $\mathcal{N} = 4$  SU(N) SYM MHV gluon scattering amplitudes at leading order in the strong 't Hooft coupling expansion can be calculated using the dual gravity (string) description. A crucial step in the calculation procedure is an application of an ordinary bosonic T-duality transformation to the four CFT coordinates of  $AdS_5 \times S^5$ . This suggestion implies that such amplitudes possess a dual conformal symmetry at strong coupling originating from the fact that  $AdS_5$  is self-dual under the T-duality. This dual symmetry has been observed also in gluon scattering amplitudes calculated in the weakly coupled gauge theory description [2, 3].

Using the AdS/CFT duality it was shown that such a symmetry is expected to be valid at all values of the 't Hooft coupling [4, 5]. This was done by proving that the aforementioned T-duality together with a novel T-duality of Grassmann-odd coordinates of the target-superspace form an exact quantum duality under which the full  $AdS_5 \times S^5$  superstring is self-dual. Since both the original background and its dual possess superconformal symmetry, it means that each of them has both the manifest superconformal symmetry and a dual one. Furthermore, [4, 5] have linked the duals of the superconformal Noether currents to the non-local currents implied by the integrability of the superstring on  $AdS_5 \times S^5$  [6].

In view of these results it is natural to inquire how ubiquitous this dual superconformal symmetry is. The aim of this paper is to consider this question by analyzing the fermionic T-duality symmetry of integrable Green-Schwarz sigma-models on AdS backgrounds with Ramond-Ramond fluxes in various dimensions. Some of these sigma-models have been constructed in [7, 8, 9, 10].

We show that sigma-models based on supercosets of PSU supergroups, such as  $AdS_2 \times S^2$  and  $AdS_3 \times S^3$  are self-dual under fermionic T-duality. Supercosets of OSp supergroups such as non-critical  $AdS_2$  and  $AdS_4$  models, and the critical  $AdS_4 \times CP^3$  background (whose coset model was constructed and explored in [11, 12, 13, 14, 15]) are not self-dual. In the OSp models we find that the Buscher procedure [16, 17] fails due to a lack of appropriate quadratic terms. This is because the Cartan-Killing bilinear form of the ortho-symplectic group is non-zero only for products of different Grassmann-odd generators. Thus, one may expect this to imply that in those cases in which a dual theory exists, the theory does not have a dual superconformal symmetry.

The paper is organized as follows. In section 2 we show that the  $AdS_p \times S^p$  ( $p = 2, 3$ ) target-spaces based on PSU cosets are self-dual under a combination of bosonic and fermionic T-duality. In section 3 we consider models based on supercosets of the ortho-symplectic supergroup, for which the Buscher procedure [16, 17] of gauging an isometry of the target-space in order to obtain the T-dual sigma-model fails. These include the non-critical superstring on  $AdS_2$  with four supersymmetries and  $AdS_4$  with eight supersymmetries, the supercoset construction of  $AdS_4 \times CP^3$ , and a model of  $AdS_2$  with eight supersymmetries. In section 4 we present a general algebraic argument to when a supercoset is expected to have a fermionic T-duality symmetry, and when it will fail to have one. In the appendices we provide details on the relevant superalgebras that are used in the paper.

## 2. $AdS_p \times S^p$ target-spaces

In this section we show that the  $AdS_p \times S^p$  ( $p = 2, 3$ ) target-spaces based on PSU supercosets are self-dual under a combination of bosonic and fermionic T-duality.

## 2.1 The $AdS_2 \times S^2$ target-space

The target superspace whose bosonic part is  $AdS_2 \times S^2$  can be realized as the coset space  $PSU(1, 1|2)/(U(1) \times U(1))^1$ . The Green-Schwarz sigma-model for supercosets with a  $\mathbb{Z}_4$  automorphism is given by the action [7]

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \text{Str} \left( J_2 \bar{J}_2 + \frac{1}{2} J_1 \bar{J}_3 - \frac{1}{2} \bar{J}_1 J_3 \right), \quad (2.1)$$

where  $J = g^{-1} \partial g$  for  $g \in G$  and  $J_i$  is the current  $J$  restricted to the invariant subspace  $\mathcal{H}_i$  of the  $\mathbb{Z}_4$  automorphism of the algebra of the group  $G$ . Using the  $psu(1, 1|2)$  algebra given in appendix D, the sigma-model (2.1) takes the form

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{1}{2} (J_P - J_K)(\bar{J}_P - \bar{J}_K) + \frac{1}{2} J_D \bar{J}_D + \frac{1}{2} J_{R_1} \bar{J}_{R_1} + \frac{1}{2} J_{R_2} \bar{J}_{R_2} - \frac{i}{2} \eta_{\alpha\beta} (J_{Q_\alpha} \bar{J}_{Q_\beta} - J_{\hat{Q}_\alpha} \bar{J}_{\hat{Q}_\beta} + J_{S_\alpha} \bar{J}_{S_\beta} - J_{\hat{S}_\alpha} \bar{J}_{\hat{S}_\beta}) \right], \quad (2.2)$$

where  $\eta_{12} = \eta_{21} = 1$  and zero otherwise. The analysis here will follow that of [4]. A general group element  $g \in PSU(1, 1|2)$  can be parameterized as

$$g = e^{xP + x'K + \theta^\alpha Q_\alpha + \xi^\alpha S_\alpha} e^B, \quad e^B \equiv e^{\hat{\theta}^\alpha \hat{Q}_\alpha + \hat{\xi}^\alpha \hat{S}_\alpha} y^D e^{\sum y^i R_i / y}. \quad (2.3)$$

We partially fix the  $\kappa$ -symmetry such that  $\xi^\alpha = 0$ , and we use the  $U(1) \times U(1)$  gauge symmetry generated by  $P + K$  and  $R_3$  to set  $x' = 0$ , thus the coset representative is

$$g = e^{xP + \theta^\alpha Q_\alpha} e^B. \quad (2.4)$$

Using the fact that the group generated by  $\{\hat{Q}, \hat{S}, D, R_i\}$  transforms  $P$  and  $Q_\alpha$  among themselves, the components of the Maurer-Cartan 1-form are

$$\begin{aligned} J_P &= [e^{-B}(dxP + d\theta^\alpha Q_\alpha)e^B]_P, & J_{Q_\alpha} &= [e^{-B}(dxP + d\theta^\alpha Q_\alpha)e^B]_{Q_\alpha}, \\ J_K &= 0, & J_{\hat{Q}_\alpha} &= [e^{-B}de^B]_{\hat{Q}_\alpha}, & J_{S_\alpha} &= 0, & J_{\hat{S}_\alpha} &= [e^{-B}de^B]_{\hat{S}_\alpha}, \\ J_D &= [e^{-B}de^B]_D, & J_{R_i} &= [e^{-B}de^B]_{R_i}. \end{aligned} \quad (2.5)$$

This sigma-model is T-dualized in the directions of the Abelian sub-algebra formed by the generators  $P$  and  $Q_\alpha$  according to the procedure of [16, 17], by introducing the gauge fields  $A, \bar{A}$  for the translation  $P$  and  $A^\alpha, \bar{A}^\alpha$  for the supercharges  $Q_\alpha$  and the corresponding Lagrange multipliers  $\tilde{x}$  and  $\tilde{\theta}_\alpha$ :

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{1}{2} [e^{-B}(AP + A^\alpha Q_\alpha)e^B]_P [e^{-B}(\bar{A}P + \bar{A}^\alpha Q_\alpha)e^B]_P - \right.$$

---

<sup>1</sup>For this to be a superstring, this superspace has to be supplemented by an additional internal CFT with the appropriate central charge. It is not clear that such a description exists using such a partial Green-Schwarz action (the hybrid model discussed in [18] might be a better option for this kind of superstring construction).

$$\begin{aligned}
& -\frac{i}{2}\eta_{\alpha\beta}[e^{-B}(AP + A^\gamma Q_\gamma)e^B]_{Q_\alpha}[e^{-B}(\bar{A}P + \bar{A}^\gamma Q_\gamma)e^B]_{Q_\beta} + \\
& +\frac{1}{2}J_D\bar{J}_D + \frac{1}{2}J_{R_1}\bar{J}_{R_1} + \frac{1}{2}J_{R_2}\bar{J}_{R_2} + \frac{i}{2}\eta_{\alpha\beta}(J_{\hat{Q}_\alpha}\bar{J}_{\hat{Q}_\beta} + J_{\hat{S}_\alpha}\bar{J}_{\hat{S}_\beta}) + \\
& +\tilde{x}(\bar{\partial}A - \partial\bar{A}) + \tilde{\theta}_\alpha(\bar{\partial}A^\alpha - \partial\bar{A}^\alpha) \Big]. \tag{2.6}
\end{aligned}$$

It is convenient to change variables to

$$A' = [e^{-B}(AP + A^\alpha Q_\alpha)e^B]_P, \quad A'^\alpha = [e^{-B}(AP + A^\beta Q_\beta)e^B]_{Q_\alpha} \tag{2.7}$$

and similarly for the right-moving gauge fields. Using the inverted relations

$$A = [e^B(A'P + A'^\alpha Q_\alpha)e^{-B}]_P, \quad A^\alpha = [e^B(A'P + A'^\beta Q_\beta)e^{-B}]_{Q_\alpha}, \tag{2.8}$$

the action in terms of the new variables reads

$$\begin{aligned}
S = \frac{R^2}{4\pi\alpha'} \int d^2z & \left[ \frac{1}{2}A'\bar{A}' - \frac{i}{2}\eta_{\alpha\beta}A'^\alpha\bar{A}'^\beta + \dots - \right. \\
& -\bar{\partial}\tilde{x}(A'[e^BPe^{-B}]_P + A'^\alpha[e^BQ_\alpha e^{-B}]_P) + \\
& +\partial\tilde{x}(\bar{A}'[e^BPe^{-B}]_P + \bar{A}'^\alpha[e^BQ_\alpha e^{-B}]_P) - \\
& -\bar{\partial}\tilde{\theta}_\alpha(A'[e^BPe^{-B}]_{Q_\alpha} + A'^\beta[e^BQ_\beta e^{-B}]_{Q_\alpha}) + \\
& \left. +\partial\tilde{\theta}_\alpha(\bar{A}'[e^BPe^{-B}]_{Q_\alpha} + \bar{A}'^\beta[e^BQ_\beta e^{-B}]_{Q_\alpha}) \right], \tag{2.9}
\end{aligned}$$

where ... denotes the spectator terms. Since the gauge fields appear quadratically in the action, one can integrate them out by substituting their equations of motion

$$\begin{aligned}
A' &= -2[e^B\partial\tilde{x}Pe^{-B}]_P - 2[e^B\bar{\partial}\tilde{\theta}_\alpha Pe^{-B}]_{Q_\alpha} = -2[e^{-B}(\partial\tilde{x}K + i\partial\tilde{\theta}_\alpha S^\alpha)e^B]_K, \\
\bar{A}' &= 2[e^B\bar{\partial}\tilde{x}Pe^{-B}]_P + 2[e^B\bar{\partial}\tilde{\theta}_\alpha Pe^{-B}]_{Q_\alpha} = 2[e^{-B}(\bar{\partial}\tilde{x}K + i\bar{\partial}\tilde{\theta}_\alpha S^\alpha)e^B]_K, \\
A'^\alpha &= 2i\eta^{\alpha\beta}([e^B\partial\tilde{x}Q_\beta e^{-B}]_P - [e^B\bar{\partial}\tilde{\theta}_\gamma Q_\beta e^{-B}]_{Q_\gamma}) = -2\eta^{\alpha\beta}[e^{-B}(\partial\tilde{x}K + i\partial\tilde{\theta}_\gamma S^\gamma)e^B]_{S^\beta}, \\
\bar{A}'^\alpha &= 2i\eta^{\alpha\beta}([e^B\bar{\partial}\tilde{x}Q_\beta e^{-B}]_P - [e^B\bar{\partial}\tilde{\theta}_\gamma Q_\beta e^{-B}]_{Q_\gamma}) = -2\eta^{\alpha\beta}[e^{-B}(\bar{\partial}\tilde{x}K + i\bar{\partial}\tilde{\theta}_\gamma S^\gamma)e^B]_{S^\beta}.
\end{aligned} \tag{2.10}$$

and obtain after rescaling  $\tilde{x} \rightarrow \frac{1}{2}\tilde{x}$  and  $\tilde{\theta}_\alpha \rightarrow \frac{1}{2}\tilde{\theta}_\alpha$

$$\begin{aligned}
S = \frac{R^2}{4\pi\alpha'} \int d^2z & \left[ \frac{1}{2}[e^{-B}(\partial\tilde{x}K + i\partial\tilde{\theta}_\alpha S^\alpha)e^B]_K[e^{-B}(\bar{\partial}\tilde{x}K + i\bar{\partial}\tilde{\theta}_\alpha S^\alpha)e^B]_K - \right. \\
& \left. -\frac{i}{2}\eta_{\alpha\beta}[e^{-B}(\partial\tilde{x}K + i\partial\tilde{\theta}_\gamma S^\gamma)e^B]_{S_\alpha}[e^{-B}(\bar{\partial}\tilde{x}K + i\bar{\partial}\tilde{\theta}_\gamma S^\gamma)e^B]_{S_\beta} + \dots \right], \tag{2.11}
\end{aligned}$$

where we used  $\epsilon_{\alpha\beta}$  to lower the spinor indices of  $\eta^{\alpha\beta}$ .

In order to show the self-duality of this background under the above T-duality, the original action has to be brought to the same form as (2.11). One can easily check by using the  $\epsilon_{\beta\gamma}\sigma_\alpha^{ij\gamma} = \epsilon_{\alpha\gamma}\sigma_\beta^{ij\gamma}$  that the  $\text{psu}(1,1|2)$  algebra admits the automorphism

$$P \leftrightarrow K, \quad D \rightarrow -D, \quad Q_\alpha \leftrightarrow S^\alpha, \quad \hat{Q}_\alpha \leftrightarrow \hat{S}_\alpha, \tag{2.12}$$

with the rest of the generators unchanged. Applying this automorphism combined with the change of variables

$$x \rightarrow \tilde{x}, \quad \theta^\alpha \rightarrow i\tilde{\theta}_\alpha, \quad \hat{\theta}^\alpha \leftrightarrow \hat{\xi}^\alpha, \quad y_i \rightarrow \frac{y_i}{y^2}, \quad (2.13)$$

to (2.5) one obtains (2.11).

In order to complete the proof of quantum mechanical equivalence we also have to show that the Jacobian functional determinant from the change of variables (2.7) is the identity. The transformation of variables was done using  $e^B$ . Since it is in a unitary subgroup of the PSU(1, 1|2) group, its super-determinant is equal to one and hence the Jacobian of the transformation is trivial and does not induce any shift of the dilaton. In order to see this explicitly, one can treat  $(A, A^\alpha)$  as a vector and write the adjoint action of the generators  $D$ ,  $R_{ij}$ ,  $\hat{Q}_\alpha$  and  $\hat{S}_\alpha$  as matrices acting on this vector

$$\begin{aligned} [\text{ad}_D] &= \begin{pmatrix} 1 & 0 \\ 0 & \frac{1}{2}\delta_\alpha^\beta \end{pmatrix}, & [\text{ad}_{R_{ij}}] &= \begin{pmatrix} 0 & 0 \\ 0 & -\frac{1}{2}\sigma_{ij\alpha}^\beta \end{pmatrix}, \\ [\text{ad}_{\hat{Q}_\alpha}] &= \begin{pmatrix} 0 & -\epsilon_{\alpha\beta} \\ 0 & 0 \end{pmatrix}, & [\text{ad}_{\hat{S}_\alpha}] &= \begin{pmatrix} 0 & 0 \\ -i\delta_\alpha^\beta & 0 \end{pmatrix}, \end{aligned} \quad (2.14)$$

which evidently have a vanishing supertrace. Hence, the super-determinant of similarity transformations with elements of the group generated by these generators is the identity.

## 2.2 The $AdS_3 \times S^3$ target-space

We construct the Green-Schwarz sigma-model on  $AdS_3 \times S^3$  using the supercoset manifold  $(\text{PSU}(1, 1|2) \times \text{PSU}(1, 1|2))/(\text{SU}(1, 1) \times \text{SU}(2))$  with 16 supersymmetry generators. Using the  $\mathbb{Z}_4$  structure (E.4) we have

$$J_2 = J_a^-(P_a - K_a) + J_D D + J_\mu R_\mu = \frac{1}{2}(J_{P_a} - J_{K_a})(P_a - K_a) + J_D D + J_\mu R_\mu \quad (2.15)$$

$$J_1 = J_{S_{\alpha\hat{\alpha}}^I + a^{IJ} Q_{\alpha\hat{\alpha}}^J} (S_{\alpha\hat{\alpha}}^I + a^{IK} Q_{\alpha\hat{\alpha}}^K) = \frac{1}{2}(J_{S_{\alpha\hat{\alpha}}^I} + a^{IJ} J_{Q_{\alpha\hat{\alpha}}^J}) (S_{\alpha\hat{\alpha}}^I + a^{IK} Q_{\alpha\hat{\alpha}}^K)$$

$$J_3 = J_{S_{\alpha\hat{\alpha}}^I - a^{IJ} Q_{\alpha\hat{\alpha}}^J} (S_{\alpha\hat{\alpha}}^I - a^{IK} Q_{\alpha\hat{\alpha}}^K) = \frac{1}{2}(J_{S_{\alpha\hat{\alpha}}^I} - a^{IJ} J_{Q_{\alpha\hat{\alpha}}^J}) (S_{\alpha\hat{\alpha}}^I - a^{IK} Q_{\alpha\hat{\alpha}}^K).$$

Thus,

$$\text{Str}(J_2 \bar{J}_2) = -\frac{1}{2}(J_{P_a} - J_{K_a})(\bar{J}_{P_b} - \bar{J}_{K_b})\eta^{ab} + J_D \bar{J}_D + J_{R_\mu} \bar{J}_{R_\mu}, \quad (2.16)$$

$$\text{Str}(J_1 \bar{J}_3) = \frac{1}{2}(\sigma^3)^{IK} \epsilon_{\alpha\beta} \epsilon_{\hat{\alpha}\hat{\beta}} (J_{S_{\alpha\hat{\alpha}}^I} + a^{IJ} J_{Q_{\alpha\hat{\alpha}}^J}) (\bar{J}_{S_{\hat{\beta}\beta}^K} - a^{KL} \bar{J}_{Q_{\hat{\beta}\beta}^L}), \quad (2.17)$$

and the action reads

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2}(J_{P_a} - J_{K_a})(\bar{J}_{P_b} - \bar{J}_{K_b})\eta_{ab} + J_D \bar{J}_D + J_{R_\mu} \bar{J}_{R_\mu} \right. \\ \left. + \frac{1}{2}(\sigma^3)^{IK} \epsilon_{\hat{\alpha}\hat{\beta}} \epsilon_{\alpha\beta} \left( J_{S_{\alpha\hat{\alpha}}^I} \bar{J}_{S_{\beta\hat{\beta}}^K} + J_{Q_{\alpha\hat{\alpha}}^I} \bar{J}_{Q_{\beta\hat{\beta}}^K} \right) \right]. \quad (2.18)$$

Next we parameterize the supergroup element

$$g = \exp(x^a P_a + \theta^{\alpha\hat{\alpha}1} Q_{\alpha\hat{\alpha}}^1) \exp(\theta^{\alpha\hat{\alpha}2} Q_{\alpha\hat{\alpha}}^2 + \xi^{\alpha\hat{\alpha}1} S_{\alpha\hat{\alpha}}^1) y^D \exp(y_\mu R_\mu / y) \\ \equiv \exp(x^a P_a + \theta^{\alpha\hat{\alpha}1} Q_{\alpha\hat{\alpha}}^1) e^B. \quad (2.19)$$

Define the currents

$$J = g^{-1} dg = j + \mathfrak{j}, \quad j = e^{-B} (dx^a P_a + d\theta^{\alpha\hat{\alpha}1} Q_{\alpha\hat{\alpha}}^1) e^B, \quad \mathfrak{j} = e^{-B} de^B. \quad (2.20)$$

Using these definitions and the algebra (E.1) we get the currents

$$J_{P_a} = j_{P_a}, \quad J_{Q_{\alpha\hat{\alpha}}^1} = j_{Q_{\alpha\hat{\alpha}}^1}, \quad J_{Q_{\alpha\hat{\alpha}}^2} = \mathfrak{j}_{Q_{\alpha\hat{\alpha}}^2}, \quad J_{S_{\alpha\hat{\alpha}}^1} = \mathfrak{j}_{S_{\alpha\hat{\alpha}}^1}, \quad J_{S_{\alpha\hat{\alpha}}^2} = 0 \quad (2.21) \\ J_{K_a} = 0, \quad J_D = \mathfrak{j}_D, \quad J_{R_\mu} = \mathfrak{j}_{R_\mu},$$

the action reads

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2} j_{P_a} \bar{j}_{P_b} \eta_{ab} + \mathfrak{j}_D \bar{\mathfrak{j}}_D + \mathfrak{j}_{R_\mu} \bar{\mathfrak{j}}_{R_\mu} \right. \\ \left. + \frac{1}{2} \epsilon_{\alpha\beta} \epsilon_{\hat{\alpha}\hat{\beta}} \left( \mathfrak{j}_{S_{\alpha\hat{\alpha}}^1} \bar{\mathfrak{j}}_{S_{\beta\hat{\beta}}^1} + j_{Q_{\alpha\hat{\alpha}}^1} \bar{j}_{Q_{\beta\hat{\beta}}^1} - \mathfrak{j}_{Q_{\alpha\hat{\alpha}}^2} \bar{\mathfrak{j}}_{Q_{\beta\hat{\beta}}^2} \right) \right]. \quad (2.22)$$

Introducing gauge fields and Lagrange multipliers we get the action

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2} A'_{P_a} \bar{A}'_{P_b} \eta_{ab} + \mathfrak{j}_D \bar{\mathfrak{j}}_D + \mathfrak{j}_{R_\mu} \bar{\mathfrak{j}}_{R_\mu} \right. \\ \left. + \frac{1}{2} \epsilon_{\alpha\beta} \epsilon_{\hat{\alpha}\hat{\beta}} \left( \mathfrak{j}_{S_{\alpha\hat{\alpha}}^1} \bar{\mathfrak{j}}_{S_{\beta\hat{\beta}}^1} + A'_{Q_{\alpha\hat{\alpha}}^1} \bar{A}'_{Q_{\beta\hat{\beta}}^1} - \mathfrak{j}_{Q_{\alpha\hat{\alpha}}^2} \bar{\mathfrak{j}}_{Q_{\beta\hat{\beta}}^2} \right) \right. \\ \left. + \tilde{x}_a (\bar{\partial} A^a - \partial \bar{A}^a) + \tilde{\theta}_{\alpha\hat{\alpha}}^1 (\bar{\partial} A_{Q_{\alpha\hat{\alpha}}^1} - \partial \bar{A}_{Q_{\alpha\hat{\alpha}}^1}) \right], \quad (2.23)$$

where

$$A' = e^{-B} (A^a P_a + A^{1\alpha\hat{\alpha}} Q_{\alpha\hat{\alpha}}^1) e^B, \quad A^{1\alpha\hat{\alpha}} \equiv A_{Q_{\alpha\hat{\alpha}}^1}, \quad A_{P_a} \equiv A^a, \quad (2.24) \\ A = e^B (A'^a P_a + A'^{1\alpha\hat{\alpha}} Q_{\alpha\hat{\alpha}}^1) e^{-B}.$$

Solving for  $A'$  we get

$$A'_a = 2(\partial \tilde{x}_b [e^B P_a e^{-B}]_{P_b} + \partial \tilde{\theta}_{\alpha\hat{\alpha}}^1 [e^B P_a e^{-B}]_{Q_{\alpha\hat{\alpha}}^1}) \quad (2.25)$$

$$\begin{aligned}
&= 2\eta_{ac}[e^{-B}(\partial\tilde{x}^b K_b + \partial\tilde{\theta}^{1\alpha\hat{\alpha}} S_{\alpha\hat{\alpha}}^2)e^B]_{K_c} \\
\bar{A}'_a &= -2(\bar{\partial}\tilde{x}_b[e^B P_a e^{-B}]_{P_b} + \bar{\partial}\tilde{\theta}^1_{\alpha\hat{\alpha}}[e^B P_a e^{-B}]_{Q^1_{\alpha\hat{\alpha}}}) \quad (2.26) \\
&= -2\eta_{ac}[e^{-B}(\bar{\partial}\tilde{x}^b K_b + \bar{\partial}\tilde{\theta}^{1\alpha\hat{\alpha}} S_{\alpha\hat{\alpha}}^2)e^B]_{K_c}
\end{aligned}$$

$$\begin{aligned}
\frac{1}{2}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}}A'_{Q^1_{\beta\hat{\beta}}} &= -(\partial\tilde{x}_b[e^B Q^1_{\alpha\hat{\alpha}} e^{-B}]_{P_b} - \partial\tilde{\theta}^1_{\beta\hat{\beta}}[e^B Q^1_{\alpha\hat{\alpha}} e^{-B}]_{Q^1_{\beta\hat{\beta}}}) \quad (2.27) \\
&= -\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}}[e^{-B}(\partial\tilde{x}^b K_b + \partial\tilde{\theta}^{1\gamma\hat{\gamma}} S_{\gamma\hat{\gamma}}^2)e^B]_{S^2_{\beta\hat{\beta}}}
\end{aligned}$$

$$\begin{aligned}
\frac{1}{2}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}}\bar{A}'_{Q^1_{\beta\hat{\beta}}} &= -(\bar{\partial}\tilde{x}_b[e^B Q^1_{\alpha\hat{\alpha}} e^{-B}]_{P_b} - \bar{\partial}\tilde{\theta}^1_{\beta\hat{\beta}}[e^B Q^1_{\alpha\hat{\alpha}} e^{-B}]_{Q^1_{\beta\hat{\beta}}}) \quad (2.28) \\
&= -\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}}[e^{-B}(\bar{\partial}\tilde{x}^b K_b + \bar{\partial}\tilde{\theta}^{1\gamma\hat{\gamma}} S_{\gamma\hat{\gamma}}^2)e^B]_{S^2_{\beta\hat{\beta}}} .
\end{aligned}$$

Thus,

$$\begin{aligned}
S &= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{1}{2}A'_{P_a}\bar{A}'_{P_b}\eta_{ab} + \mathfrak{j}_D\bar{\mathfrak{j}}_D + \mathfrak{j}_{R_\mu}\bar{\mathfrak{j}}_{R_\mu} \right. \quad (2.29) \\
&\quad \left. + \frac{1}{2}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}} \left( \mathfrak{j}_{S^1_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{S^1_{\beta\hat{\beta}}} - A'_{Q^1_{\alpha\hat{\alpha}}}\bar{A}'_{Q^1_{\beta\hat{\beta}}} - \mathfrak{j}_{Q^2_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{Q^2_{\beta\hat{\beta}}} \right) \right] \\
&= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2}4[e^{-B}(\partial\tilde{x}^b K_b + \partial\tilde{\theta}^{1\alpha\hat{\alpha}} S_{\alpha\hat{\alpha}}^2)e^B]_{K_a}[e^{-B}(\bar{\partial}\tilde{x}^b K_b + \bar{\partial}\tilde{\theta}^{1\alpha\hat{\alpha}} S_{\alpha\hat{\alpha}}^2)e^B]_{K_b}\eta^{ab} \right. \\
&\quad \left. + \mathfrak{j}_D\bar{\mathfrak{j}}_D + \mathfrak{j}_{R_\mu}\bar{\mathfrak{j}}_{R_\mu} + \frac{1}{2}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}} \left( \mathfrak{j}_{S^1_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{S^1_{\beta\hat{\beta}}} - \mathfrak{j}_{Q^2_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{Q^2_{\beta\hat{\beta}}} \right. \right. \\
&\quad \left. \left. - 4[e^{-B}(\partial\tilde{x}^b K_b + \partial\tilde{\theta}^{1\gamma\hat{\gamma}} S_{\gamma\hat{\gamma}}^2)e^B]_{S^2_{\alpha\hat{\alpha}}}[e^{-B}(\bar{\partial}\tilde{x}^b K_b + \bar{\partial}\tilde{\theta}^{1\gamma\hat{\gamma}} S_{\gamma\hat{\gamma}}^2)e^B]_{S^2_{\beta\hat{\beta}}} \right) \right] .
\end{aligned}$$

We can define  $J'$ 's

$$J' = e^{-B}(2\partial\tilde{x}^b K_b + 2\partial\tilde{\theta}^{1\gamma\hat{\gamma}} S_{\gamma\hat{\gamma}}^2)e^B , \quad (2.30)$$

and the action reads

$$\begin{aligned}
S &= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2}J'_{P_a}\bar{J}'_{P_b}\eta_{ab} + \mathfrak{j}_D\bar{\mathfrak{j}}_D + \mathfrak{j}_{R_\mu}\bar{\mathfrak{j}}_{R_\mu} \right. \quad (2.31) \\
&\quad \left. + \frac{1}{2}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}} \left( \mathfrak{j}_{S^1_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{S^1_{\beta\hat{\beta}}} - J'_{S^2_{\alpha\hat{\alpha}}}\bar{J}'_{S^2_{\beta\hat{\beta}}} - \mathfrak{j}_{Q^2_{\alpha\hat{\alpha}}}\bar{\mathfrak{j}}_{Q^2_{\beta\hat{\beta}}} \right) \right] .
\end{aligned}$$

Applying the following automorphism of the algebra

$$D \rightarrow -D, \quad K_a \leftrightarrow -\sigma_{ab}^3 P_b, \quad J_{ab} \rightarrow -J_{ab}, \quad (2.32)$$

$$R_1 \rightarrow R_1, \quad R_2 \rightarrow -R_2, \quad R_3 \rightarrow -R_3,$$

$$N_{12} \rightarrow -N_{12}, \quad N_{23} \rightarrow N_{23}, \quad N_{31} \rightarrow -N_{31},$$

$$Q^I_{\alpha\hat{\alpha}} \leftrightarrow \sigma^1_{IJ}\sigma^1_{\hat{\alpha}\hat{\beta}} S^J_{\alpha\hat{\beta}},$$

followed by the change of variables

$$x_a \rightarrow -2\sigma_{ab}^3 \tilde{x}_b, \quad \theta^1_{\alpha\hat{\alpha}} \rightarrow 2\sigma^1_{\hat{\alpha}\hat{\beta}} \tilde{\theta}^2_{\alpha\hat{\beta}}, \quad (2.33)$$

$$\theta^2_{\alpha\hat{\alpha}} \leftrightarrow \sigma^1_{\hat{\alpha}\hat{\beta}} \xi^1_{\alpha\hat{\beta}}, \quad y_1 \rightarrow y_1/y^2, \quad y_2 \rightarrow -y_2/y^2, \quad y_3 \rightarrow -y_3/y^2,$$

the action (2.31) is mapped to the original one (2.22).



### 3. Models based on the ortho-symplectic supergroup

In this section we consider models based on supercosets of the ortho-symplectic supergroup, for which the Buscher procedure [16, 17] of gauging an isometry of the target-space in order to obtain the T-dual sigma-model fails. These include the non-critical superstring on  $AdS_2$  with four supersymmetries and  $AdS_4$  with eight supersymmetries, the supercoset construction of  $AdS_4 \times \mathbb{CP}^3$  which is conjectured to be dual to superconformal Chern-Simons theory in three dimensions [19], and a model of  $AdS_2$  with eight supersymmetries.

#### 3.1 The Green-Schwarz sigma-model on $AdS_2$ with four supersymmetries

The non-critical  $AdS_2$  background with RR-flux can be realized as the supercoset  $OSp(2|2)/(SO(1,1) \times SO(2))$ . The Green-Schwarz action is of the form (2.1) (see Appendix A for the details of the algebra and the conventions that are used).

Fixing the  $SO(1,1) \times SO(2)$  gauge symmetry and  $\kappa$ -symmetry one can choose the coset representative

$$g = e^{xP+\theta Q} e^{\bar{\theta}\bar{Q}+\bar{\xi}\bar{S}} y^D . \quad (3.1)$$

The generators  $P$  and  $Q$  form an Abelian subalgebra, which we will attempt to T-dualize. In this parameterization the Maurer-Cartan current  $J$  takes the form

$$J = \frac{1}{y} (\partial x - 2\partial\theta\bar{\theta}) P + \frac{1}{y^{1/2}} \partial\theta Q + \frac{1}{y^{1/2}} [(2\partial\theta\bar{\theta} - \partial x) \bar{\xi} + \partial\bar{\theta}] \bar{Q} + y^{1/2} \partial\bar{\xi}\bar{S} + \left( \frac{\partial y}{y} - 2\partial\theta\bar{\xi} \right) D + i\partial\theta\bar{\xi} R . \quad (3.2)$$

The sigma-model can be written explicitly as

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -\frac{1}{2} (J_P + J_K) (\bar{J}_P + \bar{J}_K) + \frac{1}{2} J_D \bar{J}_D + J_Q \bar{J}_Q + J_{\bar{Q}} \bar{J}_{\bar{Q}} - J_S \bar{J}_{\bar{S}} - J_{\bar{S}} \bar{J}_S \right] . \quad (3.3)$$

The action in terms of the coordinates is then given by

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{-\partial x \bar{\partial} x + \partial y \bar{\partial} y}{2y^2} - \frac{1}{y^2} \bar{\theta} (\partial x \bar{\partial} \theta + \partial \theta \bar{\partial} x) + \frac{1}{y} \bar{\xi} (\partial y \bar{\partial} \theta + \partial \theta \bar{\partial} y) + \frac{4}{y} \bar{\theta} \bar{\xi} \partial \theta \bar{\partial} \theta + \frac{1}{y} \bar{\xi} (\partial \theta \bar{\partial} x - \partial x \bar{\partial} \theta) + \frac{1}{y} (\partial \theta \bar{\partial} \bar{\theta} + \partial \bar{\theta} \bar{\partial} \theta) \right] . \quad (3.4)$$

Note that the action is indeed quadratic in  $\theta$  so naively one should be able to T-dualize it along that coordinate.

An equivalent action can be written using two gauge fields [16, 17]

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{-A_x \bar{A}_x + \partial y \bar{\partial} y}{2y^2} - \frac{1}{y^2} \bar{\theta} (A_x \bar{A}_\theta + A_\theta \bar{A}_x) + \frac{1}{y} \bar{\xi} (\partial y \bar{A}_\theta + A_\theta \bar{\partial} y) + \right.$$

$$\begin{aligned}
& + \frac{4}{y} \bar{\theta} \bar{\xi} A_\theta \bar{A}_\theta + \frac{1}{y} \bar{\xi} (A_\theta \bar{A}_x - A_x \bar{A}_\theta) + \frac{1}{y} (A_\theta \bar{\partial} \bar{\theta} + \partial \bar{\theta} \bar{A}_\theta) + \tilde{x} (\bar{\partial} A_x - \partial \bar{A}_x) + \\
& + \tilde{\theta} (\bar{\partial} A_\theta - \partial \bar{A}_\theta) \Big] . \tag{3.5}
\end{aligned}$$

The classical equations of motion for the gauge fields are

$$\frac{1}{2y^2} A_x + \frac{1}{y^2} \bar{\theta} A_\theta - \frac{1}{y} \bar{\xi} A_\theta - \partial \tilde{x} = 0 , \tag{3.6}$$

$$\frac{1}{2y^2} \bar{A}_x + \frac{1}{y^2} \bar{\theta} \bar{A}_\theta + \frac{1}{y} \bar{\xi} \bar{A}_\theta + \bar{\partial} \tilde{x} = 0 , \tag{3.7}$$

$$\frac{1}{y^2} \bar{\theta} A_x - \frac{1}{y} \bar{\xi} \partial y - \frac{4}{y} \bar{\theta} \bar{\xi} A_\theta + \frac{1}{y} \bar{\xi} A_x - \frac{1}{y} \partial \bar{\theta} - \partial \tilde{\theta} = 0 , \tag{3.8}$$

$$\frac{1}{y^2} \bar{\theta} \bar{A}_x - \frac{1}{y} \bar{\xi} \bar{\partial} y + \frac{4}{y} \bar{\theta} \bar{\xi} \bar{A}_\theta - \frac{1}{y} \bar{\xi} \bar{A}_x + \frac{1}{y} \bar{\partial} \bar{\theta} + \bar{\partial} \tilde{\theta} = 0 . \tag{3.9}$$

Since Grassmann variables such as  $\bar{\theta}$  and  $\bar{\xi}$  have non-trivial kernels, one cannot solve for  $A_\theta$  and for  $\bar{A}_\theta$ . Solving for  $A_x$  and  $\bar{A}_x$  one gets

$$A_x = -2\bar{\theta} A_\theta + 2y\bar{\xi} A_\theta + 2y^2 \partial \tilde{x} , \tag{3.10}$$

$$\bar{A}_x = -2\bar{\theta} \bar{A}_\theta - 2y\bar{\xi} \bar{A}_\theta - 2y^2 \bar{\partial} \tilde{x} . \tag{3.11}$$

Substituting (3.10) and (3.11) in the action (this can be done since they appear only quadratically in the action, but the classical procedure should be supplemented by a functional determinant coming from the Gaussian path integration of  $A_x$  and  $\bar{A}_x$ ) yields

$$\begin{aligned}
S = \frac{R^2}{4\pi\alpha'} \int d^2z \Big[ & -2y^2 \partial \tilde{x} \bar{\partial} \tilde{x} - 2y\bar{\xi} \partial \tilde{x} \bar{A}_\theta - 2y\bar{\xi} A_\theta \bar{\partial} \tilde{x} - 2\bar{\theta} \partial \tilde{x} \bar{A}_\theta + 2\bar{\theta} A_\theta \bar{\partial} \tilde{x} + \\
& + \frac{\partial y \bar{\partial} y}{2y^2} + \frac{1}{y} \bar{\xi} (\partial y \bar{A}_\theta + A_\theta \bar{\partial} y) + \frac{1}{y} (A_\theta \bar{\partial} \bar{\theta} + \partial \bar{\theta} \bar{A}_\theta) - \bar{\partial} \tilde{\theta} A_\theta + \partial \tilde{\theta} \bar{A}_\theta \Big] . \tag{3.12}
\end{aligned}$$

Thus, after integrating out  $A_x$  and  $\bar{A}_x$  the remaining fermionic gauge fields  $A_\theta$  and  $\bar{A}_\theta$  serve as Lagrange multipliers forcing in the path integration over  $\tilde{\theta}$  that

$$\begin{aligned}
2\bar{\theta} \partial \tilde{x} - \frac{1}{y} \bar{\xi} \partial y + 2y\bar{\xi} \partial \tilde{x} - \frac{1}{y} \partial \bar{\theta} - \partial \tilde{\theta} &= 0 , \\
2\bar{\theta} \bar{\partial} \tilde{x} + \frac{1}{y} \bar{\xi} \bar{\partial} y - 2y\bar{\xi} \bar{\partial} \tilde{x} - \frac{1}{y} \bar{\partial} \bar{\theta} - \bar{\partial} \tilde{\theta} &= 0 , \tag{3.13}
\end{aligned}$$

which express the non-zero modes of  $\tilde{\theta}$  in terms of the other fields, effectively reducing the path integration over  $\tilde{\theta}$  only to its zero-modes.

This appears rather strange as the original action did include a term quadratic in  $\theta$ . A possible explanation is that the quadratic term is  $\kappa$ -symmetry-exact, and hence does not influence the equations of motion.

In order to support the above claim we do the same computation with a different gauge choice for the  $\kappa$ -symmetry. The coset representative in this gauge can be parameterized as

$$g = e^{xP+\theta Q} e^{\bar{\theta}\bar{Q}+\xi S} y^D , \quad (3.14)$$

and the Maurer-Cartan current is

$$J = \frac{1}{y}(\partial x - 2\partial\theta\bar{\theta} - \partial x\bar{\theta}\xi)P + \frac{1}{y^{1/2}}(\partial\theta - \partial x\xi + \partial\theta\bar{\theta}\xi)Q + \frac{1}{y^{1/2}}\partial\bar{\theta}\bar{Q} + y^{1/2}\partial\xi S + \left(\frac{\partial y}{y} - \partial\bar{\theta}\xi - \partial\xi\bar{\theta}\right)D - \frac{i}{2}(\partial\bar{\theta}\xi + \partial\xi\bar{\theta})R . \quad (3.15)$$

In this gauge the action is

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ \frac{(-1 + \bar{\theta}\xi)\partial x\bar{\partial}x + \partial y\bar{\partial}y}{2y^2} - \frac{1}{y^2}\bar{\theta}(\partial x\bar{\partial}\theta + \partial\theta\bar{\partial}x) - \frac{1}{y}\xi(\partial x\bar{\partial}\bar{\theta} - \partial\bar{\theta}\bar{\partial}x) + \frac{1}{2y}\xi(\partial y\bar{\partial}\bar{\theta} + \partial\bar{\theta}\bar{\partial}y) + \frac{1}{2y}\bar{\theta}(\partial y\bar{\partial}\xi + \partial\xi\bar{\partial}y) + \frac{1}{2}\bar{\theta}\xi(\partial\bar{\theta}\bar{\partial}\xi - \partial\xi\bar{\partial}\bar{\theta}) + \frac{1}{y}(1 + \bar{\theta}\xi)(\partial\theta\bar{\partial}\bar{\theta} + \partial\bar{\theta}\bar{\partial}\theta) \right] . \quad (3.16)$$

This action has no quadratic term in the Grassmann coordinate  $\theta$  so the usual procedure introduced by Buscher [16, 17] cannot be applied. This lends credence to the explanation that in the different gauge above the procedure failed because the quadratic term is  $\kappa$ -symmetry-exact.

### 3.2 The $AdS_4$ target-space

The non-critical superstring on  $AdS_4$  with eight supersymmetries can be constructed as the supercoset  $OSp(2|4)/(SO(3,1) \times SO(2))$ . Taking  $g \in OSp(2|4)$  in (2.1) one obtains the sigma-model

$$S = -\frac{R^2}{4\pi\alpha'} \int d^2z \left[ \eta_{mn}(J_{P_m} + J_{K_m})(\bar{J}_{P_n} + \bar{J}_{K_n}) + J_D\bar{J}_D + 4i\epsilon_{\alpha\beta} \left( J_{Q_\alpha}\bar{J}_{\hat{Q}_\beta} - J_{\hat{Q}_\alpha}\bar{J}_{Q_\beta} + J_{S_\alpha}\bar{J}_{\hat{S}_\beta} - J_{\hat{S}_\alpha}\bar{J}_{S_\beta} \right) \right] , \quad (3.17)$$

where the conventions for the algebra are given in Appendix B.

The general group element can be parameterized as

$$g = e^{x^m P_m + w^m K_m + \theta^\alpha Q_\alpha} e^B , \quad e^B = e^{\hat{\theta}^\alpha \hat{Q}_\alpha + \xi^\alpha S_\alpha + \hat{\xi}^\alpha \hat{S}_\alpha} y^D e^{\phi R + \omega^{mn} M_{mn}} . \quad (3.18)$$

We assume that we can partially gauge-fix the  $\kappa$ -symmetry such that  $\hat{\xi}^\alpha = 0$  and we will also fix the  $SO(3,1) \times SO(2)$  gauge symmetry by setting  $w^m = 0$ ,  $\phi = 0$  and  $\omega^{mn} = 0$  essentially picking the specific coset representative

$$g = e^{x^m P_m + \theta^\alpha Q_\alpha} e^B , \quad e^B = e^{\hat{\theta}^\alpha \hat{Q}_\alpha + \xi^\alpha S_\alpha} y^D . \quad (3.19)$$

One can check that the needed components of the Maurer-Cartan 1-form are

$$\begin{aligned}
J_{P_m} &= [e^{-B}(dx^n P_n + d\theta^\alpha Q_\alpha)e^B]_{P_m} , & J_{K_m} &= 0 , \\
J_{Q_\alpha} &= [e^{-B}(dx^m P_m + d\theta^\beta Q_\beta)e^B]_{Q_\alpha} , & J_{S_\alpha} &= [e^{-B}de^B]_{S_\alpha} , \\
J_{\hat{Q}_\alpha} &= [e^{-B}de^B]_{\hat{Q}_\alpha} , & J_{\hat{S}_\alpha} &= 0 , & J_D &= [e^{-B}de^B]_D .
\end{aligned} \tag{3.20}$$

The action then becomes

$$\begin{aligned}
S &= -\frac{R^2}{4\pi\alpha'} \int d^2z \left[ \eta_{mn} [e^{-B}(\partial x^l P_l + \partial\theta^\alpha Q_\alpha)e^B]_{P_m} [e^{-B}(\bar{\partial}x^p P_p + \bar{\partial}\theta^\beta Q_\beta)e^B]_{P_n} + \right. \\
&\quad + [e^{-B}\partial e^B]_D [e^{-B}\bar{\partial}e^B]_D + 4i\epsilon_{\alpha\beta} \left( [e^{-B}(\partial x^m P_m + \partial\theta^\gamma Q_\gamma)e^B]_{Q_\alpha} [e^{-B}\bar{\partial}e^B]_{\hat{Q}_\alpha} \right. \\
&\quad \left. \left. - [e^{-B}\partial e^B]_{\hat{Q}_\alpha} [e^{-B}(\bar{\partial}x^m P_m + \bar{\partial}\theta^\gamma Q_\gamma)e^B]_{Q_\beta} \right) \right] .
\end{aligned} \tag{3.21}$$

Replacing the partial derivatives of  $x^m$  and of  $\theta^\alpha$  by the gauge fields  $A^m$ ,  $\bar{A}^m$ ,  $A^\alpha$  and  $\bar{A}^\alpha$  and then performing the field redefinition

$$\begin{aligned}
A'^m &= [e^{-B}(A^n P_n + A^\alpha Q_\alpha)e^B]_{P_m} , \\
A'^\alpha &= [e^{-B}(A^m P_m + A^\beta Q_\beta)e^B]_{Q_\alpha}
\end{aligned} \tag{3.22}$$

with similar expressions for the right-moving sector and introducing the Lagrange multiplier terms forcing the gauge fields to be flat the new equivalent action takes the form

$$\begin{aligned}
S &= -\frac{R^2}{4\pi\alpha'} \int d^2z \left[ \eta_{mn} A'^m \bar{A}'^n + J_D \bar{J}_D + 4i\epsilon_{\alpha\beta} \left( A'^\alpha [e^{-B}\bar{\partial}e^B]_{\hat{Q}_\beta} - \right. \right. \\
&\quad \left. \left. - [e^{-B}\partial e^B]_{\hat{Q}_\alpha} \bar{A}'^\beta \right) + \tilde{x}_m (\bar{\partial}A^m - \partial\bar{A}^m) + \tilde{\theta}_\alpha (\bar{\partial}A^\alpha - \partial\bar{A}^\alpha) \right] .
\end{aligned} \tag{3.23}$$

As can be immediately seen, just as for the case of the  $AdS_2$  space discussed in section 3.1 for the (physically trivial)  $AdS_2$  the fermionic gauge fields  $A'^\alpha$  and  $\bar{A}'^\alpha$  appear only linearly in the action and hence integration over them will yield constraints rather than equations of motion.

### 3.3 The $AdS_4 \times \mathbb{C}P^3$ target-space

Using the same procedure as for the  $AdS_n \times S^m$  models we construct the Green-Schwarz sigma model action (2.1) using the bilinear forms of (C.21),

$$\begin{aligned}
S &= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2\eta_{ab}(J_{P_a} + J_{K_a})(\bar{J}_{P_b} + \bar{J}_{K_b}) \right. \\
&\quad - J_D \bar{J}_D - 2(J_{R_{kl}} \bar{J}_{R_{\dot{p}\dot{q}}} + \bar{J}_{R_{kl}} J_{R_{\dot{p}\dot{q}}})(\delta^{k\dot{p}}\delta^{l\dot{q}} - \delta^{k\dot{q}}\delta^{l\dot{p}}) \\
&\quad \left. - 2i\delta^{kl} C_{\alpha\beta} \left( J_{Q_\alpha}^l \bar{J}_{Q_\beta}^k - \bar{J}_{Q_\alpha}^l J_{Q_\beta}^k - J_{S_\alpha}^l \bar{J}_{S_\beta}^k + \bar{J}_{S_\alpha}^l J_{S_\beta}^k \right) \right] .
\end{aligned} \tag{3.24}$$

(This coset model is a Green-Schwarz superstring with kappa-symmetry partially fixed so that the broken supersymmetries are gauged away, leaving only the fermionic coordinates corresponding to unbroken supersymmetries. This partial gauge-fixing is not compatible with all the string solutions so a general analysis would seem to require the use of the complete unfixed sigma-model derived in [20]. However, as the broken supersymmetries are not isometries of the background, they cannot participate in the T-duality transformation so we expect the  $\text{OSp}(6|4)$  coset model to suffice for this purpose.)

We proceed by taking the gauge-fixed coset representative

$$g = e^{x^a P_a + \theta_l^\alpha Q_\alpha^l} e^B, \quad e^B \equiv e^{\theta_l^\alpha Q_\alpha^l + \xi_l^\alpha S_\alpha^l} y^D e^{(\Sigma y^{kl} R_{kl} + \Sigma y^{ki} R_{ki})/y}, \quad (3.25)$$

where we also fixed six out of the eight  $\kappa$ -symmetry degrees of freedom by setting  $\xi_l^\alpha = 0$ .  $P_a$  and  $\theta_l^\alpha Q_\alpha^l$  form an Abelian subalgebra.

Next, we would like to write the currents in terms of this parameterization, but in this case the commutations relation  $[Q, R] \sim \bar{Q}$  and  $\{Q, \}$   $\sim R$  prevents us from getting two kinds of currents as in (2.20), so

$$J_T = [e^{-B}(dx^a P_a + d\theta_l^\alpha Q_\alpha^l)e^B]_T + [e^{-B}de^B]_T \equiv j_T + \mathfrak{j}_T. \quad (3.26)$$

We have

$$j = j_{P_a} P_a + j_{Q_l^\alpha} Q_l^\alpha + j_{Q_i^\alpha} Q_i^\alpha + j_{R_{kl}} R_{kl} + j_{R_{ki}} R_{ki} + j_{\lambda_{\kappa l}} \lambda_{\kappa l}, \quad (3.27)$$

and

$$\mathfrak{j} = \mathfrak{j}_{Q_l^\alpha} Q_l^\alpha + \mathfrak{j}_{Q_i^\alpha} Q_i^\alpha + \mathfrak{j}_{S_l^\alpha} S_l^\alpha + \mathfrak{j}_{S_i^\alpha} S_i^\alpha + \mathfrak{j}_{R_{kl}} R_{kl} + \mathfrak{j}_{R_{ki}} R_{ki} + \mathfrak{j}_{\lambda_{\kappa l}} \lambda_{\kappa l} + \mathfrak{j}_D D + \mathfrak{j}_{M_{mn}} M_{mn}. \quad (3.28)$$

The action in terms of these currents reads

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2\eta^{ab} j_{P_a} \bar{j}_{P_b} - \mathfrak{j}_D \bar{j}_D \right. \\ \left. -2((j + \mathfrak{j})_{R_{kl}} (\bar{j} + \bar{\mathfrak{j}})_{R_{\dot{p}\dot{q}}} + (\bar{j} + \bar{\mathfrak{j}})_{R_{kl}} (j + \mathfrak{j})_{R_{\dot{p}\dot{q}}}) (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \right. \\ \left. -2i\delta^{kl} C_{\alpha\beta} \left( (j + \mathfrak{j})_{Q_\alpha^l} (\bar{j} + \bar{\mathfrak{j}})_{Q_\beta^k} - (\bar{j} + \bar{\mathfrak{j}})_{Q_\alpha^l} (j + \mathfrak{j})_{Q_\beta^k} - \mathfrak{j}_{S_\alpha^l} \mathfrak{j}_{S_\beta^k} + \mathfrak{j}_{S_\alpha^l} \mathfrak{j}_{S_\beta^k} \right) \right]. \quad (3.29)$$

We add gauge fields instead of the  $x_a$  and  $\theta_l^\alpha$  derivatives and a suitable Lagrange multiplier term. We define

$$A' = e^{-B}(A^a P_a + A_l^\alpha Q_\alpha^l) e^B, \quad (3.30)$$

(where  $A^a \equiv A_{P_a}$  and  $A_l^\alpha \equiv A_{Q_\alpha^l}$ ) and the action becomes

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2\eta_{ab} A_{P_a}' \bar{A}_{P_b}' - \mathfrak{j}_D \bar{j}_D \right. \\ \left. -2((A' + \mathfrak{j})_{R_{kl}} (\bar{A}' + \bar{\mathfrak{j}})_{R_{\dot{p}\dot{q}}} + (\bar{A}' + \bar{\mathfrak{j}})_{R_{kl}} (A' + \mathfrak{j})_{R_{\dot{p}\dot{q}}}) (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \right. \\ \left. -2i\delta^{kl} C_{\alpha\beta} \left( (A' + \mathfrak{j})_{Q_\alpha^l} (\bar{A}' + \bar{\mathfrak{j}})_{Q_\beta^k} - (\bar{A}' + \bar{\mathfrak{j}})_{Q_\alpha^l} (A' + \mathfrak{j})_{Q_\beta^k} - \mathfrak{j}_{S_\alpha^l} \mathfrak{j}_{S_\beta^k} + \mathfrak{j}_{S_\alpha^l} \mathfrak{j}_{S_\beta^k} \right) \right. \\ \left. + \tilde{x}^a (\partial \bar{A}_a - \bar{\partial} A_a) + \tilde{\theta}_l^\alpha (\partial \bar{A}_\alpha^l - \bar{\partial} A_\alpha^l) \right]. \quad (3.31)$$

### 3.3.1 Current expansion in fermions

To zeroth order in the fermions  $(\theta_l^\alpha, \theta_i^\alpha, \xi_l^\alpha)$  we get a bosonic sigma model with no fermions, which we can T-dualize,

$$S_0 = \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2y^2 \eta_{ab} A_{P_a} \bar{A}_{P_b} - j_D \bar{j}_D \right. \quad (3.32)$$

$$\left. -2 \left\{ j_{R_{kl}} \bar{j}_{R_{\dot{p}\dot{q}}} + \bar{j}_{R_{kl}} j_{R_{\dot{p}\dot{q}}} \right\} (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) + \tilde{x}^a (\partial \bar{A}_a - \bar{\partial} A_a) \right],$$

where  $j = e^{-B} de^B$ ,  $e^B = y^D e^{(\Sigma y^{kl} R_{kl} + \Sigma y^{\dot{k}\dot{l}} R_{\dot{k}\dot{l}})/y}$ .

Next, leaving only  $\theta_l^\alpha$  terms we find

$$e^{-B} (A^a P_a + A_\lambda^\alpha Q_\alpha^l) e^B = y A^a P_a + y^{1/2} A_k^\alpha \left( f_l^k \left( \frac{y^{mn}}{y}, \frac{y^{\dot{m}\dot{n}}}{y} \right) Q_\alpha^l + g_l^k \left( \frac{y^{mn}}{y}, \frac{y^{\dot{m}\dot{n}}}{y} \right) Q_\alpha^i \right), \quad (3.33)$$

thus the  $A'$ 's in terms of the  $A$ 's are

$$A'^a = y A^a, \quad A'_l{}^\alpha = y^{1/2} A_k^\alpha f_l^k \left( \frac{y^{mn}}{y}, \frac{y^{\dot{m}\dot{n}}}{y} \right), \quad A'_i{}^\alpha = y^{1/2} A_k^\alpha g_i^k \left( \frac{y^{mn}}{y}, \frac{y^{\dot{m}\dot{n}}}{y} \right). \quad (3.34)$$

Plugging these in the action we have

$$\begin{aligned} S_1 &= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2y^2 \eta_{ab} A_{P_a} \bar{A}_{P_b} - j_D \bar{j}_D - 2 \left\{ j_{R_{kl}} \bar{j}_{R_{\dot{p}\dot{q}}} + \bar{j}_{R_{kl}} j_{R_{\dot{p}\dot{q}}} \right\} (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \right. \quad (3.35) \\ &\quad \left. - 2iy \delta^{kl} C_{\alpha\beta} \left( A_n^\alpha f_l^n \bar{A}_m^\beta g_k^m - \bar{A}_n^\alpha f_l^n A_m^\beta g_k^m \right) + \tilde{x}^a (\partial \bar{A}_a - \bar{\partial} A_a) + \tilde{\theta}_l^\alpha (\partial \bar{A}_\alpha^l - \bar{\partial} A_\alpha^l) \right] \\ &= \frac{R^2}{4\pi\alpha'} \int d^2z \left[ -2y^2 \eta_{ab} A_{P_a} \bar{A}_{P_b} - j_D \bar{j}_D - 2 \left\{ j_{R_{kl}} \bar{j}_{R_{\dot{p}\dot{q}}} + \bar{j}_{R_{kl}} j_{R_{\dot{p}\dot{q}}} \right\} (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \right. \\ &\quad \left. - 2iy \delta^{kl} C_{\alpha\beta} A_n^\alpha \bar{A}_m^\beta \left( f_l^n g_k^m - f_l^m g_k^n \right) + \tilde{x}^a (\partial \bar{A}_a - \bar{\partial} A_a) + \tilde{\theta}_l^\alpha (\partial \bar{A}_\alpha^l - \bar{\partial} A_\alpha^l) \right]. \end{aligned}$$

The bosonic T-duality works as before, but now we also have a quadratic term for the fermions. The equation of motion for the fermions is

$$A_m^\alpha \delta^{lk} (f_l^n g_k^m - f_l^m g_k^n) = -\frac{i}{2y} C^{\alpha\beta} \partial \theta_\beta^n. \quad (3.36)$$

The matrix  $M^{mn} \equiv \delta^{lk} (f_l^n g_k^m - f_l^m g_k^n) = -4iy^{\dot{m}\dot{n}}/y + O(y^{\dot{m}\dot{n}} y^{kl}/y^2)$ , is an anti-symmetric three-dimensional matrix and hence has a vanishing determinant, so we cannot solve for  $A_l^\alpha$ .

Next, to first order in  $\xi_l^\alpha$  we have nontrivial  $j_{S_l^\alpha}$  and  $j_{S_i^\alpha}$ , but these terms do not mix with the  $A$ 's so we will ignore them. The  $A$ 's change as follows,

$$A_k^\beta \rightarrow A_k^\beta - iA^a (\gamma_a)_\alpha{}^\beta \xi_k^\alpha \quad (3.37)$$

$$A^a \rightarrow A^a$$

$$A_{kl} \rightarrow A_k^\beta C_{\beta\alpha} \xi_l^\alpha, \quad A^{rs} \rightarrow h_{kl}^{rs} (y^{ij}/y) A_k^\beta C_{\beta\alpha} \xi_l^\alpha.$$

Thus the change in (3.36) goes like

$$(A_m^\alpha - iA^a(\gamma_a)_\beta^\alpha \xi_m^\beta) \delta^{lk} (f_l^n g_k^m - f_l^m g_k^n) = -\frac{i}{2y} C^{\alpha\beta} \partial \theta_\beta^n + k^{\alpha n}(\xi, \mathfrak{j}_{Rkl}, y, \dots), \quad (3.38)$$

and again, we have the same singular matrix  $M^{mn}$  multiplying the gauge fields  $A$ . The second equation, which we can think of as the first order correction to the bosonic T-duality equation, is

$$2y^2 A^a + 2\delta^{kl} \xi_l^\gamma (\gamma^a C)_{\gamma\beta} (\sqrt{y} A_m^\beta g_k^m + \mathfrak{j}_k^\beta) = -\partial \tilde{x}^a. \quad (3.39)$$

We can plug the solution for  $A^m$  in (3.38), but we will just get an expression of the form

$$A_l^\beta (\delta_\beta^\alpha \delta_m^l + \phi_\beta^\alpha F_m^l) M^{mn} = -\frac{i}{2y} C^{\alpha\beta} \partial \theta_\beta^n + k^{\alpha n}(\xi, \mathfrak{j}_{Rkl}, y, \dots) + \dots \quad (3.40)$$

where  $\phi$ ,  $F$ ,  $k$  and ... are functions of the coordinates but not of  $A$ , so again we cannot solve for  $A_l^\beta$ .

Going next to first order in  $\theta_i^\alpha$  (but leaving out terms of order  $\theta_i^\alpha \xi_k^\beta$ ) we have nontrivial  $\mathfrak{j}_{Q_\alpha}^l$  and  $\mathfrak{j}_{Q_\alpha}^i$ , and also

$$A^a \rightarrow A^a + A_k^\beta \delta^{ik} (\gamma^a C)_{\beta\alpha} \theta_i^\alpha \quad (3.41)$$

with respect to (3.35), so the action we get is

$$\begin{aligned} S_2 = \frac{R^2}{4\pi\alpha'} \int d^2z \Big[ & -2y^2 \eta_{ab} (A^a + A_k^\beta \delta^{ik} (\gamma^a C)_{\beta\alpha} \theta_i^\alpha) (\bar{A}^b + \bar{A}_m^\gamma \delta^{nm} (\gamma^b C)_{\gamma\delta} \theta_n^\delta) \\ & - \mathfrak{j}_D \bar{\mathfrak{j}}_D - 2 \left\{ (h_{rs}^{kl} A_r^\beta C_{\beta\alpha} \xi_s^\alpha + \mathfrak{j}_{kl}) \bar{\mathfrak{j}}_{\dot{p}\dot{q}} + (h_{mn}^{kl} \bar{A}_m^\beta C_{\beta\alpha} \xi_n^\alpha + \bar{\mathfrak{j}}_{kl}) \mathfrak{j}_{\dot{p}\dot{q}} \right\} (\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \\ & - 2i \delta^{kl} C_{\alpha\beta} \left( y (f_l^m (A_m^\alpha - iA^a(\gamma_a)_\delta^\alpha \xi_m^\delta) + \mathfrak{j}_l^\alpha) (g_k^n (\bar{A}_n^\beta - i\bar{A}^b(\gamma_b)_\gamma^\beta \xi_n^\gamma) + \bar{\mathfrak{j}}_k^\beta) \right. \\ & \left. - y (f_l^m (\bar{A}_m^\alpha - i\bar{A}^a(\gamma_a)_\delta^\alpha \xi_m^\delta) + \bar{\mathfrak{j}}_l^\alpha) (g_k^n (A_n^\beta - iA^b(\gamma_b)_\gamma^\beta \xi_n^\gamma) + \mathfrak{j}_k^\beta) - \mathfrak{j}_{S_\alpha^l} \mathfrak{j}_{S_\beta^k} + \mathfrak{j}_{S_\alpha^k} \mathfrak{j}_{S_\beta^l} \right) \\ & \left. + \tilde{x}^a (\partial \bar{A}_a - \bar{\partial} A_a) + \tilde{\theta}_l^\alpha (\partial \bar{A}_\alpha^l - \bar{\partial} A_\alpha^l) \right]. \end{aligned} \quad (3.42)$$

The equations of motion (to first order in  $\xi$  or  $\theta$ ) are,

$$-2y A_a + A_k^\beta \delta^{ik} (\gamma_a C)_{\beta\alpha} \theta_i^\alpha \quad (3.43)$$

$$\begin{aligned} & -2\delta^{kl} y C_{\alpha\beta} \left[ g_k^n (f_l^m A_m^\alpha + \mathfrak{j}_l^\alpha) (\gamma_a)_\gamma^\beta \xi_n^\gamma - f_l^m (\gamma_a)_\delta^\alpha \xi_m^\delta (g_k^n A_n^\beta + \mathfrak{j}_k^\beta) \right] = \partial \tilde{x}_a \\ & 2i \delta^{kl} C_{\alpha\beta} y \left\{ (f_l^m (A_m^\alpha - iA^a(\gamma_a)_\delta^\alpha \xi_m^\delta) + \mathfrak{j}_l^\alpha) g_k^n - f_l^n (g_k^m (A_m^\alpha - iA^a(\gamma_a)_\gamma^\alpha \xi_m^\gamma) + \mathfrak{j}_k^\alpha) \right\} \end{aligned} \quad (3.44)$$

$$-2yA^a\delta^{nm}(\gamma_a C)_{\beta\delta}\theta_m^\delta + F_\beta^n = 0 .$$

Plugging (3.43) in (3.44) we find

$$2i\delta^{kl}C_{\alpha\beta}y\left\{(f_l^m(A_m^\alpha + i\frac{\partial\tilde{x}^a}{2y}(\gamma_a)_\delta^\alpha\xi_m^\delta) + \mathfrak{j}_l^\alpha)g_k^n - f_l^n(g_k^m(A_m^\alpha + i\frac{\partial\tilde{x}^a}{2y}(\gamma_a)_\gamma^\alpha\xi_m^\gamma) + \mathfrak{j}_k^\alpha)\right\} \\ + \partial\tilde{x}^a\delta^{nm}(\gamma_a C)_{\beta\delta}\theta_m^\delta + F_\beta^n = 0 . \quad (3.45)$$

Thus again the  $A_l^\alpha$  will multiply the same singular matrix.

More generally, the bosonic singular matrix  $M$  will get corrections with even powers of the fermionic variables,

$$M_{mn} = M_{mn}^s + M_{mn}^{2n} + M_{mn}^4 + \dots , \quad (3.46)$$

where  $M_{mn}^{2n}$  is a matrix function with a power of  $2n$  of the fermionic variables  $\theta_k^\alpha$  and/or  $\xi_k^\alpha$ , and  $M_{mn}^s$  is the purely bosonic singular matrix. In order to solve the equations of motion we will need the inverse of  $M_{mn}$ , but such an inverse does not exist, because all the matrices involving fermions should cancel each other order by order and we should get,

$$\delta_{mn} = M_{mk}M_{kn}^{-1} = M_{mk}^s(M^{-1})_{kn}^s + \dots = M_{mk}^s(M^{-1})_{kn}^s , \quad (3.47)$$

but  $M_{mn}^s$  is singular.

### 3.4 The $AdS_2$ target-space with eight supersymmetries

Similarly to section 2.1 we construct a Green-Schwarz sigma-model action on  $AdS_2$ , but using a different supercoset, namely  $PSU(1,1|2)/(U(1) \times SU(2))$ , so this time we will have eight supersymmetries rather than four. Using the  $\mathbb{Z}_4$  structure of this super-algebra given in (D.7) we construct the Green-Schwarz action

$$S = \frac{R^2}{4\pi\alpha'} \int d^2z \left( -J_P\bar{J}_K - J_K\bar{J}_P + \frac{i}{2}\epsilon_{\alpha\beta}(J_{Q_\alpha}\bar{J}_{S_\beta} - J_{\hat{Q}_\alpha}\bar{J}_{\hat{S}_\beta} - J_{S_\alpha}\bar{J}_{Q_\beta} + J_{\hat{S}_\alpha}\bar{J}_{\hat{Q}_\beta}) \right) . \quad (3.48)$$

We parameterize the coset representatives such that

$$g = e^{xP + \theta^\alpha Q_\alpha} e^B, \quad e^B = e^{\hat{\theta}^\alpha \hat{Q}_\alpha + \xi^\alpha S_\alpha + \hat{\xi}^\alpha \hat{S}_\alpha} e^{yK} . \quad (3.49)$$

If we define the following currents

$$J = g^{-1}dg = j + \mathfrak{j}, \quad j = e^{-B}(dxP + d\theta^\alpha Q_\alpha)e^B, \quad \mathfrak{j} = e^{-B}de^B, \quad (3.50)$$

Expanding  $j$  to zeroth order in the fermions in  $B$  we get,

$$j_{(0)} = dxP + s\theta^\alpha Q_\alpha + 2ydxD + id\theta^\alpha \hat{S}_\alpha - y^2dxK . \quad (3.51)$$



Thus by using the Buscher procedure of introducing gauge fields as in the examples above, to this order the action will not have quadratic terms in the fermions, so all the quadratic  $d\theta$  (or more precisely  $A^\alpha$ ) terms in the action, coming from higher orders in the fermions in  $B$ , will multiply some fermions, and we will not be able to solve the equations of motion for the fermions. The reason again is that we will have to multiply the equations of motion by a singular matrix.

## 4. A general analysis

In this section we present a general algebraic argument to when a supercoset is expected to have a fermionic T-duality symmetry, and when it will fail to have one.

Let us assume a superconformal algebra  $\mathcal{G}$  with a  $\mathbb{Z}_4$  automorphism structure with the zero grading subalgebra  $\mathcal{H}$ , with the usual conformal bosonic generators  $P_\mu$ ,  $K_\mu$ ,  $D$ ,  $J_{\mu\nu}$ , supercharges  $Q$ ,  $\bar{Q}$ , superconformal generators  $S$ ,  $\bar{S}$  and R-symmetry generators  $R_a$ . We can find an Abelian subalgebra  $\mathcal{A}$  composed of a subset of the  $P$ 's and  $Q$ 's if we can find an anti-commuting combination of supercharges. We denote by  $A, B, ..$  the indices of  $\mathcal{A}$ . Uppercase letters denote both bosonic and fermionic indices, lowercase letters bosonic ones and Greek letters fermionic ones,  $A = \{a, \alpha\}$ . We gauge fix the  $\mathcal{H}$  symmetry and parameterize the coset element as follows,

$$g = e^{x_A T^A} e^B, \quad x_A T^A \in \mathcal{A}, \quad B \in \mathcal{G} \ominus \mathcal{A}. \quad (4.1)$$

More specifically we parameterize

$$e^B = \exp(\bar{\theta}\bar{Q} + \xi S + \bar{\xi}\bar{S}) y^D \exp(y^\mu R_\mu / y), \quad (4.2)$$

where contraction of the fermions is understood ( $R_\mu$  includes only the generators in the supercoset,  $R_\mu \in \mathcal{G}/\mathcal{H}$ ). The left-invariant one-form current will split into two pieces,

$$J = e^{-B} dx_a T^a e^B + e^{-B} de^B \equiv j + \mathfrak{j}, \quad (4.3)$$

where  $\mathfrak{j}$  is independent of the coordinates associated with the Abelian subalgebra  $x_a$ . Generally,  $j$  and  $\mathfrak{j}$  can take any value in  $\mathcal{G}$ . We give special indices to these generators as follows,

$$j = j_I T^I, \quad \mathfrak{j} = \mathfrak{j}_W T^W \quad (4.4)$$

(in general  $a$  is both in  $I$  and  $W$  and  $T^I \cap T^W \neq \emptyset$ ), so we can write the general Green-Schwarz sigma model action as,

$$\begin{aligned} S = \int d^2z & \left[ j_I^{(2)\bar{(2)}} j_J^{(2)} \eta^{IJ} + \mathfrak{j}_W^{(2)\bar{(2)}} j_J^{(2)} \eta^{WJ} + j_I^{(2)\bar{(2)}} \mathfrak{j}_X^{(2)} \eta^{IX} + \mathfrak{j}_W^{(2)\bar{(2)}} \mathfrak{j}_X^{(2)} \eta^{WX} \right. \\ & \left. + \frac{1}{2} \left( j_I^{(1)\bar{(3)}} j_J^{(3)} \eta^{IJ} + \mathfrak{j}_W^{(1)\bar{(3)}} j_J^{(3)} \eta^{WJ} + j_I^{(1)\bar{(3)}} \mathfrak{j}_X^{(3)} \eta^{IX} + \mathfrak{j}_W^{(1)\bar{(3)}} \mathfrak{j}_X^{(3)} \eta^{WX} \right) \right] \end{aligned} \quad (4.5)$$

$$\begin{aligned}
& -\frac{1}{2} \left( \bar{j}_I^{(1)} j_J^{(3)} \eta^{IJ} + \bar{j}_W^{(1)} j_J^{(3)} \eta^{WJ} + \bar{j}_I^{(1)} j_X^{(3)} \eta^{IX} + \bar{j}_W^{(1)} j_X^{(3)} \eta^{WX} \right) \Big] \\
= & \int d^2z \left[ A_I'^{(2)} \bar{A}_J'^{(2)} \eta^{IJ} + j_W^{(2)} \bar{A}_J'^{(2)} \eta^{WJ} + A_I'^{(2)} \bar{j}_X^{(2)} \eta^{IX} + j_W^{(2)} \bar{j}_X^{(2)} \eta^{WX} \right. \\
& + \frac{1}{2} \left( A_I'^{(1)} \bar{A}_J'^{(3)} \eta^{IJ} + j_W^{(1)} \bar{A}_J'^{(3)} \eta^{WJ} + A_I'^{(1)} \bar{j}_X^{(3)} \eta^{IX} + j_W^{(1)} \bar{j}_X^{(3)} \eta^{WX} \right) \\
& \left. - \frac{1}{2} \left( \bar{A}_I'^{(1)} A_J'^{(3)} \eta^{IJ} + \bar{j}_W^{(1)} A_J'^{(3)} \eta^{WJ} + \bar{A}_I'^{(1)} j_X^{(3)} \eta^{IX} + \bar{j}_W^{(1)} j_X^{(3)} \eta^{WX} \right) \right. \\
& \left. + \tilde{x}^A (\partial \bar{A}_A - \bar{\partial} A_A) \right] ,
\end{aligned}$$

where  $A'$  equals  $j$  when replacing the coordinate derivatives  $dx_A$  with the gauge field  $A_A$ , and  $\tilde{x}_A$  is the Lagrange multiplier. By the structure of  $j$ , (4.3), we see that  $A'$  depends linearly on  $A$ , so quadratic terms will rise only from  $A' \bar{A}'$  interactions in the action. We also know that

$$A'^{(1)} = Q^I A'_{Q^I} + \bar{Q}^I A'_{\bar{Q}^I} + S^I A'_{S^I} + \bar{S}^I A'_{\bar{S}^I} , \quad (4.6)$$

and similarly for  $A'^{(3)}$ . Generally we encountered two case where  $J_{Q^I}$  coupled to  $\bar{J}_{\bar{Q}^I}$  or to  $\bar{J}_{\bar{Q}^I}$ , namely we have terms in the action of the form

$$c\eta^{IJ} (J_{Q^I} \bar{J}_{Q^J} - J_{\bar{Q}^I} \bar{J}_{\bar{Q}^J}), \quad \text{or} \quad c\eta^{IJ} (J_{Q^I} \bar{J}_{\bar{Q}^J} - \bar{J}_{Q^I} J_{\bar{Q}^J}) , \quad (4.7)$$

where  $c$  is some constant. Let us call these two cases, case I and case II respectively. Case I appeared usually when considering PSU based model and case II when considering ortho-symplectic based model. To zeroth order in the fermions of  $B$  we have

$$A'_i = f(y) A_i, \quad A'_{Q^\iota} = g(y) h_\iota^\kappa (y^\mu / y) A_{Q^\kappa}, \quad A'_{\bar{Q}^\iota} = g(y) \tilde{h}_\iota^\kappa (y^\mu / y) A_{\bar{Q}^\kappa}, \quad (4.8)$$

where the  $\iota, \kappa$  indices are the R-symmetry indices of  $Q, \bar{Q}$ , so  $Q$  might have more indices of transformation under the bosonic conformal group, which are the same on both sides of the equations (4.8). To this order, the equations of motion for the bosons are

$$A_I'^{(2)} \eta^{IJ} f^2 = F^J(j, \tilde{x}^A) , \quad (4.9)$$

while for the fermions we have for the two cases

$$c\eta^{\iota\kappa} g^2 h_\zeta^\kappa h_\iota^\lambda A_\lambda^{(1)} = F^\kappa(j, \tilde{x}^A), \quad \text{and} \quad c\eta^{\iota\kappa} g^2 (\tilde{h}_\zeta^\kappa h_\iota^\lambda - \tilde{h}_\zeta^\lambda h_\iota^\kappa) A_\lambda^{(1)} = G^\kappa(j, \tilde{x}^A) , \quad (4.10)$$

respectively. The matrix  $m^{\kappa\lambda} \equiv \eta^{\iota\kappa} \tilde{h}_\zeta^{[\kappa} h_\iota^{\lambda]}$  in the second equation is singular when the dimension of the representation of  $Q$  under R-symmetry is odd. That is the case for the  $AdS_4 \times CP^3$  action. The matrix  $m$  will get corrections at higher orders in the fermions of  $B$ , but these will contribute additively with even number of fermions to keep  $m$  bosonic. Let us call the full matrix  $M$  so,

$$M = m + \mathcal{O}(\chi^2) , \quad (4.11)$$

where  $\chi = \{\bar{\theta}, \xi, \bar{\xi}\}$ . The inverse matrix should have the form  $M^{-1} = m^{-1} + \mathcal{O}(\chi^2)$ , since all the fermions should cancel, so when  $m$  is singular so is  $M$ .

In case II when the R-symmetry does not mix  $Q$  with  $\bar{Q}$  (as in the  $AdS_4$  case), we don't get a quadratic term to this order. Actually in this case we will get quadratic terms in  $A_Q$  when going to higher orders, so we will get equations of motion for  $A_Q$ , but these will always multiply terms of order  $O(\chi^2)$  or higher in the fermions, so again we will have a singular matrix multiplying  $A_Q$ , and we will not be able to solve for them. The argument above does not rely on fixing any fermionic degrees of freedom (specifically we don't use  $\kappa$ -symmetry).

We saw that there can also be a case III where the WZ term gives an interaction of the form

$$J_Q \bar{J}_S + \dots, \quad (4.12)$$

and this is the case of  $AdS_2$  with eight supersymmetries realized by the supercoset  $PSU(1, 1|2)/(U(1) \times SU(2))$ . In this case, like for  $AdS_4$ ,  $A_S$  does not get corrections to zeroth order in the fermions, so the quadratic term  $\partial\theta\bar{\partial}\theta$  again will be multiplied by a singular fermionic matrix. This case has a different structure since the R-symmetry does not play any role (its generators are in  $\mathcal{H}$ ), instead we have  $K$  (special conformal transformations) that mixes  $Q$  with  $\bar{S}$  and  $\bar{Q}$  with  $S$ .

## Acknowledgments

We would like to thank N. Berkovits, P. A. Grassi, M. Magro, L. Mazzucato, V. Schomerus, D. Sorokin and S. Theisen for valuable discussions. I.A. would like to thank the KITP, UCSB for hospitality during the Fundamental Aspects of String Theory workshop. This research was supported by the German-Israeli Project cooperation (DIP H.52), the German-Israeli Fund (GIF) and supported in part by the National Science Foundation under Grant No. NSF PHY05-51164.

## A. The $osp(2|2)$ algebra

The  $osp(2|2)$  algebra is

$$\begin{aligned} [P, K] &= -2D, & [D, P] &= P, & [D, K] &= -K, & [R, D] &= 0, & [R, P] &= 0, \\ [R, K] &= 0, \\ [P, Q] &= 0, & [P, \bar{Q}] &= 0, & [K, Q] &= S, & [K, \bar{Q}] &= \bar{S}, & [D, Q] &= \frac{1}{2}Q, \\ [D, \bar{Q}] &= \frac{1}{2}\bar{Q}, & [R, Q] &= -iQ, & [R, \bar{Q}] &= i\bar{Q}, \\ [P, S] &= -Q, & [P, \bar{S}] &= -\bar{Q}, & [K, S] &= 0, & [K, \bar{S}] &= 0, & [D, S] &= -\frac{1}{2}S, \\ [D, \bar{S}] &= -\frac{1}{2}\bar{S}, & [R, S] &= -iS, & [R, \bar{S}] &= i\bar{S}, \end{aligned}$$

$$\begin{aligned}
\{Q, Q\} &= 0, & \{\bar{Q}, \bar{Q}\} &= 0, & \{Q, \bar{Q}\} &= 2P, & \{S, S\} &= 0, & \{\bar{S}, \bar{S}\} &= 0, \\
\{S, \bar{S}\} &= 2K, & \{Q, S\} &= 0, & \{\bar{Q}, \bar{S}\} &= 0, \\
\{Q, \bar{S}\} &= 2D - iR, & \{\bar{Q}, S\} &= 2D + iR.
\end{aligned} \tag{A.1}$$

This algebra admits a  $\mathbb{Z}_4$  automorphism. The automorphism relevant to our needs is the one implemented by  $\Omega X \Omega^{-1}$  for  $X \in \mathfrak{osp}(2|2)$ , where

$$\Omega = \begin{pmatrix} -i\sigma_1 & 0 \\ 0 & \mathbf{1} \end{pmatrix}. \tag{A.2}$$

Using this automorphism the algebra can be decomposed into  $\mathbb{Z}_4$ -invariant subspaces  $\mathcal{H}_k$  ( $k = 0 \dots, 3$ ) such that

$$\mathcal{H}_k = \{X \in \mathfrak{osp}(2|2) | \Omega X \Omega^{-1} = i^k X\}.$$

These subspaces are

$$\mathcal{H}_0 = \{P - K, R\}, \tag{A.3}$$

$$\mathcal{H}_1 = \{Q + S, \bar{Q} + \bar{S}\}, \tag{A.4}$$

$$\mathcal{H}_2 = \{P + K, D\}, \tag{A.5}$$

$$\mathcal{H}_3 = \{Q - S, \bar{Q} - \bar{S}\}. \tag{A.6}$$

The Cartan-Killing bilinear form is defined by  $g_{XY} = \text{Str}(XY)$ . Its non-trivial elements are

$$\begin{aligned}
\text{Str}(PK) &= \text{Str}(KP) = -1, & \text{Str}(DD) &= \frac{1}{2}, & \text{Str}(RR) &= 2, \\
\text{Str}(Q\bar{S}) &= \text{Str}(\bar{Q}S) = 2.
\end{aligned} \tag{A.7}$$

## B. The $\mathfrak{osp}(2|4)$ algebra

The definition of the  $\text{OSp}(2|4)$  group is that of [21]. The non-trivial brackets of the  $\mathfrak{osp}(2|4)$  algebra are

$$\begin{aligned}
[M_{mn}, M_{pq}] &= \eta_{mp}M_{nq} + \eta_{nq}M_{mp} - \eta_{mq}M_{np} - \eta_{np}M_{mq}, \\
[M_{mn}, P_p] &= \eta_{mp}P_n - \eta_{np}P_m, & [M_{mn}, K_p] &= \eta_{mp}K_n - \eta_{np}K_m, \\
[D, P_m] &= P_m, & [D, K_m] &= -K_m, & [P_m, K_n] &= 2\eta_{mn}D - 2M_{mn}, \\
[D, Q_\alpha] &= \frac{1}{2}Q_\alpha, & [D, \hat{Q}_\alpha] &= \frac{1}{2}\hat{Q}_\alpha, & [D, S_\alpha] &= -\frac{1}{2}S_\alpha, & [D, \hat{S}_\alpha] &= -\frac{1}{2}\hat{S}_\alpha, \\
[M_{mn}, Q_\alpha] &= \frac{1}{2}(C\gamma_{mn}C^{-1})_\alpha^\beta Q_\beta, & [M_{mn}, \hat{Q}_\alpha] &= \frac{1}{2}(C\gamma_{mn}C^{-1})_\alpha^\beta \hat{Q}_\beta, \\
[M_{mn}, S_\alpha] &= \frac{1}{2}(C\gamma_{mn}C^{-1})_\alpha^\beta S_\beta, & [M_{mn}, \hat{S}_\alpha] &= \frac{1}{2}(C\gamma_{mn}C^{-1})_\alpha^\beta \hat{S}_\beta,
\end{aligned}$$

$$\begin{aligned}
[P_m, S_\alpha] &= i(C\gamma_m C^{-1})_\alpha^\beta Q_\beta, & [P_m, \hat{S}_\alpha] &= i(C\gamma_m C^{-1})_\alpha^\beta \hat{Q}_\beta, \\
[K_m, Q_\alpha] &= -i(C\gamma_m C^{-1})_\alpha^\beta S_\beta, & [K_m, \hat{Q}_\alpha] &= -i(C\gamma_m C^{-1})_\alpha^\beta \hat{S}_\beta, \\
[R, Q_\alpha] &= iQ_\alpha, & [R, \hat{Q}_\alpha] &= -i\hat{Q}_\alpha, & [R, S_\alpha] &= iS_\alpha, & [R, \hat{S}_\alpha] &= -i\hat{S}_\alpha \\
\{Q_\alpha, \hat{Q}_\beta\} &= 4(C\gamma^m)_{\alpha\beta} P_m, & \{S_\alpha, \hat{S}_\beta\} &= -4(C\gamma^m)_{\alpha\beta} K_m, \\
\{Q_\alpha, \hat{S}_\beta\} &= -2i(C\gamma^{mn})_{\alpha\beta} M_{mn} + 4i\epsilon_{\alpha\beta} D - 4\epsilon_{\alpha\beta} R, \\
\{\hat{Q}_\alpha, S_\beta\} &= -2i(C\gamma^{mn})_{\alpha\beta} M_{mn} + 4i\epsilon_{\alpha\beta} D + 4\epsilon_{\alpha\beta} R,
\end{aligned} \tag{B.1}$$

where  $\epsilon_{12} = -\epsilon_{21} = 1$ ,  $\epsilon^{21} = -\epsilon^{12} = 1$ ,  $C_{\alpha\beta} = \epsilon_{\alpha\beta}$  and the Dirac matrices have the index structure  $\gamma^{m\alpha}_\beta$  and indices are lowered and raised using  $C$ . The anti-symmetrized Dirac-matrices are  $\gamma^{mn} = \frac{1}{2}[\gamma^m, \gamma^n]$ . The metric is  $\eta = \text{diag}(-1, 1, 1)$ ,  $m, n, p, q = 0, \dots, 2$  are space-time indices,  $\alpha, \beta = 1, 2$  are spinor indices and  $I, J = 1, 2$  are SO(2) R-symmetry indices.

The non-trivial elements of the Cartan-Killing bilinear form are

$$\begin{aligned}
\text{Str}(M_{mn}M_{pq}) &= \eta_{mp}\eta_{nq} - \eta_{mq}\eta_{np}, & \text{Str}(DD) &= -1, & \text{Str}(RR) &= -2, \\
\text{Str}(P_m K_n) &= -2\eta_{mn}, & \text{Str}(Q_\alpha \hat{S}_\beta) &= -8i\epsilon_{\alpha\beta}, & \text{Str}(\hat{Q}_\alpha S_\beta) &= -8i\epsilon_{\alpha\beta}.
\end{aligned} \tag{B.2}$$

The  $\mathbb{Z}_4$  automorphism invariant subspaces are

$$\begin{aligned}
\mathcal{H}_0 &= \{M_{mn}, P_m - K_m, R\}, \\
\mathcal{H}_1 &= \{Q_\alpha - S_\alpha, \hat{Q}_\alpha + \hat{S}_\alpha\}, \\
\mathcal{H}_2 &= \{P_m + K_m, D\}, \\
\mathcal{H}_3 &= \{Q_\alpha + S_\alpha, \hat{Q}_\alpha - \hat{S}_\alpha\}.
\end{aligned} \tag{B.3}$$

## C. The osp(6|4) algebra in $\text{so}(1, 2) \oplus \text{u}(3)$ basis

The osp(6|4) algebra's commutation relations are given by

$$[\lambda_{ki}, \lambda_{m\hat{n}}] = 2i(\delta_{mi}\lambda_{k\hat{n}} - \delta_{k\hat{n}}\lambda_{mi}) \tag{C.1}$$

$$[\lambda_{ki}, R_{mn}] = 2i(\delta_{mi}R_{kn} - \delta_{ni}R_{km}) \tag{C.2}$$

$$[R_{mn}, R_{kl}] = 0, \quad [R_{mn}, R_{ki}] = \frac{i}{2}(\delta_{mk}\lambda_{ni} - \delta_{mi}\lambda_{nk} - \delta_{nk}\lambda_{mi} + \delta_{ni}\lambda_{mk}) \tag{C.3}$$

$$[P_a, P_b] = 0, \quad [K_a, K_b] = 0, \quad [P_a, K_b] = 2\eta_{ab}D - 2M_{ab} \tag{C.4}$$

$$[M_{ab}, M_{cd}] = \eta_{ac}M_{bd} + \eta_{bd}M_{ac} - \eta_{ad}M_{bc} - \eta_{bc}M_{ad} \tag{C.5}$$

$$[M_{ab}, P_c] = \eta_{ac}P_b - \eta_{bc}P_a, \quad [M_{ab}, K_c] = \eta_{ac}K_b - \eta_{bc}K_a \tag{C.6}$$

$$[D, P_a] = P_a, \quad [D, K_a] = -K_a, \quad [D, M_{ab}] = 0 \tag{C.7}$$

$$[D, Q_\alpha^l] = \frac{1}{2}Q_\alpha^l, \quad [D, S_\alpha^l] = -\frac{1}{2}S_\alpha^l \tag{C.8}$$

$$[P_a, Q_\alpha^l] = 0, \quad [K_a, S_\alpha^l] = 0 \quad (\text{C.9})$$

$$[P_a, S_\alpha^l] = -i(\gamma_a)_\alpha^\beta Q_\beta^l, \quad [K_a, Q_\alpha^l] = i(\gamma_a)_\alpha^\beta S_\beta^l \quad (\text{C.10})$$

$$[M_{ab}, Q_\alpha^l] = -\frac{i}{2}(\gamma_{ab})_\alpha^\beta Q_\beta^l, \quad [M_{ab}, S_\alpha^l] = -\frac{i}{2}(\gamma_{ab})_\alpha^\beta S_\beta^l \quad (\text{C.11})$$

$$[R_{kl}, Q_\alpha^p] = i(\delta^{pl} Q_\alpha^k - \delta^{pk} Q_\alpha^l), \quad [R_{kl}, S_\alpha^p] = -i(\delta^{pl} S_\alpha^k - \delta^{pk} S_\alpha^l) \quad (\text{C.12})$$

$$[R_{ki}, Q_\alpha^p] = -i(\delta^{pi} Q_\alpha^k - \delta^{pk} Q_\alpha^i), \quad [R_{ki}, S_\alpha^p] = i(\delta^{pi} S_\alpha^k - \delta^{pk} S_\alpha^i) \quad (\text{C.13})$$

$$[\lambda_{ki}, Q_\alpha^p] = 2i\delta^{pi} Q_\alpha^k, \quad [\lambda_{ki}, S_\alpha^p] = 2i\delta^{pi} S_\alpha^k \quad (\text{C.14})$$

$$[\lambda_{ki}, Q_\alpha^{\dot{p}}] = -2i\delta^{\dot{p}k} Q_\alpha^i, \quad [\lambda_{ki}, S_\alpha^{\dot{p}}] = -2i\delta^{\dot{p}k} S_\alpha^i \quad (\text{C.15})$$

$$\{Q_\alpha^l, Q_\beta^k\} = 0, \quad \{Q_\alpha^l, Q_\beta^{\dot{k}}\} = -\delta^{lk}(\gamma^a C)_{\alpha\beta} P_a \quad (\text{C.16})$$

$$\{S_\alpha^l, S_\beta^k\} = 0, \quad \{S_\alpha^l, S_\beta^{\dot{k}}\} = -\delta^{lk}(\gamma^a C)_{\alpha\beta} K_a \quad (\text{C.17})$$

$$\{Q_\alpha^l, S_\beta^k\} = -C_{\alpha\beta} R_{lk}, \quad \{Q_\alpha^{\dot{l}}, S_\beta^{\dot{k}}\} = -C_{\alpha\beta} R_{\dot{l}\dot{k}} \quad (\text{C.18})$$

$$\{Q_\alpha^l, S_\beta^{\dot{k}}\} = -i\delta^{lk}(C_{\alpha\beta} D + i\frac{1}{2}(\gamma^{ab} C)_{\alpha\beta} M_{ab}) + \frac{1}{2}C_{\alpha\beta} \lambda_{lk} \quad (\text{C.19})$$

$$\{Q_\alpha^{\dot{l}}, S_\beta^k\} = i\delta^{\dot{l}k}(C_{\alpha\beta} D - i\frac{1}{2}(\gamma^{ab} C)_{\alpha\beta} M_{ab}) + \frac{1}{2}C_{\alpha\beta} \lambda_{\dot{l}k} \quad (\text{C.20})$$

The indices take the values  $k, l = 1, 2, 3$  and the same for the dotted ones — the  $\mathbf{3}$  and  $\bar{\mathbf{3}}$  of  $u(3)$ ,  $a, b = 0, 1, 2$  are the  $\mathbf{3}$  of  $so(1, 2)$  and  $\alpha, \beta, .. = 1, 2$  are the  $so(2, 1)$  spinors, and  $\eta = \text{diag}(-, +, +)$ . The generators satisfy  $R_{kl}^* = R_{ki}$  and  $\lambda_{ki} = \lambda_{ik}^*$  and  $Q_\alpha^{\dot{i}} = (Q_\alpha^i)^*$ . The  $(\gamma_a)_\alpha^\beta$  are the Dirac matrices of  $so(1, 2)$ , and  $\gamma_{ab} = \frac{i}{2}[\gamma_a, \gamma_b]$ . We raise and lower spinor indices as explained in appendix E.

The bilinear forms are given by

$$\text{Str}(R_{kl} R_{\dot{p}\dot{q}}) = -2(\delta^{k\dot{p}} \delta^{l\dot{q}} - \delta^{k\dot{q}} \delta^{l\dot{p}}) \quad (\text{C.21})$$

$$\text{Str}(Q_\alpha^l S_\beta^{\dot{k}}) = 2i\delta^{lk} C_{\alpha\beta} \quad (\text{C.22})$$

$$\text{Str}(P_a K_b) = -2\eta_{ab} \quad (\text{C.23})$$

$$\text{Str}(DD) = -1 \quad (\text{C.24})$$

$$\text{Str}(M_{ab} M_{cd}) = \eta_{ac} \eta_{bd} - \eta_{ad} \eta_{bc} \quad (\text{C.25})$$

The  $\mathbb{Z}_4$  automorphism matrix is given by

$$\Omega = \left( \begin{array}{ccc|cc} i\sigma_2 & 0 & 0 & 0 & 0 \\ 0 & i\sigma_2 & 0 & 0 & 0 \\ 0 & 0 & i\sigma_2 & 0 & 0 \\ \hline 0 & 0 & 0 & \sigma_2 & 0 \\ 0 & 0 & 0 & 0 & -\sigma_2 \end{array} \right). \quad (\text{C.26})$$

The  $\mathbb{Z}_4$  invariant subspaces of the algebra are

$$\begin{aligned}
\mathcal{H}_0 &= \{P_a - K_a, M_{ab}, \lambda_{lk}\}, \\
\mathcal{H}_1 &= \{Q_\alpha^l - S_\alpha^l, \hat{Q}_\alpha^l - \hat{S}_\alpha^l\}, \\
\mathcal{H}_2 &= \{P_a + K_a, D, R_{kl}, R_{ki}\}, \\
\mathcal{H}_3 &= \{Q_\alpha^l + S_\alpha^l, \hat{Q}_\alpha^l + \hat{S}_\alpha^l\}.
\end{aligned} \tag{C.27}$$

## D. The $\text{psu}(1, 1|2)$ algebra

The algebra of the  $\text{PSU}(1, 1|2)$  group was developed by using the definition given in [18] (which follows the definition given in [22] applied to matrices whose elements are all Grassmann-even).

The  $\text{su}(1, 1|2)$  algebra in a basis oriented towards the  $\text{SU}(2)$  R-symmetry is

$$\begin{aligned}
[D, P] &= P, \quad [D, K] = -K, \quad [P, K] = -2D, \quad [R_i, R_j] = -\epsilon_{ijk} R_k, \\
[D, Q_\alpha] &= \frac{1}{2} Q_\alpha, \quad [D, \hat{Q}_\alpha] = \frac{1}{2} \hat{Q}_\alpha, \quad [D, S_\alpha] = -\frac{1}{2} S_\alpha, \quad [D, \hat{S}_\alpha] = -\frac{1}{2} \hat{S}_\alpha, \\
[P, Q_\alpha] &= [P, \hat{Q}_\alpha] = 0, \quad [P, S_\alpha] = i \hat{Q}_\alpha, \quad [P, \hat{S}_\alpha] = i Q_\alpha, \\
[K, Q_\alpha] &= i \hat{S}_\alpha, \quad [K, \hat{Q}_\alpha] = i S_\alpha, \quad [K, S_\alpha] = [K, \hat{S}_\alpha] = 0, \\
[R_{ij}, Q_\alpha] &= -\frac{1}{2} \sigma_{ij\alpha}{}^\beta Q_\beta, \quad [R_{ij}, \hat{Q}_\alpha] = -\frac{1}{2} \sigma_{ij\alpha}{}^\beta \hat{Q}_\beta, \\
[R_{ij}, S_\alpha] &= -\frac{1}{2} \sigma_{ij\alpha}{}^\beta S_\beta, \quad [R_{ij}, \hat{S}_\alpha] = -\frac{1}{2} \sigma_{ij\alpha}{}^\beta \hat{S}_\beta, \\
\{Q_\alpha, \hat{Q}_\beta\} &= \epsilon_{\alpha\beta} P, \quad \{S_\alpha, \hat{S}_\beta\} = \epsilon_{\alpha\beta} K, \quad \{Q_\alpha, S_\beta\} = -\frac{i}{2} \epsilon_{\beta\gamma} \sigma_\alpha^{ij\gamma} R_{ij} - i \epsilon_{\alpha\beta} D, \\
\{Q_\alpha, \hat{S}_\beta\} &= \frac{i}{2} \epsilon_{\alpha\beta} \mathbf{1}, \quad \{\hat{Q}_\alpha, S_\beta\} = -\frac{i}{2} \epsilon_{\alpha\beta} \mathbf{1}, \quad \{\hat{Q}_\alpha, \hat{S}_\beta\} = \frac{i}{2} \epsilon_{\beta\gamma} \sigma_\alpha^{ij\gamma} R_{ij} + i \epsilon_{\alpha\beta} D,
\end{aligned} \tag{D.1}$$

where  $a, b = 1, 2$  are  $\text{AdS}_2$  indices,  $\alpha, \beta, \gamma = 1, 2$  are  $\text{SU}(2)$  R-symmetry indices,  $R_i$  ( $i = 1, \dots, 3$ ) are the generators of the  $\text{SU}(2)$  R-symmetry and  $R_{ij} = \epsilon_{ijk} R_k$  are defined for convenience. The antisymmetric tensors are defined such that  $\epsilon_{12} = 1$ ,  $\epsilon_{123} = 1$  and the generators of  $\text{SU}(2)$  are  $\sigma_{ij\alpha}{}^\beta = \frac{1}{2} (\sigma_{i\alpha\gamma} \sigma_j^{\gamma\beta} - \sigma_{j\alpha\gamma} \sigma_i^{\gamma\beta})$  with  $\sigma_i$  being the Pauli matrices. In order to get the  $\text{psu}(1, 1|2)$  algebra, one has to divide by the  $\text{U}(1)$  generator  $\mathbf{1}$ . The Grassmann-odd generators in the algebra above are not Hermitian but are formed as linear combinations of the original Hermitian matrices with complex coefficients. These combinations correspond to the multiplying the original Hermitian generators by the complex Killing spinors found by the requiring the supercharges and superconformal transformations to form Abelian subalgebras.

The non-trivial elements of the Cartan-Killing bilinear form are

$$\begin{aligned}
\text{Str}(PK) &= -1, \quad \text{Str}(DD) = \frac{1}{2}, \quad \text{Str}(R_i R_j) = \frac{1}{2} \delta_{ij}, \\
\text{Str}(Q_\alpha S_\beta) &= -i \epsilon_{\alpha\beta}, \quad \text{Str}(\hat{Q}_\alpha \hat{S}_\beta) = i \epsilon_{\alpha\beta}
\end{aligned} \tag{D.2}$$

The  $\mathbb{Z}_4$  automorphism is the same one as given in [18],

$$\Omega = \begin{pmatrix} \sigma_3 & 0 \\ 0 & i\sigma_3 \end{pmatrix}. \quad (\text{D.3})$$

The  $\mathbb{Z}_4$  invariant subspaces of the algebra defined as

$$\mathcal{H}_k = \{X \in \text{psu}(1, 1|2) | \Omega X \Omega^{-1} = i^k X\} \quad (\text{D.4})$$

are

$$\begin{aligned} \mathcal{H}_0 &= \{P + K, R_3\}, \\ \mathcal{H}_1 &= \{Q_2 + S_2, \hat{Q}_2 + \hat{S}_2, Q_1 - S_1, \hat{Q}_1 - \hat{S}_1\}, \\ \mathcal{H}_2 &= \{P - K, D, R_1, R_2\}, \\ \mathcal{H}_3 &= \{Q_1 + S_1, \hat{Q}_1 + \hat{S}_1, Q_2 - S_2, \hat{Q}_2 - \hat{S}_2\}. \end{aligned} \quad (\text{D.5})$$

Another  $\mathbb{Z}_4$  automorphism is given by

$$\Omega = \begin{pmatrix} \sigma_1 & 0 \\ 0 & i\mathbf{I} \end{pmatrix} \quad (\text{D.6})$$

yields the following invariant subspaces of the algebra

$$\begin{aligned} \mathcal{H}_0 &= \{D, R_i\}, \\ \mathcal{H}_1 &= \{Q_\alpha, \hat{Q}_\alpha\}, \\ \mathcal{H}_2 &= \{P, K\}, \\ \mathcal{H}_3 &= \{S_\alpha, \hat{S}_\alpha\}. \end{aligned} \quad (\text{D.7})$$

This will give us the supercoset space  $\text{PSU}(1, 1|2)/(\text{U}(1) \times \text{SU}(2))$  which is isomorphic to  $AdS_2$  with eight supersymmetries.

## E. The $\text{psu}(1, 1|2) \oplus \text{psu}(1, 1|2)$ algebra

We build the super-algebra by taking the direct sum of two  $\text{psu}(1, 1|2)$  algebras (where we take complex combinations of the odd generators), so the bosonic part is  $\text{su}(2) \oplus \text{su}(2) \simeq \text{so}(4)$  and  $\text{su}(1, 1) \oplus \text{su}(1, 1) \simeq \text{so}(2, 2)$ . The non-vanishing part of the  $\text{psu}(1, 1|2) \oplus \text{psu}(1, 1|2)$  algebra, involving odd generators, is given by

$$[D, P_a] = P_a, \quad [D, K_a] = -K_a, \quad [D, J_{ab}] = 0, \quad (\text{E.1})$$

$$[P_a, K_a] = 2(\eta_{ab}D + J_{ab}), \quad [P_c, J_{ab}] = \eta_{ac}P_b - \eta_{bc}P_a, \quad [K_c, J_{ab}] = \eta_{ac}P_b - \eta_{bc}K_a,$$

$$[R_\mu, R_\nu] = -N_{\mu\nu}, \quad [R_\rho, N_{\mu\nu}] = \delta_{\rho\mu}R_\nu - \delta_{\rho\nu}R_\mu,$$

$$[N_{\rho\sigma}, N_{\mu\nu}] = \delta_{\rho\mu}N_{\sigma\nu} + \delta_{\sigma\nu}N_{\rho\mu} - \delta_{\rho\nu}N_{\sigma\mu} - \delta_{\sigma\mu}N_{\rho\nu}$$



$$\begin{aligned}
[P_a, Q_{\alpha\hat{\alpha}}^I] &= 0, & [K_a, S_{\alpha\hat{\alpha}}^I] &= 0, & [D, Q_{\alpha\hat{\alpha}}^I] &= \frac{1}{2}Q_{\alpha\hat{\alpha}}^I, & [D, S_{\alpha\hat{\alpha}}^I] &= -\frac{1}{2}S_{\alpha\hat{\alpha}}^I, \\
[K_a, Q_{\alpha\hat{\alpha}}^I] &= i(\hat{\gamma}_a)_{\hat{\alpha}}^{\hat{\beta}}S_{\alpha\hat{\beta}}^I, & [P_a, S_{\alpha\hat{\alpha}}^I] &= i(\hat{\gamma}_a)_{\hat{\alpha}}^{\hat{\beta}}Q_{\alpha\hat{\beta}}^I, \\
[J_{01}, Q_{\alpha\hat{\alpha}}^I] &= -\frac{i}{2}(\hat{\gamma}_{01})_{\hat{\alpha}}^{\hat{\beta}}Q_{\alpha\hat{\beta}}^I, & [J_{01}, S_{\alpha\hat{\alpha}}^I] &= -\frac{i}{2}(\hat{\gamma}_{01})_{\hat{\alpha}}^{\hat{\beta}}S_{\alpha\hat{\beta}}^I, \\
[R_\mu, S_{\alpha\hat{\alpha}}^I] &= \frac{i}{2}(i\hat{\gamma}_{01})_{\hat{\alpha}}^{\hat{\beta}}(\gamma_\mu)_{\alpha}^{\beta}S_{\beta\hat{\beta}}^I, & [R_\mu, Q_{\alpha\hat{\alpha}}^I] &= -\frac{i}{2}(i\hat{\gamma}_{01})_{\hat{\alpha}}^{\hat{\beta}}(\gamma_\mu)_{\alpha}^{\beta}Q_{\beta\hat{\beta}}^I, \\
[N_{\mu\nu}, S_{\alpha\hat{\alpha}}^I] &= \frac{i}{2}\hat{C}_{\hat{\alpha}}^{\hat{\beta}}(\gamma_{\mu\nu})_{\alpha}^{\beta}S_{\beta\hat{\beta}}^I, & [N_{\mu\nu}, Q_{\alpha\hat{\alpha}}^I] &= \frac{i}{2}\hat{C}_{\hat{\alpha}}^{\hat{\beta}}(\gamma_{\mu\nu})_{\alpha}^{\beta}Q_{\beta\hat{\beta}}^I, \\
\{S_{\alpha\hat{\alpha}}^I, S_{\beta\hat{\beta}}^J\} &= \frac{i}{2}\epsilon^{IJ}C_{\alpha\beta}(\hat{\gamma}^a\hat{C})_{\hat{\alpha}\hat{\beta}}K_a, & \{Q_{\alpha\hat{\alpha}}^I, Q_{\beta\hat{\beta}}^J\} &= \frac{i}{2}\epsilon^{IJ}C_{\alpha\beta}(\hat{\gamma}^a\hat{C})_{\hat{\alpha}\hat{\beta}}P_a, \\
\{S_{\alpha\hat{\alpha}}^I, Q_{\beta\hat{\beta}}^J\} &= \frac{i}{2}\epsilon^{IJ}(C_{\alpha\beta}(i\hat{C}_{\hat{\alpha}\hat{\beta}}D - \frac{1}{2}(\hat{C}\hat{\gamma}^{ab})_{\hat{\alpha}\hat{\beta}}J_{ab}) \\
&\quad + \frac{1}{2}\hat{C}_{\hat{\alpha}\hat{\beta}}N_{\mu\nu}(\gamma^{\mu\nu}C)_{\alpha\beta} - i(\hat{\gamma}_{01}\hat{C})_{\hat{\alpha}\hat{\beta}}R_\mu(\gamma^\mu C)_{\alpha\beta})
\end{aligned}$$

where the indices go as  $I = 1, 2$ ,  $\hat{\alpha} = 1, 2$ ,  $\alpha = 1, 2$ ,  $a = 1, 2$ ,  $\mu = 1, 2, 3$  and  $C_{\alpha\beta} = \epsilon_{\alpha\beta}$ ,  $\hat{C}_{\hat{\alpha}\hat{\beta}} = \epsilon_{\hat{\alpha}\hat{\beta}}$ . The gamma matrices are defined by  $(\hat{\gamma}_a)_{\hat{\alpha}}^{\hat{\beta}} = \{i\sigma_2, -\sigma^1\}$ ,  $(\hat{\gamma}_{01})_{\hat{\alpha}}^{\hat{\beta}} = \frac{i}{2}[\hat{\gamma}_0, \hat{\gamma}_1]_{\hat{\alpha}}^{\hat{\beta}} = \{-i\sigma^3\}$ ,  $(\gamma_\mu)_{\alpha}^{\beta} = \{\sigma_1, \sigma_3, \sigma_2\}$ ,  $(\gamma_{\mu\nu})_{\alpha}^{\beta} = \frac{i}{2}[\gamma_\mu, \gamma_\nu]_{\alpha}^{\beta} = \{\sigma_2, \sigma_1, \sigma_3\}$ , where  $\sigma$  are the Pauli matrices, and  $\eta_{ab} = \text{diag}(+, -)$ . We raise and lower spinor indices using  $\psi_\alpha = \psi^\beta\epsilon_{\beta\alpha}$ ,  $\psi^\alpha = \epsilon^{\alpha\beta}\psi_\beta$ , where  $\epsilon_{12} = -\epsilon_{21} = \epsilon^{12} = -\epsilon^{21} = 1$  and the same for hatted objects. The bilinear Cartan-Killing forms can be rescaled to give

$$\text{Str}(P_a K_b) = \eta_{ab}, \quad \text{Str}(DD) = 1, \quad \text{Str}(J_{ab} J_{cd}) = \delta_{ac}\delta_{bd} - \delta_{ad}\delta_{bc} \quad (\text{E.2})$$

$$\begin{aligned}
\text{Str}(R_\mu R_\nu) &= \delta_{\mu\nu}, & \text{Str}(N_{\mu\nu} N_{\rho\sigma}) &= \delta_{\mu\rho}\delta_{\nu\sigma} - \delta_{\mu\sigma}\delta_{\nu\rho} \\
\text{Str}(Q_{\alpha\hat{\alpha}}^I S_{\beta\hat{\beta}}^J) &= \epsilon^{IJ}\epsilon_{\alpha\beta}\epsilon_{\hat{\alpha}\hat{\beta}}
\end{aligned}$$

The super-algebra has  $\mathbb{Z}_4$  grading structure using the automorphism matrix

$$\Omega = \begin{pmatrix} 0 & 0 & \sigma_3 & 0 \\ 0 & 0 & 0 & 1_{2 \times 2} \\ -i\sigma_3 & 0 & 0 & 0 \\ 0 & i1_{2 \times 2} & 0 & 0 \end{pmatrix} \quad (\text{E.3})$$

so

$$\mathcal{H}_0 = \{J_{01}, N_{\mu\nu}, P_0 - K_0, P_1 - K_1\} \quad (\text{E.4})$$

$$\mathcal{H}_1 = \{S_{\alpha\hat{\alpha}}^I + a^{IJ}Q_{\alpha\hat{\alpha}}^J\} \quad (\text{E.5})$$

$$\mathcal{H}_2 = \{R_\mu, D, P_0 + K_0, P_1 + K_1\} \quad (\text{E.6})$$

$$\mathcal{H}_3 = \{S_{\alpha\hat{\alpha}}^I - a^{IJ}Q_{\alpha\hat{\alpha}}^J\} \quad (\text{E.7})$$

where  $a^{IJ} = \sigma_1$ . So we can get the supercoset space  $\text{PSU}(1, 1|2)^2/(\text{SU}(1, 1) \times \text{SU}(2))$ , whose bosonic part is  $AdS_3 \times S^3$ . We also have another  $\mathbb{Z}_4$  grading structure with the same number of supersymmetry generators

$$\mathcal{H}_0 = \{D, J_{01}, N_{\mu\nu}, R_\mu\} \quad (\text{E.8})$$

$$\mathcal{H}_1 = \{Q_{\alpha\dot{\alpha}}^I\} \quad (\text{E.9})$$

$$\mathcal{H}_2 = \{P_a, K_a\} \quad (\text{E.10})$$

$$\mathcal{H}_3 = \{S_{\alpha\dot{\alpha}}^I\} \quad (\text{E.11})$$

which is the four-dimensional space  $B\text{DI}(2, 2) \simeq AdS_2 \times AdS_2$ , given by the supercoset  $\text{PSU}(1, 1|2)^2/(\text{SO}(2)^2 \times \text{SO}(4))$ .

## References

- [1] L. F. Alday and J. M. Maldacena, *Gluon scattering amplitudes at strong coupling*, *JHEP* **06** (2007) 064, [0705.0303].
- [2] J. M. Drummond, J. Henn, V. A. Smirnov, and E. Sokatchev, *Magic identities for conformal four-point integrals*, *JHEP* **01** (2007) 064, [hep-th/0607160].
- [3] J. M. Drummond, G. P. Korchemsky, and E. Sokatchev, *Conformal properties of four-gluon planar amplitudes and Wilson loops*, *Nucl. Phys.* **B795** (2008) 385–408, [0707.0243].
- [4] N. Berkovits and J. Maldacena, *Fermionic T-Duality, Dual Superconformal Symmetry, and the Amplitude/Wilson Loop Connection*, *JHEP* **09** (2008) 062, [0807.3196].
- [5] N. Beisert, R. Ricci, A. A. Tseytlin, and M. Wolf, *Dual Superconformal Symmetry from AdS5 x S5 Superstring Integrability*, 0807.3228.
- [6] I. Bena, J. Polchinski, and R. Roiban, *Hidden symmetries of the AdS(5) x S\*\*5 superstring*, *Phys. Rev.* **D69** (2004) 046002, [hep-th/0305116].
- [7] I. Adam, A. Dekel, L. Mazzucato, and Y. Oz, *Integrability of type II superstrings on Ramond-Ramond backgrounds in various dimensions*, *JHEP* **06** (2007) 085, [hep-th/0702083].
- [8] J.-G. Zhou, *Super 0-brane and GS superstring actions on AdS(2) x S(2)*, *Nucl. Phys.* **B559** (1999) 92–102, [hep-th/9906013].
- [9] J. Park and S.-J. Rey, *Green-Schwarz superstring on AdS(3) x S(3)*, *JHEP* **01** (1999) 001, [hep-th/9812062].
- [10] J. Rahmfeld and A. Rajaraman, *The GS string action on AdS(3) x S(3) with Ramond-Ramond charge*, *Phys. Rev.* **D60** (1999) 064014, [hep-th/9809164].

- [11] G. Arutyunov and S. Frolov, *Superstrings on  $AdS_4 \times CP^3$  as a Coset Sigma-model*, *JHEP* **09** (2008) 129, [0806.4940].
- [12] j. Stefanski, B., *Green-Schwarz action for Type IIA strings on  $AdS_4 \times CP^3$* , *Nucl. Phys.* **B808** (2009) 80–87, [0806.4948].
- [13] P. Fre and P. A. Grassi, *Pure Spinor Formalism for  $Osp(N-4)$  backgrounds*, 0807.0044.
- [14] G. Bonelli, P. A. Grassi, and H. Safaai, *Exploring Pure Spinor String Theory on  $AdS_4 \times CP^3$* , *JHEP* **10** (2008) 085, [0808.1051].
- [15] R. D’Auria, P. Fre, P. A. Grassi, and M. Trigiante, *Superstrings on  $AdS_4 \times CP^3$  from Supergravity*, 0808.1282.
- [16] T. H. Buscher, *A Symmetry of the String Background Field Equations*, *Phys. Lett.* **B194** (1987) 59.
- [17] T. H. Buscher, *Path Integral Derivation of Quantum Duality in Nonlinear Sigma Models*, *Phys. Lett.* **B201** (1988) 466.
- [18] N. Berkovits, M. Bershadsky, T. Hauer, S. Zhukov, and B. Zwiebach, *Superstring theory on  $AdS(2) \times S(2)$  as a coset supermanifold*, *Nucl. Phys.* **B567** (2000) 61–86, [hep-th/9907200].
- [19] O. Aharony, O. Bergman, D. L. Jafferis, and J. Maldacena,  *$N=6$  superconformal Chern-Simons-matter theories, M2-branes and their gravity duals*, *JHEP* **10** (2008) 091, [0806.1218].
- [20] J. Gomis, D. Sorokin, and L. Wulff, *The complete  $AdS(4) \times CP(3)$  superspace for the type IIA superstring and D-branes*, 0811.1566.
- [21] L. Frappat, P. Sorba, and A. Sciarrino, *Dictionary on Lie superalgebras*, hep-th/9607161.
- [22] V. G. Kac, *A Sketch of Lie Superalgebra Theory*, *Commun. Math. Phys.* **53** (1977) 31–64.