

Analytical Solutions of Open String Field Theory

Ehud Fuchs, Michael Kroyter

*Max-Planck-Institut für Gravitationsphysik
Albert-Einstein-Institut
14476 Golm, Germany*

Abstract

In this work we review Schnabl's construction of the tachyon vacuum solution to bosonic covariant open string field theory and the results that followed.

We survey the state of the art of string field theory research preceding this construction focusing on Sen's conjectures and the results obtained using level truncation methods.

The tachyon vacuum solution is described in various ways, in particular we describe its geometric representation using wedge states and its formal algebraic representation as a gauge solution. We also derive the form of the solution's building blocks in the oscillator representation. We show that some of Sen's conjectures can be proven analytically using this solution.

The tools used in the context of the vacuum solution can be adapted to the construction of other solutions, namely various marginal deformations. We present some of the approaches used in the construction of these solutions.

The generalization to open superstring field theory is derived in details. We start from the exposition of the problems one faces in the construction of superstring field theory. We then present the cubic and the non-polynomial versions of superstring field theory and prove their classical equivalence. Finally, the bosonic solutions are generalized to this case. In particular we focus on the (somewhat surprising) generalization of the tachyon solution to the case of a theory with no tachyons.

Key words: String Field Theory

PACS: 11.25.Sq

Email addresses: udif@aei.mpg.de (Ehud Fuchs), mikroyt@aei.mpg.de (Michael Kroyter).

Contents

1	Introduction	3
1.1	Scattering amplitudes	5
2	String field theory basics	7
3	Sen's conjectures	16
3.1	Level truncation	18
4	Algebraic primer	26
5	Schnabl's solution	33
5.1	Sen's first conjecture	36
5.2	Sen's third conjecture	39
5.3	Generalizations of the solution	41
6	Marginal deformations	43
6.1	Regular marginal deformations	44
6.2	Singular marginal deformations	47
7	Oscillator representation	52
7.1	Representation of string vertices and the continuous basis	52
7.2	Schnabl's operators and the resolution of normalization anomalies	60
8	Superstring field theory	64
8.1	Superstring variables, pictures and conventions	64
8.2	Introducing superstring field theories	75
8.3	Mapping solutions among (super)string field theories	80
8.4	Analytical solutions in superstring field theory	83
9	Outlook	88
	References	91

1 Introduction

String field theory is a field theoretical framework for string theory. As such, it can potentially be used for evaluating scattering amplitudes using Feynman diagram techniques, for studying non-perturbative effects of the theory and for exploring various string vacua by studying classical solutions.

Many attempts have been made to find analytical solutions of (open covariant bosonic) string field theory, since its introduction by Witten [1]. It took twenty years until such a solution was constructed by Schnabl [2]. This construction drew much attention and was followed by several other analytical solutions and other generalizations that improved significantly our understanding of string field theory. It is the purpose of this work to review these developments so as to make them more accessible to the general string theory community.

Previous attempts towards analytical solutions within string field theory used various tools. Among other approaches one can find:

- Conformal field theory (CFT) methods [3,4]: Defining the star product and integration in terms of CFT expressions enables the usage of many tools, gives a geometric picture to the problem, which can then be dealt with in various conformal frames. Moreover, many CFT methods are insensitive to the given background and are thus universal. Universality is an important property of solutions that represent tachyon condensation [5]. Indeed, CFT methods proved essential for the recent progress.
- Half-string formalism [6,7,8,9,10]: A fundamental entity in the construction of string field theory is the star-product, which can be cast into a form analogous to a matrix product, with the two halves of the string playing the roles of the two matrix indices. Bogoliubov transformation to the vacuum built upon the two string halves simplifies the form of the star-product. One less favorable feature of this approach (and some of the next ones) is that it is based on a given (flat) background and is therefore not universal.
- Simplifying the non-linear term in the equation of motion [11]: The equation of motion contains a single non-linear piece. As in the case of the Riccati equation, simplifying the form of the non-linear term has the potential of rendering the equation solvable. This approach led to the algebraic construction of the surface-state star-projector known as the sliver. The construction itself turned out to be equivalent to the half-string approach.
- Representing the star-product in a Moyal form [12,13,14,15,16,17,18]: A natural framework for addressing non-commutative products is the Moyal approach, which gained popularity in the string theory community following [19]. It was found that the star-product can be represented as an infinite dimensional tensor product of Moyal pairs.
- Diagonalizing the star-product [20,21]: In the oscillator basis, the star-

product is defined using some infinite dimensional matrices that can (almost) be simultaneously diagonalized. This results in the continuous basis (aka κ -basis). This approach led to some results in the matter sector. The inclusion of the ghosts, however, turns out to be more problematic.

- Vacuum string field theory [22,23,24,25,26,27,28,29]: This is not a genuine way to solve string field theory. Instead, the form of the theory expanded around the (tachyon) solution, after an unknown (and presumably singular) field redefinition, is guessed. This approach led to numerous analytical results. However, its scope of validity and its relation to the standard string field theory were not fully clarified.
- Identity string field based solutions [30,31,32,33]: The star-product contains an identity element. This state is, however, singular in several respects, namely it is not a normalizable state [34] and in the CFT description it represents surface gluing. Moreover, it is not killed by the derivation c_0 [35], implying $0 \neq c_0 |1\rangle = c_0(|1\rangle \star |1\rangle) = (c_0 |1\rangle) \star |1\rangle + |1\rangle \star (c_0 |1\rangle) = 2c_0 |1\rangle$, i.e., a contradiction. An advantage of the identity based solutions is that they are universal. On the other hand, while the identity based solutions are formally well defined, the problems with the identity string field rendered the evaluation of the action of these solutions ambiguous.
- Using other surface states to define solutions [36,37,38,39]: In order to bypass the problems with the identity string field, while keeping some of the nice properties of the solutions, one can choose to work with other star-algebra projectors, such as the butterfly and the sliver. Constructing solutions turns out to be more complicated in this case than in the case of the identity based solutions.

In this work we focus on Schnabl's solution and the developments that followed. This construction uses ideas from most of the approaches described above. Hence, we introduce some of these tools. While we tried to make this review self-contained, the reader may want to consult also older reviews focusing on other aspects of string field theory. The following is a list of related reviews.

- An introduction to string field theory emphasizing the Batalin-Vilkovisky (BV) approach and perturbation theory is the classical review by Thorn [40].
- Shorter modern general introduction to string field theory can be found in [41,42].
- The issue of tachyon condensation in string field theory and vacuum string field theory is addressed in [43].
- General and extensive reviews of tachyon condensation covering also other frameworks for addressing this subject are [44,45,46,47,48].
- A review that describes many representations of the string field algebra with some focus on string field theory as a non-commutative field theory is [49].
- The Moyal approach to string field theory is introduced in [50].
- Two papers that stress the algebraic structure of the star product are [51,52].

Let us describe the content of the various sections (by their order) and their interrelations, in order to answer the perpetual question:

How to use this review?

- (1) In the rest of the introduction we briefly describe the advance in computing scattering amplitudes in string field theory, where impressive progress has been obtained. While this is not a result regarding analytical solutions, it is important and relevant enough to be shortly described here. We refer the reader to the original works for a complete and clear exposition of this important subject.
- (2) This is an introductory section, where string field theory basics are described. Researchers familiar with its construction can skip this part.
- (3) Another introductory section, in which Sen's conjectures and their study within level truncation are described. It serves as a motivation and a historical overview and can also be skipped.
- (4) Here we present some of the tools needed for the derivation of Schnabl's solution.
- (5) In this section Schnabl's construction of the tachyon vacuum is derived using the tools of the previous section. The proofs of Sen's conjectures for this solution are then described.
- (6) This section presents some of the ways by which analytical solutions describing marginal deformations were obtained within string field theory. It relies on some basic concepts from the previous two sections.
- (7) The recent advance relies mainly on CFT technics. Still, we give an oscillator description of Schnabl's solution in this section. It relies only on some basic formulas from section 5 and the oscillator approach is explained therein. Other sections do not depend on this one and it can be skipped by readers interested in the main root of advance in the field.
- (8) Here we describe the generalization of the matter presented in previous sections to open superstring field theory. Peculiarities and derivation of superstring field theories are given. This section relies on most of the previous ones.
- (9) An outlook.

1.1 Scattering amplitudes

In string field theory scattering amplitudes can be defined even off-shell. Off-shell scattering amplitudes can be used as building blocks for loop amplitudes as well as for the evaluation of the tachyon-potential. The first calculation of off-shell amplitudes was performed in [53]. The evaluation of the quartic term in the tachyon potential immediately followed [54]. This calculation was extended to higher levels and developed in [55,56,57] and in the Moyall approach [16,18].

Scattering amplitudes were commonly calculated in the Siegel gauge. The attractive properties of the gauge choice introduced by Schnabl changed that. The Schnabl gauge has also some less attractive properties. It is based on an operator, which is not hermitian. It is singular in some sense and two Schwinger parameters should be used to describe a propagator instead of one [2]. Preliminary results in the Schnabl gauge were obtained in [58]. In this paper it was also suggested to use a hermitian version of Schnabl’s operator in order to fix the gauge.

The scattering amplitudes calculated in [58] seemed not to obey some expected symmetries. A careful study of the subject [59] showed that the propagator should be regularized. The results now were shown to obey some of the symmetries, while it was claimed that the other ones are peculiar to the Siegel gauge.

Up to this point only (off-shell) tree amplitudes were considered in the Schnabl gauge. A thorough study of scattering amplitudes using a general gauge choice based on the b -ghost, termed linear b -gauges, was performed in [60]. Regularity conditions on the gauge choice were stated and it was found that the Schnabl gauge is located on the boundary of the regular region in the space of b -coefficients. Nevertheless, no explicit problems with the results in Schnabl gauge were obtained, unlike in the case of its hermitian counterpart. Another important discovery of [60] was that Schnabl’s condition cannot be imposed at all ghost numbers. This result does not afflict previous calculations, since it is of importance only when loop amplitudes are considered, in which string fields of arbitrary ghost number appear. General b -gauge choices imply different gauge conditions at different ghost numbers. Still, the question remains, whether the Schnabl gauge can be used for the evaluation of general loop amplitudes. The disturbing (and novel) property of this gauge is that it fixes the string mid-point. Thus, it may seem that moduli space cannot be covered.

This problem was addressed in [61], where the Schnabl gauge was defined as a limit of regular b -gauges. A careful study of the limit showed that the limiting wedge states should be glued not in the standard way, but in a slant, which is responsible to a “hidden boundary at infinity”. The introduction of slanted wedges enabled the authors to show that at least at 1-loop level the moduli space is covered in the Schnabl gauge, despite the apparent problems. This seems as a strong evidence in favour of the possibility that Schnabl’s gauge can be used to an arbitrary loop level. Moreover, it was claimed that the results are technically much easier to derive than in the Siegel gauge. There is still work to be done, though. It would be desirable to study the general covering of moduli spaces, to calculate explicit scattering amplitudes and to address the Schnabl gauge using a BV formalism.

2 String field theory basics

String field theory is an approach to string theory, which is at least conceptually the most straightforward one. This approach generalizes the particle physics tools of second quantization to string theory. In particle physics second quantization amounts to taking the Schrödinger equation of a particle and reinterpret it as the equation of motion of a scalar field. It turns out that upon quantization, such a field describes just a collection of non-interacting particles of the type of the original one. To this one can add interaction terms obeying some principles, such as symmetry and renormalizability, in order to get the full field theory of interacting particles. One added advantage is that the classical equations of motion of the field can also reveal non-perturbative effects. Thus, the quantum interaction of particles and the classical, non-perturbative effects, are related.

In string theory the “first quantized” world-sheet approach already gives the form of the interaction in the form of the Polyakov path-integral. One may hope then, that the string field theory is uniquely defined by the world-sheet theory. If this is the case, one may hope to find this string field theory and then study it both for understanding and controlling string interactions and as a tool for the study of non-perturbative effects is string theory. Indeed, already in 1990, before the big advance in understanding non-perturbative string theory following the introduction of D-branes by Polchinski [62], a numerical solution to the equations of motion of string field theory was found [63]. This solution amounts to the closed string vacuum left after the removal of a D-brane. This, however, was not understood at the time and the interpretation of the solution had to wait to the introduction of the celebrated Sen’s conjectures, to be described in section 3.

There are several perturbative string theories and one should be chosen in order to construct a string field theory. There are also several approaches for quantization on the world-sheet, e.g., light-cone. So we stress at this point that unless otherwise stated, we shall be dealing with open bosonic strings and will do so in a covariant formalism, where one can hope to gain some insight about the general symmetry structure of the theory.

At the linearized level string field theory should give the equation of motion [64,65]

$$Q_B \Psi = 0, \tag{2.1}$$

where Ψ is the string field and Q_B is the BRST operator on the world-sheet of the open bosonic string. One can think of the string field as a linear combination with momentum-dependent coefficients (fields) of states of the first quantized ghost number one space. Hence, it is an off-shell generalization of

the vertex operator. In the usual flat-space it takes the form,

$$|\Psi\rangle = \int d^{26}k \left(T(k)c_1 + C(k)c_0 + A_\mu(k)\alpha_{-1}^\mu c_1 + \dots \right) |k\rangle, \quad (2.2)$$

with the usual definition

$$|k\rangle = e^{ikX(0)} |0\rangle. \quad (2.3)$$

The infinity of fields composing it are the momentum-dependent coefficients of the various modes. Each can be characterized by its level, that is its eigenvalue with respect to $L_0 + 1$ (ignoring the momentum part). Thus, the tachyon field T is of level zero, while the photon field A_μ and the (auxiliary) field C are of level one. The free equation of motion (2.1) should be accompanied by the free gauge transformation

$$\Psi \rightarrow \Psi + Q_B \Lambda. \quad (2.4)$$

Thus, the equation of motion of the string field corresponds to the condition of being closed and the gauge transformation identifies string states whose difference is exact. The gauge string field itself consists of an off-shell linear combinations of states with ghost number zero. This structure resembles that of differential forms, with Q_B playing the role of the differential, since it obeys

$$Q_B^2 = 0. \quad (2.5)$$

Note however, that unlike for forms in a finite dimensional space, here there are “forms” of an arbitrary integer degree, from minus infinity to infinity. The gauge transformation (2.4) is accordingly reducible at an infinite order, since a transformation of the form

$$\Lambda^{(n)} \rightarrow \Lambda^{(n)} + Q_B \Lambda^{(n-1)}, \quad (2.6)$$

does not contribute to the transformation of $\Lambda^{(n+1)}$ performed in the same way, due to (2.5). Here, the superscript represents the “form-degree”. Since Q_B increases the ghost number by one, the form-degree can be identified with the (first quantization) ghost number. Being a one-form, it is natural to declare that the string field is an odd element.

An appropriate linearized action is [66]

$$S_{\text{lin}} = -\frac{1}{2} \langle \Psi | Q_B | \Psi \rangle, \quad (2.7)$$

where $\langle \Psi |$ is the BPZ conjugate of $|\Psi\rangle$. Let us also explain the coefficient in front of (2.7). It is needed in order to get the canonical sign for the component fields. For the tachyon field T , which is the first field in the expansion (2.2),

one gets

$$\begin{aligned}
S &= -\frac{1}{2} \int d^{26}k d^{26}k' T(k)T(k') \langle k | c_{-1} Q c_1 | k' \rangle = \\
&\frac{1}{2} \int d^{26}k d^{26}k' T(k)T(k') \langle k | c_{-1} c_0 c_1 (1 - k'^2) | k' \rangle = \int d^{26}k \frac{1 - k^2}{2} T(k)^2.
\end{aligned} \tag{2.8}$$

Here we used the BPZ conjugation property of the c field, as well as the oscillator expansion of Q_B , neglecting in the intermediate step some terms that do not contribute to the inner product. We also defined $\alpha' = 1$ and used

$$\langle 0 | c_{-1} c_0 c_1 | 0 \rangle = \frac{1}{2} \langle \partial^2 c \partial c c \rangle = 1, \tag{2.9}$$

as the normalization convention in the ghost sector. Given our other convention, that of a mostly positive metric, the resulting action for the tachyon field T has the canonical form and we also see that it is indeed a tachyonic field. For higher level fields one may have to adjust the coefficient in the definition of the field in order to get a canonical form. The sign is, however, canonical for all fields.

The inclusion of interactions into this formalism was performed by Witten [1]. The action is given by

$$S = - \int \left(\frac{1}{2} \Psi \star Q \Psi + \frac{g_o}{3} \Psi \star \Psi \star \Psi \right). \tag{2.10}$$

Here g_o is the open string coupling constant that can (at least classically) be set to unity by a field rescaling. We shall set it to unity unless otherwise stated. From (2.10), one can derive the equation of motion¹,

$$Q \Psi + \Psi \star \Psi = 0. \tag{2.11}$$

The integration, the star product and the derivation Q are (bi-)linear operations over the space of (arbitrary ghost number) string fields \mathcal{A}

$$\int : \mathcal{A} \rightarrow \mathbb{C}, \quad \star : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}, \quad Q : \mathcal{A} \rightarrow \mathcal{A}. \tag{2.12}$$

Witten assumed that these entities obey some algebraic relations, described below. He claimed that the emergent (noncommutative geometry) structure is a natural one for describing string field theory. Regardless of this structure, the

¹ The derivation is somewhat formal at this stage. In order to be able to truly derive it, one should first specify the meaning of the operations involved and conclude that a variant of the fundamental lemma of the calculus of variations holds in this case. All that, in fact, works out well.

first requirement of interacting string field theory is that in the limit $g_o \rightarrow 0$, it reduces to the non-interacting theory (2.7). This is the case if one identifies²,

$$Q = Q_B, \quad \int \Psi_1 \star \Psi_2 = \langle \Psi_1 | \Psi_2 \rangle. \quad (2.13)$$

Henceforth, Q will represent the BRST operator Q_B . Other realizations of Q (denoted \mathcal{Q}), obeying the algebraic structure, are also of importance. A particular example is obtained when the theory (2.10) is expanded around a solution. We describe this case in section 3.

Other than linearity, Witten's axioms state that the star product is associative and Q is an odd derivation of it, namely³

$$Q^2 = 0, \quad Q(A_1 \star A_2) = (QA_1) \star A_2 + (-1)^{A_1} A_1 \star (QA_2), \quad \int QA = 0. \quad (2.14)$$

Here, $A_{1,2}$ are string fields of an arbitrary ghost number. The star product preserves the grading and the ghost number. Another axiom states that under integration the string fields behave like differential forms,

$$\int A_1 \star A_2 = (-1)^{A_1 A_2} \int A_2 \star A_1. \quad (2.15)$$

These axioms are identical to those obeyed by differential forms, except the fact that for differential forms (2.15) holds even without integration⁴. The action (2.10) can thus be considered as a generalization of the Chern-Simon action.

From the analogy with the Chern-Simon action we recognize that the action (2.10) is invariant under the infinitesimal gauge transformation

$$\delta\Psi = Q\Lambda + \Psi \star \Lambda - \Lambda \star \Psi, \quad (2.16)$$

where Λ is a ghost number zero string field. This infinitesimal gauge transformation can be exponentiated to give the finite gauge transformation,

$$\Psi \rightarrow e^{-\Lambda}(Q + \Psi)e^{\Lambda}, \quad (2.17)$$

² The definition of $\int \Psi_1 \star \Psi_2 \star \Psi_3$ is more involved. We return to this issue shortly.

³ Throughout this paper A in $(-1)^A$ represents the grading of A .

⁴ It turns out that in the bosonic case the integral is non-zero only for a string field of ghost number three. Thus, for the bosonic string the sign factor $(-1)^{\Psi_1 \Psi_2}$ equals unity. However, if the string fields themselves are allowed to carry Grassmann grading, the sign can become important. This is the case when the BV formalism is used in order to handle the gauge symmetry. Moreover, in the case of Berkovits' version of superstring field theory, to be introduced in section 8.2, the total ghost number should equal two and the minus sign is important.

where multiplication, exponentiation, as well as any other function of string fields, should be performed by expanding the relevant function in a series, while keeping the order of string fields intact and using the star product for the evaluation of all products. The 1 that one gets at zero order is realized by the identity string field, described below. No ambiguities can emerge, since the star product is the only way by which string fields can be multiplied. We continue to write it explicitly in the introductory sections, but it will be left implicit in the remaining of the paper, except in cases where its inclusion improves readability.

The gauge transformation (2.16) is reducible, since a variation of the form

$$\delta\Lambda = Q\Lambda_{-1} + \Psi \star \Lambda_{-1} - \Lambda_{-1} \star \Psi, \quad (2.18)$$

induces no change for Ψ in (2.16), provided that Ψ obeys the equation of motion (2.11), i.e., it is on-shell. Similarly,

$$\delta\Lambda_{-n} = Q\Lambda_{-(n+1)} + \Psi \star \Lambda_{-(n+1)} - \Lambda_{-(n+1)} \star \Psi, \quad (2.19)$$

will induce no change in Λ_{-n+1} . This follows from

$$\delta^2 A = (Q\Psi + \Psi \star \Psi) \star A - A \star (Q\Psi + \Psi \star \Psi), \quad (2.20)$$

where again A is a string field of an arbitrary degree. Thus, the (non-linear) gauge generator δ is nilpotent only on-shell [67], in which case the gauge system is infinitely reducible. A sensible way to deal with reducible gauge systems, as well as with gauge systems that close only on-shell, is given by the BV formalism [68,69,70,71,72] (and the reviews [73,74]).

The BV formalism was implied for quantizing string field theory in [75,76,77]⁵. It turns out that the infinity of ghost fields that should be added can be identified with string fields of (first-quantized) ghost number $n_g < 1$, while the infinity of anti-fields can be identified with string fields of ghost number $n_g > 1$. The identification is such that the ghost numbers of a field and its anti-field sum up to 3, i.e, the anti-field of the classical (ghost number one) string field carries ghost number two, the anti-field of the first (ghost number zero) ghost carries ghost number three and so on. As usual, the coefficient fields carry alternating parity. When this alternating parity is combined with the parity of the various ghost-number string states, it results in string fields of a fixed, odd, parity. These string fields can be summed to give (an odd) string field, whose ghost number is not constrained,

$$\Psi = \sum_{n=-\infty}^{\infty} \Psi_n. \quad (2.21)$$

⁵ A BRST approach to string field theory was implied in [78,79,80] using the “mid-point light-cone gauge”.

The classical master equation is obeyed by an action, which is identical in form to (2.10), only with the string field being given by (2.21), rather than by Ψ_1 . This action can be gauge fixed by the introduction of a gauge fixing fermion. A common gauge choice is the Siegel gauge

$$b_0\Psi = 0. \tag{2.22}$$

This is an extension of the Feynman gauge for photons, since this condition results in the Feynman gauge for the massless vector component of the string field, upon level truncation⁶. In this gauge, the equation of motion (2.11) reduces to

$$c_0L_0\Psi + \Psi \star \Psi = 0. \tag{2.23}$$

In order to preserve associativity, Witten suggested that the star product should implement the geometric picture of gluing the right half of the first string with the left half of the second string, while integration should amount to gluing together both halves of the same string around the middle. This suggestion is consistent with (2.13), since in this case one gets just a gluing of two strings with their orientation reversed, i.e., the BPZ inner product [84]. With these identifications of the star product, the integral and the derivation, it was shown by Giddings et. al. that the Veneziano amplitude is reproduced from the action (2.10) and that at higher order in g_o it reduces to the Polyakov path integral with correct covering of moduli space [85,86,87]. A gap in the proof of moduli space covering was filled up by Zwiebach [88].

For practical calculations it is useful to represent the star product using either CFT expressions as derived in [3,4] or using oscillator expressions in the flat background as described in [89,90,91,92,93,94]. The former method has the advantage of being universal. That is, it does not depend on a choice of a background, as long as it is composed of a bc system coupled to a $c = 26$ BCFT. Indeed, most of the recent developments in the field were achieved using this formulation. Nevertheless, the later method has also proved useful, especially, but not only, for describing phenomena peculiar to a given background. We now turn to describe the CFT approach. We describe the oscillator method and the recent related results in section 7.

In the seminal papers [3,4], LeClair, Peskin and Preitschopf defined more

⁶ One can consider also other gauges, such as the analogue of the Landau gauge. A study of gauges that interpolate between the Feynman-Siegel and the Landau gauges was carried out in [81,82]. Related discussion was given in [83]. Note, however, that there is more than one possible extension of, say the Feynman gauge, for the string field. Another such extension, the Schnabl gauge, played important role in the recent developments.




Fig. 1. The N-vertex represented on the unit disk.

general string vertices by CFT expectation values on the disk, as

$$\int \Psi_1 \star \dots \star \Psi_n = \langle f_1^{(n)} \circ \Psi_1(0), \dots, f_n^{(n)} \circ \Psi_n(0) \rangle. \quad (2.24)$$

Here, the string fields on the left hand side are interpreted as operators acting on the vacuum. In the right hand side a given set of conformal transformations acts on these operators, mapping them from the upper half plane to various points on the disk, in some given canonical representation. The CFT expectation value with this insertions gives the desired result. The formalism is general. In order to reproduce Witten's theory the conformal transformations should be fixed, such that one gets the advocated gluing. If we choose the unit disk as the canonical representative, the conformal transformations take the form

$$f_k^{(n)} = \left(\frac{1 + i\xi}{1 - i\xi} \right)^{\frac{2}{n}} e^{\frac{2\pi i k}{n}}. \quad (2.25)$$

Note that the local coordinate patch in the upper half plane, i.e., the top half-unit-disk, is transformed under these maps to wedges of angle $\frac{2\pi}{n}$ and these wedges are glued together such that the right half of the k^{th} string is glued to the left half of the $(k + 1)^{\text{th}}$ string. In this representation the symmetry of the vertex under cyclic permutation is manifest. Other coordinate systems can be more adequate for specific problems. We return to this point in section 4.

The CFT representation makes it clear that the n -vertices are multi-linear functions. In particular, the 2-vertex forms a bi-linear map, rather than a hermitian inner product, as one may expect [95]. It is possible to impose a reality condition on the string field,

$$\Psi^* = \Psi. \quad (2.26)$$

The conjugation in the above equation represents a hermitian conjugation fol-

lowed by the inversion of the orientation of the string. In terms of the operators producing the string state this conjugation corresponds to the composition of hermitian and BPZ conjugations. When this reality condition is imposed, the 2-vertex forms also a hermitian inner-product.

The CFT methods of [3,4] can be used not only for defining string vertices, but also for defining string fields of a special class, called surface states, which will be useful in the following sections. As the name implies, a surface state is associated with a given surface Σ . The association is in fact the one used to define the one-vertex, only with the function f of (2.25) being (almost) arbitrary. An arbitrary test state, Ψ formed by an operator insertion on the vacuum,

$$\Psi = \mathcal{O}(0) |0\rangle, \tag{2.27}$$

can be defined to be contracted with the surface state $\langle S|$ associated with Σ according to

$$\langle S|\Psi\rangle = \langle f \circ \mathcal{O} \rangle_{\Sigma}, \tag{2.28}$$

where the brackets represent a CFT expectation value.

In order to actually being able to evaluate the CFT expectation value, it is useful to have a canonical form for Σ , say the upper half plane. It may seem that such a choice removes all information about the surface state, but this is not the case, since the map f is still arbitrary. One can think of it in the following way. The operator inserted at the origin of the upper half plane corresponds to a string state defined on the half unit circle. The inner product corresponds to gluing this string state that evolves from zero, with another one that evolves from infinity. The half disk in which the test state evolves $|z| \leq 1$ is called the local coordinate patch. For a surface state it is possible to describe the string that evolves from infinity by a surface with open string boundary conditions instead of by an insertion at infinity. Conformal transformations, under which the theory is invariant, can be used either to fix the form of the whole disk to a canonical form, in which case the form of the local coordinate patch is non-standard, or to fix the form of the local coordinate patch to a canonical form, leaving the surface Σ arbitrary (and possibly with cuts and double covers for its image in the complex plane). The restriction that we have on the form of the surfaces is only that they should contain a local coordinate patch and leave it intact, i.e, no insertions are allowed in this region, since it is “reserved” for the test state.

The simplest surface state is the vacuum, for which the conformal map is the identity map. Using the standard local coordinate patch, while evaluating the CFT expectation value on the upper half plane, is the same as a contraction with the vacuum state. Surface states in general are given by finite deformations of the upper half plane. Infinitesimal deformations of the upper half plane are given by the Virasoro generators. Exponentiating one gets the

general form of a surface state,

$$|S\rangle = \exp\left(\sum_{n=2}^{\infty} s_n L_{-n}\right) |0\rangle, \quad (2.29)$$

where the coefficients s_n are related to the conformal transformation f in some way [96]. This form for the state implies that in the oscillator representation a surface state is a squeezed state. The converse, however, does not necessarily hold.

Another important corollary is that all these states are BRST closed,

$$Q |S\rangle = 0, \quad (2.30)$$

since the BRST charge commutes by construction with all the Virasoro generators. An important special case is

$$Q |1\rangle = 0, \quad (2.31)$$

where $|1\rangle$ is the identity string field, which is a surface state by definition, since it is generated by the conformal transformation f defining the one-vertex. The identity (2.31) is important for the consistency of Witten's axioms, since the identity string field spans a one-dimensional subspace isomorphic to the complex numbers inside the string field algebra and “constants” should be annihilated by any derivation, Q in particular.

3 Sen's conjectures

An important advance in string field theory followed the realization that it is the ideal framework for addressing Sen's conjectures [97,5]. Sen's work addressed the fate of the tachyon that is living on the bosonic D-brane⁷. According to the conjectures, the tachyon describes the instability of the D-brane. An effective tachyon potential should have a local maximum around zero, where the D-brane exists. This potential should also have a local minimum⁸. A solution in which the tachyon acquires this value as a vev (other fields also acquire vevs) represents the absence of the original D-brane⁹.

This picture suggests the following properties:

- (1) The depth of the local minimum equals the tension of the original D-brane. This reflects the energy difference between the solutions with and without the D-brane.
- (2) Other solutions exist, representing lower dimensional D-branes. These solutions are lumps from the perspective of the tachyon field.
- (3) There are no perturbative states around the tachyon solution, since perturbative states in open string field theory represent open string degrees of freedom and there are no open strings when the D-brane is absent.




Fig. 2. The tachyon potential in units of D-brane tension.

The first of Sen's conjectures states the following. Fix the value of the tachyon field to a constant and solve the equations of motion of all the other components of the string field. The action as a function of the tachyon field should be similar to the one depicted in figure 2. The local minimum of the potential represents the solution describing the condensation of the D-brane. Thus, the

⁷ Similar conjectures hold also for the open string tachyon on a non-BPS D-brane and the tachyon living on the D- \bar{D} pair [98,99,100,101,102].

⁸ The tachyon potential in the supersymmetric cases is bounded from below and the potential is symmetric around the origin. The bosonic potential is not symmetric and CFT analysis suggests that it is unbounded below.

⁹ The actual time-dependent process of tachyon condensation, was described by Sen in the "rolling tachyon" papers [103,104].

action of the solution should equal minus the tension of the D-brane (with proper normalization of the space-time volume).

The second conjecture related to studying properties of solutions that are generally space-time dependent. For a solution to represent a lump of, say, co-dimension one, it has to reduce to the vacuum solution when one moves in this one dimension away from the core, which should be thin in some sense. Analogously to the first and third conjectures, the solution should have the correct tension and carry the appropriate perturbative spectrum in order to describe a D-brane of the appropriate dimension.

In order to understand the third conjecture we have to examine the form of the action (2.10) when expanded around the solution. The expansion is obtained by substituting

$$\Psi \rightarrow \Psi_0 + \Psi, \quad (3.1)$$

into the action. As the original action is cubic it is also cubic with respect to the shifted string field. The zero order term is the constant that is used for testing the first conjecture. However, as a constant it does not contribute to the equation of motion and so can be neglected. The first order term is absent by definition, since we expand around a solution, which is an extremum of the action. The third order term cannot be changed by a shift of the field. Thus, theories that one gets by expansion around different backgrounds differ only in the form of their kinetic, i.e. second order, term. Direct evaluation gives,

$$S = - \int \left(\frac{1}{2} \Psi \star \mathcal{Q}\Psi + \frac{1}{3} \Psi \star \Psi \star \Psi \right). \quad (3.2)$$

The new kinetic operator is defined by its action on a string field with arbitrary ghost number A by,

$$\mathcal{Q}A = QA + \Psi_0 \star A - (-)^A A \star \Psi_0. \quad (3.3)$$

This form of the kinetic operator is not fixed by the ghost number one string field Ψ that appears in the action. Moreover, as far as the action is concerned, we could have written the kinetic operator as $\mathcal{Q}\Psi = Q\Psi + 2\Psi \star \Psi_0$. We have to choose the symmetric form $\mathcal{Q}\Psi = Q\Psi + \Psi \star \Psi_0 + \Psi_0 \star \Psi$ in order to be able to write the equation of motion around the solution as,

$$\mathcal{Q}\Psi + \Psi \star \Psi = 0. \quad (3.4)$$

The dependence on the parity of the form $-(-)^A$ appearing in (3.3) implies the derivation property of the new kinetic operator

$$\mathcal{Q}\mathcal{Q}A = 0 \quad \iff \quad \mathcal{Q}\Psi_0 + \Psi_0 \star \Psi_0 = 0. \quad (3.5)$$

This implies that the gauge symmetry written in terms of \mathcal{Q} takes exactly the same form as when written in terms of Q (2.16), (2.19).

Sen’s conjectures were addressed using several formalisms, such as effective theory analysis [105,106,107,108,109] and toy models such as p-adic string (field) theory [110,111,112,113] (developed in [114,115])¹⁰. However, as the conjectures involve a zero-momentum tachyon, which is quite far from being on-shell, it should be clear that a field theoretical approach is advantageous. Other than the cubic string field theory, which is the main framework described in this review, there exists also BSFT [116,117,118,119]¹¹. Analytical solutions within the framework of BSFT [120,121,122] have been derived with much less effort than those of cubic string field theory. The reason is that one can consistently truncate all the other fields and consider the action for only the tachyon field. However, it is not quite clear how should the massive (non-renormalizable) string modes be dealt within BSFT. While the analysis of tachyon condensation seems to be consistent within BSFT, it is desirable to have such a description also in the framework of cubic string field theory, which is more adequate for the study of other problems.

It turned out that within the framework of cubic string field theory the problem is technically much harder, since the tachyon field does not decouple. The main tool for studying the solution was the numerical approach called “level truncation” to which we turn next.

3.1 Level truncation

Level truncation was first introduced quite some time before Sen’s conjectures by Kostelecky and Samuel [63], following a preliminary study [54]. These authors tried to address the problem of dealing with an infinite number of coefficient fields by truncating the string field and the action to include only fields whose level is at most l . Then, they assumed (without a formal proof¹²) that by taking the limit $l \rightarrow \infty$ the correct string field theory result are obtained. They carried out their analysis up to level four and indeed, they managed to identify convergence to a solution with energy lower than that of the trivial solution. They realized that the solution is non-perturbative with respect to the open string coupling constant g_o . They even managed to notice that there is a decrease in the amount of open string degrees of freedom around the non-perturbative vacuum, although they did not have a reason to suspect that

¹⁰ We refer the reader to some of the reviews mentioned in the introduction for details regarding developments using other formalisms.

¹¹ BSFT stands for either “background-independent string field theory” or for “boundary string field theory”, as it involves the evaluation of correlation functions in a theory perturbed by the insertion of (open string) operators at the boundary.

¹² In fact, in [54] they write “Our truncation is *not* systematic”. In [63], on the other hand, the authors do give some justification and write “The approach is systematic”. The real “proof” of the power of this approach is “experimental”: it works very well.

none are left.

At that time, neither Sen's conjectures nor D-branes [62] were known. Thus, while some papers did improve much the analysis of the level-truncation results [123], it was not until Sen's conjectures that this tool (and string field theory in general) gained popularity.

String field theory has a huge amount of gauge freedom (2.16). Level truncation is usually performed with a fixed gauge. The Siegel gauge was almost exclusively chosen. Gauge fixing achieves two goals. First, the amount of fields in a given level is reduced. Then, level truncation breaks the gauge symmetry, since the gauge symmetry mixes fields of an arbitrary level. The gauge symmetry is restored in the infinite limit level. Gauge transformations induce flat directions in the potential. At a fixed level these flat directions are not exactly flat and flatness is achieved only asymptotically as the level goes to infinity. These almost flat directions have many shallow minima. Thus, without gauge fixing one finds many more solutions than one would like to find, as many of them approximate solutions that lie on the same gauge orbit. However, since gauge symmetry is only approximate in level truncation, it is generally hard to disentangle the gauge solutions from the real ones. On the other hand, gauge fixing can also induce problems, since gauge choices are generally not globally well defined. This goes under the name of Gribov ambiguities in gauge theories. The range of validity of the Siegel gauge was studied by Ellwood and Taylor [124]. They found that both the trivial and the non-perturbative vacua remain in the range where the Siegel gauge is valid, as the level is increased. Outside this region they found a complicated pattern for the boundaries of Siegel gauge validity that bifurcates as the level is increased.

One can make level truncation more efficient by using symmetries. As mentioned by Sen in [5], a set of fields that does not include the tachyon and that enter at least quadratically in the action, can be consistently set to zero when looking for the tachyon vacuum. The equations of motion of the fields that were set to zero are satisfied, as they are all at least linear with respect to these fields. This simple observation can be used in many ways to reduce the amount of fields to be considered. First, one can set to zero all states that carry momentum, since the condensation is with respect to the zero-momentum tachyon field and the action carried no momentum. A related observation is that the tachyon field is a Lorentz scalar as is also the action. At least two non-scalars are needed in order to produce a scalar (the action). Thus, only scalars should be kept in the search for the tachyon vacuum. One can also set to zero all the fields with a non-unity ghost number¹³. Further, it is enough to consider

¹³ While the gauge invariant action is a functional only of a ghost-number-one string fields, the BV treatment of the gauge symmetry introduces a string field of arbitrary ghost number (even before fixing the gauge). However, the tachyon field is of ghost number one and the fields with other ghost numbers indeed enter the action at least

only string fields that live in the “universal subspace” spanned by the ghost fields and the matter Virasoro generators¹⁴ [5]. It was shown in [35] that ghost-number one universal states (named $\mathcal{H}_{univ}^{(1)}$) are spanned by applying all possible combinations of matter-Virasoro operators and ghost-Virasoro operators on the vacuum $c_1 |0\rangle$ ¹⁵. This provided a simple systematic enumeration of the states relevant for tachyon condensation. Other symmetries with respect to which the tachyon state and the action are scalars include twist symmetry and the $SU(1, 1)$ symmetry of [125] (see also [126,127]). The twist generator is represented in $\mathcal{H}_{univ}^{(1)}$ as $(-1)^{L_0+1}$. Thus, the reduction to twist even states corresponds to using only even levels. The $SU(1, 1)$ symmetry can be applied when one works in the Siegel gauge. Then, it acts on the ghost sector with the generators

$$G = \sum_{n=1}^{\infty} (c_{-n}b_n - b_{-n}c_n), \quad X = \sum_{n=1}^{\infty} nc_{-n}c_n, \quad Y = \sum_{n=1}^{\infty} \frac{1}{n}b_{-n}b_n. \quad (3.6)$$

Here G is the ghost number, while the other generators induce rotations between the ghost and the anti-ghost modes.

Other methods used for restricting the solutions in level truncation are the use of the residual BRST symmetry [127] and the use of conservation laws [35]. Conservation laws imply relations among various coefficients of the string field [128]. More importantly, they allowed for the automatization of the calculations, which enabled going to an extremely high precision. Finally, in [129] it was found numerically that some coefficients of the string field take universal values and these values were evaluated analytically. It was suggested there that the universality property follows from (an infinity of) linear relations that the solution should obey and an example of such a relation was derived. Nonetheless, no full understanding of this universality property was achieved and some problems with it were found in [130].

quadratically. Thus, it is possible to consider only ghost-number-one string fields regardless of the question of gauge fixing.

¹⁴ The BRST charge Q is composed of these operators, so the quadratic term does not mix universal and non-universal states. For the cubic term we note that it is defined as a correlator of the three states acted upon by conformal maps. Conformal maps can be defined by exponentials of (total) Virasoro generators, which do not change the universality property and the expectation value of two universal states with one non-universal state vanishes.

¹⁵ A natural further restriction is to use only total-Virasoro generators. Acting on $c_1 |0\rangle$ this gives a set of states to which other states in $\mathcal{H}_{univ}^{(1)}$ do couple linearly. Thus, it cannot be used for further restricting the tachyon vacuum. However, the set of total-Virasoro generators acting on the ghost-number zero vacuum $|0\rangle$ forms an important subalgebra of the star product, namely that of surface states. This subalgebra plays an important role in string field theory.

3.1.1 *Sen's first conjecture*

The restrictions on the solutions and the automatization of the evaluation enabled getting to high level in the level truncation scheme, which resulted in extremely impressive results. The precision in the evaluation of the D-brane tension got far beyond the initial two percents of [131] and even beyond the 0.1% of [132]. On the other hand, it seemed that these accurate calculations introduced an overshooting of the D-brane tension [133] when the level is raised above $l = 14$. Taylor addressed this problem [134], by evaluating the tachyon potential using the methods he developed earlier in [55], as a function of the level. A technical problem in this approach is that the radius of convergence of the terms in the expansion is finite, due to a pole at a negative value of the tachyon field. This pole is related to the boundary of validity of the Siegel gauge mentioned above and is irrelevant to the problem at hand, since the tachyon vacuum is located at a positive value of the tachyon field. In order to overcome this technical problem, the Padé-approximation was used. Then, the tension as a function of the level was numerically fit to a polynomial in l^{-1} . The results suggested that the overshooting problem was merely an artifact. The (common) expectation that the approach to the correct tension would be monotonic with respect to the level was too naive. In fact, the form of the approach graph is gauge dependent and a non-monotonic approach is quite generic. In the Siegel gauge it was found that after the solution overshoots, it starts to return to the correct value around $l \approx 26$. Using a fit with the variable l^{-1} , Gaiotto and Rastelli [130] managed to show that their results also imply that around $l \approx 28$ the tension returns towards the correct value. Their final result as the level $l \rightarrow \infty$ gave the desired result with an impressive accuracy of $3 \cdot 10^{-3}\%$. The analysis was based on level truncation up to $l = 18$ with most of the restrictions described above taken into account. This resulted in more than 2000 fields to consider and over 10^{10} interaction terms. If it was not for the restriction of the coefficient fields, the amount of terms would have been larger by quite a few orders of magnitude and the evaluation, even with the strongest computers, would have been impossible to perform.

3.1.2 *Sen's third conjecture*

Sen's second conjecture states that the cohomology, at least at ghost number one, of the kinetic operator around the non-perturbative vacuum should vanish. As mentioned above, a hint in this direction appeared already in [54], although it was not understood at the time. A preliminary examination of the kinetic term following Sen's work was performed in [135]. The analysis there indicated that the kinetic term seems to vanish, at least for a few low-level scalars. A more systematic study was later performed by Ellwood and Taylor [136]. They studied the exactness of the kinetic operator at ghost number one in level truncation up to level six by calculating the projection to the

space of exact states of all scalar closed states. There are several problems with such a numerical study. First, when level truncated, the kinetic operator does not square to zero. Still, the nilpotence of \mathcal{Q} is improved as the level is increased and so this should not be a problem of principle. Second, in order to calculate the projection, an inner product in the space of string fields should be introduced and there is no canonical choice for such an inner product. This problem was handled by considering several inner products and showing that the results do not have a strong dependence on the choice. Another issue is that, at a given level l , one cannot trust the analysis regarding states whose mass squared is much higher than l . The authors considered only the states below a cut-off, $m^2 < l - 1$. It was found out that at level 6 the projection to the exact space of an arbitrary closed state below the cutoff was of length of more than 99% of the state's norm squared, regardless of the inner product chosen. For a generic (that is not closed) state, the projection gave only about 35% of the original norm squared.

A more elegant approach for investigating the triviality of the spectrum was given in [137]. While this paper also used level truncation, its main tool is general and was used also for proving Sen's conjecture using Schnabl's solution, as we describe in section 5.2. The idea is to find a ghost number -1 state A obeying¹⁶

$$\mathcal{Q}A = |1\rangle, \quad (3.7)$$

where $|1\rangle$ is the identity string field obeying¹⁷

$$|1\rangle \star \Psi = \Psi \star |1\rangle = \Psi \quad \forall \Psi, \quad (3.8)$$

and \mathcal{Q} is the kinetic operator around the tachyon vacuum Ψ_0 as defined in (3.3). The existence of A is equivalent to a strong version of Sen's conjecture. According to Sen, the cohomology of \mathcal{Q} should vanish at ghost number one, while the existence of A implies that it is zero for all ghost numbers¹⁸. The proof that the existence of A implies the vanishing of the cohomology is straightforward. Suppose that Ψ is closed, i.e., $\mathcal{Q}\Psi = 0$. Then,

$$\mathcal{Q}(A \star \Psi) = (\mathcal{Q}A) \star \Psi - A \star (\mathcal{Q}\Psi) = |1\rangle \star \Psi - 0 = \Psi, \quad (3.9)$$

and it follows that Ψ is exact. Thus, the cohomology is empty. Suppose now

¹⁶ An operator (in an operator algebra with derivation and identity) with such a property is called a homotopy operator.

¹⁷ Recall also the problems with the identity based solutions mentioned in the introduction. The subtleties mentioned there may lead one to suspect that results obtained using this state are somewhat formal and should be further examined. The authors of [137] address these issues and suggest that the level truncated identity is an "approximate identity" and that this should be enough for their purposes.

¹⁸ In [138] it was claimed that the other cohomology groups are non-empty. This result contradicts the result of [137]. We return to this point in 5.2.

that the cohomology is empty. The identity string field is closed, since

$$\mathcal{Q}|1\rangle = Q|1\rangle + \Psi \star |1\rangle - |1\rangle \star \Psi = 0, \quad (3.10)$$

where the first term vanishes trivially (2.31), while the two other terms cancel each other. Triviality of the cohomology implies that the identity is also exact, that is, a state A obeying (3.7) exists.

Now we have to identify the state A . Since it depends on the solution Ψ_0 , which was only known in level truncation, the state A was also found using level truncation. First, a measure of being close to having the desired property is introduced in the space of string fields at a given truncation level,

$$\epsilon = \frac{|\mathcal{Q}_l A_l - |1\rangle_l|}{||1\rangle_l|}. \quad (3.11)$$

Here A_l is an arbitrary state at level l , the norm is arbitrary, but following [136] it is assumed that different norms will give similar results. Also, while the norm of the state $|1\rangle$ diverges for many natural norms, the norm of $|1\rangle_l$ is finite, since the Hilbert space at any finite l is finite dimensional. Then, the minimum of ϵ is found at the level- l space. With this procedure it seems that $\epsilon \rightarrow 0$ as the level is increased (it gets to about 2% at $l = 9$). However, an examination of A_l reveals that it does not converge.

The resolution of the above problem comes from noting that A is defined only up to a gauge transformation, since the transformation

$$A \rightarrow A + \mathcal{Q}B, \quad (3.12)$$

leaves (3.7) intact. This symmetry is broken by level truncation and the gauge orbit is replaced by many isolated local minima. For every level, A_l lands arbitrary at different isolated minima. This is the reason for not having a well defined limit. Repeating the analysis for a Siegel-gauge-fixed A resulted in $\epsilon \rightarrow 0$ (about 3% at $l = 9$) as well as convergence of A_l to a well defined limit.

3.1.3 Sen's second conjecture

The examination of the conjecture concerning lump solutions is more involved than the other two. The construction of lumps involves the evaluation of non-universal terms¹⁹ also in the cubic part of the potential. This introduces not only space-time dependence but also non-locality, due to the exponentiation of momenta in the interaction term. Level truncation in such a case may become problematic, since truncation now involves also truncating the exponential of momenta to a finite order in its Taylor expansion and the infinite derivative

¹⁹ Lumps are manifestly background dependent.

theory may differ from the limit of a finite derivative theory when the number of derivatives approaches infinity [111]. Still, the study of lump solutions using level truncation produced appealing evidence in favour of Sen's conjectures [139,140,141], although the agreement was much better for lumps of a small co-dimension than for lumps with a large co-dimension. It was claimed that the exact result is the same, only the approach as a function of the level is slower for the large co-dimension lumps.

The simplest case is the truncation of the action to level zero, i.e., keeping only the tachyon field

$$\Psi = T(x)c_1 |0\rangle , \quad (3.13)$$

as we did for the non-interacting theory (in k -space (2.8)). The truncated string field action (2.10) reads,

$$S = - \int d^{26}x \left(\frac{1}{2} \partial_\mu T(x) \partial^\mu T(x) - \frac{1}{2} T(x)^2 + \frac{1}{3} K^3 \tilde{T}(x)^3 \right), \quad (3.14)$$

where we defined,

$$K = \frac{3\sqrt{3}}{4}, \quad (3.15)$$

$$\tilde{T}(x) = e^{\log K \partial_\mu \partial^\mu} T(x). \quad (3.16)$$

Since it is complicated to evaluate the derivatives in the definition of $\tilde{T}(x)$ exactly, one is led to expanding the exponent in (3.16). Level truncation now turns into a double expansion in the parameters $\log K$ and l .

When the directions in which the lump is localized are compactified, the double expansion becomes a single expansion again. This happens because the compactified momenta are quantized. The oscillators contribute to the level as before, while the momenta contribute to the level a factor of $\alpha' k^2$. From the uncompactified point of view such solutions represent an infinite array of equally spaced lumps. Another drawback of this approach is that the radius is fixed and the analysis should be repeated for different radii. On the other hand the existence of a single expansion parameter simplifies the analysis. The authors of [141] considered a co-dimension one lump for simplicity (and because the approach to the correct result was already known to be the fastest in this case). A possible subtlety of their approach is the appearance of null-states when the radius is rational in α' units. The authors avoided this complication by using only non-rational values for the compactification radius. Their result for the lump tension agreed with expectation within less than a percent at level 3. They also found a lump profile that does not change much with changes of the radius. Another check for the identification of their solution as a lump was the numerical study of the cohomology around the solution. The consistency checks of [127] were performed for these lump solutions in [142] with reasonably good results.

3.1.4 Sen's conjectures in the supersymmetric case

Level truncation was also used in the supersymmetric case²⁰. The standard open superstring theory does not include a tachyon. However, one can consider the theory that lives on a non-BPS D-brane or a D- \bar{D} pair with the appropriate conjectures as stated above.

The study of these conjectures using superstring field theory was initiated by Berkovits [143]. In this paper he used his version of superstring field theory [144], which he generalized in order to include also the GSO(-) sector living on the non-BPS D-brane, which supports a tachyon. The action was truncated so as to include the tachyon field only. Despite the non-polynomiality of the action there are only two terms that contribute within this truncation. The resulting tachyon potential is quartic and its two symmetrically located local minima correspond to about 60% of the predicted value. The results of this paper were extended in [145], where it was shown how to work with non-BPS D-branes as well as with the D- \bar{D} system. It was also shown that expanding the action to an arbitrary level gives rise to a finite number of terms, despite the non-polynomiality of the action. The minimum of the tachyon potential at level $l = 3/2$ was found to give about 85% of the brane tension. Adding one more level brought the number to about 90% [146,147].

The tachyon potential in Witten's superstring field theory [95] was studied in [148]. It was found that at least at the few lowest levels the potential is unbounded from both sides and develops singularities before any local minimum is obtained. Since there are also other problems with this theory as described in section 8.2, the study of the other superstring field theories seems to be more promising.

In the modified version of Witten's theory [149,150,151] the tachyon potential does not suffer from the above problems, at least when the non-chiral choice of Y_{-2} is being made [152]. In this paper truncation to level 1/2 was shown to reproduce already 97.5% of the tension. Truncation to level 2 gives, in the Siegel gauge, the remarkable result of 99.96% of the tension [153]. This result is, however, gauge dependent and other possible gauge choices give less impressive results [152,153].

Lump solutions of superstring field theory were constructed in [145,154]. As in the bosonic case, one gets for these lumps good agreement with the conjectured result. We are not aware of a work addressing Sen's third conjecture, regarding cohomology, in the supersymmetric case.

²⁰ For conventions and definitions of the theories involved see sections 8.1, 8.2.

4 Algebraic primer

One of the main hurdles in studying string field theory used to be the technical difficulty of working with the string field algebra. This algebra is greatly simplified by working in an appropriate coordinate system and considering the relevant basis of states.




Fig. 3. Different coordinate systems. In each coordinate system the Right Left and Middle of the string are marked. The last coordinate system is a cylinder with circumference π (the two marked lines should be identified) and the mid-point is at $i\infty$. The grayed out region is the coordinate patch where no operator insertions are allowed.

Figure 3 shows three possible coordinate systems²¹. The (τ, σ) system gives an intuitive view of a string, where $\sigma \in [0, \pi]$ is its width and $\tau \in (-\infty, \infty)$ represents the world-sheet time. The upper half plane, related to the previous one by $\xi = -\exp(\tau - i\sigma)$, is the canonical coordinate system for calculating CFT correlators. The semi-infinite cylinder C_π defined by $z = \arctan(\xi)$ turns out to be the most convenient coordinate system for string field theory, as the star-product is greatly simplified there.

We also need to transform the operators that we work with to the new coordinate system

$$\tilde{\mathcal{O}}(z) = \tan \circ \mathcal{O}(\xi). \quad (4.1)$$

For primary operators such conformal transformations have the simple form

$$\tilde{\mathcal{O}}^h(z) = f \circ \mathcal{O}^h(\xi) = f'(z)^h \mathcal{O}^h(f(z)), \quad (4.2)$$

where in our case $f(z) = \tan(z) = \xi$. We are also interested in the transformation of the mode expansion coefficients

$$\mathcal{O}^h(\xi) = \sum \mathcal{O}_n^h \xi^{-n-h}, \quad (4.3)$$

²¹ Note that in the literature one finds that left-right is defined either with respect to infinity as was done by Schnabl [2] or with respect to the origin as we do here, following Okawa [155].

which are not primary operators. Still, computing their transformation is straightforward

$$\begin{aligned}\tilde{\mathcal{O}}_n^h &= f \circ \mathcal{O}_n^h = \oint \frac{dz}{2\pi i} z^{n+h-1} \tilde{\mathcal{O}}^h(z) = \oint \frac{dz}{2\pi i} z^{n+h-1} f'(z)^h \mathcal{O}^h(f(z)) \\ &= \sum_{m=-\infty}^{\infty} \mathcal{O}_m^h \oint \frac{dz}{2\pi i} z^{n+h-1} f'(z)^h f(z)^{-m-h}.\end{aligned}\quad (4.4)$$

It is also useful to make a change of variables inside the integral

$$\tilde{\mathcal{O}}_n^h = \sum_{m=-\infty}^{\infty} \mathcal{O}_m^h \oint \frac{d\xi}{2\pi i} (f^{-1}(\xi))^{n+h-1} (f'(z(\xi)))^{h-1} \xi^{-m-h}.\quad (4.5)$$

Usually one looks at transformations that are regular at the origin, then using SL(2) transformations one can get to the canonical form

$$f(0) = 0, \quad f'(0) = 1.\quad (4.6)$$

Also, in this case the transformation of the modes simplifies a bit since the sum in (4.5) starts getting contributions only at $m = n$.

We use the above relations to calculate some modes of the energy momentum tensor

$$\mathcal{L}_{-1} = \sum L_m \oint \frac{d\xi}{2\pi i} (1 + \xi^2) \xi^{-m-2} = L_{-1} + L_1\quad (4.7)$$

$$\mathcal{L}_0 = \sum L_m \oint \frac{d\xi}{2\pi i} \arctan(\xi) (1 + \xi^2) \xi^{-m-2} = L_0 - 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{4n^2 - 1} L_{2n}.\quad (4.8)$$

These transformed Virasoro generators obey the usual Virasoro algebra

$$[\mathcal{L}_n, \mathcal{L}_m] = (n - m) \mathcal{L}_{n+m}.\quad (4.9)$$

This is so because the Virasoro algebra comes out from the OPE relations near the origin where $\tan(\xi) \sim \xi$. In general all transformations of the form (4.6) keep the algebra of local operators intact.

BPZ conjugation on the other hand is not of the form (4.6). BPZ conjugation amounts to transforming the world-sheet $(\tau, \sigma) \rightarrow (-\tau, -\sigma)$. Therefore, it can be seen as mapping an incoming string into an outgoing string. On the upper half plane the BPZ transformation is

$$I(\xi) = -\frac{1}{\xi}.\quad (4.10)$$

Using (4.5) we get the transformation rule

$$\mathcal{O}_n^{h,b} \equiv I \circ \mathcal{O}_n^h = \sum_m \mathcal{O}_m^h \oint \frac{d\xi}{2\pi i} (-1)^{h+n} \xi^{-n-m-1} = (-1)^{h+n} \mathcal{O}_{-n}^h.\quad (4.11)$$

Here, an extra minus sign came from the change of orientation of the integration contour from a contour around $\xi = \infty$ to a contour around $\xi = 0$. For the energy momentum tensor its reality implies that the hermitian conjugation of its modes obeys

$$L_n^\dagger = L_{-n}. \quad (4.12)$$

From (4.11) and (4.12) we read the relation between BPZ and hermitian conjugations for the Virasoro modes,

$$L_n^\dagger = (-1)^n L_n^b. \quad (4.13)$$

On the cylinder the BPZ transformation becomes

$$I(z) = \arctan\left(-\frac{1}{\tan z}\right) = z + \frac{\pi}{2}. \quad (4.14)$$

The simple form of the BPZ transformation on the cylinder does not translate to a simple relation between modes and their BPZ conjugates. The problem with applying (4.5) in this case is that to deform the contour from $z = \pi/2$ to $z = 0$ we need to go through the cut. Hence, the map between modes and their BPZ conjugation on the cylinder is generally quite cumbersome.

There are two equivalent ways to expand the BPZ conjugate operators. One is to apply the BPZ conjugation on the UHP operators in (4.5). The other option is to use the composed transformation $f(I(\xi))$ directly. Specifically, for the Virasoro generators on the cylinder we get

$$\mathcal{L}_n^b = - \oint \frac{d\xi}{2\pi i} (1 + \xi^2) \left(-\operatorname{arccot}(\xi)\right)^{n+1} T(\xi). \quad (4.15)$$

Here, an extra minus sign comes from orientation change as in (4.11). Hermitian conjugate Virasoro operators transform as,

$$\mathcal{L}_n^\dagger = \oint \frac{d\xi}{2\pi i} (1 + \xi^2) \left(\operatorname{arccot}(\xi)\right)^{n+1} T(\xi). \quad (4.16)$$

The last two expressions are in fact the same. This can be seen also directly from the expansion (4.8).

It is easy to see that

$$\mathcal{L}_{-1}^\dagger = \mathcal{L}_{-1} = K_1, \quad (4.17)$$

where K_1 is one of the mid-point preserving reparametrization operators [95],

$$K_n = L_n - (-1)^n L_{-n}. \quad (4.18)$$

This operator plays a dominant role in string field theory. Another important combination comes from the commutation relation

$$[\mathcal{L}_0, \mathcal{L}_0^\dagger] = \oint \frac{d\xi}{2\pi i} (1 + \xi^2) (\arctan \xi + \operatorname{arccot} \xi) T(\xi) = \mathcal{L}_0 + \mathcal{L}_0^\dagger. \quad (4.19)$$

The integrand can be written using a step function,

$$\arctan \xi + \operatorname{arccot} \xi = \frac{\pi}{2} \epsilon(\Re(\xi)) = \frac{\pi}{2} \begin{cases} -1 & \Re(\xi) < 0 \\ 1 & \Re(\xi) > 0 \end{cases}. \quad (4.20)$$

The calculation of the integral is a bit subtle, since $\operatorname{arccot}(\xi)$ has a branch cut in the range $\xi = (-i, i)$, while $\arctan(\xi)$ has a branch cut on the rest of the imaginary axis. Therefore, the contour can cross the imaginary axis only at the points $-i, i$. This is not evident when writing the integrand as a step function. The step function does suggest that we should split the contour of integration to its left (C_L) and right (C_R) parts. For each part we have the integrand of K_1 , it is therefore natural to name the two parts K_1^L and K_1^R . This gives the relation

$$\hat{\mathcal{L}}_0 \equiv \mathcal{L}_0 + \mathcal{L}_0^\dagger = \frac{\pi}{2} (-K_1^L + K_1^R). \quad (4.21)$$

$\hat{\mathcal{L}}_0$ has an interesting property of increasing the \mathcal{L}_0 -level of states. This can be seen from the commutation relation

$$[\mathcal{L}_0, \hat{\mathcal{L}}_0] = \hat{\mathcal{L}}_0. \quad (4.22)$$

For the standard Virasoro operators, only L_{-1} has such a behaviour with respect to L_0 . For \mathcal{L}_0 we have two such states, $\hat{\mathcal{L}}_0$ and \mathcal{L}_{-1} (or linear combination of these like K_1^L, K_1^R).

Notice that the definition of K_1^L and K_1^R is restricted by the cut structure. This is illustrated in figure 4. These operators have “semi-derivation” properties with respect to the star-product

$$K_1^L(\Psi_1 \Psi_2) = (K_1^L \Psi_1) \Psi_2, \quad (4.23)$$

$$K_1^R(\Psi_1 \Psi_2) = \Psi_1 (K_1^R \Psi_2). \quad (4.24)$$

Similar relations hold for the $b(z)$ ghost

$$\hat{\mathcal{B}}_0 \equiv \mathcal{B}_0 + \mathcal{B}_0^\dagger = \frac{\pi}{2} (-B_1^L + B_1^R), \quad (4.25)$$

$$B_1^L(\Psi_1 \Psi_2) = (B_1^L \Psi_1) \Psi_2, \quad (4.26)$$

$$B_1^R(\Psi_1 \Psi_2) = (-1)^{\Psi_1} \Psi_1 (B_1^R \Psi_2). \quad (4.27)$$

The commutation relation (4.19) defines an $SL(2)$ algebra. This relation only holds if the underlying CFT has vanishing central charge. Turning on a central charge in the CFT will add an infinite central charge to (4.19). A way to regularize this central charge was given in [156].




Fig. 4. Possible choices for the contour in (4.19). The natural choice in the upper half plane is the unit circle, which transforms to two infinite straight lines at $\pm\pi/4$ in the cylinder coordinates. We can squeeze the circle in the upper half plane in a way that keeps the lines in the cylinder straight. More generally, we can deform the contour as we like, as long as we do not move the points $\pm i$, which are fixed by the existence of cuts in the rest of the imaginary axis. In the cylinder coordinates, this translates to arbitrary curves connecting $\pm i\infty$.

The advantage of the cylinder coordinates is that it allows for a simple geometrical interpretation of the star product. The most basic example, is the multiplication of two vacuum states. In the upper half plane, one has to cut out the coordinate patch from the two states (see figure 3). Then, the left edge of one of the remaining half unit circles needs to be sewed to the right edge of the other. Finally, the coordinate patch will have to be reintroduced, leading to a non-trivial world-sheet. In the cylinder coordinate things are much simpler. Cutting the coordinate patch leaves us with two half-infinite strips (see again figure 3). Sewing the two half-infinite strips and the coordinate patch, which is also a half-infinite strip gives a cylinder with circumference $\frac{3\pi}{2}$.

Regardless of the choice of a coordinate system, the result of multiplying two vacuum states is the wedge state $|3\rangle$. Wedge states [35,10,24,96] are a set of surface states satisfying the abelian algebra

$$|r\rangle \star |s\rangle = |r + s - 1\rangle \quad r, s \geq 1, \quad (4.28)$$

where the wedge state $|2\rangle$ is the vacuum state $|0\rangle$ ²². The state $|3\rangle$ is the result of multiplying two vacuum states, as discussed in the previous paragraph, and $|1\rangle$ is the identity string field. In the cylinder coordinates²³, a wedge state $|r\rangle$ corresponds to a cylinder with circumference $\frac{r\pi}{2}$, with the coordinate patch

²² The equality $|0\rangle = |2\rangle$ may be a source for confusion. This should not be the case, since the wedge state $|r=0\rangle$ does not exist.

²³ Strictly speaking this is a new coordinate system. A wedge state can be defined in the standard cylinder coordinates using an operator insertion, which corresponds to rescaling (see below).

occupying a strip of width $\frac{\pi}{2}$. A different convention used for the wedge states is

$$W_\alpha \equiv |\alpha + 1\rangle \quad \Rightarrow \quad W_\alpha W_\beta = W_{\alpha+\beta}. \quad (4.29)$$

We shall be using both conventions interchangeably.

While wedge states form a very simple subalgebra, they are of limited use by themselves. Specifically, they cannot describe the solution to the equation of motion since they have ghost number zero. To overcome this limitation, we consider wedge states with insertions. The product of two such states takes in the cylinder coordinate the form,

$$\mathcal{O}(z_1^r) |r\rangle \star \mathcal{O}(z_1^s) |s\rangle = \mathcal{O}\left(z_1^r + \frac{\pi(s-1)}{4}\right) \mathcal{O}\left(z_1^s - \frac{\pi(r-1)}{4}\right) |r+s-1\rangle. \quad (4.30)$$

This wedge state multiplication is illustrated in figure 5.




Fig. 5. Multiplication of two wedge states results in a wedge state with the combined width of the two original states. Operator insertions are mapped accordingly. One can have many operator insertions and these can be anywhere on the coordinate patch.

Using wedge states with insertions, we can describe any field we want. Actually, since we are allowing an arbitrary insertion, we have more than one description per field. The simplest example to this redundancy is given by representing the wedge states themselves as²⁴

$$|n\rangle = U_n^\dagger |0\rangle = \left(\frac{2}{n}\right)^{\mathcal{L}_0^\dagger} |0\rangle, \quad (4.31)$$

where \mathcal{L}_0^\dagger is the hermitian conjugate of \mathcal{L}_0 as defined in (4.16). This expression can be understood from looking at the inner product of the wedge state with some test state

$$\langle 0 | \mathcal{O}_h(z) |n\rangle = \langle 0 | \mathcal{O}_h(z) U_n^\dagger |0\rangle = \left(\frac{2}{n}\right)^h \langle 0 | \mathcal{O}_h\left(\frac{2z}{n}\right) |0\rangle, \quad (4.32)$$

²⁴ This representation is in accord with the fact that they are surface states, in light of (2.29).

where we used the fact that \mathcal{L}_0^\dagger is the dilatation operator for bra states and assumed in the last step that \mathcal{O}_h is a primary operator of conformal weight h . This algebraic result matches the geometrical picture of starting from an operator inserted on a cylinder of width $\frac{n\pi}{2}$ and scaling it to the canonical cylinder of width π . Equivalently, wedges with insertions are written as

$$\mathcal{O}(z_n) |n\rangle = U_n^\dagger U_n \mathcal{O}(z_n) |0\rangle. \quad (4.33)$$

We can use the fact that \mathcal{L}_0 and \mathcal{L}_0^\dagger close the $SL(2)$ algebra (4.19) to get the following identities using basic group theory technics (such as using an explicit 2-dimensional representation)

$$U_r \mathcal{L}_0^\dagger U_r^{-1} = \frac{2-r}{r} \mathcal{L}_0 + \frac{2}{r} \mathcal{L}_0^\dagger, \quad (4.34)$$

$$U_r^{-1} \mathcal{L}_0^\dagger U_r = \frac{r-2}{2} \mathcal{L}_0 + \frac{r}{2} \mathcal{L}_0^\dagger, \quad (4.35)$$

$$U_r^\dagger \mathcal{L}_0 U_r^{\dagger-1} = \frac{r}{2} \mathcal{L}_0 + \frac{r-2}{r} \mathcal{L}_0^\dagger, \quad (4.36)$$

$$U_r^{\dagger-1} \mathcal{L}_0 U_r^\dagger = \frac{2}{r} \mathcal{L}_0 + \frac{2-r}{2} \mathcal{L}_0^\dagger, \quad (4.37)$$

and also

$$U_r U_s = U_{\frac{rs}{2}}, \quad (4.38)$$

$$U_r U_s^\dagger = U_{2+\frac{2}{r}(s-2)}^\dagger U_{2+\frac{2}{s}(r-2)}, \quad (4.39)$$

$$e^{\beta(\mathcal{L}_0 + \mathcal{L}_0^\dagger)} = U_{2-2\beta}^\dagger U_{2-2\beta}. \quad (4.40)$$

The building blocks that we will be working with are wedge states with local operator insertions as in (4.30) and line integrals of the $T(z)$ and $b(z)$ fields,

$$K \equiv K_1^L = - \int_{-i\infty}^{i\infty} T(z) \frac{dz}{2\pi i}, \quad B \equiv B_1^L = - \int_{-i\infty}^{i\infty} b(z) \frac{dz}{2\pi i}. \quad (4.41)$$

These line integral can be freely moved as long as the end-points are fixed and no other operator insertions are crossed.

The K operator can be used to generate wedge states from the identity state,

$$e^{\frac{\pi}{2}(n-1)K} |1\rangle = |n\rangle. \quad (4.42)$$

From this relation we learn that

$$K W_n = \frac{\pi}{2} \partial_n W_n. \quad (4.43)$$

This implies that K acts as a derivation with respect to strip length, since the factor of $\frac{\pi}{2}$ is the ratio between the length of the wedge and its number.

5 Schnabl's solution

It took almost twenty years from the time Witten wrote down the equation of motion of string field theory [1] until Schnabl found its analytical solution [2]. Prior candidate solutions were either numerical, singular in some sense or too abstract. The litmus test for a solution is its ability to reproduce the D -brane tension according to Sen's conjecture. There were some suggestion for analytical solutions before Schnabl, but none of them could pass this test. Some candidate solutions turned out to have zero action, and therefore understood to represent pure gauge solution.

Still, the simplest form to write down Schnabl's solution is as a pure gauge solution [155]. We will see that the gauge transformation is singular, and that the solution is physical. From the form of the finite gauge transformation (2.17), we can see that a pure gauge solution takes the form,

$$\Psi = e^{-\Lambda} Q e^{\Lambda}. \quad (5.1)$$

Reparametrization of Λ enables us to rewrite the above as

$$\Psi = \Gamma^{-1}(\Lambda) Q \Gamma(\Lambda), \quad (5.2)$$

with some different Λ and where $\Gamma(\Lambda)$ is some function of the form

$$\Gamma(\Lambda) = 1 + \Lambda + \mathcal{O}(\Lambda^2). \quad (5.3)$$

We refer to the choice of Γ as a scheme choice.

Schnabl's solution is most easily represented in the left scheme²⁵,

$$\Gamma(\Lambda) = 1 + \lambda \Lambda. \quad (5.4)$$

Here, λ is a parameter. It turns out that the solution is a pure gauge one for $|\lambda| < 1$, while for $\lambda = 1$, it is the desired tachyon vacuum. The solution does not converge for other (real) values of λ [2]²⁶.

The gauge string field is²⁷ [155],

$$\Lambda = Bc(0) |0\rangle, \quad (5.5)$$

²⁵ In this scheme $Q\Lambda$ is always found to the left of Λ^n , as is explicit in (5.8).

²⁶ This picture of a family of gauge solutions for $\lambda < 1$ that turn into a physical solution for $\lambda = 1$ and are not defined elsewhere, is supported by a numerical study, in which it was found that the energy density approaches a step function as a function of λ as the truncation-level is increased [157].

²⁷ An interesting property of Λ is that its \mathcal{L}_0 eigenvalue vanishes.

and the solution itself takes then the form,

$$\Psi = (1 - \lambda\Lambda)Q \frac{1}{1 - \lambda\Lambda} = Q\Lambda \frac{\lambda}{1 - \lambda\Lambda}, \quad (5.6)$$

where in the last equality we integrated by parts²⁸. It is useful to expand Ψ in λ ,

$$\Psi = \sum_{n=1}^{\infty} \lambda^n \Psi_n. \quad (5.7)$$

This gives,

$$\Psi_n = (Q\Lambda)\Lambda^{n-1}. \quad (5.8)$$

Written as in (5.6), Ψ naturally satisfies the equation of motion. Unfortunately, it seems that the action vanishes. This can be seen from

$$\partial_\lambda S(\Psi) = \langle \partial_\lambda \Psi (Q\Psi + \Psi^2) \rangle, \quad (5.9)$$

which vanishes upon using the equation of motion. Since the action is clearly zero for $\lambda = 0$, it remains zero for any value of λ . This is not surprising, since the above is equivalent to the variational method by which one gets the equation of motion in the first place.

It turns out that the argument for the vanishing of the action fails, since the action has a finite radius of convergence in λ . One clue for this singularity comes from looking at the gauge condition. Schnabl's solution does not obey the Siegel gauge (2.22), instead it obeys

$$\mathcal{B}_0\Psi = 0. \quad (5.10)$$

This is the Schnabl gauge condition. Despite the fact that Ψ obeys the gauge condition, it is (by construction) a pure gauge state. This shows that Schnabl's gauge does not fix the gauge completely. This is a bit strange, in light of the similarity between the Siegel and the Schnabl gauge choices. It may suggest that the pure gauge states obeying the condition are of a somewhat singular nature. Specifically, Ψ_1 is an exact state obeying the gauge condition,

$$\mathcal{B}_0\Psi_1 = \mathcal{B}_0Q\Lambda = (\mathcal{L}_0 - Q\mathcal{B}_0)\Lambda = 0. \quad (5.11)$$

It is instructive to try to transform Ψ_1 to the Siegel gauge, using the one-parameter family of transformations [158] (see also [60,61]),

$$\mathcal{B}_0^s = U_s \mathcal{B}_0 U_s^{-1}, \quad A_s = U_s A, \quad U_s = s^{-L_0}, \quad (5.12)$$

where $s = 1$ corresponds to Schnabl's gauge and $s = 0$ gives the Siegel gauge. The limit $s \rightarrow 0$ is well defined when U_s acts on physical states, but Λ_s is singular in the limit.

²⁸ Recall that the identity string field is annihilated by Q (2.31).

Next, we want to simplify the expanded expression for Ψ . We start with

$$\Lambda^n = Bc_1 |0\rangle \star Bc_1 |0\rangle \star \cdots \star Bc_1 |0\rangle . \quad (5.13)$$

This can be viewed as a cylinder of width $\frac{(n+1)\pi}{2}$ with alternating B line integrals and c insertions. As mentioned above, the B integral can be freely moved over the cylinder as long as the end points are fixed at $\pm i\infty$ and it does not cross any c operators. We can use the (anti)-commutation relation²⁹

$$[B, c_1] = 1 , \quad (5.14)$$

in order to move the B line integrals across the c insertions. This results in the B hitting another B that annihilates it. Hence, we are left with

$$\Lambda^n = |0\rangle^{n-1} \star Bc |0\rangle = |n\rangle \star Bc |0\rangle . \quad (5.15)$$

Next, we calculate

$$Q\Lambda = (K - BQ)c(0) |0\rangle = (1 - Bc)Kc(0) |0\rangle = (cBKc)(0) |0\rangle , \quad (5.16)$$

where we used

$$c\partial c = cKc , \quad (5.17)$$

which follows from (4.43). We conclude that

$$\Psi_n = (Q\Lambda)\Lambda^{n-1} = c |0\rangle \star BK |n-1\rangle \star c |0\rangle = \frac{d}{dn} \psi_{n-1} , \quad (5.18)$$

where we defined

$$\psi_n \equiv c |0\rangle \star B e^{\frac{\pi}{2}(n-1)K} |1\rangle \star c |0\rangle , \quad (5.19)$$

and used (4.42) to replace K with the derivative. Note, that this expression is somewhat formal for the case $n = 0$ (Ψ_1). Carefully taking the limit, one gets

$$\psi_0 = (cBc)(0) |0\rangle , \quad \psi'_0 = (cBKc)(0) |0\rangle , \quad (5.20)$$

in agreement with (5.16).

Summing (5.7) with $\lambda = 1$, we can write

$$\Psi = c |0\rangle \star \sum_{n=1}^{\infty} \left(BK e^{\frac{\pi}{2}(n-1)K} |1\rangle \star c |0\rangle \right) = c |0\rangle \star \frac{BK}{1 - e^{\frac{\pi}{2}K}} |1\rangle \star c |0\rangle . \quad (5.21)$$

It is also useful to represent the solution in a split-string form [155,159,160]. Examining the strip from left to right, one encounters a c insertion after a length of $\frac{\pi}{4}$, which is half the length of the vacuum strip (local coordinate patch not included), since the c is inserted in the middle on the vacuum state.

²⁹ Here and elsewhere in the paper, the brackets represent the graded commutator, i.e., it is the anti-commutator for two odd objects and the commutator otherwise.




Fig. 6. Schnabl's solution: The left (length $\frac{\pi}{4}$) strip length is represented in the split-string formulation by $W_{\frac{1}{2}}$. To the right of it is the c insertion. The dashed line represents the B, K line integrals as well as the varying strip length insertions. The Ψ_1 part contributes zero length, so the c insertions almost touch, but the B, K in between them make this part also non-trivial. The rest of Ψ contributes larger and larger strip lengths, up to infinity. A (gray) coordinate patch is appended to the resulting surface in the usual way.

Then, there is a varying strip length, from zero to infinity, which is summed over, with the BK line integral insertions. Finally, there is again a c insertion followed by a strip of length $\frac{\pi}{4}$. Thus, we can write

$$\Psi = W_{\frac{1}{2}} c \frac{BK}{1 - W_1} c W_{\frac{1}{2}}, \quad (5.22)$$

where free operator insertions (namely c, B, K) represent these operators acting on the identity string field, i.e., inserted on a zero-sized strip. Now, the only product appearing in the equation is the star product³⁰. The B, K insertions can be freely moved and so the expression containing them in the numerator with the denominator that represent a varying strip length is well-defined. We present Schnabl's solution schematically in figure 6.

5.1 Sen's first conjecture

We already showed in (5.9) that the action for Ψ_λ seems to vanish. Actually we showed that the action vanishes order by order in λ . This suggests that

³⁰ The string field $c|0\rangle$ for example, is written in this notation as $W_{\frac{1}{2}} c W_{\frac{1}{2}}$. In the literature one usually finds the notation $F \equiv W_{\frac{1}{2}}$.

Ψ_λ can be regularized by expanding it up to some finite order N in λ . Such a regularization will not affect terms up to λ^N in the action. Yet, at higher orders some of the terms needed for (5.9) to hold will be missing, allowing for a non-vanishing action.

Concentrating on the kinetic term in the action, we have of the order of N^2 terms with powers of λ larger than N . Therefore, for the proposed regularization to hold, it should be the case that

$$\langle \psi'_m Q \psi'_n \rangle \stackrel{?}{=} \mathcal{O}(N^{-3}) \quad m+n > N. \quad (5.23)$$

Using CFT methods we can calculate the correlator

$$\begin{aligned} \langle \psi_m Q \psi_n \rangle = \frac{1}{\pi^3} & \left(\left(\cos \left(\frac{(m-n)\pi}{m+n+2} \right) + 1 \right) \left((m+n+2) \sin \left(\frac{2\pi}{m+n+2} \right) - \pi \right) + \right. \\ & \sin^2 \left(\frac{\pi}{m+n+2} \right) \left((m+n+2)(m-n) \sin \left(\frac{(m-n)\pi}{m+n+2} \right) \right. \\ & \left. \left. - 2\pi(m+n+1) + 2\pi mn \cos \left(\frac{(m-n)\pi}{m+n+2} \right) \right) \right). \quad (5.24) \end{aligned}$$

In the limit where both n and m are large, this correlator becomes a constant. The terms in (5.23) behave in this limit (due to the two derivatives) like N^{-2} , so (5.23) does not hold and the summation produces a finite term. This demonstrates that, at least for calculating the action, cutting the series at finite N is not a consistent regularization.

We can fix this regularization by rewriting the sum using the Bernoulli numbers. The Bernoulli numbers are defined by the generating function

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} \frac{B_n}{n!} x^n. \quad (5.25)$$

The middle term in the solution (5.21) (without the B insertion) can thus be written as,

$$\sum_{n=1}^{\infty} K e^{\frac{\pi}{2}(n-1)K} = \frac{K}{1 - e^{\frac{\pi}{2}K}} = \sum_{n=1}^N K e^{\frac{\pi}{2}(n-1)K} - \left(\sum_{k=0}^{\infty} \frac{B_k}{k!} \left(\frac{\pi}{2} K \right)^k \right) e^{\frac{\pi N}{2}K}, \quad (5.26)$$

where N is arbitrary. If we cut-off the Bernoulli series at M [161], we get a regularized form for the solution (5.21),

$$\Psi^{(N,M)} = \sum_{n=0}^{N-1} \psi'_n - \left(\sum_{k=0}^M \frac{B_k}{k!} K^k \right) \psi_N, \quad (5.27)$$

where the K^k operator is inserted between the two c ghosts in ψ_{N+1} . The large

N behaviour of the kinetic term for this solution is

$$\langle \Psi^{(N)} Q \Psi^{(N)} \rangle = \langle \Psi Q \Psi \rangle + \mathcal{O}(N^{-M-1}), \quad (5.28)$$

where $\langle \Psi Q \Psi \rangle$ is the result without regularization artifacts. Similarly, for the cubic term one gets

$$\langle \Psi^{(N)} \Psi^{(N)} \Psi^{(N)} \rangle = \langle \Psi \Psi \Psi \rangle + \mathcal{O}(N^{-M-1}). \quad (5.29)$$

Therefore, it is enough to set $M = 0$, meaning that one needs only the first term in the Bernoulli series, $B_0 = 1$ ³¹. Hence, a properly regularized form of the solution is³²,

$$\Psi = \lim_{N \rightarrow \infty} \left(\sum_{n=0}^{N-1} \psi'_n - \psi_N \right). \quad (5.30)$$

To calculate the kinetic term, we replace n with xN and m with yN . For large N we get

$$\langle \psi'_x Q \psi'_y \rangle = \frac{8\pi xy}{(x+y)^6} \left(\pi xy \cos\left(\frac{\pi(x-y)}{x+y}\right) + (y^2 - x^2) \sin\left(\frac{\pi(x-y)}{x+y}\right) \right), \quad (5.31)$$

where the derivatives now are with respect to x, y and we write ψ_x instead of ψ_{Nx} . Naively, one could think that it is possible to replace the summation with an integral. Substituting (5.30) gives,

$$\begin{aligned} \langle \Psi Q \Psi \rangle &= \int_0^1 dx dy \langle \psi'_x Q \psi'_y \rangle - \int_0^1 dx \langle \psi'_x Q \psi_1 \rangle - \int_0^1 dy \langle \psi_1 Q \psi'_y \rangle + \langle \psi_1 Q \psi_1 \rangle \\ &= \langle \psi_0 Q \psi_0 \rangle = 0 \quad (\text{wrong}). \end{aligned} \quad (5.32)$$

The first two equalities are straightforward. The last equality holds, since the relevant primitive of (5.31), i.e., the one that agrees with (5.24) at infinity, is zero on the lines $x = 0$ and $y = 0$. We are getting the wrong results, since when both x and y are small, the approximation (5.31) does not hold and the summation cannot be replaced by an integral. Therefore, we calculate the sum explicitly for the case $x + y < 1$. The sum over any constant $x + y$ corresponds to a sum over terms with a $\lambda^{(x+y)N}$ coefficient. These sums vanish by the virtue of the equation of motion as discussed earlier.

³¹ It may happen that more terms are needed. This is the case for the ‘‘tachyon’’ solution in superstring field theory [161].

³² This is not the only way to regularize the solution. The simplest alternative is level truncation. The Bernoulli term (sometimes called ‘‘the phantom piece’’) has a vanishing inner product with all Fock space states in the limit $N \rightarrow \infty$. Thus, level truncation is a proper regularization of the solution. However, it is usually not adequate for the derivation of analytical results.

Therefore, to calculate the action for the kinetic term, we only need to subtract the integral over $x + y < 1$ in (5.32)

$$\langle \Psi Q \Psi \rangle = - \int_0^1 dx \int_0^{1-x} dy \langle \psi'_x Q \psi'_y \rangle = -\frac{3}{\pi^2}. \quad (5.33)$$

From the equation of motion we know that

$$\langle \Psi Q \Psi \rangle = - \langle \Psi \Psi \Psi \rangle. \quad (5.34)$$

One might want to check explicitly that this equation indeed holds [155,162]. It is a non-trivial check of the regularization. Specifically, without the Bernoulli term, the equation of motion does not hold, when contracted with the solution itself.

The energy density equals minus the action per unit volume.

$$E = -\frac{S}{V_{26}} = \frac{1}{g_o^2 V_{26}} \left(\frac{1}{2} \langle \Psi Q \Psi \rangle + \frac{1}{3} \langle \Psi \Psi \Psi \rangle \right) = \frac{\langle \Psi Q \Psi \rangle}{6g_o^2 V_{26}} = -\frac{1}{2\pi^2 g_o^2}. \quad (5.35)$$

This is exactly the tension of the D25-brane [29]. The volume factor comes from integrating over the zero-modes and was set to unity before. Lower dimensional D-branes do not have a zero-mode for the directions with the Dirichlet boundary condition, giving the correct volume factor for these D-branes.

5.2 Sen's third conjecture

As we described in 3.1.2, the cohomology of the kinetic operator around the tachyon vacuum is empty, provided that a string field A exists obeying (3.7). An exact form of such an operator was found by Ellwood and Schnabl [163] using an analogy to the numerical solution found in the Siegel gauge in [137]. The string field A takes the form,

$$A = -\frac{2}{\pi} B \int_0^1 W_r dr. \quad (5.36)$$

To show that this string field obeys (3.7), we recall (3.3) and evaluate the terms separately. First ³³,

$$QA = -\frac{2}{\pi} [Q, B] \int_0^1 W_r dr = - \int_0^1 \partial_r W_r dr = -W_1 + W_0 = |1\rangle - |0\rangle. \quad (5.37)$$

³³ Conventions here can be confusing, since $W_1 = |0\rangle$, namely the perturbative vacuum, while $W_0 = |1\rangle$, namely the identity string field.

Then, we substitute (5.22) to get,

$$\begin{aligned} \Psi A &= -\frac{2}{\pi} W_{\frac{1}{2}} c \frac{BK}{1 - W_1} cB \int_{1/2}^{3/2} W_r dr = -\frac{2}{\pi} W_{\frac{1}{2}} cBK \int_{1/2}^{\infty} W_r dr = \\ &- W_{\frac{1}{2}} cB \int_{1/2}^{\infty} \partial_r W_r dr = W_{\frac{1}{2}} cB (W_{\frac{1}{2}} - W_{\infty}) = (cB)(0) |0\rangle, \end{aligned} \quad (5.38)$$

where in the first equality we used the wedge state algebra (4.29) and in the second we commuted the ghosts, expanded the denominator and merged it with the integral. Then, we used the fact that K acts as a derivative with respect to wedge length (4.43), performed the integral and neglected the boundary term coming from an infinitely long strip, since its inner product with an arbitrary Fock space goes like N^{-3} . Similarly one gets

$$A\Psi = (Bc)(0) |0\rangle. \quad (5.39)$$

All in all one can write

$$\mathcal{Q}A = QA + \Psi A + A\Psi = |1\rangle - |0\rangle + [B, c](0) |0\rangle = |1\rangle, \quad (5.40)$$

where (5.14) was used. This ends the proof.

As mentioned in 3.1.2, the existence of A proves that the cohomology is empty not only for ghost number one string fields, but for string fields of all ghost numbers. Nonetheless, the opposite was concluded in a numerical study performed in the Siegel gauge [138]. Imbimbo, one of the authors of that work, repeated the numerical analysis, this time in the Schnabl gauge and found again, that while it seems that the cohomology at ghost number one is empty, there seem to be non-empty cohomologies at other ghost numbers [164]³⁴. This result contradicts the derivation presented above and one has to think how can they be reconciled. One option is that the numerical results get modified as higher levels are included and the analytical prove presented here holds. While this is always a possibility with numerical analysis, the results of [164] seem to be quite robust. Another option is that we were too hasty in neglecting the contribution of the terms based on infinitely long strips. This is related to the notorious problem of defining the correct space of string fields. Since it is not clear how should this space be defined³⁵, it may well be the case that either the formal steps in the Ellwood-Schnabl construction or the numerical analysis of Imbimbo, fail somehow in the relevant space. This question certainly deserves a deeper study.

³⁴ Similar conclusions for the ghost number one space were obtained in [165].

³⁵ There is no canonical positive definite norm, for example to use for the definition. We only have some criteria, such as demanding this space to form a star-algebra and existence of physical solutions in it. See also footnote 46.

5.3 Generalizations of the solution

String field theory has a huge amount of gauge symmetry. In particular, there are many solutions equivalent to the one found by Schnabl. One may wonder what is the specific property that makes this solution simpler than the other gauge equivalent ones. Deeper understanding of this point can serve as a key for finding solutions with other physical content.

One of the features that enabled the construction of Schnabl's solution, was the simplification in the form of the equation of motion in the z conformal frame. In [166], Rastelli and Zwiebach tried to pin down the properties of this conformal frame that made the string field equation of motion tractable. Instead of studying the equation of motion (2.23), a simplified ghost number zero toy model was used [129]. The equation of motion addressed was then,

$$(L_0 - 1)\Psi + \Psi^2 = 0. \quad (5.41)$$

The idea then was to consider various conformal frames and in each frame to consider the generalization of (5.41) as the equation of motion, namely

$$(L - 1)\Psi + \Psi^2 = 0. \quad (5.42)$$

Here, L represents the zero mode of the energy momentum tensor in the given conformal frame. Next, they required that L and its hermitian conjugate L^\dagger obey a generalization of (4.19),

$$[L^\dagger, L] = s(L + L^\dagger). \quad (5.43)$$

The parameter s introduced here seems to be spurious, since it is possible to absorb it by a rescaling of L . However, if one insists that the conformal frame is properly normalized, say, as in (4.6), then it becomes meaningful.

It was then suggested that the property that makes the cylinder coordinates adequate is their relation to the sliver projector. Star algebra projectors [7,167], i.e., ghost number zero string fields obeying the equation

$$\Psi^2 = \Psi, \quad (5.44)$$

were used in string field theory in various contexts. As always, one can classify projectors according to their rank. The identity string field is a projector with a maximal rank. Important role was played by rank one projectors, the most familiar ones being the sliver [35,11] and the butterfly [168,169,167].

The association of conformal frames to projectors is via the relation

$$\Psi = \lim_{\gamma \rightarrow \infty} e^{-\gamma L} |0\rangle. \quad (5.45)$$

If the limit exists, it defines a projector. In order to be adequate for defining analytical solutions, it should obey some regularity conditions, which basically imply that a left-right factorization of L is permissible. Projectors obeying these conditions were named “special projectors” and their conformal frames named “special conformal frames”. With these conditions at hand, it was shown that the sliver is the only projector with $s = 1$ in (5.43), while for $s = 2$ one has two possible projectors, namely the butterfly and a new one, named the “moth”. The solutions of (5.42) were found for $s > 1$. These solutions look like generalizations of Schnabl’s (toy) solution, in which the Bernoulli numbers were generalized in some way. Still, Schnabl’s original solution seems to be the simplest one, both algebraically and because it is based on a simple conformal transformation in which the star product is realized simply by gluing (without any other operations).

Another issue left out in [166] is the ghost number one case, as opposed to the toy model. This physical case was studied along similar lines in [170]. It was found that, while special projectors give rise to simple solutions, other projectors can also be used in principle. The map that sends solutions from one conformal frame to the other was constructed. In particular generalizations of the wedge state subalgebra leading to an arbitrary projector was found and it was established that (unlike in the toy model) the formal form of the solution is the same for all conformal frames.

Classification of the resulting solutions led to the conclusion that they all reside in a proper subspace of the universal space, in which they are supposed to be found. This enabled the authors to check whether the numerical solutions in the Siegel gauge can be related to their construction. It turns out that the vacuum solution found in the Siegel gauge does not belong to the restricted space and so cannot be obtained from a projector in the way described in [170]. Thus, while it is still believed that the numerical solution in the Siegel gauge is related to Schnabl’s solution (and its generalizations described here) by gauge transformations, these gauge transformations cannot be obtained using reparametrization.

Another generalization of Schnabl’s solution was considered in [171], where a B-field was included³⁶. It was shown that Schnabl’s solution (in its CFT form) is not modified. This is an expected result due to the universal form of the solution and Sen’s observation regarding the universality of the tachyon vacuum [5].

³⁶ The treatment of a B-field background in string theory in the language of non-commutative field theory was developed in [19]. The implementation of this idea within string field theory was studied in [172,173,174,175,176,177].

6 Marginal deformations

A boundary conformal field theory can be deformed by adding to it a (boundary) term of the form³⁷

$$\delta S_{BCFT} = \lambda \int V dx . \quad (6.1)$$

If V is a weight one vertex operator, then the resulting theory will remain conformal, at least to leading order in λ . If the theory remains conformal also for finite λ , one refers to V as being “exactly marginal”. The new theory can be described as a solution to the string field theory of the original BCFT, with the solution

$$\Psi = \lambda \Psi_1 + \mathcal{O}(\lambda^2), \quad \Psi_1 = cV(0) |0\rangle . \quad (6.2)$$

The first order term Ψ_1 trivially satisfies the linearized equation of motion (2.1)

$$Q\Psi_1 = 0 . \quad (6.3)$$

The higher order terms can be calculated, in principle, provided that the deformation is exactly marginal [179,180,181], by expanding the string field as in (5.7) and solving the recursion relations

$$Q\Psi_n = - \sum_{k=1}^{n-1} \Psi_k \Psi_{n-k} , \quad (6.4)$$

with the initial condition (6.2). Note, however, that while the perturbed BCFT is a unique theory, the solution to the recursion equations is not unique. Far from that, the huge gauge symmetry of string field theory is hidden in them. Thus, for a finite value of λ the relation is that a single BCFT corresponds to a gauge equivalence class of string field theory solutions. Moreover, one can add to Ψ_n in (6.4) any solution of (6.3). This means that an arbitrary marginal deformation can be added to the solution at any order. Adding the same marginal deformation that one starts with, results in a reparametrization of λ , while adding another one results in a solution whose physical content is different for finite values of λ . Thus, one should be cautious when constructing and interpreting these solutions.

The first investigation (using level truncation) of marginal deformations as solutions to string field theory was performed in [182]. Following this work other papers appeared [183,184,32,185,186,187,188,189,190]. The first analytical solutions of marginal deformations appeared in [191,192], where the solution describing general marginal deformations with regular OPE’s was given. The OPE whose regularity we refer to is that of V with itself. Thus, the simple

³⁷ See [178] for a general study of boundary deformations.

analytical solutions given could not describe more general deformations (of the correct dimensions) with OPE of the form

$$V(z)V(0) \sim \frac{1}{z^2} + \frac{V}{z}. \quad (6.5)$$

It is known that the inclusion of the second term in the OPE above results in a perturbation, which is not exactly marginal. The case where it is absent,

$$V(z)V(0) \sim \frac{1}{z^2}, \quad (6.6)$$

should correspond to an exactly marginal deformation. It was also considered in [191] and it was found that one has to add some counter-terms to the expressions that appear in the regular case. The first two counter-terms were given. However, it was not clear if and how could one continue with the construction of these counter-terms. A framework for addressing marginal deformations with a singular OPE was presented in [193,194]. We turn now to describe regular marginal deformations in 6.1. The case of marginal deformations with a singular OPE is presented next, in 6.2.

6.1 Regular marginal deformations

The solutions of [191,192] obtain a simple form when represented as formal gauge solutions. The key element for a gauge representation of these solutions lies in the introduction of the formal string field J [195,196], which inverts the BRST operator Q ,

$$QJ = |1\rangle. \quad (6.7)$$

This looks very similar to (3.7). The difference is that here Q is the usual BRST operator, whose cohomology is not empty and so this state should not exist. We can nevertheless write down a formal solution to (6.7),

$$J = -\frac{2}{\pi}B \int_0^{-1} W_t dt. \quad (6.8)$$

The string field J in (6.8) looks very similar to A of (5.36). The only difference is the range of integration, which is $(0, -1)$ for J and $(0, 1)$ for A . This range of integration is what causes J to be a non-legitimate string field, since the local coordinate patch is not included in this range. Recall that the local coordinate patch is reserved for the test state, with which the surface state should be contracted and so should be kept intact. A wedge state can be represented by a Gaussian wave function in the infinite dimensional space of string modes. In the limit in which the identity string field W_0 is approached, i.e., the limit in which the surface reduces to solely the local coordinate patch,

this Gaussian wave function approaches a product of delta functions in some directions and constants in the others. That is to say, the standard deviation σ in $e^{-\frac{x^2}{2\sigma}}$ approaches zero or infinity in all directions. When one tries to go beyond the identity string field to wedge states with $t < 0$, σ changes sign and the wave function diverges badly. Despite the above, one can make sense of J when it multiplies states that have enough strip length that can be removed from them. If the states multiplying J from both sides are of the form $\mathcal{O}(0)|0\rangle$ and the OPE of \mathcal{O} with itself has a zero, then (6.7) indeed holds.

Using J the form of the solution is very simple. The gauge field is given by

$$\Lambda = \lambda J\Psi_1, \quad (6.9)$$

where Ψ_1 is given by (6.2). Working in the left-scheme this results in the string field

$$\Psi = Q\Lambda \frac{1}{1-\Lambda} = \Psi_1 \frac{\lambda}{1-\lambda J\Psi_1}, \quad (6.10)$$

where we used (6.3) and (6.7). This is a solution by construction (being given in a pure gauge form), provided it is a legitimate string field despite the usage of J in its definition. We can expand the solution and write

$$\Psi_n = \Psi_1 J\Psi_1 J \cdots J\Psi_1, \quad (6.11)$$

with n appearances of Ψ_1 . Since J removes a varying amount of strip we can write the above (in split-string notations) as

$$\Psi_n = W_{\frac{1}{2}}(cV) \int_0^1 dt_1 W_{t_1}(cV) \int_0^1 dt_2 W_{t_2}(cV) \cdots \int_0^1 dt_{n-1} W_{t_{n-1}}(cV) W_{\frac{1}{2}}. \quad (6.12)$$

We see that in the lower limit of the integrations one gets a collision of two cV vertex operators. If there are singularities in the OPE it may lead to an expression that is not well behaved and to a breakdown of the equation of motion. Nevertheless, this is a perfectly well behaved solution for the regular marginal deformations.

The above method for constructing (regular) marginal deformations was used in [191,192] for the construction of the marginal deformation associated with the rolling tachyon. The coefficient of the tachyon field was evaluated as a series in e^{X_0} and it seemed that the series converges uniformly for all X_0 and represents an oscillatory tachyon with an ever growing amplitude. This is in accord with similar behaviour observed (in the Siegel gauge) in level truncation analysis [111]. This behaviour is counterintuitive and it seems to be in conflict with the exponential grow result found using BSFT³⁸. In [111] it was claimed that this behaviour is acceptable in principle for a system with infinitely many

³⁸A way to avoid this behavior was recently found by Hellerman and Schnabl [197]. They studied tachyon condensation with a tachyon perturbation evolving along a

time derivatives. Later, it was shown that the apparent contradiction between the smooth rolling of the tachyon in BSFT and the divergent oscillatory one of cubic string field theory can be attributed to the (non-local) field redefinition between the two theories [198]³⁹. Indeed, a direct evaluation of the partition function of the rolling tachyon solution of [191,192], produced a result very similar to the one obtained in BSFT [200].

Moreover, the numerical result of [198] was shown to hold analytically in [201], where a gauge equivalent family of solutions was constructed for the rolling tachyon solution. In one limit, the solution reduces to a solution based on the identity string field. This is interpreted as an IR limit, where internal structure is lost⁴⁰. Still, it may seem strange that in a fixed gauge, namely Schnabl's gauge, the rolling tachyon solution oscillates widely and does not seem to converge to the tachyon vacuum, i.e., Schnabl's solution, found in the same gauge. One possible resolution of the problem can come from the fact that the Schnabl gauge is not a complete gauge choice. An exact string field exists in this gauge, namely $Q\Lambda$ on which Schnabl's solution is based (5.5). This explanation is not very satisfactory for a couple of reasons. First, Schnabl's gauge gives almost a complete gauge fixing. Relating the apparent different behavior to the small residual gauge symmetry seems unnatural. Second, it does not explain the relation between the two solutions. So it is not clear how and in which sense does the rolling tachyon solution approach the tachyon vacuum at late time⁴¹.

Another resolution to the problem was studied by Ellwood in [201], where it was suggested that in spite of the apparent different behaviour, the rolling tachyon solution does approach the tachyon vacuum for late time. The rolling tachyon solution was expanded as an integral over strip length,

$$\Psi = \int_0^\infty \Psi_r . \tag{6.13}$$

light-like, rather than a time-like direction, including a dilaton background that introduces an effective friction term. Their solution does not suffer from any wild oscillations.

³⁹ An interesting approach for obtaining (level-truncated) solutions of tachyon condensation is to use the diffusion equation in order to relate solutions of the non-local system to analogous solutions of a local system [199]. Note, however, that due to an improper limiting procedure, one of the solutions obtained there has a cusp singularity at $t = 0$ and is actually a gluing at $t = 0$ of two genuine solutions [198]. The authors of [199] end up discarding this solution, albeit for other reasons.

⁴⁰ Related work appeared in [202].

⁴¹ Alternatively, one can try to study the rolling tachyon solution directly in the vicinity of the tachyon vacuum. A preliminary study of this question was presented in [203]. However, it is not clear how to relate the solution there neither to the tachyon vacuum of [2] nor to the rolling tachyon solution of [191,192].

The ghost part of Ψ_r is simple, but its matter dependence is complicated, since it contains contributions from all Ψ_n , as can be seen from the explicit expression (6.12). This is due to the appearance of J in (6.11). The dependence of Ψ_r on X_0 can be described by a function $F_r(X_0)$ as,

$$\Psi_r^{mat} = F_r(X_0)W_r. \quad (6.14)$$

Next, it was assumed that as $x_0 \rightarrow \infty$ a limit exists for $F_r(X_0)$ and it was further assumed that this limiting function can be written as

$$\lim_{x_0 \rightarrow \infty} F_r(X_0) = f_r(x_0). \quad (6.15)$$

With this assumption it was found that at late time the rolling tachyon solution indeed reduces to the tachyon vacuum solution. It is not quite clear how to reconcile this result with the numerical study of the tachyon coefficient described above. At any rate, this result demonstrates that the two solutions are somehow related. Further study of this point is certainly worthwhile.

6.2 Singular marginal deformations

The method of [191,192] is not easily generalized to the case of a singular OPE⁴². A different method for constructing a formal gauge field for this case was developed in [193]. There, a specific marginal deformation, namely

$$V = i\partial X^\mu, \quad (6.16)$$

for some μ , which we leave implicit henceforth, was addressed. This deformation describes a Wilson loop, for the case of a compact X . If X is not compact, this deformation describes a gauge transformation. While the method used is especially simple for the ∂X case, it can also be generalized. One root towards this generalization was sketched in [193], while another (equivalent) one was developed in [194]. We continue now with the ∂X deformation and return to the generalization later.

The observation of [193] was that in the case (6.16) one can write

$$cV(0)|0\rangle = Qx_0|0\rangle. \quad (6.17)$$

This suggests that the deformation is a pure gauge one. This is indeed the case for a non-compact X . However, for a compact X direction, the zero mode x_0 does not exist and the expression (6.17) is formal⁴³.

⁴² Some ideas regarding the origin of the singularities and the way they should be dealt with appeared in [204].

⁴³ The zero mode x_0 was similarly used in the context of closed string field theory in [205].

We therefore pretend that the compact field X has in its expansion the zero mode x_0 . Defining

$$\Lambda = i\lambda X(0) |0\rangle, \quad (6.18)$$

we get the desired first order result

$$\Psi_1 = ic\partial X(0) |0\rangle. \quad (6.19)$$

It may seem that this is the end of the story. However, it is easy to see that Ψ_2 turns out to be linear in x_0 , with higher order polynomial dependence on x_0 of Ψ_n for $n > 2$. This is not a legitimate result, since x_0 does not really exist in this case⁴⁴. The resolution of this problem comes from a redefinition of the gauge field,

$$\Lambda = \sum_{k=1}^{\infty} \lambda^k \Lambda_k, \quad (6.20)$$

with the expression (6.18) identified as the leading order term Λ_1 . Higher order terms should be chosen such that there is no x_0 -dependence of the solution, while not changing its physical content.

There are many ways to complete Λ so as to obtain an x_0 -independent solution. One particularly simple choice is to choose

$$\Lambda_n = -\frac{(-i)^n}{n!} (X^n, \underbrace{1, \dots, 1}_{n-1}). \quad (6.21)$$

Here we introduced a notation, similar to the split-string notation. A vector of length n represents the star product of n vacuum states with insertions. A vacuum state with no insertion is represented by an insertion of an identity. The fact that we are using insertions of the form X^n may seem problematic. One obvious objection is the singularities from the OPE of the X fields. This is resolved by defining the products to be normal ordered at each point. Another issue is that these fields are non-primary. That should not prevent us from using them. We only have to know how they transform under conformal transformations. For that end it is simplest to define

$$X^n \equiv \partial_k^n e^{kX} \Big|_{k=0}, \quad (6.22)$$

where e^{kX} is implicitly normal ordered. We use this definition and the relation

$$[Q, e^{kX}(z)] = \left(-k^2 \partial c e^{kX} + kc \partial X e^{kX} \right)(z), \quad (6.23)$$

⁴⁴ One can consider in principle x_0 -dependent solutions, provided that the dependence is compatible with the compactification, i.e., periodic solutions. We expect to have such form of solutions for the periodical lump system. These important solutions were not yet constructed.

in order to write

$$[Q, X^n] = (\partial_k)^n [Q, e^{kX}] \Big|_{k=0} = nc \partial X X^{n-1} - n(n-1) \partial c X^{n-2}. \quad (6.24)$$

Let us restrict ourselves to ansätze that generalize (6.21), i.e., with Λ_n being the star product of n states of the form $X^{k_i} |0\rangle$ with $\sum_{i=1}^n k_i = n$. For this class of gauge fields the x_0 -independence condition is satisfied provided the following recursion relation holds,

$$\partial_{x_0} \Lambda_n = -i\lambda \Lambda_{n-1} W_1, \quad (6.25)$$

with the initial condition for Λ_1 (6.18). The recursion relation and initial condition can be compactly summed to

$$\partial_{x_0} \Lambda = i\lambda(1 - \Lambda) W_1. \quad (6.26)$$

It can be seen that the expression (6.21) indeed obeys this equation and is therefore a solution.

Many more solutions exist with the form of the ansatz suggested. All of them are supposedly gauge equivalent, so one may ignore this degeneracy and choose to work with the simple expression (6.21). This will not be good enough, though, when one wants to impose the reality condition on string fields (2.26). The reality condition states that the string field Ψ is real with respect to the involution defined by composing hermitian and BPZ conjugations. This will be the case if Λ is pure imaginary. For our ansatz the conjugation is equivalent to inverting the orientation of the strip (while retaining the formal Grassmann ordering) and complex conjugating coefficients. The inversion of orientation implies also a minus sign for every derivative present in the expression. The first order term (6.18) is imaginary as it should, due to the i in the definition there. The higher order terms on the other hand carry no definite symmetry property and so fail to be real or imaginary.

We can trace the origin of lack of symmetry to the fact that we are using the left-scheme (6.10). This scheme is very easy to use. In particular it was easy to write a closed form expression for the x_0 -independent condition (6.26). Another simple scheme is the right scheme, which is the “mirror image” of the left scheme. In this scheme the solution is represented as

$$\Psi = \frac{1}{1 + \Lambda} Q \Lambda. \quad (6.27)$$

The gauge potential that generates an x_0 -independent solution is

$$\Lambda_n = \frac{i^n}{n!} \underbrace{(1, \dots, 1, X^n)}_{n-1}. \quad (6.28)$$

The left and right schemes are easy to use since the rational function $(1 - \Phi)^{-1}$ has a very simple variation,

$$\delta \frac{1}{1 - \Phi} = \frac{1}{1 - \Phi} \delta \Phi \frac{1}{1 - \Phi}. \quad (6.29)$$

For comparison, the variation of an exponential function can be most nicely written in terms of an integral⁴⁵,

$$\delta e^\Phi = \int_0^1 dt e^{t\Phi} \delta \Phi e^{(1-t)\Phi}. \quad (6.30)$$

Trying to work directly in more complicated schemes results in complicated reality conditions. It is nevertheless possible to define a real solution by an iteration prescription [158]. While the iteration formula is given in a closed form, the resulting expressions are quite complicated and no closed form expression (other than the iteration formula) was found.

Another way of defining real solutions was given in [194]. This relies on the gauge transformation relating the left and right schemes [193],

$$e^\Lambda = \sum_{n=0}^{\infty} \frac{(i\lambda)^n}{n!} \underbrace{(X_n - X_1)^n}_n. \quad (6.31)$$

In the above expression the n^{th} order term is built upon the wedge state $|n\rangle$. It is clear from this representation that e^Λ is x_0 -independent, which implies that this is a genuine gauge transformation. Writing the transformation explicitly one has,

$$\Psi_r = e^{-\Lambda}(\Psi_l + Q)e^\Lambda, \quad \Psi_l = e^\Lambda(\Psi_r + Q)e^{-\Lambda}, \quad (6.32)$$

where the subscripts l, r stand for “left” and “right” respectively. The real solution of [194] is based on “going half the way” between the left and right solutions,

$$\Psi = e^{-\frac{\Lambda}{2}}(\Psi_l + Q)e^{\frac{\Lambda}{2}} = e^{\frac{\Lambda}{2}}(\Psi_r + Q)e^{-\frac{\Lambda}{2}}. \quad (6.33)$$

Using the above defining expressions and the relation

$$\Psi_r = -\Psi_l^*, \quad (6.34)$$

immediately implies the reality of Ψ .

While the solution (6.34) is given in a nice closed form, it relies on the gauge transformation (6.31), whose form is misleadingly simple. This gauge transformation is not some sort of an exponent, since the size of the wedge states

⁴⁵ To prove this identity, expand the l.h.s, change one summation index and use the familiar integral representation of the Beta function.

varies. Explicitly one may write,

$$e^\Lambda = 1 + \frac{(i\lambda)^2}{2} \left((X^2, 1) + (1, X^2) - 2(X, X) \right) + \dots \quad (6.35)$$

Note, in particular, that the first order term drops out. The power of λ , the wedge size and the total X power are correlated. This will remain so in calculating any function of e^Λ such as its inverse $e^{-\Lambda}$. The exact form of $e^{-\Lambda}$ does not correspond to $\lambda \rightarrow -\lambda$. Explicit calculation gives

$$\begin{aligned} e^{-\Lambda} = & 1 - \frac{(i\lambda)^2}{2} (X_2 - X_1)^2 - \frac{(i\lambda)^3}{3!} (X_3 - X_1)^3 \\ & + \frac{(i\lambda)^4}{4!} \left(6(X_1 - X_2)^2 (X_3 - X_4)^2 - (X_1 - X_4)^4 \right) + \dots \end{aligned} \quad (6.36)$$

Here we left the wedge size implicit. Note that already the second order term has “the wrong sign” relative to a $-\lambda$ assignment, while the fourth order term is completely different from the corresponding one in e^Λ . Calculating the functions $e^{\pm\frac{\Lambda}{2}}$ needed for (6.34) results in quite complicated expressions and the practical evaluation of a real solution at a given level may turn out to be more complicated than evaluating the recursion relation. Nevertheless, one may think of several cases where the closed form of (6.34) makes it more adequate for calculations than its counterpart.

We have seen that the explicit form of real solutions is more complicated than that of the non-real solutions in the left and right schemes. These solutions are gauge equivalent in the theory without the reality condition. One may decide then, to enlarge the gauge orbits of the real solutions, so as to include these ones as well. The theory with the reality condition would then contain all the gauge orbits that contain a real representative. This procedure does not add any degrees of freedom to the theory with the reality condition, since the added solutions are all gauge equivalent to existing real ones. Thus, it seems that one can safely forget about imposing the reality condition and work with simpler non-real solutions, provided that one can prove that they are gauge equivalent to real one.

As already mentioned, the generalization of the above construction for other marginal deformations with the same OPE (6.6) can be carried out in one of two equivalent ways. In [193], it was suggested to define a formal primitive analogue to X for the other marginal deformations and demand that the final expressions are independent of it. In terms of powers of such a primitive the formal gauge solution takes exactly the same form as for the ∂X deformation. While formally this is the end of the story, when one wants to evaluate explicit coefficients, one faces the problem of normal ordering. The point is that the powers of the primitive should be normal ordered with respect to the normal ordering associated with this primitive, while for the evaluation of coefficients is performed using the normal ordering associated with ∂X . The resolution

of [194] to this problem is equivalent to noting that the fact that the final expression for Ψ is independent of the primitive implies that one can write the solution in terms of integrals of the marginal deformation itself. Thus, one can use the normal ordering of the primitive to get a fully normal ordered expression and then in this expression replace all the primitives by line integrals. This basic idea was presented in [194] in a sophisticated form, defining the normal orderings explicitly. The fact that only line integrals appear in the expression for Ψ was interpreted as suggesting that the integrated vertex operator is the building block of the string field theory solution, rather than the unintegrated one, as suggested by Sen [179,180,181].

7 Oscillator representation

In this section we discuss the recent developments in string field theory in the oscillator basis formalism. As stated already, this formalism is related to a given background and as such was not relevant to most of the recent advance in the field. Nonetheless, it is plausible that this formalism will play a more dominant role in constructing analytical solutions describing lumps.

We start in 7.1 by introducing the tools needed, i.e., the form of the string vertices in the oscillator representation and the continuous (κ) basis, as well as introducing the problem of “normalization anomaly”, which used to be a stumbling block for the derivation of analytical results in the oscillator representation. Next, in 7.2 we describe the form of Schnabl’s operators in the oscillator representation and explain how to solve the problem of normalization anomalies.

7.1 Representation of string vertices and the continuous basis

For the construction of the oscillator representation recall the form of the string field in flat background (2.1). All possible values for the coefficient fields span the (classical) string field Hilbert space⁴⁶. Our first goal is to define the

⁴⁶To have strictly a Hilbert space one should also give a positive definite inner product that will restrict the possible values of the coefficient fields. However, there is no natural candidate for such an inner product. One reason for that is that the ghost fields and the time direction induce a negative norm in the inner product of the first quantized string. On the other hand, when not restricted, star multiplication of string fields leads to “associativity anomalies” [34,206]. It is common in the literature to refer as a Hilbert space (or Fock space) to the subspace of string fields with a finite number of non-zero coefficients multiplying a finite number of oscillators acting on the vacuum. This is of course not a Hilbert space. It is also not a closed space with

string vertices in this representation.

For the description of interaction the string vertices should be defined in terms of the oscillators. The n -string vertex is a state in the n^{th} tensor power of the single string Hilbert space. In this space the string vertex is (up to zero-modes) a squeezed state in the matter as well as in the ghost sector. One can write

$$\int \Psi = \langle V_1 | \Psi \rangle, \quad (7.1)$$

$$\int \Psi_1 \star \Psi_2 = {}_{12} \langle V_2 | |\Psi_1\rangle_1 |\Psi_2\rangle_2, \quad (7.2)$$

$$\int \Psi_1 \star \Psi_2 \star \Psi_3 = {}_{123} \langle V_3 | |\Psi_1\rangle_1 |\Psi_2\rangle_2 |\Psi_3\rangle_3, \quad (7.3)$$

and so on for higher vertices if needed. Here, the subscripts next to the bras and kets refer to the Hilbert space index, since as mentioned, V_n for $n > 1$ live in a multi-string space, so for (7.2) for example, the vertex lives in the spaces one and two. The string field Ψ_1 lives in space number one and the string field Ψ_2 lives in space number two. Next, we define $|V_2\rangle$ as the inverse of $\langle V_2|$ by the relation

$${}_{12} \langle V_2 | |V_2\rangle_{23} = \mathbf{1}_{13}, \quad (7.4)$$

where

$$\mathbf{1}_{13} |\Psi\rangle_1 = |\Psi\rangle_3. \quad (7.5)$$

The vertex $|V_2\rangle$ implements (inverse) BPZ conjugation, up to a sign that may originate from the ghost zero modes.

The concrete realization of the vertices V_1 and V_2 is very simple. It can be derived from their geometrical interpretation as inducing a gluing of surfaces. In the matter sector the gluing of a string about its middle can be represented by setting to zero the coefficients of the odd modes as well as of the momenta of the even modes. In terms of oscillators this infinite product of delta functions is represented by

$$\begin{aligned} \langle V_1^m | &= \int d^{26}k \langle k | \delta(k) \exp \left(-\frac{1}{2} \sum_{n,m=1}^{\infty} a_n^\mu C_{n,m} a_m^\nu \eta_{\mu\nu} \right) \\ &= \langle 0 | \exp \left(-\frac{1}{2} \sum_{n=1}^{\infty} (-1)^n a_n^\mu a_n^\nu \eta_{\mu\nu} \right), \end{aligned} \quad (7.6)$$

where the implicitly defined matrix C is the twist matrix,

$$C = (-1)^n \delta_{n,m}, \quad (7.7)$$

and $\eta_{\mu\nu}$ is the flat 26-dimensional metric. The summation on the indices μ, ν is implicit. Henceforth, we shall refrain from writing the metric and the spatial

respect to the start product. However, as we have nothing new to say about it, we shall follow the common lore by simply ignoring this issue.

indices explicitly unless needed, as they do not play an important role in most of the following discussion. One should keep in mind though, that there are 26 coordinate dimensions. The gluing defined by the two-vertex is imposed by identifying the even modes of the two strings and the odd modes of their momentum. This results in

$$\begin{aligned} {}_{12}\langle V_2^m | &= \int d^{26}k_1 d^{26}k_2 {}_{12}\langle k_1, k_2 | \delta(k_1 + k_2) \exp\left(-\frac{1}{2} \sum_{n,m=1}^{\infty} a_n^1 C_{n,m} a_m^2\right) \\ &= \int d^{26}k {}_{12}\langle k, -k | \exp\left(-\sum_{n=1}^{\infty} (-1)^n a_n^1 a_n^2\right), \end{aligned} \quad (7.8)$$

where the superscripts on the oscillator modes represent the space-number on which they act.

The application of these simple geometric ideas to the three-vertex V_3 is technically more complicated. The easiest way to find it is first to assume the following squeezed state form

$$\begin{aligned} {}_{123}\langle V_3^m | &= \int d^{26}k_1 d^{26}k_2 d^{26}k_3 {}_{123}\langle k_1, k_2, k_3 | \delta(k_1 + k_2 + k_3) \cdot \\ &\cdot \exp\left(-\sum_{s,r=1}^3 \left(\frac{1}{2} (a^r | V^{rs} | a^s) + (a^r | V_0^{rs}) k_s + k_r V_{00}^{rs} k_s\right)\right). \end{aligned} \quad (7.9)$$

Here we introduced the notation of round brackets to represent the infinite vectors of different string modes. Now, the zero modes enter into the exponent and when singled out as above the data defining the three-vertex reduces to the matrices (in mode space) V^{rs} the vectors $|V_0^{rs}\rangle$ and the scalars V_{00}^{rs} . To find the coefficients V_{nm}^{rs} one can first note that (up to the infinite volume factor $\delta(0)$) one can write

$$V_{nm}^{rs} = -{}_{123}\langle V_3^m | a_n^{r\dagger} a_m^{s\dagger} | 0 \rangle_{123}. \quad (7.10)$$

Writing the creation operators using a contour integral of ∂X one can use the CFT expression (2.24) and the explicit form of $f_r^{(3)}$ (2.25)⁴⁷ to get to the following expression [3,4]

$$V_{nm}^{rs} = -\frac{1}{\sqrt{nm}} \oint \frac{dzdw}{(2\pi i)^2 z^n w^m} \frac{f_r'(z) f_s'(w)}{(f_r(z) - f_s(w))^2}. \quad (7.11)$$

It is easy to see in this representation that the matrices obey the symmetry properties one would expect them to have, namely V^{rs} depends only on $r - s$ modulo 3. Analogous expressions hold also for the vectors and the scalars.

A consequence of the fact that all the string vertices are given by squeezed states is that squeezed states form a subalgebra of the star-product and the

⁴⁷ We henceforth denote $f_r^{(3)}$ by f_r for simplicity.

expressions for multiplying squeezed states⁴⁸ are formally very simple. However, these formal expressions involve multiplying and inverting the infinite dimensional matrices defining the states and the vertices.

Working with infinite dimensional matrices may be complicated and one may be forced to advance using numerical tools in the spirit of level truncation. A natural route to simplify the work with matrices is to diagonalize them. This cannot be done in this case, since a direct evaluation shows that these matrices are not commutative and so cannot be simultaneously diagonalized. There is, however, a simple way out. The matrices

$$M^{rs} \equiv CV^{rs} \quad (7.12)$$

are commutative and so can be simultaneously diagonalized. This task was performed by Rastelli Sen and Zwiebach [20]. They used the fact that the three matrices M^{rr} , $M^{r(r\pm 1)}$ commute with the matrix K_1 related to the Virasoro operator \mathcal{L}_{-1} (4.7). The matrix K_1 has all real numbers as eigenvalues, without degeneration,

$$K_1 |\kappa\rangle = \kappa |\kappa\rangle. \quad (7.13)$$

The eigenvalues of the M matrices are found to be,

$$M^{rr} |\kappa\rangle = -\frac{1}{1 + 2 \cosh(\frac{\kappa\pi}{2})} |\kappa\rangle \equiv \mu(\kappa) |\kappa\rangle, \quad (7.14)$$

$$M^{r(r\pm 1)} |\kappa\rangle = \frac{1 + \exp(\pm \frac{\kappa\pi}{2})}{1 + 2 \cosh(\frac{\kappa\pi}{2})} |\kappa\rangle. \quad (7.15)$$

The eigenvectors $|\kappa\rangle$ were normalized in [21]. Given a squeezed state defined by a matrix S , one can consider the matrix $T = CS$. Squeezed states whose matrix T is diagonal in the continuous basis form a subalgebra that can be easily manipulated.

The twist matrix C also has a simple form in the continuous basis, as it only interchanges the eigenvectors $|\pm\kappa\rangle$. With the standard definition of the eigenvectors it is given in this subspace by the matrix

$$C_{-\kappa,\kappa} = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}. \quad (7.16)$$

This means that while the matrices defining the string vertex are not mutually diagonalizable and in particular are not diagonal in the continuous basis, they are block diagonal in this basis with two by two blocks. Again, block diagonal squeezed states form a subalgebra [207]. Within this subalgebra the operations

⁴⁸By squeezed states we also mean states with linear terms. In particular coherent states that do not contain the quadratic terms are also part of this subalgebra.

involving infinite matrices reduce to manipulating two by two matrices and integrating over κ .

Let $|n\rangle$ be the natural discrete basis states, i.e.,

$$|n\rangle = \underbrace{(0, \dots, 0, 1, 0, \dots)}_n. \quad (7.17)$$

The transformation between the discrete $|n\rangle$ basis and the continuous $|\kappa\rangle$ basis is given by

$$|\kappa\rangle = \sum_{n=1}^{\infty} v_n^\kappa |n\rangle, \quad |n\rangle = \int_{-\infty}^{\infty} d\kappa v_n^\kappa |\kappa\rangle. \quad (7.18)$$

The transformation coefficients v_n^κ obey the usual unitarity properties⁴⁹

$$\sum_{n=1}^{\infty} v_n^\kappa v_n^{\kappa'} = \delta(\kappa - \kappa'), \quad (7.19)$$

$$\int_{-\infty}^{\infty} d\kappa v_n^\kappa v_m^\kappa = \delta_{nm}, \quad (7.20)$$

and are defined by the generating function,

$$f_\kappa(z) = \sum_{n=1}^{\infty} \frac{v_n^\kappa}{\sqrt{n}} z^n = \frac{1}{\sqrt{\mathcal{N}(\kappa)}} \frac{1 - e^{-\kappa \tan^{-1} z}}{\kappa}, \quad \mathcal{N}(\kappa) = \frac{2}{\kappa} \sinh\left(\frac{\kappa\pi}{2}\right). \quad (7.21)$$

Transforming the creation and annihilation operators to the new basis results in the commutation relation

$$[a_\kappa, a_{\kappa'}^\dagger] = \delta(\kappa - \kappa'). \quad (7.22)$$

The continuous basis proved to be useful for dealing with many questions. In particular, it was used in [13] to formulate the continuous Moyal representation of the star algebra. Analytical calculation of tension ratio was performed in [209] while in [21] the equivalence of two definitions of the kinetic operator of vacuum string field theory was proven. Furthermore, the continuous basis is strongly related to wedge states. Of all surface states, wedge states are the only ones whose matrix T is diagonal and the so called (hybrid) butterfly states [168,169,167,207] are the only (other) ones with a block diagonal matrix [210,211]⁵⁰.

⁴⁹ Some other properties of these coefficients are collected in the appendix of [208].

⁵⁰ Surface states are necessarily represented by squeezed states in the oscillator representation. However, not all squeezed states are surface states. A simple criterion to decide whether a given (matter) squeezed state is (the matter part of) a surface state was introduced in [207]. Another criterion was later introduced that stressed integrability as the main character of surface states [212] (integrability within this

There are also some difficulties related to the continuous basis. The Virasoro generators take a very singular form when expressed in this basis. They are more singular than delta functions [13,15]. In [218] it was demonstrated that they can nevertheless be expressed by delta functions whose arguments are complex. It was explained in this paper how to deal with such expressions⁵¹.

Another problem one may encounter with the continuous basis is related to the calculation of normalization factors. It is quite common that a state has a vanishing normalization in the matter sector and an infinite one in the ghost sector or vice versa. Thus, in this type of calculations one has to introduce the ghost sector and properly regularize both sectors, in order to get a sensible result⁵². The ghost sector can be incorporated either using the bc system or its bosonized form. In the first case the string vertices are given by a zero mode product multiplying a zero ghost number squeezed state, i.e., an expression of the form $\exp(bSc) |0\rangle$. It is possible to define the continuous modes b_κ, c_κ , in a manner similar to (7.22). It turns out that the 1-vertex, i.e., the identity string field, takes a very singular form in this representation. It was not clear how overcome this problem. Therefore, the bosonized ghost approach was usually used within the continuous basis.

In the bosonized ghost formalism the field ϕ is added to the 26 spatial bosons. This field is a linear dilaton whose total central charge equals -26 . The string vertices related to this field have the same quadratic terms defining them, as the matter sector vertices, but somewhat different linear terms. When trying to evaluate the inner product of two wedge states in the oscillator representation with bosonized ghosts, erroneous results were encountered [221]. Namely, it was found that the result is not unity.

From CFT considerations it is clear that the inner product of two arbitrary surface states (wedge states in particular) should equal unity [3]⁵³,

$$\langle S|V\rangle = \langle 0| e^{\sum_{n=2}^{\infty} s_n L_n} e^{\sum_{m=2}^{\infty} v_m L_{-m}} |0\rangle = 1. \quad (7.23)$$

context was further studied in [213,214]). The two criteria were shown to be equivalent in [210], where it was also shown that the first column of the defining matrix of a squeezed state is equivalent to a unique surface state. Related study of the ghost sector was performed in [215,216,217,210].

⁵¹ Another expression for some of the Virasoro generators was given in [219].

⁵² This state of affairs does not imply by itself that the string field exists in the limit where the regularization cutoff is removed [220]. However, when dealing only with states that are well defined in the CFT no problem of principle should arise.

⁵³ Of course, in order to get a non-zero result the ghost zero modes should be saturated. Also, in the inner product an infinite factor comes from the momentum conservation delta function, which as usual represents the infinite volume of space time. In our current discussion it is assumed that all zero modes (bosonic and fermionic) were taken care of.

Here s_n, v_m are the coefficients defining the two surface states S, V according to (2.29) and the final result comes from using the Virasoro algebra with central charge zero in order to move all positive operators to the right and all negative ones to the left. Moreover, since (the integer) wedge states are the surfaces on which the string vertices are defined, the above expression is related to the descent relations of the string vertices,

$$\langle V_{n+1} | V_1 \rangle = \langle V_n | . \quad (7.24)$$

These relations should hold, without additional normalization factors, to ensure the consistency of string field theory.

Let us concentrate on evaluating the inner product of the wedge states $|1\rangle$ and $|3\rangle$. This gives the fundamental descent relation,

$$\langle V_1 | V_3 \rangle = \gamma_{13} |V_2\rangle , \quad (7.25)$$

where we introduced the normalization factor that should equal unity from CFT considerations. Squeezed state algebra reveals [11],

$$\begin{aligned} \gamma_{13} = & \det(\mathbf{1} - CV_3^{11})^{-\frac{d+1}{2}} \cdot \\ & \exp \left(\frac{9}{4} \left(-\frac{1}{2} \left(V_{30}^{g11} | (\mathbf{1} - CV_3^{11})^{-1} C | V_{30}^{g11} \right) + \left(V_{10}^g | (\mathbf{1} - V_3^{11} C)^{-1} | V_{30}^{g11} \right) \right. \right. \\ & \left. \left. - \frac{1}{2} \left(V_{10}^g | (\mathbf{1} - V_3^{11} C)^{-1} V_3^{11} | V_{10}^g \right) - \frac{1}{2} V_{300}^{g11} \right) \right). \end{aligned} \quad (7.26)$$

Here, the twist matrix comes from its part in defining the one-vertex (7.6), the factor of $d + 1 = 27$ in the determinant comes from the quadratic terms of the d space-time dimensions and from the bosonized ghost direction and the terms in the exponent all come from linear terms of the bosonized ghost, which are absent for the space-time directions. We marked here all ghost terms with a superscript g , which we omit later in order to improve readability. The first term in the exponent comes from terms in the three-vertex that are linear in momentum. This term is absent in the matter sector since the momentum there is zero. Saturation of the ghost zero mode requires $p^g = \pm \frac{3}{2}$, which results in the factor of $\frac{9}{4}$ common to all the terms in the exponent. The factor $|V_{10}^g\rangle$ is the linear term of the one-vertex. Such a factor could not exist in the matter sector, since the momentum there is zero. Both the determinant and the exponent formally diverge.

In order to be able to compare the divergence of the exponent and determinant we use the usual trick of expressing the determinant in terms of a trace of the

logarithm of the matrix,

$$\begin{aligned}
\det(\mathbf{1} - CV_3^{11}) &= \exp\left(\text{tr}\left(\log(\mathbf{1} - CV_3^{11})\right)\right) = \exp\left(\sum_{n=1}^{\infty} (n|\log(\mathbf{1} - CV_3^{11})|n)\right) \\
&= \exp\left(\sum_{n=1}^{\infty} \int d\kappa d\kappa' (n|\kappa)(\kappa|\log(\mathbf{1} - CV_3^{11})|\kappa')(\kappa'|n)\right) \quad (7.27) \\
&= \exp\left(\sum_{n=1}^{\infty} \int d\kappa (n|\kappa)\log(1 - \mu(\kappa))(\kappa|n)\right) = \exp\left(\delta(0) \int d\kappa \log(1 - \mu(\kappa))\right).
\end{aligned}$$

Here, we wrote the trace explicitly in the second equality. In the third one a unity was inserted twice in order to pass to the continuous basis. Next, we used (7.14) to get $\delta(\kappa - \kappa')$. Then, a sum over n left us with another delta function $\delta(0)$, which is the manifestation of the divergence. Similar expressions are obtained also for the exponent part of (7.26).

In order to regularized the $\delta(0)$, a cutoff on n was imposed on (7.19) for $\kappa = \kappa'$. This cutoff l , called the level⁵⁴ gives the prescription [20,221,218,222],

$$\delta(0) \rightarrow \frac{2 \log l - \psi\left(\frac{i\kappa}{2}\right) - \psi\left(-\frac{i\kappa}{2}\right)}{4\pi}, \quad (7.28)$$

where ψ is the polygamma function. It turns out that while the infinite ($\log l$) part cancels out exactly for $d = 26$, the finite part does not cancel out. Rather, it gives [223],

$$\gamma_{13} = 3^{3/8} e^{\frac{3}{2}\gamma + 36\zeta'(-1)} \left(\frac{\Gamma\left(\frac{1}{3}\right)}{\sqrt{\pi}}\right)^9 \approx 0.382948, \quad (7.29)$$

where Γ is the gamma function, γ is Euler's constant and ζ' is the derivative of the zeta function.

A possible resolution of this problem was suggested in [224,225]. According to this proposal, the normalization constant γ_{13} is not an artifact but a correct result. All string vertices should have normalization factors, i.e. $|V_n\rangle \rightarrow Z_n |V_n\rangle$. The normalization factors supposedly originate from the partition function on a surface with a conical singularity of the sort used for the definition of the vertex. The two-vertex is defined over a surface without a conical singularity and so carries a trivial normalization, $Z_2 = 1$. Hence, the descent relation holds, provided that $\gamma_{13}Z_1Z_3 = 1$. Similar relations should also hold for the other descent relations. This proposal have some problems. First, these partition

⁵⁴ Note that while resembling level truncation, this is not the same. In level truncation one keeps all fields up to a given level, here we effectively keep all fields, which are built from oscillators up to a given level, i.e., we keep an infinite number of fields. This approximation is occasionally referred to as ‘‘oscillator level truncation’’ in order to distinguish it from the usual level truncation.

functions were nowhere directly calculated and no other test was performed in order to show the consistency and universality of the proposal. More seriously, the proposal is inconsistent with (7.23).

One faces several options for an alternative clarification of the source of the anomaly. It can be related to the use of the continuous basis, to the use of the oscillator level truncation needed for defining the normalization or to the use of the bosonized form of the ghosts. In [226]⁵⁵ these possibilities were studied. It was found that when carefully treated, the discrete basis gives the same normalization constant as the continuous one, suggesting that the use of the continuous basis is not the source of the problem. Also, it was found that while the use of the ghosts in their bc form gives a different normalization factor, this factor is also not equal to unity. All that seems to suggest that oscillator level truncation is a bad regularization. Without a proper resolution of this problem it seems that one cannot really use the oscillator basis as a complete framework for the study of string field theory.

7.2 Schnabl's operators and the resolution of normalization anomalies

Schnabl's construction uses the CFT language. Despite that and regardless of the problems with the oscillator formalism, one would like to formulate his solution in the oscillator language as well. An obvious reason would be the fact that much advance in string field theory was achieved in the past in this formalism. Another reason is that Schnabl's solution is based on wedge states, which have a very simple representation in the oscillator formalism. In particular, they are diagonal in the continuous basis. Thus, this representation may turn out to be as simple as the CFT one. Alternatively, it may happen that it may teach us something about the oscillator formalism itself. As the primary building blocks of the solution are the operators $\mathcal{L}_0, \mathcal{L}_0^\dagger$, the first step towards an oscillator representation is to cast these operators into an oscillator form, preferably in the continuous basis [156].

An obvious obstacle for this program comes from the fact that Schnabl's operators are combinations of Virasoro operators, which take a very singular form in the continuous basis. Remarkably, it turns out that exactly for the case of Schnabl's operators the highly singular (complex delta function) terms conspire to vanish. The other terms add nicely to give a mild singularity, which corresponds to an almost (block) diagonal form of the operators. The defining matrices in the continuous basis are proportional to the delta function and to

⁵⁵ See also [227] and the recent papers [228,229]. In cubic superstring field theory this problem was addressed in [230].

its derivative,

$$\mathcal{L}_0 = \int_{-\infty}^{\infty} d\kappa d\kappa' \left(\left(\frac{\kappa\pi}{4} \coth\left(\frac{\kappa\pi}{2}\right) \delta(\kappa - \kappa') + \frac{\kappa + \kappa'}{2} \delta'(\kappa - \kappa') \right) a_{\kappa}^{\dagger} a_{\kappa'} \right) \quad (7.30)$$

$$+ \frac{\pi \delta(\kappa + \kappa')}{2\mathcal{N}(\kappa)} a_{\kappa} a_{\kappa'} \Big). \quad (7.31)$$

Similar results were also found for the bosonized ghost sector. However, when trying to verify the basic algebra (4.19) one seem to gets an anomaly⁵⁶,

$$[\mathcal{L}_0, \mathcal{L}_0^{\dagger}] \stackrel{?}{=} \mathcal{L}_0 + \mathcal{L}_0^{\dagger} + \frac{15}{4}. \quad (7.32)$$

This anomaly is similar to the anomaly encountered in the evaluation of the descent relations, but simpler, since it involves only linear expressions. This simple form of the normalization anomaly was used in [231] in order to reconsider the issue of regularization in the oscillator formalism.

There are some features common to good regularization schemes. First, they tend to be analytic in the regularizing parameter. Second, they tend to leave intact (or to simply deform) the main symmetries of the theory considered. In the case at hand it is clear that oscillator level truncation does not enjoy these qualities. The cutoff is performed using a step function, which is obviously not an analytic function and there is no treatment of the relevant symmetries. One may wonder, at this stage, how is that different from the usual level truncation. One difference is that for level truncation the cutoff commutes with the Virasoro algebra. Infinities may arise in relations such as (7.23) when the central charge is non-zero. Thus, the Virasoro symmetry is the relevant one in our case. While it seems that there is no regularization that keeps the Virasoro algebra intact, it is nevertheless possible to deform it in a way that does not alter the role of the central charge. Moreover, this deformation depends analytically on a parameter. The deformation is performed by defining (for $s \leq 1$)

$$L_n^s = s^{|n|} L_n, \quad (7.33)$$

which for $s \rightarrow 1$ reduces to the non-regularized case. The deformed operators satisfy the regularized Virasoro algebra

$$[L_n^s, L_m^s] = s^{|n|+|m|-|n+m|} \left((n-m)L_{n+m}^s + \frac{c}{12}(n^3 - n)\delta_{n+m} \right). \quad (7.34)$$

For the oscillators the regularization is similarly applied,

$$a_n^s = s^n a_n, \quad a_n^{\dagger s} = s^n a_n^{\dagger}. \quad (7.35)$$

⁵⁶ This is of course the result at the critical dimension. For $d \neq 26$ the central term diverges logarithmically as a function of the level l .




Fig. 7. The geometric effect of the regularization on: (a) The definition of the state. (b) The two-vertex. (c) The three-vertex.

A very simple geometric interpretation can be given to the regularization if one thinks of the positive Virasoro modes as related to bra states and of the negative ones as related to ket states. In such a case the power of the cutoff is simply the conformal weight of the state created and so the regularization simply amounts to squeezing the surface by the factor s , i.e., acting on the state by the scale transformation⁵⁷

$$f_s(\xi) = s\xi. \quad (7.36)$$

This illustrates how the singularity is resolved in yet another context. The string vertices are singular surfaces, since they induce gluing, i.e., a delta function interaction. This translates also to the singularity of the relevant surfaces in the oscillator formalism. The regularization that we are using resolves exactly this delta function. We present the effect of this regularization for the definition of a single state and for the interaction using the two- and the three-vertex in figure 7.

Having established the good formal properties of the regularization, the next task is to check whether it indeed works. The simplest possibility is to check numerically the descent relations. Indeed, a numerical evaluation of γ_{13} strongly suggests that it equals unity when using the above regularization. Next, we would like to have an expression describing the regularization in the continuous basis, where analytical results are usually easier to obtain. The problem with the level truncation regularization in the continuous basis can be traced to the manipulations performed in (7.27). There, we ended with a $\delta(0)$ that we had to regularize. However, some of the manipulations used to get to the final expression also used delta functions. It seems inconsistent to regularize one delta function and not the other. Indeed, the s -regularization is effectively performed by regularizing the identity operator inserted between various terms that contain oscillators,

$$\mathbf{1}_s \equiv \sum_{n=1}^{\infty} |n\rangle s^{2n} \langle n|. \quad (7.37)$$

Let us switch to the variable

$$\epsilon = 1 - s. \quad (7.38)$$

⁵⁷ Recall that ξ are the coordinates on the upper half plane.

Expressing $\mathbf{1}_{1-\epsilon}$ in the continuous basis gives

$$\mathbf{1}_{1-\epsilon} = \sum_{n=1}^{\infty} \int d\kappa d\kappa' |\kappa\rangle \langle \kappa| n\rangle (1-\epsilon)^{2n} \langle n| \kappa'\rangle \langle \kappa'| = \int d\kappa d\kappa' |\kappa\rangle \rho_s(\kappa, \kappa') \langle \kappa'|, \quad (7.39)$$

where we defined the regularized kernel

$$\rho_\epsilon(\kappa, \kappa') = \sum_{n=1}^{\infty} v_n^\kappa v_n^{\kappa'} (1-\epsilon)^{2n}. \quad (7.40)$$

Evaluating this kernel is neither simple nor illuminating. Thus, we simply state the final result,

$$\rho_\epsilon(\kappa, \kappa') = \frac{1}{\kappa\kappa' \sqrt{\mathcal{N}(\kappa)\mathcal{N}(\kappa')}} \left(\frac{e^{i\frac{\kappa-\kappa'}{2} \log \epsilon}}{B\left(\frac{i\kappa'}{2}, -\frac{i\kappa}{2}\right)} + \frac{e^{-i\frac{\kappa-\kappa'}{2} \log \epsilon}}{B\left(\frac{i\kappa}{2}, -\frac{i\kappa'}{2}\right)} \right), \quad (7.41)$$

where B is the Beta function. This expression is correct up to $\mathcal{O}(\epsilon)$ corrections. However, since the singularity is of the order of $\log(\epsilon)$, these corrections can be safely neglected. Using some residue calculus while consistently neglecting terms of the form $\epsilon^n \log(\epsilon)$, one can see that these kernels have some nice properties, such as

$$\int d\kappa' \rho_\epsilon(\kappa, \kappa') \rho_{\epsilon'}(\kappa', \kappa'') = \rho_{\epsilon+\epsilon'}(\kappa, \kappa''). \quad (7.42)$$

This may seem like a surprising identity. However, it is simply the (approximate) continuous basis form of the trivial identity

$$\mathbf{1}_s \mathbf{1}_{s'} = \mathbf{1}_{ss'}, \quad (7.43)$$

which is exact for arbitrary values of s, s' . It is important to note that (7.41) reduces to (7.28) in the limit $\kappa \rightarrow \kappa'$, with $\epsilon \sim l^{-1}$. Thus, (7.28) can be considered as the kernel (7.41) restricted to its diagonal and multiplied by a $\delta(\kappa - \kappa')$ factor. It is exactly this extra delta function factor that this regularization helps to resolve.

Using the result for $\rho_\epsilon(\kappa, \kappa')$ it was analytically proved in [156] that Schnabl's algebra holds in the continuous basis, without a central term, as it should. The nonlinear expressions for the inner product $\langle n|m \rangle$ ⁵⁸ were evaluated analytically as a power series in n, m and it was shown analytically that the lowest orders of the series vanish. More terms were shown to vanish using a numerical evaluation. Finally, the most potentially singular inner product, namely $\langle 1|1 \rangle$ was proven to equal unity. All that gives high credibility to the advocated regularization, thus allowing for the use of the oscillator basis as a practical and reliable formalism for string field theoretical calculations.

⁵⁸ Recall that for $n = 1$ and m an integer this is equivalent to the descent relations.

8 Superstring field theory

There are several versions of open covariant superstring field theory. All these theories are most naturally defined using the fermionized version of the superghosts [232]. We start this section by introducing the fermionization and the related concept of pictures in 8.1. Next, in 8.2 we describe the various superstring field theories. The recent results are described in 8.3, where a mapping of solutions between string field theories is studied and in 8.4 where analytical solutions of superstring field theories are described.

8.1 Superstring variables, pictures and conventions

In the RNS⁵⁹ formulation of the superstring, in addition to the world-sheet-scalar space-time-vector X^μ there is also a world-sheet-fermion space-time-vector ψ^μ . These fields are referred to as matter fields. Supersymmetry is manifest on the world-sheet. This results in the elevation of the conformal symmetry to a superconformal one. This symmetry is related to the (matter) energy-momentum tensor T_m (we work here in the $\alpha' = 2$ convention)⁶⁰,

$$T_m(z) = -\frac{1}{2}\partial X^\mu\partial X_\mu(z) - \frac{1}{2}\psi^\mu\partial\psi_\mu(z), \quad (8.1)$$

as well as to the superconformal (matter) generator G_m ,

$$G_m(z) = i\psi^\mu\partial X_\mu(z). \quad (8.2)$$

Fixing the superconformal symmetry introduces in addition to the fermionic bc ghosts with central charge $c_{bc} = -26$ also the bosonic $\beta\gamma$ superghosts with central charge $c_{\beta\gamma} = 11$. It is convenient to fermionize the superghosts. The first step in this direction consists of introducing a field ϕ in order to fermionize the superghost-number current as

$$\partial\phi(z) = \beta\gamma(z), \quad (8.3)$$

with the field ϕ satisfying

$$\phi(z)\phi(w) \sim -\log(z-w). \quad (8.4)$$

⁵⁹ We focus exclusively on the NS sector of the open superstring. The inclusion of the Ramond sector may also be possible, but is irrelevant to the results that we want to present here.

⁶⁰ Operators inserted at the same point carry implicit normal ordering.

One can see that the superghost number is shifted by q units under the action of the operator $e^{q\phi}$. Thus, the β, γ operators should be related to the operators $e^{\pm\phi}$. We define the operator $e^{q\phi}$ to carry picture number q . From the discussion above it may seem that this is just the superghost number. However, there is some freedom in the definition that we are about to explore. As a result, picture number turns out to be an independent quantum number.

One can use the Sugawara construction [233] and define the energy momentum tensor of the superghost current,

$$T_\phi(z) = -\frac{1}{2}\partial\phi\partial\phi - \partial^2\phi. \quad (8.5)$$

This definition gives the correct OPE for the energy momentum tensor and the current. Yet, it defines a system with central charge $c_\phi = 13$, so this cannot be the whole story. Also, one can check that the operators $e^{\pm\phi}$ anticommute, while the β, γ system is bosonic. The OPE's of $e^{\pm\phi}$ are also not quite what is needed. To compensate for the lost central charge and the wrong statistics one appends to ϕ the conjugate fermions η, ξ with conformal dimensions $(1, 0)$ respectively. The energy momentum tensor of this system is given by

$$T_{\eta\xi} = -\eta\partial\xi. \quad (8.6)$$

The complete fermionization formulas then read⁶¹,

$$\beta = e^{-\phi}\partial\xi, \quad \gamma = \eta e^\phi. \quad (8.7)$$

Now, we have to assign picture number to the new variables. We want all fields not in the fermionized ghost sector, including the original β, γ , to have zero picture number. This would imply the neutrality of the BRST charge Q . Thus, we assign the picture numbers as $n_p(\xi) = 1$, $n_p(\eta) = -1$. This is indeed a new quantum number. We write down for convenience the picture number, the ghost number, the conformal weight and the parity of some fields including the BRST current J_B , its “inverse” P and the picture changing operators \mathcal{X}, Y (introduced below) in table 1. The normal ordering of some of the fields

⁶¹ A co-cycle is implicitly assumed when writing $e^{\pm\phi}$, in order for these operators to anticommute also with the other anti-commuting variables of the theory.

operator	h	n_g	n_p	parity
∂X^μ	1	0	0	0
ψ^μ	$\frac{1}{2}$	0	0	1
b	2	-1	0	1
c	-1	1	0	1
η	1	1	-1	1
ξ	0	-1	1	1
$e^{q\phi}$	$-\frac{q(q+2)}{2}$	0	q	$q \bmod 2$
β	$\frac{3}{2}$	-1	0	0
γ	$-\frac{1}{2}$	1	0	0
J_B	1	1	0	1
P	0	-1	0	1
\mathcal{X}	0	0	1	0
Y	0	0	-1	0

Table 1

The conformal weight h , ghost number n_g , picture number n_p and parity of the fields relevant to us.

is given explicitly by⁶²,

$$\partial X^\mu(z)\partial X^\nu(w) = : \partial X^\mu(z)\partial X^\nu(w) : - \frac{\eta^{\mu\nu}}{(z-w)^2}, \quad (8.8a)$$

$$\psi^\mu(z)\psi^\nu(w) = : \psi^\mu(z)\psi^\nu(w) : + \frac{\eta^{\mu\nu}}{z-w}, \quad (8.8b)$$

$$b(z)c(w) = : b(z)c(w) : + \frac{1}{z-w}, \quad (8.8c)$$

$$\beta(z)\gamma(w) = : \beta(z)\gamma(w) : - \frac{1}{z-w}, \quad (8.8d)$$

$$\eta(z)\xi(w) = : \eta(z)\xi(w) : + \frac{1}{z-w}, \quad (8.8e)$$

$$e^{a\phi}(z)e^{b\phi}(w) = \frac{: e^{a\phi}(z)e^{b\phi}(w) :}{(z-w)^{ab}}. \quad (8.8f)$$

In the fermionization equations (8.7) the field ξ appears only through its derivative. That is, the zero mode ξ_0 is not needed in order to describe the β, γ system. Its inclusion would double the Hilbert space, since it is a two-level fermionic variable. It is common to refer to the Hilbert space without

⁶²Note that the order of terms in (8.7) is important for the consistency of (8.8e) and (8.8f) with (8.8d) (recall that $e^{\pm\phi}$ are anti-commuting).

the zero mode as the “small Hilbert space” and to the doubled space with the ξ_0 included as the “large Hilbert space”. In the large Hilbert space CFT correlators are normalized as

$$\left\langle \xi \partial^2 c \partial c c e^{-2\phi} \right\rangle_{\text{Large}} = 2. \quad (8.9)$$

The correlator is non-zero only for ghost number $n_g = 2$ and picture number $n_p = -1$. Since the small Hilbert space is defined by the absence of the zero mode ξ_0 , the factor of ξ is omitted in defining correlators,

$$\left\langle \partial^2 c \partial c c e^{-2\phi} \right\rangle_{\text{Small}} = 2. \quad (8.10)$$

Now, the total picture number $n_p = -2$ is needed for a non-trivial result and the ghost number should be $n_g = 3$, as in the bosonic case⁶³.

The existence of the superconformal algebra implies that a BRST charge Q should be defined. The BRST charge of a constrained system that forms a (super-)Lie algebra contains two types of terms. First, the ghost multiplied by the constraints are added. To them one has to add couplings of two ghosts and one antighost multiplied by the algebra structure constants. All that can be nicely summed up, writing the BRST charge as an integral of the BRST current J_B ,

$$Q = \frac{1}{2\pi i} \oint dz J_B(z), \quad (8.11)$$

which is defined up to a total derivative. In the case at hand, the set of constraints is encoded in the (matter) energy momentum tensor T_m , whose ghost and anti-ghost are the c and b respectively and by the superconformal (matter) generator G_m , with γ and β as the ghost and antighost. Thus, the first part of the BRST current is

$$J_B^{(0)}(z) = cT_m + \gamma G_m. \quad (8.12)$$

Adding the terms that encode the super-Virasoro algebra one gets

$$\begin{aligned} J_B(z) &= cT_m + \gamma G_m + c\partial cb - \frac{c}{2}(3\beta\partial\gamma + \partial\beta\gamma) - b\gamma^2 \\ &= c(T_m + T_{\eta\xi} + T_\phi + \partial cb) + \eta e^\phi G_m - b\eta\partial\eta e^{2\phi}. \end{aligned} \quad (8.13)$$

In the second line we converted the expression to the more convenient variables ϕ, η, ξ (recall (8.7)) and the energy-momentum tensors of the fermionized ghost systems are given by (8.5) and (8.6)⁶⁴.

⁶³ One way to think of it is to say that the left vacuum in (8.10) equals the left vacuum of (8.9), with ξ_0 acting on it.

⁶⁴ Note that normal ordering is implicit in both lines of (8.13). In order to convert, one has to use (8.8) in order to first undo the $\beta\gamma$ normal ordering and then introduce the ones of the ϕ, η, ξ systems.

An interesting property of the BRST charge in this formulation is that it defines a trivial cohomology in the large Hilbert space⁶⁵. This can be seen by defining the operator

$$P(z) = -c\xi\partial\xi e^{-2\phi}(z), \quad (8.14)$$

which, by direct inspection obeys

$$[Q, P(z)] = 1. \quad (8.15)$$

Now, let $\mathcal{O}(0)|0\rangle$ be a Q -closed state, then it is exact, since the above imply that,

$$\mathcal{O}(0)|0\rangle = Q\left(P(0)\mathcal{O}(0)|0\rangle\right). \quad (8.16)$$

One may worry that singularities may appear in the OPE of P and \mathcal{O} . However, in such a case one may modify (8.16) to

$$\mathcal{O}(0)|0\rangle = Q\left(\frac{1}{2\pi i} \oint \frac{dz}{z} P(z)\mathcal{O}(0)|0\rangle\right). \quad (8.17)$$

No singularities can remain in the above equation and its proof is as straightforward.

An important feature of superstring theory is the abundance of vertex operators adequate for the description of physical states. In fact, even for the bosonic case there is more than one vertex operator describing a given state. There, there are two variants, the integrated vertex operator⁶⁶ V_{bos} and the unintegrated vertex operator \hat{V}_{bos} . These vertex operators have to obey

$$[Q, V_{\text{bos}}] = \partial\hat{V}_{\text{bos}}, \quad (8.18)$$

$$[Q, \hat{V}_{\text{bos}}] = 0. \quad (8.19)$$

The second equation, which implies that the unintegrated vertex is closed, follows of course from the first one. Integrating the total derivative in the r.h.s of (8.18) implies that the vertex is closed in this form as well. Moreover, it follows from (8.18) that adding a (physically irrelevant) total derivative to V_{bos} corresponds to adding a (physically irrelevant) exact term to \hat{V}_{bos} . A direct evaluation gives in the bosonic case⁶⁷,

$$\hat{V}_{\text{bos}} = cV_{\text{bos}}. \quad (8.20)$$

⁶⁵ This fact is proved in [234], by constructing a similarity transformation that generates J_B from the last term of (8.13). This also serves as an elegant proof of the nilpotence of Q .

⁶⁶ By “integrated vertex operator” we actually refer to the integrand of the vertex.

⁶⁷ This is a familiar expression. However, the relation (8.18) is the fundamental one, as it can be used even in formulations with no b, c system, such as the pure-spinor formulation of superstring theory [235,236].

It is not important which form of the vertex is used, as long as there are exactly three unintegrated vertex operators to saturate the three (fermionic) zero modes of the c ghost. These zero modes are in fact the reason behind the redundancy. Their fermionic nature is the reason for having exactly two forms of each vertex operator.

It can be read from (8.10) that (on the plane) the superghosts have two zero modes (picture number) and one may expect that again this will lead to a redundant description of the vertex operators. However, as the superghosts are bosonic, one may expect an infinite redundancy instead of the two-fold redundancy related to the bc system⁶⁸. The redundancy of the bc system also exists of course, but one can treat it independently of that of the superghosts⁶⁹. Thus, each state of the superstring should correspond to one integrated vertex operator V^p and one unintegrated vertex operator \hat{V}^p at every integer value p of the picture number⁷⁰. These vertices should obey (8.18) and (8.19) separately for every values of p ,

$$[Q, V^p] = \partial \hat{V}^p, \quad (8.21)$$

$$[Q, \hat{V}^p] = 0. \quad (8.22)$$

The relation (8.20) holds now only for specific values of picture number.

Given an unintegrated vertex \hat{V}^p one can construct from it the vertex \hat{V}^{p+1} using the following definition

$$\hat{V}^{p+1} = [Q, \xi \hat{V}^p]. \quad (8.23)$$

While the new vertex operator may seem to be exact, it is such only in the

⁶⁸ In the bosonic case, the most natural description in some sense is the one where the zero mode is fixed, namely the unintegrated vertex. This is achieved by multiplication by c . In order to fix a bosonic coordinate, as we have here, one has to insert a factor of $\delta(\gamma)$. Luckily, this delta function has a simple expression after fermionization, namely $e^{-\phi}$. For this reason it is customary to refer to the -1 picture as being a natural one. One still needs, of course, also the representations of the vertex operators at other pictures.

⁶⁹ This is true for the plane. For higher genera it is no longer possible to disentangle the bosonic and fermionic zero modes and one should study the supermoduli carefully. The endeavor of D'Hoker and Phong concentrated around this issue. In particular, they showed [237] that the picture changing operators, introduced below, should be modified at genus two (see [238,239,240] for interesting recent progress and [241] for a recent review). One may hope that the framework of string field theory can be used as a simple alternative description to the complicated study of supermoduli spaces. For that end, a reliable formulation of superstring field theory is needed, in which the Ramond sector is also included.

⁷⁰ Recall that we describe the Neveu-Schwarz sector. In the Ramond sector the allowed picture numbers p are half-integers.

large Hilbert space, while being only closed in the small Hilbert space, which is the relevant one. This also shows that the only relevant part of ξ in this definition is its zero mode ξ_0 , since the rest gives rise to a truly exact part and so decouples from all calculations of scattering amplitudes. Using (8.19) one can deduce that (8.23) can equivalently be written as

$$\hat{V}^{p+1} = \mathcal{X}\hat{V}^p, \quad (8.24)$$

where the picture changing operator \mathcal{X} is given by⁷¹,

$$\mathcal{X} = [Q, \xi] = c\partial\xi + e^\phi G_m + e^{2\phi} b\partial\eta + \partial(e^{2\phi} b\eta). \quad (8.25)$$

For the integrated vertex we define

$$V^{p+1} = [Q, \xi V^p] + \partial(\xi \hat{V}^p). \quad (8.26)$$

The second, total derivative term does not contribute after integration. It should nevertheless be added both for consistency with (8.21) as well as for obtaining a vertex that resides in the small Hilbert space. This equation can be rewritten as

$$V^{p+1} = \mathcal{X}V^p + \partial\xi \hat{V}^p. \quad (8.27)$$

This procedure can be repeated ad infinitum. In the calculation of scattering amplitudes any set of representatives of the scattered states can be used, provided the total picture number is such that the picture and ghost numbers in (8.10) are saturated [232]. We illustrate the equivalence of different picture number distributions in figure 8. One can still wonder whether the repeated use of the picture changing scheme can produce singularities⁷², as it is easy to check that the OPE of \mathcal{X} with itself has a double pole. In fact this can happen. These singularities originate from the non-zero mode part of ξ that we add to ξ_0 in order to write it as a local field. Thus, such singularities correspond to terms that are exact even in the small Hilbert space⁷³. Allowing for an ϵ dependence to be left in the expressions for the vertex operators in (8.23), (8.24), (8.26) and (8.27), leads to consistent relations among the vertices, with the only problem being their non-locality. The singular terms decouple from scattering amplitudes and can simply be dropped out from the definition of the picture-changed vertex operators. Taking then the limit $\epsilon \rightarrow 0$ returns a consistent local vertex.

⁷¹The picture changing operator is usually represented by X . However, since we tend to omit the space-time index μ from the scalars X^μ , we follow the example of [44] and use \mathcal{X} instead.

⁷²Note, that the products in (8.23), (8.24), (8.26), (8.27) are simple OPE's, without any normal ordering. Had we insisted on normal ordering in these expressions, the argument described in figure 8 would have been invalidated.

⁷³For the integrated vertex they should be exact only after integration. In particular, singularities multiplying total derivatives may pop-up in this case.




Fig. 8. The equivalence of different ways to distribute picture number for an arbitrary scattering amplitude in six stages, from top to bottom:

1. We start from an arbitrary expectation value, with an arbitrary number of vertex operators (green) each one carries its own arbitrary picture number.
2. The amplitude can be evaluated also in the large Hilbert space. For that end one only has to introduce the zero mode ξ_0 . This corresponds to an insertion of the field ξ in an arbitrary place. We choose to insert it at the site of a specific vertex.
3. We now represent another vertex operator using the vertex at a picture number lower by one (red), as in (8.23). Integrated vertex operators work in the same way, since the total derivative term in (8.26) does not contribute.
4. The integration contour of the BRST current is now deformed so as to circle (with opposite orientation) the other vertices. This results in many terms, such as the first one depicted, where the current circles around a vertex without a ξ insertion. As all the vertices are closed (up to total derivatives), these terms are nullified. The only surviving term is the one where the BRST current circles the ξ insertion. An extra $(-)$ sign, coming from reversing the formal Grassmann ordering of J_B and ξ , is represented by inverting back the orientation of the integration contour. In fact, there are some more minus signs coming from considering formal Grassmann ordering, in all the stages described in this figure, but they all cancel out.
5. The vertex with the ξ insertion, surrounded by the BRST current is replaced by a vertex with picture number higher by one unit (blue).
6. The final result can be again evaluated in the small Hilbert space, by omitting the ξ insertion. The final expression is identical to the initial one, except that one vertex was “red-shifted” and another one was “blue-shifted”.

This prescription for removing the singularities amounts essentially to a sort of normal ordering. However, while this is always the usual normal ordering in (8.23) and (8.26), where only the ξ operator is inserted, it will not necessarily coincide with the usual normal ordering upon usage of (8.24) or (8.27). Moreover, these last equations are the ones that are easier to generalize for the purpose of lowering the picture, as we describe next. Again, for the Y operator, normal ordering and neglecting singular terms in the expansion may

not coincide.

Picture number can be decreased using the inverse picture changing operator,

$$Y = c\partial\xi e^{-2\phi}. \quad (8.28)$$

The operator Y is the inverse of \mathcal{X} in the sense that

$$\lim_{z \rightarrow w} \mathcal{X}(z)Y(w) = 1. \quad (8.29)$$

This relation implies,

$$\hat{V}^{p-1} = Y\hat{V}^p. \quad (8.30)$$

One can check that like \mathcal{X} , the inverse picture changing operator Y is closed but not exact. Thus, this transformation maps a (closed non-exact) vertex operator into another (closed non-exact) one, as it should⁷⁴. An interesting property of Y is that it obeys,

$$P = \xi Y. \quad (8.31)$$

We shall use this fact in 8.4.

For the integrated vertex we have to find, by analogy with (8.27), an operator Υ obeying

$$[Q, \Upsilon] = \partial Y. \quad (8.32)$$

Then, the definition

$$V^{p-1} = YV^p + \Upsilon\hat{V}^p, \quad (8.33)$$

is consistent with (8.21). Indeed, Υ is easily found to be,

$$\Upsilon = \partial\xi e^{-2\phi}. \quad (8.34)$$

A (local) primitive for Z does not exist even in the large Hilbert space. Nonetheless, one can further enlarge the Hilbert space by including such a primitive. Inverse picture changing would then be implemented using equations analogous to (8.23) and (8.26) with $p+1 \rightarrow p-1$ and the primitive of Υ replacing ξ . The arguments presented in figure 8 generalize immediately for the picture lowering procedure.

The last important feature of the RNS superstring that we have to recall is the GSO projection. This projection is based upon the operator F called the “world-sheet fermion number”, which counts the number of occurrences at a given vertex operator of the fields ψ^μ, γ, β . In the fermionized variables the fermion number of the superghosts is fully given in terms of the ϕ field,

$$[F, e^{l\phi}] = le^{l\phi}. \quad (8.35)$$

⁷⁴The remarks regarding the singularities of \mathcal{X} , hold also for Y .

The $SL(2)$ vacuum of the NS sector is defined to be F -odd,

$$F|0\rangle = -|0\rangle. \quad (8.36)$$

Since Q commutes with F , one can consider separately the sectors in the theory with $e^{i\pi F}$ being zero or unity. The former is referred to as being $GSO(+)$ while the later is $GSO(-)$. String theories contain some combinations of the $NS\pm$ and $R\pm$ sectors, subject to some consistency conditions, such as modular invariance and closure of the OPE algebra⁷⁵. Open superstring theory on a BPS D-brane has only the $(+)$ sectors. A non-BPS D-brane has both $(+)$ and $(-)$ sectors, while a $D-\bar{D}$ system has two sectors of each kind tensored with appropriate Chan-Paton factors.

8.1.1 Example: The tachyon vertex operators in various pictures

We illustrate the above discussion using the simplest vertex, namely that of the tachyon field. This vertex is often ignored, since it belongs to the $GSO(-)$ sector and so does not exist on a BPS D-brane. It does exist on a non-BPS D-branes and on other systems. Superstring field theory should be the perfect framework to study its condensation. At the end of this subsection we also describe the “ $GSO(+)$ tachyon vertex operator”.

In the natural (-1) picture, the integrated tachyon vertex is given by

$$V^{-1} = e^{-\phi} e^{ik \cdot X}. \quad (8.37)$$

Requiring that the vertex has weight one implies that

$$k^2 = 1. \quad (8.38)$$

Hence, this vertex describes a tachyon, as stated⁷⁶. Given (8.38), one obtains

$$\hat{V}^{-1} = c e^{-\phi} e^{ik \cdot X} = c V^{-1}, \quad (8.39)$$

as in the bosonic case.

Using (8.26) the picture is increased by one unit,

$$V^0 = -k \cdot \psi e^{ik \cdot X}, \quad (8.40)$$

⁷⁵ Closed string theories, which we do not consider here, can have different left and right sectors, modulo the level-matching condition.

⁷⁶ We pretend that the field X is analytic. In reality, the field X is not analytic and the vertex operator should be restricted to the boundary, where boundary normal ordering should be applied. This introduces some constants relative to the discussion here.

while for the unintegrated vertex one gets,

$$\hat{V}^0 = -(ck \cdot \psi + \eta e^\phi) e^{ik \cdot X}. \quad (8.41)$$

We see that

$$\hat{V}^0 \neq cV^0. \quad (8.42)$$

Moreover, the second term in (8.41) is what one intuitively associates with the tachyon vertex, since

$$\eta e^\phi(0) |k\rangle = \gamma(0) |k\rangle = \gamma_{\frac{1}{2}} |k\rangle. \quad (8.43)$$

Continuing this way, we can evaluate V^1 . In this case one gets among several regular terms, also a singular one,

$$V_{\text{sing}}^1 = \frac{1}{\epsilon} \partial(-e^\phi e^{ik \cdot X}). \quad (8.44)$$

However, as this term is a total derivative it can be safely neglected. For the unintegrated vertex one gets again a singular term, which equals Q acting on the expression inside the parentheses in (8.44), in accord with (8.21). A singular term is also produced upon going in the other direction (using inverse picture changing),

$$V_{\text{sing}}^{-2} = \frac{1}{\epsilon} \left(\partial(c\partial\xi e^{-3\phi} e^{ik \cdot X}) - Q(\partial\xi e^{-3\phi} e^{ik \cdot X}) \right). \quad (8.45)$$

For the singular part of \hat{V}^{-2} one gets Q acting on the expression inside the first parentheses. Again, the expressions are consistent and the singular terms can be safely dropped.

Examining (8.43) one can easily think of a “more tachyonic tachyon” living in the GSO(+) sector, namely

$$c(0) |k\rangle = c_1 |k\rangle. \quad (8.46)$$

For this operator to have zero conformal weight, as is appropriate for an unintegrated vertex operator, one has to demand $k^2 = 2$. However, even for these values of the momenta the operator fails to be closed. Thus, as is well known, there is no GSO(+) tachyon. The operator (8.46) clearly carries a zero picture. Changing the picture of an operator is a well defined procedure only for vertex operators. Still, one may ignore that and try to find the form of this operator in the natural picture, by acting on it with the inverse picture changing operator. The result is zero and there is no other operator with the same quantum numbers that can be added to it in order to remedy this result. This is not a problem, since this is not a genuine vertex operator. One may consider this operator as a peculiarity of working with zero picture number. Despite the above, this operator emerged in some recent developments, described in 8.4.

8.2 Introducing superstring field theories

The original proposal for open superstring field theory by Witten [95] is almost a straightforward generalization of the construction for the bosonic case. The main difference is the appearance of picture number, which implies that the definitions of integration and star product should be modified, in order to saturate the -2 picture number in (8.10). Also, the string field should be assigned a fixed picture number⁷⁷. Witten suggested to modify the bosonic action to

$$S = - \int \left(\frac{1}{2} \Psi \star Q \Psi + \mathcal{X} \frac{1}{3} \Psi \star \Psi \star \Psi \right), \quad (8.47)$$

where the picture changing operator \mathcal{X} is inserted at the common string midpoint. The string fields in this scheme carry picture number $n_p = -1$.

This version of superstring field theory has some problems. The picture changing operators appear in scattering amplitudes and may be inserted at the same point. This, however, produces singularities [242,243] (starting already at tree level for the four point function) that render the theory erroneous. A related issue is the collision of picture changing operators in the derivation of the g_o^2 -order gauge transformation of the string field. One can try to solve these divergences by introducing counter-terms into the theory. Wendt [242] gave the form of the fourth order counter-term that resolves both problems. Nevertheless, as also noted by him, new divergences arise at the next order. It may be possible that an infinite set of counter-terms exists that can regularize the theory (at least classically, i.e., considering only tree level interactions) to all orders. To the best of our knowledge, this avenue was never pursued. Also, as mentioned in section 3.1, Witten's theory fails to reproduce the expected results for the tachyon potential, at least for the first few levels in the level truncation scheme (without the introduction of counter-terms).

A modified version of cubic superstring field theory was constructed, using the double-step inverse picture changing operator Y_{-2} [149,150,151]. This operator is required to be closed, non-exact and obey

$$\lim_{z \rightarrow w} \mathcal{X}(z) Y_{-2}(w) = Y(w). \quad (8.48)$$

All its quantum numbers, other than the picture number, should vanish. One

⁷⁷ One could imagine constructing a string field theory without restricting the picture number. However, as vertices with different picture numbers are equivalent, a gauge symmetry should be introduced in order to avoid multiple counting. Another related option could be to start with a fixed picture number and add other picture numbers as part of the gauge fixing procedure, analogously to the way different ghost numbers appear in the bosonic case. We are not aware of any construction along these lines.

can define Y_{-2} either as a chiral or as a non-chiral (non-local) operator,

$$Y_{-2}^{chir}(z) = -e^{-2\phi(z)} - \frac{i}{5}c\partial\xi e^{-3\phi}\psi_\mu\partial X^\mu(z), \quad (8.49)$$

$$Y_{-2}^{non}(z, \bar{z}) = Y(z)Y(\bar{z}), \quad (8.50)$$

where the doubling trick is used. The non-local version is not well defined on the boundary, where picture changing operators are usually inserted. It is, nonetheless, useful for superstring field theory, in which the operator is inserted at the string mid-point, $z = i$. The Y_{-2} insertion saturates the required picture number. Hence, there is no need for further picture insertions and the string field should be assigned a zero picture number. The action reads,

$$S = - \int Y_{-2} \left(\frac{1}{2} \Psi \star Q\Psi + \frac{1}{3} \Psi \star \Psi \star \Psi \right). \quad (8.51)$$

The Y_{-2} operator can be absorbed in a redefinition of the integral. The equation of motion derived from this action is

$$Y_{-2} \left(Q\Psi + \Psi \star \Psi \right) = 0. \quad (8.52)$$

This modified form of the action does not suffer from the contact term problems. For the gauge symmetry it is clear, since the gauge parameters carry now zero picture number and picture changing operators do not occur in the gauge transformation. In the expressions for scattering amplitudes Y_{-2} will still appear. However, while each vertex now carries a factor of Y_{-2} , the propagator carries a factor of “ $\frac{1}{Y_{-2}}$ ”. These factors cancel out (at least for tree amplitudes) leading to the expected results [49].

Several objections have been raised to this modified action. One objection is that all picture changing operators have non-trivial kernels. These kernels are, however, of a somewhat exotic nature, containing only states which are localized at the string mid-point. Thus, it is not clear if this is really a problem. Also, as mentioned in footnote 69, picture changing operators should be modified at higher loop order. It may be hard to believe that one can use the picture changing operators that are adequate for the disk for describing scattering processes to all orders.

Another objection is related to the existence of two versions of Y_{-2} . The two candidates for Y_{-2} are not a-priori equivalent and it is not clear, which one should be used. Some properties of the theories for the two possible choices of Y_{-2} were studied using level truncation in [244]. The theories were found to differ when truncated to the massless level. It was claimed that the non-chiral operator is the more promising one, since the theory with the chiral operator fails to reproduce the Maxwell equations for the level zero component. Although this result can be modified when more modes are added. It was

also found out there that supersymmetry is differently realized in the two theories. Two versions of supersymmetry transformations were found for both theories⁷⁸. Though for the chiral theory, one of them suffers from singularities when the supersymmetry transformation is iterated. Later, it was noticed that the theory with the chiral operator does not respect the expected twist symmetry. The above may suggest that the theory with the non-chiral operator is the more promising one. At any rate, the ultimate test would be to check which one, if at all, reproduces the correct results, namely correct (on-shell) scattering amplitudes and the expected results regarding Sen's conjectures. Due to the criticism on the chiral theory, it was studied much less than the non-chiral theory, both in level truncation (as described in 3.1) as well as in the recent developments described below.

Another form of open superstring field theory was developed by Berkovits [144,245,246,247]. Unlike the cubic string field theories discussed so far, this theory is non-polynomial⁷⁹ and looks like a generalization of the WZW theory. This theory was constructed using the language of $N = 2$ and topological $N = 4$ string theory [253]. Nonetheless, it can be presented (and used) without going into the details of this construction.

One novel characteristic of this theory is that it is defined in the large Hilbert space. A new gauge symmetry takes care of the extra degrees of freedom that emerge due to the use of this space. Recall that in the large Hilbert space the picture number should equal minus one for a non-trivial result. Thus, it is enough to use η_0 , which acts non-trivially in the large Hilbert space, in order to saturate it. The string field can be assigned zero picture number and there is no need to use picture changing operators⁸⁰.

The action is given by

$$S = \frac{1}{2g_o^2} \int \left(e^{-\Phi} Q(e^\Phi) e^{-\Phi} \eta_0(e^\Phi) - \int_0^1 dt e^{-t\Phi} \partial_t e^{t\Phi} [e^{-t\Phi} \eta_0 e^{t\Phi}, e^{-t\Phi} Q e^{t\Phi}] \right), \quad (8.53)$$

where Φ is the string field. In the second term Φ is written as $e^{-t\Phi} \partial_t e^{t\Phi}$ in order to bring the action to a form resembling the WZW action. The equation

⁷⁸ The kernel of $Y_{-2}(i)$ was interpreted as an extra gauge symmetry. The two realizations of supersymmetry differ by picture changing operators inserted at $\pm i$ and so are presumably gauge equivalent.

⁷⁹ This theory still uses only the cubic Witten vertex, unlike the counter-terms mentioned above in the context of Witten's superstring field theory or the higher vertices that are present in closed string field theory [248], open-closed string field theory [249,250] and heterotic string field theory [251,252].

⁸⁰ From the $N = 2$ point of view η_0 and Q should be treated on an equal footing, since they are the two superconformal generators of this theory.

of motion derived from the action is

$$\eta_0(e^{-\Phi} Q e^{\Phi}) = 0. \quad (8.54)$$

This equation states that the expression inside the parentheses lies in the small Hilbert space. This can be achieved trivially by taking Φ to lie in the small Hilbert space. However, as mentioned above, there is a new gauge symmetry that removes exactly these degrees of freedom. The linearized gauge transformation is most easily given in terms of the transformation of $G \equiv e^{\Phi}$,

$$\delta G = -(Q\tilde{\Lambda})G + G(\eta_0\Lambda). \quad (8.55)$$

This can be exponentiated to give the finite gauge transformation,

$$G \rightarrow e^{-Q\tilde{\Lambda}} G e^{\eta_0\Lambda}. \quad (8.56)$$

The quantum numbers of the gauge string fields are

$$n_g(\tilde{\Lambda}) = -1, \quad n_p(\tilde{\Lambda}) = 0, \quad n_g(\Lambda) = -1, \quad n_p(\Lambda) = 1. \quad (8.57)$$

The first gauge field in (8.55)⁸¹ is analogous to the usual gauge field. The second gauge field can be used to remove any variation $\delta G = G\delta\Phi$ with Φ entirely within the small Hilbert space, by defining $\Lambda = \xi_0\Phi$, which results in $\delta G = G\eta_0\Lambda$. Introducing ξ_0 defines two copies of the small Hilbert space. Setting one of them to zero leaves us with exactly the same amount of degrees of freedom as without using ξ_0 at all. Thus, the action passes at least the preliminary test of having the correct amount of degrees of freedom. It was further described in [144] how to write a superstring field theory action that is manifestly supersymmetric after reduction to four space-time dimensions. The action passed also the important tests of describing correctly scattering amplitudes [254]. Furthermore, as described in section 3.1.4, it also reproduces the results expected according to Sen's conjectures.

Finally, we turn to describe the inclusion of the GSO(-) sector in the string field theories described above. Since the states contained in this sector differ from those in the GSO(+) sector, it is clear that a new string field should be added to describe them. The half-integer conformal weights of the GSO(-) states imply that the cyclicity property (2.15) should be modified. Let $A_{1,2}$ be two string fields with half-integer weights, then their cyclicity property reads⁸²,

$$\int A_1 \star A_2 = -(-1)^{A_1 A_2} \int A_2 \star A_1. \quad (8.58)$$

⁸¹ Many papers use a convention without the minus sign in front of this term.

⁸² If only one of them has half-integer conformal weight then the integral gives zero.

A related issue is that the $\text{GSO}(-)$ sector string field has the opposite Grassmann character as compared to the string field from the $\text{GSO}(+)$ sector⁸³. In order to solve both problems and treat the string fields from both sectors collectively one should introduce the so called “internal Chan-Paton indices” [145]. These are simply some Pauli matrices acting on the internal space that consists of the two GSO sectors. The exact prescription depends on the theory at hand⁸⁴. For Berkovits’ theory (on the non-BPS D-brane, where both sectors exist) one has to write,

$$\Phi = \Phi_+ \otimes \mathbf{1} + \Phi_- \otimes \sigma_1, \quad (8.59)$$

where the subscripts \pm refer to the GSO sector in which the string field resides. One should also redefine the operators Q and η_0 according to

$$Q \Rightarrow Q \otimes \sigma_3, \quad \eta_0 \Rightarrow \eta_0 \otimes \sigma_3, \quad (8.60)$$

and redefine the integral so as to include also a trace over the matrices of this internal space. Since $\text{tr}(\mathbf{1}) = 2$, the coefficients in front of the action should be divided by two. The gauge string fields are modified according to

$$\Lambda = \Lambda_+ \otimes \sigma_3 + \Lambda_- \otimes (i\sigma_2), \quad \tilde{\Lambda} = \tilde{\Lambda}_+ \otimes \sigma_3 + \tilde{\Lambda}_- \otimes (i\sigma_2). \quad (8.61)$$

For the cubic theories the string field is given by [255],

$$\Psi = \Psi_+ \otimes \sigma_3 + \Psi_- \otimes (i\sigma_2), \quad (8.62)$$

while the gauge string field is given by

$$\Lambda = \Lambda_+ \otimes \mathbf{1} + \Lambda_- \otimes \sigma_1. \quad (8.63)$$

The only other modification required is the assignment,

$$Y_{-2} \Rightarrow Y_{-2} \otimes \sigma_3. \quad (8.64)$$

The description of general brane configurations is almost as straightforward. We describe the D- $\bar{\text{D}}$ system in the non-polynomial theory for simplicity. In this case, one has to introduce also the proper Chan-Paton factors, each one with the appropriate type of string field. In the case at hand, the Chan-Paton

⁸³ That is, the $\text{GSO}(-)$ string field is odd rather than even in Berkovits’ formalism and even rather than odd in the formalisms based on Witten’s theory.

⁸⁴ This should be expected. While the $\text{GSO}(+)$ string field is commuting and of ghost number zero in Berkovits’ theory, it is of ghost number one and anti-commuting in the modified cubic theories. This dictates different algebraic properties to be satisfied upon the inclusion of the $\text{GSO}(-)$ sector. One might expect that the representation of the string field in one theory resembles that of the gauge field in the other one. This is indeed the case.

matrices $\mathbf{1}, \sigma_3$ support GSO(+) fields, while σ_1 and σ_2 support GSO(-) fields. Thus, we expand the string field as

$$\Phi = \Phi_+^1 \otimes \mathbf{1} \otimes \mathbf{1} + \Phi_+^2 \otimes \mathbf{1} \otimes \sigma_3 + \Phi_-^1 \otimes \sigma_1 \otimes \sigma_1 + \Phi_-^2 \otimes \sigma_1 \otimes \sigma_2. \quad (8.65)$$

Here, the middle component in the tensor product describes the inner Chan-Paton space, while the right component describes the proper Chan-Paton factor. The action is further divided by two, to account for the trace that is now taken over external as well as internal Chan-Paton indices.

8.3 Mapping solutions among (super)string field theories

The first analytical solutions of superstring field theory were constructed independently by Erler [195] and by Okawa [196,256]. The solutions represent regular marginal deformations within the framework of Berkovits' string field theory. While it was recognized that the simplest solutions found do not obey the reality condition, gauge equivalent real solutions were also constructed.

It was found in these papers that despite the apparent distinction between Witten's bosonic cubic theory and Berkovits' supersymmetric non-polynomial theory, the solutions are quite similar to the analogous bosonic ones, found in [191,192]. One may wonder whether this is merely some funny coincidence. This is not the case. In fact, given a bosonic solution in a formal gauge form, it can be used to canonically define solutions of the supersymmetric theories.

The mapping of formal gauge solutions is straightforward in the case of the modified cubic superstring field theory. Since in this case the string field has the same ghost and picture numbers as in the bosonic case, one can simply use the same gauge field and derive from it the solution. This solution may differ from the bosonic one, since now the supersymmetric BRST operator (8.13) is used.

In particular, the solutions describing singular marginal deformations can be trivially mapped to this theory (using their formal-gauge representation). Moreover, the resulting solutions, despite being different from the bosonic ones remain x_0 -independent, since the condition (6.26) still holds and only formal properties of Q were used in the derivation of x_0 -independence⁸⁵.

In the case of Berkovits' theory the string field has the same (zero) picture numbers, but the ghost number (zero) is different from that of the cubic (bosonic and supersymmetric) theories. Moreover, in the non-polynomial theory the string field has to contain ξ_0 in order to be non-trivial. Thus, for

⁸⁵ Here, we use again the terminology adequate for the photon marginal deformation, but similar logic holds also for other exactly marginal deformations.

sending solutions of the cubic theories to the non-polynomial theory, we have to consider a mapping that results in

$$\Phi_1 = \xi \mathcal{O}(z) \Psi_1, \quad (8.66)$$

for some \mathcal{O} . The quantum numbers of ξ dictate that \mathcal{O} has to have zero ghost number and conformal weight and minus one unit of picture number. Hence, it is natural to identify it as the inverse picture changing operator Y and the map in this case should take the form,

$$\Phi_1 = P(z) \Psi_1, \quad (8.67)$$

for some z and where P is given by (8.31).

The choice of z in (8.67) is important. In [257] it was found that the desirable value for z is $\pm i$. This is a natural choice, since these points are invariant under star-multiplication. The canonical nature of these points was also used in the construction of the cubic superstring field theories as described above. An important property of $P(z)$ is (8.15), which implies that the cohomology of Q is empty in the large Hilbert space. In [257] the operator P was defined as a linear combination of $P(\pm i)$, such that the normalization of (8.15) does not change,

$$P = kP(i) + (1 - k)P(-i). \quad (8.68)$$

Then, the linearized map (8.67) was trivially extended to a map of the cubic superstring field theory to the non-polynomial one⁸⁶,

$$\Phi = P\Psi. \quad (8.69)$$

The operator P is nilpotent, as can be seen from the OPE,

$$P(z)P(w) = -\frac{z-w}{12}(cc'\xi\xi'\xi''\xi'''e^{-4\phi})(w) + \mathcal{O}((z-w)^2). \quad (8.70)$$

The location of the P insertion is invariant under the star-product. This fact implies that a state Ψ carrying a P insertion but no other components with support at $\pm i$, is nilpotent with respect to the star-product. The above implies

$$e^{t\Phi} = 1 + tP\Psi. \quad (8.71)$$

For solutions of the equation of motion, the map (8.69) can be inverted. In fact, all the solutions of the equation of motion of the non-polynomial theory (8.54) can be mapped to formal gauge solutions of the cubic theory by [247,258],

$$\Psi = G^{-1}QG, \quad (8.72)$$

⁸⁶The map (8.69) is not adequate for mapping bosonic solutions to the non-polynomial theory. However, bosonic solutions that are given in a gauge form can be (trivially) mapped to the cubic theory and then to the non-polynomial one.

as (8.54) implies that Ψ lives in the small Hilbert space. The “gauge field” G is formal, since it lives in the large Hilbert space. Composing the two maps one gets the identity transformation,

$$\Psi' = G^{-1}QG = (1 - P\Psi)(\Psi - PQ\Psi) = (1 - P\Psi)(\Psi + P\Psi\Psi) = \Psi. \quad (8.73)$$

Composing the maps in the other order results in a solution of the non-polynomial theory, which is gauge equivalent to the original Φ . More generally it was proven in [257] that gauge orbits are mapped to gauge orbits under the action of both maps and all gauge orbits (of solutions) are accessible. This implies that the cohomologies around solutions agree in both theories. Moreover, it was found that the action of solutions is invariant under the maps⁸⁷, provided that the non-chiral choice is made for the Y_{-2} operator. All that seems to imply that at least on-shell the non-polynomial theory and the cubic theory with non-chiral Y_{-2} are equivalent⁸⁸. One can study some properties of a classical solution in one theory and then safely move the other theory to study some issues that are more transparent there.

Solutions of the cubic theory that are given in a formal gauge form (other than the one of (8.72)) can be written in such a form in the non-polynomial theory as well using,

$$\Lambda = \xi\Lambda_{bos}, \quad \tilde{\Lambda} = P\Lambda_{bos}, \quad (8.74)$$

where the insertion point of the ξ operator is of no importance. The formal gauge solution of the non-polynomial theory is then given by (recall (8.56)),

$$G = e^{-Q\tilde{\Lambda}}e^{\eta_0\Lambda}. \quad (8.75)$$

The solution one gets using (8.69) is identical to the one that is obtained by using (8.74) and (8.75).

The map (8.69) is easily generalized to include the cases of the non-BPS D-brane and of D-brane systems, provided we assign a factor of σ_3 also to P, ξ in the internal Chan-Paton space,

$$P \Rightarrow P \otimes \sigma_3, \quad \xi \Rightarrow \xi \otimes \sigma_3. \quad (8.76)$$

The mapping of on-shell gauge orbits to gauge orbits and the equality of the value of the action in the two theories carries over without any further modification upon imposing (8.76).

⁸⁷ For that end one has to regularize the map (8.69). Strictly speaking one should think of this map as a specific limit of maps with no support at $\pm i$. In the limiting maps the ξ and Y components of P approach the limit points at a different pace.

⁸⁸ This gives one more reason to prefer the theory with a non-chiral Y_{-2} over the theory with the chiral operator insertion.

8.4 Analytical solutions in superstring field theory

Following [195,196,256], more solutions were found. Marginal deformations with singular OPE's were studied in [158,259], within the non-polynomial theory. These works generalized the methods of [193,194] respectively. Since in the bosonic case these deformations are given in a formal gauge form, they could be written in the cubic theory immediately using the same gauge field. All that is needed to write them in the non-polynomial theory is either (8.69) or (8.74).

Currently we have only one bosonic solution at hand, other than the ones describing marginal deformations. This is Schnabl's solution, describing tachyon condensation. What happens when we consider its counterpart in superstring field theory⁸⁹ [158,161]? On the one hand, from the canonical nature of the map, one expects that the solution of the supersymmetric theory has similar physical content to the bosonic solution. On the other hand, Schnabl's solution describes tachyon condensation, while the solution in the supersymmetric theory lives, by construction⁹⁰, in the GSO(+) sector, where no tachyons are present. One may still conjecture that it describes the state without the original D-brane, despite the fact that no dynamical condensation process exist that connects these two solutions.

The most natural way to check the conjecture is by evaluating the action and calculating the cohomology around the solution. In [161], Erler studied the solution in the framework of the modified cubic theory. He found that the cohomology and action of the solution are the expected ones, namely, the cohomology vanishes and the action equals minus the tension of the original D-brane. For that end, Erler used the bosonic gauge field (5.5) with the supersymmetric BRST charge. This results in a closed form expression that differs from the bosonic solution (5.22) only slightly,

$$\Psi = \Psi_{bos} + B\gamma^2(0)|0\rangle = \Psi_{bos} + B\eta\partial\eta e^{2\phi}(0)|0\rangle. \quad (8.77)$$

The usage of the cohomology argument is straightforward, since A of (5.36) contains the operator B and so kills the extra piece, which also contains the B operator (and no c insertions). Thus, one can use the argument of [163] without any modification to prove that the cohomology vanishes.

The evaluation of the action requires some sort of regularization of the solution. This can be achieved either by level truncation (usually not adequate for analytical calculations) or by introducing the “phantom pieces” of the so-

⁸⁹ It is possible to define the solution in the supersymmetric theory, since Schnabl's solution can be given in a formal gauge form using the gauge field (5.5).

⁹⁰ The map does not involve operators of half-integer conformal weight.

lution. As noted in section 5, Erler found the amount of extra terms needed in the general case and their form. He found out that the regularization of the solution at hand should involve two phantom terms (one more than in Schnabl's case), so a properly regularize form for it is,

$$\Psi = \lim_{N \rightarrow \infty} \left(\sum_{n=0}^N \psi'_n - \psi_N - \frac{1}{2} \psi'_N \right) + B\gamma^2(0) |0\rangle. \quad (8.78)$$

The evaluation of the action is quite similar to the bosonic case and is in fact simpler. Erler showed that the solution obeys the equation of motion in the strict sense, i.e., even when contracted with the solution itself. Hence, one can use the equation of motion in order to write the action of the solution as

$$S(\Psi) = -\frac{1}{6} \int Y_{-2} \Psi Q \Psi. \quad (8.79)$$

The evaluation differs from the bosonic one in several ways. First, there is the Y_{-2} insertion in all correlators. Then, in order to get a non-zero result, the ϕ momentum should equal minus two (8.10). Since Y_{-2} has -4 charge, one has to take into account only terms whose momentum equals two, such as the γ^2 term. In particular, all the terms that contribute in the bosonic case do not contribute now. This does not imply that the Ψ_{bos} piece of the solution does not contribute, since now we use the BRST charge of the supersymmetric theory and $Q\Psi_{bos}$ contains a γ^2 piece. All in all, one has to reduce all the expression to correlators of the form

$$\left\langle Y_{-2} \int_{-i\infty}^{i\infty} dw b(w) c(y+z) c(z) \gamma^2(0) \right\rangle_{C_{\pi(x+y+z)}} = -\frac{x+y+z}{2\pi^2} y, \quad (8.80)$$

where the CFT evaluation is straightforward and can be found in the appendix of [161]⁹¹. This result is much simpler to work with than the bosonic one. Recall that in the bosonic case the building blocks for the evaluation of the action contained trigonometric functions (5.24), which made the evaluation of the sums somewhat tricky. Now, the summation is trivial.

Plugging (8.80) into the expression for the action, one sees that the bosonic piece contains terms that are not only non-zero, but even diverge in the limit $N \rightarrow \infty$. These divergences cancel against each other, leaving a total zero contribution from the bosonic piece. Similarly evaluating the contribution of the terms in the action that involve also the new pieces of the solution (the γ^2 piece and the Ψ'_N piece) results in,

$$-E(\Psi) = S(\Psi) = \frac{1}{2\pi^2}, \quad (8.81)$$

⁹¹ Note, however, that he uses a slightly different conventions.

which is the expected result for a vanishing D-brane⁹².

The evaluation of the action and the results regarding the cohomology support the physical interpretation of these solutions and imply that, at least to some extent, open string field theory can be used to describe solutions that are not continuously connected to the original theory⁹³.

One may criticize Erler's solution [161], as well as the analogous solution in Berkovits' theory [158], as describing the condensation of the wrong string field. To leading order the solution describes the condensation of the non-physical GSO(+) tachyon (8.46). It was mentioned in section 8.1.1 that this string field is not closed and so does not give rise to a vertex operator. From the perspective of string field theory, its non-closeness means that its free equation of motion is a constraint equation implying that it vanishes. It is the first of many auxiliary string fields peculiar to the zero picture. This field can nevertheless play a role within the framework of an interacting string field theory. In fact, a precursor of Erler's solution was found by Arefeva et. al. almost twenty years ago [260]. There, the modified chiral superstring field theory was truncated to level zero and a condensation of the GSO(+) tachyon was obtained as solution⁹⁴. It was claimed that the solution describes a supersymmetry-breaking vacuum. This interpretation was based on the observation that around this solution the bosonic modes become heavier, while nothing of this sort was observed for the Ramond sector states. One should take this observation with a grain of salt, due to the approximate nature of the level zero truncation.

The understanding that a solution describing the GSO(-) tachyon should also exist motivated Arefeva et. al. [261] to look for a generalization of Erler's solution. On the non-BPS D-brane Erler's solution is given by setting in (8.63),

$$\Lambda_+ = Bc(0) |0\rangle, \quad \Lambda_- = 0. \quad (8.82)$$

One can generalize that by allowing a non-zero Λ_- . In the general case the

⁹² The space-time volume is normalized to unity.

⁹³ In the case at hand the solutions are not connected as they describe states with different RR charge. One can, nevertheless, in the case of a lower dimensional D-brane, consider the continuous process of moving the D-brane to infinity. In the case of a D9-brane, which cannot be sent to infinity, one may expect the solution to have problems quantum mechanically, unless the RR charge is somehow balanced.

⁹⁴ A few years ago, Ohmori studied this solution using level truncation up to level three. He found out that a solution indeed exists for the chiral theory, but he did not find the solution for the, presumably more reliable, non-chiral version of the theory [153]. This is probably an artifact of level truncation, since Erler constructed his analytical solution just for this non-chiral theory.

solution is

$$\begin{aligned}\Psi &= (Q \otimes \sigma_3)(\Lambda_+ \otimes \mathbf{1} + \Lambda_- \otimes \sigma_1) \frac{1}{1 - (\Lambda_+ \otimes \mathbf{1} + \Lambda_- \otimes \sigma_1)} \\ &= (Q\Lambda_+\Xi_+ + Q\Lambda_-\Xi_-) \otimes \sigma_3 + (Q\Lambda_-\Xi_+ + Q\Lambda_+\Xi_-) \otimes (i\sigma_2),\end{aligned}\quad (8.83)$$

where we defined⁹⁵,

$$\Xi_{\pm} = \frac{1}{2} \left(\frac{1}{1 - (\Lambda_+ + \Lambda_-)} \pm \frac{1}{1 - (\Lambda_+ - \Lambda_-)} \right). \quad (8.84)$$

Two simple special cases are when either of Λ_{\pm} equals zero. For $\Lambda_- = 0$ one gets just the usual GSO(+) solution, while for $\Lambda_+ = 0$ one gets

$$\Psi = Q\Lambda_- \frac{\Lambda_-}{1 - \Lambda_-^2} \otimes \sigma_3 + Q\Lambda_- \frac{1}{1 - \Lambda_-^2} \otimes (i\sigma_2). \quad (8.85)$$

In this case one has a non-zero expression in both of the GSO sectors, but the GSO(+) sector starts at a higher order with respect to Λ_- .

In [261] it was suggested that the solution describing tachyon condensation should contain in the gauge field the physical, GSO(−) tachyon field. They advocated to use the gauge field,

$$\Lambda_+ = Bc(0) |0\rangle, \quad \Lambda_- = B\gamma(0) |0\rangle. \quad (8.86)$$

For this specific choice one gets

$$\Lambda_-^2 = \Lambda_- \Lambda_+ = 0, \quad (8.87)$$

which dictates that for this solution

$$\Xi_+ = \frac{1}{1 - \Lambda_+}, \quad \Xi_- = \frac{1}{1 - \Lambda_+} \Lambda_-. \quad (8.88)$$

Let us consider the slightly more general gauge field [257],

$$\Lambda_+ = Bc(0) |0\rangle, \quad \Lambda_- = \epsilon B\gamma(0) |0\rangle. \quad (8.89)$$

This is a one parameter family interpolating (8.86) ($\epsilon = 1$) and Erler's solution ($\epsilon = 0$).

The action of the whole ϵ -family agrees with that of Erler's solution. For the evaluation of the action we recall that in order to get a non-zero result, the fields should have a total power two of γ . Since Q can only increase the amount of γ 's we conclude that terms with more than two occurrences of Λ_- will not

⁹⁵ The expressions $\Lambda_+ \pm \Lambda_-$ may seem awkward, as they mix the two GSO sectors. The resulting Ξ_{\pm} are, however, standard.

contribute. Let us now consider an arbitrary ϵ -solution as being expanded around Erler's one. We can use the general expression for the action around a solution (3.2) and the fact that the ϵ -solution also obeys the equation of motion, in order to write the action of this solution as⁹⁶

$$S_\epsilon = S_E + \frac{1}{6} \int \frac{1}{2} \text{tr}(Y_{-2} \tilde{\Psi}^3). \quad (8.90)$$

Here $\tilde{\Psi}$ represents that deviation of the solution from Erler's one. For $\epsilon = 1$ the GSO(-) part of $\tilde{\Psi}$, is the term proportional to $i\sigma_2$ in (8.83), and the GSO(+) part is the second summand proportional to σ_3 in (8.83). The GSO(-) part $\tilde{\Psi}_-$ is proportional to γ , while $\tilde{\Psi}_+$ is proportional to γ^2 . At least one factor of $\tilde{\Psi}_+$ is needed for a non-zero trace. Hence, $\tilde{\Psi}^3$ contributes to the action (8.90) terms with at least four γ 's. This γ -counting works the same way for $\epsilon \neq 1$. This results in

$$S_\epsilon = S_E. \quad (8.91)$$

It was further found in [257] that the cohomology of this family of solutions is empty. These results seem to suggest that the solutions of the ϵ -family are gauge equivalent.

Writing the gauge transformation that sends Erler's solution to a general ϵ -solution is straightforward, since the solutions of this family are given as formal gauge solutions. Composing the gauge transformations one gets

$$\begin{aligned} e^{\Lambda_\epsilon^E} &= (1 - \Lambda_E) \frac{1}{1 - \Lambda_\epsilon} = 1 \otimes \mathbf{1} + \epsilon B \gamma(0) |0\rangle \otimes \sigma_1, \\ e^{-\Lambda_\epsilon^E} &= (1 - \Lambda_\epsilon) \frac{1}{1 - \Lambda_E} = 1 \otimes \mathbf{1} - \epsilon B \gamma(0) |0\rangle \otimes \sigma_1. \end{aligned} \quad (8.92)$$

These transformations form an abelian group with the simple multiplication rule,

$$e^{\Lambda_\epsilon^E} e^{\Lambda_\epsilon^E} = e^{\Lambda_\epsilon^E + \tilde{\epsilon}}. \quad (8.93)$$

In order to verify that these gauge transformations are not singular in some sense, one could try to calculate some invariants as in [262,263] and verify that the result is ϵ -independent. This check was not performed yet. The known properties of the solutions and of these gauge transformations support the idea that these transformations are genuine gauge transformations.

One is therefore lead to believe that the $\epsilon = 0$ solution can be used to describe tachyon condensation on the non-BPS D-brane. It is then very natural to assume that Erler's original solution and its counterpart in the non-polynomial theory indeed manage to describe the state without the BPS D-brane despite the fact that it supports no tachyons.

⁹⁶ Note that we added to (3.2) the action of the original solution (Erler's one) and also wrote down the trace factor needed when considering the non-BPS brane.

9 Outlook

The advance in our understanding of string field theory described in this work has the potential of turning string field theory to a practical framework for (non-perturbative) string theory research. Obviously there is still more work to be done to that end. Let us mention some of the relevant issues.

The most obvious missing ingredient is the construction of analytical solutions describing lump solutions, i.e., lower dimensional D-branes, around the tachyon vacuum solution⁹⁷. This is desirable both in its own right as well as for addressing Sen’s second conjecture that for now could not be proved. One potential complication with this construction is that it should refer to a specific BCFT, that of the lump, and is therefore not background independent. Background independence, translated into the description of solutions using general CFT methods only, played an important role in the constructions of analytical solutions. One may hope to look for general lump solutions using tools such as boundary changing operators as well as using explicit background/oscillator representations.

The, somewhat surprising, possibility of generalizing Schnabl’s solution to the case of a BPS D-brane makes one wonder: which string backgrounds are accessible as classical solutions of open string field theory around a particular background? This is an important question, whose answer can help us to estimate the relevance of string field theory to other directions of string theory research. A particular interesting question is whether it is possible to obtain a solution describing N D-branes from string field theory in the background of M D-branes with $M < N$. The opposite case, $M > N$, where some of the D-branes condense, was analytically solved in [163]. Is it possible to go to the other direction, or would one fail, since the theory does not contain “enough degrees of freedom”? The search for a two D-brane system around a single one, within level truncation, was not successful. Other than to a problem of principle, one can ascribe this failure, either to the limit of validity of the Siegel gauge, or to numerical problems resulting from “climbing up the potential”. One can hope to avoid these problems by finding an analytical solution. No such solution was found yet.

Following the treatment of solutions as formal gauge solutions and given the fact that all currently known analytical solutions can be cast in such a form, some obvious questions arise:

⁹⁷ There are also other examples of string field theory solutions for which no analytical form is known. One such example is that of solutions representing cosmological tachyon models [264]. Another interesting example includes marginal deformations on the separated D- \bar{D} pair [265].

- Is it possible to represent all string field theory solutions as formal gauge solutions?
- Given a solution how should one cast it in a gauge form?
- Is there a simple criterion to distinguish the “large gauge transformations” from the majority of pure gauge ones?
- Given a solution that is formally written in a gauge form, is there a simple and universal prescription for regularizing it?

To understand the last two questions we recall that we regularized Schnabl’s solution and the marginal deformation solutions in quite a different way and the manifestation of them being non-trivial was also quite different⁹⁸.

Another pressing issue is to recognize the physical nature of solutions, as it currently seems that it is easier to construct solutions than to interpret them. Generally speaking, one expects that a solution in string field theory corresponds to a boundary CFT. One route for recognizing the BCFT is to evaluate a large enough set of gauge invariant operators describing the solution so as to reveal its nature. Gauge invariant quantities other than the action include the invariants found in [266,168]. They are related to the 1-point disk scattering amplitudes of closed strings⁹⁹. Generalization to Berkovits’ superstring field theory was given in [271]. Ellwood [262] used these invariants in order to address the above mentioned question in the bosonic as well as in the supersymmetric theory. For specifically comparing Schnabl’s solution with the perturbative one found in the Siegel gauge, some of these gauge invariant expressions were calculated in [263], supporting the expectation that these solutions are gauge equivalent¹⁰⁰. They were further shown to coincide in a specific case with the tadpole of an open string state on the boundary state that corresponds to the closed string [272]. It would be interesting to clarify whether these invariants fully characterize a solution and whether other gauge invariant expressions can be defined and evaluated. It is also desirable to find methods for improving their calculability.

Closed strings can be described naturally using the framework of closed string field theory [248]. While recent advance in the field proved adequate for open superstring field theory, the generalization to closed string field theory was not found¹⁰¹. One of the obstacles for the generalization lies in the existence of

⁹⁸Note that evaluating the action is not always enough, since it is zero for the marginal deformations.

⁹⁹See [249] for an early incorporation of on-shell closed strings into open string field theory. Closed string amplitudes in open string field theory were also studied in [267,268]. Some interesting speculations regarding the relation between open and closed strings from a string field theoretical perspective appeared in [269,270].

¹⁰⁰The best would be of course, if one could somehow translate the solutions to Siegel gauge. We do not know how to do that yet.

¹⁰¹There was, nevertheless, quite an impressive advance in the study of tachyon

an intricate structure of many string vertices, not simply related to the star-product. Another stumbling-block comes from the ghost number of the string fields, which is two in the case of the closed string. The required saturation of six ghost zero modes in the integral implies that at the quadratic order an explicit ghost insertion is present in addition to the BRST charge. Together with the Siegel gauge this insertion implies that the string fields obey

$$b_0\Psi = \bar{b}_0\Psi = 0. \tag{9.1}$$

Modifying the gauge choice to the Schnabl gauge, without changing the form of the explicit expression leads to awkward looking formulas. It would be very desirable to examine how can the explicit ghost insertion be changed and in what way should the gauge choice be modified as compared to the open string case.

Acknowledgments

During the preparation of this work we benefited much from discussions with Ido Adam, Ofer Aharony, Sudarshan Ananth, Nathan Berkovits, Guillaume Bossard, Ian Ellwood, Stefan Fredenhagen, Sunny Itzhaki, Alexey Koshelev, Carlo Maccaferri, Nicolas Moeller, Yuji Okawa, Yaron Oz, Rob Potting, Samson Shatashvili, Martin Schnabl, Ashoke Sen, Stefan Theisen and Barton Zwiebach.

M. K. would like to thank Tel-Aviv University for hospitality. The work of M. K. was supported by a Minerva fellowship during the initiation of this project. This research was supported by the German-Israeli Project cooperation (DIP H.52).

condensation within closed string field theory [273,274,275,276,277,278,279]. The new vacuum is interpreted in this case as representing a big crunch of space time, since the metric goes to zero. Lump solutions were also found in this framework and it was claimed that they represent a lower dimensional space time. This claim is consistent with finding an approximately linear profile of the dilaton for these solutions. An important technical tool that enabled this progress is the improved numerical description of higher string vertices.

References

- [1] E. Witten, *Noncommutative geometry and string field theory*, *Nucl. Phys.* **B268** (1986) 253. (Cited on pages 3, 9 and 33.)
- [2] M. Schnabl, *Analytic solution for tachyon condensation in open string field theory*, *Adv. Theor. Math. Phys.* **10** (2006) 433–501, [[hep-th/0511286](#)]. (Cited on pages 3, 6, 26, 33 and 46.)
- [3] A. LeClair, M. E. Peskin, and C. R. Preitschopf, *String field theory on the conformal plane. 1. Kinematical principles*, *Nucl. Phys.* **B317** (1989) 411. (Cited on pages 3, 12, 14, 54 and 57.)
- [4] A. LeClair, M. E. Peskin, and C. R. Preitschopf, *String field theory on the conformal plane. 2. Generalized gluing*, *Nucl. Phys.* **B317** (1989) 464. (Cited on pages 3, 12, 14 and 54.)
- [5] A. Sen, *Universality of the tachyon potential*, *JHEP* **12** (1999) 027, [[hep-th/9911116](#)]. (Cited on pages 3, 16, 19, 20 and 42.)
- [6] J. Bordes, H.-M. Chan, L. Nellen, and S. T. Tsou, *Half string oscillator approach to string field theory*, *Nucl. Phys.* **B351** (1991) 441–473. (Cited on page 3.)
- [7] L. Rastelli, A. Sen, and B. Zwiebach, *Half strings, projectors, and multiple D-branes in vacuum string field theory*, *JHEP* **11** (2001) 035, [[hep-th/0105058](#)]. (Cited on pages 3 and 41.)
- [8] D. J. Gross and W. Taylor, *Split string field theory. I*, *JHEP* **08** (2001) 009, [[hep-th/0105059](#)]. (Cited on page 3.)
- [9] D. J. Gross and W. Taylor, *Split string field theory. II*, *JHEP* **08** (2001) 010, [[hep-th/0106036](#)]. (Cited on page 3.)
- [10] K. Furuuchi and K. Okuyama, *Comma vertex and string field algebra*, *JHEP* **09** (2001) 035, [[hep-th/0107101](#)]. (Cited on pages 3 and 30.)
- [11] V. A. Kostelecky and R. Potting, *Analytical construction of a nonperturbative vacuum for the open bosonic string*, *Phys. Rev.* **D63** (2001) 046007, [[hep-th/0008252](#)]. (Cited on pages 3, 41 and 58.)
- [12] I. Bars, *Map of Witten's $*$ to Moyal's $*$* , *Phys. Lett.* **B517** (2001) 436–444, [[hep-th/0106157](#)]. (Cited on page 3.)
- [13] M. R. Douglas, H. Liu, G. Moore, and B. Zwiebach, *Open string star as a continuous Moyal product*, *JHEP* **04** (2002) 022, [[hep-th/0202087](#)]. (Cited on pages 3, 56 and 57.)
- [14] I. Bars and Y. Matsuo, *Computing in string field theory using the Moyal star product*, *Phys. Rev.* **D66** (2002) 066003, [[hep-th/0204260](#)]. (Cited on page 3.)

- [15] T. G. Erler, *Moyal formulation of Witten's star product in the fermionic ghost sector*, [hep-th/0205107](#). (Cited on pages 3 and 57.)
- [16] I. Bars, I. Kishimoto, and Y. Matsuo, *String amplitudes from Moyal string field theory*, *Phys. Rev.* **D67** (2003) 066002, [[hep-th/0211131](#)]. (Cited on pages 3 and 5.)
- [17] I. Bars, I. Kishimoto, and Y. Matsuo, *Fermionic ghosts in Moyal string field theory*, *JHEP* **07** (2003) 027, [[hep-th/0304005](#)]. (Cited on page 3.)
- [18] I. Bars and I. Y. Park, *Improved off-shell scattering amplitudes in string field theory and new computational methods*, *Phys. Rev.* **D69** (2004) 086007, [[hep-th/0311264](#)]. (Cited on pages 3 and 5.)
- [19] N. Seiberg and E. Witten, *String theory and noncommutative geometry*, *JHEP* **09** (1999) 032, [[hep-th/9908142](#)]. (Cited on pages 3 and 42.)
- [20] L. Rastelli, A. Sen, and B. Zwiebach, *Star algebra spectroscopy*, *JHEP* **03** (2002) 029, [[hep-th/0111281](#)]. (Cited on pages 3, 55 and 59.)
- [21] K. Okuyama, *Ghost kinetic operator of vacuum string field theory*, *JHEP* **01** (2002) 027, [[hep-th/0201015](#)]. (Cited on pages 3, 55 and 56.)
- [22] L. Rastelli, A. Sen, and B. Zwiebach, *String field theory around the tachyon vacuum*, *Adv. Theor. Math. Phys.* **5** (2002) 353–392, [[hep-th/0012251](#)]. (Cited on page 4.)
- [23] L. Rastelli, A. Sen, and B. Zwiebach, *Classical solutions in string field theory around the tachyon vacuum*, *Adv. Theor. Math. Phys.* **5** (2002) 393–428, [[hep-th/0102112](#)]. (Cited on page 4.)
- [24] L. Rastelli, A. Sen, and B. Zwiebach, *Boundary CFT construction of D-branes in vacuum string field theory*, *JHEP* **11** (2001) 045, [[hep-th/0105168](#)]. (Cited on pages 4 and 30.)
- [25] L. Rastelli, A. Sen, and B. Zwiebach, *Vacuum string field theory*, [hep-th/0106010](#). (Cited on page 4.)
- [26] H. Hata and T. Kawano, *Open string states around a classical solution in vacuum string field theory*, *JHEP* **11** (2001) 038, [[hep-th/0108150](#)]. (Cited on page 4.)
- [27] L. Rastelli, A. Sen, and B. Zwiebach, *A note on a proposal for the tachyon state in vacuum string field theory*, *JHEP* **02** (2002) 034, [[hep-th/0111153](#)]. (Cited on page 4.)
- [28] H. Hata and S. Moriyama, *Reexamining classical solution and tachyon mode in vacuum string field theory*, *Nucl. Phys.* **B651** (2003) 3–25, [[hep-th/0206208](#)]. (Cited on page 4.)
- [29] Y. Okawa, *Open string states and D-brane tension from vacuum string field theory*, *JHEP* **07** (2002) 003, [[hep-th/0204012](#)]. (Cited on pages 4 and 39.)

- [30] T. Takahashi and S. Tanimoto, *Wilson lines and classical solutions in cubic open string field theory*, *Prog. Theor. Phys.* **106** (2001) 863–872, [[hep-th/0107046](#)]. (Cited on page 4.)
- [31] J. Kluson, *Exact solutions of open bosonic string field theory*, *JHEP* **04** (2002) 043, [[hep-th/0202045](#)]. (Cited on page 4.)
- [32] T. Takahashi and S. Tanimoto, *Marginal and scalar solutions in cubic open string field theory*, *JHEP* **03** (2002) 033, [[hep-th/0202133](#)]. (Cited on pages 4 and 43.)
- [33] N. Drukker, *On different actions for the vacuum of bosonic string field theory*, *JHEP* **08** (2003) 017, [[hep-th/0301079](#)]. (Cited on page 4.)
- [34] G. T. Horowitz and A. Strominger, *Translations as inner derivations and associativity anomalies in open string field theory*, *Phys. Lett.* **B185** (1987) 45. (Cited on pages 4 and 52.)
- [35] L. Rastelli and B. Zwiebach, *Tachyon potentials, star products and universality*, *JHEP* **09** (2001) 038, [[hep-th/0006240](#)]. (Cited on pages 4, 20, 30 and 41.)
- [36] Y. Okawa, *Some exact computations on the twisted butterfly state in string field theory*, *JHEP* **01** (2004) 066, [[hep-th/0310264](#)]. (Cited on page 4.)
- [37] Y. Okawa, *Solving Witten’s string field theory using the butterfly state*, *Phys. Rev.* **D69** (2004) 086001, [[hep-th/0311115](#)]. (Cited on page 4.)
- [38] H.-t. Yang, *Solving Witten’s SFT by insertion of operators on projectors*, *JHEP* **09** (2004) 002, [[hep-th/0406023](#)]. (Cited on page 4.)
- [39] N. Drukker and Y. Okawa, *Vacuum string field theory without matter-ghost factorization*, *JHEP* **06** (2005) 032, [[hep-th/0503068](#)]. (Cited on page 4.)
- [40] C. B. Thorn, *String field theory*, *Phys. Rept.* **175** (1989) 1–101. (Cited on page 4.)
- [41] L. Rastelli, *String field theory*, [hep-th/0509129](#). (Cited on page 4.)
- [42] W. Taylor, *String field theory*, [hep-th/0605202](#). (Cited on page 4.)
- [43] L. Rastelli, *Open string fields and D-branes*, *Fortsch. Phys.* **52** (2004) 302–337. (Cited on page 4.)
- [44] K. Ohmori, *A review on tachyon condensation in open string field theories*, [hep-th/0102085](#). (Cited on pages 4 and 70.)
- [45] P.-J. De Smet, *Tachyon condensation: Calculations in string field theory*, [hep-th/0109182](#). (Cited on page 4.)
- [46] W. Taylor, *Lectures on D-branes, tachyon condensation, and string field theory*, [hep-th/0301094](#). (Cited on page 4.)
- [47] W. Taylor and B. Zwiebach, *D-branes, tachyons, and string field theory*, [hep-th/0311017](#). (Cited on page 4.)

- [48] A. Sen, *Tachyon dynamics in open string theory*, *Int. J. Mod. Phys.* **A20** (2005) 5513–5656, [[hep-th/0410103](#)]. (Cited on page 4.)
- [49] I. Y. Arefeva, D. M. Belov, A. A. Giryavets, A. S. Koshelev, and P. B. Medvedev, *Noncommutative field theories and (super)string field theories*, [hep-th/0111208](#). (Cited on pages 4 and 76.)
- [50] I. Bars, *MSFT: Moyal star formulation of string field theory*, [hep-th/0211238](#). (Cited on page 4.)
- [51] B. Zwiebach, *Exploring the open string star algebra: Applications to tachyon condensation*, . Prepared for ICTP Spring School on Superstrings and Related Matters, Trieste, Italy, 18-26 Mar 2002. (Cited on page 4.)
- [52] L. Bonora, C. Maccaferri, D. Mamone, and M. Salizzoni, *Topics in string field theory*, [hep-th/0304270](#). (Cited on page 4.)
- [53] S. Samuel, *Covariant off-shell string amplitudes*, *Nucl. Phys.* **B308** (1988) 285. (Cited on page 5.)
- [54] V. A. Kostelecky and S. Samuel, *The static tachyon potential in the open bosonic string theory*, *Phys. Lett.* **B207** (1988) 169. (Cited on pages 5, 18 and 21.)
- [55] W. Taylor, *Perturbative diagrams in string field theory*, [hep-th/0207132](#). (Cited on pages 5 and 21.)
- [56] W. Taylor, *Perturbative computations in string field theory*, [hep-th/0404102](#). (Cited on page 5.)
- [57] V. Forini, G. Grignani, and G. Nardelli, *A solution to the 4-tachyon off-shell amplitude in cubic string field theory*, *JHEP* **04** (2006) 053, [[hep-th/0603206](#)]. (Cited on page 5.)
- [58] H. Fuji, S. Nakayama, and H. Suzuki, *Open string amplitudes in various gauges*, *JHEP* **01** (2007) 011, [[hep-th/0609047](#)]. (Cited on page 6.)
- [59] L. Rastelli and B. Zwiebach, *The off-shell Veneziano amplitude in Schnabl gauge*, *JHEP* **01** (2008) 018, [[0708.2591](#)]. (Cited on page 6.)
- [60] M. Kiermaier, A. Sen, and B. Zwiebach, *Linear b-gauges for open string fields*, *JHEP* **03** (2008) 050–050, [[0712.0627](#)]. (Cited on pages 6 and 34.)
- [61] M. Kiermaier and B. Zwiebach, *One-loop Riemann surfaces in Schnabl gauge*, [0805.3701](#). (Cited on pages 6 and 34.)
- [62] J. Polchinski, *Dirichlet-branes and Ramond-Ramond charges*, *Phys. Rev. Lett.* **75** (1995) 4724–4727, [[hep-th/9510017](#)]. (Cited on pages 7 and 19.)
- [63] V. A. Kostelecky and S. Samuel, *On a nonperturbative vacuum for the open bosonic string*, *Nucl. Phys.* **B336** (1990) 263. (Cited on pages 7 and 18.)
- [64] W. Siegel, *Covariantly second quantized string. 2*, *Phys. Lett.* **B151** (1984) 391. (Cited on page 7.)

- [65] W. Siegel, *Covariantly second quantized string. 3*, *Phys. Lett.* **B151s** (1984) 396. (Cited on page 7.)
- [66] A. Neveu, H. Nicolai, and P. C. West, *New symmetries and ghost structure of covariant string theories*, *Phys. Lett.* **B167** (1986) 307. (Cited on page 8.)
- [67] N. Marcus and A. Sagnotti, *String field theory and equations of motion*, *Phys. Lett.* **B178** (1986) 343. (Cited on page 11.)
- [68] J. Zinn-Justin, *Renormalization of gauge theories*, . Lectures given at Int. Summer Inst. for Theoretical Physics, Jul 29 - Aug 9, 1974, Bonn, West Germany. (Cited on page 11.)
- [69] I. A. Batalin and G. A. Vilkovisky, *Gauge algebra and quantization*, *Phys. Lett.* **B102** (1981) 27–31. (Cited on page 11.)
- [70] I. A. Batalin and G. A. Vilkovisky, *Quantization of gauge theories with linearly dependent generators*, *Phys. Rev.* **D28** (1983) 2567–2582. (Cited on page 11.)
- [71] I. A. Batalin and G. A. Vilkovisky, *Closure of the gauge algebra, generalized Lie equations and Feynman rules*, *Nucl. Phys.* **B234** (1984) 106–124. (Cited on page 11.)
- [72] B. L. Voronov and I. V. Tyutin, *Formulation of gauge theories of general form. I*, *Theor. Math. Phys.* **50** (1982) 218–225. (Cited on page 11.)
- [73] M. Henneaux and C. Teitelboim, *Quantization of gauge systems*, . Princeton, USA: Univ. Pr. (1992) 520 p. (Cited on page 11.)
- [74] J. Gomis, J. Paris, and S. Samuel, *Antibracket, antifields and gauge theory quantization*, *Phys. Rept.* **259** (1995) 1–145, [[hep-th/9412228](#)]. (Cited on page 11.)
- [75] M. Bochicchio, *Gauge fixing for the field theory of the bosonic string*, *Phys. Lett.* **B193** (1987) 31. (Cited on page 11.)
- [76] M. Bochicchio, *String field theory in the Siegel gauge*, *Phys. Lett.* **B188** (1987) 330. (Cited on page 11.)
- [77] C. B. Thorn, *Perturbation theory for quantized string fields*, *Nucl. Phys.* **B287** (1987) 61. (Cited on page 11.)
- [78] M. Maeno, *Canonical quantization of Witten’s string field theory using midpoint light cone time*, *Phys. Rev.* **D43** (1991) 4006–4014. (Cited on page 11.)
- [79] T. G. Erler and D. J. Gross, *Locality, causality, and an initial value formulation for open string field theory*, [hep-th/0406199](#). (Cited on page 11.)
- [80] T. G. Erler, *Level truncation and rolling the tachyon in the lightcone basis for open string field theory*, [hep-th/0409179](#). (Cited on page 11.)

- [81] M. Asano and M. Kato, *New covariant gauges in string field theory*, *Prog. Theor. Phys.* **117** (2007) 569–587, [[hep-th/0611189](#)]. (Cited on page 12.)
- [82] M. Asano and M. Kato, *Level truncated tachyon potential in various gauges*, *JHEP* **01** (2007) 028, [[hep-th/0611190](#)]. (Cited on page 12.)
- [83] H. Feng and W. Siegel, *Yang-Mills gauge conditions from Witten’s open string field theory*, *Phys. Rev.* **D75** (2007) 046006, [[hep-th/0611307](#)]. (Cited on page 12.)
- [84] A. A. Belavin, A. M. Polyakov, and A. B. Zamolodchikov, *Infinite conformal symmetry in two-dimensional quantum field theory*, *Nucl. Phys.* **B241** (1984) 333–380. (Cited on page 12.)
- [85] S. B. Giddings, E. J. Martinec, and E. Witten, *Modular invariance in string field theory*, *Phys. Lett.* **B176** (1986) 362. (Cited on page 12.)
- [86] S. B. Giddings and E. J. Martinec, *Conformal geometry and string field theory*, *Nucl. Phys.* **B278** (1986) 91. (Cited on page 12.)
- [87] S. B. Giddings, *The Veneziano amplitude from interacting string field theory*, *Nucl. Phys.* **B278** (1986) 242. (Cited on page 12.)
- [88] B. Zwiebach, *A proof that Witten’s open string theory gives a single cover of moduli space*, *Commun. Math. Phys.* **142** (1991) 193–216. (Cited on page 12.)
- [89] D. J. Gross and A. Jevicki, *Operator formulation of interacting string field theory*, *Nucl. Phys.* **B283** (1987) 1. (Cited on page 12.)
- [90] D. J. Gross and A. Jevicki, *Operator formulation of interacting string field theory. 2*, *Nucl. Phys.* **B287** (1987) 225. (Cited on page 12.)
- [91] D. J. Gross and A. Jevicki, *Operator formulation of interacting string field theory. 3. NSR superstring*, *Nucl. Phys.* **B293** (1987) 29. (Cited on page 12.)
- [92] S. Samuel, *The physical and ghost vertices in Witten’s string field theory*, *Phys. Lett.* **B181** (1986) 255. (Cited on page 12.)
- [93] N. Ohta, *Covariant interacting string field theory in the Fock space representation*, *Phys. Rev.* **D34** (1986) 3785–3793. (Cited on page 12.)
- [94] E. Cremmer, A. Schwimmer, and C. B. Thorn, *The vertex function in Witten’s formulation of string field theory*, *Phys. Lett.* **B179** (1986) 57. (Cited on page 12.)
- [95] E. Witten, *Interacting field theory of open superstrings*, *Nucl. Phys.* **B276** (1986) 291. (Cited on pages 13, 25, 28 and 75.)
- [96] M. Schnabl, *Wedge states in string field theory*, *JHEP* **01** (2003) 004, [[hep-th/0201095](#)]. (Cited on pages 15 and 30.)
- [97] A. Sen, *Descent relations among bosonic D-branes*, *Int. J. Mod. Phys.* **A14** (1999) 4061–4078, [[hep-th/9902105](#)]. (Cited on page 16.)

- [98] A. Sen, *Stable non-BPS bound states of BPS D-branes*, *JHEP* **08** (1998) 010, [[hep-th/9805019](#)]. (Cited on page 16.)
- [99] A. Sen, *Tachyon condensation on the brane antibrane system*, *JHEP* **08** (1998) 012, [[hep-th/9805170](#)]. (Cited on page 16.)
- [100] A. Sen, *$SO(32)$ spinors of type I and other solitons on brane-antibrane pair*, *JHEP* **09** (1998) 023, [[hep-th/9808141](#)]. (Cited on page 16.)
- [101] A. Sen, *BPS D-branes on non-supersymmetric cycles*, *JHEP* **12** (1998) 021, [[hep-th/9812031](#)]. (Cited on page 16.)
- [102] A. Sen, *Non-BPS states and branes in string theory*, [hep-th/9904207](#). (Cited on page 16.)
- [103] A. Sen, *Rolling tachyon*, *JHEP* **04** (2002) 048, [[hep-th/0203211](#)]. (Cited on page 16.)
- [104] A. Sen, *Tachyon matter*, *JHEP* **07** (2002) 065, [[hep-th/0203265](#)]. (Cited on page 16.)
- [105] A. Sen, *Supersymmetric world-volume action for non-BPS D-branes*, *JHEP* **10** (1999) 008, [[hep-th/9909062](#)]. (Cited on page 18.)
- [106] M. R. Garousi, *Tachyon couplings on non-BPS D-branes and Dirac-Born-Infeld action*, *Nucl. Phys.* **B584** (2000) 284–299, [[hep-th/0003122](#)]. (Cited on page 18.)
- [107] E. A. Bergshoeff, M. de Roo, T. C. de Wit, E. Eyras, and S. Panda, *T-duality and actions for non-BPS D-branes*, *JHEP* **05** (2000) 009, [[hep-th/0003221](#)]. (Cited on page 18.)
- [108] J. Kluson, *Proposal for non-BPS D-brane action*, *Phys. Rev.* **D62** (2000) 126003, [[hep-th/0004106](#)]. (Cited on page 18.)
- [109] A. Sen, *Field theory of tachyon matter*, *Mod. Phys. Lett.* **A17** (2002) 1797–1804, [[hep-th/0204143](#)]. (Cited on page 18.)
- [110] D. Ghoshal and A. Sen, *Tachyon condensation and brane descent relations in p -adic string theory*, *Nucl. Phys.* **B584** (2000) 300–312, [[hep-th/0003278](#)]. (Cited on page 18.)
- [111] N. Moeller and B. Zwiebach, *Dynamics with infinitely many time derivatives and rolling tachyons*, *JHEP* **10** (2002) 034, [[hep-th/0207107](#)]. (Cited on pages 18, 24 and 45.)
- [112] H.-t. Yang, *Stress tensors in p -adic string theory and truncated OSFT*, *JHEP* **11** (2002) 007, [[hep-th/0209197](#)]. (Cited on page 18.)
- [113] N. Moeller and M. Schnabl, *Tachyon condensation in open-closed p -adic string theory*, *JHEP* **01** (2004) 011, [[hep-th/0304213](#)]. (Cited on page 18.)
- [114] P. G. O. Freund and M. Olson, *Nonarchimedean strings*, *Phys. Lett.* **B199** (1987) 186. (Cited on page 18.)

- [115] L. Brekke, P. G. O. Freund, M. Olson, and E. Witten, *Nonarchimedean string dynamics*, *Nucl. Phys.* **B302** (1988) 365. (Cited on page 18.)
- [116] E. Witten, *On background independent open string field theory*, *Phys. Rev.* **D46** (1992) 5467–5473, [[hep-th/9208027](#)]. (Cited on page 18.)
- [117] E. Witten, *Some computations in background independent off-shell string theory*, *Phys. Rev.* **D47** (1993) 3405–3410, [[hep-th/9210065](#)]. (Cited on page 18.)
- [118] S. L. Shatashvili, *Comment on the background independent open string theory*, *Phys. Lett.* **B311** (1993) 83–86, [[hep-th/9303143](#)]. (Cited on page 18.)
- [119] S. L. Shatashvili, *On the problems with background independence in string theory*, *Alg. Anal.* **6** (1994) 215–226, [[hep-th/9311177](#)]. (Cited on page 18.)
- [120] A. A. Gerasimov and S. L. Shatashvili, *On exact tachyon potential in open string field theory*, *JHEP* **10** (2000) 034, [[hep-th/0009103](#)]. (Cited on page 18.)
- [121] D. Kutasov, M. Marino, and G. W. Moore, *Some exact results on tachyon condensation in string field theory*, *JHEP* **10** (2000) 045, [[hep-th/0009148](#)]. (Cited on page 18.)
- [122] D. Ghoshal and A. Sen, *Normalisation of the background independent open string field theory action*, *JHEP* **11** (2000) 021, [[hep-th/0009191](#)]. (Cited on page 18.)
- [123] V. A. Kostelecky and R. Potting, *Expectation values, Lorentz invariance, and CPT in the open bosonic string*, *Phys. Lett.* **B381** (1996) 89–96, [[hep-th/9605088](#)]. (Cited on page 19.)
- [124] I. Ellwood and W. Taylor, *Gauge invariance and tachyon condensation in open string field theory*, [hep-th/0105156](#). (Cited on page 19.)
- [125] B. Zwiebach, *Trimming the tachyon string field with $SU(1,1)$* , [hep-th/0010190](#). (Cited on page 20.)
- [126] W. Siegel and B. Zwiebach, *Gauge string fields*, *Nucl. Phys.* **B263** (1986) 105. (Cited on page 20.)
- [127] H. Hata and S. Shinohara, *BRST invariance of the non-perturbative vacuum in bosonic open string field theory*, *JHEP* **09** (2000) 035, [[hep-th/0009105](#)]. (Cited on pages 20 and 24.)
- [128] M. Schnabl, *Constraints on the tachyon condensate from anomalous symmetries*, *Phys. Lett.* **B504** (2001) 61–63, [[hep-th/0011238](#)]. (Cited on page 20.)
- [129] D. Gaiotto, L. Rastelli, A. Sen, and B. Zwiebach, *Patterns in open string field theory solutions*, *JHEP* **03** (2002) 003, [[hep-th/0201159](#)]. (Cited on pages 20 and 41.)

- [130] D. Gaiotto and L. Rastelli, *Experimental string field theory*, *JHEP* **08** (2003) 048, [[hep-th/0211012](#)]. (Cited on pages 20 and 21.)
- [131] A. Sen and B. Zwiebach, *Tachyon condensation in string field theory*, *JHEP* **03** (2000) 002, [[hep-th/9912249](#)]. (Cited on page 21.)
- [132] N. Moeller and W. Taylor, *Level truncation and the tachyon in open bosonic string field theory*, *Nucl. Phys.* **B583** (2000) 105–144, [[hep-th/0002237](#)]. (Cited on page 21.)
- [133] D. Gaiotto and L. Rastelli, *Progress in open string field theory*, (2002). Presentation by L. Rastelli at Strings 2002, Cambridge, England <http://www.damtp.cam.ac.uk/strings02/avt/rastelli/>. (Cited on page 21.)
- [134] W. Taylor, *A perturbative analysis of tachyon condensation*, *JHEP* **03** (2003) 029, [[hep-th/0208149](#)]. (Cited on page 21.)
- [135] H. Hata and S. Teraguchi, *Test of the absence of kinetic terms around the tachyon vacuum in cubic string field theory*, *JHEP* **05** (2001) 045, [[hep-th/0101162](#)]. (Cited on page 21.)
- [136] I. Ellwood and W. Taylor, *Open string field theory without open strings*, *Phys. Lett.* **B512** (2001) 181–188, [[hep-th/0103085](#)]. (Cited on pages 21 and 23.)
- [137] I. Ellwood, B. Feng, Y.-H. He, and N. Moeller, *The identity string field and the tachyon vacuum*, *JHEP* **07** (2001) 016, [[hep-th/0105024](#)]. (Cited on pages 22 and 39.)
- [138] S. Giusto and C. Imbimbo, *Physical states at the tachyonic vacuum of open string field theory*, *Nucl. Phys.* **B677** (2004) 52–86, [[hep-th/0309164](#)]. (Cited on pages 22 and 40.)
- [139] J. A. Harvey and P. Kraus, *D-branes as unstable lumps in bosonic open string field theory*, *JHEP* **04** (2000) 012, [[hep-th/0002117](#)]. (Cited on page 24.)
- [140] R. de Mello Koch, A. Jevicki, M. Mihailescu, and R. Tatar, *Lumps and p-branes in open string field theory*, *Phys. Lett.* **B482** (2000) 249–254, [[hep-th/0003031](#)]. (Cited on page 24.)
- [141] N. Moeller, A. Sen, and B. Zwiebach, *D-branes as tachyon lumps in string field theory*, *JHEP* **08** (2000) 039, [[hep-th/0005036](#)]. (Cited on page 24.)
- [142] P. Mukhopadhyay and A. Sen, *Test of Siegel gauge for the lump solution*, *JHEP* **02** (2001) 017, [[hep-th/0101014](#)]. (Cited on page 24.)
- [143] N. Berkovits, *The tachyon potential in open Neveu-Schwarz string field theory*, *JHEP* **04** (2000) 022, [[hep-th/0001084](#)]. (Cited on page 25.)
- [144] N. Berkovits, *Super-Poincaré invariant superstring field theory*, *Nucl. Phys.* **B450** (1995) 90–102, [[hep-th/9503099](#)]. (Cited on pages 25, 77 and 78.)

- [145] N. Berkovits, A. Sen, and B. Zwiebach, *Tachyon condensation in superstring field theory*, *Nucl. Phys.* **B587** (2000) 147–178, [[hep-th/0002211](#)]. (Cited on pages 25 and 79.)
- [146] P.-J. De Smet and J. Raeymaekers, *Level four approximation to the tachyon potential in superstring field theory*, *JHEP* **05** (2000) 051, [[hep-th/0003220](#)]. (Cited on page 25.)
- [147] A. Iqbal and A. Naqvi, *Tachyon condensation on a non-BPS D-brane*, [hep-th/0004015](#). (Cited on page 25.)
- [148] P.-J. De Smet and J. Raeymaekers, *The tachyon potential in Witten’s superstring field theory*, *JHEP* **08** (2000) 020, [[hep-th/0004112](#)]. (Cited on page 25.)
- [149] C. R. Preitschopf, C. B. Thorn, and S. A. Yost, *Superstring field theory*, *Nucl. Phys.* **B337** (1990) 363–433. (Cited on pages 25 and 75.)
- [150] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *New representation for string field solves the consistence problem for open superstring field*, *Nucl. Phys.* **B341** (1990) 464–498. (Cited on pages 25 and 75.)
- [151] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *Background formalism for superstring field theory*, *Phys. Lett.* **B240** (1990) 356–362. (Cited on pages 25 and 75.)
- [152] I. Y. Aref’eva, A. S. Koshelev, D. M. Belov, and P. B. Medvedev, *Tachyon condensation in cubic superstring field theory*, *Nucl. Phys.* **B638** (2002) 3–20, [[hep-th/0011117](#)]. (Cited on page 25.)
- [153] K. Ohmori, *Level-expansion analysis in NS superstring field theory revisited*, [hep-th/0305103](#). (Cited on pages 25 and 85.)
- [154] K. Ohmori, *Tachyonic kink and lump-like solutions in superstring field theory*, *JHEP* **05** (2001) 035, [[hep-th/0104230](#)]. (Cited on page 25.)
- [155] Y. Okawa, *Comments on Schnabl’s analytic solution for tachyon condensation in Witten’s open string field theory*, *JHEP* **04** (2006) 055, [[hep-th/0603159](#)]. (Cited on pages 26, 33, 35 and 39.)
- [156] E. Fuchs and M. Kroyter, *Schnabl’s \mathcal{L}_0 operator in the continuous basis*, *JHEP* **10** (2006) 067, [[hep-th/0605254](#)]. (Cited on pages 29, 60 and 63.)
- [157] T. Takahashi, *Level truncation analysis of exact solutions in open string field theory*, *JHEP* **01** (2008) 001, [[0710.5358](#)]. (Cited on page 33.)
- [158] E. Fuchs and M. Kroyter, *Marginal deformation for the photon in superstring field theory*, *JHEP* **11** (2007) 005, [[arXiv:0706.0717](#) [[hep-th](#)]]. (Cited on pages 34, 50, 83 and 85.)
- [159] T. Erler, *Split string formalism and the closed string vacuum*, *JHEP* **05** (2007) 083, [[hep-th/0611200](#)]. (Cited on page 35.)

- [160] T. Erler, *Split string formalism and the closed string vacuum. II*, *JHEP* **05** (2007) 084, [[hep-th/0612050](#)]. (Cited on page 35.)
- [161] T. Erler, *Tachyon vacuum in cubic superstring field theory*, *JHEP* **01** (2008) 013, [[0707.4591](#)]. (Cited on pages 37, 38, 83, 84 and 85.)
- [162] E. Fuchs and M. Kroyter, *On the validity of the solution of string field theory*, *JHEP* **05** (2006) 006, [[hep-th/0603195](#)]. (Cited on page 39.)
- [163] I. Ellwood and M. Schnabl, *Proof of vanishing cohomology at the tachyon vacuum*, *JHEP* **02** (2007) 096, [[hep-th/0606142](#)]. (Cited on pages 39, 83 and 88.)
- [164] C. Imbimbo, *The spectrum of open string field theory at the stable tachyonic vacuum*, *Nucl. Phys.* **B770** (2007) 155–178, [[hep-th/0611343](#)]. (Cited on page 40.)
- [165] O.-K. Kwon, B.-H. Lee, C. Park, and S.-J. Sin, *Fluctuations around the tachyon vacuum in open string field theory*, [arXiv:0709.2888](#) [[hep-th](#)]. (Cited on page 40.)
- [166] L. Rastelli and B. Zwiebach, *Solving open string field theory with special projectors*, *JHEP* **01** (2008) 020, [[hep-th/0606131](#)]. (Cited on pages 41 and 42.)
- [167] D. Gaiotto, L. Rastelli, A. Sen, and B. Zwiebach, *Star algebra projectors*, *JHEP* **04** (2002) 060, [[hep-th/0202151](#)]. (Cited on pages 41 and 56.)
- [168] D. Gaiotto, L. Rastelli, A. Sen, and B. Zwiebach, *Ghost structure and closed strings in vacuum string field theory*, [hep-th/0111129](#). (Cited on pages 41, 56 and 89.)
- [169] M. Schnabl, *Anomalous reparametrizations and butterfly states in string field theory*, *Nucl. Phys.* **B649** (2003) 101–129, [[hep-th/0202139](#)]. (Cited on pages 41 and 56.)
- [170] Y. Okawa, L. Rastelli, and B. Zwiebach, *Analytic solutions for tachyon condensation with general projectors*, [hep-th/0611110](#). (Cited on page 42.)
- [171] A. Ishida, C. Kim, Y. Kim, O.-K. Kwon, and D. D. Tolla, *Tachyon vacuum solution in open string field theory with constant B field*, [0804.4380](#). (Cited on page 42.)
- [172] F. Sugino, *Witten's open string field theory in constant B-field background*, *JHEP* **03** (2000) 017, [[hep-th/9912254](#)]. (Cited on page 42.)
- [173] T. Kawano and T. Takahashi, *Open string field theory on noncommutative space*, *Prog. Theor. Phys.* **104** (2000) 459–476, [[hep-th/9912274](#)]. (Cited on page 42.)
- [174] E. Witten, *Noncommutative tachyons and string field theory*, [hep-th/0006071](#). (Cited on page 42.)

- [175] M. Schnabl, *String field theory at large B-field and noncommutative geometry*, *JHEP* **11** (2000) 031, [[hep-th/0010034](#)]. (Cited on page 42.)
- [176] K. Furuuchi, *Non-commutative space and Chan-Paton algebra in open string field algebra*, *Nucl. Phys.* **B640** (2002) 145–170, [[hep-th/0202200](#)]. (Cited on page 42.)
- [177] C. Maccaferri, R. J. Scherer Santos, and D. D. Tolla, *Time-localized projectors in string field theory with E-field*, *Phys. Rev.* **D71** (2005) 066007, [[hep-th/0501011](#)]. (Cited on page 42.)
- [178] A. Recknagel and V. Schomerus, *Boundary deformation theory and moduli spaces of D-branes*, *Nucl. Phys.* **B545** (1999) 233–282, [[hep-th/9811237](#)]. (Cited on page 43.)
- [179] A. Sen, *On the background independence of string field theory*, *Nucl. Phys.* **B345** (1990) 551–583. (Cited on pages 43 and 52.)
- [180] A. Sen, *On the background independence of string field theory. 2. Analysis of on-shell S matrix elements*, *Nucl. Phys.* **B347** (1990) 270–318. (Cited on pages 43 and 52.)
- [181] A. Sen, *On the background independence of string field theory. 3. Explicit field redefinitions*, *Nucl. Phys.* **B391** (1993) 550–590, [[hep-th/9201041](#)]. (Cited on pages 43 and 52.)
- [182] A. Sen and B. Zwiebach, *Large marginal deformations in string field theory*, *JHEP* **10** (2000) 009, [[hep-th/0007153](#)]. (Cited on page 43.)
- [183] B. Zwiebach, *A solvable toy model for tachyon condensation in string field theory*, *JHEP* **09** (2000) 028, [[hep-th/0008227](#)]. (Cited on page 43.)
- [184] A. Iqbal and A. Naqvi, *On marginal deformations in superstring field theory*, *JHEP* **01** (2001) 040, [[hep-th/0008127](#)]. (Cited on page 43.)
- [185] J. Kluson, *Marginal deformations in the open bosonic string field theory for N D0-branes*, *Class. Quant. Grav.* **20** (2003) 827–844, [[hep-th/0203089](#)]. (Cited on page 43.)
- [186] J. Kluson, *Exact solutions in SFT and marginal deformation in BCFT*, *JHEP* **12** (2003) 050, [[hep-th/0303199](#)]. (Cited on page 43.)
- [187] F. Katsumata, T. Takahashi, and S. Zeze, *Marginal deformations and closed string couplings in open string field theory*, *JHEP* **11** (2004) 050, [[hep-th/0409249](#)]. (Cited on page 43.)
- [188] A. Sen, *Energy momentum tensor and marginal deformations in open string field theory*, *JHEP* **08** (2004) 034, [[hep-th/0403200](#)]. (Cited on page 43.)
- [189] H. Yang and B. Zwiebach, *Testing closed string field theory with marginal fields*, *JHEP* **06** (2005) 038, [[hep-th/0501142](#)]. (Cited on page 43.)

- [190] I. Kishimoto and T. Takahashi, *Marginal deformations and classical solutions in open superstring field theory*, *JHEP* **11** (2005) 051, [[hep-th/0506240](#)]. (Cited on page 43.)
- [191] M. Kiermaier, Y. Okawa, L. Rastelli, and B. Zwiebach, *Analytic solutions for marginal deformations in open string field theory*, *JHEP* **01** (2008) 028, [[hep-th/0701249](#)]. (Cited on pages 43, 44, 45, 46, 47 and 80.)
- [192] M. Schnabl, *Comments on marginal deformations in open string field theory*, *Phys. Lett.* **B654** (2007) 194–199, [[hep-th/0701248](#)]. (Cited on pages 43, 44, 45, 46, 47 and 80.)
- [193] E. Fuchs, M. Kroyter, and R. Potting, *Marginal deformations in string field theory*, *JHEP* **09** (2007) 101, [[arXiv:0704.2222](#) [[hep-th](#)]]. (Cited on pages 44, 47, 50, 51 and 83.)
- [194] M. Kiermaier and Y. Okawa, *Exact marginality in open string field theory: a general framework*, [arXiv:0707.4472](#) [[hep-th](#)]. (Cited on pages 44, 47, 50, 52 and 83.)
- [195] T. Erler, *Marginal solutions for the superstring*, *JHEP* **07** (2007) 050, [[arXiv:0704.0930](#) [[hep-th](#)]]. (Cited on pages 44, 80 and 83.)
- [196] Y. Okawa, *Analytic solutions for marginal deformations in open superstring field theory*, *JHEP* **09** (2007) 084, [[arXiv:0704.0936](#) [[hep-th](#)]]. (Cited on pages 44, 80 and 83.)
- [197] S. Hellerman and M. Schnabl, *Light-like tachyon condensation in open string field theory*, [arXiv:0803.1184](#) [[hep-th](#)]. (Cited on page 45.)
- [198] E. Coletti, I. Sigalov, and W. Taylor, *Taming the tachyon in cubic string field theory*, *JHEP* **08** (2005) 104, [[hep-th/0505031](#)]. (Cited on page 46.)
- [199] G. Calcagni and G. Nardelli, *Tachyon solutions in boundary and cubic string field theory*, [0708.0366](#). (Cited on page 46.)
- [200] N. Jokela, M. Jarvinen, E. Keski-Vakkuri, and J. Majumder, *Disk partition function and oscillatory rolling tachyons*, *J. Phys.* **A41** (2008) 015402, [[0705.1916](#)]. (Cited on page 46.)
- [201] I. Ellwood, *Rolling to the tachyon vacuum in string field theory*, *JHEP* **12** (2007) 028, [[0705.0013](#)]. (Cited on page 46.)
- [202] I. Kishimoto and Y. Michishita, *Comments on solutions for nonsingular currents in open string field theories*, *Prog. Theor. Phys.* **118** (2007) 347–369, [[0706.0409](#)]. (Cited on page 46.)
- [203] O.-K. Kwon, *Marginally deformed rolling tachyon around the tachyon vacuum in open string field theory*, [arXiv:0801.0573](#) [[hep-th](#)]. (Cited on page 46.)
- [204] B.-H. Lee, C. Park, and D. D. Tolla, *Marginal deformations as lower dimensional D-brane solutions in open string field theory*, [arXiv:0710.1342](#) [[hep-th](#)]. (Cited on page 47.)

- [205] A. Belopolsky and B. Zwiebach, *Who changes the string coupling ?*, *Nucl. Phys.* **B472** (1996) 109–138, [[hep-th/9511077](#)]. (Cited on page 47.)
- [206] I. Bars and Y. Matsuo, *Associativity anomaly in string field theory*, *Phys. Rev.* **D65** (2002) 126006, [[hep-th/0202030](#)]. (Cited on page 52.)
- [207] E. Fuchs, M. Kroyter, and A. Marcus, *Squeezed state projectors in string field theory*, *JHEP* **09** (2002) 022, [[hep-th/0207001](#)]. (Cited on pages 55 and 56.)
- [208] T. G. Erler, *A fresh look at midpoint singularities in the algebra of string fields*, *JHEP* **03** (2005) 042, [[hep-th/0304044](#)]. (Cited on page 56.)
- [209] K. Okuyama, *Ratio of tensions from vacuum string field theory*, *JHEP* **03** (2002) 050, [[hep-th/0201136](#)]. (Cited on page 56.)
- [210] E. Fuchs and M. Kroyter, *On surface states and star-subalgebras in string field theory*, *JHEP* **10** (2004) 004, [[hep-th/0409020](#)]. (Cited on pages 56 and 57.)
- [211] S. Uhlmann, *A note on kappa-diagonal surface states*, *JHEP* **11** (2004) 003, [[hep-th/0408245](#)]. (Cited on page 56.)
- [212] A. Boyarsky and O. Ruchayskiy, *Integrability in SFT and new representation of KP tau-function*, *JHEP* **03** (2003) 027, [[hep-th/0211010](#)]. (Cited on page 56.)
- [213] L. Bonora and A. S. Sorin, *Integrable structures in string field theory*, *Phys. Lett.* **B553** (2003) 317–324, [[hep-th/0211283](#)]. (Cited on page 57.)
- [214] A. Boyarsky, B. Kulik, and O. Ruchayskiy, *String field theory vertices, integrability and boundary states*, *JHEP* **11** (2003) 045, [[hep-th/0307057](#)]. (Cited on page 57.)
- [215] C. Maccaferri and D. Mamone, *Star democracy in open string field theory*, *JHEP* **09** (2003) 049, [[hep-th/0306252](#)]. (Cited on page 57.)
- [216] A. Kling and S. Uhlmann, *String field theory vertices for fermions of integral weight*, *JHEP* **07** (2003) 061, [[hep-th/0306254](#)]. (Cited on page 57.)
- [217] M. Ihl, A. Kling, and S. Uhlmann, *String field theory projectors for fermions of integral weight*, *JHEP* **03** (2004) 002, [[hep-th/0312314](#)]. (Cited on page 57.)
- [218] E. Fuchs, M. Kroyter, and A. Marcus, *Virasoro operators in the continuous basis of string field theory*, *JHEP* **11** (2002) 046, [[hep-th/0210155](#)]. (Cited on pages 57 and 59.)
- [219] D. M. Belov, *Representation of small conformal algebra in kappa-basis*, [hep-th/0210199](#). (Cited on page 57.)
- [220] G. Moore and W. Taylor, *The singular geometry of the sliver*, *JHEP* **01** (2002) 004, [[hep-th/0111069](#)]. (Cited on page 57.)

- [221] D. M. Belov and A. Konechny, *On continuous Moyal product structure in string field theory*, *JHEP* **10** (2002) 049, [[hep-th/0207174](#)]. (Cited on pages 57 and 59.)
- [222] D. M. Belov and A. Konechny, *On spectral density of Neumann matrices*, *Phys. Lett.* **B558** (2003) 111–118, [[hep-th/0210169](#)]. (Cited on page 59.)
- [223] E. Fuchs, M. Kroyter, and A. Marcus, *Continuous half-string representation of string field theory*, *JHEP* **11** (2003) 039, [[hep-th/0307148](#)]. (Cited on page 59.)
- [224] D. M. Belov and C. Lovelace, *Star products made easy*, *Phys. Rev.* **D68** (2003) 066003, [[hep-th/0304158](#)]. (Cited on page 59.)
- [225] D. M. Belov, *Witten’s ghost vertex made simple (bc and bosonized ghosts)*, *Phys. Rev.* **D69** (2004) 126001, [[hep-th/0308147](#)]. (Cited on page 59.)
- [226] E. Fuchs and M. Kroyter, *Normalization anomalies in level truncation calculations*, *JHEP* **12** (2005) 031, [[hep-th/0508010](#)]. (Cited on page 60.)
- [227] I. Y. Aref’eva, R. Gorbachev, P. B. Medvedev, and D. V. Rychkov, *Descent relations and oscillator level truncation method*, [hep-th/0606070](#). (Cited on page 60.)
- [228] L. Bonora, C. Maccaferri, R. J. Scherer Santos, and D. D. Tolla, *Ghost story. I. Wedge states in the oscillator formalism*, *JHEP* **09** (2007) 061, [[arXiv:0706.1025 \[hep-th\]](#)]. (Cited on page 60.)
- [229] L. Bonora, R. J. Scherer Santos, and D. D. Tolla, *Spectral properties of ghost Neumann matrices*, *Phys. Rev.* **D77** (2008) 106001, [[0801.2099](#)]. (Cited on page 60.)
- [230] I. Y. Aref’eva, R. V. Gorbachev, P. B. Medvedev, and D. V. Rychkov, *Descent relations in cubic superstring field theory*, *JHEP* **01** (2008) 005, [[0704.3688](#)]. (Cited on page 60.)
- [231] E. Fuchs and M. Kroyter, *Universal regularization for string field theory*, *JHEP* **02** (2007) 038, [[hep-th/0610298](#)]. (Cited on page 61.)
- [232] D. Friedan, E. J. Martinec, and S. H. Shenker, *Conformal invariance, supersymmetry and string theory*, *Nucl. Phys.* **B271** (1986) 93. (Cited on pages 64 and 70.)
- [233] H. Sugawara, *A field theory of currents*, *Phys. Rev.* **170** (1968) 1659–1662. (Cited on page 65.)
- [234] J. N. Acosta, N. Berkovits, and O. Chandia, *A note on the superstring BRST operator*, *Phys. Lett.* **B454** (1999) 247–248, [[hep-th/9902178](#)]. (Cited on page 68.)
- [235] N. Berkovits, *Super-Poincare covariant quantization of the superstring*, *JHEP* **04** (2000) 018, [[hep-th/0001035](#)]. (Cited on page 68.)

- [236] N. Berkovits, *ICTP lectures on covariant quantization of the superstring*, [hep-th/0209059](#). (Cited on page 68.)
- [237] E. D'Hoker and D. H. Phong, *Lectures on two-loop superstrings*, [hep-th/0211111](#). (Cited on page 69.)
- [238] S. L. Cacciatori, F. Dalla Piazza, and B. van Geemen, *Modular forms and three loop superstring amplitudes*, *Nucl. Phys.* **B800** (2008) 565–590, [[0801.2543](#)]. (Cited on page 69.)
- [239] S. Grushevsky, *Superstring scattering amplitudes in higher genus*, [0803.3469](#). (Cited on page 69.)
- [240] M. Matone and R. Volpato, *Superstring measure and non-renormalization of the three-point amplitude*, [0806.4370](#). (Cited on page 69.)
- [241] A. Morozov, *NSR superstring measures revisited*, *JHEP* **05** (2008) 086, [[0804.3167](#)]. (Cited on page 69.)
- [242] C. Wendt, *Scattering amplitudes and contact interactions in Witten's superstring field theory*, *Nucl. Phys.* **B314** (1989) 209. (Cited on page 75.)
- [243] I. Y. Arefeva and P. B. Medvedev, *Anomalies in Witten's field theory of the NSR string*, *Phys. Lett.* **B212** (1988) 299. (Cited on page 75.)
- [244] B. V. Urošević and A. P. Zubarev, *On the component analysis of modified superstring field theory actions*, *Phys. Lett.* **B246** (1990) 391–398. (Cited on page 76.)
- [245] N. Berkovits, *A new approach to superstring field theory*, *Fortsch. Phys.* **48** (2000) 31–36, [[hep-th/9912121](#)]. (Cited on page 77.)
- [246] N. Berkovits, *Review of open superstring field theory*, [hep-th/0105230](#). (Cited on page 77.)
- [247] N. Berkovits, *The Ramond sector of open superstring field theory*, *JHEP* **11** (2001) 047, [[hep-th/0109100](#)]. (Cited on pages 77 and 81.)
- [248] B. Zwiebach, *Closed string field theory: Quantum action and the B-V master equation*, *Nucl. Phys.* **B390** (1993) 33–152, [[hep-th/9206084](#)]. (Cited on pages 77 and 89.)
- [249] B. Zwiebach, *Interpolating string field theories*, *Mod. Phys. Lett.* **A7** (1992) 1079–1090, [[hep-th/9202015](#)]. (Cited on pages 77 and 89.)
- [250] B. Zwiebach, *Oriented open-closed string theory revisited*, *Annals Phys.* **267** (1998) 193–248, [[hep-th/9705241](#)]. (Cited on page 77.)
- [251] Y. Okawa and B. Zwiebach, *Heterotic string field theory*, *JHEP* **07** (2004) 042, [[hep-th/0406212](#)]. (Cited on page 77.)
- [252] N. Berkovits, Y. Okawa, and B. Zwiebach, *WZW-like action for heterotic string field theory*, *JHEP* **11** (2004) 038, [[hep-th/0409018](#)]. (Cited on page 77.)

- [253] N. Berkovits and C. Vafa, *$N=4$ topological strings*, *Nucl. Phys.* **B433** (1995) 123–180, [[hep-th/9407190](#)]. (Cited on page 77.)
- [254] N. Berkovits and C. T. Echevarria, *Four-point amplitude from open superstring field theory*, *Phys. Lett.* **B478** (2000) 343–350, [[hep-th/9912120](#)]. (Cited on page 78.)
- [255] I. Y. Arefeva, D. M. Belov, and A. A. Giriyavets, *Construction of the vacuum string field theory on a non-BPS brane*, *JHEP* **09** (2002) 050, [[hep-th/0201197](#)]. (Cited on page 79.)
- [256] Y. Okawa, *Real analytic solutions for marginal deformations in open superstring field theory*, *JHEP* **09** (2007) 082, [[arXiv:0704.3612 \[hep-th\]](#)]. (Cited on pages 80 and 83.)
- [257] E. Fuchs and M. Kroyter, *On the classical equivalence of superstring field theories*, [0805.4386](#). (Cited on pages 81, 82, 86 and 87.)
- [258] A. Kling, O. Lechtenfeld, A. D. Popov, and S. Uhlmann, *Solving string field equations: New uses for old tools*, *Fortsch. Phys.* **51** (2003) 775–780, [[hep-th/0212335](#)]. (Cited on page 81.)
- [259] M. Kiermaier and Y. Okawa, *General marginal deformations in open superstring field theory*, [arXiv:0708.3394 \[hep-th\]](#). (Cited on page 83.)
- [260] I. Y. Arefeva, P. B. Medvedev, and A. P. Zubarev, *Nonperturbative vacuum for superstring field theory and supersymmetry breaking*, *Mod. Phys. Lett.* **A6** (1991) 949–958. (Cited on page 85.)
- [261] I. Y. Aref’eva, R. V. Gorbachev, and P. B. Medvedev, *Tachyon solution in cubic Neveu-Schwarz string field theory*, [0804.2017](#). (Cited on pages 85 and 86.)
- [262] I. Ellwood, *The closed string tadpole in open string field theory*, [0804.1131](#). (Cited on pages 87 and 89.)
- [263] T. Kawano, I. Kishimoto, and T. Takahashi, *Gauge invariant overlaps for classical solutions in open string field theory*, *Nucl. Phys.* **B803** (2008) 135–165, [[0804.1541](#)]. (Cited on pages 87 and 89.)
- [264] I. Y. Aref’eva, L. V. Joukovskaya, and A. S. Koshelev, *Time evolution in superstring field theory on non-BPS brane. I: Rolling tachyon and energy-momentum conservation*, *JHEP* **09** (2003) 012, [[hep-th/0301137](#)]. (Cited on page 88.)
- [265] A. Bagchi and A. Sen, *Tachyon condensation on separated brane-antibrane system*, *JHEP* **05** (2008) 010, [[0801.3498](#)]. (Cited on page 88.)
- [266] A. Hashimoto and N. Itzhaki, *Observables of string field theory*, *JHEP* **01** (2002) 028, [[hep-th/0111092](#)]. (Cited on page 89.)
- [267] N. Drukker, *Closed string amplitudes from gauge fixed string field theory*, *Phys. Rev.* **D67** (2003) 126004, [[hep-th/0207266](#)]. (Cited on page 89.)

- [268] T. Takahashi and S. Zeze, *Closed string amplitudes in open string field theory*, *JHEP* **08** (2003) 020, [[hep-th/0307173](#)]. (Cited on page 89.)
- [269] L. Bonora, C. Maccaferri, R. J. Scherer Santos, and D. D. Tolla, *Bubbling AdS and vacuum string field theory*, *Nucl. Phys.* **B749** (2006) 338–357, [[hep-th/0602015](#)]. (Cited on page 89.)
- [270] L. Bonora, N. Bouatta, and C. Maccaferri, *Towards open-closed string duality: Closed strings as open string fields*, [hep-th/0609182](#). (Cited on page 89.)
- [271] Y. Michishita, *On-shell gauge invariants and field strengths in open superstring field theory*, *Nucl. Phys.* **B698** (2004) 111–131, [[hep-th/0406242](#)]. (Cited on page 89.)
- [272] T. Kawano, I. Kishimoto, and T. Takahashi, *Schnabl’s solution and boundary states in open string field theory*, [0804.4414](#). (Cited on page 89.)
- [273] H. Yang and B. Zwiebach, *Rolling closed string tachyons and the big crunch*, *JHEP* **08** (2005) 046, [[hep-th/0506076](#)]. (Cited on page 90.)
- [274] H. Yang and B. Zwiebach, *A closed string tachyon vacuum?*, *JHEP* **09** (2005) 054, [[hep-th/0506077](#)]. (Cited on page 90.)
- [275] O. Bergman and S. S. Razamat, *Toy models for closed string tachyon solitons*, *JHEP* **11** (2006) 063, [[hep-th/0607037](#)]. (Cited on page 90.)
- [276] N. Moeller and H. Yang, *The nonperturbative closed string tachyon vacuum to high level*, *JHEP* **04** (2007) 009, [[hep-th/0609208](#)]. (Cited on page 90.)
- [277] N. Moeller, *Closed bosonic string field theory at quintic order: Five-tachyon contact term and dilaton theorem*, *JHEP* **03** (2007) 043, [[hep-th/0609209](#)]. (Cited on page 90.)
- [278] N. Moeller, *Closed bosonic string field theory at quintic order. II: Marginal deformations and effective potential*, *JHEP* **09** (2007) 118, [[arXiv:0705.2102](#) [[hep-th](#)]]. (Cited on page 90.)
- [279] N. Moeller, *A tachyon lump in closed string field theory*, [0804.0697](#). (Cited on page 90.)