

Oxidation of Self-Duality to 12 Dimensions and Beyond

Chandrashekar Devchand

Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), Am Mühlenberg 1, 14476 Potsdam, Germany. E-mail: devchand@math.uni-potsdam.de

Received: 25 January 2013 / Accepted: 12 December 2013
© Springer-Verlag Berlin Heidelberg 2014

Abstract: Using (partial) curvature flows and the transitive action of subgroups of $O(d, \mathbb{Z})$ on the indices $\{1, \dots, d\}$ of the components of the Yang–Mills curvature in an orthonormal basis, we obtain a nested system of equations in successively higher dimensions d , each implying the Yang–Mills equations on d -dimensional Riemannian manifolds possessing special geometric structures. This ‘matryoshka’ of self-duality equations contains the familiar self-duality equations on Riemannian fourfold as well as their generalisations on complex Kähler threefold and on seven- and eight-dimensional manifolds with G_2 and $\text{Spin}(7)$ holonomy. The matryoshka allows enlargement (‘oxidation’) to a remarkable system in 12 dimensions, invariant under $\text{Sp}(3)$. There are hints that the underlying geometry is related to the sextonions, a six-dimensional algebra between the quaternions and octonions.

1. Introduction

Many interesting examples of special geometric structures on d -dimensional Riemannian manifolds (M, g) are provided by certain G -invariant covariantly constant (parallel) p -forms $\varphi \in \Lambda^p T^*M$, where $G = \text{Hol}$, the restricted holonomy group of M . If $p < d$, then G is clearly a proper subgroup of $\text{SO}(d)$, since in the generic rotationally invariant case, only the volume form is invariant.

For Riemannian manifolds which are locally neither a product of lower dimensional spaces nor a symmetric space, Berger’s list [1] provides the most interesting examples of restricted holonomy groups. These include $U(n) \subset \text{SO}(2n)$, which leaves the Kähler two-form ω on a $2n$ -dimensional Kähler manifold invariant. The $SU(n)$ Calabi-Yau specialisation has, in addition, an invariant complex n -form, the holomorphic volume form. The group $\text{Sp}(n) \subset \text{SO}(4n)$, $d = 4n$, of $n \times n$ matrices with quaternion elements satisfying $A^\dagger A = 1$, has three invariant Kähler two-forms ω_α , combinable in a two-form, $\omega = \omega_1 i + \omega_2 j + \omega_3 k$, taking values in the imaginary quaternions. These characterise hyper-Kähler geometry. The quaternionic Kähler generalisation has $\text{Hol} = \text{Sp}(n) \cdot$

$\text{Sp}(1) \subset \text{SO}(4n)$, with the three Kähler forms existing only locally. Globally, they define an invariant parallel four-form $\sum \omega_\alpha \wedge \omega_\alpha$. The two exceptional $d = 7$ and 8 geometries with $\text{Hol} = G_2$ and $\text{Spin}(7)$ have, respectively, an invariant three- and four-form. In all these cases, the geometric information can be equally well encoded uniformly in an invariant four-form: the two-forms afford squaring and the three-form in seven dimensions has a Hodge-dual four-form. The Lie group inclusions

$$\text{Sp}(n) \subset \text{SU}(2n) \subset \text{U}(2n) \subset \text{SO}(4n)$$

imply corresponding inclusions of geometries: hyperkähler manifolds are Calabi-Yau manifolds, the latter are Kähler, which in turn are orientable. The two exceptional cases are also part of lower dimensional sets of inclusions:

$$\begin{aligned} \text{U}(2) &\subset \text{Sp}(2) \subset \text{SU}(4) \subset \text{Spin}(7) \subset \text{SO}(8) \\ \text{SU}(3) &\subset G_2 \subset \text{Spin}(7) \subset \text{SO}(8). \end{aligned}$$

The respective invariant tensors can be obtained by successive reductions of the $4n$ -dimensional volume form. For instance, the $\text{Spin}(7)$ invariant four-form in eight dimensions contracted with an arbitrary vector yields the G_2 -invariant three-form in the orthogonal seven-dimensional space. Similarly, the latter yields an $\text{SU}(3)$ -invariant two-form on projection to the complex threefold orthogonal to an arbitrary vector.

For Riemannian manifolds (M, g) admitting a G -structure, a principle subbundle of the frame bundle of M , with structure group $G \subset \text{GL}(d, \mathbb{R})$, the tangent space at every point admits an isomorphism with \mathbb{R}^d . For every point $p \in M$ there exists a choice of local coordinates with p as the origin in which the Riemannian metric takes the euclidean form $d^2s = g_{ij}dx^i dx^j = \sum_i dx^i dx^i$ and the special geometric structure φ in these coordinates is the constant G -invariant form

$$\varphi = \sum_{(i_1, \dots, i_p) \in \mathcal{J}^+} dx^{i_1, \dots, i_p}. \quad (1)$$

where $dx^{i_1, \dots, i_p} := dx^{i_1} \wedge \dots \wedge dx^{i_p}$ and \mathcal{J}^+ is a set of oriented subsets $\{i_1, \dots, i_p\} \subset \{1, \dots, d\}$ with $\varphi_{i_1 \dots i_p} = 1$. Differential forms like φ have been called *special democratic forms* [2,3]. They are ‘special’ in the sense that they have components $\varphi_{\mu_1 \dots \mu_p}$ equal to $+1, -1$ or 0 in some orthonormal basis, just like the volume form $\text{vol}_d = dx^1 \wedge dx^2 \wedge \dots \wedge dx^d =: dx^{12 \dots d}$ on a Euclidean vector space. More precisely, a p -form φ is called *special* if it lies in the $\text{SO}(d, \mathbb{R})$ -orbit of

$$\varphi = \sum_{1 \leq \mu_1 < \dots < \mu_p \leq d} \varphi_{\mu_1 \dots \mu_p} dx^{\mu_1 \dots \mu_p} \quad (2)$$

with components $\varphi_{\mu_1 \dots \mu_p} \in \{-1, 0, 1\}$. There are clearly only a finite number of orbits of special p -forms parametrised by the components $\varphi_{\mu_1 \dots \mu_p} \in \{-1, 0, 1\}$ under $\text{SO}(d, \mathbb{R})$ or $\text{O}(d, \mathbb{R})$. Distinct sets of components may give rise to special p -forms in the same orbit, because the subgroups $\text{SO}(d, \mathbb{Z}) \subset \text{SO}(d, \mathbb{R})$ or $\text{O}(d, \mathbb{Z}) \subset \text{O}(d, \mathbb{R})$ map the special form φ in Eq. (1) into a special form parametrised by different components. These groups are isomorphic to the semidirect product of the permutation group S_d acting naturally on $d-1$ or d copies of \mathbb{Z}_2 , namely $\text{SO}(d, \mathbb{Z}) \cong S_d \times \mathbb{Z}_2^{d-1}$ or $\text{O}(d, \mathbb{Z}) \cong S_d \times \mathbb{Z}_2^d$. Thus, special p -forms which appear to be different may nevertheless be in the same orbit under $\text{SO}(d, \mathbb{R})$ or $\text{O}(d, \mathbb{R})$. The orbit of a special p -form may always be labelled by a choice of a representative (1).

A special p -form φ is called democratic if its set of nonzero components $\{\varphi_{i_1 \dots i_p}\}$ is symmetric under the transitive action of a subgroup of $O(d, \mathbb{Z})$ on the indices $\{1, \dots, d\}$. The action of an element $(\sigma, \eta_1, \dots, \eta_d) \in S_d \times \mathbb{Z}_2^d$, on the components of φ being given by

$$\varphi_{i_1 \dots i_p} \mapsto \eta_{i_1} \dots \eta_{i_p} \varphi_{\sigma(i_1) \dots \sigma(i_p)}, \quad (3)$$

where $\eta_i^2 = 1$, $i = 1, \dots, d$. So for a democratic form, no choice of indices is privileged. We refer to [2,3] for further details. It was shown in [2] that knowledge of the above symmetry groups allows an enlargement (oxidation) of the base space; the symmetries may be used to remix the sets of indices $\{(i_1 \dots i_p)\}$ of the nonzero components amongst a larger set of indices $\{1, \dots, D\}$, $D > d$, thus defining special democratic P -forms in D dimensions from special democratic p -forms in d dimensions for successively higher $P \geq p$ and $D \geq d$. In this paper, we consider two such oxidation maps:¹

(a) Oxidation through remixing

This is a map $\varphi \in \Lambda^p \mathbb{R}^d \rightarrow \Lambda^p \mathbb{R}^D \ni \Phi$ defining a special democratic p -form Φ in $D > d$ dimensions in terms of the components of a special p -form φ in d -dimensions thus:

$$\varphi = \sum_{(i_1, \dots, i_p) \in \mathcal{J}^+} dx^{i_1, \dots, i_p} \mapsto \Phi = \sum_{\sigma \in H \subset S_D} \sum_{(i_1, \dots, i_p) \in \mathcal{J}^+} dx^{\sigma(i_1) \dots \sigma(i_p)}, \quad (4)$$

where H is some subgroup of the symmetric group S_D acting on the D indices.

(b) Oxidation through heat flow

Alternatively, for $D = d+q$ the nonzero components of a special democratic $P = p+q$ -form are given by a map $\varphi \in \Lambda^p \mathbb{R}^d \rightarrow \Lambda^{p+q} \mathbb{R}^{d+q} \ni \Phi$ defined by

$$\varphi = \sum_{(i_1, \dots, i_p) \in \mathcal{J}^+} dx^{i_1, \dots, i_p} \mapsto \Phi = \sum_{\sigma \in H \subset S_D} \sum_{(i_1, \dots, i_p) \in \mathcal{J}^+} dx^{\sigma(i_1) \dots \sigma(i_p) \sigma(d+1) \dots \sigma(D)}. \quad (5)$$

Using these mappings, a nested structure of special forms in successively higher dimensions emerges. This is reminiscent of a matryoshka (матрёшка), a set of nested Russian dolls, traditionally carved in wood, where the inner surface of each doll is basically a copy of the outer surface of the previous doll; but the outer surface can then vary somewhat, depending on the geometry of the bulk.

A remarkable nested structure of special democratic forms was displayed in [2], which included a $U(3)$ -invariant two-form in six dimensions, a G_2 -invariant three-form in seven dimensions, and a $Spin(7)$ -invariant four-form in eight dimensions; corresponding to the embeddings $SU(3) \subset G_2 \subset Spin(7)$ mentioned above. It was also shown that this matryoshka with three dolls fits into even larger dolls and interesting properties of a special democratic six-form in ten dimensions were presented.

Motivated by the discussion in [2] of nested special democratic forms, we shall presently show that there exists a corresponding matryoshka of self-duality equations in successively higher dimensions; each implying the Yang–Mills equations, just as

¹ The notion of using the inverse of dimensional reduction as a means of searching for higher dimensional parents of lower dimensional theories goes back to early work on supergravity by Julia [4].

four-dimensional self-duality [5]. Successive sets of equations are ‘oxidised’ to higher dimensions and ‘reduced’ to lower dimensions by enhancing or restricting the permutation symmetries on the sets of indices of special geometric tensors. Remarkably, the simplest case of the mapping (5), with $q = D - d = 1$ corresponds to equations for (partial) curvature flows for the vector potentials, hence ‘Oxidation through heat flow’. Solutions of the lower dimensional equations then provide initial values for the flow into the extra dimension, the flow to the next doll of the matryoshka. We shall display oxidations up to $d = 16$. The representation theory underlying the 12 dimensional system seems to be related to a mathematical curiosity, the algebra of the *sextonions* [6,7], a six-dimensional algebra between quaternions and octonions. This algebra gives rise to a new row in Freudenthal’s magic chart, corresponding to a (non-simple) Lie algebra between \mathfrak{e}_7 and \mathfrak{e}_8 , which has been called $\mathfrak{e}_{7\frac{1}{2}}$ [7].

2. Generalised Duality for Gauge Fields in $d > 4$

Generalisations of the four-dimensional self-duality equations to higher dimensions were introduced some time ago in [5], where it was shown that restrictions of the Yang–Mills curvature two-form F to an eigenspace of a four-form T , implies the Yang–Mills equations. In a standard orthonormal basis of T^*M these take the form,

$$\frac{1}{2}g^{km}g^{ln}T_{ijkl}F_{mn} = \lambda F_{ij}, \quad i, j, \dots = 1, \dots, d. \quad (6)$$

Here T_{mnpq} is a covariantly constant tensor, g^{pr} the inverse metric tensor and $F = dA + A \wedge A$ is the curvature of a connection $D = d + A$ on a Riemannian d -fold (M, g) with values in the Lie algebra of a real gauge group contained in $\text{GL}(n, \mathbb{R})$. These partial-flatness conditions on the curvature are first order equations for the vector potentials A , so they are more amenable to solution than the second order Yang–Mills equations. Indeed, many special solutions are known (see e.g. [9–11]). The usefulness of the linear curvature constraints (6) follows from the observation [5]:

Theorem 1. *For nonzero eigenvalues λ , the conditions (6) imply the Yang–Mills equations $g^{ij}D_i F_{jk} = 0$. Thus, potentials A satisfying these first order equations automatically satisfy the Yang–Mills equations.*

This result follows in virtue of the Bianchi identities $D_{[i}F_{mn]} \equiv 0$. In [5], constant four-forms T in flat euclidean spaces were considered, but it is clear that, more generally [12], it suffices for the consistency condition

$$g^{km}g^{ln}(g^{ip}\nabla_p T_{ijkl})F_{mn} = 0 \quad (7)$$

to hold, which follows if T is co-closed, $g^{ip}\nabla_p T_{ijkl} = 0$. The latter in turn follows if T is parallel (i.e. covariantly constant) with respect to the Levi-Civita connection ∇ . In dimensions $d > 4$, the four-form T clearly breaks the d -dimensional rotational invariance of the Yang–Mills equations. Examples of four-forms and the corresponding partial-flatness conditions (6) invariant under various subgroups $G \subset \text{SO}(d)$ were studied in [5] for dimensions $4 < d \leq 8$. In particular, interesting examples invariant under $(\text{SU}(n) \otimes \text{U}(1))/\mathbb{Z}_2$ and $\text{SU}(n)$, G_2 and $\text{Spin}(7)$, in dimensions $d = 2n, 7, 8$ were constructed. The example of $\text{Sp}(n) \otimes \text{Sp}(1)/\mathbb{Z}_2$ was discussed shortly thereafter in [13, 14]. The above groups are precisely the holonomy groups of Calabi-Yau, quaternionic Kähler and exceptional holonomy manifolds, so remarkably, the generalisations of self-duality

for most of Berger's special holonomy manifolds [1] were unwittingly constructed before the subject acquired widespread differential geometric interest (e.g. [12, 15–20]). On all the above manifolds, there exists a ∇ -parallel four-form, so the above-mentioned consistency condition on T is satisfied.

On a d -dimensional Riemannian manifold (M, g) the volume form vol^g , associated to the metric and orientation, is given in local coordinates by $\text{vol}^g = \sqrt{\det(g)} dx^1 \wedge \dots \wedge dx^d$. The Hodge star operator on the space of p -forms $*$: $\Lambda^p T^*M \rightarrow \Lambda^{d-p} T^*M$ is defined by $\alpha \wedge * \beta = (\alpha, \beta) \text{vol}^g$, where $\alpha, \beta \in \Lambda^p T^*M$ and (α, β) is the natural inner product induced on p -forms by the metric; $(\alpha, \beta) = \alpha_{i_1 \dots i_p} \beta_{j_1 \dots j_p} g^{i_1 j_1} \dots g^{i_p j_p}$. Given the existence of a G -invariant four-form T on M , the space of two-forms naturally decomposes into its T -eigenspaces, $\Lambda^2 T^*M = \bigoplus_{\lambda \in \sigma_T} V_\lambda$, where σ_T is the spectrum of T and the eigenspaces V_λ are G -modules. The endomorphism defined by the four-form T on the space of two-forms is traceless, so $\sum_{\lambda \in \sigma_T} d_\lambda \lambda = 0$; $d_\lambda := \dim V_\lambda$.

The Yang–Mills equations are the Euler–Lagrange equations for the Yang–Mills functional defined on the space of vector potentials,

$$S = \|F\|^2 = \langle F, F \rangle := \int_M \text{Tr} (F, F) \text{vol}^g = \int_M \text{Tr} F \wedge *F. \quad (8)$$

The topological functional associated with a co-closed four-form T yields a Chern–Simons-like form on the boundary of M ,

$$Q = \int_M \text{Tr} *T \wedge F \wedge F = \int_M \text{Tr} d(*T \wedge (A \wedge dA + \frac{2}{3} A \wedge A \wedge A)). \quad (9)$$

This functional was used in [21] to construct examples of higher dimensional analogues of topological field theories; its relation to the topology of the bundle for some specific examples has been discussed, for instance, in [18, 20]. The curvature decomposes into its T -eigenspaces, $F = \sum_{\lambda \in \sigma_T} F_\lambda$, where the components $F_\lambda \in V_\lambda$ satisfy (6) and are mutually orthogonal with respect to the L^2 inner product in (8); i.e. $\langle F_\lambda, F_\mu \rangle = 0$ for $\lambda \neq \mu$. Equation (6) may be expressed in terms of the projection operator $P: \Lambda^2 T^*M \rightarrow (V_\lambda)^\perp$ to the orthogonal complement of the eigenspace V_λ (see e.g. [22]),

$$F = F_\lambda \Leftrightarrow P_{ij}^I F^{ij} = 0, \quad I = 1, \dots, \bar{d}_\lambda, \quad (10)$$

where the number of equations, \bar{d}_λ , is the codimension of the eigenspace V_λ with eigenvalue λ . Here, the projector P is the analogue of the 't Hooft tensor in four dimensions and we lower (raise) indices using the (inverse) Riemannian metric. The most interesting sets of equations (10) correspond to the projections to the largest eigenspace of T ; these being the least overdetermined systems. We choose the convenient orientation and normalisation of the four-form T in which this largest eigenspace has eigenvalue $\lambda = -1$, corresponding in four dimensions to anti-self-duality.

Now, the eigenspace decomposition of F , together with Eq. (6), implies that

$$*T \wedge F = \sum_{\lambda \in \sigma_T} *T \wedge F_\lambda = \sum_{\lambda \in \sigma_T} \lambda * F_\lambda.$$

So the topological functional in (9) has the spectral decomposition

$$Q = \int_M \text{Tr} *T \wedge F \wedge F = \sum_{\lambda \in \sigma_T} \lambda * F_\lambda \wedge F_\lambda = \sum_{\lambda \in \sigma_T} \lambda \|F_\lambda\|^2. \quad (11)$$

Similarly, the Yang–Mills action takes the form $S = \sum_{\lambda \in \sigma_T} \|F_\lambda\|^2$. Picking out a specific sector of the theory with eigenvalue μ , we may write

$$Q = \mu \|F_\mu\|^2 + \sum_{\lambda \in \sigma_T \setminus \{\mu\}} \lambda \|F_\lambda\|^2. \quad (12)$$

It follows that

$$S = \|F_\mu\|^2 + \sum_{\lambda \in \sigma_T \setminus \{\mu\}} \|F_\lambda\|^2 = \frac{1}{\mu} Q + \sum_{\lambda \in \sigma_T} \left(1 - \frac{\lambda}{\mu}\right) \|F_\lambda\|^2. \quad (13)$$

Thus, if $(1 - \frac{\lambda}{\mu}) \geq 0$ for all $\lambda \in \sigma_T$, then the action is bounded below:

$$S = \|F\|^2 \geq \frac{1}{\mu} Q. \quad (14)$$

This L^2 -bound on the curvature is saturated precisely when F is projected to the eigenspace with eigenvalue μ . Vector potentials satisfying $F(A) = F_\mu(A)$ thus minimise the functional S , with the minimal action taking the value $\|F_\mu(A)\|^2$. The bound (14) shows that manifolds with a co-closed four-form T differ essentially from spheres. It is known that a connection over $M = S^d$, $d \geq 5$, with sufficiently small L^2 -norm is necessarily flat [23,24]. This reflects the argument in [25] that in flat spaces of dimension $d \geq 5$, the only finite action Yang–Mills solutions are pure-gauge with $F = 0$. In the case of more general compact manifolds M , curvature estimates have been discussed in [26,27]. Using these results, Tian [20] has proven the existence of finite action solutions to (6) on manifolds M with a co-closed four-form T satisfying certain conditions, in particular when T is a certain calibrating form [28]. Further, in certain cases, the moduli space of equivalence classes of solutions to (6) has also been shown to be a compact manifold [20].

Note that if $a := (1 - \frac{\lambda}{\mu})$ does not have the same sign for every $\lambda \neq \mu$, so that the *difference* of two sums of squares (the terms with $a > 0$ and those with $a < 0$) gives the action plus a topological term, then the equation $F(A) = F_\mu(A)$ determines saddle points of the Yang–Mills functional. This yields a theory with no finite lower bound to the action and solutions A are necessarily unstable under those infinitesimal deformations δA which contribute to curvature components F_λ , where $\lambda/\mu > 1$. For such solutions the second variation of the action functional $\delta^2 S / \delta A^2$ is not non-negative. Particular examples of such systems of equations for saddle points of the Yang–Mills action have been known for some time [29]. The quantum field theoretic significance of negative eigenvalues of the second variation of the action has been discussed for instance in [30,31].

In even dimensions, with $d = 2n$, if the manifold M admits a complex structure J , this provides, at any point p in M , a linear map $J_p: T_p^* M \rightarrow T_p^* M$ under which the complexification $T_p^* M \otimes_{\mathbb{R}} \mathbb{C}$ splits into the eigenspaces $T_p^{(1,0)} M$ and $T_p^{(0,1)} M$, both of which are isomorphic to \mathbb{C}^n . This allows the choice of complex coordinates (z^1, \dots, z^n) and $(z^{\bar{1}}, \dots, z^{\bar{n}})$. The complex (1,0)- and (0,1)-forms $\{dz^\alpha\}$ and $\{dz^{\bar{\alpha}}\}$, for $\alpha, \bar{\alpha} = 1, \dots, n$, then provide bases for $T_p^{(1,0)} M$ and $T_p^{(0,1)} M$ respectively. Imposing the reality conditions $dz^{\bar{\alpha}} = \overline{dz^\alpha}$, we may recover $\mathbb{R}^{2n} \simeq \mathbb{C}^n$. The curvature two-form in this basis has components $F_{\alpha\beta}, F_{\alpha\bar{\beta}}, F_{\bar{\alpha}\bar{\beta}} = \overline{F_{\alpha\beta}}$ and the Riemannian metric locally takes the

hermitian form $d^2s = g_{\alpha\bar{\beta}}dz^\alpha dz^\beta = dz^\alpha dz_\alpha = \sum_\alpha dz^\alpha dz^{\bar{\alpha}}$ and the complex $(n, 0)$ volume form is given by $\Omega = dz^{\alpha_1 \dots \alpha_n}$. In the complex setting, Eq. (6) is a G -invariant equation, where the structure group G is a subgroup of $\text{GL}(n, \mathbb{C}) \subset \text{GL}(2n, \mathbb{R})$. For the particularly important $\lambda = -1$ case, we shall use the following complex variant of Theorem 1.

Theorem 2. *On a Riemannian complex n -fold (M^{2n}, g) , with hermitian metric $g = g_{\alpha\bar{\beta}}dz^\alpha dz^\beta$ and $(4,0)$ -form Φ , the linear curvature constraints,*

$$F_{\alpha\beta} + \frac{1}{2}g^{\gamma\bar{\eta}}g^{\delta\bar{\kappa}}\Phi_{\alpha\beta\gamma\delta}F_{\bar{\eta}\bar{\kappa}} = 0, \quad (15)$$

$$g^{\alpha\bar{\beta}}F_{\alpha\bar{\beta}} = 0, \quad (16)$$

$$g^{\gamma\bar{\eta}}g^{\delta\bar{\kappa}}(g^{\alpha\bar{\rho}}\nabla_{\bar{\rho}}\Phi_{\alpha\beta\gamma\delta})F_{\bar{\eta}\bar{\kappa}} = 0, \quad (17)$$

imply the Yang–Mills equations $g^{\alpha\bar{\rho}}D_{\bar{\rho}}F_{\alpha\beta} = g^{\alpha\bar{\rho}}D_{\bar{\rho}}F_{\alpha\bar{\beta}} = 0$.

Proof. Using (15) we have

$$g^{\alpha\bar{\rho}}D_{\bar{\rho}}F_{\alpha\beta} = D^\alpha F_{\alpha\beta} = -\frac{1}{2}(\nabla^\alpha\Phi_{\alpha\beta\gamma\delta})F^{\gamma\delta} - \frac{1}{2}\Phi_{\alpha\beta\gamma\delta}D^\alpha F^{\gamma\delta} = 0, \quad (18)$$

the first term being the left side of (17) and the second vanishes in virtue of the Bianchi identity $D_{\bar{\alpha}}F_{\bar{\gamma}\bar{\delta}} + \text{cyclic permutations} = 0$. Similarly, using the Bianchi identity between $D_{\bar{\rho}}$, $D_{\bar{\beta}}$ and D_a we have, $D_{\bar{\rho}}F_{\alpha\bar{\beta}} = D_{\bar{\beta}}F_{\alpha\bar{\rho}} + D_\alpha F_{\bar{\alpha}\bar{\rho}}$. On contracting with $g^{\alpha\bar{\rho}}$, the second term on the right yields the complex conjugate of the left side of (18) and the first term contains the trace of the $(1,1)$ -part of the curvature, which vanishes by Eq. (16). \square

Already in [5], it was noticed that the lower dimensional cases, including four-dimensional self-duality, the six-dimensional $\text{SU}(3)\otimes\text{U}(1)/\mathbb{Z}_2$ -invariant equations and the seven-dimensional G_2 -invariant equations, were reductions of the eight-dimensional $\text{Spin}(7)$ -invariant set of equations. In the present paper, we show that using the results of [2] these equations also admit a systematic ‘oxidation’ to higher dimensions starting from the lower dimensional ones.

We consider two types of oxidation. The first is based on the map (4) and uses cyclic permutations to remix the index sets appearing in the lower dimensional equations amongst a larger set of indices. The second oxidation method is based on the heat flow for some appropriate partial curvature. This is related to the $D - d = 1$ case of (5). More specifically, if in $(d-1)$ -dimensions, there exists a special set of $d-1$ curvature constraints $f_{ijk}F^{jk} = 0$, $i = 1, \dots, d-1$, where f is some appropriate tensor, then we can consider the corresponding partial curvature flow

$$\dot{A}_i = f_{ijk}F^{jk}, \quad i = 1, \dots, d. \quad (19)$$

Identifying the parameter of the variation or ‘time’ of the flow with a d th independent variable x^d , the left hand side is the $A_d = 0$ ‘temporal’ gauge form of the curvature components F_{di} , so that the flow equations (19) are in fact linear curvature constraints of the form

$$F_{di} = f_{ijk}F^{jk}. \quad (20)$$

Remarkably, in many interesting cases, these constraints may be reformulated in the form (6), thus implying the Yang–Mills equations. The idea of choosing such a temporal gauge to obtain a flow equation is not new. For instance, both Nahm’s equations for magnetic monopoles [32] and the generalisations to higher dimensions of Euler’s equations for a spinning top [33], arise from the imposition of precisely such a gauge choice on equations of the form (6). Conversely, the flow equations can be written as gauge covariant equations in one dimension higher by gauge un-fixing the component of the gauge potential in the direction of the flow. This idea has also been used by Tao [34] in the context of the full second order Yang–Mills gradient flow $\dot{A}_k = g^{ij} D_i F_{jk}$.

As we shall see, the juxtaposition of the two oxidation methods above yields the advertised matryoshka of self-duality equations, starting from zero curvature in $d = 2$ and including the familiar four-dimensional self-duality, as well as its generalisations to six, seven and eight dimensions mentioned above. Remarkably, the matryoshka affords enlargement to even higher dimensions. We discuss an interesting 12 dimensional extension and display its oxidation to 14 and 16 dimensions.

3. The Matryoshka of Self-Duality Equations

Let us begin in two dimensions with the flatness condition $F_{12} = 0$ for the sole component of the curvature two-form. In the complex setting, the curvature only has a (1,1)-part, $F_{z\bar{z}}$, where we use complex coordinates $z = x^1 + ix^2$, $\bar{z} = x^1 - ix^2$. The flatness condition means that the curvature is in the kernel of the volume form. We therefore have,

$$\epsilon_{ij} F^{ij} = 0 \Leftrightarrow F_{12} = 0 \Leftrightarrow F_{z\bar{z}} = 0. \quad (21)$$

Both real and complex forms of the equations are locally rotationally invariant, since their respective invariance algebras $\mathfrak{so}(2)$ and $\mathfrak{u}(1)$ are isomorphic. The rich properties of the solutions of these equations on Riemann surfaces have been investigated by Atiyah and Bott [35].

We oxidise the equation $F_{12} = 0$ to a system in three dimensions by acting on the indices by all permutations generated by the cycle $\sigma = (1\ 2\ 3) \in S_3$, so as to obtain a system of equations invariant under these permutations:

$$\{F_{12} = 0\} \longrightarrow \{F_{\sigma^p(1)\sigma^p(2)} = 0; \sigma = (1\ 2\ 3), p = 1, 2\}. \quad (22)$$

This of course yields flatness in three dimensions; the curvature lies in the kernel of the three-dimensional volume form,

$$\epsilon_{ijk} F^{jk} = 0 \Leftrightarrow F_{12} = F_{23} = F_{31} = 0. \quad (23)$$

Since this is a set of three equations for the three vector potentials A_i , $i = 1, 2, 3$, it allows us to write the Yang–Mills curvature flow

$$\frac{\partial}{\partial x^4} A_i(x^i, x^4) = \frac{1}{2} \epsilon_{ijk} F^{jk}, \quad i = 1, 2, 3, \quad (24)$$

with initial (at $x^4 = 0$) flat connection $A_i(x^i, 0)$ satisfying (23). This evolution equation is the gradient flow of the Chern–Simons functional [36] on the space of connections

$$S_{CS} = \int_{M^3} \text{Tr} \left(\frac{1}{2} A dA + \frac{1}{3} A^3 \right) = \int_{M^3} \text{Tr} \left(\frac{1}{2} A_i \partial_j A_k + \frac{1}{3} A_i A_j A_k \right) dx^{ijk}, \quad (25)$$

where $dx^{ijk} = dx^i \wedge dx^j \wedge dx^k$, the volume form. In his canonical quantisation of this theory, Witten [36] considered the threefold to be of the form $M^3 = \Sigma \times \mathbb{R}^1$, where the data on the two-dimensional boundary Σ , a Riemann surface, satisfied the equations (21).

Now applying an x^4 -dependent gauge transformation to the vector potentials

$$A_a(x^i, 0) \mapsto g^{-1}(x^i, x^4)A_a(x^i, 0)g(x^i, x^4) + g^{-1}(x^i, x^4)\partial_a g(x^i, x^4), \quad a = 1, \dots, 4, \quad (26)$$

where $A_4(x^i, 0) = 0$, yields a pure-gauge form for the fourth vector potential, $A_4 = g^{-1}\partial_4 g$. The non-gauge covariant equation (24) now takes the gauge covariant form of the four dimensional SO(4)-invariant anti-self-duality equations

$$F_{ab} + \frac{1}{2}\epsilon_{abcd}F^{cd} = 0, \quad a, b, c, d = 1, \dots, 4, \quad (27)$$

a set of three equations for the four vector potentials. (The self-duality equations emerge on reversing the x^4 -direction of the flow.)

Using a manifestly u(2)-covariant notation for Yang's complex coordinates ($z^\alpha, z^{\bar{\alpha}} := \bar{z}^\alpha, \alpha, \bar{\alpha} = 1, 2$), these equations take the form [c.f. (21)] [37],

$$\Omega_{\alpha\beta}F^{\alpha\beta} = 0 \quad \Leftrightarrow \quad F_{\bar{1}\bar{2}} = 0 \quad (28)$$

$$g^{\alpha\bar{\beta}}F_{\alpha\bar{\beta}} = 0 \quad \Leftrightarrow \quad F_{1\bar{1}} + F_{2\bar{2}} = 0. \quad (29)$$

This is a system consisting of one complex and one real equation, leaving as the sole non-zero part, the trace-free part of the (1,1)-curvature. The U(2)-invariant metric on $\mathbb{C}^2 \simeq \mathbb{R}^4$ is given by $g_{\alpha\bar{\beta}}dz^\alpha dz^{\bar{\beta}} = dz^1 dz^{\bar{1}} + dz^2 dz^{\bar{2}} =: dz^\alpha dz^{\bar{\alpha}}$ and the symplectic (2,0) volume form, invariant under SU(2), by $\Omega = \Omega_{\alpha\beta}dz^\alpha \wedge dz^\beta = dz^1 \wedge dz^2 =: dz^{12}$.

Now, complexifying all the data by dropping all reality conditions (see for instance the discussion in [12]), we obtain the additional equation $F_{\alpha\beta} = 0$, which allows us to choose the holomorphic gauge $A_\alpha = 0$. The Eq. (29) then takes the form of a conservation law [38]

$$g^{\alpha\bar{\beta}}\partial_\alpha A_{\bar{\beta}} = \partial^{\bar{\beta}} A_{\bar{\beta}} = 0, \quad \alpha, \bar{\beta} = 1, 2, \quad (30)$$

which has local solution $A_{\bar{\beta}} = \Omega_{\bar{\alpha}\bar{\beta}}\partial^{\bar{\alpha}} f$, where $\bar{\Omega} = \Omega_{\bar{\alpha}\bar{\beta}}dz^{\bar{\alpha}} \wedge dz^{\bar{\beta}} = dz^{\bar{1}\bar{2}}$ is the symplectic (0,2)-form. The remaining equation in (28) then takes the form of Leznov's wave equation [39]

$$\square f + \frac{1}{2}\Omega^{\alpha\beta}[\partial_\alpha f, \partial_\beta f] = 0, \quad (31)$$

with Laplacian $\square = g^{\alpha\bar{\beta}}\partial_\alpha \partial_{\bar{\beta}} = \partial^{\bar{\alpha}}\partial_{\bar{\alpha}}$. Solutions provide stationary points of the Leznov functional

$$S_L = \int_{M^c} \text{Tr} \left(\frac{1}{2}f\square f + \frac{1}{3}\Omega^{\alpha\beta}f\partial_\alpha f\partial_\beta f \right). \quad (32)$$

whose gradient flow has the standard heat equation form,

$$\frac{\partial}{\partial t} f = \square f + \frac{1}{2}\Omega^{\alpha\beta}[\partial_\alpha f, \partial_\beta f]. \quad (33)$$

Here t is the parameter of the infinitesimal variation, so stationary points correspond to functions $f(t, z, \bar{z})$ independent of t , i.e. solutions of Eq.(31). In the case of Eq.(33), the left-hand-side does not allow interpretation as a (gauge-fixed) component of the curvature.

In all the above cases, in dimensions $d = 1, \dots, 4$, the equations are fully $\text{SO}(d)$ -invariant. The special geometric structures characterising these equations are thus precisely the volume forms, which are trivially special democratic forms. The oxidised volume form in d -dimensions $\text{vol}_d = dx^{1\dots d}$ may be obtained from lower dimensional volume forms by taking successive wedge products with the additional basis one-forms, $\text{vol}_d = \text{vol}_{d-1} \wedge dx^d$.

4. From Four to Eight Dimensions

4.1. Permutation to $d = 6$. To proceed to higher dimensions, we now consider the complex version (28), (29) of the four-dimensional equations. Following the previous mapping from two to three dimensions (22), we now oxidise these equations from \mathbb{C}^2 to \mathbb{C}^3 by requiring invariance under the cyclic permutations generated by $\sigma = (1\ 2\ 3) \in S_3$, where the indices are now complex;

$$\{F_{\bar{1}\bar{2}} = 0\} \longrightarrow \{F_{\sigma^p(\bar{i})\sigma^p(\bar{j})} = 0; \sigma = (1\ 2\ 3), p = 1, 2\}. \quad (34)$$

This yields the system [c.f. (23)]

$$\Omega_{\alpha\beta\gamma} F^{\beta\gamma} = 0 \Leftrightarrow \{F_{\bar{1}\bar{2}} = F_{\bar{2}\bar{3}} = F_{\bar{3}\bar{1}} = 0\} \quad (35)$$

$$g^{\alpha\bar{\beta}} F_{\alpha\bar{\beta}} = 0 \Leftrightarrow F_{1\bar{1}} + F_{2\bar{2}} + F_{3\bar{3}} = 0, \quad \alpha, \bar{\alpha} = 1, 2, 3, \quad (36)$$

a set of three complex and one real equation. Here $g_{\alpha\bar{\beta}} dz^\alpha d\bar{z}^\beta$ is the $\text{U}(3)$ -invariant hermitian metric and $\Omega = dz^1 \wedge dz^2 \wedge dz^3 = dz^{123}$, the complex (3,0) volume form. These equations were obtained in [5] as $\text{SU}(3) \otimes \text{U}(1) / \mathbb{Z}_2$ -invariant curvature constraints which imply the second order Yang–Mills equations. They later made an appearance in work by Donaldson [40], Uhlenbeck and Yau [15] as the equations for holomorphic connections on three (complex) dimensional Kähler manifolds, g being the Kähler metric.

In the six real coordinates, $x^\alpha := \text{Re } z^\alpha$, $x^{\alpha+3} = \text{Im } z^\alpha$, $\alpha = 1, 2, 3$, the equations take the form (6), with the special democratic four-form (see [5])

$$T_{(6)} = dx^{1425} + dx^{1436} + dx^{2536}. \quad (37)$$

This is invariant under the group S_3 of permutations of the 3 ordered pairs ($\{1, 4\}$, $\{2, 3\}$, $\{4, 5\}$), or, equivalently, the symmetries generated by the permutation $\sigma = (123)(456) \in S_6$. The stabiliser of $T_{(6)}$ in $\text{SO}(6)$ is the group $\text{SU}(3) \times \text{U}(1) / \mathbb{Z}_2$ and under this, the space of two-forms has the following decomposition into eigenspaces of $T_{(6)}$ [5]:

$$\Lambda^2 \mathbb{R}^6 = (\mathfrak{su}(3)_0, \lambda = -1) \oplus (V_2^3 \oplus \bar{V}_{-2}^3, \lambda = 1) \oplus (\mathbb{R}\omega_0, \lambda = 2), \quad (38)$$

where (V_q^n, λ) is the n -dimensional irreducible representation of $\text{SU}(3)$, the index q denotes the $\text{U}(1)$ charge, λ the eigenvalue of $T_{(6)}$ and $\omega_0 = g_{\alpha\bar{\beta}} dz^\alpha \wedge d\bar{z}^\beta$ is the invariant metric form associated with g . Two-forms parallel to ω_0 are contained in the $\lambda = 2$

eigenspace. Under the action of $T_{(6)}$ the curvature tensor therefore decomposes into $T_{(6)}$ -eigenspaces according to

$$F = (F_{\gamma\bar{\delta}} - \frac{1}{3}g_{\gamma\bar{\delta}}F_0, \lambda = -1) \oplus (F_{\alpha\beta} \oplus F_{\bar{\alpha}\bar{\beta}}, \lambda = 1) \oplus (F_0, \lambda = 2), \quad (39)$$

where F_0 denotes the trace $g^{\alpha\bar{\beta}}F_{\alpha\bar{\beta}}$. The set of seven equations (35), (36) thus projects the curvature to the eight-dimensional $\mathfrak{su}(3)$ part, the $\lambda = -1$ eigenspace. We note that solutions of the set of nine equations projecting to the six-dimensional $\lambda = 1$ eigenspace are saddle points of the Yang–Mills action in accordance with Eq. (13).

Analogously to (29), complexifying the Yang–Mills fields, Eq. (36), in the holomorphic gauge $A_\alpha = 0$, $\alpha = 1, 2, 3$, can be locally solved in terms of three prepotentials taking values in the complexification of the gauge group:

$$A_{\bar{\alpha}} = \Omega_{\bar{\alpha}\bar{\beta}\bar{\gamma}}\partial^{\bar{\gamma}}f^{\bar{\beta}}. \quad (40)$$

The remaining conditions (35) provide extrema of the Chern–Simons action

$$\begin{aligned} S &= \int_{M_{\mathbb{C}}} \text{Tr} (\bar{A}\partial\bar{A} + \bar{A}^3) \wedge *\bar{\Omega} \\ &= \int_{M_{\mathbb{C}}} \text{Tr} (\frac{1}{2}A_{\bar{\alpha}}\partial_{\bar{\beta}}A_{\bar{\gamma}} + \frac{1}{3}A_{\bar{\alpha}}A_{\bar{\beta}}A_{\bar{\gamma}}) dz^{\bar{\alpha}\bar{\beta}\bar{\gamma}}. \end{aligned} \quad (41)$$

Inserting (40) in (35) yields a wave equation analogous to (31) for the triplet of complex prepotentials f_β ,

$$\partial^{\beta}\partial_{[\alpha}f_{\beta]} + \frac{1}{2}\Omega^{\beta\delta\eta}[\partial_{\delta}f_{\eta}, \partial_{[\alpha}f_{\beta]}] = 0. \quad (42)$$

The associated gradient flow for the functional (41) takes the heat equation form,

$$\frac{\partial}{\partial t}f_{\alpha} = \partial^{\beta}\partial_{[\alpha}f_{\beta]} + \frac{1}{2}\Omega^{\beta\delta\eta}[\partial_{\delta}f_{\eta}, \partial_{[\alpha}f_{\beta]}]. \quad (43)$$

The reduction of (35), (36) to the missing $d = 5$ case involves choosing a constant unit vector in \mathbb{R}^6 and projecting to the five-dimensional space orthogonal to it. Without loss of generality, we may simply choose one of the basis vectors, say e_6 , effectively deleting the variables x^6 and yielding an $\text{SO}(4)$ -invariant four-form $T = dx^{1245}$. The corresponding equations (see [5]) are an embedding of four-dimensional self-duality (27) in five dimensional space. A five dimensional reduction of the Chern–Simons action (41) and corresponding flow equations were discussed some time ago by Nair and Schiff [41].

4.2. Flow to $d = 7$ and $d = 8$. Since the three complex equations (35) have an action (41), we may write down the partial curvature flow, for the three complex potentials A_α , now depending on seven variables ($z^\alpha, z^{\bar{\alpha}}, x^7$):

$$\frac{\partial}{\partial x^7}A_\alpha = \Omega_{\alpha\beta\gamma}F^{\beta\gamma}, \quad \alpha = 1, 2, 3. \quad (44)$$

This being the gradient flow for the functional (41). Now, analogously to the four-dimensional case (26), an x^7 -dependent gauge transformation yields the fully gauge covariant form of this partial curvature flow

$$F_{7\alpha} = \Omega_{\alpha\beta\gamma}F^{\beta\gamma} \Leftrightarrow \{F_{71} = F_{2\bar{3}}, F_{72} = F_{3\bar{1}}, F_{73} = F_{1\bar{2}}\}. \quad (45)$$

Here $\partial/\partial x^7$ denotes the real vector field (the ‘time’ of the flow) and $\alpha, \beta, \gamma = 1, 2, 3$ are complex indices. The three complex equations (45) together with the real equation,

$$g^{\alpha\bar{\beta}} F_{\alpha\bar{\beta}} = F_{1\bar{1}} + F_{2\bar{2}} + F_{3\bar{3}} = 0, \quad (46)$$

imply the Yang–Mills equations in seven dimensions. Choosing real coordinates (x^1, \dots, x^7) , these equations they take the manifestly G_2 -invariant form [5]

$$\psi_{ijk} F^{jk} = 0, \quad i, j, k = 1, \dots, 7. \quad (47)$$

Here ψ is the G_2 -invariant Cayley three form whose components ψ_{ijk} provide structure constants of the algebra of imaginary octonions. Choosing the first six real coordinates as the real and imaginary parts of the complex coordinates as follows, $z^\alpha = x^\alpha + i x^{\alpha+3}$, $\alpha = 1, 2, 3$, we obtain,

$$\psi = dx^{367} + dx^{257} + dx^{147} + dx^{465} + dx^{243} + dx^{135} + dx^{162}. \quad (48)$$

Its four-form dual is given by

$$\varphi := *\psi = dx^{1245} + dx^{1346} + dx^{2356} + dx^{7123} + dx^{1567} + dx^{7246} + dx^{3457}, \quad (49)$$

in terms of which the equations (47) take the form,

$$F_{ij} + \frac{1}{2} \varphi_{ijkl} F^{kl} = 0, \quad i, j, k = 1, \dots, 7, \quad (50)$$

which projects the curvature to the $\lambda = -1$ eigenspace of φ ; the eigenspace decomposition of the space of two-forms being [5]

$$\Lambda^2 \mathbb{R}^7 = (\mathfrak{g}_2, \lambda = -1) \oplus (\mathbb{R}^7, \lambda = 3). \quad (51)$$

Since the system (47) consists of seven equations for seven potentials and has the Chern–Simons type action

$$S_{CS} = \int_{M^7} \text{Tr} (AdA + A^3) \wedge *\psi = \int_{M^7} \text{Tr} \left(\frac{1}{2} A_i \partial_j A_k + \frac{1}{3} A_i A_j A_k \right) \psi^{ijk}, \quad (52)$$

we can immediately write down the corresponding partial curvature flow in eight dimensions analogous to (24):

$$\frac{\partial}{\partial x^8} A_i = \frac{1}{2} \psi_{ijk} F^{jk}, \quad i = 1, \dots, 7. \quad (53)$$

This is the temporal gauge ($A_8 = 0$) form of the Spin(7)-invariant equations in eight dimensions, which were discovered in [5] and shown there to arise as the projection of the curvature form to the $\lambda = -1$ eigenspace of the Spin(7)-invariant four-form ϕ ,

$$F_{ab} + \frac{1}{2} \phi_{abcd} F^{cd} = 0, \quad a, b, c, d = 1, \dots, 8, \quad (54)$$

where in terms of the seven dimensional forms ψ, φ in (47) and (50) the four-form ϕ in eight dimensions is given by $\phi = dx^8 \wedge \psi + \varphi$. The decomposition $\Lambda^2 \mathbb{R}^8$ into eigenspaces of this four-form is given by

$$\Lambda^2 \mathbb{R}^8 = (\mathfrak{spin}_7, \lambda = -1) \oplus (\mathbb{R}^7, \lambda = 3). \quad (55)$$

In complex coordinates, $z^\alpha = x^\alpha + ix^{\alpha+4}$, $\alpha = 1, 2, 3, 4$, the Eq. (54) take the form (see [5]) incorporating (45),

$$\begin{aligned} F_{\alpha\beta} + \frac{1}{2}\Omega_{\alpha\beta\gamma\delta} F^{\gamma\delta} = 0 &\Leftrightarrow \{F_{41} = F_{\bar{2}\bar{3}}, F_{42} = F_{\bar{3}\bar{1}}, F_{43} = F_{\bar{1}\bar{2}}\} \\ g^{\alpha\bar{\beta}} F_{\alpha\bar{\beta}} = 0 &\Leftrightarrow F_{1\bar{1}} + F_{2\bar{2}} + F_{3\bar{3}} + F_{4\bar{4}} = 0, \end{aligned} \quad (56)$$

where g is the $U(4)$ -invariant hermitian metric on $\mathbb{C}^4 \simeq \mathbb{R}^8$ and $\Omega = dz^{1234}$ is the $SU(4)$ -invariant volume form in \mathbb{C}^4 . In the complex ‘temporal’ gauge, $A_4 = 0$, the three complex equations in (56) therefore take the form of a partial curvature flow with complex flow parameter z^4 ,

$$\frac{\partial}{\partial z^4} A_\alpha = \frac{1}{2}\Omega_{\alpha\beta\gamma} F^{\beta\gamma} \quad (57)$$

$$\frac{\partial}{\partial z^4} A_{\bar{4}} = g_3^{\alpha\bar{\beta}} F_{\alpha\bar{\beta}} \quad (58)$$

where $\Omega_{\alpha\beta\gamma}$, $g_3^{\alpha\bar{\beta}}$ are the volume form and inverse metric of the complex 3-space orthogonal to the complex vector field $\partial/\partial z^4$. The Eq. (57) thus gives the complex variation of the Chern–Simons action (41).

All the above duality equations in dimensions up to eight are more or less well-known [5]. Our main result is that the pattern of successive dimensional oxidation actually continues to higher dimensions. Proceeding further, we see that a particularly interesting 12-dimensional system results.

5. Self-Duality in 12 Dimensions

Following the method of oxidising the duality equations from \mathbb{R}^4 to \mathbb{R}^6 , we now extend the system (56) in \mathbb{C}^4 to \mathbb{C}^6 by juxtaposing two additional complex variables z^5, z^6 and then remixing the six complex indices by requiring symmetry under permutations generated by $\sigma = (135)(246) \in S_6$. We thus obtain the equations,

$$g^{\alpha\bar{\beta}} F_{\alpha\bar{\beta}} = F_{1\bar{1}} + F_{2\bar{2}} + F_{3\bar{3}} + F_{4\bar{4}} + F_{5\bar{5}} + F_{6\bar{6}} = 0, \quad (59)$$

together with

$$F_{12} + F_{\bar{3}\bar{4}} + F_{\bar{5}\bar{6}} = 0, \quad F_{34} + F_{\bar{5}\bar{6}} + F_{\bar{1}\bar{2}} = 0, \quad F_{56} + F_{\bar{1}\bar{2}} + F_{\bar{3}\bar{4}} = 0 \quad (60)$$

$$F_{13} + F_{\bar{4}\bar{2}} = 0, \quad F_{14} + F_{\bar{2}\bar{3}} = 0, \quad F_{15} + F_{\bar{6}\bar{2}} = 0$$

$$F_{16} + F_{\bar{2}\bar{5}} = 0, \quad F_{35} + F_{\bar{6}\bar{4}} = 0, \quad F_{36} + F_{\bar{4}\bar{5}} = 0. \quad (61)$$

These equations imply the 12-dimensional Yang–Mills equations! The proof follows from Theorem 2 and the observation that these equations allow expression in the form (15), (16), with the (4,0)-form Φ taking the form

$$\Phi = dz^{1234} + dz^{1256} + dz^{3456}. \quad (62)$$

This four-form is thus given by $\Phi = \omega^2$, where ω is the symplectic (2,0)-form

$$\omega = dz^{12} + dz^{34} + dz^{56} \in \Lambda^2 \mathbb{C}^6. \quad (63)$$

This is analogous to the \mathbb{R}^6 case, except that now everything is complex. The (4,0)-form Φ is manifestly invariant under the action of $\text{Sp}(3) \subset \text{SU}(6) \subset \text{Spin}(12)$.

The three conditions in (60) are equivalent to the four real equations,

$$\begin{aligned} \text{Im}(F_{\bar{1}\bar{2}}) &= \text{Im}(F_{\bar{3}\bar{4}}) = \text{Im}(F_{\bar{5}\bar{6}}) = 0, \\ \text{Re}(\omega^{\bar{\alpha}\bar{\beta}} F_{\bar{\alpha}\bar{\beta}}) &= \text{Re}(F_{\bar{1}\bar{2}} + F_{\bar{3}\bar{4}} + F_{\bar{5}\bar{6}}) = 0, \end{aligned} \quad (64)$$

where the symplectic (0,2)-form $\bar{\omega} = \omega_{\bar{\alpha}\bar{\beta}} dz^{\bar{\alpha}} \wedge dz^{\bar{\beta}} = dz^{\bar{1}\bar{2}} + dz^{\bar{3}\bar{4}} + dz^{\bar{5}\bar{6}}$ and $\omega^{\bar{\alpha}\bar{\beta}} \omega_{\bar{\beta}\bar{\gamma}} = \delta_{\bar{\gamma}}^{\bar{\alpha}}$. The system of equations thus consists of five real equations, (59) and (64), together with six complex equations (61), a total of 17 real equations.

The entire system (59),(60),(61) in real coordinates for 12-dimensional euclidean space given by $x^i = \text{Re } z^i$, $x^{i+6} = \text{Im } z^i$, $i = 1, \dots, 6$, takes the following form. Here we denote the indices 10,11,12 by 0, a , b respectively.

$$F_{12} + F_{34} + F_{56} + F_{87} + F_{09} + F_{ba} = 0 \quad (65)$$

$$F_{17} + F_{28} + F_{39} + F_{40} + F_{5a} + F_{6b} = 0 \quad (66)$$

$$F_{13} + F_{42} + F_{97} + F_{80} = 0$$

$$F_{14} + F_{23} + F_{07} + F_{98} = 0$$

$$F_{15} + F_{62} + F_{a7} + F_{8b} = 0$$

$$F_{16} + F_{25} + F_{b7} + F_{a8} = 0$$

$$F_{35} + F_{64} + F_{a9} + F_{0b} = 0$$

$$F_{36} + F_{45} + F_{b9} + F_{a0} = 0$$

$$F_{19} + F_{73} + F_{84} + F_{20} = 0$$

$$F_{10} + F_{74} + F_{92} + F_{38} = 0$$

$$F_{1a} + F_{75} + F_{86} + F_{2b} = 0$$

$$F_{1b} + F_{76} + F_{a2} + F_{58} = 0$$

$$F_{3a} + F_{95} + F_{06} + F_{4b} = 0$$

$$F_{3b} + F_{96} + F_{a4} + F_{50} = 0 \quad (67)$$

$$F_{18} + F_{72} = 0$$

$$F_{30} + F_{94} = 0$$

$$F_{5b} + F_{a6} = 0 \quad (68)$$

These equations have the familiar form (6), with the four-form $T_{(12)} \in \Lambda^4 \mathbb{R}^{12}$ being given by the special democratic form

$$\begin{aligned} T_{(12)} = & dx^{1234} + dx^{1256} + dx^{1287} + dx^{1209} + dx^{12ba} + dx^{1397} + dx^{1380} \\ & + dx^{1407} + dx^{1498} + dx^{15a7} + dx^{158b} + dx^{16b7} + dx^{16a8} + dx^{2307} \\ & + dx^{2398} + dx^{2479} + dx^{2408} + dx^{25b7} + dx^{25a8} + dx^{267a} + dx^{26b8} \\ & + dx^{3456} + dx^{3487} + dx^{3409} + dx^{34ba} + dx^{35a9} + dx^{35b0} + dx^{36b9} \\ & + dx^{36a0} + dx^{45b9} + dx^{45a0} + dx^{469a} + dx^{46b0} + dx^{5687} + dx^{5609} \\ & + dx^{56ba} + dx^{7890} + dx^{78ab} + dx^{90ab}, \end{aligned} \quad (69)$$

which has a set of 39 non-zero components. The characteristic polynomial of this $\text{Sp}(3)$ -invariant four-form, acting on the space of two-forms has been calculated using Maple.

The eigenspace decomposition of the space of two-forms in terms of $\mathrm{Sp}(3)$ representations (see e.g. [42,43]) is given by

$$\begin{aligned} \Lambda^2 \mathbb{R}^{12} = & (\mathfrak{sp}_3 \oplus V^{14}(\pi_2) \oplus V^{14}(\pi_2), \lambda = -1) \oplus (V^{14}(\pi_2), \lambda = 3) \\ & \oplus (\mathbb{C}\omega, \lambda = 5) \oplus (\mathbb{R}\omega_0, \lambda = -3). \end{aligned} \quad (70)$$

Here, ω is the symplectic form (63) and ω_0 the metric form $\omega_0 = g_{\alpha\bar{\beta}} dz^\alpha \wedge dz^{\bar{\beta}}$. $V^{14}(\pi_2)$ denotes the 14-dimensional representation with highest weight π_2 , the second fundamental weight of \mathfrak{sp}_3 . The four-form $T_{(12)}$ is in fact one of six $\mathrm{Sp}(3)$ -invariant four-forms in 12 dimensions. The 17 equations (65)–(67) project the curvature two-form to the 49-dimensional eigenspace with eigenvalue $\lambda = -1$. The other eigenspaces have rather small dimensions compared with $\dim(\Lambda^2 \mathbb{R}^{12}) = 66$. We therefore expect the corresponding solutions to be rather trivial. $\mathrm{Sp}(3)$, the stabiliser of the four-form $T_{(12)}$ is a maximal subgroup of $\mathrm{SU}(6)$.

The similarity of the equations (59), (60), and (61) to the three and six dimensional systems in \mathbb{R}^3 and $\mathbb{C}^3 \simeq \mathbb{R}^6$ discussed above suggests that this is the counterpart in three dimensional quaternionic space $\mathbb{H}^3 \simeq \mathbb{C}^6$. The imaginary quaternion units satisfy $i^2 = j^2 = k^2 = -1$ and $ij = -ji = k$, together with the relations which result on cyclically permuting (i, j, k) . We consider \mathbb{C} to be an \mathbb{R} -vector space spanned by $(1, i)$ and \mathbb{H} a \mathbb{C} -vector space spanned by $(1, j)$. Scalar multiplication of $z \in \mathbb{C}$ with the quaternionic basis element j satisfies $zj = j\bar{z}$, so quaternions may be written in the form

$$q := z + j\bar{w} = z + wj, \quad q \in \mathbb{H}, \quad z, w \in \mathbb{C}. \quad (71)$$

The conjugate quaternion is then given by

$$\bar{q} := \bar{z} - wj = \bar{z} - j\bar{w}, \quad q \in \mathbb{H}, \quad z, w \in \mathbb{C}. \quad (72)$$

The conjugate imaginary units are clearly given by $\bar{i} = -i$, $\bar{j} = -j$, $\bar{k} = -k$. Quaternions being noncommutative, conjugation is an involutive antiautomorphism, i.e. $\bar{\bar{q}} = q$ and $\overline{q_1 q_2} = \bar{q}_2 \bar{q}_1$. There exist related involutive automorphisms given by conjugation with the quaternion units,

$$\begin{aligned} \mathrm{id} : q &\mapsto q = z + wj, \\ \alpha : q &\mapsto -iqi = z - wj, \\ \beta : q &\mapsto -jqj = \bar{z} + \bar{w}j, \\ \gamma : q &\mapsto -kqk = \bar{z} - \bar{w}j, \end{aligned} \quad (73)$$

in terms of which the real and imaginary parts of q can be expressed as linear combinations of $q, \alpha(q), \beta(q), \gamma(q)$ (see e.g. [44]).

Now, let M be a three quaternionic-dimensional (i.e. 12 real-dimensional) space. In a local coordinate frame $T_p M \simeq \mathbb{H}^3 \simeq \mathbb{C}^6$. We define three quaternionic coordinates q^A , $A = 1, 2, 3$, in terms of pairs of the complex coordinates $z^\alpha := x^\alpha + ix^{\alpha+6}$, $\alpha = 1, \dots, 6$ used above,

$$\begin{aligned} q^1 &:= z^1 + z^2 j = x^1 + ix^7 + jx^2 + kx^8, \\ q^2 &:= z^3 + z^4 j = x^3 + ix^9 + jx^4 + kx^0, \\ q^3 &:= z^5 + z^6 j = x^5 + ix^a + jx^6 + kx^b \end{aligned} \quad (74)$$

and we denote the conjugate coordinates as $\overline{q^A} = q^{\bar{A}}$.

For any two quaternionic vector fields Q_1, Q_2 the curvature components $F(Q_1, \gamma(Q_2))$ and $F(Q_2, \gamma(Q_1))$ have the same content in terms of real curvature components, since γ is an involutive automorphism. We now denote the basis vectors of the coordinate vector fields on M by $Q_A := \partial/\partial q^A$, their quaternionic conjugates by $Q_{\bar{A}} := \overline{Q_A} = \partial/\partial q^{\bar{A}}$ and their α, β, γ -conjugates by $Q_{\alpha(A)} := \alpha(Q_A)$, etc. The hermitian metric in local quaternionic coordinates is given by $d^2s = g_{A\bar{B}}dq^A dq^{\bar{B}} = dq^1 dq^{\bar{1}} + dq^2 dq^{\bar{2}} + dq^3 dq^{\bar{3}}$.

Proposition 1. *On a three quaternionic dimensional Riemannian manifold, the following eight quaternionic curvature constraints are equivalent to the system (65), (66), (67) and (68) of self-duality equations in 12 dimensions:*

$$g^{A\bar{B}} F(Q_{\bar{B}}, Q_{\alpha(A)}) = \sum_{A=1}^3 F(Q_{\bar{A}}, Q_{\alpha(A)}) = 0 \quad (75)$$

$$g^{A\bar{B}} F(Q_{\bar{B}}, Q_{\beta(A)}) = \sum_{A=1}^3 F(Q_{\bar{A}}, Q_{\beta(A)}) = 0 \quad (76)$$

$$F(Q_{\bar{1}}, Q_{\gamma(2)}) = F(Q_{\bar{2}}, Q_{\gamma(3)}) = F(Q_{\bar{3}}, Q_{\gamma(1)}) = 0 \quad (77)$$

$$F(Q_{\bar{1}}, Q_{\gamma(1)}) = F(Q_{\bar{2}}, Q_{\gamma(2)}) = F(Q_{\bar{3}}, Q_{\gamma(3)}) = 0. \quad (78)$$

Proof. The equivalence to the 17 equations (65)–(68), or equivalently to the complex form (59)–(61) follows from a direct expansion of the quaternionic vector fields in the basis (1, j). \square

6. Flowing to 14 Dimensions

The similarity between the three quaternionic equations in (77), the three complex equations in (35) and the 3 real equations in (23) immediately suggests that in analogy to the flows (24) and (44), we may write down flows for the three quaternionic partial curvatures in (77) into a further complex direction, with coordinate z^7 . We write, in $M = M_{\mathbb{H}}^3 \times \mathbb{C}$ with coordinates (q^1, q^2, q^3, z^7) , in analogy with (57) and (58),

$$\begin{aligned} \frac{\partial}{\partial z^7} A(Q_1) &= -F(Q_{\bar{2}}, Q_{\gamma(3)}) \\ \frac{\partial}{\partial z^7} A(Q_2) &= -F(Q_{\bar{3}}, Q_{\gamma(1)}) \\ \frac{\partial}{\partial z^7} A(Q_3) &= -F(Q_{\bar{1}}, Q_{\gamma(2)}) \\ \frac{\partial}{\partial z^7} A(Z_{\bar{7}}) &= \sum_{A=1}^3 F(Q_{\bar{A}}, Q_{\alpha(A)}) \end{aligned} \quad (79)$$

together with (76), considered as an equation in 14 dimensions,

$$\sum_{A=1}^3 F(Q_{\bar{A}}, Q_{\beta(A)}) = 0. \quad (80)$$

Writing the quaternionic vector fields Q_A , $A = 1, \dots, 3$ in terms of complex vector fields Z_α , $\alpha = 1, \dots, 6$ according to the choice in (74) and unravelling the $A(Z_7) = 0$ gauge, we obtain the system

$$\begin{aligned}
 F(Z_7, Z_1 + Z_2j) + F(\overline{Z_3} - j\overline{Z_4}, \overline{Z_5} - \overline{Z_6}j) &= 0 \\
 F(Z_7, Z_3 + Z_4j) + F(\overline{Z_5} - j\overline{Z_6}, \overline{Z_1} - \overline{Z_2}j) &= 0 \\
 F(Z_7, Z_5 + Z_6j) + F(\overline{Z_1} - j\overline{Z_2}, \overline{Z_3} - \overline{Z_4}j) &= 0 \\
 F(Z_7, \overline{Z_7}) - \sum_{\alpha=1}^3 F(\overline{Z_{2\alpha-1}} - j\overline{Z_{2\alpha}}, Z_{2\alpha-1} - j\overline{Z_{2\alpha}}) &= 0 \\
 \sum_{\alpha=1}^3 F(\overline{Z_{2\alpha-1}} - j\overline{Z_{2\alpha}}, \overline{Z_{2\alpha-1}} + j\overline{Z_{2\alpha}}) &= 0
 \end{aligned} \tag{81}$$

Expanding the quaternionic vector fields in the basis $(1, j)$, we obtain equations on \mathbb{C}^7 , which are contained in the system

$$F_{\alpha\beta} + \frac{1}{2}\Phi_{\alpha\beta\gamma\delta} F_{\overline{\gamma\delta}} = 0 \tag{82}$$

$$g^{\alpha\overline{\beta}} F_{\alpha\overline{\beta}} = 0 \tag{83}$$

with Φ given by the G_2^C -invariant (4,0)-form

$$\Phi = dz^{1234} + dz^{1256} + dz^{3456} + dz^{1375} + dz^{1467} + dz^{2367} + dz^{2457}. \tag{84}$$

By Theorem 2 we therefore have a system of equations which implies the Yang–Mills equations in 14 dimensions.

Unlike the previous analogous cases, the equations (82) are not equivalent to the set (81). The former set contains more equations than the latter. More precisely, (82) includes, for instance, the three equations

$$F_{71} + F_{\overline{35}} + F_{\overline{64}} = F_{71} + F_{\overline{35}} + F_{64} = F_{71} + F_{35} + F_{\overline{64}} = 0. \tag{85}$$

Under the G_2^C -invariant four-form Φ , both real and imaginary parts of $F_{\alpha\beta}$ split into their 7- and 14-dimensional irreducible parts. The equations of the form (85) imply that under (82) the real part is projected to the 14-dimensional piece (seven equations) and the imaginary part is zero (21 equations). The real form of the system (82), (83) is given by the set of 29 equations,

$$\begin{aligned}
 F_{18} + F_{29} + F_{30} + F_{4a} + F_{5b} + F_{6c} + F_{7d} &= 0 \\
 F_{12} + F_{34} + F_{56} - F_{89} - F_{0a} - F_{bc} &= 0 \\
 F_{13} - F_{24} - F_{80} + F_{9a} + F_{bd} + F_{75} &= 0 \\
 F_{14} + F_{23} - F_{8a} - F_{90} - F_{cd} - F_{76} &= 0 \\
 F_{15} - F_{26} - F_{8b} + F_{9c} - F_{0d} - F_{73} &= 0 \\
 F_{16} + F_{25} - F_{8c} - F_{9b} + F_{ad} + F_{74} &= 0 \\
 F_{17} - F_{35} + F_{46} - F_{8d} + F_{0b} - F_{ac} &= 0 \\
 F_{72} - F_{36} - F_{45} + F_{9d} + F_{0c} + F_{ab} &= 0 \\
 F_{78} - F_{1d} = F_{79} - F_{2d} = F_{70} - F_{3d} &= 0
 \end{aligned} \tag{86}$$

$$\begin{aligned}
F_{7a} - F_{4d} &= F_{7b} - F_{5d} = F_{7c} - F_{6d} = 0 \\
F_{19} - F_{28} &= F_{10} - F_{38} = F_{1a} - F_{48} = 0 \\
F_{1b} - F_{58} &= F_{20} - F_{39} = F_{2a} - F_{49} = 0 \\
F_{3a} - F_{40} &= F_{2b} - F_{59} = F_{3b} - F_{50} = 0 \\
F_{4b} - F_{5a} &= F_{1c} - F_{68} = F_{2c} - F_{69} = 0 \\
F_{3c} - F_{60} &= F_{4c} - F_{6a} = F_{5c} - F_{6b} = 0.
\end{aligned} \tag{87}$$

These 29 equations correspond to the $\lambda = -1$ eigenspace of the special democratic four-form given by

$$\begin{aligned}
T_{(14)} = & dx^{1234} + dx^{1256} + dx^{1298} + dx^{12a0} + dx^{12cb} + dx^{1375} + dx^{13bd} \\
& + dx^{1308} + dx^{139a} + dx^{14a8} + dx^{1409} + dx^{1467} + dx^{14dc} + dx^{15b8} \\
& + dx^{159c} + dx^{15d0} + dx^{16c8} + dx^{16b9} + dx^{1ad6} + dx^{170b} + dx^{1a7c} \\
& + dx^{2367} + dx^{23a8} + dx^{2309} + dx^{23dc} + dx^{2457} + dx^{2480} + dx^{24a9} \\
& + dx^{24db} + dx^{25c8} + dx^{25b9} + dx^{2ad5} + dx^{268b} + dx^{26c9} + dx^{20d6} \\
& + dx^{207c} + dx^{2a7b} + dx^{3456} + dx^{3498} + dx^{34a0} + dx^{34cb} + dx^{35b0} \\
& + dx^{35ac} + dx^{835d} + dx^{93d6} + dx^{36c0} + dx^{36ba} + dx^{83b7} + dx^{937c} \\
& + dx^{94d5} + dx^{45c0} + dx^{45ba} + dx^{460b} + dx^{46ca} + dx^{84d6} + dx^{847c} \\
& + dx^{947b} + dx^{5698} + dx^{56a0} + dx^{56cb} + dx^{9a75} + dx^{8a76} + dx^{9076} \\
& + dx^{890a} + dx^{89bc} + dx^{8057} + dx^{80db} + dx^{8acd} + dx^{90cd} + dx^{9abd} \\
& + dx^{0abc} + dx^{187d} + dx^{297d} + dx^{307d} + dx^{4a7d} + dx^{5b7d} + dx^{6c7d}.
\end{aligned}$$

Its characteristic polynomial is given by

$$\chi(T_{(14)}) = (\lambda + 1)^{62}(\lambda - 3)^{14}(\lambda + 3)^7(\lambda - 5)^7(\lambda - 6) \tag{88}$$

and the above 29 equations correspond to the projection to 62-dimensional $\lambda = -1$ eigenspace.

Deleting all terms containing the 14th index d from the above equations yields the 13-dimensional reduction, corresponding to a flow along a real parameter rather than the complex one chosen in (79). This is also a set of 29 equations, projecting the curvature to the 49-dimensional $\lambda = -1$ eigenspace of the corresponding reduction of the four-form $T_{(14)}$. The reduced four-form has characteristic polynomial

$$\chi(T_{(13)}) = (\lambda + 1)^{49}(\lambda - 3)^8(\lambda + 3)(\lambda - 5)^2(\lambda - 4)^6(\lambda^2 + \lambda - 4)^6. \tag{89}$$

7. Oxidation to 16 Dimensions

Analogously to the oxidations (53), (57) and (58) to eight real dimensions, we may oxidise the system (82), (83) in \mathbb{C}^7 to one in \mathbb{C}^8 by taking $g_{\alpha\bar{\beta}}$ to be the \mathbb{C}^8 -metric and the (4,0)-form Φ to be given by the $\text{Spin}(7)^{\mathbb{C}}$ -invariant,

$$\Phi = dz^{1234} + dz^{1256} + dz^{1278} + dz^{3456} + dz^{3478} + dz^{5678} + dz^{1368} \tag{90}$$

$$+ dz^{1375} + dz^{1467} + dz^{1458} + dz^{2367} + dz^{2457} + dz^{2358} + dz^{2486}. \tag{91}$$

The corresponding system includes the flow equations based on (81),

$$\begin{aligned}
 F(Z_8, Z_{1j} - Z_2) + F(Z_7, Z_1 + Z_{2j}) + F(\overline{Z_3} - j\overline{Z_4}, \overline{Z_5} - \overline{Z_6}j) &= 0 \\
 F(Z_8, Z_{3j} - Z_4) + F(Z_7, Z_3 + Z_{4j}) + F(\overline{Z_5} - j\overline{Z_6}, \overline{Z_1} - \overline{Z_2}j) &= 0 \\
 F(Z_8, Z_{5j} - Z_6) + F(Z_7, Z_5 + Z_{6j}) + F(\overline{Z_1} - j\overline{Z_2}, \overline{Z_3} - \overline{Z_4}j) &= 0 \\
 F(Z_8, \overline{Z_8}) + F(Z_7, \overline{Z_7}) - \sum_{\alpha=1}^3 F(\overline{Z_{2\alpha-1}} - j\overline{Z_{2\alpha}}, Z_{2\alpha-1} - j\overline{Z_{2\alpha}}) &= 0 \\
 \sum_{\alpha=1}^4 F(\overline{Z_{2\alpha-1}} - j\overline{Z_{2\alpha}}, \overline{Z_{2\alpha-1}} + j\overline{Z_{2\alpha}}) &= 0.
 \end{aligned} \tag{92}$$

The real form of the full system of equations with (4,0)-form Φ given in (91) is given by

$$\begin{aligned}
 F_{12} + F_{34} + F_{56} + F_{78} - F_{90} - F_{ab} - F_{cd} - F_{ef} &= 0 \\
 F_{13} - F_{24} - F_{57} + F_{68} - F_{9a} + F_{0b} + F_{ce} - F_{df} &= 0 \\
 F_{14} + F_{23} + F_{58} + F_{67} - F_{9b} - F_{0a} - F_{cf} - F_{de} &= 0 \\
 F_{15} - F_{26} + F_{37} - F_{48} - F_{9c} + F_{0d} - F_{ae} + F_{bf} &= 0 \\
 F_{16} + F_{25} - F_{38} - F_{47} - F_{9d} - F_{0c} + F_{af} + F_{be} &= 0 \\
 F_{17} - F_{28} - F_{35} + F_{46} - F_{9e} + F_{0f} + F_{ac} - F_{bd} &= 0 \\
 F_{18} + F_{27} + F_{36} + F_{45} - F_{9f} - F_{0e} - F_{ad} - F_{bc} &= 0 \\
 F_{19} + F_{20} + F_{3a} + F_{4b} + F_{5c} + F_{6d} + F_{7e} + F_{8f} &= 0 \\
 F_{79} - F_{1e} = F_{80} - F_{2f} = F_{5a} - F_{3c} = F_{6b} - F_{4d} &= 0 \\
 F_{70} - F_{2e} = F_{3d} - F_{6a} = F_{4c} - F_{5b} = F_{1f} - F_{89} &= 0 \\
 F_{69} - F_{1d} = F_{2c} - F_{50} = F_{3f} - F_{8a} = F_{7b} - F_{4e} &= 0 \\
 F_{1c} - F_{59} = F_{2d} - F_{60} = F_{3e} - F_{7a} = F_{8b} - F_{4f} &= 0 \\
 F_{1b} - F_{49} = F_{5f} - F_{8c} = F_{2a} - F_{30} = F_{6e} - F_{7d} &= 0 \\
 F_{6f} - F_{8d} = F_{5e} - F_{7c} = F_{40} - F_{2b} = F_{39} - F_{1a} &= 0 \\
 F_{7f} - F_{8e} = F_{5d} - F_{6c} = F_{10} - F_{29} = F_{3b} - F_{4a} &= 0.
 \end{aligned} \tag{93}$$

The corresponding four-form $T_{(16)} \in \Lambda^4 \mathbb{R}^{16}$ has characteristic polynomial

$$\chi(T_{(16)}) = (\lambda + 1)^{84}(\lambda - 3)^{21}(\lambda - 7)^8(\lambda + 5)^7, \tag{94}$$

so the above 36 equations correspond to the vanishing of the imaginary part of $F_{\alpha\beta}$ (28 equations), the seven-dimensional irreducible piece of the real part of $F_{\alpha\beta}$ and the singlet trace condition on the (1,1)-curvature.

Deleting all terms containing f , the 16th index, from the above equations yields 36 equations in 15 dimensions which projects the curvature to the 69-dimensional $\lambda = -1$ eigenspace of the corresponding four-form $T_{(15)}$, which has characteristic polynomial

$$\chi(T_{(15)}) = (\lambda + 1)^{69}(\lambda - 6)^8(\lambda - 3)^{14}(\lambda^2 + 3\lambda - 6)^7. \tag{95}$$

8. The Reductions to $8 < d < 12$

We now briefly comment on some reductions of the above 12-dimensional system to the lower dimensions which were missed out in the discussion above.

$d = 11$

Deleting all terms containing dx^b in (65), (66) and (67) yields a set of 17 equations in 11-dimensions. The correspondingly reduced four-form $T_{(11)} := T_{(12)}|_{dx^b=0}$ has characteristic polynomial

$$\chi(T_{(11)}) = (\lambda + 1)^{38}(\lambda - 2)^8(\lambda - 3)^5(\lambda - 4)^2(\lambda^2 + \lambda - 4). \quad (96)$$

$d = 10$

Reducing the above 11-dimensional four-form further to the 10-dimensional hypersurface defined, for instance, by $x^6 = 0$ yields a four-form with characteristic polynomial

$$\chi(T_{(10)}) = (\lambda + 1)^{30}(\lambda - 1)^8(\lambda - 3)^6(\lambda - 4). \quad (97)$$

The $\lambda = -1$ eigenspace corresponds to a set of 15 equations amongst the 45 curvature components. This case is the complex counterpart of the $d = 5$ case discussed at the end of Sect. 4.1. In \mathbb{C}^5 , these equations take the form (15), (16) with $\alpha, \beta = 1, \dots, 5$ and the complex (4,0)-form given by the contraction of the (5,0) volume form with a constant unit (0,1)-vector. This (4,0)-form is the $SU(4)$ -invariant volume form in the four-dimensional complex space orthogonal to this vector. Choosing, this vector, for instance in the direction of the z^5 -axis, we obtain $\Phi = dz^{1234}$, yielding the following equations on \mathbb{C}^5

$$\begin{aligned} F_{1\bar{1}} + F_{2\bar{2}} + F_{3\bar{3}} + F_{4\bar{4}} + F_{5\bar{5}} &= 0 \\ F_{12} + F_{3\bar{4}} &= F_{13} + F_{4\bar{2}} = F_{14} + F_{2\bar{3}} = 0 \\ F_{15} &= F_{25} = F_{35} = F_{45} = 0. \end{aligned} \quad (98)$$

$d = 9$

The most symmetric reduction of (98) to nine-dimensions, making z^5 real, is a trivial embedding of the $Spin(7)$ -invariant set of equations (56) in nine dimensions.

9. Some Open Questions

An intriguing open problem is the relation of the 12-dimensional system to sextonions and to the ‘missing row’ of the Freudenthal magic square related to $E_{7\frac{1}{2}}$ (see [7, 13]).

It remains to be seen whether interesting solutions to the higher dimensional equations presented here can be found. The simplest solutions are embeddings of known solutions in dimensions $d \leq 8$ in the higher dimensional theories. Do such solutions allow oxidation to nontrivial solutions in higher dimensions? In particular, when the duality equations describe (partial) curvature flows, to what extent do solutions in the bulk arise from solutions on the initial value surface (boundary) of the flow? For instance, can the known four-dimensional solutions of the self-duality equations (27) be seen as arising from a flow which has a flat 3D connection as its initial value, or do the

known solutions of the eight-dimensional Spin(7)-invariant equation [9, 10, 46] arise as solutions of the flow equation (53) from solutions (e.g. [45]) of the G_2 -invariant equation (47) on the initial value sevenfold?

Acknowledgements. This work has benefitted a great deal from innumerable discussions over the last 30 years with Jean Nuyts. I acknowledge useful discussions with Andrea Spiro and Gregor Weingart, as well as partial funding from the SFB 647 “Raum-Zeit-Materie” of the Deutsche Forschungsgemeinschaft. I should like to thank Hermann Nicolai and the Albert-Einstein-Institut for hospitality.

References

1. Besse, A.L.: Einstein Manifolds. Springer, Berlin (1987)
2. Devchand, C., Nuyts, J., Weingart, G.: Matryoshka of special democratic forms. *Commun. Math. Phys.* **293**, 545–562, arXiv:0812.3012 (2010)
3. Devchand, C., Nuyts, J., Weingart, G.: Special graphs. *Int. J. Geom. Meth. Mod. Phys.* **3**, 1011–1018 (2006) arXiv:math/0604558.
4. Julia, B.: Group disintegrations. In: Rocek, M., Hawking, S. (eds.) *Superspace and Supergravity*. CUP, Cambridge (1980)
5. Corrigan, E., Devchand, C., Fairlie, D.B., Nuyts, J.: First order equations for gauge fields in spaces of dimension greater than four. *Nucl. Phys. B* **214**, 452–464 (1983)
6. Westbury, B.W.: Sextonions and the magic square. *J. London Math. Soc.* **73**(2), 455–474 (2006). arXiv:math/0411428
7. Landsberg J.M., Manivel L.: The sextonions and $E_{7\frac{1}{2}}$. *Adv. Math.* **201**, 143–179 (2006). arXiv:math/0402157
8. Kleinfeld, E.: On extensions of quaternions. *Indian J. Math.* **9**, 443–446 (1967)
9. Fairlie, D.B., Nuyts, J.: Spherically symmetric solutions of gauge theories in eight-dimensions. *J. Phys. A* **17**, 2867–2872 (1984)
10. Fubini, S., Nicolai, H.: The octonionic instanton. *Phys. Lett. B* **155**, 369–372 (1985)
11. Brihaye, Y., Devchand, C., Nuyts, J.: Selfduality for eight-dimensional gauge theories. *Phys. Rev. D* **32**, 990–994 (1985)
12. Alekseevsky, D.V., Cortes, V., Devchand, C.: Yang–Mills connections over manifolds with Grassmann structure. *J. Math. Phys.* **44**, 6047–6076 (2003). arXiv:math/0209124
13. Ward, R.S.: Completely solvable gauge field equations in dimension greater than four. *Nucl. Phys. B* **236**, 381–396 (1984)
14. Corrigan, E., Goddard, P., Kent, A.: Some comments on the ADHM construction in $4k$ dimensions. *Commun. Math. Phys.* **100**, 1–13 (1985)
15. Uhlenbeck, K., Yau, S.T.: On the existence of Hermitian–Yang–Mills connections in stable vector bundles. *Commun. Pure Appl. Math.* **39**, S257–S293 (1986)
16. Capria, M.M., Salamon, S.M.: Yang–Mills fields on quaternionic spaces. *Nonlinearity* **1**, 517–530 (1988)
17. Nitta, T.: Vector bundles over quaternionic Kähler manifolds. *Tohoku Math. J.* **40**, 425–440 (1988)
18. Itoh, M., Nakajima, H.: Yang–Mills connections and Einstein–Hermitian metrics. In: *Kähler metric and moduli spaces*, Adv. Stud. Pure Math., vol. 18, pp. 395–457. Academic Press, Boston (1990)
19. Donaldson, S.K., Thomas, R.P.: Gauge theory in higher dimensions. In: Huggett, S.A., et al (eds.) *The geometric universe: science, geometry, and the work of Roger Penrose*, pp. 31–47. Oxford Univ. Press, Oxford (1998)
20. Tian, G.: Gauge theory and calibrated geometry I. *Ann. Math. (2)* **151**, 193–268 (2000)
21. Baulieu, L., Kanno, H., Singer, I.M.: Special quantum field theories In eight and other dimensions. *Commun. Math. Phys.* **194**, 149–175 (1998). arXiv:hep-th/9704167v2
22. Devchand, C., Nuyts, J.: Super-selfduality for Yang–Mills fields in dimensions greater than four. *JHEP* **0112**, 020 (2001). arXiv:hep-th/0109072
23. Nakajima, H.: Removable singularities for Yang–Mills connections in higher dimensions. *J. Fac. Sci. Univ. Tokyo Sect. IA Math.* **34**, 299–307 (1987)
24. Bourguignon, J.P., Lawson, H.B., Simons, J.: Stability and gap phenomena for Yang–Mills fields. *Differential geometry of Yang–Mills fields*. Proc. Nat. Acad. Sci. **76**, 1550 (1979)
25. Jaffe, A., Taubes, C.: *Vortices and Monopoles*. pp. 32 Birkhauser, Boston (1980)
26. Price, P.: A monotonicity formula for Yang–Mills fields. *Manuscr. Math.* **43**, 131–166 (1983)
27. Nakajima, H.: Compactness of the moduli space of Yang–Mills connections in higher dimensions. *J. Math. Soc. Japan* **40**, 383–392 (1988)
28. Harvey, R., Lawson, H.B.: Calibrated geometries. *Acta Math.* **148**, 47–157 (1982)

29. Devchand, C., Fairlie, D.B.: Bogomolny equations and dimensional reduction. *Phys. Lett. B* **141**, 73 (1984)
30. Coleman, S.R.: Quantum Tunneling and Negative Eigenvalues. *Nucl. Phys. B* **298**, 178–186 (1988)
31. Witten, E.: Supersymmetry and Morse theory. *J. Diff. Geom.* **17**, 661–692 (1982)
32. Corrigan, E., Goddard, P.: Construction of instanton and monopole solutions and reciprocity. *Ann. Phys.* **154**, 253–279 (1984)
33. Fairlie, D.B., Ueno, T.: Higher dimensional generalizations of the Euler top equations. *Phys. Lett A* **240**, 132–136 (1998). arXiv:hep-th/9710079
34. Tao, T.: Geometric renormalization of large energy wave maps, Journées équations aux dérivées partielles, Exp. No. **11**, 1–32 (2004). arXiv:math/0411354v1
35. Atiyah, M.F., Bott, R.: The Yang–Mills equations over Riemann surfaces. *Phil. Trans. Roy. Soc. Lond. A* **308**, 523–615 (1982)
36. Witten, E.: Quantum field theory and the Jones polynomial. *Commun. Math. Phys.* **121**, 351–399 (1989)
37. Yang, C.N.: Condition of self-duality for SU(2) gauge fields on euclidean four-dimensional space. *Phys. Rev. Lett.* **38**, 1377–1379 (1977)
38. Brihaye, Y., Fairlie, D.B., Nuyts, J., Yates, R.G.: Properties of the self dual equations for an SU(n) gauge theory. *J. Math. Phys.* **19**, 2528–2532 (1978)
39. Leznov, A.N.: Equivalence of four-dimensional self-duality equations and the continuum analog of the principal chiral field problem. *Teor. Mat. Fiz.* **73**, 302–307 [*Theor. Math. Phys.* **73** 1233–1237 (1988)] (1987)
40. Donaldson, S.K.: Anti self-dual Yang–Mills connections over complex algebraic surfaces and stable vector bundles. *Proc. Lond. Math. Soc.* **50**, 1–26 (1985)
41. Nair, V.P., Schiff, J.: Kähler Chern–Simons theory and symmetries of anti-self-dual gauge fields. *Nucl. Phys. B* **371**, 329–352 (1992)
42. McKay, W.G., Patera, J.: Tables of dimensions, indices and branching rules for representations of simple Lie algebras. Marcel Dekker, New York (1981)
43. Onishchik, A.L., Vinberg, E.B.: Lie groups and algebraic groups. Springer, Berlin (1990)
44. Sudbery, A.: Quaternionic analysis. *Math. Proc. Cambridge Philos. Soc.* **85**(2), 199–224 (1979)
45. Gunaydin, M., Nicolai, H.: Seven-dimensional octonionic Yang–Mills instanton and its extension to an heterotic string soliton. *Phys. Lett. B* **351**, 169–172 (1995). arXiv:hep-th/9502009
46. Harland, D., Ivanova, T.A., Lechtenfeld, O., Popov, A.D.: Yang–Mills flows on nearly Kähler manifolds and G_2 -instantons. *Commun. Math. Phys.* **300**, 185–204 (2010). arXiv:0909.2730

Communicated by P. T. Chruściel