

Recent Progress in Group Field Theory

Daniele Oriti

*Max Planck Institute for Gravitational Physics (Albert Einstein Institute)
Am Mühlenberg 1 D-14476 Golm, Germany, EU*

Abstract. We introduce the key ideas behind the group field theory approach to quantum gravity, and the basic elements of its formalism. We also briefly report on some recent results obtained in this approach, concerning both the mathematical definition of these models, and possible avenues towards extracting interesting physics from them.

Keywords: quantum gravity, loop quantum gravity, spin foam models, matrix models, non-commutative geometry, simplicial quantum gravity

PACS: 04.60.Pp, 04.60.Gw, 04.60.Nc, 11.10.Nx

The field of background independent quantum gravity is progressing fast [1]. Not only new research directions are being developed, and new important developments are taking place in existing approaches, but some of these approaches are converging to one another, leading to further progress. The group field theory formalism [4, 2, 3] can be understood in several different ways. It is a generalization matrix models for 2d quantum gravity [5]. It is an important part, nowadays, of the loop quantum gravity and spin foam approach to the quantization of 4d gravity [7, 6]. It is a point of convergence of loop quantum gravity and of simplicial quantum gravity approaches, like quantum Regge calculus and dynamical triangulations [2]. Recently, tools from non-commutative geometry have been introduced as well in the formalism, which, thanks to them, started to make tentative contact with quantum gravity phenomenology.

In this paper we introduce the general idea behind the GFT formalism, and some basic elements of the same (for more detailed introduction, we refer to [2, 3, 4]), then report briefly on some recent results.

THE GROUP FIELD THEORY FORMALISM

Group field theories are an attempt to define quantum gravity in terms of *combinatorially non-local quantum field theories on group manifolds*, related to the Lorentz or rotation group. Let us motivate briefly the three main elements in this characterization.

A good theory of quantum gravity should be background-independent, as it should explain origin and properties of *spacetime* itself, while we know how to formulate quantum field theories only on fixed backgrounds. Therefore, if an (almost) ordinary QFT it should be, quantum gravity can only be a QFT on some auxiliary, internal or “higher-level” space. What background (non-dynamical) structures in GR could provide such space. One is the internal, local symmetry group of the theory, i.e. the Lorentz group. This gives the primary motivation for using the Lorentz group (and related) in GFT. Another background structure is the configuration space of GR: the (meta-)space

of (spatial) geometries on a given (spatial) topology, coined “superspace” by Wheeler. We will see that these two possibilities for the field domain in GFT actually coincide, because, as we have learned from loop quantum gravity, the set of possible geometries can be characterized in terms of (Lorentz) group elements. The other key ingredient of GFTs is their *combinatorial non-locality*. Consider a point particle in 0+1 dimensions, with action $S(X) = \frac{1}{2}X^2 + \frac{\lambda}{3}X^3$. This action defines a trivial dynamics (for a trivial system), of course. What interests us here, however, is the combinatorial structure of its “Feynman diagrams”, i.e. the graphs that can be used as a convenient book-keeping tool in computing the corresponding partition function $Z = \int dX e^{-S_\lambda(X)}$ perturbatively in λ . These are simple 3-valent (because of the order of the “interaction”) graphs.




FIGURE 1. A Feynman graph for a point particle and the corresponding field theory

The fact that the Feynman diagrams of the theory are simple graphs like the above follows from 1) the *point-like* nature of the particle, and 2) the *locality* of the corresponding interaction, encoded in the identification of X variables in the interaction term.

The same structure of diagrams is maintained, because the local nature of the interaction and the point-like nature of the corresponding quanta are maintained, also when moving to a field theory setting, i.e. going from the above particle dynamics to the corresponding field theory (still dynamically rather trivial), governed by the action: $S(\phi) = \frac{1}{2} \int dx \phi(x)^2 + \frac{\lambda}{3} \int dx \phi(x)^3$. The associated Feynman amplitudes have integrations over position or momentum variables, but still, the combinatorics of the diagrams is the same.

Now we move up in combinatorial dimension. Instead of point particles, let us consider 1d objects, that could be represented graphically by a line, with two end points. We label these two end points with two indices i, j , and we represent the fundamental 1d objects by (e.g. orthogonal) *matrices* M_{ij} . We want to move up in dimension also in the corresponding Feynman diagrams. We want diagrams corresponding to 2-dimensional structures. For this, we have to drop the assumption of *locality*. We define an action for M , given by $S(M) = \frac{1}{2} M_{ij}M_{ji} + \frac{\lambda}{3} M_{ij}M_{jk}M_{ki}$. The racing of indices i, j, k in the kinetic and vertex term represent identification of the points labeled by the same indices. This graphical representation of the Feynman diagrams used in evaluating the partition function $Z = \int \mathcal{D}M_{ij} e^{-S(M)}$ gives 2-dimensional simplicial complexes of arbitrary topology, because obtained by arbitrary gluing of lines to form triangles (in the interaction vertex) and of these triangles to one another along common lines (as dictated by the propagator).

Matrix models [5] have been quite successful in describing 2d quantum gravity.

There is no obstruction to keep moving upward in combinatorial dimension. We can move from 1d objects represented by matrices, with indices labeling the end points of the line, to 2d (closed) objects, represented by tensors, with indices labeling the boundary edges of the same 2d objects. The simplest combinatorial choice is that of triangles (2d simplices) and thus of tensors T_{ijk} , with indices (representing the edges of the triangles)




FIGURE 2. A (piece of) Feynman diagram for a matrix model, of which we give both direct and dual (simplicial) representation; the two parallel lines of propagation correspond to the two indices of the matrix; the extra line on the bottom indicates identification of the two edges of the triangles.

traced out in the interaction term in such a way as to represent 3d objects, tetrahedra (3-simplices) bounded by such triangles.




FIGURE 3. A (piece of) Feynman diagram for a tensor model, of which we give both direct and dual (simplicial) representation; the three parallel lines of propagation (dual to the three edges in the triangles) correspond to the three indices of the tensor.

The process of combinatorial generalization can be continued to tensor models whose Feynman diagrams are d -dimensional simplicial complexes. Moreover, while maintaining the combinatorial structure of the theory, we can generalize matrix and tensor models in the direction of adding degrees of freedom, i.e. defining corresponding field theories. The tensor indices will be replaced by variables taking values in appropriate domain spaces. Choosing these spaces to be group manifolds, we obtain group field theories.

The GFT field is a \mathbb{C} -valued function of D group elements $\phi(g_1, \dots, g_d)$, for a group G , usually the $SO(d-1, 1)$ Lorentz group (or $SO(d)$ for Riemannian gravity). It is interpreted as the fundamental building block of quantum space, a second quantized $(d-1)$ -simplex, and each argument corresponds to one of its boundary $(d-2)$ -faces. One imposes invariance under diagonal action of the group G : $\phi(g_1, \dots, g_d) = \phi(g_1 g, \dots, g_d g)$ to give closure of the d $(d-2)$ -faces to form a $(d-1)$ -simplex. The mode expansion gives:

$$\phi(g_i) = \sum_{J_i, \Lambda, k_i} \phi_{k_i}^{J_i \Lambda} \prod_i D_{k_i l_i}^{J_i} (g_i) C_{l_1 \dots l_d}^{J_1 \dots J_d \Lambda},$$

with the J 's labeling representations of G , the k 's being vector indices in the representation spaces, and the C 's being group intertwiners, labeled by Λ . So, dually, the same field quanta can be interpreted and represented as *spin network* vertices for the group G . The GFT amplitudes will justify a geometric interpretation of the group variables as parallel transport of the gravity connection along elementary paths dual to the $(d-2)$ -faces, and of the representations J as quantum numbers for volume operators for the same $(d-2)$ -faces.

The combinatorics of the field arguments in the action

$$S = \frac{1}{2} \int dg_i d\tilde{g}_i \phi(g_i) \mathcal{K}(g_i \tilde{g}_i^{-1}) \phi(\tilde{g}_i) + \frac{\lambda}{(d+1)!} \int dg_{ij} \phi(g_{1j}) \dots \phi(g_{d+1j}) \mathcal{V}(g_{ij} g_{ji}^{-1}),$$




FIGURE 4. The 3d example: the field, in group and representation picture, and its dual graphical representations as triangle and as spin network vertex

is constructed so to obtain d -dimensional complexes Γ as Feynman diagrams. It describes the interaction of $d+1$ ($d-1$)-simplices to form a d -simplex by gluing them along their ($d-2$)-faces (arguments of the fields). The interaction is specified by the choice of function \mathcal{V} . The kinetic term involves two fields each representing a given ($d-1$)-simplex seen from one of the two d -simplices (interaction vertices) sharing it. Thus the function \mathcal{K} specifies how the geometric degrees of freedom are propagated across simplices.




FIGURE 5. The 3d example: the simplicial representation of the interaction and of the propagation

The resulting complexes/triangulations can have arbitrary topology, because each simplicial complex is built up by arbitrary gluings of d -simplices to one another.

In representation space, each Feynman diagram is given by a spin foam (a 2-complex with faces f labeled by representation variables), and each Feynman amplitude by a spin foam model: $Z(\Gamma) = \sum_{\{J_f\}} A(\{J_f\})$. These models had been introduced in the loop quantum gravity approach to define the covariant dynamics of states of the quantum gravitational field, i.e. spin networks, and in fact transition amplitudes between given spin network states on the boundary is obtained in GFT as the expectation value of field operators having the same combinatorial structure of the two spin networks. The same representation variables in loop quantum gravity, have a geometric interpretation as quantum numbers of geometric operators, thus each of these Feynman amplitudes defines a sum-over-histories for discrete quantum gravity on the specific triangulation dual to the Feynman diagram. They can be understood, therefore, as a new implementation of the idea of simplicial gravity path integrals, like quantum Regge calculus and dynamical triangulations. To summarize: GFT incorporates many ingredients of other approaches to quantum gravity. Boundary states are spin networks, as in LQG, but they have a dual description as simplicial spaces, as in simplicial quantum gravity. Their dynamics is expressed as a covariant sum over quantum geometries as in spin foam models, or a discrete gravity path integral, as in quantum Regge calculus, but involves also a sum over inequivalent triangulations, as in the dynamical triangulations approach.

As an explicit example, consider the Boulatov GFT model for 3d quantum gravity:

$$S_{3d}[\phi] = \frac{1}{2} \int [dg]^3 \phi(g_1, g_2, g_3) \phi(g_3, g_2, g_1) + \\ - \frac{\lambda}{4!} \int [dg]^6 \phi(g_1, g_2, g_3) \phi(g_3, g_4, g_5) \phi(g_5, g_2, g_6) \phi(g_6, g_4, g_1) \quad (1)$$

for a real field $\phi : \text{SU}(2)^3 \rightarrow \mathbb{R}$ invariant under the diagonal right action of $\text{SU}(2)$: $\phi(g_1, g_2, g_3) = \phi(g_1 g, g_2 g, g_3 g) \forall g \in \text{SU}(2)$.

The Feynman diagrams are dual to 3d triangulations, and the Feynman amplitudes are

$$Z(\Gamma) = \prod_{L \in \Gamma} \int dh_L \prod_f \delta\left(\prod_{L \in \partial f} h_L\right)$$

with a single delta function on the group $\delta(\prod_{L \in \partial f} h_L)$ for each 2-cell f in the Feynman diagram, dual to a single edge e of the simplicial complex, with argument given by the product of group elements h_L each associated to a link in the boundary of the 2-cell.

The *same* amplitude can be given in representation space:

$$Z(\Gamma) = \left(\prod_f \sum_{j_f} \right) \prod_f (2j_f + 1) \prod_v \left\{ \begin{matrix} j_1 & j_2 & j_3 \\ j_4 & j_5 & j_6 \end{matrix} \right\},$$

where the symbol for the vertex amplitude is the well-know $6j$ -symbol from the recoupling theory of angular momentum This is the Ponzano-Regge spin foam model.

At this stage the connection with simplicial path integrals is rather indirect, but still very clear. Consider the continuum 3d gravity action:

$$S(e, \omega) = \int_{\mathcal{M}} \text{tr}(e \wedge F(\omega)),$$

with variables the triad 1-form $e^I(x) \in \mathfrak{su}(2)$ and the 1-form connection $\omega^J(x) \in \mathfrak{su}(2)$, with curvature $F(\omega)$. Introducing the simplicial complex Δ and its topological dual cellular complex, we discretize the triad in terms of Lie algebra elements associated to the edges of the simplicial complex as $E_e = E_f = \int_e e(x) = E^i J_i \in \mathfrak{su}(2)$, and the connection in terms of parallel transports along links of the dual complex as $g_L = e^{\int_L \omega} \in \text{SU}(2)$. The discrete curvature is given by: $G_f = G_e = \prod_{L \in \partial f} g_L = e^{\int_f F} \in \text{SU}(2)$, and a discrete counterpart of the continuum action is: $S(E, g) = \sum_{f \in \Gamma} \text{tr}(E_f G_f)$.

The path integral for this discrete theory on the simplicial complex Δ is *equal* to the previously computed GFT Feynman amplitude $Z(\Gamma)$ for the diagram Γ dual to Δ :

$$Z(\Gamma)_{GFT} = \prod_f \int_{\mathfrak{su}(2)} dE_f \prod_L \int_{\text{SU}(2)} dg_L e^{i \sum_f \text{tr}(E_f G_f)} = \prod_L \int_{\text{SU}(2)} dg_L \prod_f \delta(G_f),$$

which, as we have seen, can be also expressed as a function of group representations j_e .

From all this, we also learn that spin foam models can be seen as a way of re-writing simplicial gravity path integrals expressed in connection (group) variables.

SOME RECENT RESULTS

Group field theory renormalization and the sum over topologies

An area of recent developments [10, 11] has been the application of quantum field theory techniques to GFTs, to gain a better understanding and control over its perturbative expansion, using tools from renormalization theory.

GFTs define a sum over simplicial complexes 1) of arbitrary topology and 2) that correspond, in general, to pseudo-manifolds, i.e. contain conical singularities at the vertices. The issue of controlling the sum over topologies, and of identifying an approximation in which simple topologies dominate, has an analogue in the context of matrix models [5]. In matrix models, it has been shown that, in the so-called large- N limit, diagrams of trivial topology (S^2 in the compact case) dominate the perturbative sum. The issue of controlling the relative weight of manifolds and pseudo-manifolds in the perturbative sum arises instead only in dimensions $D > 2$ and it has represented an obstacle to the development of tensor models. A third issue is to identify and control the divergences that arise in this perturbative expansion, which are of two types: a) divergences in the sum over (pre-)geometric data (group elements or group representations) for each amplitude associated to a given simplicial complex; b) the divergence of the entire sum over simplicial complexes.

The work of [10] makes the first steps toward solving these three issues, leaving aside 3b), starting a systematic study of GFT renormalization, in the context of the Boulatov model for 3d (Riemannian) quantum gravity, that we have discussed above. The divergences of this model are related to the topology of the bubbles (3-dimensional cells), dual to vertices of the simplicial complex, in the Feynman diagrams, but it is difficult to establish which diagrams need renormalization in full generality, mainly due to the very complicated topological structure of 3D simplicial complexes, after a scale is introduced in the theory by an explicit cut-off in the spectral decomposition of the propagator. What is achieved in [10] is the following:

- a detailed algorithm is given for identifying bubbles (3-cells) in the Feynman diagrams of the model, together with their boundary triangulations, which in turn can be used to identify the topology (genus) of the same boundary.
- using this algorithm, the authors are able to identify a subclass of Feynman diagrams which allow for a complete contraction procedure, and thus the ones that allow for an almost standard renormalization; moreover, this class of graphs, dubbed “type 1” is shown to be a natural generalization of the 2d planar graphs of matrix models, thus suggesting that they can play a similar role in GFTs to that of planar diagrams in matrix models. Notice also that the existence of such contraction procedure can be seen as a sort of generalised locality property
- for this class of diagrams, an exact power counting of divergences is proven, according to which their divergence is of the order:

$$A_{\Gamma} = \left(\delta^{\Lambda}(I) \right)^{|\mathcal{B}_{\Gamma}|-1}$$

where $|\mathcal{B}_\Gamma|$ is the number of bubbles in the diagram Γ , and $\delta^\Lambda(I)$ is the delta function on the group, with cut-off Λ , evaluated at the identity I .

On the basis of these results, of the experience gained with explicit evaluation of Feynman amplitudes, and of a better understanding of the combinatorial structure of the Feynman diagrams, it is then possible to put forward two main conjectures, obviously confirmed in all examples considered: 1) that *all* “type 1” diagrams correspond to *manifolds of trivial topology*; and 2) that an appropriate generalization of the usual scaling limit (large- N) of matrix models to these GFT would lead to the relative suppression of all the “non-type 1” diagrams, and thus leave us with: only type 1 diagrams in need for renormalization, and only manifolds of trivial topology in the theory.

A different perspective on GFT divergences is taken in [11], which also tackles the difficult issue of the summability of the entire perturbative sum (thus including the sum over topologies). The authors consider both the Boulatov model and a modification of the same proposed in [12], obtained adding a second interaction term in the action:

$$+\frac{\lambda}{4!} \delta \prod_{i=1}^6 \int dg_i [\phi(g_1, g_2, g_3) \phi(g_3, g_4, g_5) \phi(g_4, g_2, g_6) \phi(g_6, g_5, g_1)]. \quad (2)$$

The new term corresponds to the only other possible way of gluing 4 triangles to form a closed surface. This mild modification gives a Borel summable partition function [12]. This shows that a control over the sum over topologies and a non-perturbative definition of the corresponding GFT is feasible.

For both the Boulatov model and the modified one, the authors of [11] establish general perturbative bounds on amplitudes using powerful constructive techniques, rather than focusing on explicit power counting or Feynman evaluations. They find that, using the same regularization used in [10], the amplitudes of the Boulatov model for a diagram with n vertices, are bounded, with cut-off Λ , by $K^n \Lambda^{6+3n/2}$, for some arbitrary positive constant K , while the modified model has amplitudes bounded by $K^n \Lambda^{6+3n}$, and that both bounds can be saturated. This result shows that the Freidel-Louapre modification (BFL), even though Borel summable, is perturbatively more divergent than the original model.

The second main result of [11] relies again on constructive field theory techniques. A cactus expansion of the BFL model is obtained, and used to prove the Borel summability of the free energy of the model and to define its Borel sum. We can expect more applications of these techniques to other GFT models, also in higher dimensions.

Emergent non-commutative matter fields from group field theories

The last set of results we want to mention are interesting steps in the direction of bridging the gap between the microscopic GFT description of quantum space (and the language of spin networks, simplices, spin foams, etc) and macroscopic continuum physics, including usual General Relativity and quantum field theories for matter. In fact, this problem is faced by *all* discrete approaches to quantum gravity[1].

One would expect [9] a generic continuum spacetime to be formed by zillions of Planck size building blocks, and thus to be, from the GFT point of view, a many-particle

system whose microscopic theory is given by some fundamental GFT action. This suggests us to look for ideas and techniques from statistical field theory and condensed matter theory, and to try to apply/re-interpret them in a GFT context.

Condensed matter theory also provides examples of systems in which the collective behaviour of the microscopic constituents, in some hydrodynamic approximation, gives rise to effective emergent geometries from the collective variables themselves [13]. What happens is that the collective parameters describing the fluid and its dynamics can be recast as the component functions of an *effective metric field*, and that the effective dynamics of perturbations (quasi-particles, themselves collective excitations of the fundamental constituents of the fluid) takes the form of matter field theories in curved spacetimes, on the same effective metrics. Inspired by these results, we ask: assuming that a given GFT model (Lagrangian) describes the microscopic dynamics of (the fundamental building blocks of) a *discrete quantum spacetime*, and that some solution of the corresponding fundamental equations can be interpreted as identifying a given quantum spacetime configuration, 1) can we obtain an effective macroscopic *continuum* field theory for matter fields from it? and if so, 2) what is the effective spacetime and geometry that these emergent matter fields see? As it turns out, it is possible [8] to apply the same procedure to GFT models and obtain rather straightforwardly effective continuum field theories for matter fields. The effective matter field theories that we obtain most easily are QFTs on non-commutative spaces of Lie algebra type.

The basic point is the use of the same natural duality between Lie algebra and corresponding Lie group, interpreted as the non-commutative version of the usual duality between coordinate and momentum space. More precisely, if we have a non-commutative spacetime of Lie algebra type $[X_\mu, X_\nu] = C_{\mu\nu}^\lambda X_\lambda$, the corresponding momentum space is naturally identified with the corresponding Lie group, in such a way that the non-commutative coordinates X_μ act on it as (Lie) derivatives (as we expect in the quantum setting). The link with GFTs is then obvious: in momentum space the field theory on such non-commutative spacetime will be given, by definition, by some sort of group field theory. The task will then be to derive the relevant field theories from matter from interesting GFT models of quantum spacetime.

In 3 spacetime dimensions the results obtained recently [14] concern the euclidean signature and a non-commutative spacetime given by the $su(2)$ Lie algebra, and we start again from the Boulatov GFT model. We look [14] at two-dimensional variations $\delta\phi(g_1, g_2, g_3) \equiv \psi(g_1 g_3^{-1})$ around a class of GFT classical solutions (they can be interpreted as quantum flat space on some a priori non-trivial topology) given by:

$$\phi^{(0)}(g_1, g_2, g_3) = \sqrt{\frac{3!}{\lambda}} \int dg \delta(g_1 g) F(g_2 g) \delta(g_3 g), \quad F: G \rightarrow \mathbb{R}, \quad \int F^2 = 1 \quad (3)$$

The effective action is then:

$$\begin{aligned} S_{eff}[\psi] &\equiv S_{3d}[\phi^{(0)} + \psi] - S_{3d}[\phi^{(0)}] = \frac{1}{2} \int \psi(g) \mathcal{K}(g) \psi(g^{-1}) - \\ &- \frac{\mu}{3!} \int [dg]^3 \psi(g_1) \psi(g_2) \psi(g_3) \delta(g_1 g_2 g_3) - \frac{\lambda}{4!} \int [dg]^4 \psi(g_1) \dots \psi(g_4) \delta(g_1 \dots g_4), \end{aligned}$$

with the kinetic term and the 3-valent coupling given in term of F :

$$\mathcal{H}(g) = 1 - 2 \left(\int F \right)^2 - \int dh F(h) F(hg), \quad \frac{\mu}{3!} = \sqrt{\frac{\lambda}{3!}} \int F.$$

with $F(g)$ assumed to be invariant under conjugation $F(g) = F(hgh^{-1})$. This is a non-commutative quantum field theory invariant under the quantum double of $SU(2)$ (which provides a quantum deformation of the Poincaré group $ISU(2)$).

Expanding F in group characters: $F(g) = \sum_{j \in \mathbb{N}/2} F_j \chi_j(g)$, where $j \in \mathbb{N}$ label irreducible representations of $SU(2)$, the kinetic term reads:

$$\mathcal{H}(g) = 1 - 3F_0^2 - \sum_{j \geq 0} \frac{F_j^2}{d_j} \chi_j(g) = \sum_{j \geq 0} F_j^2 \left(1 - \frac{\chi_j(g)}{d_j} \right) - 2F_0^2 \equiv Q^2(g) - M^2. \quad (4)$$

We can interpret $Q^2(g) \geq 0$ as a generalized ‘‘Laplacian’’, and F_0^2 as a ‘‘gravitational’’ mass M^2 . For the simple classical solution $F(g) = a + \sqrt{1 - a^2} \chi_1(g)$, we obtain

$$\mathcal{H}(g) = \frac{4}{3} (1 - a^2) \bar{p}^2 - 2a^2. \quad (5)$$

Similar results have also been obtained in the 4d context [15, 8]. It has been shown that from GFT models (indirectly) related to 4d quantum gravity, it is possible to derive effective non-commutative matter field theories of ‘‘deformed special relativity’’ type, based on momentum group given by $AN(3) \approx SO(4, 1)/SO(3, 1)$ and a non-commutative κ -Minkowski spacetime: $[x_0, x_i] = i\kappa x_i$; these field theories form the basis for much current work in the area of quantum gravity phenomenology [16].

Work in this direction, therefore, including these recent and preliminary results, is a step in bridging the gap between the fundamental discrete theory of spacetime we have at hand, and a continuum description of spacetime, and getting closer to possible quantum gravity phenomenology, thus bringing this class of models a bit closer to experimental falsifiability. Let us also notice that, contrary to the situation in analog gravity models in condensed matter, we have here models which are non-geometric and far from usual geometrodynamics in their formalism, but which at the same time are expected to encode quantum geometric information and indeed to determine, in particular in their classical solutions, a (quantum and therefore classical) geometry for spacetime [2]. We are, in other words, far beyond a pure analogy.

CONCLUSIONS

We have introduced the key ideas behind the group field theory approach to quantum gravity, and to the microscopics of quantum space, and the basic elements of its formalism. We have also briefly reported on some recent results obtained in this approach, concerning both the mathematical definition of these models, and possible avenues towards extracting interesting physics from them. From our outline it should be clear that, while much more work is certainly needed in this area of research, the new direction toward quantum gravity that group field theories provide is exciting and full of potential.

ACKNOWLEDGMENTS

We thank the organizers of the Max Born symposium, in particular Jurek Kowalski-Glikman, and all the attendees, for a very stimulating and enjoyable meeting.

REFERENCES

1. D. Oriti, ed., *Approaches to Quantum Gravity*, Cambridge University Press, Cambridge (2009)
2. D. Oriti, in [1], [arXiv: gr-qc/0607032]
3. D. Oriti, in *Quantum Gravity*, B. Fauser, J. Tolksdorf and E. Zeidler, eds., Birkhaeuser, Basel, (2007), [arXiv: gr-qc/0512103]
4. L. Freidel, Int.J.Phys. **44**, 1769-1783, (2005) [arXiv: hep-th/0505016]
5. F. David, Nucl. Phys. B257, **45** (1985); P. Ginsparg, 'Matrix models of 2-d gravity', [arXiv: hep-th/9112013]
6. D. Oriti, Rept. Prog. Phys. **64**, 1489, (2001), [arXiv: gr-qc/0106091]; A. Perez, 'Spin foam models for quantum gravity', Class. Quant. Grav. **20**, R43, (2003), [arXiv: gr-qc/0301113]
7. C. Rovelli, *Quantum Gravity*, Cambridge University Press, Cambridge (2006)
8. D. Oriti, J. Phys. Conf. Ser. **174**, 012047 (2009) [arXiv:0903.3970]
9. D. Oriti, Proceedings of Science, [arXiv:0710.3276]
10. L. Freidel, R. Gurau and D. Oriti, Phys. Rev. D **80**, 044007 (2009), [arXiv:0905.3772]
11. J. Magnen, K. Noui, V. Rivasseau and M. Smerlak, Class. Quant. Grav. **26**, 185012 (2009), [arXiv:0906.5477]
12. L. Freidel and D. Louapre, Phys. Rev. D **68**, 104004 (2003) [arXiv: hep-th/0211026]
13. C. Barcelo, S. Liberati, and M. Visser, Living Rev. Rel. **8**, 12 (2005) [arXiv: gr-qc/0505065]
14. W. Fairbairn and E. Livine, Class. Quant. Grav. **24**, 5277 (2007) [arXiv:gr-qc/0702125]
15. F. Girelli, E. Livine and D. Oriti, *4d deformed special relativity from group field theories*, [arXiv: 0903.3475]
16. G. Amelino-Camelia, Lect. Notes Phys. **669**, 59-100, [arXiv:gr-qc/0412136]