

SECOND QUANTIZED MATHIEU MOONSHINE

Daniel Persson

Fundamental Physics, Chalmers University of Technology,
412 96, Gothenburg, Sweden
`daniel.persson@chalmers.se`

Roberto Volpato

Max-Planck-Institut für Gravitationsphysik,
Am Mühlenberg 1, 14476 Golm, Germany
`roberto.volpato@aei.mpg.de`

Abstract

We study the second quantized version of the twisted twining genera of generalized Mathieu moonshine, and verify that they give rise to Siegel modular forms with infinite product representations. Most of these forms are expected to have an interpretation as twisted partition functions counting 1/4 BPS dyons in type II superstring theory on $K3 \times T^2$ or in heterotic CHL-models. We show that all these Siegel modular forms, independently of their possible physical interpretation, satisfy an “S-duality” transformation and a “wall-crossing formula”. The latter reproduces all the eta-products of an older version of generalized Mathieu moonshine proposed by Mason in the '90s. Surprisingly, some of the Siegel modular forms we find coincide with the multiplicative (Borcherds) lifts of Jacobi forms in umbral moonshine.

Contents

1	Introduction and Summary	3
1.1	Background	4
1.2	Summary of results	6
1.3	Outline	9
2	Twisted-twining partition functions	10
2.1	Definition and basic properties	10
2.2	Modular properties and the α -twist	11
2.3	(Generalized) Mathieu moonshine	13
2.4	Geometric perspective	16
3	Second quantization of Mathieu moonshine	17
3.1	Symmetric Orbifolds and Hecke Operators	17
3.1.1	Symmetric Orbifolds	17
3.1.2	Hecke operators	19
3.2	Twisted equivariant Hecke operators	20
3.2.1	Main definition	20
3.2.2	Central extensions	21
3.2.3	Alternative definition	23
3.3	Second-Quantized Twisted Twining Genera	23
3.3.1	Definition	23
3.3.2	Infinite product representation	25
3.3.3	Multiplicative versus additive lift	27
4	Wall-crossing and Mason's generalized moonshine for M_{24}	28
4.1	Mason's generalized M_{24} -moonshine	29
4.2	Connecting the two moonshines via wall-crossing	30
5	Automorphic Properties	32
5.1	Relabeling and S-duality	33
5.1.1	Orbifolds and relabeling	34
5.1.2	S -duality invariance	37
5.2	Automorphic properties	40
5.3	Examples	43

6	Conclusions	48
A	Some group cohomology	49
B	Modular properties of twisted twining partition functions	52
B.1	$SL(2, \mathbb{Z}) \times G$ - action on \mathcal{P}_G	52
B.2	Twisted action on \mathcal{P}_G : generalized permutation	53
B.3	Modular properties of twisted-twining partition functions	54
B.4	Cohomological obstructions	55
C	Definition of twisted equivariant Hecke operators	56
C.1	Twisted equivariant Hecke operator	56
C.2	Definition of slash operator for $\text{Mat}(\mathbb{Z}) \times G$	57
C.3	Proof of consistency	59
D	Some technical proofs	62
D.1	Proof of (3.53).	62
D.2	Central extensions and special choices of the cocycle	64
D.2.1	Case $\lambda = 1$ (trivial multiplier)	65
D.2.2	Case $\lambda \neq 1$ (non-trivial multiplier)	66
D.3	S-duality for g in classes $2B, 3B, 4C, 6B, 12B$	67
E	Tables	72

1 Introduction and Summary

Mathieu moonshine [1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16] and its generalized version [17, 18] pertain to the association of a class of weak Jacobi forms $\phi_{g,h}(\tau, z)$, called twisted twining genera, to each commuting pair of elements (g, h) in M_{24} . It has been verified that these functions satisfy all the requirements of Norton's generalized moonshine conjectures [19]; in particular they decompose into (projective) graded characters of the centralizer of g in M_{24} . Many of these Jacobi forms arise as supersymmetric indices in certain non-linear sigma models with target space K3; in particular, $\phi_{e,e}$ is the K3 elliptic genus. This obvious idea that M_{24} might be the symmetry group of some of these non-linear sigma models has been ruled out quite quickly [8, 10, 11] and the most recent works on the subject [20, 21, 22] seek for an explanation of Mathieu moonshine in the framework

of K3 compactifications of full-fledged superstring theory rather than within conformal field theory.

In this paper we adopt a similar viewpoint, and construct the second-quantized* version $\Psi_{g,h}$ of the twisted twining genera. From a mathematical perspective, the functions $\Psi_{g,h}$ are obtained from the Jacobi forms $\phi_{g,h}$ via a certain twisted equivariant version of Borchers multiplicative lift. Physically, many of these functions can be interpreted as twisted supersymmetric indices counting 1/4 BPS states in type II superstring theory compactified on $K3 \times T^2$ and in CHL models [23, 24, 25, 26]. A similar construction for the twining genera of the ordinary Mathieu moonshine was first proposed by Cheng [2].

We show that these multiplicative lifts are Siegel modular forms, generalizing results for the case $(g, h) = (e, h)$ [2, 27, 28]. In some cases we can identify $\Psi_{g,h}$ with known Siegel modular forms which have previously appeared in the context of umbral moonshine [29]. Furthermore, they satisfy a “wall-crossing formula”, which reproduces all of Mason’s generalized eta-products $\eta_{g,h}$ for M_{24} [30, 31, 32, 33]. This therefore establishes the link between the two existing versions of generalized moonshine for M_{24} .

1.1 Background

The first hint of Mathieu moonshine was found by Eguchi, Ooguri and Tachikawa [1], who noticed a connection between the elliptic genus of K3 and the largest Mathieu group M_{24} . The elliptic genus of K3 $\phi_{K3}(\tau, z)$ is the unique (up to normalization) weak Jacobi form of weight 0 and index 1 and it can be defined as a supersymmetric index non-linear sigma models with target space K3. It therefore is natural to consider a decomposition of $\phi_{K3}(\tau, z)$ into characters of the $\mathcal{N} = 4$ superconformal algebra. The authors of [1] noticed that the coefficients in this decomposition are sums of dimensions of irreducible representations of M_{24} . This observation led to the Mathieu moonshine conjecture: each conjugacy class $[g]$ of M_{24} there should exist a weak Jacobi form $\phi_g : \mathbb{H} \times \mathbb{C} \rightarrow \mathbb{C}$ such that the Fourier coefficients of ϕ_g are the characters of these M_{24} -representations evaluated at g . In particular, for the identity element e of M_{24} , $\phi_e(\tau, z)$ coincides with the elliptic genus of K3. In subsequent work [2, 3, 4, 5], all the functions ϕ_g , dubbed “twining elliptic genera”, were found and substantial evidence was given for the validity of Mathieu moonshine, namely the existence of a graded M_{24} -module such that the ϕ_g ’s are its graded characters. This conjecture has now been proven rigorously by Gannon [12].

The story outlined above is of course in close analogy with Monstrous moonshine [34,

*The terminology “second-quantized” originates in the work of Dijkgraaf, Moore, Verlinde, Verlinde [49].

35, 36] which assigns modular functions $T_g : \mathbb{H} \rightarrow \mathbb{C}$ (McKay-Thompson series) with each conjugacy class $[g]$ of the Fischer-Griess Monster group \mathbb{M} , the largest of the finite sporadic simple groups. After the initial conjecture [34], Norton proposed his generalized Monstrous moonshine [19], which assigns modular functions $f(g, h; \tau)$ to each commuting pair $g, h \in \mathbb{M}$. For fixed $g \in \mathbb{M}$, these generalized moonshine functions should then have (possibly rational) Fourier coefficients that correspond to projective characters of the centralizer $C_{\mathbb{M}}(g) = \{k \in \mathbb{M} \mid gk = kg\}$. Dixon-Ginsparg-Harvey subsequently suggested that Norton's functions $f(g, h; \tau)$ naturally arise in string theory as the path integral on a torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ with boundary conditions twisted by (g, h) along the (a, b) -cycles. Although the full generalized Monstrous moonshine conjecture is still open, considerable progress has been made toward proving it [37, 38, 39, 40, 41, 42, 43].

In earlier work [17, 18], we gave substantial evidence that Norton's generalization also holds, with small modifications, for Mathieu moonshine. We found that for each commuting pair $g, h \in M_{24}$ there exists a weak Jacobi form $\phi_{g,h}(\tau, z)$, dubbed *twisted twining genus*, whose Fourier coefficients are characters of a projective representation of $C_{M_{24}}(g)$. Inspired by orbifolds of holomorphic CFT's [44, 45, 46, 47], we further showed that the modular properties of the functions $\phi_{g,h}$ are controlled by a cohomology class $[\alpha]$ in the third cohomology group $H^3(M_{24}, U(1))$, as was anticipated in [48] (see also [12]).

A different kind of generalized moonshine for M_{24} had in fact already been established in old work by Mason [30, 32, 33]. Mason associated to each commuting pair g, h in M_{24} a so called multiplicative eta product $\eta_{g,h}(\tau)$, based on the action of M_{24} on 24 chiral free bosons. This leads to the natural question: *Is there a relation between the recently discovered Mathieu moonshine, pertaining to weak Jacobi forms, and Mason's M_{24} -moonshine involving eta-products?*

For the special case of commuting pairs $(g, h) = (e, h)$, where e is the identity element of M_{24} , this was given an affirmative answer by Cheng in [2]. Cheng's idea was to generalize the known fact that the elliptic genus of K3, $\phi_e(\tau, z)$, exhibits an exponential Borcherds lift to the unique weight 10 Siegel modular form for $Sp(4; \mathbb{Z})$, referred to as the "Igusa cusp form" and commonly denoted by Φ_{10} . She proposed that to all twining genera $\phi_h(\tau, z)$ one should have a similar lift of the form

$$\Phi_h(\sigma, \tau, z) = pqy \prod_{(n,m,\ell) > 0} \exp \left[- \sum_{k=1}^{\infty} \frac{c_{h^k}(4mn - \ell^2)}{k} (p^m q^n y^\ell)^k \right], \quad (1.1)$$

where $q = e^{2\pi i \tau}$, $y = e^{2\pi i z}$, $p = e^{2\pi i \sigma}$, and c_g denotes the Fourier coefficients of $\phi_g(\tau, z)$:

$$\phi_g(\tau, z) = \sum_{m \geq 0, \ell \in \mathbb{Z}} c_g(m, \ell) q^m y^\ell. \quad (1.2)$$

It was conjectured in [27] that Φ_h is automorphic with respect to a subgroup $\Gamma_h^{(2)}$ of $Sp(4; \mathbb{Z})$. This was then proven by Raum [28] for most of the conjugacy classes $[h] \subset M_{24}$.

A central point for us is that $\Phi_h(\sigma, \tau, z)$ has a double pole at $z = 0$ and in the limit one finds

$$\lim_{z \rightarrow 0} \frac{\Phi_h(\sigma, \tau, z)}{(2\pi iz)^2} = \eta_h(\tau) \eta_h(\sigma), \quad (1.3)$$

where $\eta_h = \eta_{e,h}$ are the eta-products of Mason's M_{24} -moonshine [30]. Hence, the process of taking the multiplicative lift of ϕ_h followed by studying the limiting behavior as $z \rightarrow 0$ provides a link between the two moonshines.

1.2 Summary of results

In this paper we answer the question above in the general case of commuting pairs (g, h) in M_{24} . In other words, we establish a link between the generalized Mathieu moonshine proposed in [17, 18] and the generalized eta-products of Mason. To describe our results, we recall the notion of “second quantized elliptic genus” as defined in [49]. Suppose $\phi_X(\tau, z)$ is the elliptic genus of some Calabi-Yau manifold X . Then one defines the second quantized genus Ψ_X as the generating function of the elliptic genus $\phi_{S^n X}$ for the n :th symmetric product $S^n X$: $\Psi_X = \sum_{n \geq 0} p^n \phi_{S^n X}$. In [49], the following remarkable formula is proved

$$\Psi_X(\sigma, \tau, z) = \exp \left[\sum_{n=1}^{\infty} p^n T_n \phi_X(\tau, z) \right], \quad (1.4)$$

where $p = e^{2\pi i \sigma}$, and T_n is the standard Hecke operator acting on Jacobi forms. Subsequently, Gritsenko showed [50] that by multiplying the inverse Ψ_X^{-1} by a certain factor $A_X(\sigma, \tau, z)$ (the “Hodge anomaly”) one obtains a Siegel modular form

$$\Phi_X(\sigma, \tau, z) = \frac{A_X(\sigma, \tau, z)}{\Psi_X(\sigma, \tau, z)}, \quad (1.5)$$

of weight $c_X(0, 0)/2$ for a subgroup $\Gamma_X^{(2)} \subset Sp(4; \mathbb{Z})$, where $c_X(m, n)$ are the Fourier coefficients of ϕ_X .

Inspired by these results, we define for each commuting pair (g, h) in M_{24} the associated *second quantized twisted twining genus* $\Psi_{g,h}$ via a generalization of the formula (1.4):

$$\Psi_{g,h}(\sigma, \tau, z) = \exp \left[\sum_{n=1}^{\infty} p^n \mathcal{T}_n^\alpha \phi_{g,h}(\tau, z) \right], \quad (1.6)$$

where \mathcal{T}_n^α is now a certain twisted equivariant Hecke operator (for the precise definition see section (3.2)) which reduces to the T_n in (1.4) in the special case $(g, h) = (e, e)$. Notice that,

when $g = e$, the presence of a non trivial 3-cocycle α governing the modular properties of the twisted-twining genera can be safely ignored (see [17] and section 2.2). This leads to the simpler definition of the multiplicative lift adopted in [27]. However, for the general case considered in this paper, the technical subtleties associated with non-trivial 3-cocycles cannot be avoided.

We show that upon multiplying the second quantized twisted twining genera by a correction $A_{g,h}(\sigma, \tau, z)$ (see eq. (3.61) for the explicit form of $A_{g,h}$) we obtain a class of Siegel modular forms

$$\Phi_{g,h}(\sigma, \tau, z) = \frac{A_{g,h}(\sigma, \tau, z)}{\Psi_{g,h}(\sigma, \tau, z)}, \quad (1.7)$$

for certain discrete subgroups $\Gamma_{g,h}^{(2)} \subset Sp(4; \mathbb{R})$, containing the invariance subgroups $\Gamma_{g,h} \subset SL(2, \mathbb{Z})$ of the twisted twining genera $\phi_{g,h}$. The most noticeable new automorphic property is the ‘S-duality’ transformation

$$\Phi_{g,h}(\sigma, \tau, z) = \Phi_{g,h'}\left(\frac{\tau}{N\lambda}, N\lambda\sigma, z\right), \quad (1.8)$$

where N is the order of g and λ is the length of the shortest cycle of g in the 24-dimensional permutation representation. Notice that $h, h' \in C_{M_{24}}(g)$ are not necessarily in the same conjugacy class. This is related with the ‘relabeling’ phenomenon described in [17]. In particular, as described in section 5.1.1, when g is an element of an M_{23} subgroup, then h' is the symmetry induced by h in the g -orbifold theory. We proved (1.8) for at least one generator g in each group $\langle g, h \rangle$, but we conjecture that similar relations hold for all commuting pairs g, h .

In section 4 we prove the following “wall-crossing formula”

$$\lim_{z \rightarrow 0} \frac{\Phi_{g,h}(\sigma, \tau, z)}{(2\pi iz)^2} = \eta_{g,h}(\tau) \eta_{g,h'}(N\lambda\sigma), \quad (1.9)$$

which generalizes (1.3). This shows that the limit $z \rightarrow 0$ of $\Phi_{g,h}(\sigma, \tau, z)$ reproduces all of Mason’s generalized eta-products $\eta_{g,h}$ [32, 31], thus providing the desired link between the two moonshines. See figure 1.2 for a pictorial overview of the relation between the various modular objects.

When (g, h) lies in the conjugacy classes $(2A, 2A)$, $(3A, 3A)$ and $(4B, 4B)$ we can identify $\Phi_{g,h}$ with known Siegel modular forms:

$$\begin{aligned} (2A, 2A) &: \Phi_{g,h} = (\Delta_2)^2 = \Phi^{(3)}, \\ (3A, 3A) &: \Phi_{g,h} = (\Delta_1)^2 = \Phi^{(4)}, \\ (4B, 4B) &: \Phi_{g,h} = (\Delta_{1/2})^2 = \Phi^{(5)}, \end{aligned} \quad (1.10)$$

where the first equality identifies them with the squares of the weight k Siegel modular forms Δ_k obtained by Gritsenko and Nikulin [51], and the second equality with the umbral Siegel modular forms $\Phi^{(\ell)}$ of lambency $\ell = 3, 4, 5$ [29]. Thus we obtain a surprising connection between generalized Mathieu moonshine and umbral moonshine, which clearly deserves further investigation.[†]




Figure 1: Pictorial description of the relation between generalized Mathieu moonshine, involving the twisted twining genera $\phi_{g,h}$, and Mason’s generalized eta-products $\eta_{g,h}$. Starting from $\phi_{g,h}$ one constructs a second-quantized twisted twining genus $\Psi_{g,h}$, whose reciprocal becomes a Siegel modular form $\Phi_{g,h}$ after multiplying by a factor $A_{g,h}$. This is the content of the multiplicative (automorphic) lift indicated in the leftmost arrow. At the level of the Siegel modular form $\Phi_{g,h}$ one then takes the limit $z \rightarrow 0$ which produces the generalized eta-products $\eta_{g,h}(\tau)\eta_{g,h'}(\sigma)$. Here, h' is possibly in a different class from h , as a consequence of the ‘relabeling phenomenon’ described in [17]. See section 4.2 for more details.

Finally, we would like to comment further on the possible physical interpretation of the functions $\Phi_{g,h}$. For $\Phi_{e,e} = \Phi_{10}$, it is of course well-known that the reciprocal $1/\Phi_{10}$ is the generating function of 1/4-BPS dyons in heterotic string theory on T^6 [52], and

[†]We thank Miranda Cheng for a discussion that pointed us in this direction.

in this case the decomposition (1.3) becomes $1/\Phi_{10} \sim \eta(\tau)^{-24}\eta(\rho)^{-24}$, where each factor on the right hand side is identified with the partition function of 1/2 BPS-states. More generally, $1/\Phi_h$ should correspond to the partition function of “twisted dyons”, and the limit (1.3) reflects the wall-crossing phenomenon when a 1/4 BPS-state splits into two 1/2 BPS-states. Similarly we expect that in general $1/\Phi_{g,h}$ should be the generating function of certain “twisted dyons” in CHL orbifolds T^6/\mathbb{Z}_N of heterotic string theory [23, 24, 25, 26]. In particular, this should be true whenever g, h can be interpreted as a pair of commuting symmetries of some K3 surface and the orbifold by g is consistent. Given the results of [8], the first condition amounts to the group $\langle g, h \rangle$ having at least four orbits in the standard 24-dimensional permutation representation of M_{24} , while the second condition simply requires g to belong to some M_{23} subgroup of M_{24} . From the tables in appendix E, it is easy to check that the first condition is satisfied by all 34 groups except the groups numbered 22, 30, 31, 32. As for the remaining groups, the second condition is satisfied by at least one generator g , except for groups 4, 5, 6, 13, 14.

1.3 Outline

Our paper is organized as follows. In section 2 we recall some relevant facts about holomorphic orbifolds, focussing on properties of the associated twisted twining partition functions. We explain the important role played by the third cohomology group $H^3(G, U(1))$, where G is a finite automorphism group of the holomorphic CFT. In particular, this cohomology group controls the modular properties of the twisted twining partition functions. After discussing these properties in a general context we restrict to the relevant case of $G = M_{24}$ for which we recall the main results of [17, 18]. Section 3 introduces some aspects of symmetric products of holomorphic CFT’s and the connection with Hecke operators. In section 3.2 we then introduce the key notion of twisted equivariant Hecke operators that will play a crucial role in what follows. In section 3.3 we use the twisted equivariant Hecke operators to define second-quantized twisted twining genera $\Psi_{g,h}$ and we compute their infinite product expansions. The connection with Mason’s eta-products is then made in section 4 by evaluating $\Psi_{g,h}$ in the limit $z \rightarrow 0$. The modular properties of the second-quantized twisted twining genera are analyzed in section 5.2. Finally, in the concluding section 6 we summarize our results and give some ideas for future research. Various background material, technical calculations and tables have been relegated to the appendices.

2 Twisted-twining partition functions

In this section we introduce and analyze twisted twining partition functions associated with the orbifold of a conformal field theory (CFT) \mathcal{C} by a finite group G of symmetries. When \mathcal{C} is a bosonic CFT, these are modular functions $Z_{g,h}(\tau)$ on the upper-half plane \mathbb{H} , associated with each commuting pair $g, h \in G$. For superconformal field theories the analogous functions are the twisted twining genera $\phi_{g,h}(\tau, z)$, which are Jacobi forms on $\mathbb{H} \times \mathbb{C}$. Our main interest is the case when \mathcal{C} is a putative $\mathcal{N} = (4, 4)$ superconformal field theory on a K3-surface with $G = M_{24}$, but we shall keep the discussion general whenever possible.

2.1 Definition and basic properties

Consider a two-dimensional conformal field theory \mathcal{C} and let G be its symmetry group, i.e. the group of linear automorphisms of its space of states \mathcal{H} that preserves the OPE of the corresponding fields and fixes (at least) the left- and right-moving Virasoro algebra. Given such data, one can construct a new conformal field theory by considering the orbifold of \mathcal{C} by (a subgroup of) G . The first step in the orbifold construction consists in introducing, for each $g \in G$, the g -twisted fields Φ_g , generating a g -twisted space of states \mathcal{H}_g . The defining property of the g -twisted fields is that every field of \mathcal{C} has a non-trivial monodromy g when it is moved along a (sufficiently small) closed path encircling the twisted field Φ_g . The orbifold theory is defined by including all twisted sectors in the spectrum and then restricting to the G -invariant part. The orbifold theory is a consistent CFT provided certain conditions (in particular, the level-matching condition), assuring the locality of the OPE of twisted fields, are satisfied.

Even if the consistency conditions for the orbifold theory are not satisfied, it makes sense to consider the g -twisted sector \mathcal{H}_g as a module over the Virasoro algebra (or more generally, over the G -invariant part of the chiral algebra of \mathcal{C}). Let us consider the case where \mathcal{C} is a holomorphic CFT, so that it coincides with its chiral algebra, which is a self-dual vertex operator algebra. Then, under suitable assumptions, there exists a unique irreducible g -twisted sector \mathcal{H}_g for each $g \in G$ [38]. The action of a generic element $h \in G$ on \mathcal{C} induces a linear map

$$\rho_g(h) : \mathcal{H}_g \rightarrow \mathcal{H}_{h^{-1}gh} , \quad (2.1)$$

from the g -twisted to the $h^{-1}gh$ -twisted sector. In particular, when h commutes with g , it

defines an endomorphism of \mathcal{H}_g , so that \mathcal{H}_g carries a representation ρ_g of the centralizer

$$C_G(g) := \{h \in G : gh = hg\} , \quad (2.2)$$

of g in G . It should be noticed that the representation ρ_g of $C_G(g)$ is in general only projective. This fact will play a crucial role in what follows.

Let us define a g -twisted h -twining partition function on the upper half-plane \mathbb{H}_+ by

$$Z_{g,h}(\tau) = \text{Tr}_{\mathcal{H}_g}(\rho_g(h)q^{L_0 - \frac{c}{24}}) , \quad q := e^{2\pi i\tau}, \quad g, h \in G, \quad gh = hg . \quad (2.3)$$

When g and h are the identity, this reduces to the usual torus partition function for \mathcal{C} . On general physical grounds, one expects the twisted-twining partition function $Z_{g,h}$ to be given by a path-integral on an elliptic curve $E_\tau \cong \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ with modular parameter τ , where the fields have monodromies g and h around the generators $-1, \tau$ of the first homology group $H_1(E_\tau, \mathbb{Z}) \cong \mathbb{Z} + \tau\mathbb{Z}$. Furthermore, each twisted-twining partition function $Z_{g,h}$ is a holomorphic function of the modular parameter τ and is expected to be a modular function under the subgroup $\Gamma_{g,h} \subset SL(2, \mathbb{Z})$, that preserves the monodromies of the fields around the non-trivial cycles of the torus.

Analogous properties are expected for the twisted-twining genera in superconformal field theories with an extended (at least $\mathcal{N} = (2, 2)$) superconformal algebra and a group G of symmetries preserving such an algebra. The twisted-twining genera are defined as traces

$$\phi_{g,h}(\tau, z) = \text{Tr}_{\mathcal{H}_g^{RR}}(hq^{L_0 - \frac{c}{24}}\bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}}(-1)^{F+\bar{F}}y^{J_0}) , \quad q := e^{2\pi i\tau}, \quad y := e^{2\pi iz}, \quad (2.4)$$

over the Ramond-Ramond (RR) g -twisted sector \mathcal{H}_g^{RR} . Here J_0 is the zero mode of a $\mathfrak{u}(1)$ current algebra contained in the left $\mathcal{N} = 2$ superconformal algebra (normalized so that the charges are integral) and $(-1)^{F+\bar{F}}$ is the total fermion number. This trace is computed by a path-integral with the same g - and h -twisted periodicity conditions for both bosonic and fermionic fields, with the insertion of an operator y^{J_0} . Although these conformal field theories are not chiral, the twisted-twining partition functions are holomorphic with respect to both τ and z and are expected to be Jacobi forms of weight 0 and index $\frac{c}{6}$ under the same subgroup $\Gamma_{g,h} \subset SL(2, \mathbb{Z})$ as above.

2.2 Modular properties and the α -twist

We mentioned above that for a holomorphic CFT \mathcal{C} with automorphism group G , the twisted sectors \mathcal{H}_g , carry projective representations ρ_g of the centralizer $C_G(g)$. This

implies that ρ_g satisfies

$$\rho(h)_g \rho_g(h') = c_g(h, h') \rho_g(hh'), \quad h, h' \in C_G(g), \quad (2.5)$$

where $c_g(h, h')$ is a phase, or, more precisely, a 2-cocycle representing a class $[c_g]$ in $H^2(C_G(g), U(1))$. As a consequence, the twisted twining partition function $Z_{g,h}$ is not an honest class function on G , but rather satisfies

$$Z_{g,h}(\tau) = \frac{c_g(h, k)}{c_g(k, k^{-1}hk)} Z_{k^{-1}gk, k^{-1}hk}(\tau), \quad (2.6)$$

which, as we shall see, puts strong cohomological constraints on $Z_{g,h}$. The same formula also applies to the twisted twining elliptic genera $\phi_{g,h}$.

The role of the cohomology group $H^2(C_G(g), U(1))$ can in fact be derived from a more fundamental property of \mathcal{C} , namely that consistent orbifolds of a holomorphic CFT are classified by the third cohomology $H^3(G, U(1))$. Concretely, this implies there exists a $U(1)$ -valued 3-cocycle $\alpha(g, h, k)$, with $(g, h, k) \in G \times G \times G$, representing a class $[\alpha]$ in $H^3(G, U(1))$, which in turn determines the class $[c_g] \in H^2(C_G(g), U(1))$ via the formula [46, 47]

$$c_g(x, y) := \frac{\alpha(g, x, y) \alpha(x, y, (xy)^{-1}g(xy))}{\alpha(x, x^{-1}gx, y)}, \quad g, x, y \in G. \quad (2.7)$$

Thus, α characterizes the projective representations ρ_g . The 3-cocycle α is normalized, i.e. $\alpha(g, h, k) = 1$ whenever at least one of its arguments g, h, k is the identity. The cohomology group $H^3(G, U(1))$ also controls the modular properties of the twisted twining partition functions $Z_{g,h}$, in the sense that under a modular transformation $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ they satisfy

$$\epsilon_{g,h} \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right) Z_{g^d h^{-c}, g^{-b} h^a} \left(\frac{a\tau + b}{c\tau + d} \right) = Z_{g^a h^c, g^b h^d}(\tau), \quad (2.8)$$

where $\epsilon_{g,h} \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} \right)$ is a $U(1)$ -valued multiplier system[‡] for $SL(2, \mathbb{Z})$ that can be explicitly constructed from the 3-cocycle $\alpha(g, h, k)$ (see appendix B). In [18, 17] these properties were used to constrain and determine all of the twisted twining elliptic genera $\phi_{g,h}$ for g, h a commuting pair in M_{24} .

Given the rather complicated modular action in (2.8) it turns out to be very convenient to introduce a certain slash operator which combines the action of $SL(2, \mathbb{Z})$ on $(\tau; g, h)$ with the conjugation by G in (2.6). In addition this must incorporate the multiplier induced by the cocycle twist. To define the slash operator we must first define the following twisted modular action on commuting pairs of elements (g, h) in G :

$$(\gamma, k) : (g, h) \mapsto (kgk^{-1}, khk^{-1})\gamma^{-1} = (kg^d h^{-c} k^{-1}, kg^{-b} h^a k^{-1}). \quad (2.9)$$

[‡]This was called $\chi_{g,h}$ in [18].

Here, $(\gamma, k) = \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, k\right) \in SL(2, \mathbb{Z}) \times G$.

When acting on twisted twining partition functions $Z_{g,h}$ the α -twisted equivariant slash operator is defined as follows

$$Z_{g,h}(\tau)|_{\alpha}(\gamma, k) := \epsilon_{g,h}(\gamma, k) Z_{(\gamma,k) \cdot (g,h)}(\gamma \cdot \tau), \quad (\gamma, k) \in SL(2, \mathbb{Z}) \times G, \quad (2.10)$$

where $\epsilon_{g,h}(\gamma, k) \in U(1)$ is a phase which depends on the choice of 3-cocycle α representing a class $[\alpha] \in H^3(G, U(1))$ (see appendix B.2 for more details). When $k = e$ this phase reduces to the phase $\epsilon_{g,h}(\gamma, e) = \epsilon_{g,h}(\gamma)$ in (2.8), while for $\gamma = 1$ we have that $\epsilon_{g,h}(1, k)$ is identified with the phase in (2.6).[§]

The class $[\alpha] \in H^3(G, U(1))$ is part of the defining data of a holomorphic CFT \mathcal{C} with finite automorphism group G . In general the twisted-twining partition functions $Z_{g,h}(\tau)$ associated with \mathcal{C} are invariant under the α -twisted slash operator

$$Z_{g,h}(\tau)|_{\alpha}(\gamma, k) = Z_{g,h}(\tau). \quad (2.11)$$

Different choices of representative α in the class $[\alpha]$ are related to each other by rescalings of $Z_{g,h}(\tau)$ by overall (g, h) -dependent phases. More precisely, if two normalized 3-cocycles α and α' differ by a 3-coboundary $\partial\beta$, i.e.

$$\alpha(g_1, g_2, g_3) = \alpha'(g_1, g_2, g_3) \frac{\beta(g_1, g_2 g_3) \beta(g_2, g_3)}{\beta(g_1 g_2, g_3) \beta(g_1, g_2)}, \quad (2.12)$$

for some $\beta : G \times G \rightarrow U(1)$, with $\beta(e, g) = \beta(g, e) = 1$ for all $g \in G$, then the corresponding twisted-twining partition functions $Z_{g,h}$ and $Z'_{g,h}$ are related by

$$Z'_{g,h}(\tau, z) = \frac{\beta(g, h)}{\beta(h, g)} Z_{g,h}(\tau, z). \quad (2.13)$$

We also define the α -twisted slash operators on the twisted twining genera $\phi_{g,h}$ by the natural generalization of (2.10) to Jacobi forms of weight 0 and index 1:

$$\phi_{g,h}(\tau, z)|_{\alpha}(\gamma, k) := \epsilon_{g,h}(\gamma, k) e^{-\frac{2\pi i m c z^2}{c\tau + d}} \phi_{k g^d h^{-c} k^{-1}, k g^{-b} h^a k^{-1}}(\gamma \cdot \tau, \frac{z}{c\tau + d}). \quad (2.14)$$

2.3 (Generalized) Mathieu moonshine

The elliptic genus of K3 is defined in physics as a refined partition function of certain conformal field theories with $\mathcal{N} = (4, 4)$ superconformal algebra at central charge $c = 6$, namely non-linear sigma models with target space K3. The only states contributing to this

[§]This phase was called $\xi_{g,h}$ in [18].

partition function are the Ramond-Ramond right-moving ground states, i.e. with $\bar{L}_0 - \frac{\bar{c}}{24} = 0$. Similarly, the properties of the twining genera in the Mathieu moonshine conjecture are the ones expected by ‘twining’ this refined partition function by some symmetry of the model that preserves the $\mathcal{N} = (4, 4)$ algebra. The connection between non-linear sigma models on K3 and Mathieu moonshine is however unclear [8, 11]

It is useful to formulate the Mathieu moonshine conjecture (now a theorem, thanks to [12]) in terms of an abstract representation \mathcal{H} of the (holomorphic) $\mathcal{N} = 4$ superconformal algebra at central charge $c = 6$, which is also a module for the Mathieu group M_{24} . Heuristically, \mathcal{H} can be interpreted as the spectrum of R-R right-moving ground states in a generic K3 model, although the meaning of the M_{24} action is not clear from this point of view. The module \mathcal{H} also admits a \mathbb{Z}_2 -grading by a ‘right-moving fermion number’ $(-1)^{\bar{F}}$, which is preserved both by the $\mathcal{N} = 4$ superconformal algebra and by the action of M_{24} . The precise statement of the conjecture is as follows.

Theorem (Mathieu moonshine). *There exists unitary Ramond representation \mathcal{H} of the $\mathcal{N} = 4$ superconformal algebra at central charge $c = 6$, graded by the ‘right-moving fermion number’ $(-1)^{\bar{F}} \in \mathbb{Z}_2$, that carries a non-trivial action of the Mathieu M_{24} commuting with both the $\mathcal{N} = 4$ algebra and the \mathbb{Z}_2 -grading. Furthermore, for each $g \in M_{24}$, the twining genus*

$$\phi_g(\tau, z) := \text{Tr}_{\mathcal{H}}(g q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) , \quad g \in M_{24} \quad (2.15)$$

is a weak Jacobi form of weight 0 and index 1 (possibly with multiplier) under $\Gamma_0(N)$, where N is the order of g . In particular, ϕ_e is the elliptic genus of K3.

A list of Jacobi forms ϕ_g , $g \in M_{24}$, satisfying the expected modular properties was proposed in [2]–[5] and the existence of the corresponding module \mathcal{H} was proved in [12]. The module \mathcal{H} is not uniquely determined by this description, since one can always adjoin to \mathcal{H} a pair of isomorphic representations of the $\mathcal{N} = 4$ algebra and M_{24} with opposite right-moving fermion number, so that they do not contribute to any twisted twining genera. However, there is a ‘minimal’ module that contains no such pair of representations and this is uniquely determined by the twining genera ϕ_g . Remarkably, in this minimal module, the only $\mathcal{N} = 4$ representations with negative $(-1)^{\bar{F}}$ are the BPS ones. It is believed that the module \mathcal{H} and the Jacobi forms ϕ_g of [2]–[5] are the unique solutions of the Mathieu moonshine conjecture, although this has not been proved yet.

In the same spirit, generalized Mathieu moonshine can be expressed in terms of the existence of twisted modules.

Conjecture (Generalized Mathieu moonshine). *For each $g \in M_{24}$, there exists a \mathbb{Z}_2 -graded unitary Ramond representation \mathcal{H}_g of the $\mathcal{N} = 4$ algebra at central charge $c = 6$,*

that carries a projective representation $\rho_g: C_{M_{24}}(g) \rightarrow GL(\mathcal{H}_g)$ of the centralizer of g in M_{24} , with group law

$$\rho_g(h)\rho_g(k) = c_g(h, k)\rho_g(hk) , \quad h, k \in C_{M_{24}}(g) , \quad (2.16)$$

where $c_g(h, k)$ is defined by (2.7) in terms of a normalized 3-cocycle α representing a primitive class in $H^3(M_{24}, U(1)) \cong \mathbb{Z}_{12}$. The operators $\rho_g(h)$ commute with the $\mathcal{N} = 4$ algebra and with the right-moving fermion number $(-1)^{\bar{F}}$, and we have

$$\rho_g(g) = e^{2\pi i(L_0 - \frac{c}{24})} . \quad (2.17)$$

Furthermore, for each pair of commuting elements $g, h \in M_{24}$, the function

$$\phi_{g,h}(\tau, z) := \text{Tr}_{\mathcal{H}_g}(\rho_g(h) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) , \quad (2.18)$$

satisfies the following properties:

(A) For $g = e$, where e is the identity element of M_{24} , the functions $\phi_{e,h}$ coincide with the twining genera ϕ_h constructed in [2]–[5]. In particular, $\phi_{e,e}$ is the elliptic genus of $K3$.

(B) Elliptic properties:

$$\phi_{g,h}(\tau, z + \ell\tau + \ell') = e^{-2\pi i(\ell^2\tau + 2\ell z)} \phi_{g,h}(\tau, z) \quad \ell, \ell' \in \mathbb{Z} . \quad (2.19)$$

(C) Invariance under the α -twisted slash operator (see eq.(2.14))

$$\phi_{g,h}(\tau, z)|_{\alpha}(\gamma, k) = \phi_{g,h}(\tau, z) , \quad (\gamma, k) \in SL(2, \mathbb{Z}) \times M_{24} . \quad (2.20)$$

In particular, each $\phi_{g,h}$ is an even weak Jacobi form of weight 0 and index 1 under a group $\Gamma_{g,h} \subseteq PSL(2, \mathbb{Z})$, with multiplier $\chi_{g,h}$ depending on α .

These conditions were presented in [17] where all the functions $\phi_{g,h}$ were explicitly found, and strong numerical evidence was given that they decompose into projective characters of $C_{M_{24}}(g)$ with respect to the $\mathcal{N} = 4$ algebra. It was also proven in [17] that there exists a unique cohomology class $[\alpha] \in H^3(M_{24}, U(1))$ for which (A)-(C) can be satisfied and that, for each choice of a normalized representative α , the functions $\phi_{g,h}$ satisfying (A)-(C) are uniquely determined. If two normalized 3-cocycles α and α' differ by a 3-coboundary $\partial\beta$, as in (2.12), then the corresponding twisted-twining genera $\phi_{g,h}$ and $\phi'_{g,h}$ are related as in (2.13)

$$\phi'_{g,h}(\tau, z) = \frac{\beta(g, h)}{\beta(h, g)} \phi_{g,h}(\tau, z) . \quad (2.21)$$

The explicit expressions for $\phi_{g,h}$ derived in [17] are collected in a table in appendix E. Although the existence of the modules \mathcal{H}_g matching (2.18) is not proven yet for g different from the identity, strong evidence in this direction has been given [17]. In the rest of the paper we will assume that the conjecture holds. As in the ordinary moonshine case, one can choose minimal modules \mathcal{H}_g , such that the only states with negative $(-1)^{\bar{F}}$ are contained in irreducible BPS representations of $\mathcal{N} = 4$. In fact, the results of [17] suggest that such states only appear in the untwisted sector $\mathcal{H}_{g=e}$.

2.4 Geometric perspective

Before we close this section we shall offer a more geometric perspective on the twisted twining genera which will be useful later on. In [53], Ganter showed that the natural home for the Norton series $f(g, h; \tau)$ [19] is the equivariant elliptic cohomology developed in [54]. We shall here give a short review of Ganter's perspective, adapted to the case of Jacobi forms on $\mathbb{H}_+ \times \mathbb{C}$, as opposed to modular forms on \mathbb{H}_+ .

Let \mathcal{P} be the set of pairs of commuting elements of M_{24}

$$\mathcal{P} := \{(g, h) \in M_{24} \mid gh = hg\} , \quad (2.22)$$

and $\bar{\mathcal{P}}$ the set of conjugacy classes of such pairs, i.e. the quotient of \mathcal{P} with respect to $(g, h) \sim (k^{-1}gk, k^{-1}hk)$, for any $k \in M_{24}$. The set $\bar{\mathcal{P}}$ can be identified with the set of isomorphism classes of principal M_{24} -bundles over the elliptic curve $E_\tau = \mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$. One can then consider the associated moduli space[¶] of principal M_{24} -bundles on E_τ

$$\mathcal{M} = \mathcal{P} \times (\mathbb{H}_+ \times \mathbb{C}) / M_{24} \times (SL(2, \mathbb{Z}) \ltimes \mathbb{Z}^2) . \quad (2.23)$$

For each pair $(g, h) \in \mathcal{P}$ there exists an $(SL(2, \mathbb{Z}) \ltimes \mathbb{Z}^2)$ -equivariant line bundle

$$\begin{array}{c} \mathcal{L}_{g,h}^\alpha \\ \downarrow \\ \mathcal{M} \end{array} \quad (2.24)$$

which is twisted by the 3-cocycle $\alpha \in H^3(M_{24}, U(1))$ [53]. Thus, for fixed (g, h) we can think of the twisted twining genus $\phi_{g,h}$ as a section of $\mathcal{L}_{g,h}^\alpha$. The twist by α in the bundle $\mathcal{L}_{g,h}^\alpha$ accounts for the α -dependent multiplier phases occurring in the modular transformations of the twisted twining genera. In the cases when α describes a trivial class in $H^3(M_{24}, U(1))$,

[¶]This should really be a moduli *stack* but we ignore this technical point; see [53] for a more precise description.

the bundle $\mathcal{L}_{g,h}^\alpha$ is canonically trivialized and the associated $\phi_{g,h}$ exhibits no multiplier phase. The interpretation of the twisted twining genera as sections of $\mathcal{L}_{g,h}^\alpha$ will in particular play a role in section 3.3 when we discuss twisted equivariant Hecke operators.

3 Second quantization of Mathieu moonshine

In this section we construct the second quantized twisted twining genera $\Psi_{g,h}$ as the exponentiated generating function of certain twisted equivariant Hecke operators acting on $\phi_{g,h}$. These Hecke operators have a natural interpretation in terms of symmetric orbifolds and we therefore begin by reviewing some relevant facts about symmetric orbifold theory. Having defined the functions $\Psi_{g,h}$ we also show that they have infinite product expansions.

3.1 Symmetric Orbifolds and Hecke Operators

3.1.1 Symmetric Orbifolds

Given a holomorphic CFT, or self-dual Vertex Operator Algebra (VOA), \mathcal{C} with a group $G \subseteq \text{Aut}(\mathcal{C})$ of automorphisms, let us consider the theory $\mathcal{C}^{\otimes N}$ obtained by taking the tensor product of N copies of \mathcal{C} , for some $N \geq 1$. The group of automorphisms of $\mathcal{C}^{\otimes N}$ contains the direct product of the symmetric group S_N and the group G acting diagonally on all copies of \mathcal{C} .^{||} The twisted twining partition function $Z_{g,h}^{\mathcal{C}^{\otimes N}}$ of the theory $\mathcal{C}^{\otimes N}$, associated to a pair of commuting elements (g, h) of the diagonal group G , is simply the N :th-power of the twisted twining partition function $Z_{g,h}$ in the original theory \mathcal{C} :

$$Z_{g,h}^{\mathcal{C}^{\otimes N}}(\tau) = Z_{g,h}(\tau)^N. \quad (3.1)$$

In other words, the path integral on the torus with boundary conditions g, h in the product theory $\mathcal{C}^{\otimes N}$ is simply the product of N copies of the path integral in the original theory with the same boundary conditions. In particular, if $Z_{g,h}$ is invariant under the α -twisted slash operator for some 3-cocycle α , then $Z_{g,h}^{\mathcal{C}^{\otimes N}}$ is invariant under the α^N -twisted slash operator.

The N :th symmetric orbifold $S^N \mathcal{C}$ of \mathcal{C} is defined as the orbifold of $\mathcal{C}^{\otimes N}$ by S_N , and its group of automorphisms contains the diagonal G . Therefore, one can define the twisted twining partition functions

$$Z_{g,h}^{(N)}(\tau) \equiv Z_{g,h}^{S^N \mathcal{C}}(\tau) \quad (3.2)$$

^{||}Actually, $\text{Aut}(\mathcal{C}^{\otimes N})$ contains the wreath product $S_N \wr G := S_N \ltimes (G \times \dots \times G)$, i.e. the semidirect product of S_N and the direct product $G^N := G \times \dots \times G$ of N copies of G , with S_N acting on G^N by the obvious permutation.

in each symmetric product. These partition functions can be computed using the usual formulae for orbifold CFTs.

The orbifold formula for $Z_{g,h}^{(N)}$ can be nicely understood in the geometric setting of section 2.4. Recall that the twisted twining partition function $Z_{g,h}(\tau)$ of a VOA \mathcal{C} with $\text{Aut}(\mathcal{C}) = G$ is a section of a (α -twisted) line bundle $\mathcal{L}_{g,h}^\alpha$ over the moduli space \mathcal{M}_G of G -bundles over the elliptic curve $E = \mathbb{C}/\Lambda$ with $\Lambda = \mathbb{Z} + \tau\mathbb{Z}$. In this picture g and h correspond to the monodromies around the cycles -1 and τ of the elliptic curve.

As shown in [49], $Z_{g,h}^{(N)}$ can be expressed as a sum of contributions from all isomorphism classes of unramified N -fold coverings $\Upsilon : E' \rightarrow E$, namely

$$Z_{g,h}^{(N)}(\tau) \sim \sum_{\substack{\text{isomorphism classes of} \\ N\text{-fold coverings} \\ \Upsilon : E' \rightarrow E}} \Upsilon^* Z_{g,h}(\tau) , \quad (3.3)$$

up to a suitable normalization. Here, $\Upsilon^* Z_{g,h}$ is simply the partition function associated with the pull-back $\Upsilon^* \mathcal{L}_{g,h}^\alpha \rightarrow E'$ of the line bundle $\mathcal{L}_{g,h}^\alpha$.

Up to isomorphisms, we can consider coverings that preserve the base-point of the elliptic curve. Any *connected* unramified base-point preserving N -fold cover of $E = \mathbb{C}/\Lambda$ is given by $E' \cong \mathbb{C}/\Lambda' \rightarrow E$, $z \mapsto z$, where Λ' is a sublattice of index N in Λ . In turn, the sublattices of index N are given by $\Lambda' = M\Lambda$ for any M in the set

$$\text{Mat}_N(\mathbb{Z}) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{Z}, ad - bc = N \right\} \quad (3.4)$$

of integral 2×2 matrices M of determinant N . In particular, when $N = 1$, the covers $\Upsilon : E' \rightarrow E$ are actually isomorphisms and are classified by $\text{Mat}_1(\mathbb{Z}) \equiv SL(2, \mathbb{Z})$. Let Υ_γ denote the isomorphism associated with a specific choice of $\gamma \in SL(2, \mathbb{Z})$. The pull-back $\Upsilon_\gamma^* Z_{g,h}$ then coincides with the slash operator

$$\Upsilon_\gamma^* Z_{g,h}(\tau) = Z_{g,h}(\tau)|_\alpha(\gamma, e) , \quad \gamma \in SL(2, \mathbb{Z}) . \quad (3.5)$$

For generic $N > 0$, the isomorphism classes of coverings are in one to one correspondence with the cosets

$$SL(2, \mathbb{Z}) \backslash \text{Mat}_N(\mathbb{Z}) , \quad (3.6)$$

and a set of coset representatives is given by

$$\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} , \quad ad = N, \quad a, d > 0, \quad 0 \leq b < d . \quad (3.7)$$

Therefore, the contribution to the partition function $Z_{g,h}^{(N)}(\tau)$ from the *connected* N -fold coverings is given by

$$T_N Z_{g,h}(\tau) := \frac{1}{N} \sum_{M \in SL_2(\mathbb{Z}) \setminus \text{Mat}_N(\mathbb{Z})} \Upsilon_M^* Z_{g,h}(\tau) \equiv \frac{1}{N} \sum_{M \in SL_2(\mathbb{Z}) \setminus \text{Mat}_N(\mathbb{Z})} Z_{g,h}(\tau)|_\alpha(M, e) , \quad (3.8)$$

where $|_\alpha(M, e)$ is a suitable generalization of the slash operator to $\text{Mat}_N(\mathbb{Z})$, that will be discussed in the following sections. The linear operators T_N defined by (3.8) are (α -twisted) equivariant Hecke operators, that will be studied in more detail in the remainder of this section.

3.1.2 Hecke operators

Hecke operators are linear operators acting on modular forms. Specifically, for each $N \geq 1$ there is an operator T_N that acts on a weight k modular form $f(\tau)$ and produces another modular form $T_N f(\tau)$ of the same weight. On modular functions, i.e. weight $k = 0$ modular forms, the Hecke operator is defined by

$$T_N f(\tau) := \frac{1}{N} \sum_{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \setminus \text{Mat}_N(\mathbb{Z})} f\left(\frac{a\tau + b}{c\tau + d}\right). \quad (3.9)$$

These operators have a natural generalization to Jacobi forms given by [55] (restricting to the weight 0, index m case)

$$T_N \phi(\tau, z) := \frac{1}{N} \sum_{\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \setminus \text{Mat}_N(\mathbb{Z})} e^{\frac{-mNcz^2}{c\tau + d}} \phi\left(\frac{a\tau + b}{c\tau + d}, \frac{Nz}{c\tau + d}\right). \quad (3.10)$$

In fact, on Jacobi forms there exists an additional Hecke operator U_N with the simple action

$$U_N \phi(\tau, z) := \phi(\tau, Nz). \quad (3.11)$$

These operators map Jacobi forms of weight 0 and index m to Jacobi forms of weight 0 and index Nm and N^2m , respectively. They also form a Hecke algebra, with the relations

$$T_m \cdot T_n = T_{mn} , \quad \text{for } \gcd(m, n) = 1 , \quad (3.12)$$

$$U_m \cdot U_n = U_{mn} , \quad (3.13)$$

$$T_m \cdot U_n = U_n \cdot T_m , \quad (3.14)$$

$$T_p \cdot T_{p^m} = T_{p^{m+1}} + \frac{1}{p} T_{p^{m-1}} \cdot U_p , \quad \text{for } p \text{ prime} . \quad (3.15)$$

It is easy to see that a set of representatives for the quotient $SL(2, \mathbb{Z}) \backslash \text{Mat}_N(\mathbb{Z})$ can always be chosen of the form (3.7). This allows us to write the action of T_N on $\phi(\tau, z)$ as follows

$$T_N \phi(\tau, z) = \frac{1}{N} \sum_{a, d > 0, ad=N} \sum_{b=0}^{d-1} \phi\left(\frac{a\tau + b}{d}, az\right). \quad (3.16)$$

3.2 Twisted equivariant Hecke operators

In order to define the exponential lift of our twisted twining genera $\phi_{g,h}$ we must find a set of Hecke operators that are compatible with the properties (A)-(C) in section 2.3. This means that the Hecke action should be compatible, i.e. equivariant, with respect to the M_{24} -action on the pair (g, h) , and it should incorporate the α -twist in the modular transformation. Equivariant versions of Hecke operators acting on modular forms was proposed by Ganter [53] in the context of generalized Monstrous moonshine, and we shall see that this result also applies here after some minor modifications.

3.2.1 Main definition

The standard Hecke-operators T_N map weak Jacobi forms of weight 0, index 1 to weak Jacobi forms of weight 0, index N . In the generalized setting the analogous statement implies that we should have equivariant Hecke operators \mathcal{T}_N that map sections of $\mathcal{L}_{g,h}^\alpha$ to sections of the product bundle $(\mathcal{L}_{g,h}^\alpha)^{\otimes N}$, where $\mathcal{L}_{g,h}^\alpha \rightarrow \mathcal{M}$ is the line bundle discussed in section 2.4. In fact, because of the cocycle-dependent multiplier phases arising in the modular transformations of the $\phi_{g,h}$ we need to introduce a certain α -twisted version \mathcal{T}_N^α of the equivariant Hecke operator. More explicitly, if the sections of $\mathcal{L}_{g,h}^\alpha$ correspond to holomorphic functions $\phi_{g,h}$ on the covering space $\mathcal{P} \times \mathbb{H}_+ \times \mathbb{C}$ of \mathcal{M} , with an automorphy factor determined by a 3-cocycle α , then $\mathcal{T}_N^\alpha \phi_{g,h}$ should correspond to holomorphic functions whose automorphy factor derives from the N :th power α^N of the 3-cocycle.

Since the detailed analysis is rather technical, we will postpone it to appendix C and just quote the explicit formula for the action of the Hecke operators on a twisted-twining character $\phi_{g,h}$

$$\mathcal{T}_N^\alpha \phi_{g,h}(\tau, z) := \frac{1}{N} \sum_{\substack{a, d > 0, \\ ad=N}} \sum_{b=0}^{d-1} \epsilon_{g,h} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \phi_{g^d, g^{-b}h^a} \left(\frac{a\tau + b}{d}, az \right), \quad (3.17)$$

$$\mathcal{U}_N^\alpha \phi_{g,h}(\tau, z) := \epsilon_{g,h} \begin{pmatrix} N & 0 \\ 0 & N \end{pmatrix} \phi_{g^N, h^N}(\tau, Nz), \quad (3.18)$$

where

$$\epsilon_{g,h} \left(\begin{smallmatrix} a & b \\ 0 & d \end{smallmatrix} \right) := \frac{\prod_{i=1}^{a-1} c_g(h, h^i)^d}{\prod_{j=1}^{d-1} c_{g^{-b}h^a}(g, g^j) \prod_{k=1}^b c_g(g, g^{-k}h^a)^d} , \quad (3.19)$$

and $c_g(x, y)$, $g, x, y \in G$, depends on the cocycle α that determines the automorphy factor of $\phi_{g,h}$ via (2.7).

As discussed in appendix C, these Hecke operators satisfy the Hecke algebra (3.12)–(3.15). The explicit expressions for $\mathcal{T}_N^\alpha \phi_{g,h}$, for $N = 1, 2, 3, 4$, in the case of Mathieu moonshine are collected in appendix E.

Notice that the phases $\epsilon_{g,h} \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix} \right)$ are trivial whenever g is the identity or when the restriction of the cocycle α to the group $\langle g, h \rangle$ is trivial. An example where non-trivial phases appear is given by $\mathcal{T}_2^\alpha \phi_{g,g}$, where g is in class $2B$ of M_{24} . These phases can be understood by first considering

$$\mathcal{T}_2^\alpha \phi_{e,g}(\tau, z) = \frac{1}{2} \left(\phi_{e,e}(2\tau, 2z) + \phi_{e,g}\left(\frac{\tau}{2}, z\right) + \phi_{e,g}\left(\frac{\tau+1}{2}, z\right) \right) , \quad (3.20)$$

and then imposing the relations

$$\mathcal{T}_2^\alpha \phi_{g,e}(\tau, z) = e^{\frac{-4\pi i z^2}{\tau}} \mathcal{T}_2^\alpha \phi_{e,g}\left(-\frac{1}{\tau}, \frac{z}{\tau}\right) , \quad \mathcal{T}_2^\alpha \phi_{g,g}(\tau, z) = \mathcal{T}_2^\alpha \phi_{g,e}(\tau+1, z) . \quad (3.21)$$

Using the modular properties of $\phi_{e,g}$, we obtain

$$\mathcal{T}_2^\alpha \phi_{g,e}(\tau, z) = \frac{1}{2} \left(\phi_{g,e}(2\tau, 2z) + \phi_{e,e}\left(\frac{\tau}{2}, z\right) + \phi_{e,g}\left(\frac{\tau+1}{2}, z\right) \right) , \quad (3.22)$$

$$\mathcal{T}_2^\alpha \phi_{g,g}(\tau, z) = \frac{1}{2} \left(-\phi_{g,e}(2\tau, 2z) + \phi_{e,g}\left(\frac{\tau}{2}, z\right) + \phi_{e,e}\left(\frac{\tau+1}{2}, z\right) \right) , \quad (3.23)$$

where the minus sign in front of $\phi_{g,e}(2\tau, 2z)$ in the latter equation is due to the non-trivial multiplier system of the twining genus $\phi_{e,g}$.

3.2.2 Central extensions

Rather than considering the projective representation ρ_g of the centralizer $C_{M_{24}}(g)$, it is often useful to work with a linear representation of a central extension

$$1 \rightarrow U(1) \rightarrow C_{M_{24}}^\alpha(g) \rightarrow C_{M_{24}}(g) \rightarrow 1 . \quad (3.24)$$

The group $C_{M_{24}}^\alpha(g)$ can be constructed explicitly in terms of generators and relations. Let $q(x) := e^{2\pi i x}$, $x \in \mathbb{R}/\mathbb{Z}$, be the generator of the central $U(1)$ factor and consider one generator h_α for each $h \in C_{M_{24}}(g)$. Then, $C_{M_{24}}^\alpha(g)$ is generated by all such elements subject to the relations

$$h_\alpha k_\alpha = q(\mu_g(h, k)) (hk)_\alpha , \quad q(x) h_\alpha = h_\alpha q(x) \quad (3.25)$$

for all $h, k \in C_{M_{24}}(g)$. Here, $\mu_g(h, k) \in \mathbb{R}/\mathbb{Z}$ is defined by

$$e^{2\pi i \mu_g(h, k)} = c_g(h, k) , \quad (3.26)$$

where $c_g(h, k)$ is the 2-cocycle determining the projective representation ρ_g as in (2.5). This definition provides also a canonical lift

$$h \mapsto h_\alpha , \quad (3.27)$$

from $C_{M_{24}}(g)$ to its central extension $C_{M_{24}}^\alpha(g)$. By construction, the g -twisted sector carries a genuine representation $\tilde{\rho}_g$ of $C_{M_{24}}^\alpha(g)$, with

$$\tilde{\rho}_g(q(x)) = e^{2\pi i x} , \quad \tilde{\rho}_g(h_\alpha) = \rho_g(h) , \quad (3.28)$$

for all $h \in C_{M_{24}}(g)$, $x \in \mathbb{R}/\mathbb{Z}$. A less trivial observation is that also the g^r -twisted sectors, for all $r \in \mathbb{Z}_{\geq 0}$, carry a genuine representation $\tilde{\rho}_{g,r}$ of $C_{M_{24}}^\alpha(g)$, defined by**

$$\tilde{\rho}_{g,r}(q(x)) := e^{2\pi i r x} , \quad \tilde{\rho}_{g,r}(h_\alpha) \equiv \tilde{\rho}_{g,r}(h) := \frac{\rho_{g^r}(h)}{f_{g,r}(h)} , \quad (3.29)$$

for all $h \in C_{M_{24}}(g) \subseteq C_{M_{24}}(g^r)$, $x \in \mathbb{R}/\mathbb{Z}$, $r \in \mathbb{Z}_{\geq 0}$, where

$$f_{g,r}(h) = \prod_{i=1}^{r-1} c_h(g, g^i) . \quad (3.30)$$

The fact that $\tilde{\rho}_{g,r}$ are well defined representations of $C_{M_{24}}^\alpha(g)$ is an immediate consequence of the identity

$$c_{g^r}(h, k) = \frac{f_{g,r}(h)f_{g,r}(k)}{f_{g,r}(hk)} c_g(h, k)^r , \quad h, k \in C(g) , \quad (3.31)$$

which follows from the definition of c_g in terms of α and repeated applications of the cocycle condition for α . Notice that $f_{g,1}(h) = 1$ for all $h \in C_{M_{24}}(g)$, so that $\tilde{\rho}_g \equiv \tilde{\rho}_{g,1}$.

If α and α' are different cocycle representatives of the same cohomology class $[\alpha]$, then the central extensions are isomorphic $C_{M_{24}}^\alpha(g) \cong C_{M_{24}}^{\alpha'}(g)$, although the corresponding lifts h_α and $h_{\alpha'}$ are different. In particular, for special choices of the representative cocycle α , it is sufficient to consider a central extension of $C_{M_{24}}(g)$ by a finite subgroup of $U(1)$. With slight abuse of notation, we will denote also these finite central extensions by $C_{M_{24}}^\alpha(g)$. See appendix D.2 for more details on these special choices for α .

**We denote by the same symbol $\tilde{\rho}_{g,r}$ also the projective representation of $C_{M_{24}}(g)$ on \mathcal{H}_{g^r} with $\tilde{\rho}_{g,r}(h) := \rho_{g^r}(h)/f_{g,r}(h)$

3.2.3 Alternative definition

The α -twisted equivariant Hecke operators are more easily defined in terms of the central extension $C_{M_{24}}^\alpha(g)$ described in section 3.2.2. Set

$$\phi_{g_\alpha^r, h_\alpha}(\tau, z) := \text{Tr}_{\mathcal{H}_{g^r}}(\tilde{\rho}_{g,r}(h_\alpha) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) . \quad (3.32)$$

so that

$$\phi_{g_\alpha^d, g_\alpha^{-b} h_\alpha}(\tau, z) = \text{Tr}_{\mathcal{H}_{g^d}}(\tilde{\rho}_{g,d}(g)^{-b} \tilde{\rho}_{g,d}(h)^a q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) \quad (3.33)$$

$$= \frac{\prod_{i=1}^{a-1} c_g(h, h^i)^d}{\prod_{k=1}^b c_g(g, g^{-k} h^a)^d} \text{Tr}_{\mathcal{H}_{g^d}}(\tilde{\rho}_{g,d}(g^{-b} h^a) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) \quad (3.34)$$

$$= \frac{\prod_{i=1}^{a-1} c_g(h, h^i)^d}{\prod_{j=1}^{d-1} c_{g^{-b} h^a}(g, g^j) \prod_{k=1}^b c_g(g, g^{-k} h^a)^d} \text{Tr}_{\mathcal{H}_{g^d}}(\rho_{g^d}(g^{-b} h^a) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) \quad (3.35)$$

$$= \epsilon_{g,h} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \phi_{g^d, g^{-b} h^a}(\tau, z) . \quad (3.36)$$

Therefore, we can reinterpret the α -twisted Hecke operators \mathcal{T}_N^α acting on the twisted-twining genera $\phi_{g,h}$ as (untwisted) equivariant operators acting on the twisted-twining genera $\phi_{g_\alpha, h_\alpha}$

$$\mathcal{T}_L^\alpha \phi_{g,h}(\tau, z) = \frac{1}{L} \sum_{ad=L} \sum_{b=0}^{d-1} \phi_{g_\alpha^d, g_\alpha^{-b} h_\alpha^a} \left(\frac{a\tau + b}{d}, az \right) =: \mathcal{T}_L \phi_{g_\alpha, h_\alpha}(\tau, z) . \quad (3.37)$$

This form of the Hecke operators will turn out to be very useful for some of the calculations in sections 3.3 and 4.

3.3 Second-Quantized Twisted Twining Genera

3.3.1 Definition

For any Calabi-Yau manifold X , one can define its *second-quantized* elliptic genus Ψ_X as the exponentiated generating function of the orbifold elliptic genus $\phi_{S^N X}(\tau, z)$ of the symmetric products $S^N X$ [49]. In fact, ref. [49] gave three equivalent expressions for Ψ_X :

$$\Psi_X(\sigma, \tau, z) = \sum_{N=0}^{\infty} p^N \phi_{S^N X}(\tau, z) = \exp \left[\sum_{L=1}^{\infty} p^L (T_L \phi_X)(\tau, z) \right] = \prod_{\substack{n>0, m \geq 0, \\ \ell \in \mathbb{Z}}} (1 - p^n q^m y^\ell)^{c(mn - \ell^2)}, \quad (3.38)$$

where we have set

$$q = e^{2\pi i \tau}, \quad y = e^{2\pi i z}, \quad p = e^{2\pi i \rho}, \quad (3.39)$$

and $c(mn - \ell^2)$ are the Fourier coefficients of the elliptic genus of X :

$$\phi_X(\tau, z) = \sum_{m \geq 0, \ell \in \mathbb{Z}} c(m, \ell) q^m y^\ell. \quad (3.40)$$

Here, T_N are standard Hecke operators defined in (3.16).

It was shown in [50] that $\Phi \equiv A_X \Psi_X$, where A_X is a simple correction factor (“Hodge anomaly”) that is determined by the Hodge numbers of X , transforms as a Siegel modular form for some congruence subgroup of $Sp(4; \mathbb{Z})$.

Taking the formula (3.38) as a starting point, we now wish to define the *second quantized twisted twining genera* as follows

$$\Psi_{g,h}(\sigma, \tau, z) := \exp \left[\sum_{L=1}^{\infty} p^L (\mathcal{T}_L^\alpha \phi_{g,h})(\tau, z) \right], \quad (3.41)$$

where \mathcal{T}_L^α is the twisted equivariant Hecke operator defined in (3.17). We will later show that after including a correction factor $A_{g,h}$ the functions $\Psi_{g,h}$ transform as Siegel modular forms for some subgroup $\Gamma_{g,h}^{(2)} \subset Sp(4; \mathbb{Z})$ that contains the invariance group $\Gamma_{g,h} \subset SL(2, \mathbb{Z})$ of the twisted twining genera $\phi_{g,h}$.

Notice that the definition (3.41) depends on the choice of a normalized 3-cocycle α . If α and α' differ by a 3-coboundary $\partial\beta$ as in (A.5), the corresponding twisted-twining genera are related as

$$\mathcal{T}_L^{\alpha'} \phi'_{g,h}(\tau, z) = e^{2\pi i \nu_{g,h} L} \mathcal{T}_L^\alpha \phi_{g,h}(\tau, z), \quad (3.42)$$

where $\nu \in \mathbb{R}/\mathbb{Z}$ is defined as

$$e^{2\pi i \nu_{g,h}} = \frac{\beta(g, h)}{\beta(h, g)}, \quad (3.43)$$

so that

$$\Psi'_{g,h}(\sigma, \tau, z) = \exp \left(\sum_{L=1}^{\infty} p^L \mathcal{T}_L^{\alpha'} \phi'_{g,h}(\tau, z) \right) \quad (3.44)$$

$$= \exp \left(\sum_{L=1}^{\infty} (e^{2\pi i \nu_{g,h}} p)^L \mathcal{T}_L^\alpha \phi_{g,h}(\tau, z) \right) = \Psi_{g,h}(\sigma + \nu_{g,h}, \tau, z). \quad (3.45)$$

Therefore, a different choice for the cocycle representative α simply amounts to a redefinition $\sigma \rightarrow \sigma + \nu_{g,h}$ of the variable σ .

3.3.2 Infinite product representation

We shall now derive infinite product representations for the second quantized twisted-twining genera $\Psi_{g,h}$ in terms of the Fourier coefficients of $\phi_{g,h}$.

Let us consider a generic commuting pair of elements $g, h \in M_{24}$, with the cocycle α inducing a possibly non-trivial multiplier for $\phi_{g,h}$. Let $N = o(g)$ be the order of g and λ the length of the shortest cycle of g in the 24-dimensional permutation representation. Then, the lift g_α of g to the central extension $C_{M_{24}}^\alpha(g)$, as defined in section 3.2.2, has order $N\lambda$ (see appendix D.2). For any $h \in C_{M_{24}}(g)$ the associated twisted-twining genus $\phi_{g,h}$ has a Fourier expansion of the form

$$\phi_{g,h}(\tau, z) = \sum_{n=0}^{\infty} \sum_{\ell \in \mathbb{Z}} c_{g,h}\left(\frac{n}{N\lambda}, \ell\right) q^{\frac{n}{N\lambda}} y^\ell, \quad (3.46)$$

where, in particular, $c_{g,h}\left(\frac{n}{N\lambda}, \ell\right) = 0$ unless $n \equiv -1 \pmod{\lambda}$. Let $M = o(h_\alpha)$ be the order of the lift h_α of h to the central extension $C_{M_{24}}^\alpha(g)$.^{††} The logarithm of the second quantized twisted-twining genus of $\phi_{g,h}$ is

$$\log \Psi_{g,h}(\sigma, \tau, z) = \sum_{L=1}^{\infty} p^L \mathcal{T}_L^\alpha \phi_{g,h}(\tau, z) \quad (3.47)$$

$$= \sum_{a,d=1}^{\infty} \frac{1}{ad} \sum_{b=0}^{d-1} \sum_{\ell \in \mathbb{Z}} \sum_{n=0}^{\infty} \epsilon_{g,h} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} c_{g^d, g^{-b}h^a}\left(\frac{n}{N\lambda}, \ell\right) e^{\frac{2\pi i b n}{N\lambda d}} q^{\frac{an}{N\lambda d}} y^{a\ell} p^{ad} \quad (3.48)$$

$$= \sum_{a,d=1}^{\infty} \frac{1}{ad} \sum_{\ell \in \mathbb{Z}} \sum_{n=0}^{\infty} \sum_{b=0}^{d-1} c_{g_\alpha^d, g_\alpha^{-b}h_\alpha^a}\left(\frac{n}{N\lambda}, \ell\right) e^{\frac{2\pi i b n}{N\lambda d}} q^{\frac{an}{N\lambda d}} y^{a\ell} p^{ad} \quad (3.49)$$

$$= \sum_{a,d=1}^{\infty} \frac{1}{a} \sum_{t,k=0}^{M-1} \frac{e^{\frac{2\pi i t(a-k)}{M}}}{M} \sum_{\ell \in \mathbb{Z}} \sum_{n=0}^{\infty} \frac{1}{d} \sum_{b=0}^{d-1} c_{g_\alpha^d, g_\alpha^{-b}h_\alpha^k}\left(\frac{n}{N\lambda}, \ell\right) e^{\frac{2\pi i b n}{N\lambda d}} q^{\frac{an}{N\lambda d}} y^{a\ell} p^{ad} \quad (3.50)$$

$$= \sum_{a,d=1}^{\infty} \frac{1}{a} \sum_{t,k=0}^{M-1} \frac{e^{\frac{2\pi i t(a-k)}{M}}}{M} \sum_{\ell \in \mathbb{Z}} F_{g,h}(a, d, k, \ell) y^{a\ell} p^{ad}, \quad (3.51)$$

where

$$F_{g,h}(a, d, k, \ell) := \sum_{n=0}^{\infty} \frac{1}{d} \sum_{b=0}^{d-1} e^{\frac{2\pi i b n}{N\lambda d}} \epsilon_{g,h} \begin{pmatrix} k & b \\ 0 & d \end{pmatrix} c_{g^d, g^{-b}h^k}\left(\frac{n}{N\lambda}, \ell\right) q^{\frac{an}{N\lambda}}. \quad (3.52)$$

In appendix D.1 we show that this sum can be rewritten as follows

$$F_{g,h}(a, d, k, \ell) = \sum_{m=0}^{\infty} \frac{1}{N\lambda} \sum_{b=0}^{\lambda N-1} e^{\frac{2\pi i b m}{N\lambda}} \epsilon_{g,h} \begin{pmatrix} k & b \\ 0 & d \end{pmatrix} c_{g^d, g^{-b}h^k}\left(\frac{md}{N\lambda}, \ell\right) q^{\frac{am}{N\lambda}}. \quad (3.53)$$

^{††}We assume that α is chosen in such a way that the order is finite.

By plugging this expression into (3.51) we obtain

$$\log \Psi_{g,h}(\sigma, \tau, z) = \sum_{d=1}^{\infty} \sum_{m=0}^{\infty} \sum_{\ell \in \mathbb{Z}} \sum_{t=0}^{M-1} \hat{c}_{g,h}(d, m, \ell, t) \sum_{a=1}^{\infty} \frac{1}{a} \left(e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell} p^d \right)^a \quad (3.54)$$

$$= - \sum_{d=1}^{\infty} \sum_{m=0}^{\infty} \sum_{\ell \in \mathbb{Z}} \sum_{t=0}^{M-1} \hat{c}_{g,h}(d, m, \ell, t) \log(1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell} p^d), \quad (3.55)$$

where

$$\hat{c}_{g,h}(d, m, \ell, t) := \sum_{k=0}^{M-1} \sum_{b=0}^{\lambda N-1} \frac{e^{-\frac{2\pi i t k}{M}} e^{\frac{2\pi i b m}{\lambda N}}}{M} \epsilon_{g,h} \left(\begin{smallmatrix} k & b \\ 0 & d \end{smallmatrix} \right) c_{g^d, g^{-b} h^k} \left(\frac{md}{N\lambda}, \ell \right) \quad (3.56)$$

$$= \sum_{k=0}^{M-1} \sum_{b=0}^{\lambda N-1} \frac{e^{-\frac{2\pi i t k}{M}} e^{\frac{2\pi i b m}{\lambda N}}}{M} \text{Tr}_{\mathcal{H}_{g^d}(\frac{md}{N\lambda}, \ell)} (\tilde{\rho}_{g,d}(g)^{-b} \tilde{\rho}_{g,d}(h)^k (-1)^{F+\bar{F}}) . \quad (3.57)$$

Thus, for the inverse of the second quantized twisted twining genus we obtain the infinite product expression

$$\frac{1}{\Psi_{g,h}(\sigma, \tau, z)} = \exp \left(- \sum_{L=1}^{\infty} p^L \mathcal{T}_L^{\alpha} \phi_{g,h}(\tau, z) \right) = \prod_{d=1}^{\infty} \prod_{m=0}^{\infty} \prod_{\ell \in \mathbb{Z}} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell} p^d)^{\hat{c}_{g,h}(d, m, \ell, t)} . \quad (3.58)$$

Note that (3.57) makes sense also for $d = 0$, so that the infinite product in d and m can be symmetrized to obtain

$$\Phi_{g,h}(\sigma, \tau, z) := p q^{\frac{1}{N\lambda}} y \prod_{(d, m, \ell) > 0} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell} p^d)^{\hat{c}_{g,h}(d, m, \ell, t)}, \quad (3.59)$$

where the first product runs over

$$d, m \in \mathbb{Z}_{\geq 0} \quad \text{and} \quad \begin{cases} \ell \in \mathbb{Z}, \ell < 0, & \text{if } m = 0 = d, \\ \ell \in \mathbb{Z}, & \text{otherwise.} \end{cases} \quad (3.60)$$

One can rewrite $\Phi_{g,h}$ in terms of the original $\Psi_{g,h}$ as

$$\Phi_{g,h}(\sigma, \tau, z) = \frac{p \psi_{g,h}(\tau, z)}{\Psi_{g,h}(\sigma, \tau, z)}, \quad (3.61)$$

where $\psi_{g,h}$ includes the $d = 0$ factors in (3.59) and takes the form

$$\psi_{g,h}(\tau, z) = q^{\frac{1}{N\lambda}} y \prod_{t=0}^{M-1} \left(\prod_{\ell < 0} (1 - e^{\frac{2\pi i t}{M}} y^{\ell})^{\hat{c}_{g,h}(0, 0, \ell, t)} \right) \left(\prod_{\ell \in \mathbb{Z}} \prod_{m=1}^{\infty} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell})^{\hat{c}_{g,h}(0, m, \ell, t)} \right) . \quad (3.62)$$

The numerator in (3.61) is the analogue of the “Hodge anomaly” in [50].

Infinite products of the form (3.59) were studied in [36], where it was shown that they converge for $(\begin{smallmatrix} \sigma & z \\ z & \tau \end{smallmatrix})$ in a suitable domain of the Siegel upper half-space of 2×2 complex symmetric matrices with positive definite imaginary part

$$\mathbb{H}_2 := \left\{ Z \in \text{Mat}_2(\mathbb{C}) \mid Z = Z^t, \text{Im} Z > 0 \right\}. \quad (3.63)$$

Furthermore, these products can be analytically continued to meromorphic functions on the whole \mathbb{H}_2 , with zeroes and (possibly) poles along the rational quadratic divisor [56].

The prefactor $pq^{\frac{1}{N\lambda}}y$ in (3.59) has been chosen in such a way that $\Phi_{g,h}$ are Siegel modular forms under certain discrete subgroups of $Sp(4, \mathbb{R})$. The automorphic properties of $\Phi_{g,h}$ will be discussed in section 5.2.

3.3.3 Multiplicative versus additive lift

The construction of a Siegel modular form $\Phi(\sigma, \tau, z)$ from a (weak) Jacobi form $\phi(\tau, z)$ via an infinite product representation, as exemplified by (3.59), is generally referred to as a *multiplicative (automorphic) lift*. The Jacobi form ϕ is said to be the *multiplicative seed* of the lift (see, for instance, [56, 50, 51, 57]), and we write

$$\Phi = \text{Mult}[\phi]. \quad (3.64)$$

In some cases one can also obtain the Siegel modular form using a different procedure, known as the *additive lift*^{††}. In this case the Siegel modular form is constructed as a certain generating function (without exponentiation) of Hecke operators acting on a different Jacobi form ψ , which is then called the *additive seed*. For $\psi(\tau, z)$ a Jacobi form of weight k we write $\tilde{\psi}(\sigma, \tau, z) = p\psi(\tau, z)$ and define the additive lift as

$$\Phi = \text{Add}[\psi] := \sum_{m \geq 1} m^{2-k} (T_m^- \tilde{\psi})(\sigma, \tau, z), \quad (3.65)$$

where T_m^- is a certain Hecke operator; see, e.g. [59] for the precise definition and properties of the right hand side.

As an example, consider the case of the Igusa cusp form Φ_{10} (corresponding to $\Phi_{e,e}$). The multiplicative seed is the K3 elliptic genus $\phi_{e,e} = \phi_{0,1}$ which is the unique weak Jacobi form of weight 0 and index 1. The multiplicative lift yields an infinite product formula for Φ_{10} [58]:

$$\Phi_{10} = \text{Mult}[\phi_{0,1}] = pqy \prod_{(d,m,\ell) > 0} (1 - q^m y^\ell p^d)^{c(d,m,\ell)} \quad (3.66)$$

^{††}This generalizes the *Saito-Kurokawa-Maass lift* defined in [55].

where $c(d, m, \ell)$ are the Fourier coefficients of $\phi_{0,1}$. This is indeed obtained from (3.59) by restricting to $(g, h) = (e, e)$. As explained in [51] one can also obtain this Siegel modular form via an additive lift from the seed $\phi_{10,1}$ (the unique weak Jacobi form of weight 10 and index 1):

$$\Phi_{10} = \text{Add}[\phi_{10,1}]. \quad (3.67)$$

We now observe that the additive seed $\phi_{10,1}$ can be expressed as $\vartheta(\tau, z)^2 \eta(\tau)^{18}$ which is precisely the symmetrization factor $\psi_{g,h}$ in (3.62) when restricting to $(g, h) = (e, e)$. This is in fact a general feature that holds whenever a Siegel modular form can be realized both as an additive lift as well as a multiplicative lift. It is a consequence of the fact that the additive seed ψ is the first Fourier-Jacobi coefficient in the expansion of Φ and it is precisely this coefficient which appears as the prefactor in the symmetrization of the infinite product. For purposes we thus expect that for all (g, h) for which an additive lift exists we should have the following equalities

$$\Phi_{g,h} = \text{Mult}[\phi_{g,h}] = \text{Add}[\psi_{g,h}]. \quad (3.68)$$

In section 5.3 we indeed verify this for a number of examples. For $(g, h) = (e, h)$ modified versions of these additive lifts have also been considered in [60].

4 Wall-crossing and Mason's generalized moonshine for M_{24}

Already in 1990, Mason proposed an M_{24} version of Norton's generalized moonshine conjecture [31, 32]. In this section, we will establish the connection between Mason's generalized moonshine for M_{24} and the recent Mathieu moonshine involving the $K3$ elliptic genus. This involves taking the multiplicative lift $\phi_{g,h} \rightarrow \Phi_{g,h}$, defined in section 3, after which the limit $z \rightarrow 0$ reproduces the generalized eta-products $\eta_{g,h}$ constructed by Mason [32]. Thus, the exponential lift $\Phi_{g,h}$ links Mason's generalized M_{24} -moonshine to the $M_{24}/K3$ -moonshine considered here. This connection was first suggested in [2] for the special case of the twining genera $\phi_{1,h} = \phi_h$. A pictorial overview of this relation is given in figure 1.2. We begin by reviewing Mason's construction of the generalized eta-products $\eta_{g,h}$ and explain their interpretation in terms of partition functions of twisted chiral bosons. Physically, the Siegel modular forms $\Phi_{g,h}$ have interpretations as generating functions of twisted dyons in (CHL) orbifolds of $\mathcal{N} = 4$ string theory, and the eta-products emerge as a result of a wall-crossing formula, generalizing the one written down in [2] for $\Phi_{e,h}$.

4.1 Mason's generalized M_{24} -moonshine

For each commuting pair $(g, h) \in M_{24}$, Mason associates a modular function on the upper half-plane satisfying the requirements posed by Norton in [19]. The starting point of Mason was the 24-dimensional permutation representation of M_{24} in which each element can be associated to a *cycle shape*, which describes the element as a product of cycles of permutations. For instance, in this representation the identity element is represented by the cycle shape 1^{24} , corresponding to the product of 24 identity permutations, while the elements in class $2A$ are represented by $1^8 2^8$, corresponding to the product of 8 identity permutations followed by 8 consecutive order 2 permutations.

Mason's functions were all given in terms of so called *eta-products*, namely products of Dedekind eta-functions. Suppose an order M element $h \in M_{24}$ has cycle shape

$$\prod_{\ell|M} \ell^{i(\ell)} , \quad (4.1)$$

on the 24-dimensional representation of M_{24} , for some integers $i(\ell)$. Then one can associate an eta-product to h by

$$\eta_h(\tau) := \prod_{\ell|M} \eta(\ell\tau)^{i(\ell)} . \quad (4.2)$$

Mason found a generalization of such eta-products associated to “generalized cycle shapes” labelled by commuting pairs (g, h) in M_{24} . We shall denote these generalized moonshine functions by $\eta_{g,h}(\tau)$.^{§§} Let $g, h \in M_{24}$ be a pair of commuting elements of order $o(g) = N$ and $o(h) = M$ and consider their action in the standard 24-dimensional representation V of M_{24} . Let v_1, \dots, v_{24} be a basis of simultaneous eigenvectors for g and h , relative to the eigenvalues $(e^{\frac{2\pi i r_i}{N}}, e^{\frac{2\pi i t_i}{M}})$, $i = 1, \dots, 24$, with $1 \leq r_i \leq N$ and $1 \leq t_i \leq M$. Then, the eta-product $\eta_{g,h}$ is defined as

$$\eta_{g,h}(\tau) = q^{\frac{1}{N\lambda}} \prod_{i=1}^{24} \prod_{n=0}^{\infty} (1 - e^{\frac{2\pi i t_i}{M}} q^{\frac{r_i}{N} + n}) , \quad (4.3)$$

where λ is the length of the shortest cycle of g . In particular, the eta-products η_h of equation (4.2) correspond to $(g, h) = (e, h)$. The products $\eta_{g,h}$ (or rather their inverse) can be interpreted as h -twining partition functions for 24 g -twisted chiral free bosons.

As proven in [31], the eta products $\eta_{g,h}$ satisfy the modular transformations

$$v_{g,h} \left(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix} \right) (c\tau + d)^{-w} \eta_{g^d h^{-c}, g^{-b} h^a} \left(\frac{a\tau + b}{c\tau + d} \right) = \eta_{g,h}(\tau) , \quad (4.4)$$

^{§§}In [32] these functions were denoted by $f(g, h; \tau)$.

for some $v_{g,h} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in U(1)$. The full list of Mason's generalized eta-products and the corresponding weights w can be found in Table 1. Furthermore, by definition, $\eta_{g,h}$ are invariant under conjugation in M_{24}

$$\eta_{kgk^{-1},khk^{-1}}(\tau) = \eta_{g,h}(\tau) . \quad (4.5)$$

4.2 Connecting the two moonshines via wall-crossing

The connection between the $K3/M_{24}$ -moonshine for twining genera $\phi_h(\tau, z)$ and the M_{24} -moonshine associated with the functions $\eta_h(\tau)$ was pointed out in [2]. We now want to use our results above to establish this link also between the two different *generalized* moonshines for M_{24} .

As already mentioned, infinite products of the form (3.59) have been considered by Borcherds [56], where it was shown that they converge in some region of the Siegel upper half-space \mathbb{H}_2 and they can be analytically continued to meromorphic functions on the whole \mathbb{H}_2 . Furthermore, the zeroes and poles of $\Phi_{g,h}$ can only be located at rational quadratic divisors [36, 56, 61], namely

$$H_{d,m,\ell,v} = \left\{ \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} \mid \frac{m}{N\lambda}\tau + \ell z + d\sigma + \frac{v}{M} = 0 \right\} , \quad (4.6)$$

where $m, d, v, \ell \in \mathbb{Z}$ are coprime integers ($\gcd(m, d, v, \ell) = 1$) that satisfy a positive discriminant condition

$$\ell^2 - \frac{4md}{N\lambda} > 0 . \quad (4.7)$$

This generalizes the rational quadratic divisor found in [24] in the context of twisted dyon counting in CHL-models.

The multiplicity of the zero or pole of $\Phi_{g,h}$ at the divisor $H_{m,d,\ell,v}$ is given by $\hat{c}_{g,h}(d, m, \ell, v)$. In particular, $\Phi_{g,h}$ is holomorphic if and only if $\hat{c}_{g,h}(d, m, \ell, v)$ is non-negative at every rational quadratic divisor. As will be shown in section 5, $\Phi_{g,h}$ is an automorphic form for a discrete subgroup $\Gamma_{g,h}^{(2)} \subset Sp(4, \mathbb{R})$. Therefore, one only needs to consider the distinct orbits of rational quadratic divisors under the action of $\Gamma_{g,h}^{(2)}$ to determine the full divisor of $\Phi_{g,h}$.

A special role is played by (the modular orbit of) the divisor $H_{0,0,-1,0}$, corresponding to the locus $z = 0$ in \mathbb{H}_2 . Since $c_{e,g}(0, -1) = 2$ for all $g \in M_{24}$, we have

$$\hat{c}_{g,h}(0, 0, -1, 0) = \frac{1}{MN\lambda} \sum_{k=0}^{M-1} \sum_{b=0}^{\lambda N-1} c_{e,g^{-b}h^k}(0, -1) = 2 . \quad (4.8)$$

It follows that every $\Phi_{g,h}$ has a double zero at this divisor. Using the relation

$$\sum_{\ell \in \mathbb{Z}} c_{g,h}(r, \ell) = 0, \quad \text{for } r > 0, \quad (4.9)$$

together with the relations

$$\text{Tr}_{\mathcal{H}_e(0, \pm 1)}(g^{-b} h^k (-1)^{F+\bar{F}}) = 2, \quad (4.10)$$

$$\text{Tr}_{\mathcal{H}_e(0, \pm \ell)}(g^{-b} h^k (-1)^{F+\bar{F}}) = 0 \quad \text{for } \ell > 1, \quad (4.11)$$

$$\sum_{\ell \in \mathbb{Z}} \text{Tr}_{\mathcal{H}_e(0, \ell)}(g^{-b} h^k (-1)^{F+\bar{F}}) = \text{Tr}_{\mathbf{24}}(g^{-b} h^k), \quad (4.12)$$

we obtain

$$\begin{aligned} \lim_{z \rightarrow 0} \frac{\Phi_{g,h}(\sigma, \tau, z)}{(2\pi i z)^2} &= \lim_{z \rightarrow 0} \frac{y(1-y^{-1})^2}{(2\pi i z)^2} \left(q^{\frac{1}{N\lambda}} \prod_{t=1}^M \prod_{m=1}^{\infty} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}})^{\sum_{\ell \in \mathbb{Z}} \hat{c}_{g,h}(m, 0, \ell, t)} \right) \\ &\quad \times \left(p \prod_{t=1}^M \prod_{d=1}^{\infty} (1 - e^{\frac{2\pi i t}{M}} p^d)^{\sum_{\ell \in \mathbb{Z}} \hat{c}_{g,h}(0, d, \ell, t)} \right). \end{aligned} \quad (4.13)$$

Next we use the relations (4.10)-(4.12) to find an expression for $\psi_{g,h}$ in (3.62) in terms of $\eta_{g,h}$ and known modular objects. Using the aforementioned equations we obtain

$$\begin{aligned} \psi_{g,h}(\tau, z) &= q^{\frac{1}{N\lambda}} y \prod_{t=0}^{M-1} \left((1 - e^{\frac{2\pi i t}{M}} y^{-1})^{\hat{c}_{g,h}(0, 0, -1, t)} \prod_{m=1}^{\infty} \prod_{\ell \in \mathbb{Z}} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}} y^{\ell})^{\hat{c}_{g,h}(0, m, \ell, t)} \right) \\ &= q^{\frac{1}{N\lambda}} y (1 - y^{-1})^2 \prod_{n=1}^{\infty} \frac{(1 - q^n y)^2 (1 - q^n y^{-1})^2}{(1 - q^n)^4} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}})^{\sum_{\ell \in \mathbb{Z}} \hat{c}_{g,h}(0, m, \ell, t)}. \end{aligned} \quad (4.14)$$

To proceed we note the identity

$$\begin{aligned} &q^{\frac{1}{N\lambda}} \prod_{t=0}^{M-1} \prod_{m=1}^{\infty} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}})^{\sum_{k=0}^{M-1} \sum_{b=0}^{N\lambda-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} e^{\frac{2\pi i b m}{N\lambda}} \text{Tr}_{\mathbf{24}}(g^{-b} h^k)} \\ &= q^{\frac{1}{N\lambda}} \prod_{t=0}^{M-1} \prod_{m=1}^{\infty} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N\lambda}})^{\sum_{i=1}^{24} \frac{1}{M} \sum_{k=0}^{M-1} \frac{1}{N\lambda} \sum_{b=0}^{N\lambda-1} e^{-\frac{2\pi i (t-t_i) k}{M}} e^{\frac{2\pi i b (m-\lambda r_i)}{N\lambda}}} \\ &= q^{\frac{1}{N\lambda}} \prod_{i=1}^{24} \prod_{n=0}^{\infty} (1 - e^{\frac{2\pi i t_i}{M}} q^{n+\frac{r_i}{N}}) = \eta_{g,h}(\tau), \end{aligned} \quad (4.15)$$

where $1 \leq r_i \leq N$ and $1 \leq t_i \leq M$ are such that $(e^{\frac{2\pi i r_i}{N}}, e^{\frac{2\pi i t_i}{M}})$, $i = 1, \dots, 24$, are the (g, h) -eigenvalues of a basis of simultaneous eigenvectors for g and h in the 24-dimensional

representation of M_{24} . Using this relation in (4.14) we obtain

$$\psi_{g,h}(\tau, z) = -\frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^6} \eta_{g,h}(\tau) = \phi_{-2,1}(\tau, z) \eta_{g,h}(\tau) , \quad (4.16)$$

where $\phi_{-2,1}$ is the standard weak Jacobi forms of weight -2 and index 1 for $SL(2, \mathbb{Z})$.

To complete the analysis of the limit in (4.13) we note that

$$\lim_{z \rightarrow 0} \frac{y(1-y^{-1})^2}{(2\pi iz)^2} \left(q^{\frac{1}{N\lambda}} \prod_{t=1}^M \prod_{m=1}^{\infty} (1 - e^{\frac{2\pi it}{M}} q^{\frac{m}{N\lambda}})^{\sum_{\ell \in \mathbb{Z}} \hat{c}_{g,h}(m, 0, \ell, t)} \right) = \lim_{z \rightarrow 0} \frac{\psi_{g,h}(\tau, z)}{(2\pi iz)^2} = \eta_{g,h}(\tau) . \quad (4.17)$$

For the pair $(g, h) = (e, e)$, the function $\Phi_{e,e}(\sigma, \tau, z)$ is invariant under the exchange $\sigma \leftrightarrow \tau$. From a physical viewpoint, this is a consequence of S-duality of type II superstring compactified on $K3 \times T^2$. In section 5.2, we will prove that $\Phi_{g,h}$ satisfy analogous transformations

$$\Phi_{g,h}(\sigma, \tau, z) = \Phi_{g,h'}\left(\frac{\tau}{N\lambda}, N\lambda\sigma, z\right) , \quad (4.18)$$

where $h' \in C_{M_{24}}(g)$ is not necessarily in the same conjugacy class as h . Using this identity, we conclude that

$$\lim_{z \rightarrow 0} \frac{\Phi_{g,h}(\sigma, \tau, z)}{(2\pi iz)^2} = \eta_{g,h}(\tau) \eta_{g,h'}(N\lambda\sigma) . \quad (4.19)$$

This equation has a physical interpretation as a wall-crossing formula. Indeed, $\Phi_{g,h}^{-1}$ corresponds to the generating function for the h -twisted degeneracies of $1/4$ BPS states in a CHL model. More precisely, these degeneracies are the Fourier coefficients of the automorphic form $\Phi_{g,h}^{-1}$ and the region where the Fourier expansion is performed depends on the moduli. These multiplicities jump as one crosses the pole at $z = 0$; the physical interpretation is that some $1/4$ BPS dyon corresponding to a bound state of $1/2$ BPS configurations becomes unstable in a certain region of the moduli space and thus disappears from the spectrum. Indeed, the mismatch $\eta_{g,h}(\tau)^{-1} \eta_{g,h'}(N\lambda\sigma)^{-1}$ between the Fourier coefficients at the two sides of the pole represents the degeneracy of a bound state of two $1/2$ BPS states.

5 Automorphic Properties

In this final section we analyze the modular transformation properties of $\Phi_{g,h}(\sigma, \tau, z)$ with respect to discrete subgroups of $Sp(4, \mathbb{R})$. We show that they are Siegel modular forms and in some cases we are able to identify them with previously known objects. Our proof of modularity is done in two steps. The crucial first step is to determine the transformation properties of $\Phi_{g,h}(\sigma, \tau, z)$ with respect to the interchange $\sigma \leftrightarrow \tau$. We refer to this as ‘‘S-duality’’ since in the cases when $\Phi_{g,h}$ has an interpretation as a partition function in an

$\mathcal{N} = 4$ string theory this corresponds precisely to the S-duality that exchanges electric and magnetic charges (see, e.g., [52, 62, 63, 23, 24]). It turns out that this transformation involves the subtle concept of “relabeling” of the group elements (g, h) introduced in [17]. By a careful analysis of this relabeling phenomenon we establish the S-duality symmetry in section 5.1. To complete the analysis of the automorphic properties of $\Phi_{g,h}(\sigma, \tau, z)$ we must also verify the transformation properties with respect to the remaining generators of the relevant modular subgroups of $Sp(4, \mathbb{R})$. This is rather straightforward since it essentially follows from the modularity of the seed functions $\phi_{g,h}(\tau, z)$ and $\psi_{g,h}(\tau, z)$. We present this analysis in section 5.2, and in section 5.3 we also investigate some examples in detail.

5.1 Relabeling and S-duality

In this section, we will show that the functions $\Phi_{g,h}(\sigma, \tau, z)$ defined by the infinite product (3.59) satisfy some ‘S-duality’ identities that exchange σ and τ . This property, together with the modular properties of the twisted-twining genera $\phi_{g,h}$ proved in [17], will be sufficient to prove that all $\Phi_{g,h}$ are automorphic functions under some subgroup of $Sp(4, \mathbb{R})$ (see section 5.2).

In the case where g belongs to some M_{23} subgroup of M_{24} , the main step in the derivation of this S-duality for $\Phi_{g,h}$ is the relabeling phenomenon [17]. This can be understood by considering the example of a holomorphic vertex operator algebra \mathcal{C} with automorphism group G . Given an element $g \in G$, the orbifold of \mathcal{C} by $\langle g \rangle$ is constructed by introducing the twisted representations of \mathcal{C} and then restricting to the g -invariant sector. All twisted sectors carry a representation of some central extension of the centralizer $C_G(g)$ of g in G . This induces an action of $C_G(g)$ by automorphisms on the orbifold theory.

Now suppose that this orbifold defines a consistent VOA isomorphic to the original \mathcal{C} . Thus, the automorphism group G' of the orbifold theory is isomorphic to G , so that the central extension of $C_G(g)$ (or rather of its quotient group acting faithfully on the orbifold) must be a subgroup of $G' \cong G$. It might happen that some automorphism $h \in C_G(g)$ of the original theory and the corresponding automorphism induced on the orbifold theory belong to different conjugacy classes of G and G' . In particular, the characters of h in the original theory and of the induced automorphism in the orbifold theory might be different. In this case, we say that the class of h has been relabeled.

Roughly speaking, a similar phenomenon occurs for the Mathieu moonshine. In particular, we will see that S-duality relates a function $\Phi_{g,h}$ to a function $\Phi_{g,h'}$, where h' belongs to the ‘relabelled’ class.

5.1.1 Orbifolds and relabeling

Given a conformal field theory with a symmetry g , one can construct an orbifold CFT by introducing the g^r -twisted sectors and then restricting to the g -invariant states. The space obtained in this way is the spectrum of a consistent CFT, provided that the level-matching condition is satisfied.

An analogous construction can be considered for the abstract modules of the generalized Mathieu moonshine with respect to some $g \in M_{24}$ of order N , by considering the g -invariant subspace of the direct sum of all twisted sectors

$$\mathcal{H}' = \oplus_{r=0}^{N-1} \left(\mathcal{H}_{g^r}^{\langle \tilde{\rho}_{g,r}(g) \rangle} \right). \quad (5.1)$$

The level-matching condition is equivalent to the requirement that the restriction of the cohomology class $[\alpha]$ is trivial in $H^3(\langle g \rangle, U(1))$, so we will consider only this case. If \mathcal{H} can be interpreted as the spectrum of R-R right-moving ground states in a non-linear sigma model on K3 with a symmetry g , then \mathcal{H}' corresponds to the spectrum of right-moving ground states in the g -orbifold theory. The orbifold is a consistent $\mathcal{N} = (4, 4)$ SCFT with central charge 6, which turns out to be again a non-linear sigma model on K3, since its elliptic genus is

$$\text{Tr}_{\mathcal{H}'} \left(q^{L_0 - \frac{c}{24}} \bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}} y^{J_0^3} (-1)^{F+\bar{F}} \right) = \sum_{r,s=0}^{N-1} \frac{1}{N} \text{Tr}_{\mathcal{H}_{g^r}} \left(\tilde{\rho}_{g,r}(g)^s q^{L_0 - \frac{c}{24}} \bar{q}^{\bar{L}_0 - \frac{\bar{c}}{24}} y^{J_0^3} (-1)^{F+\bar{F}} \right) \quad (5.2)$$

$$= \frac{1}{N} \sum_{r,s=1} \phi_{g^r, g^s}(\tau, z) = \phi_{e,e}(\tau, z). \quad (5.3)$$

Here, the twisted-twining genera ϕ_{g^r, g^s} are relative to a cocycle α satisfying the conditions (D.18) and (D.23), in particular with trivial restriction to $\langle g \rangle$. The identity (5.3) can be verified case by case for all $g \in M_{24}$ satisfying the level-matching condition.

By (5.3), the space \mathcal{H}' defined by (5.1) is isomorphic to \mathcal{H} as a module over $\mathcal{N} = 4$ superconformal algebra; as such, one can define a representation of M_{24} over \mathcal{H}' satisfying the properties of Mathieu moonshine. Furthermore, \mathcal{H}' also carries a representation ρ' of $C_{M_{24}}^\alpha(g)/\langle g_\alpha \rangle$ given by the restriction of $\oplus_{r=0}^{N-1} \tilde{\rho}_{g,r}$ to \mathcal{H}' . It can be proved that, as an abstract group, $C_{M_{24}}^\alpha(g)/\langle g_\alpha \rangle$ is isomorphic to a subgroup of M_{24} [17]. More precisely, Q is identified with an element g' in the same conjugacy class as g and the image of $C_{M_{24}}^\alpha(g)/\langle g \rangle$

is the centralizer $C_{M_{24}}(g')$ of g' . Furthermore, for each $h \in C_{M_{24}}(g)$ the twining-genus

$$\begin{aligned}\phi_{e,g'^n h'}(\tau, z) &:= \text{Tr}_{\mathcal{H}'}(\rho'(g')^n \rho'(h) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) \\ &= \frac{1}{N} \sum_{r,s=0}^{N-1} \text{Tr}_{\mathcal{H}_{g^r}}(\tilde{\rho}_{g,r}(Q)^n \tilde{\rho}_{g,r}(g)^{-s} \tilde{\rho}_{g,r}(h) q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) \\ &= \frac{1}{N} \sum_{r,s=0}^{N-1} e^{\frac{2\pi i n r}{N}} \epsilon_{g,h} \left(\frac{1}{0} \frac{s}{r} \right) \phi_{g^r, g^{-s} h}(\tau, z) ,\end{aligned}\tag{5.4}$$

is the Mathieu moonshine twining genus associated with an element $g'^n h' \in C_{M_{24}}(g') \subseteq M_{24}$. The identities (5.4) can be verified directly, given the explicit knowledge of the twisted-twining genera. Thus, the representation ρ' can be thought of as the restriction to $C_{M_{24}}(g')$ of the moonshine representation of M_{24} over \mathcal{H}' . From this construction, it is clear that one can simply identify g and g' as elements of an abstract M_{24} group, as well as their centralizers in M_{24} . Although we will make this identification in the following sections, for the moment being it is convenient to consider g and g' as distinct elements.

Since \mathcal{H}' satisfies the properties of Mathieu moonshine, we can introduce the twisted sectors $\mathcal{H}'_{g'^n}$ that carry projective representations $\rho'_{g'^n}$ of $C_{M_{24}}(g')$, such that the twisted-twining genera

$$\phi_{g'^n, h'}(\tau, z) = \text{Tr}_{\mathcal{H}'_{g'^n}}(\rho'_{g'^n}(h') q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) ,\tag{5.5}$$

satisfy the generalized Mathieu moonshine conjecture with respect to some 3-cocycle α' . As in the previous section, we can choose the cocycle α' to satisfy the analogue of (D.18) and (D.23) and define a corresponding central extension

$$1 \rightarrow \langle Q' \rangle \rightarrow C^{\alpha'}(g') \rightarrow C_{M_{24}}(g') \rightarrow 1 ,\tag{5.6}$$

and the representations $\tilde{\rho}'_{g',n}$. By (5.4), we have

$$\text{Tr}_{\mathcal{H}'_{g'^n}}(q^{L_0 - \frac{c}{24}} y^{J_0^3} (-1)^{F+\bar{F}}) = \phi_{g'^n, e}(\tau, z) = \phi_{e, g'^{-n}}(-\frac{1}{\tau}, \frac{z}{\tau})\tag{5.7}$$

$$= \frac{1}{N} \sum_{r,s=0}^{N-1} e^{-\frac{2\pi i n r}{N}} \phi_{g^r, g^{-s}}(-\frac{1}{\tau}, \frac{z}{\tau}) = \frac{1}{N} \sum_{r,s=0}^{N-1} e^{-\frac{2\pi i n r}{N}} \phi_{g^s, g^r}(\tau, z) ,\tag{5.8}$$

so that the g'^n -twisted sector $\mathcal{H}'_{g'^n}$ can be identified as an $\mathcal{N} = 4$ module with the g -eigenspace of $\oplus_{r=0}^{N-1} \mathcal{H}_{g^r}$ with eigenvalue $e^{\frac{2\pi i n}{N}}$. If we exchange the roles of g and g' , we obtain an isomorphism of $\mathcal{N} = 4$ modules

$$\mathcal{H}'_{g'^n, r} \cong \mathcal{H}_{g^r, n} ,\tag{5.9}$$

where $\mathcal{H}_{g^r,n}$ is the g -eigenspace of \mathcal{H}_{g^r} relative to the eigenvalue n and $\mathcal{H}'_{g'^n,r}$ is the analogous eigenspace for g' . It is then natural to conjecture that this isomorphism is equivariant with respect to the action of $C_{M_{24}}^\alpha(g)$

Conjecture. *For all choices of cocycles α and α' satisfying (D.18) and (D.23), there is an isomorphism $\varphi : C_{M_{24}}^\alpha(g) \rightarrow C_{M_{24}}^{\alpha'}(g')$ such that $\varphi(g) = Q'$, $\varphi(Q) = g'$, and for all $k \in C_{M_{24}}^\alpha(g)$, $m, r \in \mathbb{Z}$,*

$$\mathrm{Tr}_{\mathcal{H}'_{g'^m,r}}(\tilde{\rho}'_{g',m}(\varphi(k))q^{L_0-\frac{c}{24}}y^{J_0^3}(-1)^{F+\bar{F}}) = \mathrm{Tr}_{\mathcal{H}_{g^r,m}}(\tilde{\rho}_{g,r}(k)q^{L_0-\frac{c}{24}}y^{J_0^3}(-1)^{F+\bar{F}}). \quad (5.10)$$

Notice that if such an isomorphism exists for a certain choice of α and α' , then for any other choices it can be obtained by composing φ with (D.14). It is useful to rephrase the conjecture in terms of identities among twisted-twining genera. Recall that any element of $C_{M_{24}}^\alpha(g)$ can be written as $Q^x h_\alpha$, for some $x \in \mathbb{Z}/N\mathbb{Z}$, $h \in C_{M_{24}}(g)$ and any element of $C_{M_{24}}^{\alpha'}(g')$ can be written as $Q'^y h'_{\alpha'}$, for some $y \in \mathbb{Z}/N\mathbb{Z}$, $h' \in C_{M_{24}}(g')$. The conjecture implies that for any $k = Q^x h_\alpha \in C_{M_{24}}^\alpha(g)$ there is an element $\varphi(k) = Q'^y h'_{\alpha'} \in C_{M_{24}}^{\alpha'}(g')$ such that for all $r, m \in \mathbb{Z}$

$$\sum_{b=0}^{N-1} \frac{e^{\frac{2\pi i r(b-x)}{N}}}{N} \epsilon'_{g',h'} \left(\begin{smallmatrix} 1 & b \\ 0 & m \end{smallmatrix} \right) \phi_{g'^m, g'^{-b} h'}(\tau, z) = \sum_{s=0}^{N-1} \frac{e^{\frac{2\pi i m(s-y)}{N}}}{N} \epsilon_{g,h} \left(\begin{smallmatrix} 1 & s \\ 0 & r \end{smallmatrix} \right) \phi_{g^r, g^{-s} h}(\tau, z). \quad (5.11)$$

In particular, for $m \equiv 0 \pmod{N}$, we have $\epsilon_{g',h'} \left(\begin{smallmatrix} 1 & b \\ 0 & m \end{smallmatrix} \right) = 1$, and by taking the sum of these identities over $0 \leq r < N$, we reobtain (5.4).

Obviously, since g and g' are elements in the same conjugacy classes of M_{24} , there exist also isomorphisms $C_{M_{24}}^\alpha(g) \cong C_{M_{24}}^{\alpha'}(g')$ that map g to g' and Q to Q' . In particular, since g' is defined up to conjugation in M_{24} , we can simply choose $g' = g$ so that there is a natural identification of the centralizers $C_{M_{24}}(g) = C_{M_{24}}(g')$ and, upon choosing the same cocycles $\alpha = \alpha'$, also of their central extensions $C_{M_{24}}^\alpha(g) = C_{M_{24}}^{\alpha'}(g')$. With these conventions, the conjecture is equivalent to the existence of an outer automorphism of $C_{M_{24}}^\alpha(g)$ that exchanges g and Q and the representations $\mathcal{H}_{g^m,r}$ and $\mathcal{H}_{g^r,m}$.

The existence of automorphisms that exchange g and Q is easy to check for the elements g of order higher than 4 (thus, excluding the classes 2A and 4B of M_{24}). Indeed, in all these cases the central extension of $C_{M_{24}}(g)$ has the form $C_{M_{24}}^\alpha(g) \cong \langle Q \rangle \times (\langle Qg \rangle.G)$. This group clearly admits an automorphism that fixes the factor $\langle Qg \rangle.G$ and exchanges Q and g . For the classes 2A and 4B, the existence of the automorphism φ has been verified in [17] with the aid of the software GAP. In order for the conjecture to be proved, one needs to

show that, for each $h \in C_{M_{24}}(g)$, the image $\varphi(h_\alpha) = Q^y h'_\alpha$ of the lift $h_\alpha \in C_{M_{24}}^\alpha(g)$ satisfies

$$e^{\frac{2\pi i m y}{N}} \sum_{b=0}^{N-1} \frac{e^{\frac{2\pi i r b}{N}}}{N} \epsilon_{g,h'} \begin{pmatrix} 1 & b \\ 0 & m \end{pmatrix} \phi_{g^m, g^{-b} h'}(\tau, z) = \sum_{s=0}^{N-1} \frac{e^{\frac{2\pi i m s}{N}}}{N} \epsilon_{g,h} \begin{pmatrix} 1 & s \\ 0 & r \end{pmatrix} \phi_{g^r, g^{-s} h}(\tau, z) , \quad (5.12)$$

for all $m, r \in \mathbb{Z}$. If g is not in one of the classes $2A$ or $4B$, then these identities are satisfied with

$$h' = h , \quad y = 0 , \quad (5.13)$$

for all $h \in C_{M_{24}}(g) \setminus \langle g \rangle$, as well as $\varphi(g^i) = Q^i$. When g is in one of the classes $2A$ or $4B$, the identities is satisfied for suitable automorphisms φ of $C_{M_{24}}^\alpha(g)$. Tables 2 and 3 report the conjugacy class of h' in $C_{M_{24}}(g)$ for each $C_{M_{24}}(g)$ -conjugacy class of h . Notice that, in general, h and h' might belong to different conjugacy classes of M_{24} ; this phenomenon has been dubbed ‘relabeling’ in [17]. When h and h' are in different conjugacy classes of $C_{M_{24}}(g)$, we can choose the cocycle α in such a way that $y = 0$. When h and h' are conjugated in $C_{M_{24}}(g)$, then y can be set to 0 if and only if $\varphi(h_\alpha)$ is conjugated with h_α in the central extension $C_{M_{24}}^\alpha(g)$. This is always the case, except for g in M_{24} -class $4B$ and h in one of the classes $2B_3$ or $4B_1$ of $C_{M_{24}}(g)$. In both these cases, the pair g, h generates group 20 in the list of appendix E and $\varphi(h_\alpha)$ is conjugated with $Q^2 h_\alpha$, i.e. $y = 2$.

5.1.2 S-duality invariance

In this section, we will prove that all functions $\Phi_{g,h}(\sigma, \tau, z)$ defined in terms of the infinite product (3.59), satisfy identities of the form

$$\Phi_{g,h}(\sigma, \tau, z) \sim \Phi_{g,h'}\left(\frac{\tau}{N\lambda} + x, N\lambda\sigma, z\right) , \quad (5.14)$$

for some suitable $h' \in C_{M_{24}}(g)$ and real x that depends on the cocycle α . Here, \sim denotes equality up to a phase, that depends on the cocycle α . More precisely, in this section we will prove identities of the form (5.14) for all pairs $(g, h) \in \mathcal{P}$, except when g is in one of the classes $4A$, $10A$, $12A$ and $21AB$. For the latter classes, the identities (5.14) can be derived through a complete analysis of the automorphic properties of $\Phi_{g,h}$, as discussed in section 5.2.

When the function $\Phi_{g,h}$ admits a physical interpretation as the generating functions for twisted multiplicities of $1/4$ BPS states in a CHL model, the identity (5.14) corresponds to S-duality exchanging electric and magnetic charges in the low energy effective action.

Case 1: $g \in M_{23}$

Let g be an element of $M_{23} \subset M_{24}$ of order N . The restriction of $[\alpha]$ to $H^3(\langle g \rangle, U(1))$ is trivial, so that $\lambda = 1$ and we can choose a representative α satisfying (D.18) and (D.23).

Let us identify $g = g'$ and the centralizers $C_{M_{24}}(g) = C_{M_{24}}(g')$ and consider the same cocycle $\alpha = \alpha'$. The conjecture (5.10) implies that for any $h \in C_{M_{24}}(g)$, $d, m \in \mathbb{Z}$, there are $h' \in C_{M_{24}}(g')$ and $y \in \mathbb{Z}/N\mathbb{Z}$ such that

$$\sum_{s=0}^{N-1} \frac{e^{\frac{2\pi i ds}{N}}}{N} \epsilon_{g,h} \left(\begin{smallmatrix} k & s \\ 0 & m \end{smallmatrix} \right) \phi_{g^m, g^{-s} h'^k}(\tau, z) = \sum_{b=0}^{N-1} \frac{e^{\frac{2\pi i m(b-ky)}{N}}}{N} \epsilon_{g,h} \left(\begin{smallmatrix} k & b \\ 0 & d \end{smallmatrix} \right) \phi_{g^d, g^{-b} h^k}(\tau, z) , \quad (5.15)$$

that generalizes the identity of the previous section. Thus,

$$\hat{c}_{g,h}(d, m, \ell, t) = \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i tk}{M}}}{M} \sum_{b=0}^{N-1} \frac{e^{\frac{2\pi i bm}{N}}}{N} \epsilon_{g,h} \left(\begin{smallmatrix} k & b \\ 0 & d \end{smallmatrix} \right) c_{g^d, g^{-b} h^k} \left(\frac{md}{N}, \ell \right) \quad (5.16)$$

$$= \sum_{k=0}^{M-1} e^{\frac{2\pi i k m y}{N}} \frac{e^{-\frac{2\pi i tk}{M}}}{M} \sum_{s=0}^{N-1} \frac{e^{\frac{2\pi i ds}{N}}}{N} \epsilon_{g,h'} \left(\begin{smallmatrix} k & s \\ 0 & m \end{smallmatrix} \right) c_{g^m, g^{-s} h'^k} \left(\frac{md}{N}, \ell \right) \quad (5.17)$$

$$= \hat{c}_{g,h'}(m, d, \ell, t + my \frac{M}{N}) . \quad (5.18)$$

If h and h' are in different conjugacy classes of $C_{M_{24}}(g)$, we can choose the cocycle α in such a way that $y = 0$, so that

$$\Phi_{g,h}(\sigma, \tau, z) = p q^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h}(d,m,\ell,t)} \quad (5.19)$$

$$= p q^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h'}(m,d,\ell,t)} \quad (5.20)$$

$$= \Phi_{g,h'} \left(\frac{\tau}{N}, N\sigma, z \right) . \quad (5.21)$$

When h and h' are in the same conjugacy class of $C_{M_{24}}(g)$, then y cannot be eliminated, in general. This corresponds to the case when $\varphi(h_\alpha)$ is conjugated with $Q^y h_\alpha$ in the central extension $C_{M_{24}}(g)$. In particular, since $\varphi(h_\alpha)^M = \varphi(h_\alpha^M) = e$, this implies that Q^y has order M , i.e. there is an integer x such that

$$yM = xN . \quad (5.22)$$

Thus,

$$\Phi_{g,h}(\sigma, \tau, z) = pq^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h}(d,m,\ell,t)} \quad (5.23)$$

$$= pq^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t=0}^{M-1} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t+mx)} \quad (5.24)$$

$$= pq^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t'=0}^{M-1} (1 - e^{\frac{2\pi i t'}{M}} e^{-\frac{2\pi i x m}{M}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t)} \quad (5.25)$$

$$= pq^{\frac{1}{N}} y \prod_{(d,m,\ell)} \prod_{t'=0}^{M-1} (1 - e^{\frac{2\pi i t'}{M}} e^{-\frac{2\pi i y m}{N}} q^{\frac{m}{N}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t)} \quad (5.26)$$

$$= e^{\frac{2\pi i y}{N}} \Phi_{g,h}\left(\frac{\tau - y}{N}, N\sigma, z\right). \quad (5.27)$$

In fact, $y \neq 0$ only when g is in class 4B and h in one of the classes 2B₃ or 4B₁ of $C_{M_{24}}(g)$. In both cases, the pair g, h generates group 20 in the list of appendix E and $y = 2$. For such pairs (g, h) , however, since $T_L^\alpha \phi_{g,h} = 0$ for L odd, the function $\Phi_{g,h}$ satisfies the identity

$$\Phi_{g,h}(\sigma, \tau, z) = -\Phi_{g,h}\left(\sigma + \frac{1}{2}, \tau, z\right). \quad (5.28)$$

Using this identity, we conclude

$$\Phi_{g,h}(\sigma, \tau, z) = \Phi_{g,h}\left(\frac{\tau}{N}, N\sigma, z\right). \quad (5.29)$$

Case 2: g in one of the classes 2B, 3B, 4C, 6B, 12B

The calculations in this case are rather technical and have therefore been relegated to appendix D.3. Below we summarize the results.

For a suitable choice of the cocycle α , we have

$$\Phi_{g,h}(\sigma, \tau, z) = \Phi_{g,h^{-1}}\left(\frac{\tau}{N^2}, N^2\sigma, z\right), \quad (5.30)$$

if h and h^{-1} are not conjugated within $C_{M_{24}}(g)$, and

$$\Phi_{g,h}(\sigma, \tau, z) = \Phi_{g,h}\left(\frac{\tau}{N^2}, N^2\sigma, z\right), \quad (5.31)$$

otherwise. In particular, by (D.68) and (3.42) the latter holds for a choice of the cocycle α such that

$$\phi_{g,h}(\tau, z) = \phi_{g,h}^*(\tau, z), \quad (5.32)$$

i.e. such that all Fourier coefficients of $\phi_{g,h}$ are real.

5.2 Automorphic properties

Using the results of the previous sections, we can now prove that the functions $\Phi_{g,h}(\sigma, \tau, z)$, defined by the analytic continuation of the infinite product (3.59) to the Siegel upper half-space \mathbb{H}_2 are automorphic forms under certain subgroups of $Sp(4, \mathbb{R})$ acting by

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \cdot Z = (AZ + B)(CZ + D)^{-1}, \quad \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in Sp(4, \mathbb{R}) \quad (5.33)$$

where

$$Z = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix}. \quad (5.34)$$

Recall that the function $\Phi_{g,h}$ can be written as

$$\Phi_{g,h}(\sigma, \tau, z) = p\psi_{g,h}(\tau, z) \exp\left[-\sum_{N=1}^{\infty} p^L(\mathcal{T}_L^\alpha \phi_{g,h})(\tau, z)\right], \quad (5.35)$$

where the right-hand side converges on a suitable domain in \mathbb{H}_2 .

Let us consider the action of various generators of $Sp(4, \mathbb{R})$ on $\Phi_{g,h}$.

- The Heisenberg subgroup $H(\mathbb{Z})$ of $Sp(4, \mathbb{Z})$ is defined as

$$H(\mathbb{Z}) = \{[\lambda, \mu, \kappa] := \begin{pmatrix} 1 & 0 & 0 & \mu \\ \lambda & 1 & \mu & \kappa \\ 0 & 0 & 1 & -\lambda \\ 0 & 0 & 0 & 1 \end{pmatrix}, \lambda, \mu, \kappa \in \mathbb{Z}\}, \quad (5.36)$$

and acts by

$$[\lambda, \mu, \kappa] \cdot (\sigma, \tau, z) = (\sigma + \kappa + 2\lambda z + \lambda^2 \tau, \tau, z + \mu + \lambda \tau). \quad (5.37)$$

By (5.35), using the elliptic properties of the Jacobi forms $\psi_{g,h}$ and $\mathcal{T}_L^\alpha \phi_{g,h}$, we conclude easily that every $\Phi_{g,h}$ is invariant under $H(\mathbb{Z})$.

- For many commuting pairs $g, h \in M_{24}$, the genera $\mathcal{T}_L^\alpha \phi_{g,h}$ vanish unless L is an integer multiple of some $r \equiv r_{g,h} \in \mathbb{Z}$. Therefore, $\Phi_{g,h}$ is invariant up to a multiplier under $\sigma \mapsto \sigma + \frac{1}{r}$, i.e.

$$\Phi_{g,h}\left([0, 0, \frac{\kappa}{r}] \cdot (\sigma, \tau, z)\right) = e^{-\frac{2\pi i \kappa'}{r}} \Phi_{g,h}(\sigma, \tau, z), \quad (5.38)$$

where $[0, 0, \frac{\kappa}{r}] \cdot (\sigma, \tau, z) = (\sigma + \frac{\kappa'}{r}, \tau, z)$. In particular, $r_{g,h} = 1$ when g, h generate a cyclic group and for groups 13, 23, 24 and 27, $r = 3$ for groups 33 and 34, $r = 4$ for

groups 25 and 26 and $r = 2$ in all the other cases. Notice that $r_{g,h}$ always divides $N\lambda$.

More generally, the transformation $[0, 0, \frac{\kappa}{N\lambda}]$ relates the functions $\Phi_{g,h}$ relative to two distinct choices of the cocycle α , that differ from one each other by a $N\lambda$ -root of unity.

- The twisted-twining genera $\mathcal{T}_L^\alpha \phi_{g,h}$ are Jacobi forms of weight 0 and index L under a group $\Gamma_{g,h} \subset SL(2, \mathbb{Z})$ (see the tables in appendix E), up to a multiplier $\chi_{g,h}^L$. As noticed above, for each group $\langle g, h \rangle$, there is an integer $r \equiv r_{g,h}$ such that $\mathcal{T}_L^\alpha \phi_{g,h}$ vanishes unless $L|r$. Therefore, only the power $\chi_{g,h}^r$ needs to be a well-defined character of $\Gamma_{g,h}$.

On the other hand, the properties (4.4) and (4.5) imply that the eta products $\eta_{g,h}$ are modular forms of weight w and multiplier $v_{g,h}$ under the same group $\Gamma_{g,h}$ associated with $\phi_{g,h}$. Equivalently, each $\psi_{g,h}$ is a weak Jacobi form of weight $w - 2$, index 1 and multiplier $v_{g,h}$ under $\Gamma_{g,h}$.

For each $(\gamma, k) \in SL(2, \mathbb{Z}) \times M_{24}$, let us choose $\mu_{g,h}(\gamma, k) \in \mathbb{R}/\mathbb{Z}$ such that

$$e^{2\pi i r_{g,h} \mu_{g,h}(\gamma, k)} = \left(\epsilon_{g,h}(\gamma, k) \right)^{r_{g,h}}, \quad \gamma \in SL(2, \mathbb{Z}) . \quad (5.39)$$

Then for any $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ and $k \in M_{24}$, we have

$$v_{g,h}(\gamma) \Phi_{(\gamma, k) \cdot (g, h)}(\xi(\gamma) \cdot Z) = \det(c\tau + d)^{w-2} \Phi_{g,h}(\sigma + \mu_{g,h}(\gamma, k), \tau, z) , \quad (5.40)$$

where $Z = \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix}$ and

$$\xi \begin{pmatrix} a & b \\ c & d \end{pmatrix} := \begin{pmatrix} a & 0 & b & 0 \\ 0 & 1 & 0 & 0 \\ c & 0 & d & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} , \quad (5.41)$$

so that

$$\xi(\gamma) \cdot \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} = \begin{pmatrix} \frac{a\tau+b}{c\tau+d} & \frac{z}{c\sigma+d} \\ \frac{z}{c\sigma+d} & \sigma - \frac{z^2}{c\sigma+d} \end{pmatrix} . \quad (5.42)$$

In particular, by taking $(\gamma, k) \in M_{24}$ stabilizing (g, h) , we obtain

$$v'_{g,h}(\gamma, k) \Phi_{g,h}([0, 0, -\mu_{g,h}(\gamma, k)] \cdot (\xi(\gamma) \cdot Z)) = \det(c\tau + d)^{w-2} \Phi_{g,h}(\sigma, \tau, z) , \quad (5.43)$$

where $\gamma \in \Gamma_{g,h}$ and

$$v'_{g,h}(\gamma, k) = v_{g,h}(\gamma) e^{-2\pi i \mu_{g,h}(\gamma, k)} . \quad (5.44)$$

- Finally, for any pair (g, h) of commuting elements of M_{24} (with the exception of g in the classes 4A, 10A, 12A, 21AB), we have proved that

$$\Phi_{g,h}(Z) = \Phi_{g,h'}(V_{N\lambda} \cdot Z) , \quad (5.45)$$

for a suitable $h' \in C_{M_{24}}(g)$ and a suitable choice of the cocycle α , where

$$V_t = \frac{1}{\sqrt{t}} \begin{pmatrix} 0 & t & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & t & 0 \end{pmatrix} , \quad (5.46)$$

acts by

$$V_t \cdot \begin{pmatrix} \tau & z \\ z & \sigma \end{pmatrix} = \begin{pmatrix} \frac{\tau}{t} & z \\ z & t\sigma \end{pmatrix} . \quad (5.47)$$

Notice that in each of the 55 conjugacy classes of abelian subgroups $\langle g, h \rangle \subset M_{24}$, an identity of the form (5.45) has been proved for at least one pair of generators.

As discussed in [51, 61], for any integers $N, t > 0$, the elements

$$V_t, \quad \xi(\gamma) , \quad \gamma \in \Gamma_0(N) \quad [\lambda, \mu, \kappa/t] , \quad \lambda, \mu, \kappa \in \mathbb{Z} , \quad (5.48)$$

generate the group $\Gamma_t^+(N) = \langle \Gamma_t(N), V_t \rangle \subset Sp(4, \mathbb{R})$ which is a normal double extension of the paramodular group

$$\Gamma_t(N) = \left\{ \begin{pmatrix} * & t* & * & * \\ * & * & * & t^{-1}* \\ N* & Nt* & * & * \\ Nt* & Nt* & t* & * \end{pmatrix} \in Sp(4, \mathbb{Q}), \quad * \in \mathbb{Z} \right\} . \quad (5.49)$$

From the discussion above, it follows that every $\Phi_{g,h}$ is a modular function under some finite index subgroup $\Gamma_{g,h}^{(2)}$ of a paramodular group $\Gamma_t \equiv \Gamma_t(1)$, for some suitable t . The image of $\Phi_{g,h}$ under the action of a generic element of Γ_t is expected to be again a function $\Phi_{g',h'}$ for some (possibly different) commuting pair $g', h' \in M_{24}$, and defined with respect to a suitable choice of the cocycle α .

In section 5.3, we will discuss the automorphic properties of some of these functions $\Phi_{g,h}$. We leave the detailed description of the groups $\Gamma_{g,h}^{(2)}$ for all pairs g, h to a future work.

5.3 Examples

Having established that the $\Phi_{g,h}$ are indeed Siegel modular forms we now wish to analyze some specific examples in detail.

$\Phi_{2A,2A_2,3,5}$ (Groups 1, 2, 3)

Groups 1, 2 and 3 are $\mathbb{Z}_2 \times \mathbb{Z}_2$ groups that contain three elements in class 2A of M_{24} . The corresponding $\Phi_{g,h}$ are identical

$$\Phi_{2A,2A_2}(\sigma, \tau, z) = \Phi_{2A,2A_3}(\sigma, \tau, z) = \Phi_{2A,2A_5}(\sigma, \tau, z) . \quad (5.50)$$

The 3-cocycle α can be chosen to have trivial restriction to these groups. The Hecke transforms of the twisted-twining genera $\mathcal{T}_L \phi_{g,h}$ vanish unless L is a multiple of $r = 2$; in this case, they are Jacobi forms under $\Gamma_{g,h} = SL(2, \mathbb{Z})$ with trivial multiplier $\chi_{g,h}^2 = 1$. The eta products for these elements are given by

$$\eta_{g,h}(\tau) = \eta(\tau)^{12} , \quad (5.51)$$

and are modular forms of weight $w = 6$ under $SL(2, \mathbb{Z})$ with multiplier (see eq. (5.40))

$$v_{g,h}(T) = -1 \quad v_{g,h}(S) = -1 . \quad (5.52)$$

Furthermore, from the results of section 5.1.2 we deduce that, since $g \in M_{23}$ (with $N = 2$ and $\lambda = 1$), $\Phi_{g,h}$ satisfies

$$\Phi_{2A,2A_2}(\sigma, \tau, z) = \Phi_{2A,2A_3}\left(\frac{\tau}{2}, 2\sigma, z\right) , \quad \Phi_{2A,2A_5}(\sigma, \tau, z) = \Phi_{2A,2A_5}\left(\frac{\tau}{2}, 2\sigma, z\right) . \quad (5.53)$$

Thus, $\Phi_{g,h}$ is a modular form of weight $w - 2 = 4$ with a multiplier $v_{g,h}$ given above under the subgroup of $Sp(4, \mathbb{R})$ generated by the Heisenberg group $H(\mathbb{Z})$, by $SL(2, \mathbb{Z})$ and under V_2 . These elements of $Sp(4, \mathbb{R})$ generate the paramodular group $\Gamma_2^+(1)$. Furthermore, $\Phi_{g,h}$ has a double zero at the rational quadratic divisor $z \rightarrow 0$ and at all modular images of this divisor. This allows to identify $\Phi_{g,h}$ as

$$\Phi_{g,h}(Z) = \Delta_2(Z)^2 , \quad (5.54)$$

where Δ_2 is the modular form of weight 2 under $\Gamma_2^+(1)$ defined by Gritsenko and Nikulin in [51]. Notice that in this case the function $\psi_{g,h}$ in (4.16) is given by

$$\psi_{g,h}(\tau, z) = -\vartheta_1(\tau, z)^2 \eta(\tau)^6 , \quad (5.55)$$

which is the additive seed for $\Delta_2(Z)$ [51]. Hence we conclude that in this case

$$\Phi_{g,h} = \text{Add}[\psi_{g,h}] , \quad (5.56)$$

as claimed in section 3.3.3. The Siegel modular form $(\Delta_2)^2$ also appears in the context of umbral moonshine, where it corresponds to the Siegel modular form $\Phi^{(3)}(Z)$ (see section 2.6 of [29]). In this context $\Phi^{(3)}$ arises as the multiplicative lift of the umbral Jacobi form $Z^{(3)}(\tau, z)$ of weight 0 and index 2. It is therefore interesting to ask how this relates to our construction of $\Phi_{g,h}$ as the multiplicative lift (3.59). By comparing Fourier coefficients one can verify that we have the identity (see the table in appendix E for the result of the Hecke action)

$$\mathcal{T}_2 \phi_{g,h}(\tau, z) = Z^{(3)}(\tau, z). \quad (5.57)$$

However, this does not necessarily imply that the two lifts are the same, since a priori our multiplicative lift in (3.59) differ from that in [29] since we are constructing it from the seed function $\phi_{g,h}$, which is a weak Jacobi form of weight 0 and index 1, using the equivariant Hecke operator \mathcal{T}_N (the twist by the 3-cocycle α is trivial in this case). Even so, by virtue of (5.56), in the case at hand it turns out that the lifts do coincide and we thus have

$$\text{Mult}[\phi_{g,h}] = \Phi^{(3)}. \quad (5.58)$$

$\Phi_{2A,2A_4}$ (Group 7)

The function $\Phi_{2A,2A_4}$ of group 7 satisfies

$$\Phi_{2A,2A_4}(V_2 \cdot Z) = \Phi_{2A,4B_1}(Z), \quad (5.59)$$

where h' in class $4B_1$ of $C_{M_{24}}(g)$ is such that $h'^2 = g$. Therefore, the right hand side is a $Sp(4, \mathbb{Z})$ transformation of $\Phi_{e,4B}$, which is a Siegel modular form under $\Gamma_1(4) = \Gamma_0^{(2)}(4)$ [27, 28].

$\Phi_{2A,2B_{1,2}}$ (Groups 8,9)

The functions $\Phi_{g,h}$ for groups 8 and 9 are identical

$$\Phi_{2A,2B_1}(Z) = \Phi_{2A,2B_2}(Z). \quad (5.60)$$

The twisted-twining genera $\mathcal{T}_L^\alpha \phi_{g,h}$ vanish unless L is a multiple of $r = 2$; in this case, they are Jacobi forms under $\Gamma_{g,h} = \Gamma_0(2)$ with trivial multiplier. The eta products are given by

$$\eta_{g,h}(\tau) = \eta(\tau)^4 \eta(2\tau)^4, \quad (5.61)$$

and are modular forms of weight $w = 4$ under $\Gamma_0(2)$ with multiplier

$$v_{g,h}\left(\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right) = -1 \quad v_{g,h}\left(\begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}\right) = -1. \quad (5.62)$$

Therefore, $\Phi_{g,h}$ transforms as a modular form of weight $w - 2 = 2$ under $\xi(\gamma)$, $\gamma \in \Gamma_0(2)$, with multiplier $v_{g,h}(\gamma)$. Furthermore, for a suitable choice of the cocycle α , we have

$$\Phi_{2A,2B_{1,2}}(V_2 \cdot Z) = \Phi_{2A,2B_{1,2}}(Z) . \quad (5.63)$$

We conclude $\Phi_{g,h}$ is a modular form of weight 2 under the paramodular group $\Gamma_2(2)$. Assuming that it is holomorphic, then it can be identified as

$$\Phi_{g,h}(Z) = Q_1(Z)^2 , \quad (5.64)$$

where Q_1 is the modular form of weight 1 defined by Gritsenko and Clery in [61]. This conjecture is also supported by the fact that in this case $\psi_{g,h}$ is given by

$$\psi_{g,h}(\tau, z) = -\frac{\vartheta_1(\tau, z)^2}{\eta(\tau)^2} \eta(2\tau)^4, \quad (5.65)$$

which is the square of the additive seed for Q_1 (see eq. (16) in [61]).

$\Phi_{2A,4B_4}$ (Group 12)

The function $\Phi_{2A,4B_4}$ of group 12 satisfies

$$\Phi_{2A,4B_4}(V_2 \cdot Z) = \Phi_{2A,8A_1}(Z) , \quad (5.66)$$

where h' in class $8A_1$ of $C_{M_{24}}(g)$ is such that $h'^4 = g$. Therefore, the right hand side is a $Sp(4, \mathbb{Z})$ transformation of $\Phi_{e,8A}$, which is a Siegel modular form under $\Gamma_1(8) = \Gamma_0^{(2)}(8)$ [28].

$\Phi_{2A,4B_{2,3,5}}$ (Groups 17,18,19)

The functions $\Phi_{g,h}$ for groups 17, 18, 19 are identical

$$\Phi_{2A,4B_2}(Z) = \Phi_{2A,4B_3}(Z) = \Phi_{2A,4B_5}(Z) . \quad (5.67)$$

The Jacobi forms $\mathcal{T}_L^\alpha \phi_{g,h}$ for these groups have exactly the same modular properties as the ones for groups 8 and 9; the eta products are also the same. Therefore, $\Phi_{g,h}$ is a modular function of weight 2 under the paramodular group $\Gamma_2(2)$.

$\Phi_{4B,4B_{4,7}}$ (Groups 25,26)

The functions $\Phi_{g,h}$ for groups 25 and 26 are identical

$$\Phi_{4B,4B_4}(Z) = \Phi_{4B,4B_7}(Z) . \quad (5.68)$$

The 3-cocycle α can be chosen to be trivial when restricted to $\langle g, h \rangle$. The Hecke-transformed twisted-twining genera $\mathcal{T}_L \phi_{g,h}$ vanish unless L is a multiple of $r = 4$; in this case, they are Jacobi forms under $\Gamma_{g,h} = SL(2, \mathbb{Z})$ with trivial multiplier. The eta products are given by

$$\eta_{g,h}(\tau) = \eta(\tau)^6, \quad (5.69)$$

and are modular forms of weight $w = 3$ under $SL(2, \mathbb{Z})$ with multiplier

$$v_{g,h}\left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right) = -i \quad v_{g,h}\left(\begin{smallmatrix} 1 & 0 \\ 2 & 1 \end{smallmatrix}\right) = -i. \quad (5.70)$$

Furthermore,

$$\Phi_{4B,4B_4}(V_4 \cdot Z) = \Phi_{4B,4B_7}(Z). \quad (5.71)$$

We conclude $\Phi_{g,h}$ is a modular function of weight $w - 2 = 1$ under the paramodular group $\Gamma_4^+(1)$ and the multiplicity at the rational quadratic divisor $z \rightarrow 0$ is non-negative. It follows that

$$\Phi_{g,h}(Z) = \Delta_{1/2}(Z)^2, \quad (5.72)$$

where $\Delta_{1/2}$ is the modular form of weight $1/2$ defined in [51]. Also in this case the additive lift matches since

$$\psi_{g,h}(\tau, z) = -\vartheta_1(\tau, z)^2 \quad (5.73)$$

is the square of the additive seed for $\Delta_{1/2}(Z)$, and hence $\text{Add}[\psi_{g,h}] = (\Delta_{1/2}(Z))^2$.

The function $(\Delta_{1/2}(Z))^2$ also corresponds to the umbral Siegel modular form $\Phi^{(5)}(Z)$, which is the multiplicative lift of the umbral Jacobi form $Z^{(5)}(\tau, z)$ of weight 0 and index 4 [29]. The relation between $Z^{(5)}(\tau, z)$ and the twisted twining genus $\phi_{g,h}$ is now:

$$(\mathcal{T}_4 \phi_{g,h})(\tau, z) = Z^{(4)}(\tau, z). \quad (5.74)$$

Again, it is not a priori clear that the equivariant multiplicative lift of $\phi_{g,h}$, defined by (3.59), will coincide with the ordinary multiplicative Borcherds lift of $Z^{(5)}(\tau, z)$ considered in [29]. However, in the case at hand they do:

$$\text{Mult}[\phi_{g,h}] = \Phi^{(5)}. \quad (5.75)$$

$\Phi_{3A,3A_3}$ (Group 33)

Consider the function $\Phi_{3A,3A_3}$ of groups 33. The twisted-twining genera $\mathcal{T}_L^\alpha \phi_{g,h}$ vanish unless L is a multiple of $r = 3$; in this case, they are Jacobi forms under $\Gamma_{g,h} = SL(2, \mathbb{Z})$ with trivial multiplier. The eta product is given by

$$\eta_{g,h}(\tau) = \eta(\tau)^8, \quad (5.76)$$

and is a modular form of weight $w = 4$ under $SL(2, \mathbb{Z})$ with multiplier

$$v_{g,h}\left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right) = e^{-\frac{2\pi i}{3}} \quad v_{g,h}\left(\begin{smallmatrix} 0 & -1 \\ 0 & 1 \end{smallmatrix}\right) = 1 . \quad (5.77)$$

Furthermore,

$$\Phi_{3A,3A_3}(V_3 \cdot Z) = \Phi_{3A,3A_3}(Z) . \quad (5.78)$$

We conclude $\Phi_{g,h}$ is a modular function of weight $w - 2 = 2$ under the paramodular group $\Gamma_3^+(1)$ and the analysis of the multiplicities at its divisors shows that it is holomorphic with a double zero at $z = 0$. It follows that

$$\Phi_{3A,3A_3}(Z) = \Delta_1(Z)^2 , \quad (5.79)$$

where Δ_1 is the modular form of weight 1 defined in [51]. Also here we find that

$$\psi_{g,h}(\tau, z) = -\vartheta_1(\tau, z)^2 \eta(\tau)^2 \quad (5.80)$$

is the square of the additive seed for $\Delta_1(Z)$. The function $\Delta_1(Z)^2$ coincides with the umbral Siegel modular form $\Phi^{(4)}(Z)$, which is the multiplicative lift of the umbral Jacobi form $Z^{(4)}(\tau, z)$ of weight 0 and index 3. In this case we have that, for a suitable choice of cocycle α ,

$$(\mathcal{T}_3^\alpha \phi_{g,h})(\tau, z) = Z^{(3)}(\tau, z), \quad (5.81)$$

and the equivariant and multiplicative lifts coincide:

$$\text{Mult}[\phi_{g,h}] = \Phi^{(4)}. \quad (5.82)$$

$\Phi_{3A,3B_1}$ (Group 34)

Consider the function $\Phi_{3A,3B_1}$ of group 34. The twisted-twining genera $\mathcal{T}_L^\alpha \phi_{g,h}$ vanish unless L is a multiple of $r = 3$; in this case, they are Jacobi forms under $\Gamma_{g,h} = \Gamma_0(3)$ with trivial multiplier. The eta product is given by

$$\eta_{g,h}(\tau) = \eta(\tau)^2 \eta(3\tau)^2 , \quad (5.83)$$

and is a modular form of weight $w = 2$ under $SL(2, \mathbb{Z})$ with multiplier

$$v_{g,h}\left(\begin{smallmatrix} 1 & 1 \\ 0 & 1 \end{smallmatrix}\right) = e^{-\frac{2\pi i}{3}} \quad v_{g,h}\left(\begin{smallmatrix} 1 & 0 \\ 3 & 1 \end{smallmatrix}\right) = e^{\frac{2\pi i}{3}} . \quad (5.84)$$

Furthermore,

$$\Phi_{3A,3B_1}(V_3 \cdot Z) = \Phi_{3A,3B_1}(Z) . \quad (5.85)$$

We conclude $\Phi_{g,h}$ is a modular function of weight $w - 2 = 0$ under the paramodular group $\Gamma_3^+(3)$. Since the weight vanishes, it must necessarily be meromorphic.

6 Conclusions

In this paper we have proposed a second quantized version of (generalized) Mathieu moonshine, involving a class of Siegel modular forms $\Phi_{g,h}$ for discrete subgroups $\Gamma_{g,h}^{(2)} \subset Sp(4, \mathbb{R})$, constructed from a multiplicative lift of the twisted twining genera $\phi_{g,h}$. For certain pairs of conjugacy classes of M_{24} we were able to identify $\Phi_{g,h}$ with known Siegel modular forms. It would be interesting to extend these results and perform a more detailed investigation of all the modular groups $\Gamma_{g,h}^{(2)}$ and determine whether the remaining $\Phi_{g,h}$ coincide with known objects, or perhaps constitute new examples of Siegel modular forms. In [28], Raum proved modularity for most of the cases $\Phi_{e,h}$ and found that not all of them are of a standard Borcherds product type, but in fact correspond to certain rescaled products of Borcherds modular forms. One would like to extend this analysis to determine whether the $\Phi_{g,h}$ for $g \neq e$ also contain such rescaled Borcherds products, or some generalization thereof.

As already mentioned in the introduction, an interesting by-product of our analysis is the fact that some of the Siegel modular forms $\Phi_{g,h}$ coincide with multiplicative lifts of the umbral Jacobi forms analyzed in [29]. This might be a simple consequence of the constraints from modularity, but it might also indicate some deeper relation between umbral moonshine and generalized Mathieu moonshine which would be interesting to uncover. Could it be that some of the other products $\Phi_{g,h}$ also coincide with lifts of the more general D_n or E_n Niemeier-umbral Jacobi forms in [64]?

It is natural to wonder whether our products $\Phi_{g,h}$ have interpretations in terms of denominator formulas of some generalized Kac-Moody algebras (GKM). It is well-known that the Igusa cusp form $\Phi_{10} = \Phi_{e,e}$ constitutes one side of the denominator formula for a rank 3 GKM-algebra of hyperbolic type [58] (the other side corresponds to the additive lift in (3.67)). In the terminology of Borcherds [36], the other functions $\Phi_{e,h}$ then corresponds to twisted denominator formulas for the same algebra. On the other hand, by analogy with generalized Monstrous moonshine [41, 43], we would expect that the functions $\Phi_{g,e}$ give denominator formulas for a class of rank 3 GKM-algebras. For some of the elements g of small order these algebras have indeed been constructed in [65, 66, 67, 68] in the context of CHL-models (see also [58, 59, 51] for earlier mathematical results). In this context the prefactor $pq^{\frac{1}{N\lambda}}y$ in (3.59) should have an interpretation as the exponential of the Weyl vector ρ of the algebra. If true one would expect that for fixed g , the associated twisted denominator formulas $\Phi_{g,h}$ all have the same prefactor involving the Weyl vector of the original GKM-algebra determined by the class $[g]$. In other words, the prefactor should be independent of the twining element h , and this is indeed what we find. In fact, the Weyl vectors extracted from $pq^{\frac{1}{N\lambda}}y$ reduces to the ones in [65, 66] when the length of the

shortest cycle λ equals one.

As stressed in the introduction, we think that our results could have immediate applications to the understanding of dyon counting in CHL-orbifolds. In particular, most of the functions $\Phi_{g,h}$ are expected to have interpretations as partition functions of twisted dyons, and we hope to investigate this relation in more detail in a future publication.

Acknowledgments

We are especially grateful to Matthias Gaberdiel for collaboration in the initial stages of this work, and for many helpful discussions. We are also grateful to Geoff Mason for several useful discussions on M_{24} moonshine for eta-products, and for sending us a printed copy of [31]; to Miranda Cheng for a discussion that drew our attention to the connection with umbral moonshine; and to Martin Raum for helpful discussions on infinite product representations of Siegel modular forms. In addition we thank Scott Carnahan, Thomas Creutzig, Terry Gannon, Suresh Govindarajan, Jeff Harvey, Stefan Hohenegger, Gerald Höhn, Jim Lepowsky, Greg Moore, Sameer Murthy, Anne Taormina and Katrin Wendland for inspiring discussions during the course of this work. We also thank the Simons Center and the organizers of the programme “Mock Modular Forms, Moonshine, and String Theory” for providing a stimulating research environment, and all the participants for many inspiring discussions. D.P. thanks the Albert Einstein Institute in Golm, and R.V. thanks the Department of Fundamental Physics at Chalmers for hospitality while this work was carried out.

A Some group cohomology

In this appendix, we summarize some general results about group cohomology. See [69] for more details.

For a finite group G , a 2-cochain $\beta : G \times G \rightarrow U(1)$ is closed (and hence defines a cocycle) provided it satisfies

$$\beta(g_1, g_2 g_3) \beta(g_2, g_3) = \beta(g_1 g_2, g_3) \beta(g_1, g_2) \quad (\text{A.1})$$

for $g_1, g_2, g_3 \in G$. The second cohomology $H^2(G, U(1))$ then consists of the closed 2-cochains, modulo the ambiguity

$$\beta(g_1, g_2) \rightarrow \beta(g_1, g_2) \frac{\gamma(g_1) \gamma(g_2)}{\gamma(g_1 g_2)}, \quad (\text{A.2})$$

where $\gamma : G \rightarrow U(1)$ is an arbitrary 1-cochain, i.e. an arbitrary function $\gamma : G \rightarrow U(1)$.

A 3-cochain α ,

$$\alpha : G \times G \times G \rightarrow U(1) \quad (\text{A.3})$$

is closed provided it satisfies

$$\alpha(g_1, g_2, g_3) \alpha(g_1, g_2 g_3, g_4) \alpha(g_2, g_3, g_4) = \alpha(g_1 g_2, g_3, g_4) \alpha(g_1, g_2, g_3 g_4) . \quad (\text{A.4})$$

In the cohomology group $H^3(G, U(1))$ closed 3-cochains are then identified modulo

$$\alpha(g_1, g_2, g_3) \rightarrow \alpha(g_1, g_2, g_3) \frac{\beta(g_1, g_2 g_3) \beta(g_2, g_3)}{\beta(g_1 g_2, g_3) \beta(g_1, g_2)} . \quad (\text{A.5})$$

Note that the multiplying factor is trivial if β is closed, i.e. if it satisfies the 2-cocycle condition (A.1). In particular, for each cocycle α and element $x \in G$, the 3-cocycle α_x , defined by

$$\alpha_x(g, h, k) := \alpha(x^{-1} g x, x^{-1} h x, x^{-1} k x) , \quad (\text{A.6})$$

differs from α just by a 3-coboundary [46]

$$\alpha_x(g_1, g_2, g_3) = \alpha(g_1, g_2, g_3) \frac{\eta_x(g_1, g_2 g_3) \eta_x(g_2, g_3)}{\eta_x(g_1 g_2, g_3) \eta_x(g_1, g_2)} , \quad (\text{A.7})$$

where

$$\eta_z(x, y) := \frac{\alpha(x, y, z) \alpha(z, z^{-1} x z, z^{-1} y z)}{\alpha(x, z, z^{-1} y z)} . \quad (\text{A.8})$$

Given a 3-cocycle α , we can define, for any $h \in G$, a map $c_h : G \times G \rightarrow U(1)$ via

$$c_g(h_1, h_2) = \frac{\alpha(g, h_1, h_2) \alpha(h_1, h_2, (h_1 h_2)^{-1} g (h_1 h_2))}{\alpha(h_1, h_1^{-1} g h_1, h_2)} . \quad (\text{A.9})$$

It is shown in [44] that c_g defines a 2-cocycle of the stabilizer subgroup $C_G(g) \subseteq G$ (i.e. the subgroup of all elements h_1, h_2 which commute with g). When $h_1, h_2 \in C_G(g)$, we have the simplified expression

$$c_g(h_1, h_2) = \frac{\alpha(g, h_1, h_2) \alpha(h_1, h_2, g)}{\alpha(h_1, g, h_2)} , \quad h_1, h_2 \in C_G(g) . \quad (\text{A.10})$$

It is straightforward to check this by inserting (A.9) into (A.1), and making repeated use of the 3-cocycle condition (A.4) (twice on each side of the equality sign) together with the fact that g commutes with h_1, h_2, h_3 .

Under the ‘gauge transformation’ (A.5), c_g transforms as

$$c_g(h_1, h_2) \rightarrow \tilde{c}_g(h_1, h_2) := c_g(h_1, h_2) \frac{\gamma_g(h_1) \gamma_g(h_2)}{\gamma_g(h_1 h_2)} , \quad h_1, h_2 \in C_G(g) , \quad (\text{A.11})$$

where we defined the 1-cochain γ_g by

$$\gamma_g(h) \equiv \frac{\beta(h, g)}{\beta(g, h)} . \quad (\text{A.12})$$

This is indeed of the form (A.2), and hence, for all $g \in G$, c_g defines a map

$$c_g : H^3(G, U(1)) \rightarrow H^2(C_G(g), U(1)) . \quad (\text{A.13})$$

In fact, if c_g is the 2-cocycle associated to a projective representation ρ_g of $C_G(g)$, i.e.

$$\rho_g(h_1)\rho_g(h_2) = c_g(h_1, h_2)\rho_g(h_1h_2) , \quad (\text{A.14})$$

then \tilde{c}_g is the 2-cocycle associated with the projectively equivalent representation

$$\tilde{\rho}_g(h) := \gamma_g(h)\rho_g(h) . \quad (\text{A.15})$$

In the context of holomorphic CFTs, the transformation (A.5) corresponds to a redefinition (A.15) of the projective representations ρ_g of the centralizer $C_G(g)$ over the twisted sector \mathcal{H}_g , which induces the analogous transformation of the twisted twining partition functions

$$Z_{g,h} \rightarrow \tilde{Z}_{g,h} = \gamma_g(h)Z_{g,h} . \quad (\text{A.16})$$

Indeed, the new partition functions $\tilde{Z}_{g,h}$ satisfy the expected modular properties with respect to the new cocycle $\tilde{\alpha}$.

In particular, in the case (A.7) of conjugation by $x \in G$, we have $\beta \equiv \eta_x$, so that

$$\gamma_g(h) = \frac{\eta_x(h, g)}{\eta_x(g, h)} = \frac{c_g(x, x^{-1}hx)}{c_g(h, x)} , \quad h \in C_g(G) , \quad (\text{A.17})$$

where the latter equality follows from [45, 46]. From this identity, one recovers

$$Z_{(\mathbb{I}, x^{-1}) \cdot (g, h)}(\tau) \equiv Z_{x^{-1}gx, x^{-1}hx}(\tau) = \frac{c_g(x, x^{-1}hx)}{c_g(h, x)} Z_{g,h}(\tau) , \quad (\text{A.18})$$

from which the formula (B.8) for $\epsilon_{g,h}(\mathbb{I}, x)$ follows.

One further useful identity is [46]

$$\frac{c_{x_1x_2}(z_1, z_2)}{c_{x_1}(z_1, z_2)c_{x_2}(z_1, z_2)} = \frac{c_{z_1}(x_1, x_2)c_{z_2}(x_1, x_2)}{c_{z_1z_2}(x_1, x_2)} , \quad (\text{A.19})$$

that holds for *pairwise commuting* $x_1, x_2, z_1, z_2 \in G$.

For most applications, only the restriction of a 3-cocycle α to subgroups of the form $\langle g, h \rangle \cong \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}$ is needed. For such groups, we have $\mathcal{H}^3(\mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}, U(1)) = \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2} \times \mathbb{Z}_{\text{lcm}(N_1, N_2)}$ and a set of normalized representatives for the generators are [70]

$$\alpha_{v_i}(g^{a_1} h^{a_2}, g^{b_1} h^{b_2}, g^{c_1} h^{c_2}) := e^{\frac{2\pi i v_i a_i}{N_i^2} ([b_i]_{N_i} + [c_i]_{N_i} - [b_i + c_i]_{N_i})} \quad v_i \in \mathbb{Z}/N_i\mathbb{Z}, \quad i = 1, 2, \quad (\text{A.20})$$

$$\alpha_{v_{12}}(g^{a_1} h^{a_2}, g^{b_1} h^{b_2}, g^{c_1} h^{c_2}) := e^{\frac{2\pi i v_{12} a_1}{N_1 N_2} ([b_2]_{N_2} + [c_2]_{N_2} - [b_2 + c_2]_{N_2})} \quad v_{12} \in \mathbb{Z}/\text{lcm}(N_1, N_2)\mathbb{Z}, \quad (\text{A.21})$$

where $[\cdot]_x : \mathbb{Z} \rightarrow \{0, \dots, x-1\}$ denotes the reduction modulo x . Notice that, with this choice for the generators, we have the simplified formula

$$c_x(y, z) = \alpha(x, y, z), \quad x, y, z \in \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}. \quad (\text{A.22})$$

B Modular properties of twisted twining partition functions

In this appendix we include some details on the modular properties of twisted twining partition functions. In particular, we analyze the combined action of $SL(2, \mathbb{Z})$ and G on the set of commuting pairs (g, h) in G . Here we must also take into account the presence of a non-trivial 3-cocycle α which leads to a certain twisted action. We introduce a convenient “twisted equivariant slash-operator” that simplifies many expressions since it combines (twisted) $SL(2, \mathbb{Z})$ -equivariance with G -equivariance. Finally, we discuss a reformulated version of the cohomological obstructions found in [17].

B.1 $SL(2, \mathbb{Z}) \times G$ - action on \mathcal{P}_G

It is clear from (2.8) that modular transformations act on $\tau \in \mathbb{H}$ as well as on the set of commuting pairs $(g, h) \in G \times G$. In addition G acts on itself by conjugation and thereby on the set of pairs g, h . In order to determine the subgroups $\Gamma_{g, h} \subset SL(2, \mathbb{Z})$ under which $Z_{g, h}$ (and $\phi_{g, h}$) are invariant for fixed $g, h \in G$, we must classify the orbits of the combined $SL(2, \mathbb{Z}) \times G$ -action on the commuting pair (g, h) . We are mainly interested in the case $G = M_{24}$ but we only make this specification at the end of the subsection.

For any finite group G , let $\mathcal{P}_G \subset G \times G$ be the set of commuting pairs of elements:

$$\mathcal{P}_G = \{(g, h) \in G \times G \mid gh = hg\}. \quad (\text{B.1})$$

The group $SL(2, \mathbb{Z}) \times G$ has a left action as a permutation over this set by

$$\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, k \right) \cdot (g, h) := (k g k^{-1}, k h k^{-1}) \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = (k g^d h^{-c} k^{-1}, k g^{-b} h^a k^{-1}), \quad (\text{B.2})$$

where $(g, h) \in \mathcal{P}_G$, $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ and $k \in G$. This action can be extended to an action of $GL(2, \mathbb{Z}) \times G$ in the obvious way.

B.2 Twisted action on \mathcal{P}_G : generalized permutation

As explained in section 2.2, in order to include the possibility of non-trivial multipliers in the modular properties of the twisted twining genera one needs to consider the “twisting” of the action (B.2) by a 3-cocycle α , representing a cohomology class in $H^3(G, U(1))$. This α -twisted action is a generalized permutation

$$\mathbb{C}\mathcal{P}_G \times SL(2, \mathbb{Z}) \times G \rightarrow \mathbb{C}\mathcal{P}_G, \quad (\text{B.3})$$

on the complex vector space $\mathbb{C}\mathcal{P}_G$ freely generated by the elements of \mathcal{P}_G . We define this action by the formula

$$(\gamma, k)_\alpha : (g, h) \mapsto \epsilon_{g,h}(\gamma, k) (k g k^{-1}, k h k^{-1}) \gamma^{-1}. \quad (\text{B.4})$$

Here, $(\gamma, k) \in SL(2, \mathbb{Z}) \times G$ and $\epsilon_{g,h}(\gamma, k) \in U(1)$ is a phase which depends on the choice of 3-cocycle α . More precisely, in terms of the 2-cocycle $c_g(h_1, h_2)$ in (2.7) the phases $\epsilon_{g,h}(\gamma, k)$ are defined as

$$\epsilon_{g,h}(\gamma_1 \gamma_2, k_1 k_2) := \epsilon_{(\gamma_2, k_2) \cdot (g, h)}(\gamma_1, k_1) \epsilon_{g,h}(\gamma_2, k_2), \quad (\text{B.5})$$

$$\epsilon_{g,h}(T, e) := \frac{1}{c_g(g, g^{-1}h)}, \quad (\text{B.6})$$

$$\epsilon_{g,h}(S, e) := c_g(h^{-1}, h), \quad (\text{B.7})$$

$$\epsilon_{g,h}(\mathbb{I}, k) := \frac{c_g(h, k^{-1})}{c_g(k^{-1}, k h k^{-1})}, \quad k \in G, \quad (\text{B.8})$$

where

$$S := \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad T := \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \quad (\text{B.9})$$

are the generators of $SL(2, \mathbb{Z})$.

B.3 Modular properties of twisted-twining partition functions

We now wish to analyze the modular properties of the twisted twining partition function $Z_{g,h}$ in more detail. To this end we define the following equivariant slash operator

$$f(g, h; \tau)|(\gamma, k) := f((\gamma, k) \cdot (g, h); \gamma \cdot \tau) , \quad (\gamma, k) \in SL(2, \mathbb{Z}) \times G , \quad (\text{B.10})$$

acting on modular functions $f : \mathcal{P}_G \times \mathbb{H}_+ \rightarrow \mathbb{C}$. We can think of the twisted-twining partition functions $Z_{g,h}(\tau)$ as functions on $\mathcal{P}_G \times \mathbb{H}_+$ that are equivariant under $SL(2, \mathbb{Z}) \times G$, up to a multiplier. In the simplest case when the multiplier system $\chi_{g,h}$ in (2.8) is trivial, this property can be expressed as follows in terms of the slash operator:

$$Z_{g,h}(\tau)|(\gamma, k) = Z_{g,h}(\tau) . \quad (\text{B.11})$$

In particular, for each fixed $(g, h) \in \mathcal{P}_G$, the function $Z_{g,h}(\tau)$ is a modular function under some subgroup $\Gamma_{g,h} \in PSL(2, \mathbb{Z})$, i.e.

$$Z_{g,h}(\gamma \cdot \tau) = Z_{g,h}(\tau) , \quad \gamma \in \Gamma_{g,h} \subseteq PSL(2, \mathbb{Z}) . \quad (\text{B.12})$$

The group $\Gamma_{g,h}$ is the image $\pi(\tilde{\Gamma}_{g,h})$ of the stabilizer

$$\tilde{\Gamma}_{g,h} := \{(\gamma, k) \in SL(2, \mathbb{Z}) \times G \mid (g, h) \cdot (\gamma, k) = (g, h)\} , \quad (\text{B.13})$$

under the homomorphism

$$\pi : SL(2, \mathbb{Z}) \times G \rightarrow PSL(2, \mathbb{Z}) . \quad (\text{B.14})$$

Similar properties hold for the twisted-twining genera $\phi_{g,h}(\tau, z)$ of an $\mathcal{N} = (4, 4)$ superconformal algebra: these are expected to be Jacobi forms of weight zero and index 1 with respect to the same groups $\Gamma_{g,h}$. For this reason, we need to extend the definition of the slash operator to an action on the space of functions $\psi : \mathcal{P}_G \times \mathbb{H}_+ \times \mathbb{C} \rightarrow \mathbb{C}$. Specifically, for Jacobi forms of weight 0 and index m with respect to some $\Gamma \subset SL(2, \mathbb{Z})$ we define

$$\psi(g, h; \tau, z)|(\gamma, k) := e^{-\frac{2\pi i m c z^2}{c\tau + d}} \psi\left(k g^d h^{-c} k^{-1}, k g^{-b} h^a k^{-1}; \gamma \cdot \tau, \frac{z}{c\tau + d}\right) , \quad (\text{B.15})$$

where $(\gamma, k) = \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, k\right) \in SL(2, \mathbb{Z}) \times G$. Note that for theories with $\mathcal{N} = (4, 4)$ superconformal symmetry, these genera are expected to be *even* functions of z , so that the central element $C = S^2 \in SL(2, \mathbb{Z})$ acts trivially and it makes sense to consider the action on such functions of the quotient $PSL(2, \mathbb{Z}) = SL(2, \mathbb{Z})/\mathbb{Z}_2$.

We are now ready to incorporate the α -twist in the modular properties of the twisted twining genera. To this end we define the α -twisted generalization of the equivariant slash operator:

$$f(g, h, \tau)|_\alpha(\gamma, k) := \epsilon_{g,h}(\gamma, k) f((\gamma, k) \cdot (g, h); \gamma \cdot \tau) , \quad (\gamma, k) \in SL(2, \mathbb{Z}) \times G , \quad (\text{B.16})$$

for some 3-cocycle α representing a class $[\alpha] \in H^3(G, U(1))$. Similarly, we define the α -twisted slash operators on Jacobi forms of weight 0 and index m by

$$\psi(g, h; \tau, z)|_\alpha(\gamma, k) := \epsilon_{g,h}(\gamma, k) e^{-\frac{2\pi i m c z^2}{c\tau + d}} \psi\left(k g^d h^{-c} k^{-1}, k g^{-b} h^a k^{-1}; \gamma \cdot \tau, \frac{z}{c\tau + d}\right) , \quad (\text{B.17})$$

where $(\gamma, k) = \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, k\right) \in SL(2, \mathbb{Z}) \times G$.

B.4 Cohomological obstructions

When α represents a non-trivial class in $H^3(G, U(1))$, the partition functions $Z_{g,h}(\tau)$ are modular functions under $\Gamma_{g,h} \subseteq PSL(2, \mathbb{Z})$ only up to some multiplier $\chi_{g,h}$, which depends on α (see section 2.2). In fact, by (B.13) and (B.8), the restriction of $\epsilon_{g,h}$ to $\tilde{\Gamma}_{g,h}$ is a group homomorphism $\tilde{\Gamma}_{g,h} \rightarrow U(1)$. If

$$\epsilon_{g,h}(\gamma, k) = 1 , \quad (\text{B.18})$$

for all (γ, k) in

$$\tilde{\Gamma}_{g,h} \cap \ker \pi = \{(e, k) \in SL(2, \mathbb{Z}) \times G \mid (k^{-1} g k, k^{-1} h k) = (g, h)\} \quad (\text{B.19})$$

$$\cup \{(S^2, k) \in SL(2, \mathbb{Z}) \times G \mid (k^{-1} g k, k^{-1} h k) = (g^{-1}, h^{-1})\} , \quad (\text{B.20})$$

then the restriction $\epsilon_{g,h} : \tilde{\Gamma}_{g,h} \rightarrow U(1)$ induces a well-defined homomorphism $\chi_{g,h} : \Gamma_{g,h} \rightarrow U(1)$ on the image $\Gamma_{g,h} = \pi(\tilde{\Gamma}_{g,h})$. Explicitly, for each $\gamma \in \Gamma_{g,h} \subseteq PSL(2, \mathbb{Z})$, one can choose a lift $(\gamma, k) \in \tilde{\Gamma}_{g,h} \subseteq SL(2, \mathbb{Z}) \times G$ and set

$$\chi_{g,h}(\gamma) := \epsilon_{g,h}(\gamma, k) , \quad \text{for } \gamma = \pi(\gamma, k) \in \Gamma_{g,h} . \quad (\text{B.21})$$

By (B.18), the definition is independent of the lift.

On the contrary, if (B.18) is not satisfied for some $(\gamma, k) \in \tilde{\Gamma}_{g,h} \cap \ker \pi$, then eq.(2.11) implies that $Z_{g,h}(\tau)$ must vanish identically. In this case, we say that the twisted-twining partition function is *obstructed*. Following the discussion in [17, 18], we can distinguish between two kinds of obstructions:

- (1) We say that, for a certain $(g, h) \in \mathcal{P}_G$, there is an obstruction of the *first kind* if there is some element of the form $(e, k) \in \tilde{\Gamma}_{g,h} \cap \ker \pi$ for which (B.18) is not satisfied.

- (2) We say that there is an obstruction of the *second kind* if (B.18) is satisfied for all elements of the form $(e, k) \in \tilde{\Gamma}_{g,h} \cap \ker \pi$, but is violated by some element of the form $(S^2, k) \in \tilde{\Gamma}_{g,h} \cap \ker \pi$. By (B.8), this implies that (B.18) is false for *all* elements of the form $(S^2, k) \in \tilde{\Gamma}_{g,h} \cap \ker \pi$.

Although not phrased this way in [17] these obstructions are equivalent to the ones given there.

C Definition of twisted equivariant Hecke operators

In this appendix, we shall discuss the definition and properties of the α -twisted equivariant Hecke operators.

C.1 Twisted equivariant Hecke operator

Let us denote by $\text{Mat}(\mathbb{Z})$ the ring of 2×2 integral matrices with positive determinant, graded by the determinant,

$$\text{Mat}(\mathbb{Z}) = \bigcup_{N>0} \text{Mat}_N(\mathbb{Z}) , \quad (\text{C.1})$$

where $\text{Mat}_N(\mathbb{Z})$ was defined in (3.4). For any $u \in \text{Mat}_N(\mathbb{Z})$ denote by $u^\vee \in \text{Mat}_N(\mathbb{Z})$ the dual

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^\vee := N \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{-1} = \begin{pmatrix} d & -b \\ -c & a \end{pmatrix} , \quad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{Mat}_N(\mathbb{Z}) . \quad (\text{C.2})$$

Note that $\text{Mat}_1(2, \mathbb{Z}) \equiv SL(2, \mathbb{Z})$ is the group of invertible elements of $\text{Mat}(\mathbb{Z})$ and that each $\text{Mat}_N(\mathbb{Z})$ is a bimodule over $SL(2, \mathbb{Z})$.

In order to define the α -twisted equivariant Hecke operators \mathcal{T}_N^α , we need to extend the action (B.3) of $SL(2, \mathbb{Z}) \times G$ on \mathbb{CP}_G to an action

$$\mathbb{CP}_G \times \text{Mat}(\mathbb{Z}) \times G \rightarrow \mathbb{CP}_G , \quad (\text{C.3})$$

by

$$(u, k)_\alpha : (g, h) \mapsto \epsilon_{g,h}(u, k)(kgk^{-1}, khk^{-1})u^\vee , \quad (\text{C.4})$$

for a suitable $\epsilon_{g,h}(u, k) \in U(1)$, that reduces to the one discussed in section B.2 when $u \in \text{Mat}_1(\mathbb{Z}) = SL(2, \mathbb{Z})$. Furthermore, the interpretation of $\phi_{g,h}$ as a section of $\mathcal{L}_{g,h}^\alpha$ suggests that the following composition law should be imposed

$$(u_1 u_2, k_1 k_2)_\alpha \cdot (g, h) = (u_1, k_1)_\alpha \cdot ((u_2, k_2)_{\alpha^N} \cdot (g, h)) , \quad (\text{C.5})$$

where $(u_1, k_1) \in \text{Mat}_N(\mathbb{Z}) \times G$ and $(u_2, k_2) \in \text{Mat}(\mathbb{Z}) \times G$. In terms of the phases $\epsilon_{g,h}$, this condition reads

$$\epsilon_{g,h}((u_1, k_1) \cdots (u_n, k_n)) = \prod_{i=1}^n \epsilon_{(u_{i+1}, k_{i+1}) \cdots (u_n, k_n) \cdot (g,h)}(u_i, k_i)^{|u_1 \cdots u_{i-1}|}, \quad (\text{C.6})$$

where $|u_i| := \det u_i$.

Observe that $\text{Mat}(\mathbb{Z})$, as a multiplicative semigroup, is generated by $SL(2, \mathbb{Z})$ together with the matrices of the form $\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}$ for p prime. Therefore, it is sufficient to specify the phases $\epsilon(\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}, e)$ for all primes p and any other phase $\epsilon_{g,h}(u, k)$, $(u, k) \in \text{Mat}(\mathbb{Z}) \times G$, is then determined by (C.6). The α -twisted slash operator (2.14) can be trivially extended to an action with respect to $\gamma \in \text{Mat}_N(\mathbb{Z})$, in which case it maps a Jacobi form ϕ of weight 0 and index m to another Jacobi form $\phi|_\alpha(\gamma, k)$ of weight 0 and index Nm . Notice that, by (C.6), the twisted slash operators satisfy the composition relation

$$|_\alpha(u_1 u_2, k_1 k_2) = |_\alpha(u_1, k_1)|_{\alpha^N}(u_2, k_2) \quad (\text{C.7})$$

We can now define the α -twisted equivariant Hecke operators acting on the space of $SL(2, \mathbb{Z}) \times G$ -equivariant Jacobi forms $\phi_{g,h}$ by

$$\mathcal{T}_N^\alpha \phi_{g,h}(\tau, z) := \frac{1}{N} \sum_{u \in SL(2, \mathbb{Z}) \setminus \text{Mat}_N(\mathbb{Z})} \phi_{g,h}(\tau, z)|_\alpha(u, e), \quad (\text{C.8})$$

$$\mathcal{U}_N^\alpha \phi_{g,h}(\tau, z) := \phi_{g,h}(\tau, z)|_\alpha\left(\begin{pmatrix} N & 0 \\ 0 & N \end{pmatrix}, e\right). \quad (\text{C.9})$$

One may check that these operators satisfy the Hecke algebra (3.12)–(3.15). Furthermore, \mathcal{T}_N^α (respectively, \mathcal{U}_N^α) maps the system of α -twisted $SL(2, \mathbb{Z}) \times G$ -equivariant (weak) Jacobi forms of weight 0 and index m to a system of α^N -twisted (respectively, α^{N^2} -twisted) $SL(2, \mathbb{Z}) \times G$ -equivariant (weak) Jacobi forms of weight 0 and index Nm (respectively, index N^2m). The proofs of these properties are completely analogous to the case where α is trivial; in fact, these properties follow directly from the definition of the Hecke operators in terms of slash operators satisfying a composition law of the form (C.7).

C.2 Definition of slash operator for $\text{Mat}(\mathbb{Z}) \times G$

In this subsection, we will describe the extension of the α -twisted slash operators to $\text{Mat}(\mathbb{Z}) \times G$. As stressed in section 3.2, it is sufficient to define such operators for matrices of the form $\begin{pmatrix} N & 0 \\ 0 & 1 \end{pmatrix}$, since, together with $SL(2, \mathbb{Z})$ they generate the whole $\text{Mat}(\mathbb{Z})$. We will make an ansatz for the phase $\epsilon_{g,h}(\begin{pmatrix} N & 0 \\ 0 & 1 \end{pmatrix}, e)$ and verify that it extends consistently to the whole $\text{Mat}(\mathbb{Z})$.

Our ansatz for the phase $\epsilon_{g,h}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e)$ is based on the interpretation of the corresponding slash operator within the theory of symmetric orbifolds of conformal field theories. We recall that the twisted-twining partition function $Z_{g,h}(\tau)$ in a holomorphic CFT is defined as a trace $\text{Tr}_{\mathcal{H}_g}(\rho_g(h)q^{L_0 - \frac{c}{24}})$, where \mathcal{H}_g is the g -twisted sector and ρ_g is the (possibly projective) representation of the centralizer $C_G(g)$ on \mathcal{H}_g . This partition function can be computed by a path integral on a torus $\mathbb{C}/(\mathbb{Z} + \tau\mathbb{Z})$ where the fields are required to have monodromies g and h along the cycles -1 and τ , respectively. The function

$$Z_{g,h}(\tau)|_{\alpha}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e) = \Upsilon_{(\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix})}^* Z_{g,h} , \quad (\text{C.10})$$

is associated with the N -fold covering $\mathbb{C}/(\mathbb{Z} + N\tau\mathbb{Z})$ and has a natural interpretation as a trace

$$Z_{g,h}(\tau)|_{\alpha}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e) = \text{Tr}_{\mathcal{H}_g}((\rho_g(h)q^{L_0 - \frac{c}{24}})^N) . \quad (\text{C.11})$$

Using the product law (2.5) for the projective representation ρ_g , we obtain

$$Z_{g,h}(\tau)|_{\alpha}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e) = \prod_{i=1}^{N-1} c_g(h, h^i) \text{Tr}_{\mathcal{H}_g}(\rho_g(h^N)q^{NL_0 - \frac{Nc}{24}}) = \prod_{i=1}^{N-1} c_g(h, h^i) Z_{g,h^N}(N\tau) . \quad (\text{C.12})$$

This formula suggests the definition

$$\epsilon_{g,h}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e) = \prod_{i=1}^{N-1} c_g(h, h^i) , \quad (\text{C.13})$$

so that

$$f(g, h; \tau)|_{\alpha}((\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}), e) = \prod_{i=1}^{N-1} c_g(h, h^i) f\left((g, h)(\begin{smallmatrix} 1 & 0 \\ 0 & N \end{smallmatrix}); (\begin{smallmatrix} N & 0 \\ 0 & 1 \end{smallmatrix}) \cdot \tau\right) , \quad (\text{C.14})$$

Any integer matrix of determinant N can be written as a product of elements in $SL(2, \mathbb{Z})$ and matrices of the form $(\begin{smallmatrix} p & 0 \\ 0 & 1 \end{smallmatrix})$ for prime p . For each $(u, k) \in \text{Mat}(\mathbb{Z}) \times G$, given a representation of u as a word $u = a_1 \cdots a_n$ in the generators a_1, \dots, a_n of $\text{Mat}(\mathbb{Z})$, one can use the composition law

$$f(g, h; \tau)|_{\alpha}(u_1 u_2, k_1 k_2) = f(g, h; \tau)|_{\alpha}(u_1, k_1) |_{\alpha|u_1|}(u_2, k_2) , \quad (\text{C.15})$$

where $|u| := \det u$, to define the slash operator

$$f(g, h; \tau)|_{\alpha}(a_1 \cdots a_n, k) := f(g, h; \tau)|_{\alpha}(\mathbb{I}, k) |_{\alpha}(a_1, e) |_{\alpha|a_1|}(a_2, e) \cdots |_{\alpha|a_1 \cdots a_{n-1}|}(a_n, e) . \quad (\text{C.16})$$

One needs to check that this definition is consistent, i.e. that the operator $|_\alpha(u, k)$ does not depend on representation of $(u, k) \in \text{Mat}(\mathbb{Z}) \times G$ as a word in the generators and that the composition law is respected. The outline of the proof is given in the next subsection.

In particular, by (C.15), the identity

$$\begin{pmatrix} 1 & 0 \\ 0 & N \end{pmatrix} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} N & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (\text{C.17})$$

yields

$$\epsilon_{g,h}\left(\begin{pmatrix} 1 & 0 \\ 0 & N \end{pmatrix}, e\right) = \frac{c_g(h^{-1}, h)^N \prod_{i=1}^{N-1} c_{h^{-1}}(g, g^i)}{c_{g^N}(h^{-1}, h)} = \frac{1}{\prod_{i=1}^{N-1} c_h(g, g^i)}, \quad (\text{C.18})$$

where in the last step we use (A.19), while from the identity

$$\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & d \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^b \begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \quad (\text{C.19})$$

we obtain the general formula

$$\epsilon_{g,h}\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}, e\right) := \epsilon_{g,h}\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}, e\right) = \frac{\prod_{i=1}^{a-1} c_g(h, h^i)^d}{\prod_{j=1}^{d-1} c_{g^{-b}h^a}(g, g^j) \prod_{k=1}^b c_g(g, g^{-k}h^a)^d}. \quad (\text{C.20})$$

Since for any coset of $SL(2, \mathbb{Z}) \setminus \text{Mat}(\mathbb{Z})$ we can choose a representative of the form $\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}$, this formula is sufficient to determine the Hecke operators \mathcal{T}_N^α and \mathcal{U}_N^α for all N . In particular, if the restriction of the 3-cocycle α to the group $\langle g, h \rangle \cong \mathbb{Z}_{N_1} \times \mathbb{Z}_{N_2}$ is given by $\alpha = \alpha_{v_1} \alpha_{v_2} \alpha_{v_{12}}$, in terms of the generators (A.20)–(A.21), then $(N_1, N_2 > 1)$

$$\epsilon_{g,h}\left(\begin{pmatrix} a & b \\ 0 & d \end{pmatrix}, e\right) = e^{\frac{2\pi i}{N_1} \left(v_1 b \sum_{j=1}^{d-1} \delta^{(N_1)}(j+1) - v_1 d \sum_{j=1}^b \delta^{(N_1)}(1-j) + v_{12} d \sum_{j=1}^{a-1} \delta^{(N_2)}(j+1) \right)}. \quad (\text{C.21})$$

C.3 Proof of consistency

In the rest of this section, we will show that our proposal for the twisted equivariant Hecke operators satisfies several consistency conditions. As a first consistency, let us consider the effect of a ‘gauge transformation’ (A.5), under which the normalized 3-cocycle α is multiplied by a 3-coboundary $\partial\beta$ (with $\beta(e, g) = \beta(g, e) = 1$ to keep the normalization). Formally, this transformation corresponds to a different choice of trivialization of the pull-back $\pi^* \mathcal{L}_{g,h}^\alpha$, where $\pi : \mathcal{P} \times \mathbb{H}_+ \times \mathbb{C} \rightarrow \mathcal{M}$ is the covering map. As stressed in the previous appendix A, for a holomorphic CFTs \mathcal{C} , the transformation (A.5) corresponds to a redefinition (A.15) of the projective representations ρ_g of the centralizer $C_G(g)$ over the

twisted sector \mathcal{H}_g and induces the transformation (A.16) of the partition functions. For the tensor product $\mathcal{C}^{\otimes N}$ and the symmetric product $S^N \mathcal{C}$ theories, the twisted twining partition functions $Z_{g,h}$ are defined traces over the representation induced by the (symmetrized) N -tensor product of ρ_g , so that the appropriate transformation is

$$Z_{g,h}^{(N)} \mapsto \tilde{Z}_{g,h}^{(N)} := \gamma_g(h)^N Z_{g,h}^{(N)} . \quad (\text{C.22})$$

Analogous properties hold for the twisted twining genera $\phi_{g,h}^{(N)}$ in superconformal field theories. Using the identities $\gamma_e(g) = \gamma_g(e) = 1$ and $\gamma_g(h) = \gamma_h(g)^{-1}$, it is easy to check that, under (A.15) and (A.11),

$$(\gamma_g(h)f(g, h; \tau))|_{\tilde{\alpha}}((\begin{smallmatrix} a & b \\ 0 & d \end{smallmatrix})\gamma, e) = \gamma_g(h)^N \left(f(g, h; \tau)|_{\alpha}((\begin{smallmatrix} a & b \\ 0 & d \end{smallmatrix})\gamma, e) \right) , \quad (\text{C.23})$$

where $N = ad$, and similar properties hold for Jacobi forms. As a consequence, we obtain

$$\mathcal{T}_N^\alpha \tilde{Z}_{g,h} = \gamma_g(h)^N \mathcal{T}_N^\alpha Z_{g,h} , \quad (\text{C.24})$$

and

$$\mathcal{T}_N^\alpha \tilde{\phi}_{g,h} = \gamma_g(h)^N \mathcal{T}_N^\alpha \phi_{g,h} , \quad \mathcal{U}_N^\alpha \tilde{\phi}_{g,h} = \gamma_g(h)^{N^2} \mathcal{U}_N^\alpha \phi_{g,h} , \quad (\text{C.25})$$

which indeed reproduce (C.22) and the analogous formula for twisted twining genera. More generally, given *any* representation of a matrix $u \in \text{Mat}_N(\mathbb{Z})$ as a word in the generators of $\text{Mat}(\mathbb{Z})$, it is easy to check that (C.23) holds for the slash operator $|_{\alpha}(u, k)$, i.e

$$\gamma_g(h)\tilde{\epsilon}_{g,h}(u, k) = \gamma_g(h)^N \epsilon_{g,h}(u, k) . \quad (\text{C.26})$$

An easy corollary of this property is

$$f(g, h; \tau)|_{\alpha}(u, k)|_{\alpha^N}(\mathbb{I}, x) = f(g, h; \tau)|_{\alpha}(\mathbb{I}, x)|_{\alpha}(u, k) , \quad (u, k) \in \text{Mat}_N(\mathbb{Z}) \times G, \ x \in G , \quad (\text{C.27})$$

which follows by considering (C.26) for the transformation (A.7). As a consequence of (C.27), one only needs to prove consistency of the slash operator for elements of the form (u, e) .

A set of generators of the multiplicative semigroup $\text{Mat}(\mathbb{Z})$ is given by the generators S, T of $SL(2, \mathbb{Z})$ (see eq.(B.9)), together with all matrices $(\begin{smallmatrix} p & 0 \\ 0 & 1 \end{smallmatrix})$ for p prime. To check the consistency of the definition of the slash operator, one needs to verify that, for each relation $a_1 \cdots a_n = b_1 \cdots b_m$ among the generators of $\text{Mat}(\mathbb{Z})$, the following identities hold for all $(g, h) \in \mathcal{P}_G$

$$f(g, h; \tau)|_{\alpha}(a_1, e) \cdots |_{\alpha^{|a_1 \cdots a_{n-1}|}}(a_n, e) = f(g, h; \tau)|_{\alpha}(b_1, e) \cdots |_{\alpha^{|b_1 \cdots b_{m-1}|}}(b_m, e) . \quad (\text{C.28})$$

This is equivalent to checking the following identities for the phases

$$\prod_{i=1}^n \epsilon_{(g,h)a_{i+1}^\vee \dots a_n^\vee}(a_i, e)^{|a_1 \dots a_{i-1}|} = \prod_{i=1}^m \epsilon_{(g,h)b_{i+1}^\vee \dots b_m^\vee}(b_i, e)^{|b_1 \dots b_{i-1}|} . \quad (\text{C.29})$$

For the relations $S^4 = 1$ and $(ST)^3 = S^2$ within $SL(2, \mathbb{Z})$, these conditions are a consequence of the identities considered in [45, 46]; therefore, we will only consider the independent relations within $\text{Mat}_N(\mathbb{Z})$ for $N > 1$.

For each prime power p , the relations within $\text{Mat}_p(\mathbb{Z})$ can be obtained from the relations for $SL(2, \mathbb{Z})$, together with relations of the form

$$\gamma_1 \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \gamma_2 , \quad (\text{C.30})$$

for suitable $\gamma_1, \gamma_2 \in SL(2, \mathbb{Z})$. It is easy to check that such a relation exists if and only if γ_2 is an element of $\Gamma_0(p)$, which is generated by T , S^2 and $ST^pS = \begin{pmatrix} 1 & 0 \\ -p & 1 \end{pmatrix}$. Therefore, the only independent relations within $\text{Mat}_p(\mathbb{Z})$ are

$$T^p \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} T , \quad (\text{C.31})$$

$$S^2 \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} S^2 , \quad (\text{C.32})$$

and

$$STS \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} ST^pS . \quad (\text{C.33})$$

The only remaining relations in $\text{Mat}(\mathbb{Z})$ are of the form

$$\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \gamma_1 \begin{pmatrix} p' & 0 \\ 0 & 1 \end{pmatrix} = \gamma_2 \begin{pmatrix} p' & 0 \\ 0 & 1 \end{pmatrix} \gamma_3 \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \gamma_4 , \quad (\text{C.34})$$

for all pairs of distinct primes p, p' and suitable $\gamma_1, \gamma_2, \gamma_3, \gamma_4 \in SL(2, \mathbb{Z})$. In fact, it is easier to consider formula (C.20) as a definition of $\epsilon_{g,h}((\begin{smallmatrix} a & b \\ 0 & d \end{smallmatrix}), e)$ and verify the identities (C.29) for the relations

$$\begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \begin{pmatrix} pa & b \\ 0 & d \end{pmatrix} , \quad (\text{C.35})$$

$$\begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} = \begin{pmatrix} pa & pb \\ 0 & d \end{pmatrix} , \quad (\text{C.36})$$

since the consistency conditions for the relations (C.34) then follow. The consistency conditions are therefore

$$\epsilon_{g,h^p} \begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \epsilon_{g,g^{-1}h} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}^p \quad (\text{C.37})$$

$$\epsilon_{h^{-p},g} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \epsilon_{g,h^p} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} = \epsilon_{g^{-1},h^{-1}} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \epsilon_{h^{-1},p} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^p \epsilon_{g,h} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}^p \quad (\text{C.38})$$

$$\epsilon_{g^p,h} \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} = \epsilon_{g,g^{-p}h} \begin{pmatrix} 1 & 0 \\ 0 & p \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} 1 & p \\ 0 & 1 \end{pmatrix}^p \quad (\text{C.39})$$

$$\epsilon_{g,h^p} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix}^{ad} = \epsilon_{g,h} \begin{pmatrix} pa & b \\ 0 & d \end{pmatrix} \quad (\text{C.40})$$

$$\epsilon_{g^d,g^{-b}h^a} \begin{pmatrix} p & 0 \\ 0 & 1 \end{pmatrix} \epsilon_{g,h} \begin{pmatrix} a & b \\ 0 & d \end{pmatrix}^p = \epsilon_{g,h} \begin{pmatrix} pa & pb \\ 0 & d \end{pmatrix} . \quad (\text{C.41})$$

The proof is a tedious but straightforward computation, consisting of a repeated use of the cocycle conditions. For example, the proof of (C.40):

$$\frac{\prod_{i=1}^{a-1} c_g(h, h^{ip})^d}{\prod_{j=1}^{d-1} c_{g^{-b}h^{ap}}(g, g^j) \prod_{k=1}^b c_g(g, g^{-k}h^{ap})^d} \prod_{i=1}^{p-1} c_g(h, h^i)^{ad} \frac{\prod_{j=1}^{d-1} c_{g^{-b}h^{ap}}(g, g^j) \prod_{k=1}^b c_g(g, g^{-k}h^{ap})^d}{\prod_{i=1}^{pa-1} c_g(h, h^i)^d} \quad (\text{C.42})$$

$$= \left(\frac{\prod_{i=1}^{a-1} c_g(h, h^{ip}) \prod_{i=1}^{p-1} c_g(h, h^i)^a}{\prod_{i=1}^{pa-1} c_g(h, h^i)} \right)^d = 1 \quad (\text{C.43})$$

where the last equality follows from the 2-cocycle condition on c_g (the easiest way to check this is to consider a projective representation ρ_g associated with c_g , and to rewrite this expression as a ratio of two phases, both corresponding to the mismatch between $\rho_g(h)^{pa}$ and $\rho_g(h^{pa})$).

D Some technical proofs

D.1 Proof of (3.53).

Let us prove (3.53). The sum $F_{g,h}(a, d, k, \ell)$ can be written as

$$F_{g,h}(a, d, k, \ell) := \sum_{n=0}^{\infty} \frac{1}{d} \sum_{b=0}^{d-1} e^{\frac{2\pi i b n}{N\lambda d}} c_{g_{\alpha}^d, g_{\alpha}^{-b}h_{\alpha}^k} \left(\frac{n}{N\lambda}, \ell \right) q^{\frac{na}{N\lambda d}} , \quad (\text{D.1})$$

in terms of the Fourier coefficients of the twisted-twining genera $\phi_{g_{\alpha}^d, g_{\alpha}^{-b}h_{\alpha}^k}$ introduced in section 3.2.2. Notice that the functions $\phi_{g_{\alpha}, h_{\alpha}}$ satisfy

$$\phi_{g_{\alpha}^r, g_{\alpha}^{-r}h_{\alpha}}(\tau + 1, z) = \phi_{g_{\alpha}, h_{\alpha}}(\tau, z) , \quad (\text{D.2})$$

so that

$$c_{g_{\alpha}^d, g_{\alpha}^d h_{\alpha}}(r, \ell) = e^{\frac{2\pi i r}{N\lambda}} c_{g_{\alpha}, h_{\alpha}}(r, \ell) , \quad (\text{D.3})$$

for all $h_\alpha \in C_{M_{24}}^\alpha(g)$. Set

$$e := \gcd(d, N\lambda) , \quad f := \frac{d}{e} , \quad (\text{D.4})$$

and note that

$$\gcd(f, \frac{N\lambda}{e}) = 1 , \quad (\text{D.5})$$

otherwise $e \gcd(f, \frac{N\lambda}{e})$ would be a common divisor of d and $N\lambda$ greater than e . The order of g_α^d is $o(g_\alpha^{ef}) = o(g_\alpha^e) = N\lambda/e$, so that $c_{g_\alpha^d, g_\alpha^{-b} h_\alpha^k}(\frac{n}{N\lambda}, \ell) = 0$ unless $e|n$. Thus, we can set $r := n/e$ and obtain

$$F_{g,h}(a, d, k, \ell) = \sum_{r=0}^{\infty} \frac{1}{d} \sum_{b=0}^{d-1} e^{\frac{2\pi i b r e}{N\lambda d}} c_{g_\alpha^d, g_\alpha^{-b} h_\alpha^k}(\frac{re}{N\lambda}, \ell) q^{\frac{ar}{N\lambda f}} . \quad (\text{D.6})$$

Notice that, by (D.3), the general term of the sum over b is periodic under $b \rightarrow b + d$. We can set $b := se + b'$ and replace the sum over $b \in \mathbb{Z}/d\mathbb{Z}$ by a sum over $b' \in \mathbb{Z}/e\mathbb{Z}$ and $s \in \mathbb{Z}/f\mathbb{Z}$

$$F_{g,h}(a, d, k, \ell) = \sum_{r=0}^{\infty} \frac{1}{e} \sum_{b'=0}^{e-1} \frac{1}{f} \sum_{s=0}^{f-1} e^{\frac{2\pi i (b'+se)r}{N\lambda f}} c_{g_\alpha^d, g_\alpha^{-b'} g_\alpha^{-se} h_\alpha^k}(\frac{re}{N\lambda}, \ell) q^{\frac{ar}{N\lambda f}} . \quad (\text{D.7})$$

Thus, the general term of the sum over b' is periodic under $b' \rightarrow b' + e$ and we can sum over $N\lambda/e$ periods and divide by $N\lambda/e$

$$F_{g,h}(a, d, k, \ell) = \sum_{r=0}^{\infty} \frac{1}{N\lambda} \sum_{b'=0}^{N\lambda-1} \frac{1}{f} \sum_{s=0}^{f-1} e^{\frac{2\pi i (b'+se)r}{N\lambda f}} c_{g_\alpha^d, g_\alpha^{-b'} g_\alpha^{-se} h_\alpha^k}(\frac{re}{N\lambda}, \ell) q^{\frac{ar}{N\lambda f}} . \quad (\text{D.8})$$

By (D.5), there are integers x, y such that

$$xf + y\frac{N\lambda}{e} = 1 \quad (\text{D.9})$$

so that $se = sxd + syN\lambda$ and we obtain

$$F_{g,h}(a, d, k, \ell) = \sum_{r=0}^{\infty} \frac{1}{N\lambda} \sum_{b'=0}^{N\lambda-1} \frac{1}{f} \sum_{s=0}^{f-1} e^{\frac{2\pi i b' r}{N\lambda f}} e^{\frac{2\pi i s (xd+yN\lambda)r}{N\lambda f}} c_{g_\alpha^d, g_\alpha^{-b'} g_\alpha^{-sdx} h_\alpha^k}(\frac{re}{N\lambda}, \ell) q^{\frac{ar}{N\lambda f}} \quad (\text{D.10})$$

$$= \sum_{r=0}^{\infty} \sum_{b'=0}^{N\lambda-1} \frac{e^{\frac{2\pi i b' r}{N\lambda f}}}{N\lambda} \sum_{s=0}^{f-1} \frac{e^{\frac{2\pi i s y r}{f}}}{f} c_{g_\alpha^d, g_\alpha^{-b'} h_\alpha^k}(\frac{re}{N\lambda}, \ell) q^{\frac{ar}{N\lambda f}} , \quad (\text{D.11})$$

where in the last step we used (D.3). Since by (D.9) $\gcd(y, f) = 1$, the sum over s just a projection

$$\sum_{s=0}^{f-1} \frac{e^{\frac{2\pi i s y r}{f}}}{f} = \begin{cases} 1 & \text{if } f|r , \\ 0 & \text{otherwise} , \end{cases} \quad (\text{D.12})$$

so that, by setting $r = mf$, we obtain

$$F_{g,h}(a, d, k, \ell) = \sum_{m=0}^{\infty} \sum_{b'=0}^{N\lambda-1} \frac{e^{\frac{2\pi i b' m}{N\lambda}}}{N\lambda} c_{g_{\alpha}^d, g_{\alpha}^{-b'} h_{\alpha}^k} \left(\frac{md}{N\lambda}, \ell \right) q^{\frac{am}{N\lambda}}, \quad (\text{D.13})$$

that is equivalent to (3.53).

D.2 Central extensions and special choices of the cocycle

In section 3.2.2 a central extension $C_{M_{24}}^{\alpha}(g)$ of the centralizer $C_{M_{24}}(g)$ is defined, together with the representations $\tilde{\rho}_{g,r}$ on the g^r -twisted sector \mathcal{H}_{g^r} , for all $r \in \mathbb{Z}_{\geq 0}$. In this section, we will describe some special choices of the cocycle α , for which these representations are particularly simple.

If two 3-cocycles α, α' differ by a coboundary $\partial\beta$, then there is an isomorphism $C_{M_{24}}^{\alpha'}(g) \xrightarrow{\cong} C_{M_{24}}^{\alpha}(g)$ that relates the canonical lifts

$$h_{\alpha'} \mapsto h_{\alpha} q(\nu_g(h)), \quad (\text{D.14})$$

where $\nu_g(h) \in \mathbb{R}/\mathbb{Z}$ is defined by $e^{2\pi i \nu_g(h)} = \frac{\beta(g, h)}{\beta(h, g)}$.

Under the shift $\alpha \rightarrow \alpha'$ by a coboundary $\partial\beta$, the cocycle c_g transforms as

$$c_g(h, k) \rightarrow \frac{\beta(g, h)}{\beta(h, g)} \frac{\beta(g, k)}{\beta(k, g)} \frac{\beta(hk, g)}{\beta(g, hk)} c_g(h, k), \quad h, k \in C_{M_{24}}(g). \quad (\text{D.15})$$

Correspondingly, the phases $f_{g,r}(h)$ transform as

$$f_{g,r}(h) \rightarrow f_{g,r}(h) \frac{\beta(h, g)^r}{\beta(g, h)^r} \frac{\beta(g^r, h)}{\beta(h, g^r)}, \quad h \in C_{M_{24}}(g), \quad (\text{D.16})$$

and, in particular,

$$f_{g,N}(h) \rightarrow f_{g,N}(h) \frac{\beta(h, g)^N}{\beta(g, h)^N}, \quad h \in C_{M_{24}}(g). \quad (\text{D.17})$$

Therefore, we can choose $\beta(g, h)/\beta(h, g)$ for each $h \in C_{M_{24}}(g)$, $h \notin \langle g \rangle$, in such a way that

$$f_{g,N}(h) = 1, \quad \forall h \in C_{M_{24}}(g) \setminus \langle g \rangle. \quad (\text{D.18})$$

This condition determines $\beta(g, h)/\beta(h, g)$ for all $h \in C_{M_{24}}(g) \setminus \langle g \rangle$ up to N -th roots of unity. Notice that, in general,

$$\tilde{\rho}_{g,r+N}(h) = \frac{\rho_{g^r}(h)}{f_{g,r}(h) f_{g,N}(h)} = \frac{\tilde{\rho}_{g^r}(h)}{f_{g,N}(h)}, \quad h \in C_{M_{24}}(g), \quad (\text{D.19})$$

so that, by imposing (D.18), we have

$$\tilde{\rho}_{g,r+N}(h) = \tilde{\rho}_{g^r}(h) , \quad h \in C_{M_{24}}(g) \setminus \langle g \rangle . \quad (\text{D.20})$$

On the other hand, $f_{g,N}(g)$ depends only on the cohomology class $[\alpha]$. Since the restriction of $[\alpha]^N$ to $H^3(\langle g \rangle, U(1)) \cong \mathbb{Z}_N$ is the trivial class, it follows that $f_{g,N}(g)$ must be a N -th root of unity. In fact, for M_{24} , we have

$$f_{g,N}(g) = e^{-\frac{2\pi i}{\lambda}} , \quad (\text{D.21})$$

where $\lambda|N$ is the length of the shortest cycle of g in the 24-dimensional permutation representation. It is also related to the spectrum of $L_0 - \frac{c}{24}$ in the g -twisted sector, which takes values in $-\frac{1}{\lambda N} + \frac{1}{N}\mathbb{Z}$ and to the presence of a non-trivial multiplier system for the twining genus $\phi_{e,g}$. It is useful to distinguish the cases when $\lambda = 1$ and $\lambda \neq 1$.

D.2.1 Case $\lambda = 1$ (trivial multiplier)

This case occurs whenever g is an element of some M_{23} subgroup of M_{24} , i.e. when g belongs to the classes

$$2A, 3A, 4B, 5A, 6A, 7A, 7B, 8A, 11A, 14A, 14B, 15A, 15B, 23A, 23B . \quad (\text{D.22})$$

In this case the restriction of $[\alpha]$ to $H^3(\langle g \rangle, U(1))$ is the trivial class and one can choose $\beta(g^i, g^j)$ in such a way that

$$c_{g^i}(g^j, g^k) = 1 . \quad (\text{D.23})$$

In particular,

$$f_{g,r}(g^i) = 1 , \quad (\text{D.24})$$

for all r, i . By specializing (3.31) to the case $r = N$, we obtain

$$c_g(h, k)^N = c_{g^N}(h, k) \frac{f_{g,N}(hk)}{f_{g,N}(h)f_{g,N}(k)} = 1 , \quad h, k \in C(g) , \quad (\text{D.25})$$

so that $c_g(h, k)$ is an N -th root of unity for all $h, k \in C_{M_{24}}(g)$, i.e. there is $\mu_g(h, k) \in \mathbb{Z}/N\mathbb{Z}$ such that

$$c_g(h, k) = e^{\frac{2\pi i \mu_g(h, k)}{N}} . \quad (\text{D.26})$$

With this choice of cocycle, the central extension $C_{M_{24}}^\alpha(g)$ of section 3.2.2 can be chosen to be finite

$$1 \rightarrow \langle Q \rangle \cong \mathbb{Z}_N \rightarrow C_{M_{24}}^\alpha(g) \rightarrow C_{M_{24}}(g) \rightarrow 1 . \quad (\text{D.27})$$

where the central element Q is related to the $U(1)$ -generator $q(x)$ in section 3.2.2 by

$$Q = q(1/N) . \quad (\text{D.28})$$

D.2.2 Case $\lambda \neq 1$ (non-trivial multiplier)

This case occurs when g is in one of the M_{24} -classes

$$2B, 3B, 4A, 4C, 6B, 10A, 12A, 12B, 21A, 21B . \quad (\text{D.29})$$

Since $H^3(\langle g \rangle, U(1)) \cong \mathbb{Z}_N$, we can choose the cocycle in such a way that

$$c_{g^i}(g^j, g^k)^N = 1 , \quad (\text{D.30})$$

for all $i, j, k \in \mathbb{Z}$. This condition fixes $\beta(g^i, g^j)$ up to N -th roots of unity. For $h, k \in C_{M_{24}}(g)$, with $h, k, hk \notin \langle g \rangle$, we can apply again (3.31) with $r = N$ and obtain

$$c_g(h, k)^N = 1 , \quad h, k, hk \in C_{M_{24}}(g) \setminus \langle g \rangle . \quad (\text{D.31})$$

If α satisfies (D.18) and (D.30), then by (3.31) we have

$$f_{g,N}(g^i) c_g(g^i, h)^N = 1 , \quad \frac{f_{g,N}(g^i) f_{g,N}(g^j)}{f_{g,N}(g^{i+j})} = 1 , \quad (\text{D.32})$$

so that

$$f_{g,N}(g^n) = f_{g,N}(g)^n = e^{-\frac{2\pi i n}{\lambda}} , \quad (\text{D.33})$$

and

$$c_g(g^i, h)^{N\lambda} = 1 . \quad (\text{D.34})$$

Therefore, $c_g(h, k)$ is a $N\lambda$ roots of unity for all $h, k \in C_{M_{24}}(g)$ and we can define $\mu_g(h, k) \in \mathbb{Z}/N\lambda\mathbb{Z}$ such that

$$c_g(h, k) = e^{\frac{2\pi i \mu_g(h, k)}{N\lambda}} . \quad (\text{D.35})$$

The discussion is similar to the case $\lambda = 1$. With this choice of cocycle, the central extension $C_{M_{24}}^\alpha(g)$ of section 3.2.2 can be chosen to be finite

$$1 \rightarrow \langle q(\frac{1}{N\lambda}) \rangle \cong \mathbb{Z}_{N\lambda} \rightarrow C_{M_{24}}^\alpha(g) \rightarrow C_{M_{24}}(g) \rightarrow 1 . \quad (\text{D.36})$$

It is convenient to define also a central element Q of order N by

$$Q = q(1/N) , \quad (\text{D.37})$$

so that the g^r -twisted sector is an eigenspace of Q with eigenvalue $e^{\frac{2\pi i r}{N}}$.

D.3 S-duality for g in classes $2B, 3B, 4C, 6B, 12B$

Here we prove the S-duality transformation property of $\Phi_{g,h}$ for g in the classes $2B, 3B, 4C, 6B, 12B$. Let g be an element of M_{24} in one of these classes and let N be its order. In all these cases, the restriction of the class $[\alpha]$ to $H^3(\langle g \rangle, U(1))$ is non-trivial and has order $\lambda = N$. Therefore,

$$\hat{c}_{g,h}(d, m, \ell, t) = \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b m}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^d}(\frac{md}{N^2}, \ell)} (\tilde{\rho}_{g,d}(g)^{-b} \tilde{\rho}_{g,d}(h)^k (-1)^{F+\bar{F}}) . \quad (\text{D.38})$$

In the g^r -twisted sector, for r and N coprime, the spectrum of $L_0 - \frac{c}{24}$ takes values in $-\frac{1}{N^2} + \frac{1}{N}\mathbb{Z}$. More generally, if $\gcd(r, N) = e$, then in the g^r -twisted sector the spectrum of $L_0 - \frac{c}{24}$ takes values in $-\frac{1}{(N/e)^2} + \frac{1}{N/e}\mathbb{Z}$. Notice that $\hat{c}_{g,h}(d, m, \ell, t)$ is a sum of traces in the g^d -twisted sector at level $\frac{md}{N^2}$, so that $\hat{c}_{g,h}(d, m, \ell, t) = 0$ unless $\frac{md}{N^2} \in -\frac{1}{(N/e)^2} + \frac{1}{N/e}\mathbb{Z}$, where $e = \gcd(N, d)$. It is easy to verify that this condition is equivalent to

$$m \equiv -d \pmod{N} , \quad (\text{D.39})$$

where we used the property

$$\gcd(x, N) = 1 \quad \Leftrightarrow \quad x^2 \equiv 1 \pmod{N} , \quad (\text{D.40})$$

which holds for all N that divide 24.

We can choose the cocycle α in such a way that

$$f_{g,N}(h) \equiv \prod_{i=1}^{N-1} c_h(g, g^i) = 1 , \quad h \in C_{M_{24}}(g) \setminus \langle g \rangle , \quad (\text{D.41})$$

while $f_{g,N}(g^k)$ depends only on the cohomology class $[\alpha]$ and equals

$$f_{g,N}(g^k) = e^{-\frac{2\pi i k}{N}} . \quad (\text{D.42})$$

With this choice of cocycle, we have

$$\tilde{\rho}_{g,x+N}(h) = \frac{\rho_{g^x}(h)}{f_{g,N}(h)} = \tilde{\rho}_{g,x}(h) , \quad h \in C_{M_{24}}(g) \setminus \langle g \rangle , \quad (\text{D.43})$$

while

$$\tilde{\rho}_{g,x+N}(g) = \frac{\tilde{\rho}_{g,x}(g)}{f_{g,N}(g)} = e^{\frac{2\pi i}{N}} \tilde{\rho}_{g,x}(g) . \quad (\text{D.44})$$

For $d \equiv -m \equiv 0 \pmod{N}$, we have

$$\hat{c}_{g,h}(d = Nu, m = Nv, \ell, t) = \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b v}{N}}}{N^2} \text{Tr}_{\mathcal{H}_e(uv, \ell)} (\tilde{\rho}_{g, Nu}(g)^{-b} \tilde{\rho}_{g, Nu}(h)^k (-1)^{F+\bar{F}}) \quad (\text{D.45})$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(v-u)}{N}}}{N^2} \text{Tr}_{\mathcal{H}_e(uv, \ell)} (g^{-b} h^k (-1)^{F+\bar{F}}) . \quad (\text{D.46})$$

Since the (untwisted) twining genera are invariant under charge conjugation, i.e.

$$\phi_{e, g^{-b} h^k}(\tau, z) = \phi_{e, g^b h^{-k}}(\tau, z) , \quad (\text{D.47})$$

we obtain

$$\begin{aligned} \hat{c}_{g,h}(Nu, Nv, \ell, t) &= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(v-u)}{N}}}{N^2} \text{Tr}_{\mathcal{H}_e(uv, \ell)} (g^b h^{-k} (-1)^{F+\bar{F}}) \\ &= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(u-v)}{N}}}{N^2} \text{Tr}_{\mathcal{H}_e(uv, \ell)} (g^{-b} h^{-k} (-1)^{F+\bar{F}}) = \hat{c}_{g, h^{-1}}(Nv, Nu, \ell, t) . \end{aligned} \quad (\text{D.48})$$

Similarly, for $d \equiv x \equiv -m \pmod{N}$, with $x = 1, \dots, N-1$, we obtain

$$\begin{aligned} \hat{c}_{g,h}(d = Nu + x, m = Nv + N - x, \ell, t) &= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(Nv + N - x - Nu)}{N^2}}}{N^2} \\ &\quad \text{Tr}_{\mathcal{H}_{g^x}(\frac{md}{16}, \ell)} (\tilde{\rho}_{g,x}(g)^{-b} \tilde{\rho}_{g,x}(h)^k (-1)^{F+\bar{F}}) . \end{aligned} \quad (\text{D.49})$$

Using the relation

$$\phi_{g,h}(\tau, z) = \frac{1}{c_h(g, g^{-1}) c_{g^{-1}}(h, h^{-1})} \phi_{g^{-1}, h^{-1}}(\tau, z) , \quad (\text{D.50})$$

and by

$$\frac{\rho_{g^{-1}}(h^{-1})}{c_{g^{-1}}(h, h^{-1})} = \rho_{g^{-1}}(h)^{-1} , \quad (\text{D.51})$$

we obtain

$$\text{Tr}_{H_{g^x}(n, \ell)} (\tilde{\rho}_{g,x}(h)) = \frac{\text{Tr}_{\mathcal{H}_{g^x}(n, \ell)} (\rho_{g^x}(h))}{\prod_{i=1}^{x-1} c_h(g, g^i)} = \frac{\text{Tr}_{\mathcal{H}_{g^{-x}}(n, \ell)} (\rho_{g^{-x}}(h)^{-1})}{c_h(g^x, g^{-x}) \prod_{i=1}^{x-1} c_h(g, g^i)} \quad (\text{D.52})$$

$$= \frac{\text{Tr}_{\mathcal{H}_{g^{N-x}}(n, \ell)} (\tilde{\rho}_{g, N-x}(h)^{-1})}{c_h(g^x, g^{N-x}) \prod_{i=1}^{x-1} c_h(g, g^i) \prod_{j=1}^{N-x-1} c_h(g, g^j)} = \frac{\text{Tr}_{\mathcal{H}_{g^{N-x}}(n, \ell)} (\tilde{\rho}_{g, N-x}(h)^{-1})}{f_{g, N}(h)} , \quad (\text{D.53})$$

where the last equality follows from

$$c_h(g^x, g^{N-x}) \prod_{i=1}^{x-1} c_h(g, g^i) = \prod_{i=N-x}^N c_h(g, g^i) , \quad (\text{D.54})$$

that in turn is a consequence of the 2-cocycle condition for c_g . It is easy to verify that this identity is compatible with the product in the central extension, i.e.

$$\text{Tr}_{H_{g^x}(n, \ell)}(\tilde{\rho}_{g,x}(h) \tilde{\rho}_{g,x}(k)) = \frac{\text{Tr}_{\mathcal{H}_{g^{N-x}}(n, \ell)}(\tilde{\rho}_{g, N-x}(k)^{-1} \tilde{\rho}_{g, N-x}(h)^{-1})}{f_{g, N}(k) f_{g, N}(h)} . \quad (\text{D.55})$$

Using this relation, we obtain

$$\hat{c}_{g,h}(d = Nu + x, m = Nv + N - x, \ell, t) \quad (\text{D.56})$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(Nv-x-Nu)}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^{N-x}}(\frac{md}{N^2}, \ell)}(\tilde{\rho}_{g, N-x}(g)^b \tilde{\rho}_{g, N-x}(h)^{-k} (-1)^{F+\bar{F}}) \quad (\text{D.57})$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(Nu+x-Nv)}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^{N-x}}(\frac{md}{N^2}, \ell)}(\tilde{\rho}_{g, N-x}(g)^{-b} \tilde{\rho}_{g, N-x}(h)^{-k} (-1)^{F+\bar{F}}) \quad (\text{D.58})$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b(Nu+x)}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^{N-x}}(\frac{md}{N^2}, \ell)}(\tilde{\rho}_{g, Nv+N-x}(g)^{-b} \tilde{\rho}_{g, Nv+N-x}(h)^{-k} (-1)^{F+\bar{F}}) \quad (\text{D.59})$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b d}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^m}(\frac{md}{N^2}, \ell)}(\tilde{\rho}_{g, m}(g)^{-b} \tilde{\rho}_{g, m}(h)^{-k} (-1)^{F+\bar{F}}) . \quad (\text{D.60})$$

Finally, we notice that

$$\tilde{\rho}_{g,r}(h)^{-k} = c_g(h, h^{-1})^{-rk} \tilde{\rho}_{g,r}(h^{-1})^k , \quad (\text{D.61})$$

where $c_g(h, h^{-1})$ is a N -th root of unity thanks to our choice of cocycle. We recall that imposing the condition $f_{g,N}(h) = 1$ still leaves the possibility of modifying the cocycle α by a coboundary $\partial\beta$ with $\beta^N = 1$. Under such a modification, $c_g(h, h^{-1})$ transforms as

$$c_g(h, h^{-1}) \rightarrow \frac{\beta(g, h)}{\beta(h, g)} \frac{\beta(g, h^{-1})}{\beta(h^{-1}, g)} c_g(h, h^{-1}) , \quad (\text{D.62})$$

We distinguish two cases. If h^{-1} and h are not conjugated within $C_{M_{24}}(g)$, then one can choose the cocycle α in such a way that

$$c_g(h, h^{-1}) = 1 . \quad (\text{D.63})$$

With this choice, we obtain

$$\hat{c}_{g,h}(d, m, \ell, t) = \hat{c}_{g,h^{-1}}(m, d, \ell, t) , \quad (\text{D.64})$$

so that

$$\Phi_{g,h}(\sigma, \tau, z) = pq^{\frac{1}{N^2}} y \prod_{t=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h}(d,m,\ell,t)} \quad (\text{D.65})$$

$$= pq^{\frac{1}{N^2}} y \prod_{t=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h^{-1}}(m,d,\ell,t)} \quad (\text{D.66})$$

$$= \Phi_{g,h^{-1}}\left(\frac{\tau}{N^2}, N^2\sigma, z\right) . \quad (\text{D.67})$$

In most cases, however, there is some $w \in C_{M_{24}}(g)$ such that $h^{-1} = w^{-1}hw$. This property implies a ‘reality condition’ on the Fourier coefficients of $\phi_{g,h}(\tau, z)$, namely

$$\phi_{g,h}(\tau, z) = \frac{c_g(h, w)c_g(h, h^{-1})}{c_g(w, h^{-1})} \phi_{g,h}^*(\tau, z) = e^{-\frac{2\pi i n}{N}} \phi_{g,h}^*(\tau, z) , \quad (\text{D.68})$$

where

$$\phi_{g,h}^*(\tau, z) := \overline{\phi_{g,h}(-\bar{\tau}, -\bar{z})} = \sum_{r \in \mathbb{Q}} \sum_{\ell \in \mathbb{Z}} \overline{c_{g,h}(r, \ell)} q^r y^\ell , \quad (\text{D.69})$$

is the Jacobi form with complex conjugate Fourier coefficients. Here, $n \in \mathbb{Z}/N\mathbb{Z}$ is such that

$$e^{-\frac{2\pi i n}{N}} = \frac{c_g(h, w)c_g(h, h^{-1})}{c_g(w, h^{-1})} , \quad (\text{D.70})$$

where we used the fact that, with our choice of cocycle, the right-hand side is a N -th root of unity. Thus, (D.68) implies that all Fourier coefficients of $\phi_{g,h}$ have the same argument. At the level of the central extension $C_{M_{24}}^\alpha(g)$, the relation $hw = wh^{-1}$ leads to a relation among the lifts

$$h_\alpha^{-1} = \frac{c_g(w, h^{-1})}{c_g(h, w)c_g(h, h^{-1})} w_\alpha^{-1} h_\alpha w_\alpha = w_\alpha^{-1} Q^n h_\alpha w_\alpha . \quad (\text{D.71})$$

Since $Q^n h_\alpha$ is conjugated with h_α^{-1} within $C_{M_{24}}^\alpha(g)$, it must have the same order M , so that $nM \equiv 0 \pmod{N}$ or, equivalently,

$$nM = sN , \quad (\text{D.72})$$

for some $s \in \mathbb{Z}/M\mathbb{Z}$. Therefore,

$$\hat{c}_{g,h}(d, m, \ell, t) \quad (D.73)$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b d}{N^2}}}{N^2} \text{Tr}_{\mathcal{H}_{g^m}(\frac{md}{N^2}, \ell)} (\tilde{\rho}_{g,m}(g)^{-b} \tilde{\rho}_{g,m}(h)^k \tilde{\rho}_{g,m}(Q^n)^k (-1)^{F+\bar{F}}) \quad (D.74)$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i t k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b d}{N^2}}}{N^2} e^{\frac{2\pi i n m k}{N}} \text{Tr}_{\mathcal{H}_{g^m}(\frac{md}{N^2}, \ell)} (\tilde{\rho}_{g,m}(g)^{-b} \tilde{\rho}_{g,m}(h)^k (-1)^{F+\bar{F}}) \quad (D.75)$$

$$= \sum_{k=0}^{M-1} \frac{e^{-\frac{2\pi i (t-sm) k}{M}}}{M} \sum_{b=1}^{N^2} \frac{e^{\frac{2\pi i b d}{N^2}}}{N^2} e^{\frac{2\pi i n m k}{N^2}} \text{Tr}_{\mathcal{H}_{g^m}(\frac{md}{N^2}, \ell)} (\tilde{\rho}_{g,m}(g)^{-b} \tilde{\rho}_{g,m}(h)^k (-1)^{F+\bar{F}}) \quad (D.76)$$

$$= \hat{c}_{g,h}(m, d, \ell, t - sm) . \quad (D.77)$$

Using this identity, we obtain

$$\Phi_{g,h}(\sigma, \tau, z) = pq^{\frac{1}{N^2}} y \prod_{t=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h}(d,m,\ell,t)} \quad (D.78)$$

$$= pq^{\frac{1}{N^2}} y \prod_{t=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i t}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t-sm)} \quad (D.79)$$

$$= pq^{\frac{1}{N^2}} y \prod_{t'=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i sm}{M}} e^{\frac{2\pi i t'}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t')} \quad (D.80)$$

$$= pq^{\frac{1}{N^2}} y \prod_{t'=1}^M \prod_{(d,m,\ell)} (1 - e^{\frac{2\pi i n m}{N}} e^{\frac{2\pi i t'}{M}} q^{\frac{m}{N^2}} y^\ell p^d)^{\hat{c}_{g,h}(m,d,\ell,t')} \quad (D.81)$$

$$= e^{-\frac{2\pi i n}{N}} \Phi_{g,h}\left(\frac{\tau}{N^2} + \frac{n}{N}, N^2\sigma, z\right) \quad (D.82)$$

$$= e^{\frac{2\pi i n}{N}} \Phi_{g,h}\left(\frac{\tau}{N^2}, N^2\sigma - Nn, z\right) , \quad (D.83)$$

where we used the property that the exponent $\hat{c}_{g,h}(m, d, \ell, t')$ is non-zero only for $d \equiv -m \pmod{N^2}$, so that

$$e^{-\frac{2\pi i x}{N^2}} \Phi_{g,h}\left(\sigma + \frac{x}{N^2}, \tau, z\right) = e^{\frac{2\pi i x}{N^2}} \Phi_{g,h}(\sigma, \tau - x, z) , \quad x \in \mathbb{Z} . \quad (D.84)$$

Let us set $\nu = \frac{r}{N^2}$, for some $r \in \mathbb{Z}$, so that

$$\Phi'_{g,h}(\sigma, \tau, z) = e^{-\frac{2\pi i r}{N^2}} \Phi_{g,h}\left(\sigma + \frac{r}{N^2}, \tau, z\right) = e^{\frac{2\pi i (Nn-r)}{N^2}} \Phi_{g,h}\left(\frac{\tau}{N^2}, N^2\sigma + r - Nn, z\right) \quad (D.85)$$

$$= e^{\frac{2\pi i (r-Nn)}{N^2}} \Phi_{g,h}\left(\frac{\tau}{N^2} + \frac{Nn-r}{N^2}, N^2\sigma, z\right) = e^{\frac{2\pi i (2r-Nn)}{N^2}} \Phi'_{g,h}\left(\frac{\tau}{N^2} + \frac{Nn-2r}{N^2}, N^2\sigma, z\right) . \quad (D.86)$$

By choosing $r = \frac{n(N+N^2)}{2}$, we finally obtain

$$\Phi'_{g,h}(\sigma, \tau, z) = \Phi'_{g,h}\left(\frac{\tau}{N^2} - n, N^2\sigma, z\right) = \Phi'_{g,h}\left(\frac{\tau}{N^2}, N^2\sigma, z\right). \quad (\text{D.87})$$

E Tables

In the following tables, we collect information about the 34 non-cyclic abelian subgroups $\langle g, h \rangle$ of M_{24} and the corresponding twisted-twining genera. There we have described their structure as an abelian group, i.e. as $\mathbb{Z}_m \times \mathbb{Z}_n$, the M_{24} classes of all its elements (excluding the identity), the orders of the centralizer $C(g, h)$ and its index $|N(g, h)|/|C(g, h)|$ in the normalizer of $\langle g, h \rangle$ in M_{24} , and the lengths of the orbits of $\langle g, h \rangle \subset M_{24}$ when acting as a group of permutations of 24 objects.

Furthermore, we give a classification of all modular groups $\Gamma_{g,h} \subset PSL(2, \mathbb{Z})$ and twisted twining genera $\phi_{g,h}$ for commuting pairs $g, h \in M_{24}$.

Group 27 is the only case where the pairs (g, h) and (g^{-1}, h^{-1}) are not conjugated within M_{24} . Thus, in this case, charge conjugation gives the identities $\phi_{2B,8A_1} = \phi_{2B,8A_2}$, $\phi_{8A,2B_1} = \phi_{8A,2B_2}$, and so on, and the respective functions are denoted in the following tables by $\phi_{2B,8A_{1,2}}$, $\phi_{8A,2B_{1,2}}$, etcetera.

Most of the modular groups $\Gamma_{g,h}$ are of the form $\Gamma(1) = SL(2, \mathbb{Z})$ or

$$\Gamma_0(N) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid c \equiv 0 \pmod{N} \right\}, \quad (\text{E.1})$$

or conjugates of $\Gamma_0(N)$ in $SL(2, \mathbb{Z})$. The exceptions are the group in case 32, where

$$\Gamma_{2B,10A} = \bigcup_{i \in \mathbb{Z}/3\mathbb{Z}, j \in \mathbb{Z}/4\mathbb{Z}} \begin{pmatrix} 1 & 1 \\ -5 & -4 \end{pmatrix}^i \begin{pmatrix} -3 & -1 \\ 10 & 3 \end{pmatrix}^j \Gamma_{2,10}, \quad (\text{E.2})$$

is a subgroup of index 12 in $SL(2, \mathbb{Z})$ and

$$\Gamma_{2,10} := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid a \equiv 1, b \equiv 0 \pmod{2}, c \equiv 0, d \equiv 1 \pmod{10} \right\}, \quad (\text{E.3})$$

is the group of elements $\gamma \in SL(2, \mathbb{Z})$ such that $(g, h) \cdot \gamma = (g, h)$; the group in case 12, with

$$\Gamma_{2A,4A} = \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid b \equiv 0 \pmod{2}, c \equiv 0 \pmod{4} \right\}, \quad (\text{E.4})$$

which is a conjugate of $\Gamma_0(8)$ in $SL(2, \mathbb{R})$; and the group in case 22, with

$$\Gamma_{4A,4C} = \left\langle \begin{pmatrix} -1 & 1 \\ -2 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 0 \\ 4 & 1 \end{pmatrix}, \begin{pmatrix} 3 & -2 \\ -4 & 3 \end{pmatrix} \right\rangle. \quad (\text{E.5})$$

Finally, for each of the 34 groups we provide the explicit expressions of the twisted-twining genera $\phi_{g,h}$ and $\mathcal{T}_N^\alpha \phi_{g,h}$, $N = 2, 3, 4$, for a pair (g, h) of generators and for a choice of cocycle α satisfying (D.18) and (D.23) (when g is in $M_{23} \subset M_{24}$) or (D.30) (otherwise).

#	Structure	Elements	$ C(g, h) $	$\frac{ N(g, h) }{ C(g, h) }$	Orbits on 24	Max subgr.	Functions	$\Gamma_{g, h}$
1.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)^3$	1536	6	$1^8 \cdot 4^4$		$\phi_{2A, 2A_2}$	$\Gamma(1)$
2.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)^3$	1536	6	2^{12}		$\phi_{2A, 2A_3}$	$\Gamma(1)$
3.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)^3$	128	6	$1^4 \cdot 2^6 \cdot 4^2$		$\phi_{2A, 2A_5}$	$\Gamma(1)$
4.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2B)^3$	3840	6	4^6		$\phi_{2B, 2B_2}$	$\Gamma(1)$
5.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2B)^3$	96	6	4^6		$\phi_{2B, 2B_4}$	$\Gamma(1)$
6.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2B)^3$	64	6	4^6		$\phi_{2B, 2B_6}$	$\Gamma(1)$
7.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)^2(2B)$	256	2	$2^8 \cdot 4^2$		$\phi_{2B, 2A_1}, \phi_{2A, 2B_3}, \phi_{2A, 2A_4}$	$\Gamma_0(2)$
8.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)(2B)^2$	512	2	$2^4 \cdot 4^4$		$\phi_{2A, 2B_1}, \phi_{2B, 2A_2}, \phi_{2B, 2B_1}$	$\Gamma_0(2)$
9.	$\mathbb{Z}_2 \times \mathbb{Z}_2$	$(2A)(2B)^2$	128	2	$2^4 \cdot 4^4$		$\phi_{2A, 2B_2}, \phi_{2B, 2A_3}, \phi_{2B, 2B_5}$	$\Gamma_0(2)$
10.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^3(4A)^4$	64	8	$2^4 \cdot 8^2$	1	$\phi_{2A, 4A_2}, \phi_{4A, 2A_2}, \phi_{4A, 4A_3}$	$\Gamma_0(2)$
11.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^3(4A)^4$	64	8	4^6	2	$\phi_{2A, 4A_3}, \phi_{4A, 2A_3}, \phi_{4A, 4A_7}$	$\Gamma_0(2)$
12.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^2(2B)$ $(4A)^2(4B)^2$	32	2	$2^2 \cdot 4^3 \cdot 8^1$	7	$\phi_{2A, 4A_4}, \phi_{2A, 4B_4}, \phi_{2B, 4A_1},$ $\phi_{2B, 4B_1}, \phi_{4A, 2A_1}, \phi_{4A, 2B_1},$ $\phi_{4B, 2A_4}, \phi_{4B, 2B_2}, \phi_{4B, 4A_1},$ $\phi_{4B, 4A_2}, \phi_{4A, 4B_3}, \phi_{4A, 4B_4}$	$\Gamma_{2A, 4A}$
13.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4A)^4$	64	8	$4^2 \cdot 8^2$	8	$\phi_{2B, 4A_2}, \phi_{4A, 2B_3}, \phi_{4A, 4A_5}$	$\Gamma_0(2)$
14.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4A)^4$	32	8	$4^2 \cdot 8^2$	8	$\phi_{2B, 4A_3}, \phi_{4A, 2B_2}, \phi_{4A, 4A_2}$	$\Gamma_0(2)$
15.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4C)^4$	32	4	$4^2 \cdot 8^2$	8	$\phi_{2A, 4C_1}, \phi_{4C, 2A_2}, \phi_{4C, 2B_3},$ $\phi_{2B, 4C_2}, \phi_{4C, 4C_4}, \phi_{4C, 4C_6}$	$\Gamma_0(4)$
16.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4C)^4$	16	4	$4^2 \cdot 8^2$	9	$\phi_{2A, 4C_2}, \phi_{4C, 2A_1}, \phi_{4C, 2B_2},$ $\phi_{2B, 4C_3}, \phi_{4C, 4C_3}, \phi_{4C, 4C_5}$	$\Gamma_0(4)$
17.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^3(4B)^4$	64	8	$1^4 \cdot 2^2 \cdot 8^2$	1	$\phi_{2A, 4B_2}, \phi_{4B, 2A_2}, \phi_{4B, 4B_8}$	$\Gamma_0(2)$
18.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^3(4B)^4$	64	8	$2^4 \cdot 4^4$	2	$\phi_{2A, 4B_3}, \phi_{4B, 2A_5}, \phi_{4B, 4B_9}$	$\Gamma_0(2)$
19.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)^3(4B)^4$	16	8	$1^2 \cdot 2^3 \cdot 4^2 \cdot 8^1$	3	$\phi_{2A, 4B_5}, \phi_{4B, 2A_3}, \phi_{4B, 4B_3}$	$\Gamma_0(2)$

#	(g, h)	$\phi_{g,h}$	$2\mathcal{T}_2^\alpha \phi_{g,h}$	$3\mathcal{T}_3^\alpha \phi_{g,h}$	$4\mathcal{T}_4^\alpha \phi_{g,h}$
1.	$(2A, 2A_2)$				$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2A}\left(\frac{2\tau+1}{2}, 2z\right)$
2.	$(2A, 2A_3)$	0	$\phi_{2A,e}(2\tau, 2z) + \phi_{e,2A}\left(\frac{\tau}{2}, z\right) + \phi_{e,2A}\left(\frac{\tau+1}{2}, z\right)$	0	$+\phi_{e,e}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,2A}\left(\frac{\tau+i}{4}, z\right)$
3.	$(2A, 2A_5)$				
4.	$(2B, 2B_2)$				$\phi_{2B,e}(4\tau, 4z) - \phi_{e,2B}\left(\frac{2\tau+1}{2}, 2z\right)$
5.	$(2B, 2B_4)$	0	$-\phi_{2B,e}(2\tau, 2z) + \phi_{e,2B}\left(\frac{\tau}{2}, z\right) - \phi_{e,2B}\left(\frac{\tau+1}{2}, z\right)$	0	$+\phi_{e,e}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,2B}\left(\frac{\tau+i}{4}, z\right)$
6.	$(2B, 2B_6)$				
7.	$(2A, 2A_4)$	0	$-\phi_{2A,e}(2\tau, 2z) + \phi_{e,2A}\left(\frac{\tau}{2}, z\right) + \phi_{e,2B}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2A}\left(\frac{2\tau+1}{2}, 2z\right) + \phi_{e,e}(\tau, 2z)$ $+ \sum_{i=0}^1 \left(\phi_{e,2A}\left(\frac{\tau+2i}{4}, z\right) + \phi_{e,2B}\left(\frac{\tau+2i+1}{4}, z\right)\right)$
8.	$(2A, 2B_1)$	0	$\phi_{2A,e}(2\tau, 2z) + \phi_{e,2B}\left(\frac{\tau}{2}, z\right) + \phi_{e,2B}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2A}\left(\frac{2\tau+1}{2}, 2z\right)$
9.	$(2A, 2B_2)$				$+\phi_{e,e}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,2B}\left(\frac{\tau+i}{4}, z\right)$
10.	$(2A, 4A_2)$	0	$\phi_{e,4A}\left(\frac{\tau}{2}, z\right) + \phi_{e,4A}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2A}\left(\frac{2\tau+1}{2}, 2z\right)$
11.	$(2A, 4A_3)$				$+\phi_{e,2A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4A}\left(\frac{\tau+i}{4}, z\right)$
12.	$(2A, 4B_4)$	0	$\phi_{e,4A}\left(\frac{\tau+1}{2}, z\right) + \phi_{e,4B}\left(\frac{\tau}{2}, z\right)$	0	$-\phi_{2A,e}(4\tau, 4z) + \phi_{e,2B}\left(\frac{2\tau+1}{2}, 2z\right) + \phi_{e,2A}(\tau, 2z)$ $+ \sum_{i=0}^1 \left(\phi_{e,4B}\left(\frac{\tau+2i}{4}, z\right) + \phi_{e,4A}\left(\frac{\tau+2i+1}{4}, z\right)\right)$
13.	$(2B, 4A_2)$	$4\frac{\eta(2\tau)^2}{\eta(\tau)^4}\vartheta_1(\tau, z)^2$	$\phi_{e,4A}\left(\frac{\tau}{2}, z\right) - \phi_{e,4A}\left(\frac{\tau+1}{2}, z\right)$	$\phi_{2B,4A_2}(3\tau, 3z) + \phi_{2B,4A_2}\left(\frac{\tau}{3}, z\right)$ $+i\phi_{2B,4A_2}\left(\frac{\tau+1}{3}, z\right) - \phi_{2B,4A_2}\left(\frac{\tau+2}{3}, z\right)$	$\phi_{2B,e}(4\tau, 4z) - \phi_{e,2B}\left(\frac{2\tau+1}{2}, 2z\right)$ $+\phi_{e,2A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4A}\left(\frac{\tau+i}{4}, z\right)$
14.	$(2B, 4A_3)$	0	$\phi_{e,4A}\left(\frac{\tau}{2}, z\right) - \phi_{e,4A}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2B,e}(4\tau, 4z) - \phi_{e,2B}\left(\frac{2\tau+1}{2}, 2z\right)$ $+\phi_{e,2A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4A}\left(\frac{\tau+i}{4}, z\right)$
15.	$(2A, 4C_1)$	0	$\phi_{e,4C}\left(\frac{\tau}{2}, z\right) + \phi_{e,4C}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2B}\left(\frac{2\tau+1}{2}, 2z\right)$
16.	$(2A, 4C_2)$				$+\phi_{e,2B}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4C}\left(\frac{\tau+i}{4}, z\right)$
17.	$(2A, 4B_2)$				
18.	$(2A, 4B_3)$	0	$\phi_{e,4B}\left(\frac{\tau}{2}, z\right) + \phi_{e,4B}\left(\frac{\tau+1}{2}, z\right)$	0	$\phi_{2A,e}(4\tau, 4z) + \phi_{e,2A}\left(\frac{2\tau+1}{2}, 2z\right)$
19.	$(2A, 4B_5)$				$+\phi_{e,2A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4B}\left(\frac{\tau+i}{4}, z\right)$

#	Structure	Elements	$ C(g, h) $	$\frac{ N(g, h) }{ C(g, h) }$	Orbits on 24	Max subgr.	Functions	$\Gamma_{g, h}$
20.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4B)^4$	64	8	$2^4 \cdot 8^2$	8	$\phi_{2B, 4B_2}, \phi_{4B, 2B_3}, \phi_{4B, 4B_1}$	$\Gamma_0(2)$
21.	$\mathbb{Z}_2 \times \mathbb{Z}_4$	$(2A)(2B)^2(4B)^4$	16	8	$2^4 \cdot 8^2$	9	$\phi_{2B, 4B_3}, \phi_{4B, 2B_1}, \phi_{4B, 4B_5}$	$\Gamma_0(2)$
22.	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$\frac{(2A)(2B)^2}{(4A)^4(4C)^8}$	16	16	$8^1 \cdot 16^1$	13, 15, 15	$\phi_{4A, 4C_1}, \phi_{4A, 4C_2}, \phi_{4C, 4A_1},$ $\phi_{4C, 4A_2}, \phi_{4C, 4C_7}, \phi_{4C, 4C_8}$	$\Gamma_{4A, 4C}$
23.	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$(2A)^3(4A)^8(4B)^4$	16	32	$2^2 \cdot 4^1 \cdot 16^1$	10, 10, 17	$\phi_{4B, 4A_3}, \phi_{4A, 4B_1}, \phi_{4A, 4A_1}$	$\Gamma_0(2)$
24.	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$(2A)^3(4A)^8(4B)^4$	16	32	$4^2 \cdot 8^2$	11, 11, 18	$\phi_{4B, 4A_4}, \phi_{4A, 4B_2}, \phi_{4A, 4A_4}$	$\Gamma_0(2)$
25.	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$(2A)^3(4B)^{12}$	16	96	$1^4 \cdot 4^1 \cdot 16^1$	17, 17, 17	$\phi_{4B, 4B_4}$	$\Gamma(1)$
26.	$\mathbb{Z}_4 \times \mathbb{Z}_4$	$(2A)^3(4B)^{12}$	16	96	4^6	18, 18, 18	$\phi_{4B, 4B_7}$	$\Gamma(1)$
27.	$\mathbb{Z}_2 \times \mathbb{Z}_8$	$\frac{(2A)(2B)^2}{(4B)^4(8A)^8}$	16	8	$2^2 \cdot 4^1 \cdot 16^1$	20	$\phi_{2B, 8A_{1,2}}, \phi_{8A, 2B_{1,2}}, \phi_{4B, 8A_{2,3}},$ $\phi_{8A, 4B_{1,3}}, \phi_{8A, 8A_{2,8}}, \phi_{8A, 8A_{6,7}}$	$\Gamma_0(4)$
28.	$\mathbb{Z}_2 \times \mathbb{Z}_6$	$(2A)^3(3A)^2(6A)^6$	12	12	$1^2 \cdot 3^2 \cdot 4^1 \cdot 12^1$	1	$\phi_{2A, 6A_2}, \phi_{6A, 2A_1},$ $\phi_{6A, 6A_1}, \phi_{6A, 6A_2}$	$\Gamma_0(3)$
29.	$\mathbb{Z}_2 \times \mathbb{Z}_6$	$(2A)^3(3A)^2(6A)^6$	12	12	$2^3 \cdot 6^3$	2	$\phi_{2A, 6A_3}, \phi_{6A, 2A_2},$ $\phi_{6A, 6A_3}, \phi_{6A, 6A_4}$	$\Gamma_0(3)$
30.	$\mathbb{Z}_2 \times \mathbb{Z}_6$	$(2B)^3(3B)^2(6B)^6$	12	12	12^2	4	$\phi_{2B, 6B_2}, \phi_{6B, 2B_1},$ $\phi_{6B, 6B_2}, \phi_{6B, 6B_5}$	$\Gamma_0(3)$
31.	$\mathbb{Z}_2 \times \mathbb{Z}_6$	$(2B)^3(3B)^2(6B)^6$	12	12	12^2	5	$\phi_{2B, 6B_3}, \phi_{6B, 2B_2},$ $\phi_{6B, 6B_1}, \phi_{6B, 6B_3}$	$\Gamma_0(3)$
32.	$\mathbb{Z}_2 \times \mathbb{Z}_{10}$	$(2B)^3(5A)^4(10A)^{12}$	20	12	$4^1 \cdot 20^1$	4	$\phi_{2B, 10A_1}, \phi_{2B, 10A_3}, \phi_{10A, 10A_1},$ $\phi_{10A, 10A_2}, \phi_{10A, 10A_3}, \phi_{10A, 10A_5},$ $\phi_{10A, 10A_7}, \phi_{10A, 10A_9}, \phi_{10A, 10A_{10}},$ $\phi_{10A, 10A_{11}}, \phi_{10A, 2B_2}, \phi_{10A, 2B_3}$	$\Gamma_{2B, 10A}$
33.	$\mathbb{Z}_3 \times \mathbb{Z}_3$	$(3A)^8$	9	48	$1^3 \cdot 3^4 \cdot 9^1$		$\phi_{3A, 3A_3}$	$\Gamma(1)$
34.	$\mathbb{Z}_3 \times \mathbb{Z}_3$	$(3A)^2(3B)^6$	9	12	$3^2 \cdot 9^2$		$\phi_{3A, 3B_1}, \phi_{3B, 3A_1},$ $\phi_{3B, 3B_3}, \phi_{3B, 3B_4}$	$\Gamma_0(3)$

#	(g, h)	$\phi_{g,h}$	$2\mathcal{T}_2^\alpha \phi_{g,h}$	$3\mathcal{T}_3^\alpha \phi_{g,h}$	$4\mathcal{T}_4^\alpha \phi_{g,h}$
20.	$(2B, 4B_2)$	0	$\phi_{e,4B} \left(\frac{\tau}{2}, z \right) - \phi_{e,4B} \left(\frac{\tau+1}{2}, z \right)$	0	$\phi_{2B,e}(4\tau, 4z) - \phi_{e,2B} \left(\frac{2\tau+1}{2}, 2z \right)$
21.	$(2B, 4B_3)$				$+\phi_{e,2A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,4B} \left(\frac{\tau+i}{4}, z \right)$
22.	$(4A, 4C_1)$	0	$\phi_{4A,2B_3}(2\tau, 2z)$	0	$-\phi_{4A,e}(4\tau, 4z) + \phi_{2A,4A_1} \left(\frac{2\tau+1}{2}, 2z \right)$
23.	$(4B, 4A_3)$	$2\sqrt{2} \frac{\eta(2\tau)^2}{\eta(\tau)^4} \vartheta_1(\tau, z)^2$	0	$\phi_{4B,4A_3}(3\tau, 3z) + \phi_{4B,4A_3} \left(\frac{\tau}{3}, z \right)$	$\phi_{4B,e}(4\tau, 4z) + \phi_{2A,4B_1} \left(\frac{2\tau+1}{2}, 2z \right)$
24.	$(4B, 4A_4)$			$+i\phi_{4B,4A_3} \left(\frac{\tau+1}{3}, z \right) - \phi_{4B,4A_3} \left(\frac{\tau+2}{3}, z \right)$	$+\sum_{i=0}^3 \phi_{e,4A} \left(\frac{\tau+i}{4}, z \right)$
25.	$(4B, 4B_4)$	0	0	0	$\phi_{4B,e}(4\tau, 4z) + \phi_{2A,4B_1} \left(\frac{2\tau+1}{2}, 2z \right)$
26.	$(4B, 4B_7)$				$+\sum_{i=0}^3 \phi_{e,4B} \left(\frac{\tau+i}{4}, z \right)$
27.	$(2B, 8A_1)$	$2\frac{\eta(2\tau)^2}{\eta(\tau)^4} \vartheta_1(\tau, z)^2$	$\phi_{e,8A} \left(\frac{\tau}{2}, z \right) - \phi_{e,8A} \left(\frac{\tau+1}{2}, z \right)$	$\phi_{2B,8A_{1,2}}(3\tau, 3z) + \phi_{2B,8A_{1,2}} \left(\frac{\tau}{3}, z \right)$ $+i\phi_{2B,8A_{1,2}} \left(\frac{\tau+1}{3}, z \right) - \phi_{2B,8A_{1,2}} \left(\frac{\tau+2}{3}, z \right)$	$-\phi_{e,4B} \left(\frac{2\tau+1}{2}, 2z \right) + \phi_{e,4B}(\tau, 2z)$ $+\sum_{i=0}^3 \phi_{e,8A} \left(\frac{\tau+i}{4}, z \right)$
28.	$(2A, 6A_2)$	0	$\phi_{2A,3A_1}(2\tau, 2z) + \phi_{e,6A} \left(\frac{\tau}{2}, z \right) + \phi_{e,6A} \left(\frac{\tau+1}{2}, z \right)$	0	$\phi_{2A,3A}(4\tau, 4z) + \phi_{e,6A} \left(\frac{2\tau+1}{2}, 2z \right)$
29.	$(2A, 6A_3)$				$+\phi_{e,3A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,6A} \left(\frac{\tau+i}{4}, z \right)$
30.	$(2B, 6B_2)$	0	$\phi_{2B,3B_1}(2\tau, 2z) + \phi_{e,6B} \left(\frac{\tau}{2}, z \right) - \phi_{e,6B} \left(\frac{\tau+1}{2}, z \right)$	0	$\phi_{2B,3B}(4\tau, 4z) - \phi_{e,6B} \left(\frac{2\tau+1}{2}, 2z \right)$
31.	$(2B, 6B_3)$				$+\phi_{e,3B}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,6B} \left(\frac{\tau+i}{4}, z \right)$
32.	$(2B, 10A_1)$	0	$\phi_{2B,5A_1}(2\tau, 2z) + \phi_{e,10A} \left(\frac{\tau}{2}, z \right) - \phi_{e,10A} \left(\frac{\tau+1}{2}, z \right)$	0	$\phi_{2B,5A}(4\tau, 4z) - \phi_{e,10A} \left(\frac{2\tau+1}{2}, 2z \right)$ $+\phi_{e,5A}(\tau, 2z) + \sum_{i=0}^3 \phi_{e,10A} \left(\frac{\tau+i}{4}, z \right)$
33.	$(3A, 3A_3)$	0	0	$\phi_{3A,e}(3\tau, 3z) + \sum_{k=0}^2 \phi_{e,3A} \left(\frac{\tau+k}{3}, z \right)$	0
34.	$(3A, 3B_1)$	0	0	$\phi_{3A,e}(3\tau, 3z) + \sum_{k=0}^2 \phi_{e,3B} \left(\frac{\tau+k}{3}, z \right)$	0

Group	(g, h)	$\eta_{g,h}$	w	Group	(g, h)	$\eta_{g,h}$	w
1, 2, 3	$(2A, 2A_{2,3,5})$	2^{12}	6	20, 21	$(2B, 4B_{2,3}), (4B, 2B_{3,1}), (4B, 4B_{1,5}),$	4^6	3
4, 5, 6	$(2B, 2B_{2,4,6})$	4^6	3	22	$(4A, 4C_{1,2}), (4C, 4A_{1,2})$	$8 \cdot 16$	1
	$(2A, 2A_4)$	$1^4 \cdot 2^2 \cdot 4^4$			$(4C, 4C_{7,8})$	$16^4/8 \cdot 32$	
7	$(2A, 2B_3)$	$2^{14}/1^4$	5	23, 24	$(4B, 4A_{3,4}), (4A, 4B_{1,2})$	$4^2 \cdot 8^2$	2
	$(2B, 2A_1)$	$4^{14}/8^4$			$(4A, 4A_{1,4})$	$8^8/4^2 \cdot 16^2$	
8, 9	$(2A, 2B_{1,2}), (2B, 2A_{2,3})$	$2^4 4^4$	4	25, 26	$(4B, 4B_{4,7})$	4^6	3
	$(2B, 2B_{1,5})$	$4^{16}/2^4 8^4$			$(2B, 8A_{1,2}), (8A, 2B_{1,2})$	$4^2 \cdot 8^2$	
10, 11	$(2A, 4A_{2,3}), (4A, 2A_{2,3})$	4^6	3	27	$(4B, 8A_{2,3}), (8A, 4B_{1,3})$	$4^2 \cdot 8^2$	2
	$(4A, 4A_{3,7})$	$8^{18}/4^6 \cdot 16^6$			$(8A, 8A_{2,6,7,8})$	$8^8/4^2 \cdot 16^2$	
	$(2A, 4B_4)$	$1^2 \cdot 2 \cdot 4 \cdot 8^2$		28, 29	$(2A, 6A_{2,3}), (6A, 2A_{1,2})$	$2^3 \cdot 6^3$	3
	$(2A, 4A_4)$	$2^7 \cdot 8^2/1^2 \cdot 4$			$(6A, 6A_{1,2,3,4})$	irrational	
12	$(2B, 4B_1)$	$2^2 \cdot 8^7/4 \cdot 16^2$	3	30, 31	$(2B, 6B_{2,3}), (6B, 2B_{1,2})$	12^2	1
	$(2B, 4A_1)$	$4^5 \cdot 8^5/2^2 \cdot 16^2$			$(6B, 6B_{2,5,1,3})$	irrational	
	$(4B, 4A_2)$	irrational		32	$(2B, 10A_{1,3}), (10A, 2B_{2,3})$	$4 \cdot 20$	1
13, 14	$(2B, 4A_{2,3}), (4A, 2B_{3,2})$	$4^2 \cdot 8^2$	2		$(10A, 10A)$	irrational	
	$(4A, 4A_{5,2})$	$8^8/4^2 \cdot 16^2$		33	$(3A, 3A_3)$	3^8	4
	$(2A, 4C_{1,2}), (4C, 2A_{2,1})$	$4^2 \cdot 8^2$			$(3A, 3B_1), (3B, 3A_1)$	$3^2 \cdot 9^2$	2
15, 16	$(2B, 4C_{2,3}), (4C, 2B_{3,2})$	$8^8/4^2 \cdot 16^2$	2	34	$(3B, 3B_{3,4})$	irrational	
	$(4C, 4C_{4,6,3,5})$	irrational					
17, 18, 19	$(2A, 4B_{2,3,5}), (4B, 2A_{2,5,3})$	$2^4 \cdot 4^4$	4				
	$(4B, 4B_{8,9,3})$	$4^{16}/2^4 \cdot 8^4$					

Table 1: The eta products of [32]. For each pair g, h , a cycle shape of the form $\prod_{\ell} \ell^{i(\ell)}$ is given: the corresponding eta-product is $\eta_{g,h}(\tau) = \prod_{\ell} \eta(\ell\tau/N\lambda)^{i(\ell)}$, where N is the order of g and λ is the length of the shortest cycle. In the last column, w is the modular weight.

$\langle g, h \rangle$	h	$\langle g, h' \rangle$	h'	$\langle g, h \rangle$	h	$\langle g, h' \rangle$	h'
$\mathbb{Z}_2 A$	$1A$	$\mathbb{Z}_2 A$	$1A$	$\mathbb{Z}_{14} AB$	$14A_1$	$\mathbb{Z}_{14} AB$	$14B_1$
$\mathbb{Z}_2 A$	Qg	$\mathbb{Z}_2 A$	Qg	$\mathbb{Z}_{14} AB$	$14B_1$	$\mathbb{Z}_{14} AB$	$14A_1$
$\mathbb{Z}_2 A$	g	$\mathbb{Z}_2 A$	Q	1	$2A_2$	2	$2A_3$
$\mathbb{Z}_2 A$	Q	$\mathbb{Z}_2 A$	g	2	$2A_3$	1	$2A_2$
$\mathbb{Z}_4 A$	$4A_1$	$\mathbb{Z}_4 A$	$4A_1$	3	$2A_5$	3	$2A_5$
$\mathbb{Z}_4 A$	$4A_1$	$\mathbb{Z}_4 A$	$4A_1$	7	$2A_4$	$\mathbb{Z}_4 B$	$4B_1$
$\mathbb{Z}_4 B$	$4B_1$	7	$2A_4$	7	$2B_3$	$\mathbb{Z}_4 B_1$	$4B$
$\mathbb{Z}_4 B$	$4B_1$	7	$2B_3$	8	$2B_1$	8	$2B_1$
$\mathbb{Z}_6 A$	$3A_1$	$\mathbb{Z}_6 A$	$3A_1$	9	$2B_2$	9	$2B_2$
$\mathbb{Z}_6 A$	$3A_1$	$\mathbb{Z}_6 A$	$6A_1$	10	$4A_2$	11	$4A_3$
$\mathbb{Z}_6 A$	$6A_1$	$\mathbb{Z}_6 A$	$3A_1$	11	$4A_3$	10	$4A_2$
$\mathbb{Z}_6 A$	$6A_1$	$\mathbb{Z}_6 A$	$6A_1$	12	$4A_4$	$\mathbb{Z}_8 A$	$8A_1$
$\mathbb{Z}_8 A$	$8A_1$	12	$4B_4$	12	$4B_4$	$\mathbb{Z}_8 A$	$8A_1$
$\mathbb{Z}_8 A$	$8A_1$	12	$4A_4$	15	$4C_1$	15	$4C_1$
$\mathbb{Z}_{12} A$	$12A_1$	$\mathbb{Z}_{12} A$	$12A_1$	16	$4C_2$	16	$4C_2$
$\mathbb{Z}_{12} A$	$12A_1$	$\mathbb{Z}_{12} A$	$12A_1$	17	$4B_2$	18	$4B_3$
$\mathbb{Z}_{14} AB$	$7A_1$	$\mathbb{Z}_{14} AB$	$7B_1$	18	$4B_3$	17	$4B_2$
$\mathbb{Z}_{14} AB$	$7A_1$	$\mathbb{Z}_{14} AB$	$14A_1$	19	$4B_5$	19	$4B_5$
$\mathbb{Z}_{14} AB$	$7B_1$	$\mathbb{Z}_{14} AB$	$7A_1$	28	$6A_2$	29	$6A_3$
$\mathbb{Z}_{14} AB$	$7B_1$	$\mathbb{Z}_{14} AB$	$14B_1$	28	$6A_2$	29	$6A_3$
$\mathbb{Z}_{14} AB$	$14A_1$	$\mathbb{Z}_{14} AB$	$7A_1$	29	$6A_3$	28	$6A_2$
$\mathbb{Z}_{14} AB$	$14B_1$	$\mathbb{Z}_{14} AB$	$7B_1$	29	$6A_3$	28	$6A_2$

Table 2: Relabeling for g in class 2A. Each line corresponds to a conjugacy class in the central extension $C_{M_{24}}^\alpha(g)$. For each such class, we report the corresponding class in $C_{M_{24}}(g)$ with representative h , the group $\langle g, h \rangle$ generated by g and h , the ‘reabeled’ class in $C_{M_{24}}(g)$ with representative h' and the group $\langle g, h' \rangle$ generated by g and h' .

$\langle g, h \rangle$	h	$\langle g, h' \rangle$	h'	$\langle g, h \rangle$	h	$\langle g, h' \rangle$	h'	$\langle g, h \rangle$	h	$\langle g, h' \rangle$	h'
\mathbb{Z}_4B	$1A$	\mathbb{Z}_4B	$1A$	12	$2A_4$	\mathbb{Z}_8A	$8A_4$	21	$4B_1$	21	$2B_3$
\mathbb{Z}_4B	Q^2	\mathbb{Z}_4B	g^2	12	$4A_1$	\mathbb{Z}_8A	$8A_1$	21	$4B_1$	21	$4B_3$
\mathbb{Z}_4B	Q	\mathbb{Z}_4B	g	12	$4A_1$	\mathbb{Z}_8A	$8A_4$	23	$4A_3$	24	$4A_4$
\mathbb{Z}_4B	Q^3	\mathbb{Z}_4B	g^3	12	$4A_2$	\mathbb{Z}_8A	$8A_1$	23	$4A_3$	24	$4A_4$
\mathbb{Z}_4B	g^2	\mathbb{Z}_4B	Q^2	12	$4A_2$	\mathbb{Z}_8A	$8A_4$	23	$4A_3$	24	$4A_4$
\mathbb{Z}_4B	g^2Q^2	\mathbb{Z}_4B	g^2Q^2	17	$2A_2$	18	$2A_5$	23	$4A_3$	24	$4A_4$
\mathbb{Z}_4B	g^2Q	\mathbb{Z}_4B	gQ^2	17	$2A_2$	18	$4B_9$	24	$4A_4$	23	$4A_3$
\mathbb{Z}_4B	g^2Q^3	\mathbb{Z}_4B	g^3Q^2	17	$4B_8$	18	$2A_5$	24	$4A_4$	23	$4A_3$
\mathbb{Z}_4B	gQ^2	\mathbb{Z}_4B	g^2Q	17	$4B_8$	18	$4B_9$	24	$4A_4$	23	$4A_3$
\mathbb{Z}_4B	g	\mathbb{Z}_4B	Q	18	$2A_5$	17	$2A_2$	24	$4A_4$	23	$4A_3$
\mathbb{Z}_4B	gQ	\mathbb{Z}_4B	gQ	18	$2A_5$	17	$4B_8$	25	$4B_4$	26	$4B_7$
\mathbb{Z}_4B	gQ^3	\mathbb{Z}_4B	g^3Q	18	$4B_9$	17	$2A_2$	25	$4B_4$	26	$4B_7$
\mathbb{Z}_4B	g^3Q^2	\mathbb{Z}_4B	g^2Q^3	18	$4B_9$	17	$4B_8$	25	$4B_4$	26	$4B_7$
\mathbb{Z}_4B	g^3	\mathbb{Z}_4B	Q^3	19	$2A_3$	19	$2A_3$	25	$4B_4$	26	$4B_7$
\mathbb{Z}_4B	g^3Q	\mathbb{Z}_4B	gQ^3	19	$2A_3$	19	$4B_3$	26	$4B_7$	25	$4B_4$
\mathbb{Z}_4B	g^3Q^3	\mathbb{Z}_4B	g^3Q^3	19	$4B_3$	19	$2A_3$	26	$4B_7$	25	$4B_4$
\mathbb{Z}_8A	$8A_1$	12	$2B_2$	19	$4B_3$	19	$4B_3$	26	$4B_7$	25	$4B_4$
\mathbb{Z}_8A	$8A_1$	12	$2A_4$	20	$2B_3$	20	$2B_3$	26	$4B_7$	25	$4B_4$
\mathbb{Z}_8A	$8A_1$	12	$4A_1$	20	$2B_3$	20	$2B_3$	27	$8A_2$	27	$8A_2$
\mathbb{Z}_8A	$8A_1$	12	$4A_2$	20	$2B_3$	20	$4B_1$	27	$8A_2$	27	$8A_2$
\mathbb{Z}_8A	$8A_4$	12	$2B_2$	20	$2B_3$	20	$4B_1$	27	$8A_3$	27	$8A_3$
\mathbb{Z}_8A	$8A_4$	12	$2A_4$	20	$4B_1$	20	$2B_3$	27	$8A_3$	27	$8A_3$
\mathbb{Z}_8A	$8A_4$	12	$4A_1$	20	$4B_1$	20	$2B_3$	27	$8A_3$	27	$8A_3$
\mathbb{Z}_8A	$8A_4$	12	$4A_2$	20	$4B_1$	20	$4B_1$	27	$8A_3$	27	$8A_3$
12	$2B_2$	\mathbb{Z}_8A	$8A_1$	20	$4B_1$	20	$4B_1$	27	$8A_2$	27	$8A_2$
12	$2B_2$	\mathbb{Z}_8A	$8A_4$	21	$2B_3$	21	$2B_3$	27	$8A_2$	27	$8A_2$
12	$2A_4$	\mathbb{Z}_8A	$8A_1$	21	$2B_3$	21	$4B_1$				

Table 3: Relabeling for g in class 4B.

References

- [1] T. Eguchi, H. Ooguri and Y. Tachikawa, “Notes on the K3 Surface and the Mathieu group M_{24} ,” *Exper. Math.* **20**, 91 (2011) [[arXiv:1004.0956](#) [[hep-th](#)]].
- [2] M.C.N. Cheng, “K3 Surfaces, N=4 dyons, and the Mathieu group M_{24} ,” *Commun. Number Theory Phys.* **4**, 623 (2010) [[arXiv:1005.5415](#) [[hep-th](#)]].
- [3] M.R. Gaberdiel, S. Hohenegger and R. Volpato, “Mathieu twining characters for K3,” *JHEP* **1009**, 058 (2010) [[arXiv:1006.0221](#) [[hep-th](#)]].
- [4] M.R. Gaberdiel, S. Hohenegger and R. Volpato, “Mathieu Moonshine in the elliptic genus of K3,” *JHEP* **1010**, 062 (2010) [[arXiv:1008.3778](#) [[hep-th](#)]].
- [5] T. Eguchi and K. Hikami, “Note on twisted elliptic genus of K3 surface,” *Phys. Lett. B* **694**, 446 (2011) [[arXiv:1008.4924](#) [[hep-th](#)]].
- [6] A. Taormina and K. Wendland, “The Symmetries of the tetrahedral Kummer surface in the Mathieu group M_{24} ,” [arXiv:1008.0954](#) [[hep-th](#)].
- [7] M.C.N. Cheng and J.F.R. Duncan, “On Rademacher sums, the largest Mathieu group, and the holographic modularity of Moonshine,” *Commun. Number Theory Phys.* **6** (3), 697–758 (2012) [[arXiv:1110.3859](#) [[math.RT](#)]].
- [8] M.R. Gaberdiel, S. Hohenegger and R. Volpato, “Symmetries of K3 sigma models,” *Commun.Num.Theor.Phys.* **6** 1–50 (2012) [[arXiv:1106.4315](#) [[hep-th](#)]].
- [9] A. Taormina and K. Wendland, “The overarching finite symmetry group of Kummer surfaces in the Mathieu group M_{24} ,” *JHEP* **1308** (2013) 125 [[arXiv:1107.3834](#) [[hep-th](#)]].
- [10] R. Volpato, “Mathieu Moonshine and symmetries of K3 sigma models,” *Fortsch.Phys.* **60**, 1112–1117 (2012) [[arXiv:1201.6172](#)].
- [11] M.R. Gaberdiel and R. Volpato, “Mathieu moonshine and orbifold K3s,” [arXiv:1206.5143](#) [[hep-th](#)].
- [12] T. Gannon, “Much ado about Mathieu,” [arXiv:1211.5531](#) [[math.RT](#)].
- [13] A. Taormina and K. Wendland, “Symmetry-surfing the moduli space of Kummer K3s,” [arXiv:1303.2931](#) [[hep-th](#)].

- [14] A. Taormina and K. Wendland, “A twist in the M24 moonshine story,” arXiv:1303.3221 [hep-th].
- [15] M. R. Gaberdiel, A. Taormina, R. Volpato and K. Wendland, “A K3 sigma model with $\mathbb{Z}_2^8 : M_{20}$ symmetry,” arXiv:1309.4127 [hep-th].
- [16] T. Creutzig and G. Hoehn, “Mathieu Moonshine and the Geometry of K3 Surfaces,” arXiv:1309.2671 [math.QA].
- [17] M. R. Gaberdiel, D. Persson, H. Ronellenfitsch and R. Volpato, “Generalized Mathieu Moonshine,” Commun. Num. Theor. Phys. **7**, 145-223 (2013) arXiv:1211.7074 [hep-th].
- [18] M. R. Gaberdiel, D. Persson and R. Volpato, “Generalized Moonshine and Holomorphic Orbifolds,” arXiv:1302.5425 [hep-th].
- [19] S. Norton, “Generalized moonshine”, Proc. Sympos. Pure Math. **47**, 209-210, Amer. Math. Soc., Providence, RI (1987).
- [20] M. C. N. Cheng, X. Dong, J. Duncan, J. Harvey, S. Kachru and T. Wrase, “Mathieu Moonshine and N=2 String Compactifications,” JHEP **1309**, 030 (2013) [arXiv:1306.4981 [hep-th]].
- [21] S. Harrison, S. Kachru and N. M. Paquette, “Twining Genera of (0,4) Supersymmetric Sigma Models on K3,” arXiv:1309.0510 [hep-th].
- [22] J. A. Harvey and S. Murthy, “Moonshine in Fivebrane Spacetimes,” arXiv:1307.7717 [hep-th].
- [23] A. Sen, “A Twist in the Dyon Partition Function,” JHEP **1005** (2010) 028 [arXiv:0911.1563 [hep-th]].
- [24] A. Sen, “Discrete Information from CHL Black Holes,” JHEP **1011** (2010) 138 [arXiv:1002.3857 [hep-th]].
- [25] S. Govindarajan, “BKM Lie superalgebras from counting twisted CHL dyons,” JHEP **1105** (2011) 089 [arXiv:1006.3472 [hep-th]].
- [26] S. Govindarajan, “Unravelling Mathieu Moonshine,” Nucl. Phys. B **864** (2012) 823 [arXiv:1106.5715 [hep-th]].

- [27] M.C.N. Cheng and J.F.R. Duncan, “The largest Mathieu group and (mock) automorphic forms,” Proc. Sympos. Pure Math. **85**, 53–82 (2012) [arXiv:1201.4140 \[math.RT\]](#).
- [28] M. Raum, “Powers of M_{24} -twisted Siegel product expansions are modular,” [arXiv:1208.3453v2 \[math.NT\]](#).
- [29] M. C. N. Cheng, J. F. R. Duncan and J. A. Harvey, “Umbral Moonshine,” [arXiv:1204.2779 \[math.RT\]](#).
- [30] G. Mason, “ M_{24} and certain automorphic forms”, Contemp. Math. **45**, p. 223-244, (1985).
- [31] G. Mason, “Elliptic systems and the eta-function,” Notas Soc. Mat. Chile **8**, 37 (1989).
- [32] G. Mason, “On a system of elliptic modular forms attached to the largest Mathieu group”, Nagoya Math. J. **118**, 177 (1990).
- [33] G. Mason, “ G -elliptic systems and the genus zero problem for M_{24} ”, Bull. Amer. Math. Soc. **25**, no. 1, p. 45-53, (1991).
- [34] J.H. Conway and S.P. Norton, “Monstrous Moonshine”, Bull. London Math. Soc. **11**, 308 (1979).
- [35] I. Frenkel, J. Lepowsky and A. Meurman, “Vertex operator algebras and the Monster”, Pure and Applied Mathematics **134**, Academic Press, Inc., Boston, MA, (1988).
- [36] R.E. Borcherds, “Monstrous moonshine and monstrous Lie superalgebras” Invent. Math. **109**, 405 (1992).
- [37] M.P. Tuite, “Generalized moonshine and Abelian orbifold constructions,” Contemp. Math. **193**, 353–368 (1996) [arXiv:hep-th/9412036](#).
- [38] C.-y. Dong, H.-s. Li and G. Mason, “Modular invariance of trace functions in orbifold theory,” Commun. Math. Phys. **214**, 1 (2000) [[arXiv:q-alg/9703016](#)].
- [39] G. Höhn, “Generalized moonshine for the baby Monster,” Habilitation thesis, Mathematisches Institut, Universität Freiburg (2003), <http://www.math.ksu.edu/~gerald/papers/baby8.ps>.
- [40] S. Carnahan, “Generalized Moonshine I: Genus zero functions,” Algebra Number Theory **4** (6), 649–679 (2010) [arXiv:0812.3440 \[math.RT\]](#).

- [41] S. Carnahan, “Generalized Moonshine II: Borchers products,” *Duke Math. J.* **161** (5), 893–950 (2012) [arXiv:0908.4223](#) [[math.RT](#)].
- [42] S. Carnahan, “Generalized Moonshine III: Equivariant intertwining operators,” to appear.
- [43] S. Carnahan, “Generalized Moonshine IV: Monstrous Lie algebras,” [arXiv:1208.6254](#) [[math.RT](#)].
- [44] R. Dijkgraaf and E. Witten, “Topological gauge theories and group cohomology,” *Commun. Math. Phys.* **129**, 393 (1990).
- [45] P. Roche, V. Pasquier and R. Dijkgraaf, “Quasi-Hopf algebras, group cohomology and orbifold models,” *Nucl. Phys. Proc. Suppl.* **18B**, 60 (1990).
- [46] P. Bantay, “Orbifolds and Hopf algebras,” *Phys. Lett. B* **245**, 477 (1990).
- [47] A. Coste, T. Gannon and P. Ruelle, “Finite group modular data,” *Nucl. Phys. B* **581**, 679 (2000) [[arXiv:hep-th/0001158](#)].
- [48] T. Gannon, talk at ‘Mathieu Moonshine’, ETH Zürich, July 2011.
- [49] R. Dijkgraaf, G.W. Moore, E.P. Verlinde and H.L. Verlinde, “Elliptic genera of symmetric products and second quantized strings,” *Commun. Math. Phys.* **185**, 197 (1997) [[arXiv:hep-th/9608096](#)].
- [50] V. Gritsenko, “Elliptic genus of Calabi-Yau manifolds and Jacobi and Siegel modular forms,” *St. Petersburg Math. J.* **11** (5), 781–804 (2000) [arXiv:math/9906190](#).
- [51] V. A. Gritsenko and V. V. Nikulin, “Automorphic Forms and Lorentzian Kac-Moody Algebras. II”, *Internat. J. Math.* **9** (2), 201–275 (1998) [[arXiv:alg-geom/9611028](#)].
- [52] R. Dijkgraaf, E.P. Verlinde and H.L. Verlinde, “Counting dyons in N=4 string theory,” *Nucl. Phys. B* **484**, 543 (1997) [[arXiv:hep-th/9607026](#)].
- [53] N. Ganter, “Hecke operators in equivariant elliptic cohomology and generalized moonshine”, *CRM Proc. Lecture Notes* **47**, 173, Amer. Math. Soc., Providence, RI (2009) [[arXiv:0706.2898](#) [[math.AT](#)]].
- [54] V. Ginzburg, M. Kapranov, E. Vasserot, “Elliptic algebras and equivariant elliptic cohomology I.” [[arXiv:q-alg/9505012](#)]

- [55] M. Eichler and D. Zagier, *The Theory of Jacobi Forms*, Birkhäuser (1985).
- [56] R. Borcherds, “Automorphic forms on $O_{s+2,2}(\mathbf{R})$ and infinite products,” *Invent. Math.* **120** (1), 161 (1995).
- [57] A. Dabholkar and S. Nampuri, “Spectrum of dyons and black holes in CHL orbifolds using Borcherds lift,” *JHEP* **0711** (2007) 077 [[hep-th/0603066](#)].
- [58] V. A. Gritsenko and V. V. Nikulin, “Siegel automorphic form corrections of some Lorentzian Kac–Moody Lie algebras”, *Amer. J. Math.* **119** (1), 181–224 (1997) [[arXiv:alg-geom/9504006](#)].
- [59] V. A. Gritsenko and V. V. Nikulin, “The Igusa modular form and the ”simplest” Lorentzian Kac-Moody algebras”, *Sb. Math.* **187** (11), 1601–1641 (1996) [[arXiv:alg-geom/9603010](#)].
- [60] T. Eguchi and K. Hikami, “Twisted Elliptic Genus for K3 and Borcherds Product,” *Lett. Math. Phys.* **102**, 203 (2012) [[arXiv:1112.5928](#) [[hep-th](#)]].
- [61] F. Clery and V. A. Gritsenko, “The Siegel modular forms of genus 2 with the simplest divisor”, *Proc. Lond. Math. Soc.* **102** (6), 1024–1052 (2011) [[arXiv:0812.3962](#) [[math.NT](#)]].
- [62] J.R. David, D.P. Jatkar and A. Sen, “Product representation of Dyon partition function in CHL models,” *JHEP* **0606**, 064 (2006) [[arXiv:hep-th/0602254](#)].
- [63] J.R. David, D.P. Jatkar and A. Sen, “Dyon spectrum in generic $N=4$ supersymmetric $Z(N)$ orbifolds,” *JHEP* **0701**, 016 (2007) [[arXiv:hep-th/0609109](#)].
- [64] M. C. N. Cheng, J. F. R. Duncan and J. A. Harvey, “Umbral Moonshine and the Niemeier Lattices,” [arXiv:1307.5793](#) [[math.RT](#)].
- [65] M.C.N. Cheng and E.P. Verlinde, “Wall crossing, discrete attractor flow, and Borcherds algebra,” *SIGMA* **4**, 068 (2008) [[arXiv:0806.2337](#) [[hep-th](#)]].
- [66] M. C. N. Cheng and A. Dabholkar, “Borcherds-Kac-Moody Symmetry of $N=4$ Dyons,” *Commun. Num. Theor. Phys.* **3** (2009) 59 [[arXiv:0809.4258](#) [[hep-th](#)]].
- [67] S. Govindarajan and K. Gopala Krishna, “Generalized Kac-Moody Algebras from CHL dyons,” *JHEP* **0904** (2009) 032 [[arXiv:0807.4451](#) [[hep-th](#)]].

- [68] S. Govindarajan and K. Gopala Krishna, “BKM Lie superalgebras from dyon spectra in $Z(N)$ CHL orbifolds for composite N ,” JHEP **1005** (2010) 014 [[arXiv:0907.1410](#) [[hep-th](#)]].
- [69] G. Karpilovsky, “Group Representations: Volume 2,” North-Holland Mathematical Studies **177**, Elsevier (1993).
- [70] M.D.F. de Wild Propitius, “Topological interactions in broken gauge theories,” [arXiv:hep-th/9511195](#).