

Auxiliary Material for:

The use of radiocarbon to constrain current and future soil organic matter turnover and transport in a temperate forest

Maarten C. Braakhekke, Christian Beer, Marion Schrumpf, Altug Ekici, Bernhard Ahrens, Marcel R. Hoosbeek, Bart Kruijt, Pavel Kabat, and Markus Reichstein

¹Max Planck Institute for Biogeochemistry, Jena Germany

²Earth System Science and Climate Change Chairgroup, Wageningen University, Wageningen, The Netherlands

Journal of Geophysical Research – Biogeosciences, 2013

Introduction

This PDF file contains auxiliary figures and tables for the study.

List of Figures

1	Measured soil temperature at Hainich	2
2	Measured soil moisture at Hainich	2
3	Atmospheric radiocarbon content used in the simulations.	3
4	Correlation matrix of the posterior parameter distribution	4
5	Measured lead-210 fractions and corresponding model results	5
6	Measured and modeled total annual heterotrophic respiration.	5
7	Measured effective decomposition rate coefficients and corresponding model results	5
8	Marginal posterior distributions for a calibration with uninformative (uniform) priors for setup woC14	6
9	Marginal posterior distributions for a calibration with uninformative (uniform) priors for setup wC14	7
10	Organic carbon measurements for 2009 and corresponding model results based on JSBACH forcing .	8
11	Marginal prior and posterior distributions for the sub-dominant mode	9
12	Modeled organic carbon stocks and profile for the sub-dominant mode	10
13	Modeled $\Delta^{14}\text{C}$ for the sub-dominant mode	11
14	Modeled organic carbon transport fluxes for the sub-dominant mode	12
15	Modeled C stocks for projection simulations until 2100 for the sub-dominant mode	13
16	Modeled heterotrophic respiration for projection simulations until 2100 for the sub-dominant mode .	14

List of Tables

1	Model driving data and not-estimated parameters.	14
2	Properties of the marginal posterior distributions	15