

First simulations of multi-ELM cycles in tokamak X-point plasmas

F.Orain,¹ M.Bécoulet,¹ G.T.A.Huijsmans,² G.Dif-Pradalier,¹ M.Hoelzl,³ J.Morales,¹
X.Garbet,¹ E.Nardon,¹ S.Pamela,⁴ C.Passeron,¹ G.Latu,¹ A.Fil,¹ and P.Cahyna⁵

¹CEA/DSM/IRFM, Cadarache, 13108 Saint-Paul-Lez-Durance, France

²ITER Organization, route de Vinon, 13115 Saint-Paul-Lez-Durance, France

³Max-Planck-Institut für Plasmaphysik, Garching, Germany

⁴CCFE, Culham Science Centre, Oxon, OX14 3DB, UK

⁵IPP.CR, Prague, Czech Republic

(Dated: May 13, 2014)

The full dynamics of a multi-ELM cycle is modeled for the first time in realistic tokamak X-point geometry with the non-linear reduced MHD code JOEAK. The diamagnetic rotation is found to be instrumental to stabilize the plasma after an ELM crash and to model the cyclic reconstruction and collapse of the plasma pressure profile. ELM relaxations are cyclically initiated each time the pedestal gradient crosses a triggering threshold. Diamagnetic drifts are also found to yield a near-symmetric ELM power deposition on the inner and outer divertor target plates, consistent with experimental measurements.

Introduction – Edge Localized Modes (ELMs) are magnetohydrodynamic (MHD) instabilities occurring in fusion plasmas in high confinement regime (H-mode) of tokamaks and sometimes in stellarators, and present similarities with the solar flares occurring in astrophysical plasmas [1]. As the ELMs represent a particular concern for the lifetime of the divertor in ITER [2], their dynamics has been widely studied experimentally and theoretically [3–6]. In X-point tokamak plasmas, MHD instabilities develop due to a large edge pressure gradient (so-called ballooning modes) or a large edge current (so-called peeling-tearing modes), triggering the formation of ELM filaments. The ELMs consist in the quasi-periodic ejection of these filaments in H-mode regimes across the transport barrier at the plasma edge (also called pedestal).

So far, non-linear MHD computations with codes such as M3D [7], BOUT++ [8, 9], NIMROD [10] and JOEAK [11–13] have been able to reproduce a single ELM crash in simulations, in realistic tokamak geometry. Other works using a model including 2-fluid diamagnetic effects [14, 15] have shown transport bursts comparable to ELM relaxations in simplified cylindrical geometry. This Letter presents the first simulations of a multi-ELM cycle in realistic toroidal geometry, using the non-linear reduced MHD code JOEAK [11] recently extended to add the 2-fluid diamagnetic effects [16]. The diamagnetic effects are known to have a stabilizing effect on plasma instabilities. They are shown to prevent the magnetic field from reconnecting and/or to reduce the growth rate of ideal and resistive instabilities in both astrophysical [17] and laboratory plasmas [18–20]. In particular, in tokamaks, they are evidenced to be a key parameter for simulating cycles of sawtooth crashes [21]. We show here that the diamagnetic rotation is also a key parameter for modeling the ELM cycling dynamics: as the diamagnetic effects stabilize peeling-ballooning instabilities after an ELM crash, they allow for the pedestal to rebuild, thus leading to an ELM cycle. This approach is different and complementary to that in Ref [22] in which it is an externally

imposed $E \times B$ stabilization in the plasma edge that allows for oscillation relaxations.




FIG. 1: Magnetic and kinetic energies of the mode $n = 8$ without diamagnetic effects. The time in x-axis is normalized to the Alfvén time t_A .

Multi-ELM cycle simulations – Experimentally, ELM relaxations consist in a cyclical phenomenon, resulting in the quasiperiodic deposition of energy on the divertor target plates. Modeling this phenomenon requires to reproduce this cyclical behavior. Besides, the simulation of ELM cycles rather than a single ELM crash which depends on the initial unstable pressure profile involves different physical mechanisms. Indeed, after the first ELM crash, memory of the choice for the initial state is lost. The phase coherence between modes that determines the ELM instability growth has reorganized and is now consistently determined by the dynamics rather than imposed through the choice of initial conditions. And so is the energy content within the bath of toroidal harmonics. In this respect, an ELM crash starting from such a coherent state is significantly different from a first initial relaxation triggered by the choice of an initial state. This Letter discusses the specificities of the multiple ELM dynamics starting from self-consistent inter-ELM states

with respect to the usual single ELM relaxation from an initial chosen condition.

As we consider long runs of ELM-cycle simulations ($> 10^4$ Alfvén times $\sim 10^{-2}s$), we have to deal with very different time scales: the diffusive time scale $\tau_D \sim a^2/D_\perp \sim 10^{-1} - 1s$ (D_\perp being the perpendicular diffusion coefficient), the timescale of the pedestal reconstruction after an ELM ($\sim 10^{-3} - 10^{-1}s$) and the ELM crash timescale ($\sim 10^{-4}s$). Thus the presence of heat and particle sources – needed to rebuild the pedestal after an ELM – as much as diffusion propagating the heat and particles, is essential in the simulations. The heat source is chosen to follow the initial temperature profile and the particle source is taken constant over the plasma. Heat and particle diffusion coefficients are reduced in the pedestal in order to preserve a transport barrier at the plasma edge and a $3cm$ -large pedestal. The injected heat and particle sources determine the pedestal reconstruction timescale and thus the simulated ELM frequency.

In this Letter, ELM cycles are modeled for JET-like plasma parameters and geometry, similar to Ref [16]: major radius $R_0 = 3m$, minor radius $a = 1m$, toroidal magnetic field $B_t = 2.9T$ and safety factor $q_{95} \sim 3$. Established H-mode experimental profiles are taken initially, with central electron density $n_{e,0} = 6 \times 10^{19}m^{-3}$ and central temperature $T_{e,0} = 5keV$. The pedestal density and temperature are $n_{e,ped} = 3.8 \times 10^{19}m^{-3}$ and $T_{e,ped} = 2.5keV$. Because of computational restrictions, the central resistivity is taken $\eta_0 = 2.5 \times 10^{-7}\Omega.m$, which is 2 orders of magnitude larger than the Spitzer resistivity $\eta_{Spitzer} = 2.5 \times 10^{-9}\Omega.m$. The resistivity profile follows a $T^{-3/2}$ radial dependence. A JET realistic diamagnetic velocity is taken for both species s (ions and electrons): $\vec{V}_s^* = Z_s(R/R_0)^2\tau_{IC}/\rho \cdot \vec{b} \times \nabla P_s$. Z_s and P_s are respectively the charge number and the scalar pressure of the species s , R is the horizontal coordinate, ρ is the mass density and \vec{b} is the magnetic field normalized to B_t . The realistic value of the diamagnetic parameter (inverse of the ion cyclotron frequency) $\tau_{IC} = m_i/eB_t \sim 7 \times 10^{-9}s$ is taken, and ion and electron temperatures are supposed to be equal. The full description of the model is given on Ref [16]. Compared to Ref [16], the toroidal source of rotation and the neoclassical friction are not included in the modeling to focus only on the impact of the diamagnetic flow on the ELM dynamics.

A simulation is run as follows: once the axisymmetric ($n = 0$) equilibrium flows are established (mainly due to diamagnetic drifts and sheath conditions [16]), the toroidal modes of perturbation $n = 2, 4, 6$ and 8 are added in simulation. Linear runs show that the $n = 8$ mode is the most unstable mode in this configuration, as the larger $n > 9$ modes are stabilized by the diamagnetic effects.

In ELM simulations without diamagnetic effects, the ballooning and/or peeling-tearing instabilities at the onset on the ELM crash generate the stochasticization of the edge magnetic field. Yet after the ELM crash, the MHD activity is not fully stabilized and the edge mag-

netic field remains stochastic: the residual MHD activity still generates an enhanced edge transport that does not allow for the pedestal reconstruction. Therefore, a second ELM crash cannot be obtained as the plasma remains below the peeling-ballooning stability limit. An example is given for a single $n = 8$ ELM (only the $n = 0$ and $n = 8$ modes are included in the simulation) modeled without taking into account the diamagnetic effects (Fig.1): during the ELM crash, characterized by a large peak of kinetic and magnetic energy of the $n = 8$ mode (Fig.1), the plasma edge is stochasticized in a large extent: the region for normalized flux $\psi_{norm} > 0.85$ is fully reconnected (Fig.2 (a) and (c)). After the crash, the instability does not completely vanish (Fig.1) and the magnetic field remains partially reconnected: large magnetic islands subsist for $\psi_{norm} > 0.85$ and an ergodic layer remains for $\psi_{norm} > 0.95$ (Fig.2 (b) and (d), given for $t \sim 7 \times 10^3 t_A$). Thus the edge transport remains high due to both the stochastic layer enhancing the heat parallel diffusivity and the $E \times B$ convection of particles.




FIG. 2: Magnetic field topology in the $n = 8$ ELM simulation without diamagnetic effect, given at the peak of the ELM crash (a-c) and after the ELM crash at $t \sim 7 \times 10^3 t_A$ (b-d). The figures (a) and (b) show the magnetic topology in the poloidal plane, and the figure (c) [respectively (d)] is a zoom of the plasma edge of the figure (a) [respectively (b)] in (ψ, θ) coordinates for a normalized flux $\psi \geq 0.8$. Note the full edge reconnection at the peak of the crash, and the remaining partial reconnection after the crash.

However, the addition of the diamagnetic rotation in modeling has two major effects: first, the amplitude of the ELM crash is smaller due to the diamagnetic stabilization, and second the $n = 8$ mode is completely damped after the crash. The plasma is actually stabilized by the diamagnetic effects after the crash, but is not destabilized again until the pedestal pressure gradi-




FIG. 3: Kinetic energy of the modes $n = 2 - 8$ in the multi-harmonic $n = 0 : 2 : 8$ simulation with diamagnetic effects.




FIG. 4: Pedestal pressure gradient normalized over flux surfaces before (dash line), during (dash-dot line) and after (full line) the ELM crash.

ent that increases due to the injected heat source reaches the ELM-triggering threshold. Then a second ELM crash occurs, and so forth. The simulation of a multi-ELM cycle (multi-harmonic simulation $n = 0$ to 8 by step 2) is presented in Fig.3. The ELM energy is one order of magnitude smaller than in Fig.1 due to the diamagnetic stabilization. The ELM cycle can be decomposed into two parts: transient ELMs – from $t = 3200t_A$ to $t = 6200t_A$ – are followed by a quasiperiodic repetitive regime. The system evolves from a state depending on the initial conditions towards a quasiperiodic cycling regime which mainly depends on the injected heat and particles and on the diamagnetic frequency. The three first transient ELMs are largely dominated by the most unstable mode (first $n = 8$ and then $n = 6$) and there is only a few coupling between modes. However, after 4 ELMs, we get a quasi-periodic regime driven by a bunch of $n = 2 - 8$ modes strongly nonlinearly coupled with each other. Each ELM in this regime is ini-




FIG. 5: Peeling-Ballooning diagram: evolution of the pressure gradient as a function of the mean pedestal current in quasi-periodic regime. The time between two points is $20t_A$, and each color represents one ELM.

tiated by the growth of the $n = 6$ mode directly followed by the coupled growth of the other modes. After a crash, because of the quick refilling of the pedestal by the heat and particle sources, the plasma is barely stabilized by the diamagnetic effects before being destabilized again by the increase in pressure gradient: this leads to a large frequency ($\sim 3kHz$) of small ELMs. Each ELM in the quasi-periodic regime is characterized by the collapse of the edge pressure gradient, presented in Fig.4. This pedestal relaxation occurs when a threshold in pressure gradient – corresponding to the ELM triggering threshold – is reached. The Peeling-Ballooning diagram (maximal pressure gradient as a function of the mean pedestal current Fig.5) shows that these ELM are constituted of three steps: the linear growth of the instability as the pressure gradient increases, then the ELM crash – occurring at approximately the same threshold for all the ELMs in quasi-periodic regime – and last the relaxation of the density and temperature profiles. The increase of the pressure gradient in the pedestal due to the heat source is going together with the increase of the rotational shear at the plasma edge, due to the increase in diamagnetic velocity proportional to the pressure gradient. At the ELM crash, an $n = 0$ flow is induced by Maxwell stress [23], increasing the edge rotational shear. The plasma filaments are then sheared by this ($n = 0$) flow and expelled out of the plasma.

Near-symmetric deposition on divertor – After a small delay ($\sim 100t_A$) corresponding to the ion parallel time $\tau_{||,i} = \pi qR/c_s$ (where c_s is the sound speed), the energy and particles due to the ELM filaments reach the divertor target plates at the sound speed. The particles are convected by $E \times B$ and diamagnetic drifts while the energy is mainly transported by parallel conduction. Although the power deposition due to the four first ELMs is very variable, the integrated peak power deposited by an ELM on the divertor (Fig.6) is approximately the same for all




FIG. 6: Integrated power reaching the inner (dash line) and outer (full line) divertor plates.

the ELMs in quasi-periodic regime ($\sim 5 - 6 MW$ on the outer divertor, $\sim 2 - 3 MW$ on the inner divertor). This again shows that the first ELMs before reorganization are singular. Note that in quasiperiodic regime, the peak power deposition on the inner divertor is slightly delayed (by $\sim 50t_A$) compared to the outer deposition: because of the ballooning character of the instabilities, the filaments are mostly expelled from the Low Field Side, so the filaments first hit the outer target.

Furthermore, the deposition on the inner and outer divertor targets is near-symmetric in this simulation: the integrated deposited power has the same order of magnitude on both targets, even though roughly two-times more power is deposited on the outer divertor target. Experimentally, the peak power deposition is either symmetric on inner/outer targets, or the inner target receives twice more power [24, 25] after a crash. In previous modeling performed without diamagnetic drifts, the outer target received almost all the ELM power deposition, which was contrary to the experimental observations. An $n = 2-8$ ELM simulation is performed with realistic JET parameters corresponding to the shot #77329 described in [26] for two different cases: one without including flows in the model (full line in Fig.7) and one with diamagnetic effects, neoclassical friction and toroidal source of rotation included (dash line in Fig.7). In the case without flows, almost all the heat flux generated by the ELM filaments is deposited on the outer divertor, whereas in the case where flows are included in the model, the deposited heat flux is near-symmetric on the inner and outer divertor targets (Fig.7). Therefore, even though the filaments are mostly expelled from the outer region and the heat transport is mainly diffusive in direction of the outer region, the particle convection towards the inner region is enhanced by the diamagnetic flow, leading to a near-equal repartition of the power (proportional to density and temperature) on inner and outer divertor plates. Hence plasma flows and in particular the bi-fluid diamagnetic drifts allow for simulating more realistically the ELM cyclical dynamics up to the deposition on di-

vertor.

Discussion and conclusions – Even though computational restrictions impose to choose in the simulation a plasma resistivity that is two orders of magnitude larger than the realistic value, the modeling of these resistive ELMs represents a significant step forward in the understanding of the cycling dynamics. The addition of the bi-fluid diamagnetic rotation in MHD models allows for the stabilization of the plasma after an ELM relaxation. The plasma reorganizes and memory of the initial chosen profiles is lost. The phasing and the energy repartition between modes is consistently determined. The steepening of the pressure profile generated by the pedestal reconstruction destabilizes again the edge plasma until the edge pressure gradient reaches the ELM-triggering threshold: a new ELM crash then occurs. Similar coupling between modes, similar maximal pressure gradient reached when the crash occurs and similar power deposition on the divertor plates are cyclically recovered for all the ELMs in the quasiperiodic regime. These differ much from the first transient ELMs, pointing out the importance of simulating cycles rather than a single ELM crash. In addition, the inclusion of the diamagnetic drifts induces a near-symmetric ELM power deposition on the inner/outer divertor target plates, which is in closer agreement with experimental measurements.




FIG. 7: Radial distribution of the heat flux (in GW/m^2) reaching the inner and outer divertor target plates: without flows included in simulation (full line) or with diamagnetic, neoclassical and toroidal flows (dashed line).

Acknowledgments – This work has benefited from financial support from the National French Research Program (ANR): ANEMOS (2011) and E2T2(2010). *Disclaimer:* The views and opinions expressed herein do not necessarily reflect those of the ITER Organization. This work was granted access to the HPC resources of Aix-Marseille Université financed by the project Equip@Meso (ANR-10-EQPX-29-01). A part of this work was carried out using the HELIOS supercomputer system (IFERC-CSC), Aomori, Japan, under the Broader Approach collaboration, implemented by Fusion for Energy and JAEA, and using the CURIE supercomputer, operated into the TGCC by CEA, France, in the framework of GENCI and PRACE projects.

-
- [1] W. Fundamenski, *et al.* Plasma Physics and Controlled Fusion **49(5)**, R43 (2007), URL <http://stacks.iop.org/0741-3335/49/i=5/a=R01>.
- [2] A. Loarte, *et al.* Plasma Physics and Controlled Fusion **45(9)**, 1549 (2003), URL <http://stacks.iop.org/0741-3335/45/i=9/a=302>.
- [3] P. Gohil, *et al.* Phys. Rev. Lett. **61**, 1603 (1988), URL <http://link.aps.org/doi/10.1103/PhysRevLett.61.1603>.
- [4] H. Zohm. Plasma Physics and Controlled Fusion **38(2)**, 105 (1996), URL <http://stacks.iop.org/0741-3335/38/i=2/a=001>.
- [5] A. Kirk, *et al.* (MAST Team). Phys. Rev. Lett. **92**, 245002 (2004), URL <http://link.aps.org/doi/10.1103/PhysRevLett.92.245002>.
- [6] H. R. Wilson *et al.* Phys. Rev. Lett. **92**, 175006 (2004), URL <http://link.aps.org/doi/10.1103/PhysRevLett.92.175006>.
- [7] L. E. Sugiyama *et al.* Physics of Plasmas **17(6)**, 062505 (2010), URL <http://scitation.aip.org/content/aip/journal/pop/17/6/10.1063/1.3449301>.
- [8] X. Xu, *et al.* Nuclear Fusion **51(10)**, 103040 (2011), URL <http://stacks.iop.org/0029-5515/51/i=10/a=103040>.
- [9] W. Xi, P. *et al.* Phys. Rev. Lett. **112**, 085001 (2014), URL <http://link.aps.org/doi/10.1103/PhysRevLett.112.085001>.
- [10] C. R. Sovinec, *et al.* Journal of Physics: Conference Series **78(1)**, 012070 (2007), URL <http://stacks.iop.org/1742-6596/78/i=1/a=012070>.
- [11] G. T. A. Huysmans, *et al.* Plasma Physics and Controlled Fusion **51(12)**, 124012 (2009), URL <http://stacks.iop.org/0741-3335/51/i=12/a=124012>.
- [12] I. Krebs, *et al.* Physics of Plasmas (1994-present) **20(8)**, 082506 (2013), URL <http://scitation.aip.org/content/aip/journal/pop/20/8/10.1063/1.4817953>.
- [13] S. Pamela, *et al.* Plasma Physics and Controlled Fusion **55**, 095001 (13 pp.) (2013), URL <http://iopscience.iop.org/0741-3335/55/9/095001/>.
- [14] R. G. Kleva *et al.* Physics of Plasmas (1994-present) **14(1)**, 012303 (2007), URL <http://scitation.aip.org/content/aip/journal/pop/14/1/10.1063/1.2424560>.
- [15] A. Thyagaraja, *et al.* Physics of Plasmas (1994-present) **17(4)**, 042507 (2010), URL <http://scitation.aip.org/content/aip/journal/pop/17/4/10.1063/1.3381074>.
- [16] F. Orain, *et al.* Physics of Plasmas **20(10)**, 102510 (2013), URL <http://scitation.aip.org/content/aip/journal/pop/20/10/10.1063/1.4824820>.
- [17] M. Swisdak, *et al.* The Astrophysical Journal **710(2)**, 1769 (2010), URL <http://stacks.iop.org/0004-637X/710/i=2/a=1769>.
- [18] J. F. Drake, *et al.* Physics of Fluids (1958-1988) **26(9)** (1983).
- [19] P. H. Diamond, *et al.* Physics of Fluids (1958-1988) **28(4)** (1985).
- [20] G. T. A. Huysmans, *et al.* Physics of Plasmas (1994-present) **8(10)** (2001).
- [21] F. D. Halpern, *et al.* Physics of Plasmas (1994-present) **18(10)**, 102501 (2011), URL <http://scitation.aip.org/content/aip/journal/pop/18/10/10.1063/1.3646305>.
- [22] P. Beyer, *et al.* Phys. Rev. Lett. **94**, 105001 (2005), URL <http://link.aps.org/doi/10.1103/PhysRevLett.94.105001>.
- [23] G. Huysmans *et al.* Nuclear Fusion **47(7)**, 659 (2007), URL <http://stacks.iop.org/0029-5515/47/i=7/a=016>.
- [24] R. Pitts, *et al.* Nuclear Fusion **47(11)**, 1437 (2007), URL <http://stacks.iop.org/0029-5515/47/i=11/a=005>.
- [25] T. Eich, *et al.* (ASDEX Upgrade Team). Phys. Rev. Lett. **91**, 195003 (2003), URL <http://link.aps.org/doi/10.1103/PhysRevLett.91.195003>.
- [26] M. Bécoulet, *et al.* Nuc. Fus. **52(5)**, 054003. 5th International Workshop on Stochasticity in Fusion Plasmas (SFP), Julich, GERMANY, APR 11-14, 2011 (2012), URL <http://iopscience.iop.org/0029-5515/52/5/054003/>.