

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Trends and drivers of regional sources and sinks of carbon dioxide over the past two decades

S. Sitch¹, P. Friedlingstein¹, N. Gruber², S. D. Jones³, G. Murray-Tortarolo¹,
A. Ahlström⁴, S. C. Doney⁵, H. Graven², C. Heinze^{6,7,8}, C. Huntingford⁹,
S. Levis¹⁰, P. E. Levy¹¹, M. Lomas¹², B. Poulter¹³, N. Viovy¹³, S. Zaehle¹⁴,
N. Zeng¹⁵, A. Arneeth¹⁶, G. Bonan¹⁰, L. Bopp¹³, J. G. Canadell¹⁷, F. Chevallier¹³,
P. Ciais¹³, R. Ellis⁹, M. Gloor¹⁸, P. Peylin¹³, S. Piao¹⁹, C. Le Quéré³, B. Smith⁴,
Z. Zhu^{20,21}, and R. Myneni²²

¹University of Exeter, Exeter EX4 4QF, UK

²Institute of Biogeochemistry and Pollutant Dynamics, ETH Zurich, Zürich, Switzerland

³Tyndall Centre for Climate Change Research, University of East Anglia, Norwich NR4 7TJ, UK

⁴Lund University, Sölvegatan 12, 223 62 Lund, Sweden

⁵Marine Chemistry and Geochemistry Department, Woods Hole Oceanographic Institution, 266 Woods Hole Road, Woods Hole, MA 02543, USA

⁶Geophysical Institute, University of Bergen, Bergen, Norway

⁷Bjerknes Centre for Climate Research, Bergen, Norway

⁸Uni Climate, Uni Research AS, Allégaten 70, 5007 Bergen, Norway

20113

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

⁹Centre for Ecology and Hydrology, Benson Lane, Wallingford OX10 8BB, UK

¹⁰National Center for Atmospheric Research, Boulder, Colorado, USA

¹¹Centre for Ecology and Hydrology, Bush Estate, Penicuik, Midlothian EH26 0QB, UK

¹²Department of Animal and Plant Sciences, University of Sheffield, Sheffield S10 2TN, UK

¹³Laboratoire des Sciences du Climat et de l'Environnement, CEA CNRS UVSQ, 91191 Gif-sur-Yvette, France

¹⁴Biogeochemical Integration Department, Max Planck Institute for Biogeochemistry, P.O. Box 10 01 64, 07701 Jena, Germany

¹⁵Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD 20740, USA

¹⁶Karlsruhe Institute of Technology, Garmisch-Partenkirchen, Germany

¹⁷Global Carbon Project, CSIRO Marine and Atmospheric Research, Canberra, Australia

¹⁸University of Leeds, School of Geography, Woodhouse Lane, LS9 2JT, Leeds, UK

¹⁹College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

²⁰State Key Laboratory of Remote Sensing Science, Institute of Remote Sensing and Digital Earth, Chinese Academy of Sciences, Beijing, 100101, China

²¹Center for Applications of Spatial Information Technologies in Public Health, Beijing, 100101, China

²²Department of Geography and Environment, Boston University, 675 Commonwealth Avenue, Boston, MA 02215, USA

Received: 21 November 2013 – Accepted: 26 November 2013

– Published: 23 December 2013

Correspondence to: S. Sitch (s.a.sitch@exeter.ac.uk)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Abstract

The land and ocean absorb on average over half of the anthropogenic emissions of carbon dioxide (CO₂) every year. These CO₂ “sinks” are modulated by climate change and variability. Here we use a suite of nine Dynamic Global Vegetation Models (DGVMs) and four Ocean Biogeochemical General Circulation Models (OBGCMs) to quantify the global and regional climate and atmospheric CO₂ – driven trends in land and oceanic CO₂ exchanges with the atmosphere over the period 1990–2009, attribute these trends to underlying processes, and quantify the uncertainty and level of model agreement. The models were forced with reconstructed climate fields and observed global atmospheric CO₂; Land Use and Land Cover Changes are not included for the DGVMs. Over the period 1990–2009, the DGVMs simulate a mean global land carbon sink of $-2.4 \pm 0.7 \text{ Pg C yr}^{-1}$ with a small significant trend of $-0.06 \pm 0.03 \text{ Pg C yr}^{-2}$ (increasing sink). Over the more limited period 1990–2004, the ocean models simulate a mean ocean sink of $-2.2 \pm 0.2 \text{ Pg C yr}^{-1}$ with a trend in the net C uptake that is indistinguishable from zero ($-0.01 \pm 0.02 \text{ Pg C yr}^{-2}$). The two ocean models that extended the simulations until 2009 suggest a slightly stronger, but still small trend of $-0.02 \pm 0.01 \text{ Pg C yr}^{-2}$. Trends from land and ocean models compare favourably to the land greenness trends from remote sensing, atmospheric inversion results, and the residual land sink required to close the global carbon budget. Trends in the land sink are driven by increasing net primary production (NPP) whose statistically significant trend of $0.22 \pm 0.08 \text{ Pg C yr}^{-2}$ exceeds a significant trend in heterotrophic respiration of $0.16 \pm 0.05 \text{ Pg C yr}^{-2}$ – primarily as a consequence of wide-spread CO₂ fertilisation of plant production. Most of the land-based trend in simulated net carbon uptake originates from natural ecosystems in the tropics ($-0.04 \pm 0.01 \text{ Pg C yr}^{-2}$), with almost no trend over the northern land region, where recent warming and reduced rainfall offsets the positive impact of elevated atmospheric CO₂ on carbon storage. The small uptake trend in the ocean models emerges because climate variability and change, and in particular increasing sea surface temperatures, tend to counteract the trend in ocean

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



torial simulations to elucidate the processes that drive trends in the sources and sinks of atmospheric CO₂.

This study has two major aims. First, to quantify the trends in the carbon exchange over the period 1990–2009, associated with changes in climate and atmospheric CO₂ concentration, for three land regions and seven ocean regions (Fig. 1): Northern, Tropics, and Southern land, and North Pacific, North Atlantic, Tropical Indo-Pacific, Tropical Atlantic, South Indo-Pacific, South Atlantic, Southern Ocean. Second, to determine which factors and processes among those included in the models are driving the modeled/observed trends in the regional land/ocean to atmosphere net CO₂ fluxes. For the land models these include the CO₂ fertilization effect on productivity and storage, and climate effects on productivity, respiration and climate-caused natural disturbances (see Table S1 for details represented in individual models). A particular focus is on the impacts of climate variation and change on land ecosystems at the regional scale as extreme climate events have occurred over the 1990–2009 period across many regions of the World, e.g. including North America (South West USA, 2000–2002), Europe (2003), Amazonia (2005), and eastern Australia (2001–2008), raising considerable attention in the ecological community on the consequences of recent climate variability on ecosystem structure and function (Allen et al., 2010) and the carbon cycle (Ciais et al., 2005; Van der Molen et al., 2011; Reichstein et al., 2013). Consideration of Land Use Change (LUC) on regional trends is beyond the scope of the present study. There are large uncertainties in the global LULCC flux (Le Quéré et al., 2009), and in comparison the net land use flux represents only approximately 2–3% of the large land-atmosphere fluxes associated with net primary productivity and heterotrophic respiration. The response of these large fluxes to climate variability and CO₂ are the focus of this study. In addition the net LU flux for the period 1990–2009 will be influenced by earlier LULCC (i.e. legacy fluxes), confounding the analysis. Other companion papers investigate ecosystem response to interannual and seasonal timescales (Piao et al., 2012), and the carbon balance for individual land and ocean

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



regions over the period 1990–2009 (see RECCAP special issue, Canadell et al., 2013, http://www.biogeosciences.net/special_issue107.html).

Trends and variability in the air–sea CO₂ fluxes simulated by the employed OBGCMs are driven by the increase in atmospheric CO₂ and by variability and change in ocean temperature, circulation, winds, and biology largely governed by climate variability. The air–sea CO₂ flux arising from the increase in atmospheric CO₂ is often referred to as the flux of anthropogenic CO₂, while the remainder, induced by changes in the natural cycling of carbon in the ocean–atmosphere system is called the “natural” CO₂ component (e.g., Gruber et al., 2009). Although this conceptual separation has its limits (McNeill and Matear, 2013), it provides for a powerful way to understand how different forcings affect the net ocean sink.

DGVM results are compared with estimates of the Residual Land Sink and with remote sensing products indicating trends of greening and browning in the northern region. Regional sources and sink trends are attributed to processes based on factorial simulations.

2 Methods

2.1 Dynamic Global Vegetation Models

Following the studies of Le Quéré et al. (2009) and Sitch et al. (2008), a consortium of Dynamic Global Vegetation Model (DGVM) groups set up a project to investigate further the spatial trends in land–atmosphere flux and agreed to perform a factorial set of DGVM simulations over the historical period, 1901–2009. These simulations have contributed to the RECCAP activity (Canadell et al., 2011, 2013). There are now a variety of DGVMs with origins in different research communities that typically contain alternative parameterisations and a diverse inclusion of processes (Prentice et al., 2007; Piao et al., 2013). DGVMs have emerged from the Land Surface Modelling (LSM), forest ecology, global biogeography, and global biogeochemical modelling communities.

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



inal global mean gas transfer velocity of Broecker et al. (1985) was too high (Sweeney et al., 2007; Müller et al., 2008).

2.3 Datasets

2.3.1 Land

Climate forcing is based on a merged product of Climate Research Unit (CRU) observed monthly 0.5° climatology (v3.0, 1901–2009; New et al., 2000) and the high temporal fidelity NCEP reanalysis forcing. The merged product has a 0.5° spatial and 6 hourly temporal resolution. A coarse resolution $3.75^\circ \times 2.5^\circ$ version at monthly time scales was also provided (see Table 1 for spatial resolution of individual DGVMs).

Global atmospheric CO_2 was derived from ice core and NOAA monitoring station data, and provided at annual resolution over the period, 1860–2009. As land use change was not simulated in these model experiments, models assume a constant land use throughout the simulation period. Atmospheric nitrogen deposition data for CLM4CN and OCN was sourced from Jean-Francois Lamarque (personal communication, 2013) and Dentener et al. (2006), respectively.

Gridded fields of Leaf Area Index (LAI) are used in the evaluation of DGVM northern greening trends. These LAI datasets based on remote sensing data were generated from the AVHRR GIMMS NDVI3g using an Artificial Neural Network derived model (Zhu et al., 2012). The dataset has a temporal resolution of 15 days over the period 1981–2010, and a spatial resolution of $1/12^\circ$.

2.3.2 Ocean

Unlike how the land models simulations were set up, no common climatic forcing dataset was used for the ocean model simulations. In fact, some models even provided results with different climatic forcings. Models were forced by the NCEP climatic data (Kalnay et al., 1996) in its original form, or in the modified CORE (Common Ocean-ice

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO_2 over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Reference Experiments – Corrected Normal Year Forcing (CORE-CNYF; Large and Yeager, 2004) form.

2.3.3 Atmospheric inversion

5 Simulated trends are compared with those from version 11.2 of the CO₂ inversion product from the Monitoring Atmospheric Composition and Climate – Interim Implementation (MACC-II) service (<http://www.gmes-atmosphere.eu/>). It covers years 1979–2011 and a previous release has been documented by Chevallier et al. (2010). It uses a climatological prior without inter-annual variability, except for fossil fuel.

2.4 Experimental design

10 2.4.1 Land

The land models were forced over the 1901–2009 period with changing CO₂, climate and fixed present-day land use according to the following simulations:

S_L1: changing CO₂ only (time-invariant present-day land use mask, fixed pre-industrial climate).

15 S_L2: changing CO₂ and climate (time-invariant present-day land use mask).

In both cases, DGVMs spin-up were initially performed, with pre-industrial CO₂, and climate. For DGVMs including the N cycle, N deposition was a time-variant forcing in both simulations, such that the difference between S_L2 and S_L1 includes the synergistic effects of N deposition on CO₂ fertilisation (Zaehle et al., 2010).

20 Figure 2 shows the historical changes in climate, atmospheric CO₂ concentration, nitrogen deposition over the period 1990–2009 used to force the DGVMs. A summary of DGVM characteristics is given in Table 1. A more detailed description of DGVM process representations is given in Table A1.

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

2.4.2 Ocean

The ocean models have employed two different strategies for creating the initial conditions for the experiments. The first strategy, followed by CCSM-ETH, CCSM-WHOI and BER, involved first a several century long spinup with climatological forcing and with atmospheric CO₂ held constant at its pre-industrial value, bringing these models very close to a climatological steady-state for preindustrial conditions (in some models ~ 1750; in others ~ 1850). In the second step, the models are then integrated forward in time through the historical period until 1948, with atmospheric CO₂ prescribed to follow the observed trend and a climatological forcing. The length of the spinup varies between a few hundred years to several thousand years, resulting in differing global integrated drift fluxes, although their magnitudes are substantially smaller than 0.05 PgCyr⁻¹ with essentially no rate of change. The second strategy, followed by the University of East Anglia, was to initialize the model with reconstructed initial conditions in 1920, and then also run it forward in time until 1948 with prescribed atmospheric CO₂, repeating the daily forcing conditions of a single year (1980). The modelled export production was tuned to obtain an ocean CO₂ sink of 2.2 PgCyr⁻¹ in the 1990s. This second method offers the advantage that the model's carbon fields remain closer to the observations compared to the long spinup approach, but it comes at the cost of generating a drift that affects the mean conditions and to a lesser extent the trend. Tests with the model runs of Le Quéré et al. (2010) suggests the drift in the mean CO₂ sink is about 0.5 PgCyr⁻¹ and the drift in the trend is about 0.005 PgCyr⁻² globally, and is largest in the Southern Ocean. In these runs, both S_O1 and S_O2 are affected by the same drift, and their differences thus removes the drift. From ~ 1950 onward, the models performed two separate simulations:

S_O1: CO₂ only, i.e., atmospheric CO₂ increases, but models are forced with climatological atmospheric boundary conditions (referred to as ACO2 in the RECCAP archive).

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



S_O2: CO₂ and climate, i.e., as S_O1, but models are forced with “realistic” year-to-year variability in atmospheric boundary conditions (ANTH).

The CCSM-based models performed an additional experiment to better separate between the fluxes of natural and anthropogenic CO₂:

5 S_O3: pre-industrial and climate, i.e., atmospheric CO₂ is fixed at its pre-industrial level, but atmospheric boundary conditions vary as in S_O2 (PIND).

From these simulations, only the results from 1990 through 2009 were analysed. Only the UEA and CCSM-WHOI models made available results for the S_O1 and S_O2 simulations for the entire analysis time. The results for the BER model for 2009 are
10 incomplete and the CCSM-ETH simulations extend only to 2007. In order to maintain a sufficiently large set of models, we decided to focus our analysis primarily on the 1990 through 2004 period, but include occasionally also the results through 2009, with the important caveat that the latter are based only on two models.

2.5 Output variables

15 2.5.1 Land

Output variables include those associated with the carbon, hydrological and energy balance over the period 1901–2009. Variables were considered either “Level 1”, and essential, or “Level 2”, desirable, for additional analysis/studies. Output variables were typically at either the monthly or annual resolution, and at the spatial resolution of the
20 individual model application. In this study we analyse a sub-set of the DGVM output, and primarily focus on the carbon cycle variables, Net Primary Productivity (NPP), Heterotrophic Respiration (RH), and the Net Biome Productivity (NBP), and Leaf Area Index (LAI), a measure of vegetation greenness. The land to atmosphere net CO₂ flux is equal in magnitude to the NBP but has the opposite sign, i.e. we adopt the sign
25 convention where a negative value for the net CO₂ flux represents a carbon sink. Here

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



we define NBP as:

$$\text{Land to atmosphere net CO}_2 \text{ flux} = -\text{NBP} = \text{RH} + \text{Wildfire Flux} + \text{Riverine C Flux} \\ + \text{harvest} - \text{NPP}$$

5 DGVMs typically do not represent all these processes; a list for each individual DGVM is given in Table 1. DGVM results for simulation S2 are compared against the global Residual Land Sink (RLS), calculated as the annual anthropogenic CO₂ emissions (fossil fuel, cement manufacture and land use C flux) minus the annual CO₂ growth rate and model mean ocean C sink as given by Friedlingstein et al. (2010). The ocean
10 uptake is from the same OGGCMs as the ones used here, and the land use C flux is based on a book-keeping approach from Houghton (2010).

The regional analysis will focus on 3 large regions: northern, tropical and southern land regions (Fig. 1). Within these regions, trends at a finer spatial resolution, from multi-grid cell to the sub-region are analysed. The Northern land region is divided by continent into sub-regions, North America, Europe and North Asia, and into ecoregions, Temperate and Boreal North America and Asia, and the Tundra biome. Tropical Land is divided into four regions: Tropical South America Forests, Tropical Asia, Equatorial Africa, North Africa Savanna. Likewise Southern Land is divided into four sub-regions: South America Savanna, Temperate South America, Southern Africa, Australia and New Zealand.
20

The comparison of DGVM simulated trends in the northern growing season against satellite-derived NDVI observations was based on 8 models (JULES, LPJ, LPJ-GUESS, NCAR-CLM4, ORCHIDEE, OCN, SDGVM, VEGAS), which provided Leaf Area Index outputs (LAI). The means and trends in the onset, offset and length of growing season were computed. Growing season variables were calculated using the methodology of Murray-Tortarolo et al. (2013). Leaf onset is defined as the day (t) when LAI begins to increase above a critical threshold (CT), defined as:
25

$$\text{CT} = \text{LAI}_{\min} + 0.2 \cdot (\text{LAI}_{\max} - \text{LAI}_{\min})$$

20125

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



where LAI_{\min} and LAI_{\max} represent the minimum and maximum LAI over the annual cycle. Similarly, leaf senescence, or offset, is defined as the day when LAI decreases below the CT. The length of the growing season in days is calculated as the offset minus the onset. This calculation was made for each gridcell above 30° N (i.e. northern extra-tropics) from the models and the satellite data. In addition, any gridcell where LAI varied by less than 0.5 over the annual cycle from the satellite data was considered to be predominantly evergreen (e.g. Boreal Forest), and not considered in the analysis. We also masked out regions where LAI decreases in the summer (drought deciduous vegetation). In addition, when the growing season spans over the end of year (e.g. Mediterranean and some pixels particularly on the southern margin of the domain), we include the first 3 months of the second year in our analysis. Means and trends were calculated using a linear model over the period 1990–2009.

2.5.2 Ocean

The modelling groups provided output on a monthly basis for the years 1990 through 2004 and 2009, respectively, at two levels of priority. Tier one data included the surface ocean fields of the air–sea CO_2 flux, oceanic pCO_2 , dissolved inorganic carbon (DIC), alkalinity (Alk), temperature (T), salinity (S), and mixed layer depth. The second tier data included the biological export at 100 m, the vertically integrated net primary production, and the surface chlorophyll a concentration. Some models supplied also three-dimensional climatological fields of DIC, Alk, temperature, and salinity.

To determine the different factors contributing to the modelled trends and variations, we undertook two (linear) separations:

The contribution of climate variability on the ocean carbon cycle: $X_{\text{var}} = X(S_{O2}) - X(S_{O1})$, where X is any state variable or flux, where the expression in parentheses represents the results of the corresponding simulation, and where X_{var} represents the impact of climate change and variability on the ocean carbon cycle.

The contribution of anthropogenic CO_2 : $X_{\text{ant}} = X(S_{O2}) - X(S_{O3})$.

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO_2 over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

For each of the integrations, but particularly for the changing CO₂ and climate simulation S_O2, we analysed the factors contributing to the temporal change in the air–sea CO₂ flux F by a linear Taylor expansion (see e.g., Lovenduski et al. (2007) and Doney et al., 2009a):

$$\Delta F = \frac{\partial F}{\partial ws} \cdot \Delta ws + \frac{\partial F}{\partial T} \cdot \Delta T + \frac{\partial F}{\partial ice} \cdot \Delta ice + \frac{\partial F}{\partial sDIC} \cdot \Delta sDIC + \frac{\partial F}{\partial sAlk} \cdot \Delta sAlk + \frac{\partial F}{\partial FS} \cdot \Delta S$$

where ws is the wind speed, ice is the sea–ice fraction, $sDIC$ and $sAlk$ are the salinity normalized DIC and Alk concentrations, and where $\frac{\partial F}{\partial FS}$ is the change in the air–sea CO₂ flux in response to freshwater fluxes. This latter term includes not only the sensitivity of oceanic pCO_2 to changes in salinity, but also the dilution effects of freshwater on DIC and Alk (see Doney et al., 2009a for details). The partial derivatives were computed directly from the model equations for the mean conditions in each region. The changes in the driving components were derived from the trend computed via a linear regression of the model results and then multiplied by the length of the timeseries.

3 Results

3.1 Global trends

3.1.1 Land

The ensemble mean global land to atmosphere net carbon dioxide flux from S_L2 is $-2.38 \pm 0.72 \text{ PgCyr}^{-1}$ over the period 1990–2009 ($P = 0.04$) (Figs. 3 and A1, Table 2). The numbers behind \pm signs are the standard deviation, calculated as statistic of the 20 yr means for 9 DGVMs. This compares to the global RLS of $-2.45 \pm 1.17 \text{ PgCyr}^{-1}$, inferred as a residual from other terms in the global carbon budget by Friedlingstein et al. (2010) over the same period. All DGVMs agree on an increasing land sink with

20127

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

a net flux trend over this period ranging between -0.02 and -0.11 PgCyr^{-2} , corresponding to DGVMs, OCN and Hyland, respectively (Table 2). DGVMs simulate an increase in the land C sink with an ensemble mean trend of $-0.06 \pm 0.03 \text{ PgCyr}^{-2}$ ($P < 0.05$) over the period 1990 to 2009 (Table 2), in response to changes in climate and atmospheric CO₂ content. The two DGVMs with a fully coupled carbon and nitrogen cycle (CN), OCN and CLM4CN, still simulate an increase in the land sink, but with a trend below the ensemble mean, at -0.02 ($P = 0.6$) and -0.05 PgCyr^{-2} ($P = 0.06$), respectively. DGVMs suggest the increase in global land sink between 1990 and 2009 is driven by increases in simulated global NPP (Fig. 2).

DGVMs simulate an ensemble mean global NPP of $62.9 \pm 8.73 \text{ PgCyr}^{-1}$ over the period 1990 and 2009 (Table 2). All DGVMs simulate an increase in NPP over this period, with an ensemble mean DGVM trend in NPP of $0.22 \pm 0.08 \text{ PgCyr}^{-2}$ ($P = 0.00$) (Table 2). Also models with a higher NPP trend also produce a higher NBP trend (Fig. A2). VEGAS, and the two CN models, CLM4CN and OCN, simulate the smallest positive trends in global NPP of 0.11, 0.15 and 0.16 PgCyr^{-2} , respectively (Table 2). The ensemble mean NPP trend of $0.22 \pm 0.08 \text{ PgCyr}^{-2}$ ($P = 0.00$) from simulation, S_L2 (CO₂ and climate forcing) contrasts with an ensemble trend of $0.19 \pm 0.08 \text{ PgCyr}^{-2}$ ($P = 0.00$), and $0.03 \pm 0.05 \text{ PgCyr}^{-2}$ ($P = 0.24$) over the same period for the S_L1 (CO₂ only) and S_L2–S_L1 (the climate effect), respectively (Tables A2 and A3). These results suggest that the simulated increase in global NPP is mainly in response to increasing atmospheric CO₂ (direct CO₂ fertilization of photosynthesis, in addition to the indirect benefits from an improved water balance in water-limited ecosystems due to the physiological effects of CO₂ on water use efficiency). Again, VEGAS, CLM4CN and OCN, simulate the smallest positive trends in NPP among the DGVMs in response to elevated CO₂ forcing (Table A2). This suggests that the potential CO₂ fertilization effect may be already strongly limited by present day nitrogen availability in some ecosystems (Vitousek and Howarth, 1991). There is more uncertainty among models on the impact of climate changes on global NPP, with only 2 models simulating a significant positive trend (Table A3).

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

DGVMs simulate an ensemble mean global RH of $57.5 \pm 9.8 \text{ PgCyr}^{-1}$ over the period 1990 and 2009 (Table 2). All DGVMs simulate an increase in RH for S2 (CO₂ and climate), with an ensemble mean trend of $0.16 \pm 0.05 \text{ PgCyr}^{-2}$ ($P = 0.00$) over the period 1990 to 2009 (Table 2). This is lower than the trend in global NPP, thus explaining the trend towards increasing net land carbon uptake. This is unsurprising as there is a lagged response in increases in RH relative to NPP, reflecting the turnover time of the newly incorporated plant material. The ensemble mean trend in RH is $0.12 \pm 0.06 \text{ PgCyr}^{-2}$ ($P = 0.00$) and $0.04 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.09$) over the same period for the S1 (CO₂ only) and S_L2-S_L1 (the climate effect), respectively (Tables A2, A3), implying the dominant effect on RH is increased substrate for microbial respiration, with the additional litter input into soils, as a consequence of enhanced NPP, rather than enhanced rates of microbial decomposition with rising temperatures. Nevertheless, the simulated mean residence time (MRT = Soil Carbon/RH) of soil organic matter decreases, in response to warming, which is especially pronounced in high latitude regions (Fig. A3). The differences in land–atmosphere flux trends for the CN models, OCN (-0.02 PgCyr^{-2}) and CLM4CN (-0.05 PgCyr^{-2}) are largely due their differences in RH trends at 0.14 and 0.11 PgCyr^{-2} , respectively, rather than differential responses of simulated NPP to elevated CO₂ (Table 2).

Only 4 DGVMs provided wildfire fluxes (CLM4CN, LPJ, LPJ-GUESS, SDGVM). No significant trends in the global wildfire flux were reported by any of the DGVMs. Only two DGVMs (CLM4CN and LPJ) simulated small significant trends of opposite sign in the wildfire flux to the 90 % confidence interval for S_L1 (CO₂ only), possibly reflecting the counteracting effects of CO₂ on plant water use, leading to more moist land surface (reduced flammability), and elevated productivity (increased fuel loads).

3.1.2 Ocean

The global ocean is simulated to have acted as a very substantial sink for atmospheric CO₂, but one that has increased only slightly over the last two decades (see also

discussion in Wanninkhof et al., 2013). The mean ocean sink in the 4 models (CCSM-ETH, CCSM-WHOI, UEA and BER), increased from $\sim -2.0 \text{ PgCyr}^{-1}$ in the early 1990s to $\sim -2.1 \text{ PgCyr}^{-1}$ during the first five years of the 21st century (Fig. 4). The overall sink is largely a consequence of the increase in atmospheric CO_2 (i.e., it corresponds mostly to the uptake flux of anthropogenic CO_2), but it includes a substantial perturbation flux stemming from the impact of climate variability and change on the ocean carbon cycle.

This is confirmed if we separate the mean and variable components by using our factorial experiments, i.e., by using S_O1 results to identify the ocean uptake in the absence of climate variability and change, and the difference between S_O2 and S_O1 as measure of the impact of climate change. This separation reveals that in the absence of climate variability and change, the global ocean uptake would have increased from about $-1.98 \pm 0.04 \text{ PgCyr}^{-1}$ for the 1990–1994 period to $-2.3 \pm 0.09 \text{ PgCyr}^{-1}$ for 2000–2004 (for the two models CCSM-WHOI and UEA that provided S_O1 results up to 2009, the uptake flux would have increased from -1.99 PgCyr^{-1} to -2.56 PgCyr^{-1} for 2005–2009). This global net uptake flux and its substantial trend in time (-0.03 PgCyr^{-2} for 1990–2004, and -0.04 PgCyr^{-2} for 1990–2010) is entirely driven by the increase of atmospheric CO_2 and is – integrated globally – numerically equivalent to the ocean uptake flux of anthropogenic CO_2 . Climate variability and change modified these fluxes, and particularly the trend in these models. The four models suggest an enhancement of the uptake in the early 1990s (1990–1994) of about -0.2 PgCyr^{-1} , turning into a reduction of the uptake in the subsequent period (1995–1999), followed by a further reduction in the 2000–2004 period of $\sim +0.1 \text{ PgCyr}^{-1}$. This trend toward reduced uptake in response to climate variability and change of $+0.03 \text{ PgCyr}^{-2}$ nearly completely compensates for the anthropogenic CO_2 driven increase in uptake, causing the overall uptake of CO_2 to have a nearly flat trend over the 1990 through 2004 period of $< 0.01 \text{ PgCyr}^{-2}$. The same tendencies are found for the two models that extend over the entire 1990 through 2009 period: climate change and variability reduces in these models the CO_2 driven trend of -0.04 PgCyr^{-2} by more than $+0.02 \text{ PgCyr}^{-2}$, to around -0.02 PgCyr^{-2} .

Trends and drivers of regional sources and sinks of CO_2 over the past two decades

S. Sitch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Consideration of the different factors affecting the ocean carbon sink following our Taylor expansion, we find increasing sea-surface temperature to be globally one of the most important drivers for the positive trends (reduced sinks) induced by climate change and variability. Over the 1990 through 2004 period, the surface ocean warmed, on average, by $0.004\text{ }^{\circ}\text{C yr}^{-1}$ ($0.005\text{ }^{\circ}\text{C yr}^{-1}$ from 1990 through 2009). Isochemically, this leads to an increase of the oceanic $p\text{CO}_2$ of $\sim 0.06\text{ }\mu\text{atm yr}^{-1}$, which appears small. However, it needs to be compared with the trend in the global mean air–sea $p\text{CO}_2$ difference of about $\sim 0.1\text{ }\mu\text{atm yr}^{-1}$ that is required in order to generate a trend in the ocean uptake of -0.03 PgC yr^{-2} (see e.g., Matsumoto and Gruber, 2005; Sarmiento and Gruber, 2006).

3.2 Regional trends

3.2.1 Land

Spatial correlations between trends in drivers and carbon fluxes (NBP, NPP, RH) were significant for precipitation (0.36, 0.5, 0.48 respectively; $P < 0.05$), but not for temperature (see Table A11), underlining the sensitivity of carbon cycle models to precipitation changes at the multi-year to decadal timescale.

3.2.2 Northern land

All DGVMs agree on a land C sink over the northern land region, with a mean land–atmosphere flux of $-1.03\pm 0.30\text{ PgC yr}^{-1}$, over the period 1990–2009 (Fig. A4, Table 2). The ensemble mean land–atmosphere flux trend is near zero for this region between 1990 and 2009 (Fig. A5). Although 4 of 9 DGVMs simulate reductions in the land sink over the northern land region, none of the trends from the 9 DGVMs are significant at the 95% confidence level (Table 2). Of particular interest are sub-regions with a simulated positive land–atmosphere flux trend (Fig. 5), implying a diminishing sink of atmospheric CO_2 , or an increasing source of CO_2 to the atmosphere. At least 6 mod-

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitsh et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

els out of 9 agree on a decreasing regional land sink across some areas in Temperate North America, Eastern Europe, North-East China and Mongolia (Fig. 5). These net flux trends largely correspond to regions with negative trends in precipitation (Fig. 6).

Over the northern region, DGVMs simulate an ensemble mean NPP of $24.1 \pm 4.48 \text{ PgCyr}^{-1}$, which represents almost 40 % of the global total (Table 2). All DGVMs simulate an increase in northern NPP over this period, with a trend in NPP of $0.06 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.00$) (Table 2). However, enhanced productivity in the northern land region accounts for only around 29 % of the simulated global trend in NPP. The ensemble mean NPP trend of $0.06 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.00$) from simulation S2 (CO₂ and climate forcing) compares to a trend of $0.07 \pm 0.03 \text{ PgCyr}^{-2}$ ($P = 0.00$), and $-0.00 \pm 0.04 \text{ PgCyr}^{-2}$ ($P = 0.85$) for the S1 (CO₂ only) and S2–S1 (the climate effect), respectively (Tables A2, A3). All DGVMs simulate a positive trend in NPP in response to elevated CO₂ across the northern land region and trends are all significant to the 95 % confidence level with the exception of CLM4CN ($P = 0.21$). Large areas in Temperate North America and Asia have experienced warming combined with reductions in precipitation over the period 1990–2009 (Fig. 5).

The ensemble mean NPP for North America, Europe and North Asia are $7.78 \pm 1.38 \text{ PgCyr}^{-1}$, $5.08 \pm 1.40 \text{ PgCyr}^{-1}$, and $11.20 \pm 1.99 \text{ PgCyr}^{-1}$, respectively (Table A5). Despite the two-fold difference in mean NPP, NPP trends for North America, Europe and North Asia are similar, at $0.021 \pm 0.008 \text{ PgCyr}^{-2}$ ($P = 0.02$), $0.018 \pm 0.006 \text{ PgCyr}^{-2}$ ($P = 0.00$) and $0.024 \pm 0.015 \text{ PgCyr}^{-2}$ ($P = 0.01$), respectively (Table A5). DGVMs simulate larger mean NPP in Temperate compared to Boreal regions with mean NPP in Temperate North America and Asia of $4.22 \pm 0.83 \text{ PgCyr}^{-1}$ and $7.09 \pm 1.03 \text{ PgCyr}^{-1}$, respectively, compared to $3.57 \pm 1.21 \text{ PgCyr}^{-1}$ and $4.12 \pm 1.46 \text{ PgCyr}^{-1}$ in Boreal North America and Asia, respectively (Table A5). DGVMs simulate a significant positive trend in Boreal North America and Boreal Asia of $0.014 \pm 0.007 \text{ PgCyr}^{-2}$ ($P = 0.02$) and $0.018 \pm 0.006 \text{ PgCyr}^{-2}$ ($P = 0.01$), respectively. DGVM NPP trends in both Temperate North America and Asia are smaller than those for Boreal regions but are not significant at the 95 % confidence level (Table A5).

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



In response to warming, models simulate an earlier onset (ensemble mean model trend = $-0.078 \pm 0.131 \text{ dayyr}^{-1}$) and delayed termination of the growing season ($0.217 \pm 0.097 \text{ dayyr}^{-1}$) based on LAI, and thus a trend towards a longer growing season in the northern extra-tropics ($0.295 \pm 0.228 \text{ dayyr}^{-1}$) (Fig. 7). This is in broad agreement with observed greening trends: onset = -0.11 dayyr^{-1} ; offset = 0.252 dayyr^{-1} and growing season length = 0.361 dayyr^{-1}). There is less agreement among models on reproducing the observed browning trends in some regions of the boreal forest.

DGVMs simulate an ensemble mean RH of $21.8 \pm 4.6 \text{ PgCyr}^{-1}$ across the northern land region (Table 2). All DGVMs simulate an increase in northern RH over the period 1990–2009, with a significant trend in RH of $0.063 \pm 0.02 \text{ PgCyr}^{-1}$ ($P = 0.00$) (Table 2). DGVMs simulate larger mean RH in Temperate compared to Boreal regions with RH in Temperate North America and Asia of $3.78 \pm 0.93 \text{ PgCyr}^{-1}$ and $6.33 \pm 1.13 \text{ PgCyr}^{-1}$, respectively, compared to $3.26 \pm 1.08 \text{ PgCyr}^{-1}$ and $3.78 \pm 1.35 \text{ PgCyr}^{-1}$ in Boreal North America and Asia, respectively (Table A6). DGVMs simulate a significant positive trend in Boreal North America and Boreal Asia of $0.012 \pm 0.003 \text{ PgCyr}^{-2}$ ($P = 0.02$) and $0.015 \pm 0.004 \text{ PgCyr}^{-2}$ ($P = 0.01$), respectively (Table A6). DGVM RH trends in both Temperate North America and Asia are smaller at $0.008 \pm 0.005 \text{ PgCyr}^{-2}$ ($P = 0.01$) and $0.010 \pm 0.009 \text{ PgCyr}^{-2}$ ($P = 0.08$), respectively than those for boreal regions and both significant at the 90 % confidence level. This is because of relatively smaller increases in substrate (i.e. NPP), in temperate regions, and greater warming in boreal regions stimulating microbial decomposition, reducing mean residence time of carbon in soils (MRT = Soil Carbon/RH; see Fig. A3).

No significant trends in the wildfire flux were reported by any of the DGVMs for the northern land region. However DGVMs agree on simulating a small negative trend in wildfire flux across Boreal North America and Tundra, with ensemble mean trends of $-0.002 \pm 0.003 \text{ PgCyr}^{-2}$ ($P = 0.01$) and $-0.001 \pm 0.002 \text{ PgCyr}^{-2}$ ($P = 0.08$), respectively at the 90 % confidence level.

3.2.3 Tropical Land

All DGVMs simulate a land C sink over the last two decades, in response to climate variability and changes in atmospheric CO₂ concentration, with an ensemble mean land–atmosphere flux of $-0.96 \pm 0.43 \text{ PgCyr}^{-1}$ over the Tropical land region (Table 2, Fig. A4). All DGVMs simulate an increasing land sink, with an ensemble mean trend of $-0.04 \pm 0.01 \text{ PgCyr}^{-2}$ ($P = 0.04$) for this region (Table 2, Fig. A5). The ensemble mean NBP trend over the tropical land region is significant ($P < 0.05$) and represents 65 % of the increase in land sink over the last two decades. DGVMs simulate a significant land–atmosphere flux trend (i.e. increasing sink) of $-0.016 \pm 0.007 \text{ PgCyr}^{-2}$ ($P = 0.00$) and $-0.008 \pm 0.006 \text{ PgCyr}^{-2}$ ($P = 0.05$) across Tropical Asia and Equatorial Africa, respectively (Table A4). Land–atmosphere flux trends for the Tropical South American forest and North Africa Savanna regions are not significant (Table A4).

DGVMs simulate an ensemble mean NPP of $27.0 \pm 4.29 \text{ PgCyr}^{-1}$ averaged over the tropical region representing 42 % of the global total (Table 2). All DGVMs simulate a significant increase in tropical NPP over this period, with an ensemble mean trend in NPP of $0.10 \pm 0.03 \text{ PgCyr}^{-2}$ ($P = 0.00$) for S2 (Table 2). This compares to a trend of $0.09 \pm 0.03 \text{ PgCyr}^{-2}$ ($P = 0.00$), and $0.02 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.33$) over the same period for the S1 (CO₂ only) and S2–S1 (the climate effect), respectively (Tables A2 and A3). Again the simulated trend in NPP is dominated by the simulated response of ecosystems to elevated atmospheric CO₂ content. DGVMs simulate positive NPP trends of $0.038 \pm 0.015 \text{ PgCyr}^{-2}$ ($P = 0.00$), $0.031 \pm 0.011 \text{ PgCyr}^{-2}$ ($P = 0.00$), $0.024 \pm 0.01 \text{ PgCyr}^{-2}$ ($P = 0.00$), $0.01 \pm 0.008 \text{ PgCyr}^{-2}$ ($P = 0.07$), across Tropical South America forests, Tropical Asia, Equatorial Africa, and North Africa savanna, respectively (Table A5). Nevertheless there are some areas in Tropical South America and in the North Africa Savanna regions with negative trends in NPP (Fig. 6).

DGVMs simulate an ensemble mean RH of $24.49 \pm 4.75 \text{ PgCyr}^{-1}$ over the Tropical land region (Table 2). All DGVMs simulate an increase in RH over the period 1990–2009, with a significant trend in the ensemble mean RH of $0.065 \pm 0.025 \text{ PgCyr}^{-2}$ ($P =$

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

0.00). This can be attributed largely to the response of ecosystems to elevated CO₂; the ensemble mean trend in RH from S1 is $0.059 \pm 0.022 \text{ PgCyr}^{-2}$ ($P = 0.00$) (Table A2).

No significant trends in the wildfire flux were reported by any of the DGVMs for the tropical land region. However DGVMs agree on simulating a negative trend in wildfire flux across Equatorial Africa and Tropical Asia, with an ensemble means trends of $-0.004 \pm 0.002 \text{ PgCyr}^{-2}$ ($P = 0.00$) and $-0.003 \pm 0.003 \text{ PgCyr}^{-2}$ ($P = 0.03$).

3.2.4 Southern land

All DGVMs agree on a net land sink over the southern land during the last two decades, with an ensemble mean land–atmosphere flux of $-0.38 \pm 0.29 \text{ PgCyr}^{-1}$ (Table 2, Fig. A4). The ensemble mean land–atmosphere flux trend is $-0.02 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.20$) for this region (Fig. A5). Although all DGVMs simulate an increase in the land sink over the southern extra-tropics, only trends for HYL and ORC are significant at the 95 % confidence level (Table 2). Ensemble mean land–atmosphere flux trends are significant for Temperate South America, Australia and New Zealand, and Southern Africa regions at $0.005 \pm 0.005 \text{ PgCyr}^{-2}$ ($P = 0.05$), $-0.003 \pm 0.004 \text{ PgCyr}^{-2}$ ($P = 0.68$), and $-0.022 \pm 0.011 \text{ PgCyr}^{-2}$ ($P = 0.01$), respectively (Table A4). For Southern Africa, all DGVMs simulate an increase in the land sink in response to climate changes (S_L2–S_L1) over this period (5 out of 9 are significant to the 90 % confidence level), with an ensemble mean land–atmosphere flux of $-0.018 \pm 0.01 \text{ PgCyr}^{-2}$ ($P = 0.02$) (Table A7). This compares with a land–atmosphere flux trend of $-0.004 \pm 0.002 \text{ PgCyr}^{-2}$ ($P = 0.08$) in response to CO₂ forcing only (Table A7). This is likely related to an increase in precipitation over this region (Fig. 6).

In contrast, the simulated decrease in land sink for Temperate South America is associated with a decrease in precipitation over 1990–2009. All DGVMs, except HYL, simulate a decrease in land uptake in this region for the S_L2–S_L1 simulation (however only 4 out of 9 models are significant at the 90 % confidence level), and an ensemble mean land–atmosphere flux trend is $0.005 \pm 0.005 \text{ PgCyr}^{-2}$ ($P = 0.06$) (Table A8).

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There is no significant trend for the ensemble mean land sink for S_L1 over Temperate South America.

DGVMs simulate an ensemble mean NPP of $12.3 \pm 3.53 \text{ PgCyr}^{-1}$ over the southern extra-tropics, which represents $\sim 20\%$ of the global total (Table 2). All DGVMs simulate an increase in NPP over this period, with a significant ensemble mean trend of $0.05 \pm 0.03 \text{ PgCyr}^{-2}$ ($P = 0.01$), i.e. the southern land region accounts for around 25% of the simulated global trend in NPP. This compares to ensemble mean trends of $0.03 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.00$), and $0.02 \pm 0.01 \text{ PgCyr}^{-2}$ ($P = 0.34$) over the same period for the S1 (CO₂ only) and S2–S1 (the climate effect), respectively (Tables A2, A3). Southern Africa is the only southern sub-region with a significant trend of $0.041 \pm 0.018 \text{ PgCyr}^{-2}$ ($P = 0.00$) (Table A5). This large positive trend in NPP is due to a positive response of plant production to both CO₂ (NPP S1 = $0.014 \pm 0.007 \text{ PgCyr}^{-2}$, $P = 0.00$), and climate (NPP S2–S1 = $0.027 \pm 0.012 \text{ PgCyr}^{-2}$, $P = 0.01$) (Table A7). An increase in NPP over Southern Africa is likely in response to increases in precipitation over the last two decades (Fig. 5), whereas a decrease in NPP over Temperate South America is likely due to a decrease in precipitation.

DGVMs simulate an ensemble mean RH of $11.20 \pm 3.60 \text{ PgCyr}^{-1}$ over the Southern land region (Table 2). All DGVMs simulate an increase in RH over the period 1990–2009, with a significant trend in the ensemble mean RH of $0.03 \pm 0.02 \text{ PgCyr}^{-2}$ ($P = 0.00$). This is only partly explained by the response of ecosystems to elevated CO₂; over Southern Africa the ensemble mean trend in RH from S1 is $0.02 \pm 0.01 \text{ PgCyr}^{-2}$ ($P = 0.00$), and a climate induced positive trend in RH, $0.01 \pm 0.00 \text{ PgCyr}^{-1}$ ($P = 0.00$) (Tables A2 and A7).

No significant trends in the wildfire flux were reported by any of the DGVMs for the southern land region. However DGVMs agree on simulating a negative trend in wildfire flux across Southern Africa, with an ensemble mean trend of $-0.006 \pm 0.002 \text{ PgCyr}^{-2}$ ($P = 0.03$).

3.2.5 Qualitative change in processes

A qualitative assessment of the differential responses of the underlying land processes to changes in environmental conditions, and their contribution to the sink-source land-atmosphere flux trends are shown in Fig. 8. Many regions are simulated to have a negative land-atmosphere flux trend, with increases in NPP leading increases in RH. However the locations with positive trends over the period 1990 to 2009, i.e. the red colours in Fig. 8, generally fall into two categories: First, regions where models simulate a positive trend in NPP, but an even larger positive trend in RH (eastern Europe, South East USA, Amazonia, South China, North America Tundra). Warming is likely to enhance both NPP and RH in high latitude ecosystems, but primarily RH in low latitudes. Reduced precipitation may partially or fully offset the benefits of elevated atmospheric CO₂ abundance on NPP, and the response of RH to changes in precipitation is not obvious, as this depends on the initial soil moisture status. This is because microbial activity increases with increasing soil moisture at low moisture levels, before reaching a maximum activity, and then begins to decline as water fills the soil pore spaces and oxygen becomes more limiting to respiration. Locations in eastern USA, southern Asia, northern boreal China, south-eastern South America, western and southern Australia are simulated to have negative NPP trends over the last two decades, as a result of reduced rainfall, and there is a less negative trend in RH, possibly due to a reduction in microbial respiration rates with increased soil dryness. The warming and drying in Inner Asia (North East China and Mongolia) and southern Australia is simulated to reduce the rate of microbial decomposition in these regions (Fig. A3), which partly opposes the NPP-driven lagged decrease of RH. The source trend in eastern Europe is simulated as a combination of a negative trend in NPP, as a result of a combination of elevated temperatures and reduced precipitation (i.e. soil drying), and a positive trend in RH, despite reduced plant litter input.

3.2.6 Ocean

3.2.7 Regional fluxes

The large-scale distribution of the modeled mean surface fluxes consists of strong outgassing in the tropical regions, especially in the Pacific, and broad regions of uptake in the mid latitudes, with a few regions in the high latitudes of particularly high uptake, such as the North Atlantic (Fig. 9). This pattern is largely the result of the exchange flux of natural CO₂ that balances globally to a near zero flux, but exhibits regionally strong variations (Gruber et al., 2009). Superimposed on this natural CO₂ flux pattern is the uptake of anthropogenic CO₂, which leads to uptake everywhere, but with substantial regional differences. Large anthropogenic CO₂ uptake fluxes occur in the regions of surface ocean divergence, such as the equatorial Pacific and particularly the Southern Ocean (Sarmiento et al., 1992; Mikaloff Fletcher et al., 2006). This is a result of the divergence causing waters to come from below to the surface which have not been exposed to the atmosphere for while, thereby permitting them to take up a substantial amount of anthropogenic CO₂. This reduces the outgassing that typically characterizes these regions as a result of these upwelling waters bringing with them also high carbon loads from the remineralization of organic matter.

Over the analysis period, the air–sea CO₂ fluxes exhibit only a remarkably small trend in most places with some regions increasing in uptake, while others show a positive flux anomaly, i.e., lesser uptake. Thus the small global trend in ocean uptake over the 1990 through 2004 analysis period is a result of also the individual regions having relatively modest trends.

3.2.8 Process-decomposition

The regional flux trends are, however, much smaller than expected from an ocean with constant circulation that is only responding to increasing atmospheric CO₂ and hence would tend to increase its uptake of anthropogenic CO₂ through time (Fig. 10). In the

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



absence of climate variability and change, all regions would have flux density trends of more than $-0.05 \text{ gCm}^{-2} \text{ yr}^{-2}$, with some regions, such as the Southern Ocean exceeding $-0.15 \text{ gCm}^{-2} \text{ yr}^{-2}$. But climate variability and change compensate these negative trends in every single region by increasing them by $+0.04 \text{ gCm}^{-2} \text{ yr}^{-2}$ or more (with the exception of the South Pacific), such that the overall trends fluctuate from region to region around zero (Fig. 10). The largest reductions in trends are simulated to occur in the North and Equatorial Pacific and in the North Atlantic, where they even cause a change in the sign of the overall trend. A similar, although slightly more moderate pattern is seen if the analysis is undertaken for the entire 1990 through 2009 period on the basis of two models only. The most important difference is found in the North Atlantic, where the climate variability impact is substantially smaller, and not offsetting the anthropogenic CO_2 trend when analyzed for 1990–2009.

The mechanisms driving the oceanic flux trends differ between the analyzed regions. In some regions, surface ocean warming dominates and hence reduces or even cancels increasing ocean anthropogenic CO_2 uptake, as is the case globally (Roy et al., 2011). In other regions, wind changes dominate and yet in other regions, flux trends reflect changes in DIC and Alk, possibly a result of changes in ocean circulation, mixing, and biological productivity.

4 Discussion

4.1 Land

DGVMs simulate an increase in land carbon uptake over the period 1990–2009. The result agrees with earlier findings of Sarmiento et al. (2010), who suggested a large increase in the RLS between the periods 1960–1988 and 1989–2009 (Table A9). The ensemble mean land-atmosphere flux increased by -1.11 PgCyr^{-1} for the same period, compared to the estimated RLS increase of -0.88 PgCyr^{-1} from Sarmiento et al. (2010). The DGVM ensemble trends in land uptake for the globe,

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO_2 over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

northern, tropical and southern land regions of -0.06 ± 0.03 , 0.00 ± 0.01 , -0.04 ± 0.01 , -0.02 ± 0.02 PgCyr⁻², respectively, compare favorably with the inversion estimates of -0.06 ± 0.04 , -0.01 ± 0.01 , -0.04 ± 0.02 , -0.01 ± 0.01 PgCyr⁻², over the period, 1990–2009. Although encouraging, these results should be interpreted with caution because the inversion accounts for any trend in the land use change flux over this period, whereas DGVMs had fixed land use. For this reason we do not compare results at scales finer than the zonal one.

There is empirical evidence for a large increase in biomass in intact forest in the tropical South America and Africa (Pan et al., 2011; Baker et al., 2004; Lewis et al., 2009a, b), which is consistent with the DGVM projections presented here. Subsequently, Lewis et al. (2009b) found broad agreement between biomass trends from observations and from a suite of carbon cycle models applied with 20th century forcing of climate and atmospheric CO₂ content, using a similar protocol to the current analysis. This suggests a large possible role of CO₂ fertilization in stimulating plant productivity in tropical ecosystems. DGVMs suggest a large component of the uptake trend is associated with a positive NPP response to elevated CO₂, which is broadly consistent with the enhancement of forest production due to CO₂ observed in Free Air Carbon Enrichment (FACE) experiments (Norby et al., 2005), albeit they are largely located in temperate forest ecosystems. However, recent studies have highlighted the role of nitrogen in limiting the long term CO₂ response (Canadell et al., 2007; Norby et al., 2010) in these ecosystems. The long-term plant response to elevated CO₂ is likely affected by nutrients and its impact on plant C allocation (Zaehle et al., 2014), however only two out of the nine models used here (CLM4CN and OCN) include interactive nutrient cycling (see DGVM characteristics, Table A1).

In contrast to the large trend in net C uptake across the tropics, DGVMs simulate no statistically significant trend over the northern land. In particular, trends in NPP over Temperate regions are smaller than those in Boreal regions, and are also not significant. Many temperate areas have experienced a decrease in rainfall between 1990 and 2009, and suffered periods of prolonged and severe drought. Examples include

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

the drought in western USA 2000–2004 (McDowell et al., 2008; Anderreg et al., 2012), and the 2003 summer heatwave in Europe (Ciais et al., 2005). These are likely to have had a detrimental effect on the land C uptake (Ciais et al., 2005). This is particularly relevant as climate models agree on a future warming and reduced summer precipitation over continental mid-latitudes (IPCC, 2007).

Modelling ecosystem structure and function in water stressed environments remains a challenge for DGVMs (Morales et al., 2005; Keenan et al., 2009). In general, there is a need for a greater understanding of the mechanisms behind drought-induced plant mortality (Allen et al., 2010; McDowell et al., 2011), and changes in plant water use (De Kauwe et al., 2013). Incorporating hydraulic failure under drought stress improved the ability of the LPJ-GUESS model to simulate the distribution and productivity of xeric and arid vegetation types on a global scale (Hickler et al., 2006); however, the latter is not included in most models. In general, an improvement in the description of vegetation is needed with the adoption of more plant trait information now available (Kattge et al., 2011).

Satellite observations suggest a general greening trend in high latitudes with an earlier onset and longer growing season in high latitude ecosystems, which is reproduced by the DGVMs. Observations suggest a greening Tundra and a slower greening and possible browning in some regions of the boreal forest (Tucker et al., 2001; Bhatt et al., 2010), especially in North America (Beck and Goetz, 2011). In tundra ecosystems, an earlier onset is attributed to warming and earlier snowmelt. In these ecosystems radiation melts snow in the early spring, and the start of growing season corresponds to near peak in radiation. Thus any temperature induced early snowmelt (McDonald et al., 2004; Sitch et al., 2007) is likely to enhance plant production. Warming may not have such a great effect on the offset of the growing season in Arctic tundra ecosystems as this may be driven primarily by radiation. DGVMs simulate a significant positive trend in NPP Boreal North America and Boreal Asia and Tundra. Nitrogen limitation is also likely to constrain the carbon cycle at high latitudes. Only 2 out of 9 DGVMs studied

here included a fully interactive carbon and nitrogen cycle, and it was not possible to quantify N-limitations effects on regional trends in this study.

DGVMs simulate decreasing NPP across North East China and Mongolia, contributing to the overall decreasing land uptake trend, in response to recent climate. In a regional study, Poulter et al. (2013) investigated the differential response of cool semi-arid ecosystems to recent warming and drying trends across Mongolia and Northern China, using multiple sources of evidence, including, LPJ DGVM, FPAR remotely sensed data (derived from GIMMS NDVI3g) and tree-ring widths. They found coherent pattern of high-precipitation sensitivity across data sources, which showed some areas with warming-induced springtime greening and drought-induced summertime browning, and limitations to NPP explained mainly by soil moisture.

Browning is a consequence of regional drought, wildfire and insect outbreak, and their interaction, especially in North America (Beck and Goetz, 2011). Disturbance plays a key role in the ecology of many global ecosystems. For example, wildfire plays a dominant role in the carbon balance of boreal forest in central Canada and other regions (Bond-Lamberty et al., 2007), and insect outbreaks like the mountain pine beetle epidemic 2000–2006; British Columbia, Canada, resulted in the transition of forests from a small carbon sink to a source (Kurz et al., 2008). In general disturbance and forest management are inadequately represented by the current generation of DGVMs, even though several models include simple prognostic wildfire schemes (Table A1), while some are starting to include other disturbance types such as insect attacks (Jönsson et al., 2012) and windthrow (Lagergren et al., 2012). The improvement of DGVMs to include representations of globally and regionally important disturbance types and their response to changing environmental conditions is a priority.

4.2 Ocean

The investigated BOGCMs consistently simulate an ocean characterized by a substantial uptake of CO₂ from the atmosphere, but with a global integrated trend in the two recent decades that is substantially smaller than that expected based on the in-

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



crease in atmospheric CO₂. Results based on the predictions from ocean inversion and ocean Green function methods (Mikaloff Fletcher et al., 2006; Gruber et al., 2009; Khatiwala et al., 2010) suggest an increase in ocean uptake with a trend of the order of -0.04 Pg C yr⁻² over the analysis period (see also Wanninkhof et al., 2013). These latter methods assume constant circulation, while our simulations here include the impact of climate variability and change.

Our analyses reveal indeed that recent climate variability and change has caused the ocean carbon cycle to take up less CO₂ from the atmosphere than expected on the basis of the increase in atmospheric CO₂, i.e., it reduces the efficiency of the ocean carbon sink. Globally, we demonstrated that this efficiency reduction is primarily a result of ocean warming, while regionally, many more processes (e.g., wind changes, alkalinity/DIC concentrations changes) are at play.

Is this reduction in uptake efficiency over the analysis period the first sign of a positive feedback between global warming and the ocean carbon cycle – or alternatively, could it be due to natural decadal-scale variability in air–sea CO₂ fluxes? Without a formal attribution study, it is not possible to provide a firm answer. We suspect that the majority of the trend in the efficiency is due to “natural” decadal-scale variability, however, largely based on the results of McKinley et al. (2011) and Fay and McKinley (2013) who showed that whereas trends in oceanic pCO₂ (and air–sea CO₂ fluxes) are variable on a decadal time-scale, they converge towards atmospheric pCO₂ trends when analyzed over a longer 30 yr period for most global regions. Nevertheless, they also show that in the permanently-stratified subtropical gyre of the North Atlantic, warming (partly driven by anthropogenic climate change) has started to reduce ocean uptake in recent years. In the Southern Ocean, where Le Quéré et al. (2007) and Lovenduski et al. (2008) used models to suggest a reduction in ocean carbon uptake efficiency over the past 25 yr in response to increasing Southern Ocean winds, Fay and McKinley (2013) concluded that the data are insufficient to draw any conclusions.

We should note that the associated uncertainties remain large. Of particular concern is the moderate success of the models to simulate the time-mean ocean sinks and

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

their long-term seasonal cycle (e.g. McKinley et al., 2006). Furthermore, some of the models underestimate the oceanic uptake of transient tracers such as anthropogenic radiocarbon (see e.g., Graven et al., 2012). Such a reduction in the oceanic uptake efficiency is also not suggested by independent measures of oceanic CO₂ uptake, such as the atmospheric O₂/N₂ method (Manning and Keeling, 2006; Ishidoya et al., 2012), albeit the large uncertainties in these estimates make the determination of trends in uptake highly uncertain.

4.3 Reducing uncertainty in regional sinks

In order to better quantify the regional carbon cycle and its trends, DGVM and ocean carbon cycle models need to improve both process representations and model evaluation and benchmarking (Luo et al., 2012). There is a need for up to date global climate and land use and cover change datasets to force the DGVMs, as well as a deeper investigation of the quality and differences between the different reanalysis products used to force ocean carbon cycle models. Also techniques such as detection and attribution can be applied to elucidate trends in the regional carbon cycle and their drivers.

4.3.1 Model benchmarking

There is a critical need for comprehensive model benchmarking, as a first step to attempt to reduce model uncertainty. Several prototype carbon cycle benchmarking schemes have been developed (Randerson et al., 2009; Cadule et al., 2010). A more in depth evaluation and community benchmarking set needs to be agreed and implemented, which also evaluates models for their implicit land response timescales (especially relevant in the discussion on future tipping elements and non-linear future responses) and for the simulated carbon, water and nutrient cycles. New emerging frameworks now exist (Blyth et al., 2011; Abramowitz, 2012; Luo et al., 2012; Dalmonch and Zaehle, 2013).

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



4.3.2 Model resolution

5 Simulated ocean carbon dynamics may be sensitive to horizontal resolution, particularly as model resolution improves sufficiently to adequately capture mesoscale eddies. Mesoscale turbulence influences the ocean carbon cycle in a variety of ways, and the present eddy parameterizations may not adequately capture the full range of effects and the responses to climate variability and change. For example, mesoscale processes are thought to modulate biological productivity by altering the supply of limiting nutrients (Falkowski et al., 1991; McGillicuddy et al., 1998; Gruber et al., 2011). A particularly crucial issue involves the wind-driven overturning circulation in the Southern Ocean, where non-eddy resolving models indicate a strong sensitivity of the overturning circulation and ocean carbon uptake to surface wind stress (Le Quere et al., 2007; Lovenduski et al., 2008). Some eddy-resolving models in contrast suggest that enhanced wind stress is dissipated by increased eddy activity, leading to only a small increase in overturning (Boning et al., 2008), though more recent results indicate a larger response (Gent and Danabasoglu, 2011).

4.3.3 Model structure

20 There is a need for improved representation of ecological processes in land and ocean models, e.g. nutrient cycling (N, P), disturbance (wildfire, wind-throw, insects), land use and land cover change in land models and better representation of the key functional diversity in ocean biogeochemical models. DGVMs need to represent land use and land cover changes, forest management and forest age, to improve estimates of the regional and global land carbon budget. There are recent developments to include nutrient dynamics, mostly nitrogen, into global land biosphere models (as reviewed by Zaehle and Dalmonech, 2011). Too few model simulations are available to date to allow for an ensemble model trend assessment. However, a few general trends appear robust: As evident from Table 2, C–N models generally show less of a response to increasing atmospheric CO₂ due to nitrogen limitation of plant production. N dynam-

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

ics further alter the climate–carbon relationship, which tend to reduce the C loss from temperate and boreal terrestrial ecosystems due to warming; however, with a considerable degree of uncertainty (Thornton et al., 2009; Sokolov et al., 2008; Zaehle et al., 2010). Changes in the nitrogen cycle due to anthropogenic reactive nitrogen additions (both fertiliser to croplands and N deposition on forests and natural grasslands) further modify the terrestrial net C balance and contribute with -0.2 to -0.5 PgCyr⁻¹ to current land sink, with a geographic distribution closely related to that of anthropogenic reactive nitrogen deposition (Zaehle and Dalmonech, 2011). Zaehle et al. (2011), using the OCN model, have estimated the 1995–2005 trend in land uptake due to N deposition as -1.1 ± 1.7 TgCyr⁻², with strong regional differences, depending on the regional trends in air pollution and reactive N loading of the atmosphere and the nitrogen status of the ecosystems, which are generally lower in less responsive in ecosystems close to nitrogen saturation highly polluted regions.

There are several additional land processes that have not been considered in this current multi-model analysis. These include the effects of aerosols and tropospheric ozone on the carbon cycle. Unlike a global forcing agent such as CO₂, the effects of air pollutants (aerosols, NO_x, and O₃) with their shorter atmospheric lifetimes are at the regional scale. Aerosol induced changes in radiation quantity and quality (i.e. the ratio of diffuse to direct) affect plant productivity and the land sink (Mercado et al., 2009). From around 1960 onwards until the 1980s radiation levels declined across industrialized regions, a phenomenon called “global dimming”, followed by a recent brightening in Europe and North America with the adoption of air pollution legislation. Reductions in acid rain have been found to greatly influence trends in riverine DOC, vegetation health, and rates of soil organic matter decomposition. Tropospheric ozone is known to be toxic to plants and lead to reductions in plant productivity, and reduce the efficiency of the land carbon sink (Sitch et al., 2007; Anav et al., 2011). Drivers of the land carbon sink related to air pollution, e.g. N deposition, acid precipitation, diffuse and direct radiation, and surface O₃ have varied markedly in space and time over recent decades.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Similar gaps need to be addressed in ocean biogeochemical models. The ecosystem modules in the current generation of BOGCMs lack the ability to assess many of the suggested mechanisms by which climate and ocean acidification could alter marine biogeochemistry and ocean carbon storage. Proposed biological processes that could influence ocean carbon uptake and release involve, for example, decoupling of carbon and macro-nutrient cycling, changes in micro-nutrient limitation, variations in elemental stoichiometry in organic matter, and changes in the vertical depth-scale for the respiration of sinking organic carbon particles (e.g., Boyd and Doney, 2003; Sarmiento and Gruber, 2006). Some advances have been made with the incorporation of dynamic iron cycling and iron limitation, multiple plankton groups, calcification, and nitrogen fixation (Le Quéré et al., 2005). However, the evaluation of these aspects of the models is currently hindered by both data and process-level information limitations.

4.3.4 Climate and land use and cover datasets

In addition to model structure, the choice of climate forcing and model initial conditions can also contribute to differences between the simulated terrestrial carbon sink. At regional scales, differences in land cover can introduce $\sim 10\%$ uncertainty in simulated regional-scale GPP (Jung et al., 2007; Quaife et al., 2008) and about a 3.5% uncertainty for global NPP. Climate forcing uncertainty tends to have larger effects on carbon flux uncertainty than land cover (Hicke, 2005; Poulter et al., 2011) with up to 25% differences in GPP reported over Europe (Jung et al., 2007) and a 10% difference for global NPP (Poulter et al., 2011). Climate forcing uncertainty and land cover (i.e., PFT distributions) can alter long-term trends in NBP and inter-annual variability of carbon fluxes to climate (Poulter et al., 2011). The DGVMs applied here did not consider LULCC. This is an active area of research; models need a consistent implementation of LULCC. Uncertainties in the simulated net land use flux are associated with assumptions on the implementation of LULCC gridded maps (e.g. whether conversion to cropland in a grid-cell is taken preferentially from grassland, forest, or from both), simulated biomass estimates, and subsequent decomposition rates. However DGVMs

offer the exciting prospect to disentangle the component fluxes associated with land use (e.g. direct emissions, and legacy fluxes), and to separate the environmental and direct human impacts on the net LU flux.

5 Conclusions

5 Land models suggest an increase in the global land net C uptake over the period 1990–2009, mainly driven by trends in NPP, in response to changes in climate and atmospheric CO₂ concentration. Over the same period, ocean models suggest a negligible increase in net ocean C uptake; a result of ocean warming counteracting the expected increase in ocean uptake driven by the increase in atmospheric CO₂. At the
10 sub-regional level, trends vary both in sign and magnitude, particularly over land. Areas in Temperate North America, eastern Europe, North-East China, show a decreasing regional land sink trend, due to regional drying, suggesting a possibility for a transition to a net carbon source in the future if drying continues or drought become more severe and/or frequent. In the ocean, the trends tend to be more homogeneous, but the underlying dynamics differ greatly, ranging from ocean warming, to winds, and to changes in
15 circulation/mixing and ocean productivity, making simple extrapolations into the future difficult.

Our conclusions need to be viewed with several important caveats: Few land models include a prognostic representation of wildfire and no land model represents other disturbances and their interactions, indicating the model response to warming and drought
20 may be conservative in some regions. In addition, only a few models include a fully coupled carbon–nitrogen cycle. Ocean models tend to be too coarse in resolution to properly represent important scales of motions and mixing, such as eddies and other mesoscale processes, and coastal boundary processes. Furthermore, their representation of ocean ecosystem processes and their sensitivity to climate change and other
25 stressors (e.g. ocean acidification, deoxygenation, etc.; Gruber, 2011; Boyd, 2011) is rather simplistic.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



There is a need for detailed model evaluation and benchmarking, to reduce the uncertainty in the sinks in the land and ocean and particularly in how these sinks have changed in the past and how they may change in the future. For land ecosystems, a concerted effort is needed in the DGVM community to incorporate nutrient cycling, and land use and land cover change. For the oceans, models need to improve their representation of unresolved physical transport and mixing processes, and ecosystem models need to evolve to better characterize their response to global change.

Supplementary material related to this article is available online at
[http://www.biogeosciences-discuss.net/10/20113/2013/
bgd-10-20113-2013-supplement.zip](http://www.biogeosciences-discuss.net/10/20113/2013/bgd-10-20113-2013-supplement.zip).

Acknowledgements. N. Gruber and C. Heinze acknowledge financial support by the European Commission through the EU FP7 projects CARBOCHANGE (grant no. 264879) and GEOCARBON (grant no. 283080). N. Gruber was in addition supported through ETH Zurich. SCD acknowledges support from the US National Science Foundation (NSF AGS-1048827). P. Friedlingstein, A. Arneeth and S. Zaehle acknowledge support by the European Commission through the EU FP7 projects EMBRACE (grant no. 282672), and Greencycles II (grant no. 238366). A. Ahlström and B. Smith acknowledge funding through the Mistra Swedish Research Programme for Climate, Impacts and Adaptation (SWECIA). This study is a contribution to the Lund Centre for Studies of Carbon Cycle and Climate Interactions (LUCCI) and the Strategic Research Area Modelling the Regional and Global Earth System (MERGE). This is a contribution to the Bjerknes Centre for Climate Research (BCCR) and core project BIOFEEDBACK of the Centre for Climate Dynamics (SKD) at BCCR. The authors wish to thank Jonathan Barichivich for discussions on greening trends.

References

Abramowitz, G.: Towards a public, standardized, diagnostic benchmarking system for land surface models, *Geosci. Model Dev.*, 5, 819–827, doi:10.5194/gmd-5-819-2012, 2012.

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Allen, M. R. and Stott, P. A.: Estimating signal amplitudes in optimal fingerprinting: Part I. Theory, *Clim. Dynam.*, 21, 477–491, 2003.
- Allen, M. R. and Tett, S. F. B.: Checking for model consistency in optimal fingerprinting, *Clim. Dynam.*, 15, 419–434, 1999.
- 5 Allen, C. D., Macalady, A., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Gonzales, P., Hogg, T., Rigling, A., Breshears, D., Fensham, R., Zhang, Z., Kitzberger, T., Lim, J., Castro, J., Allard, G., Running, S., Semerci, A., and Cobb, N.: Climate-induced forest mortality: a global overview of emerging risks, *Forest Ecol. Manag.*, 259, 660–684, doi:10.1016/j.foreco.2009.09.001, 2010.
- 10 Anav, A., Menut, L., Khvorostyanov, D., and Viovy, N.: Impact of tropospheric ozone on the Euro-Mediterranean vegetation, *Glob. Change Biol.*, 17, 2342–2359, 2011.
- Andres, R. J., Boden, T. A., Bréon, F.-M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A., Marland, G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A synthesis of carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845–1871, doi:10.5194/bg-9-1845-2012, 2012.
- 15 Angert, A., Biraud, S., Bonfils, C., Henning, C. C., Buermann, W., Pinzon, J., Tucker, C. J., and Fung, I.: Drier summers cancel out the CO₂ uptake enhancement induced by warmer springs, *P. Natl. Acad. Sci. USA*, 102, 10823–10827, 2005.
- Assmann, K. M., Bentsen, M., Segsneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model, *Geosci. Model Dev.*, 3, 143–167, doi:10.5194/gmd-3-143-2010, 2010.
- 20 Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A cross-calibrated multiplatform ocean surface wind velocity product for meteorological and oceanographic applications, *B. Am. Meteorol. Soc.*, 92, 157–174, doi:10.1175/2010BAMS2946.1, 2011.
- 25 Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochem. Cy.*, 20, GB2017, doi:10.1029/2005GB002591, 2006.
- Bacastow, R. B.: Modulation of atmospheric carbon dioxide by the Southern Oscillation, *Nature*, 261, 116–118, 1976.
- Baker, T. R., Phillips, O. L., Malhi, Y., Almeida, S., Arroyo, L., Di Fiore, A., Erwin, T., Higuchi, N., Killeen, T. J., Laurance, S. G., Laurance, W. F., Lewis, S. L., Monteagudo, A., Neill, D. A., Nunez Vargas, P., Pitman, N. C. A., Silva, J. N. M., and Vasquez Martinez, R.: Increasing biomass in Amazonian forest plots, *Philos. T. Roy. Soc. B*, 359, 353–365, 2004.
- 30

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon dioxide uptake by land and oceans during the past 50 years, *Nature*, 488, 70–72, doi:10.1038/nature11299, 2012.

Battle, M., Bender, M. L., Tans, P. P., White, J. W. C., Ellis, J. T., Conway, T., and Francey, R. J.: Global carbon sinks and their variability inferred from atmospheric O₂ and δ¹³C, *Science*, 287, 2467–2469, 2000.

Beaulieu, C., Sarmiento, J. L., Mikaloff Fletcher, S. E., Chen, J., and Medvigy, D.: Identification and characterization of abrupt changes in the land uptake of carbon, *Global Biogeochem. Cy.*, 26, GB1007, doi:10.1029/2010GB004024, 2012.

Beck, S. A. and Goetz, S. J.: Satellite observations of high northern latitude vegetation productivity changes between 1982 and 2008: ecological variability and regional differences, *Environ. Res. Lett.*, 6, 045501, doi:10.1088/1748-9326/6/4/045501, 2011.

Bhatt, U. S., Walker, D. A., Reynolds, M. K., Comiso, J. C., Epstein, H. E., Jia, G., Gens, R., and Pinzon, J. E., Tucker, C. J., Tweedie, C. E., and Webber, P. J.: Circumpolar Arctic tundra vegetation change is linked to sea ice decline, *Earth Interact.*, 14, 1–20, 2010.

Buermann, W., Lintner, B. R., Koven, C. D., Angert, A., Pinzon, J. E., Tucker, C. J., and Fung, I. Y.: The changing carbon cycle at Mauna Loa observatory, *P. Natl. Acad. Sci. USA*, 104, 4249–4254, 2007.

Blyth, E., Clark, D. B., Ellis, R., Huntingford, C., Los, S., Pryor, M., Best, M., and Sitch, S.: A comprehensive set of benchmark tests for a land surface model of simultaneous fluxes of water and carbon at both the global and seasonal scale, *Geosci. Model Dev.*, 4, 255–269, doi:10.5194/gmd-4-255-2011, 2011.

Bonan, G. B. and Levis, S.: Quantifying carbon-nitrogen feedbacks in the Community Land Model (CLM4), *Geophys. Res. Lett.*, 37, L07401, doi:10.1029/2010GL042430, 2010.

Böning, C. W., Dispert, A., Visbeck, M., Rintoul, S. R., and Schwarzkopf, F. U.: The response of the Antarctic Circumpolar Current to recent climate change, *Nat. Geosci.*, 1, 864–869, 2008.

Bousquet, P., Peylin, P., Ciais, P., Friedlingstein, P., Lequere, C., and Tans, P.: Interannual CO₂ sources and sinks as deduced by inversion of atmospheric CO₂ data, *Science*, 290, 1342–1346, 2000.

Boyd, P. W.: Beyond ocean acidification, *Nat. Geosci.*, 4, 273–274, 2011.

Boyd, P. and Doney, S. C.: The impact of climate change and feedback process on the ocean carbon cycle, in: *Ocean Biogeochemistry*, edited by: Fasham, M., Springer, 157–193, 2003.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Broecker, W. S., Peng, T.-H., Ostlund, G., and Stuiver, M.: The distribution of bomb radiocarbon in the ocean, *J. Geophys. Res.*, 90, 6953–6970, 1985.
- Buitenhuis, E. T., Rivkin, R. B., Sailley, S., and Le Quéré, C.: Biogeochemical fluxes through microzooplankton, *Global Biogeochem. Cy.*, 24, GB4015, doi:10.1029/2009GB003601, 2010.
- 5 Cadule, P., Friedlingstein, P., Bopp, L., Sitch, S., Jones, C. D., Ciais, P., Piao, S. L., and Peylin, P.: Benchmarking coupled climate-carbon models against long-term atmospheric CO₂ measurements, *Global Biogeochem. Cy.*, 24, GB2016, doi:10.1029/2009GB003556, 2010.
- Canadell, J.: Saturation of the Terrestrial Carbon Sink, 2007.
- Canadell, J. G., Pataki, D., Gifford, R., Houghton, R. A., Lou, Y., Raupach, M. R., Smith, P., and Steffen, W.: Saturation of the terrestrial carbon sink, in: *Terrestrial Ecosystems in a Changing World*, edited by: Canadell, J. G., Pataki, D., and Pitelka, L., The IGBP Series, Springer-Verlag, Berlin Heidelberg, 59–78, 2007.
- 10 Canadell, J. G., Ciais, P., Gurney, K., Le Quéré C., Piao, S., Raupach, M. R., and Sabine, C. L.: An international effort to quantify regional carbon fluxes, *EOS*, 92, 81–82, 2011.
- 15 Canadell, J. G., Ciais, P., Sabine, C., and Joos, F. (Eds.): REgional Carbon Cycle Assessment and Processes (RECCAP), Special Issue, *Biogeosciences*, http://www.biogeosciences.net/special_issue107.html, 2013.
- Chevallier, F., Ciais, P., Conway, T. J., Aalto, T., Anderson, B. E., Bousquet, P., Brunke, E. G., Ciattaglia, L., Esaki, Y., Frölich, M., Gomez, A., Gomez-Pelaez, A. J., Haszpra, L., Krummel, P. B., Langenfelds, R. L., Leuenberger, M., Machida, T., Maignan, F., Matsueda, H., Morguí, J. A., Mukai, H., Nakazawa, T., Peylin, P., Ramonet, M., Rivier, L., Sawa, Y., Schmidt, M., Steele, L. P., Vay, S. A., Vermeulen, A. T., Wofsy, S., and Worthy, D.: CO₂ surface fluxes at grid point scale estimated from a global 21-year reanalysis of atmospheric measurements, *J. Geophys. Res.*, 115, D21307, doi:10.1029/2010JD013887, 2010.
- 20 Ciais, P., Tans, P. P., Trolier, M., White, J. W. C., and Francey, R. J.: A large Northern Hemisphere terrestrial CO₂ sink indicated by the ¹³C/¹²C ratio of atmospheric CO₂, *Science*, 269, 1098–1102, doi:10.1126/science.269.5227.1098, 1995.
- 25 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, doi:10.5194/gmd-4-701-2011, 2011.
- 30

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Conway, T. J., Lang, P. M., and Masarie, K. A.: Atmospheric Carbon Dioxide Dry Air Mole Fractions from the NOAA ESRL Carbon Cycle Cooperative Global Air Sampling Network, 1968–2010, Version: 2011-06-21, available at: <ftp://ftp.cmdl.noaa.gov/ccg/co2/flask/event/>, 2011.

5 Cox, P. M.: Description of the “TRIFFID” dynamic global vegetation model, Hadley Centre, Technical Note 24, 2001.

Dalmonech, D. and Zaehle, S.: Towards a more objective evaluation of modelled land-carbon trends using atmospheric CO₂ and satellite-based vegetation activity observations, *Biogeosciences*, 10, 4189–4210, doi:10.5194/bg-10-4189-2013, 2013.

10 Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P.,
15 Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Q. J. Roy. Meteor. Soc.*, 137, 553–597, 2011.

Defries, R. S., Houghton, R. A., Hansen, M. C., Field, C. B., Skole, D., and Townshend, J.: Carbon emissions from tropical deforestation and regrowth based on satellite observations from the 1980s and 1990s, *P. Natl. Acad. Sci. USA*, 99, 14256–14261, 2002.

20 Denman, K., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., da Silva Dias, P. L., Wofsy, S. C., and Zhang, X.: Couplings Between Changes in the Climate System and Biogeochemistry, in: IPCC, *Climate Change 2007: The Physical Science Basis*, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on
25 Climate Change, edited by: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K., Tignor, M., and Miller, H., Cambridge University Press, Cambridge, United Kingdom and United States, 2007.

Dentener, F., Stevenson, D., Ellingsen, K., van Noije, T., Schultz, M., Amann, M., Atherton, C., Bell, N., Bergmann, D., Bey, I., Bouwman, L., Butler, T., Cofala, J., Collins, B., Drevet, J., Doherty, R., Eickhout, B., Eskes, H., Fiore, A., Gauss, M., Hauglustaine, D., Horowitz, L., Isaksen, I. S. A., Josse, B., Lawrence, M., Krol, M., Lamarque, J. F., Montanaro, V., Muller, J. F.,
30 Peuch, V. H., Pitari, G., Pyle, J., Rast, S., Rodriguez, J., Sanderson, M., Savage, N. H., Shindell, D., Strahan, S., Szopa, S., Sudo, K., Van Dingenen, R., Wild, O., and Zeng, G.:

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

The 5 global atmospheric environment for the next generation, *Environ. Sci. Technol.*, 40, 3586–3594, doi:10.1021/es0523845, 2006.

Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air–sea CO₂ fluxes: physical climate and atmospheric dust, *Deep-Sea Res. Pt. II*, 56, 640–655, 2009a.

Doney, S. C., Lima, I., Moore, J. K., Lindsay, K., Behrenfeld, M. J., Westberry, T. K., Mahowald, N., Glover, D. M., and Takahashi, T.: Skill metrics for confronting global upper ocean ecosystem-biogeochemistry models against field and remote sensing data, *J. Mar. Syst.*, 76, 95–112, 2009b.

Falkowski, P. G., Ziemann, D., Kolber, Z., and Bienfang, P. K.: Role of eddy pumping in enhancing primary production in the ocean, *Nature*, 352, 55–58, 1991.

Fan, S., Gloor, M., Mahlmam, J., Pacala, S., Sarmiento, J., Takahashi, T., and Tans, P.: A large terrestrial carbon sink in North America implied by atmospheric and oceanic carbon dioxide data and models, *Science*, 282, 442–446, 1998.

Fay, A. R. and McKinley, G. A.: Global trends in surface ocean pCO₂ from in situ data, *Global Biogeochem. Cy.*, 27, 541–557, doi:10.1002/gbc.20051, 2013.

Friedlingstein, P., Houghton, R. A., Marland, G., Hackler, J., Boden, T. A., Conway, T. J., Canadell, J. G., Raupach, M. R., Ciais, P., and Le Quere, C.: Update on CO₂ emissions, *Nat. Geosci.*, 3, 811–812, 2011.

Galbraith, D., Levy, P. E., Sitch, S., Huntingford, C., Cox, P., Williams, M., and Meir, P.: Multiple mechanisms of Amazonian Forest biomass losses in three Dynamic Global Vegetation Models under climate change, *New Phytol.*, 187, 647–665, 2010.

Gedney, N., Cox, P. M., Betts, R. A., Boucher, O., Huntingford, C., and Stott, P. A.: Detection of a direct carbon dioxide effect in continental river runoff records, *Nature*, 439, 835–838, 2006.

Gent, P. R. and Danabasoglu, G.: Response to increasing Southern Hemisphere winds in CCSM4, *J. Climate*, 24, 4992–4998, 2011.

Graven, H. D., Gruber, N., Key, R., Khatiwala, S., and Giraud, X.: Changing controls on oceanic radiocarbon: new insights on shallow-to-deep ocean exchange and anthropogenic CO₂ uptake, *J. Geophys. Res.*, 117, C10005, doi:10.1029/2012JC008074, 2012.

Gruber, N.: Warming up, turning sour, losing breath: ocean biogeochemistry under global change, *Phil. Trans. R. Soc. A*, 369, 1980–1996, doi:10.1098/rsta.2011.0003, 2011.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

- Gruber, N., Gloor, M., Mikaloff Fletcher, S. E., Doney, S. C., Dutkiewicz, S., Follows, M. J., Gerber, M., Jacobson, A. R., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., Sarmiento, J. L., and Takahashi, T.: Oceanic sources, sinks, and transport of atmospheric CO₂, *Global Biogeochem. Cy.*, 23, GB1005, doi:10.1029/2008GB003349, 2009.
- 5 Gruber, N., Lachkar, Z., Frenzel, H., Marchesiello, P., Munnich, M., McWilliams, J., Nagai, T., and Plattner, G.-K.: Eddy-induced reduction of biological production in Eastern Boundary Upwelling Systems, *Nat. Geosci.*, 4, 787–792, doi:10.1038/ngeo1273, 2011.
- Guan, D., Liu, Z., Geng, Y., Lindner, S., and Hubacek, K.: The gigatonne gap in China's carbon dioxide inventories, *Nature Climate Change*, 2, 672–675 doi:10.1038/nclimate1560, 2012.
- 10 Houghton, R. A.: How well do we know the flux of CO₂ from land-use change?, *Tellus B*, 62, 337–351, doi:10.1111/j.1600-0889.2010.00473.x, 2010.
- Hicke, J. A.: NCEP and GISS solar radiation data sets available for ecosystem modeling: description, differences, and impacts on net primary production, *Global Biogeochem. Cy.*, 19, GB2006, doi:10.1029/2004GB002391, 2005.
- 15 Hickler, T., Prentice, I. C., Smith, B., Sykes, M. T., and Zaehle, S.: Implementing plant hydraulic architecture within the LPJ Dynamic Global Vegetation Model, *Global Ecol. Biogeogr.*, 15, 567–577, 2006.
- Huntingford, C., Stott, P. A., Allen, M. R., and Lambert, F. H.: Incorporating model uncertainty into attribution of observed temperature change, *Geophys. Res. Lett.*, 33, L05710, doi:10.1029/2005GL024831, 2006.
- 20 Jain, A. K. and Yang, X.: Modeling the effects of two different land cover change data sets on the carbon stocks of plants and soils in concert with CO₂ and climate change, *Global Biogeochem. Cy.*, 19, GB2015, doi:10.1029/2004GB002349, 2005.
- Janssens, I. A., Freibauer, A., Ciais, P., Smith, P., Nabuurs, G.-J., Folberth, G., Schlamadinger, B., Hutjes, R. W. A., Ceulemans, R., Schulze, E.-D., Valentini, R., and Dolman, A. J.: Europe's terrestrial biosphere absorbs 7 to 12 % of European anthropogenic CO₂ emissions, *Science*, 300, 1538–1542, 2003.
- 25 Jeansson, E., Olsen, A., Eldevik, T., Skjelvan, I., Omar, A. M., Lauvset, S. K., Nilsen, J. E. Ø., Bellerby, R. G. J., Johannessen, T., and Falck, E.: The Nordic Seas carbon budget: sources, sinks, and uncertainties, *Global Biogeochem. Cy.*, 25, GB4010, doi:10.1029/2010GB003961, 2011.
- 30

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Jönsson, A. M., Schröder, M., Lagergren, F., Anderbrandt, O. and Smith, B.: Guess the impact of *Ips typographus* – an ecosystem modelling approach for simulating spruce bark beetle outbreaks, *Agr. Forest Meteorol.*, 166–167, 188–200, 2012.

Jung, M., Vetter, M., Herold, M., Churkina, G., Reichstein, M., Zaehle, S., Ciais, P., Viovy, N., Bondeau, A., Chen, Y., Trusilova, K., Feser, F., and Heimann, M.: Uncertainties of modelling gross primary productivity over Europe: a systematic study on the effects of using different drivers and terrestrial biosphere models, *Global Biogeochem. Cy.*, 21, GB4021, doi:10.1029/2006GB002915, 2007.

Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M., Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M., Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K. C., Ropelewski, C., Wang, J., Jenne, R., and Joseph, D.: The NCEP/NCAR 40-year reanalysis project, *B. Am. Meteorol. Soc.*, 77, 437–471, 1996.

Kattge, J., Díaz, S., Lavorel, S., Prentice, I. C., et al.: TRY – a global database of plant traits, *Glob. Change Biol.*, 17, 2905–2935, 2011.

Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F., Sabine, C. L., and Sarmiento, J. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences Discuss.*, 9, 8931–8988, doi:10.5194/bgd-9-8931-2012, 2012.

Keeling, C. D. and Whorf, T. P.: Atmospheric CO₂ records from sites in the SIO sampling network, In *Trends: A compendium of data on global change*, Carbon Dioxide Information Analysis Center, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tenn., USA, 2005.

Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdahl Jr, C. A., Guenther, P. R., Waterman, L. S., and Chin, J. F. S.: Atmospheric carbon dioxide variations at Mauna Loa observatory, Hawaii, *Tellus*, 28, 538–551, 1976.

Keeling, C. D., Whorf, T. P., Wahlen, M., and van der Plicht, J.: Interannual extremes in the rate of rise of atmospheric carbon dioxide since 1980, *Nature*, 375, 666–670, 1995.

Keenan, T., García, R., Friend, A. D., Zaehle, S., Gracia, C., and Sabate, S.: Improved understanding of drought controls on seasonal variation in Mediterranean forest canopy CO₂ and water fluxes through combined in situ measurements and ecosystem modelling, *Biogeosciences*, 6, 1423–1444, doi:10.5194/bg-6-1423-2009, 2009.

Krinner, G., Viovy, N., and de Noblet-Ducoudré, N., Ogée, J., Polcher, J., Friedlingstein, P., Ciais, P., Sitch, S., and Prentice, I. C.: A dynamic global vegetation model for stud-

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



ies of the coupled atmosphere-biosphere system, *Global Biogeochem. Cy.*, 19, GB1015, doi:10.1029/2003GB002199, 2005.

Lagergren, F., Jönsson, A. M., Blennow, K., and Smith, B.: Implementing storm damage in a dynamic vegetation model for regional applications in Sweden, *Ecol. Model.*, 247, 71–82, 2012.

Lambert, F. H., Gillett, N. P., Stone, D. A., and Huntingford, C.: Attribution studies of observed land precipitation changes with nine coupled model, *Geophys. Res. Lett.*, 32, L18704, doi:10.1029/2005GL023654, 2005.

Large, W. G. and Yeager, S.: Diurnal to decadal global forcing for ocean and sea-ice models: the data sets and flux climatologies, NCAR Tech. Rep. TN-460_STR, 105 pp., 2004.

Lawrence, D. M., Oleson, K. W., Flanner, M. G., Thornton, P. E., Swenson, S. C., Lawrence, P. J., Zeng, X., Yang, Z.-L., Levis, S., Sakaguchi, K., Bonan, G. B., and Slater, A. G.: Parameterization improvements and functional and structural advances in version 4 of the Community Land Model, *J. Adv. Model. Earth Syst.*, 3, M03001, doi:10.1029/2011MS000045, 2011.

Le Quéré, C., Harrison, S. P., Colin Prentice, I., Buitenhuis, E. T., Aumont, O., Bopp, L., Claustre, H., Cotrim Da Cunha, L., Geider, R., Giraud, X., Klaas, C., Kohfeld, K. E., Legendre, L., Manizza, M., Platt, T., Rivkin, R. B., Sathyendranath, S., Uitz, J., Watson, A. J., and Wolf-Gladrow, D.: Ecosystem dynamics based on plankton functional types for global ocean biogeochemistry models, *Glob. Change Biol.*, 11, 2016–2040, doi:10.1111/j.1365-2486.2005.1004.x, 2005.

Le Quéré C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R. A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House, J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metz, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2, 831–836, 2009.

Le Quéré, C., Takahashi, T., Buitenhuis, C. E., Rödenbeck, C., and Sutherland, S.: Impact of climate change and variability on the global oceanic sink of CO₂, *Global Biogeochem. Cy.*, 24, GB4007, doi:10.1029/2009GB003599, 2010.

Levy, P. E., Cannell, M. G. R., and Friend, A. D.: Modelling the impact of future changes in climate, CO₂ concentration and land use on natural ecosystems and the terrestrial carbon sink, *Global Environ. Chang.*, 14, 21–30, 2004.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Lewis, S. L., Lopez-Gonzalez, G., Sonké, B., Affum-Baffoe, K., Baker, T. R., Ojo, L. O., Phillips, O. L., Reitsma, J. M., White, L., Comiskey, J. A., Djuikouo K. M.-N., Ewango, C. E. N., Feldpausch, T. R., Hamilton, A. C., Gloor, M., Hart, T., Hladik, A., Lloyd, J., Lovett, J. C., Makana, J.-R., Malhi, Y., Mbago, F. M., Ndangalasi, H. J., Peacock, J., Peh, K. S. H., Sheil, D., Sunderland, T., Swaine, M. D., Taplin, J., Taylor, D., Thomas, S. C., Votere, R., and Wöll, H.: Increasing carbon storage in intact African tropical forests, *Nature*, 457, 1003–1006, 2009a.
- Lewis, S. L., Lloyd, J., Sitch, S., Mitchard, E. T. A., and Laurance, W. F.: Changing ecology of tropical forests: evidence and drivers, *Annu. Rev. Ecol. Evol. Syst.*, 40, 529–49, 2009b.
- Lovenduski, N. S., Gruber, N., Doney, S. C., and Lima, I. D.: Enhanced CO₂ outgassing in the Southern Ocean from a positive phase of the Southern Annular Mode, *Global Biogeochem. Cy.*, 21, GB2026, doi:10.1029/2006GB002900, 2007.
- Lovenduski, N. S., Gruber, N., and Doney, S. C.: Toward a mechanistic understanding of the decadal trends in the Southern Ocean carbon sink, *Global Biogeochem. Cy.*, 22, 1–9, 2008.
- Luo, Y. Q., Randerson, J. T., Abramowitz, G., Bacour, C., Blyth, E., Carvalhais, N., Ciais, P., Dalmonech, D., Fisher, J. B., Fisher, R., Friedlingstein, P., Hibbard, K., Hoffman, F., Huntzinger, D., Jones, C. D., Koven, C., Lawrence, D., Li, D. J., Mahecha, M., Niu, S. L., Norby, R., Piao, S. L., Qi, X., Peylin, P., Prentice, I. C., Riley, W., Reichstein, M., Schwalm, C., Wang, Y. P., Xia, J. Y., Zaehle, S., and Zhou, X. H.: A framework for benchmarking land models, *Biogeosciences*, 9, 3857–3874, doi:10.5194/bg-9-3857-2012, 2012.
- Matear, R. J. and Lenton, A.: Impact of historical climate change on the Southern Ocean carbon cycle, *J. Climate*, 21, 5820–5834, doi:10.1175/2008JCLI2194.1, 2008.
- Matsumoto, K. and Gruber, N.: How accurate is the estimation of anthropogenic carbon in the ocean? An evaluation of the DC* method, *Global Biogeochem. Cy.*, 19, GB3014, doi:10.1029/2004GB002397, 2005.
- McDowell, N. G.: Mechanisms linking drought, hydraulics, carbon metabolism, and vegetation mortality, *Plant Physiol.*, 155, 1051–1059, 2011.
- McDowell, N. G., Pockman, W. T., Allen, C. D., Breshears, D. D., Cobb, N., Kolb, T., Plaut, J., Sperry, J., West, A., Williams, D. G., and Yepez, E. A.: Mechanisms of plant survival and mortality during drought: why do some plants survive while others succumb to drought?, *New Phytol.*, 178, 719–739, 2008.
- McGillicuddy, D. J., Robinson, Jr., A. R., Siegel, D. A., Jannasch, H. W., Johnson, R., Dickey, T. D., McNeil, J., Michaels, A. F., and Knap, A. H.: Influence of mesoscale eddies on new production in the Sargasso Sea, *Nature*, 394, 263–266, 1998.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- McNeil, B. I. and Matear, R. J.: The non-steady state oceanic CO₂ signal: its importance, magnitude and a novel way to detect it, *Biogeosciences*, 10, 2219–2228, doi:10.5194/bg-10-2219-2013, 2013.
- McKinley, G. A., Takahashi, T., Buitenhuis, E., Chai, F., Christian, J. R., Doney, S. C., Jiang, M.-S., Lindsay, K., Moore, J. K., Le Quéré, C., Lima, I., Murtugudde, R., Shi, L., and Wetzel, P.: North Pacific carbon cycle response to climate variability on seasonal to decadal timescales, *J. Geophys. Res.*, 111, 1–22, 2006.
- McKinley, G. A., Fay, A. R., Takahashi, T., and Metzl, N.: Convergence of atmospheric and North Atlantic carbon dioxide trends on multidecadal timescales, *Nat. Geosci.*, 4, 606–610, 2011.
- Mercado, L. M., Bellouin, N., Sitch, S., Boucher, O., Huntingford, C., and Cox, P. M.: Impact of changes in diffuse radiation on the global land carbon sink, *Nature*, 458, 1014–1017, doi:10.1038/nature07949, 2009.
- Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento, J. L.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the ocean, *Global Biogeochem. Cy.*, 20, GB2002, doi:10.1029/2005GB002530, 2006.
- Myneni, R. B., Keeling, C. D., Tucker, C. J., Asrar, G., and Nemani, R. R.: Increased plant growth in the northern high latitudes from 1981 to 1991, *Nature*, 386, 698–702, doi:10.1038/386698a0, 1997.
- Nemani, R. R., Keeling, C. D., Hashimoto, H., Jolly, W. M., Piper, S. C., Tucker, C. J., Myneni, R. B., and Running, S. W.: Climate-driven increases in global terrestrial net primary production from 1982 to 1999, *Science*, 300, 1560–1563, 2003.
- New, M. G., Hulme, M., and Jones, P. D.: Representing twentieth-century space-climate variability, Part II, Development of 1901–1996 monthly grids of terrestrial surface climate, *J. Climate*, 13, 2217–2238, 2000.
- Norby, R. J., DeLucia, E. H., Gielen, B., Calfapietra, C., Giardina, C. P., King, J. S., Ledford, J., McCarthy, H. R., Moore, D. J. P., Ceulemans, R., De Angelis, P., Finzi, A. C., Karnosky, D. F., Kubiske, M. E., Lukac, M., Pregitzer, K. S., Scarascia-Mugnozza, G. E., Schlesinger, W. H., and Oren, R.: Forest response to elevated CO₂ is conserved across a broad range of productivity, *P. Natl. Acad. Sci. USA*, 102, 18052–18056, 2005.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Norby, R. J., Warren, J. M., Iversen, C. M., Medlyn, B. E., and McMurtrie, R. E.: CO₂ enhancement of forest productivity constrained by limited nitrogen availability, *P. Natl. Acad. Sci. USA*, 107, 19368–19373, 2010.
- Oleson, K. W., Lawrence, D. M., Bonan, G. B., Flanner, M. G., Kluzek, E., Lawrence, P. J., Levis, S., Swenson, S. C., Thornton, P. E., Dai, A., Decker, M., Dickinson, R., Feddes, J., Heald, C. L., Hoffman, F., Lamarque, J.-F., Mahowald, N., Niu, G.-Y., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stockli, R., Wang, A., Yang, Z.-L., Zeng, X., and Zeng, X.: Technical description of version 4.0 of the Community Land Model (CLM), NCAR Technical Note NCAR/TN-478+STR, 257 pp., 2010.
- Olivier, J. G. J., Janssens-Maenhout, G., and Peters, J. A. H. W.: Trends in global CO₂ emissions, 2012 report, Background Studies, PBL Netherlands Environmental Assessment Agency, 40 pp., 2012.
- Pacala, S. W., Hurtt, G. C., Baker, D., Peylin, P., Houghton, R. A., Birdsey, R. A., Heath, L., Sundquist, E. T., Stallard, R. F., Ciais, P., Moorcroft, P., Caspersen, J. P., Shevliakova, E., Moore, B., Kohlmaier, G., Holland, E., Gloor, M., Harmon, M. E., Fan, S.-M., Sarmiento, J. L., Goodale, C. L., Schimel, D., and Field, C. B.: Consistent land- and atmosphere-based US carbon sink estimates, *Science*, 292, 2316–2320, 2001.
- Pan, Y., Birdsey, R. A., Fang, J., Houghton, R., Kauppi, P. E., Kurz, W. A., Phillips, O. L., Shvidenko, A., Lewis, S. L., Canadell, J. G., Ciais, P., Jackson, R. B., Pacala, S., McGuire, A. D., Piao, S., Rautiainen, A., Sitch, S., Hayes, D., and Watson, C.: A large and persistent carbon sink in the World's forests, 1990–2007, *Science*, 333, 6045, doi:10.1126/science.1201609, 2011.
- Phillips, O. L., Malhi, Y., Higuchi, N., Laurance, W. F., Núñez, P. V., Vásquez, R. M., Laurance, S. G., Ferreira, L. V., Stern, M., Brown, S., and Grace, J.: Changes in the carbon balance of tropical forests: evidence from long-term plots, *Science*, 282, 439–442, 1998.
- Piao, S. L., Friedlingstein, P., Ciais, P., Zhou, L. M., and Chen, A. P.: Effect of climate and CO₂ changes on the greening of the Northern Hemisphere over the past two decades, *Geophys. Res. Lett.*, 33, L23402, doi:10.1029/2006GL028205, 2006.
- Piao, S., Friedlingstein, P., Ciais, P., Viovy, N., and Demarty, J.: Growing season extension and its impact on terrestrial carbon cycle in the Northern Hemisphere over the past 2 decades, *Global Biogeochem. Cy.*, 21, GB3018, doi:10.1029/2006GB002888, 2007.
- Piao, S. L., Fang, J. Y., Ciais, P., Peylin, P., Huang, Y., Sitch, S., and Wang, T.: The carbon balance of terrestrial ecosystems in China, *Nature*, 458, 1009–1014, 2009a.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Piao, S. L., Ciais, P., de Noblet-Ducoudré, P., Cadule, P., Viovy, N., and Wang, T.: Spatiotemporal patterns of terrestrial carbon cycle during the 20th century, *Global Biogeochem. Cy.*, 23, GB4026, doi:10.1029/2008GB003339, 2009b.
- Piao, S., Sitch, S., Ciais, P., Friedlingstein, P., Peylin, P., Wang, X., Ahlström, A., Anav, A., Canadell, J. G., Cong, N., Huntingford, C., Jung, M., Levis, S., Levy, P. E., Li, J., Lin, X., Lomas, M. R., Lu, M., Luo, Y., Ma, Y., Myneni, R. B., Poulter, B., Sun, Z., Wang, T., Viovy, N., Zaehle, S., and Zeng, N.: Evaluation of terrestrial carbon cycle models for their response to climate variability and to CO₂ trends, *Glob. Change Biol.*, 19, 2117–2132, doi:10.1111/gcb.12187, 2013.
- Poulter, B., Frank, D. C., Hodson, E. L., and Zimmermann, N. E.: Impacts of land cover and climate data selection on understanding terrestrial carbon dynamics and the CO₂ airborne fraction, *Biogeosciences*, 8, 2027–2036, doi:10.5194/bg-8-2027-2011, 2011.
- Poulter, B., Pederson, N., Liu, H., Zhu, Z., D'Arrigo, R., Ciais, P., Davi, N., Frank, D., Leland, C., Myneni, R., Piao, S., and Wang, T.: Recent trends in Inner Asian forest dynamics to temperature and precipitation indicate high sensitivity to climate change, *Agr. Forest Meteorol.*, 178–179, 31–45, 2013.
- Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Kheshgi, H. S., Le Quere, C., Scholes, R. J., Wallace, D. W. R., Archer, D., Ashmore, M. R., Aumont, O., Baker, D., Battle, M., Bender, M., Bopp, L. P., Bousquet, P., Caldeira, K., Ciais, P., Cox, P. M., Cramer, W., Dentener, F., Enting, I. G., Field, C. B., Friedlingstein, P., Holland, E. A., Houghton, R. A., House, J. I., Ishida, A., Jain, A. K., Janssens, I. A., Joos, F., Kaminski, T., Keeling, C. D., Keeling, R. F., Kicklighter, D. W., Kohfeld, K. E., Knorr, W., Law, R., Lenton, T., Lindsay, K., Maier-Reimer, E., Manning, A. C., Matear, R. J., McGuire, A. D., Melillo, J. M., Meyer, R., Mund, M., Orr, J. C., Piper, S., Plattner, K., Rayner, P. J., Sitch, S., Slater, R., Taguchi, S., Tans, P. P., Tian, H. Q., Weirig, M. F., Whorf, T., and Yool, A.: The carbon cycle and atmospheric carbon dioxide, in: *Climate Change, 2001: The Scientific Basis*, edited by: Houghton, J. T., Cambridge Univ. Press, New York, 2001.
- Prentice, I. C., Bondeau, A., Cramer, W., Harrison, S. P., Hickler, T., Lucht, W., Sitch, S., Smith, B., and Sykes, M. T.: *Terrestrial Ecosystems in a Changing world*, IGBP Book Series, edited by: Canadell, J., Pitelka, L., and Pataki, D., Springer, Heidelberg, Germany, 336 pp., 2007.
- Quaife, T., Quegan, S., Disney, M., Lewis, P., Lomas, M., and Woodward, F. I.: Impact of land cover uncertainties on estimates of biospheric carbon fluxes, *Global Biogeochem. Cy.*, 22, GB4016, doi:10.1029/2007GB003097, 2008.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

- Reichstein, M., Bahn, M., Ciais, P., Frank, D., Mahecha, M. D., Seneviratne, S. I., Zscheischler, J., Beer, C., Buchmann, N., Frank, D. C., Papale, D., Rammig, A., Smith, P., Thonicke, K., van der Velde, M., Vicca, S., Walz, A., and Wattenbach, M.: Climate extremes and the carbon cycle, *Nature*, 500, 287–295, 2013.
- 5 Roy, T., Bopp, L., Gehlen, M., Schneider, B., Cadule, P., Frölicher, T. L., Segschneider, J., Tjiputra, J., Heinze, C., and Joos, F.: Regional impacts of climate change and atmospheric CO₂ on future ocean carbon uptake: a multimodel linear feedback analysis, *J. Climate*, 24, 2300–2318, 2011.
- Sabine, C. S., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R.,
10 Wong, C. S., Wallace, D. W. R., Tillbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for anthropogenic CO₂, *Science*, 305, 367–371, 2004.
- Sarmiento, J. L. and Gruber, N.: Sinks for anthropogenic carbon, *Physics Today*, 55, 30–36, 2002.
- Sarmiento, J. L. and Gruber, N.: *Ocean Biogeochemical Dynamics*, Princeton Univ. Press, 526
15 pp., 2006.
- Sarmiento, J. L., Gloor, M., Gruber, N., Beaulieu, C., Jacobson, A. R., Mikaloff Fletcher, S. E., Pacala, S., and Rodgers, K.: Trends and regional distributions of land and ocean carbon sinks, *Biogeosciences*, 7, 2351–2367, doi:10.5194/bg-7-2351-2010, 2010.
- Schuster, U., McKinley, G. A., Bates, N., Chevallier, F., Doney, S. C., Fay, A. R., González-Dávila, M., Gruber, N., Jones, S., Krijnen, J., Landschützer, P., Lefèvre, N., Manizza, M.,
20 Mathis, J., Metzl, N., Olsen, A., Rios, A. F., Rödenbeck, C., Santana-Casiano, J. M., Takahashi, T., Wanninkhof, R., and Watson, A. J.: An assessment of the Atlantic and Arctic sea-
air CO₂ fluxes, 1990–2009, *Biogeosciences*, 10, 607–627, doi:10.5194/bg-10-607-2013, 2013.
- Sitch, S., Smith, B., Prentice, I. C., Arneeth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S.,
25 Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic vegetation model, *Glob. Change Biol.*, 9, 161–185, 2003.
- Sitch, S., Cox, P. M., Collins, W. J., and Huntingford, C.: Indirect radiative forcing of climate change through ozone effects on the land-carbon sink, *Nature*, 448, 791–795, doi:10.1038/nature06059, 2007.
- 30 Sitch, S., Huntingford, C., Gedney, N., Levy, P. E., Lomas, M., Piao, S., Betts, R., Ciais, P., Cox, P., Friedlingstein, P., Jones, C. D., Prentice, I. C., and Woodward, F. I.: Evaluation of the terrestrial carbon cycle, future plant geography, and climate-carbon cycle feed-

backs using 5 Dynamic Global Vegetation Models (DGVMs), *Glob. Change Biol.*, 14, 1–25, doi:10.1111/j.1365-2486.2008.01626.x, 2008.

Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches within European climate space, *Global Ecol. Biogeogr.*, 10, 621–637, 2001.

Stephens, B. B., Gurney, K. R., Tans, P. P., Sweeney, C., Peters, W., Bruhwiler, L., Ciais, P., Ramonet, M., Bousquet, P., Nakazawa, T., Aoki, S., Machida, T., Inoue, G., Vinnichenko, N., Lloyd, J., Jordan, A., Heimann, M., Shibistova, Langenfelds, R. L., Steele, L. P., Francey, R. J., and Denning, A. S.: Weak northern and strong tropical land carbon uptake from vertical profiles of atmospheric CO₂, *Science*, 316, 1732–1735, 2007.

Sweeney, C., Gloor, E., Jacobson, A. R., Key, R. M., McKinley, G., Sarmiento, J. L., and Wankminkhof, R.: Constraining global air–sea gas exchange for CO₂ with recent bomb ¹⁴C measurements, *Global Biogeochem. Cy.*, 21, GB2015, doi:10.1029/2006GB002784, 2007.

Tans, P. P., Fung, I. Y., and Takahashi, T.: Observational constraints on the global atmospheric CO₂ budget, *Science*, 247, 1431–1439, doi:10.1126/science.247.4949.1431, 1990.

Thornton, P. E. and Rosenbloom, N. A.: Ecosystem model spin-up: estimating steady state conditions in a coupled terrestrial carbon and nitrogen cycle model, *Ecol. Model.*, 189, 25–48, 2005.

Thornton, P. E., Lamarque, J. F., Rosenbloom, N. A., and Mahowald, N. M.: Influence of carbon-nitrogen cycle coupling on land model response to CO₂ fertilization and climate variability, *Global Biogeochem. Cy.*, 21, GB4018, doi:10.1029/2006GB002868, 2007.

Thornton, P. E., Doney, S. C., Lindsay, K., Moore, J. K., Mahowald, N., Randerson, J. T., Fung, I., Lamarque, J.-F., Feddes, J. J., and Lee, Y.-H.: Carbon-nitrogen interactions regulate climate-carbon cycle feedbacks: results from an atmosphere-ocean general circulation model, *Biogeosciences*, 6, 2099–2120, doi:10.5194/bg-6-2099-2009, 2009.

Tjiputra, J. F., Assmann, K., and Heinze, C.: Anthropogenic carbon dynamics in the changing ocean, *Ocean Sci.*, 6, 605–614, doi:10.5194/os-6-605-2010, 2010.

Tucker, C. J., Slayback, D. A., Pinzon, J. E., Los, S. O., Myneni, R. B. and Taylor, M. G.: Higher northern latitude normalized difference vegetation index and growing season trends from 1982 to 1999, *Int. J. Biometeorol.*, 45, 184–190, 2001.

Van der Molen, M. K., Dolman, A. J., Ciais, P., Eglin, T., Gobron, N., Law, B. E., Meir, P., Peters, W., Phillips, O. L., Reichstein, M., Chen, T., Dekker, S. C., Doubková, M., Friedl, M. A., Jung, M., van den Hurk, B. J. J. M., de Jeu, R. A. M., Kruijt, B., Ohta, T., Rebel, K. T.,

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Plummer, S., Seneviratne, S. I., Sitch, S., Teuling, A. J., van der Werf, G. R., and Wang, G.: Drought and ecosystem carbon cycling, *Agr. Forest Meteorol.*, 151, 765–773, doi:10.1016/j.agrformet.2011.01.018, 2011.

Vitousek, P. M. and Howarth, R. W.: Nitrogen limitation on land and in the sea: how can it occur?, *Biogeochemistry*, 13, 87–115, 1991.

Wanninkhof, R.: Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, 97, 7373–7382, 1992.

Wanninkhof, R., Park, G.-H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C., McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983–2000, doi:10.5194/bg-10-1983-2013, 2013.

Woodward, F. I. and Lomas, M. R.: Vegetation-dynamics – simulating responses to climate change, *Biolog. Rev.*, 79, 643–670, 2004.

Woodward, F. I., Smith, T. M., and Emanuel, W. R.: A global land primary productivity and phytogeography model, *Global Biogeochem. Cy.*, 9, 471–490, 1995.

Zaehle, S. and Dalmonch, D.: Carbon–nitrogen interactions on land at global scales: current understanding in modeling climate biosphere feedbacks, *Curr. Op. Environ. Sustain.*, 3, 311–320, 2011.

Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model, I: Model description, site-scale evaluation and sensitivity to parameter estimates, *Global Biogeochem. Cy.*, 24, GB1005, doi:10.1029/2009GB003521, 2010.

Zaehle, S., Medlyn, B. E., De Kauwe, M. G., Walker, A. P., Dietze, M. C., Hickler, T., Luo, Y., Wang, Y.-P., El-Masri, B., Thornton, P., Jain, A., Wang, S., Warlind, D., Weng, E., Parton, W., Iversen, C. M., Gallet-Budynek, A., McCarthy, H., Finzi, A., Hanson, P. J., Prentice, I. C., Oren, R., and Norby, R. J.: Evaluation of eleven terrestrial carbon-nitrogen cycle models against observations from two temperate Free-Air CO₂ Enrichment Studies, *New Phytol.*, submitted, 2013.

Zhang, X. B., Zwiers, F. W., Hegerl, G. C., Lambert, F. H., Gillett, N. P., Solomon, S., Stott, P. A., and Nozawa, T.: Detection of human influence on twentieth-century precipitation trends, *Nature*, 448, 461–465, doi:10.1038/nature06025, 2007.

Zeng, N.: Glacial-interglacial atmospheric CO₂ change – the glacial burial hypothesis, *Adv. Atmos. Sci.*, 20, 677–673, 2003.

BGD

10, 20113–20177, 2013

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Table 1. Characteristics of the 9 Dynamic Global Vegetation Models.

Model Name	Abbreviation	Spatial resolution	Land Surface Model	Full Nitrogen Cycle	River Export Flux	Fire simulation	Harvest/ grazing flux	Source
Community Land Model 4CN	CLM4CN	0.5° × 0.5°	Yes	Yes	No	Yes	No	Oleson et al. (2010); Lawrence et al. (2011)
Hyland	HYL	3.75° × 2.5°	No	No	No	No	Yes	Friend et al. (1997); Levy et al. (2004)
Lund-Potsdam-Jena	LPJ	0.5° × 0.5°	No	No	No	Yes	Yes	Sitch et al. (2003)
LPJ-GUESS	LPJ-GUESS	0.5° × 0.5°	No	No	No	Yes	No	Smith et al. (2001)
ORCHIDEE-CN	OCN	3.75° × 2.5°	Yes	Yes	No	No	Yes	Zaehle and Friend (2010); Zaehle et al. (2010)
ORCHIDEE	ORC	0.5° × 0.5°	Yes	No	No	No	No	Krinner et al. (2005)
Sheffield-DGVM	SDGVM	3.75° × 2.5°	No	No	Yes	Yes	No	Woodward et al. (1995)
TRIFFID	TRI	3.75° × 2.5°	Yes	No	No	No	No	Cox (2001)
VEGAS	VEGAS	0.5° × 0.5°	Yes	No	Yes	Yes	Yes	Zeng et al. (2005)

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Table 2. Mean and trends in NPP, RH, land–atmosphere flux as simulated by individual DGVMs and the Ensemble mean.

MODEL	NPP (PgCyr ⁻¹)	Trend (PgCyr ⁻²)	P value	RH (PgCyr ⁻¹)	Trend (PgCyr ⁻²)	P value	Land sink (PgCyr ⁻¹)	Trend (PgCyr ⁻²)	P value
Global_Land									
CLM4CN	51.508	0.148	0.000	47.668	0.106	0.000	-1.459	-0.052	0.059
HYLAND	73.422	0.319	0.000	68.835	0.203	0.000	-3.466	-0.109	0.000
LPJ	59.306	0.216	0.000	47.612	0.117	0.000	-2.251	-0.068	0.061
LPJ-GUESS	62.506	0.174	0.000	55.448	0.145	0.000	-1.802	-0.043	0.346
OCN	53.941	0.155	0.000	50.611	0.135	0.000	-2.272	-0.015	0.568
ORCHIDEE	75.516	0.293	0.000	72.037	0.208	0.000	-3.479	-0.086	0.046
SDGVM	60.965	0.240	0.000	53.778	0.190	0.000	-2.127	-0.044	0.170
TRIFFID	71.929	0.305	0.000	69.167	0.244	0.000	-2.762	-0.061	0.265
VEGAS	57.308	0.113	0.006	51.930	0.092	0.000	-1.783	-0.018	0.551
ENSEMBLE	62.934	0.218	0.000	57.454	0.160	0.000	-2.378	-0.055	0.048
Std	8.729	0.076		9.791	0.053		0.721	0.030	
Northern_Land									
CLM4CN	17.523	0.043	0.003	16.215	0.036	0.000	-0.670	-0.007	0.612
HYLAND	19.139	0.098	0.000	17.591	0.080	0.000	-0.876	-0.014	0.311
LPJ	24.566	0.079	0.001	19.578	0.062	0.006	-1.168	-0.006	0.735
LPJ-GUESS	28.484	0.039	0.085	25.883	0.067	0.009	-0.634	0.023	0.521
OCN	21.008	0.044	0.035	19.264	0.047	0.008	-1.117	0.007	0.632
ORCHIDEE	30.337	0.070	0.007	29.112	0.063	0.000	-1.226	-0.006	0.740
SDGVM	25.144	0.063	0.006	22.598	0.065	0.006	-0.828	0.004	0.762
TRIFFID	28.476	0.088	0.009	27.006	0.103	0.001	-1.470	0.016	0.455
VEGAS	21.895	0.048	0.012	18.914	0.043	0.001	-1.322	-0.000	0.968
ENSEMBLE	24.064	0.063	0.001	21.796	0.063	0.001	-1.034	0.002	0.865
Std	4.484	0.022		4.562	0.020		0.295	0.012	
Tropical_Land									
CLM4CN	26.400	0.090	0.000	24.464	0.058	0.000	-0.692	-0.039	0.110
HYLAND	34.489	0.112	0.000	32.695	0.067	0.000	-1.560	-0.044	0.001
LPJ	25.830	0.100	0.001	21.224	0.035	0.001	-0.817	-0.049	0.031
LPJ-GUESS	21.922	0.078	0.000	19.332	0.051	0.000	-0.785	-0.036	0.038
OCN	22.750	0.084	0.000	21.476	0.065	0.000	-0.982	-0.017	0.210
ORCHIDEE	31.313	0.151	0.000	29.640	0.108	0.000	-1.673	-0.043	0.084
SDGVM	23.505	0.118	0.000	20.677	0.075	0.000	-0.984	-0.038	0.030
TRIFFID	29.801	0.141	0.000	28.925	0.096	0.000	-0.876	-0.045	0.218
VEGAS	23.472	0.041	0.061	21.994	0.033	0.004	-0.278	-0.010	0.527
ENSEMBLE	26.609	0.102	0.000	24.492	0.065	0.000	-0.961	-0.036	0.045
Std	4.350	0.034		4.752	0.025		0.428	0.013	
Southern_Land									
CLM4CN	7.617	0.014	0.187	7.017	0.011	0.036	-0.098	-0.005	0.719
HYLAND	19.875	0.109	0.000	18.623	0.056	0.000	-1.035	-0.051	0.000
LPJ	8.940	0.037	0.074	6.833	0.021	0.004	-0.267	-0.013	0.355
LPJ-GUESS	12.124	0.058	0.003	10.255	0.026	0.001	-0.385	-0.031	0.192
OCN	10.222	0.027	0.165	9.909	0.023	0.053	-0.174	-0.004	0.744
ORCHIDEE	13.884	0.073	0.002	13.304	0.037	0.000	-0.581	-0.036	0.027
SDGVM	12.358	0.059	0.034	10.539	0.050	0.000	-0.317	-0.010	0.701
TRIFFID	13.707	0.077	0.020	13.290	0.045	0.000	-0.417	-0.032	0.269
VEGAS	11.971	0.024	0.382	11.049	0.016	0.140	-0.182	-0.009	0.656
ENSEMBLE	12.300	0.053	0.011	11.202	0.032	0.000	-0.384	-0.021	0.196
Std	3.528	0.031		3.597	0.016		0.285	0.017	

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

Title Page

Abstract Introduction

Conclusions References

Tables Figures

⏪ ⏩

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 1. Land and ocean regions. The three land regions: Northern land, Tropical land and Southern land. Northern land comprises: Boreal North America (navy blue), Europe (light blue), Boreal Asia (blue), Temperate North America (pale red), and Temperate Asia (red). Tropical land comprises: Tropical South America Forests (sea green), Northern Africa (sand), Equatorial Africa (green), and Tropical Asia (dark green). Southern land comprises: South America Savanna (pale green), Temperate South America (violet), Southern Africa (orange), Australia and New Zealand (yellow). Ocean regions comprise: North Pacific (dark red), Equatorial Pacific (orange-red), South Pacific (orange), North Atlantic (orchid), Equatorial/South Atlantic (slate blue), Indian Ocean (thistle), and Southern Ocean (sky blue), Arctic Ocean and Antarctic (white).

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#) [Introduction](#)

[Conclusions](#) [References](#)

[Tables](#) [Figures](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

[Back](#) [Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[◀](#)

[▶](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)




Fig. 2. Global trends in environmental driving variables: Panel 1: temperature; Panel 2: precipitation, Panel 3: atmospheric [CO₂], Panel 4: N deposition.

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 3. Global trends in ensemble land model responses. Panel 1: DGVM mean model land sink and standard deviation (grey lines), Component fluxes, NPP (Panel 2), and RH (= RH + wildfire + Riverine C flux) (Panel 3). Remotely sensed trends in annual mean NDVI (crosses), a measure of vegetation greenness, and a linear regression through the data points (bold line) (Panel 4)

Title Page

Abstract Introduction

Conclusions References

Tables Figures

◀ ▶

◀ ▶

Back Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 4. Global trends in ensemble ocean model fluxes. Black line: results from simulation S_O2 with variable “climate” and increasing CO₂. Red line: results from Simulation S_O1 with constant “climate” and increasing CO₂. The dashed lines indicate the \pm uncertainty bands given by the 4 models that contribute to the ensemble mean.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 5. Gridded maps of average land sink over the period 1990–2009 for the ensemble mean (top left); model agreement with stippling representing agreement for 66 % of DGVMs (top middle panel); standard deviation across DGVMs. The bottom left panel shows the trend in land sink across the ensemble, and model agreement; stippling representing agreement of at least 66 % of the DGVMs (bottom middle), and the standard deviation of the trend.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 6. Multi-panel gridded maps of trends in land climate drivers and process responses. Panel 1: trend in temperature; Panel 2: precipitation; Panel 3: land sink; Panel 4: NPP; Panel 5: Rh (= RH + wildfire + Riverine C flux).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 7. Ensemble-mean trends in the onset (top), offset (middle) and length of growing season in days for the ensemble mean (left) compared with satellite derived estimates (right).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 8. Qualitative change in processes over the period, 1990–2009. Negative trend in land–atmosphere CO₂ flux: enhanced NPP > enhanced RH (= RH + wildfire + Riverine C flux) (pale blue); enhanced NPP, reduced RH (turquoise); reduced NPP < reduced RH (dark blue). Positive trend in land–atmosphere CO₂ flux: enhanced NPP < enhanced RH (dark red); reduced NPP, enhanced RH (red); reduced NPP > reduced RH (pink).

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 9. Gridded maps of the ensemble mean air–sea CO₂ flux over the period 1990–2004 (top left); standard deviation of the mean flux across the four BOGCMs. The bottom left panel shows the trend in the net flux across the ensemble, while the bottom right panel shows the standard deviation of the trend.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

Trends and drivers of regional sources and sinks of CO₂ over the past two decades

S. Sitch et al.




Fig. 10. Regional ocean flux trends from 1990 through 2004 for the standard case, i.e., variable climate and increasing CO₂ (simulation S_O2), and for the constant climate case (simulation S_O1).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)

[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)
