

# Signature of a universal statistical description for drift-wave plasma turbulence

Johan Anderson and Pavlos Xanthopoulos  
Max-Planck-Institut für Plasmaphysik, IPP-Euratom Association,  
Teilinstitut Greifswald, D-17491 Greifswald, Germany

This Letter provides a theoretical interpretation of numerically generated probability density functions (PDFs) of intermittent plasma transport events. Specifically, nonlinear gyrokinetic simulations of ion-temperature-gradient turbulence produce time series of heat flux which exhibit manifestly non-Gaussian PDFs with enhanced tails. It is demonstrated that, after the removal of autocorrelations, the numerical PDFs can be matched with predictions from a fluid theoretical setup, based on the instanton method. This result points to a universality in the modeling of intermittent stochastic process, offering predictive capability.

Stochastic physical processes are most often observed to be unimodal with exponential tails [1], a feature which is also attributed to fluctuations in magnetically confined plasmas [2]- [7]. These fluctuations are intermittent events manifesting a patchy spatial and bursty temporal structure, pertaining to radially propagating coherent structures like blobs or avaloids [8], and have been suggested to carry a significant fraction of the total transport [9]. Therefore, a comprehensive predictive theory is called for, in order to understand and, subsequently, improve properties related to intermittency. A major goal would be, for instance, the control of edge heat flux loads, which depend on the instant amplitude of fluctuations, as opposed to the mean load, which can be calculated by quasilinear theory.

In terms of mathematical description, the likelihood of intermittent events related to plasma turbulence is expressed by probability density functions (PDFs), which usually deviate significantly from the Gaussian distribution. Along these lines, there have been attempts to characterize the statistical properties of the PDFs, based on phenomenological premises [10, 11] or numerical investigations [12, 13] alone. Here, however, we carry out a direct comparison between first-principles analytical modeling and numerical simulations. As will be evident in the sequel, although the two approaches express the same physics, they nevertheless greatly differ in their theoretical backgrounds. A key finding of this work is that the intermittent process in the context of drift-wave turbulence appears to be independent of the specific modeling framework, opening the way to the prediction of its salient features.

The main part of this Letter consists in providing a theoretical interpretation of the PDFs of radial heat flux derived by nonlinear, local, gyrokinetic simulations of drift wave turbulence in tokamaks. Our paradigm in this work is Ion-Temperature-Gradient (ITG) turbulence with adiabatic electrons [14]. The simulations have been carried out with the GENE code [15] in a simple large aspect ratio, circular tokamak geometry. In particular, we calculate the turbulent ion radial heat flux,  $Q = \left\langle v_r \int \left( \frac{1}{2} m_i v_{\parallel}^2 + \mu B \right) F_i \right\rangle$ , where  $v_r$  is the radial  $\mathbf{E} \times \mathbf{B}$  velocity,  $v_{\parallel}$  the parallel velocity,  $\mu$  the magnetic

moment,  $m_i$  the ion mass,  $F_i$  the perturbed ion gyrocenter distribution function and  $B$  the modulus of the magnetic field. The brackets denote spatial averaging over the entire simulation domain. In order to perform a reliable comparison, we produce a series of different cases, by varying the magnetic shear  $\hat{s} = \frac{r}{q} \frac{dq}{dr}$  ( $q$  is the safety factor, i.e., the number of toroidal turns of the magnetic field for each poloidal turn), which appears both in the gyrokinetic simulations and the theoretical model. There is nothing special about the selection of this parameter, and a similar scan could be performed over, for example, the ion temperature or density gradient. Here, we have set the (normalized) ion temperature gradient  $R/L_{T_i} = 9$ , the density gradient  $R/L_n = 2$ , and the ion-to-electron temperature ratio  $\tau = 1$ .

For each case, the time evolution of the ion heat flux is considered as a time series, to which we apply standard Box-Jenkins modeling [16]. This mathematical procedure effectively removes deterministic autocorrelations from the system, allowing for the statistical interpretation of the residual part, which a posteriori turns out to be relevant for comparison with the analytical theory. In our setup, it turns out that an ARIMA(3,1,0) model accurately describes the stochastic procedure, in that, one can express the (differenced) heat flux time trace in the form

$$Q_{t+1} = a_1 Q_t + a_2 Q_{t-1} + a_3 Q_{t-2} + Q_{res}(t) \quad (1)$$

where the fitted coefficients  $a_1, a_2, a_3$  describe the deterministic component and  $Q_{res}(t)$  is the residual part (noise). The reliability of the produced autoregressive model was routinely verified via portmanteau testing and overfitting.

It is systematically observed that the residual PDFs are manifestly non-Gaussian with elevated tails, as shown for instance in Fig. 1, for the case  $\hat{s} = 1.0$ , where the sample residuals from a GENE simulation are tested against the Gaussian distribution, via a normal quantile-quantile plot. Furthermore, two significant statistical quantities, namely the variance  $\sigma^2 = \langle Q^2 \rangle - \langle Q \rangle^2$  and the kurtosis  $K = \langle Q^4 \rangle / \sigma^4$  (here, the brackets denote averaging over the statistical sample), during the magnetic shear scan are summarized in Table 1. Evidently, the decrease

of the variance and the rapid increase in kurtosis for increasing magnetic shear renders a Gaussian description improper. We note in passing, that the values of  $\hat{s}$  were selected such that the resolution of the simulation domain remains constant, thus avoiding numerical artifacts.

In the sequel, it will be shown that the aforementioned PDF tails can be satisfactorily predicted, starting from a fluid model consisting of a continuity and an energy equation [17], and using a nonperturbative statistical technique, called the instanton method, which has been adopted from Quantum Field Theory and then modified to classical statistical physics for Burgers turbulence and in the passive scalar model of Kraichnan [18]. In this context, the key element is to identify the bursty or intermittent event with the appearance of a coherent structure (e.g., streamer).




FIG. 1: Normal quantile-quantile plot showing strong deviation of the residual PDF from Gaussian statistics, in view of the enhanced tails representing about half of the sample.

TABLE I: Simulation Results

$\hat{s}$	Variance	Kurtosis
0.25	5.962e-4	3.529
0.40	5.253e-4	3.517
0.50	6.049e-4	3.527
0.60	5.874e-4	3.715
0.75	4.403e-4	4.324
1.00	8.253e-5	14.413

In the following paragraph, we briefly outline the implementation of the instanton method. For more details, the reader is referred to the existing literature [20]- [23]. The PDF tail is first formally expressed in terms of a path integral by utilizing the Gaussian statistics of the forcing in the continuity equation in a similar spirit as in [9]. An optimum path will then be associated with the creation of a modon (among all possible paths or functional values) and the action ( $S_\lambda$ , below) is evaluated using the saddle-point method on the effective action in the limit  $\lambda \rightarrow \infty$ .

The instanton is localized in time, existing during the formation of the modon. The saddle-point solution of the dynamical variable  $\phi(\vec{x}, t)$  of the form  $\phi(\vec{x}, t) = F(t)\psi(\vec{x})$  is called an instanton if  $F(t) = 0$  at  $t = -\infty$  and  $F(t) \neq 0$  at  $t = 0$ . Note that, the function  $\psi(\vec{x})$  here represents the spatial form of the coherent structure. Thus, the intermittent character of the transport consisting of bursty events can be described by the creation of modons. The probability density function of the heat flux  $Q$  can be defined as

$$P(Q) = \langle \delta(v_r n T_i(\vec{r} = \vec{x}_0) - Q) \rangle = \int d\lambda e^{i\lambda Q} I_\lambda, \quad (2)$$

where

$$I_\lambda = \langle \exp(-i\lambda v_r n T_i(\vec{x} = \vec{x}_0)) \rangle. \quad (3)$$

Here,  $v_r$  is the radial drift velocity and  $T_i$  the ion temperature. The integrand can then be rewritten in the form of a path-integral as

$$I_\lambda = \int \mathcal{D}\phi \mathcal{D}\bar{\phi} e^{-S_\lambda}. \quad (4)$$

Although  $\bar{\phi}$  appears to be simply a convenient mathematical tool, it does have a useful physical meaning that should be noted; it arises from the uncertainty in the value of  $\phi$  due to the stochastic forcing. That is, the dynamical system with a stochastic forcing should be extended to a larger space involving this conjugate variable, whereby  $\phi$  and  $\bar{\phi}$  constitute an uncertainty relation. Furthermore,  $\bar{\phi}$  acts as a mediator between the observables (heat flux) and instantons (physical variables) through stochastic forcing. In Eq. 4, the integral in  $\lambda$  is computed using the saddle-point method where it is shown that the limit  $\lambda \rightarrow \infty$  corresponds to  $Q \rightarrow \infty$ , representing the tail part of the distribution. Based on the assumption that the total PDF can be characterized by an exponential form, the expression

$$P(Q) = \frac{1}{Nb} \exp\{-b|Q - \mu|^{3/2}\}, \quad (5)$$

$$b = b_0 \left( \frac{R}{L_n} + 2\langle g_i \rangle \beta - U - \langle k_\perp^2 \rangle \left( U + \frac{R}{L_n} \right) \right), \quad (6)$$

$$\beta = 2 + \frac{2}{3} \frac{R/L_n - U}{U + 10/3\tau \langle g_i \rangle}. \quad (7)$$

is found [23], where the heat flux  $Q$  plays the role of the stochastic variable, with  $P(Q)$  determining its statistical properties. Several auxiliary definitions are also utilized; the normalization constant  $N$ ; the gradient scale lengths  $L_f = -(d \ln f / dr)^{-1}$ ; the normalized modon speed  $\tilde{U} = RU/L_n = R/L_n$  and temperature ratio  $\tau = T_i/T_e$ ;  $g_i = \omega_d / \omega_\star = \frac{2L_n}{R} (\cos(\xi) + \hat{s}\xi \sin(\xi))$  where  $\omega_d$  is the curvature drift frequency and  $\omega_\star$  is the diamagnetic drift frequency;  $k_\perp^2 = k_y^2 (1 + \hat{s}^2 \xi^2)$  is the perpendicular wave number; the brackets denote averaging along the field line, e.g. for an arbitrary scalar function  $f$ ,  $\langle f \rangle = \int_{-\pi}^{\pi} d\xi \phi f \phi / \int_{-\pi}^{\pi} d\xi \phi^2$  where the eigenfunctions  $\phi(\xi)$  are extracted from the GENE simulations and

$\mu = \langle Q \rangle$  is the mean value of the heat flux. The coefficient  $b_0$  is a free parameter and represents the strength of the forcing in the continuity equation. Note that the proposed PDF is close enough to a Gaussian distribution to match the bulk of the PDF while retaining the enhanced tails. Furthermore, the exponential form of the PDF will be the same for momentum flux with a modified coefficient  $b$ , which is in agreement with findings in recent experiments [24].

In Fig. 2 ( $\hat{s} = 0.25$ ), the PDF of heat flux  $Q$  (i.e. stemming from the *raw* timetrace of the simulation) (blue line) is shown. Although a simple visual inspection precludes the Gaussian distribution, the analytically predicted instanton method from Eq. 5 (red line) does a rather poor job modeling the simulation result. This is, however, contrasted to the PDF using the weak non-linear model (dashed-dotted black line) where reasonably good agreement is found similar to what has been reported earlier [6, 9]. Note that all PDFs shown are normalized such that the total probability is unity. The weak non-linear model (a Laplace distribution) has the form  $P(Q) = e^{-|Q-\mu|/b}/(2b)$  where  $b$  is determined by the variance. It is important to stress, however, that it can be predicted by the instanton model only when the non-linearities are neglected [25]. The Laplace distribution can easily be obtained by assuming that the background fluctuating fields (i.e. the potential and temperature) are close to Gaussian distributions whereas the resulting distribution for the heat flux is non-Gaussian with an unknown correlation factor. This fact points to the possibility that the total PDF does not reliably represent the statistics of the intermittent process. Along these lines, the well-studied K-S scaling, based on the total PDF, does not prove to be insightful [26].




FIG. 2: The numerically estimated PDF of heat flux (dashed blue line) in comparison with the analytically predicted instanton method (red line) and the weak non-linear model (dashed-dotted black line) at magnetic shear  $\hat{s} = 0.25$ .

Figs. 3 and 4 single out the comparison of the numerically estimated residual PDFs against the instanton prediction in Fig. 1. Here, the PDF of  $Q_{res}$  from

the simulation (blue line) and the analytically predicted PDF from the instanton method (dashed-dotted black line) demonstrate a significantly better agreement both at the tails and the center of the PDF, as compared to the best fit (using the first two moments) Gaussian distribution (dashed red line) at magnetic shears  $\hat{s} = 0.25$  and  $\hat{s} = 1.0$ , respectively. The coefficient  $b_0$  in Eq. 6 is determined from the simulations at the point  $\hat{s} = 0.5$  and for the weak non-linear model a value of the variance of  $8.00e-4$  was used for better agreement. Note that the weak model does not capture the form of the residual PDF while surprisingly good agreement is found for the instanton model.




FIG. 3: Comparison of the numerically estimated PDF (blue line) against the analytically predicted instanton model (dashed-dotted black line) and weak non-linear model (dashed green line) compared to the best fit Gaussian (dashed red line) for  $\hat{s} = 0.25$ .




FIG. 4: Comparison of the numerically estimated PDF (blue line) against the analytically prediction (dashed dotted black line) and the best fit Gaussian (dashed red line) for  $\hat{s} = 1.0$ .

Since the quality of the agreement in the previous figures relies heavily on the exponential coefficient  $b$  in Eq.

5, this is displayed for the whole magnetic shear scan in Fig. 5, verifying that good agreement is expected for any value of  $\hat{s}$ . Here, we note that the abrupt bending of the curve for larger values of shear is solely attributed to FLR effects.




FIG. 5: The coefficient  $b$  as a function of magnetic shear  $\hat{s}$ . The simulation results (red squares) are compared to the analytical fluid model prediction (black stars) and the same with suppressed FLR effects (blue diamonds).

In conclusion, we have presented a first quantitative comparison between a first-principles theoretical model of drift wave turbulence with self-consistent non-linear simulations. Numerical PDFs of heat flux were generated with the gyrokinetic code GENE in the framework of toroidal ITG turbulence, and subsequently processed

with Box-Jenkins modeling, in order to remove deterministic autocorrelations, thus retaining their stochastic parts only. These PDFs have been shown to agree very well with analytical predictions based on a fluid model, on applying the instanton method. Specifically, we were able to quantitatively confirm the exponential form of the PDFs, therefore adding the important element of predictive strength to the existing phenomenological approaches. More importantly, a strong indication of universality in the description of drift wave turbulence has emerged.

In future publications we will address the emergent universal scalings of the PDFs of potential, density and temperature where the theory predicts different tails [25]. A study of potential or density fluctuations opens up opportunities for comparison with experimentally measured PDFs. Still within the scope of ITG turbulence, we will include effects from kinetic electrons, in order to test the robustness of the exponential scaling. A quite interesting complementary work would involve incorporating results from nonlinear fluid simulations, based on Braginskii-like equations, and relate them to the already presented findings. Finally, our present setup provides a testbed for the investigation of the impact of zonal flows on the statistics, as reported elsewhere [12]. Work addressing these avenues has been initiated.

#### Acknowledgments

The GENE simulations have been performed at the Jülich Supercomputing Center (JSC). The Authors are grateful to Prof. P. Helander and Prof. F. Jenko for many fruitful discussions.

- 
- [1] S. T. Bramwell, *Nature* **5**, 443 (2009)
  - [2] S. Zweben, J. A. Boedo, O. Grulke, C. Hidalgo, B. LaBombard, R. J. Maqueda, P. Scarin and J. L. Terry, *Plasma Phys. Contr. Fusion* **49**, S1 (2007)
  - [3] P. A. Politzer, *Phys. Rev. Lett.* **84**, 1192 (2000)
  - [4] P. Beyer, S. Benkadda, X. Garbet and P. H. Diamond, *Phys. Rev. Lett.* **85**, 4892 (2000)
  - [5] J. F. Drake, P. N. Guzdar and A. B. Hassam, *Phys. Rev. Lett.* **61**, 2205 (1988)
  - [6] B. A. Carreras, B. van Milligen, C. Hidalgo, R. Balbin, E. Sanches, I. Garcia-Cortes, M. A. Pedrosa, J. Bleuel and M. Endler, *Phys. Rev. Lett.* **83**, 3653 (1999)
  - [7] F. Sattin, P. Scarin, M. Agostini, R. Cavazzana, G. Seriani, M. Spolaore and N. Vianello, *Plasma Phys. Contr. Fusion* **48** 1033 (2006)
  - [8] G. Y. Antar, S. I. Krashenninnikov, P. Devynck, R. P. Doerner, E. M. Hollmann, J. A. Boedo, S. C. Luckhardt and R. W. Conn, *Phys. Rev. Lett.* **87**, 065001 (2001)
  - [9] B. A. Carreras, C. Hidalgo, E. Sanches, M. A. Pedrosa, R. Balbin, I. Garcia-Cortes, B. van Milligen, D. E. Newman and V. E. Lynch, *Phys. Plasmas* **3**, 2664 (1996)
  - [10] I. Sandberg, S. Benkadda, X. Garbet, G. Ropokis, K. Hizandis and D. del-Castillo-Negrete, *Phys. Rev. Lett.* **103**, 165001 (2009)
  - [11] J. A. Krommes, *Phys. Plasmas* **15**, 030703 (2008)
  - [12] R. Sánchez, D. E. Newman, J.-N. Leboeuf, B. A. Carreras and V. K. Decyk, *Phys. Plasmas* **16**, 055905 (2009)
  - [13] G. Dif-Pradalier, P. H. Diamond, V. Grandgirard, Y. Sarazin, J. Abiteboul, X. Garbet, Ph. Ghendrih, A. Strugarek, S. Ku and C. S. Chang, *Phys. Rev. E* **82**, 025401 (2010)
  - [14] W. Horton, *Rev. Mod. Phys.* **71**, 735 (1999)
  - [15] F. Jenko, W. Dorland, M. Kotschenreuther and B. N. Rogers, *Phys. Plasmas* **7**, 1904 (2000)
  - [16] G. Box, G. Jenkins, G. Reinsel, *Time series analysis; Forecasting and control*, Prentice Hall, 1994
  - [17] J. Anderson, H. Nordman, R. Singh and J. Weiland, *Phys. Plasmas* **9**, 4500 (2002)
  - [18] E. Balkovsky and V. Lebedev, *Phys. Rev. E* **58**, 5776 (1998)
  - [19] J. Zinn-Justin, *Field Theory and Critical Phenomena* (Clarendon, Oxford, 1989) p. 57
  - [20] V. Gurarie and A. Migdal, *Phys. Rev. E* **54**, 4908 (1996)
  - [21] G. Falkovich, I. Kolokolov, V. Lebedev and A. Migdal, *Phys. Rev. E* **54**, 4896 (1996)
  - [22] E. Kim and P. H. Diamond, *Phys. Rev. Lett.* **88**, 225002 (2002)
  - [23] J. Anderson and E. Kim, *Phys. Plasmas* **15**, 082312

- (2008)
- [24] Z. Yan, G. R. Tynan, J. H. Yu, C. Holland, S. H. Muller and M. Xu, *Bull. Am. Phys. Soc.* **52**, 74 (2007)
- [25] E. Kim and J. Anderson, *Phys. Plasmas* **15**, 114506 (2008)
- (2008)
- [26] F. Sattin, M. Agostini, R. Cavazzana, G. Serianni, P. Scarin and N. Vianello, *Phys. Scripta* **79**, 045006 (2009)