

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.

This discussion paper is/has been under review for the journal Biogeosciences (BG).
Please refer to the corresponding final paper in BG if available.

Organic matter quality of deep permafrost carbon – a study from Arctic Siberia

J. Strauss¹, L. Schirrmeister¹, K. Mangelsdorf², L. Eichhorn³, S. Wetterich¹, and U. Herzschuh¹

¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Periglacial Research Unit Potsdam, Telegrafenberg A 43, Potsdam, Germany

²Helmholtz Centre Potsdam GFZ German Research Centre for Geosciences, Telegrafenberg, 14473 Potsdam, Germany

³International Max Planck Research School for Global Biogeochemical Cycles (Max-Planck-Institute for Biogeochemistry and Friedrich Schiller University), Burgweg 11, 07749 Jena, Germany

Received: 5 August 2014 – Accepted: 2 November 2014 – Published: 21 November 2014

Correspondence to: J. Strauss (jens.strauss@awi.de)

Published by Copernicus Publications on behalf of the European Geosciences Union.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

The organic carbon (OC) pool accumulated in Arctic permafrost (perennially frozen ground) equals the carbon stored in the recent atmosphere. To give an idea of how Yedoma region permafrost could respond under future climatic warming, we conducted a study to quantify the organic matter quality for future decomposition of late Pleistocene (Yedoma) and Holocene (thermokarst) deposits on the Buor Khaya Peninsula, northeast Siberia. The objective of this study was to develop a stratigraphic classified organic matter quality characterization. For this purpose the degree of organic matter decomposition was estimated by using a multiproxy approach. We applied sedimentological (grain-size analyses, bulk density, ice content) and geochemical parameters (total OC, stable carbon isotopes ($\delta^{13}\text{C}$), carbon : nitrogen (C/N) ratios) as well as lipid biomarkers (*n*-alkanes, *n*-fatty acids, hopanes, triterpenoids, and biomarker proxies/indices: average chain length, carbon preference index (CPI), and higher plant fatty acid index (HPFA)). Our results show that the Yedoma and thermokarst organic matter qualities exhibit no obvious degradation – depth trend. The C/N, $\delta^{13}\text{C}$, and hop-17(21)-ene values and the HPFA index show a better quality of the organic matter stored in thermokarst deposits compared to Yedoma deposits, but the CPI points in the other direction. As the ranges of the proxies mostly overlap, we interpret this as to indicate similar quality for both kind of deposits with perhaps slightly better thermokarst organic matter quality. Supported by principal component analyses, the sediment parameters and quality proxies of Yedoma and thermokarst deposits could not be clearly separated from each other. This lack of clear quality differences revealed that the organic matter vulnerability is heterogeneous, independent from radiocarbon age and depends on different decomposition trajectories and the previous decomposition and preservation history. Elucidating this was one of the major novelties of our multiproxy study. With the addition of biomarker data, it was possible to show that permafrost organic matter degradation likely occurs via a combination of (uncompleted) degradation cycles or a cascade of degradation steps rather than as a linear function of age or sediment

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



facies. We conclude that the amount of organic matter in the studied sediments is high for mineral soils and of good quality and therefore susceptible to future decomposition. The missing depth trends reveal that permafrost acts like a giant freezer, preserving the constant quality of ancient organic matter independently from its age. When undecomposed Yedoma organic matter is mobilized via thermokarst processes, the fate of this carbon depends largely on the environmental conditions; the carbon could be preserved in an undecomposed state till refreezing occurs. If recent input has occurred, thermokarst organic matter could be of a better quality than that found in Yedoma deposits.

1 Introduction

During the late Quaternary, the rate of organic matter decomposition in the Arctic has been slower than plant growth, sedimentation, and freezing rates. Thus, a large pool of organic carbon (OC) accumulated in the Arctic and was deeply sequestered in the permafrost. Hugelius et al. (2014) estimates an OC storage of 1300 Gt for the circum-Arctic permafrost region with ~ 850 Gt OC sequestered in permafrost. This is approximately the carbon stored in the recent atmosphere (Dlugokencky and Tans, 2014). During warming and permafrost thawing, this formerly cryo-sequestered OC gradually entered the recent biogeochemical cycle by microbial turnover. By thawing and microbial activity, the permafrost deposits can turn from a carbon sink to a source (Schuur et al., 2009), releasing greenhouse gases such as carbon dioxide and methane to the atmosphere. Besides the near-surface carbon pool representing the uppermost 3m below surface, and because of rapid permafrost thaw like thermokarst and thermoerosion, deep OC pools, especially those held in ice-rich permafrost deposits in the Yedoma region, are of great significance for current concerns about the effects of global warming. According to Strauss et al. (2013) the Yedoma region is defined as the area of potential distribution of late Pleistocene ice-rich and organic-rich silty deposits (Yedoma) covering large areas in Siberia and Alaska. Estimates of OC stored in the

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Yedoma region amount to 83 +61/ – 57 Gt for late Pleistocene Yedoma deposits (ages shown in Table 1). Due to Holocene warming, subsequent ground ice melt and surface subsidence, thermokarst basins formed and were partly occupied by lakes. Holocene thermokarst deposits (ages shown in Table 1) contain 128 +99/ – 96 Gt organic carbon. In total, the Yedoma region extends to an area of about 1 387 000 km² of which about 70 % is already affected by permafrost degradation (thermokarst) (Strauss et al., 2013). Kuhry et al. (2009) and Schirrmeister et al. (2011a) showed that Yedoma deposits accumulated at fast rates, implying a short time for the organic matter to decay before it became locked into a perennially-frozen state. Therefore, the organic matter availability for microorganisms is expected to be excellent, resulting in great vulnerability to warming ground conditions (Mu et al., 2014). To elucidate how Yedoma region permafrost could respond under conditions of future climatic warming, we studied the organic matter quality of Yedoma and its Holocene degradation features (called thermokarst deposits) on Buor Khaya Peninsula, Eastern Laptev Sea. As mentioned above, Strauss et al. (2013) found that thermokarst deposits contain the quantitatively more important carbon pool, but the unsolved question is this: Is the thermokarst organic matter pool as degradable as the frozen late Pleistocene Yedoma, or has the most labile carbon already been emitted due to thermokarst degradation processes? In both kinds of deposits the OC was deeply (deeper than 3 m) incorporated into permafrost (Schirrmeister et al., 2013; Strauss et al., 2013). As shown by models and extrapolation from recent observations, the more southern portions of Yedoma deposits thawed during the last deglaciation, resulting in large emissions of greenhouse gases to the atmosphere (Walter et al., 2007a; Ciais et al., 2012; Walter Anthony et al., 2014). Recent ground warming has been observed in the permafrost zone (Romanovsky et al., 2010), and incubation experiments reveal that permafrost warming is accompanied by a substantial outgassing of greenhouse gases (Lee et al., 2012; Knoblauch et al., 2013; Schädel et al., 2014). As an illustration of the important influence of ground temperature on organic matter quality, a higher respiration rate at greater depth close to the permafrost table (Mangelsdorf et al., 2009; Waldrop et al., 2010) was found inside the

seasonally-thawed active layer and interpreted as a greater lability of the organic matter close to the perennally frozen ground. Focusing on permafrost deposits in the Laptev Sea region, which includes our Buor Khaya study site, Schirrmeister et al. (2011a) characterize the Yedoma region permafrost organic matter as weakly decomposed.

5 Biomarkers are used for paleoenvironmental reconstruction of terrestrial permafrost (Andersson et al., 2011) or characterization of permafrost organic matter degradation (Andersson and Meyers, 2012; Vonk et al., 2013; Routh et al., 2014). In our study we estimate molecular markers (*n*-alkanes, *n*-fatty acids, hopanes, and triterpenoids) and use biomarker proxies/indices (absolute lipid concentration, average chain length
10 (ACL), carbon preference index (CPI), hop-17(21)-ene, higher plant fatty acid (HPFA) index, and an Oleanen ratio) to test whether they are useful mirrors of organic matter decomposition, i.e. organic matter quality in permafrost deposits. Rather established methods, both cryolithological (grain size analyses, bulk density, ice content) and biogeochemical (total organic carbon (TOC_{wt%}), stable carbon isotope ratios ($\delta^{13}\text{C}$ in
15 TOC), total nitrogen (TN), and TOC_{wt%}/TN (C/N) ratios), are applied to our sample set. Finally, principal components analysis (PCA) highlights the relationships between different organic matter degradation proxies. Because the future feedback from the Yedoma region permafrost OC to climate forcing is driven by both (1) the pool size, estimated to be ~ 211 Gt (Strauss et al., 2013), and (2) the quality of OC stored in the
20 studied deposits, the objective of this study is the development of a stratigraphically differentiated organic matter quality characterization using sample material representative of widespread Yedoma and thermokarst permafrost.

2 Material and methods

2.1 Study area

25 The Buor Khaya Peninsula study site (71°34' N, 132°12' E) is located in the north-eastern part of Siberia (Fig. 1). Buor Khaya Peninsula is framed by the Laptev Sea,

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



a shallow epicontinental part of the Arctic Ocean, and geologically by two rift structures (Drachev et al., 1998). Buor Khaya is underlain by continuous permafrost with ground temperatures of less than -11°C (Drozhdov et al., 2005). The permafrost thicknesses is estimated to be between 450 and 650 m (Romanovskii et al., 2004). Stratigraphically, outcrops from two sediment units are distinguished and studied, (1) ice-rich permafrost, called Yedoma deposits, and (2) deposits in permafrost rapid thaw features, generalized as thermokarst deposits. Three profiles of thermokarst deposits (in a thermokarst basin: Buo-01 and Buo-05; initial thermokarst on top of a Yedoma hill: Buo-03) and two profiles of Yedoma deposits (Buo-02, Buo-04) were studied and sampled. Figure 1 shows an overview of the sampled profiles and their position relative to each other.

2.2 Field work

Field studies were undertaken in summer 2010 at outcrops situated at the western coast of the Buor Khaya Peninsula. The sediment of the profiles and sub-profiles, exposed at the cliff wall or partly in thermokarst mounds in thaw slumps, were dug by spades and cleaned with hacks. The cryolithology, sediment characteristics, and visible organic matter in the sediments of the chosen sequences were surveyed and described. Moreover, the profiles were photographed and sketched. Sub-profiles were stacked together to create composite profiles. Sampling positions in neighboring sub-profiles were correlated by height estimation using measuring tape. The upper edge of each profile was calibrated with tacheometer measurements (Günther et al., 2012). In the field laboratory all sample volumes were measured with a balance and Archimedes principle, and the absolute ice content was determined by drying the sample. In total, 91 samples were taken and kept cool for transport to laboratories for further analysis. Detailed sampling positions for each profile are shown in Strauss and Schirrmmeister (2011).

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3 Indicators of organic matter quality

To validate and to extend the sedimentological approach used, and to estimate the organic matter quality, lipid biomarkers were measured to estimate the degree of organic matter degradation. For biomarker studies we used a “fingerprint” approach by focusing on identifiable markers related to organic matter quality. Below, the utilized geochemical indicators and biomarkers are described.

2.3.1 Grain-size analyses

Grain sizes were analyzed using a laser particle sizer (LS 200, Beckmann–Coulter) between 0.375 and 1000 μm (Figs. 2 and S1). Grain-size calculations were done after Folk and Ward (1957) using Gradostat v8 (Blott and Pye, 2001). A detailed description of this analytical techniques is given in the Supplement (Supplement Sect. 1.1).

2.3.2 Elemental composition

To determine the total elemental carbon and total nitrogen (TN) content, the samples were measured by a carbon-nitrogen-sulphur analyzer (Vario EL III, Elementar). $\text{TOC}_{\text{wt}\%}$ was measured with a TOC analyzer (Vario Max C, Elementar). The volumetric TOC content ($\text{TOC}_{\text{kg m}^{-3}}$) was calculated according to Strauss et al. (2013). A detailed description of this techniques is given in the Supplement (Supplement Sect. 1.2).

The C/N ratio has been used as a general indicator of the degree of organic matter decomposition (Stevenson, 1994). Based on the assumption that organic matter components are degraded selectively, degradation modifies elemental compositions and hence C/N in deposits. Because a decrease in the C/N ratio has been observed in aerated deposits with microbial immobilization of TN (nitrogen stays in the system) accompanied by the re-mineralization of TOC (Sollins et al., 1984) and CO_2 emission, this ratio is used in the following way: The higher the C/N ratio, the lower the degree of decomposition. This assumption implies a similar source signal.

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.3 Bulk density and volumetric carbon content

BD was calculated using Eq. (1).

$$\text{BD} [10^3 \text{ kg m}^{-3}] = \frac{\text{sample dry weight} [10^3 \text{ kg}]}{\text{sample volume} [\text{m}^3]} \quad (1)$$

Estimating the BD is required to convert the measured-weight-based $\text{TOC}_{\text{wt}\%}$ content per sample to a volume-based value. Thus, the $\text{TOC}_{\text{kg m}^{-3}}$ was calculated according to Eq. (2):

$$\text{TOC}_{\text{kg m}^{-3}} = \text{BD} [10^3 \text{ kg m}^{-3}] \times \frac{\text{TOC}_{\text{wt}\%}}{100} \quad (2)$$

2.3.4 Carbon isotope studies

Stable TOC carbon isotopes were determined with a Finnigan MAT Delta-S mass spectrometer combined with a FLASH elemental analyzer and a CONFLO III gas mixing system. A detailed method is given in the Supplement (Supplement Sect. 1.4). The stable carbon isotopes of OC reflect (1) initial contribution from different plant species and plant components, and (2) subsequent degradation processes (Gundelwein et al., 2007). Assuming constant photosynthetic isotope fractionation in source plants in the region (C_3 plants are ubiquitous in the Arctic, Tieszen, 1973), we use $\delta^{13}\text{C}$ ratios as a degradation proxy. After Heyer et al. (1976), decomposition discriminates against the lighter isotope (^{12}C), resulting in more negative $\delta^{13}\text{C}$ ratios. Thus, this proxy is used in the following way: Lower (more negative) $\delta^{13}\text{C}$ values are connected to less degraded material, while higher (less negative) $\delta^{13}\text{C}$ values reflect greater decomposition.

Agers were determined by radiocarbon dating of selected macroscopic plant remains performed at the Poznań Radiocarbon Laboratory, Poland (Goslar et al., 2004). The presented radiocarbon ages are uncalibrated ages; Table 1 includes calibrated ages as well. Radiocarbon ages are given in year before present (a BP).

15952

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.3.5 Lipid biomarkers

To look more closely at the molecular composition, we used specific lipid biomarkers. Molecular fossils or biomarkers were studied by chromatography methods coupled with mass spectrometers. Characteristic fractions like *n*-alkanes, *n*-fatty acids, sterols, and hopanes were isolated. Because the $\text{TOC}_{\text{wt}\%}$ in the profiles is not equally distributed, we calculated and visualized the biomarker concentration as $\mu\text{g g TOC}_{\text{wt}\%}^{-1}$ and $\mu\text{g g Sediment}^{-1}$ ($\mu\text{g g Sed}^{-1}$). For the results, we focus on $\mu\text{g g TOC}_{\text{wt}\%}^{-1}$.

Extraction and fraction separation

For lipid biomarker analyses 2–12 g of ground sediment was weighed in an extraction cell with an accelerated solvent extractor (ASE 200, Dionex). Samples were extracted with dichloromethane/methanol (99 : 1). Each sample was held in a static phase for 20 min at 75 °C (after 5 min heating, no preheating) at a pressure of 5 MPa. Afterwards, the dissolved compounds were concentrated with a Turbo Vap (Zymark) closed cell concentrator and further dried by evaporating the solvent in a stream of nitrogen gas. After that, internal standards (5 α -androstane for the aliphatic fraction, ethylpyrene for the aromatic fraction, 5 α -androstan-17-on for nitrogen-, sulfur-, and oxygen- (NSO-) containing compounds, and erucic acid for the NSO fatty acid fraction) were added. The amount of internal standards depended on the $\text{TOC}_{\text{wt}\%}$ content (< 10wt%: 8 μg ; > 10 to \leq 25wt%: 20 μg ; > 25wt%: 50 μg). After the removal of the *n*-hexane-insoluble fraction (by the addition of a large excess of *n*-hexane, called “asphaltene” precipitation), the hexane-soluble portion of the extract was separated by medium-pressure liquid chromatography (MPLC; Radke et al., 1980) into fractions of different polarity (aliphatic and aromatic hydrocarbons as well as polar hetero (NSO) components). Afterwards, the NSO fraction was split into a fatty acids and an alcohol fraction using a KOH-impregnated silica gel column (Schulte et al., 2000).

For this study, the focus was placed on the aliphatic fraction (containing *n*-alkanes and triterpenoid compounds) and the NSO fraction (containing *n*-fatty acids). The frac-

tions were measured by gas chromatography–mass spectrometry (GC–MS). All compounds of interest were identified using the Xcalibur software (Thermo Fisher Scientific).

GC-MS measurement and compound quantification

5 The *n*-alkanes, *n*-alcohols, hopenes (hop-17(21)-ene), and other triterpenoids (β -amyryn (olean-12-en-3 β -ol), Olean-12-ene, and Olean-13(18)-ene) were measured with a GC-MS system (GC: Trace GC Ultra; MS: DSQ, both Thermo Fisher Scientific). Prior to the measurements, the *n*-fatty acids were methylated with diazomethane and the alcohols were silylated with *N*-methyl-*N*-trimethylsilyltrifluoroacetamide (MSTFA).
10 The GC was equipped with a programmable temperature vaporization (PTV) injector system (starting temperature of 50 °C; heating rate of 10 °C s⁻¹ to 300 °C; isothermal holding time of 10 min; operated in splitless mode) and a fused silica capillary column (SGE BPX5, 50 m length, 0.22 mm inner diameter, 0.25 μ m film thickness). For the measurements the GC oven was programmed with a starting temperature of 50 °C, a heating rate of 3 °C min⁻¹ to 310 °C, and an isothermal holding time of 30 min. Helium
15 with a constant flow rate of 1 mL min⁻¹ was used as a carrier gas. For the *n*-fatty acid fraction a different temperature program (starting temperature of 50 °C, 1 min isotherm, heating rate of 3 °C min⁻¹ to 350 °C, isothermal holding time 25 min) was used. For compound identification, the gas chromatograph was linked to a mass spectrometer,
20 which was operated in electron impact ionization mode at 70 eV. The temperature of the ion source was set to 230 °C. Full scan mass spectra were recorded from *m/z* 50 to 600 Da at a scan rate of 2.5 scans s⁻¹. For the *n*-fatty acids fraction the scan rate was *m/z* 50 to 650 Da.

25 Quantification of *n*-alkanes, *n*-fatty acids, and β -amyryn was done in the GC-MS total ion current chromatogram by relating the peak area of the target compound to the peak area of an internal standard of known concentration. Other triterpenoids like Olean-12-ene, Olean-13(18)-ene, and hopene were quantified using the *m/z* 191 mass trace relative to the peak area of the β,β -diploptene (in the *m/z* 191 mass trace), the con-

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



centration of which was calculated in the total ion current chromatogram relative to the internal standard.

2.3.6 Biomarker proxies/indices

Absolute lipid concentration

- 5 The absolute lipid concentration is used as rough estimator of organic matter quality in the following sense: The higher the concentration, the better the conservation of the lipid, and the better the quality of the organic matter.

Carbon preference index

10 The CPI was introduced by Bray and Evans (1961) as the ratio of odd- to neighboring even-numbered alkanes, which is a measure of the alteration of organic matter. Here we use the improved formula after Marzi et al. (1993). In addition, we also applied the CPI for fatty acids in which even-numbered fatty acids predominate over adjacent odd n -fatty acids (Glombitza et al., 2009).

$$\text{CPI} = \frac{\left(\sum_{i=n}^m C_{2i+1} \right) + \left(\sum_{i=n+1}^{m+1} C_{2i+1} \right)}{2 \times \left(\sum_{i=n+1}^{m+1} C_{2i} \right)} \quad (3)$$

15 n : starting dominating chain length/2; m : ending dominating chain length/2; i : index (carbon number); C : concentration.

The CPI is used as a degradation/alteration proxy by quantifying the odd/even (n -alkanes, Fig. S2) or even/odd (n -fatty acids, Fig. S3) of the carbon chains (Bray and Evans, 1961; Glombitza et al., 2009). A low CPI means mature/degraded organic matter (e.g. CPI of crude oil ~ 1).

20

tion of the sample and of standard quantification was $< 5\%$. Because acetate can act as excellent feedstock for microbes (Smith and Mah, 1980; Vieth et al., 2008) and it has been shown that acetate was rapidly consumed in the presence of oxygen and nitrate (Kuesel and Drake, 1995), we use this parameter in the following way. If there is an acetate concentration of $> 1 \text{ mgL}^{-1}$, the deposit, including its acetate, was less available for degradation, resulting in good organic matter quality.

2.4 Principal component analysis

Multivariate statistical techniques, like the PCA used here, allow the analysis of multiple variables in order to investigate connections between the different degradation proxies. Prior to the PCA, concentration data were transformed using a $\log(x+1)$ transformation, and TOC ($\text{wt}\%$ and kg m^{-3}) data were transformed using a square root transformation. We performed three PCA runs. First, a PCA of the sediment parameters was implemented to infer differences between Yedoma and thermokarst deposits. Second, a PCA of biomarker proxies was performed. For this purpose, other characteristics were added as supplementary variables (TOC $_{\text{wt}\%}$, TOC $_{\text{kg m}^{-3}}$, C/N, $\delta^{13}\text{C}$, grain size, BD, ice content, and depth) without inclusion in the PCA calculation. These supplementary variables have no influence on the PCA and were plotted afterwards in the PCA biplot. Third, a PCA was conducted on samples of the major odd n -alkanes to infer possible changes of the source organisms with the same supplementary variables as described above to relate the different biomarker proxies to each other. Computations were performed using the “vegan” package of R software (Oksanen, 2013).

3 Results

Stratigraphically, there are two types of deposition units at the study site. The first unit is composed of Yedoma deposits. The second unit represents thermokarst deposits resulting from thermal degradation of Yedoma. Grain-size distributions (Figs. 2 and S1)

BGD

11, 15945–15989, 2014

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and PCA of sediments illustrate that thermokarst deposits are made up of degraded Yedoma sediments. After Gubin and Veremeeva (2010) and Zanina et al. (2011) the Yedoma deposits soil types are mainly less-developed cryopedoliths containing more-developed paleocryosol parts (Figs. 1 and 2, labeled and grey-shaded areas).

3.1 Organic matter quality of Yedoma deposits

3.1.1 Sedimentological and biogeochemical proxies

The radiocarbon ages (Table 1, Fig. 3) of the Yedoma deposits range from infinite ages (> 55 000 a BP) at the very bottom to 30 100 a BP at the uppermost sampled Yedoma unit. This is comparable to other Yedoma sequences in the region (Schirrmeister et al., 2011b). The mean grain sizes show a decreasing trend in the Buo-04 lower Yedoma profile, from 28 μm at the bottom to 11 μm in the upper part of Buo-04-A. The Buo-02 Yedoma profile shows no trend, but exhibits a more heterogeneous mean-grain size including three maxima at 22.5 m.a.s.l. (32 μm), 23.7 m.a.s.l. (34 μm), and 25.5 m.a.s.l. (33 μm). Nevertheless, all Yedoma deposit samples are classified as poorly-sorted medium-to-coarse silts with a stable low clay fraction (< 15 %).

The $\text{TOC}_{\text{wt}\%}$ contents vary from 0.2 wt% at 5 m.a.s.l. to 24.0 wt% in a peaty paleocryosol horizon at 24 m.a.s.l. (Fig. 3). The mean $\text{TOC}_{\text{wt}\%}$ content is 2.4 wt% (median 0.97 wt%). Calculating the $\text{TOC}_{\text{kg m}^{-3}}$ according to Strauss et al. (2013) by utilizing the BD (between 0.1 and 1.5 10^3 kg m^{-3} ($10^3 \text{ kg m}^{-3} = \text{g cm}^{-3}$)) and ice content (without ice wedges; 21 to 90 vol%), the Yedoma sediments contain from 3 to 46 kg C m^{-3} with a mean of 14 kg C m^{-3} (median 9 kg C m^{-3}). The maxima correspond to the peaty horizons with large $\text{TOC}_{\text{wt}\%}$ contents and a low BD. Within the paleocryosol horizons, located at 6.8, 24.0 to 24.5, 24.8, and 27.8 to 28.9 m.a.s.l., maxima in the C/N ratio are observable. The C/N range in these horizons is 8 to 31. In the cryopedolith profile parts the C/N maximum is reached at the lowermost Buo-04-C sub-profile (17.7 and 16.7). The C/N of the rest of the Yedoma profile falls between 4.1 (at Buo-02-C, 23.7 m.a.s.l.) and 14.3 (below the paleocryosol at 23.5 m.a.s.l.)

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The $\delta^{13}\text{C}$ of the Yedoma deposits ranges between -29.0 and -24.7‰ . The minima fit well to the maxima of the C/N ratio in the paleocryosol horizons at 6.8, 24.0 to 24.5, 24.8, and 27.8 to 28.9 ma.s.l. The minimum C/N of the Buo-02-C sub-profile corresponds approximately to the $\delta^{13}\text{C}$ maximum (-25.0 to -24.7‰).

3.1.2 Biomarker proxies/indices

Every radiocarbon-dated sample and additional samples were used for biomarker analysis. In total 25 biomarker samples were analyzed.

A series of long-chain *n*-alkanes that exhibit a strong odd-carbon preference ranging from *n*-C₂₁ to *n*-C₃₃ are recognized in all Yedoma samples (Fig. S2). Moreover, the *n*-alkanes show a unimodal distribution maximizing at the C₂₇, C₂₉, or C₃₁ *n*-alkane (Fig. S2). The *n*-fatty acids show strong even-over-odd carbon number predominance and a bimodal distribution ranging from C₁₄ to C₃₀ (Fig. S3). The maxima are generally located at *n*-C₁₆ in the lower carbon number range and at *n*-C₂₄ in the higher carbon number range. Total *n*-alkanes and *n*-fatty acids concentrations related to TOC_{wt%} and sediment weight show a homogeneous pattern similar to that of the TOC_{wt%} and C/N values. The *n*-alkane concentration ranges from 3 to 75 $\mu\text{g g Sed}^{-1}$ (mean 20 $\mu\text{g g Sed}^{-1}$) and from 387 to 1715 $\mu\text{g TOC}_{\text{wt}\%}^{-1}$ (mean 1132 $\mu\text{g TOC}_{\text{wt}\%}^{-1}$). The *n*-fatty acids range from 4 to 306 $\mu\text{g g Sed}^{-1}$ (mean 51 $\mu\text{g g Sed}^{-1}$) and from 475 to 4669 $\mu\text{g TOC}^{-1}$ (mean 2196 $\mu\text{g TOC}_{\text{wt}\%}^{-1}$).

This Yedoma series shows distinct preference between even and odd carbon. The mean CPI values of the *n*-alkanes (12.2, ranging between 8.3 and 15.9) are higher than the CPI values of the *n*-fatty acids (4.9, ranging between 3.8 and 7.6). Because *n*-fatty acids are functional compounds (including a functional group, e.g. a carboxyl group), their degradation rates are much higher compared to those of *n*-alkanes (Poynter and Eglinton, 1990). This statement is also based on the assumption of similar sources. The ACL of the *n*-alkanes and *n*-fatty acids is very stable at around 28.4 (range 27.6 to 29.2) and 25.0 (range 23.8 to 25.6), respectively.

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Higher hop-17(21)-ene concentrations are used as an indicator for lower organic matter degradation state. In the lower Yedoma profile the hop-17(21)-ene ranges from $0.0 \mu\text{ggTOC}^{-1}$ at the lowermost and uppermost samples (4.3 and 18.5 m.a.s.l.) to the overall maximum at the Buor Khaya site ($19.4 \mu\text{ggTOC}^{-1}$) at 9.1 m.a.s.l. At Buo-02, the hop-17(21)-ene concentration is lower compared to the other Yedoma profile with a mean of $1.9 \mu\text{ggTOC}_{\text{wt}\%}^{-1}$ and a maximum of $7.7 \mu\text{ggTOC}_{\text{wt}\%}^{-1}$ in the potentially Holocene-contaminated uppermost sample. The HPFA ratio for the Yedoma samples is very stable around the mean value of 0.50 (median 0.54) with a minimum at 18.5 m.a.s.l. (0.15) and a maximum at the uppermost sample (0.69) at 29.7 m.a.s.l. For Yedoma, the Oleanen ratio is 0.0 (except a ratio of 10.0 at the uppermost sample). The acetate content of the Yedoma sample is between 0.6 and 57.5mgL^{-1} with a mean of 6.7mgL^{-1} (median 1.2mgL^{-1}).

3.2 Organic matter quality of thermokarst deposits

3.2.1 Sedimentological and biogeochemical proxies

The radiocarbon dating shows Holocene ages between 8140 and 3665 a BP (Fig. 4, Table 1). The lowermost Buo-05-C profile shows an age inversion for the two samples, (0.3 and 2.2 m.a.s.l.). The mean grain size at Buo-05 from the bottom to 6.7 m.a.s.l. is $13 \mu\text{m}$. Above, the mean grain size increases to $19 \mu\text{m}$. The Buo-05 clay fraction is stable at a low level ($< 15\%$). The Buo-01 profile shows a very scattered grain size ranging from 4 to $44 \mu\text{m}$ mean grain size. For the whole dataset, there is a maximum in the clay fraction (35%) in the peat horizon at 8.7 m.a.s.l. Buo-03 shows a slight decrease from 18 to $11 \mu\text{m}$. All thermokarst deposits are classified as (very) poorly-sorted silts. Similar to the Yedoma deposits, the BD of the thermokarst deposits is between 0.1 and $1.5 \cdot 10^3 \text{kgm}^{-3}$ and the ice content (without the ice wedges) is 23 to 87 wt%.

The mean $\text{TOC}_{\text{wt}\%}$ contents of the thermokarst deposits, 4.7 wt% (median 1.7 wt%), are higher compared to Yedoma deposits, varying between 0.2 and 43.0 wt%. Minimum

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and maximum $\text{TOC}_{\text{wt}\%}$ both occur at Buo-01 and exhibit the same scatter as in the grain sizes. $\text{TOC}_{\text{kg m}^{-3}}$ ranges between 2.8 and 93.5 kg C m^{-3} (mean 24 kg C m^{-3} , median 19 kg C m^{-3}).

At Buo-05 the C/N ratio is stable around 9 to 10, except for a paleocryosol horizon at 9.2 m a.s.l. that shows a value of 22. At Buo-01, the C/N ratio below the paleocryosol horizon is remarkably low, between 2 and 9, followed by the overall maximum in the peaty horizon with a ratio of 34. The Buo-03 cryopedolith samples show C/N ratios around 10, while the paleocryosol samples exhibit C/N ratios from 16 to 19. The $\delta^{13}\text{C}$ values range between -29.5 and -25.0‰ , with minima corresponding to the C/N maxima at the paleocryosol horizons (anti-correlated to the C/N, Fig. 5a).

3.2.2 Biomarker proxies/indices

The absolute lipid concentration of *n*-alkanes are in the same range but slightly higher compared to the Yedoma profiles. The *n*-alkane average is $1275.7 \mu\text{g g TOC}_{\text{wt}\%}^{-1}$ (median $1260.1 \mu\text{g g TOC}_{\text{wt}\%}^{-1}$), ranging from 599.7 (8.7 m a.s.l.) to $1907.2 \mu\text{g g TOC}_{\text{wt}\%}^{-1}$ (29.5 m a.s.l.). The *n*-fatty acids average is nearly double that found in the Yedoma samples. On average, $4096.1 \mu\text{g g TOC}_{\text{wt}\%}^{-1}$ (median $3805.7 \mu\text{g g TOC}_{\text{wt}\%}^{-1}$) are stored in the thermokarst deposits of Buor Khaya, ranging from 554.5 (uppermost Buo-01 sample) to 11 013.3 (uppermost Buo-03 sample) $\mu\text{g g TOC}_{\text{wt}\%}^{-1}$.

A series of long-chain *n*-alkanes were recognized in all thermokarst samples with a strong odd carbon number preference ranging from *n*- C_{21} to *n*- C_{33} . Nearly all samples show a unimodal distribution of *n*-alkanes maximized at C_{27} , C_{29} , or C_{31} (Fig. S2). Sample Buo-03-A-03 alone does not fit into this scheme because it maximizes at *n*- C_{25} . Compared to Yedoma samples, the short-chain fraction $< n\text{-C}_{27}$ is more pronounced (Fig. S2). The *n*-fatty acids show strong even-carbon-number preference and a bimodal distribution between *n*- C_{14} and *n*- C_{30} (Fig. S3), but the *n*- C_{16} is less pronounced than in the Yedoma deposits. An exception to this is found in sample Buo-01-A-02, where

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the C₁₆ monomer reaches the overall maximum of the distribution. Apart from that, the maxima are generally located at the C₂₄ *n*-fatty acid.

The *n*-alkane CPI of thermokarst averages 9.6 (median 9.3) and is lower compared to the Yedoma deposits, although the CPI values are in the same range (between 7.0 and 15.3). The CPI of the fatty acids ranges from 4.0 to 9.0 (mean 5.3, median 4.9). The ACL of *n*-alkanes and fatty acids reveal a homogeneous signal between 27.2 and 29.2 (mean 28.3) for *n*-alkanes and 23.6 to 25.6 (mean 24.8) for *n*-fatty acids.

Except for the maximum value of 16.1 µg g TOC_{wt%}⁻¹ at 8.7 ma.s.l., the hop-17(21)-ene concentration at Buo-05 is quite stable, between 0.1 and 4.9 µg g TOC_{wt%}⁻¹. Buo-01 paleocryosol values are 0.9 (8.7 ma.s.l.) and 8.4 at the lowermost sample (7.8 ma.s.l.). For Buo-03 the hop-17(21)-ene concentration ranges from 5 µg g TOC_{wt%}⁻¹ up to 8 µg g TOC_{wt%}⁻¹.

The HPFA ratio for the Buo-05 thermokarst samples is high, between 0.6 and 0.8; only the uppermost sample (9.3 ma.s.l.) shows a significantly lower value of 0.2. The Buo-01 profile decreases from 0.7 at the lowest sample to 0.2 at the top. Buo-03 shows high parameter values of 0.8 and 0.9. The Oleanen ratio for the thermokarst deposits ranges between 0 (Buo-01) and 13.8 (Buo-03). The overall mean Oleanen ratio in thermokarst is 3.7 (median 2.2), which is remarkably higher compared to the Yedoma deposits.

The acetate content of the thermokarst samples is between 0.4 and 109.4 mg L⁻¹ with a mean of 23.5 mg L⁻¹ (median 2.8 mg L⁻¹). Large acetate contents are found especially in the middle part of Buo-05, from 3.4 (74.1 mg L⁻¹) to 6.1 ma.s.l. (109.4 mg L⁻¹), and in the uppermost Buo-03 sample (35.3 mg L⁻¹).

3.3 Principal component analyses

The first PCA diagram (Fig. 6a) shows that thermokarst sediments, especially at Buo-05, could not be separated from Yedoma deposits. This diagram, including the first two principal components, explains 79 % (pc1 57 %, pc2 22 %) of the total data set vari-

BGD

11, 15945–15989, 2014

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



ance. The second PCA diagram (Fig. 6b) illustrates that biomarker quality estimators in Yedoma samples have slightly lower variability because they cluster in an area at $pc1$ and $pc2 > 0$, while the thermokarst samples do not cluster. In this diagram 53 % of the data set variance is explained. Moreover, this PCA shows that there is good consistency between the CPI_{alkane} quality estimator and the C/N ratio (Fig. 6b). The PCA of the n -alkane chain length (Fig. 6c) shows that the best separating variables for thermokarst are the shorter-chain n -alkanes (C_{17} , C_{19} , and C_{21}), contrary to C_{29} for the majority of the Yedoma samples. The $pc1$ explains 39 % and $pc2$ explains 29 % (total 68 %) of the data set variance.

4 Discussion

The Buor Khaya Peninsula is a typical Yedoma hill – thermokarst basin Yedoma region landscape (Strauss et al., 2013). The Yedoma deposits cover ~ 15 % of the peninsula (Günther et al., 2013), which is less than the Yedoma region mean of 30 %, but inside the overall range of Yedoma deposit coverage (Grosse et al., 2013; Strauss et al., 2013). Thus, the current study of Yedoma and thermokarst deposits is representative for an area covered by similar permafrost deposits of late Pleistocene and Holocene age.

4.1 Sediment facies

The grain-size distribution curves (Figs. 2 and S1) indicate a constant deposition environment for the Yedoma sequences. According to Strauss et al. (2012), there have been stable deposition conditions during Yedoma accumulation; this hypothesis is supported by the data presented here. The three thermokarst profiles include three different kinds of thermokarst deposits. Buo-05 is dominated by a lake facies containing valves of two freshwater ostracod taxa: *Cytherissa lacustris* and *Cypria* sp. Moreover, shells have been found in Buo-05 (Strauss and Schirrmmeister, 2011). An ice wedge

BGD

11, 15945–15989, 2014

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



is located next to Buo-01, which points to sub-aerial conditions like a polygon mire. Buo-03 is interpreted as initial thermokarst on top of a Yedoma hill. Thus, the grain-size distributions of Buo-05 and Buo-01 reveal that the thermokarst is granulometrically composed of the same material as Yedoma. The grain size distributions in Buo-03 paint a different picture. This distribution is likely caused by the early state of thermokarst development dominated by peat aggradation. This peat can act like a selective sediment trap influencing the grain-size distributions, e.g. by producing a less distinct coarse silt-fine sand peak.

4.2 Organic matter degradation

The organic matter proxies of Yedoma deposits are less variable than those of thermokarst deposits (Buo-01 and 03). Except for the paleocryosols, the cryopedolith parts of the Yedoma and the Buo-05 thermokarst profile reveal a rather homogenous picture (Figs. 3 and 4). Constant grain-size distributions, less $\text{TOC}_{\text{wt}\%}$, and smaller absolute lipid concentration scattering reveal that the OC stored in the Yedoma deposits has likely been kept perennially frozen since incorporation. The organic matter signatures (Figs. 4, S2, and S3) as well as the grain-size distributions (Figs. 2 and S1) of thermokarst deposits, especially in Buo-01 and Buo-03, show broader variations. This is caused by a more complex degradation and re-deposition history due to reworking. The degradation markers of organic matter found in the paleocryosol parts of all profiles reveal a less-degraded state, indicating that the organic matter in these portions is the best preserved.

The mean $\text{TOC}_{\text{wt}\%}$ content for Yedoma deposits is comparable to other sites (Fig. S5) in the Yedoma region (Schirrmeister et al., 2011b, 2013). Intense accumulation and frozen preservation of plant remains (14 kgCm^{-3} for Yedoma and 24 kgCm^{-3} for thermokarst deposits) is caused by syngenetic permafrost formation in polygonal tundra landscapes over long periods in the Quaternary (Schirrmeister et al., 2013). But comparing the studied deposits to the overall Yedoma region mean (19 kgCm^{-3} for Yedoma deposits and 33 kgCm^{-3} (disregarding wedge-ice content) for thermokarst

deposits, Strauss et al., 2013) on Buor Khaya Peninsula reveals that both deposit types contain less OC. Nevertheless, these numbers show that these deposits comprise a large pool of dormant carbon, which could be reactivated due to permafrost thawing. Moreover, thermokarst deposits seem to be the quantitatively more important OC pool (Yedoma : thermokarst carbon ratio $\sim 2 : 3$). The higher carbon inventory in thermokarst deposits is partially related to a concentration effect for reworked Yedoma OC due to thaw subsidence progression including ground ice loss plus input of Holocene OC. Together with ecosystem recovery, thermokarst basins can act as a local sink for portions of the carbon released from thawing permafrost deposits (van Huissteden and Dolman, 2012). Nevertheless, at the same time thermokarst lakes also promote intense organic matter degradation including methane production in the anaerobic environments of organic-rich lake sediments and unfrozen deposits (Walter et al., 2007b). To answer this question arisen in the introduction, if the thermokarst organic matter pool is as degradable as the frozen late Pleistocene Yedoma (or has the most labile carbon already been emitted due to thermokarst degradation processes?), we visualized the stratigraphically differentiated main proxies in Fig. 7.

In our study the C/N does not reveal a clear picture. The average values are relatively close together for all profiles (Fig. 7b). Nevertheless, the medians and means hint at a lower degradation state/better organic matter quality of thermokarst deposits (especially Buo-03 and Buo-03). Moreover, in both Yedoma and thermokarst deposits the same pattern is visible: A positive linear relationship exists between $\text{TOC}_{\text{wt}\%}$ and C/N ratios (Fig. 5b). In soil science literature it is agreed that the elemental composition of organic matter is affected by the degree of humification and microbial activities that metabolize the organic matter (Kumada, 1987). Ongoing organic matter decomposition will release stored C to the atmosphere and N to the soil (Weintraub and Schimel, 2005), resulting in a lower C/N ratio for more-degraded deposits (Gundelwein et al., 2007). This was found in (sub-) arctic peat deposits and soils, where the C/N ratio decreases with depth (Kuhry and Vitt, 1996; McKane et al., 1997; Ping et al., 1998). Because a high TN content can promote stabilization of organic matter at late stages

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



compounds of shorter chain length (Höfle et al., 2013), we confirm the interpretation of good organic matter quality in both Yedoma and thermokarst deposits. At first view, the hop-17(21)-ene (Fig. 7e) concentration does not show a distinct quality difference between both kinds of deposits, because the Buo-04 Yedoma profile contains hopene concentrations in the same range as those found in thermokarst deposits. However, if we focus on the median values, the Yedoma deposits again appear to be slightly more strongly degraded than the thermokarst deposits. With the exception of Buo-01, the HPFA index (Fig. 7f) also suggests slightly lower degradation and better organic matter quality in the thermokarst deposit profiles (Buo-05 and Buo-03). Our HPFA index, introduced based on Poynter's (1989) HPA index which was tested in the Arctic environment by Routh et al. (2014), is an appropriate indicator of the relative amount of the labile fatty acids that remain in a sample. The uppermost samples just below the surface at Buo-04, Buo-05, and Buo-01 with lower HPFA values are clearly an exception and suggest the entrainment of higher proportions of material influenced by Holocene degradation. This is likely caused by the fairly recent influence of an active layer or transient layer and warmer permafrost temperatures. The Oleanen ratio shows a separation of Yedoma and thermokarst deposits, but this ratio is dominated by numerous 0.0 measurements in the Yedoma deposits. These results might be caused not only by transformation of β -amyrin to Olean-12-ene (by losing the hydroxyl group) or to Olean-13(18)-ene (by losing the double bond), but also by unknown processes in the Yedoma deposits. Thus, because of sparse data, we interpret this proxy as potentially hinting at slightly better Yedoma organic matter quality.

Summing up Fig. 7, thermokarst organic matter is partly less degraded and of better quality compared to the organic matter sequestered in Yedoma deposits (as revealed by measurements of C/N, $\delta^{13}\text{C}$, hop-17(21)-ene, and the HPFA index). The CPI points in the other direction and describes the Yedoma organic matter as better preserved (Fig. 7). As the interquartile ranges show an overlap for most proxies, we see no significant differences (although better thermokarst organic matter quality is indicated).

BGD

11, 15945–15989, 2014

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The PCA confirms the picture of little difference between the organic matter quality of the Yedoma and the thermokarst samples. Especially Fig. 6a, supported by Fig. 2, reveals that Yedoma and thermokarst are composed of similar sediments. The Buo-05 thermokarst profile is very similar to both Yedoma profiles. The PCA of the degradation proxies (Fig. 6b) also shows no clusters, but exhibits slightly better separation between both kinds of deposits. Figure 6b reveals that the C/N ratio, the $\delta^{13}\text{C}$ ratio, and the CPI are correlated. This is also separately illustrated in Fig. 5a and c. Thus, these proxies seem to confirm each other. The PCA of the *n*-alkane chain length points to a potential dominance of longer chain alkanes in Yedoma and shorter chain alkanes in thermokarst, indicating better quality of Yedoma samples (Höfle et al., 2013). Exceptions are the Buo-05-A-01 and Buo-03-A-03 thermokarst samples which point in the same direction as the *n*-C₃₅ concentration.

The abovementioned overlap of the interquartile range (Fig. 7) and especially the PCA of the biomarkers (Fig. 6b and c) show that the organic matter degradation/decomposition vulnerability is heterogeneous and depends on different decomposition trajectories and differing former decomposition/incorporation histories. This is likely shown in both Yedoma and thermokarst deposits, covering the whole range of degradation proxy values (Fig. 7b, c and e). To elucidate this was one of the benefits of the applied multiproxy approach. With the addition of biomarker data, it is possible to show that the permafrost organic matter degradation is not a linear function of age or sediment facies, but likely a combination of (interrupted) degradation cycles and a cascade of degradation steps. In particular, the reasonably good organic matter quality of thermokarst deposits reveals that the sediment degradation processes do not necessarily degrade the organic matter. Potentially, the loss of labile OC during thermokarst processes was compensated for by high rates of Holocene OC accumulation in e.g. lake sediments. Nutrient release from thawing permafrost could have stimulated lake productivity, whereas decomposition was slow because of low lake temperatures, resulting in cold anoxic lake environments (Boike et al., 2013; Walter Anthony et al.,

describing the Yedoma organic matter as being better preserved. We do not see any conflict between these two determinations, because the interquartile ranges overlap for most proxies. We interpret this to indicate similar quality in both kind of deposits, with perhaps slightly better thermokarst organic matter quality. For a modelling approach, this conclusion could be extrapolated to the Laptev Sea Region as the studied deposits are akin to other Yedoma and thermokarst deposits of the northeast Siberian Arctic (Schirrmeister et al., 2011a).

The fate of mobilized Yedoma deposit OC depends largely on the environmental conditions that exist during the thermokarst processes and in the resulting thermokarst basin. When the conditions are good for organic matter preservation, for example cold (slightly above 0 °C) or anoxic (lake) conditions, and reworked fossil organic matter can rapidly refreeze to permafrost, good-quality organic matter can be maintained and inputs could compensate for losses due to degradation.

In conclusion, we found that a combination of classical sedimentological proxies and biomarker ratios is useful for getting closer to an understanding of the complex history of organic matter delivery and degradation in permafrost, as well as the future fate of organic matter when it is exposed due to active layer deepening and further thermokarst development.

**The Supplement related to this article is available online at
doi:10.5194/bgd-11-15945-2014-supplement.**

Author contributions. J. Strauss, L. Schirrmeister, and S. Wetterich sampled and coordinated all sediment sampling at the Buor Khaya field campaign in 2010. K. Mangelsdorf supported the biomarker analysis and interpretation. J. Strauss carried out the laboratory analyses, except for one profile, which was analyzed by L. Eichhorn. U. Herzschuh designed the statistical analyses. J. Strauss planned and wrote the publication with input from all co-authors.

Acknowledgements. We acknowledge support of this research by the German Federal Ministry of Education and Research (the “System Laptev Sea” and “CarboPerm” projects).

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We also thank the Russian and German partners who were involved in the “Eastern Laptev Sea – Buor Khaya Peninsula 2010” expedition. J. Strauss was supported by a grant by the Studienstiftung des deutschen Volkes (German National Academic Foundation).

5 The service charges for this open access publication have been covered by a Research Centre of the Helmholtz Association.

References

- 10 Andersson, R. A. and Meyers, P. A.: Effect of climate change on delivery and degradation of lipid biomarkers in a holocene peat sequence in the eastern european russian arctic, *Org. Geochem.*, 53, 63–72, doi:10.1016/j.orggeochem.2012.05.002, 2012.
- Andersson, R. A., Kuhry, P., Meyers, P., Zebühr, Y., Crill, P., and Mörrth, M.: Impacts of paleohydrological changes on *n*-alkane biomarker compositions of a holocene peat sequence in the eastern european russian arctic, *Org. Geochem.*, 42, 1065–1075, doi:10.1016/j.orggeochem.2011.06.020, 2011.
- 15 Berg, B.: Litter decomposition and organic matter turnover in northern forest soils, *Forest Ecol. Manag.*, 133, 13–22, doi:10.1016/S0378-1127(99)00294-7, 2000.
- Blott, S. J. and Pye, K.: Gradistat: a grain size distribution and statistics package for the analysis of unconsolidated sediments, *Earth Surf. Proc. Land.*, 26, 1237–1248, doi:10.1002/esp.261, 2001.
- 20 Boike, J., Kattenstroth, B., Abramova, K., Bornemann, N., Chetverova, A., Fedorova, I., Fröb, K., Grigoriev, M., Grüber, M., Kutzbach, L., Langer, M., Minke, M., Muster, S., Piel, K., Pfeiffer, E.-M., Stoof, G., Westermann, S., Wischnewski, K., Wille, C., and Hubberten, H.-W.: Baseline characteristics of climate, permafrost and land cover from a new permafrost observatory in the Lena River Delta, Siberia (1998–2011), *Biogeosciences*, 10, 2105–2128, doi:10.5194/bg-10-2105-2013, 2013.
- 25 Bray, E. E. and Evans, E. D.: Distribution of *n*-paraffins as a clue to recognition of source beds, *Geochim. Cosmochim. Ac.*, 22, 2–15, doi:10.1016/0016-7037(61)90069-2, 1961.
- Ciais, P., Tagliabue, A., Cuntz, M., Bopp, L., Scholze, M., Hoffmann, G., Lourantou, A., Harrison, S. P., Prentice, I. C., Kelley, D. I., Koven, C., and Piao, S. L.: Large inert carbon
- 30

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

pool in the terrestrial biosphere during the last glacial maximum, *Nat. Geosci.*, 5, 74–79, doi:10.1038/ngeo1324, 2012.

Dlugokencky, E. and Tans, P.: Trends in atmospheric carbon dioxide, available at: <http://www.esrl.noaa.gov/gmd/ccgg/trends/global.html> (last access: 23 March 2014), 2014.

Drachev, S. S., Savostin, L. A., Groshev, V. G., and Bruni, I. E.: Structure and geology of the continental shelf of the laptev sea, *Eastern Russian Arctic*, *Tectonophysics*, 298, 357–393, doi:10.1016/S0040-1951(98)00159-0, 1998.

Drozдов, D. S., Rivkin, F. M., Rachold, V., Ananjeva-Malkova, G. V., Ivanova, N. V., Chehina, I. V., Koreisha, M. M., Korostelev, Y. V., and Melnikov, E. S.: Electronic atlas of the russian arctic coastal zone, *Geo-Mar. Lett.*, 25, 81–88, doi:10.1007/s00367-004-0189-7, 2005.

Folk, R. L. and Ward, W. C.: Brazos river bar: a study in the significance of grain size parameters, *J. Sediment. Petrol.*, 27, 3–26, 1957.

Gaglioti, B. V., Mann, D. H., Jones, B. M., Pohlman, J. W., Kunz, M. L., and Wooller, M. J.: Radiocarbon age-offsets in an arctic lake reveal the long-term response of permafrost carbon to climate change, *J. Geophys. Res.-Biogeo.*, 119, 2014JG002688, doi:10.1002/2014jg002688, 2014.

Glombitza, C., Mangelsdorf, K., and Horsfield, B.: Maturation related changes in the distribution of ester bound fatty acids and alcohols in a coal series from the new zealand coal band covering diagenetic to catagenetic coalification levels, *Org. Geochem.*, 40, 1063–1073, doi:10.1016/j.orggeochem.2009.07.008, 2009.

Goslar, T., Czernik, J., and Goslar, E.: Low-energy ¹⁴C AMS in Poznań radiocarbon laboratory, Poland, *Nucl. Instrum. Meth. B*, 223–224, 5–11, doi:10.1016/j.nimb.2004.04.005, 2004.

Grosse, G., Harden, J., Turetsky, M. R., McGuire, A. D., Camill, P., Tarnocai, C., Frolking, S., Schuur, E. A. G., Jorgenson, T., Marchenko, S., Romanovsky, V., Wickland, K. P., French, N., Waldrop, M. P., Bourgeau-Chavez, L., and Striegl, R. G.: Vulnerability of high-latitude soil organic carbon in north america to disturbance, *J. Geophys. Res.*, 116, G00K06, doi:10.1029/2010JG001507, 2011a.

Grosse, G., Romanovsky, V., Jorgenson, T., Anthony, K. W., Brown, J., and Overduin, P. P.: Vulnerability and feedbacks of permafrost to climate change, *EOS T. Am. Geophys. Un.*, 92, 73–74, doi:10.1029/2011eo090001, 2011b.

Grosse, G., Robinson, J. E., Bryant, R., Taylor, M. D., Harper, W., DeMasi, A., Kyker-Snowman, E., Veremeeva, A., Schirrmeister, L., and Harden, J.: Distribution of Late Pleis-

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



tocene Ice-Rich Syngenetic Permafrost of the Yedoma Suite in East and Central Siberia, Russia, US Geological Survey Open File Report, 1078, U.S. Geological Survey, Reston, USA, 37 pp., 2013.

Gubin, S. V. and Veremeeva, A. A.: Parent materials enriched in organic matter in the northeast of Russia, *Eurasian Soil Sci.*, 43, 1238–1243, doi:10.1134/s1064229310110062, 2010.

Gundelwein, A., Muller-Lupp, T., Sommerkorn, M., Haupt, E. T. K., Pfeiffer, E., and Wiechmann, H.: Carbon in tundra soils in the lake Labaz region of Arctic Siberia, *Eur. J. Soil Sci.*, 58, 1164–1174, 2007.

Günther, F., Overduin, P. P., Sandakov, A., Grosse, G., and Grigoriev, M. N.: Thermo-erosion along the yedoma coast of the buor khaya peninsula, laptev sea, east siberia, in: *Proceedings of the Tenth International Conference on Permafrost, Volume 1: International Contributions*, Salekhard, Russia, 25–29 June 2012, 137–142, 2012.

Günther, F., Overduin, P. P., Sandakov, A. V., Grosse, G., and Grigoriev, M. N.: Short- and long-term thermo-erosion of ice-rich permafrost coasts in the Laptev Sea region, *Biogeosciences*, 10, 4297–4318, doi:10.5194/bg-10-4297-2013, 2013.

Heyer, J., Hübner, H., and Maaß, I.: Isotopenfraktionierung des Kohlenstoffs bei der mikrobiellen Methanbildung, *Isot. Environ. Healt. S.*, 12, 202–205, doi:10.1080/10256017608543912, 1976.

Hodgkins, S. B., Tfaily, M. M., McCalley, C. K., Logan, T. A., Crill, P. M., Saleska, S. R., Rich, V. I., and Chanton, J. P.: Changes in peat chemistry associated with permafrost thaw increase greenhouse gas production, *P. Natl. Acad. Sci. USA*, 111, 5819–5824, doi:10.1073/pnas.1314641111, 2014.

Höfle, S., Rethemeyer, J., Mueller, C. W., and John, S.: Organic matter composition and stabilization in a polygonal tundra soil of the Lena Delta, *Biogeosciences*, 10, 3145–3158, doi:10.5194/bg-10-3145-2013, 2013.

Hugelius, G., Routh, J., Kuhry, P., and Crill, P.: Mapping the degree of decomposition and thaw remobilization potential of soil organic matter in discontinuous permafrost terrain, *J. Geophys. Res.-Biogeo.*, 117, G02030, doi:10.1029/2011jg001873, 2012.

Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirmermeister, L., Grosse, G., Michaelson, G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Improved estimates show large circumpolar stocks of permafrost carbon while quantifying substantial uncertainty ranges and identifying

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



remaining data gaps, *Biogeosciences Discuss.*, 11, 4771–4822, doi:10.5194/bgd-11-4771-2014, 2014.

Jenkinson, D. S., Poulton, P. R., and Bryant, C.: The turnover of organic carbon in subsoils. Part 1. Natural and bomb radiocarbon in soil profiles from the rothamsted long-term field experiments, *Eur. J. Soil Sci.*, 59, 391–399, doi:10.1111/j.1365-2389.2008.01025.x, 2008.

Jones, B. M., Grosse, G., Arp, C. D., Jones, M. C., Walter Anthony, K. M., and Romanovsky, V. E.: Modern thermokarst lake dynamics in the continuous permafrost zone, northern Seward peninsula, Alaska, *J. Geophys. Res.*, 116, G00M03, doi:10.1029/2011JG001666, 2011.

Knoblauch, C., Beer, C., Sosnin, A., Wagner, D., and Pfeiffer, E.-M.: Predicting long-term carbon mineralization and trace gas production from thawing permafrost of northeast siberia, *Glob. Change Biol.*, 19, 1160–1172, doi:10.1111/gcb.12116, 2013.

Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.: Permafrost carbon-climate feedbacks accelerate global warming, *P. Natl. Acad. Sci. USA*, 108, 14769–14774, doi:10.1073/pnas.1103910108, 2011.

Kuesel, K. and Drake, H. L.: Effects of environmental parameters on the formation and turnover of acetate by forest soils, *Appl. Environ. Microb.*, 61, 3667–3675, 1995.

Kuhry, P. and Vitt, D. H.: Fossil carbon/nitrogen ratios as a measure of peat decomposition, *Ecology*, 77, 271–275, doi:10.2307/2265676, 1996.

Kuhry, P., Ping, C.-L., Schuur, E. A. G., Tarnocai, C., and Zimov, S.: Report from the international permafrost association: carbon pools in permafrost regions, *Permafrost Periglac.*, 20, 229–234, doi:10.1002/ppp.648, 2009.

Kumada, K.: *Chemistry of soil organic matter*, Developments in Soil Science, Elsevier/Japan Scientific Societies Press, Amsterdam, 1987.

Lee, H., Schuur, E. A. G., Inglett, K. S., Lavoie, M., and Chanton, J. P.: The rate of permafrost carbon release under aerobic and anaerobic conditions and its potential effects on climate, *Glob. Change Biol.*, 18, 515–527, doi:10.1111/j.1365-2486.2011.02519.x, 2012.

Li, J., Luo, Y., Natali, S., Schuur, E. A. G., Xia, J., Kowalczyk, E., and Wang, Y.: Modeling permafrost thaw and ecosystem carbon cycle under annual and seasonal warming at an arctic tundra site in alaska, *J. Geophys. Res.-Biogeo.*, 119, 2013JG002569, doi:10.1002/2013jg002569, 2014.

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Luo, Q., Yu, S., Liu, Y., Zhang, Y., Han, H., Qi, L., and Zhong, N.: Existence and implications of hop-17(21)-enes in the lower cretaceous of the Saihantala sag, Erlian basin, China, *Pet. Sci.*, 9, 154–160, doi:10.1007/s12182-012-0195-8, 2012.
- Mangelsdorf, K., Finsel, E., Liebner, S., and Wagner, D.: Temperature adaptation of microbial communities in different horizons of siberian permafrost-affected soils from the lena delta, *Chem. Erde-Geochem.*, 69, 169–182, doi:10.1016/j.chemer.2009.02.001, 2009.
- Mann, P. J., Sobczak, W. V., LaRue, M. M., Bulygina, E., Davydova, A., Vonk, J. E., Schade, J., Davydov, S., Zimov, N., Holmes, R. M., and Spencer, R. G. M.: Evidence for key enzymatic controls on metabolism of arctic river organic matter, *Glob. Change Biol.*, 20, 1089–1100, doi:10.1111/gcb.12416, 2014.
- Marzi, R., Torkelson, B. E., and Olson, R. K.: A revised carbon preference index, *Org. Geochem.*, 20, 1303–1306, doi:10.1016/0146-6380(93)90016-5, 1993.
- McKane, R. B., Rastetter, E. B., Shaver, G. R., Nadelhoffer, K. J., Giblin, A. E., Laundre, J. A., and Chapin, F. S.: Climatic effects on tundra carbon storage inferred from experimental data and a model, *Ecology*, 78, 1170–1187, doi:10.1890/0012-9658(1997)078[1170:ceotcs]2.0.co;2, 1997.
- Mu, C., Zhang, T., Schuster, P. F., Schaefer, K., Wickland, K. P., Repert, D. A., Liu, L., Schaefer, T., and Cheng, G.: Carbon and geochemical properties of cryosols on the north slope of alaska, *Cold Reg. Sci. Technol.*, 100, 59–67, doi:10.1016/j.coldregions.2014.01.001, 2014.
- Oksanen, J.: *Multivariate Analysis of Ecological Communities in R: Vegan Tutorial*, University of Oulu, Oulu, 43 pp., 2013.
- Ping, C. L., Bockheim, J. G., Kimble, J. M., Michaelson, G. J., and Walker, D. A.: Characteristics of cryogenic soils along a latitudinal transect in arctic alaska, *J. Geophys. Res.-Atmos.*, 103, 28917–28928, doi:10.1029/98jd02024, 1998.
- Poynter, J.: *Molecular Stratigraphy: the Recognition of Palaeoclimatic Signals in Organic Geochemical Data*, Ph.D. thesis, School of Chemistry, University of Bristol, Bristol, 324 pp., 1989.
- Poynter, J. and Eglinton, G.: 14. Molecular composition of three sediments from hole 717c: the Bengal fan, *Proceedings of the Ocean Drilling Program: Scientific results*, 116, 155–161, 1990.
- Radke, M., Willsch, H., and Welte, D. H.: Preparative hydrocarbon group type determination by automated medium pressure liquid chromatography, *Anal. Chem.*, 52, 406–411, doi:10.1021/ac50053a009, 1980.

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Romanovskii, N. N., Hubberten, H. W., Gavrillov, A. V., Tumskoy, V. E., and Kholodov, A. L.: Permafrost of the east Siberian Arctic shelf and coastal lowlands, *Quaternary Sci. Rev.*, 23, 1359–1369, doi:10.1016/j.quascirev.2003.12.014, 2004.
- Romanovsky, V. E., Smith, S. L., and Christiansen, H. H.: Permafrost thermal state in the polar Northern Hemisphere during the international polar year 2007–2009: a synthesis, *Permafrost Periglac.*, 21, 106–116, doi:10.1002/ppp.689, 2010.
- Routh, J., Hugelius, G., Kuhry, P., Filley, T., Tillman, P. K., Becher, M., and Crill, P.: Multi-proxy study of soil organic matter dynamics in permafrost peat deposits reveal vulnerability to climate change in the european russian arctic, *Chem. Geol.*, 368, 104–117, doi:10.1016/j.chemgeo.2013.12.022, 2014.
- Rumpel, C. and Kögel-Knabner, I.: Deep soil organic matter – a key but poorly understood component of terrestrial C cycle, *Plant Soil.*, 338, 143–158, doi:10.1007/s11104-010-0391-5, 2011.
- Schädel, C., Schuur, E. A. G., Bracho, R., Elberling, B., Knoblauch, C., Lee, H., Luo, Y. Q., Shaver, G. R., and Turetsky, M. R.: Circumpolar assessment of permafrost C quality and its vulnerability over time using long-term incubation data, *Glob. Change Biol.*, 20, 641–652, doi:10.1111/gcb.12417, 2014.
- Schaphoff, S., Heyder, U., Ostberg, S., Gerten, D., Heinke, J., and Lucht, W.: Contribution of permafrost soils to the global carbon budget, *Environ. Res. Lett.*, 8, 014026, doi:10.1088/1748-9326/8/1/014026, 2013.
- Schirrmester, L., Grosse, G., Wetterich, S., Overduin, P. P., Strauss, J., Schuur, E. A. G., and Hubberten, H.-W.: Fossil organic matter characteristics in permafrost deposits of the north-east siberian arctic, *J. Geophys. Res.*, 116, G00M02, doi:10.1029/2011jg001647, 2011a.
- Schirrmester, L., Kunitsky, V., Grosse, G., Wetterich, S., Meyer, H., Schwamborn, G., Babiy, O., Derevyagin, A., and Siegert, C.: Sedimentary characteristics and origin of the late pleistocene ice complex on north-east siberian arctic coastal lowlands and islands – a review, *Quatern. Int.*, 241, 3–25, doi:10.1016/j.quaint.2010.04.004, 2011b.
- Schirrmester, L., Froese, D. G., Tumskoy, V., Grosse, G., and Wetterich, S.: Yedoma: Late pleistocene ice-rich syngenetic permafrost of beringia, in: *Encyclopedia of Quaternary Sciences, Quaternary Glaciation | Cold Regions Landforms*, 2nd edn., edited by: Elias, S. A., Elsevier, Amsterdam, 2013.

Organic matter quality of deep permafrost carbon

J. Strauss et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)



[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



- Schneider von Deimling, T., Meinshausen, M., Levermann, A., Huber, V., Frieler, K., Lawrence, D. M., and Brovkin, V.: Estimating the near-surface permafrost-carbon feedback on global warming, *Biogeosciences*, 9, 649–665, doi:10.5194/bg-9-649-2012, 2012.
- Schulte, S., Mangelsdorf, K., and Rullkötter, J.: Organic matter preservation on the pakistan continental margin as revealed by biomarker geochemistry, *Org. Geochem.*, 31, 1005–1022, doi:10.1016/S0146-6380(00)00108-X, 2000.
- Schuur, E. A. G. and Abbott, B.: High risk of permafrost thaw, *Nature*, 480, 32–33, doi:10.1038/480032a, 2011.
- Schuur, E. A. G., Vogel, J. G., Crummer, K. G., Lee, H., Sickman, J. O., and Osterkamp, T. E.: The effect of permafrost thaw on old carbon release and net carbon exchange from tundra, *Nature*, 459, 556–559, doi:10.1038/nature08031, 2009.
- Smith, M. R. and Mah, R. A.: Acetate as sole carbon and energy source for growth of methanosarcina strain 227, *Appl. Environ. Microb.*, 39, 993–999, 1980.
- Sollins, P., Spycher, G., and Glassman, C. A.: Net nitrogen mineralization from light- and heavy-fraction forest soil organic matter, *Soil Biol. Biochem.*, 16, 31–37, doi:10.1016/0038-0717(84)90122-6, 1984.
- Stevenson, F. J.: *Humus Chemistry: Genesis, Composition, Reactions*, John Wiley & Sons, New York, 1994.
- Strauss, J. and Schirrmeister, L.: Permafrost sequences of buor khaya peninsula, in: *Reports on Polar and Marine Research – Russian–German Cooperation System Laptev Sea: the Expedition Eastern Laptev Sea-Buor Khaya Peninsula 2010*, edited by: Wetterich, S., Overduin, P. P., and Grigoriev, M., Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany, 35–50, 2011.
- Strauss, J., Schirrmeister, L., Wetterich, S., Borchers, A., and Davydov, S. P.: Grain-size properties and organic-carbon stock of yedoma ice complex permafrost from the Kolyma lowland, northeastern Siberia, *Global Biogeochem. Cy.*, 26, GB3003, doi:10.1029/2011GB004104, 2012.
- Strauss, J., Schirrmeister, L., Grosse, G., Wetterich, S., Ulrich, M., Herzsich, U., and Hubberten, H.-W.: The deep permafrost carbon pool of the yedoma region in siberia and alaska, *Geophys. Res. Lett.*, 40, 2013GL058088, doi:10.1002/2013GL058088, 2013.
- Stuiver, M., Reimer, P., and Reimer, R.: *Calib 6.0*, ¹⁴C CHRONO Centre, Queens University Belfast, Belfast, 2010.

Organic matter quality of deep permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Tieszen, L.: Photosynthesis and respiration in arctic tundra grasses: field light intensity and temperature responses, *Arctic Alpine Res.*, 5, 239–251, 1973.
- Ulrich, M., Grosse, G., Strauss, J., and Schirrmeyer, L.: Quantifying wedge-ice volumes in yedoma and thermokarst basin deposits, *Permafrost Periglac.*, 25, 151–161, doi:10.1002/ppp.1810, 2014.
- van Huissteden, J. and Dolman, A. J.: Soil carbon in the arctic and the permafrost carbon feedback, *Current Opinion in Environmental Sustainability*, 4, 545–551, doi:10.1016/j.cosust.2012.09.008, 2012.
- Vieth, A., Mangelsdorf, K., Sykes, R., and Horsfield, B.: Water extraction of coals – potential for estimating low molecular weight organic acids as carbon feedstock for the deep terrestrial biosphere, *Org. Geochem.*, 39, 985–991, doi:10.1016/j.orggeochem.2008.02.012, 2008.
- Vonk, J. E., Sánchez-García, L., Semiletov, I., Dudarev, O., Eglinton, T., Andersson, A., and Gustafsson, Ö.: Molecular and radiocarbon constraints on sources and degradation of terrestrial organic carbon along the Kolyma paleoriver transect, East Siberian Sea, *Biogeosciences*, 7, 3153–3166, doi:10.5194/bg-7-3153-2010, 2010.
- Vonk, J. E., Mann, P. J., Davydov, S., Davydova, A., Spencer, R. G. M., Schade, J., Sobczak, W. V., Zimov, N., Zimov, S., Bulygina, E., Eglinton, T. I., and Holmes, R. M.: High biolability of ancient permafrost carbon upon thaw, *Geophys. Res. Lett.*, 40, 2689–2693, doi:10.1002/grl.50348, 2013.
- Waldrop, M. P., Wickland, K. P., White III, R., Berhe, A. A., Harden, J. W., and Romanovsky, V. E.: Molecular investigations into a globally important carbon pool: permafrost-protected carbon in alaskan soils, *Glob. Change Biol.*, 16, 2543–2554, doi:10.1111/j.1365-2486.2009.02141.x, 2010.
- Walter Anthony, K. M., Zimov, S. A., Grosse, G., Jones, M. C., Anthony, P. M., Chapin III, F. S., Finlay, J. C., Mack, M. C., Davydov, S., Frenzel, P., and Frolking, S.: A shift of thermokarst lakes from carbon sources to sinks during the holocene epoch, *Nature*, 511, 452–456, doi:10.1038/nature13560, 2014.
- Walter, K. M., Edwards, M. E., Grosse, G., Zimov, S. A., and Chapin, F. S.: Thermokarst lakes as a source of atmospheric ch₄ during the last deglaciation, *Science*, 318, 633–636, 2007a.
- Walter, K. M., Smith, L. C., and Chapin, S. F.: Methane bubbling from northern lakes: present and future contributions to the global methane budget, *Philos. T. R. Soc. A*, 365, 1657–1676, doi:10.1098/rsta.2007.2036, 2007b.

Weintraub, M. N. and Schimel, J. P.: Nitrogen cycling and the spread of shrubs control changes in the carbon balance of arctic tundra ecosystems, *BioScience*, 55, 408–415, doi:10.1641/0006-3568(2005)055[0408:ncatso]2.0.co;2, 2005.

- 5 Zanina, O. G., Gubin, S. V., Kuzmina, S. A., Maximovich, S. V., and Lopatina, D. A.: Late-pleistocene (mis 3–2) palaeoenvironments as recorded by sediments, palaeosols, and ground-squirrel nests at duvanny yar, kolyma lowland, northeast siberia, *Quaternary Sci. Rev.*, 30, 2107–2123, doi:10.1016/j.quascirev.2011.01.021, 2011.

BGD

11, 15945–15989, 2014

**Organic matter
quality of deep
permafrost carbon**

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Organic matter
quality of deep
permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 1. Radiocarbon AMS dating on plant macro remains. Calibrations were done by using Calib 6.0 software and the IntCal09 calibration curve (Stuiver et al., 2010). Depth is given in meter below surface level (m b.s.l.) and height in meter above sea level (m a.s.l.). Age is given as year before present (a BP). Poz: Poznań Radiocarbon Laboratory, Poland.

| Lab. no. | Sample name | Depth [m b.s.l.] | Height [m a.s.l.] | Radiocarbon ages [a BP] | ± | Calibrated ages 2σ 95.4 % [a BP] | ± | |
|-----------|-------------|------------------|-------------------|-------------------------|------|----------------------------------|------|----------------------|
| Poz-42080 | Buo-03-A-03 | 1.3 | 28.7 | 4760 | 40 | 5519 | 70 | thermokarst deposits |
| Poz-42072 | Buo-01-A-02 | 0.7 | 8.7 | 3665 | 35 | 3990 | 100 | |
| Poz-42073 | Buo-01-A-04 | 1.8 | 7.6 | 8140 | 50 | 9075 | 78 | |
| Poz-42086 | Buo-05-A-04 | 0.8 | 8.7 | 5990 | 40 | 6837 | 103 | |
| Poz-42087 | Buo-05-B-10 | 3.4 | 6.1 | 8000 | 80 | 8817 | 215 | |
| Poz-42088 | Buo-05-B-19 | 6.1 | 3.4 | 7940 | 50 | 8811 | 122 | |
| Poz-42090 | Buo-05-C-23 | 7.3 | 2.2 | 5280 | 35 | 6059 | 74 | |
| Poz-42091 | Buo-05-C-29 | 9.2 | 0.3 | 6710 | 90 | 7566 | 138 | |
| Poz-42074 | Buo-02-A-03 | 0.7 | 29.3 | 30 100 | 300 | 34 613 | 596 | |
| Poz-42075 | Buo-02-B-09 | 3.5 | 26.5 | 34 650 | 550 | 39 813 | 1242 | |
| Poz-42076 | Buo-02-B-12 | 5 | 25 | 41 500 | 1500 | 45 312 | 2649 | |
| Poz-42077 | Buo-02-D-20 | 5.5 | 24.5 | 45 000 | 2000 | 47 614 | 2386 | |
| Poz-42078 | Buo-02-D-23 | 7 | 23 | 43 000 | 1500 | 46 830 | 2678 | |
| Poz-42081 | Buo-04-A-02 | 1.5 | 17.1 | 49 000 | 3000 | | | |
| Poz-42082 | Buo-04-A-08 | 5 | 13.6 | > 48 000 | | | | |
| Poz-42083 | Buo-04-B-10 | 8.5 | 9.1 | > 55 000 | | | | |
| Poz-42084 | Buo-04-C-16 | 10.5 | 8 | > 49 000 | | | | |
| Poz-42085 | Buo-04-C-20 | 11.7 | 6.8 | > 55 000 | | | | |

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.




Figure 1. Location of the Buor Khaya Peninsula and the study area. The square black box in the globe inset indicates the area shown in the map below. The profile diagram and the photographs below it show the profiles and their positions relative to each other. Modified after Strauss and Schirrmeister (2011), pictures taken by J. Strauss.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Organic matter quality of deep permafrost carbon

J. Strauss et al.




Figure 2. Three-dimensional grain-size distributions of (a) Yedoma and (b) thermokarst profiles. To avoid an overlap of Buoc-05 and Buoc-01 in (b), the altitude axis was adapted and does not ascend consistently. A two-dimensional grain-size plot is shown in Fig. S1.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.




Figure 3. Summary of sedimentological, biogeochemical, and biomarker parameters for the Buo-04 and Buo-02 Yedoma profiles. All diagrams are drawn in such a way as to show more degraded samples on the left and less degraded samples on the right side. Thus, the axis of $\delta^{13}\text{C}$ values and the Oleanen ratio are descending. In the text, the paleocryosol parts are reported with altitude measurements from the lowest to the highest sample of each paleocryosol. The grey shaded areas are for visualization, not for exact height estimations of the paleocryosols.

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




Figure 4. Summary of sedimentological, biogeochemical, and biomarker parameters for the Buo-05, Buo-01, and Buo-03 thermokarst profiles. The grain-size colors and the *n*-alkane and *n*-fatty acid symbols are explained in Fig. 3. All diagrams are drawn in such a way as to show more degraded samples on the left and less degraded samples on the right side (descending axis of $\delta^{13}\text{C}$ values and Oleanen ratio). In the text, the paleocryosol parts are reported with altitude measurements from the lowest to the highest sample of each paleocryosol. The grey shaded areas are for visualization, not for exact height estimations of the paleocryosols.

Organic matter
quality of deep
permafrost carbon

J. Strauss et al.




Figure 5. Scatter plots of selected degradation markers. Yedoma deposits are shown as black triangles, thermokarst deposits as grey diamonds. Regression equations, the r^2 , and the sample number (n) are inserted as texts.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Organic matter
quality of deep
permafrost carbon

J. Strauss et al.




Figure 7. Conceptual scheme of the organic matter degradation state, estimated using the different applied proxies with boxplots for each studied profile. The boxplots show the studied profiles separately, with Yedoma deposits (white boxes) shown below the thermokarst deposits (grey boxes). The whiskers illustrate the data range, and the box ends indicate the 25th and the 75th quartile (interquartile range). The vertical lines inside each box show the median (= 50th quartile). All diagrams are drawn in such a way as to show more degraded samples on the left and less degraded samples on the right side. Thus, the axis of $\delta^{13}\text{C}$ values is descending.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures



Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

