

Sharp asymptotics for Einstein- λ -dust flows

Helmut Friedrich
Max-Planck-Institut für Gravitationsphysik
Am Mühlenberg 1
14476 Golm, Germany

January 19, 2016

Abstract

We consider the Einstein-dust equations with positive cosmological constant λ on manifolds with time slices diffeomorphic to an orientable, compact 3-manifold S . It is shown that the set of standard Cauchy data for the Einstein- λ -dust equations on S contains an open (in terms of suitable Sobolev norms) subset of data that develop into solutions which admit at future time-like infinity a space-like conformal boundary \mathcal{J}^+ that is C^∞ if the data are of class C^∞ and of correspondingly lower smoothness otherwise. As a particular case follows a strong stability result for FLRW solutions. The solutions can conveniently be characterized in terms of their asymptotic end data induced on \mathcal{J}^+ , only a linear equation must be solved to construct such data. In the case where the energy density $\hat{\rho}$ is everywhere positive such data can be constructed without solving any differential equation at all.

1 Introduction

It has been known for a while that among the solutions to Einstein's vacuum field equations $\hat{R}_{\mu\nu} = \lambda \hat{g}_{\mu\nu}$ with positive cosmological constant λ on manifolds with space-sections diffeomorphic to an orientable, compact 3-manifold S there is an open (in terms of Sobolev norms on Cauchy data) subset of solutions which are future asymptotically simple in the sense of Penrose [17], i.e. the solutions admit the construction of a conformal boundary \mathcal{J}^+ at their infinite time-like future which is C^∞ if the solutions are C^∞ and is of correspondingly lower smoothness otherwise (see [11] for more details and references). This property generalises to the Einstein- λ equations coupled to conformally covariant matter field equations with trace free energy momentum tensor. In [6] this has been discussed in detail for the Maxwell and the Yang-Mills equations, where a procedure has been laid out which applies, possibly with some modifications in specific cases, to other such field equations (see [16] for a recent example).

Matter fields with energy momentum tensors which are not trace free were generally expected to lead to difficulties in the construction of reasonably smooth conformal boundaries. (The emphasis here is on results about the evolution problem, we are not talking about geometric studies near conformal boundaries which postulate properties of energy momentum tensors convenient for their analysis). It has recently been observed, however, that this need not be the true [10].

In the case of the Einstein-Klein-Gordon equations the conformal field equations with suitably transformed matter field imply evolutions system which are hyperbolic, irrespective of the sign of the conformal factor, if the mass and the cosmological constants are related by the equation $m^2 = \frac{2}{3}\lambda$. If this condition is imposed a fairly direct calculation shows that the equation for the rescaled scalar field becomes regular where the conformal factor goes to zero. However, that the conformal equations for the geometric fields become regular in this limit is far from immediate and, as in the case discussed in the following, came as a surprise after various attempts to cast the singular equations into a form which would allow one to draw conclusions about the precise asymptotic behaviour of the solutions in the presence of singularities.

Leaving aside the questions about the significance of this particular result, the present article is concerned with the analysis of another matter model with non-vanishing trace of the energy momentum tensor. We study in detail the future asymptotic behaviour of solutions to the Einstein- λ -dust equations.

In a recent article Hadžić and Speck have shown that the FLRW solutions to the Einstein- λ -dust equations with underlying manifolds of the form $\mathbb{R} \times \mathbb{T}^3$ are future stable, i.e. slightly perturbed FLRW data on \mathbb{T}^3 develop into solutions to the Einstein- λ -dust equations whose causal geodesics are future complete [13]. The authors use the method proposed in [3] to control the evolution of a general wave gauge in terms of its gauge source functions. As emphasized in [3], it is clear that (under fairly weak smoothness assumptions) any coordinate system can in principle be controlled in terms of its gauge source functions and suitable initial data. But finding gauge source functions which are useful in a specific problem is quite a delicate matter. The authors manage to identify gauge source functions which allow them to derive estimates that give control on the long time evolution of their solutions (see [19] for another such case).

It is, however, quite a different question whether the gauge so established lends itself to analyzing the asymptotic behaviour of solutions in detail and to deciding, for instance, whether the differentiable as well as the conformal structure of the solutions admit simultaneously extensions of some smoothness to (future) time-like infinity as required by asymptotic simplicity.

FLRW solutions are known to be future asymptotically simple (see section 6.2). This may be expected to be just an artifact of the high symmetry requirements which imply local conformal flatness and hypersurface orthogonality of the flow field. The present study grew out of attempts to understand what may go wrong under more general assumptions and what kind of obstruction to the asymptotic smoothness of the conformal structure may possibly arise from the presence of a non-vanishing energy density $\hat{\rho}$.

In the article [8] have been derived hyperbolic evolution equations from the Einstein-dust equation in a geometric gauge based of the flow field. The following analysis may be seen as a conformal version of this discussion. After presenting the Einstein- λ -dust equations in section 2, we derive in section 3 the conformal field equations and suitably transformed matter field equations. It turns out that two equations of the system are singular in the sense that there occur factors of the form Ω^{-1} on the right hand side, where Ω is the conformal factor which is positive on the *physical* solution space-time and relates the *physical* metric $\hat{g}_{\mu\nu}$ there to the *conformal* metric $g_{\mu\nu}$ by $g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}$. Since things are to be arranged such that $\Omega \rightarrow 0$ at future time-like infinity, where we want to understand the precise nature of the solutions, there arise problems. One of the singularities, namely the one in the transformed (geodesic) flow field equation, was to be expected. Much more serious is a singularity in the equation for the rescaled conformal Weyl tensor $W^\mu{}_{\nu\lambda\rho} = \Omega^{-1} C^\mu{}_{\nu\lambda\rho}[g]$, which plays a central role in the system. The singularities carry, however, interesting geometric information. They imply that the (so far formally defined) set $\{\Omega = 0\}$ can only define a smooth conformal boundary of the solution space-time if the flow lines approach this set orthogonally. Thus, if one wants to approach the problem in terms of estimates, one has to aim for sufficient control to be able to define simultaneously a conformal boundary at time-like infinity, if admitted by the solution at all, and correspondingly control the behaviour of the flow lines.

In the present article we try to exploit the conformal properties of the system in the most direct way. In section 4 it is shown that due to the specific form of the energy momentum tensor for dust the geodesics tangent to the flow field can be identified after a parameter transformation with curves underlying certain conformal geodesics. Since conformal geodesics are invariants of the conformal structure, this opens the possibility to define a gauge which extends regular across the conformal boundary $\mathcal{J}^+ = \{\Omega = 0\}$ if the latter can indeed be attached in a smooth way to the solution manifold (on which $\Omega > 0$, of course). It turns out that this gauge implies a certain *regularising relation* which proves useful in three different contexts. Its first important merit is to render the conformal field equations regular.

In section 5 it is shown that the conformal field equations imply a *hyperbolic reduced system of evolution equations* which can make sense up to and beyond the conformal boundary at time-like infinity (if it exists). This system is not obtained immediately. The regularizing relation leads to a system which is hyperbolic where $\Omega > 0$ but becomes singular where $\Omega \rightarrow 0$. A further regularization is performed to obtain a system which is

hyperbolic independent of the sign of the conformal factor.

In section 7 is derived a subsidiary system which implies that solutions to the hyperbolic evolution system for data that satisfy the constraints on a given Cauchy hypersurface (with respect to the metric provided by the evolution system) will satisfy in fact the complete system of conformal field equations. This closes the hyperbolic reduction argument.

To obtain complete information on the class of future asymptotically simple solutions to the Einstein- λ -dust solutions we characterize in Lemma 6.1 the possible *asymptotic end data* which may be prescribed on the conformal boundary $\mathcal{J}^+ = \{\Omega = 0\}$ (assumed to be 3-dimensional, orientable, compact) of a solution that admits the construction of such a boundary with sufficient smoothness. As observed already in [4] in the vacuum case, the constraints reduce on \mathcal{J}^+ to a linear system of equations. Remarkably, there is a case where the problem of solving the constraints simplifies even further. In the case where the density $\hat{\rho}$ is positive everywhere certain fields can be prescribed completely freely on \mathcal{J}^+ and the rest follows by algebra and taking derivatives. There is no need to solve any differential equation at all (but see the remarks following Lemma 6.1).

The reduced system of evolution equations is used in section 8 to derive our main results. Being based on hyperbolic equations, a completely detailed statement of the results should give information about Sobolev norms. Since we only use properties of symmetric hyperbolic systems which can be found in the literature at various places and because we are mainly interested in solutions of class C^∞ , we refrain from listing Sobolev indices. We would consider these only be of interest if the weakest possible smoothness assumptions were needed in the context of some concrete problems.

Theorem 1.1 *Let S be a smooth, orientable, compact 3-manifold, assume $\lambda > 0$, and denote by $\mathcal{A}_{\lambda,S}$ the set of standard Cauchy data on S to the Einstein- λ -dust equations with energy density $\hat{\rho} \geq 0$. Then*

- (i) *There is an open (with respect to suitable Sobolev norms) subset $\mathcal{B}_{\lambda,S}$ of data in $\mathcal{A}_{\lambda,S}$ which develop into solutions that admit the construction of conformal boundaries in their infinite time-like future which are of class C^∞ if the data are of class C^∞ and of correspondingly lower differentiability if the data are of lower differentiability.*
- (ii) *The solutions which develop from data in $\mathcal{B}_{\lambda,S}$ are completely parametrized by the asymptotic end data on S (specified in Lemma 6.1) which correspond to the data induced on the future conformal boundaries \mathcal{J}^+ of the solutions.*

The case of the Nariai solution, an explicit, geodesically complete solution to the Einstein- λ -dust equations with $\hat{\rho} = 0$ that admits not even a patch of a smooth conformal boundary (see [11]), shows that our reduced evolution system is by itself not sufficient to ensure the existence of a smooth conformal boundary. Some extra information on the Cauchy data is required.

Because the FLRW solutions do admit a smooth conformal future boundary one could consider data close to FLRW data. Following instead the arguments introduced in [5] and [6], a much larger class of suitable reference solutions (which includes the FLRW solutions) will be constructed in section 8 by solving a backward Cauchy problem for the reduced equations with asymptotic end data that are given on a 3-manifold S which in the end

will represent the future conformal boundary $\mathcal{J}^+ = \{\Omega = 0\}$ of the physical space-time defined on the set $\{\Omega > 0\}$.

In a second step we consider the ‘physical’ standard Cauchy data that are induced by one of these solutions on a ‘physical’ Cauchy hypersurface. It is shown that under sufficiently small perturbations of these data the resulting solutions are *strongly stable* in the sense that the smooth extensibility of their conformal structures at future time-like infinity is preserved. This makes use of the fact that a future asymptotically simple solution admits a conformal representation that extends as a smooth solution to the conformal Einstein- λ -dust equations beyond the conformal boundary into a domain where $\Omega < 0$. The strong stability result follows then as a consequence of the well known Cauchy stability property of hyperbolic equations and the fact that the equation themselves ensure that the set of points where $\Omega = 0$ defines a smooth space-like hypersurface.

Though they lead to the same sets of solutions in the end, it is of interest to distinguish the two different ways of looking at the solutions. In the construction of the reference solutions some features of asymptotic simplicity are *built in from the start* by using asymptotic end data. In the stability result, however, asymptotic simplicity for the perturbed solution is *deduced* as a consequence of the conformal properties of the equations and the reference solution.

In contrast to the approach of [13], which concentrates on deriving suitable estimates, the emphasis is put in this article on the analysis of the field equations and the explicit use of their conformal properties. While the conformal equations may lead to serious difficulties when the conformal structure of the solutions is intrinsically not well behaved at time-like infinity, they give results which are sharp and complete if the conformal structure extends smoothly and only the standard energy estimates for symmetric hyperbolic systems are needed.

Moreover, the information obtained on the equations is in that case of considerable practical interest. The reduced evolution system provides the possibility to calculate numerically - on a finite grid - future complete solutions to Einstein’s field equations, including the details of their asymptotic behaviour. In the Einstein- λ case this has been successfully demonstrated by the work of Beyer (see [2] and the references given there).

Besides the one analysed in [10] this is the second example that illustrates that even in cases in which the energy momentum tensor is not trace free the conformal field equations with $\lambda > 0$ and suitably rescaled matter fields can imply hyperbolic evolution equations that are well defined up to and beyond the future time-like infinity of the physical solutions. The two cases are quite different but the results suggest that the analysis of the asymptotic conformal structure in the presence of matter fields can be more useful than expected.

The possibility to extend solutions to the conformal field equations into a domain in which $\Omega < 0$, where they define another solution to the original equations (see section 8), has been used here only as a technical device in the stability argument leading to Theorem 1.1. Whether it is of any significance in the context of Penrose’s proposal of *conformal cyclic cosmologies* [18] is a question not discussed here.

2 The Einstein- λ -dust system

The Einstein-Euler system with cosmological constant λ consists of the Einstein equations

$$\hat{R}_{\mu\nu} - \frac{1}{2} \hat{R} \hat{g}_{\mu\nu} + \lambda \hat{g}_{\mu\nu} = \kappa \hat{T}_{\mu\nu}, \quad (2.1)$$

for a Lorentz metric $\hat{g}_{\mu\nu}$ on a four-dimensional manifold \hat{M} with an energy momentum tensor of a simple ideal fluid

$$\hat{T}_{\mu\nu} = (\hat{\rho} + \hat{p}) \hat{U}_\mu \hat{U}_\nu + \hat{p} \hat{g}_{\mu\nu}. \quad (2.2)$$

Here \hat{U}^μ is the future directed time-like flow vector field, normalized so that $\hat{U}_\mu \hat{U}^\mu = -1$, and $\hat{\rho}$ and \hat{p} denote the total energy density and the pressure as measured by an observer moving with the fluid. The equations require the relation $\hat{\nabla}^\mu \hat{T}_{\mu\nu} = 0$, which is equivalent to the system consisting of the equations

$$(\hat{\rho} + \hat{p}) \hat{U}^\mu \hat{\nabla}_\mu \hat{U}_\nu + \{\hat{U}_\nu \hat{U}^\mu \hat{\nabla}_\mu + \hat{\nabla}_\nu\} \hat{p} = 0, \quad (2.3)$$

$$\hat{U}^\mu \hat{\nabla}_\mu \hat{\rho} + (\hat{\rho} + \hat{p}) \hat{\nabla}_\mu \hat{U}^\mu = 0. \quad (2.4)$$

These equations must be implemented by an equation of state.

In the following we set $\kappa = 1$, assume $\lambda > 0$, and consider solutions on manifolds diffeomorphic to $\hat{M} = \mathbb{R} \times S$ where S is a compact (without boundary), orientable 3-manifold which specifies the topology of the time slices. We will be interested in the case where $\hat{p} = 0$ throughout, referred to as *pressure free matter* or, shortly, as *dust*. It is supposed that $\hat{\rho}$ does not vanish identically and satisfies

$$\hat{\rho} \geq 0 \quad \text{on} \quad \hat{M}. \quad (2.5)$$

Equation (2.3) reduces then to $\hat{\rho} \hat{U}^\mu \hat{\nabla}_\mu \hat{U}^\nu = 0$. This will be satisfied without condition on \hat{U}^μ on sets where $\hat{\rho} = 0$ and implies that the flow is geodesic where $\hat{\rho} \neq 0$. We require \hat{U}^μ to be geodesic everywhere. The system to be considered consists then of (2.1),

$$\hat{T}_{\mu\nu} = \hat{\rho} \hat{U}_\mu \hat{U}_\nu, \quad (2.6)$$

$$\hat{U}^\mu \hat{\nabla}_\mu \hat{U}^\nu = 0, \quad \hat{U}_\mu \hat{U}^\mu = -1, \quad (2.7)$$

$$\hat{\nabla}_\mu (\hat{\rho} \hat{U}^\mu) = 0. \quad (2.8)$$

Let \hat{S} be a hypersurface in \hat{M} which is space-like for $\hat{g}_{\mu\nu}$ and denote by \hat{n}^μ the future directed normal of \hat{S} normalized by $\hat{n}_\mu \hat{n}^\mu = -1$. Let coordinates x^μ be given near \hat{S} so that $\hat{S} = \{x^0 = 0\}$ and the x^α , $\alpha, \beta = 1, 2, 3$, are local coordinates on \hat{S} . Denote by $\hat{h}_{\alpha\beta}$, $\hat{\kappa}_{\alpha\beta}$ the first and the second fundamental form induced on \hat{S} by $\hat{g}_{\mu\nu}$ and by $\hat{h}_\mu{}^\nu = \hat{g}_\mu{}^\nu + \hat{n}_\mu \hat{n}^\nu$ the orthogonal projector onto the tangent spaces of \hat{S} . Equations (2.7), (2.8) are evolution equations for \hat{U}^μ and $\hat{\rho}$. Equation (2.1) induces with (2.6) on \hat{S} the constraints

$$0 = R[\hat{h}] - \hat{\kappa}_{\alpha\beta} \hat{\kappa}^{\alpha\beta} + (\hat{\kappa}_\alpha{}^\alpha)^2 - 2\lambda - 2\hat{n}^\mu \hat{n}^\nu \hat{T}_{\mu\nu},$$

$$0 = \hat{D}_\beta \hat{\kappa}_\alpha{}^\beta - \hat{D}_\alpha \hat{\kappa}_\beta{}^\beta - \hat{n}^\mu \hat{h}_\alpha{}^\nu \hat{T}_{\mu\nu}.$$

Setting $a = -\hat{n}^\mu \hat{U}_\mu > 0$, $\hat{u}_\mu = \hat{h}_\mu{}^\nu \hat{U}_\nu$, so that

$$\hat{U}_\mu = a \hat{n}_\mu + \hat{u}_\mu \quad \text{with} \quad -1 = -a^2 + \hat{u}_\beta \hat{u}^\beta \quad \text{where} \quad \hat{u}_\beta \hat{u}^\beta = \hat{h}^{\beta\gamma} \hat{u}_\beta \hat{u}_\gamma,$$

the constraints take the form

$$0 = R[\hat{h}] - \hat{\kappa}_{\alpha\beta} \hat{\kappa}^{\alpha\beta} + (\hat{\kappa}_\alpha{}^\alpha)^2 - 2\lambda - 2\hat{\rho}(1 + \hat{u}_\alpha \hat{u}^\alpha), \quad (2.9)$$

$$0 = \hat{D}_\beta \hat{\kappa}_\alpha{}^\beta - \hat{D}_\alpha \hat{\kappa}_\beta{}^\beta + \hat{\rho} \sqrt{1 + \hat{u}_\beta \hat{u}^\beta} \hat{u}_\alpha. \quad (2.10)$$

It has been shown in [8] how to derive from equations (2.1), (2.6), (2.7), (2.8) a symmetric hyperbolic evolution system of equations for all unknowns in a gauge based on the flow vector field \hat{U} . Given $\lambda > 0$ and a sufficiently smooth initial data set

$$(\hat{S}, \hat{h}_{\alpha\beta}, \hat{\kappa}_{\alpha\beta}, \hat{u}^\alpha, \hat{\rho}), \quad (2.11)$$

satisfying (2.9), (2.10) with $\hat{h}_{\alpha\beta}$ a Riemannian metric and $\hat{\rho} \geq 0$, the evolution system can be used to construct a globally hyperbolic solution $(\hat{M}, \hat{g}_{\mu\nu}, \hat{U}^\mu, \hat{\rho})$ to the Einstein-dust equations with cosmological constant λ into which the initial data set is isometrically embedded so that \hat{S} represents after an identification a space-like Cauchy hypersurface for $(\hat{M}, \hat{g}_{\mu\nu})$. The manifold \hat{M} will then be ruled by the geodesics tangent to \hat{U}^μ . The ODE's

$$\hat{U}^\mu \hat{\nabla}_\mu \hat{\rho} + \hat{\rho} \hat{\nabla}_\mu \hat{U}^\mu = 0,$$

along the geodesics tangent to \hat{U}^μ ensure that $\hat{\rho} > 0$ or $= 0$ along a given geodesic, depending on whether this relation is satisfied at the point where the geodesic intersects \hat{S} . Thus $\hat{\rho} \geq 0$ will hold on \hat{M} .

For smooth initial data the evolution system given in [8] provides a smooth solution in coordinates $x^0 = t$, x^a so that $\langle dx^a, \hat{U} \rangle = 0$, $\langle dt, \hat{U} \rangle = 1$, whence $\hat{U} = \partial_t$. The initial hypersurface is given by $\hat{S} = \{t = t_*\}$ for some fixed value t_* , the metric is of the form

$$\hat{g} = -(a dt)^2 + h_{\alpha\beta} (\hat{u}^\alpha dt + dx^\alpha) (\hat{u}^\beta dt + dx^\beta) \quad \text{on} \quad \hat{M}, \quad (2.12)$$

the future directed \hat{g} -unit normal to \hat{S} is given by

$$\hat{n}^\mu = \frac{1}{a} (\delta^\mu - \hat{u}^\mu) \quad \text{with shift vector field } \hat{u}^\mu \text{ so that } \hat{u}^0 = 0, \quad (2.13)$$

and the lapse function a satisfies $-1 = \hat{g}(\hat{U}, \hat{U}) = -a^2 + h_{\alpha\beta} \hat{u}^\alpha \hat{u}^\beta$. If \hat{U} is hypersurface orthogonal we can assume that $a = 1$, $\hat{u}^\alpha = 0$ and the coordinates define a Gauss system. This will not necessarily be assumed in this article.

The questions to be analyzed in the following asks whether there exist a reasonably large set of data for which the solutions can be extended to become future complete, so that t takes values in $[t_*, \infty[$, and whether these solutions allow us to give a sharp and detailed description of the asymptotic behaviour of the conformal structure in the expanding direction, where $t \rightarrow \infty$.

3 The metric conformal field equations

Let Ω denote a positive *conformal factor* on \hat{M} and $g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}$ the *rescaled metric*. We shall in the following consider the tensor fields

$$\Omega, \quad s = \frac{1}{4} \nabla_\mu \nabla^\mu \Omega + \frac{1}{24} \Omega R[g], \quad L_{\mu\nu} = \frac{1}{2} \left(R_{\mu\nu}[g] - \frac{1}{6} R[g] g_{\mu\nu} \right), \quad (3.1)$$

$$W^\mu{}_{\eta\nu\lambda} = \Omega^{-1} C^\mu{}_{\eta\nu\lambda}[g], \quad (3.2)$$

where ∇_μ denotes the Levi-Civita connection of g and the last two fields denote the Schouten and the rescaled conformal Weyl tensor of $g_{\mu\nu}$ respectively. Moreover, we shall consider the *conformal matter fields*

$$U_\mu = \Omega \hat{U}_\mu, \quad \rho = \Omega^{-3} \hat{\rho}.$$

The vector fields $U^\mu = g^{\mu\nu} U_\nu$ and $\hat{U}^\mu = \hat{g}^{\mu\nu} \hat{U}_\nu$ are then related by

$$U^\mu = \Omega^{-1} \hat{U}^\mu \quad \text{so that} \quad g(U, U) = \hat{g}(\hat{U}, \hat{U}) = -1.$$

The tensor fields above satisfy the system of *conformal field equations* (see [6], [10])

$$6 \Omega s - 3 \nabla_\eta \Omega \nabla^\eta \Omega - \lambda = -\frac{1}{4} \hat{T}, \quad (3.3)$$

$$\nabla_\mu \nabla_\nu \Omega + \Omega L_{\mu\nu} - s g_{\mu\nu} = \frac{1}{2} \Omega T_{\mu\nu}^*, \quad (3.4)$$

$$\nabla_\mu s + \nabla^\eta \Omega L_{\eta\mu} = \frac{1}{2} \nabla^\eta \Omega T_{\eta\mu}^* - \frac{1}{24 \Omega} \nabla_\mu \hat{T}, \quad (3.5)$$

$$\nabla_\nu L_{\lambda\eta} - \nabla_\lambda L_{\nu\eta} - \nabla_\mu \Omega W^\mu{}_{\eta\nu\lambda} = 2 \hat{\nabla}_{[\nu} \hat{L}_{\lambda]\eta}, \quad (3.6)$$

$$\nabla_\mu W^\mu{}_{\eta\nu\lambda} = 2 \Omega^{-1} \hat{\nabla}_{[\nu} \hat{L}_{\lambda]\eta}. \quad (3.7)$$

The right hand sides are determined by the trace

$$\hat{T} = \hat{g}^{\eta\mu} \hat{T}_{\eta\mu} = -\hat{\rho} = -\Omega^3 \rho, \quad (3.8)$$

and the trace free part

$$T_{\eta\mu}^* = \hat{\rho} \left(\hat{U}_\eta \hat{U}_\mu + \frac{1}{4} \hat{g}_{\eta\mu} \right) = \Omega \rho \left(U_\eta U_\mu + \frac{1}{4} g_{\eta\mu} \right), \quad (3.9)$$

of the energy momentum tensor (2.6) and the *physical Schouten tensor* $\hat{L}_{\mu\nu}$, which takes with our energy momentum tensor, the field equations, and the rescaled fields the form

$$\hat{L}_{\mu\nu} = \frac{1}{6} (\hat{\rho} + \lambda) \hat{g}_{\mu\nu} + \frac{1}{2} \hat{\rho} \hat{U}_\mu \hat{U}_\nu = \frac{1}{6} \lambda \hat{g}_{\mu\nu} + \Omega \rho \left(\frac{1}{2} U_\mu U_\nu + \frac{1}{6} g_{\mu\nu} \right). \quad (3.10)$$

Taking into account the transformation law of the connection coefficients under conformal rescaling this gives

$$\begin{aligned}
2 \hat{\nabla}_{[\nu} \hat{L}_{\lambda]\eta} &= \hat{\nabla}_{[\nu} \hat{\rho} \hat{U}_{\lambda]} \hat{U}_\eta + \frac{1}{3} \hat{\nabla}_{[\nu} \hat{\rho} \hat{g}_{\lambda]\eta} + \hat{\rho} (\hat{\nabla}_{[\nu} \hat{U}_{\lambda]} \hat{U}_\eta + \hat{U}_{[\lambda} \hat{\nabla}_{\nu]} \hat{U}_\eta) \\
&= \Omega \left(\rho (\nabla_{[\nu} U_{\lambda]} U_\eta + U_{[\lambda} \nabla_{\nu]} U_\eta) + \nabla_{[\nu} \rho U_{\lambda]} U_\eta + \frac{1}{3} \nabla_{[\nu} \rho g_{\lambda]\eta} \right) \\
&\quad + \rho (\nabla_{[\nu} \Omega g_{\lambda]\eta} + 2 \nabla_{[\nu} \Omega U_{\lambda]} U_\eta + U_{[\nu} g_{\lambda]\eta} g^{\pi\delta} \nabla_\pi \Omega U_\delta).
\end{aligned}$$

Finally, the geodesic equation (2.7) translates into

$$\nabla_U U^\mu = \frac{1}{\Omega} (-g(U, U) g^\mu{}_\rho + U^\mu U_\rho) \nabla^\rho \Omega. \quad (3.11)$$

while equation (2.8) for the density $\hat{\rho}$ gives

$$\nabla_U \rho + \rho \nabla_\mu U^\mu = 0. \quad (3.12)$$

We express the equations in terms of a frame field $e_k = e^\mu{}_k \partial_{x^\mu}$, $k = 0, 1, 2, 3$, which has a time-like vector field given by

$$e_0 = U,$$

and which is orthonormal, so that $g_{jk} \equiv g(e_j, e_k) = \eta_{jk} = \text{diag}(-1, 1, 1, 1)$. The space-like frame fields are given by the e_a , where $a, b, c = 1, 2, 3$ denote spatial indices to which the summation convention applies. The metric is given by

$$g = \eta_{jk} \sigma^j \sigma^k,$$

where σ^j denotes the field of 1-forms dual to e_k so that their coefficients in the coordinates x^μ satisfy $\sigma^j{}_\mu e^\mu{}_k = \delta^j{}_k$.

The connection coefficients, defined by $\nabla_j e_k \equiv \nabla_{e_j} e_k = \Gamma_j{}^l{}_k e_l$, satisfy $\Gamma_{jlk} = -\Gamma_{jkl}$ with $\Gamma_{jlk} = \Gamma_j{}^i{}_k g_{li}$, because $\nabla_i g_{jk} = 0$. The covariant derivative of a tensor field $X^\mu{}_\nu$, given in the frame by $X^i{}_j$, takes the form

$$\nabla_k X^i{}_j = X^i{}_{j,\mu} e^\mu{}_k + \Gamma_k{}^i{}_l X^l{}_j - \Gamma_k{}^i{}_l X^i{}_j.$$

For the covariant version of U , i.e. $U_j = -\delta^0{}_j$, equation (3.11) implies the form

$$\nabla_k U_l = \Gamma_k{}^0{}_l = \delta^0{}_k \Omega^{-1} (\nabla_l \Omega + U_l \nabla_0 \Omega) + \delta^a{}_k \delta^b{}_l \chi_{ab}. \quad (3.13)$$

If U is hypersurface orthogonal and if \hat{S} were chosen to be orthogonal to U so that the vector fields e_a define an orthonormal frame on \hat{S} , the field χ_{ab} would represent the second fundamental form induced by g on the slice \hat{S} whence $\chi_{ab} = \chi_{(ab)}$. In general hypersurface orthogonality will not be assumed here. We shall write $g^{ab} \chi_{ab} = \chi_a{}^a$.

The metric coefficients and the connection coefficients satisfy the *first structural equations*

$$e^\mu{}_{i,\nu} e^\nu{}_j - e^\mu{}_{j,\nu} e^\nu{}_i = (\Gamma_j{}^k{}_i - \Gamma_i{}^k{}_j) e^\mu{}_k, \quad (3.14)$$

which ensures the connection to be torsion free, and the *second structural equations*

$$\begin{aligned} \Gamma_l^i{}_{j,\mu} e^\mu{}_k - \Gamma_k^i{}_{j,\mu} e^\mu{}_l + 2\Gamma_{[k}^i{}_{p]}\Gamma_{l]p}{}^j - 2\Gamma_{[k}^p{}_{l]}\Gamma_p^i{}_{j} & \quad (3.15) \\ & = \Omega W^i{}_{jkl} + 2\{g^i{}_{[k}L_{l]j} + L^i{}_{[k}g_{l]j}\}, \end{aligned}$$

which relates the coefficients (and thus the metric $g_{\mu\nu}$) to the unknowns in the conformal field equations. The conformal field equations read now

$$6\Omega s - 3\nabla_i\Omega\nabla^i\Omega - \lambda = \frac{1}{4}\Omega^3\rho, \quad (3.16)$$

$$\nabla_j\nabla_k\Omega + \Omega L_{jk} - s g_{jk} = \frac{1}{2}\Omega^2\rho\left(U_j U_k + \frac{1}{4}g_{jk}\right), \quad (3.17)$$

$$\nabla_k s + \nabla^i\Omega L_{ik} = \frac{1}{2}\Omega\rho\nabla^i\Omega\left(U_i U_k + \frac{1}{4}g_{ik}\right) + \frac{1}{8}\Omega\rho\nabla_k\Omega + \frac{1}{24}\Omega^2\nabla_k\rho, \quad (3.18)$$

$$\begin{aligned} \nabla_k L_{lj} - \nabla_l L_{kj} - \nabla_i\Omega W^i{}_{jkl} & \quad (3.19) \\ & = \Omega\left(\rho(\nabla_{[k}U_{l]}U_j + U_{[l}\nabla_{k]}U_j) + \nabla_{[k}\rho U_{l]}U_j + \frac{1}{3}\nabla_{[k}\rho g_{l]j}\right) \\ & \quad + \rho(\nabla_{[k}\Omega g_{l]j} + 2\nabla_{[k}\Omega U_{l]}U_j + U_{[k}g_{l]j}g^{pq}\nabla_p\Omega U_q), \end{aligned}$$

$$\begin{aligned} \nabla_i W^i{}_{jkl} & \quad (3.20) \\ & = \rho(\nabla_{[k}U_{l]}U_j + U_{[l}\nabla_{k]}U_j) + \nabla_{[k}\rho U_{l]}U_j + \frac{1}{3}\nabla_{[k}\rho g_{l]j} + \frac{1}{\Omega}\rho Z_{jkl} \end{aligned}$$

with

$$Z_{jkl} = \nabla_{[k}\Omega g_{l]j} + 2\nabla_{[k}\Omega U_{l]}U_j + U_{[k}g_{l]j}g^{pq}\nabla_p\Omega U_q.$$

The matter equations are given by

$$\nabla_U U^k = \frac{1}{\Omega}(g^k{}_i + U^k U_i)\nabla^i\Omega, \quad (3.21)$$

$$\nabla_U \rho + \rho\chi_a{}^a = 0. \quad (3.22)$$

Equations (3.14) to (3.22) establish a system of differential equations for the unknowns

$$e^\mu{}_k, \quad \Gamma_i{}^j{}_k, \quad \Omega, \quad s, \quad L_{jk}, \quad W^i{}_{jkl}, \quad U^k, \quad \rho, \quad (3.23)$$

which is (apart from subtleties which may arise in cases of low differentiability) equivalent to the system (2.1), (2.6), (2.7), (2.8) in domains where $\Omega > 0$.

If the system is to be used to solve Cauchy problems with data given on a space-like hypersurface \hat{S} , one has to restrict the available gauge freedom. We shall follow the procedure of [6] and [10], where the conformal freedom is removed by considering the Ricci

scalar $R = R[g]$ in a suitable neighborhood of \hat{S} as a prescribed function of the space-time coordinates and by prescribing suitable initial data for Ω and $\nabla_i \Omega$ on \hat{S} . The coordinates $\tau = x^0$ and x^a are chosen near \hat{S} so that $\tau = \tau_*$ on \hat{S} and $\langle U, dx^a \rangle = 0$, $\langle U, d\tau \rangle = 1$, whence

$$U^\mu = e^\mu{}_{0} = \delta^\mu{}_{0} \quad \text{near } \hat{S}.$$

Apart from a parameter transformation $t = t(\tau)$ these coordinates coincide with the ones considered in (2.12). Precise conditions on the vector fields e_a orthogonal to U will be stated later.

Our main interest is the question whether there exist solutions to the system above on the domain where $\Omega > 0$ which admit a meaningful (i.e. sufficiently smooth) limit to a boundary where $\Omega \rightarrow 0$. In that case we write $\{\Omega = 0\} = \mathcal{J}^+$, and refer to this set as the future *conformal boundary* of the solution. By equation (3.16) the limit of $\nabla^i \Omega$ will then define a time-like normal to the set \mathcal{J}^+ so that the latter will define a space-like hypersurface. It represents (future) time-like and null infinity for the ‘physical’ space-time on which $\Omega > 0$.

There arises an obvious problem with the differential system above. The right hand sides of equations (3.20) and (3.21) are formally singular where $\Omega \rightarrow 0$. This problem will be analyzed in the next section. Here we just point out its geometric nature.

If the fields entering equation (3.21) have limits as $\Omega \rightarrow 0$ the term in brackets on the right hand side of (3.21) defines a projection operator with kernel generated by the unit vector U . The right hand side of (3.21) can only admit a limit as $\Omega \rightarrow 0$ if the gradient of Ω is in the kernel of that operator and thus proportional to U , whence

The solutions can only admit a reasonably smooth conformal boundary \mathcal{J}^+ if the geodesics generated by \hat{U} approach \mathcal{J}^+ orthogonally.

Remarkably, the singularity of equation (3.20) is of a similar geometric nature. If we want to keep the freedom to have non-vanishing conformal densities ρ on \mathcal{J}^+ , the right hand side of (3.20) can only assume a limit if $Z_{jkl} \rightarrow 0$ at \mathcal{J}^+ . Since this implies that $U^j Z_{jkl} = -\nabla_{[k} \Omega U_{l]} \rightarrow 0$, which implies in turn that $Z_{jkl} \rightarrow 0$, the conclusion above follows again.

4 The regularizing relation

A conformal geodesic in a given space-time (\hat{M}, \hat{g}) is a curve $x^\mu(\sigma)$ together with a 1-form field $b_\nu(\sigma)$ which satisfy the system of *conformal geodesic equations*

$$\hat{\nabla}_V V^\mu + S(b)_{\lambda}{}^{\mu}{}_{\rho} V^\lambda V^\rho = 0,$$

$$\hat{\nabla}_V b_\nu - \frac{1}{2} b_\mu S(b)_{\lambda}{}^{\mu}{}_{\nu} V^\lambda - \hat{L}_{\lambda\nu} V^\lambda = 0,$$

where $S(b)_{\lambda}{}^{\mu}{}_{\rho} = \delta_{\lambda}{}^{\mu} b_{\rho} + \delta_{\rho}{}^{\mu} b_{\lambda} - \hat{g}_{\lambda\rho} \hat{g}^{\mu\nu} b_{\nu}$ and $V^\mu(\sigma) = \frac{dx^\mu}{d\sigma}$ denotes the tangent vector of the curve. Sometimes it will be convenient to write these equations in the form

$$\hat{\nabla}_V V + 2 \langle b, V \rangle V - \hat{g}(V, V) b = 0, \quad (4.1)$$

$$\hat{\nabla}_V b - \langle b, V \rangle b + \frac{1}{2} \hat{g}(b, b) V - \hat{L}(V, \cdot) = 0, \quad (4.2)$$

where the index position should be clear from the above.

For a conformal geodesic the initial data at a given point consist of its tangent vector and its 1-form at that point. On a given space-time there exist thus more conformal geodesics than metric geodesics. Moreover, there exists in general no particular relation between conformal and metric geodesics. The problem of interest here is, however, very special in this respect.

Lemma 4.1 *Let (\hat{M}, \hat{g}) be a solution to the Einstein-dust system (2.1), (2.6), (2.7), (2.8). Then the geodesics tangential to the vector field \hat{U} coincide after a reparameterization with the curves underlying certain conformal geodesics.*

Proof: Suppose $\bar{x}^\mu(t)$ is a \hat{g} -geodesic with $\frac{d\bar{x}^\mu}{dt} = \hat{U}^\mu(\bar{x}(t))$ and $(x^\mu(\sigma), b_\nu(\sigma))$ a conformal geodesics with $V^\mu(\sigma) = \frac{dx^\mu}{d\sigma}$. Then there exists a parameter transformation $t = t(\sigma)$ so that $\frac{dt}{d\sigma} > 0$ and $x^\mu(\sigma) = \bar{x}^\mu(t(\sigma))$ if and only if

$$V^\mu(\sigma) = \omega(\sigma)^{-1} \hat{U}^\mu(\bar{x}(t(\sigma))) \quad \text{with} \quad \omega^{-1} = \frac{dt}{d\sigma} > 0, \quad \hat{g}(V, V) = -\omega^{-2}. \quad (4.3)$$

For $x^\mu(\sigma)$ to be up to a reparameterization a geodesic we need to have a relation

$$b_\mu = \alpha V_\mu, \quad (4.4)$$

with some function $\alpha = \alpha(\sigma)$ so that (4.1) reads

$$\hat{\nabla}_V V^\mu + \alpha \hat{g}(V, V) V^\mu = 0. \quad (4.5)$$

It follows then that $2\omega^{-3} \hat{\nabla}_V \omega = \hat{\nabla}_V (\hat{g}(V, V)) = -2\alpha\omega^{-4}$, whence

$$\alpha = -\omega \hat{\nabla}_V \omega. \quad (4.6)$$

Basic for our result is that relations (3.10) and (4.3) give along $x^\mu(\sigma)$

$$V^\nu \hat{L}_{\nu\mu} = \frac{1}{6} (\lambda - 2\hat{\rho}) V_\mu, \quad \text{with} \quad \hat{\rho} = \hat{\rho}(\bar{x}^\mu(t(\sigma))).$$

Inserting this and (4.4) into (4.2) and observing (4.5), (4.6) gives the equation

$$\omega \frac{d^2\omega}{d\sigma^2} - \frac{1}{2} \left(\frac{d\omega}{d\sigma} \right)^2 + \frac{1}{6} (\lambda - 2\hat{\rho}(\bar{x}^\mu(t(\sigma)))) = 0,$$

which provides with the relation

$$\frac{dt}{d\sigma} = \frac{1}{\omega}, \quad (4.7)$$

a system of ODE's for $\omega = \omega(\sigma)$ and $t = t(\sigma)$ along $x^\mu(\sigma) = \bar{x}^\mu(t(\sigma))$. Prescribing arbitrary initial data $t|_{\sigma_*} = t_*$, $\omega|_{\sigma_*}$, and $\frac{d\omega}{d\sigma}|_{\sigma_*}$ with $\omega_* > 0$ at the point $x^\mu(\sigma_*) = \bar{x}^\mu(t_*)$ it can be solved. A straight forward calculation then shows that

$$V^\mu(\sigma) = \frac{1}{\omega} \hat{U}^\mu(\bar{x}(t(\sigma))), \quad b_\nu(\sigma) = -\frac{d\omega}{d\sigma} \hat{U}^\mu(\bar{x}(t(\sigma))),$$

do indeed satisfy equations (4.1) and (4.2). \square

It will later be important to note that the freedom to prescribe the initial data for ω gives the freedom to prescribe α arbitrarily at a given point.

Conformal geodesics are of interest in the present context because the curves underlying *conformal geodesics are conformal invariants of a given conformal structure*: If $g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}$, where Ω is a conformal factor as considered above and $x^\mu(\sigma)$, $b_\lambda(\sigma)$ satisfy the conformal geodesic equations with respect to $\hat{g}_{\mu\nu}$, then $x^\mu(\sigma)$, $f_\nu(\sigma)$ with

$$f_\nu(\sigma) = b_\nu(\sigma) - \Omega^{-1} \nabla_\nu \Omega|_{x(\sigma)}, \quad (4.8)$$

satisfy the conformal geodesics equations

$$\nabla_V V + 2 \langle f, V \rangle V - g(V, V) f = 0, \quad (4.9)$$

$$\nabla_V f - \langle f, V \rangle f + \frac{1}{2} g(f, f) V - L(V, \cdot) = 0, \quad (4.10)$$

with respect to $g_{\mu\nu}$, where ∇ and L denote the Levi-Civita connection and the Schouten tensor of $g_{\mu\nu}$ (for this and further properties of conformal geodesics we refer to [7], [9]). If $g(V, V) = -\theta^{-2}$ with $\theta > 0$ at a given point, equation (4.9) gives

$$\nabla_V \theta = \theta \langle V, f \rangle,$$

which shows that θ will stay positive and $x^\mu(\sigma)$ will be time-like as long as V and f remain sufficiently smooth. Equations (4.9), (4.10) do not see the relation $g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}$. Thus, if (\hat{M}, \hat{g}) admits a smooth conformal boundary \mathcal{J}^+ , one can arrange time-like conformal geodesics to extend smoothly to \mathcal{J}^+ with finite and non-vanishing tangent vector.

In the following we shall assume V to be a conformal geodesic vector field which is related, as in (4.3), to the \hat{g} -geodesic vector field \hat{U} by

$$V^\mu = \omega^{-1} \hat{U}^\mu. \quad (4.11)$$

With the notation above we have then

$$\theta V^\mu = U^\mu = \Omega^{-1} \hat{U}^\mu,$$

and thus

$$\theta = \frac{\omega}{\Omega}, \quad \nabla_U \theta = \theta \langle U, f \rangle. \quad (4.12)$$

Since θ stays smooth and positive if U crosses the conformal boundary this has the remarkable consequence, used already in [7], that ω goes to zero precisely where Ω does.

In terms of U equation (4.9) takes the form

$$\nabla_U U + \langle U, f \rangle U - g(U, U) f = 0. \quad (4.13)$$

Replacing in (4.10) the field V by $U = \theta V$ renders that equation in the form

$$\nabla_U f - \langle U, f \rangle f + \frac{1}{2} g(f, f) U - L(U, \cdot) = 0. \quad (4.14)$$

This version of the conformal geodesic equations will be assumed from now on. The only effect of the transition is a reparametrization of $x^\mu(\sigma) \rightarrow x^\mu(\tau)$, $f_\nu(\sigma) \rightarrow f_\nu(\tau)$ where σ is replaced by a function $\sigma(\tau)$ so that

$$\frac{d\tau}{d\sigma} = \frac{1}{\theta(x(\sigma))}. \quad (4.15)$$

In the following the parameter τ will be used.

With (4.12) and the relations obtained in the proof of Lemma 4.1 we get

$$\begin{aligned} f_\mu &= b_\mu - \Omega^{-1} \nabla_\mu \Omega = -\omega \nabla_V \omega \hat{g}_{\mu\nu} V^\nu - \Omega^{-1} \nabla_\mu \Omega \\ &= -(\theta \Omega) \theta^{-1} \nabla_U (\theta \Omega) \Omega^{-2} g_{\mu\nu} \theta^{-1} U^\nu - \Omega^{-1} \nabla_\mu \Omega \\ &= -(\theta^{-1} \nabla_U \theta + \Omega^{-1} \nabla_U \Omega) U_\mu - \Omega^{-1} \nabla_\mu \Omega, \\ &= -(\langle U, f \rangle + \Omega^{-1} \nabla_U \Omega) U_\mu - \Omega^{-1} \nabla_\mu \Omega, \end{aligned}$$

and thus the *regularising relation*

$$\nabla_\mu \Omega = -(\nabla_U \Omega + \Omega \langle U, f \rangle) U_\mu - \Omega f_\mu. \quad (4.16)$$

This relation will play a critical role. It will be used later to obtain a hyperbolic system of evolution equations which extends in a regular way to the set $\{\Omega = 0\}$ and it will be used to set up a subsidiary system to show that constraints and gauge conditions are preserved by the evolution system. Here it is used to remove the singularities in equations (3.20) and (3.21). In fact, replacing in Z_{jkl} the term $\nabla_k \Omega$ by the right hand side of (4.16), we get (3.20) in the form

$$\begin{aligned} \nabla_i W^i{}_{jkl} &= \nabla_{[k} \rho U_{l]} U_j + \frac{1}{3} \nabla_{[k} \rho g_{l]j} \\ &+ \rho (\nabla_{[k} U_{l]} U_j + U_{[l} \nabla_{k]} U_j - f_{[k} g_{l]j} - 2 f_{[k} U_{l]} U_j - U_{[k} g_{l]j} U^i f_i). \end{aligned} \quad (4.17)$$

Using (4.16) to replace $\nabla_k \Omega$ on the right hand side of (3.21), the equation takes the form

$$\nabla_0 U^k + f^k + U^k U_i f^i = 0, \quad (4.18)$$

which is just (4.13) again. Equation (3.13) is then replaced by the formally regular version

$$\nabla_k U_l = \Gamma_k{}^0{}_l = (-\delta^0{}_k f_b + \delta^a{}_k \chi_{ab}) \delta^b{}_l. \quad (4.19)$$

Finally we note that given sufficient asymptotic smoothness and an arrangement such that $\Omega(x(\tau)) \rightarrow 0$ for some finite value of τ , the relation

$$\frac{dt}{d\tau} = \frac{1}{\Omega(x(\tau))}, \quad (4.20)$$

which follows from (4.7), (4.12), (4.15) implies with (3.3) that $t \rightarrow \infty$ as $\Omega(x(\tau)) \rightarrow 0$.

5 The hyperbolic reduced equations

To extract from our equations a hyperbolic system we need to complete the gauge conditions for the g -orthonormal frame field e_k satisfying $e_0 = U$. The reduction procedure of the Einstein-dust system in [8] employs a frame that is \hat{g} -parallelly transported in the direction of \hat{U} . Since the field U is not geodesic with respect to g this cannot be done here. We use instead a frame whose vector fields X satisfy the Fermi transport law

$$0 = \mathbb{F}_U X \equiv \nabla_U X - g(X, \nabla_U U) U + g(X, U) \nabla_U U,$$

which has the properties: $\mathbb{F}_U U = 0$ and if $\mathbb{F}_U X = 0$, $\mathbb{F}_U Y = 0$ then $\nabla_U(g(X, Y)) = 0$.

On a given space-like hypersurface transverse to the flow line of U we thus choose smooth fields e_k with $e_0 = U$ such that $g_{jk} = g(e_j, e_k) = \eta_{jk}$ and extend the e_a away from the hypersurface by the requirement that $\mathbb{F}_U e_a = 0$. The smooth orthonormal frame field so obtained is then closely related to the frame considered in [8]. In fact, if \hat{e}_k is a \hat{g} -orthonormal frame such that $\hat{e}_0 = \hat{U}$ and $\hat{\nabla}_{\hat{U}} \hat{e}_k = 0$, then $e_k = \Omega^{-1} \hat{e}_k$ is a $g_{\mu\nu} = \Omega^2 \hat{g}_{\mu\nu}$ -orthonormal frame with $e_0 = U$ and $\mathbb{F}_U e_a = 0$.

As a consequence of relation $\mathbb{F}_U e_k = 0$ the connection coefficients satisfy

$$\Gamma_0^a{}_b = 0. \quad (5.1)$$

The transport equation for the flow field U is given by (4.13). The coefficients $U^\mu = e^\mu{}_0 = \delta^\mu{}_0$ have been fixed by our choice of coordinates, however, and equation (4.18) reduces to the relation

$$\Gamma_0^a{}_0 = -f^a = -g^{ab} f_b \quad \text{resp.} \quad \Gamma_0^0{}_a = -f_a, \quad (5.2)$$

between the connection coefficients and the acceleration of U . The remaining not necessarily vanishing connection coefficients are then given by

$$\Gamma_a^b{}_c \quad \text{and} \quad \Gamma_a^0{}_b = \nabla_a U_b = g(\nabla_{e_a} e_0, e_b) \equiv \chi_{ab} \quad \text{resp.} \quad \Gamma_a^b{}_0 = \chi_a^b = \chi_{ac} g^{cb}. \quad (5.3)$$

In the case in which U resp. \hat{U} is hypersurface orthogonal, the field χ_{ab} is symmetric and represents the second fundamental form while the $\Gamma_a^b{}_c$ are the connection coefficients of the intrinsic connection induced on the hypersurfaces orthogonal to U in the frame e_a .

We shall now derive the reduced equations for the remaining frame and connection coefficients. With our gauge conditions and the connection coefficients above the first structural equations (3.14) induce the evolution equations

$$e^\mu{}_{a,0} = -f_a \delta^\mu{}_0 - \chi_a^b e^\mu{}_b, \quad (5.4)$$

for the fields $e^\mu{}_a$.

The second structural equations (3.15) induce the evolution equations

$$\Gamma_c^a{}_b,0 = f^a \chi_{cb} - \chi_c^a f_b - \chi_c^d \Gamma_d^a{}_b + \Omega W^a{}_{b0c} - g^a{}_c L_{0b} + L^a{}_0 g_{cb}, \quad (5.5)$$

$$\chi_{ab,0} + D_a f_b = f_a f_b - \chi_a^c \chi_{cb} - \Omega W_{0b0a} + L_{ab} - L_{00} g_{ab}, \quad (5.6)$$

for $\Gamma_c^a{}_b$ and χ_{ab} , where we set

$$D_a f_b = f_{b,\mu} e^\mu{}_a - \Gamma_a^c{}_b f_c.$$

No equation is implied for $\Gamma_0^0{}_a = -f_a$ by (3.15). Such an equation is provided, however, by (4.14), which takes in our gauge the explicit form

$$f_{0,0} = -\frac{1}{2} f_j f^j + L_{00}, \quad (5.7)$$

$$f_{a,0} = L_{0a}. \quad (5.8)$$

At this stage arises a problem. We are aiming for a system that is symmetric hyperbolic. The principal part of the coupled system

$$\chi_{ab,0} + D_a f_b = \dots, \quad f_{a,0} = \dots,$$

does not satisfy the required symmetry condition. One might think of proceeding as follows. The structural equations (3.15) imply after a contraction an analogue of Codacci's equation, which takes with the convention $D_c \chi_{ab} \equiv \chi_{ab,\mu} e^\mu{}_c - \Gamma_c{}^d{}_a \chi_{db} - \Gamma_c{}^d{}_b \chi_{ad}$ the form

$$D^a \chi_{ab} - D_b(\chi_a{}^a) = \dots,$$

(where the index position in the first term has to be respected because χ_{ab} is not necessarily symmetric). By adding a suitable multiple of this equation to the second of the equations above one could hope to obtain a symmetric system. A careful analysis shows, however, that this does not work. We skip the details.

Help is again provided by (4.16). By this relation the field

$$N_k = \nabla_k \Omega + (\nabla_U \Omega + \Omega \langle U, f \rangle) U_k + \Omega f_k,$$

vanishes in our gauge. While $N_0 = N_k U^k = 0$ identically, the equation $N_a = 0$ with

$$N_a = \Omega f_a + \nabla_a \Omega,$$

has non-trivial content. The relation

$$\begin{aligned} \nabla_j N_k &= \nabla_j \nabla_k \Omega + \nabla_j (\nabla_U \Omega + \Omega \langle U, f \rangle) U_k \\ &+ (\nabla_U \Omega + \Omega \langle U, f \rangle) \nabla_j U_k + \nabla_j \Omega f_k + \Omega \nabla_j f_k, \end{aligned}$$

implies in our gauge

$$\begin{aligned} \nabla_a N_b - N_a f_b &= \nabla_a \nabla_b \Omega + (\nabla_U \Omega + \Omega \langle U, f \rangle) \chi_{ab} - \Omega f_a f_b + \Omega \nabla_a f_b \\ &= \nabla_a \nabla_b \Omega + (\nabla_U \Omega + \Omega \langle U, f \rangle) \chi_{ab} - \Omega f_a f_b + \Omega (D_a f_b - \chi_{ab} f_0). \\ &= \nabla_a \nabla_b \Omega + \nabla_U \Omega \chi_{ab} - \Omega f_a f_b + \Omega D_a f_b, \end{aligned}$$

which gives with (3.17)

$$\nabla_a N_b - N_a f_b = \nabla_U \Omega \chi_{ab} + s g_{ab} + \Omega (D_a f_b - f_a f_b - L_{ab} + \frac{1}{8} \Omega \rho g_{ab}). \quad (5.9)$$

Solving the equation $\nabla_a f_b - N_a f_b = 0$ for $D_a f_b$ and using the resulting expression to replace that term in the evolution equation for χ_{ab} , gives the latter in the form

$$\chi_{ab,0} - \Omega^{-1} (\nabla_U \Omega \chi_{ab} + s g_{ab}) = -\chi_a{}^c \chi_{cb} - \Omega W_{0a0b} - L_{00} g_{ab}. \quad (5.10)$$

With the reduced equations obtained so far and the ones that follow below this gives again a symmetric hyperbolic system where $\Omega \neq 0$.

Let us assume that the solution admits a smooth conformal boundary $\mathcal{J}^+ = \{\Omega = 0\}$. To obtain a system which extends in a regular fashion to \mathcal{J}^+ we recall that this would require that $e_0 = U$ approaches \mathcal{J}^+ orthogonally. With (3.16) this would imply that

$$\nabla_U \Omega \rightarrow -\nu < 0 \quad \text{as} \quad \Omega \rightarrow 0, \quad \text{where} \quad \nu \equiv \sqrt{-\frac{\lambda}{3}},$$

and thus $\nabla_U \Omega < 0$ also in a neighborhood of \mathcal{J}^+ . In the discussion of the conformal constraints on \mathcal{J}^+ in the next section we shall see that the conformal gauge can be chosen such that s and χ_{ab} vanish at \mathcal{J}^+ . If data on a ‘physical’ initial hypersurface are evolved in the direction of \mathcal{J}^+ it is, however, difficult to decide how the conformal gauge must be chosen such that these fields will vanish at \mathcal{J}^+ . This suggests to introduce regularizing unknowns which are derived from fields which go to zero at \mathcal{J}^+ in any conformal gauge. Such unknowns are suggested by the equation $\nabla_a f_b - N_a f_b = 0$. In fact, the fields

$$\zeta_{ab} \equiv \frac{\chi_{ab} - \frac{1}{3} g_{ab} \chi_c{}^c}{\Omega}, \quad \xi \equiv \frac{\nabla_U \Omega \chi_c{}^c + 3s}{\Omega}, \quad (5.11)$$

satisfy for $\Omega \neq 0$ and $\nabla_U \Omega \neq 0$ by (5.9)

$$\zeta_{ab} = -(\nabla_U \Omega)^{-1} \left(D_a f_b - f_a f_b - L_{ab} - \frac{1}{3} (D_c f^c - f_c f^c - L_c{}^c) g_{ab} \right), \quad (5.12)$$

and

$$\xi = -D_a f^a + f_a f^a + L_a{}^a - \frac{3}{8} \Omega \rho, \quad (5.13)$$

and can thus be expected to extend smoothly to \mathcal{J}^+ . The original unknown will be recovered from the new ones by

$$\chi_{ab} = \Omega \zeta_{ab} + \frac{1}{3} (\nabla_U \Omega)^{-1} (\Omega \xi - 3s) g_{ab}, \quad (5.14)$$

which will certainly be well defined on neighbourhoods of \mathcal{J}^+ where $\nabla_U \Omega \neq 0$. This will suffice for our purpose because we can use equation (5.10) where $\Omega \neq 0$.

The equations we have obtained so far imply equations for the unknowns (5.11) that are regular where $\nabla_U \Omega \neq 0$. Indeed, a direct calculation gives with (5.10) the equation

$$\zeta_{ab,0} = -\Omega (\zeta_a{}^c \zeta_{cb} - \frac{1}{3} \zeta^{cd} \zeta_{dc} g_{ab}) - \frac{2}{3} (\nabla_U \Omega)^{-1} (\Omega \xi - 3s) \zeta_{ab} - W_{0a0b}. \quad (5.15)$$

From (3.17) follows

$$\Omega_{,00} - \Gamma_0{}^a{}_0 \nabla_a \Omega = \nabla_0 \nabla_0 \Omega = -\Omega L_{00} - s + \frac{3}{8} \Omega^2 \rho,$$

and thus with $0 = N_a = \Omega f_a + \nabla_a \Omega$

$$\Omega_{,00} = \Omega f_a f^a - \Omega L_{00} - s + \frac{3}{8} \Omega^2 \rho.$$

Equation (3.18) gives with (3.22) and $N_a = 0$

$$s_{,0} = \nabla_U \Omega L_{00} + \Omega f^a L_{a0} - \frac{1}{4} \rho \Omega \nabla_U \Omega - \frac{1}{24} \rho \Omega^2 \chi.$$

With these two equations relation (5.10) implies

$$\begin{aligned} \xi_{,0} = (\nabla_U \Omega)^{-1} (\Omega \xi - 3s) & \left(-\frac{1}{3} \xi + f_a f^a - L_{00} + \frac{1}{4} \rho \Omega \right) \\ & - \nabla_U \Omega \Omega \zeta_{cd} \zeta^{dc} + 3 f^a L_{a0} - \frac{3}{4} \rho \nabla_U \Omega. \end{aligned} \quad (5.16)$$

This completes the evolution system for the metric and the connection coefficients.

To deal with equations of first order we introduce

$$\Sigma_k = \nabla_k \Omega,$$

as an unknown and use (3.17) to get the evolution equations

$$\nabla_0 \Omega = \Sigma_0, \quad (5.17)$$

$$\nabla_0 \Sigma_k = -\Omega L_{0k} + s g_{0k} + \frac{1}{2} \Omega^2 \rho \left(U_0 U_k + \frac{1}{4} g_{0k} \right). \quad (5.18)$$

From (3.18) we get

$$\nabla_0 s = -\nabla^i \Omega L_{i0} = \frac{1}{2} \Omega \rho \nabla^i \Omega \left(U_i U_0 + \frac{1}{4} g_{i0} \right) + \frac{1}{8} \Omega \rho \nabla_0 \Omega + \frac{1}{24} \Omega^2 \nabla_0 \rho. \quad (5.19)$$

As mentioned above, the Ricci scalar $R = R[g]$ of $g_{\mu\nu}$ will play the role of a conformal gauge source function and thus be prescribed as an explicit function of the coordinates near the initial hypersurface. Because of the relation

$$-L_{00} + g^{ab} L_{ab} = L_j{}^j = \frac{1}{6} R, \quad (5.20)$$

it suffices to derive an evolution system for the components L_{0a} , L_{ab} , $a, b = 1, 2, 3$, of the Schouten tensor. To simplify the equations we set

$$K_{jkl} = \nabla_i \Omega W^i{}_{jkl} \quad (5.21)$$

$$\begin{aligned} & + \Omega \left(\rho (\nabla_{[k} U_{l]} U_j + U_{[l} \nabla_{k]} U_j) + \nabla_{[k} \rho U_{l]} U_j + \frac{1}{3} \nabla_{[k} \rho g_{l]j} \right) \\ & + \rho (\nabla_{[k} \Omega g_{l]j} + 2 \nabla_{[k} \Omega U_{l]} U_j + U_{[k} g_{l]j} g^{pq} \nabla_p \Omega U_q), \end{aligned}$$

so that (3.19) takes the form

$$\nabla_k L_{lj} - \nabla_l L_{kj} = K_{jkl}.$$

It implies by contraction

$$\nabla_0 L_{l0} - g^{bc} \nabla_b L_{lc} = \frac{1}{6} \nabla_l R + K^j_{jl}.$$

These equations are used to define the evolution system

$$\nabla_0 L_{0a} - h^{bc} \nabla_b L_{ac} = \frac{1}{6} \nabla_a R + K^j_{ja}, \quad a = 1, 2, 3, \quad (5.22)$$

$$\nabla_0 L_{aa} - \nabla_a L_{0a} = K_{a0a}, \quad a = 1, 2, 3, \quad (5.23)$$

$$2 \nabla_0 L_{ab} - \nabla_a L_{0b} - \nabla_b L_{0a} = K_{a0b} + K_{b0a}, \quad a, b = 1, 2, 3, a \neq b. \quad (5.24)$$

for the set of unknowns

$$L_{01}, \quad L_{02}, \quad L_{03}, \quad L_{11}, \quad L_{12}, \quad L_{13}, \quad L_{22}, \quad L_{23}, \quad L_{33}.$$

For given right hand sides the system will then be symmetric hyperbolic on a neighborhood of an initial hypersurface on which $e_0^\mu = \delta^\mu_0$ and on which e^0_a is sufficiently small. Moreover, we find with our gauge conditions

$$K^j_{ja} = -\frac{1}{2} \rho (\Omega f_a + \nabla_a \Omega),$$

$$K_{a0b} = \nabla_i \Omega W^i_{a0b} + \frac{1}{2} \Omega \left(\rho \chi_{ba} + \frac{1}{3} \nabla_U \rho g_{ab} \right),$$

and thus the important fact that on the right hand sides of the evolution system above only that derivative of ρ occurs which can be removed by using the equation (3.22), i.e.

$$\nabla_U \rho + \rho \chi_a^a = 0. \quad (5.25)$$

This equation is assumed, of course, to be part of the reduced system.

The following extraction of an evolution system for the rescaled conformal Weyl tensor from equation (4.17) is close to the procedure to obtain evolution equations for the conformal Weyl tensor discussed in [8], [12], to which we refer for more details. Let

$$h^j_k = g^j_k + U^j U_k, \quad l^j_k = g^j_k + 2U^j U_k,$$

denote the projection operator which maps the tangent spaces onto their subspaces U^\perp orthogonal to U and the reflection operator which maps U onto $-U$ and induces the identity on U^\perp and consider the totally antisymmetric tensor densities

$$\epsilon_{ijkl} = \epsilon_{[ijkl]} \quad \text{with} \quad \epsilon_{0123} = 1 \quad \text{and} \quad \epsilon_{jkl} = U^i \epsilon_{ijkl}.$$

Further, define the U -electric part w_{jl} and the U -magnetic part w_{jl}^* of W^i_{jkl} by setting

$$w_{jl} = W_{ipkq} U^i h^p_j U^k h^q_l, \quad w_{jl}^* = \frac{1}{2} W_{ipmn} \epsilon^{mn}_{kq} U^i h^p_j U^k h^q_l,$$

so that these symmetric trace free fields are given in our gauge essentially by their ‘spatial’ components w_{ab} and w_{ab}^* .

It will be convenient to write equation (4.17) in the form $F_{jkl} = 0$ with

$$F_{jkl} = \nabla_i W^i{}_{jkl} - \nabla_{[k} \rho U_{l]} U_j - \frac{1}{3} \nabla_{[k} \rho g_{l]j} \quad (5.26)$$

$$-\rho (\nabla_{[k} U_{l]} U_j + U_{[l} \nabla_{k]} U_j - f_{[k} g_{l]j} - 2 f_{[k} U_{l]} U_j - U_{[k} g_{l]j} U^i f_i).$$

Inserting the representation

$$W_{ijkl} = 2 (l_{i[k} w_{l]j} - l_{j[k} w_{l]i} - U_{[k} w_{l]p}^* \epsilon^p{}_{ij} - U_{[i} w_{j]p}^* \epsilon^p{}_{kl}),$$

of the rescaled conformal Weyl tensor into the equations

$$0 = P_{ij} \equiv -F_{pkq} h^p{}_{(i} U^k h^q{}_{j)} + \frac{1}{3} h_{ij} h^{kl} F_{pmq} h^p{}_{k} U^m h^q{}_{l}, \quad (5.27)$$

$$0 = Q_{ij} \equiv -\frac{1}{2} F_{mpq} h^m{}_{(i} \epsilon_j)^{pq}, \quad (5.28)$$

the latter take the explicit form

$$w_{ab,0} + D_c w_{d(b}^* \epsilon_a)^{cd} = \chi_{(a}{}^c w_{b)c} + 2 \chi^c{}_{(a} w_{b)c} - 2 \chi_c{}^c w_{ab} \quad (5.29)$$

$$-h_{ab} \chi^{cd} w_{cd} - 2 a_c w_{d(b} \epsilon_b)^{cd} - \frac{1}{6} \rho (3 \chi_{(ab)} - h_{ab} \chi_c{}^c),$$

$$\begin{aligned} w_{ab,0}^* - D_c w_{d(b} \epsilon_a)^{cd} &= \chi^c{}_{(a} w_{b)c}^* - \chi_c{}^c w_{ab}^* \\ &+ 2 a_c w_{d(a} \epsilon_b)^{cd} + \chi_{cd} w_{ef} \epsilon_{(i}{}^{ce} \epsilon_j)^{df}, \end{aligned} \quad (5.30)$$

where we set, as before,

$$D_a w_{bc} = w_{bc,\mu} e^\mu{}_a - \Gamma_a{}^d{}_b w_{dc} - \Gamma_a{}^d{}_c w_{bd},$$

etc. (The slight differences with the analogues equations in [8], [12] result from the use of the relation $\mathcal{L}_U w_{ij} = w_{ij,0} + 2 \chi_{(i}{}^k w_{k)j}$ for w_{ab} and w_{ab}^* .) For given right hand side equations (5.29) and (5.30) represent a symmetric hyperbolic system for w_{ab} and w_{ab}^* if it is ignored that these fields are trace free. Their trace-freeness will be taken care of by the construction of the initial data and then be preserved by the equations. Again it is important that no derivatives of the field ρ occur on the right hand sides.

If on the right hand sides the field $\nabla_k \Omega$ is replaced by Σ_k , $\nabla_0 \rho$ is removed by using (5.25), χ_{ab} is replaced by (5.14), and L_{00} is removed where it occurs (also in expressions like $\nabla_a L_{0b} = L_{0b,\mu} e^\mu{}_a - \Gamma_a{}^k{}_0 L_{kb} - \Gamma_a{}^k{}_b L_{0k}$) by using (5.20), then equations (5.4), (5.5), (5.7), (5.8), (5.15), (5.16), (5.17), (5.18), (5.19), (5.22), (5.23), (5.24), (5.25), (5.29), (5.30) represent, irrespectively of the sign of Ω , for suitably chosen initial data a quasi-linear symmetric hyperbolic evolution system for the unknowns

$$e^\mu{}_a, \Gamma_c{}^a{}_b, f_k, \zeta_{ab}, \xi, \Omega, \Sigma_k, s, L_{0a}, L_{ab}, \rho, w_{ab}, w_{ab}^*,$$

where $\nabla_0 \Omega \neq 0$. Where $\Omega \neq 0$ such an evolution system can be obtained by replacing ζ_{ab} and ξ by χ_{ab} and using directly equation (5.10). The characteristics of the systems so obtained are time-like or null with respect to the solution metric, i.e. the metric $g_{\mu\nu}$ that satisfies $g_{\mu\nu} e^\mu{}_j e^\nu{}_k = \eta_{jk}$.

6 Asymptotic end data

In section 8 we shall discuss the natural question how initial data for the reduced field equations are derived from solutions to the constraints (2.9, (2.10) induced by the Einstein- λ -dust system on ‘physical’ initial hypersurfaces. The nature of the argument employed in 8 suggests, however, to consider first asymptotic data.

For solutions to Einstein’s field equations with a positive cosmological constant which admit a smooth conformal boundary \mathcal{J}^+ it has been observed in the vacuum case [4], in the case of matter models involving conformally covariant matter models with $\hat{g}^{\mu\nu} \hat{T}_{\mu\nu} = 0$ [6], and also in the case of a matter model with $\hat{g}^{\mu\nu} \hat{T}_{\mu\nu} \neq 0$ [10] that the problem of providing initial data simplifies considerably if solutions to the constraints are constructed on that boundary. There is no need any longer to consider non-linear elliptic equations. Assuming that the solutions admit a smooth conformal boundary $\mathcal{J}^+ = \{\Omega = 0\}$, it will be shown in this section that the constraints induced on \mathcal{J}^+ by the conformal equations in the Einstein-dust case with a positive cosmological constant lead to the same simplification. Moreover, in the particular case where $\rho > 0$ on \mathcal{J}^+ they simplify even further. The solutions to the conformal Einstein-dust constraints can then in principle be constructed without solving any differential equation at all.

To construct the *asymptotic end data* on a 3-manifold which will later acquire the status of a smooth conformal boundary, let S be a smooth, orientable, compact (though the latter is not really needed in the following discussion) 3-manifold. Assume that it represents a smooth conformal boundary \mathcal{J}^+ of an Einstein dust solution with cosmological constant $\lambda > 0$. The conformal constraints induced on it must then be considered with an induced metric which is Riemannian and a conformal factor Ω which vanishes on S . As seen earlier, the future directed conformal flow field U must be orthogonal to S . The conformal field equations will be considered in a frame e_k , $k = 0, 1, 2, 3$, on S so that $e_0 = U$ and the e_a , $a = 1, 2, 3$, represent a frame on S for the induced metric

$$h_{ab} = g_{ab} = g(e_a, e_b) = \text{diag}(1, 1, 1),$$

on S . The connection coefficients defined by g in the frame e_k are given again by $\nabla_k e_j = \Gamma_k^l{}_j e_l$. As before $h^j{}_k = g_j^k + U_j U^k$ denotes the orthogonal projector onto S . By assumption we have $\Omega > 0$ in the past and < 0 in the future of S and thus $e_0(\Omega) < 0$ on S . Because e_0 is orthogonal to S the field

$$\chi_{ab} = \Gamma_a^0{}_b = g(\nabla_{e_a} e_0, e_b),$$

represents the second fundamental form induced on S and is thus symmetric, while the $\Gamma_a^b{}_c$ define the connection coefficients on S in the frame e_a of the Levi-Civita connection D defined by the intrinsic metric h_{ab} .

The electric part $w_{jl} = W_{ipkq} U^i U^k h^p{}_j h^q{}_l$ of the rescaled conformal Weyl tensor is then represented by $w_{ab} = W_{0a0b}$ and $w_{ab}^* = \frac{1}{2} W_{0acd} \epsilon_b{}^{cd}$ represents its magnetic part $w_{jl}^* = \frac{1}{2} W_{ipmn} \epsilon^{mn}{}_{kq} U^i U^k h^p{}_j h^q{}_l$, where ϵ_{ijkl} and ϵ_{jkl} are defined as before.

With these assumptions equation (3.16) reduces to the condition

$$\nabla_0 \Omega = -\nu, \quad \nabla^0 \Omega = \nu \quad \text{on } S, \quad \text{where } \nu = \sqrt{\lambda/3} > 0. \quad (6.1)$$

Equation (3.17) reduces on S to $\nabla_i \nabla_j \Omega = s g_{ij}$. The only non-trivial condition implied by this relation is a restriction on the second fundamental form

$$\nu \chi_{ab} = s h_{ab} \quad \text{on } S. \quad (6.2)$$

Equation (3.18) implies the constraint

$$\nabla_a s + \nu L_{0a} = 0 \quad \text{on } S. \quad (6.3)$$

Under the conformal gauge transformation $g \rightarrow \bar{g} = \theta^2 g$, $\Omega \rightarrow \bar{\Omega} = \theta \Omega$ with smooth $\theta > 0$ the function s transforms as $s \rightarrow \bar{s} = \theta s + g^{\rho\delta} \nabla_\rho \Omega \nabla_\nu \theta$. This shows that for given $\theta > 0$ on S the derivative $\nabla_\mu \theta$ can be determined on S such that \bar{s} coincides on S with any prescribed function. The function s could be carried along as a free function in the following equations but for simplicity the choice that

$$s = 0, \quad \chi_{ab} = 0, \quad \nabla_i \nabla_j \Omega = 0, \quad L_{0a} = L_{a0} = 0 \quad \text{on } S, \quad (6.4)$$

will be assumed, which still leaves the freedom to rescale the metric on S . It should be observed, however, that the gauge above may not be satisfied if a solution is evolved into S from the domain where $\Omega > 0$. In that case the more general relations like (6.2) and (6.3) must be considered.

Because the conformal Weyl tensor $\Omega W^i{}_{jkl}$ vanishes on S , the curvature tensor of g is determined there by its Schouten tensor L_{jk} . Because the second fundamental form vanishes on S , the orthogonal projection of the curvature tensor of g onto S coincides by Gauss' theorem with the curvature tensor of h , i.e. $R_{abcd}[g] = R_{abcd}[h]$. It follows that the decomposition of $R_{abcd}[g]$ in terms $g_{ab} = h_{ab}$ and the components $L_{ab}[g]$ of its Schouten tensor is formally identically with the decomposition of $R_{abcd}[h]$ in terms h_{ab} and its Schouten tensor $l_{ab}[h] = R_{ab}[h] - \frac{1}{4} R[h] h_{ab}$. This implies that

$$L_{ab}[g] = l_{ab}[h],$$

which can be calculated from h_{ab} . The component L_{00} then follows from $\frac{1}{6} R[g] = L_j{}^j$ as

$$L_{00} = -\frac{1}{6} R[g] + h^{ab} L_{ab},$$

once the conformal gauge source function $R[g]$ has been prescribed.

Equation (3.19) induces the constraint $\nabla_a L_{bc} - \nabla_b L_{ac} = -\nu W^0{}_{cab}$ on S . Because the second fundamental form on S vanishes, it can be written in the form

$$w_{ab}^* = \frac{1}{\nu} \epsilon_a{}^{cd} D_c l_{db}. \quad (6.5)$$

The equation says that the magnetic part of the rescaled conformal Weyl tensor is given on S up to a factor by the (dualized) Cotton tensor of h . Equation (3.19) induces the further constraint $\nabla_a L_{b0} - \nabla_b L_{a0} = 0$ on S . This is satisfied as a consequence of (6.4).

With F_{jkl} given by (5.26), the constraints induced on S by equation (4.17) are given by (see [8], [12])

$$0 = P_k \equiv F_{jpl} U^j h^p{}_k U^l, \quad 0 = Q_k \equiv -\frac{1}{2} F_{jpq} U^j \epsilon_k{}^{pq}. \quad (6.6)$$

They can be written more explicitly in the form

$$D^a w_{ac} = \frac{1}{3} D_c \rho - \rho f_c, \quad (6.7)$$

which is a genuine constraint, and

$$D^a w_{ab}^* = 0, \quad (6.8)$$

which is, consistent with (6.5), the differential identity satisfied by the Cotton tensor and imposes thus no additional restriction.

The 1-form f_a characterizes the deviation of U from hypersurface orthogonality (see the datum \hat{u}^α in (2.11) and the following discussion of hypersurface orthogonal flows) and can be prescribed freely on S . The value of f_0 only affects the gauge. It can be prescribed freely and we assume that $f_0 = 0$ on S .

The initial data for ζ_{ab} and ξ which follow from (5.12) and (5.13) are then given on S by

$$\zeta_{ab} = \nu^{-1} \left(D_a f_b - f_a f_b - L_{ab} - \frac{1}{3} (D_c f^c - f_c f^c - L_c{}^c) g_{ab} \right), \quad (6.9)$$

and

$$\xi = -D_a f^a + f_a f^a + L_a{}^a. \quad (6.10)$$

The observations above can be summarized in terms of local coordinates x^α , $\alpha = 1, 2, 3$, on S as follows.

Lemma 6.1 *Any smooth initial data set for the reduced equations is determined on the set $S = \{\Omega = 0\}$ uniquely by a Riemannian metric $h_{\alpha\beta}$, the density $\rho \geq 0$, the acceleration f_α and a symmetric, h -trace free tensor field $w_{\alpha\beta}$, which are arbitrary up to the relation*

$$D^\alpha w_{\alpha\beta} = \frac{1}{3} D_\beta \rho - \rho f_\beta \quad \text{on } S, \quad (6.11)$$

where D denotes the Levi-Civita operator defined by $h_{\alpha\beta}$.

As in the cases mentioned in the beginning there is no need to solve an analogue of the Hamiltonian constraint. The Riemannian space $(S, h_{\alpha\beta})$ is not subject to any further restriction. The situation even simplifies for the class of data with $\rho > 0$ on S . In that case $h_{\alpha\beta}$, $\rho > 0$, and $w_{\alpha\beta}$ can be prescribed completely freely and f_β is then determined by reading (6.11) as its defining equation. It should be pointed out, however, that if f_α is required to satisfy some extra conditions, as in the hypersurface orthogonal case discussed below, equation (6.11) must be read as a differential equation. The situation can then be discussed by the well known splitting techniques used in the discussion of the standard constraints [1].

The gauge requirement $s|_{\{\Omega=0\}} = 0$ leaves the conformal gauge freedom

$$\Omega \rightarrow \Omega' = \theta \Omega, \quad g_{\mu\nu} \rightarrow g'_{\mu\nu} = \theta^2 g_{\mu\nu},$$

with smooth functions $\theta > 0$ that are arbitrary on S . If n^μ denotes the future directed unit normal to S the conformal gauge transformation above implies associated transformations

$$h_{\alpha\beta} \rightarrow h'_{\alpha\beta} = \theta^2 g_{\alpha\beta}, \quad n^\mu \rightarrow n'^\mu = \theta^{-1} n^\mu, \quad U^\mu \rightarrow U'^\mu = \theta^{-1} U^\mu, \quad \rho \rightarrow \rho' = \theta^{-3} \rho,$$

and, by the transformation law for the 1-forms associated with conformal geodesics,

$$f_\alpha \rightarrow f'_\alpha = f_\alpha - \theta^{-1} D_\alpha \theta. \quad (6.12)$$

If n is extended as unit vector field into \hat{M} , the relation $g_{\alpha\mu} W^\mu{}_{\nu\beta\rho} n^\nu n^\rho = \Omega^{-1} g_{\alpha\mu} C^\mu{}_{\nu\beta\rho} n^\nu n^\rho$ makes sense and suggests on S for $w_{\alpha\beta}$ the transformation law

$$w_{\alpha\beta} \rightarrow w'_{\alpha\beta} = \theta^{-1} w_{\alpha\beta}.$$

It follows then

$$h'^{\alpha\beta} D'_\alpha w'_{\rho\gamma} = \theta^{-3} h^{\alpha\beta} D_\alpha w_{\rho\gamma},$$

whence

$$D'^\alpha w'_{\alpha\beta} - \frac{1}{3} D'_\beta \rho' + \rho' f'_\beta = \theta^{-3} (D_\alpha w^\alpha{}_\beta - \frac{1}{3} D_\beta \rho + \rho f_\beta),$$

so that the constraints are preserved.

6.1 Hypersurface orthogonal flows

Obviously, the vector field \hat{U}^μ is hypersurface orthogonal where $\Omega \neq 0$ if and only if this is true for $U^\mu = \Omega^{-1} \hat{U}^\mu$. Formally this follows from the relation $\hat{U}_{[\rho} \hat{\nabla}_\mu \hat{U}_{\nu]} = \Omega^{-2} U_{[\rho} \nabla_\mu U_{\nu]}$. In our gauge the hypersurface orthogonality condition $\hat{U}_{[\rho} \hat{\nabla}_\mu \hat{U}_{\nu]} = 0$ is equivalent to

$$0 = \nabla_{[a} U_{b]} = \chi_{[ab]}. \quad (6.13)$$

From (5.6) we get with $\sigma_{ab} = \chi_{(ab)}$ along the flow lines of U^μ the ODE

$$\chi_{[ab],0} + D_{[a} f_{b]} = \sigma_a{}^c \chi_{[cb]} - \sigma_b{}^c \chi_{[ca]}.$$

It follows that $D_{[a} f_{b]} = 0$ if U^μ is hypersurface orthogonal. If the solution admitted a smooth conformal extension, so that $\chi_{[ab]} = 0$ on \mathcal{J}^+ , we could conclude from the equation above that $\chi_{[ab]} = 0$ if we knew that $D_{[a} f_{b]} = 0$. With the gauge condition $\nabla_a N_b - N_a f_b = 0$ equation (5.9) gives, however, only the relation

$$0 = \Omega_{,0} \chi_{[ab]} + \Omega D_{[a} f_{b]}.$$

But this combines with the equation above to give

$$(\Omega^{-1} \chi_{[ab]})_{,0} = \sigma_a{}^c (\Omega^{-1} \chi_{[cb]}) - \sigma_b{}^c (\Omega^{-1} \chi_{[ca]}).$$

It follows that $\chi_{[ab]} = 0$ along a given integral curve of U^μ if it vanishes at a point of it where $\Omega \neq 0$. On the other hand, the relation above shows that $\Omega^{-1} \chi_{[ab]}$ assumes the limit $(\nabla_0 \Omega)^{-1} D_{[a} f_{b]}$ on \mathcal{J}^+ , which vanishes where the integral curves of U^μ meet \mathcal{J}^+ and only if $D_{[a} f_{b]} = 0$ there. Observing the discussion of the conformal gauge freedom in the construction of data on the conformal boundary, in particular (6.12), we conclude:

Lemma 6.2 *Let be given a solution to the Einstein-dust system (2.1), (2.6), (2.7), (2.8) that admits a smooth conformal boundary \mathcal{J}^+ . Then the field U^μ is hypersurface orthogonal if and only if the initial data for the conformal field equations induced on \mathcal{J}^+ in the gauge above are such that*

$$D_{[a} f_{b]} = 0 \quad \text{on } \mathcal{J}^+.$$

If this condition is satisfied and the field f_a can be given on \mathcal{J}^+ as the differential of a function f , then the conformal gauge can be chosen so that $f_a = 0$ on \mathcal{J}^+ .

6.2 FLRW-type solutions

In the following we discuss the FLRW solutions along the lines of the previous sections. The FLRW-type solutions to (2.1), (2.6), (2.7), (2.8) on $\hat{M} = \mathbb{R} \times S$ with $S = \mathbb{S}^3, \mathbb{T}^3$ or \mathbb{H}_*^3 (a suitable factor space of hyperbolic 3-space) are of the form

$$\hat{g} = -dt^2 + a^2 k, \quad \hat{U} = \partial_t, \quad \hat{\rho} = \hat{\rho}(t) \geq 0,$$

with a function $a = a(t) > 0$ and a 3-metric of constant curvature which is given in local coordinates x^α , $\alpha, \beta, \dots = 1, 2, 3$, on S by $k = k_{\alpha\beta} dx^\alpha dx^\beta$, so that $R_{\alpha\beta\gamma\delta}[k] = 2\epsilon k_{\alpha[\gamma} k_{\beta]\delta}$ where $\epsilon = 1, 0$ or -1 . Rescaling the fields with a conformal factor $\Omega = \Omega(t)$

$$\hat{g} \rightarrow g = \Omega^2 \hat{g}, \quad \hat{U} \rightarrow U = \Omega^{-1} \hat{U}, \quad \hat{\rho} \rightarrow \rho = \Omega^{-3} \hat{\rho},$$

and introducing a coordinate $x^0 = \tau(t)$ so that $\langle U, d\tau \rangle = 1$, the conformal version of the metric above takes the form

$$g = -d\tau^2 + l^2 k, \quad U = \partial_\tau, \quad \rho = \rho(\tau),$$

with some function $l = l(\tau) > 0$. The non-vanishing Christoffel symbols and the second fundamental form $\chi_{\alpha\beta}$ of the slices $\{\tau = \text{const.}\}$ are then given by

$$\chi_{\alpha\beta} = \Gamma_{\alpha}{}^0{}_{\beta}[g] = ll' k_{\alpha\beta}, \quad \Gamma_0{}^\alpha{}_\gamma[g] = \Gamma_\gamma{}^\alpha{}_0[g] = \frac{1}{l} l' k^\alpha{}_\gamma, \quad \Gamma_\beta{}^\alpha{}_\gamma[g] = \Gamma_\beta{}^\alpha{}_\gamma[k],$$

where $' = \frac{d}{d\tau}$. The Ricci scalar and the Schouten tensor are given by

$$R[g] = \frac{6}{l^2} (\epsilon + ll'' + (l')^2),$$

$$L_{00}[g] = \frac{1}{2l^2} (\epsilon - 2ll'' + (l')^2), \quad L_{\alpha 0}[g] = L_{0\alpha}[g] = 0, \quad L_{\alpha\beta}[g] = \frac{1}{2} (\epsilon + (l')^2) k_{\alpha\beta}.$$

Choosing the conformal gauge function as $R[g] = 6\epsilon$ on \hat{M} , the function l must satisfy $ll'' + (l')^2 + \epsilon(1 - l^2) = 0$. Using the remaining conformal gauge freedom to achieve $l = 1$, $l' = 0$ on a slice $\{\tau = \text{const.}\}$, it follows that $l = 1$. The only non-vanishing Christoffel symbols are then given by $\Gamma_\beta{}^\alpha{}_\gamma[g] = \Gamma_\beta{}^\alpha{}_\gamma[k]$ and

$$L_{00} = \frac{\epsilon}{2}, \quad L_{\alpha 0} = L_{0\alpha} = 0, \quad L_{\alpha\beta} = \frac{\epsilon}{2} k_{\alpha\beta}.$$

Where $\Omega > 0$ the physical field is then given by

$$\hat{g} = \Omega^{-2} g = -dt^2 + a^2 d\omega^2, \quad (6.14)$$

$$a(t) = \frac{1}{\Omega(\tau(t))}, \quad \frac{dt}{d\tau} = \frac{1}{\Omega(\tau)}. \quad (6.15)$$

The high symmetry assumptions leads to a simplification of the conformal field equations. There do not occur singularities any longer in the equations. In fact, because U is g -geodesic and hypersurface orthogonal and $\Omega = \Omega(\tau)$, the singularity in (3.11) is gone. Because the line element g is locally conformally flat it follows that $W^\mu{}_{\nu\rho\kappa} = 0$ and thus $\hat{\nabla}_{[\nu} \hat{L}_{\lambda]\rho} = 0$ by (3.7). Moreover, it follows by (3.6) that $\nabla_{[\nu} L_{\lambda]\rho} = 0$.

It will be assumed in the following that the conformal time coordinate τ vanishes on a set $\{\Omega = 0\}$ and that $\nabla_U \Omega = \Omega' < 0$ there. Equations (3.3) and (3.8) then imply

$$\Omega'(0) = -\nu = -\sqrt{\lambda/3} < 0.$$

Equation (3.12) reduces because of $\nabla_\mu U^\mu = \chi_c{}^c = 0$ to $\rho' = 0$, so that

$$\rho = \rho_* = \text{const.} > 0,$$

equations (3.4) and (3.9) imply $s = \frac{\epsilon}{2} \Omega - \frac{1}{8} \rho_* \Omega^2$, $\Omega'' + \epsilon \Omega - \frac{1}{2} \rho_* \Omega^2 = 0$ and equations (3.5), (3.8), (3.9) give $s' = \frac{\epsilon}{2} \Omega' - \frac{1}{4} \rho_* \Omega \Omega'$, which is satisfied by the function s given above. The equations for s are redundant under the given assumptions. So we are left with the initial value problems

$$\Omega'' + \epsilon \Omega - \frac{1}{2} \rho_* \Omega^2 = 0, \quad \Omega(0) = 0, \quad \Omega'(0) = -\nu,$$

which clearly have a smooth solutions near $\{\tau = 0\} = \mathcal{J}^+$. Where $\Omega' \neq 0$ (thus in particular near \mathcal{J}^+ .) the ODE is equivalent to $(3\Omega'^2 + 3\epsilon\Omega^2 - \rho_*\Omega^3)' = 0$, which implies with the boundary conditions

$$3\Omega'^2 + 3\epsilon\Omega^2 - \rho_*\Omega^3 = \lambda. \quad (6.16)$$

The decreasing solutions to this equation cover all the expanding ends of the FRW-type solutions. With (6.15) the usual (physical) equations (see [14]) for $a(t)$ are implied by (6.16).

7 The subsidiary system

To show that solutions to the reduced equations for data which satisfy the constraints do indeed satisfy the complete set of conformal field equations, it has to be shown that the *zero quantities* N_j and

$$T_i{}^j{}_k, \quad \Delta^i{}_{jkl}, \quad A, \quad B_j, \quad C_{jl}, \quad D_j, \quad H_{jkl}, \quad F_{jkl}, \quad (7.1)$$

vanish as a consequence of the reduced equations and the given initial data. Here

$$N_j \equiv \Omega f_j + U^k \Sigma_k U_j + \Sigma_j + \Omega U^k f_k U_j,$$

$$T_i{}^k{}_j e_k \equiv -[e_i, e_j] + (\Gamma_i{}^l{}_j - \Gamma_j{}^l{}_i) e_l,$$

$$\Delta^i{}_{jkl} \equiv R^i{}_{jkl} - \Omega W^i{}_{jkl} - 2 \{g^i{}_{[k} L_{l]j} + L^i{}_{[k} g_{l]j}\}, \quad (7.2)$$

with

$$R^i{}_{jkl} = \Gamma_l{}^i{}_{j,\mu} e^\mu{}_k - \Gamma_k{}^i{}_{j,\mu} e^\mu{}_l + 2\Gamma_{[k}{}^i{}_{p} \Gamma_{l]pj} - 2\Gamma_{[k}{}^p{}_{l]} \Gamma_p{}^i{}_j,$$

$$A \equiv 6\Omega s - 3\Sigma_i \Sigma^i - \lambda - \frac{1}{4}\Omega^3 \rho,$$

$$B_k \equiv \nabla_k \Omega - \Sigma_k$$

$$C_{jk} \equiv \nabla_j \Sigma_k + \Omega L_{jk} - s g_{jk} - \frac{1}{2}\Omega^2 \rho \left(U_j U_k + \frac{1}{4} g_{jk} \right),$$

$$D_k \equiv \nabla_k s + \Sigma^i L_{ik} - \frac{1}{2}\Omega \rho \Sigma^i \left(U_i U_k + \frac{1}{4} g_{ik} \right) - \frac{1}{8}\Omega \rho \Sigma_k - \frac{1}{24}\Omega^2 \nabla_k \rho,$$

$$H_{jkl} \equiv \nabla_k L_{lj} - \nabla_l L_{kj} - K_{jkl},$$

$$F_{jkl} \equiv \nabla_i W^i{}_{jkl} - M_{jkl},$$

where

$$M_{jkl} = \nabla_{[k} \rho U_{l]} U_j + \frac{1}{3} \nabla_{[k} \rho g_{l]j} \quad (7.3)$$

$$+ \rho (\nabla_{[k} U_{l]} U_j + U_{[l} \nabla_{k]} U_j - f_{[k} g_{l]j} - 2 f_{[k} U_{l]} U_j - U_{[k} g_{l]j} U^i f_i),$$

$$K_{jkl} = \Sigma_i W^i{}_{jkl} + \Omega M_{jkl}. \quad (7.4)$$

Some of these quantities vanish trivially because of symmetries, gauge conditions, or the reduced equations. The latter comprise equations (4.13), (4.14), (5.25) and

$$U^i T_i{}^k{}_j = 0, \quad U^k \Delta^i{}_{jkl} = 0, \quad U^j B_j = 0, \quad U^j C_{jl} = 0, \quad U^j D_j = 0, \quad (7.5)$$

$$H^j{}_{ja} = 0, \quad H_{a0b} + H_{b0a} = 0, \quad a, b = 1, 2, 3, \quad P_{ij} = 0, \quad Q_{ij} = 0, \quad (7.6)$$

The zero quantities not in this list correspond to constraints or gauge conditions. Concerning the second of equations (7.5) we refer to the remarks below.

In the following we shall use the covariant derivative operator ∇_j defined by the connection coefficients $\Gamma_i{}^j{}_k$ that satisfy the gauge conditions and the reduced equations. This operator is metric in the sense that $\nabla_i g_{jk} = 0$ but, as seen from the first of conditions (7.5), it is not known a priori whether the connection is torsion free. In the following

arguments will be needed the commutators of covariant derivatives, which are for a function ϕ and a vector field X^i in the case of a general metric connection of the form

$$\begin{aligned}(\nabla_i \nabla_j - \nabla_j \nabla_i) \phi &= -T_i^l{}_j \nabla_l \phi \\ (\nabla_i \nabla_j - \nabla_j \nabla_i) X^k &= R^k{}_{lij} X^l - T_i^l{}_j \nabla_l X^j.\end{aligned}$$

To avoid carrying along various non-illuminating terms involving components of the torsion tensor we shall refer to such terms in an equation often in the form $\dots + P(T)$, where the dots indicate the equation of interest and $P(T)$ is a generic symbol for a polynomial in the components of the torsion tensor that satisfies $P(0) = 0$. The equation above will then take the form

$$(\nabla_i \nabla_j - \nabla_j \nabla_i) X^k = R^k{}_{lij} X^l + P(T).$$

The other zero quantities in the list (7.1) will be kept explicitly in an equation if needed to indicate how the calculations goes, otherwise the equations will be written in the form $\dots + P(Z)$, where the dots indicate the members of interest and $P(Z)$ is a polynomial in the components of the zero quantities (that may occasionally absorb a $P(T)$) with smooth coefficients that satisfies $P(0) = 0$.

The regular system has been obtained from the original version of the conformal field equations by using the gauge requirements $N_j = 0$ and $\nabla_a N_b = 0$. It needs to be shown that they are preserved by the reduced equations to establish that the original version of the conformal field equations is satisfied. They are needed in particular to show that the equations for ζ_{ab} and ξ imply the equations $U^i \Delta^0{}_{aib} = 0$, $U^i \Delta^a{}_{0ib} = 0$. The zero quantity N_j plays a particular role because its vanishing follows directly from the reduced equations and the initial conditions.

If $N_k = 0$ on a hypersurface transverse to the flow lines of U^k (which will, for instance, be the case if data are prescribed on $\{\Omega = 0\}$), this relation is preserved along the flow lines of U as a consequence of the reduced equations.

In fact, equations (4.13) and (4.14) imply

$$\nabla_U N_i = U^k U^l C_{kl} U_i + U^k C_{ki} + U^k B_k (f_i + U^l f_l U_i) - U_i f^k N_k,$$

which reduces with (7.5) to the linear homogeneous ODE

$$\nabla_U N_i = -U_i f^k N_k, \tag{7.7}$$

along the flow lines of U . From this the assertion follows. Since the solution to the reduced equations is ruled by the flow lines it follows also that $\nabla_i N_j = 0$ on the solution.

It can thus be assumed that $N_j = 0$, $\nabla_a N_b = 0$ so that we have indeed $U^i \Delta^0{}_{aib} = 0$ and the equivalent equation $U^i \Delta^a{}_{0ib} = 0$ as written in (7.5).

The subsequent discussion follows to some extent the derivation of subsidiary systems in earlier work on the conformal field equations. It will be convenient to use for the covariant derivative of a given tensor field $X_{ij}{}^k$ the notation

$$\nabla_l X_{ij}{}^k = e_l(X_{ij}{}^k) + (\Gamma X)_{lij}{}^k,$$

so that $X_{ij}{}^k \rightarrow (\Gamma X)_{lij}{}^k$ denotes a purely algebraic linear operator which does not involve derivatives.

The connection defined by the $\Gamma_i{}^j{}_k$ and the associated torsion and curvature tensor satisfy the first Bianchi identity

$$\sum_{(jkl)} \nabla_j T_k{}^i{}_l = \sum_{(jkl)} (R^i{}_{jkl} + T_j{}^m{}_k T_l{}^i{}_m),$$

where $\sum_{(jkl)}$ denotes the sum over the cyclic permutation of the indices jkl . Setting here $j = 0$, observing that the symmetries of $C^i{}_{jkl} = \Omega W^i{}_{jkl}$ and L_{kl} imply $\sum_{(jkl)} R^i{}_{jkl} = \sum_{(jkl)} \Delta^i{}_{jkl}$ and taking into account the reduced equations, we get from this the equation

$$\nabla_0 T_k{}^i{}_l = -(\Gamma T)_{l0}{}^i{}_k + (\Gamma T)_{k0}{}^i{}_l + 3 \sum_{(0kl)} (\Delta^i{}_{0kl} + T_0{}^m{}_k T_l{}^i{}_m) = P(Z). \quad (7.8)$$

To obtain an equation of the desired type for $\Delta^i{}_{jkl}$ we show that the right hand side of the identity

$$\nabla_j \Delta^i{}_{mkl} + \nabla_l \Delta^i{}_{mjk} + \nabla_k \Delta^i{}_{mlj} = \frac{1}{2} \epsilon_{njkl} \epsilon^{npqr} \nabla_p \Delta^i{}_{mqr}.$$

can be written as a linear expression in the zero quantities. We write (7.2) in the form

$$R^i{}_{jkl} = \Delta^i{}_{jkl} + \Omega W^i{}_{jkl} + G^i{}_{jkl} + E^i{}_{jkl},$$

with

$$G^i{}_{jkl} = L g^i{}_{[k} g_{l]j}, \quad L = L_i{}^i, \quad E^i{}_{jkl} = 2 \{g^i{}_{[k} L^*_{l]j} + L^{*i}{}_{[k} g_{l]j}\}, \quad L^*_{lj} = L_{lj} - \frac{1}{4} L g_{lj},$$

and use the second Bianchi identity

$$\sum_{(jkl)} \nabla_j R^i{}_{mkl} = - \sum_{(jkl)} R^i{}_{mpj} T_k{}^p{}_l, \quad (7.9)$$

to obtain

$$\begin{aligned} \epsilon^{njkl} \nabla_j \Delta_{imkl} &= -\epsilon^{njkl} (\nabla_j \Omega W_{imkl} + \Omega \nabla_j W_{imkl} \\ &\quad + \nabla_j G_{imkl} + \nabla_j E_{imkl} + R_{impj} T_k{}^p{}_l). \end{aligned}$$

The well known facts that the left and right duals of W_{ijkl} and G_{ijkl} coincide respectively while the left dual of E_{ijkl} differs from its right dual by a sign then imply with the reduced equations

$$\begin{aligned} \epsilon_n{}^{ijkl} \nabla_j \Delta_{imkl} &= \epsilon_{im}{}^{kl} (\nabla_j \Omega W^j{}_{nkl} + \Omega \nabla_j W^j{}_{nkl} \\ &\quad + \nabla_j G^j{}_{nkl} - \nabla_j E^j{}_{nkl}) - \epsilon_n{}^{ijkl} R_{impj} T_k{}^p{}_l \\ &= \epsilon_{im}{}^{kl} (\nabla_j \Omega W^j{}_{nkl} + \Omega (F_{nkl} + M_{nkl})) \\ &\quad - 2 \nabla_{[k} L g_{l]n} - 2 \nabla_{[k} L_{l]n} - 2 \nabla_j L^j{}_{[k} g_{l]n}) - \epsilon_n{}^{ijkl} R_{impj} T_k{}^p{}_l \end{aligned}$$

$$\begin{aligned}
&= \epsilon_{im}{}^{kl} (\nabla_j \Omega W^j{}_{nkl} + \Omega (F_{nkl} + M_{nkl}) \\
&\quad - H_{nkl} - \Sigma_i W^i{}_{nkl} - \Omega M_{nkl} - 2 H^j{}_{j[k} g_{l]n}) - \epsilon_n{}^{jkl} R_{impj} T_k{}^p{}{}_l.
\end{aligned}$$

In the last step it has been used that $K^j{}_{jl} = 0$. This follows because the tensor $W^i{}_{jkl}$ has vanishing contractions and because equations (4.13) and (5.25), which are satisfied as members of the reduced system, imply that $M^j{}_{jl} = 0$. Using again the reduced equations we finally get

$$\begin{aligned}
&\nabla_0 \Delta^i{}_{mkl} = -(\Gamma \Delta)_l{}^i{}_{m0k} + (\Gamma \Delta)_k{}^i{}_{m0l} \tag{7.10} \\
&-\frac{1}{2} \epsilon^n{}_{0kl} \{ \epsilon^i{}_{m}{}^{kl} (B_p W^p{}_{nkl} + \Omega F_{nkl} - H_{nkl} - 2 H^p{}_{pk} g_{ln}) - \epsilon_n{}^{jkl} R^i{}_{mp0} T_k{}^p{}{}_l \} = P(Z).
\end{aligned}$$

A direct calculation gives for the quantity

$$A = 6 \Omega s - 3 \Sigma_i \Sigma^i - \lambda - \frac{1}{4} \Omega^3 \rho, \tag{7.11}$$

the relation

$$\nabla_j A = 6 \Omega D_j - 6 \Sigma^i C_{ji} + (6s - \frac{3}{4} \Omega^2 \rho) B_j.$$

On the initial slice, where the zero quantities on the right hand side vanish by the construction of the initial data, this relation reduces to $\nabla_j A = 0$. This implies that $A = 0$ on that slice if it holds at one point of it. In the case of ‘physical’ data (i. e. $\Omega = 1$) the condition $A = 0$ reduces to $0 = 4 \hat{A} = \hat{R} - 4 \lambda - \hat{\rho}$, which will be satisfied by the construction of the physical data. Using the freedom to prescribe Ω and its time derivative on the initial slice the condition $A = 0$ can also be achieved in the transition to conformal data. We recall that the relation $A = 0$ served to determine the value of Σ_j in our discussion of the conformal data on $\{\Omega = 0\}$. With the reduced equations the relation above implies that

$$\nabla_U A = 0.$$

We can thus assume that $A = 0$ on the solution manifold.

A straightforward but lengthy calculation shows that the fields

$$Z_{jk}^B = \nabla_{[j} B_{k]}, \quad Z_{jkl}^C = \nabla_{[j} C_{k]l}, \quad Z_{jk}^D = \nabla_{[j} D_{k]},$$

can be expressed as linear (homogeneous) functions of the zero quantities with smooth coefficients. Taking into account the reduced equation $U^j B_j = 0$, $U^j C_{jl} = 0$, $U^j D_j = 0$ one gets

$$U^j \nabla_j B_k = 2 U^j Z_{jk}^B + U^j \nabla_k B_j = 2 U^j Z_{jk}^B + \nabla_k (U^j B_j) - (\nabla_k U^j) B_j = P(Z).$$

Similar calculations give

$$U^j \nabla_j B_k = P(Z), \quad U^j \nabla_j C_{kl} = P(Z), \quad U^j \nabla_j D_k = P(Z). \tag{7.12}$$

The remaining subsidiary equations are obtained by analyzing the expressions

$$\nabla_{[l} H^i{}_{jk]} \quad \nabla^j F_{jkl},$$

from two different points of view. As a preparation we observe the algebraic relations

$$M_{jkl} = -M_{jkl}, \quad M_{[jkl]} = 0, \quad M^j{}_{jl} = 0. \quad (7.13)$$

The first of them follow immediately from the definition while, as pointed out above, the last one follows as a consequence of the reduced equations (4.13) and (5.25). These relations imply

$$F_{jkl} = -F_{jkl}, \quad F_{[jkl]} = 0, \quad F^j{}_{jl} = 0, \quad (7.14)$$

and also

$$K^j{}_{jl} = 0. \quad (7.15)$$

Moreover, a straightforward though fairly lengthy calculation which makes repeatedly use of the reduced equations, shows that

$$\nabla^j M_{jkl} = P(Z), \quad (7.16)$$

and

$$\begin{aligned} \nabla_l K^l{}_{jk} &= \nabla_l \Sigma_i W^{il}{}_{jk} + \Sigma_i \nabla_l W^{il}{}_{jk} + \nabla_l \Omega M^l{}_{jk} + \Omega \nabla_l M^l{}_{jk} \\ &= C_{li} W^{il}{}_{jk} + B_l M^l{}_{jk} - \Sigma_l F^l{}_{jk} + \Omega \nabla_l M^l{}_{jk} = P(Z). \end{aligned}$$

From this follows the relation

$$\nabla_{[l} H^l{}_{jk]} = \Delta^l{}_{p[lj} L_k]^p - \nabla_{[l} K^l{}_{jk]} + P(T) = P(Z). \quad (7.17)$$

Similar calculations, which use that the left and right duals of the conformal Weyl tensor coincide, gives

$$\begin{aligned} \epsilon^{qljk} \nabla_l H^p{}_{jk} &= \epsilon^{qljk} (\nabla_l \nabla_j L_k{}^p - \nabla_l (\Sigma_n W^{np}{}_{jk} + \Omega M^p{}_{jk})) \\ &= \epsilon^{qljk} (\Delta^p{}_{nlj} L_k{}^n + W^p{}_{nlj} C_k{}^n - B_l M^p{}_{jk}) + \epsilon^{npik} \Sigma_n F^q{}_{jk} \\ &+ \frac{1}{2} \rho \Omega^2 \epsilon^{qljk} W^p{}_{njk} U^n U_l - 2 \Sigma_l M^{(q}{}_{jk} \epsilon^{p)ljk} - \Omega \nabla_l M^p{}_{jk} \epsilon^{pljk}. \end{aligned}$$

From equations (7.13), (7.16) follows that

$$\epsilon_{pqmn} \nabla_l M^p{}_{jk} \epsilon^{qljk} = P(Z).$$

Solving the equation $N_l = 0$ for Σ_l and inserting this into the equation above, we thus finally get

$$\begin{aligned} \epsilon^{qljk} \nabla_l H^p{}_{jk} &= \frac{1}{2} \rho \Omega^2 \epsilon^{qljk} W^p{}_{njk} U^n U_l \\ &- 2 \Sigma_l M^{(q}{}_{jk} \epsilon^{p)ljk} - \Omega \nabla_l M^{(p}{}_{jk} \epsilon^{q)ljk} + P(Z). \\ \epsilon^{qljk} \nabla_l H^p{}_{jk} &= \frac{1}{2} \rho \Omega^2 \epsilon^{qljk} W^p{}_{njk} U^n U_l + 2 \nabla_U \Omega U_l M^{(q}{}_{jk} \epsilon^{p)ljk} \\ &+ \Omega \{ 2 (f_l + \langle U, f \rangle U_l) M^{(q}{}_{jk} \epsilon^{p)ljk} - \nabla_l M^{(p}{}_{jk} \epsilon^{q)ljk} \} + P(Z). \end{aligned}$$

A direct calculation shows now that

$$\epsilon^{0ljk} \nabla_l H^0_{jk} = P(Z), \quad \epsilon^{aljk} \nabla_l H^b_{jk} = P(Z), \quad a, b = 1, 2, 3. \quad (7.18)$$

After solving the 9 reduced equations for the components L_{0a} , L_{ab} , they resume their original form if $1/6 R$ is replaced again by $L_j{}^j$. To show that they imply for suitably given initial data the full set $H_{jkl} = 0$, it needs to be shown that

$$H_{abc} = 0, \quad H_{0ab} = 0, \quad a \neq b.$$

In fact, the equation $0 = H^j{}_{ja} = -H_{00a} + g^{cd} H_{cad}$ implies then that $H_{00a} = 0$ and with the identities

$$H_{jkl} = -H_{jlk} \text{ and } \epsilon^{ijkl} H_{jkl} = 0, \text{ i.e. } \epsilon^{abc} H_{abc} = 0 \text{ and } H_{0ab} + H_{b0a} + H_{ab0} = 0, \quad a \neq b,$$

and the reduced equation $H_{a0b} + H_{b0a} = 0$ it follows then that

$$0 = H_{0ab} = -H_{b0a} + H_{a0b} = 2 H_{a0b} \quad a \neq b,$$

which exhaust the remaining cases.

We derive now the equations for the zero quantities above. The reduced equation $H^j{}_{ja} = 0$ implies that $\nabla_k H^l{}_{la} = (\Gamma H)_k{}^i{}_{la} = P(Z)$. Observing this in equations (7.17) we an equation of the form

$$\nabla_0 H_{0ab} - g_{cd} \nabla_c H_{dab} = P(Z). \quad (7.19)$$

On the other hand we have by (7.18)

$$\nabla_0 H_{dab} + \nabla_b H_{d0a} - \nabla_a H_{d0b} = 3 \nabla_{[0} H_{|d|ab]} = P(Z)$$

and

$$\nabla_d H_{0ab} + \nabla_b H_{0da} + \nabla_a H_{0bd} = 3 \nabla_{[d} H_{|0|ab]} = P(Z)$$

(where indices with a modulus sign are exempt from the anti-symmetrization). Observing the relations $H_{a0b} = -H_{b0a}$ and $2 H_{c0d} = H_{0cd}$ implied by the reduced equations, one gets from this an equation of the form

$$2 \nabla_0 H_{dab} - \nabla_d H_{0ab} = P(Z). \quad (7.20)$$

Equations (7.19), (7.20) constitute a system of equations for the unknowns H_{0ab} and H_{abc} which is, for given right hand sides, symmetric hyperbolic.

The properties (7.14) imply in particular the relation $F^a{}_{0a} = F^i{}_{0i} = 0$. The field P_{ij} and Q_{kl} introduced in (5.27 and (5.28) are thus completely represented by

$$P_{ab} = -F_{(a|0|b)}, \quad Q_{ab} = -\frac{1}{2} F_{(a}{}^{cd} \epsilon_{b)cd}.$$

To discuss the remaining content of the field F_{jkl} we recall the definitions

$$P_a = F_{0a0}, \quad Q_b = -\frac{1}{2} F_{0cd} \epsilon_b{}^{cd},$$

given in the discussion of the constraints. These fields exhaust the information in F_{0a0} and F_{0bc} . Because F_{a0b} is trace free it remains to control its anti-symmetric part. The relation $F_{[jkl]} = 0$ gives

$$-Q_c \epsilon^c{}_{ab} = \frac{1}{2} F_{0de} \epsilon_c{}^{de} \epsilon^c{}_{ab} = F_{0ab} = F_{a0b} - F_{b0a},$$

whence

$$F_{a0b} = -P_{ab} - \frac{1}{2} \epsilon_{abc} Q^c.$$

Because $F_{abc} \epsilon^{abc} = 0$, the field $F_{acd} \epsilon_b{}^{cd}$ is trace free. Contracting its anti-symmetric part suitably twice with epsilons and using that gives $F^j{}_{jl} = 0$ gives

$$F_{[a}{}^{cd} \epsilon_{b]cd} = -F_d{}^{dc} \epsilon_{cab} = -F_{00c} \epsilon^c{}_{ab} = P_c \epsilon^c{}_{ab},$$

and thus

$$F_{abc} = \frac{1}{2} Q_{ad} \epsilon_{bc}{}^d - h_{a[b} P_{c]}.$$

Observing now the reduced equations $P_{ab} = 0$ and $Q_{ab} = 0$, the remaining content of F_{jkl} is then described by the formula

$$F_{jkl} = 3U_j P_{[k} U_{l]} - g_{j[k} P_{l]} + Q_i (U_j \epsilon^i{}_{kl} - \epsilon^i{}_{j[k} U_{l]}).$$

Inserting this into $\nabla^j F_{jkl}$ and projecting suitably gives his

$$(\nabla_U P_l) h^l{}_i + \frac{1}{2} \epsilon_i{}^{kj} \nabla_k Q_j = \nabla^j F_{jkl} U^k h^l{}_i + P(Z),$$

$$(\nabla_U Q_l) h^l{}_i - \frac{1}{2} \epsilon_i{}^{kj} \nabla_k P_j = \frac{1}{2} \nabla^j F_{jkl} \epsilon_i{}^{kl} + P(Z).$$

Working then out $\nabla^j F_{jkl}$ explicitly and observing (7.16) one finally gets equations of the form

$$P_{a,0} + \frac{1}{2} \epsilon_a{}^{bc} D_b Q_c = P(Z), \tag{7.21}$$

$$Q_{a,0} - \frac{1}{2} \epsilon_a{}^{bc} D_b P_c = P(Z). \tag{7.22}$$

For given right hand sides this is a symmetric hyperbolic system for the fields P_a and Q_a .

We have seen above that solutions to the reduced equations for suitably arranged initial data satisfy $N_j = 0$ and $A = 0$. Equations (7.8), (7.10), (7.12), (7.19), (7.20), (7.21), (7.22) constitute a system of differential equations for those of the remaining components of the zero quantities (7.1) which do not vanish already because of gauge conditions or the reduced equation. The system is symmetric hyperbolic and has characteristics which are time-like or null with respect to the metric $g_{\mu\nu}$ that is supplied by the reduced system.

It follows that a solution to the reduced system for data that satisfy the conformal constraints on the initial slice satisfies on the domain of dependence of the initial slice the gauge conditions and the complete set of conformal Einstein- λ -dust equations.

8 Existence and strong future stability

In this section the properties of the conformal field equations derived above and standard results about quasi-linear symmetric hyperbolic systems will be used to draw conclusions on the global structure of solutions to the Einstein- λ -dust equations. Since we are mainly interested in C^∞ solutions and not in the weakest possible smoothness assumptions on the data we refrain from specifying Sobolev norms. We refer to [6] for details of the patching arguments in the context of Cauchy stability and for some relevant PDE reference.

8.1 Existence of asymptotically simple solutions

To construct solutions to the Einstein-dust equations with positive cosmological constant λ that admit a smooth conformal boundary in their infinite future we consider Cauchy problems for the reduced field equations on $\mathbb{R} \times S$ where data are prescribed on the submanifold $\{0\} \times S$. We identify the latter diffeomorphically with the manifold S underlying a given *asymptotic end data set* as considered in section 6. The *conformal time* variable τ in the reduced field equations will correspond to the factor \mathbb{R} above and it will be assumed that $\tau = 0$ on S . The conformal gauge source function represented by the Ricci scalar $R[g]$ of the conformal metric g to be constructed will be required to vanish and it is assumed that the condition $R[g] = 0$ is also underlying the construction of the given asymptotic end data. A fixed gauge source function will in general only work well for some limited time. For our purpose this will suffice, however, because it will be arranged that a finite interval of the conformal time τ will cover an interval of *physical* time of infinite extent.

Since S is compact and may have complicated topology, we use the fact that the hyperbolicity of the reduced equation allows us to obtain a solution on a neighborhood of $S \sim \{0\} \times S$ in $\mathbb{R} \times S$ by patching together local solutions. Compactness implies that S can be covered by a finite number of open subsets V_A , $A = 1, 2, \dots, k$, of S which carry smooth local coordinates x^α , $\alpha = 1, 2, 3$, and a smooth frame field e_a , $a = 1, 2, 3$, that satisfies $h_{ab} \equiv h(e_a, e_b) = \delta_{ab}$, where h denotes the 3-metric on S supplied by the asymptotic end data. It can be assumed that their exist shrinkings V'_A with compact closure \overline{V}'_A in V_A so that the V'_A still define an open covering and the boundary of V'_A in V_A is smooth. Standard results on symmetric hyperbolic systems then imply the existence of smooth solutions to the reduced field equations on open neighbourhoods \mathcal{D}_A of V'_A in $\mathbb{R} \times S$ which imply on V'_A the data induced on V'_A by the asymptotic end data on S in the gauge chosen on V_A . It can be assumed that the solution extends smoothly to the closure of \mathcal{D}_A in $\mathbb{R} \times S$ with $\det(e^\mu{}_k) \neq 0$ so that \mathcal{D}_A acquires a boundary which consists of (i) smooth hypersurfaces \mathcal{H}_A^\pm in the future/past of \mathcal{D}_A which are null with respect to the solution metric g and approach $\overline{V}'_A \setminus V'_A$ in their past/future, (ii) the intersection of $\overline{\mathcal{D}}_A$ with hypersurfaces $\{\tau = \tau_\pm\}$ in $\mathbb{R} \times S$ defined by some constants $\tau_- < 0 < \tau_+$ (which can be chosen to be the same for all V'_A), and (iii) the three 2-dimensional edges diffeomorphic to $\overline{V}'_A \setminus V'_A$ where these hypersurfaces approach each other. It can be assumed that the solution on \mathcal{D}_A is globally hyperbolic with respect to metric g . The subsidiary system then implies that the full set of conformal Einstein- λ -dust equations is satisfied on $\overline{\mathcal{D}}_A$.

If $p \in V'_A \cap V'_B$, there exists an open neighborhood $V_p \subset V'_A \cap V'_B$ of p so that solutions are given in the domain of dependence $\mathcal{D}_{A,p}$ of V_p in \mathcal{D}_A as well as in the domain of

dependence $\mathcal{D}_{B,p}$ of V_p in \mathcal{D}_B . On V_p these two solutions can be related to each other because the coordinate and frame transformations which relate the data induced on V_p by the data on V'_A and the data on V'_B respectively are known explicitly. Because the gauge inherent in the reduced equations is evolved by invariant propagation laws along the invariantly defined flow lines of the flow field U , the coordinate and frame transformations extend, independent of τ , and allow us to relate the solution on $\mathcal{D}_{A,p}$ isometrically to the solution on $\mathcal{D}_{B,p}$. By extending the argument it follows that the solution induced on the domain of dependence of $V'_A \cap V'_B$ in \mathcal{D}_A can be identified isometrically with the solution induced on the domain of dependence of $V'_A \cap V'_B$ in \mathcal{D}_B .

By patching together the local solutions, we obtain a smooth, globally hyperbolic solution to the conformal Einstein- λ -dust equations on a subset of the form $M = [\tau_*, \tau_{**}] \times S$ of $\mathbb{R} \times S$ with constants $\tau_* < 0 < \tau_{**}$ so that the conformal factor obtained on M satisfies $\Omega > 0$ on $\hat{M} = [\tau_*, 0[\times S$ while $\Omega < 0$ on $\check{M} =]0, \tau_{**}] \times S$.

The hypersurfaces $S_\sigma = \{\tau = \sigma = \text{const.}\}$ with $\tau_* \leq \sigma \leq \tau_{**}$ can be required to be space-like. In fact, with the co-normal to $\{\tau = \text{const.}\}$ given by $n_\mu = -a \tau_{,\mu}$ the future directed normal is given by

$$n^\mu = \frac{-a g^{\mu 0}}{\sqrt{|a^2 g^{00}|}} = -\frac{\eta^{jk} e^\mu_j e^0_k}{\sqrt{|\eta^{jk} e^0_j e^0_k|}} = \frac{\delta^\mu_0 - \eta^{ab} e^\mu_a e^0_b}{\sqrt{1 - \eta^{ab} e^0_a e^0_b}},$$

and the condition $n_\mu n^\mu = -1$ implies the expression

$$a = \frac{1}{\sqrt{1 - \eta^{ab} e^0_a e^0_b}}. \quad (8.1)$$

Moreover,

$$n^\mu = a (\delta^\mu_0 - \eta^{ab} e^\mu_a e^0_b) = \frac{1}{a} (U^\mu - u^\alpha \delta^\mu_\alpha) \quad \text{with} \quad u^\alpha = a^2 \eta^{ab} e^\alpha_a e^0_b. \quad (8.2)$$

We thus require that

$$e^0_a e^0_b \eta^{ab} \leq \text{const.} < 1 \quad \text{on} \quad M, \quad (8.3)$$

which can be achieved with suitable choices of τ_* and τ_{**} because $e^0_a = 0$ on S_0 . The hypersurfaces S_σ will then be Cauchy hypersurfaces for $(M, g_{\mu\nu})$. To simplify things, so that we only need to consider the regularized reduced equations involving the unknowns ζ_{ab} and ξ , it will also be assumed that $\Omega_{,\tau} < 0$ on M , which makes sense because $\Omega_{,\tau} = -\nu$ on S_0 .

The metric $g_{\mu\nu}$, the conformal factor Ω , the flow field U and the density function ρ are then such that the ‘physical’ fields

$$\hat{g}_{\mu\nu} = \Omega^{-2} g_{\mu\nu}, \quad \hat{U}_\mu = \Omega^{-1} U_\mu, \quad \hat{\rho} = \Omega^3 \rho \quad (8.4)$$

define a solution to the Einstein- λ -dust equations on the manifold \hat{M} with $\hat{\rho} \geq 0$ on \hat{M} . Extending smoothly to S_{τ_*} , this solution admits an extension into the past of S_{τ_*} but we are not interested here in controlling something like a maximal globally hyperbolic

solution. What is important for us is that the set $\mathcal{J}^+ \equiv S_0 = \{\Omega = 0\}$ defines for the solution $(\hat{M}, \hat{g}_{\mu\nu})$ a smooth conformal boundary at future time-like infinity.

Equations (3.14) to (3.22) are invariant under the transformation which implies the map

$$\Omega \rightarrow -\Omega, \quad \nabla_k \Omega \rightarrow -\nabla_k \Omega, \quad s \rightarrow -s, \quad W^i{}_{jkl} \rightarrow -W^i{}_{jkl}, \quad \rho \rightarrow -\rho, \quad \nabla_k \rho \rightarrow -\nabla_k \rho,$$

but leaves the fields $e^\mu{}_k$, $\Gamma_i{}^j{}_k$, L_{jk} , and U^k unchanged. It follows that after performing this transition on M and restricting to \check{M} gives us another solution to the Einstein- λ -dust equations on the manifold \check{M} . It follows, however, that then $\hat{\rho} \leq 0$ on \check{M} . For this solution the set $\{\Omega = 0\}$ defines a smooth conformal boundary in the infinite past. In this article we shall not be interested in this solution any further.

Two facts have been used above to obtain solutions whose conformal structures extend smoothly across future time-like infinity so as to define there smooth conformal boundaries: (i) The Einstein- λ -dust equations admit conformal representations which imply with suitable gauge conditions systems of evolution equations that are hyperbolic irrespective of the sign of the conformal factor Ω , (ii) some requirements needed to ensure the existence of smooth conformal extensions *are put in by hand* by starting from asymptotic end data.

The case of the Nariai solution, an explicit, geodesically complete solution to the Einstein- λ -dust equations with $\hat{\rho} = 0$, shows that that the property (i) is by itself not sufficient to ensure the existence of a smooth conformal boundary (see [11]). This raises the question whether the use of asymptotic end data may result in the construction of a very restricted class of solutions.

The following argument, introduced in the vacuum case in [5] and used in the presence of conformally invariant matter fields in [6], shows that the existence of smooth asymptotic conformal structures is in fact a fairly general feature of solutions to the Einstein- λ -dust equations. The smooth extensibility of the conformal structure across future time-like infinity will be *derived* as a consequence of the property (i) of the Einstein- λ -dust equations and the existence of a given reference solution that admits a smooth asymptotic structure.

8.2 Strong future stability of the solutions

Let

$$\Delta = (e^\mu{}_k, \Gamma_i{}^j{}_k, \zeta_{ab}, \xi, f_k, \Omega, \nabla_i \Omega, s, L_{jk}, W^i{}_{jkl}, U^k, \rho), \quad (8.5)$$

be one of the solutions constructed above. The associated physical fields $\hat{g}_{\mu\nu} = \Omega^{-2} g_{\mu\nu}$, $\hat{U}^\mu = \Omega U^\mu$, $\hat{\rho} = \Omega^3 \rho$ then induce on the Cauchy hypersurface $S' \equiv S_{\tau_*}$ with local coordinates x^α , $\alpha = 1, 2, 3$, standard Cauchy data $\hat{\delta} = (\hat{h}_{\alpha\beta}, \hat{k}_{\alpha\beta}, \hat{u}^\alpha, \hat{\rho})$, i.e. a solution to the constraints (2.9) and (2.10), where \hat{u}^α denotes the orthogonal projection of \hat{U}^μ onto S' .

As a first step towards showing that the asymptotic simplicity of the solution above is preserved under sufficiently small perturbations of the data $\hat{\delta}$, any given standard Cauchy data set on S' needs to be transformed into a suitable Cauchy data set for the conformal field equations. This involves several transformations and a suitable handling of the gauge freedom which will be discussed now by showing how the restriction of Δ to S' is obtained from $\hat{\delta}$.

Conformal data $\delta = (h_{\alpha\beta}, \kappa_{\alpha\beta}, u^\alpha, \rho)$ on S' are obtained from the standard data $\hat{\delta}$ by using the functions $\Omega > 0$ and $\nabla_U \Omega < 0$ on S' to define

$$h_{\alpha\beta} = \Omega^2 \hat{h}_{\alpha\beta}, \quad u^\alpha = \Omega^{-1} \hat{u}^\alpha, \quad \rho = \Omega^{-3} \hat{\rho},$$

and, using the transformation law of second fundamental forms under conformal rescalings,

$$\kappa_{\alpha\beta} = \Omega (\hat{\kappa}_{\alpha\beta} + \hat{h}_{\alpha\beta} \nabla_n \Omega).$$

Here n denotes the future directed unit normal to S' with respect to g , which is related to the flow vector field U and its projection u onto S' (that represents the shift vector field on S' , see the ADM representation of g below) by the relation

$$n = \frac{1}{a} (U - u) \quad \text{with} \quad a = \sqrt{1 + h_{\alpha\beta} u^\alpha u^\beta},$$

where the expression for the positive lapse function a is obtained from

$$-1 = g(U, U) = a^2 g(n, n) + g(u, u) = -a^2 + h_{\alpha\beta} u^\alpha u^\beta.$$

It follows that

$$\nabla_n \Omega = \frac{1}{a} (\nabla_U \Omega - \Omega_{,\alpha} u^\alpha),$$

can be calculated from the data given above.

When starting from arbitrarily given standard Cauchy data $\hat{\delta}$ the functions $\Omega > 0$ and $\nabla_U \Omega < 0$ are not given but represent part of the conformal gauge freedom. Suitable choices will be discussed later.

As a second step it will be convenient to derive all the unknowns entering the conformal field equations in a g -orthonormal frame c_k on S' which is adapted to S' in the sense that $c_0 = n$. This frame, which is not needed in the final process, is introduced because it simplifies various discussions. In a third step all the data will be expressed on S' in terms of the g -orthonormal frame e_k satisfying $e_0 = U$.

To remove the gauge freedom in the transition $c_k \rightarrow e_k$, we prescribe a specific field of Lorentz transformations K^i_j on S' which map the g -orthonormal frame field e_k with $e_0 = U$ onto a smooth g -orthonormal frame $c_j = K^i_j e_i$ field with $c_0 = n$ by setting

$$K^i_j = \begin{pmatrix} K^0_0 & K^0_b \\ K^a_0 & K^a_b \end{pmatrix} = \begin{pmatrix} -g(c_0, e_0) & g(c_0, e_b) \\ \eta^{ad} g(c_0, e_d) & \delta^a_b + \frac{1}{1-g(c_0, e_0)} \eta^{ad} g(c_0, e_d) g(c_0, e_b) \end{pmatrix}. \quad (8.6)$$

In terms of the frame coefficients e^μ_k given by the solution Δ this reads

$$K^i_j = \begin{pmatrix} a & -a e^0_b \\ -a \eta^{ac} e^0_c & \delta^a_b + \frac{a^2}{1+a} \eta^{ac} e^0_c e^0_b \end{pmatrix}.$$

It follows that indeed

$$K^i_0 e_i = K^0_0 e_0 + K^a_0 e_a = -g(c_0, e_0) e_0 + \eta^{ad} g(c_0, e_d) e_a = g(c_0, e_i) \eta^{ij} e_j = c_0.$$

In the following considerations (8.1) and (8.2) will be useful. A direct calculation verifies that $\eta_{ij} K^i{}_k K^j{}_l = \eta_{kl}$.

The coefficients of the frame c_k are given in the coordinates x^μ by

$$c^\mu{}_k = \begin{pmatrix} c^0{}_0 & , & 0 \\ c^\alpha{}_0 & , & c^\alpha{}_b \end{pmatrix} = \begin{pmatrix} \frac{1}{a} & , & 0 \\ -\frac{1}{a} u^\alpha & , & e^\alpha{}_b + \frac{1}{1+a} u^\alpha e^0{}_b \end{pmatrix},$$

and the coefficients of the 1-forms μ^k that satisfy $c^\mu{}_k \mu^k{}_\nu = \delta^\mu{}_\nu$ are so that

$$\mu^k{}_\nu = \begin{pmatrix} \mu^0{}_0 & , & 0 \\ \mu^a{}_0 & , & \mu^a{}_\beta \end{pmatrix},$$

with

$$\mu^0{}_0 = a, \quad u^\alpha = (e^\alpha{}_b + \frac{1}{1+a} u^\alpha e^0{}_b) \mu^b{}_0,$$

$$c^\alpha{}_b \mu^b{}_\beta = (e^\alpha{}_b + \frac{1}{1+a} u^\alpha e^0{}_b) \mu^b{}_\beta = \delta^\alpha{}_\beta,$$

whence

$$\mu^a{}_\alpha (e^\alpha{}_b + \frac{1}{1+a} u^\alpha e^0{}_b) = \delta^a{}_b, \quad \mu^a{}_\alpha u^\alpha = \mu^a{}_0.$$

The comparison of

$$g = \eta_{jk} \mu^j \mu^k = -a^2 d\tau^2 + \eta_{ab} \mu^a{}_\alpha \mu^b{}_\beta (u^\alpha d\tau + dx^\alpha) (u^\beta d\tau + dx^\beta),$$

with the ADM representation

$$g = -(a d\tau)^2 + h_{\alpha\beta} (u^\alpha d\tau + dx^\alpha) (u^\beta d\tau + dx^\beta)$$

gives then

$$h_{\alpha\beta} = \eta_{ab} \mu^a{}_\alpha \mu^b{}_\beta, \quad a = \sqrt{1 + u^\alpha u^\beta h_{\alpha\beta}}.$$

With the frame c_j defined above we set

$$M^i{}_j = \begin{pmatrix} -g(e_0, c_0) & , & g(e_0, c_b) \\ \eta^{ad} g(e_0, c_d) & , & \delta^a{}_b + \frac{1}{1-g(e_0, c_0)} \eta^{ad} g(e_0, c_d) g(e_0, c_b) \end{pmatrix}. \quad (8.7)$$

In terms of the frame coefficients c_k^μ this can be written

$$\begin{aligned} M^i{}_j &= \begin{pmatrix} M^0{}_0 & , & M^0{}_b \\ M^a{}_0 & , & M^a{}_b \end{pmatrix} = \begin{pmatrix} a & , & u_\alpha c^\alpha{}_b \\ \eta^{ac} u_\alpha c^\alpha{}_c & , & \delta^a{}_b + \frac{1}{1+a} \eta^{ac} u_\alpha c^\alpha{}_c u_\beta c^\beta{}_b \end{pmatrix} \\ &= \begin{pmatrix} a & , & a e^0{}_b \\ a \eta^{ac} e^0{}_c & , & \delta^a{}_b + \frac{a^2}{1+a} \eta^{ac} e^0{}_c e^0{}_b \end{pmatrix}. \end{aligned}$$

A direct calculation shows that $M^i{}_j K^j{}_k = \delta^i{}_k$ and $e_k = M^j{}_k c_j$.

Because the fields c_a , $a = 1, 2, 3$ are tangential to S' we can set

$$h'_{ab} = h_{\alpha\beta} c^\alpha{}_a c^\beta{}_b = \eta_{ab}, \quad \kappa'_{ab} = \kappa_{\alpha\beta} c^\alpha{}_a c^\beta{}_b$$

where here and in the following a prime is used to indicate when a tensor field is given in terms of the frame c_k . Directional derivatives with respect to c_k will also indicated by a prime, so that $\nabla'_k = \nabla_{c_k}$ etc.

When the data for the conformal field equations are to be constructed by starting from standard Cauchy data, the frame e_k is not available. Instead, the frame c_k has to be chosen first and e_k will then be obtained by applying M^i_k . The field c_0 is uniquely determined as the future directed unit normal to S' but the frame c_a tangent to S' is only determined up to rotations. In the stability argument given below this freedom will have to be removed in a specific way.

Connection coefficients with respect to the frame c_k satisfying the relation $\nabla_{c_i} c_k = \gamma_i^j c_j$ with respect to the Levi-Civita connection ∇ given by g can only be defined if the frame is defined near S' . It will be convenient to extend the frame by the requirement $\nabla_{c_0} c_k = 0$ and to define coordinates $v = x^{0'}$ and $x^{\alpha'}$ near S' so that $x^{\mu'} = x^\mu$ on S' and $\langle c_0, dv \rangle = 1$ and $\langle c_0, x^{\alpha'} \rangle = 0$. The coordinates $x^{\mu'}$ are then Gauss coordinates based on S' and the coefficients $c^{\mu'}_k$ satisfy $c^{\mu'}_0 = \delta^{\mu'}_0$ and $c^{0'}_a = 0$ so that $c_0 = \partial_v$. The coordinates x^μ and $x^{\mu'}$ satisfy

$$\begin{aligned} \frac{\partial x^0}{\partial x^{0'}} &= \langle n, d\tau \rangle = \frac{1}{a} \langle U - u, d\tau \rangle = \frac{1}{a}, & \frac{\partial x^0}{\partial x^{\alpha'}} &= 0, \\ \frac{\partial x^\alpha}{\partial x^{0'}} &= \frac{1}{a} \langle U - u, dx^\alpha \rangle = -\frac{1}{a} u^\alpha, & \frac{\partial x^\alpha}{\partial x^{\alpha'}} &= \delta^\alpha_{\alpha'} \quad \text{on } S', \end{aligned}$$

so that the relation $e^{\mu'}_k = M^j_k c^{\mu'}_j$ can be used to determine on S'

$$e^\mu_k = \frac{\partial x^\mu}{\partial x^{\mu'}} c^{\mu'}_l M^l_k.$$

The connection coefficients with respect to c_k can now be defined. They satisfy

$$\gamma_0^j{}_k = 0, \quad \gamma_a^0{}_b = \kappa'_{ab} = \kappa'_{ba}, \quad \gamma_a^c{}_0 = \kappa'_{ab} h^{bc}, \quad \gamma_a^d{}_b c_d = D_{c_a} c_b \quad \text{on } S',$$

where D denotes the Levi-Civita connection of the metric h on S' .

The connection coefficients in the frame c_k are related to the connection coefficients in the frame e_k by

$$\begin{aligned} \Gamma_i^j{}_k &= K^j_n \left(M^n_{k,\mu'} e^{\mu'}_i + \gamma_l^n{}_p M^l_i M^p_k \right) \\ &= K^j_n \left(M^n_{k,0'} e^{0'}_i + M^n_{k,\alpha'} e^{\alpha'}_i + \gamma_l^n{}_p M^l_i M^p_k \right). \end{aligned}$$

Apart from $M^n_{k,0'}$, which can only be determined by taking into account the evolution equations for the frame e_k , all the other terms in the expression above can be calculated from the data available so far. The relation $e^{\mu'}_k = M^j_k c^{\mu'}_j$ implies

$$e^{\mu'}_{k,0'} = M^j_{k,0'} c^{\mu'}_j + M^j_k c^{\mu'}_{j,0'}.$$

The first structural equation with respect to the frame c_k gives

$$c^{\mu'}_{j,0'} = \delta^{\mu'}_{\alpha'} \delta^a_j c^{\alpha'}_{a,0'} = -\delta^{\mu'}_{\alpha'} \delta^a_j \gamma_a^b{}_0 c^{\alpha'}_b = -\delta^{\mu'}_{\alpha'} \delta^a_j \kappa'_{ac} h^{bc} c^{\alpha'}_b \quad \text{on } S'.$$

The field $e_0 = U = U'^k c_k$, given on S' by $U = a c_0 + u'^a c_a$ with $u'^a = \mu^a{}_{\alpha'} u^{\alpha'}$, must thus satisfy by (4.13)

$$\begin{aligned} 0 &= U'^k{}_{,\mu'} c^{\mu'}{}_l U'^l + U'^l U'^j \gamma_l{}^k{}_j + \langle U, f \rangle U'^k + f'^k \\ &= a U'^k{}_{,0'} + U'^k{}_{,\alpha'} u^{\alpha'} + U'^l U'^j \gamma_l{}^k{}_j + \langle U, f \rangle U'^k + f'^k \quad \text{on } S'. \end{aligned}$$

The fields $e_a = e'^k{}_a c_k$ must satisfy $\mathbb{F}_U e_a = 0$, which implies with (4.13)

$$0 = a e'^k{}_{c,0'} + e'^k{}_{c,\alpha'} u^{\alpha'} + U'^i e'^j{}_c \gamma_i{}^k{}_j + f'_l e'^l{}_c U'^k - U'_l e'^l{}_c f'^k \quad \text{on } S'.$$

These relations determine $c^{\mu'}{}_{j,0'}$, $e^{\mu'}{}_{k,0'}$ whence $M^j{}_{k,0'}$ and $\Gamma_i{}^j{}_k$ uniquely from the given data on S' once f'_k is given there.

Our gauge requires that the tensorial field

$$N'_k = \nabla'_k \Omega + (\nabla_U \Omega + \Omega \langle U, f \rangle) U'_k + \Omega f'_k,$$

vanishes on S' . The condition that its orthogonal projection N'_a into S' vanishes gives

$$f'_a = -\frac{1}{\Omega} \{ \nabla'_a \Omega + (\nabla_U \Omega + \Omega \langle U, f \rangle) u'_a \} \quad \text{on } S'.$$

If this is satisfied it follows with $U_k = U'_i M^i{}_k$, $N_k = N'_i M^i{}_k$

$$0 = U^k N_k = U'^k N'_k = a n'^k N'_k,$$

and thus together $N'_k = 0$. The relation

$$f'_0 = n'^k f'_k = \frac{1}{a} (\langle U, f \rangle - u'^a f'_a),$$

shows that f'_k is determined from the data given on S' only up to $f_0 = \langle U, f \rangle$. This is consistent with the fact remarked on earlier that the quantity f_0 is pure gauge and can be chosen arbitrarily. With a suitable choice of f_0 (made in a specific way later) we can the set $f_k = f'_j M^j{}_k$.

The Einstein equations and the conformal rescaling of the density imply $R[\hat{g}] = 4\lambda + \Omega^3 \rho$. With this the conformal transformation law of the Ricci scalar gives

$$\nabla_\mu \nabla^\mu \Omega + \frac{1}{6} R[g] \Omega = \frac{2}{\Omega} \nabla_\mu \Omega \nabla^\mu \Omega + \frac{1}{6\Omega} R[\hat{g}] = \frac{2}{\Omega} \nabla_\mu \Omega \nabla^\mu \Omega + \frac{1}{6\Omega} (4\lambda + \Omega^3 \rho).$$

With the gauge condition $R[g] = 0$ we thus set

$$4s = \nabla'_k \nabla'^k \Omega = \frac{2}{\Omega} \nabla'_i \Omega \nabla'^i \Omega + \frac{1}{6\Omega} (4\lambda + \Omega^3 \rho).$$

The second equation determines $\partial_v^2 \Omega = c_0(c_0 \Omega)$ in terms of known data because

$$\nabla'_k \nabla'^k \Omega = -\nabla'_0 \nabla'_0 \Omega + \eta^{ab} \nabla'_a \nabla'_b \Omega = -c_0(c_0 \Omega) + \eta^{ab} (D'_a D'_b \Omega - \kappa'_{ab} \nabla_n \Omega) \quad \text{on } S'.$$

Thus s and $\nabla'_j \nabla'_k \Omega$ are determined on S' from known data and the scalar equation (3.16) is satisfied there. Given s and $\chi_{ab} = \Gamma_a^0 b$, the fields ζ_{ab} and ξ are then defined on S' by (5.11).

The conformal transformation law of the Schouten tensor, the field equations, and the conformal rescalings of the flow vector field and the density give

$$\begin{aligned} L_{\mu\nu} &= \hat{L}_{\mu\nu} - \frac{1}{\Omega} \nabla_\mu \nabla_\nu \Omega + \frac{1}{2\Omega^2} \nabla_\rho \Omega \nabla^\rho \Omega g_{\mu\nu} \\ &= \frac{1}{6} \lambda \Omega^{-2} g_{\mu\nu} + \Omega \rho \left(\frac{1}{2} U_\mu U_\nu + \frac{1}{6} g_{\mu\nu} \right) - \frac{1}{\Omega} \nabla_\mu \nabla_\nu \Omega + \frac{1}{2\Omega^2} \nabla_\rho \Omega \nabla^\rho \Omega g_{\mu\nu}, \end{aligned}$$

and we set

$$L'_{ij} = \frac{1}{6} \lambda \Omega^{-2} g'_{ij} + \Omega \rho \left(\frac{1}{2} U'_i U'_j + \frac{1}{6} g'_{ij} \right) - \frac{1}{\Omega} \nabla'_i \nabla'_j \Omega + \frac{1}{2\Omega^2} \nabla'_l \Omega \nabla'^l \Omega g'_{ij} \quad \text{on } S'.$$

By the way $\nabla'_0 \nabla'_0 \Omega$ has been determined above it follows that $g'^{ik} L'_{ik} = \frac{1}{6} R[g] = 0$. The appropriate data on S' for the reduced field equations are then given by $L'_{jk} = L'_{il} M^i_j M^l_k$.

To determine the rescaled conformal Weyl tensor we observe that the Gauss and the Codazzi equation with respect to S' read in terms of the frame c_k

$$\begin{aligned} R'_{abcd}[g] &= R'_{abcd}[h] + \kappa'_{ac} \kappa'_{bd} - \kappa'_{ad} \kappa'_{bc}, \\ n'^k R'_{kabc}[g] &= D'_c \kappa'_{ba} - D'_c \kappa'_{da}, \end{aligned}$$

where the fields on the right hand sides can be determined from the data available so far. With L'_{jk} as given above, the general relation

$$R'_{ijkl}[g] = 2 \{ g'_{i[k} L'_{l]j} + L'_{i[k} g'_{l]j} \} + C'_{ijkl},$$

then allows us to calculate the components $C'_{abcd}[g]$ and $n'^k C'_{kabc}[g]$ of the conformal Weyl tensor. The conformal Weyl tensor admits the decomposition

$$C'_{ijkl} = 2 \left(k'_{i[k} e'_{l]j} - k'_{j[k} e'_{l]i} + n'_{[k} m'_{l]m} \epsilon'^m{}_{ij} + n'_{[i} m'_{j]m} \epsilon'^m{}_{kl} \right).$$

where $h'_{jk} = g'_{jk} + n'_j n'_k$ and $k'_{jk} = g'_{jk} + 2n'_j n'_k$ and $e'_{ik} = h'^m{}_i h'^n{}_k C'_{mjnl} n'^j n'^l$ and $m'_{ik} = h'^m{}_i h'^n{}_k C'^*_{mjnl} n'^j n'^l$ with $C'^*_{ijkl} = \frac{1}{2} C'_{ijmn} \epsilon'^{mn}{}_{kl}$ denote the electric and magnetic part of the conformal Weyl tensor *with respect to* n in the frame c_k respectively. It holds $e'_{ij} = e'_{ji}$, $e'_{ij} n'^j = 0$, $e'^i{}_i = 0$ and similar relations hold for m'_{ij} .

It follows that

$$C'_{abcd} = 2 (h'_{a[c} e'_{d]b} + e'_{a[c} h'_{d]b}) \quad \text{whence} \quad e'_{bd} = h'^{ac} C'_{abcd},$$

and

$$n'^k C'_{kbcd} = 2 (n'_{[i} m'_{j]m} \epsilon'^m{}_{kl}) \quad \text{whence} \quad m'_{ab} = -\frac{1}{2} n'^k C'_{kbcd} \epsilon'_b{}^{cd}.$$

The tensors C'_{ijkl} and $W'_{ijkl} = \Omega^{-1} C'_{ijkl}$ whence $W_{ijkl} = W'_{mnpq} M^m{}_i M^n{}_j M^p{}_k M^q{}_l$ can thus be determined from the given data and thus also U -electric and -magnetic parts w_{ij} and w^*_{kl} of W_{ijkl} which enter the reduced conformal field equations.

The conformal field equations and their unknowns are derived from the Einstein equations by conformal rescalings, the use of various differential identities, and the use of the frame formalism. This leaves a coordinate, frame, and conformal gauge freedom which is controlled by suitable initial data and propagation laws for the coordinates, the frame field, and the conformal factor (controlled here implicitly by the requirement $R[g] = 0$). Following this procedure it follows from the discussion above how to derive from a given smooth solution $\hat{\delta} = (\hat{h}_{\alpha\beta}, \hat{\kappa}_{\alpha\beta}, \hat{u}^\alpha, \hat{\rho})$ to the constraints (2.9) and (2.10) and given smooth gauge dependent fields

$$\Omega > 0, \quad \nabla_U \Omega < 0, \quad f_0 = \langle U, f \rangle, \quad \text{and a smooth } h\text{-orthonormal field } c_a \text{ on } S', \quad (8.8)$$

the unknowns $\Delta'_{S'}$ on S' of the conformal field equations in the frame c_k and also the unknowns

$$\Delta_{S'} = (e^\mu{}_k, \Gamma_i{}^j{}_k, \zeta_{ab}, \xi, f_k, \Omega, \nabla_j \Omega, s, L_{jk}, W^i{}_{jkl}, U^k, \rho), \quad (8.9)$$

in the frame e_k on S' .

Written in terms of the frame c_k and the frame coefficients $c^{\mu'}{}_k$ as defined above, the conformal field equations allow us to derive from the data $\Delta'_{S'}$ a formal expansion type solution in terms of the coordinate v so that the complete set of conformal field equations is satisfied at all orders. The constraints are satisfied because of differential identities and the fact that the data $\hat{\delta}$ satisfy the ‘physical’ constraints.

A similar formal expansion is obtained in terms of the coordinate τ if the equations and the data are expressed in terms of the frame e_k . In this case the expansion coefficients are seen, however, to be the coefficients of a Taylor expansion of an actual smooth solution to the conformal field equations because the equations comprise the hyperbolic system of reduced conformal field equations.

The life time of the solution in the given gauge depends, of course, on the data (8.9) and in particular on the choice of the free fields in (8.8). Suppose

$$\Delta^*(\tau) = (e^{*\mu}{}_k, \Gamma_i^{*j}{}_k, \zeta_{ab}^*, \xi^*, f_k^*, \Omega^*, \nabla_j \Omega^*, s^*, L_{jk}^*, W^{*i}{}_{jkl}, U^{*k}, \rho^*), \quad (8.10)$$

is one of the solutions to the conformal field equations considered in the previous subsection. It exists and is smooth for $\tau_* \leq \tau \leq \tau_{**}$ with $\Omega^* \rightarrow 0$ as $\tau \rightarrow 0$ so that S_0 represents the conformal boundary at future time-like infinity for the physical solution associated with $\Delta^*(\tau)$. Denote by $\Delta_{S'}^* = \Delta^*(\tau_*)$ the data for the reduced equations on S' and by $\hat{\delta}^* = (\hat{h}_{\alpha\beta}^*, \hat{\kappa}_{\alpha\beta}^*, \hat{u}^{\alpha*}, \hat{\rho}^*)$ the physical data induced by this solution on S' . Let $\hat{\delta} = (S', \hat{h}_{\alpha\beta}, \hat{\kappa}_{\alpha\beta}, \hat{u}^\alpha, \hat{\rho})$ denote a smooth solution to the constraints (2.9) and (2.10), $\Delta_{S'}$ the corresponding initial data on S' for the reduced conformal field equations as considered in (8.8), and $\Delta(\tau)$, where $\tau \in [\tau_*, \tau_* + \tau^*[$ with some $\tau^* > 0$, the solution to the conformal field equations determined by these data.

To compare the life times of the solutions $\Delta^*(\tau)$ and $\Delta(\tau)$ the corresponding gauge conditions must be comparable. It will be assumed that the data $\Delta_{S'}$ have been con-

structured such that

$$\Omega = \Omega^*, \quad \nabla_U \Omega = \nabla_U \Omega^*, \quad f_0 = f_0^* \quad \text{on } S'.$$

Let $h_{\alpha\beta}^* = \Omega^{*2} \hat{h}_{\alpha\beta}^*$, and $h_{\alpha\beta} = \Omega^2 \hat{h}_{\alpha\beta} = \Omega^{*2} \hat{h}_{\alpha\beta}$ denote the metric induced on S' by the solution $\Delta^*(\tau)$ and $\Delta(\tau)$ respectively. As discussed above, the frame e_k^* given by the data $\Delta_{S'}^*$, can be used to define a field of Lorentz transformation K^{*j}_i on S' so that the relation $c_k^* = K^{*j}_k c_j^*$ defines a frame field on S' for which c_0^* is normal to S' . The fields c_a^* , $a = 1, 2, 3$, then define an h^* -orthonormal frame field on S' . It will be assumed in the following that the h -orthonormal field c_a has been chosen so that $c_a = c_c^* \alpha^c_a$ with a 3×3 matrix α^c_a that satisfies $\alpha^1_1 > 0$, $\alpha^2_2 > 0$, $\alpha^3_3 > 0$, and $\alpha^c_a = 0$ if $a < c$. The frame c_a so defined is smooth and fixed uniquely so that $\alpha^c_a \rightarrow \delta^c_a$ precisely if $c_a \rightarrow c_a^*$.

The point of these choices is that the space-time conditions $R[g^*] = 0$ and $R[g] = 0$ combine with these gauge conditions on S' to ensure that $\|\hat{\delta} - \hat{\delta}^*\| \rightarrow 0$ if and only if $\|\Delta_{S'} - \Delta_{S'}^*\| \rightarrow 0$, where the norms are meant to indicate Sobolev norms on S' which are chosen corresponding to the differentiability order of the fields involved.

We can invoke now the Cauchy stability property which holds for hyperbolic equations to conclude that for data $\hat{\delta}$ sufficiently close to $\hat{\delta}^*$ or, equivalently, for data $\Delta_{S'}$ sufficiently close to $\Delta_{S'}^*$, the solution $\Delta(\tau)$ of the conformal field equations that develops from the data $\Delta_{S'}$ also exists in the interval $\tau_* \leq \tau \leq \tau_{**}$ and the conformal factor Ω supplied by $\Delta(\tau)$ is negative on $S_{\tau_{**}}$ [15]. This conclusion may require repeated patchings (see [6]).

There exists then a map $S \ni q \rightarrow \tau(q) \in]\tau_*, \tau_{**}[$ so that $\Omega(\tau(q), q) = 0$ for $q \in S$ and $\Omega(\tau, q) > 0$ if $\tau_* \leq \tau < \tau(q)$. Equation (3.16) then implies that on the subset $\mathcal{J}^+ = \{(\tau(q), q), q \in S\}$ of $\mathbb{R} \times S$ the gradient $\nabla^i \Omega$ is time-like for the metric g supplied by $\Delta(\tau)$. It follows that \mathcal{J}^+ defines a smooth space-like hypersurface which represents a conformal boundary in the infinite future of the set $\hat{M} = \{(\tau, q) \in \mathbb{R} \times S \mid \tau_* \leq \tau < \tau(q)\}$ on which the fields $\hat{g}_{\mu\nu} = \Omega^{-2} g_{\mu\nu}$, $\hat{U}_\mu = \Omega^{-1} U^\mu$, $\hat{\rho}' = \Omega^3 \rho$ define a smooth solution to the Einstein- λ -dust equations. The smooth asymptotic end data induced by its conformal extension $\Delta(\tau)$ on $\mathcal{J}^+ \sim S$ belongs then to the class of conformal end data considered in section 6. Combining the results of the last two subsection we obtain Theorem 1.1.

Acknowledgements: I would like to thank the relativity group at Cordoba in Argentina, where this work was begun, for hospitality and discussions.

References

- [1] R. Bartnik, J. Isenberg. The constraint equations. In: P. T. Chruściel, H. Friedrich (eds.): *The Einstein equations and the large scale behaviour of gravitational fields*. Birkhäuser, Basel, 2004.
- [2] F. Beyer. Investigations of solutions of Einstein's field equations close to lambda-Taub-NUT. *Class. Quantum Gravity* 25 (2008) 235005.
- [3] H. Friedrich. On the hyperbolicity of Einstein's and other gauge field equations. *Commun. Math. Phys.* 100 (1985) 525 - 543.

- [4] H. Friedrich. Existence and structure of past asymptotically simple solution of Einstein's field equations with positive cosmological constant. *J. Geom. Phys.* 3 (1986) 101 - 117.
- [5] H. Friedrich. On the existence of n-geodesically complete or future complete solutions of Einstein's field equations with smooth asymptotic structure. *Commun. Math. Phys.* 107 (1986), 587 - 609.
- [6] H. Friedrich. On the global existence and the asymptotic behaviour of solutions to the Einstein-Maxwell-Yang-Mills equations. *J. Differential Geometry* 34 (1991) 275 - 345.
- [7] H. Friedrich. Einstein equations and conformal structure: existence of anti-de Sitter-type space-times. *J. Geom. Phys.* 17 (1995) 125–184.
- [8] Friedrich, H. (1998) Evolution equations for gravitating ideal fluid bodies in general relativity. *Phys. Rev. D* 57, 2317–2322.
- [9] H. Friedrich. Conformal geodesics on vacuum space-times. *Commun. Math. Phys.* 235 (2003) 513 - 543.
- [10] H. Friedrich. Smooth non-zero rest-mass evolution across time-like infinity. *Ann. Henri Poincaré* 16 (2015) 2215 - 2238.
- [11] H. Friedrich. Geometric asymptotics and beyond. In: L.Bieri, S.-T. Yau (eds), *Surveys in Differential Geometry*, Vol.20. International Press, Boston, 2015. arXiv:1411.3854
- [12] H. Friedrich, A. Rendall. The Cauchy Problem for the Einstein Equations. In: B. Schmidt (ed.): *Einstein's Field Equations and Their Physical Implications*. Springer, Lecture Notes in Physics, Berlin 2000.
- [13] M. Hadžić, J. Speck. The global future stability of the FLRW solutions to the Dust–Einstein system with a positive cosmological constant. *J. Hyperbolic Differential Equations* 12 (2015) 87.
- [14] S. Hawking, G. Ellis. *The large scale structure of space-time*. Cambridge University Press, 1973.
- [15] T. Kato. The Cauchy problem for quasi-linear symmetric hyperbolic systems. *Arch. Ration. Mech. Anal.*, 58 (1975) 181 - 205.
- [16] C. Lübbe, J. A. Valiente Kroon. A conformal approach to the analysis of the non-linear stability of radiation cosmologies. *Ann. Phys.* 328 (2013) 1.
- [17] R. Penrose. Zero rest-mass fields including gravitation: asymptotic behaviour. *Proc. Roy. Soc. Lond. A* 284 (1965) 159 - 203.
- [18] R. Penrose. *Cycles of Time*. Vintage, London, 2011.
- [19] H. Ringström. Future stability of the Einstein-non-linear scalar field system. *Invent. math.* 173 (2008) 123 - 208.