

**Global carbon budget
2013**

C. Le Quéré et al.

Global carbon budget 2013

C. Le Quéré¹, G. P. Peters², R. J. Andres³, R. M. Andrew², T. Boden³, P. Ciais⁴, P. Friedlingstein⁵, R. A. Houghton⁶, G. Marland⁷, R. Moriarty¹, S. Sitch⁸, P. Tans⁹, A. Arneth¹⁰, A. Arvanitis¹⁰, D. C. E. Bakker¹¹, L. Bopp⁴, J. G. Canadell¹², L. P. Chini¹³, S. C. Doney¹⁴, A. Harper¹⁵, I. Harris¹⁶, J. I. House¹⁷, A. K. Jain¹⁸, S. D. Jones¹, E. Kato¹⁹, R. F. Keeling²⁰, K. Klein Goldewijk²¹, A. Körtzinger²², C. Koven²³, N. Lefèvre²⁴, A. Omar²⁵, T. Ono²⁶, G.-H. Park²⁷, B. Pfeil^{25,28}, B. Poulter⁴, M. R. Raupach¹², P. Regnier²⁹, C. Rödenbeck³⁰, S. Saito³¹, J. Schwinger^{25,28}, J. Segsneider³², B. D. Stocker³³, B. Tilbrook³⁴, S. van Heuven³⁵, N. Viovy⁴, R. Wanninkhof³⁶, A. Wiltshire³⁷, S. Zaehle³⁰, and C. Yue⁴

¹Tyndall Centre for Climate Change Research, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK

²Center for International Climate and Environmental Research – Oslo (CICERO), Norway

³Carbon Dioxide Information Analysis Center (CDIAC), Oak Ridge National Laboratory, Oak Ridge, Tennessee, USA

⁴Laboratoire des Sciences du Climat et de l'Environnement, CEA-CNRS-UVSQ, CE Orme des Merisiers, 91191 Gif sur Yvette Cedex, France

⁵College of Engineering, Mathematics and Physical Sciences, University of Exeter, UK

⁶Woods Hole Research Centre (WHRC), Falmouth, Massachusetts 02540, USA

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- ⁷Research Institute for Environment, Energy, and Economics, Appalachian State University, Boone, North Carolina 28608, USA
- ⁸College of Life and Environmental Sciences University of Exeter, Exeter, Devon, EX4 4RJ, UK
- ⁹National Oceanic & Atmosphere Administration, Earth System Research Laboratory (NOAA/ESRL), Boulder, Colorado 80305, USA
- ¹⁰Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research/ Atmospheric Environmental Research, 82467 Garmisch-Partenkirchen, Germany
- ¹¹School of Environmental Sciences, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK
- ¹²Global Carbon Project, CSIRO Marine and Atmospheric Research, Canberra, Australia
- ¹³Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA
- ¹⁴Woods Hole Oceanographic Institution (WHOI), Woods Hole, MA 02543, USA
- ¹⁵College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, Devon, EX4 4RJ, UK
- ¹⁶Climatic Research Unit, University of East Anglia, Norwich Research Park, Norwich, NR4 7TJ, UK
- ¹⁷Cabot Institute, Dept. of Geography, University of Bristol, Bristol, Bristol BS8 1TH, UK
- ¹⁸Department of Atmospheric Sciences, University of Illinois, Urbana, IL 61821, USA
- ¹⁹Center for Global Environmental Research (CGER), National Institute for Environmental Studies (NIES), 16-2 Onogawa, Tsukuba, Ibaraki 305-8506, Japan
- ²⁰University of California, San Diego, Scripps Institution of Oceanography, La Jolla, California 92093-0244, USA
- ²¹PBL Netherlands Environmental Assessment Agency, The Hague/Bilthoven and Utrecht University, Utrecht, the Netherlands
- ²²GEOMAR Helmholtz Centre for Ocean Research, Wischhofstr. 1–3, 24148 Kiel, Germany
- ²³Earth Sciences Division, Lawrence Berkeley National Lab, 1 Cyclotron Road, Berkeley, California 94720, USA
- ²⁴IRD LOCEAN, Paris, France
- ²⁵Bjerknes Centre for Climate Research, Bergen, Norway
- ²⁶Fisheries Research Agency, 2-3-3 Minato Mirai, Nishi-ku, Yokohama, Kanagawa 220-6115, Japan

Global carbon budget 2013

C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


²⁷ East Sea Research Institute, Korea Institute of Ocean Science & Technology (KIOST), Ulsan, 767-813, South Korea

²⁸ Geophysical Institute, University of Bergen, Bergen, Norway

²⁹ Department of Earth & Environmental Sciences, CP160/02, Université Libre de Bruxelles, 1050 Brussels, Belgium

³⁰ Max-Planck-Institut für Biogeochemie, P.O. Box 600164, Hans-Knöll-Str. 10, 07745 Jena, Germany

³¹ Marine Division, Global Environment and Marine Department, Japan Meteorological Agency, 1-3-4 Otemachi, Chiyoda-ku, Tokyo 100-8122, Japan

³² Max Planck Institute for Meteorology, Bundesstr. 53, 20146 Hamburg, Germany

³³ Physics Institute, and Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland

³⁴ CSIRO Marine and Atmospheric Research and Antarctic Climate and Ecosystems Co-operative Research Centre, Hobart, Australia

³⁵ Centre for Isotope Research, University of Groningen, Groningen, the Netherlands

³⁶ NOAA/AOML, Miami, Florida, FL 33149, USA

³⁷ Met Office Hadley Centre, FitzRoy Road, Exeter, Devon, EX1 3PB, UK

Received: 7 November 2013 – Accepted: 8 November 2013 – Published: 19 November 2013

Correspondence to: C. Le Quéré (c.lequere@uea.ac.uk)

Published by Copernicus Publications.

ESSDD

6, 689–760, 2013

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Abstract

Accurate assessment of anthropogenic carbon dioxide (CO₂) emissions and their redistribution among the atmosphere, ocean, and terrestrial biosphere is important to better understand the global carbon cycle, support the development of climate policies, and project future climate change. Here we describe datasets and a methodology to quantify all major components of the global carbon budget, including their uncertainties, based on the combination of a range of data, algorithms, statistics and model estimates and their interpretation by a broad scientific community. We discuss changes compared to previous estimates consistency within and among components, alongside methodology and data limitations. CO₂ emissions from fossil-fuel combustion and cement production (E_{FF}) are based on energy statistics, while emissions from Land-Use Change (E_{LUC}), including deforestation, are based on combined evidence from land-cover change data, fire activity in regions undergoing deforestation, and models. The global atmospheric CO₂ concentration is measured directly and its rate of growth (G_{ATM}) is computed from the annual changes in concentration. The mean ocean CO₂ sink (S_{OCEAN}) is based on observations from the 1990s, while the annual anomalies and trends are estimated with ocean models. The variability in S_{OCEAN} is evaluated for the first time in this budget with data products based on surveys of ocean CO₂ measurements. The global residual terrestrial CO₂ sink (S_{LAND}) is estimated by the difference of the other terms of the global carbon budget and compared to results of Dynamic Global Vegetation Models. All uncertainties are reported as ± 1 sigma, reflecting the current capacity to characterise the annual estimates of each component of the global carbon budget. For the last decade available (2003–2012), E_{FF} was $8.6 \pm 0.4 \text{ GtC yr}^{-1}$, E_{LUC} $0.8 \pm 0.5 \text{ GtC yr}^{-1}$, G_{ATM} $4.3 \pm 0.1 \text{ GtC yr}^{-1}$, S_{OCEAN} $2.6 \pm 0.5 \text{ GtC yr}^{-1}$, and S_{LAND} $2.6 \pm 0.8 \text{ GtC yr}^{-1}$. For year 2012 alone, E_{FF} grew to $9.7 \pm 0.5 \text{ GtC yr}^{-1}$, 2.2% above 2011, reflecting a continued trend in these emissions; G_{ATM} was $5.2 \pm 0.2 \text{ GtC yr}^{-1}$, S_{OCEAN} was $2.9 \pm 0.5 \text{ GtC yr}^{-1}$, and assuming and E_{LUC} of $0.9 \pm 0.5 \text{ GtC yr}^{-1}$ (based on 2001–2010 average), S_{LAND} was $2.5 \pm 0.9 \text{ GtC yr}^{-1}$. G_{ATM} was high in 2012 compared to

ESSDD

6, 689–760, 2013

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the 2003–2012 average, almost entirely reflecting the high E_{FF} . The global atmospheric CO_2 concentration reached 392.52 ± 0.10 ppm on average over 2012. We estimate that E_{FF} will increase by 2.1 % (1.1–3.1 %) to 9.9 ± 0.5 GtC in 2013, 61 % above emissions in 1990, based on projections of World Gross Domestic Product and recent changes in the carbon intensity of the economy. With this projection, cumulative emissions of CO_2 will reach about 550 ± 60 GtC for 1870–2013, 70 % from E_{FF} (390 ± 20 GtC) and 30 % from E_{LUC} (160 ± 55 GtC). This paper is intended to provide a baseline to keep track of annual carbon budgets in the future.

All data presented here can be downloaded from the Carbon Dioxide Information Analysis Center (doi:10.3334/CDIAC/GCP_2013_v1.1).

1 Introduction

The concentration of carbon dioxide (CO_2) in the atmosphere has increased from approximately 277 parts per million (ppm) in 1750 (Joos and Spahni, 2008), the beginning of the Industrial Era, to 392.52 in 2012 (Dlugokencky and Tans, 2013). Daily averages went above 400 ppm for the first time at Mauna Loa station in May 2013 (Scripps, 2013). This station holds the longest running record of direct measurements of atmospheric CO_2 concentration (Tans and Keeling, 2013). The atmospheric CO_2 increase above preindustrial levels was caused initially by the release of carbon to the atmosphere from deforestation and other land-use change activities (Ciais et al., 2013). Emissions from fossil-fuel combustion started before the Industrial Era and became the dominant source of anthropogenic emissions to the atmosphere from around 1920 until present. Anthropogenic emissions occur on top of an active natural carbon cycle that circulates carbon between the atmosphere, ocean, and terrestrial biosphere reservoirs on time scales from days to millennia, while geologic reservoirs have even longer timescales (Archer et al., 2009).

The global carbon budget presented here refers to the mean, variations, and trends in the anthropogenic perturbation of CO_2 in the atmosphere, referenced to the

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



beginning of the Industrial Era. It quantifies the input of CO₂ to the atmosphere by emissions from human activities, the growth of CO₂ in the atmosphere, and the resulting changes in land and ocean carbon fluxes in response to increasing atmospheric CO₂ levels, climate change and climate variability, and other anthropogenic and natural changes. An understanding of this perturbation budget over time and the underlying variability and trends of the natural carbon cycle are necessary to understand and quantify climate-carbon feedbacks.

The components of the CO₂ budget that are reported in this paper include separate estimates for (1) the CO₂ emissions from fossil-fuel combustion and cement production (E_{FF}), (2) the CO₂ emissions resulting from deliberate human activities on land leading to Land-Use Change (LUC; E_{LUC}), (3) the growth rate of CO₂ in the atmosphere (G_{ATM}), and the uptake of CO₂ by the “CO₂ sinks” in (4) the ocean (S_{OCEAN}) and (5) on land (S_{LAND}). The CO₂ sinks as defined here include the response of the land and ocean to elevated CO₂ and changes in climate and other environmental conditions. The global emissions and their partitioning among the atmosphere, ocean and land are in balance:

$$E_{FF} + E_{LUC} = G_{ATM} + S_{OCEAN} + S_{LAND} \quad (1)$$

G_{ATM} is usually reported in ppm, and we convert to units of carbon mass using 1 ppm = 2.120 GtC (Joos et al., 2013) (Table 1). We also include a quantification of E_{FF} by country, both computed with territorial and consumption based accounting (see Methods).

Equation (1) partly omits two kinds of processes. The first is the net input of CO₂ to the atmosphere from the chemical oxidation of reactive carbon-containing gases from sources other than fossil-fuels (e.g. landfills, industrial processes, etc), primarily methane (CH₄), carbon monoxide (CO), and volatile organic compounds such as isoprene and terpene. The second is the anthropogenic perturbation to carbon cycling in terrestrial freshwaters, estuaries, and coastal areas, that modify lateral fluxes transported from land ecosystems to the open ocean, the evasion CO₂ flux from rivers, lakes

and estuaries to the atmosphere, and the net air-sea anthropogenic CO₂ flux of coastal areas (Regnier et al., 2013). These flows are omitted in absence of annual information on the natural versus anthropogenic terms of these loops of the carbon cycle, and they are discussed in Sect. 2.6. The inclusion of these fluxes of anthropogenic CO₂ would affect the estimates of, and partitioning between, S_{LAND} and S_{OCEAN} in Eq. (1) in complementary ways, but would not affect the other terms in Eq. (1).

The CO₂ budget has been assessed by the Intergovernmental Panel on Climate Change (IPCC) in all assessment reports (Ciais et al., 2013; Denman et al., 2007; Prentice et al., 2001; Schimel et al., 1995; Watson et al., 1990), and by others (e.g. Balantyne et al., 2012). These assessments included budget estimates for the decades of the 1980s, 1990s (Denman et al., 2007) and, most recently, the period 2002–2011 (Ciais et al., 2013). The IPCC methodology has been adapted and used by the Global Carbon Project (GCP, www.globalcarbonproject.org), who have coordinated a cooperative community effort for the annual publication of global carbon budgets up to year 2005 (Raupach et al., 2007; including fossil emissions only), year 2006 (Canadell et al., 2007), year 2007 (published online; GCP, 2007), year 2008 (Le Quéré et al., 2009), year 2009 (Friedlingstein et al., 2010), year 2010 (Peters et al., 2012b), and most recently, year 2011 (Le Quéré et al., 2013; Peters et al., 2013). Each of these papers updated previous estimates with the latest available information for the entire time series. From 2008, these publications projected fossil-fuel emissions for one additional year using the projected World Gross Domestic Product and estimated improvements in the carbon intensity of the economy.

We adopt a range of ± 1 standard deviation (σ) to report the uncertainties in our annual estimates, representing a likelihood of 68% that the true value lies within the provided range if the errors have a Gaussian distribution. This choice reflects the difficulty of characterising the uncertainty in the CO₂ fluxes between the atmosphere and the ocean and land reservoirs individually, particularly on an annual basis, as well as the difficulty to update the CO₂ emissions from LUC. A 68% likelihood provides an indication of our current capability to quantify each term and its uncertainty given

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the available information. For comparison, the Fifth Assessment Report of the IPCC (AR5) generally reported 90 % likelihood for large datasets whose uncertainty is well characterised, or for long time intervals less affected by year-to-year variability. Our 68 % uncertainty value is near the 66 % that the IPCC characterises as “likely” for values falling into the $\pm 1\sigma$ interval. The uncertainties reported here combine statistical analysis of the underlying data and expert judgement of the likelihood of results lying outside this range. The limitations of current information are discussed in the paper.

All units are presented in gigatonnes of carbon (GtC, 10^{15} gC), which is the same as petagrams of carbon (PgC; Table 1). Units of gigatonnes of CO_2 (or billion tonnes of CO_2) used in policy are equal to 3.664 multiplied by the value in units of GtC.

This paper provides a detailed description of the datasets and methodology used to compute the global carbon budget estimates for the period preindustrial (1750) to 2012 and in more detail for the period 1959 to 2012. We also provide decadal averages starting in 1960 including the last decade (2003–2012), results for the year 2012, and a projection of E_{FF} for year 2013. Finally we provide the total or cumulative emissions from fossil-fuels and land-use change since year 1750, the pre-industrial period, and since year 1870, the reference year for the cumulative carbon estimate used by the Fifth Assessment Report of the Intergovernmental Panel on Climate Change based on the availability of global temperature data (Stocker et al., 2013). It is intended that this paper will be updated every year using the format of “living data”, to help keep track of new versions of the budget that result from new data, revision of data, and changes in methodology. Additional materials associated with the release of each new version will be posted at the Global Carbon Project (GCP) website (<http://www.globalcarbonproject.org/carbonbudget>). With this approach, we aim to provide transparency and traceability in reporting indicators and drivers of climate change.

2 Methods

The original measurements and data used to complete the global carbon budget are generated by multiple organizations and research groups around the world. The effort presented here is thus mainly one of synthesis, where results from individual groups are collated, analysed and evaluated for consistency. We facilitate access to original data with the understanding that that primary datasets will be referenced in future work (see Table 2 for “How to cite” the datasets). Descriptions of the measurements, models, and methodologies follow below and in depth descriptions of each component are described elsewhere (e.g. Andres et al., 2012; Houghton et al., 2012).

This is the second revised version of the “global carbon budget”. It is an update of Le Quéré et al. (2013), including data until year 2012 and a projection for fossil-fuel emissions for year 2013. The main changes from Le Quéré et al. (2013) are: (1) we have introduced a new section (Sect. 2.6) that describes and quantifies the main missing processes; (2) we have introduced data-products to assess the interannual variability in the ocean CO₂ sink; (3) we have introduced a confidence level to characterise the annual estimates from each term based on the type, amount, quality and consistency of the evidence as defined by the IPCC (Stocker et al., 2013); and (4) we now also update the cumulative CO₂ emissions.

2.1 CO₂ emissions from fossil-fuel combustion and cement production (E_{FF})

2.1.1 Fossil-fuel and cement emissions and their uncertainty

The calculation of global and national CO₂ emissions from fossil-fuel combustion, including gas flaring and cement production (E_{FF}), relies primarily on energy consumption data, specifically data on hydrocarbon fuels, collated and archived by several organisations (Andres et al., 2012). These include the Carbon Dioxide Information Analysis Center (CDIAC), the International Energy Agency (IEA), the United Nations (UN), and the United States Department of Energy (DoE) Energy Information

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Administration (EIA). We use the emissions estimated by the CDIAC (Boden et al., 2013). The CDIAC emission estimates also extend back in time to 1751 with consistent and well-documented emissions from all fossil-fuel combustion, cement production, and gas flaring for all countries and their uncertainty (Andres et al., 1999, 2012); this makes the dataset a unique resource for research of the carbon cycle during the fossil-fuel era. During the period 1959–2010, the emissions are based primarily on energy data provided by the UN Statistics Division (Table 3; UN, 2013a, b). For the most recent two years (2011 and 2012) when the UN statistics are not yet available, we generate preliminary estimates based on the BP annual energy review for extrapolation of emissions in 2011 and 2012 (BP, 2013). BP's sources for energy statistics overlap with those of the UN data, but are compiled more rapidly using a smaller group of mostly developed countries and assumptions for missing data. We use the BP values only for the year-to-year rate of change, because the rates of change are less uncertain than the absolute values and to avoid discontinuities in the time-series when linking the UN-based energy data (up to 2010) with the BP energy data (2011 and 2012). These preliminary estimates are replaced by the more complete CDIAC data based on UN statistics when they become available. Past experience shows that projections based on the BP rate of change provide reliable estimates for the two most recent years when full data are not yet available from the UN (see Sect. 3.2 and Supplementary Information from Peters et al., 2013).

When necessary, fuel masses/volumes are converted to fuel energy content using coefficients provided by the UN and then to CO₂ emissions using conversion factors that take into account the relationship between carbon content and heat content of the different fuel types (coal, oil, gas, gas flaring) and the combustion efficiency (to account, for example, for soot left in the combustor or fuel otherwise lost or discharged without oxidation). Most data on energy consumption and fuel quality are available at the level of countries. In general, CO₂ emissions for equivalent primary energy consumption are about 30 % higher for coal compared to oil, and 70 % higher for coal compared to natural gas (Marland et al., 2007). All estimated fossil-fuel emissions are based on the

ESSDD

6, 689–760, 2013

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



mass flows of carbon and assume that the fossil carbon emitted as CO or CH₄, will soon be oxidized to CO₂ in the atmosphere and hence count the carbon mass with CO₂ emissions.

Emissions from cement production are based on cement data from the US Geological Survey (van Oss, 2013) up to year 2010, and from preliminary data for 2011 and 2012 where available (US Geological Survey, 2013). Some fraction of the CaO and MgO in cement is returned to the carbonate form during cement weathering but this is generally regarded to be small and is ignored here.

Emission estimates from gas flaring are calculated in a similar manner as those from solid, liquid, and gaseous fuels, and rely on the UN Energy Statistics to supply the amount of flared or vented fuel. For emission years 2011 and 2012, flaring is assumed constant from the emission year 2010 UN-based data. The basic data on gas flaring report atmospheric losses during petroleum production and processing, have large uncertainty and do not distinguish between gas that is flared as CO₂ or vented as CH₄. Fugitive emissions of CH₄ from the so-called upstream sector (coal mining and natural gas distribution, for examples) are not included in the accounts of CO₂ emissions except to the extent that they are captured in the UN energy data and counted as gas “flared or lost”.

The published CDIAC dataset has 250 countries and regions included. This expanded list includes countries that no longer exist, such as the USSR or East Pakistan. For the budget, we reduce the list to 219 countries by reallocating emissions to the currently defined territories. This involved both aggregation and disaggregation, and does not change global emissions. Examples of aggregation include merging East and West Germany to the currently defined Germany. Examples of disaggregation include real-locating the emissions from former USSR to the resulting independent countries. For disaggregation, we use the emission shares when the current territory first appeared. Most recent two years, 2011 and 2012, the BP statistics are more aggregated, but we retain the detail in CDIAC by applying the same growth rates to individual countries in CDIAC as in the aggregated regions in the BP dataset.

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Estimates of CO₂ emissions show that the global total of emissions is not equal to the sum of emissions from all countries. This is largely attributable to emissions that occur in international territory, in particular the combustion of fuels used in international shipping and aviation (bunker fuels), where the emissions are included in the global totals but are not attributed to individual countries. In practice, the emissions from international bunker fuels are calculated based on where the fuels were loaded, but they are not included with national emissions estimates. Smaller differences occur because globally the sum of imports in all countries is not equal to the sum of exports and because of differing treatment of oxidation of non-fuel uses of hydrocarbons (e.g. as solvents, lubricants, feedstocks, etc.), and changes in stocks.

The uncertainty of the annual fossil-fuel and cement emissions for the globe has been estimated at $\pm 5\%$ (scaled down from the published $\pm 10\%$ at ± 2 sigma to the use of ± 1 sigma bounds reported here; Andres et al., 2012). This includes an assessment of uncertainties in the amounts of fuel consumed, the carbon contents of fuels, and the combustion efficiency. While in the budget we consider a fixed uncertainty of $\pm 5\%$ for all years, in reality the uncertainty, as a percentage of the emissions, is growing with time because of the larger share of global emissions from non-Annex B countries with less precise statistical systems (Marland et al., 2009). For example, the uncertainty in Chinese emissions has been estimated at around $\pm 10\%$ (for ± 1 sigma; Gregg et al., 2008). Generally, emissions from mature economies with good statistical bases have an uncertainty of only a few per cent (Marland, 2008). Further research is needed before we can quantify the time evolution of the uncertainty. We assign a medium confidence to the results presented here because they are based on indirect estimates of emissions using energy data (Durant et al., 2010).

2.1.2 Emissions embodied in goods and services

National emissions inventories take a territorial (production) perspective and “include greenhouse gas emissions and removals taking place within national territory and offshore areas over which the country has jurisdiction” (Rypdal et al., 2006). That is,

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



emissions are allocated to the country where and when the emissions actually occur. The territorial emission inventory of an individual country does not include the emissions from the production of goods and services produced in other countries (e.g. food and clothes) that are used for consumption. Consumption-based emission inventories for an individual country allocate global emissions to products that are consumed within a country, and are conceptually calculated as the territorial emissions less the territorial emissions to produce exported products plus the emissions in other countries to produce imported products (Consumption = Territorial – Exports + Imports). The difference between the territorial- and consumption-based emission inventories is the net transfer (exports minus imports) of emissions from the production of internationally traded products. Consumption-based emission inventories (e.g. Davis and Caldeira, 2010) provide additional information to territorial inventories that can be used to understand emission drivers (Hertwich and Peters, 2009), quantify emission leakages between countries (Peters et al., 2011b) and potentially design more effective and efficient climate policy (Peters and Hertwich, 2008).

We estimate consumption-based emissions by enumerating the global supply chain using a global model of the economic relationships between sectors within and between every country (Andrew and Peters, 2013; Peters et al., 2011a). Due to availability of the input data, detailed estimates are made for the years 1997, 2001, 2004, and 2007 (using the methodology of Peters et al., 2011b) using economic and trade data from the Global Trade and Analysis Project version 8.1 (GTAP; Narayanan et al., 2013). The results cover 57 sectors and 134 countries and regions. The results are extended into an annual time-series from 1990 to the latest year of the fossil-fuel emissions or GDP data (2011 in this budget), using GDP data by expenditure in current USD (from the UN National Accounts main Aggregates database; UN, 2013c) and time series of trade data from GTAP (based on the methodology in Peters et al., 2011b).

The consumption-based emission inventories in this carbon budget incorporate several improvements over previous versions (Le Quéré et al., 2013; Peters et al., 2011b, 2012b). The detailed estimates for 2004 and 2007 and time series approximation from

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



1990–2011 are based on an updated version of the GTAP database (Narayanan et al., 2013). We estimate the sector level CO₂ emissions using our own calculations based on the GTAP data and methodology, include flaring and cement emissions from CDIAC, and then scale the national totals to match the CDIAC estimates from the most recent carbon budget. We do not include international transportation in our estimates of national totals, but include them in the global total. The time-series of trade data provided by GTAP covers the period 1995–2009 and our methodology uses the trade shares of this dataset. For the period 1990–1994 we assume the trade shares of 1995, while in 2010 and 2011 we assume the trade shares of 2008 since 2009 was heavily affected by the global financial crisis. We identified errors in the trade shares of Taiwan in 2008 and 2009, so the trade shares for 2008–2010 are based on the 2007 trade shares.

We do not provide an uncertainty estimate for these emissions, but based on model comparisons and sensitivity analysis, they are unlikely to be larger than for the territorial emission estimates (Peters et al., 2012a). Uncertainty is expected to increase for more detailed results, and to decrease with aggregation (Peters et al., 2011b; e.g. the results for Annex B will be more accurate than the sector results for an individual country).

The consumption-based emissions consider the carbon emitted to the atmosphere in the production of products, but not the trade in fossil fuels (coal, oil, gas). It is also possible to account for the carbon trade in fossil fuels (Davis et al., 2011), but we do not present that data here. Peters et al. (2012a) additionally consider trade in biomass.

The consumption data do not contribute to the global average terms in Eq. (1), but are relevant to the anthropogenic carbon cycle as they reflect the movement of carbon across the Earth's surface in response to human needs (both physical and economic). Furthermore, if national and international climate policies continue to develop in an unharmonised way, then the trends reflected in these data will need to be accommodated by those developing policies.

2.1.3 Growth rate in emissions

We report the annual growth rate in emissions for adjacent years (in percent per year) by calculating the difference between the two years and then comparing to the emissions in the first year: $\left[\frac{E_{FF}(t_{0+1}) - E_{FF}(t_0)}{E_{FF}(t_0)} \right] \times 100 / (1 \text{ yr})$. This is the simplest method to characterise a one-year growth compared to the previous year and is in widespread usage. We do not apply a leap-year adjustment, which could affect the growth rate by about $0.3\% \text{ yr}^{-1}$ ($1/365.25$).

The growth rate of E_{FF} over time periods of greater than one year can be re-written using its logarithm equivalent as follows:

$$\frac{1}{E_{FF}} \frac{dE_{FF}}{dt} = \frac{d(\ln E_{FF})}{dt} \quad (2)$$

Here we calculate growth rates in emissions for multi-year periods (e.g. a decade) by fitting a linear trend to $\ln(E_{FF})$ in Eq. (2), reported in percent per year. We fit the logarithm of E_{FF} rather than E_{FF} directly because this method ensures that computed growth rates satisfy Eq. (6). This method differs from previous papers (Canadell et al., 2007; Le Quéré et al., 2009; Raupach et al., 2007) who computed the fit to E_{FF} and divided by average E_{FF} directly, but the difference is very small ($<0.05\%$) in the case of E_{FF} .

2.1.4 Emissions projections using GDP

Energy statistics are normally available around June for the previous year. We use the close relationship between the growth in world Gross Domestic Product (GDP) and the growth in global emissions (Raupach et al., 2007) to project emissions for the current year. This is based on the so-called Kaya (also called IPAT) identity, whereby E_{FF} is decomposed by the product of GDP and the fossil-fuel carbon intensity of the economy

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(I_{FF}) as follows:

$$E_{FF} = \text{GDP} \times I_{FF} \quad (3)$$

Such product-rule decomposition identities imply that the growth rates of the multiplied quantities are additive. Taking a time derivative of Eq. (3) gives:

$$5 \quad \frac{dE_{FF}}{dt} = \frac{d(\text{GDP} \times I_{FF})}{dt} \quad (4)$$

and applying the rules of calculus:

$$\frac{dE_{FF}}{dt} = \frac{d\text{GDP}}{dt} \times I_{FF} + \text{GDP} \times \frac{dI_{FF}}{dt} \quad (5)$$

finally, dividing Eq. (4) by Eq. (2) gives:

$$\frac{1}{E_{FF}} \frac{dE_{FF}}{dt} = \frac{1}{\text{GDP}} \frac{d\text{GDP}}{dt} + \frac{1}{I_{FF}} \frac{dI_{FF}}{dt} \quad (6)$$

10 where the left hand term is the relative growth rate of E_{FF} , and the right hand terms are the relative growth rates of GDP and I_{FF} , respectively, which can simply be added linearly to give overall growth rate. The growth rates are reported in percent by multiplying each term by 100. As preliminary estimates of annual change in GDP are made well before the end of a calendar year, making assumptions on the growth rate of I_{FF}
15 allows us to make projections of the annual change in CO₂ emissions well before the end of a calendar year.

2.2 CO₂ emissions from land use, land-use change and forestry (E_{LUC})

LUC emissions reported in the 2013 carbon budget (E_{LUC}) include CO₂ fluxes from deforestation, afforestation, logging (forest degradation and harvest activity), shifting

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



cultivation (cycle of cutting forest for agriculture then abandoning), regrowth of forests following wood harvest or abandonment of agriculture, fire-based peatland emissions and other land management practices (Table 4). Some of these processes lead to emissions of CO₂ to the atmosphere, while others lead to CO₂ sinks. E_{LUC} is the net sum of all processes considered. Our annual estimate for 1959–2010 is from a bookkeeping method (Sect. 2.2.1) primarily based on forest area change and biomass data from the Forest Resource Assessment (FRA) of the Food and Agriculture Organisation (FAO) published at intervals of five years, as revised by Houghton and Hackler (2013). E_{LUC} for year 2011 is based upon E_{LUC} averaged over 2001–2010 from the bookkeeping method and E_{LUC} anomalies based on fire emissions in deforested areas, as in Le Quéré et al. (2013) (Sect. 2.2.2). Fire emissions were not available for year 2012. E_{LUC} for 2012 is thus assigned the mean of 2001–2010 (last decade where the bookkeeping method is available) and a low confidence. In addition, we use results from eight Dynamic Global Vegetation Models (see Sect. 2.2.3 and Table 5) that calculate net LUC CO₂ emissions in response to observed land cover change prescribed to each model, to help quantify the uncertainty in E_{LUC} , and to explore the consistency of our understanding. Compared to E_{LUC} provided in Le Quéré et al. (2013; Sect. 2.2.2), the revised estimate of the bookkeeping method directly includes emissions from peat and is used through year 2010. The three methods are described below, and differences are discussed in Sect. 3.2.

2.2.1 Bookkeeping method

LUC CO₂ emissions are calculated by a bookkeeping model approach (Houghton, 2003) that keeps track of the carbon stored in vegetation and soils before deforestation or other land-use change, and the changes in forest age classes, or cohorts, of disturbed lands after land-use change. It tracks the CO₂ emitted to the atmosphere over time due to decay of soil and vegetation carbon in different pools, including wood products pools after logging and deforestation. It also tracks the regrowth of vegetation and build-up of soil carbon pools following land-use change. It considers transitions

**Global carbon budget
2013**

 C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


between forests, pastures and cropland, shifting cultivation, degradation of forests where a fraction of the trees is removed, abandonment of agricultural land, and forest management such as logging and fire management. In addition to tracking logging debris on the forest floor, the bookkeeping model tracks the fate of carbon contained in harvested wood products that is eventually emitted back to the atmosphere as CO₂, although a detailed treatment of the lifetime in each product pool is not performed (Earles et al., 2012). Harvested wood products are partitioned into three pools with different turnover times. All fuel-wood is assumed to be burned in the year of harvest (1.0 yr⁻¹). Pulp and paper products are oxidized at a rate of 0.1 yr⁻¹. Timber is assumed to be oxidized at a rate of 0.01 yr⁻¹, and elemental carbon decays at 0.001 yr⁻¹. The general assumptions about partitioning wood products among these pools are based on national harvest data.

Here we use the updated estimate of Houghton and Hackler (2013) based on better and more recent data in three ways. First, regions outside the tropics were updated, for the first time since 1990 in some cases. As a result, nearly all regions outside the tropics show a small net sink from LUC after ~1980. Second, emissions of carbon from draining and burning of peatlands in SE Asia were included. The practice is not thought to have been important before around 1985. Finally, a greater number of forest types that could be deforested or logged were included, which improved the accounting for variations in biomass density associated with any land use (Houghton and Hackler, 2013). The revised estimate produces higher emissions during 1850–1950 compared to Houghton et al. (2012) due to the revision of historical deforestation in Australia and New Zealand based on Bradshaw (2012).

The primary land cover change and biomass data for the bookkeeping model analysis is the Forest Resource Assessment of the FAO published at intervals of five years (FAO, 2010), which is based on countries' self-reporting of statistics on forest cover change and management partially combined with satellite data in more recent assessments (Table 3). Changes in land cover other than forest are based on annual, national changes in cropland and pasture areas reported by the FAO Statistics Division

(FAOSTAT, 2010). The LUC data set is non-spatial and aggregated by regions. The carbon stocks on land (biomass and soils), and their response functions subsequent to LUC, are based on averages per land cover type, per biome and per region. Similar results were obtained using forest biomass carbon density based on satellite data (Baccini et al., 2012). The bookkeeping model does not include land ecosystems' transient response to changes in climate, atmospheric CO₂ and other environmental factors, but the growth/decay curves are based on contemporary data that will implicitly reflect the effects of CO₂ and climate at that time. Results from the bookkeeping method are available from 1850 to 2010.

2.2.2 Fire-based method

LUC associated CO₂ emissions calculated from satellite-based fire activity in deforested areas (van der Werf et al., 2010) provide information that is complementary to the bookkeeping approach. They do not provide a direct estimate of E_{LUC} as they do not include non-combustion processes such as respiration, wood harvest, wood products or forest regrowth, or legacy emissions such as decomposition from on-ground debris or soils are missed by this method. They do however provide insight on the year-to-year variations in E_{LUC} that result from the interactions between climate and human activity (e.g. there is more burning and clearing of forests in dry years) that are not as well represented by other methods. The “deforestation fire emissions” assume an important role of fire in removing biomass in the deforestation process, and thus can be used to infer direct CO₂ emissions from deforestation using satellite-derived data on fire activity in regions with active deforestation. The method requires information on the fraction of total area burned associated with deforestation versus other types of fires, and can be merged with information on biomass stocks and the fraction of the biomass lost in a deforestation fire to estimate CO₂ emissions. The satellite-based fire emissions are limited to the tropics, where fires result mainly from human activities. Tropical deforestation is the largest and most variable single contributor to E_{LUC} .

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global carbon budget
2013**

 C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Burned area from (Giglio et al., 2010) is merged with active fire retrievals to mimic more sophisticated assessments of deforestation rates in the pan-tropics (van der Werf et al., 2010). This information is used as input data in a modified version of the satellite-driven CASA biogeochemical model to estimate carbon emissions, keeping track of what fraction was due to deforestation (van der Werf et al., 2010). The CASA model uses different assumptions to compute decay functions compared to the bookkeeping model, and does not include historical emissions or regrowth from land-use change prior to the availability of satellite data. Comparing coincident CO emissions and their atmospheric fate with satellite-derived CO concentrations allows for some validation of this approach (e.g. van der Werf et al., 2008). Results from the fire-based method are available from 1997 to 2011 only. Le Quéré et al. (2013) used annual estimates from the Global Fire Emissions Database (GFED3; available from <http://www.globalfiredata.org>) to quantify the mean emissions over five years with the satellite-based method to distribute these emissions annually up to 2011. It thus assumed that all land management activities apart from deforestation do not vary significantly on a year-to-year basis.

2.2.3 Dynamic Global Vegetation Models (DGVMs)

LUC CO₂ emissions have been estimated using an ensemble of eight DGVMs, coordinated by the project “Trends and drivers of the regional-scale sources and sinks of carbon dioxide (Trendy)”. These DGVMs were forced with historical changes in land cover distribution, climate, atmospheric CO₂ concentration, and N deposition. As further described below, each historical DGVM simulation was repeated with a time-invariant pre-industrial land cover distribution, allowing to estimate, by difference with the first simulation, the dynamic evolution of biomass and soil carbon pools in response to prescribed land cover change (Tables 4 and 5). All DGVMs represent deforestation and (to some extent) regrowth, the most important components of E_{LUC} , but they do not represent all processes resulting directly from human activities on land (Table 4). DGVMs represent processes of vegetation establishment, growth, mortality and decomposition associated with natural cycles and include the vegetation and soil response to

increasing atmospheric CO₂ levels, to climate variability and change, in addition to atmospheric N deposition in the presence of nitrogen limitation (in four models; Table 4). The DGVMs are independent from the other budget terms except for their use of atmospheric CO₂ concentration to calculate the fertilization effect of CO₂ on primary production.

The DGVMs used a consistent land-use change dataset (Hurtt et al., 2011), which provided annual, half-degree, fractional data on cropland, pasture, primary vegetation and secondary vegetation, as well as all underlying transitions between land-use states, including wood harvest and shifting cultivation. This dataset used the HYDE (Klein Goldewijk et al., 2011) spatially gridded maps of cropland, pasture, and ice/water fractions of each grid cell as an input. The HYDE data is based on annual FAO statistics of change in agricultural area (FAOSTAT, 2010). For the year 2012, the HYDE dataset was extrapolated from 2011, based on the trend in agricultural area over the previous 5 yr. The HYDE dataset is independent from the data set used in the bookkeeping method (Houghton, 2003 and updates), which is based on forest area change statistics (FAO, 2010). Although the land-use change datasets indicate whether land-use changes occur on forested or non-forested land, the changes in agricultural areas are then implemented differently within each model (for instance, an increased cropland fraction in a grid cell can either be at the expense of grassland, or forest, the latter resulting in deforestation; land cover fractions of the non-agricultural land differ between models). Similarly, model-specific assumptions are also applied for the conversion of wood harvest mass or area and other product pools into carbon in some models (Table 4).

The DGVM model runs were forced by observed monthly temperature, precipitation, and cloud cover fields, provided on a 0.5° × 0.5° grid and updated to 2012 by the Climatic Research Unit (Harris et al., 2013a, b). The forcings include both observed climate change and change in atmospheric CO₂ (Dlugokencky and Tans, 2013), and in some models N deposition (Lamarque et al., 2010). E_{LUC} is diagnosed in each model by the difference between a model simulation with prescribed historical land cover change and

**Global carbon budget
2013**

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

a simulation with constant, pre-industrial land cover distribution. Both simulations were driven by changing atmospheric CO₂, climate, and in some models N deposition over the period 1860–2012. Using the difference between these two DGVM simulations to diagnose E_{LUC} is not entirely consistent with the definition of E_{LUC} in the bookkeeping model (Gasser and Ciais, 2013). The DGVM approach should produce systematically higher E_{LUC} emissions than the bookkeeping approach if all the parameters of the two approaches were the same. Here, given the different input data of DGVMs and bookkeeping, this systematic difference cannot be quantified.

2.2.4 Uncertainty assessment for E_{LUC}

Differences between the bookkeeping, fire-based and DGVM methods originate from three main sources: the land cover change data set, different approaches in models, and in the different processes represented (Table 4). We examine the results from the eight DGVM models and of the bookkeeping methods to assess the uncertainty in E_{LUC} . The standard deviation across models in each year ranged from 0.3 to 0.9 GtC yr⁻¹, with an average of 0.5 GtC yr⁻¹ from 1960 to 2012 (Table 6). The multi-model mean is higher than the bookkeeping estimate used in the budget with a mean absolute difference of 0.3 GtC for 1960 to 2010. The multi-model mean and bookkeeping method differ by less than 0.5 GtC yr⁻¹ over 80 % of the time. Based on this comparison, we assess that an uncertainty of ± 0.5 GtC yr⁻¹ provides a semi-quantitative measure of uncertainty for annual emissions, and reflects our best value judgment that there is at least 68 % chance (± 1 sigma) that the true LUC emission lies within the given range, for the range of processes considered here. This is consistent with the analysis of Houghton et al. (2012), which partly reflects improvements in data on forest area change using satellite data, and partly more complete understanding and representation of processes in models.

The uncertainties in the decadal mean estimates from the DGVM ensemble are likely correlated between decades. They come from (1) system boundaries (e.g. not counting forest degradation in some models); (2) definition issues when calculating E_{LUC} from

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



the difference of simulations with and without LUC, which cause a bias compared to the bookkeeping estimates that makes decadal uncertainty estimates perfectly correlated (Gasser and Ciais, 2013); (3) common and uncertain land cover change input data which cause a bias, though if a different input dataset is used each decade, decadal fluxes from DGVMs may be partly decorrelated; (4) model structural errors (e.g. errors in biomass stocks), which cause bias that correlate decadal estimates. In addition, errors arising from uncertain DGVM parameter values would be random but they are not accounted for in this study, since no DGVM provided an ensemble of runs with perturbed parameters.

Prior to 1959, the uncertainty in E_{LUC} is taken as $\pm 33\%$, which is the ratio of uncertainty to mean from the 1960s (Table 6), the first decade available. This ratio is consistent with the mean standard deviation of DGVMs LUC emissions over 1870–1958 (0.37 GtC) over the multi-model mean (1.1 GtC).

2.3 Atmospheric CO₂ growth rate (G_{ATM})

Global atmospheric CO₂ growth rate estimates

The atmospheric CO₂ growth rate is provided by the US National Oceanic and Atmospheric Administration Earth System Research Laboratory (Dlugokencky and Tans, 2013), which is updated from Ballantyne et al. (2012). For the 1959–1980 period, the global growth rate is based on measurements of atmospheric CO₂ concentration averaged from the Mauna Loa and South Pole stations, as observed by the CO₂ Program at Scripps Institution of Oceanography (Keeling et al., 1976). For the 1980–2012 time period, the global growth rate is based on the average of multiple stations selected from the marine boundary layer sites (Ballantyne et al., 2012), after fitting each station with a smoothed curve as a function of time, and averaging by latitude band (Masarie and Tans, 1995). The annual growth rate is estimated from atmospheric CO₂ concentration by taking the average of the most recent December–January months corrected for the average seasonal cycle and subtracting this same average one year earlier. The growth

rate in units of ppm yr^{-1} is converted to fluxes by multiplying by a factor of 2.120 GtC per ppm (Joos et al., 2013) for comparison with the other components.

The uncertainty around the annual growth rate based on the multiple stations dataset ranges between 0.11 and 0.72 GtC yr^{-1} , with a mean of 0.60 GtC yr^{-1} for 1959–1980 and 0.19 GtC yr^{-1} for 1980–2012, when a larger set of stations were available. It is based on the number of available stations, and thus takes into account both the measurement errors and data gaps at each station. This uncertainty is larger than the uncertainty of $\pm 0.1 \text{ GtC yr}^{-1}$ reported for decadal mean growth rate by the IPCC because errors in annual growth rate are strongly anti-correlated in consecutive years leading to smaller errors for longer time scales. The decadal change is computed from the difference in concentration ten years apart based on measurement error of 0.35 ppm (based on offsets between NOAA/ESRL measurements and those of the World Meteorological Organization World Data Center for Greenhouse Gases (NOAA/ESRL, 2013) for the start and end points (the decadal change uncertainty is the $\sqrt{2 \times (0.35 \text{ ppm})^2}$)/10 yr assuming that each yearly measurement error is independent). This uncertainty is also used in Table 6.

The contribution of anthropogenic CO and CH₄ is neglected from the global anthropogenic CO₂ budget (see Sect. 2.6.1). We assign a high confidence to the annual estimates of G_{ATM} because they are based on direct measurements.

In order to estimate the total carbon accumulated in the atmosphere since 1750 or 1870, we use an atmospheric CO₂ concentration of $277 \pm 3 \text{ ppm}$ or $288 \pm 3 \text{ ppm}$, respectively, based on a cubic spline fit to ice core data (Joos and Spahni, 2008). The uncertainty of $\pm 3 \text{ ppm}$ (converted to $\pm 1\sigma$) is taken directly from the IPCC's assessment (Ciais et al., 2013). Typical uncertainties in the atmospheric growth rate from ice core data are $\pm 1\text{--}1.5 \text{ GtC per decade}$ as evaluated from the Law Dome data (Etheridge et al., 1996) for individual 20-year intervals over the period from 1870 to 1960 (Bruno and Joos, 1997).

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



2.4 Ocean CO₂ sink

The global ocean CO₂ sink is based on a combination of a mean CO₂ sink estimates for the 1990s from observations, and a trend in the ocean CO₂ sink for 1959–2012 from six global ocean biogeochemistry models. Data products that estimate the annual CO₂ sink are beginning to emerge. These are used here for the first time to provide a qualitative assessment of confidence in the reported results.

2.4.1 Data-based estimates

A mean ocean CO₂ sink of $2.2 \pm 0.4 \text{ GtC yr}^{-1}$ for the 1990s was estimated by the IPCC (Denman et al., 2007) based on three data-based methods: direct ocean/land CO₂ sink partitioning from observed atmospheric O₂/N₂ concentration trends (Manning and Keeling, 2006), an oceanic inversion method constrained by ocean biogeochemistry data (Mikaloff Fletcher et al., 2006), and a method based on penetration time scale for CFCs (McNeil et al., 2003). This is comparable with the sink of $2.0 \pm 0.5 \text{ GtC yr}^{-1}$ estimated by Khatiwala et al. (2013) for the 1990s, and with the sink of 1.9 to 2.5 estimated from a range of methods for the period 1990–2009 (Wanninkhof et al., 2013), with uncertainties ranging from ± 0.3 to $\pm 0.7 \text{ GtC yr}^{-1}$.

The interannual variability in the ocean CO₂ sink was estimated for 1990–2011 by Rödenbeck et al. (2013 updated version “oc_v1.1”) using an inversion method based on observed oceanic partial pressure of CO₂ ($p\text{CO}_2$) derived from the Surface Ocean Carbon Atlas (SOCAT v2; Bakker et al., 2013; Pfeil et al., 2013). The interannual variability in ocean CO₂ was also estimated with an update of Park et al. (2010) based on regional correlations between surface temperature and $p\text{CO}_2$, changes in surface temperature observed by satellite, and wind speed estimates also from satellite data for 1990–2009 (Atlas et al., 2011). This estimate provides a data-based assessment of the interannual variability combined with a model-based assessment of the trend and mean in S_{OCEAN} .

ESSDD

6, 689–760, 2013

Global carbon budget
2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



We use the data-based results of Khatiwala et al. (2009) updated by Khatiwala et al. (2013) to estimate the cumulative carbon accumulated in the ocean during 1765–1958 (60.2 GtC) and 1870–1958 (47.5 GtC), and assume an oceanic uptake of 0.4 GtC for 1750–1765 where no data are available based on the mean uptake during 1765–1770. The estimate of Khatiwala et al. (2009) is based on regional disequilibrium between surface $p\text{CO}_2$ and atmospheric CO_2 , and a Green's function utilizing transient ocean tracers like CFCs and ^{14}C to ascribe changes through time. It does not include changes associated with changes in ocean circulation, temperature and climate, but these are thought to be small over the time period considered here (Ciais et al., 2013). The uncertainty in cumulative uptake of ± 20 GtC (converted to $\pm 1\sigma$) is taken directly from the IPCC's review of the literature (Rhein et al., 2013), or about $\pm 30\%$ for the annual values (Khatiwala et al., 2009).

2.4.2 Global Ocean Biogeochemistry models

The trend in the ocean CO_2 sink for 1959–2012 is computed using a combination of six global ocean biogeochemistry models (Table 5). The models represent the physical, chemical and biological processes that influence the surface ocean concentration of CO_2 and thus the air-sea CO_2 flux. The models are forced by meteorological reanalysis data and atmospheric CO_2 concentration available for the entire time period. They compute the air-sea flux of CO_2 over grid boxes of 1 to 4 degrees in latitude and longitude. The ocean CO_2 sink for each model is normalised to the observations, by dividing the annual model values by their observed average over 1990–1999, and multiplying this by the observational-based estimate of 2.2 GtC yr^{-1} . The ocean CO_2 sink for each year (t) is therefore:

$$S_{\text{OCEAN}}(t) = \frac{1}{n} \sum_{m=1}^{m=n} \frac{S_{\text{OCEAN}}^m(t)}{S_{\text{OCEAN}}^m(1990-1999)} \times 2.2 \quad (7)$$

where n is the number of models. This normalisation ensures that the ocean CO₂ sink for the global carbon budget is based on observations, and that the trends and annual values in CO₂ sinks are consistent with model estimates. The normalisation based on a ratio assumes that if models over or underestimate the sink in the 1990s, it is primarily due to the process of diffusion which depends on the gradient of CO₂. Thus a ratio is more appropriate than an offset as it takes into account the time-dependence of CO₂ gradients in the ocean. We use the four models published in Le Quéré et al. (2009), including updates of Aumont and Bopp (2006), Doney et al. (2009), Buitenhuis et al. (2010), and Galbraith et al. (2010) and further model estimate updated from Assman (2010) and Ilyina et al. (2013; Table 5). All models are available to 2012 except Galbraith et al. (2010), which is available to 2008. The mean ocean CO₂ sink from the six uncorrected models for 1990–1999 ranges between 1.3 and 2.6 GtC yr⁻¹, with a multi model mean of 1.93 GtC yr⁻¹ and a standard deviation across models of 0.46 GtC yr⁻¹.

2.4.3 Uncertainty assessment for S_{OCEAN}

The uncertainty around the mean CO₂ sink was already quantified for the 1990s (see Sect. 2.4.1). To quantify the uncertainty around annual values, we examine the standard deviation of the normalised model ensemble. We further use information from the two data-based products to assess the confidence level. The standard deviation of the ocean model ensemble averages to 0.13 GtC yr⁻¹ during 1980–2010 (with a maximum of 0.22), but it increases as the model ensemble goes back in time, with a standard deviation of 0.29 GtC yr⁻¹ across models in the 1960s. We estimate that the uncertainty in the annual ocean CO₂ sink is about ± 0.5 GtC yr⁻¹ from the quadratic sum of the data uncertainty of ± 0.4 GtC yr⁻¹ and standard deviation across models of up to ± 0.3 GtC yr⁻¹, reflecting both the uncertainty in the mean sink and in the interannual variability as assessed by models.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

The interannual variability of the ocean fluxes of the two data-based estimates for 1990–2009 (when they overlap) is $\pm 0.34 \text{ GtC yr}^{-1}$ (Rödenbeck et al., 2013) and $\pm 0.14 \text{ GtC yr}^{-1}$ (Park et al., 2010), which compares well to the interannual variability of $\pm 0.20 \text{ GtC yr}^{-1}$ estimated here based on Eq. (7). The phase is generally consistent between estimates, with a higher ocean CO_2 sink during El Niño events. The two data-based estimates correlate with the ocean CO_2 sink estimated here with the same correlation of $r = 0.59$, but with a mutual correlation between data-based estimates of 0.30 only. A comparison of variability in regional fluxes also shows generally consistent patterns in amplitude, although not everywhere in phase (not shown). We assess a medium confidence level to the annual ocean CO_2 sink and its uncertainty because the interannual variability in the model and data-based estimates are both generally small and consistent in time, and the mean CO_2 sink is based on observations.

2.5 Terrestrial CO_2 sink

The difference between the fossil-fuel (E_{FF}) and LUC net emissions (E_{LUC}), the growth rate in atmospheric CO_2 concentration (G_{ATM}) and the ocean CO_2 sink (S_{OCEAN}) is attributable to the net sink of CO_2 in terrestrial vegetation and soils (S_{LAND}), within the given uncertainties. Thus, this sink can be estimated either as the residual of the other terms in the mass balance budget but also directly calculated using DGVMs. The residual land sink (S_{LAND}) is in part due to the fertilising effect of rising atmospheric CO_2 on plant growth, N deposition and climate change effects such as prolonged growing seasons in northern temperate areas. S_{LAND} does not include gross land sinks directly resulting from LUC (e.g. regrowth of vegetation) as these are estimated as part of the net land use flux (E_{LUC}). System boundaries make it difficult to attribute exactly CO_2 fluxes on land between S_{LAND} and E_{LUC} (Erb et al., 2013), and by design most of the uncertainties in our method are allocated to S_{LAND} for those processes that are poorly known or represented in models.

2.5.1 Residual of the budget

For 1959–2012, the terrestrial carbon sink was estimated from the residual of the other budget terms by rearranging Eq. (1):

$$S_{\text{LAND}} = E_{\text{FF}} + E_{\text{LUC}} - (G_{\text{ATM}} + S_{\text{OCEAN}}) \quad (8)$$

- 5 The uncertainty in S_{LAND} is estimated annually from the quadratic sum of the uncertainty in the right-hand terms assuming the errors are not correlated. The uncertainty averages to $\pm 0.8 \text{ GtC yr}^{-1}$ over 1959–2012 (Table 6). S_{LAND} estimated from the residual of the budget will include, by definition, all the missing processes and potential biases in the other component of Eq. (8).

10 2.5.2 DGVMs

A comparison of the residual calculation of S_{LAND} in Eq. (8) with the same DGVMs used to estimate E_{LUC} in Sect. 2.2.3, but here excluding the effects of changes in land cover (using a constant pre-industrial land cover distribution), provides an independent estimate of the consistency of S_{LAND} with our understanding of the functioning of the terrestrial vegetation in response to CO_2 and climate variability (Table 6). As described in Sect. 2.2.3, the DGVMs include all climate variability and CO_2 effects over land, but do not include reductions in CO_2 sink capacity associated with human activity directly affecting changes in vegetation cover and management, which by design is allocated to E_{LUC} . This effect has been estimated to have led to a reduction in the terrestrial sink by 0.5 GtC yr^{-1} since 1750 (Gitz and Ciais, 2003). The models estimate the mean and variability of S_{LAND} based on atmospheric CO_2 and climate, and thus both terms can be compared to the budget residual.

25 The multi-model mean of $2.7 \pm 1.0 \text{ GtC yr}^{-1}$ for the period 2003–2012 agrees well with the value computed from the budget residual (Table 6). The standard deviation of the annual CO_2 sink across the eight DGVMs ranges from ± 0.4 to $\pm 1.3 \text{ GtC yr}^{-1}$, with a mean standard deviation of $\pm 0.9 \text{ GtC yr}^{-1}$ for the period 1959 to 2012. The model mean

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



correlates with the budget residual with $r = 0.71$, compared to correlations of $r = 0.47$ to $r = 0.71$ (median of 0.63) by individual models. The standard deviation is similar to that of the five model ensembles presented in Le Quéré et al. (2009), but the correlation is improved compared to $r = 0.54$ obtained in that later study. The DGVM results confirm that the sum of our knowledge on annual CO_2 emissions and their partitioning is plausible (see Discussion), and they enable the attribution of the fluxes to the underlying processes and provide a breakdown of the regional contributions (not shown here). However as the standard deviation across the DGVM models (of $\pm 0.9 \text{ GtC yr}^{-1}$) is of the same magnitude as the combined uncertainty due to the other components (E_{FF} , E_{LUC} , G_{ATM} , S_{OCEAN} ; Table 6), the DGVMs do not provide further constraints on the terrestrial CO_2 sink compared to the residual of the budget (Eq. 8). We assess a medium confidence level to the annual land CO_2 sink and its uncertainty because the estimates from the residual budget and DGVMs match well within the given uncertainty, and the estimates based on the residual budget are primarily dependent on E_{FF} and G_{ATM} , both of which are well constrained. The sum of E_{LUC} and S_{LAND} is better constrained than their individual components.

2.6 Processes not included in the global carbon budget

2.6.1 Contribution of anthropogenic CO and CH_4 to the global anthropogenic CO_2 budget

Anthropogenic emissions of CO and CH_4 to the atmosphere are eventually oxidized to CO_2 and thus are part of the anthropogenic CO_2 budget. These contributions are omitted in Eq. (1), but a first attempt is made in this section to estimate their magnitude, and identify the sources of uncertainty. The anthropogenic CO emissions that are part of the anthropogenic CO_2 budget are from incomplete fossil-fuel burning for CO and deforestation fires. The only anthropogenic emissions of fossil CH_4 that matter for the anthropogenic CO_2 budget are the fugitive emissions of coal, oil and gas upstream

sectors (see below). These emissions of CO and CH₄ contribute a net addition of fossil carbon to the atmosphere.

In our estimate of E_{FF} we assume that all the fuel burned is emitted as CO₂, thus CO emissions and their atmospheric oxidation into CO₂ within a few months are already counted implicitly in E_{FF} and should not be counted twice (same for E_{LUC} and CO deforestation fires). Anthropogenic emissions of fossil CH₄ are not included in the fossil-fuel CO₂ emissions E_{FF} , because these mainly fugitive emissions are not included in fuel inventories, but they contribute to the carbon budget after CH₄ gets oxidized into CO₂ (lifetime 12.4 yr; Prather et al., 2012). These anthropogenic fossil CH₄ emissions are estimated to be 0.07 GtC yr⁻¹ [0.06–0.09] (after Kirschke et al., 2013) and are assumed to be oxidized into CO₂ with a lifetime of 12.4 yr. After one year, 92% of these emissions thus remain in the atmosphere as CH₄ and contribute to the observed CH₄ global growth rate but not to the CO₂ growth rate, whereas the rest (8%) get oxidized into CO₂, and contribute to the CO₂ growth rate. Given that anthropogenic fossil-fuel CH₄ emissions represent a fraction of 15% of the total global CH₄ source (Kirschke et al., 2013) we assumed that a fraction of 0.15 times 0.92 of the observed global growth rate of CH₄ of 6 Tg C yr⁻¹ during 2000–2009 is due to fossil CH₄ sources. Therefore, annual fossil-fuel CH₄ emissions contribute 0.8 Tg C yr⁻¹ to the CH₄ growth rate and 0.8 Tg C yr⁻¹ (units of C in CO₂ form) to the CO₂ growth rate. Summing up the effect of fossil-fuel CH₄ emissions from each previous year during the past 10 yr, a fraction of which is oxidized into CO₂ in the current year, this defines a contribution of 5 Tg C yr⁻¹ to the CO₂ growth rate, or about 0.1%. Thus the effect of anthropogenic fossil CH₄ emissions and their oxidation to anthropogenic CO₂ in the atmosphere can be assessed to have a negligible effect on the observed CO₂ growth rate, although they do contribute significantly to the global CH₄ growth rate.

Other anthropogenic biogenic sources of CO and CH₄ from wildfires, biomass, and CH₄ wetlands are assumed to be balanced by annual CO₂ uptake by photosynthesis on continental and long time-scale (e.g. decadal or longer).

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

2.6.2 Anthropogenic carbon fluxes in the land to ocean continuum

The approach used to determine the global carbon budget considers only anthropogenic CO₂ emissions and their partitioning among the atmosphere, ocean and land. In the analysis, the land and ocean reservoirs that take up anthropogenic CO₂ from the atmosphere are conceived as independent carbon storage repositories. This approach thus omits that carbon is continuously displaced along the land-ocean aquatic continuum (LOAC) comprising freshwaters, estuaries and coastal areas. Carbon is transferred both in inorganic (bicarbonates and dissolved CO₂), and organic (dissolved and particulate organic carbon) forms along this continuum. During its journey from upland terrestrial ecosystems to the oceans, carbon is not only transferred laterally, but is also sequestered in e.g. lake and coastal sediments (Krumins et al., 2013; Tranvik et al., 2009) or released back to the atmosphere, mainly as respired CO₂ (Aufdenkampe et al., 2011; Battin et al., 2009; Cole et al., 2007; Laruelle et al., 2010; Regnier et al., 2013), and to a much lesser extent, as CH₄ (Bastviken et al., 2011; Borges and Abril, 2011). A significant fraction of this lateral carbon flux is entirely “natural” and is thus a steady state component of the pre-industrial carbon cycle that can be ignored in the current analysis. The remaining fraction is anthropogenic carbon entrained into the lateral transport loop of the LOAC, a perturbation that is relevant for the global carbon budget presented here.

The recent synthesis by Regnier et al. (2013) is the first attempt to estimate the anthropogenic component of LOAC carbon fluxes and their significance for the global carbon budget. The results of their analysis can be summarized in three points of relevance to the budget. First, only a portion of the anthropogenic CO₂ taken up by land ecosystems is sequestered in soil and biomass pools, as $1 \pm 0.5 \text{ GtC yr}^{-1}$ is exported to the LOAC. This flux is comparable to the C released to the atmosphere by LUC (Table 6). Second, the exported anthropogenic C is both stored ($0.55 \pm 0.3 \text{ GtC yr}^{-1}$) and released back to the atmosphere as CO₂ ($0.35 \pm 0.2 \text{ GtC yr}^{-1}$), the magnitude of these fluxes resulting from the combined effects of freshwaters, estuaries and coastal seas.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global carbon budget
2013**

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Third, a small fraction of anthropogenic carbon displaced by the LOAC accumulates in the open ocean ($0.1 \pm > 0.05 \text{ GtC yr}^{-1}$). The anthropogenic perturbation of the carbon fluxes from land to ocean do not question the method used in Sect. 2.5 to define the ocean sink and residual land sink (Table 4). However, it does point to the need to account for the fate of anthropogenic carbon once it is removed from the atmosphere by land ecosystems (summarized in Fig. 1). In theory, direct estimates of changes of ocean inorganic carbon inventory over time would see the land flux of anthropogenic carbon and would thus have a bias relative to air-sea flux estimates and tracer based reconstructions. However currently the value is small enough that it is not noticeable relative to the errors in the individual techniques.

Of greater importance is the finding that the residual land sink calculated in a budget which accounts for the LOAC ($2.95 \pm 0.9 \text{ GtC yr}^{-1}$) is larger than the value of $2.6 \pm 0.8 \text{ GtC yr}^{-1}$ reported in Table 6, because this flux is partially offset by the net source of CO_2 to the atmosphere of $0.35 \pm 0.3 \text{ GtC yr}^{-1}$ from rivers, estuaries and coastal seas. In addition, because anthropogenic CO_2 taken up by land ecosystems is exported to the LOAC, the annual land carbon storage change (1.15 GtC yr^{-1}) is notably smaller than the net CO_2 uptake by land ecosystems calculated in the GCP budget (1.8 GtC yr^{-1}), a significant fraction of the displaced carbon (0.65 GtC yr^{-1}) being stored in freshwater and coastal sediments (0.55 GtC yr^{-1}), and to a lesser extent, in the open ocean (0.1 GtC yr^{-1}).

All estimates of LOAC are given with low confidence, because they originate from a single source. The carbon budget presented here implicitly incorporates the fluxes from the LOAC with S_{LAND} . We do not attempt to separate these fluxes because the uncertainties in either estimate are too large, and there is insufficient information available to estimate the LOAC fluxes on an annual basis.

3 Results

3.1 Global carbon budget averaged over decades and its variability

The global carbon budget averaged over the last decade (2003–2012) is shown in Fig. 1. For this time period, 92 % of the total emissions ($E_{\text{FF}} + E_{\text{LUC}}$) were caused by fossil-fuel combustion and cement production, and 8 % by land-use change. The total emissions were partitioned among the atmosphere (45 %), ocean (27 %) and land (27 %). All components except land-use change emissions have grown since 1959 (Figs. 2 and 3), with important interannual variability in the atmospheric growth rate caused primarily by variability in the land CO_2 sink (Fig. 3), and some decadal variability in all terms (Table 7).

Global CO_2 emissions from fossil-fuel combustion and cement production have increased every decade from an average of $3.1 \pm 0.2 \text{ GtC yr}^{-1}$ in the 1960s to an average of $8.6 \pm 0.4 \text{ GtC yr}^{-1}$ during 2003–2012 (Table 7 and Fig. 4). The growth rate in these emissions decreased between the 1960s and the 1990s, from $4.5 \% \text{ yr}^{-1}$ in the 1960s, $2.7 \% \text{ yr}^{-1}$ in the 1970s, $2.0 \% \text{ yr}^{-1}$ in the 1980s, $1.1 \% \text{ yr}^{-1}$ in the 1990s, and increased again since year 2000 at an average of $3.1 \% \text{ yr}^{-1}$ for 2003–2012, the growth rate was 2.7 %. In contrast, CO_2 emissions from LUC have remained constant at around $1.4 \pm 0.5 \text{ GtC yr}^{-1}$ during 1960–1999, and decreased to $0.8 \pm 0.5 \text{ GtC yr}^{-1}$ during 2003–2012. The E_{LUC} estimates from the bookkeeping method and from the DGVM models are remarkably similar, except for the 1990s where the DGVM emissions are about 1.5 times larger than those of the bookkeeping method (Table 6 and Fig. 5). The decreased emissions from LUC since 2000 is also reproduced by the DGVMs (Fig. 5).

The growth rate in atmospheric CO_2 increased from $1.7 \pm 0.1 \text{ GtC yr}^{-1}$ in the 1960s to $4.3 \pm 0.1 \text{ GtC yr}^{-1}$ during 2003–2012 with important decadal variations (Table 7). The ocean CO_2 sink increased from $1.1 \pm 0.5 \text{ GtC yr}^{-1}$ in the 1960s to $2.6 \pm 0.5 \text{ GtC yr}^{-1}$ during 2003–2012, with decadal variations of the order of a few tenths of GtC yr^{-1} . The low uptake anomaly around year 2000 originates from multiple regions in all models

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



(western Equatorial Pacific, Southern Ocean and North Atlantic), and is caused by climate variability. The land CO₂ sink increased from 1.8 ± 0.9 GtC yr⁻¹ in the 1960s to 2.6 ± 0.8 GtC yr⁻¹ during 2003–2012, with important decadal variations of 1–2 GtC yr⁻¹. The high uptake anomaly around year 1991 is thought to be caused by the effect of the volcanic eruption of Mount Pinatubo and is not generally reproduced by the DGVMs (Fig. 5). The larger land CO₂ sink during 2003–2012 is reproduced by the DGVMs in response to combined atmospheric CO₂ increase and climate change and variability, fully consistent with the budget residual (Table 6). Both ocean and land CO₂ sinks increased roughly in line with the atmospheric increase, but with large decadal variability on land.

3.2 Global carbon budget for year 2012 and emissions projection for 2013

Global CO₂ emissions from fossil-fuel combustion and cement production reached 9.7 ± 0.5 GtC in 2012 (Fig. 4; see also Peters et al., 2013), 2.2 % higher than the emissions in 2011. This compares to our estimate of 2.6 % yr⁻¹ made last year (Peters et al., 2013), based on an estimated GDP growth of 3.3 % yr⁻¹ and improvement in *I*_{FF} of -0.7 % yr⁻¹ (Table 8) The latest estimate of GDP growth for 2012 was 3.2 % yr⁻¹ (IMF, 2013) and hence *I*_{FF} improved -1.0 % yr⁻¹, slightly better than our prediction. The 2012 emissions were distributed among coal (43 %), oil (33 %), gas (18 %), cement (5.3 %) and gas flaring (0.6 %). These first four categories increased by 2.8 %, 1.2 %, 2.5 %, and 2.5 % respectively over the previous year. Due to lack of data gas flaring in 2012 is assumed equal to 2011.

Using Eq. (6), we estimate that these global CO₂ emissions in 2013 will reach 9.9 ± 0.5 GtC, or 2.1 % above 2012 levels (likely range of 1.1–3.1 %), and that emissions in 2013 will thus be 61 % above emissions in 1990. The expected value is computed using the world GDP projection of 2.9 % made by the IMF (2013) and a growth rate for *I*_{FF} of -0.8 % yr⁻¹ which is the average from the previous 10 yr. The *I*_{FF} is based on GDP in constant PPP from the IEA (2012) up to 2010 (IEA/OECD, 2012) and

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

**Global carbon budget
2013**

 C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

5 extended using the IMF growth rates of 3.9% in 2011 and 3.2% in 2012. The uncertainty range is based on an uncertainty of 0.6% for GDP growth (the range in IMF estimates of 2013 GDP growth published in January, April, July, and October 2013, 3.5%, 3.3%, 3.1%, and 2.9%, respectively) and the range in I_{FF} due to short term trends of $-0.4\% \text{ yr}^{-1}$ (2008–2012) and medium term trends of $-1.2\% \text{ yr}^{-1}$ (1990–2012); the combined uncertainty range is therefore 1.1% (2.9–0.6–1.2; low GDP growth, large I_{FF} improvements) and 3.1% (2.9+0.6–0.4; high GDP growth, small I_{FF} improvements). Projections made in the previous global carbon budgets compared well to the actual CO_2 emissions for that year (Table 8 and Fig. 6) and were useful to capture the current state of the fossil-fuel emissions (see also Peters et al., 2013).

10 In 2012, global CO_2 emissions were dominated by emissions from China (27%), the USA (14%), the EU (28 member states; 10%), and India (6%) compared to the global total including bunker fuels. These five regions account for 63% of global emissions. Growth rates for these countries from 2011 to 2012 were 5.9% (China), -3.7% (USA), -1.3% (EU28), and 7.7% (India). The countries contributing most to the 2012 change in emissions were China (71% increase), USA (26% decrease), India (21% increase), Japan (11% increase), and Australia (6% decrease). The per-capita CO_2 emissions in 2012 were $1.4 \text{ tC person}^{-1} \text{ yr}^{-1}$ for the globe, and 4.4, 1.9, 1.9 and $0.5 \text{ tC person}^{-1} \text{ yr}^{-1}$ for the USA, China, the EU and India, respectively (Fig. 4e).

20 Territorial-based emissions in Annex B countries have remained stable from 1990–2011, while consumption-based emissions have grown at $0.5\% \text{ yr}^{-1}$ (Fig. 4c). In non-Annex B countries territorial-based emissions have grown at $4.3\% \text{ yr}^{-1}$, while consumption-based emissions have grown at $4.0\% \text{ yr}^{-1}$. In 1990, 62% of global territorial-based emissions were emitted in Annex B countries (34% in non-Annex B, and 4% in bunker fuels used for international shipping and aviation), while in 2011 this had reduced to 38% (56% in non-Annex B, and 6% in bunkers). In terms of consumption-based emissions this split was 63% in 1990 and 43% in 2011 (33 to 51% in non-Annex B). The difference between territorial-based and consumption-based emissions (the net emission transfer via international trade) from non-Annex

Global carbon budget
2013

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

B to Annex B countries has increased from 0.05 GtC yr⁻¹ in 1990 to 0.46 GtC in 2011 (Fig. 4), with an average annual growth rate of 12% yr⁻¹. The increase in net emission transfers of 0.41 GtC from 1990–2011 compares with the emission reduction of 0.21 GtC in Annex B countries. These results clearly show a growing net emission transfer via international trade from non-Annex B to Annex B countries. In 2011, the biggest emitters from a consumption-based perspective were China (22 % of the global total), USA (17 %), EU28 (14 %), and India (5 %).

Based on DGVMs only, the global CO₂ emissions from land-use change activities are estimated as 0.6 ± 0.7 GtC in 2012, lower than the 2003–2012 average of 1.1 ± 0.5 GtC yr⁻¹. However, although the decadal mean generally agreed, the estimated annual variability was not consistent between the bookkeeping method and the DGVMs (Fig. 5a). This could be partly due to the design of the DGVM experiments, which use flux differences between simulations with and without land cover change, and thus may overestimate variability due to fires in regions where the contemporary land cover is smaller than pre-industrial cover. For this reason we assign a mean value to E_{LUC} for year 2012 based on the 2001–2010 average.

Atmospheric CO₂ growth rate was 5.2 ± 0.2 GtC in 2012 (2.43 ± 0.09 ppm; Fig. 3; Dlugokencky and Tans, 2013). This is significantly above the 2003–2012 average of 4.3 ± 0.1 GtC yr⁻¹, though the interannual variability in atmospheric growth rate is large.

The ocean CO₂ sink was 2.9 ± 0.5 GtC yr⁻¹ in 2012, an increase of 0.2 GtC yr⁻¹ over 2011. This is larger than the 2003–2012 average of 2.6 ± 0.5 GtC yr⁻¹. All models produce an increase in the ocean CO₂ sink in 2012 compared to 2011 except for MICOM-HAMOCC, which shows a very small decrease in the sink.

The terrestrial CO₂ sink calculated as the residual from the carbon budget was 2.5 ± 0.9 GtC in 2012, well below the 4.1 ± 0.9 GtC in 2011, which was a La Niña year, but near the 2003–2012 average of 2.6 ± 0.8 GtC (Fig. 3). The DGVMs model mean suggests a lower terrestrial CO₂ sink in 2012 of 1.7 ± 1.2 GtC (Table 6), possibly from weak El Niño conditions in the Northern Hemisphere spring of year 2012. The DGVMs thus estimate internally consistent land fluxes over 2012, with both E_{LUC} and S_{LAND}

being weaker than those of the carbon budget. Internal consistency is an emerging property of the models, not an a-priori constraint as is the residual calculation of S_{LAND} . These results thus suggest that constraints from DGVMs may provide sufficient information to be directly incorporated in the budget calculations in the future.

3.3 Cumulative emissions

Cumulative emissions for 1870–2012 were 380 ± 20 GtC for E_{FF} , and 160 ± 55 GtC for E_{LUC} based on the bookkeeping method of Houghton and Hackler (2013) for 1870–2010, with an extension to 2012 based on methods described in Sect. 2.2 (Table 9). The cumulative emissions are rounded to the nearest 5 GtC. The total cumulative emissions for 1870–2012 are 540 ± 60 GtC. These emissions were partitioned among the atmosphere (220 ± 5 GtC) based on atmospheric measurements in ice cores of 288 ppm (Sect. 2.3.1; Joos and Spahni, 2008) and recent direct measurements of 392.52 ppm (Dlugokencky and Tans, 2013), ocean (150 ± 20 GtC) using Khatiwala et al. (2013) prior to 1959 and Table 7 otherwise, and the land (170 ± 65 GtC) by difference.

Cumulative emissions for the early period 1750–1869 were 3 GtC for E_{FF} , and about 45 GtC for E_{LUC} of which 15 GtC were emitted in the period 1850–1870 (Houghton and Hackler, 2013) and 30 GtC were emitted in the period 1750–1850 based on the average of four publications (22 GtC by Pongratz et al., 2009; 15 GtC by van Minnen et al., 2009; 64 GtC by Shevliakova et al., 2009 and 24 GtC by Zaehle et al., 2011). The growth in atmospheric CO_2 during that time was about 25 GtC, and the ocean uptake about 15 GtC, implying a land uptake of 10 GtC. These numbers have large relative uncertainties but balance within the limits of our understanding.

Cumulative emissions for 1750–2012 based on the sum of the two periods above were 385 ± 20 GtC for E_{FF} , and 205 ± 70 GtC for E_{LUC} , for a total of 590 ± 75 GtC, partitioned among the atmosphere (245 ± 5 GtC), ocean (165 ± 20 GtC), and the land (180 ± 75 GtC).

Cumulative emissions through to year 2013 can be estimated based on the 2013 projections of E_{FF} (Sect. 3.2), the largest contributor, and assuming a constant E_{LUC} .

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



For 1870–2013, these are 550 ± 60 GtC for total emissions, with about 70 % contribution from E_{FF} (390 ± 20 GtC) and about 30 % contribution from E_{FF} (160 ± 55 GtC). Cumulative emissions since year 1870 are higher than the emissions of 515 [445 to 585] GtC reported in the IPCC (Stocker et al., 2013) because they include and additional 21 GtC from emissions in 2012 and 2013 (mostly from E_{FF}), and an additional 20 GtC from the revisions of E_{LUC} in the early century (see Sect. 2.2.1). The uncertainty presented here ($\pm 1\sigma$) is smaller than the range of 90 % used by IPCC, but both estimates overlap within their uncertainty ranges.

4 Discussion

Each year when the global carbon budget is published, each component for all previous years is updated to take into account corrections that are due to further scrutiny and verification of the underlying data in the primary input data sets. The updates have generally been relatively small and focused on the most recent years, except for LUC, where they are more significant but still generally within the provided uncertainty range (Fig. 6). The difficulty in accessing land cover change data to estimate E_{LUC} is the key problem to providing continuous records of emissions in this sector. Revisions in E_{LUC} for the 2008/2009 budget was the result of the release of FAO 2010, which contained a major update to forest cover change for the period 2000–2005 and provided the data for the following 5 yr to 2010 (Fig. 6b). The differences this year could be attributable to both the different data and the different methods. Updates were at most 0.24 GtC yr⁻¹ for the fossil-fuel and cement emissions, 0.19 GtC yr⁻¹ for the atmospheric growth rate, 0.20 GtC yr⁻¹ for the ocean CO₂ sink, all within the reported uncertainty. The update for the residual land CO₂ sink was also large (Fig. 6e), with maximum value of 0.71 GtC yr⁻¹, directly reflecting the revision in other terms of the budget, but still within the reported uncertainty.

Our capacity to separate the CO₂ budget components can be evaluated by comparing the land CO₂ sink estimated with the budget residual (S_{LAND}), which includes errors

and biases from all components, with the land CO₂ sink estimates by the DGVM ensemble, which are based on our understanding of processes of how the land responds to increasing CO₂, climate change and variability. The two estimates are generally close (Fig. 5), both for the mean and for the interannual variability. The DGVM mean correlate with the budget residual with $r = 0.71$ (Sect. 2.5.2; Fig. 5). The DGVMs produce a decadal mean and standard deviation across eight models of $2.6 \pm 0.8 \text{ GtC yr}^{-1}$ for the period 2000–2009, nearly the same as the estimate produced with the budget residual (Table 6). Finally the fact that the DGVMs provide an internally consistent split between E_{LUC} and S_{LAND} for year 2012 suggests that they could inform the annual budget analysis more extensively as the effort evolves. Analysis of regional CO₂ budgets would provide further information to quantify and improve our estimates, as has been undertaken by the REgional Carbon Cycle Assessment and Processes (Canadell et al., 2012–2013).

Annual estimates of each component of the global carbon budgets have their limitations, some of which could be improved with better data and/or better understanding of carbon dynamics. The primary limitations involve resolving fluxes on annual time scales and providing updated estimates for recent years for which data-based estimates are not yet available or only beginning to emerge. Of the various terms in the global budget, only the fossil-fuel burning and atmospheric growth rate terms are based primarily on empirical inputs supporting annual estimates in this carbon budget. The data on fossil-fuel consumption and cement production are based on survey data in all countries. The other terms can be provided on an annual basis only through the use of models. While these models represent the current state of the art, they provide only estimates of actual changes. For example, the decadal trends in ocean uptake and the interannual variations associated with El Niño/La Niña (ENSO) are not directly constrained by observations, although many of the processes controlling these trends are sufficiently well known that the model-based trends still have value as benchmarks for further validation. Data-based products for the ocean CO₂ sink provide new ways to evaluate the model results, and could be used directly as data become more rapidly

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

available and methods for creating such products improve. Estimates of land-use emissions and their year-to-year variability have even larger uncertainty, and much of the underlying data are not available as an annual update. Efforts are underway to work with annually available satellite area change data or FAO reported data in combination with fire data and modelling to provide annual updates for future budgets. The best resolved changes are in atmospheric growth (G_{ATM}), fossil-fuel emissions (E_{FF}), and by difference, the change in the sum of the remaining terms ($S_{OCEAN} + S_{LAND} - E_{LUC}$). The variations from year-to-year in these remaining terms are largely model-based at this time. Further efforts to increase the availability and use of annual data for estimating the remaining terms with annual to decadal resolution are especially needed.

Our approach also depends on the reliability of the energy and land-cover change statistics provided at the country level, and are thus potentially subject to biases. Thus it is critical to develop multiple ways to estimate the carbon balance at the global and regional level, including from the inversion of atmospheric CO₂ concentration, the use of other oceanic and atmospheric tracers, and the compilation of emissions using alternative statistics (e.g. sectors). Multiple approaches ranging from global to regional scale would greatly help increase confidence and reduce uncertainty in CO₂ emissions and their fate.

5 Conclusions

The estimation of global CO₂ emissions and sinks is a major effort by the carbon cycle research community that requires a combination of measurements and compilation of statistical estimates and results from models. The delivery of an annual CO₂ budget serves two purposes. First, there is a large demand for up-to-date information on the state of the anthropogenic perturbation of the climate system and its underpinning causes. A broad stakeholder community relies on the datasets associated with the annual CO₂ budget including scientists, policy makers, businesses, journalists, and the broader society increasingly engaged in adapting to and mitigating human-driven

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



**Global carbon budget
2013**

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

climate change. Second, over the last decade we have seen unprecedented changes in the human and biophysical environments (e.g. increase in the growth of fossil-fuel emissions, ocean temperatures, and strength of the land sink), which call for more frequent assessments of the state of the Planet, and by implication a better understanding of the future evolution of the carbon cycle, and the requirements for climate change mitigation and adaptation. Both the ocean and the land surface presently remove a large fraction of anthropogenic emissions. Any significant change in the function of carbon sinks is of great importance to climate policymaking, as they affect the excess carbon dioxide remaining in the atmosphere and therefore the compatible emissions for any climate stabilization target. Better constraints of carbon cycle models against contemporary datasets raises the capacity for the models to become more accurate at future projections.

This all requires more frequent, robust, and transparent datasets and methods that can be scrutinized and replicated. After eight annual releases from the GCP, the effort is growing and the traceability of the methods has become increasingly complex. Here, we have documented in detail the datasets and methods used to compile the annual updates of the global carbon budget, explained the rationale for the choices made, the limitations of the information, and finally highlighted need for additional information where gaps exist.

This paper via “living data” will help to keep track of new budget updates. The evolution over time of the CO₂ budget is now a key indicator of the anthropogenic perturbation of the climate system, and its annual delivery joins a set of climate indicators to monitor the evolution of human-induced climate change, such as the annual updates on the global surface temperature, sea level rise, minimum Arctic sea ice extent and others.

6 Data access

The data presented here is made available in the belief that their wide dissemination will lead to greater understanding and new scientific insights of how the carbon cycle works, how humans are altering it, and how we can mitigate the resulting human-driven climate change. The free availability of these data (doi:10.3334/CDIAC/GCP_2013_v1.1) does not constitute permission for publication of the data. For research projects, if the data are essential to the work, or if an important result or conclusion depends on the data, co-authorship may need to be considered. Full contact details and information on how to cite the data are given at the top of each page in the accompanying database.

The accompanying database includes an Excel file organised in the following spreadsheets:

1. Summary
2. The global carbon budget (1959–2012).
3. Global CO₂ emissions from fossil-fuel combustion and cement production by fuel type, and the per-capita emissions (1959–2012).
4. Territorial-based (e.g. as reported to the UN Framework Convention on Climate Change) country CO₂ emissions from fossil-fuel combustion and cement production (1959–2012).
5. Consumption-based country CO₂ emissions from fossil-fuel combustion and cement production and emissions transfer from the international trade of goods and services (1990–2011).
6. Emissions transfers (Consumption minus territorial emissions) (1990–2011).
7. CO₂ emissions from land-use change from the individual methods and models (1959–2012).

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



8. Ocean CO₂ sink from the individual ocean models (1959–2012).
9. Terrestrial residual CO₂ sink from the DGVMs (1959–2012).
10. Country definitions.

7 Data Provenance and Structure

5 All data sources and individual components of the global carbon budget 2013 are documented throughout Sect. 2 and summarised in Tables 2, 3 and 5.

Acknowledgements. We thank all people and institutions that provided data used in this carbon budget, P. Cadule, J. C. Enright, Ghattas, and G. Hurtt for support to the model simulations, F. Joos and S. Khatiwala for providing historical data, and M. Heimann and G. McKinley for their reviews and suggestions. The observations and modelling analysis were possible thanks to the generous support from the author's institutions and funding from multiple agencies around the world. The UK Natural Environment Research Council (NERC) provided funding to C. Le Quéré and the GCP through their International Opportunities Fund specifically to support this publication (project NE/103002X/1). R. J. Andres and T. A. Boden were supported by US Department of Energy, Office of Science, Biological and Environmental Research (BER) programs and performed at Oak Ridge National Laboratory (ORNL) under US Department of Energy contract DE-AC05-00OR22725. G. P. Peters and R. M. Andrews were supported by the Norwegian Research Council (221355). A. Arneeth, A. Omar, C. Le Quéré, J. Schwinger, P. Ciais, P. Friedlingstein, P. Regnier, S. Sitch and S. Zaehle were supported by the EU FP7 for funding through projects GEOCarbon (283080), COMBINE (226520), CARBOCHANGE (264879), EM-BRACE (GA282672), and LUC4C (GA603542). A. Harper was supported by the NERC Joint Weather and Climate Research Programme. A. K. Jain was supported by the US National Science Foundation (NSF AGS 12-43071) and the US Department of Energy, Office of Science and BER programs (DOE DE-SC0006706). B. D. Stocker was supported by the Swiss National Science Foundation. A. Wiltshire was supported by the Joint UK DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). E. Kato was supported by the Environment Research and Technology Development Fund (S-10) of the Ministry of Environment of Japan. J. G. Canadell and M. R. Raupach are supported by the Australian Climate Change Science

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Program. J. I. House was supported by a Leverhulme Research Fellowship. S. C. Doney was supported by the US National Science Foundation (NSF AGS-1048827).

References

- Andres, R. J., Fielding, D. J., Marland, G., Boden, T. A., Kumar, N., and Kearney, A. T.: Carbon dioxide emissions from fossil fuel use, 1751–1950, *Tellus*, 51, 759–765, 1999.
- Andres, R. J., Boden, T. A., Bréon, F.-M., Ciais, P., Davis, S., Erickson, D., Gregg, J. S., Jacobson, A., Marland, G., Miller, J., Oda, T., Olivier, J. G. J., Raupach, M. R., Rayner, P., and Treanton, K.: A synthesis of carbon dioxide emissions from fossil-fuel combustion, *Biogeosciences*, 9, 1845–1871, doi:10.5194/bg-9-1845-2012, 2012.
- Andrew, R. M. and Peters, G. P.: A multi-region input-output table based on the Global Trade Analysis Project Database (GTAP-MRIO), *Economic Systems Research*, 2013, 99–121, 2013.
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K. M., K., Munhoven, G., Montenegro, A., and Tokos, K.: Atmospheric Lifetime of Fossil Fuel Carbon Dioxide, *Annu. Rev. Earth Pl. Sc.*, 37, 117–134, 2009.
- Assmann, K. M., Bentsen, M., Segschneider, J., and Heinze, C.: An isopycnic ocean carbon cycle model, *Geosci. Model Dev.*, 3, 143–167, doi:10.5194/gmd-3-143-2010, 2010.
- Atlas, R., Hoffman, R. N., Ardizzone, J., Leidner, S. M., Jusem, J. C., Smith, D. K., and Gombos, D.: A cross-calibrated, multiplatform ocean surface wind velocity product for meteorological and oceanographic applications, *B. Am. Meteor. Soc.*, 92, 157–174, 2011.
- Aufdenkampe, A. K., Mayorga, E., Raymond, P. A., Melack, J. M., Doney, S. C., Alin, S. R., Aalto, R. E., and Yoo, K.: Riverine coupling of biogeochemical cycles between land, oceans and atmosphere, *Frontiers Ecology Environ.*, 9, 53–60, 2011.
- Aumont, O. and Bopp, L.: Globalizing results from ocean in situ iron fertilization studies, *Global Biogeochem. Cy.*, 20, GB2017, doi:10.1029/2005GB002591, 2006.
- Baccini, A., Goetz, S. J., Walker, W. S., Laporte, N. T., Sun, M., Sulla-Menashe, D., Hackler, J., Beck, P. S. A., Dubayah, R., Friedl, M. A., Samanta, S., and Houghton, R. A.: Estimated carbon dioxide emissions from tropical deforestation improved by carbon-density maps, *Nature Clim. Change*, 2, 182–186, 2012.

**Global carbon budget
2013**

 C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S., Kozyr, A., Nojiri, Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl, J., Barbero, L., Bates, N., Boutin, J., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen, L., Chierici, M., Currie, K., de Baar, H. J. W., Evans, W., Feely, R. A., Fransson, A., Gao, Z., Hales, B., Hardman-Mountford, N., Hoppema, M., Huang, W.-J., Hunt, C. W., Huss, B., Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S. D., Jutterström, S., Kitidis, V., Körtzinger, A., Landschutzer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat, L., Metz, N., Murata, A., Newberger, T., Ono, T., Park, G.-H., Paterson, K., Pierrot, D., Ríos, A. F., Sabine, C. L., Saito, S., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R., Skjelvan, I., Steinhoff, T., Sullivan, K., Sun, H., Sutton, A. J., Suzuki, T., Sweeney, C., Takahashi, T., Tjiputra, J., Tsurushima, N., van Heuven, S. M. A. C., Vandemark, D., Vlahos, P., Wallace, D. W. R., Wanninkhof, R., and Watson, A. J.: An update to the Surface Ocean CO₂ Atlas (SOCAT version 2), *Earth Syst. Sci. Data Discuss.*, 6, 465–512, doi:10.5194/essdd-6-465-2013, 2013.

Ballantyne, A. P., Alden, C. B., Miller, J. B., Tans, P. P., and White, J. W. C.: Increase in observed net carbon dioxide uptake by land and oceans during the last 50 years, *Nature*, 488, 70–72, 2012.

Bastviken, D., Tranvik, L. J., Downing, J. A., Crill, P. M., and Enrich-Prast, A.: Fresh-water methane emissions offset the continental carbon sink, *Science*, 331, 6013, doi:10.1126/science.1196808, 2011.

Battin, T. J., Luysaert, S., Kaplan, L. A., Aufdenkampe, A. K., Richter, A., and Tranvik, L. J.: The boundless carbon cycle, *Nat. Geosci.*, 2009, 598–600, 2009.

Best, M. J., Pryor, M., Clark, D. B., Rooney, G. G., Essery, R. L. H., Ménard, C. B., Edwards, J. M., Hendry, M. A., Porson, A., Gedney, N., Mercado, L. M., Sitch, S., Blyth, E., Boucher, O., Cox, P. M., Grimmond, C. S. B., and Harding, R. J.: The Joint UK Land Environment Simulator (JULES), model description – Part 1: Energy and water fluxes, *Geosci. Model Dev.*, 4, 677–699, doi:10.5194/gmd-4-677-2011, 2011.

Boden, T. A., Marland, G., and Andres, R. J.: Global, Regional, and National Fossil-Fuel CO₂ Emissions, Oak Ridge National Laboratory, US Department of Energy, Oak Ridge, Tenn., USA, http://cdiac.ornl.gov/trends/emis/meth_reg.html, 2013.

Borges, A. V. and Abril, G.: Carbon dioxide and methane dynamics in estuaries, in: *Treatise on Estuarine and Coastal Science*, edited by: Wolanski, E. and McLusky, D. S., Academic Press, 2011.

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- BP: Statistical Review of World Energy 2013, <http://www.bp.com/en/global/corporate/about-bp/statistical-review-of-world-energy-2013.html>, last access: August 2013.
- Bradshaw, C. J. A.: Little left to lose: deforestation and forest degradation in Australia since European colonization, *J. Plant Ecol.*, 5, 109–120, 2012.
- 5 Bruno, M. and Joos, F.: Terrestrial carbon storage during the past 200 years: A monte carlo analysis of CO₂ data from ice core and atmospheric measurements, *Global Biogeochem. Cy.*, 11, 111–124, 1997.
- Buitenhuis, E. T., Rivkin, R. B., Sailley, S., and Le Quéré, C.: Biogeochemical fluxes through microzooplankton, *Global Biogeochem. Cy.*, 24, GB4015, doi:10.1029/2009GB003601, 2010.
- 10 Canadell, J. G., Le Quéré, C., Raupach, M. R., Field, C. B., Buitenhuis, E. T., Ciais, P., Conway, T. J., Gillett, N. P., Houghton, R. A., and Marland, G.: Contributions to accelerating atmospheric CO₂ growth from economic activity, carbon intensity, and efficiency of natural sinks, *P. Natl. Acad. Sci. USA*, 104, 18866–18870, 2007.
- Canadell, J., Ciais, P., Sabine, C., and Joos, F.: REgional Carbon Cycle Assessment and Processes (RECCAP), http://www.biogeosciences.net/special_issue107.html, 2012–2013.
- 15 Ciais, P., Sabine, C., Govindasamy, B., Bopp, L., Brovkin, V., Canadell, J., Chhabra, A., DeFries, R., Galloway, J., Heimann, M., Jones, C., Le Quéré, C., Myneni, R., Piao, S., and Thornton, P.: Chapter 6: Carbon and Other Biogeochemical Cycles, in: *Climate Change 2013 The Physical Science Basis*, edited by: Stocker, T., Qin, D., and Plattner, G.-K., Cambridge University Press, Cambridge, 2013.
- 20 Clark, D. B., Mercado, L. M., Sitch, S., Jones, C. D., Gedney, N., Best, M. J., Pryor, M., Rooney, G. G., Essery, R. L. H., Blyth, E., Boucher, O., Harding, R. J., Huntingford, C., and Cox, P. M.: The Joint UK Land Environment Simulator (JULES), model description – Part 2: Carbon fluxes and vegetation dynamics, *Geosci. Model Dev.*, 4, 701–722, doi:10.5194/gmd-4-701-2011, 2011.
- 25 Cole, J. J., Prairie, Y. T., Caraco, N. F., McDowell, W. H., Tranvik, L. J., Striegl, R. G., Duarte, C. M., Kortelainen, P., Downing, J. A., Middelburg, J. J., and Melack, J.: Plumbing the global carbon cycle: Integrating inland waters into the terrestrial carbon budget, *Ecosystems*, 2007, 171–184, 2007.
- 30 Davis, S. J. and Caldeira, K.: Consumption-based accounting of CO₂ emissions, *P. Natl. Acad. Sci.*, 107, 5687–5692, 2010.
- Davis, S. J., Peters, G. P., and Caldeira, K.: The supply chain of CO₂ emissions, *P. Natl. Acad. Sci.*, 108, 18554–18559, 2011.

**Global carbon budget
2013**

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Denman, K. L., Brasseur, G., Chidthaisong, A., Ciais, P., Cox, P. M., Dickinson, R. E., Hauglustaine, D., Heinze, C., Holland, E., Jacob, D., Lohmann, U., Ramachandran, S., Leite da Silva Dias, P., Wofsy, S. C., and Zhang, X.: Couplings Between Changes in the Climate System and Biogeochemistry, Intergovernmental Panel on Climate Change, ISBN: 978-0-521-70596-7, 499–587, 2007.

5 Dlugokencky, E. and Tans, P.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL), <http://www.esrl.noaa.gov/gmd/ccgg/trends>, last access: 8 October 2013.

10 Doney, S. C., Lima, I., Feely, R. A., Glover, D. M., Lindsay, K., Mahowald, N., Moore, J. K., and Wanninkhof, R.: Mechanisms governing interannual variability in upper-ocean inorganic carbon system and air–sea CO₂ fluxes: Physical climate and atmospheric dust, *Deep-Sea Res. Pt. II*, 56, 640–655, 2009.

Durant, A. J., Le Quéré, C., Hope, C., and Friend, A. D.: Economic value of improved quantification in global sources and sinks of carbon dioxide, *Phil. Trans. A*, 269, 1967–1979, 2010.

15 Earles, J. M., Yeh, S., and Skog, K. E.: Timing of carbon emissions from global forest clearance, *Nature Climate Change*, 2, 682–685, 2012.

El-Masri, B., Barman, R., Meiyappan, P., Song, Y., Liang, M., and Jain, A. K.: Carbon dynamics in the Amazonian Basin: Integration of eddy covariance and ecophysiological data with a land surface model, *Agr. Forest Meteorol.*, 182–183, 156–167, 2013.

20 Erb, K.-H., Kastner, T., Luysaert, S., Houghton, R. A., Kuemmerle, T., Olofsson, P., and Haberl, H.: Bias in the attribution of forest carbon sinks, *Nature Climate Change*, 3, 854–856, 2013.

Etheridge, D. M., Steele, L. P., Langenfelds, R. L., and Francey, R. J.: Natural and anthropogenic changes in atmospheric CO₂ over the last 1000 years from air in Antarctic ice and firn, *J. Geophys. Res.*, 101, 4115–4128, 1996.

25 FAO: Global Forest Resource Assessment 2010, 378 pp., 2010.

FAOSTAT: Food and Agriculture Organization Statistics Division, <http://faostat3.fao.org/faostat-gateway/go/to/home/E>, 2010.

Friedlingstein, P., Houghton, R. A., Marland, G., Hackler, J., Boden, T. A., Conway, T. J., Canadell, J. G., Raupach, M. R., Ciais, P., and Le Quéré, C.: Update on CO₂ emissions, *Nat. Geosci.*, 3, 811–812, 2010.

30 Galbraith, E. D., Gnanadesikan, A., Dunne, J. P., and Hiscock, M. R.: Regional impacts of iron-light colimitation in a global biogeochemical model, *Biogeosciences*, 7, 1043–1064, doi:10.5194/bg-7-1043-2010, 2010.

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Gasser, T. and Ciais, P.: A theoretical framework for the net land-to-atmosphere CO₂ flux and its implications in the definition of “emissions from land-use change”, *Earth Syst. Dynam.*, 4, 171–186, doi:10.5194/esd-4-171-2013, 2013.

GCP: The Global Carbon Budget 2007, http://lgmwebweb.env.uea.ac.uk/lequere/co2/2007/carbon_budget_2007.htm, last access: November 2013.

Giglio, L., Randerson, J. T., van der Werf, G. R., Kasibhatla, P. S., Collatz, G. J., Morton, D. C., and DeFries, R. S.: Assessing variability and long-term trends in burned area by merging multiple satellite fire products, *Biogeosciences*, 7, 1171–1186, doi:10.5194/bg-7-1171-2010, 2010.

Gitz, V. and Ciais, P.: Amplifying effects of land-use change on future atmospheric CO₂ levels, *Global Biogeochem. Cy.*, 17, 1024, doi:10.1029/2002GB001963, 2003.

Gregg, J. S., Andres, R. J., and Marland, G.: China: Emissions pattern of the world leader in CO₂ emissions from fossil fuel consumption and cement production, *Geophys. Res. Lett.*, 35, L08806, doi:10.1029/2007GL032887, 2008.

Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: Updated high-resolution grids of monthly climatic observations, *Int. J. Climatol.*, doi:10.1002/joc.3711, 2013a.

Harris, I., Jones, P. D., Osborn, T. J., and Lister, D. H.: CRU TS3.21: Climatic Research Unit (CRU) Time-Series (TS) Version 3.21 of High Resolution Gridded Data of Monthly-by-month Variation in Climate (Jan. 1901–Dec. 2012), doi:10.5285/D0E1585D-3417-485F-87AE-4FCECF10A992, 2013b.

Hertwich, E. G. and Peters, G. P.: Carbon Footprint of Nations: A Global, Trade-Linked Analysis, *Environ. Sci. Technol.*, 43, 6414–6420, 2009.

Houghton, R. A.: Revised estimates of the annual net flux of carbon to the atmosphere from changes in land use and land management 1850–2000, *Tellus B*, 55, 378–390, 2003.

Houghton, R. A. and Hackler, J. L.: Annual Flux of Carbon from Land Use and Land-Cover Change 1850 to 2010, *Global Biogeochem. Cy.*, in review, 2013.

Houghton, R. A., House, J. I., Pongratz, J., van der Werf, G. R., DeFries, R. S., Hansen, M. C., Le Quéré, C., and Ramankutty, N.: Carbon emissions from land use and land-cover change, *Biogeosciences*, 9, 5125–5142, doi:10.5194/bg-9-5125-2012, 2012.

Hurttt, G. C., Chini, L. P., Froliking, S., Betts, R. A., Feddema, J., Fischer, G., Fisk, J. P., Hibbard, K., Houghton, R. A., Janetos, A., Jones, C. D., Kindermann, G., Kinoshita, T., Klein Goldewijk, K., Riahi, K., Shevliakova, E., Smith, S., Stehfest, E., Thomson, A., Thornton, P., van Vuuren, D. P., and Wang, Y. P.: Harmonization of land-use scenarios for the period 1500–2100: 600

**Global carbon budget
2013**

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands, *Climatic Change*, 109, 117–161, 2011.

IEA/OECD: CO₂ emissions from fuel combustion, Paris, 2012.

Ilyina, T., Six, K., Segsneider, J., Maier-Reimer, E., Li, H., and Núñez-Riboni, I.: The global ocean biogeochemistry model HAMOCC: Model architecture and performance as component of the MPI-Earth System Model in different CMIP5 experimental realizations, *Journal of Advances in Modeling Earth Systems*, 5, 287–315, doi:10.1002/jame.20017, 2013.

IMF: World Economic Outlook of the International Monetary Fund, <http://www.imf.org/external/ns/cs.aspx?id=29>, last access: 10 October 2013.

Ito, A. and Inatomi, M.: Use of a process-based model for assessing the methane budgets of global terrestrial ecosystems and evaluation of uncertainty, *Biogeosciences*, 9, 759–773, doi:10.5194/bg-9-759-2012, 2012.

Jacobson, A. R., Mikaloff Fletcher, S. E., Gruber, N., Sarmiento, J. L., and Gloor, M.: A joint atmosphere-ocean inversion for surface fluxes of carbon dioxide: 1. Methods and global-scale fluxes, *Global Biogeochem. Cy.*, 21, GB1019, doi:10.1029/2005GB002556, 2007.

Jain, A. K., Meiyappan, P., Song, Y., and House, J. I.: CO₂ Emissions from Land-Use Change Affected More by Nitrogen Cycle, than by the Choice of Land Cover Data, *Glob. Change Biol.*, 19, 2893–2906, doi:10.1111/gcb.12207, 2013.

Joos, F. and Spahni, R.: Rates of change in natural and anthropogenic radiative forcing over the past 20,000 years, *P. Natl. Acad. Sci.*, 105, 1425–1430, 2008.

Joos, F., Roth, R., Fuglestedt, J. S., Peters, G. P., Enting, I. G., von Bloh, W., Brovkin, V., Burke, E. J., Eby, M., Edwards, N. R., Friedrich, T., Frölicher, T. L., Halloran, P. R., Holden, P. B., Jones, C., Kleinen, T., Mackenzie, F. T., Matsumoto, K., Meinshausen, M., Plattner, G.-K., Reisinger, A., Segsneider, J., Shaffer, G., Steinacher, M., Strassmann, K., Tanaka, K., Timmermann, A., and Weaver, A. J.: Carbon dioxide and climate impulse response functions for the computation of greenhouse gas metrics: a multi-model analysis, *Atmos. Chem. Phys.*, 13, 2793–2825, doi:10.5194/acp-13-2793-2013, 2013.

Kato, E., Kinoshita, T., Ito, A., Kawamiya, M., and Yamagata, Y.: Evaluation of spatially explicit emission scenario of land-use change and biomass burning using a process-based biogeochemical model, *J. Land Use Sci.*, 8, 104–122, 2013.

Keeling, C. D., Bacastow, R. B., Bainbridge, A. E., Ekdhal, C. A., Guenther, P. R., and Waterman, L. S.: Atmospheric carbon dioxide variations at Mauna Loa Observatory, Hawaii, *Tellus*, 28, 538–551, 1976.

**Global carbon budget
2013**

C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- Khatiwala, S., Primeau, F., and Hall, T.: Reconstruction of the history of anthropogenic CO₂ concentrations in the ocean, *Nature*, 462, 346–350, 2009.
- Khatiwala, S., Tanhua, T., Mikaloff Fletcher, S., Gerber, M., Doney, S. C., Graven, H. D., Gruber, N., McKinley, G. A., Murata, A., Ríos, A. F., and Sabine, C. L.: Global ocean storage of anthropogenic carbon, *Biogeosciences*, 10, 2169–2191, doi:10.5194/bg-10-2169-2013, 2013.
- Kirschke, S., Bousquet, P., Ciais, P., Saunois, M., Canadell, J. G., Dlugokencky, E. J., Bergamaschi, P., Bergmann, D., Blake, D. R., Bruhwiler, L., Cameron Smith, P., Castaldi, S., Chevallier, F., Feng, L., Fraser, A., Heimann, M., Hodson, E. L., Houweling, S., Josse, B., Fraser, P. J., Krummel, P. B., Lamarque, J., Langenfelds, R. L., Le Quéré, C., Naik, V., O'Doherty, S., Palmer, P. I., Pison, I., Plummer, D., Poulter, B., Prinn, R. G., Rigby, M., Ringeval, B., Santini, M., Schmidt, M., Shindell, D. T., Simpson, I. J., Spahni, R., Steele, L. P., Strode, S. A., Sudo, K., Szopa, S., van der Werf, G. R., Voulgarakis, A., van Weele, M., Weiss, R. F., Williams, J. E., and Zeng, G.: Three decades of global methane sources and sinks, *Nat. Geosci.*, 6, 813–823, doi:10.1038/ngeo1955, 2013.
- Klein Goldewijk, K., Beusen, A., van Drecht, G., and de Vos, M.: The HYDE 3.1 spatially explicit database of human-induced global land-use change over the past 12,000 years, *Global Ecol. Biogeogr.*, 20, 73–86, 2011.
- Krumins, V., Gehlen, M., Arndt, S., Van Cappellen, P., and Regnier, P.: Dissolved inorganic carbon and alkalinity fluxes from coastal marine sediments: model estimates for different shelf environments and sensitivity to global change, *Biogeosciences*, 10, 371–398, doi:10.5194/bg-10-371-2013, 2013.
- Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass burning emissions of reactive gases and aerosols: methodology and application, *Atmos. Chem. Phys.*, 10, 7017–7039, doi:10.5194/acp-10-7017-2010, 2010.
- Laruelle, G. G., Durr, H. H., Slomp, C. P., and Borges, A. V.: Evaluation of sinks and sources of CO₂ in the global coastal ocean using a spatially-explicit typology of estuaries and continental shelves, *Geophys. Res. Lett.*, 37, L15607, doi:10.1029/2010GL043691, 2010.
- Le Quéré, C.: Closing the global budget for CO₂, *Global Change*, 74, 28–31, 2009.
- Le Quéré, C., Raupach, M. R., Canadell, J. G., Marland, G., Bopp, L., Ciais, P., Conway, T. J., Doney, S. C., Feely, R. A., Foster, P., Friedlingstein, P., Gurney, K., Houghton, R. A., House,

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



J. I., Huntingford, C., Levy, P. E., Lomas, M. R., Majkut, J., Metz, N., Ometto, J. P., Peters, G. P., Prentice, I. C., Randerson, J. T., Running, S. W., Sarmiento, J. L., Schuster, U., Sitch, S., Takahashi, T., Viovy, N., van der Werf, G. R., and Woodward, F. I.: Trends in the sources and sinks of carbon dioxide, *Nat. Geosci.*, 2, 831–836, 2009.

5 Le Quéré, C., Andres, R. J., Boden, T., Conway, T., Houghton, R. A., House, J. I., Marland, G., Peters, G. P., van der Werf, G. R., Ahlström, A., Andrew, R. M., Bopp, L., Canadell, J. G., Ciais, P., Doney, S. C., Enright, C., Friedlingstein, P., Huntingford, C., Jain, A. K., Jourdain, C., Kato, E., Keeling, R. F., Klein Goldewijk, K., Levis, S., Levy, P., Lomas, M., Poulter, B., Raupach, M. R., Schwinger, J., Sitch, S., Stocker, B. D., Viovy, N., Zaehle, S., and Zeng, N.:
10 The global carbon budget 1959–2011, *Earth Syst. Sci. Data*, 5, 165–185, doi:10.5194/essd-5-165-2013, 2013.

Manning, A. C. and Keeling, R. F.: Global oceanic and land biotic carbon sinks from the Scripps atmospheric oxygen flask sampling network, *Tellus B*, 58, 95–116, 2006.

Marland, G.: Uncertainties in accounting for CO₂ from fossil fuels, *J. Ind. Ecol.*, 12, 136–139,
15 2008.

Marland, G., Andres, R. J., Blasing, T. J., Boden, T. A., Broniak, C. T., Gregg, J. S., Losey, L. M., and Treanton, K.: Energy, industry and waste management activities: An introduction to CO₂ emissions from fossil fuels, in: A report by the US Climate Change Science Program and the Subcommittee on Global Change Research, in *The First State of the Carbon Cycle Report (SOCCR): The North American Carbon Budget and Implications for the Global Carbon Cycle*, edited by: King, A. W., Dilling, L., Zimmerman, G. P., Fairman, D. M., Houghton, R. A., Marland, G., Rose, A. Z., and Wilbanks, T. J., Asheville, NC, 2007.

Marland, G., Hamal, K., and Jonas, M.: How Uncertain Are Estimates of CO₂ Emissions?, *J. Ind. Ecol.*, 13, 4–7, 2009.

25 Masarie, K. A. and Tans, P. P.: Extension and integratio of atmospheric carbon dioxide data into a globally consistent measurement record, *J. Geophys. Res.-Atmos.*, 100, 11593–11610, 1995.

McNeil, B. I., Matear, R. J., Key, R. M., Bullister, J. L., and Sarmiento, J. L.: Anthropogenic CO₂ uptake by the ocean based on the global chlorofluorocarbon data set, *Science*, 299,
30 235–239, 2003.

Mikaloff Fletcher, S. E., Gruber, N., Jacobson, A. R., Doney, S. C., Dutkiewicz, S., Gerber, M., Follows, M., Joos, F., Lindsay, K., Menemenlis, D., Mouchet, A., Müller, S. A., and Sarmiento,

**Global carbon budget
2013**

C. Le Quéré et al.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


- J. L.: Inverse estimates of anthropogenic CO₂ uptake, transport, and storage by the oceans, *Global Biogeochem. Cy.*, 20, GB2002, doi:10.1029/2005GB002530, 2006.
- Narayanan, B., Aguiar, A., and McDougall, R.: Global Trade, Assistance, and Production: The GTAP 8 Data Base, <https://www.gtap.agecon.purdue.edu/databases/v8/default.asp>, last access: September 2013.
- NOAA/ESRL: NOAA/ESRL calculation of global means, http://www.esrl.noaa.gov/gmd/ccgg/about/global_means.html, last access: August 2013.
- Oleson, K., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., Subin, Z., Swenson, S., Thornton, P., Bozbiyik, A., Fisher, R., Heald, C., Kluzek, E., Lamarque, J., Lawrence, P., Leung, L., Lipscomb, W., Muszala, S., Ricciuto, D., Sacks, W., Tang, J., and Yang, Z.: Technical Description of version 4.5 of the Community Land Model (CLM), NCAR, 2013.
- Park, G. H., Wanninkhof, R., Doney, S. C., Takahashi, T., Lee, K., Feely, R. A., Sabine, C. L., Trinanes, J., and Lima, I. D.: Variability of global net sea-air CO₂ fluxes over the last three decades using empirical relationships, *Tellus B*, 62, 352–368, 2010.
- Peters, G. P. and Hertwich, E. G.: Post-Kyoto Greenhouse Gas Inventories: Production versus Consumption, *Climatic Change*, 85, 51–66, 2008.
- Peters, G. P., Andrew, R., and Lennos, J.: Constructing a multi-regional input-output table using the GTAP database, *Economic Systems Research*, 23, 131–152, 2011a.
- Peters, G. P., Minx, J. C., Weber, C. L., and Edenhofer, O.: Growth in emission transfers via international trade from 1990 to 2008, *P. Natl. Acad. Sci. USA*, 108, 8903–8908, 2011b.
- Peters, G. P., Davis, S. J., and Andrew, R.: A synthesis of carbon in international trade, *Biogeosciences*, 9, 3247–3276, doi:10.5194/bg-9-3247-2012, 2012a.
- Peters, G. P., Marland, G., Le Quéré, C., Boden, T. A., Canadell, J. G., and Raupach, M. R.: Correspondence: Rapid growth in CO₂ emissions after the 2008–2009 global financial crisis, *Nature Climate Change*, 2, 2–4, 2012b.
- Peters, G. P., Andrew, R. M., Boden, T., Canadell, J. G., Ciais, P., Le Quéré, C., Marland, G., Raupach, M. R., and Wilson, C.: The challenge to keep global warming below 2 °C, *Nature Clim. Change*, 3, 4–6, 2013.
- Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metz, N., Sabine, C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F. P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-Mountford, N., Heinze, C., Hood,

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S. D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J., Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125–143, doi:10.5194/essd-5-125-2013, 2013.

Pongratz, J., Reick, C. H., Raddatz, T., and Claussen, M.: Effects of anthropogenic land cover change on the carbon cycle of the last millennium, *Global Biogeochem. Cy.*, 23, GB4001, doi:10.1029/2009GB003488, 2009.

Prentice, I. C., Farquhar, G. D., Fasham, M. J. R., Goulden, M. L., Heimann, M., Jaramillo, V. J., Kheshgi, H. S., Le Quéré, C., Scholes, R. J., and Wallace, D. W. R.: The Carbon Cycle and Atmospheric Carbon Dioxide, in: *Climate Change 2001: The Scientific Basis. Contribution of Working Group I to the Third Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Houghton, J. T., Ding, Y., Griggs, D. J., Noguer, M., van der Linden, P. J., Dai, X., Maskell, K., and Johnson, C. A., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2001.

Raupach, M. R., Marland, G., Ciais, P., Le Quéré, C., Canadell, J. G., Klepper, G., and Field, C. B.: Global and regional drivers of accelerating CO₂ emissions, *P. Natl. Acad. Sci. USA*, 104, 10288–10293, 2007.

Regnier, P., Friedlingstein, P., Ciais, P., Mackenzie, F. T., Gruber, N., Janssens, I. A., Laruelle, G. G., Lauerwald, R., Luysaert, S., Andersson, A. J., Arndt, S., Arnosti, C., Borges, A. V., Dale, A. W., Gallego-Sala, A., Goddérís, Y., Goossens, N., Hartmann, J., Heinze, C., Ilyina, T., Joos, F., La Rowe, D. E., Leifeld, J., Meysman, F. J. R., Munhoven, G., Raymond, P. A., Spahni, R., Suntharalingam, P., and Thullner M.: Anthropogenic perturbation of the carbon fluxes from land to ocean, *Nat. Geosci.*, 6, 597–607, 2013.

Rhein, M., Rintoul, S. R., Aoki, S., Campos, E., Chambers, D., Feely, R. A., Gulev, S., Johnson, G. C., Josey, S. A., Kostianoy, A., Mauritzen, C., Roemmich, D., Talley, L. D., and Wang, F.: Chapter 3: Observations: Ocean, in: *Climate Change 2013 The Physical Science Basis*, Cambridge University Press, 2013.

**Global carbon budget
2013**

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Rödenbeck, C., Keeling, R. F., Bakker, D. C. E., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-ocean $p\text{CO}_2$ and sea-air CO_2 flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Sci.*, 9, 193–216, doi:10.5194/os-9-193-2013, 2013.

Rypdal, K., Paciomik, N., Eggleston, S., Goodwin, J., Irving, W., Penman, J., and Woodfield, M.: Chapter 1 Introduction to the 2006 Guidelines, in: 2006 IPCC Guidelines for National Greenhouse Gas Inventories, edited by: Eggleston, S., Buendia, L., Miwa, K., Ngara, T., and Tanabe, K., Institute for Global Environmental Strategies (IGES), Hayama, Kanagawa, Japan, 2006.

Schimel, D., Alves, D., Enting, I., Heimann, M., Joos, F., Raynaud, D., Wigley, T., Prater, M., Derwent, R., Ehhalt, D., Fraser, P., Sanhueza, E., Zhou, X., Jonas, P., Charlson, R., Rodhe, H., Sadasivan, S., Shine, K. P., Fouquart, Y., Ramaswamy, V., Solomon, S., Srinivasan, J., Albritton, D., Derwent, R., Isaksen, I., Lal, M., and Wuebbles, D.: Radiative Forcing of Climate Change, in: *Climate Change 1995 The Science of Climate Change. Contribution of Working Group I to the Second Assessment Report of the Intergovernmental Panel on Climate Change*, edited by: Houghton, J. T., Meira Rilho, L. G., Callander, B. A., Harris, N., Kattenberg, A., and Maskell, K., Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 1995.

Scripps: The Keeling Curve, <http://keelingcurve.ucsd.edu/>, last access: 7 November 2013.

Shevliakova, E., Pacala, S., Malyshev, S., Hurtt, G., Milly, P., Caspersen, J., Sentman, L., Fisk, J., Wirth, C., and Crevoisier, C.: Carbon cycling under 300 years of land use change: Importance of the secondary vegetation sink, *Global Biogeochem. Cy.*, 23, GB2022, doi:10.1029/2007GB003176, 2009.

Sitch, S., Smith, B., Prentice, I. C., Arneeth, A., Bondeau, A., Cramer, W., Kaplan, J. O., Levis, S., Lucht, W., Sykes, M. T., Thonicke, K., and Venevsky, S.: Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model, *Glob. Change Biol.*, 9, 161–185, 2003.

Smith, B., Prentice, I. C., and Sykes, M. T.: Representation of vegetation dynamics in the modelling of terrestrial ecosystems: comparing two contrasting approaches with European climate space, *Global Ecol. Biogeogr.*, 10, 621–637, doi:10.1046/j.1466-822X.2001.t01-1-00256.x, 2001.

Stocker, B. D., Strassmann, K., and Joos, F.: Sensitivity of Holocene atmospheric CO_2 and the modern carbon budget to early human land use: analyses with a process-based model, *Biogeosciences*, 8, 69–88, doi:10.5194/bg-8-69-2011, 2011.

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- Stocker, T., Qin, D., and Plattner, G.-K.: Climate Change 2013 The Physical Science Basis, Cambridge University Press, 2013.
- Tans, P. and Keeling, R. F.: Trends in atmospheric carbon dioxide, National Oceanic & Atmospheric Administration, Earth System Research Laboratory (NOAA/ESRL) & Scripps Institution of Oceanography, <http://www.esrl.noaa.gov/gmd/ccgg/trends/> and <http://scrippsco2.ucsd.edu/>, last access: 7 October 2013.
- Tjiputra, J. F., Roelandt, C., Bentsen, M., Lawrence, D. M., Lorentzen, T., Schwinger, J., Seland, Ø., and Heinze, C.: Evaluation of the carbon cycle components in the Norwegian Earth System Model (NorESM), *Geosci. Model Dev.*, 6, 301–325, doi:10.5194/gmd-6-301-2013, 2013.
- Tranvik, L. J., Downing, J. A., Cotner, J. B., Loiselle, S. A., Striegl, R. G., Ballatore, T. J., Dillon, P., Finlay, K., Fortino, K., Knoll, L. B., Kortelainen, P. L., Kutser, T., Larsen, S., Laurion, I., Leech, D. M., McCallister, S. L., McKnight, D. M., Melack, J. M., Overholt, E., Porter, J. A., Prairie, Y., Renwick, W. H., Roland, F., Sherman, B. S., Schindler, D. W., Sobek, S., Tremblay, A., Vanni, M. J., Verschoor, A. M., von Wachenfeldt, E., and Weyhenmeyer, G. A.: Lakes and reservoirs as regulators of carbon cycling and climate, *Limnol. Oceanogr.*, 54, 2298–2314, 2009.
- UN: United Nations Statistics Division – Energy Statistics, <http://unstats.un.org/unsd/energy/>, last access: August 2013a.
- UN: United Nations Statistics Division – Industry Statistics, <http://unstats.un.org/unsd/industry/default.asp>, last access: August 2013b.
- UN: United Nations Statistics Division – National Accounts Main Aggregates Database, <http://unstats.un.org/unsd/snaama/Introduction.asp>, last access: August 2013c.
- US Geological Survey: Mineral Commodities Summaries, US Geological Survey, 2013.
- van der Werf, G. R., Dempewolf, J., Trigg, S. N., Randerson, J. T., Kasibhatla, P., Giglio, L., Murdiyarso, D., Peters, W., Morton, D. C., Collatz, G. J., Dolman, A. J., and DeFries, R. S.: Climate regulation of fire emissions and deforestation in equatorial Asia, *P. Natl. Acad. Sci. USA*, 15, 20350–20355, 2008.
- van der Werf, G. R., Randerson, J. T., Giglio, L., Collatz, G. J., Mu, M., Kasibhatla, P. S., Morton, D. C., DeFries, R. S., Jin, Y., and van Leeuwen, T. T.: Global fire emissions and the contribution of deforestation, savanna, forest, agricultural, and peat fires (1997–2009), *Atmos. Chem. Phys.*, 10, 11707–11735, doi:10.5194/acp-10-11707-2010, 2010.

Global carbon budget 2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



- van Minnen, J. G., Goldewijk, K. K., Stehfest, E., Eickhout, B., van Drecht, G., and Leemans, R.: The importance of three centuries of land-use change for the global and regional terrestrial carbon cycle, *Climatic Change*, 97, 123–144, 2009.
- van Oss, H. G.: Cement, US Geological Survey, June, 2013.
- 5 Wanninkhof, R., Park, G. -H., Takahashi, T., Sweeney, C., Feely, R., Nojiri, Y., Gruber, N., Doney, S. C., McKinley, G. A., Lenton, A., Le Quéré, C., Heinze, C., Schwinger, J., Graven, H., and Khatiwala, S.: Global ocean carbon uptake: magnitude, variability and trends, *Biogeosciences*, 10, 1983–2000, doi:10.5194/bg-10-1983-2013, 2013.
- 10 Watson, R. T., Rodhe, H., Oeschger, H., and Siegenthaler, U.: Greenhouse Gases and Aerosols, in: *Climate Change: The IPCC Scientific Assessment*. Intergovernmental Panel on Climate Change (IPCC), edited by: Houghton, J. T., Jenkins, G. J., and Ephraums, J. J., Cambridge University Press, Cambridge, 1990.
- Zaehle, S. and Friend, A. D.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 1. Model description, site-scale evaluation, and sensitivity to parameter estimates, *Global Biogeochem. Cy.*, 24, GB1005, doi:10.1029/2009GB003521, 2010.
- 15 Zaehle, S., Friend, A. D., Friedlingstein, P., Dentener, F., Peylin, P., and Schulz, M.: Carbon and nitrogen cycle dynamics in the O-CN land surface model: 2. Role of the nitrogen cycle in the historical terrestrial carbon balance, *Global Biogeochem. Cy.*, 24, GB1006, doi:10.1029/2009GB003522, 2010.
- 20 Zaehle, S., Ciais, P., Friend, A. D., and Prieur, V.: Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions, *Nat. Geosci.*, 4, 601–605, 2011.

Global carbon budget 2013

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 1. Factors used to convert carbon in various units (by convention, Unit 1 = Unit 2 × conversion).

Unit 1	Unit 2	Conversion	Source
GtC (gigatonnes of Carbon)	ppm (parts per million)	2.120	Joos et al. (2013)
GtC (gigatonnes of Carbon)	PgC (petagrammes of Carbon)	1	SI unit conversion
GtCO ₂ (gigatonnes of Carbon Dioxide)	GtC (gigatonnes of Carbon)	3.664	44/12 in mass equivalent
GtC (gigatonnes of Carbon)	MtC (megatonnes of Carbon)	1000	SI unit conversion

Global carbon budget 2013

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 2. How to cite the individual components of the global carbon budget presented here.

Component	Primary reference
Territorial fossil-fuel and cement emissions (E_{FF}) global, by fuel type, and by country	Boden et al. (2013; CDIAC: cdiac.ornl.gov/trends/emis/meth_reg.html)
Consumption-based fossil-fuel and cement emissions (E_{FF}) by country (consumption)	Peters et al. (2011b) updated as described in this paper
Land-use change emissions (E_{LUC})	Houghton and Hackler (2013)
Atmospheric CO ₂ growth rate	Dlugokencky and Tans (2013; NOAA/ESRL: www.esrl.noaa.gov/gmd/ccgg/trends/)
Ocean and land CO ₂ sinks (S_{OCEAN} and S_{LAND})	this paper for S_{OCEAN} and S_{LAND} and references in Table 5 for individual models

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)**Table 3.** Data sources used to compute each component of the global carbon budget.

Component	Process	Data source	Data reference
E_{FF}	Fossil-fuel combustion and gas flaring	UN Statistics Division to 2010 BP for 2011–2012	UN (2013a, b) BP (2013)
	Cement production	US Geological Survey	van Oss (2013) US Geological Survey (2013)
E_{LUC}	Land cover change (deforestation, afforestation, and forest regrowth)	Forest Resource Assessment (FRA) of the Food and Agriculture Organisation (FAO)	FAO (2010)
	Wood harvest	FAO Statistics Division	FAOSTAT (2010)
	Shifting agriculture	FAO FRA and Statistics Division	FAO (2010) FAOSTAT (2010)
G_{ATM}	Change in atmospheric CO ₂ concentration	1959–1980: CO ₂ Program at Scripps Institution of Oceanography and other research groups	Keeling et al. (1976)
		1980–2011: US National Oceanic and Atmospheric Administration Earth System Research Laboratory	Dlugokencky and Tans (2013) and Ballantyne et al. (2012)
S_{OCEAN}	Uptake of anthropogenic CO ₂	1990–1999 average: indirect estimates based on CFCs, atmospheric O ₂ , and other tracer observations	Manning and Keeling (2006), McNeil et al. (2003) and Mikaloff Fletcher et al. (2006) as assessed by the IPCC Denman et al. (2007)
	Impact of increasing atmospheric CO ₂ , and climate change and variability	Ocean models	Table 5
S_{LAND}	Response of land vegetation to: Increasing atmospheric CO ₂ concentration Climate change and variability Other environmental changes	Budget residual	

Global carbon budget 2013

C. Le Quéré et al.

Table 4. Comparison of the processes included in the E_{LUC} of the global carbon budget and the DGVMs. See Table 5 for model references. All models include deforestation, afforestation, and forest regrowth after abandonment of agriculture.

	Bookkeeping	CLM4.5BGC	ISAM	JULES	LPJ-GUESS	LPJ	LPX-Bern	ORCHIDEE-CN	VISIT
Wood harvest and forest degradation	yes	yes	yes	no	no	no	no	no	yes*
Shifting cultivation	yes	yes	no	no	no	no	no	no	yes
Cropland harvest	yes	yes	no	no	yes	no	yes	yes	yes
Peat fires	no	yes	no	no	no	no	no	no	no
Fire simulation and/or suppression	for US only	yes	no	no	yes	yes	yes	no	yes
Climate change and variability	no	yes	yes	yes	yes	yes	yes	yes	yes
CO ₂ fertilisation	no	yes	yes	yes	yes	yes	yes	yes	yes
Nitrogen fertilisation and/or dynamics	no	yes	yes	no	no	no	yes	yes	no

* Wood stems are harvested according to the land-use data.

[Title Page](#)
[Abstract](#)
[Instruments](#)
[Data Provenance & Structure](#)
[Tables](#)
[Figures](#)
[◀](#)
[▶](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


Table 5. References for the process models and data products included in Fig. 3.

Model/Data name	Reference
Dynamic Global Vegetation Models providing E_{LUC} and S_{LAND}	
CLM4.5BGC ^a	Oleson et al. (2013)
ISAM	Jain et al. (2013) ^b
JULES ^c	Clarke et al. (2011) ^d
LPJ-GUESS	Smith et al. (2001)
LPJ ^e	Sitch et al. (2003)
LPX-Bern	Stocker et al. (2011)
ORCHIDEE-CN	Zaehle and Friend (2010) ^f
VISIT	Kato et al. (2013) ^g
Ocean Biogeochemistry Models providing S_{OCEAN}	
NEMO-PlankTOM5	Buitenhuis et al. (2010) ^h
LSCE	Aumont and Bopp (2006)
CCSM-BEC	Doney et al. (2009)
MICOM-HAMOCC	Assmann et al. (2010) ⁱ
MPI-MET	Ilyina et al. (2013)
BLING ^j	Galbraith (2009)
Ocean CO ₂ Data Products informing ocean analysis	
Park	Park et al. (2010) ^k
Rödenbeck	Rödenbeck et al. (2013) ^l

^a Community Land Model 4.5, ^b see also El-Masri et al. (2013), ^c Joint UK Land Environment Simulator, ^d see also Best et al. (2011), ^e Lund-Potsdam-Jena, ^f see also Zaehle et al. (2010), ^g see also Ito and Inatomi (2012), ^h with no nutrient restoring below the mixed layer depth, ⁱ with updates to the physical model as described in Tjiputra et al. (2013), ^j available to year 2008 only, ^k using winds from Atlas et al. (2011), ^l updated version “oc_v1.1”

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Global carbon budget
2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 6. Comparison of results from the bookkeeping model and budget residuals with results from the DGVMs for the periods 1960–1969, 1970–1979, 1980–1989, 1990–1999, 2000–2009 and the last decade available. All values are in GtC yr^{-1} . The DGVM uncertainties represents ± 1 sigma of results from the eight individual models.

	mean (GtC yr^{-1})						
	1960–1969	1970–1979	1980–1989	1990–1999	2000–2009	2003–2012	2012
Land-use change emissions (E_{LUC})							
Bookkeeping method	1.5 ± 0.5	1.3 ± 0.5	1.2 ± 0.5	1.4 ± 0.5	0.9 ± 0.5	$0.8 \pm 0.5^*$	$0.9 \pm 0.5^*$
DGVMs	1.6 ± 0.5	1.5 ± 0.4	1.6 ± 0.4	2.1 ± 0.8	1.2 ± 0.6	1.1 ± 0.6	0.6 ± 0.8
Residual terrestrial sink (S_{LAND})							
Budget residual	1.8 ± 0.7	1.8 ± 0.8	1.4 ± 0.8	2.5 ± 0.7	2.3 ± 0.8	2.6 ± 0.8	2.5 ± 0.9
DGVMs	1.2 ± 0.8	2.1 ± 0.8	1.6 ± 0.9	2.1 ± 0.9	2.6 ± 0.9	2.7 ± 1.0	1.7 ± 1.2

* E_{LUC} for 2012 is assigned the mean of 2001–2010 as the estimate based on the bookkeeping method was not available for that year.

Global carbon budget
2013

C. Le Quéré et al.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Table 7. Decadal mean in the five components of the anthropogenic CO₂ budget for the periods 1960–1969, 1970–1979, 1980–1989, 1990–1999, 2000–2009 and the last decade available. All values are in GtC yr⁻¹. All uncertainties are reported as ±1 sigma.

	mean (GtC yr ⁻¹)						
	1960–1969	1970–1979	1980–1989	1990–1999	2000–2009	2003–2012	2012
Emissions							
Fossil-fuel combustion and cement production (E_{FF})	3.1 ± 0.2	4.7 ± 0.2	5.5 ± 0.3	6.4 ± 0.3	7.8 ± 0.4	8.6 ± 0.4	9.7 ± 0.5
Land-Use Change emissions (E_{LUC})	1.5 ± 0.5	1.3 ± 0.5	1.2 ± 0.5	1.4 ± 0.5	0.9 ± 0.5	0.8 ± 0.5*	0.9 ± 0.5*
Partitioning							
Atmospheric growth rate (G_{ATM})	1.7 ± 0.1	2.8 ± 0.1	3.4 ± 0.1	3.1 ± 0.1	4.0 ± 0.1	4.3 ± 0.1	5.2 ± 0.2
Ocean sink (S_{OCEAN})	1.1 ± 0.5	1.5 ± 0.5	1.9 ± 0.5	2.2 ± 0.4	2.4 ± 0.5	2.6 ± 0.5	2.9 ± 0.5
Residual terrestrial sink (S_{LAND})	1.8 ± 0.7	1.8 ± 0.8	1.4 ± 0.8	2.5 ± 0.7	2.3 ± 0.8	2.6 ± 0.8	2.5 ± 0.9

* E_{LUC} for 2012 is assigned the mean of 2001–2010 as the estimate based on the bookkeeping method was not available for that year.

Global carbon budget 2013

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 8. Actual CO₂ emissions from fossil-fuel combustion and cement production (E_{FF}) compared to projections made the previous year based on world GDP (IMF October 2013) and the fossil-fuel intensity of GDP (I_{FF}) based on subtracting the CO₂ and GDP growth rates. The “Actual” values are the latest estimate available and the “Projected” value for 2013 refer to those presented in this paper.

Component	2009 ^a		2010 ^b		2011 ^c		2012 ^d		2013 Projected
	Projected	Actual	Projected	Actual	Projected	Actual	Projected	Actual	
E_{FF}	−2.8%	−0.5%	> 3%	4.9%	3.1 ± 1.5%	3.2%	2.6 (1.9–3.5)%	2.2%	2.1%
GDP	−1.1%	−0.4%	4.8%	5.2%	4.0%	3.9%	3.3%	3.2%	2.9%
I_{FF}	−1.7%	−0.9%	> −1.7%	−0.3%	−0.9 ± 1.5%	−0.7%	−0.7%	−1.0%	−0.8%

^aLe Quéré et al. (2009), ^bFriedlingstein et al. (2010), ^cPeters et al. (2013), ^dLe Quéré et al. (2013)

Global carbon budget 2013

C. Le Quéré et al.

[Title Page](#)[Abstract](#)[Instruments](#)[Data Provenance & Structure](#)[Tables](#)[Figures](#)[I◀](#)[▶I](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

Table 9. Cumulative CO₂ emissions for the periods 1750–2012, 1870–2012 and 1870–2013 in GtC. All uncertainties are reported as ± 1 sigma. All values are rounded to nearest 5 GtC as in Stocker et al. (2013), reflecting the limits of our capacity to constrain cumulative estimates.

	1750–2012 (GtC)	1870–2012 (GtC)	1870–2013 (GtC)
Emissions			
Fossil-fuel combustion and cement production (E_{FF})	385 ± 20	380 ± 20	$390 \pm 20^*$
Land-Use Change emissions (E_{LUC})	205 ± 70	160 ± 55	$160 \pm 55^*$
Total emissions	590 ± 75	540 ± 60	$550 \pm 60^*$
Partitioning			
Atmospheric growth rate (G_{ATM})	245 ± 5	220 ± 5	
Ocean sink (S_{OCEAN})	165 ± 20	150 ± 20	
Residual terrestrial sink (S_{LAND})	180 ± 75	170 ± 65	

* The extension to year 2013 uses the emissions projections for 2013 of 9.9 GtC (Sect. 3.2) and assumes a constant E_{LUC} flux as in 2012 (Sect. 2.2).




Fig. 1. Schematic representation of the overall perturbation of the global carbon cycle caused by anthropogenic activities, averaged globally for the decade 2003–2012. The arrows represent emission from fossil-fuel burning and cement production (E_{FF}); emissions from deforestation and other land-use change (E_{LUC}); and the growth of carbon in the atmosphere (G_{ATM}) and the uptake of carbon by the “sinks” in the ocean (S_{OCEAN}) and land (S_{LAND}) reservoirs. All fluxes are in units of $GtC\ yr^{-1}$, with uncertainties reported as ± 1 sigma (68 % confidence that the real value lies within the given interval) as described in the text. This Figure is an update of one prepared by the International Geosphere Biosphere Programme for the GCP, first presented in Le Quéré (2009).




Fig. 2. Combined components of the global carbon budget illustrated in Fig. 1 as a function of time, for (top) emissions from fossil-fuel combustion and cement production (E_{FF} ; grey) and emissions from land-use change (E_{LUC} ; brown), and (bottom) their partitioning among the atmosphere (G_{ATM} ; light blue), land (S_{LAND} ; green) and oceans (S_{OCEAN} ; dark blue). All time-series are in GtC yr^{-1} . G_{ATM} and S_{OCEAN} (and by construction also S_{LAND}) prior to 1959 are based on different methods and shown as a dashed line. The primary data sources are for: fossil-fuel and cement emissions from Boden et al. (2013), with uncertainty of about $\pm 5\%$ ($\pm 1\sigma$); land-use change emissions from Houghton and Hackler (2013) with uncertainties of about $\pm 30\%$; atmospheric growth rate prior to 1959 is from Joos and Spahni (2008) with uncertainties of about $\pm 1-1.5 \text{ GtC decade}^{-1}$ or $\pm 0.1-0.15 \text{ GtC yr}^{-1}$ (Bruno and Joos, 1997), and from Dlugokencky and Tans (2013) from 1959 with uncertainties of about $\pm 0.2 \text{ GtC yr}^{-1}$; ocean sink prior to 1959 is from Khatiwala et al. (2013) with uncertainty of about $\pm 30\%$, and from this study from 1959 with uncertainties of about $\pm 0.5 \text{ GtC yr}^{-1}$; residual land sink is obtained by difference (Eq. 8), resulting in uncertainties of about $\pm 50\%$ prior to 1959 and $\pm 0.8 \text{ GtC yr}^{-1}$ after that. See the text for more details of each component and their uncertainties.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 3. Components of the global carbon budget and their uncertainties as a function of time, presented individually for **(a)** emissions from fossil-fuel combustion and cement production (E_{FF}), **(b)** emissions from land-use change (E_{LUC}), **(c)** atmospheric CO_2 growth rate (G_{ATM}), **(d)** the ocean CO_2 sink (S_{OCEAN} , positive indicates a flux from the atmosphere to the ocean), and **(e)** the land CO_2 sink (S_{LAND} , positive indicates a flux from the atmosphere to the land). All time-series are in $GtC\ yr^{-1}$ with the uncertainty bounds representing ± 1 sigma in shaded colour. Data sources are as in Fig. 2. The black dots in panels **(a)**, **(b)** and **(e)** show the values extrapolated from original data as explained in the text.

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 4. CO₂ emissions from fossil-fuel combustion and cement production for **(a)** the globe, including an uncertainty of $\pm 5\%$ (grey shading), the emissions extrapolated using BP energy statistics (black dots) and the emissions projection for year 2012 based on GDP projection (red dot), **(b)** global emissions by fuel type, including coal (red), oil (black), gas (blue), and cement (purple), and excluding gas flaring which is small (0.7% in 2011), **(c)** territorial (full line) and consumption (dashed line) emissions for the countries listed in the Annex B of the Kyoto Protocol (blue lines; mostly advanced economies with emissions limitations) versus non-Annex B countries (red lines), also shown are the emissions transfer from non-Annex B to Annex B countries (black line) **(d)** territorial CO₂ emissions for the top three country emitters (USA – purple; China – red; India – green) and for the European Union (EU; blue for the 28 member states of the EU in 2012), and **(e)** per-capita emissions for the top three country emitters and the EU (all colours as in panel **d**) and the world (black). In panels **(b)** to **(e)**, the dots show the years where the emissions were extrapolated using BP energy statistics. All time-series are in GtC yr⁻¹ except the per-capita emissions (panel **e**), which are in tonnes of carbon per person per year (tC person⁻¹ yr⁻¹). All territorial emissions are primarily from Boden et al. (2013) as detailed in the text; consumption-based emissions are updated from Peters et al. (2011a).




Fig. 5. Comparison between the CO₂ budget values estimated here (full black line), and other methods and models (Table 5; coloured lines) for **(a)** CO₂ emissions from land-use change showing individual DGVM model results (purple) and the multi model mean (dash black line), and fire-based results (orange), **(b)** land CO₂ sink (S_{LAND}) showing individual DGVM model results (green) and multi model mean (dash black line), and **(c)** ocean CO₂ sink (S_{OCEAN}) showing individual models before normalisation (blue lines), and the two data-based products (red line for Rödenbeck et al., 2012 and purple line for Park et al., 2010). Both data-based products were corrected for the pre-industrial ocean source of CO₂ by adding a sink of 0.45 GtC yr⁻¹ (Jacobson et al., 2007), to make them comparable to S_{OCEAN} .

Title Page

Abstract

Instruments

Data Provenance & Structure

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion






Fig. 6. Comparison of global carbon budget components released annually by GCP since 2005. CO₂ emissions from both **(a)** fossil-fuel combustion and cement production (E_{FF}), and **(b)** land-use change (E_{LUC}), and their partitioning among **(c)** the atmosphere (G_{ATM}), **(d)** the ocean (S_{OCEAN}), and **(e)** the land (S_{LAND}). See legend for the corresponding years, with the 2006 carbon budget from Raupach et al. (2007); 2007 from Canadell et al. (2007); to 2008 published online only; 2009 from Le Quéré et al. (2009); 2010 from Friedlingstein et al. (2010); 2011 from Peters et al. (2012b); 2012 from Le Quéré et al. (2013); and this year's budget (2013). The budget year corresponds to the year of the budget was first release. All values are in GtC yr⁻¹.