SUPPORTING INFORMATION

The Loading Capacity *versus* the Enzyme Activity in

New Anisotropic and Spherical Calcium Carbonate

Microparticles

Senem Donatan^{1,‡}, Alexey Yashchenok^{1,4,‡}*, Nazimuddin Khan^{2,‡}, Bogdan Parakhonskiy^{3,4,5‡,*}, Melissa Cocquyt⁵, Bat-El Pinchasik¹, Dmitry Khalenkow⁵, Helmuth Möhwald¹, Manfred Konrad², Andre Skirtach^{1,5}

¹ Department of Interfaces, Max Planck Institute of Colloids and Interfaces, Golm/Potsdam, D-14476, Germany, ² Enzyme Biochemistry Group, Max Planck Institute for Biophysical Chemistry, Göttingen, D-37077, Germany, ³A.V. Shubnikov Institute of Crystallography RAS, 119333, Moscow, Russia

⁴Remote Controlled Theranostic Systems Lab, Institute of Nanostructres and Biosystems, Saratov State

University, 410012 Saratov, Russia

⁵Department of Molecular Biotechnology, NB-Photonics Group, Ghent University, Ghent 9000, Belgium

⁶ Department of Physics at Interfaces, Max Planck Institute for Polymer

*Corresponding authors: Bogdan Parakhonskiy: bogdan.parakhonskiy@ugent.be; Department of

Molecular Biotechnology, NB-Photonics Group, Ghent University, Ghent 9000, Belgium

Alexey Yashchenok: yashenokam@gmail.com Max Planck Institute of Colloids and Interfaces, Golm,
Germany; Remote Controlled Theranostic Systems Lab, Institute of Nanostructres and Biosystems,
Saratov State University, 410012 Saratov, Russia




Figure S1. Powder X-ray diffraction analysis of rhomboidal, ellipsoidal, and spherical vaterite particles. C= calcite, V= vaterite.