

SUPPORTING INFORMATION

On the interaction between digitonin and cholesterol in Langmuir monolayers

Kamil Wojciechowski^{1*}, Marta Orczyk¹, Thomas Gutberlet², Gerald Brezesinski³, Thomas Geue⁴,
Philippe Fontaine⁵

¹*Faculty of Chemistry, Warsaw University of Technology, Noakowskiego 3, 00-664 Warsaw, Poland*

²*Helmholtz-Zentrum Berlin für Materialien und Energie GmbH, Hahn-Meitner-Platz 1, 14109 Berlin,
Germany*

³*Max Planck Institute of Colloids and Interfaces, Am Mühlenberg 1, Potsdam, Germany*

⁴*Laboratory for Neutron Scattering, Paul Scherrer Institute, WHGA/110, 5232 Villigen – PSI,
Switzerland*

⁵*Synchrotron SOLEIL, L'Orme des Merisiers, Saint Aubin, BP48, Gif-sur-Yvette Cedex, France*

*corresponding author kamil.wojciechowski@ch.pw.edu.pl




Fig. S1. GIXD patterns for DPPC, cholesterol and their mixed monolayers compressed to $\Pi_0 = 32.5$ mN/m on water.




Fig. S2. GIXD patterns for DPPC, cholesterol and their mixed monolayers compressed to $\Pi_0 = 32.5$ mN/m on digitonin (10⁻⁴ M)