

Search for structure in the $B_s^0\pi^\pm$ invariant mass spectrum

The LHCb collaboration[†]

Abstract

The $B_s^0\pi^\pm$ invariant mass distribution is investigated in order to search for possible exotic meson states. The analysis is based on a data sample recorded with the LHCb detector corresponding to 3 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV. No significant excess is found, and upper limits are set on the production rate of the claimed $X(5568)$ state within the LHCb acceptance. Upper limits are also set as a function of the mass and width of a possible exotic meson decaying to the $B_s^0\pi^\pm$ final state. The same limits also apply to a possible exotic meson decaying through the chain $B_s^{*0}\pi^\pm$, $B_s^{*0} \rightarrow B_s^0\gamma$ where the photon is excluded from the reconstructed decays.

Published in Phys. Rev. Lett. 117 (2016) 152003

© CERN on behalf of the LHCb collaboration, licence CC-BY-4.0.

[†]Authors are listed at the end of this Letter.

Interest in exotic hadrons has recently intensified, with a wealth of experimental data becoming available [1, 2]. All the well-established exotic states contain a heavy quark-antiquark ($c\bar{c}$ or $b\bar{b}$) pair together with additional light particle content. However, the D0 collaboration has reported evidence [3] of a narrow structure, referred to as the $X(5568)$, in the $B_s^0\pi^\pm$ spectrum produced in $p\bar{p}$ collisions at centre-of-mass energy $\sqrt{s} = 1.96$ TeV. The claimed $X(5568)$ state, if confirmed, would differ from any of the previous observations, as it must have constituent quarks with four different flavours (b, s, u, d). As such, it would be unique among observed exotic hadrons in having its mass dominated by a single constituent quark rather than by a quark-antiquark pair. This could provide a crucial piece of information to help understand how exotic hadrons are bound; specifically, whether they are dominantly tightly bound (often referred to as “tetraquarks” and “pentaquarks”) or loosely bound meson-meson or meson-baryon molecules.

In this Letter, results are presented from a search for an exotic meson, denoted X , decaying to $B_s^0\pi^\pm$ in a data sample corresponding to 3 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV recorded by LHCb. The search is performed by scanning over the mass and width of the purported state, with dedicated fits for parameters corresponding to those of the claimed $X(5568)$ state. The B_s^0 mesons are reconstructed in decays to $D_s^-\pi^+$ and $J/\psi\phi$ final states to obtain a B_s^0 yield approximately 20 times larger than that used by the D0 collaboration. The inclusion of charge-conjugate processes is implied throughout the Letter. The analysis techniques follow closely those developed for studies of the B^+K^- [4], $B^+\pi^-$ and $B^0\pi^+$ [5] spectra. As in previous analyses, the charged pion which is combined with the B_s^0 meson in order to form the $B_s^0\pi^\pm$ candidate is referred to as the “companion pion”.

The LHCb detector [6, 7] is a single-arm forward spectrometer covering the pseudorapidity range $2 < \eta < 5$, designed for the study of particles containing b or c quarks. The detector includes a high-precision tracking system consisting of a silicon-strip vertex detector surrounding the pp interaction region, a large-area silicon-strip detector located upstream of a dipole magnet with a bending power of about 4 Tm, and three stations of silicon-strip detectors and straw drift tubes placed downstream of the magnet. The tracking system provides a measurement of momentum, p , of charged particles with a relative uncertainty that varies from 0.5% at low momentum to 1.0% at 200 GeV (units in which $c = \hbar = 1$ are used throughout). The minimum distance of a track to a primary vertex (PV), the impact parameter, is measured with a resolution of $(15 + 29/p_T)\mu\text{m}$, where p_T is the component of the momentum transverse to the beam, in GeV. Different types of charged hadrons are distinguished using information from two ring-imaging Cherenkov detectors. Photons, electrons and hadrons are identified by a calorimeter system consisting of scintillating-pad and preshower detectors, an electromagnetic calorimeter and a hadronic calorimeter. Muons are identified by a system composed of alternating layers of iron and multiwire proportional chambers. The online event selection is performed by a trigger, which consists of a hardware stage, based on information from the calorimeter and muon systems, followed by a software stage, which applies a full event reconstruction.

Simulations of pp collisions are generated using PYTHIA [8] with a specific LHCb configuration [9]. Decays of hadronic particles are described by EVTGEN [10], in which final-state radiation is generated using PHOTOS [11]. The interaction of the generated particles with the detector, and its response, are implemented using the GEANT4 toolkit [12] as described in Ref. [13].




Figure 1: Selected candidates for (left) $B_s^0 \rightarrow D_s^- \pi^+$ and (right) $B_s^0 \rightarrow J/\psi \phi$ decays, with $p_T(B_s^0) > 5$ GeV, where the B_s^0 signal window requirements of $|m(D_s^- \pi^+) - 5367 \text{ MeV}| < 30 \text{ MeV}$ and $|m(J/\psi \phi) - 5367 \text{ MeV}| < 13 \text{ MeV}$ are indicated by dotted lines. Results of the fits described in the text are superimposed with the total fit result shown as a red line, the signal component as an unfilled area, the combinatorial background component as a dark blue area and additional background contributions as a light green area.

Candidate B_s^0 mesons are reconstructed through the decays $B_s^0 \rightarrow D_s^- \pi^+$ with $D_s^- \rightarrow K^+ K^- \pi^-$, and $B_s^0 \rightarrow J/\psi \phi$ with $J/\psi \rightarrow \mu^+ \mu^-$ and $\phi \rightarrow K^+ K^-$. Particle identification, track quality and impact parameter requirements are imposed on all final-state particles. Both B_s^0 and intermediate particle (D_s^- and J/ψ) candidates are required to have good vertex quality and to have invariant mass close to the known values [14]. Specific backgrounds due to other b -hadron decays are removed with appropriate vetoes. A requirement is imposed on the multiplicity of tracks originating from the PV associated with the B_s^0 candidate; this requirement is about 90% efficient on B_s^0 signal and significantly reduces background due to random $B_s^0 \pi^\pm$ combinations. To further reduce background, the p_T of the B_s^0 candidate, $p_T(B_s^0)$, is required to be greater than 5 GeV. Results are also obtained with this requirement increased to 10 or 15 GeV, to be more sensitive to scenarios in which the X state is predominantly produced from hard processes. The definition of the fiducial acceptance is completed with the requirements $p_T(B_s^0) < 50$ GeV and $2.0 < y < 4.5$, where y is the rapidity of the B_s^0 candidate.

The signals in the two B_s^0 decay modes are shown in Fig. 1. To estimate the B_s^0 yields, the data are fitted with functions that include a signal component, described by a double Gaussian function with a shared mean, and a combinatorial background component, described by a polynomial function. Backgrounds from $B_s^0 \rightarrow D_s^\mp K^\pm$ decays in the $D_s^- \pi^+$ sample and from $\Lambda_b^0 \rightarrow J/\psi p K^-$ decays in the $J/\psi \phi$ sample, where a final-state hadron is misidentified, are modelled using empirical shapes derived from simulated samples. An additional component, modelled with a Gaussian function, is included to account for possible $B^0 \rightarrow J/\psi K^+ K^-$ decays [15] in the $J/\psi \phi$ sample. The results of these fits are reported in Table 1. The signal-to-background ratio in the B_s^0 signal windows is about 10 for the $D_s^- \pi^+$ sample and above 50 for the $J/\psi \phi$ sample.

The B_s^0 candidates are combined with each track originating from the associated PV that gives a good quality $B_s^0 \pi^\pm$ vertex and that has $p_T > 500$ MeV. A loose pion identification requirement is imposed in order to suppress possible backgrounds involving misidentified particles. In case multiple candidates are obtained in the same event, all are retained. Mass and vertex constraints are imposed [16] in the calculation of the $B_s^0 \pi^\pm$ invariant mass.

In order to obtain quantitative results on the contributions from resonant structures in

the data, the $B_s^0\pi^\pm$ mass distributions are fitted with a function containing components for signal and background. The signal shape is an S-wave Breit–Wigner function multiplied by a function that accounts for the variation of the efficiency with $B_s^0\pi^\pm$ mass. The efficiency function, determined from simulation, plateaus at high $B_s^0\pi^\pm$ mass and falls near the threshold to a value that depends on $p_T(B_s^0)$. The resolution is better than 1 MeV and does not affect the results. The background is modelled with a polynomial function. It is verified that this function gives a good description of backgrounds composed of either a real or a fake B_s^0 decay combined with a random pion, as determined from simulation or from data in B_s^0 candidate mass sideband regions, respectively.

For each choice of signal mass and width parameters, a binned maximum likelihood fit to the $B_s^0\pi^\pm$ candidate mass spectrum is used to determine the signal and background yields and the parameters of the polynomial shape that describes the background. The two B_s^0 decay modes are fitted simultaneously. The results of the fit where the mass and width are fixed according to the central values obtained by the D0 collaboration, $m = 5567.8 \pm 2.9$ (stat) $_{-1.9}^{+0.9}$ (syst) MeV and $\Gamma = 21.9 \pm 6.4$ (stat) $_{-2.5}^{+5.0}$ (syst) MeV [3], are shown in Fig. 2 for both B_s^0 decay modes combined. The $X(5568)$ yield is not significant for any minimum $p_T(B_s^0)$ requirement. In each case the change in negative log-likelihood between fits including or not including the signal component is less than 2 units for two additional free parameters corresponding to the yields in the two B_s^0 decay modes. The results of the fits are summarised in Table 1.

The yields N obtained from the fits are used to measure the ratio of cross-sections

$$\rho_X^{\text{LHCb}} \equiv \frac{\sigma(pp \rightarrow X + \text{anything}) \times \mathcal{B}(X \rightarrow B_s^0\pi^\pm)}{\sigma(pp \rightarrow B_s^0 + \text{anything})}, \quad (1)$$

$$= \frac{N(X)}{N(B_s^0)} \times \frac{1}{\epsilon^{\text{rel}}(X)}, \quad (2)$$

where the cross-sections σ are for promptly produced particles within the LHCb acceptance. Since $\sigma(pp \rightarrow B_s^0 + \text{anything})$ in the LHCb acceptance has been previously measured [17], any result for ρ_X^{LHCb} can be scaled to give a result for $\sigma(pp \rightarrow X + \text{anything}) \times \mathcal{B}(X \rightarrow B_s^0\pi^\pm)$ in the LHCb acceptance. The relative efficiency $\epsilon^{\text{rel}}(X) = \frac{\epsilon(X)}{\epsilon(B_s^0)}$ accounts for the reconstruction and selection efficiency of the companion pion as well as the requirement that it is within the LHCb acceptance. These effects are determined from simulation, weighted to reproduce the measured differential B_s^0 production spectrum [17], together with a data-driven evaluation [18] of the efficiency of the particle identification requirement on the companion pion. In the simulation the X state is assumed to be spinless; it has been verified that the systematic uncertainty associated with this choice is negligible. The quantities used to evaluate ρ_X^{LHCb} are summarised in Table 1.

Systematic uncertainties are assigned due to possible biases in the evaluation of $N(X)$, $N(B_s^0)$ and $\epsilon^{\text{rel}}(X)$. The signal shape is modified by varying the efficiency function, and separately by changing the assumed angular momentum in the relativistic Breit–Wigner function from S-wave to P-wave. In each case, the changes in $N(X)$ are assigned as the associated systematic uncertainties. Uncertainties associated with the determination of $N(B_s^0)$ arise due to the size of the B_s^0 sample and the estimation of the background in the signal region. In addition to the limited size of the simulation sample, uncertainties associated with $\epsilon^{\text{rel}}(X)$ arise due to the precision with which the companion pion reconstruction and particle identification efficiencies are known [18, 19]. The uncertainties from




Figure 2: Results of the fit to the $B_s^0 \pi^\pm$ mass distribution for candidates (both B_s^0 modes combined) with minimum $p_T(B_s^0)$ of (top) 5 GeV, (middle) 10 GeV and (bottom) 15 GeV. The component for the claimed $X(5568)$ state is included in the fit but is not significant. The distributions of the normalised residuals, or “pulls”, displayed underneath the main figures show good agreement between the fit functions and the data.

Table 1: Yields, N , of B_s^0 and $X(5568)$ candidates obtained from the fits to the B_s^0 and $B_s^0\pi^\pm$ candidate mass distributions, with statistical uncertainties only. The values reported for $N(B_s^0)$ are those inside the B_s^0 signal window. The reported values for $X(5568)$ are obtained from fits with signal mass and width parameters fixed to those determined by the D0 collaboration. Relative efficiencies $\epsilon^{\text{rel}}(X)$ of the B_s^0 and $X(5568)$ candidate selection criteria are also given. The reported uncertainties on the relative efficiencies are only statistical, due to the finite size of the simulated samples.

		$B_s^0 \rightarrow D_s^- \pi^+$	$B_s^0 \rightarrow J/\psi \phi$	Sum
$N(B_s^0)/10^3$	$p_T(B_s^0) > 5 \text{ GeV}$	62.2 ± 0.3	43.6 ± 0.2	105.8 ± 0.4
	$p_T(B_s^0) > 10 \text{ GeV}$	28.4 ± 0.2	13.2 ± 0.1	41.6 ± 0.2
	$p_T(B_s^0) > 15 \text{ GeV}$	8.8 ± 0.1	3.7 ± 0.1	12.5 ± 0.1
$N(X)$	$p_T(B_s^0) > 5 \text{ GeV}$	3 ± 64	-33 ± 43	-30 ± 77
	$p_T(B_s^0) > 10 \text{ GeV}$	75 ± 52	12 ± 33	87 ± 62
	$p_T(B_s^0) > 15 \text{ GeV}$	14 ± 31	-10 ± 17	4 ± 35
$\epsilon^{\text{rel}}(X)$	$p_T(B_s^0) > 5 \text{ GeV}$	0.127 ± 0.002	0.093 ± 0.001	—
	$p_T(B_s^0) > 10 \text{ GeV}$	0.213 ± 0.003	0.206 ± 0.002	—
	$p_T(B_s^0) > 15 \text{ GeV}$	0.289 ± 0.005	0.290 ± 0.004	—

different sources are combined in quadrature and give a total that is much smaller than the statistical uncertainty. To obtain results that can be compared to those for the claimed $X(5568)$ state reported by the D0 collaboration, additional systematic uncertainties are assigned from the changes in the results for ρ_X^{LHCb} when the mass and width parameters are varied independently within $\pm 1\sigma$ ranges from their central values. These are the dominant sources of systematic uncertainty.

To cross-check the results, candidates are selected with criteria similar to those used in the observation of $B_c^+ \rightarrow B_s^0 \pi^+$ decays [20], with consistent results. In addition, $B^0 \rightarrow D^- \pi^+$ decays are used to create $B^0 \pi^+$ combinations, and the results on the excited B states of Ref. [5] are reproduced.

The values of ρ_X^{LHCb} for the two B_s^0 decay modes are consistent and are therefore combined in a weighted average. In the average, systematic uncertainties are taken to be uncorrelated between the two B_s^0 decay modes. An exception is made when obtaining results corresponding to the claimed $X(5568)$ state, where the uncertainty due to the limited precision of the reported mass and width values [3] is treated as correlated between the two modes. These results are

$$\begin{aligned}
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 5 \text{ GeV}) &= -0.003 \pm 0.006 \pm 0.002, \\
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 10 \text{ GeV}) &= 0.010 \pm 0.007 \pm 0.005, \\
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 15 \text{ GeV}) &= 0.000 \pm 0.010 \pm 0.006,
\end{aligned}$$

where the first uncertainty is statistical and the second is systematic. Since the signal is not significant, upper limits on ρ_X^{LHCb} are obtained by integration of the likelihood in the positive region to find the value that contains the fraction of the integral corresponding to the required confidence level (CL). The upper limits at 90 (95) % CL are found to be

$$\begin{aligned}
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 5 \text{ GeV}) &< 0.011 (0.012), \\
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 10 \text{ GeV}) &< 0.021 (0.024), \\
\rho_X^{\text{LHCb}}(p_T(B_s^0) > 15 \text{ GeV}) &< 0.018 (0.020).
\end{aligned}$$

No significant signal for a $B_s^0\pi^\pm$ resonance is seen at any value of mass and width in the range considered. To obtain limits on ρ_X^{LHCb} for different values of these parameters, fits are performed for widths (Γ) of 10 to 50 MeV in 10 MeV steps. For each width, the mass is scanned in steps of $\Gamma/2$, starting one unit of width above the kinematic threshold and ending approximately one and a half units of width below 6000 MeV. The upper edge of the range is chosen because an exotic state with higher mass would be expected to give a clearer signature in the B^0K^\pm final state [21]. The results are obtained in the same way as described above, and converted into upper limits that are shown in Fig. 3. The upper limits are weaker when a broader width is assumed, due to the larger amount of background under the putative peak. The limits also become weaker when there is an excess of events in the signal region, although all such excesses are consistent with being statistical fluctuations. The method used to set the upper limits smooths out any negative fluctuations.

In summary, a search for the claimed $X(5568)$ state has been carried out using a data sample corresponding to 3 fb^{-1} of pp collision data at $\sqrt{s} = 7$ and 8 TeV recorded by LHCb. No significant excess is found and thus the existence of the $X(5568)$ state is not confirmed. Upper limits are set on the relative production rate of the claimed state in the LHCb acceptance. Limits are also set as a function of the mass and width of a possible exotic meson decaying to the $B_s^0\pi^\pm$ final state. The same limits also apply to a possible exotic meson decaying through the chain $B_s^{*0}\pi^\pm$, $B_s^{*0} \rightarrow B_s^0\gamma$ where the photon is excluded from the reconstructed decays.

Acknowledgements

We express our gratitude to our colleagues in the CERN accelerator departments for the excellent performance of the LHC. We thank the technical and administrative staff at the LHCb institutes. We acknowledge support from CERN and from the national agencies: CAPES, CNPq, FAPERJ and FINEP (Brazil); NSFC (China); CNRS/IN2P3 (France); BMBF, DFG and MPG (Germany); INFN (Italy); FOM and NWO (The Netherlands); MNiSW and NCN (Poland); MEN/IFA (Romania); MinES and FASO (Russia); MinECo (Spain); SNSF and SER (Switzerland); NASU (Ukraine); STFC (United Kingdom); NSF (USA). We acknowledge the computing resources that are provided by CERN, IN2P3 (France), KIT and DESY (Germany), INFN (Italy), SURF (The Netherlands), PIC (Spain), GridPP (United Kingdom), RRCKI and Yandex LLC (Russia), CSCS (Switzerland), IFIN-HH (Romania), CBPF (Brazil), PL-GRID (Poland) and OSC (USA). We are indebted to the communities behind the multiple open source software packages on which we depend. Individual groups or members have received support from AvH Foundation (Germany), EPLANET, Marie Skłodowska-Curie Actions and ERC (European Union), Conseil Général de Haute-Savoie, Labex ENIGMASS and OCEVU, Région Auvergne (France), RFBR and Yandex LLC (Russia), GVA, XuntaGal and GENCAT (Spain), Herchel Smith Fund, The Royal Society, Royal Commission for the Exhibition of 1851 and the Leverhulme Trust (United Kingdom).




Figure 3: Upper limits (ULs) at 90 % confidence level (CL) as functions of the mass and width of a purported exotic state X decaying to $B_s^0\pi^\pm$ with minimum $p_T(B_s^0)$ of (top) 5 GeV, (middle) 10 GeV and (bottom) 15 GeV. The same limits also apply to a possible exotic meson decaying through the chain $B_s^{*0}\pi^\pm$, $B_s^{*0} \rightarrow B_s^0\gamma$ where the photon is excluded from the reconstructed decays. In the latter case the nominal mass difference $m(B_s^{*0}) - m(B_s^0) = 48.6^{+1.8}_{-1.6}$ MeV [14] has to be added to the values on the x-axis to get the mass of the exotic meson under investigation.

References

- [1] S. L. Olsen, *A new hadron spectroscopy*, Front. Phys. **10** (2015) 101401, [arXiv:1411.7738](#).
- [2] H.-Y. Cheng, *Multiquark hadrons*, The Universe **3** (2015), no. 3 33, [INSPIRE:Record 1452624].
- [3] D0 collaboration, V. M. Abazov *et al.*, *Evidence for a $B_s^0\pi^\pm$ state*, Phys. Rev. Lett. **117** (2016) 022003, [arXiv:1602.07588](#).
- [4] LHCb collaboration, R. Aaij *et al.*, *First observation of the decay $B_{s2}^*(5840)^0 \rightarrow B^{*+}K^-$ and studies of excited B_s^0 mesons*, Phys. Rev. Lett. **110** (2013) 151803, [arXiv:1211.5994](#).
- [5] LHCb collaboration, R. Aaij *et al.*, *Precise measurements of the properties of the $B_1(5721)^{0,+}$ and $B_2^*(5747)^{0,+}$ states and observation of structure at higher invariant mass in the $B^+\pi^-$ and $B^0\pi^+$ spectra*, JHEP **04** (2015) 024, [arXiv:1502.02638](#).
- [6] LHCb collaboration, A. A. Alves Jr. *et al.*, *The LHCb detector at the LHC*, JINST **3** (2008) S08005.
- [7] LHCb collaboration, R. Aaij *et al.*, *LHCb detector performance*, Int. J. Mod. Phys. **A30** (2015) 1530022, [arXiv:1412.6352](#).
- [8] T. Sjöstrand, S. Mrenna, and P. Skands, *PYTHIA 6.4 physics and manual*, JHEP **05** (2006) 026, [arXiv:hep-ph/0603175](#); T. Sjöstrand, S. Mrenna, and P. Skands, *A brief introduction to PYTHIA 8.1*, Comput. Phys. Commun. **178** (2008) 852, [arXiv:0710.3820](#).
- [9] I. Belyaev *et al.*, *Handling of the generation of primary events in Gauss, the LHCb simulation framework*, J. Phys. Conf. Ser. **331** (2011) 032047.
- [10] D. J. Lange, *The EvtGen particle decay simulation package*, Nucl. Instrum. Meth. **A462** (2001) 152.
- [11] P. Golonka and Z. Was, *PHOTOS Monte Carlo: A precision tool for QED corrections in Z and W decays*, Eur. Phys. J. **C45** (2006) 97, [arXiv:hep-ph/0506026](#).
- [12] Geant4 collaboration, J. Allison *et al.*, *Geant4 developments and applications*, IEEE Trans. Nucl. Sci. **53** (2006) 270; Geant4 collaboration, S. Agostinelli *et al.*, *Geant4: A simulation toolkit*, Nucl. Instrum. Meth. **A506** (2003) 250.
- [13] M. Clemencic *et al.*, *The LHCb simulation application, Gauss: Design, evolution and experience*, J. Phys. Conf. Ser. **331** (2011) 032023.
- [14] Particle Data Group, K. A. Olive *et al.*, *Review of particle physics*, Chin. Phys. **C38** (2014) 090001, and 2015 update.
- [15] LHCb collaboration, R. Aaij *et al.*, *Amplitude analysis and branching fraction measurement of $\bar{B}_s^0 \rightarrow J/\psi K^+ K^-$* , Phys. Rev. **D87** (2013) 072004, [arXiv:1302.1213](#).

- [16] W. D. Hulsbergen, *Decay chain fitting with a Kalman filter*, Nucl. Instrum. Meth. **A552** (2005) 566, [arXiv:physics/0503191](#).
- [17] LHCb collaboration, R. Aaij *et al.*, *Measurement of B meson production cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV*, JHEP **08** (2013) 117, [arXiv:1306.3663](#).
- [18] M. Adinolfi *et al.*, *Performance of the LHCb RICH detector at the LHC*, Eur. Phys. J. **C73** (2013) 2431, [arXiv:1211.6759](#).
- [19] LHCb collaboration, R. Aaij *et al.*, *Measurement of the track reconstruction efficiency at LHCb*, JINST **10** (2015) P02007, [arXiv:1408.1251](#).
- [20] LHCb collaboration, R. Aaij *et al.*, *Observation of the decay $B_c^+ \rightarrow B_s^0 \pi^+$* , Phys. Rev. Lett. **111** (2013) 181801, [arXiv:1308.4544](#).
- [21] A. Esposito, A. Pilloni, and A. D. Polosa, *Hybridized tetraquarks*, Phys. Lett. **B758** (2016) 292, [arXiv:1603.07667](#).

LHCb collaboration

R. Aaij⁴⁰, B. Adeva³⁹, M. Adinolfi⁴⁸, Z. Ajaltouni⁵, S. Akar⁶, J. Albrecht¹⁰, F. Alessio⁴⁰, M. Alexander⁵³, S. Ali⁴³, G. Alkhazov³¹, P. Alvarez Cartelle⁵⁵, A.A. Alves Jr⁵⁹, S. Amato², S. Amerio²³, Y. Amhis⁷, L. An⁴¹, L. Anderlini¹⁸, G. Andreassi⁴¹, M. Andreotti^{17,g}, J.E. Andrews⁶⁰, R.B. Appleby⁵⁶, O. Aquines Gutierrez¹¹, F. Archilli¹, P. d'Argent¹², J. Arnau Romeu⁶, A. Artamonov³⁷, M. Artuso⁶¹, E. Aslanides⁶, G. Auriemma²⁶, M. Baalouch⁵, I. Babuschkin⁵⁶, S. Bachmann¹², J.J. Back⁵⁰, A. Badalov³⁸, C. Baesso⁶², S. Baker⁵⁵, W. Baldini¹⁷, R.J. Barlow⁵⁶, C. Barschel⁴⁰, S. Barsuk⁷, W. Barter⁴⁰, V. Batozskaya²⁹, B. Batsukh⁶¹, V. Battista⁴¹, A. Bay⁴¹, L. Beaucourt⁴, J. Beddow⁵³, F. Bedeschi²⁴, I. Bediaga¹, L.J. Bel⁴³, V. Bellee⁴¹, N. Belloli^{21,i}, K. Belous³⁷, I. Belyaev³², E. Ben-Haim⁸, G. Bencivenni¹⁹, S. Benson⁴⁰, J. Benton⁴⁸, A. Berezhnoy³³, R. Bernet⁴², A. Bertolin²³, F. Betti¹⁵, M.-O. Bettler⁴⁰, M. van Beuzekom⁴³, I. Bezshyiko⁴², S. Bifani⁴⁷, P. Billoir⁸, T. Bird⁵⁶, A. Birnkraut¹⁰, A. Bitadze⁵⁶, A. Bizzeti^{18,u}, T. Blake⁵⁰, F. Blanc⁴¹, J. Blouw¹¹, S. Blusk⁶¹, V. Bocci²⁶, T. Boettcher⁵⁸, A. Bondar³⁶, N. Bondar^{31,40}, W. Bonivento¹⁶, A. Borgheresi^{21,i}, S. Borghi⁵⁶, M. Borisyak³⁵, M. Borsato³⁹, F. Bossu⁷, M. Boubdir⁹, T.J.V. Bowcock⁵⁴, E. Bowen⁴², C. Bozzi^{17,40}, S. Braun¹², M. Britsch¹², T. Britton⁶¹, J. Brodzicka⁵⁶, E. Buchanan⁴⁸, C. Burr⁵⁶, A. Bursche², J. Buytaert⁴⁰, S. Cadeddu¹⁶, R. Calabrese^{17,g}, M. Calvi^{21,i}, M. Calvo Gomez^{38,m}, A. Camboni³⁸, P. Campana¹⁹, D. Campora Perez⁴⁰, D.H. Campora Perez⁴⁰, L. Capriotti⁵⁶, A. Carbone^{15,e}, G. Carboni^{25,j}, R. Cardinale^{20,h}, A. Cardini¹⁶, P. Carniti^{21,i}, L. Carson⁵², K. Carvalho Akiba², G. Casse⁵⁴, L. Cassina^{21,i}, L. Castillo Garcia⁴¹, M. Cattaneo⁴⁰, Ch. Cauet¹⁰, G. Cavallero²⁰, R. Cenci^{24,t}, M. Charles⁸, Ph. Charpentier⁴⁰, G. Chatzikonstantinidis⁴⁷, M. Chefdeville⁴, S. Chen⁵⁶, S.-F. Cheung⁵⁷, V. Chobanova³⁹, M. Chrzaszcz^{42,27}, X. Cid Vidal³⁹, G. Ciezarek⁴³, P.E.L. Clarke⁵², M. Clemencic⁴⁰, H.V. Cliff⁴⁹, J. Closier⁴⁰, V. Coco⁵⁹, J. Cogan⁶, E. Cogneras⁵, V. Cogoni^{16,40,f}, L. Cojocariu³⁰, G. Collazuol^{23,o}, P. Collins⁴⁰, A. Comerma-Montells¹², A. Contu⁴⁰, A. Cook⁴⁸, S. Coquereau⁸, G. Corti⁴⁰, M. Corvo^{17,g}, C.M. Costa Sobral⁵⁰, B. Couturier⁴⁰, G.A. Cowan⁵², D.C. Craik⁵², A. Crocombe⁵⁰, M. Cruz Torres⁶², S. Cunliffe⁵⁵, R. Currie⁵⁵, C. D'Ambrosio⁴⁰, F. Da Cunha Marinho², E. Dall'Occo⁴³, J. Dalseno⁴⁸, P.N.Y. David⁴³, A. Davis⁵⁹, O. De Aguiar Francisco², K. De Bruyn⁶, S. De Capua⁵⁶, M. De Cian¹², J.M. De Miranda¹, L. De Paula², M. De Serio^{14,d}, P. De Simone¹⁹, C.-T. Dean⁵³, D. Decamp⁴, M. Deckenhoff¹⁰, L. Del Buono⁸, M. Demmer¹⁰, D. Derkach³⁵, O. Deschamps⁵, F. Dettori⁴⁰, B. Dey²², A. Di Canto⁴⁰, H. Dijkstra⁴⁰, F. Dordei⁴⁰, M. Dorigo⁴¹, A. Dosil Suárez³⁹, A. Dovbnya⁴⁵, K. Dreimanis⁵⁴, L. Dufour⁴³, G. Dujany⁵⁶, K. Dungs⁴⁰, P. Durante⁴⁰, R. Dzhelyadin³⁷, A. Dziurda⁴⁰, A. Dzyuba³¹, N. Déleage⁴, S. Easo⁵¹, M. Ebert⁵², U. Egede⁵⁵, V. Egorychev³², S. Eidelman³⁶, S. Eisenhardt⁵², U. Eitschberger¹⁰, R. Ekelhof¹⁰, L. Eklund⁵³, Ch. Elsasser⁴², S. Ely⁶¹, S. Esen¹², H.M. Evans⁴⁹, T. Evans⁵⁷, A. Falabella¹⁵, N. Farley⁴⁷, S. Farry⁵⁴, R. Fay⁵⁴, D. Fazzini^{21,i}, D. Ferguson⁵², V. Fernandez Albor³⁹, A. Fernandez Prieto³⁹, F. Ferrari^{15,40}, F. Ferreira Rodrigues¹, M. Ferro-Luzzi⁴⁰, S. Filippov³⁴, R.A. Fini¹⁴, M. Fiore^{17,g}, M. Fiorini^{17,g}, M. Firlej²⁸, C. Fitzpatrick⁴¹, T. Fiutowski²⁸, F. Fleuret^{7,b}, K. Fohl⁴⁰, M. Fontana^{16,40}, F. Fontanelli^{20,h}, D.C. Forshaw⁶¹, R. Forty⁴⁰, V. Franco Lima⁵⁴, M. Frank⁴⁰, C. Frei⁴⁰, J. Fu^{22,q}, E. Furfaro^{25,j}, C. Färber⁴⁰, A. Gallas Torreira³⁹, D. Galli^{15,e}, S. Gallorini²³, S. Gambetta⁵², M. Gandelman², P. Gandini⁵⁷, Y. Gao³, L.M. Garcia Martin⁶⁸, J. García Pardiñas³⁹, J. Garra Tico⁴⁹, L. Garrido³⁸, P.J. Garsed⁴⁹, D. Gascon³⁸, C. Gaspar⁴⁰, L. Gavardi¹⁰, G. Gazzoni⁵, D. Gerick¹², E. Gersabeck¹², M. Gersabeck⁵⁶, T. Gershon⁵⁰, Ph. Ghez⁴, S. Gianì⁴¹, V. Gibson⁴⁹, O.G. Girard⁴¹, L. Giubega³⁰, K. Gizdov⁵², V.V. Gligorov⁸, D. Golubkov³², A. Golutvin^{55,40}, A. Gomes^{1,a}, I.V. Gorelov³³, C. Gotti^{21,i}, M. Grabalosa Gándara⁵, R. Graciani Diaz³⁸, L.A. Granado Cardoso⁴⁰, E. Graugés³⁸, E. Graverini⁴², G. Graziani¹⁸, A. Grecu³⁰, P. Griffith⁴⁷, L. Grillo^{21,40}, B.R. Gruberg Cazon⁵⁷, O. Grünberg⁶⁶, E. Gushchin³⁴, Yu. Guz³⁷, T. Gys⁴⁰, C. Göbel⁶², T. Hadavizadeh⁵⁷, C. Hadjivasiliou⁵, G. Haefeli⁴¹, C. Haen⁴⁰,

S.C. Haines⁴⁹, S. Hall⁵⁵, B. Hamilton⁶⁰, X. Han¹², S. Hansmann-Menzemer¹², N. Harnew⁵⁷,
 S.T. Harnew⁴⁸, J. Harrison⁵⁶, M. Hatch⁴⁰, J. He⁶³, T. Head⁴¹, A. Heister⁹, K. Hennessy⁵⁴,
 P. Henrard⁵, L. Henry⁸, J.A. Hernando Morata³⁹, E. van Herwijnen⁴⁰, M. Heß⁶⁶, A. Hicheur²,
 D. Hill⁵⁷, C. Hombach⁵⁶, W. Hulsbergen⁴³, T. Humair⁵⁵, M. Hushchyn³⁵, N. Hussain⁵⁷,
 D. Hutchcroft⁵⁴, M. Idzik²⁸, P. Ilten⁵⁸, R. Jacobsson⁴⁰, A. Jaeger¹², J. Jalocha⁵⁷, E. Jans⁴³,
 A. Jawahery⁶⁰, F. Jiang³, M. John⁵⁷, D. Johnson⁴⁰, C.R. Jones⁴⁹, C. Joram⁴⁰, B. Jost⁴⁰,
 N. Jurik⁶¹, S. Kandybei⁴⁵, W. Kanso⁶, M. Karacson⁴⁰, J.M. Kariuki⁴⁸, S. Karodia⁵³,
 M. Kecke¹², M. Kelsey⁶¹, I.R. Kenyon⁴⁷, M. Kenzie⁴⁹, T. Ketel⁴⁴, E. Khairullin³⁵,
 B. Khanji^{21,40,i}, C. Khurewathanakul⁴¹, T. Kirn⁹, S. Klaver⁵⁶, K. Klimaszewski²⁹, S. Koliiev⁴⁶,
 M. Kolpin¹², I. Komarov⁴¹, R.F. Koopman⁴⁴, P. Koppenburg⁴³, A. Kozachuk³³, M. Kozeiha⁵,
 L. Kravchuk³⁴, K. Kreplin¹², M. Krepes⁵⁰, P. Krokovny³⁶, F. Kruse¹⁰, W. Krzemien²⁹,
 W. Kucewicz^{27,l}, M. Kucharczyk²⁷, V. Kudryavtsev³⁶, A.K. Kuonen⁴¹, K. Kurek²⁹,
 T. Kvaratskheliya^{32,40}, D. Lacarrere⁴⁰, G. Lafferty⁵⁶, A. Lai¹⁶, D. Lambert⁵², G. Lanfranchi¹⁹,
 C. Langenbruch⁹, T. Latham⁵⁰, C. Lazzeroni⁴⁷, R. Le Gac⁶, J. van Leerdam⁴³, J.-P. Lees⁴,
 A. Leflat^{33,40}, J. Lefrançois⁷, R. Lefèvre⁵, F. Lemaitre⁴⁰, E. Lemos Cid³⁹, O. Leroy⁶, T. Lesiak²⁷,
 B. Leverington¹², Y. Li⁷, T. Likhomanenko^{35,67}, R. Lindner⁴⁰, C. Linn⁴⁰, F. Lionetto⁴²,
 B. Liu¹⁶, X. Liu³, D. Loh⁵⁰, I. Longstaff⁵³, J.H. Lopes², D. Lucchesi^{23,o}, M. Lucio Martinez³⁹,
 H. Luo⁵², A. Lupato²³, E. Luppi^{17,g}, O. Lupton⁵⁷, A. Lusiani²⁴, X. Lyu⁶³, F. Machefert⁷,
 F. Maciuc³⁰, O. Maev³¹, K. Maguire⁵⁶, S. Malde⁵⁷, A. Malinin⁶⁷, T. Maltsev³⁶, G. Manca⁷,
 G. Mancinelli⁶, P. Manning⁶¹, J. Maratas^{5,v}, J.F. Marchand⁴, U. Marconi¹⁵, C. Marin Benito³⁸,
 P. Marino^{24,t}, J. Marks¹², G. Martellotti²⁶, M. Martin⁶, M. Martinelli⁴¹, D. Martinez Santos³⁹,
 F. Martinez Vidal⁶⁸, D. Martins Tostes², L.M. Massacrier⁷, A. Massafferri¹, R. Matev⁴⁰,
 A. Mathad⁵⁰, Z. Mathe⁴⁰, C. Matteuzzi²¹, A. Mauri⁴², B. Maurin⁴¹, A. Mazurov⁴⁷,
 M. McCann⁵⁵, J. McCarthy⁴⁷, A. McNab⁵⁶, R. McNulty¹³, B. Meadows⁵⁹, F. Meier¹⁰,
 M. Meissner¹², D. Melnychuk²⁹, M. Merk⁴³, A. Merli^{22,q}, E. Michielin²³, D.A. Milanes⁶⁵,
 M.-N. Minard⁴, D.S. Mitzel¹², A. Mogini⁸, J. Molina Rodriguez⁶², I.A. Monroy⁶⁵, S. Monteil⁵,
 M. Morandin²³, P. Morawski²⁸, A. Mordà⁶, M.J. Morello^{24,t}, J. Moron²⁸, A.B. Morris⁵²,
 R. Mountain⁶¹, F. Muheim⁵², M. Mulder⁴³, M. Mussini¹⁵, D. Müller⁵⁶, J. Müller¹⁰, K. Müller⁴²,
 V. Müller¹⁰, P. Naik⁴⁸, T. Nakada⁴¹, R. Nandakumar⁵¹, A. Nandi⁵⁷, I. Nasteva², M. Needham⁵²,
 N. Neri²², S. Neubert¹², N. Neufeld⁴⁰, M. Neuner¹², A.D. Nguyen⁴¹, C. Nguyen-Mau^{41,n},
 S. Nieswand⁹, R. Niet¹⁰, N. Nikitin³³, T. Nikodem¹², A. Novoselov³⁷, D.P. O'Hanlon⁵⁰,
 A. Oblakowska-Mucha²⁸, V. Obraztsov³⁷, S. Ogilvy¹⁹, R. Oldeman⁴⁹, C.J.G. Onderwater⁶⁹,
 J.M. Otalora Goicochea², A. Otto⁴⁰, P. Owen⁴², A. Oyanguren⁶⁸, P.R. Pais⁴¹, A. Palano^{14,d},
 F. Palombo^{22,q}, M. Palutan¹⁹, J. Panman⁴⁰, A. Papanestis⁵¹, M. Pappagallo^{14,d},
 L.L. Pappalardo^{17,g}, W. Parker⁶⁰, C. Parkes⁵⁶, G. Passaleva¹⁸, A. Pastore^{14,d}, G.D. Patel⁵⁴,
 M. Patel⁵⁵, C. Patrignani^{15,e}, A. Pearce^{56,51}, A. Pellegrino⁴³, G. Penso²⁶, M. Pepe Altarelli⁴⁰,
 S. Perazzini⁴⁰, P. Perret⁵, L. Pescatore⁴⁷, K. Petridis⁴⁸, A. Petrolini^{20,h}, A. Petrov⁶⁷,
 M. Petruzzo^{22,q}, E. Picatoste Olloqui³⁸, B. Pietrzyk⁴, M. Pikies²⁷, D. Pinci²⁶, A. Pistone²⁰,
 A. Piucci¹², S. Playfer⁵², M. Plo Casasus³⁹, T. Poikela⁴⁰, F. Polci⁸, A. Poluektov^{50,36},
 I. Polyakov⁶¹, E. Polycarpo², G.J. Pomery⁴⁸, A. Popov³⁷, D. Popov^{11,40}, B. Popovici³⁰,
 S. Poslavskii³⁷, C. Potterat², E. Price⁴⁸, J.D. Price⁵⁴, J. Prisciandaro³⁹, A. Pritchard⁵⁴,
 C. Prouve⁴⁸, V. Pugatch⁴⁶, A. Puig Navarro⁴¹, G. Punzi^{24,p}, W. Qian⁵⁷, R. Quagliani^{7,48},
 B. Rachwal²⁷, J.H. Rademacker⁴⁸, M. Rama²⁴, M. Ramos Pernas³⁹, M.S. Rangel², I. Raniuk⁴⁵,
 G. Raven⁴⁴, F. Redi⁵⁵, S. Reichert¹⁰, A.C. dos Reis¹, C. Remon Alepuz⁶⁸, V. Renaudin⁷,
 S. Ricciardi⁵¹, S. Richards⁴⁸, M. Rihl⁴⁰, K. Rinnert⁵⁴, V. Rives Molina³⁸, P. Robbe^{7,40},
 A.B. Rodrigues¹, E. Rodrigues⁵⁹, J.A. Rodriguez Lopez⁶⁵, P. Rodriguez Perez⁵⁶,
 A. Rogozhnikov³⁵, S. Roiser⁴⁰, V. Romanovskiy³⁷, A. Romero Vidal³⁹, J.W. Ronayne¹³,
 M. Rotondo¹⁹, M.S. Rudolph⁶¹, T. Ruf⁴⁰, P. Ruiz Valls⁶⁸, J.J. Saborido Silva³⁹, E. Sadykhov³²,
 N. Sagidova³¹, B. Saitta^{16,f}, V. Salustino Guimaraes², C. Sanchez Mayordomo⁶⁸,
 B. Sanmartin Sedes³⁹, R. Santacesaria²⁶, C. Santamarina Rios³⁹, M. Santimaria¹⁹,

E. Santovetti^{25,j}, A. Sarti^{19,k}, C. Satriano^{26,s}, A. Satta²⁵, D.M. Saunders⁴⁸, D. Savrina^{32,33}, S. Schael⁹, M. Schellenberg¹⁰, M. Schiller⁴⁰, H. Schindler⁴⁰, M. Schlupp¹⁰, M. Schmelling¹¹, T. Schmelzer¹⁰, B. Schmidt⁴⁰, O. Schneider⁴¹, A. Schopper⁴⁰, K. Schubert¹⁰, M. Schubiger⁴¹, M.-H. Schune⁷, R. Schwemmer⁴⁰, B. Sciascia¹⁹, A. Sciubba^{26,k}, A. Semennikov³², A. Sergi⁴⁷, N. Serra⁴², J. Serrano⁶, L. Sestini²³, P. Seyfert²¹, M. Shapkin³⁷, I. Shapoval⁴⁵, Y. Shcheglov³¹, T. Shears⁵⁴, L. Shekhtman³⁶, V. Shevchenko⁶⁷, A. Shires¹⁰, B.G. Siddi^{17,40}, R. Silva Coutinho⁴², L. Silva de Oliveira², G. Simi^{23,o}, S. Simone^{14,d}, M. Sirendi⁴⁹, N. Skidmore⁴⁸, T. Skwarnicki⁶¹, E. Smith⁵⁵, I.T. Smith⁵², J. Smith⁴⁹, M. Smith⁵⁵, H. Snoek⁴³, M.D. Sokoloff⁵⁹, F.J.P. Soler⁵³, B. Souza De Paula², B. Spaan¹⁰, P. Spradlin⁵³, S. Sridharan⁴⁰, F. Stagni⁴⁰, M. Stahl¹², S. Stahl⁴⁰, P. Stefko⁴¹, S. Stefkova⁵⁵, O. Steinkamp⁴², S. Stemmler¹², O. Stenyakin³⁷, S. Stevenson⁵⁷, S. Stoica³⁰, S. Stone⁶¹, B. Storaci⁴², S. Stracka^{24,t}, M. Straticiuc³⁰, U. Straumann⁴², L. Sun⁵⁹, W. Sutcliffe⁵⁵, K. Swientek²⁸, V. Syropoulos⁴⁴, M. Szczekowski²⁹, T. Szumlak²⁸, S. T'Jampens⁴, A. Tayduganov⁶, T. Tekampe¹⁰, G. Tellarini^{17,g}, F. Teubert⁴⁰, E. Thomas⁴⁰, J. van Tilburg⁴³, M.J. Tilley⁵⁵, V. Tisserand⁴, M. Tobin⁴¹, S. Tolc⁴⁹, L. Tomassetti^{17,g}, D. Tonelli⁴⁰, S. Topp-Joergensen⁵⁷, F. Toriello⁶¹, E. Tournefier⁴, S. Tourneur⁴¹, K. Trabelsi⁴¹, M. Traill⁵³, M.T. Tran⁴¹, M. Tresch⁴², A. Trisovic⁴⁰, A. Tsaregorodtsev⁶, P. Tsopelas⁴³, A. Tully⁴⁹, N. Tuning⁴³, A. Ukleja²⁹, A. Ustyuzhanin^{35,67}, U. Uwer¹², C. Vacca^{16,f}, V. Vagnoni^{15,40}, S. Valat⁴⁰, G. Valenti¹⁵, A. Vallier⁷, R. Vazquez Gomez¹⁹, P. Vazquez Regueiro³⁹, S. Vecchi¹⁷, M. van Veghel⁴³, J.J. Velthuis⁴⁸, M. Veltri^{18,r}, G. Veneziano⁴¹, A. Venkateswaran⁶¹, M. Vernet⁵, M. Vesterinen¹², B. Viaud⁷, D. Vieira¹, M. Vieites Diaz³⁹, X. Vilasis-Cardona^{38,m}, V. Volkov³³, A. Vollhardt⁴², B. Voneki⁴⁰, D. Voong⁴⁸, A. Vorobyev³¹, V. Vorobyev³⁶, C. Voß⁶⁶, J.A. de Vries⁴³, C. Vázquez Sierra³⁹, R. Waldi⁶⁶, C. Wallace⁵⁰, R. Wallace¹³, J. Walsh²⁴, J. Wang⁶¹, D.R. Ward⁴⁹, H.M. Wark⁵⁴, N.K. Watson⁴⁷, D. Websdale⁵⁵, A. Weiden⁴², M. Whitehead⁴⁰, J. Wicht⁵⁰, G. Wilkinson^{57,40}, M. Wilkinson⁶¹, M. Williams⁴⁰, M.P. Williams⁴⁷, M. Williams⁵⁸, T. Williams⁴⁷, F.F. Wilson⁵¹, J. Wimberley⁶⁰, J. Wishahi¹⁰, W. Wislicki²⁹, M. Witek²⁷, G. Wormser⁷, S.A. Wotton⁴⁹, K. Wraight⁵³, S. Wright⁴⁹, K. Wyllie⁴⁰, Y. Xie⁶⁴, Z. Xing⁶¹, Z. Xu⁴¹, Z. Yang³, H. Yin⁶⁴, J. Yu⁶⁴, X. Yuan³⁶, O. Yushchenko³⁷, K.A. Zarebski⁴⁷, M. Zavertyaev^{11,c}, L. Zhang³, Y. Zhang⁷, Y. Zhang⁶³, A. Zhelezov¹², Y. Zheng⁶³, A. Zhokhov³², X. Zhu³, V. Zhukov⁹, S. Zucchelli¹⁵.

¹Centro Brasileiro de Pesquisas Físicas (CBPF), Rio de Janeiro, Brazil

²Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

³Center for High Energy Physics, Tsinghua University, Beijing, China

⁴LAPP, Université Savoie Mont-Blanc, CNRS/IN2P3, Annecy-Le-Vieux, France

⁵Clermont Université, Université Blaise Pascal, CNRS/IN2P3, LPC, Clermont-Ferrand, France

⁶CPPM, Aix-Marseille Université, CNRS/IN2P3, Marseille, France

⁷LAL, Université Paris-Sud, CNRS/IN2P3, Orsay, France

⁸LPNHE, Université Pierre et Marie Curie, Université Paris Diderot, CNRS/IN2P3, Paris, France

⁹I. Physikalisches Institut, RWTH Aachen University, Aachen, Germany

¹⁰Fakultät Physik, Technische Universität Dortmund, Dortmund, Germany

¹¹Max-Planck-Institut für Kernphysik (MPIK), Heidelberg, Germany

¹²Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany

¹³School of Physics, University College Dublin, Dublin, Ireland

¹⁴Sezione INFN di Bari, Bari, Italy

¹⁵Sezione INFN di Bologna, Bologna, Italy

¹⁶Sezione INFN di Cagliari, Cagliari, Italy

¹⁷Sezione INFN di Ferrara, Ferrara, Italy

¹⁸Sezione INFN di Firenze, Firenze, Italy

¹⁹Laboratori Nazionali dell'INFN di Frascati, Frascati, Italy

²⁰Sezione INFN di Genova, Genova, Italy

²¹Sezione INFN di Milano Bicocca, Milano, Italy

²²Sezione INFN di Milano, Milano, Italy

²³Sezione INFN di Padova, Padova, Italy

- ²⁴ *Sezione INFN di Pisa, Pisa, Italy*
- ²⁵ *Sezione INFN di Roma Tor Vergata, Roma, Italy*
- ²⁶ *Sezione INFN di Roma La Sapienza, Roma, Italy*
- ²⁷ *Henryk Niewodniczanski Institute of Nuclear Physics Polish Academy of Sciences, Kraków, Poland*
- ²⁸ *AGH - University of Science and Technology, Faculty of Physics and Applied Computer Science, Kraków, Poland*
- ²⁹ *National Center for Nuclear Research (NCBJ), Warsaw, Poland*
- ³⁰ *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest-Magurele, Romania*
- ³¹ *Petersburg Nuclear Physics Institute (PNPI), Gatchina, Russia*
- ³² *Institute of Theoretical and Experimental Physics (ITEP), Moscow, Russia*
- ³³ *Institute of Nuclear Physics, Moscow State University (SINP MSU), Moscow, Russia*
- ³⁴ *Institute for Nuclear Research of the Russian Academy of Sciences (INR RAN), Moscow, Russia*
- ³⁵ *Yandex School of Data Analysis, Moscow, Russia*
- ³⁶ *Budker Institute of Nuclear Physics (SB RAS) and Novosibirsk State University, Novosibirsk, Russia*
- ³⁷ *Institute for High Energy Physics (IHEP), Protvino, Russia*
- ³⁸ *ICCUB, Universitat de Barcelona, Barcelona, Spain*
- ³⁹ *Universidad de Santiago de Compostela, Santiago de Compostela, Spain*
- ⁴⁰ *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- ⁴¹ *Ecole Polytechnique Fédérale de Lausanne (EPFL), Lausanne, Switzerland*
- ⁴² *Physik-Institut, Universität Zürich, Zürich, Switzerland*
- ⁴³ *Nikhef National Institute for Subatomic Physics, Amsterdam, The Netherlands*
- ⁴⁴ *Nikhef National Institute for Subatomic Physics and VU University Amsterdam, Amsterdam, The Netherlands*
- ⁴⁵ *NSC Kharkiv Institute of Physics and Technology (NSC KIPT), Kharkiv, Ukraine*
- ⁴⁶ *Institute for Nuclear Research of the National Academy of Sciences (KINR), Kyiv, Ukraine*
- ⁴⁷ *University of Birmingham, Birmingham, United Kingdom*
- ⁴⁸ *H.H. Wills Physics Laboratory, University of Bristol, Bristol, United Kingdom*
- ⁴⁹ *Cavendish Laboratory, University of Cambridge, Cambridge, United Kingdom*
- ⁵⁰ *Department of Physics, University of Warwick, Coventry, United Kingdom*
- ⁵¹ *STFC Rutherford Appleton Laboratory, Didcot, United Kingdom*
- ⁵² *School of Physics and Astronomy, University of Edinburgh, Edinburgh, United Kingdom*
- ⁵³ *School of Physics and Astronomy, University of Glasgow, Glasgow, United Kingdom*
- ⁵⁴ *Oliver Lodge Laboratory, University of Liverpool, Liverpool, United Kingdom*
- ⁵⁵ *Imperial College London, London, United Kingdom*
- ⁵⁶ *School of Physics and Astronomy, University of Manchester, Manchester, United Kingdom*
- ⁵⁷ *Department of Physics, University of Oxford, Oxford, United Kingdom*
- ⁵⁸ *Massachusetts Institute of Technology, Cambridge, MA, United States*
- ⁵⁹ *University of Cincinnati, Cincinnati, OH, United States*
- ⁶⁰ *University of Maryland, College Park, MD, United States*
- ⁶¹ *Syracuse University, Syracuse, NY, United States*
- ⁶² *Pontifícia Universidade Católica do Rio de Janeiro (PUC-Rio), Rio de Janeiro, Brazil, associated to ²*
- ⁶³ *University of Chinese Academy of Sciences, Beijing, China, associated to ³*
- ⁶⁴ *Institute of Particle Physics, Central China Normal University, Wuhan, Hubei, China, associated to ³*
- ⁶⁵ *Departamento de Física, Universidad Nacional de Colombia, Bogota, Colombia, associated to ⁸*
- ⁶⁶ *Institut für Physik, Universität Rostock, Rostock, Germany, associated to ¹²*
- ⁶⁷ *National Research Centre Kurchatov Institute, Moscow, Russia, associated to ³²*
- ⁶⁸ *Instituto de Física Corpuscular (IFIC), Universitat de Valencia-CSIC, Valencia, Spain, associated to ³⁸*
- ⁶⁹ *Van Swinderen Institute, University of Groningen, Groningen, The Netherlands, associated to ⁴³*

^a *Universidade Federal do Triângulo Mineiro (UFTM), Uberaba-MG, Brazil*

^b *Laboratoire Leprince-Ringuet, Palaiseau, France*

^c *P.N. Lebedev Physical Institute, Russian Academy of Science (LPI RAS), Moscow, Russia*

^d *Università di Bari, Bari, Italy*

^e *Università di Bologna, Bologna, Italy*

^f *Università di Cagliari, Cagliari, Italy*

^g *Università di Ferrara, Ferrara, Italy*

^h *Università di Genova, Genova, Italy*

ⁱ *Università di Milano Bicocca, Milano, Italy*

^j *Università di Roma Tor Vergata, Roma, Italy*

^k *Università di Roma La Sapienza, Roma, Italy*

^l *AGH - University of Science and Technology, Faculty of Computer Science, Electronics and Telecommunications, Kraków, Poland*

^m *LIFAEELS, La Salle, Universitat Ramon Llull, Barcelona, Spain*

ⁿ *Hanoi University of Science, Hanoi, Viet Nam*

^o *Università di Padova, Padova, Italy*

^p *Università di Pisa, Pisa, Italy*

^q *Università degli Studi di Milano, Milano, Italy*

^r *Università di Urbino, Urbino, Italy*

^s *Università della Basilicata, Potenza, Italy*

^t *Scuola Normale Superiore, Pisa, Italy*

^u *Università di Modena e Reggio Emilia, Modena, Italy*

^v *Iligan Institute of Technology (IIT), Iligan, Philippines*