

Coulomb explosion imaging of small organic molecules at LCLS

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2012 J. Phys.: Conf. Ser. 388 022108

(<http://iopscience.iop.org/1742-6596/388/2/022108>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 134.76.223.157

This content was downloaded on 02/12/2016 at 11:13

Please note that [terms and conditions apply](#).

You may also be interested in:

[Inner-shell multiple ionization of polyatomic molecules with an intense x-ray free-electron laser studied by coincident ion momentum imaging](#)

B Erk, D Rolles, L Foucar et al.

[Probing ultrafast electronic and molecular dynamics with free-electron lasers](#)

L Fang, T Osipov, B F Murphy et al.

[Exploring few-photon, few-electron reactions at FLASH](#)

A Rudenko, Y H Jiang, M Kurka et al.

[Femtosecond x-ray photoelectron diffraction on gas-phase dibromobenzene molecules](#)

D Rolles, R Boll, M Adolph et al.

[Field-free molecular alignment probed by FLASH](#)

P Johnsson, A Rouzée, W Siu et al.

[Strongly aligned gas-phase molecules at free-electron lasers](#)

Thomas Kierspel, Joss Wiese, Terry Mullins et al.

[Stereochemical configuration and selective excitation of the chiral molecule halothane](#)

Martin Pitzer, Gregor Kastirke, Phillip Burzynski et al.

[AMO science at the FLASH and European XFEL free-electron laser facilities](#)

J Feldhaus, M Krikunova, M Meyer et al.

[Towards imaging of ultrafast molecular dynamics using FELs](#)

A Rouzée, P Johnsson, L Rading et al.

Coulomb explosion imaging of small organic molecules at LCLS

Benjamin Erk^{*1,2}, Artem Rudenko^{1,2}, Daniel Rolles^{1,3}, Benedikt Rudek^{1,2}, Lutz Foucar^{1,3},
Sascha Epp^{1,2}, Max Cryle³, Ilme Schlichting^{1,3}, Christoph Bostedt⁴, Sebastian Schorb⁴, John Bozek⁴,
Arnaud Rouzee⁵, Axel Hundertmark⁵, Tatiana Marchenko⁶, Mark Simon⁶, Frank Filsinger⁷,
Lauge Christensen⁸, Shankar De⁸, Sebastian Trippel⁹, Shinichi Wada⁴, Kiyoshi Ueda¹⁰,
Claus Dieter Schroeter², Joachim Ullrich^{1,2}

¹ Max Planck Advanced Study Group at CFEL, 22607 Hamburg, Germany, ² Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany, ³ Max-Planck-Institut für Medizinische Forschung, 69120 Heidelberg, Germany, ⁴ LCLS, SLAC National Accelerator Laboratory, Menlo Park, CA 94025, USA, ⁵ Max-Born Institute, 12489, Berlin, Germany, ⁶ Laboratoire de Chimie Physique-Matière et Rayonnement. Université Pierre et Marie Curie, Paris, France, ⁷ Fritz-Haber-Institut, 14195 Berlin, Germany, ⁸ Department of Physics and Astronomy, Aarhus University, Denmark, ⁹ Center for Free-Electron Laser Science, DESY, 22607 Hamburg, Germany, ¹⁰ IMRAM, Tohoku University, Sendai, Japan

Synopsis Fragmentation of small organic molecules by intense few-femtosecond X-ray free-electron laser pulses has been studied using Coulomb explosion imaging. By measuring kinetic energies and emission angles of the ionic fragments in coincidence, we disentangle different fragmentation pathways, for certain cases can reconstruct molecular geometry at the moment of explosion, and show how it depends on LCLS pulse duration.

The development of free-electron lasers (FEL) delivering intense ultrashort coherent X-ray pulses paves the way towards structural investigations of matter with atomic spatial and femtosecond time resolution. One of the basic underlying processes for all of these studies is an inner-shell (multi-) photon absorption followed by the Auger or radiative decay and structural rearrangement (atomic motion and charge redistribution), often proceeding on comparable femtosecond time scale.

Here we have studied multiple fragmentation of small organic molecules upon multiphoton inner-shell ionization by intense ($> 10^{16}$ W/cm²) short (3-200 fs) 2 keV pulses provided by the Linac Coherent Light Source (LCLS) at Stanford. In order to increase and localize X-ray absorption, we used methylselenol (CH₃SeH), ethylselenol (C₂H₅SeH) and phenylselenol (C₆H₅SeH) compounds containing high-Z selenium atom as a substitute for naturally occurring oxygen. The experiment was conducted in the CFEL-ASG Multi-Purpose (CAMP) end station [1] installed at the AMO beamline of LCLS. By measuring kinetic energies and emission angles of the ionic fragments in coincidence, we can separate different fragmentation pathways and, for certain cases, reconstruct molecular geometry (bond lengths and angles) at the moment of explosion. In particular, in case of methylselenol proton energies and emission angles in the molecular frame can be defined for each of the identified pairs of heavy ion fragments (shown in Fig. 1). Comparing these data

for a given final state and different pulse durations, we observe lower proton energies for longer pulses. This indicates molecular deformation during the pulse, which has direct implications for the studies of radiation damage. Overall, the results yield unique information on the structural rearrangement the molecule undergoes upon few-photon absorption, in particular, pointing to the ultrafast charge redistribution within the molecule.




Figure 1. Photoion-photoion coincidence spectrum measured after Coulomb explosion of methylselenol by 3 fs LCLS pulses at 2 keV photon energy.

References

[1] L. Strüder *et al* 2010 *Nucl. Instr. and Meth. A* **614** 483

* E-mail: Benjamin.Erk@asg.mpg.de