



Historical carbon dioxide emissions caused by land-use changes are possibly larger than assumed

Accepted version

This is the peer reviewed version of the following article:

Arneth, A., Sitch, S., Pongratz, J., Stocker, B. D., Ciais, P., Poulter, B., et al. (2017). Historical carbon dioxide emissions caused by land-use changes are possibly larger than assumed. *Nature Geoscience*, 10(2), 79-84. [doi:10.1038/ngeo288](https://doi.org/10.1038/ngeo288)

Conditions of use:

Articles published within the Springer Nature group of companies which are made available through academic repositories remain subject to copyright. The following restrictions on use of such articles apply:

Academic research only

1. Archived content may only be used for academic research. Any content downloaded for text based experiments should be destroyed when the experiment is complete.

Use must not be for Commercial Purposes

2. Archived content may not be used for purposes that are intended for or directed towards commercial advantage or monetary compensation by means of sale, resale, licence, loan, transfer or any other form of commercial exploitation ("Commercial Purposes"). Wholesale re-publishing is prohibited

Wholesale re-publishing is prohibited

3. Archived content may not be published verbatim in whole or in part, whether or not this is done for Commercial Purposes, either in print or online.

4. This restriction does not apply to reproducing normal quotations with an appropriate citation. In the case of text-mining, individual words, concepts and quotes up to 100 words per matching sentence may be used, whereas longer paragraphs of text and images cannot (without specific permission from Springer Nature).

Moral rights

5. All use must be fully attributed. Attribution must take the form of a link—using the article DOI—to the published article on the journal's website.

6. All use must ensure that the authors' moral right to the integrity of their work is not compromised.

Third party content

7. Where content in the document is identified as belonging to a third party, it is the obligation of the user to ensure that any use complies with copyright policies of the owner.

Use at own risk

8. Any use of Springer Nature content is at your own risk and Springer Nature accepts no liability arising from such use

Published in: Nature Geoscience

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 647204



1 **Historical carbon dioxide emissions due to land use changes possibly larger than assumed**

2 A Arneth (1), S Sitch (2), J Pongratz (3), B Stocker (4,5), P Ciais (6), B Poulter (7), A Bayer
3 (1), A Bondeau (8), L Calle (7), L. Chini (9), T Gasser (6), M Fader (8,10), P Friedlingstein (11), E
4 Kato (12), W Li (6), M Lindeskog (13), J E M S Nabel (3), TAM Pugh (1, 14), E Robertson (15), N
5 Viovy (6), C Yue (6), S Zaehle (16)

6

7

8 (1) Karlsruhe Institute of Technology, Dept. Atmospheric Environmental Research, Kreuzeckbahnstr.
9 19, 82467 Garmisch-Partenkirchen, Germany

10 (2) College of Life and Environmental Sciences, University of Exeter, Exeter, EX4 4RJ, UK

11 (3) Max Planck Institute for Meteorology, Bundesstr. 53, 20146 Hamburg, Germany

12 (4) Department of Life Sciences and Grantham Institute for Climate Change, Imperial College London,
13 Silwood Park, Ascot, SL5 7PY, UK

14 (5) Institute for Atmospheric and Climate Science, ETH Zürich, Universitätstrasse 16, 8092 Zürich,
15 Switzerland

16 (6) IPSL – LSCE, CEA CNRS UVSQ, Centre d'Etudes Orme des Merisiers, 91191 Gif sur Yvette France

17 (7) Institute on Ecosystems and Department of Ecology, Montana State University, Bozeman, MT
18 59717

19 (8) Institut Méditerranéen de Biodiversité et d'Ecologie marine et continentale, Aix-Marseille

20 Université, CNRS, IRD, Avignon Université, Technopôle Arbois-Méditerranée, Bâtiment Villemin, BP
21 80, 13545 Aix-en-Provence CEDEX 04, France

22 (9) Department of Geographical Sciences, University of Maryland, College Park, MD 20742, USA

- 23 (10) International Centre for Water Resources and Global Change, hosted by the German Federal
24 Institute of Hydrology. Am Mainzer Tor 1, 56068 Koblenz, Germany
- 25 (11) College of Engineering, Mathematics and Physical Sciences, University of Exeter, Exeter, EX4
26 4QE, UK
- 27 (12) The Institute of Applied Energy, Minato, Tokyo 105-0003, Japan
- 28 (13) Dept of Physical Geography and Ecosystem Science, Sölvegatan 12, Lund University,
29 22362 Lund, Sweden
- 30 (14) School of Geography, Earth & Environmental Sciences and Birmingham Institute of Forest
31 Research, University of Birmingham, Birmingham, B15 2TT, United Kingdom
- 32 (15) Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, UK
- 33 (16) Max Planck Institute for Biogeochemistry, 07701 Jena, Germany
- 34
- 35
- 36

37 **The terrestrial biosphere absorbs about 20% of fossil fuel CO₂ emissions. The overall**
38 **magnitude of this sink is constrained by the difference between emissions, the rate of**
39 **increase in atmospheric CO₂ concentrations and the ocean sink. However, the land sink**
40 **is actually composed of two largely counteracting fluxes that are poorly quantified: fluxes**
41 **from land-use change and CO₂ uptake by terrestrial ecosystems. Dynamic global**
42 **vegetation model simulations suggest that CO₂ emissions from land-use change have been**
43 **substantially underestimated because processes such as tree harvesting and land-clearing**
44 **from shifting cultivation have not been considered. Since the overall terrestrial sink is**
45 **constrained, a larger net flux as a result of land-use change implies that terrestrial uptake**
46 **of CO₂ is also larger, and that terrestrial ecosystems might have greater potential to**
47 **sequester carbon in the future. Consequently, reforestation projects and efforts to avoid**
48 **further deforestation could represent important mitigation pathways, with co-benefits for**
49 **biodiversity. It is unclear whether a larger land carbon sink can be reconciled with our**
50 **current understanding of terrestrial carbon cycling. In light of our possible**
51 **underestimation of the historical residual terrestrial carbon sink and associated**
52 **uncertainties, we argue that projections of future terrestrial carbon uptake and losses are**
53 **more uncertain than ever.**

54

55 The net atmosphere-to-land carbon flux (F_L) is typically inferred as the difference between relatively
56 well-constrained terms of the global carbon cycle: fossil fuel and cement emissions, oceanic carbon
57 uptake and atmospheric growth rate of CO₂ (see Textbox) ¹. In contrast, very large uncertainties exist in
58 how much anthropogenic land-use and land-cover change (F_{LULCC}) contributes to F_L , which propagates
59 into large uncertainties in the estimation of the ‘residual’ F_{RL} (see Box). The lack of confidence in
60 separating F_L into its component fluxes diminishes the predictive capacity for terrestrial carbon cycle

61 projections into the future. It restricts our ability to estimate the capacity of land ecosystems to continue
62 to mitigate climate change, and to assess land management options for land-based mitigation policies.

63 As land-use change emissions and the residual sink are spatially closely enmeshed, global-scale
64 observational constraints do not exist for estimating F_{LULCC} or F_{RL} separately. Dynamic Global
65 Vegetation Models (DGVMs) have over recent years been used to infer the magnitude and spatial
66 distribution of F_{LULCC} as well as of F_{RL} , while F_{LULCC} has traditionally been also derived from data-
67 driven approaches such as the bookkeeping method¹⁻³ (see Box). Although large, for some sources of
68 uncertainties in F_{LULCC} (such as differences in baseline years used for calculation, how environmental
69 effects have been considered, or assumptions about wood products) there is no good reason to believe
70 that these would introduce a systematic under- or overestimation⁴⁻⁶. However, until recently, most
71 processes related to land management and the subgrid-scale dynamics of land-use change have been
72 ignored in large-scale assessments of the terrestrial carbon balance, and we argue here that including
73 these missing processes might systematically increase the magnitude of F_{LULCC} . In turn, an upward
74 revision of F_{LULCC} implies through the global budget the existence of a substantially higher F_{RL} and
75 raises the question whether a larger F_{RL} is plausible given our understanding of the response of
76 ecosystems to changing environmental conditions.

77 **Gross land-cover transitions such as shifting cultivation (SC)**

78 Opposing changes in different land-use types can take place simultaneously within a region (see
79 methods, and Supplementary Figure), e.g. an area is converted from natural to managed land, whereas
80 an equal area within the same region might be abandoned or reforested, equating to a net zero land-
81 cover change. The magnitude of these bi-directional changes depends on the size of the area
82 investigated. Over thousands of km², the typical resolution of DGVMs, ignoring sub-grid changes can
83 have a substantial effect on the simulated carbon cycle, since accounting for the gross changes (e.g., the
84 parallel conversion to, and abandonment of, agricultural land in the same grid-cell) includes (rapid)
85 carbon losses from deforestation, (slow) loss from post-deforestation soil legacy effects, and (slow)

86 uptake in areas of regrowth. In sum this leads to younger mean stand-age, smaller biomass pools and
87 thus higher F_{LULCC} compared to net area-change simulations.

88 Gross area transitions are fundamental to LULCC dynamics in areas of shifting cultivation in the
89 tropics⁷, but also occur elsewhere⁸. Gross forest loss far exceeding net area loss can be demonstrated
90 from remote-sensing products globally⁹, although these products in themselves cannot distinguish
91 effects of logging from natural disturbance events such as fire or storms. Secondary forests in the tropics
92 can return to biomass carbon stocks comparable to old-growth forest within 5-6 decades¹⁰, but the same
93 is not the case for soil carbon. Also, fallow lengths in shifting cultivation systems tends to be shorter,
94 and show a decreasing trend in many regions¹¹. These dynamics result in the degraded vegetation and
95 reduced soil carbon stocks commonly observed in disturbed forest land¹².

96 **Wood harvest (*WH*)**

97 Until recently, global DGVM studies that accounted for LULCC concentrated on the representation of
98 conversion of natural lands to croplands and pastures, while areas under forest cover were represented
99 as natural forest, and hence by each model's dynamics of establishment, growth and mortality. Two
100 thirds to three quarters of global forests have been affected by human use, mainly harvest, as a source
101 of firewood, roundwood and secondary products, or for recreational purposes¹³. Between 1700-2000 an
102 estimated 86 PgC has been removed globally from forests due to wood harvest¹⁴. Wood harvest leads
103 to reduced carbon density on average in managed forests¹⁵ and can ultimately result in degradation in
104 the absence of sustainable management strategies. Furthermore, the harvest of wood can reduce litter
105 input, which lowers soil pools¹³. The effect of bringing a natural forest under any harvesting regime will
106 be net CO₂ emissions to the atmosphere, its time-dependency depending on harvest intensity and
107 frequency, regrowth, and by the fate and residence time of the wood products.

108 **Grazing and crop harvest (*GH*) and cropland management (*MC*)**

109 Management is not only fundamental for the carbon balance of forests, but also for pasture and
110 cropland. As with forests, accounting for management processes on arable lands has only recently been
111 included in DGVMs (see methods). Regular grazing and harvesting (*GH*), and more realistic crop

112 management processes (*MC*) such as flexible sowing and harvesting, or tillage, will enhance F_{LULCC} ¹⁶.
113 Over decadal timescales, conversion of forest to cropland has been observed to reduce soil carbon pools
114 by around 40% ¹⁷, resulting from reduced vegetation litter soil inputs and enhanced soil respiration in
115 response to tillage, although the effect and magnitude of the latter is being debated ¹⁸. Conversion to
116 pasture often has either little effect, or may even increase soil carbon ¹⁷.

117 **Impacts of land management processes on the carbon cycle**

118 The few DGVM studies published that account for the management of land more realistically ^{16,19-21}
119 consistently suggest a systematically larger F_{LULCC} over the historical period compared to estimates that
120 ignored these processes, with important implications for our understanding of the terrestrial carbon cycle
121 and its role for historical (and future) climate change. In order to assess if results from these initial
122 experiments hold despite differences among models, we compile here results from a wider set of
123 DGVMs (and one DGVM “emulator”, see methods and Supplementary Table 1), adopting the approach
124 described in ². F_{LULCC} was calculated as the difference between a simulation in which CO₂ and climate
125 were varied over the historical period, at constant (pre-industrial) land use, and one in which land use
126 was varied as well.

127 When accounting for shifting cultivation and wood harvest, F_{LULCC} was systematically enhanced
128 (Fig. 1). Shifting-cultivation, assuming that no shade-trees remain in cultivated areas, results in
129 increased cumulative F_{LULCC} over the period 1901-2014 on average by 35 ± 18 PgC (Fig. 1;
130 Supplementary Table 2). While three DGVMs had demonstrated this effect previously¹⁹⁻²¹, an upward
131 shift of F_{LULCC} was also found in the other models that performed additional *SC* simulations for this
132 study. Including wood harvest caused F_{LULCC} to increase over the same time period by a similar
133 magnitude to *SC*, 30 ± 21 PgC. Trends in wood-harvest-related F_{LULCC} over time differed between
134 models (Fig. 1) likely due to different rates of post-harvest regrowth, and assumptions about residence
135 time in different pools²². Including the harvest of crops and the grazing of pastures also resulted in larger
136 F_{LULCC} , since carbon harvested or grazed is consumed and released as CO₂ rapidly instead of decaying
137 slowly as litter and soil organic matter. Beyond harvest, accounting for more realistic cropland

138 management such as tillage processes also showed, with one exception (in which tillage effects were
139 not modelled, see methods) an enhancement of F_{LULCC} emissions.

140 When ignoring the additional land-use processes investigated here, average F_{LULCC} is 119 ± 50 PgC
141 (Supplementary Table 2). Adding effects of *SC*, *WH*, *GH* and *MC* enhance land-use change emissions
142 by, on average, 20-30% each (Fig. 2; Supplementary Table), with individually large uncertainties. The
143 total effects on F_{LULCC} are difficult to judge as models do not yet account for all land-use dynamics. For
144 instance, shifting cultivation and wood harvest effects are expected to enhance F_{LULCC} additively as there
145 is little overlap in the input dataset used by DGVMs regarding the areas that are assumed to be under
146 shifting cultivation, and areas where wood harvest occurs⁷. But in the case of accounting for harvest
147 and other management on arable lands and pastures, carbon cycle interactions with *SC* and *WH* cannot
148 be excluded because subsequent transitions could occur in a grid location, between primary vegetation
149 and cropland, pastures or secondary forests. The overall enhancement of F_{LULCC} therefore will need to
150 be explored with model frameworks that include all dynamic land-use change processes. DGVMs
151 currently contributing to the annual update of the global carbon budget account for some of the processes
152 examined here, but as yet not at all comprehensively, and we thus expect DGVM-based F_{LULCC} to
153 increase substantially compared to results reported in¹. As a consequence the discrepancy to book-
154 keeping estimates of F_{LULCC} will become larger, although results in²³ call for a broader range of book-
155 keeping approaches as well.

156 **Implications for the historical residual land sink**

157 In order to match F_L in the global carbon budget (Box) for the historical period a substantially larger
158 F_{LULCC} would need to be balanced by a corresponding increase in F_{RL} , which could be either due to
159 underestimated historical increase in GPP and vegetation biomass, overestimated heterotrophic carbon
160 loss, or both. The question arises if such a discrepancy is credible in light of today's understanding. For
161 instance, by compiling a number of observations Pan et al.²⁴ suggested a forest sink that is in line with
162 total carbon budget estimates¹. However, their study excluded savannahs, grasslands, and woodlands
163 and in semi-arid regions alone C uptake was estimated to be about 20% of the terrestrial sink (plus

164 around another 30% from other non-forested ecosystems), which also dominate the recent positive trend
165 in C uptake ²⁵. Reconstructing the Austrian historical forest sink from inventory data also suggested a
166 much larger residual sink, compared with (bookkeeping) model results ²⁶.

167 The response of photosynthesis to increasing CO₂ could underlie more than half of today's land carbon
168 sink ²⁷. Several recent lines of observation-based evidence suggest that GPP may have undergone much
169 stronger enhancement over the last century than currently calculated by DGVMs. These studies include
170 isotopic analysis of herbarium plant samples, of stable oxygen isotope ratios in atmospheric CO₂, and
171 accounting for the effect of leaf mesophyll resistance to CO₂ ²⁸⁻³⁰. Ciais et al. ³¹ inferred a pre-industrial
172 GPP of 80 PgC a⁻¹ based on measurements of oxygen isotopes in ice-core air, indicative for a 33%
173 difference to the often-used present-day GPP benchmark of ca. 120 PgC a⁻¹ ³² and independently
174 consistent with the 35% increase suggested by ²⁸. In contrast, the participating DGVMs in this study
175 show an average increase of GPP by only 15% between the first and last ten years of the simulation (not
176 shown).

177 Whether or not enhancements in GPP translate into increased carbon storage depends on other factors
178 such as nutrient and water supply, seen for instance in the mixed trends in stem growth found in forest
179 inventories ^{33,34}. Much work remains to better understand the response of ecosystem carbon storage to
180 increasing atmospheric CO₂ concentration ³⁵. Ultimately, enhanced growth will only result in increasing
181 carbon pools if turnover time does not change at the same rate ²². Besides GPP and heterotrophic
182 ecosystem respiration (ER), lateral carbon flows play an important role in the ecosystem carbon sink.
183 Recent syntheses that combined a range of observations, inventories of carbon stock changes, trade
184 flows and transport in waterways, estimated dissolved organic carbon losses to account for a flux of >
185 1.0 PgC a⁻¹, with an unknown historical trend ^{36,37}. The fate of this carbon is highly uncertain, but its
186 inclusion would enhance the calculated residual sink via an additional loss term (eqn. 1, textbox). Taken
187 together, a number of candidates for underestimated F_{RL} in today's models are plausible, and a
188 combination of the above listed processes likely. It remains to be seen whether a larger F_{LULCC} can be
189 supported by observation-based estimates. Several lines of evidence suggest that a common low-bias in

190 the historic F_{LULCC} could affect all DGVMs, and the challenge of resolving the many open issues will
191 stay with us for some years to come.

192 **Unknowns in historical LULCC reconstructions**

193 Patterns and historical trends of deforestation, cropland and pasture management or wood harvest are
194 uncertain. Land use reconstructions differ substantially in terms of the time, location and rate of LULCC
195 (see ³⁸ and reference therein). The DGVM and climate science community has mostly relied on the
196 LUH1 data-set by Hurtt et al. ⁷, chiefly because it provides the needed seamless time-series from the
197 historical period into future projections at the spatial resolution required by DGVMs. Clearly such a
198 globally applicable, gridded data-set must necessarily include simplifications. For instance, the assumed
199 uniform 15-year turnover in tropical shifting cultivation systems⁷ cannot account for the known variation
200 between a few years and one to two decades, or trends towards shorter fallow periods in some regions
201 (see ¹¹ and references therein), while there is also an increasing proportion of permanent agriculture.
202 Likewise, not only the amount of wood harvest but also the type of forestry (coppice, clear-cut, selective
203 logging, fuel-wood) will vary greatly in time and space, which is difficult to hindcast ^{39,40}.

204 In upcoming revisions to LUH1 (LUH-2, <http://luh.umd.edu/data.shtml>), forest-cover gross
205 transitions are now constrained by the remote sensing information⁹, and have overall been re-estimated
206 (Fig. 3). Whether or not this will result in reduced *SC* carbon loss estimates in recent decades remains
207 to be seen. At the same time, these historical estimates consider large gross transitions of land-cover
208 change only for tropical regions even though there is good reason to believe that bi-directional changes
209 occur elsewhere⁴¹. For Europe alone, a recent assessment that is relatively impartial to spatial resolution
210 estimated twice the area having undergone land-use transitions since 1900 when accounting for gross
211 *vs.* net area changes⁸. This leads to substantial increase in the calculated historical European F_{LULCC} ,
212 both in a bookkeeping-model and DGVM-based study⁴². Historical land carbon cycle estimates
213 therefore are not only highly uncertain due to missing LULCC processes, but equally so due to the
214 LULCC reconstructions *per se*. However, for a given reconstruction, accounting for additional processes

215 discussed here will always introduce a unidirectional enhancement in F_{LULCC} compared to ignoring these
216 processes.

217 **Implications for the future land carbon mitigation potential**

218 Our calculated increases in F_{LULCC} , in absence of a clear understanding of the processes underlying F_{RL} ,
219 notably strengthen the existing arguments to avoid further deforestation (and all ecosystem degradation)
220 – an important aspect of climate change mitigation, with considerable co-benefits to biodiversity and a
221 broad range of ecosystem service supply. One could also conjecture whether or not a larger historical
222 carbon loss through LULCC would imply a larger potential to sequester carbon through reforestation,
223 than thought so far. However, assessments of mitigation potentials must consider the often relatively
224 slow carbon gain in re-growing forests (compared to the rapid, large loss during deforestation), in
225 particular the sluggish replenishment of long-term soil carbon storage^{43,44}. What is more, trees grow
226 now, and will in future, under very different environmental conditions compared to the past. A warmer
227 climate increases mineralisation rates and hence enhances nutrient supply to plant growth, supporting
228 the CO₂ fertilisation effect, but also stimulates heterotrophic decay of existing soil carbon and/or flow
229 of dissolved carbon, with as yet no agreement about the net effects^{3,45}. Re-growing forests might also
230 in future be more prone to fire risk, and other episodic events such as wind-throw or insect outbreaks^{46,47},
231 crucial ecosystem features not yet represented well in models⁴⁸. This question of “permanence” has
232 been an important point of discussion at conferences under the UNFCCC, and also endangers the success
233 of payment-for-ecosystem-services schemes that target conservation measures, since it is unclear how
234 an increasing risk of losing carbon-uptake potential can be accounted for^{49,50}.

235 Given that we may be greatly underestimating the present-day F_{RL} , and therefore missing or
236 underestimating the importance of key driving mechanisms, projections of future terrestrial carbon
237 uptake and losses appear more fraught with uncertainty than ever. In the light of the findings summarised
238 here, this poses not only a major challenge when judging mitigation efforts, but also for the next
239 generation of DGVMs and Earth System models to assess the future global carbon budget. Future work

240 therefore needs to concentrate on representing the interactions between physiological responses to
241 environmental change in ecosystems with improved representations of human land management.

242

243

244

245 **References**

246

- 247 1 Le Quere, C. *et al.* Global Carbon Budget 2015. *Earth Sys. Sci. Data* 7, 349-396 (2015).
- 248 2 Sitch, S. *et al.* Recent trends and drivers of regional sources and sinks of carbon dioxide.
249 *Biogeosciences* 12, 653-679 (2015).
- 250 3 Ciais, P. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working*
251 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.*
252 (eds T.F. Stocker *et al.*) (Cambridge University Press, 2013).
- 253 4 Pongratz, J., Reick, C., Houghton, R. A. & House, J. I. Terminology as a key uncertainty in net
254 land use flux estimates. *Earth Syst. Dyn.* 5, 177-195 (2013).
- 255 5 Gasser, T. & Ciais, P. A theoretical framework for the net land-to-atmosphere CO₂ flux and its
256 implications in the definitions of "emissions from land-use change". *Earth Syst. Dyn.* 4, 171-
257 186 (2013).
- 258 6 Houghton, R. A. *et al.* Carbon emissions from land use and land-cover change. *Biogeosciences*
259 9, 5125-5142 (2012).
- 260 7 Hurtt, G. C. *et al.* Harmonization of land-use scenarios for the period 1500-2100: 600 years of
261 global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Clim.*
262 *Change* 109, 117-161 (2011).
- 263 8 Fuchs, R., Herold, M., Verburg, P. H., Clevers, J. G. P. W. & Eberle, J. Gross changes in
264 reconstructions of historic land cover/use for Europe between 1900 and 2010. *Glob. Change*
265 *Biol.* 21, 299-313 (2015).
- 266 9 Hansen, M. C., Stehman, S. V. & Potapov, P. V. Quantification of global gross forest cover
267 loss. *Proc. Nat. Acad. Sci.* 107, 8650-8655 (2010).

- 268 10 Poorter, L. *et al.* Biomass resilience of Neotropical secondary forests. *Nature* 530, 211-214
269 (2016).
- 270 11 van Vliet, N. *et al.* Trends, drivers and impacts of changes in swidden cultivation in tropical
271 forest-agriculture frontiers: A global assessment. *Glob. Env. Change* 22, 418-429 (2012).
- 272 12 Grace, J., Mitchard, E. & Gloor, E. Perturbations in the carbon budget of the tropics. *Glob.*
273 *Change Biol.* 20, 3238-3255 (2014).
- 274 13 Erb, K.-H. *et al.* Land management: data availability and process understanding for global
275 change studies. *Glob. Change Biol.*, gcb.13443 (2016).
- 276 14 Hurtt, G. C. *et al.* The underpinnings of land-use history: three centuries of global gridded land-
277 use transitions, wood-harvest activity, and resulting secondary lands. *Glob. Change Biol.*, 12,
278 1208-1229 (2006).
- 279 15 Noormets, A. *et al.* Effects of forest management on productivity and carbon sequestration: A
280 review and hypothesis. *For. Ecol. Manag.* 355, 124-140 (2015).
- 281 16 Pugh, T. A. M. *et al.* Carbon emission from land-use change is substantially enhanced by
282 agricultural management. *Env. Res. Lett.*, 124008 (2015).
- 283 17 Guo, L. B. & Gifford, R. M. Soil carbon stocks and land use change: a meta analysis. *Glob.*
284 *Change Biol.* 8, 345-360 (2002).
- 285 18 Powlson, D. S. *et al.* Limited potential of no-till agriculture for climate change mitigation. *Nat.*
286 *Clim. Change* 4, 678-683 (2014).
- 287 19 Shevliakova, E. *et al.* Carbon cycling under 300 years of land use change: Importance of the
288 secondary vegetation sink. *Glob. Biogeochem. Cycles* 23, GB2022 (2009).
- 289 20 Stocker, B. D., Feissli, F., Strassmann, K. M., Spahni, R. & Joos, F. Past and future carbon
290 fluxes from land use change, shifting cultivation and wood harvest. *Tellus B* 66, 23188 (2014).

- 291 21 Wilkenskjeld, S., Kloster, S., Pongratz, J., Raddatz, T. & Reick, C. H. Comparing the influence
292 of net and gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-
293 ESM. *Biogeosciences* 11, 4817-4828 (2014).
- 294 22 Friend, A. D. *et al.* Carbon residence time dominates uncertainty in terrestrial vegetation
295 responses to future climate and atmospheric CO₂. *Proc. Nat. Acad. Sci.* 111, 3280-3285 (2014).
- 296 23 Hansis, E., Davis, S. J. & Pongratz, J. Relevance of methodological choices for accounting of
297 land use change carbon fluxes. *Glob. Biogeochem. Cycles* 29, 1230-1246 (2015).
- 298 24 Pan, Y. *et al.* A Large and Persistent Carbon Sink in the World's Forests. *Science* 333, 988-993
299 (2011).
- 300 25 Ahlström, A. *et al.* The dominant role of semi-arid ecosystems in the trend and variability of the
301 land CO₂ sink. *Science* 348, 895-899 (2015).
- 302 26 Erb, K.-H. *et al.* Bias in attributing of forest carbon sinks. *Nat. Clim. Change* 3, 854-856 (2013).
- 303 27 Schimel, D., Stephens, B. B. & Fisher, J. B. Effect of increasing CO₂ on the terrestrial carbon
304 cycle. *Proc. Nat. Acad. Sci.* 112, 436-441 (2015).
- 305 28 Ehlers, I. *et al.* Detecting long-term metabolic shifts using isotopomers: CO₂-driven
306 suppression of photorespiration in C-3 plants over the 20th century. *Proc. Nat. Acad. Sci.* 112,
307 15585-15590 (2015).
- 308 29 Sun, Y. *et al.* Impact of mesophyll diffusion on estimated global land CO₂ fertilization.
309 *Proceedings of the National Academy of Sciences*, doi:10.1073/pnas.1418075111 (2014).
- 310 30 Welp, L. R. *et al.* Interannual variability in the oxygen isotopes of atmospheric CO₂ driven by
311 El Nino. *Nature* 477, 579-582 (2011).
- 312 31 Ciais, P. *et al.* Large inert carbon pool in the terrestrial biosphere during the Last Glacial
313 Maximum. *Nat. Geosc.* 5, 74-79 (2012).

- 314 32 Beer, C. *et al.* Terrestrial Gross Carbon Dioxide Uptake: Global Distribution and Covariation
315 with Climate. *Science* 329, 834-838 (2010).
- 316 33 McMahon, S. M., Geoffrey G Parker, and Dawn R Miller. 2010. . Evidence for a recent increase
317 in forest growth. *Proc. Nat. Acad. Sci* 107, 3611–3615 (2010).
- 318 34 van der Sleen, P. *et al.* No growth stimulation of tropical trees by 150 Years of CO₂ fertilization
319 but water-use efficiency increased. *Nat. Geosc.* 8, 24–28 (2015).
- 320 35 Pugh, T. A. M., Muller, C., Arneth, A., Haverd, V. & Smith, B. Key knowledge and data gaps
321 in modelling the influence of CO₂ concentration on the terrestrial carbon sink. *J. Plant Phys.*
322 203, 3-15 (2016).
- 323 36 Raymond, P. A. *et al.* Global carbon dioxide emissions from inland waters. *Nature* 503, 355-
324 359 (2013).
- 325 37 Regnier, P. *et al.* Anthropogenic perturbation of the carbon fluxes from land to ocean. *Nat.*
326 *Geosc.* 6, 597-607, doi:10.1038/ngeo1830 (2013).
- 327 38 Prestele, R. *et al.* Hotspots of uncertainty in land use and land cover change projections: a global
328 scale model comparison. *Glob. Change Biol.*, gcb.13337 (2016).
- 329 39 Bais, A. L. S., Lauk, C., Kastner, T. & Erb, K. Global patterns and trends of wood harvest and
330 use between 1990 and 2010. *Ecol. Econ.* 119, 326-337 (2015).
- 331 40 McGrath, M. J. *et al.* Reconstructing European forest management from 1600 to 2010.
332 *Biogeosciences* 12, 4291-4316 (2015).
- 333 41 Richter, D. D. & Houghton, R. A. Gross CO₂ fluxes from land-use change: implications for
334 reducing global emissions and increasing sinks. *Carb. Manag.* 2, 41-47 (2011).
- 335 42 Bayer, A. D., Lindeskog, M., Pugh, T. A. M., Fuchs, R. & Arneth, A. Uncertainties in the land
336 use flux resulting from land use change reconstructions and gross land transitions. *Earth Syst.*
337 *Dyn. Discuss.* (2016).

- 338 43 Korner, C. Slow in, rapid out - Carbon flux studies and Kyoto targets. *Science* 300, 1242-1243
339 (2003).
- 340 44 Krause, A., Pugh, T. A. M., Bayer, A. D., Lindeskog, M. & Arneith, A. Impacts of land-use
341 history on the recovery of ecosystems after agricultural abandonment. *Earth Syst. Dyn.* 7, 745-
342 766 (2016).
- 343 45 Zaehle, S., Jones, C. D., Houlton, B., Lamarque, J.-F. & Robertson, E. Nitrogen Availability
344 Reduces CMIP5 Projections of Twenty-First-Century Land Carbon Uptake. *J. Clim.* 28, 2494-
345 2511 (2015).
- 346 46 Seidl, R., Schelhaas, M. J., Rammer, W. & Verkerk, P. J. Increasing forest disturbances in
347 Europe and their impact on carbon storage. *Nat. Clim. Change* 4, 806-810 (2014).
- 348 47 Hantson, S. *et al.* The status and challenge of global fire modelling. *Biogeosciences* 13, 3359-
349 3375 (2016).
- 350 48 Running, S. W. Ecosystem disturbance, carbon, and climate. *Science* 321, 652-653 (2008).
- 351 49 Galik, C. S., Murray, B. C., Mitchell, S. & Cottle, P. Alternative approaches for addressing non-
352 permanence in carbon projects: an application to afforestation and reforestation under the Clean
353 Development Mechanism. *Mitigation and Adaptation Strategies for Global Change* 21, 101-
354 118 (2016).
- 355 50 Friess, D. A., Phelps, J., Garmendia, E. & Gomez-Baggethun, E. Payments for Ecosystem
356 Services (PES) in the face of external biophysical stressors. *Glob. Env. Change* 30, 31-42
357 (2015).

358

359 **Corresponding Author**

360 Correspondence and request for materials should be addressed to Almut Arneith, Almut.arneth@kit.edu

361

362 **Acknowledgements**

363 AA, ADB and TAMP acknowledge support from EU FP7 grants LUC4C (grant no. 603542) and
364 OPERAS (grant no.308393), and the Helmholtz Association in its ATMO programme and its impulse
365 and networking fund. MF, WL, CY and SS were also funded by LUC4C. JP and JEMSN were supported
366 by the German Research Foundation's Emmy Noether Program (PO 1751/1-1). EK was supported by
367 the ERTDF (S-10) from the Ministry of the Environment, Japan. ER was funded by LUC4C and by the
368 Joint UK DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). SZ has received
369 funding from the European Research Council (ERC) under the European Union's Horizon 2020 research
370 and innovation programme (grant agreement no. 647204; QUINCY). BDS is supported by the Swiss
371 National Science Foundation and FP7 funding through project EMBRACE (282672). PC received
372 support from the ERC SyG project IMBALANCE-P 'Effects of phosphorus limitations on Life, Earth
373 system and Society' Grant agreement no.: 610028.'

374

375 **Author contributions**

376 AA, SS, JP, BS conceived the study. BP, LC, AB, MF, EK, JEMN, ADB, ML, TAMP, ER, TG, NV,
377 CY, SZ made changes to model code and provided simulation results. AA and SS analysed results. BS,
378 PC, WL provided Fig. 3. AA wrote the first draft, all authors commented on the draft and discussion of
379 results.

380

381

382 **Textbox: Calculations of global terrestrial carbon uptake and removal**

383 The net atmosphere-to-land carbon flux (F_L) is generally inferred as the difference between other terms
384 of the global carbon cycle perturbation,

385
$$F_L = F_{FFC} - F_O - \frac{dA_{CO_2}}{dt} \quad (1)$$

386 where F_{FFC} are fossil fuel and cement emissions, F_O is the atmosphere-ocean carbon exchange (currently
387 an uptake) and $\frac{dA_{CO_2}}{dt}$ is the atmospheric growth rate of CO_2 (1). F_{FFC} and $\frac{dA_{CO_2}}{dt}$ are well known, and
388 the estimate of the decadal global ocean carbon sink is bounded by a range of observations ¹ such that
389 the net land carbon flux is relatively well constrained. By contrast, there is much less confidence in
390 separating F_L into a carbon flux from anthropogenic land use and land cover change (F_{LULCC}), and a
391 ‘residual’ carbon flux to the land (F_{RL} ; (2)) which is typically calculated as the difference from the other
392 carbon-cycle components:

$$393 \quad F_L = F_{RL} - F_{LULCC} \quad (2)$$

394 F_{LULCC} and F_{LR} are both made up of source and sink fluxes. Uncertainties in F_{LULCC} and F_{RL} are around
395 35% - 40% over the period 1870-2014 (when expressed as % of the cumulative mean absolute values),
396 compared to 13% for the cumulative ocean sink and 5% for fossil fuel burning and cement emissions¹.

397 F_{LULCC} has been modelled by the bookkeeping method (combining data-driven representative carbon
398 stocks trajectories and/or –for the satellite period– remote-sensing information on carbon density for
399 different biomes, with estimates of land-cover change), or by dynamic global vegetation models
400 (DGVMs; calculating carbon density of ecosystems with process-based algorithms; see methods).
401 DGVMs can also be used to calculate explicitly the magnitude and spatial distribution of F_{RL} ^{1,2} instead
402 of deducing its global value as a difference between F_L and F_{LULCC} as done in global budget analyses.
403 The bookkeeping approach has the advantage that carbon densities and carbon response functions that
404 describe the temporal evolution and fate of carbon after a LULCC disturbance can be based directly on
405 observational evidence ^{6,23}, but has to assume that local observations can be extrapolated to
406 regions/countries or biomes, thus partly ignoring spatial edaphic and climatic gradients of carbon stocks.
407 The DGVM-based simulations have the advantage to account for environmental effects on carbon stocks
408 through time, and account for spatial heterogeneity, but are poorly constrained by data. DGVMs and
409 bookkeeping models have similarly large degree of uncertainties ¹.

410

411 **Figure captions**

412

413 Figure 1: Difference in LULCC emission flux (Δ_{FLULCC}) due to individual processes. Coloured lines
414 represent different models, grey symbols and hairlines are average \pm one standard deviation.

415 a: wood harvest; b: shifting cultivation; c: harvest (using the grass functional type); d: full crop
416 representation

417

418 Figure 2: Response ratio of cumulative $F_{LULCC,1}$ and $F_{LULCC,0}$. See also Supplementary Table 1 and
419 methods for individual processes and models.

420

421 Figure 3: Comparison of net (a) and gross (b) forest / natural land change (in Million km²) between
422 different LULCC data sets. Changes in LUH1 data ⁷ represents the change of natural land because there
423 is no separate forest type in LUH1 while change in the other data sets indicates the forest change.

424

425

426

427 **Methods (and references for methods)**

428 1) General simulation set-up

429 Carbon fluxes from land-use change are derived as the difference between a simulation with historically
430 varying observed climate, atmospheric CO₂ concentration and land-cover change (S3) and one in which
431 land-cover change was held constant (S2)^{1,2}. Land-cover changes were taken from HYDE³ or LUH1⁴.
432 In S2, land-cover distribution was fixed. Gridded historical estimates of gross-transitions (shifting
433 cultivation in the tropics; *SC*) and wood harvesting (*WH*) were taken from⁴.

434 Spin up used repeated climate from the first decades of the 20th century, and constant CO₂ concentration
435 and land-cover distribution (for details, see section 2). Upon achieving steady-state, land-cover
436 distribution and CO₂ concentration were allowed to evolve transiently, whilst transient climate evolution
437 began at 1901. Atmospheric CO₂ concentration was taken from ice core data until ca. mid-20th century,
438 when atmospheric measurements became available². A “baseline” carbon flux related to land-use change
439 ($F_{LULCC,0}$; see Supplementary Table 1) is defined as excluding gross transitions and wood harvest, and
440 using the grass plant functional type to represent crop areas. Data in this Perspective article were from
441 previously published work, supplemented by from additional, new simulations. In cases where more
442 than one of the processes that are under investigation here were assessed by one model several S3
443 experiments were provided. While spin-up and model configurations differed between models, for S2
444 and S3 simulations of any one individual model the set-up was the same, which allows to identify the
445 effect of adding the individual processes. Section (2) provides a brief summary of relevant aspects of
446 models and simulation protocol, in particular where they differ from their previously published versions.

447

448 2) Individual models

449 2.1 JULES

450 Here, to implement crop harvest, four additional PFTs were added: C3 crops, C4 crops, C3 pasture and
451 C4 pasture, with identical parameter sets as the C3 and C4 grass PFTs. Lotka-Volterra equations⁵ are

452 used three times to calculate the vegetation distribution in natural areas, crop and pasture areas, with the
453 calculations in each area being independent of the others. Crop harvest is represented by diverting 30%
454 of crop litter to the fast product pool instead of to the soil; the fast product pool has a rapid decay
455 timescale of 1 year. Pasture is not harvested.

456 The model is forced by crop and pasture area from the Hyde 3.2 dataset ² and by CRU-NCEP climate^{1,2},
457 both at 1.875x1.25 degrees, using an hourly time-step, and updating vegetation distribution every ten
458 days. 1080 years of spin-up were run by fixing crop and pasture areas at 1860 levels and by repeating
459 1901-1920 climate and CO₂ concentrations.

460 2.2 JSBACH

461 The JSBACH version used here is similar to the version in ². S3 experiments include gross land-use
462 transitions and wood harvest ⁶. $F_{LULCC,0}$ in Supplementary Table 2 were calculated by subtracting the
463 individual contributions of these processes. Net transitions are derived from the gross transition
464 implementation, but by minimizing land conversions ⁶. Wood harvest ⁴ is taken not only from forest
465 PFTs but also shrubs and natural grasslands are harvested. Upon harvest, 20% of the carbon is
466 immediately released to the atmosphere; the rest is transferred into the litter and subject to soil dynamics.
467 JSBACH simulations were conducted at 1.9°x1.9° forced with remapped 1° LUH1 data from 1860-
468 2014 and daily climate calculated from the 6-hourly 0.5° CRU-NCEP product ² for the years 1901-2014.
469 The initial state in 1860 is based on a spin-up with 1860 CO₂ concentrations (286.42 ppm), cycling
470 (detrended) 1901-1921 climate and constant 1860 LUH1 wood harvest amounts. From 1860 annual CO₂
471 forcing was used, and after 1901 climate was taken from CRU-NCEP. In the no-harvest simulation the
472 1860 wood harvest amounts were applied throughout the whole simulated period.

473 2.3 LPJ-GUESS

474 SC: For implementing shifting cultivation, recommendations followed those by ⁴, with rotation periods
475 of 15 years. Simulations used the coupled carbon-nitrogen version of the model ⁷⁻⁸ Spin-up used constant
476 1701 land-cover and CO₂ concentration, and 1901-1930 recycled climate. Upon steady-state land-cover
477 and CO₂ were allowed to change from 1701, and climate from 1901 onwards⁹. When land is cleared,
21

478 76% of woody biomass and 71% of leaf biomass is removed and oxidised within one year, with a further
479 21 % of woody biomass assigned to a product pool with 25 year turnover time ⁹. Upon abandonment a
480 secondary forest stand is created and recolonization of natural vegetation takes place from a state of bare
481 soil. With forest rotation, young stands (above a minimum age of 15 years) are preferentially converted.

482 *GH/MC*: Simulations are taken from ⁸, using the carbon-only version of the model. 68% of deforested
483 woody biomass and 75% of leaf biomass is oxidised within one year, with a further 30% of woody
484 biomass going to the product pool. In the *GH* case, 50% of the above-ground biomass are annually
485 removed from the ecosystem. In *MC*, 90% of the harvestable organs and an additional 75% of above-
486 ground crop residues are removed each year. Simulations ran from 1850 to 2012, with 1850 land-cover
487 and CO₂ concentrations, and recycled climate (1901-1930) being used for spin-up.

488 All LPJ-GUESS simulations used CRU TS 3.23 climate ¹⁰.

489 2.4 LPJ

490 Compared to previous versions, the model now uses the World Harmonization Soils Database version
491 1.2 for soil texture and Cosby equations ¹¹ to estimate soil water holding capacity. Further developments
492 allow for gross land-use transitions and wood harvest to be prescribed. Changes include (1) the primary
493 grid-cell fraction only decreases in size; (2) secondary grid-cell fractions can decrease or increase in size
494 by combining with other secondary forest fractions, recently abandoned land, or fractions with recent
495 wood harvest; (3) deforestation results in an immediate flux to the atmosphere equal to 100% of
496 heartwood biomass and 50% of sapwood biomass; root biomass enters belowground litter pools, while
497 100% leaf and 50% of sapwood biomass becomes part of aboveground litter.

498 Wood harvest demand ⁴ on primary or secondary lands was met by the biomass in tree sapwood and
499 heartwood only. Only whole trees were harvested (i.e., tree-density was reduced); wood from
500 deforestation was not included to meet wood harvest demand. 100% of leaf biomass and 40% of the
501 sapwood and heartwood enters the aboveground litter, and 100% of root biomass enters the belowground
502 litter pools; 60% of sapwood and heartwood are assumed to go into a product pool. Of these, 55% go to
503 the 1-year product pool (emitted in the same year), 35% go to the 10-year product pool (emitted at rate

504 10% per year) and 10% go to the 100-year product pool (emitted at rate 1% per year). These delayed
505 pool-emission fluxes are part of the LULCC fluxes. After harvest, the harvested fraction is mixed with
506 existing secondary forest fraction, or a secondary fraction is created if none exists, while fully
507 conserving biomass. For simulations with shifting cultivation, grid-cell fractions that underwent land-
508 use change were not mixed with existing managed lands or secondary fractions until all land-use
509 transitions had occurred.

510 Simulations were performed using monthly CRU ¹⁰ (TS3.23) climate at 0.5° degrees, and finished in
511 year 2013. Spin-up was done using recycled 1901-20 climate, and using 1860 land-cover and CO₂. Upon
512 steady-state, land cover and CO₂ varied after 1860 and climate varied after 1900.

513 2.5 LPJmL

514 The LPJmL version used was as described in ¹²⁻¹⁴. In the baseline scenario all crops were simulated as
515 a mixture of C3 and C4 managed grasslands, 50% of the aboveground biomass is transferred to the
516 harvest compartment and assumed to be respired in the same year. Climate data was 1901-2014 CRU
517 TS v. 3.23 monthly datasets and land-use patterns from the HYDE 3.2 dataset. Simulations were
518 performed at 0.5° spatial resolution. Model spin-up used recycled climate data from 1901-1920, and
519 with land use patterns and CO₂ concentrations fixed to the 1860 value. Simulations from 1861-2014
520 were done with varying annual CO₂ concentration values, and varying land use patterns according to
521 the HYDE dataset, and with transient climate from 1901 until 2014.

522 2.6 LPX

523 Land-use change, including shifting cultivation and wood harvesting, is implemented as
524 described in¹⁵, using the full land-use transition and wood harvesting data provided ⁴. Wood
525 (heartwood and sapwood) removed by harvesting and land conversion is diverted to products
526 pools with turnover rates of 2 years (37.5%) and 20 years (37.5%). The rest, including slash
527 from roots and leaves is respired within the same year.

528 Simulation results shown here are based on employing the GCP 2015 protocol and input data².
529 LPX includes interactive C and N cycling with N deposition and N fertiliser inputs

530 ¹⁶. Simulations with shifting cultivation and wood harvesting were spun up to equilibrium under
531 land-use transitions and wood harvesting of year 1500 ¹⁵. Varying land-use transitions and wood
532 harvesting was included from 1500 onwards, with CO₂ and N deposition of year 1860 and
533 recycled climate from CRU TS 3.23, years 1901-1931. All simulations are done on a 1 x 1
534 degree spatial resolution and make use of monthly climate input. Original GCP standard input
535 files were aggregated to 1 x 1 degrees conserving area-weighted means (climate input) or
536 absolute area of cropland and pasture (land use input).

537 2.7 OCN

538 The OCN version used here is applied as in the framework of the annual carbon budget ². OCN includes
539 interactive C and N cycling with N deposition and N fertiliser inputs ¹⁷. Wood harvest was implemented
540 by first satisfying the prescribed wood extraction rate from wood production due to land-use change,
541 and then removing additional biomass proportionally from forested tiles. Wood (heartwood and
542 sapwood) removed by harvesting and land conversion is diverted to products pools with turnover rates
543 of 1 years (59.7%), 10 years (40.2% for tropical, and 29.9% for extratropical trees) and 100 years (10.4
544 % for extratropical trees)¹⁸. The remainder enters the litter pools. In case OCN's forest growth rate did
545 not suffice to meet the prescribed wood extraction rate, harvesting was limited to 5% of the total stand
546 biomass and assumed to stop if the stand biomass density fell below 1 kg C m⁻². These limits were set
547 to account for offsets in annual wood production between OCN's predicted biomass growth and the
548 assumptions in the Hurtt et al. database ⁴. These limits may lead to lower than prescribed wood harvest
549 rates in low productive areas. An additional run was performed with keeping wood harvest constant at
550 1860s level.

551 Simulations with wood harvesting were spun up to equilibrium using harvesting of the year 1860 ².
552 Varying land-use transitions or wood harvesting was included from 1860 onwards, with CO₂ and N
553 deposition of year 1860 and recycled climate from CRU-NCEP, years 1901-1931. All simulations are
554 done on a 1 x 1 degree spatial resolution and make use of daily climate input, which is disaggregated to
555 half-hourly values by means of a weather generator ¹⁹. Original GCP standard input files were

556 aggregated to 1 x 1 degrees conserving area-weighted means (climate input) or absolute area of cropland
557 and pasture (land use input).

558

559 2.8 ORCHIDEE

560 *WH*: Developments to the version included in ² include annual wood harvest, the total wood harvested
561 of a grid cell is removed from above-ground biomass of the different forest PFTs proportional (i) to its
562 fraction in the gridcell and (ii) also to its relative biomass among forest PFTs. This results in harvesting
563 more wood in biomass-rich forests. In cases of inconsistencies between the Orchidee and Hurtt forest
564 fraction, and to avoid forest being degraded from excessive harvest we assume that no more than 20%
565 of the total forest biomass of a gridcell can be harvested in one year. Hence the biomass actually
566 harvested each year can be slightly lower than prescribed ⁴. The harvested biomass enters 3 pools of 1,
567 10 and 100 residence years respectively (and is part of F_{LULCC}). Model runs were done at 0.5°x0.5°
568 resolution. Spin-up used recycled climate of 1901-1910. CO₂ concentration, land-cover and wood-
569 harvest we those of the year 1860. The model was run until the change in mean total carbon of 98% of
570 grid-points over a ten-year spin-up period was < 0.05%.

571 *SC*: Land cover transition matrices are upscaled from 0.5° LUH1 data ⁴ so no transition information is
572 lost in the low-resolution run. The minimum bi-directional fluxes between two land cover types in LUH1
573 were treated as shifting cultivation. The model was forced with CRU-NCEP forcing (v5.3.2), re-gridded
574 to 5° resolution from the original 0.5° resolution. Spin-up simulation used recycled climate data for
575 1901-1910 with atmospheric CO₂ held at 1750 level, and land cover fixed at 1500. Transient runs started
576 from 1501 until 2014, with CO₂ varying from 1750 and climate varying from 1901. In the transient run
577 for the control simulation, land cover is held constant at 1500; for the *SC* run, land cover varies by
578 applying annual land use transition matrices of shifting cultivation. All runs have been performed with
579 outputs on annual temporal resolution but forcing data is with 6-hourly.

580 2.9 OSCAR

581 A complete description of OSCAR v2.2 is provided by ²⁰. OSCAR is not a DGVM, but a compact Earth
582 system model calibrated on complex models. Here, it is used in an offline setup in which the terrestrial
583 carbon-cycle module is driven by exogenous changes in atmospheric CO₂ (IPCC AR5 WG1 Annex 2),
584 climate (CRU TS v. 3.23), and land-use and land cover (HYDE 3.2).

585 The global terrestrial biosphere is disaggregated into 9 regions (detailed by ²¹) and subdivided into 5
586 biomes (bare soil, forest, shrubland+grassland, cropland, pasture). The carbon-cycle in each of these 45
587 subparts is represented by a three-box model whose parameters are calibrated on DGVMs. The
588 preindustrial equilibrium (carbon densities and fluxes) is calibrated on TRENDY v2 models ¹. The
589 transient response of NPP, heterotrophic respiration and wildfires to CO₂ and/or climate is calibrated on
590 CMIP5 models ²². The impact of land-use and land-cover change on the terrestrial carbon-cycle is
591 modelled using a book-keeping approach. Coefficients used to allocate biomass after land-use or land-
592 cover change are based on ²³.

593 Since OSCAR v2.2 is meant to be used in a probabilistic setup we made an ensemble of 2400 simulations
594 in which the parameters (e.g. preindustrial equilibrium, transient responses, allocation coefficients) are
595 drawn randomly from the pool of available parameterizations. See ²⁰ for more details. The resulting
596 “OSCAR” values discussed and shown in the main text are the median of this ensemble.

597 2.10 VISIT

598 Implementation of climate, land-use change (gross transitions, *SC*) and wood harvest (*WH*) has not
599 changed from ². Land-use, land-use change, and wood harvest data for 1860-2014 were from LUH1 ⁴.
600 For *WH*, the amount of harvested biomass prescribed in ⁴ were transferred from simulated stem biomass
601 to 1-year product pool (emitted in entirety in same year of wood harvest), 10-year product pool, and
602 100-year product pool in a same manner as in the cleared biomass with land-use change described in ²⁴.
603 Non-harvested part of biomass were remain in the ecosystem. The fluxes from wood harvest pools are
604 included in the NBP calculations.

605 Climate data was 1901-2014 monthly CRU TS v. 3.23 and all simulations were conducted with 0.5°
606 spatial resolution. The model spin-up was performed recycling climate data from 1901-1920, and with
26

607 land use patterns and CO₂ concentrations fixed to the 1860 value. Simulations from 1860-2014 were
608 done with varying annual CO₂ concentration values, varying land use patterns according to LUH1,
609 recycling the climate from 1901-1920 in the period 1860-1900, and with transient climate from 1901
610 until 2014.

611

612 3) Data in Figure 3

613 Data for net forest change from FAO ²⁵ is calculated as the difference of forest area between 2000 and
614 2010 in each region. The same data were also used in the Houghton et al. bookkeeping model ²⁶. The
615 net forest change from Hansen et al. ²⁷ is based on satellite observations, and is their difference between
616 gross forest gain and gross forest loss during 2000-2012. Because the LUH1 data set ⁴ only has one type
617 of natural vegetation, and does not separate natural forest from natural grassland, the change in Figure
618 3 represents the total change of natural land. In Figure 3b, for LUH1 the gross loss includes transitions
619 from primary/secondary vegetation to cropland / pasture, while the gross gain is the sum of transitions
620 from cropland and pasture to secondary land. With grasslands and forests treated as separate land-cover
621 types in LUH2 (<http://luh.umd.edu/>), the change includes transitions from primary / secondary forest to
622 cropland / pasture (gross loss) and transitions from cropland / pasture to secondary forest (gross gain).
623 The net change for LUH1 or LUH2 is the difference between gross loss and gross gain. To be consistent
624 with ²⁷, the period calculated for LUH1 and LUH2 is also from 2000 to 2012.

625

626 Data and code availability

627 The data that support the findings of this study are available upon request, for access please contact
628 almut.arneth@kit.edu and s.a.sitch@exeter.ac.uk. We are unable to make the computer code of each of
629 the models associated with this paper freely available because in many cases the code is still under
630 development. However, individual groups are open to share code upon request, in case of interest please
631 contact the co-authors for specific models.

632 Access for LUH1 & LUH2 is under <http://luh.umd.edu/data.shtml>; the HYDE data are accessible via
633 <http://themasites.pbl.nl/tridion/en/themasites/hyde/download/index-2.html>

634

635 References

636 1 Sitch, S. *et al.* Recent trends and drivers of regional sources and sinks of carbon dioxide.
637 *Biogeosciences* **12**, 653-679 (2015).

638 2 Le Quere, C. *et al.* Global Carbon Budget 2015. *Earth System Science Data* **7**, 349-396 (2015).

639 3 Klein Goldewijk, L., Beusen, A., van Drecht, G. & de Vos, M. The HYDE 3.1 spatially explicit
640 database of human-induced global land use change over the past 12,000 years. *Globl Ecol.*
641 *Biogeogr.* **20**, 73–86 (2011).

642 4 Hurtt, G. C. *et al.* Harmonization of land-use scenarios for the period 1500-2100: 600 years of
643 global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Clim.*
644 *Change* **109**, 117-161 (2011).

645 5 Clark, D. B. *et al.* The Joint UK Land Environment Simulator (JULES), model description –
646 Part 2: Carbon fluxes and vegetation dynamics. *Geosci. Model Dev.* **4**, 701-722 (2011).

647 6 Wilkenskield, S., Kloster, S., Pongratz, J., Raddatz, T. & Reick, C. H. Comparing the influence
648 of net and gross anthropogenic land-use and land-cover changes on the carbon cycle in the MPI-
649 ESM. *Biogeosciences* **11**, 4817-4828 (2014).

650 7 Smith, B. *et al.* Implications of incorporating N cycling and N limitations on primary production
651 in an individual-based dynamic vegetation model. *Biogeosciences* **11**, 2027-2054 (2014).

652 8 Pugh, T. A. M. *et al.* Carbon emission from land-use change is substantially enhanced by
653 agricultural management. *Environmental Research Letters*, 124008 (2015).

- 654 9 Bayer, A. D., Lindeskog, M., Pugh, T. A. M., Fuchs, R. & Arneth, A. Uncertainties in the land
655 use flux resulting from land use change reconstructions and gross land transitions. *Earth Syst.*
656 *Dyn. Disc.* (2016).
- 657 10 Jones, P. & Harris, I. University of East Anglia Climatic Research Unit, CRU TS3. 21: Climatic
658 Research Unit (CRU) Time-Series (TS) Version 3.21 of High Resolution Gridded Data of
659 Month-by-month Variation in Climate (Jan. 1901—Dec. 2012). *NCAS British Atmospheric*
660 *Data Centre* (2013).
- 661 11 Cosby, B. J., Hornberger, G. M., Clapp, R. B. & Ginn, T. R. A STATISTICAL
662 EXPLORATION OF THE RELATIONSHIPS OF SOIL-MOISTURE CHARACTERISTICS
663 TO THE PHYSICAL-PROPERTIES OF SOILS. *Water Resources Res.* **20**, 682-690 (1984).
- 664 12 Bondeau, A. *et al.* Modelling the role of agriculture for the 20th century global terrestrial carbon
665 balance. *Glob. Change Biol.* **13**, 679-706 (2007).
- 666 13 Fader, M., von Bloh, W., Shi, S., Bondeau, A. & Cramer, W. Modelling Mediterranean agro-
667 ecosystems by including agricultural trees in the LPJmL model. *Geosc. Model Dev.* **8**, 3545-
668 3561 (2015).
- 669 14 Waha, K., van Bussel, L. G. J., Müller, C. & Bondeau, A. Climate-driven simulation of global
670 crop sowing dates. *Glob. Ecol. Biogeogr.* **12**, 247–259 (2012).
- 671 15 Stocker, B. D., Feissli, F., Strassmann, K. M., Spahni, R. & Joos, F. Past and future carbon
672 fluxes from land use change, shifting cultivation and wood harvest. *Tellus B*, **66**, 23188 (2014).
- 673 16 Stocker, B. D. *et al.* Multiple greenhouse-gas feedbacks from the land biosphere under future
674 climate change scenarios. *Nat. Clim. Change* **3**, 666-672 (2013).
- 675 17 Zaehle, S., Ciais, P., Friend, A. D. & Prieur, V. Carbon benefits of anthropogenic reactive
676 nitrogen offset by nitrous oxide emissions. *Nat. Geosc.* **4**, 601-605 (2011).

- 677 18 McGuire, A. D. *et al.* Carbon balance of the terrestrial biosphere in the twentieth century:
678 Analysis of CO₂, climate and land use effects with four process-based ecosystem models. *Glob.*
679 *Biogeochem. Cycles* **15**, 183-206 (2001).
- 680 19 Krinner, G., Ciais, P., Viovy, N. & Friedlingstein, P. A simple parameterization of nitrogen
681 limitation on primary productivity for global vegetation models. *Biogeosciences Discussions* **2**,
682 1243-1282 (2005).
- 683 20 Gasser, T. *et al.* The compact Earth system model OSCAR v2.2: description and first results.
684 *Geosc. Model Dev.* **submitted** (2016).
- 685 21 Houghton, R. A. & Hackler, J. L. Carbon flux to the atmosphere from land-use changes: 1850
686 to 1990. (Carbon Dioxide Information Analysis Center, Oak Ridge, Tennessee, 2001).
- 687 22 Arora, V. K. *et al.* Carbon-Concentration and Carbon-Climate Feedbacks in CMIP5 Earth
688 System Models. *J. Clim.* **26**, 5289-5314 (2013).
- 689 23 Mason, E. J., Yeh, S. & Skog, K. E. Timing of carbon emissions from global forest clearance.
690 *Nature Clim. Change* **2**, 682-685 (2012).
- 691 24 Kato, E., Kinoshita, T., Ito, A., Kawamiya, M. & Yamagata, Y. Evaluation of spatially explicit
692 emission scenario of land-use change and biomass burning using a process-based
693 biogeochemical model. *J. Land Use Sc.* **8**, 104-122 (2013).
- 694 25 FAO. Global Forest Resources Assessment 2010. (2010).
- 695 26 Houghton, R. A. *et al.* Carbon emissions from land use and land-cover change. *Biogeosciences*
696 **9**, 5125-5142 (2012).
- 697 27 Hansen, M. C. *et al.* High-Resolution Global Maps of 21st-Century Forest Cover Change.
698 *Science* **342**, 850-853 (2013).

699

700





