



Erratum

Erratum to “Synthesis and characteristic properties of novel Ba(II)Fe(III) layered double hydroxides” [Applied Clay Science 48 (2010) 214–217]

Dávid Srankó^a, Attila Pallagi^{a,b}, Ernő Kuzmann^c, Sophie E. Canton^d, Monika Walczak^d, András Sági^e,
Ákos Kukovecz^e, Zoltán Kónya^e, Pál Sipos^{a,*}, István Pálinkó^{b,*}

^a Department of Inorganic and Analytical Chemistry, University of Szeged, Dóm tér 7, Szeged, H-6720, Hungary

^b Department of Organic Chemistry, University of Szeged, Dóm tér 8, Szeged, H-6720, Hungary

^c Laboratory of Nuclear Chemistry, Eötvös Lóránd University, Pázmány Péter sétány 1/A, Budapest, H-1117, Hungary

^d Chemical Physics Department, Chemical Centre, Lund University, PO Box 124, Lund, SE-221 00, Sweden

^e Department of Applied and Environmental Chemistry, University of Szeged, Rerrich B. tér 1, H-6720 Szeged, Hungary

Figs. 6 and 7 appeared erroneously in the printed version (the energy scale of the various curves are horizontally shifted in a big way and therefore these graphs are misaligned). The figures correctly are as follows:




Fig. 6. Fe–K-edge XANES spectra of the solid substances obtained from solutions with varying concentrations of NaOH.




Fig. 7. Fe–K-edge X-ray absorption spectra on BaFe-LDH-10 samples after calcination at various temperatures.

DOI of original article: [10.1016/j.clay.2009.11.028](https://doi.org/10.1016/j.clay.2009.11.028).

* Corresponding authors. Tel.: +36 62 544 288; fax: +36 62544 200.

E-mail addresses: sipos@chem.u-szeged.hu (P. Sipos), palinko@chem.u-szeged.hu (I. Pálinkó).