



Functional diversity of microbial communities in pristine aquifers inferred by PLFA- and sequencing-based approaches

Valérie F. Schwab^{1,2}, Martina Herrmann^{3,4}, Vanessa-Nina Roth⁵, Gerd Gleixner⁵, Robert Lehmann¹, Georg Pohnert², Susan Trumbore⁵, Kirsten Küsel^{3,4}, and Kai U. Totsche¹

¹Friedrich Schiller University, Institute of Geosciences, Jena, Germany

²Friedrich Schiller University, Institute for Inorganic and Analytical Chemistry, Jena, Germany

³Friedrich Schiller University, Institute of Ecology, Jena, Germany

⁴German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Leipzig, Germany

⁵Max-Planck-Institute for Biogeochemistry, Jena, Germany

Correspondence to: Valérie F. Schwab (vf.schwab@uni-jena.de)

Received: 17 October 2016 – Discussion started: 3 November 2016

Revised: 23 March 2017 – Accepted: 4 April 2017 – Published: 31 May 2017

Abstract. Microorganisms in groundwater play an important role in aquifer biogeochemical cycles and water quality. However, the mechanisms linking the functional diversity of microbial populations and the groundwater physicochemistry are still not well understood due to the complexity of interactions between surface and subsurface. Within the framework of Hainich (north-western Thuringia, central Germany) Critical Zone Exploratory of the Collaborative Research Centre AquaDiva, we used the relative abundances of phospholipid-derived fatty acids (PLFAs) to link specific biochemical markers within the microbial communities to the spatio-temporal changes of the groundwater physicochemistry. The functional diversities of the microbial communities were mainly correlated with groundwater chemistry, including dissolved O₂, Fe_t and NH₄⁺ concentrations. Abundances of PLFAs derived from eukaryotes and potential nitrite-oxidizing bacteria (11Me16:0 as biomarker for *Nitrospira moscoviensis*) were high at sites with elevated O₂ concentration where groundwater recharge supplies bioavailable substrates. In anoxic groundwaters more rich in Fe_t, PLFAs abundant in sulfate-reducing bacteria (SRB), iron-reducing bacteria and fungi increased with Fe_t and HCO₃⁻ concentrations, suggesting the occurrence of active iron reduction and the possible role of fungi in mediating iron solubilization and transport in those aquifer domains. In more NH₄⁺-rich anoxic groundwaters, anammox bacteria and SRB-derived PLFAs increased with NH₄⁺ concentration, further evidencing the dependence of the anammox process on ammonium

concentration and potential links between SRB and anammox bacteria. Additional support of the PLFA-based bacterial communities was found in DNA- and RNA-based Illumina MiSeq amplicon sequencing of bacterial 16S rRNA genes, which showed high predominance of nitrite-oxidizing bacteria *Nitrospira*, e.g. *Nitrospira moscoviensis*, in oxic aquifer zones and of anammox bacteria in more NH₄⁺-rich anoxic groundwater. Higher relative abundances of sequence reads in the RNA-based datasets affiliated with iron-reducing bacteria in more Fe_t-rich groundwater supported the occurrence of active dissimilatory iron reduction. The functional diversity of the microbial communities in the biogeochemically distinct groundwater assemblages can be largely attributed to the redox conditions linked to changes in bioavailable substrates and input of substrates with the seepage. Our results demonstrate the power of complementary information derived from PLFA-based and sequencing-based approaches.

1 Introduction

Continental and marine subsurface environments represent the largest habitat on Earth for microbial life and therefore are of primary importance for energy fluxes on a global scale (Edwards et al., 2012). In terrestrial ecosystems, complex interactions between the surface and subsurface compartments (including aquifers), such as groundwater recharge and rainfall event-driven flow, influence the availability of

O₂, and the nature and abundance of bioavailable organic matter (OM; Benner et al., 1995; Kalbus et al., 2006). Young and shallow groundwaters tend to maintain the chemical characteristics of surface, i.e. higher O₂ levels and greater amounts of bioavailable substrates (e.g. labile OM) which support aerobic heterotrophic microbial activity (Landmeyer et al., 1996). In contrast, deep and especially fossil groundwaters tend to reflect the chemistry of the aquifer materials. They have typically lower concentrations of O₂ and bioavailable substrates which cause facultative anaerobes to switch to terminal electron acceptors with lower energy yield such as NO₃⁻, MnO₂, FeOOH and SO₄²⁻ (Chapelle and Lovley, 1992). In pristine aquifers, a low amount of OM typically results in a higher amount of terminal electron acceptors than electron donors (Chapelle, 2001). As many chemo-lithoautotrophs can use a variety of compounds to meet their energy needs in the dark subsurface, an increasing number of studies report the importance of chemo-lithoautotrophy in groundwater (Stevens and McKinley, 1995; Emerson et al., 2015; Herrmann et al., 2015). However, it is still not well understood how the composition and function of microbial communities in groundwaters depend on hydrology, chemistry and the relationship to groundwater recharge dynamics.

There are a number of ways to assess the composition and function of microbial communities in groundwaters. Intact polar lipids, mainly phospholipids, are important constituents of bacterial and eukaryotic cell membranes. They consist of a polar head group linked to a glycerol backbone with two fatty acids esterified to it. Because various phospholipid-derived fatty acids (PLFAs) are indicative of specific types or groups of bacteria in soil (e.g. Frostegård and Bååth, 1996; Frostegård et al., 2011) and aquifers (Green and Scow, 2000), PLFA-based studies are recognised as a valuable approach to infer the presence of specific microbial groups and to show trends in the spatial distribution of active microbial populations related to specific substrate utilization patterns in environments (e.g. Torsvik and Øvreås, 2002; Schneider et al., 2012). PLFAs that are commonly associated with a group or genus of bacteria are listed in Table 1. Additionally, the stable carbon isotope ratios ($\delta^{13}\text{C}$ values) of PLFAs reflect a combination of the source of microbial carbon and kinetic isotope fractionation effects associated with the carbon assimilation pathway (e.g. heterotrophy, autotrophy, methanotrophy; Teece et al., 1999; Zhang et al., 2003; Londry et al., 2004). Although a wide range of carbon isotope effects have been measured, in general autotrophs are expected to have PLFA $\delta^{13}\text{C}$ values more negative than heterotrophs (Blair et al., 1985; Teece et al., 1999; van der Meer et al., 2001; Zhang et al., 2003; Londry et al., 2004; Schouten et al., 2004). In particular, large isotope effects have been associated with anammox bacteria that have PLFA $\delta^{13}\text{C}$ values as much as 47‰ more negative than the dissolved inorganic carbon (DIC) source (Schouten et al., 2004).

Despite PLFAs being widely used in microbial ecology, their potential to assess changes in microbial communities

still remains the topic of many research efforts. One of the major limitations of PLFA-based studies is the definitive identification of the lipid sources, since many PLFAs commonly associated with a group or genus of bacteria (Table 1) may also be found, albeit in smaller amounts, in cell membranes of other organisms (Frostegård et al., 2011). A few PLFAs are highly specific, for example ladderanes are characteristic membrane constituents of anammox bacteria (Sinninghe Damsté et al., 2002, 2005) and have commonly been used to infer the presence of active anammox bacteria in diverse environments (Kuypers et al., 2003; Jaeschke et al., 2009). As these organisms are capable of anaerobically oxidizing ammonium with nitrite to molecular N₂, they play an essential role in N removal from marine (Dalsgaard et al., 2003; Burgin and Hamilton, 2007) and lacustrine environments (Yoshinaga et al., 2011). Yet their role in aquifer environments is only starting to be considered (Humbert et al., 2009). Another important limitation may reside in the PLFA extraction technique. Heinzlmann et al. (2014) showed that the proposed method to separate the glycolipids and phospholipids is incomplete and results in a significant proportion of glycolipids, betaine lipids and sulfoquinovosyl diacylglycerols (SQDGs) in the phospholipids fraction. Therefore, PLFA fractions might also contain fatty acids derived from glycolipids, betaine lipids, and to some extent SQDGs, and thus might not only reflect the active biomass.

In an attempt to overcome some of these limitations, we combined a detailed multivariate statistical analysis of PLFAs with PLFA $\delta^{13}\text{C}$ values, and DNA- and RNA-based Illumina MiSeq amplicon sequencing of bacterial 16S rRNA genes in groundwaters with different redox conditions and water chemistry (Kohlhepp et al., 2016). This approach allows for parallel study of microbial community composition and specific substrate consumption by evidencing specific PLFAs that respond significantly to changes in the groundwater chemistry. Microbial community structure and potential function assessed by PLFAs were confirmed by Illumina MiSeq amplicon sequencing targeting 16S rRNA genes and transcripts, providing a more detailed insight into bacterial community structure and taxonomic affiliation (Kozich et al., 2013). We showed that such a PLFA-based study has particular relevance and importance when trying to understand how micro-organisms in groundwater interact with their environment. This study provides baselines for future studies investigating the impact of changes in surface conditions on microorganism in carbonate-rock aquifer ecosystems.


2 Sampling and methods

2.1 Study site

The sampled groundwater wells are part of the monitoring well transect of the Hainich Critical Zone Exploratory (CZE: north-western Thuringia, central Germany) of the Collabo-

Table 1. Lipid markers proposed to represent a group or genus of microorganisms.

Lipid marker	Interpretation	References
Branched PLFAs	Gram-positive bacteria	Harwood and Russell (1984)
Mono-saturated PLFAs	Gram-negative bacteria	Wilkinson (1988)
10Me18:0	Actinomycetes	Zogg et al. (1997); Zelles et al. (1997)
18:1 ω 9c, 18:2 ω 6c, 18:3 ω 6c, 18:3 ω 3c	Fungi	Frostegard and Bååth (1996)
10Me16:0, cy18:0(ω 7,8)	<i>Desulfobacter</i>	Dowling et al. (1986)
11Me16:0, 16:1 ω 11	<i>Nitrospira moscoviensis</i>	Lipski et al. (2001)
i17:1 ω 7c, i15:1 ω 7c, i19:1 ω 7c	<i>Desulfovibrio</i>	Edlund et al. (1985), Kohring et al. (1994)
17:1 ω 6, 15:1	<i>Desulfobulbus</i>	Parkes and Calder (1985), Macalady et al. (2000)
i17:1 ω 5, 10Me18:1 ω 6, 11Me18:1 ω 6	<i>Thiobacillus</i>	Kerger et al. (1986, 1987)
20:2 ω 6, 20:3 ω 6, 20:4 ω 6, 22:5, 22:6	Fungi, <i>Protozoa</i> , Algae	Kennedy et al. (1993), Olsson (1999), White (1988); Volkman et al. (1989)
[3]-ladderane, [5]-ladderane	Anammox	Sinninghe Damsté et al. (2002, 2005)

**Figure 1.** Schematic geological cross section of the Hainich monitoring well transect (without karst features). The wells sampled for this study are numbered in black. The black colours in the wells indicate screen sections and accessed depths of the aquifer assemblages. Abbreviations: mm: Middle Muschelkalk; mo: Upper Muschelkalk; moTK: Trochitenkalk formation; moM & CB: Meissner formation with Cycloides-Bank; moW: Warburg formation; ku: Lower Keuper.

ative Research Centre (CRC) AquaDiva. AquaDiva aims to determine how deep signals of surface environmental conditions can be traced into the critical zone (Küsel et al., 2016). The wells access two distinct aquifer assemblages in marine sediments of the Upper Muschelkalk (mo) lithostratigraphic subgroup (Germanic Triassic, Middle Triassic epoch) at different depths and locations (Fig. 1). Wells in the hilltop recharge areas (H1, H2) were not sampled, due to very low groundwater levels or desaturation. Aquifers predominantly receive surface recharge in their outcrop areas at the eastern Hainich hillslope. The lower aquifer assemblage (subsequently referred to as HTL) represents one aquifer hosted in the Trochitenkalk formation (moTK), whereas the upper aquifer assemblage (referred to as HTU) comprises several aquifers and aquitards of the Meissner formation. The HTL, sampled at depths ranging from 41 to 88 m below the surface, is rich in O₂, whereas the upper aquifer found at depths from 12 to 50 m below the surface, is anoxic to sub-oxic. Both aquifer assemblages are found in alternating sequences of limestones and marlstones that are partly karstified (Kohlhepp et al., 2016). More details on the CZE and well constructions can be found in Küsel et al. (2016) and Kohlhepp et al. (2016).

2.2 Groundwater sampling

Groundwater was sampled for chemical analyses and colloidal and/or particulate organic matter in June, September and December of 2014 (Table 2) during regular sampling campaigns within the coordinated joint monitoring program of the CRC. Groundwater samples were collected at locations H3, H4, and H5 (i.e. the lower topographic positions of the well transect, Fig. 1). Wells H3.2, H4.2, H4.3, H5.2 and H5.3 reach into the HTU, while wells H3.1, H4.1 and H5.1 access the HTL aquifer (Fig. 1). The wells were originally drilled between 2009 and 2011, and were specifically designed for sampling groundwater (micro-) organisms and particles. Prior to sampling, stagnant water (at least three well volumes) was pumped out and discarded until the physico-chemical parameters pH, dissolved O₂ concentration, redox potential and specific electrical conductivity remained constant. Subsequently, ~1000 L of groundwater were filtered on-site using a submersible pump (Grundfos SQE 5-70, Grundfos, Denmark) connected to a stainless steel filter device (diameter 293 mm, Millipore USA) equipped with a removable pre-combusted (5 h at 500 °C) glass fiber filter (Sterlitech, USA) of fine porosity (0.3 µm), allowing

Table 2. Well depths, sampling dates and stratigraphic units of the monitoring wells studied.

Well name	Aquifer assemblage	Well depth* (m)	Sampling dates	Stratigraphic unit	Notes
H3.1	HTL	42.7–46.7	June 2014	moTK	well almost dry. Pumped only 100 L
H3.2	HTU	15–22	June, September 2014	moM	well dry in December 2014
H4.1	HTL	44.5–47.5	June, September 2014	moTK	well not accessible in December 2014
H4.2	HTU	8.5–11.5	June, September 2014	moM	well not accessible in December 2014
H4.3	HTU	8.5–12.5	June, September 2014	moM	well not accessible in December 2014
H5.1	HTL	84–88	June, September, December 2014	moTK	
H5.2	HTU	65–69	June, September, December 2014	moM	
H5.3	HTU	47–50	June, September, December 2014	moM	

* Depth of well screen section below surface. HTL: Hainich transect lower aquifer assemblage. HTU: Hainich transect upper aquifer assemblage. moTK: Upper Muschelkalk, Trochitenkalk formation. moM: Upper Muschelkalk, Meissner formation.

a water flow of ca. 20 L min⁻¹. Filters with the collected particulates were carefully removed, immediately transferred to dry ice and stored at -80 °C until analysis. Groundwater extraction temperature, redox potential, specific electrical conductivity, pH and dissolved O₂ concentration were monitored continuously during pumping in a flow-through cell equipped with the probes TetraCon 925, FDO 925, SenTix 980, ORP 900 and Multi 3430 IDS meter (WTW GmbH, Germany).

During the sampling campaign of June 2014, groundwater was additionally sampled for nucleic acid extraction. The groundwater was transferred to sterile glass bottles and kept at 4 °C. Within a few hours after sampling, 5–6 L of groundwater was filtered through 0.2 µm pore size polyethersulfone (PES) filters (Pall Corporation, USA), and 2 L were filtered through 0.2 µm pore size polycarbonate filters (Nuclepore, Whatman, United Kingdom) for extraction of DNA and RNA, respectively. Filters were immediately transferred to dry ice and stored at -80 °C until nucleic acid extraction.

2.3 Groundwater chemistry analyses

Concentration of the major anions (SO₄²⁻, Cl⁻, NO₃⁻ and PO₄³⁻; PES filter < 0.45 µm) were determined according to DIN EN ISO 10304-1 (2009) using an ion chromatograph (DX-120, Dionex, USA; equipped with an IonPac AS11-HC column and an IonPac AG11-HC pre-column). The redox-sensitive parameters (Fe²⁺, NO₂⁻, NH₄⁺) were determined by colorimetry (DR 890, Hach Company, USA) according to manufacturer's protocol following APHA (1981) and Rardon et al. (1966). The concentration of dissolved organic carbon (DOC) and DIC (filter < 0.45 µm) were determined by high-temperature catalytic oxidation (multi 18 N/C 2100S, Analytik Jena, Germany) according to DIN EN 1484 (1997). Total S (S_t), Mn (Mn_t) and iron (Fe_t) were analysed by ICP-OES (725 ES, Varian, Agilent, USA) according to DIN EN ISO 11885 (2009). The acid and base neutralizing capacity (ANC, BNC) by acid–base endpoint titration was determined according to DIN 38409-7 (2005). The ap-

proximated concentrations of HCO₃⁻ and CO₂⁻ were converted from ANC_{4,3} and BNC_{8,2} by simple replacement (cCO₂⁻ (mmol L⁻¹) = BNC_{8,2} (mmol L⁻¹); cHCO₃⁻ (mmol L⁻¹) = BNC_{4,3} (mmol L⁻¹)), assuming that other buffering species than those are negligible, in the nearly pH-neutral waters (Wisotzky, 2011).

2.4 PLFA extraction and pre-treatment

To minimize external contamination, all material (including filters) and glass in contact with the samples during extraction and purification were baked at 500 °C for 5 h to remove organic contaminants. Only trace levels of 16:0 fatty acid methyl ester (FAME) have been detected in blank extracts. PLFAs were extracted from filters using a method slightly modified from Bligh and Dyer (1959) and Seifert et al. (2013). The filters were cut into small pieces and extracted in a phase solution of chloroform–methanol (2 : 1; v/v) with a 0.005 M phosphate buffer. The solution was rotated and shaken for 4 h. Chloroform and water (1 : 1; v/v) were then added to the mixture. After shaking, the chloroform phase, containing the Bligh–Dyer extract (BDE), was separated from the water–MeOH phase and concentrated by a rotary evaporator. The BDE was then partitioned into the conventionally defined neutral lipid, glycolipid and phospholipid fractions by chromatography (solid phase extraction: SPE, 6 mL column) on pre-activated silica gel (Merck silica mesh 230–400, 2 g pre-activated for 1 h at 100 °C) using chloroform (12 mL), acetone (12 mL) and methanol (48 mL), respectively. The phospholipids were converted to FAMES using mild-alkaline hydrolysis and methylation (White et al., 1979). The different fatty acids were then separated using NH₂ column (Chromabond 3 mL, 500 mg) with 3 mL of hexane–DCM (3 : 1; v/v) for eluting the unsubstituted FAMES; 3 mL of DCM–ethylacetate (9 : 1; v/v) for the hydroxy FAME and 6 mL of 2 % acetic acid in methanol for unsaponifiable lipids. To quantify the recovery, the standard, 1,2-dinonadecanoyl-sn-glycero-3-phosphatidyl-choline (Avanti Polar Lipids, Inc. USA) was

added on clean pre-combusted glass filters that were treated exactly the same as the samples following the above protocol. The formed C19:0 FAME was quantified to calculate a mean recovery of 82 %. To test the efficiency of the separation between the glycolipids and the phospholipids, the glycolipid standard digalactosyl diglyceride (Sigma Aldrich) and the phospholipid standard 1,2-dinonadecanoyl-sn-glycero-3-phosphatidyl-choline were run through the SPE column using the above protocol. The absence of phospholipid-derived FA (C19:0) in the glycolipid fraction and glycolipid-derived fatty acids (C17:2) in the phospholipid fractions points to an efficient separation and thus a major origin of the studied FAME from phospholipid head groups.

2.5 Nucleic acid extraction, amplicon sequencing, and sequence analysis

DNA was extracted from the PES filters using the Power Soil DNA extraction kit (Mo Bio, CA, USA) following the manufacturer's instructions. RNA was extracted from polycarbonate filters using the Power Water RNA Isolation Kit (Mo Bio, CA, USA). Traces of co-extracted genomic DNA were removed using Turbo DNA free (Thermo Fisher Scientific, Germany), and reverse transcription to cDNA was performed using ArrayScript Reverse Transcriptase (Thermo Fisher Scientific) as described previously (Herrmann et al., 2012). DNA and cDNA obtained from the groundwater samples from PNK51 were shipped to LGC Genomic GmbH (Berlin, Germany) for Illumina MiSeq amplicon sequencing of the V3–V5 region of 16S rRNA genes and transcripts, using the primer combination Bakt_341F and Bakt_805R (Herrmann et al., 2011). Sequence analysis was performed using Mothur v. 1.36 (Schloss et al., 2009), following the MiSeq SOP (http://www.mothur.org/wiki/MiSeq_SOP; Kozich et al., 2013). Quality-trimmed sequence reads were aligned to the SILVA reference database (v 119; Quast et al., 2013). Potential chimeric sequences were detected and removed using the UCHIME algorithm implemented in Mothur. Taxonomic classification of sequence reads was based on the SILVA reference database (v 119). To facilitate comparisons across samples, sequence-read numbers per sample were normalized to the smallest number of sequence reads obtained across all samples using the subsample command implemented in Mothur. Raw data from 16S rRNA amplicon Illumina sequencing were submitted to the European Nucleotide Archive database under the study accession number PRJEB14968 and sample accession numbers ERS1270616 to ERS1270631.

2.6 Gas chromatography (GC) and gas chromatography–mass spectrometry

A total of 10 % of the PLFA extracts were used for peak identification and relative quantification using a gas chromatograph (Trace 1310 GC) coupled to a triple quadrupole

mass spectrometer (TSQ-8000; Thermo Fisher, Bremen, Germany) at the Friedrich Schiller University Jena's Institute of Inorganic and Analytical Chemistry (Germany). The GC was equipped with a TG 5silms capillary column (60 m, 0.25 mm, 0.25 μm film thickness). Helium was used as carrier gas at a constant flow of 1.2 mL min⁻¹. The GC oven was programmed to have an initial temperature of 70 °C (hold 1 min) and a heating rate of 2 °C min⁻¹ until 250 °C followed by a heating rate of 50 °C min⁻¹ until 280 °C and held for 5 min. The PTV injector was operated in splitless mode at an initial temperature of 70 °C. Upon injection, the injector was heated to 280 °C at a programmed rate of 14.5 °C s⁻¹ and held at this temperature for 2 min. Afterwards, the PTV was heated to 320 °C and held for 5 min. FAMES were quantified relative to an internal standard nonadecanoic acid-methyl ester (19:0) added prior to GC analysis. FAMES were identified based on the mass spectra and on retention time of standards. Standard nomenclature is used to describe PLFAs. The number before the colon refers to the total number of C atoms; the number(s) following the colon refers to the number of double bonds and their location (after the “ ω ”) in the fatty acid molecule. The prefixes “me”, “cy”, “i” and “a” refer to the methyl group, cyclopropane groups, and iso- and anteiso-branched fatty acids, respectively.

2.7 PLFA distribution and statistical analyses

The 47 PLFAs, expressed in percentage, were investigated in the different wells (Supplement Table S1). The sum of the PLFAs considered to be predominantly of bacterial origin (BactPLFA; *i*15:0, *a*15:0, 15:0, 16:1 ω 7, 16:0, *cy*17:0, 18:1 ω 7, 18:0 and *cy*19:0) was used as an index of the bacterial biomass (Bossio and Scow, 1998; Frostegård and Bååth, 1996). The fungal biomass (FunPLFA) was estimated from the sum of the relative abundance of the 18:2 ω 6c (Bååth et al., 1995), 18:3 ω 6c (Hamman et al., 2007) and 18:1 ω 9c (Myers et al., 2001); these were all significantly correlated with each other. Gram-positive (G+) bacteria were represented by the sum of PLFAs: *i*12:0, *i*13:0, *a*15:0 and *i*15:0 (Kaur et al., 2005). Gram-negative (G-) bacteria included 16:1 ω 7c, *cy*17:0, 18:1 ω 7c and *cy*19:0 (Kaur et al., 2005). The ratios of FunPLFA / BactPLFA and G+ / G- were calculated from the above PLFAs.

The PLFA data and 29 environmental parameters were used for principal component analysis (PCA) and redundancy analysis (RDA) using Canoco for Windows, version 5 (Microcomputer Power, Ithaca, New York, United States). Before regression, the data were centred and standardized. We used PCA to emphasise strong variations and similarities of the PLFA distributions between the wells and to identify patterns in the dataset. RDA is used to determine PLFA variations and similarities (response variables) that can be significantly explained by different environmental parameters (explanatory variables). This technique helps to identify the environmental parameters that have the highest effects on the

PLFA distribution, i.e. on the microbial communities in the different wells.

Additionally, we used variation partitioning analyses with conditional effects to determine the variations in PLFA composition between the different wells that can be explained significantly by the preselected environmental variables. To visualise the PLFAs acting significantly with the environmental variables (predictor), we used PLFA environmental-variable *t*-value biplots (Šmilauer and Lepš, 2014). These plots can be used to approximate the *t*-value of the regression between a particular PLFA and an environmental variable. The PLFAs are represented by arrows projecting from the origin. Those with a preference for higher values of the environmental variable are enclosed by a red (indicating positive relationship) circle. Inversely, those with preference for low values of the corresponding environmental variable have their arrow tips enclosed by a blue (indicating negative relationship) circle.

2.8 Compound-specific stable isotope carbon measurements

The carbon stable isotope composition of pre-purified PLFAs was determined using a GC–C–IRMS system (Deltaplus XL, Finnigan MAT, Bremen, Germany) at the Max Planck Institute for Biogeochemistry, Jena. Analyses were performed using 50 % of the total amount of PLFA extracts. The gas chromatograph (HP5890 GC, Agilent Technologies, Palo Alto USA) was equipped with a DB1-ms column (60 m, 0.25 mm internal diameter, 0.52 µm film thickness; Agilent). The injector at 280 °C was operated in splitless mode with a constant flow of 1 mL min⁻¹. The oven temperature was maintained for 1 min at 70 °C, heated with 4 °C min⁻¹ to 300 °C and held for 15 min, then heated by 30 °C min⁻¹ to 330 °C and held 3 min. Isotope values, expressed in the delta notation (‰), were calculated with ISODAT version software relative to the reference CO₂. Offset correction factor was determined on a daily basis using a reference mixture of *n*-alkanes (*n*-C₁₇ to *n*-C₃₃) of known isotopic composition. The carbon isotopic composition of the reference *n*-alkanes was determined off-line using a thermal conversion elemental analyser (TCEA) (Thermo Fisher, Bremen, Germany) interfaced to the DELTA V PLUS IRMS system via a Conflo III combustion interface (Thermo Fisher, Bremen, Germany; Werner and Brand, 2001). The contribution of the methyl carbon derived from the methanol after mild-alkaline hydrolysis and methylation of the PLFAs to the FAME was removed by isotopic mass balance, with $\delta^{13}\text{C}_{\text{PLFA}} = [(N_{\text{PLFA}} + 1) \times \delta^{13}\text{C}_{\text{FAME}} - \delta^{13}\text{C}_{\text{MeOH}}] / N_{\text{PLFA}}$, where *N* is the number of carbon atoms in the PLFA and $\delta^{13}\text{C}_{\text{FAME}}$ stands for the measured values of the methylated PLFAs (Kramer and Gleixner, 2006). The carbon isotope composition of MeOH used for derivatization ($\delta^{13}\text{C}$ value = -31.13 ± 0.03 ‰) was determined off-line using a thermal conversion elemental analyzer (TC/EA) (Thermo Fisher, Bremen, Germany) inter-

faced to the DELTA V PLUS irMS system via a Conflo III combustion interface (Thermo Fisher, Bremen, Germany).

3 Results


3.1 Groundwater physico-chemistry

The deeper aquifer assemblage, HTL (wells H3.1, H4.1 and H5.1), had higher mean concentration of O₂ (3.7 ± 1.0 mg L⁻¹) than the shallow aquifer assemblage, HTU (wells H4.2, H4.3, H5.2 and H5.3). Groundwater extracted from HTU wells were anoxic, with O₂ < 0.02 mg L⁻¹ (Supplement Table S2 and Fig. 2), except for well H3.2 that had mean O₂ = 2.4 ± 0.7 mg L⁻¹. No significant differences in the content of DOC (mean = 2.3 ± 1.0 mg L⁻¹) were measured between the different aquifers. The HTL had higher mean concentration of sulfate (183.5 ± 110.9 mg L⁻¹) than the anoxic HTU (76.4 ± 3.1 mg L⁻¹). The highest concentrations of nitrate were measured in the well H3.2 (30.0 ± 3.3 mg L⁻¹) of the HTU. Higher mean concentrations of total iron (Fe_t = 0.1 ± 0.08 mg L⁻¹), TIC (86.6 ± 7.0 mg L⁻¹) and HCO₃⁻ (4.69 ± 0.07 mg L⁻¹), the latter measured as acid-neutralizing capacity (Wisotzky, 2011), were found in the anoxic groundwater of the wells H4.2 and H4.3 than of the wells H5.2 and H5.3, which had mean Fe_t = 0.01 ± 0.00 mg L⁻¹, TIC = 75.6 ± 5.4 mg L⁻¹ and HCO₃⁻ = 4.02 ± 0.2 mg L⁻¹ (Fig. 2). Inversely, mean concentrations of total sulfur (S_t = 26.1 ± 4.9 mg L⁻¹), sulfate (76.7 ± 14.8 mg L⁻¹) and ammonium (0.62 ± 0.16 mg L⁻¹) were higher in the anoxic groundwater of the wells H5.2 and H5.3 than of the wells H4.2 and H4.3 that had mean S_t = 12.3 ± 0.5 mg L⁻¹, SO₄²⁻ = 37.6 ± 2.0 mg L⁻¹ and NH₄⁺ = 0.13 ± 0.06 mg L⁻¹ (Fig. 2 and Supplement Table S2).

The PCA analyses, using the physico-chemical parameters of the groundwater, separate the wells into three main groups (Fig. 3), with 73.6 % of the variability explained by the first three principal components (PCs): PC1, 32.8 %; PC2, 23.8 %; and PC3, 16.9 %. The conductivity, redox potential and the concentrations of Ca²⁺, SO₄²⁻, S_t and O₂ positively correlated (response > 0.5) with PC1 separating the oxic to sub-oxic wells H5.1, H4.1, H3.1 and H3.2 from the anoxic wells H4.2–H4.3 and H5.2–H5.3. The concentrations of NH₄⁺, K⁺ and Mg²⁺ inversely correlated (response < 0.5) with PC1, separating wells H5.2–H5.3 from the others. The Fe_t, TIC and HCO₃⁻ positively correlated along PC2 and mainly separated the anoxic wells between location H4 and H5. Groundwaters in location H5 have lower Fe_t, TIC and HCO₃⁻ concentrations but higher NH₄⁺ and K⁺ concentrations, whereas higher Fe_t, TIC and HCO₃⁻ concentrations but lower NH₄⁺ and K⁺ concentrations were measured at location H4.


Table 3. FunPLFA / BactPLFA, G− / G+ and cy17:0 / 16:1 ω 7c ratios averaged in the upper aquifer (HTU) and lower aquifer (HTL) as well as in the anoxic groundwater at locations H4 and H5.

	FunPLFA	SD	G−	SD	G+	SD	FunPLFA / BactPLFA	SD	G+ / G−	SD	cy17:0 / 16 ω 7c	SD
HTL	7.8	3.3	27.2	7.1	8.9	2.6	0.2	0.1	0.4	0.2	0.2	0.1
HTU	8.2	9.2	29.4	8.0	11.4	4.4	0.2	0.2	0.4	0.2	0.0	0.0
H4.2–H4.3	19.0	7.8	25.5	4.5	16.3	1.5	0.5	0.2	0.7	0.1	0.0	0.0
H5.2–H5.3	1.9	2.0	34.0	8.4	9.5	3.2	0.1	0.1	0.3	0.1	0.0	0.0

**Figure 2.** Variations of the chemical compositions of the groundwaters relevant for the discussion. HTL and HTU refer to the wells of the lower and upper aquifer assemblage, respectively. Chemical compositions of the groundwater of the wells H4.2–H4.3 and H5.2–H5.3 of the HTU are given separately for comparison.

3.2 PLFA distribution and statistical analyses

The 16:1 ω 7c (mean $22.2 \pm 8.9\%$), 16:0 (mean $13.4 \pm 2.3\%$) and 18:1 ω 7c (mean $5.2 \pm 2.6\%$), common in most bacteria, were the most abundant PLFAs in both aquifer assemblages (Supplement Table S1). The PLFAs 10Me16:0 (mean $7.8 \pm 5.6\%$), 17:1 ω 6c (mean $1.2 \pm 1.0\%$) and i17:1 (mean

**Figure 3.** Principal component analysis (PCA) of the groundwater physico-chemical compositions. Vectors indicate the steepest increase of the respective physico-chemical parameter. The different wells are represented by dots with different colours: blue for oxidic groundwater, yellow for sub-oxidic groundwater, dark red and violet for anoxic groundwater richer in Fe_t and NH_4^+ . Note the separation between the lower and upper aquifer (HTL and HTU, respectively) and the anoxic wells at location H4.2–H4.3 and H5.2–H5.3.

$0.8 \pm 0.7\%$), derived from Deltaproteobacteria mainly encompassing SRB, iron-reducing or iron-oxidizing bacteria, were dominant only in the anoxic groundwater, whereas the 11Me16:0 (mean $4.2 \pm 4.7\%$) was found in high relative abundance in the oxidic groundwaters. The [3]- and [5]-ladderane PLFAs specific to anammox bacteria were found in the anoxic wells H5.2 and H5.3 and the sub-oxidic well H3.2 in a relative abundance of up to 5.0%. The highest fungal biomass, based on the FunPLFA ratios (Table 3), was observed in the anoxic wells H4.2 and H4.3 (mean 19.0 ± 7.8), whereas the lowest was observed in the anoxic wells H5.2 and H5.3 (mean 1.9 ± 2). The Gram-negative (G−) bacteria were more abundant than Gram-positive bacteria (G+) in both HTU and HTL (Table 3: mean G+ / G− ratio = 0.4 ± 0.2). The highest values of the G+ / G− ratios were in the anoxic wells H4.2 and H4.3 (mean 0.7 ± 0.1).

A PCA analysis explained 56.5% of the PLFA variation. PC1 and PC2, respectively explaining 29.1 and 15.9% of




Figure 4. Principal component analysis (PCA) of PLFAs composition. The different wells are represented by dots with different colours: blue for oxic groundwater, yellow for sub-oxic or oxic groundwater, dark red and violet for anoxic groundwater richer in Fe_t and NH_4^+ . Note the separation between the lower and upper aquifer (HTL and HTU, respectively) and the anoxic wells at locations H4.2–H4.3 and H5.2–H5.3.

overall variability (Fig. 4), separated the wells into the same three groups evidenced by the PCA analysis of the groundwater chemistry (Fig. 3). The wells of the HTU assemblage were separated along PC1; wells from sites H4.2–H4.3 separated from those of the sites H5.2–H5.3. Along PC2, the wells were separated between the oxic (wells H3.1, H4.1 and H5.1), sub-oxic (well H3.2) and anoxic groundwaters (H4.2–H4.3 and H5.2–H5.3). The RDA analyses showed that O_2 , Fe_t and NH_4^+ concentrations or O_2 , HCO_3^- and NH_4^+ concentrations explained the greatest proportion (39.9 %) of the PLFA variability (Fig. 5). Well grouping obtained using the RDA analysis was consistent with the results of the PCA. The first RDA axis (21.5 %) separated the anoxic wells of the upper aquifer according to Fe_t or HCO_3^- (wells H4.2–H4.3) and NH_4^+ (wells H5.2–H5.3) concentration. The second RDA axis (14.6 %) separated suboxic to oxic (mainly lower aquifer) from anoxic groundwater (upper aquifer assemblage). In the following discussion, the wells are separated according to the PCA and RDA analyses into these three main groups.

To identify the individual effects of O_2 , Fe_t and NH_4^+ on the explained PLFA variation, we used variation partitioning with conditional effects implemented in Canoco 5 (Heikkinen et al., 2004; Roth et al., 2015). Because these environmental variables were the most significant factors, their combined variation was set to explain 100 % of total PLFA variation in each RDA plot. In our case, the following fractions explained the PLFA distribution by effect of O_2 alone: $a = 19.7\%$, effect of NH_4^+ alone: $b = 22.0\%$, effect of Fe_t alone: $c = 13.4\%$, by combined effects of O_2 and NH_4^+ : $d = 22.3\%$, by combined effects of Fe_t and NH_4^+ : $e = 29.2\%$, by combined effect of O_2




Figure 5. Redundancy analysis (RDA) of PLFAs, used as species, and the most significant environmental parameters O_2 , NH_4^+ and Fe_t that explained 39.9 % of the variability. The different wells are represented by dots with different colours: blue for oxic groundwater, yellow for sub-oxic groundwater, dark red and violet for anoxic groundwater richer in Fe_t and NH_4^+ .

and Fe_t ; and $f = 25.9\%$. The fraction g (-32.4%) explained the combined effect of the three environmental variables (Fig. 6). The PLFA environmental-variable O_2 t -plot (Fig. 6a) showed that the relative abundance of Me15:0, 16:1 ω 11c, cy17:0, 11Me16:0, 18:1, 22:5 and 22:6 increased significantly with O_2 concentration and the relative abundance of 10Me12:0, i 13:0, a 15:0, 17:1 and [5]-ladderane decreased with O_2 concentration. The PLFA environmental-variable Fe_t t -values biplot (Fig. 6b) showed that 10Me12:0, 17:1, 18:1 ω 9c, 18:1 ω 7c and 12:0 relative abundance increased with Fe_t concentration, whereas 10Me16:0, i 17:1, [3]-ladderane and [5]-ladderane relative abundance decreased. Inversely, the PLFA environmental-variable NH_4^+ t -values biplot (Fig. 6c) showed that 10Me16:0, 17:1, [3]-ladderane and [5]-ladderane relative abundance increased with NH_4^+ concentration, whereas 10Me12:0, 12:0, 18:1 ω 9c, 18:1 ω 7c and 17:1 relative abundance decreased.

3.3 PLFA $\delta^{13}\text{C}$ values

The PLFA $\delta^{13}\text{C}$ values for individual compounds ranged from -26 to -68.8% (Supplement Table S3, Fig. 7 and Table 4). The most negative mean $\delta^{13}\text{C}$ values were found in the anoxic groundwater from locations H5.2 and H5.3 ($-48.0 \pm 10.5\%$ and $-45.9 \pm 11.7\%$, respectively) and in the suboxic groundwater at the location H3.2




Figure 6. Variation-partitioning *t*-value biplots showing the PLFAs significantly correlated with the environmental variables (a) O₂, (b) Fe_t and (c) NH₄⁺. Results of the significance test of the variation partitioning are shown in the associated table. The PLFAs are represented by arrows projecting from the origin. Concentration changes, between sampling data of a particular PLFA, are significantly related to concentration changes in the environmental variables, when the arrow tip of those PLFA is enclosed within circles. The arrow tip is enclosed within the red circle for positive correlation and inversely within the blue circle for negative correlation.

Table 4. PLFA δ¹³C values averaged for the sampled month in the different wells.

PLFA	H3.2	SD	H4.2	SD	H4.3	SD	H5.2	SD	H5.3	SD	H3.1	SD	H4.1	SD	H5.1	SD
<i>i</i> 13:0	-39.2	3.5	-42.9		-37.7	0.8	-53.6	1.6	-51.2	10.3	-38.8		-40.7	1.8	-40.4	3.4
<i>a</i> 15:0	-40.3	0.7	-40.8		-35.9	0.3	-45.4	1.4	-43.0	2.8			-40.5	0.8	-37.1	1.5
<i>i</i> 15:0	-43.7	0.0	-47.4		-36.3	0.8	-55.0	3.3	-56.8	12.7			-46.0	0.8	-42.1	2.5
16:17 ω c	-47.4	0.6	-37.9	1.6	-36.0	2.3	-44.9	2.3	-44.3	1.3	-42.1		-40.0	3.8	-40.2	2.1
16:1 ω 11c	-44.4	2.4	-40.5	1.8	-36.0	4.0	-36.0	1.7	-34.6	4.1	-39.4		-27.8	2.4	-28.1	0.7
C16:0	-45.2	3.0	-34.8	2.6	-34.4	1.2	-45.3	2.4	-42.9	3.4	-36.4		-34.3	1.0	-35.4	2.2
10Me16:0	-49.9	0.9			-42.7		-57.6	3.3	-54.6	7.4	-42.2				-40.0	2.2
11Me16:0	-30.5										-28.3		-26.5	0.6	-25.7	2.5
cy17:0											-33.2				-26.3	1.6
<i>i</i> 17:1	-45.1						-42.6	2.0	-46.1							
18:1 ω 9c	-34.0	0.5	-30.2	0.1	-30.9	3.5	-36.7	1.8	-33.4	0.0	-32.1		-31.4	0.0	-31.0	1.4
18:1 ω 7c	-42.5	0.3	-32.5	0.3	-32.2	1.8	-39.6	2.3	-40.2	2.1	-32.4		-34.9	2.5	-34.6	1.2
18:1 ω 5c			-33.5		-30.0				-31.7		-35.1		-31.6			
18:0	-35.5	4.4	-42.1	1.0	-36.7	1.3	-36.1		-33.5		-32.9		-32.9		-31.5	1.7
cy19:0	-45.2															
20:4	-42.3												-33.1	1.0	-34.7	0.9
20:5	-41.0												-33.4	0.0		
[3]-lad	-62.9						-64.0	0.3	-63.7	0.8						
[5]-lad	-68.8						-67.0	0.9	-67.7	1.5						
mean	-44.6		-38.3	1.2	-35.4		-48.0		-46.0		-35.7		-34.9		-34.4	
SD	9.4		5.4	1.0	3.5		10.5		11.4		4.5		5.5		5.5	

($-45.4 \pm 9.0\%$) and coincided with the presence of the [5]- and [3]-ladderane. In those wells, the *i*13:0 ($-52.4 \pm 2.0\%$), *i*15:0 ($-55.6 \pm 2.0\%$), 10Me16:0 ($-56.1 \pm 2.1\%$) and *i*17:1 ($-44.3 \pm 2.0\%$) were slightly ^{13}C -enriched compared to both [5]- and [3]-ladderane ($-65.6 \pm 2.0\%$). More positive mean PLFA $\delta^{13}\text{C}$ values were measured in the anoxic wells H4.2 and H4.3 ($-36.8 \pm 2.1\%$) and in the oxic wells H5.1, H4.1 and H3.1 ($-35.3 \pm 1.1\%$). In those wells, the $\delta^{13}\text{C}$ values of the *i*13:0, *i*15:0 and 10Me16:0 were in the same range as the other PLFAs (Fig. 7). The most positive $\delta^{13}\text{C}$ values were measured for 16:1 ω 11c and 11Me16:0 in the oxic wells H5.1 and H4.1 (mean $-28.2 \pm 2.5\%$) and for 18:1 ω 9c (mean $-30.2 \pm 2.3\%$) in the anoxic wells H4.2 and H4.3.

3.4 Bacterial community composition based on 16S rRNA gene sequences

Based on Illumina sequencing of DNA-based 16S rRNA gene amplicons, bacterial communities were largely dominated by members of the phylum Nitrospirae and of Candidate Division OD1, followed by Delta- and Betaproteobacteria, Planctomycetes, and Alpha- and Gammaproteobacteria (Fig. 8a). Members of the Nitrospirae were especially abundant in the groundwater of the anoxic wells H5.2 and H5.3 as well as the oxic wells H4.1 and H5.1, while this phylum only contributed a minor fraction to the groundwater of the anoxic wells H4.2 and H4.3 and the oxic wells H3.1 and H3.2 (Fig. 8a). In addition, we performed sequencing of 16S rRNA amplicons derived from the extracted RNA to get insight into which taxonomic groups harbour protein synthesis potential, as proposed by Blazewicz et al. (2013). RNA-based community analysis targeting 16S rRNA sequences has traditionally been used as an approximation of the currently active fraction of the microbial community. However, this interpretation is critical since many cells may retain high ribosome contents even in a dormant state (Filion et al., 2009; Sukenik et al., 2012), and thus rRNA content of cells does not necessarily indicate current metabolic activity, especially in low-nutrient environments such as groundwater (reviewed in Blazewicz et al., 2013). Here, we used this approach to investigate whether key microbial groups identified by PLFA-based analysis were supported to be metabolically active or have the potential to resume metabolic activities based on the detection of the corresponding 16S rRNA gene sequences on the RNA level. In general, members of the Candidate Division OD1 formed only a minor part of the community obtained by RNA-based amplicon sequencing, while members of the phyla Nitrospirae, Planctomycetes and Proteobacteria showed the largest relative abundances (Fig. 8b). Members of the phylum Nitrospirae were especially highly represented in the RNA-based analyses of wells H3.2, H4.1, H5.2 and H5.3. Among the Proteobacteria, Deltaproteobacteria were more frequently represented in the RNA-based analysis of communities of wells H3.1, H3.2, H5.2 and H5.3, while Al-

phaproteobacteria showed a higher relative abundance in the groundwater of wells H4.2, H4.3 and H5.1 (Fig. 8b).

Bacterial phyla and classes may harbour organisms with a high diversity of different metabolisms. Therefore, as some source-specific PLFAs displayed strong relationships with the environmental variables O_2 , NH_4^+ and Fe_t , we specifically focused on groups potentially involved in iron oxidation and reduction, sulfate reduction, anammox, and nitrite oxidation. Here, relative fractions of reads assigned to bacterial genera known to be involved in either of these processes were summed up to get an estimation of the potential for these processes within the microbial community with both DNA- and RNA-based analyses. On the level of DNA-based sequencing, bacteria involved in iron oxidation accounted for 0.25 to 6.2 % of the sequence reads across sites (Fig. 9a), while they accounted for 0.24 to 2.8 % on the level of the RNA-based analyses with the highest relative fraction of bacteria potentially involved in iron oxidation at wells H5.1 and H5.3 (Fig. 9b). Differences across sites and aquifers were more pronounced for bacteria involved in iron reduction, which were accounted for by 0.16 to 3.7 % of the sequence reads on the DNA level but for 0.15 to 20.4 % on the RNA level, with the highest number of sequence reads affiliated with known iron reducers in the groundwater of well H4.3 (Fig. 9b). Bacteria related to the genera *Acidiferrobacter*, *Gallionella* and *Sideroxydans* were the most frequent genera among the known iron oxidizers, while members of the genera *Albidiferax* and *Ferribacterium* dominated the iron-reducing groups. Bacterial groups potentially involved in sulfur reduction included the genera *Desulfacinum*, *Desulfovibrio*, *Desulfosporosinus* and *Desulfatiferula* as the most frequent groups and accounted for 0.2 to 2.8 % of the sequence reads on the DNA level and 0.4 to 10.4 % on the RNA level, with the maximum in the anoxic well H4.2 (Fig. 9). Anammox bacteria mostly represented by the Candidatus genera *Brocadia* and *Kuenenia* accounted for 0.6 to 3.0 % of the sequence reads on the DNA level and for 1.1 to 16.8 % on the RNA level, with the highest fractions in the groundwater of the wells H3.1, H5.1, H5.2 and H5.3 (Fig. 9). Finally, we observed large fractions of potential nitrite oxidizers mostly related to the genus *Nitrospira*, with the vast majority of the *Nitrospira*-affiliated reads especially in the lower aquifer assemblage showing a high sequence similarity to the 16S rRNA gene sequence of *Nitrospira moscoviensis* (96–99 %). Moreover, reads associated with the genus *Nitrospira* may also include potential comammox organisms (Pinto et al., 2016). Relative fractions of sequence reads affiliated with this genus on the DNA and RNA level were highest in the oxic groundwater of the well H4.1 and lowest in the anoxic groundwater of wells H4.2 and H5.2 (Fig. 9). Since nitrifiers such as *Nitrospira* are known to retain a high ribosome content even if cells are not active (Morgenroth et al., 2000), these results do not necessarily indicate high nitrite oxidation activity at the time point of sampling but point to nitrite ox-




Figure 7. PLFA $\delta^{13}\text{C}$ values averaged in the different wells for June, September and December. The dotted and full lines represent the $\delta^{13}\text{C}$ values of 11Me16:0 and ladderanes associated with nitrite-oxidizing bacteria (e.g. *Nitrospira moscoviensis*) and anammox bacteria, respectively.




Figure 8. Taxonomic classification (phylum level) of the bacterial communities in the groundwater of the eight different wells from the upper aquifer assemblages (HTU) or the lower aquifer (HTL). (a) Bacterial communities based on sequencing of 16S rRNA genes from extracted genomic DNA. (b) Bacterial communities based on sequencing of RNA-derived 16S rRNA amplicons.

idizers forming a large fraction of the microbial community with protein synthesis potential.

4 Discussion

4.1 PLFA distribution

The PCA of PLFAs indicated that the oxic, suboxic and anoxic groundwaters had distinct bacterial communities, with the anoxic groundwater additionally differentiated into two distinct bacterial communities (Fig. 4). Of the environmental variables tested, the variation partitioning showed that




Figure 9. Fractions of sequence reads affiliated with iron oxidizing or iron reducing bacteria, sulfate-reducing bacteria, anammox bacteria, and nitrite oxidizers (*Nitrospira moscoviensis*-related and others) within the bacterial community (a) DNA-based analysis, (b) RNA-based analysis. Only those sequence reads were considered which were unambiguously classified to described taxa on the genus level.

NH_4^+ , O_2 and Fe_i concentration explained 22.0, 19.7 and 13.4 % of the PLFA variations, respectively (Fig. 6), and differentiated between those three bacterial communities. Variation partitioning analyses revealed, along those environmental variables, clusters of covarying PLFAs that may originate from the same functional group of organisms or closely affiliated organisms that react similarly to certain environmental conditions. While the ladderanes are unequivocally attributed to anammox bacteria (Sinninghe Damsté et al., 2002, 2005), the other PLFAs are not exclusive to a phylogenetic or functional microbial group, which complicates their use in understanding the role of microbes in environments. The t -

value biplots of variation partitioning analyses evidenced the PLFAs that significantly correlated with the environmental variables O₂ (Fig. 6a), Fe_t (Fig. 6b) and NH₄⁺ (Fig. 6c), and provided better insights into the functional diversity of active microorganisms in the subdivided groundwaters. Additional supports of the bacterial community structure, assessed by the PLFA patterns, were found in the 16S rRNA-based results. Although a large fraction of the microbial community remains poorly classified and thus precludes the knowledge of the metabolic capacities, high sequence similarity to genera known to be involved in iron oxidation or reduction, sulfate reduction, anammox, and nitrite oxidation allowed an estimation of the fraction of the microbial population potentially involved in these processes. By combining the PLFA-based and sequencing-based approaches, we aimed, here, to compensate for biases introduced by PCR as well as for the limited phylogenetic resolution of PLFA-based analysis. This combined approach resulted in highly supported evidence of some key microbial players and associated biogeochemical processes in physico-chemical distinct aquifer assemblages of the aquifer transect.

4.1.1 PLFA cluster in oxic to suboxic groundwater (wells 5.1, 4.1, 3.1 and 3.2)

A cluster of the covarying 20:4, 20:5, 22:5 and 22:6 PLFAs has to our knowledge heretofore never been observed in groundwater. Associations of those PLFAs have been commonly found in eukaryotes such as microalgae (Volkman et al., 1989; Shinmen et al., 1989; Kennedy et al., 1993; Olsson, 1999; Qi et al., 2004) and protozoans (White, 1988), etc. Protozoa act as detritivores and are expected to be key predators in the microbial loop feeding on different subsets of the bacterial communities and other protozoa (Brad et al., 2008; Akob and Küsel, 2011). Consistently, sessile and free-swimming suspension-feeding flagellates (e.g. *Spumella* sp.), mobile naked amoebae, and ciliates could be detected in this aquifer, with a cultivable protist abundance of up to 8000 cells L⁻¹ (Risse-Buhl et al., 2013). The 18S rRNA gene sequences also revealed high relative fractions of *Spumella*-like Stramenopiles, and sequences affiliated with fungi and metazoan grazers. DNA-based pyro-tag sequencing of fungal internal transcribed spacer sequences revealed a fungi community structure dominated by Ascomycota and Basidiomycota (Nawaz et al., 2016), with the majority of the observed fungal groups being involved in ectomycorrhizal symbioses. In general, the abundance of micro-eukaryotes in pristine groundwater is estimated to be low, because they are limited in nutrients and space and are unable to cope with oxygen limitations (Akob and Küsel, 2011). Consistently, they are commonly found in higher concentrations in OM-rich contaminated groundwaters (Ludvigsen et al., 1997). In pristine aquifers, the origin of those eukaryotic organisms is difficult to determine as they may be autochthonous, allochthonous or both. In the studied sites, the close relation of eukaryotic

PLFA biomarkers with O₂ concentrations (Fig. 6a) suggests their association with recharging groundwater within larger conduits prone to faster water flow. Freshly introduced surface OC and O₂ could fuel the heterotrophic bacterial growth in groundwater. This may subsequently stimulate protists that selectively graze on the prokaryotic biomass and result in the observed relationship between the eukaryotic PLFAs and the O₂ concentration. It is possible to speculate that some surface microorganisms would also survive the transport from surface to the aquifer (Dibbern et al., 2014), especially if the transport is fast. In this case, high cy17:0-to-16:1 ω 7c ratios (Table 3) may show physiological stress due to change of the environmental conditions within the Gram-negative communities (Balkwill et al., 1998).

The 16:1 ω 11c and particularly the 11Me16:0 are major components of *Nitrospira moscoviensis* (Lipski et al., 2001) cell membrane, an obligatory chemo-lithoautotrophic nitrite-oxidizing bacterium (Ehrich et al., 1995). In the oxic groundwater, the occurrence of 16S rRNA gene sequence reads that are closely related to *Nitrospira moscoviensis* (Herrmann et al., 2015) further supports the potential of 11Me16:0 as a biomarker for *Nitrospira moscoviensis* and confirms previous assumptions about an important role of nitrite oxidizers within the autotrophic community of the lower aquifers (Herrmann et al., 2015). The correlation of 11Me16:0 and 16:1 ω 11c with O₂ (Fig. 6a) indicated the occurrence of active nitrification in oxic zones of the aquifers, in agreement with observation of experiments (Satoh et al., 2003). *Nitrospira* use the reverse tricarboxylic acid cycle as the pathway for CO₂ fixation (Lücker et al., 2010), which leads to small ¹³C fractionation (2–6‰) between biomass and CO₂ (van der Meer et al., 1998). The ¹³C-enrichment of 11Me16:0 and 16:1 ω 11c relative to the other PLFAs (up to 18‰ in well H4.1) thus supports a major *Nitrospira* contribution to those PLFAs found in oxic groundwaters (Fig. 7).

4.1.2 PLFA cluster in anoxic, more Fe_t-rich groundwater (wells H4.2 and H4.3)

In groundwater, the concentration of dissolved iron is often inversely related to oxygen, as O₂ in water will chemically oxidize iron that will precipitate as insoluble iron-hydroxides at neutral pH. In the wells H4.2 and H4.3, the increase of the PLFAs 10Me12:0, 12:0, 17:1, 18:1 ω 7c and 18:1 ω 9c with concentrations of Fe_t, Fe₂⁺ and HCO₃⁻ (Figs. 5 and 6b) and the DNA- and RNA-based analyses (Fig. 9) suggested degradation of OM by anaerobic iron-reducing bacteria (IRB). Because many IRB are highly versatile, i.e. they can use different metal substrates as electron acceptors coupled to the oxidation of the OM (Coleman et al., 1993; Lovley et al., 1993; Holmes et al., 2004), specific PLFAs linked to the reduction of iron in anoxic environments are poorly described. The two most studied genera of IRB are *Geobacter* and *Shewanella*, which contain most of those PLFAs (Coleman et al., 1993; Lovley et al., 1993; Hedrick et al., 2009). However, none of

these PLFAs are specific to a certain genus or species. The 17:1 is generally related to anaerobic SRB (Dowling et al., 1986) as *Desulfobulbus* (Parkes and Calder, 1985; Macalady et al., 2000) but also occurs in dissimilatory IRB as *Shewanella* (Coleman et al., 1993). The ability of some sulfate reducers to reduce iron rather than sulfate has long been recognized in groundwater (Coleman et al., 1993).

The 18:1 ω 9c is common and abundant in fungi (Frostegård and Bååth, 1996; Kaiser et al., 2010), but may also occur in micro-algae (Arts et al., 2001) and Gram-negative bacteria (Kandeler, 2007). The 18:1 ω 9c, 18:2 ω 6,9 and 18:3 ω 6 are typically used as fungi biomarkers in soil (Frostegård and Bååth, 1996; Ruzicka et al., 2000; Bååth and Anderson, 2003) and more particularly for saprotrophs (Etingoff, 2014). The correlations between 18:1 ω 9c, 18:2 ω 6,9 and 18:3 ω 6 suggested a major fungal origin of those PLFAs in the studied groundwaters. In soil, fungi are well known for their role in accelerating weathering and solubilisation of iron-containing minerals by excreting organic acids including phenolic compounds, siderophores and protons (Arrieta and Grez, 1971; Landeweert et al., 2001). By forming dense hyphae tunnelling in soils and shallow rocks, fungi mediate and facilitate iron transport in plants and increase iron availability in the environment (van Schöll et al., 2008). Therefore, several studies have linked enhanced rates of iron cycling to the presence of fungal biomass (Gadd, 2010). Moreover, in a recent study, it is been shown that rhizoplanes are important root channels for preferential vertical transport from soil to seepage area of soil colloids including microbes (Dibbern et al., 2014). Limitation of ferric iron may restrain the growth and activity of IRB in the subsurface (O'Neil et al., 2008). In the groundwater of wells H4.2 and H4.3, the close relation of 18:1 ω 9c and 18:2 ω 6,9 with Fe₂ concentration (Fig. 6b) suggested that fungal biomass may, by mediating and facilitating the transport of different types of organic and/or inorganic particles and colloids, play a key role in iron bioavailability and thus sustain IRB growth and activity.

4.1.3 PLFA cluster in anoxic, more NH₄⁺-rich groundwater (wells H5.2, H5.3 and H3.2)

To our knowledge, this is the first time phospholipid [3]-ladderane and [5]-ladderane, which attest the presence of viable or recently degraded anammox bacteria (Jaeschke et al., 2009), have been identified in groundwater. The occurrence of anammox bacteria in those groundwaters is consistent with the DNA- and RNA-based analyses (Fig. 9) and coincided with higher concentrations of ammonium (Fig. 2). The difference between DIC and ladderane $\delta^{13}\text{C}$ values of 55‰ was within the range previously reported for anammox in the Black Sea (Schouten et al., 2004), further suggesting that autotrophic carbon fixation pathways within the diverse group of anaerobic ammonium-oxidizing bacteria are similar. In the sub-oxic (well H3.2) and anoxic groundwaters

(well H5.2 and H5.3), the increasing concentration of ladderane lipids derived from anammox bacteria with decreasing O₂ concentration (Fig. 6a) agrees well with the reported high sensitivity of the anammox process to O₂ (Kalvelage et al., 2011). Denitrification and anammox are the dominant nitrogen loss pathways in aquatic ecosystems (Burgin and Hamilton, 2008; Koeve and Kähler, 2010). The occurrence of lipids derived from anammox bacteria in those groundwaters indicates that the anammox process may be critically important in the nitrogen loss from this part of the aquifer assemblage.

High amounts of 10Me16:0 are typically found in SRB (Dowling et al., 1986; Vainshtein et al., 1992; Kohring et al., 1994) but also occur in anammox bacteria (Sinninghe Damsté et al., 2002). Anammox bacteria strongly fractionate against ¹³C, producing ladderane lipids which are ¹³C-depleted by 47‰ compared to the inorganic carbon source (Schouten et al., 2004). Relative to ladderanes, SRB-derived lipids are expected to be ¹³C-enriched as cultured SRB under heterotrophic and autotrophic growth fractionated against ¹³C by up to 27‰ (Londry et al., 2004). Therefore, the ¹³C-enrichment of 10Me16:0 (up to 19‰) relative to the ladderanes supported major SRB contribution to the 10Me16:0 found in these groundwaters. The i13:0, i15:0 and i17:1 are typically, as with 10Me16:0, associated with SRB (Edlund et al., 1985; Kohring et al., 1994). In those groundwaters, similar $\delta^{13}\text{C}$ values, in the −44 to −56‰ range, also supported a common SRB origin of those PLFAs.

Variation partitioning analyses showed that the concentrations of [3]-ladderane, [5]-ladderane, 10Me16:0 and i17:1 correlated with NH₄⁺ concentration (Fig. 6c). Many studies in other aquatic environments showed that the relative importance of the anammox process is directly related to the availability of NH₄⁺ (Dalsgaard and Thamdrup, 2002; Kuypers et al., 2003). Commonly, the breakdown of OM via ammonification or dissimilatory nitrate reduction to ammonia is presumed to be the major source of NH₄⁺ for anammox (Kartal et al., 2007). However, the recent discovery of comammox organisms capable of complete nitrification underlines the complexity of the nitrogen cycle and the variability of ammonium sources for anammox (van Kessel et al., 2015). The availability of OM is known as an additional important factor influencing the anammox process. Higher anammox activity has been observed in OM-poor environments and interpreted as a decrease in competition for NO₂⁻ by heterotrophic denitrifiers (Hu et al., 2011). Consistently, high anammox activity was observed in redox zones associated with sulfate reduction or sulfur oxidation (Mills et al., 2006; Canfield et al., 2010; Prokopenko et al., 2013; Wenk et al., 2013). In the groundwater of the wells H5.2 and H5.3, the occurrence of anammox bacteria and SRB thus supported low groundwater–surface interactions which likely threatened the availability of generically favourable electron acceptors and labile OM.

5 Conclusion

In this study, we used constrained ordination to show environmental variables that significantly correlated with PLFA relative abundances in groundwater of distinct carbonate-rock aquifer assemblages. This technique shows that the active subsurface microbial communities were mainly affected by variations in dissolved O_2 , Fe_I and NH_4^+ concentrations. Variation partitioning identified PLFA-based microbial functional groups that were directly supported by results of DNA- and RNA-based amplicon sequencing targeting bacterial 16S rRNA genes. Higher O_2 concentration resulted in increased eukaryotic biomass and higher relative fractions of nitrite oxidizing bacteria (e.g. *Nitrospira moscoviensis*) but impeded anammox bacteria, sulfate-reducing bacteria and iron-reducing bacteria. In anoxic groundwater, concomitant increase of total iron (Fe_I), HCO_3^- and PLFAs abundant in Gram-negative bacteria and fungi suggested the occurrence of active dissimilatory iron reduction and a possible role of fungi in mediating iron solubilisation and transport in those aquifer assemblages. The relative abundance of PLFA derived from anammox bacteria correlated with NH_4^+ concentrations, showing the dependence of the anammox process on the availability of NH_4^+ . Our study shows that different relationships among the microbial community structures, estimated based on both the PLFA patterns and 16S rRNA gene-targeted next generation sequencing, reflected changes in the physiological strategies of microorganisms related to a decrease in substrate bioavailability and redox potential of the groundwater.

Data availability. All data are accessible in the Supplement.

The Supplement related to this article is available online at doi:10.5194/bg-14-2697-2017-supplement.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. The work has been funded by the Deutsche Forschungsgemeinschaft (DFG) CRC 1076 “AquaDiva”. Field work permits were issued by the responsible state environmental offices of Thuringia. We thank Heiko Minkmar and Falko Gutmann for sampling and on-site measurements, Christine Hess for scientific coordination and the Hainich National Park administration for help and support. Patricia Geesink is acknowledged for assistance in DNA and RNA extractions. Illumina MiSeq amplicon sequencing was financially supported by the German Center for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, funded by the Deutsche Forschungsgemeinschaft (FZT 118). The authors would like to thank two anonymous referees and the Associate Editor Marcel van der Meer for their helpful comments in Biogeosciences Discussions.

Edited by: M. van der Meer

Reviewed by: two anonymous referees

References

- Akob, D. M. and Küsel, K.: Where microorganisms meet rocks in the Earth's Critical Zone, *Biogeosciences*, 8, 3531–3543, doi:10.5194/bg-8-3531-2011, 2011.
- American Public Health Association: Standard methods for the examination of water and wastewater: selected analytical methods approved and cited by the United States Environmental Protection Agency, 1981.
- Arrieta, L. and Grez, R.: Solubilization of iron-containing minerals by soil microorganisms, *Appl. Microbiol.*, 22, 487–490, 1971.
- Arts, M. T., Ackman, R. G., and Holub, B. J.: “Essential fatty acids” in aquatic ecosystems: a crucial link between diet and human health and evolution, *Can. J. Fish. Aquat. Sci.*, 58, 122–137, doi:10.1139/f00-224, 2001.
- Bååth, E. and Anderson, T.-H.: Comparison of soil fungal/bacterial ratios in a pH gradient using physiological and PLFA-based techniques, *Soil Biol. Biochem.*, 35, 955–963, doi:10.1016/S0038-0717(03)00154-8, 2003.
- Bååth, E., Frostegård, Å., Pennanen, T., and Fritze, H.: Microbial community structure and pH response in relation to soil organic matter quality in wood-ash fertilized, clear-cut or burned coniferous forest soils, *Soil Biol. Biochem.*, 27, 229–240, doi:10.1016/0038-0717(94)00140-V, 1995.
- Balkwill, D. L., Murphy, E. M., Fair, D. M., Ringelberg, D. B., and White, D. C.: Microbial communities in high and low recharge environments: implications for microbial transport in the vadose zone, *Microbiol. Ecol.*, 35, 156–171, doi:10.1007/s002489900070, 1998.
- Benner, S. G., Smart, E. W., and Moore, J. N.: Metal behavior during surface-groundwater interaction, Silver Bow Creek, Montana, *Environ. Sci. Technol.*, 29, 1789–1795, doi:10.1021/es00007a015, 1995.
- Blair, N., Leu, A., Muñoz, E., Olsen, J., Kwong, E., and Des Marais, D.: Carbon isotopic fractionation in heterotrophic microbial metabolism, *Appl. Environ. Microbiol.*, 50, 996–1001, 1985.
- Blazewicz, S. J., Barnard, R. L., Daly, R. A., and Firestone, M. K.: Evaluating rRNA as an indicator of microbial activity in environmental communities: limitations and uses, *ISME J.*, 7, 2061–2068, doi:10.1038/ismej.2013.102, 2013.
- Bligh, E. G. and Dyer, W. J.: A rapid method of total lipid extraction and purification, *Can. J. Biochem. Physiol.*, 37, 911–917, doi:10.1139/o59-099, 1959.
- Bossio, D. A. and Scow, K. M.: Impacts of carbon and flooding on soil microbial communities: phospholipid fatty acid profiles and substrate utilization patterns, *Microbiol. Ecol.*, 35, 265–278, doi:10.1007/s002489900082, 1998.
- Brad, T., Braster, M., van Breukelen, B. M., van Straalen, N. M., and Röling, W. F. M.: Eukaryotic diversity in an anaerobic aquifer polluted with landfill leachate, *Appl. Environ. Microbiol.*, 74, 3959–3968, doi:10.1128/AEM.02820-07, 2008.
- Burgin, A. and Hamilton, S. K.: NO_3^- -driven SO_4^{2-} production in freshwater ecosystems: implications for N and S cycling, *Ecosystems*, 11, 908–922, doi:10.1007/s10021-008-9169-5, 2008.

- Burgin, A. J. and Hamilton, S. K.: Have we overemphasized the role of denitrification in aquatic ecosystems? A review of nitrate removal pathways, *Front. Ecol. Environ.*, 5, 89–96, doi:10.1890/1540-9295(2007)5[89:HWOTRO]2.0.CO;2, 2007.
- Canfield, D. E., Stewart, F. J., Thamdrup, B., Brabandere, L. D., Dalsgaard, T., Delong, E. F., Revsbech, N. P., and Ulloa, O.: A cryptic sulfur cycle in oxygen-minimum-zone waters of the Chilean coast, *Science*, 330, 1375–1378, doi:10.1126/science.1196889, 2010.
- Chapelle, F. H.: *Ground-water microbiology and geochemistry*, John Wiley & Sons, 2001.
- Chapelle, F. H. and Lovley, D. R.: Competitive exclusion of sulfate reduction by Fe(III)-reducing bacteria: A mechanism for producing discrete zones of high-iron ground water, *Groundwater*, 30, 29–36, doi:10.1111/j.1745-6584.1992.tb00808.x, 1992.
- Coleman, M. L., Hedrick, D. B., Lovley, D. R., White, D. C., and Pye, K.: Reduction of Fe(III) in sediments by sulphate-reducing bacteria, *Nature*, 361, 436–438, doi:10.1038/361436a0, 1993.
- Dalsgaard, T. and Thamdrup, B.: Factors controlling anaerobic ammonium oxidation with nitrite in marine sediments, *Appl. Environ. Microbiol.*, 68, 3802–3808, doi:10.1128/AEM.68.8.3802-3808.2002, 2002.
- Dalsgaard, T., Canfield, D. E., Petersen, J., Thamdrup, B., and Acuña-González, J.: N₂ production by the anammox reaction in the anoxic water column of Golfo Dulce, Costa Rica, *Nature*, 422, 606–608, doi:10.1038/nature01526, 2003.
- Dibbern, D., Schmalwasser, A., Lueders, T., and Totsche, K. U.: Selective transport of plant root-associated bacterial populations in agricultural soils upon snowmelt, *Soil Biol. Biochem.*, 69, 187–196, doi:10.1016/j.soilbio.2013.10.040, 2014.
- DIN 38409-7: Deutsche Einheitsverfahren zur Wasser-, Abwasser- und Schlammuntersuchung – Summarische Wirkungs- und Stoffkenngrößen (Gruppe H) – Teil 7: Bestimmung der Säure- und Basekapazität (H7), in: Normung, D.D.I.f., Beuth Verlag GmbH, Berlin, 2005.
- DIN EN 1484: Wasseranalytik – Anleitungen zur Bestimmung des gesamten organischen Kohlenstoffs (TOC) und des gelösten organischen Kohlenstoffs (DOC), in: Normung, D.D.I.f., Beuth Verlag GmbH, Berlin, 1997.
- DIN EN ISO 10304-1: Wasserbeschaffenheit – Bestimmung von gelösten Anionen mittels Flüssigkeits-Ionenchromatographie – Teil 1: Bestimmung von Bromid, Chlorid, Fluorid, Nitrat, Nitrit, Phosphat und Sulfat, in: Normung, D.D.I.f., Beuth Verlag GmbH, Berlin, p. 24, 2009.
- DIN EN ISO 11885: Wasserbeschaffenheit – Bestimmung von ausgewählten Elementen durch induktiv gekoppelte Plasma-Atom-Emissionsspektrometrie (ICP-OES), in: Normung, D.D.I.f., Beuth Verlag GmbH, Berlin, 2009.
- Dowling, N. J. E., Widdel, F., and White, D. C.: Phospholipid ester-linked fatty acid biomarkers of acetate-oxidizing sulphate-reducers and other sulphide-forming bacteria, *Microbiol.*, 132, 1815–1825, doi:10.1099/00221287-132-7-1815, 1986.
- Edlund, A., Nichols, P. D., Roffey, R., and White, D. C.: Extractable and lipopolysaccharide fatty acid and hydroxy acid profiles from *Desulfovibrio* species, *J. Lipid Res.*, 26, 982–988, 1985.
- Edwards, K. J., Becker, K., and Colwell, F.: The deep, dark energy biosphere: intraterrestrial life on earth, *Annu. Rev. Earth Pl. Sc.*, 40, 551–568, doi:10.1146/annurev-earth-042711-105500, 2012.
- Ehrich, S., Behrens, D., Lebedeva, E., Ludwig, W., and Bock, E.: A new obligately chemolithoautotrophic, nitrite-oxidizing bacterium, *Nitrospira moscoviensis* sp. nov. and its phylogenetic relationship, *Arch. Microbiol.*, 164, 16–23, doi:10.1007/BF02568729, 1995.
- Emerson, J. B., Thomas, B. C., Alvarez, W., and Banfield, J. F.: Metagenomic analysis of a high carbon dioxide subsurface microbial community populated by chemolithoautotrophs and bacteria and archaea from candidate phyla, *Environ. Microbiol.*, 18, 1462–2920, doi:10.1111/1462-2920.12817, 2015.
- Etingoff, K.: *Organic agricultural practices: Alternatives to conventional agricultural systems*, CRC Press, Toronto, Canada, 2014.
- Filion, G., Laflamme, C., Turgeon, N., Ho, J., and Duchaine, C.: Permeabilization and hybridization protocols for rapid detection of *Bacillus* spores using fluorescence in situ hybridization, *J. Microbiol. Meth.*, 77, 29–36, doi:10.1016/j.mimet.2008.12.009, 2009.
- Frostegård, A. and Bååth, E.: The use of phospholipid fatty acid analysis to estimate bacterial and fungal biomass in soil, *Biol. Fert. Soils*, 22, 59–65, doi:10.1007/BF00384433, 1996.
- Frostegård, Å., Tunlid, A., and Bååth, E.: Use and misuse of PLFA measurements in soils, *Soil Biol. Biochem.*, 43, 1621–1625, doi:10.1016/j.soilbio.2010.11.021, 2011.
- Gadd, G. M.: Metals, minerals and microbes: geomicrobiology and bioremediation, *Microbiology*, 156, 609–643, doi:10.1099/mic.0.037143-0, 2010.
- Green, C. T. and Scow, K. M.: Analysis of phospholipid fatty acids (PLFA) to characterize microbial communities in aquifers, *Hydrogeol. J.*, 8, 126–141, doi:10.1007/s100400050013, 2000.
- Hamman, S. T., Burke, I. C., and Stromberger, M. E.: Relationships between microbial community structure and soil environmental conditions in a recently burned system, *Soil Biol. Biochem.*, 39, 1703–1711, doi:10.1016/j.soilbio.2007.01.018, 2007.
- Harwood, D. L. and Russell, N. J.: *Lipids in plants and microbes*, George Allen and Unwin, London, 1984.
- Hedrick, D. B., Peacock, A. D., Lovley, D. R., Woodard, T. L., Nevin, K. P., Long, P. E., and White, D. C.: Polar lipid fatty acids, LPS-hydroxy fatty acids, and respiratory quinones of three Geobacter strains, and variation with electron acceptor, *J. Ind. Microbiol. Biot.*, 36, 205–209, doi:10.1007/s10295-008-0486-7, 2009.
- Heikkinen, R. K., Luoto, M., Virkkala, R., and Rainio, K.: Effects of habitat cover, landscape structure and spatial variables on the abundance of birds in an agricultural–forest mosaic, *J. Appl. Ecol.*, 41, 824–835, doi:10.1111/j.0021-8901.2004.00938.x, 2004.
- Heinzelmann, S. M., Bale, N. J., Hopmans, E. C., Sinninghe Damsté, J. S., Schouten, S., and van der Meer, M. T. J.: Critical Assessment of glyco- and phospholipid separation by using silica chromatography, *Appl. Environ. Microbiol.*, 80, 360–365, doi:10.1128/AEM.02817-13, 2014.
- Herlemann, D. P., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J. J., and Andersson, A. F.: Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea, *ISME J.*, 5, 1571–1579, doi:10.1038/ismej.2011.41, 2011.
- Herrmann, M., Hädrich, A., and Küsel, K.: Predominance of thaumarchaeal ammonia oxidizer abundance and transcriptional activity in an acidic fen, *Environ. Microbiol.*, 14, 3013–3025, doi:10.1111/j.1462-2920.2012.02882.x, 2012.

- Herrmann, M., Rusznyák, A., Akob, D. M., Schulze, I., Opitz, S., Totsche, K. U., and Küsel, K.: Large Fractions of CO₂-fixing microorganisms in pristine limestone aquifers appear to be involved in the oxidation of reduced sulfur and nitrogen compounds, *Appl. Environ. Microbiol.*, 81, 2384–2394, doi:10.1128/AEM.03269-14, 2015.
- Holmes, D. E., Bond, D. R., and Lovley, D. R.: Electron Transfer by *Desulfobulbus propionicus* to Fe(III) and Graphite Electrodes, *Appl. Environ. Microbiol.*, 70, 1234–1237, doi:10.1128/AEM.70.2.1234-1237.2004, 2004.
- Hu, B. L., Shen, L. D., Xu, X. Y., and Zheng, P.: Anaerobic ammonium oxidation (anammox) in different natural ecosystems, *Biochem. Soc. Trans.*, 39, 1811–1816, doi:10.1042/BST20110711, 2011.
- Humbert, S., Tarnawski, S., Fromin, N., Mallet, M.-P., Aragno, M., and Zopfi, J.: Molecular detection of anammox bacteria in terrestrial ecosystems: distribution and diversity, *ISME J.*, 4, 450–454, doi:10.1038/ismej.2009.125, 2009.
- Jaeschke, A., Rooks, C., Trimmer, M., Nicholls, J. C., Hopmans, E. C., Schouten, S., and Sinninghe Damsté, J. S.: Comparison of ladderane phospholipid and core lipids as indicators for anaerobic ammonium oxidation (anammox) in marine sediments, *Geochim. Cosmochim. Ac.*, 73, 2077–2088, doi:10.1016/j.gca.2009.01.013, 2009.
- Kaiser, C., Frank, A., Wild, B., Koranda, M., and Richter, A.: Negligible contribution from roots to soil-borne phospholipid fatty acid fungal biomarkers 18:2 ω 6,9 and 18:1 ω 9, *Soil Biol. Biochem.*, 42, 1650–1652, doi:10.1016/j.soilbio.2010.05.019, 2010.
- Kalbus, E., Reinstorf, F., and Schirmer, M.: Measuring methods for groundwater – surface water interactions: a review, *Hydrol. Earth Syst. Sci.*, 10, 873–887, doi:10.5194/hess-10-873-2006, 2006.
- Kalvelage, T., Jensen, M. M., Contreras, S., Revsbech, N. P., Lam, P., Günter, M., LaRoche, J., Lavik, G., and Kuypers, M. M.: Oxygen sensitivity of anammox and coupled N-cycle processes in oxygen minimum zones, *PLoS ONE*, 6, e29299, doi:10.1371/journal.pone.0029299, 2011.
- Kandeler, E.: Physiological and biochemical methods for studying soil biota and their function, in: *Soil Microbiology, Ecology, and Biochemistry*, edited by: Eldor, A. P., Academic Press Elsevier, 53–83, 2007.
- Kartal, B., Kuypers, M. M. M., Lavik, G., Schalk, J., Op den Camp, H. J. M., Jetten, M. S. M., and Strous, M.: Anammox bacteria disguised as denitrifiers: nitrate reduction to dinitrogen gas via nitrite and ammonium, *Environ. Microbiol.*, 9, 635–642, doi:10.1111/j.1462-2920.2006.01183.x, 2007.
- Kaur, A., Chaudhary, A., Kaur, A., Choudhary, R., and Kaushik, R.: Phospholipid fatty acid-A bioindicator of environment monitoring and assessment in soil ecosystem, *Curr. Sci.*, 89, 1103–1112, 2005.
- Kennedy, M., Reader, S., and Davies, R.: Fatty acid production characteristics of fungi with particular emphasis on gamma linolenic acid production, *Biotechnol. and Bioeng.*, 42, 625–634, doi:10.1002/bit.260420511, 1993.
- Kerger, B. D., Nichols, P. D., Antworth, C. P., Sand, W., Bock, E., Cox, J. C., Langworthy, T. A., and White, D. C.: Signature fatty acids in the polar lipids of acid-producing *Thiobacillus* spp.: methoxy, cyclopropyl, alpha-hydroxycyclopropyl and branched and normal monoenoic fatty acids, *FEMS Microbiol. Ecol.*, 38, 67–78, doi:10.1016/0378-1097(86)90144-8, 1986.
- Kerger, B. D., Nichols, P. D., Sand, W., Bock, E., and White, D. C.: Association of acid-producing thiobacilli with degradation of concrete: analysis by “signature” fatty acids from the polar lipids and lipopolysaccharide, *J. Ind. Microbiol.*, 2, 63–70, doi:10.1007/BF01569504, 1987.
- Koeve, W. and Kähler, P.: Heterotrophic denitrification vs. autotrophic anammox – quantifying collateral effects on the oceanic carbon cycle, *Biogeosciences*, 7, 2327–2337, doi:10.5194/bg-7-2327-2010, 2010.
- Kohlhepp, B., Lehmann, R., Seeber, P., Küsel, K., Trumbore, S. E., and Totsche, K. U.: Pedological and hydrogeological setting and subsurface flow structure of the carbonate-rock CZE Hainich in western Thuringia, Germany, *Hydrol. Earth Syst. Sci. Discuss.*, doi:10.5194/hess-2016-374, in review, 2016.
- Kohring, L. L., Ringelberg, D. B., Devereux, R., Stahl, D. A., Mittelman, M. W., and White, D. C.: Comparison of phylogenetic relationships based on phospholipid fatty acid profiles and ribosomal RNA sequence similarities among dissimilatory sulfate-reducing bacteria, *FEMS Microbiol. Lett.*, 119, 303–308, doi:10.1111/j.1574-6968.1994.tb06905.x, 1994.
- Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D.: Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence data on the MiSeq Illumina sequencing platform, *Appl. Environ. Microbiol.*, 79, 5112–5120, doi:10.1128/AEM.01043-13, 2013.
- Kramer, C. and Gleixner, G.: Variable use of plant- and soil-derived carbon by microorganisms in agricultural soils, *Soil Biol. Biochem.*, 38, 3267–3278, doi:10.1016/j.soilbio.2006.04.006, 2006.
- Küsel, K., Totsche, K. U., Trumbore, S. E., Lehmann, R., Steinhäuser, C., and Herrmann, M.: How deep can surface signals be traced in the critical zone? Merging biodiversity with biogeochemistry research in a central German Muschelkalk landscape, *Front. Earth Sci.*, 4, 32, doi:10.3389/feart.2016.00032, 2016.
- Kuypers, M. M. M., Sliemers, A. O., Lavik, G., Schmid, M., Jorgensen, B. B., Kuenen, J. G., Sinninghe Damsté, J. S., Strous, M., and Jetten, M. S. M.: Anaerobic ammonium oxidation by anammox bacteria in the Black Sea, *Nature*, 422, 608–611, doi:10.1038/nature01472, 2003.
- Landeweert, R., Hoffland, E., Finlay, R. D., Kuypers, T. W., and van Breemen, N.: Linking plants to rocks: ectomycorrhizal fungi mobilize nutrients from minerals, *Trends Ecol. Evol.*, 16, 248–254, doi:10.1016/S0169-5347(01)02122-X, 2001.
- Landmeyer, J. E., Vroblesky, D. A., and Chapelle, F. H.: Stable carbon isotope evidence of biodegradation zonation in a shallow jet-fuel contaminated aquifer, *Environ. Sci. Technol.*, 30, 1120–1128, doi:10.1021/es950325t, 1996.
- Lipski, A., Spieck, E., Makolla, A., and Altendorf, K.: Fatty acid profiles of nitrite-oxidizing bacteria reflect their phylogenetic heterogeneity, *Syst. Appl. Microbiol.*, 24, 377–384, doi:10.1078/0723-2020-00049, 2001.
- Londry, K. L., Jahnke, L. L., and Des Marais, D. J.: Stable carbon isotope ratios of lipid biomarkers of sulfate-reducing bacteria, *Appl. Environ. Microbiol.*, 70, 745–751, doi:10.1128/AEM.70.2.745-751.2004, 2004.
- Lovley, D. R., Giovannoni, S. J., White, D. C., Champine, J. E., Phillips, E., Gorby, Y. A., and Goodwin, S.: *Geobacter metal-*

- lireducens* gen. nov. sp. nov., a microorganism capable of coupling the complete oxidation of organic compounds to the reduction of iron and other metals, *Arch. Microbiol.*, 159, 336–344, doi:10.1007/BF00290916, 1993.
- Lücker, S., Wagner, M., Maixner, F., Pelletier, E., Koch, H., Vacherie, B., Rattei, T., Sinninghe Damsté, J. S., Spieck, E., Le Paslier, D., and Daims, H.: A *Nitrospira* metagenome illuminates the physiology and evolution of globally important nitrite-oxidizing bacteria, *P. Natl. Acad. Sci. USA.*, 107, 13479–13484, doi:10.1073/pnas.1003860107, 2010.
- Ludvigsen, L., Albrechtsen, H. J., Holst, H., and Christensen, T. H.: Correlating phospholipid fatty acids (PLFA) in a landfill leachate polluted aquifer with biogeochemical factors by multivariate statistical methods, *FEMS Microbiol. Rev.*, 20, 447–460, doi:10.1111/j.1574-6976.1997.tb00329.x, 1997.
- Macalady, J., Mack, E., Nelson, D., and Scow, K.: Sediment microbial community structure and mercury methylation in mercury-polluted Clear Lake, California, *Appl. Environ. Microbiol.*, 66, 1479–1488, doi:10.1128/AEM.66.4.1479-1488.2000, 2000.
- Mills, C. T., Dias, R. F., Graham, D., and Mandernack, K. W.: Determination of phospholipid fatty acid structures and stable carbon isotope compositions of deep-sea sediments of the Northwest Pacific, ODP site 1179, *Mar. Chem.*, 98, 197–209, doi:10.1016/j.marchem.2005.10.001, 2006.
- Morgenroth, E., Obermayer, A., Arnold, E., Brühl, A., Wagner, M., and Wilderer, P.: Effect of long-term idle periods on the performance of sequencing batch reactors, *Water Sci. Technol.*, 41, 105–113, 2000.
- Myers, R. T., Zak, D. R., White, D. C., and Peacock, A.: Landscape-level patterns of microbial community composition and substrate use in upland forest ecosystems, *Soil Sci. Soc. Am. J.*, 65, 359–367, doi:10.2136/sssaj2001.652359x, 2001.
- Nawaz, A., Purahong, W., Lehmann, R., Herrmann, M., Küsel, K., Totsche, K. U., Buscot, F., and Wubet, T.: Superimposed pristine limestone aquifers with marked hydrochemical differences exhibit distinct fungal communities, *Front. Microbiol.*, 7, 666, doi:10.3389/fmicb.2016.00666, 2016.
- Olsson, P.: Signature fatty acids provide tools for determination of the distribution and interactions of mycorrhizal fungi in soil, *FEMS Microbiol. Ecol.*, 29, 303–310, doi:10.1016/S0168-6496(99)00021-5, 1999.
- O’Neil, R. A., Holmes, D. E., Coppi, M. V., Adams, L. A., Larrahondo, M. J., Ward, J. E., Nevin, K. P., Woodard, T. L., Vriomis, H. A., N’Guessan, A. L., and Lovley, D. R.: Gene transcript analysis of assimilatory iron limitation in *Geobacteraceae* during groundwater bioremediation, *Environ. Microbiol.*, 10, 1218–1230, doi:10.1111/j.1462-2920.2007.01537.x, 2008.
- Parkes, R. J. and Graham Calder, A.: The cellular fatty acids of three strains of *Desulfobulbus*, a propionate-utilising sulphate-reducing bacterium, *FEMS Microbiol. Lett.*, 31, 361–363, doi:10.1016/0378-1097(85)90032-1, 1985.
- Pinto, A. J., Marcus, D. N., Ijaz, U. Z., Bautista-de Iose Santos, Q. M., Dick, G. J., and Raskin, L.: Metagenomic evidence for the presence of Comammox *Nitrospira*-like bacteria in a drinking water system, *mSphere*, 1, doi:10.1128/mSphere.00054-15, 2016.
- Prokopenko, M. G., Hirst, M. B., De Brabandere, L., Lawrence, D. J. P., Berelson, W. M., Granger, J., Chang, B. X., Dawson, S., Crane III, E. J., Chong, L., Thamdru, B., Townsend-Small, A., and Sigman, D. M.: Nitrogen losses in anoxic marine sediments driven by *Thioploca*-anammox bacterial consortia, *Nature*, 500, 194–198, doi:10.1038/nature12365, 2013.
- Qi, B., Fraser, T., Mugford, S., Dobson, G., Sayanova, O., Butler, J., Napier, J. A., Stobart, A. K., and Lazarus, C. M.: Production of very long chain polyunsaturated omega-3 and omega-6 fatty acids in plants, *Nat. Biotech.*, 22, 739–745, doi:10.1038/nbt972, 2004.
- Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T., Yarza, P., Peplies, J., and Glöckner, F. O.: The SILVA ribosomal RNA gene database project: improved data processing and web-based tools, *Nucleic Acids Res.*, 41, D590–D596, doi:10.1093/nar/gks1219, 2013.
- Reardon, J., Foreman, J., and Searcy, R.: New reactants for the colorimetric determination of ammonia, *Clin. Chim. Acta*, 14, 203–205, doi:10.1016/0009-8981(66)90120-3, 1966.
- Risse-Buhl, U., Herrmann, M., Lange, P., Akob, D. M., Pizani, N., Schönborn, W., Totsche, K. U., and Küsel, K.: Phagotrophic protist diversity in the groundwater of a karstified aquifer—morphological and molecular analysis, *J. Eukaryot. Microbiol.*, 60, 467–479, doi:10.1111/jeu.12054, 2013.
- Roth, V.-N., Dittmar, T., Gaupp, R., and Gleixner, G.: The molecular composition of dissolved organic matter in forest soils as a function of pH and temperature, *PLoS ONE*, 10, e0119188, doi:10.1371/journal.pone.0119188, 2015.
- Ruzicka, S., Edgerton, D., Norman, M., and Hill, T.: The utility of ergosterol as a bioindicator of fungi in temperate soils, *Soil Biol. Biochem.*, 32, 989–1005, doi:10.1016/S0038-0717(00)00009-2, 2000.
- Satoh, H., Nakamura, Y., Ono, H., and Okabe, S.: Effect of oxygen concentration on nitrification and denitrification in single activated sludge flocs, *Biotechnol. Bioeng.*, 83, 604–607, doi:10.1002/bit.10717, 2003.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., Lesniewski, R. A., Oakley, B. B., Parks, D. H., Robinson, C. J., Sahl, J. W., Stres, B., Thallinger, G. G., Van Horn, D. J., and Weber, C. F.: Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities, *Appl. Environ. Microbiol.*, 75, 7537–7541, doi:10.1128/AEM.01541-09, 2009.
- Schneider, T., Keiblinger, K. M., Schmid, E., Sterflinger-Gleixner, K., Ellersdorfer, G., Roschitzki, B., Richter, A., Eberl, L., Zechmeister-Boltenstern, S., and Riedel, K.: Who is who in litter decomposition and quest; metaproteomics reveals major microbial players and their biogeochemical functions, *ISME J.*, 6, 1749–1762, doi:10.1038/ismej.2012.1, 2012.
- Schouten, S., Strous, M., Kuypers, M. M. M., Rijpstra, W. I. C., Baas, M., Schubert, C. J., Jetten, M. S. M., and Sinninghe Damsté, J. S.: Stable carbon isotopic fractionations associated with inorganic carbon fixation by anaerobic ammonium-oxidizing bacteria, *Appl. Environ. Microbiol.*, 70, 3785–3788, doi:10.1128/AEM.70.6.3785-3788.2004, 2004.
- Seifert, A.-G., Trumbore, S., Xu, X., Zhang, D., and Gleixner, G.: Variable effects of plant colonization on black slate uptake into microbial PLFAs, *Geochim. Cosmochim. Ac.*, 106, 391–403, doi:10.1016/j.gca.2012.12.011, 2013.
- Shinmen, Y., Shimizu, S., Akimoto, K., Kawashima, H., and Yamada, H.: Production of arachidonic acid by

- Mortierella fungi, *Appl. Microbiol. Biotechnol.*, 31, 11–16, doi:10.1007/BF02932833, 1989.
- Sinninghe Damsté, J. S., Strous, M., Rijpstra, W. I. C., Hopmans, E. C., Geenevasen, J. A. J., van Duin, A. C. T., van Niftrik, L. A., and Jetten, M. S. M.: Linearly concatenated cyclobutane lipids form a dense bacterial membrane, *Nature*, 419, 708–712, doi:10.1038/nature01128, 2002.
- Sinninghe Damsté, J. S., Rijpstra, W. I. C., Geenevasen, J. A. J., Strous, M., and Jetten, M. S. M.: Structural identification of ladderane and other membrane lipids of planctomycetes capable of anaerobic ammonium oxidation (anammox), *FEBS J.*, 272, 4270–4283, doi:10.1111/j.1742-4658.2005.04842.x, 2005.
- Šmilauer, P. and Lepš, J.: Multivariate analysis of ecological data using CANOCO 5, Cambridge University Press, 2014.
- Stevens, T. O. and McKinley, J. P.: Lithoautotrophic microbial ecosystems in deep basalt aquifers, *Science*, 270, 450, doi:10.1126/science.270.5235.450, 1995.
- Sukenik, A., Kaplan-Levy, R. N., Welch, J. M., and Post, A. F.: Massive multiplication of genome and ribosomes in dormant cells (akinetes) of *Aphanizomenon ovalisporum* (Cyanobacteria), *ISME J.*, 6, 670–679, doi:10.1038/ismej.2011.2012.
- Teece, M. A., Fogel, M. L., Dollhopf, M. E., and Neelson, K. H.: Isotopic fractionation associated with biosynthesis of fatty acids by a marine bacterium under oxic and anoxic conditions, *Org. Geochem.*, 30, 1571–1579, doi:10.1016/S0146-6380(99)00108-4, 1999.
- Torsvik, V. and Øvreås, L.: Microbial diversity and function in soil: from genes to ecosystems, *Curr. Opin. Microbiol.*, 5, 240–245, doi:10.1016/S1369-5274(02)00324-7, 2002.
- Vainshtein, M., Hippe, H., and Kroppenstedt, R. M.: Cellular fatty acid composition of *Desulfovibrio species* and its use in classification of sulfate-reducing bacteria, *Syst. Appl. Microbiol.*, 15, 554–566, doi:10.1016/S0723-2020(11)80115-3, 1992.
- van der Meer, M. T. J., Schouten, S., and Sinninghe Damsté, J. S.: The effect of the reversed tricarboxylic acid cycle on the ¹³C contents of bacterial lipids, *Org. Geochem.*, 28, 527–533, doi:10.1016/S0146-6380(98)00024-2, 1998.
- van der Meer, M. T. J., Schouten, S., van Dongen, B. E., Rijpstra, W. I. C., Fuchs, G., Sinninghe Damsté, J. S., de Leeuw, J. W., and Ward, D. M.: Biosynthetic controls on the ¹³C contents of organic components in the photoautotrophic bacterium *chloroflexus aurantiacus*, *J. Biol. Chem.*, 276, 10971–10976, doi:10.1074/jbc.M009701200, 2001.
- van Kessel, M. A. H. J., Speth, D. R., Albertsen, M., Nielsen, P. H., Op den Camp, H. J. M., Kartal, B., Jetten, M. S. M., and Lüscher, S.: Complete nitrification by a single microorganism, *Nature*, 528, 555–559, doi:10.1038/nature16459, 2015.
- van Schöll, L., Kuypers, T., Smits, M., Landeweert, R., Hoffland, E., and Breemen, N.: Rock-eating mycorrhizas: their role in plant nutrition and biogeochemical cycles, *Plant Soil*, 303, 35–47, doi:10.1007/s11104-007-9513-0, 2008.
- Volkman, J. K., Jeffrey, S. W., Nichols, P. D., Rogers, G. I., and Garland, C. D.: Fatty acid and lipid composition of 10 species of microalgae used in mariculture, *J. Exp. Mar. Biol. Ecol.*, 128, 219–240, doi:10.1016/0022-0981(89)90029-4, 1989.
- Wenk, C. B., Brees, J., Zopf, J., Veronesi, M., Bourbonnais, A., Schubert, C. J., Niemann, H., and Lehmann, M. F.: Anaerobic ammonium oxidation (anammox) bacteria and sulfide-dependent denitrifiers coexist in the water column of a meromictic south-alpine lake, *ASLO*, 58, 1–12, doi:10.4319/lo.2013.58.1.0001, 2013.
- White, D. C.: Validation of quantitative analysis for microbial biomass, community structure, and metabolic activity, *Adv. Limnol.*, 31, 1–18, 1988.
- White, D. C., Davis, W. M., Nickels, J. S., King, J. D., and Robbie, R. J.: Determination of the sedimentary microbial biomass by extractable lipid phosphate, *Oecologia*, 40, 51–52, doi:10.1007/BF00388810, 1979.
- Wilkinson, S. G.: Gram-negative bacteria, in: *Microbial lipids*, edited by: Ratledge, C. and Wilkinson, S. G., Vol. 1. Academic Press, London, 299–488, 1988.
- Wisotzky, F.: *Angewandte Grundwasserchemie, Hydrogeologie und hydrogeochemische Modellierung*, Springer, Berlin, 2011.
- Yoshinaga, I., Amano, T., Yamagishi, T., Okada, K., Ueda, S., Sako, Y., and Suwa, Y.: Distribution and diversity of anaerobic ammonium oxidation (Anammox) bacteria in the sediment of a eutrophic freshwater lake, Lake Kitaura, Japan, *Microbiol. Environ.*, 26, 189–197, doi:10.1264/jsme2.ME10184, 2011.
- Zelles, L., Palojarvi, A., Kandeler, E., von Lütow, M., Winter, K., and Bai, Q. Y.: Changes in soil microbial properties and phospholipid fatty acid fractions after chloroform fumigation, *Soil Biol. Biochem.*, 29, 1325–1336, doi:10.1016/S0038-0717(97)00062-X, 1997.
- Zhang, C. L., Li, Y., Ye, Q., Fong, J., Peacock, A. D., Blunt, E., Fang, J., Lovley, D. R., and White, D. C.: Carbon isotope signatures of fatty acids in *Geobacter metallireducens* and *Shewanella algae*, *Chem. Geol.*, 195, 17–28, doi:10.1016/S0009-2541(02)00386-8, 2003.
- Zogg, G. P., Zak, D. R., Ringelberg, D. B., White, D. C., MacDonald, N. W., and Pregitzer, K. S.: Compositional and functional shifts in microbial communities due to soil warming, *Soil Sci. Soc. Am. J.*, 61, 475–481, doi:10.2136/sssaj1997.03615995006100020015x, 1997.