

**Physiology of a marine *Beggiatoa* strain and the
accompanying organism *Pseudovibrio* sp. –
a facultatively oligotrophic bacterium**

Dissertation

zur Erlangung des Doktorgrades eines
Doktors der Naturwissenschaften (Dr. rer. nat.)

dem Fachbereich Biologie/Chemie der
Universität Bremen

vorgelegt von

Anne Schwedt

Bremen, September 2011

Diese Arbeit wurde von November 2007 bis September 2011 in der Abteilung Mikrobiologie (Gruppe Ökophysiologie) am Max-Planck-Institut für Marine Mikrobiologie in Bremen angefertigt.

1. Gutachterin: Dr. Heide Schulz-Vogt

Universität Bremen

Max-Planck-Institut für Marine Mikrobiologie, Bremen

2. Gutachter: Prof. Dr. Ulrich Fischer

Universität Bremen

Tag des Promotionskolloquiums: 31. Oktober 2011

Table of contents

Summary	6
Zusammenfassung	7
<hr/>	
Chapter 1 – General introduction	10
Aims of the study	24
<hr/>	
Chapter 2 – Physiology and mat formation of a marine <i>Beggiatoa</i> culture	25
2.1 Sulfur respiration in a marine chemolithoautotrophic <i>Beggiatoa</i> strain	27
2.2 Coordinated movement of <i>Beggiatoa</i> filaments in oxygen-sulfide gradients and the effect of blue/green light	43
<hr/>	
Chapter 3 – Co-cultivation of a marine <i>Beggiatoa</i> strain and <i>Pseudovibrio</i> sp.	47
3.1 A chemolithoautotrophic <i>Beggiatoa</i> strain requiring the presence of a <i>Pseudovibrio</i> sp. for cultivation	49
3.2 The <i>Pseudovibrio</i> genus contains metabolically versatile and symbiotically interacting bacteria	53
<hr/>	
Chapter 4 – Isolation and cultivation of <i>Pseudovibrio</i> sp. and other facultatively oligotrophic bacteria	55
4.1 Substrate use of <i>Pseudovibrio</i> sp. growing in extremely oligotrophic seawater	57

Table of contents

4.2 Facultatively oligotrophic bacteria isolated from the habitat of large sulfide-oxidizers 77

Chapter 5 – Concluding remarks 88

Conclusions 98

Outlook 99

References 101

List of abbreviations 114

Appendix 115

Acknowledgements 145

Summary

The oceans cover large parts of the earth's surface and play an important role in the cycling of elements. The large filamentous sulfide-oxidizing bacteria are capable of forming huge microbial mats at the oxic-anoxic interface of the sediment surface, where they oxidize sulfide using either oxygen or nitrate as electron acceptor. Thereby, they can strongly influence and connect the different nutrient cycles. The water column above is populated by planktonic bacteria, which account for a large fraction of biomass on earth. Consequently, these organisms also strongly influence the turnover of nutrients in the oceans.

The first part of this thesis (Chapter 2) addresses the physiology and mat formation processes of the large sulfide-oxidizers belonging to the genus *Beggiatoa*. Until now, it was assumed that nitrate as an alternative electron acceptor is crucial for the migration of marine *Beggiatoa* spp. into deeper anoxic sediment layers. We found that a subpopulation of the investigated *Beggiatoa* filaments actively migrates into anoxic, sulfidic layers as a reaction to high sulfide fluxes without the presence of nitrate. Our experiments show that the reason for this so far unknown migration behavior seems to be excessive storage of elemental sulfur and organic carbon due to high sulfide fluxes, which leads to filaments extremely filled with storage compounds that tend to break easily at this stage. By moving into anoxic regions, aerobic sulfide oxidation is stopped and storage space is emptied by reducing the stored sulfur with carbon reserve compounds.

The investigated sulfide-oxidizer (*Beggiatoa* sp.) depends on the presence of a small heterotrophic bacterium (*Pseudovibrio* sp.). This association is investigated in the second part of this thesis (Chapter 3). The associated *Pseudovibrio* sp. mainly populates the oxic part of the gradient co-culture. This suggests that these bacteria are mainly required for the oxic growth of the *Beggiatoa* sp. and might protect them from oxidative stress, as *Beggiatoa* spp. are typically known to lack the gene encoding for the enzyme catalase. Supporting this hypothesis, we found that the genome of the accompanying *Pseudovibrio* sp. possesses several genes for enzymes involved in the protection against reactive oxygen species.

In contrast to the large *Beggiatoa* sp., the associated *Pseudovibrio* sp. is able to grow in pure culture. Besides heterotrophic growth on organic-rich media, the bacteria are also able to grow under extremely oligotrophic (nutrient-poor) conditions. A detailed analysis of the substrate use under oligotrophic conditions revealed that *Pseudovibrio* sp. grows on organic

contaminations preferentially containing nitrogen (Chapter 4). Interestingly, we could isolate further facultatively oligotrophic bacteria from water overlaying Namibian sediments, which are known to inhabit many different large sulfide-oxidizers.

Zusammenfassung

Die Ozeane bedecken große Teile der Erdoberfläche und spielen somit eine wichtige Rolle in Bezug auf die Kreisläufe der Elemente. Große, filamentöse, sulfidoxidierende Bakterien können enorme mikrobielle Matten auf der Sedimentoberfläche bilden. Diese Bakterien oxidieren das aufsteigende Sulfid mit Sauerstoff oder Nitrat als Elektronenakzeptor, wodurch sie die verschiedenen Nährstoffkreisläufe der Ozeane beeinflussen und verbinden. In der darüber liegenden Wassersäule befinden sich planktonische Bakterien, welche durch die enorme Größe der Ozeane einen erheblichen Anteil der Biomasse auf der Erde darstellen. Folglich wird auch der Umsatz der Nährstoffe im Ozean stark von diesen Organismen beeinflusst.

Der erste Teil dieser Dissertation (Kapitel 2) befasst sich mit der Physiologie und Mattenbildung der großen, sulfidoxidierenden Bakterien aus dem Genus *Beggiatoa*. Bisher wurde angenommen, dass das Vorhandensein von Nitrat als alternativer Elektronenakzeptor essenziell für die Migration von *Beggiatoa* sp. in anoxische Sedimentschichten sei. Wir konnten in unserer Studie zeigen, dass eine Subpopulation der untersuchten *Beggiatoa* Filamente ohne zur Verfügung stehendes Nitrat aktiv anoxische, sulfidische Bereiche aufsuchen kann. Der Grund für dieses bislang unbekanntes Migrationsverhalten scheint die übermäßige Speicherung an internem Schwefel und Kohlenstoff zu sein, welche als Folge von einem hohen Sulfidflux auftritt. Die erhöhte Speicherung führt dazu, dass die Filamente sehr mit Speicherstoffen angefüllt sind und dadurch leicht brechen. Die aerobe Sulfidoxidation kann unterbrochen werden, indem die Filamente sich in anoxische Bereiche bewegen, wo sie den internen Schwefel mit intern gespeichertem Kohlenstoff reduzieren.

Das Wachstum der untersuchten Sulfidoxidierer (*Beggiatoa* sp.) ist abhängig von der Anwesenheit von kleinen heterotrophen Bakterien (*Pseudovibrio* sp.). Diese Assoziation wurde im zweiten Teil dieser Dissertation untersucht (Kapitel 3). Die assoziierten Bakterien (*Pseudovibrio* sp.) sind vorwiegend im oxischen Bereich der Co-Kultur zu finden, was vermuten lässt, dass sie besonders für das aerobe Wachstum von *Beggiatoa* sp. erforderlich sind. Da *Beggiatoa* spp. typischerweise nicht über das Gen für das Enzym Katalase verfügen,

ist es möglich, dass die assoziierten Bakterien ihre Partner vor oxidativem Stress schützen. Diese Vermutung wird dadurch unterstützt, dass wir im Genom des Begleitorganismus (*Pseudovibrio* sp.) diverse Gene für Enzyme gefunden haben, die vor reaktiven Sauerstoffspezies schützen.

Im Gegensatz zu den großen Sulfidoxidierern (*Beggiatoa* sp.) können die assoziierten Bakterien (*Pseudovibrio* sp.) in Reinkultur leben. Neben heterotrophem Wachstum auf kohlenstoffhaltigen Medien, können die Bakterien unter extrem oligotrophen (nährstoffarmen) Bedingungen wachsen. Eine detaillierte Analyse der Substrate, die unter diesen nährstoffarmen Bedingungen benutzt werden, hat gezeigt, dass *Pseudovibrio* sp. auf stickstoffhaltigen, organischen Kontaminationen wachsen kann (Kapitel 4). Interessanterweise konnten wir weitere fakultativ oligotrophe Bakterien aus dem Wasser über Namibischen Sedimenten isolieren. Namibische Sedimente sind bekannt für ihre Vielzahl an verschiedenen Sulfidoxidierern.

„Science is built up of facts, as a house is built of stones; but an accumulation of facts is no more a science than a heap of stones is a house.“

~Jules Henri Poincaré (1854-1912)

Chapter 1

General introduction

Marine element cycles

Nutrients are chemical compounds that are required for the metabolism of living organisms and have to be taken up from the environment. Bacterial nutrition includes both organic and inorganic molecules. The turnover of the individual elements in these nutrients is referred to as ‘element cycling’.

The marine carbon cycle

Carbon is the major element of cellular material (Battley, 1995). In the model organism *Escherichia coli*, for instance, the amount of cellular carbon accounts for 48 to 59% of the dry weight (Battley, 1995; Norland *et al.*, 1995). The production of new organic material, also referred to as primary production, takes place in the ocean mainly via photosynthesis. In this process, carbon dioxide from the atmosphere is fixed to form new organic matter (Figure 1.1) using the energy from sunlight. Primary production is the main source of dissolved organic carbon (DOC) in the open ocean, which occurs within the euphotic zone (Hansell *et al.*, 2009). An additional source of DOC is terrestrial organic carbon that is transported into the ocean by rivers and serves likewise as a fixed carbon source for marine microorganisms (Schlünz and Schneider, 2000), but accounts for only a minor fraction. The rate of primary production in the ocean surface waters generally controls the flux of organic matter towards the sediment (Suess, 1980; Jørgensen, 1983). Sinking to the bottom of the ocean, the fixed organic material is degraded and transformed by microorganisms and chemical processes.




Figure 1.1: The oceanic carbon cycle. Carbon dioxide from the atmosphere is fixed into organic carbon which can sink down to the seafloor as particulate organic matter (POM). The labile dissolved organic matter (LDOM) can be respired to CO_2 and the recalcitrant dissolved organic matter (RDOM) is inert to bacterial breakdown. (Image redrawn from Jiao *et al.*, 2010 and references therein)

The organic matter in the ocean can be divided in particulate organic matter (POM) and dissolved organic matter (DOM). Part of the POM pool sinks down to the seafloor where it can be stored for long periods of time (Figure 1.1, Ducklow *et al.*, 2001). The DOM pool consists of labile dissolved organic matter (LDOM) and recalcitrant dissolved organic matter (RDOM). The LDOM fraction can partly be transformed by microorganisms, thereby, LDOM is oxidized by heterotrophic microorganisms within days forming again carbon dioxide. Molecules like amino acids and monosaccharides as part of the LDOM fraction can easily be utilized by the marine bacterioplankton (Bauer *et al.*, 1992; Cherrier *et al.*, 1996; Kirchman *et al.*, 2001) and make up 75% of the DOC that is consumed by marine microorganisms in the upper layers of the ocean (Cherrier and Bauer, 2004). The RDOM, on the other hand, is assumed to be resistant to biological degradation and can be stored in the ocean for millennia (Figure 1.1, Bauer *et al.*, 1992; Kirchman *et al.*, 2001; Hopkinson and Vallino, 2005; Jiao *et al.*, 2010). The composition of dissolved organic matter in the ocean is highly diverse and DOM can consist of thousands of different organic compounds of which only few (<10%) have yet been identified with specific molecular formulas (Koch *et al.*, 2005; Hertkorn *et al.*, 2006; Dittmar and Paeng, 2009).

The marine sulfur cycle

Sulfur makes up only about 1% of the cellular dry weight (Battley, 1995), however, it is essential for the formation of amino acids (cysteine, methionine) and vitamins (biotin). In most marine environments, sulfur is not a limiting factor due to the high sulfate concentration of 28 mmol L^{-1} in seawater (Volkov and Rozanov, 1983). In the marine environment, sulfur can be found in varying oxidation states ranging between $[-2]$ and $[+6]$ (Figure 1.2). The potential to transform between the different oxidation states represents the importance of this element as it can serve as an electron donor or acceptor in various key redox reactions.




Figure 1.2: Different oxidation states of the element sulfur ranging from $[+6]$ to $[-2]$. (Image adapted from Chameides and Perdue, 1997)

In marine sediments, alternative electron acceptors, like sulfate, are present below the oxygen penetration depth. In anoxic layers, sulfate is used by microorganisms to oxidize organic and inorganic electron donors while reducing sulfate to sulfide. In coastal marine sediment from Aarhus Bay (Denmark) sulfate reduction takes place below 4 cm depth, which was concluded from hydrogensulfide (H_2S) production (Jørgensen and Nelson, 2004). These anoxic sediment layers are, therefore, characterized by an upwards directed sulfide flux. When sulfide reaches the oxic-anoxic interface and reacts with oxygen it gets oxidized back to sulfur or sulfate either chemically or biologically. The biological oxidation mediated by bacteria, for example of the genus *Beggiatoa*, was shown to be three times faster than the chemical oxidation (Nelson *et al.*, 1986a). Due to the formation of large bacterial mats in certain habitats, the sulfide-oxidizing bacteria *Beggiatoa* spp. have a huge potential to oxidize large amounts of the upwards diffusing sulfide in these areas (Jørgensen, 1977), thereby strongly influencing the marine sulfur cycle.

The marine nitrogen cycle

Nitrogen, as a component of proteins and nucleic acids, is a fundamental molecule of life and cellular material consists to about 15% of nitrogen (Battley, 1995). The major nitrogen reservoir is the atmosphere, consisting of 78% nitrogen in the form of N_2 gas (Fiadeiro, 1983). Only few microorganisms have the ability to fix the atmospheric N_2 and make it available also for other organisms. Nitrogen fixation is an energy consuming process since N_2 is triple-bonded and has to be cleaved during the fixation process. Thereby, nitrogen gets reduced and is present in organisms in the most reduced form, the particulate organic nitrogen (PON, Figure 1.3). The PON can be remineralized to ammonia. Nitrifying microorganisms are able to oxidize ammonia aerobically to nitrate over nitrite, which is a process mediated by two metabolically different groups of bacteria. The formed nitrate can be used as electron acceptor in anaerobic environments (Figure 1.3), for example by the large sulfur bacteria of the genus *Beggiatoa*. Thereby, nitrate is reduced back to ammonia (dissimilatory nitrate reduction to ammonia = DNRA) or to gaseous nitrogen compounds (denitrification). Denitrification removes fixed nitrogen from the system, because the gaseous end-product N_2 needs to be fixed again by microorganisms to make it biologically available. Besides denitrification, fixed nitrogen can also be removed from the system by anaerobic ammonium oxidation (anammox). During this process, ammonia is anaerobically oxidized to N_2 using nitrite as electron acceptor (Strous *et al.*, 1999).




Figure 1.3: The marine nitrogen cycle. Nitrogen from the atmosphere is fixed into particulate organic nitrogen (PON) which can be remineralized to ammonia. Ammonia can be either oxidized aerobically to nitrate or anaerobically with nitrite (anammox) producing N_2 and removing fixed nitrogen from the system. Nitrate can also be reduced to gaseous nitrogen compounds (denitrification) that leave the system. (Image based in part on Arrigo, 2005; and is reproduced from Francis *et al.*, 2007)

Connection of marine element cycles

The cycling of the elements ranges from the turnover of single molecules to entire pathways occurring in living cells, thereby connecting all element cycles. The element cycling of individual cells does eventually influence the entire ecosystem on a broad scale (Bolin *et al.*, 1983). Microorganisms are composed of many different elements, such as carbon, nitrogen, sulfur, phosphorus, oxygen, hydrogen and many microelements like iron or magnesium (Battley, 1995). As a consequence, the new production or decomposition of biomass will automatically connect the different element cycles.

The marine element cycles are, furthermore, connected by the diverse metabolisms of bacteria. Redox reactions always combine the reduction of an electron acceptor with the oxidation of an electron donor. In nearly all cases, electron acceptor and donor are composed of different elements. Denitrification, which is the reduction of nitrate (NO_3^-) to molecular nitrogen (N_2 , N-cycle), for example can be coupled to the oxidation of organic carbon compounds (C-cycle) or the oxidation of reduced inorganic sulfur compounds (S-cycle). Additionally, both organic carbon and inorganic reduced sulfur compounds can also be oxidized using oxygen (O-cycle) as an electron acceptor. This is only an excerpt of many metabolic pathways connecting the cycling of the single elements, including different electron donors (e.g. sulfide, hydrogen, organic material) and electron acceptors (e.g. oxygen, nitrate, sulfate).

In marine habitats, the mineralization of organic matter, such as dead organic material consisting of many different elements, is an important process combining nutrient cycles. In pelagic regions, this mainly occurs in the water column by the metabolic activity of free-living bacteria (Azam and Hodson, 1977; Tabor and Neihof, 1982; Ishida *et al.*, 1989). There, nutrient hotspots exist, such as marine snow particles that contain high amounts of organic matter. Bacteria densely aggregate on these particles (e.g. Smith *et al.*, 1992; Azam and Malfatti, 2007 and references therein) and can achieve high growth rates (e.g. Alldredge *et al.*, 1986; Kjørboe and Jackson, 2001). In contrast, organic matter remineralization in shallow waters, such as fjords or continental shelves, takes mainly place in the sediment. Thus, depending on the water depth, these are the substantial regions for nutrient cycling in the marine environment (Jørgensen, 1983). The connection of nutrient cycles in marine sediments (reviewed in Jørgensen, 1983) involves a cascade of transformation processes. Aerobic degradation of organic material in shallow marine sediments takes place within a thin layer at the sediment surface, where the oxidation of organic matter to carbon dioxide occurs. Below

this oxic zone, anaerobic processes take place that successively oxidize the residual organic matter via different metabolic pathways by diverse microorganisms. From the top sediment layers to the deeper regions, the electron acceptor used is determined by its energy yield per mole carbon being oxidized. From top to bottom, the preferred electron acceptor gradually decreasing from oxygen to carbon dioxide via nitrate, iron, manganese and sulfate, combining the C-cycle to the N-, Fe-, Mn- and S-cycle (Jørgensen, 1983). Most importantly in the anoxic regions are, therefore, the highly abundant inorganic nitrogen and sulfur compounds, which are concomitantly reduced to N₂ and H₂S. Reduced substances, such as sulfide and methane that are produced in deep sediment layers diffuse upwards and become oxidized to form sulfate and carbon dioxide, thereby closing the cycling of elements (Jørgensen, 1983).

Sulfide-oxidizing bacteria of the genus *Beggiatoa*

More than two centuries ago, bacteria of the genus *Beggiatoa* were discovered (Vaucher, 1803). They were originally described as *Oscillatoria alba* because they feature a similar filamentous morphology as the cyanobacteria of the genus *Oscillatoria*, but have a whitish appearance instead of the blue-green pigments (Figure 1.4). About 40 years later, these colorless sulfur bacteria were reclassified as *Beggiatoa alba*, named after the Italian scientist F. S. Beggiato (Trevisan, 1842). Based on their morphology, different filamentous sulfur bacteria were assigned to the genus *Beggiatoa*. Several species were differentiated on the basis of filament diameter size classes ranging between 1–55 µm (Vaucher, 1803; Trevisan, 1842; Hinze, 1901; Klas, 1937). However, only a small number of 16S rDNA sequences were available until recently, which made it difficult to phylogenetically classify the large sulfur bacteria. It was even found that filaments with a similar morphology belong to phylogenetically different genera (Ahmad *et al.*, 1999; Ahmad *et al.*, 2006). In a single-cell 16S rDNA gene sequencing approach of large sulfur bacteria, Salman *et al.* (2011) strongly extended the amount of available sequences and proposed based on phylogenetic analysis new candidatus genera names for the members of the family *Beggiatoaceae*. According to this reclassification, the genus *Beggiatoa* contains aerobic or microaerophilic filamentous bacteria with a diameter of 1–9 µm.




Figure 1.4: Bright field micrographs of filamentous bacteria of the genera (A) *Oscillatoria* and (B) *Beggiatoa* from a freshwater enrichment culture. (Image reproduced from Bondarev, 2007)

Mat-formation and physiology of *Beggiatoa* spp.

Filaments of the genus *Beggiatoa* can be several centimeters long and move by gliding. Pores on the surface of *Beggiatoa* filaments are arranged as spirals and are assumed to be involved in the gliding motility by the excretion of slime (Larkin and Henk, 1996). This spatial flexibility allows the *Beggiatoa* filaments to position themselves in the chemical microenvironment of sediments. As a consequence, *Beggiatoa* are able to form mats in different habitats, such as sulfidic marine and freshwater sediments (Winogradsky, 1887; Jørgensen, 1977; Nelson and Castenholz, 1982; McHatton *et al.*, 1996), activated sludge (Farquhar and Boyle, 1971), hot vents (Nelson *et al.*, 1989), cold seeps (Barry *et al.*, 1996) and in hypersaline lakes (Hinck *et al.*, 2007).

Beggiatoa filaments usually form a distinct mat in the transition zone of oxygen and sulfide (Winogradsky, 1887; Keil, 1912; Jørgensen, 1977). *Beggiatoa* spp. oxidize the upwards diffusing sulfide, via elemental sulfur to sulfate using oxygen as electron acceptor (Winogradsky, 1887; Nelson and Castenholz, 1981). The consumption of oxygen and sulfide by the bacteria steepens the gradients of oxygen and sulfide and narrows the transition zone to a few micrometer (Figure 1.5, Nelson *et al.*, 1986a).




Figure 1.5: H_2S and O_2 microprofiles in (A) an uninoculated control medium and (B) an inoculated *Beggiatoa* culture. In the uninoculated medium, O_2 and H_2S gradients overlap, whereas in the culture the bacteria form a mat between the opposing gradients (shaded area) and steepen the gradients by aerobic sulfide oxidation and raise the overlapping zone to 2.5 mm. (Image reproduced from Nelson *et al.*, 1986a)

Simulating the natural habitat of the *Beggiatoa*, agar-based oxygen-sulfide gradient media are used to cultivate these large sulfide-oxidizers (Nelson *et al.*, 1982; Nelson and Jannasch, 1983). The formation of a distinct mat of *Beggiatoa* filaments between their electron acceptor and donor depends on different parameters. Besides the fact that both oxygen and sulfide are essential for the growth of the bacteria, each of these substances is also toxic if present in higher concentrations. Exceeding 5% air saturation, oxygen induces a phobic reaction of *Beggiatoa* filaments (Møller *et al.*, 1985). In contrast, long-lasting depletion of oxygen causes filaments to move into the direction of the oxygen source (Winogradsky, 1887; Møller *et al.*, 1985). The concentration of oxygen, therefore, defines the upper border of the *Beggiatoa* mat. The lower border of the *Beggiatoa* layer is defined by the sulfide flux from below. With increasing sulfide flux the *Beggiatoa* filaments position themselves at higher layers in the agar-based gradient culture tubes (Figure 1.6, Nelson and Jannasch, 1983) and if sulfide exceeds a critical concentration, the filaments die (Winogradsky, 1887). Additionally, a phobic reaction of *Beggiatoa* filaments towards light was observed and thus light might also influence the gliding direction and consequently the position of the *Beggiatoa* mat (Winogradsky, 1887; Nelson and Castenholz, 1982; Møller *et al.*, 1985).




Figure 1.6: Position of *Beggiatoa* cell layers (mats) in culture tubes with different sulfide concentrations in the bottom agar plug. With increasing sulfide, the filaments form a mat located higher in the culture tube. (Image reproduced from Nelson and Jannasch, 1983)

Elemental sulfur, which is the intermediate of sulfide oxidation, can be stored inside the *Beggiatoa* cells (Winogradsky, 1887) and leads to the whitish appearance of the filaments. Using electron microscopy, it was shown that the sulfur globules in the cells are surrounded by the cytoplasmic membrane and are located in the periplasm (Figure 1.7 A, Strohl *et al.*, 1982). The intracellular sulfur can serve as an electron donor and be further oxidized to sulfate when sulfide gets limited in the environment (Winogradsky, 1887). In addition to the storage of sulfur, *Beggiatoa* have the ability to store polyhydroxyalkanoates (PHA, sometimes specifically denoted as poly- β -hydroxybutyric acid [PHB]) in the cytoplasm of the cell (Figure 1.7 A, Pringsheim, 1964; Strohl and Larkin, 1978; Strohl *et al.*, 1982). The amount of PHA in the cell can account for up to 50% of the dry weight of the cell (Güde *et al.*, 1981). Furthermore, an accumulation of polyphosphate in *Beggiatoa* cells was shown by transmission electron microscopy and different staining methods (Figure 1.7 C, Maier and Murray, 1965; Strohl and Larkin, 1978; de Albuquerque *et al.*, 2010; Brock and Schulz-Vogt, 2011).

About two decades ago, extremely large marine filamentous sulfur bacteria (116–122 μm in diameter) containing a central vacuole were found and identified as *Beggiatoa* spp. based on morphological similarities to these organisms (Figure 1.7 B, Nelson *et al.*, 1989). Few years later, the storage of nitrate, an alternative electron acceptor, was detected within the vacuoles of these large filaments (McHatton *et al.*, 1996). It was proposed that the oxidation of sulfide can be coupled to either DNRA (Sayama, 2001; Sayama *et al.*, 2005) or denitrification (Sweerts *et al.*, 1990). The storage of nitrate allows the filaments to inhabit deeper anoxic sediment layers. Carrying nitrate down into anoxic sediment layers and use it for sulfide oxidation can lead to the separation of oxygen and sulfide gradients over several centimeters (Mußmann *et al.*, 2003; Sayama *et al.*, 2005; Kamp *et al.*, 2006). This life strategy enables large, vacuolated sulfur bacteria like *Beggiatoa* spp. to outcompete non-vacuolated, non-

motile sulfide-oxidizers in anaerobic environments. Close relatives of *Beggiatoa*, like bacteria belonging to the candidate genus “*candidatus* Marithioploca”, also use and store nitrate and even show a positive chemotactic response towards nitrate (Huettel *et al.*, 1996; reclassified by Salman *et al.*, 2011). Thus, the orientation and mat formation of the vacuolated nitrate-storing sulfur bacteria may also be influenced by the nitrate flux.

Studying the physiology of *Beggiatoa*, Winogradsky (1887) developed the concept of chemolithotrophy. He observed that the growth of *Beggiatoa* was dependent on reduced inorganic sulfur compounds but not on the presence of organic compounds. The utilization of CO₂ as a sole carbon source was later confirmed by isotope-labeling studies (Nelson and Jannasch, 1983). Besides these chemolithoautotrophic strains, many chemoorganoheterotrophic *Beggiatoa* strains were isolated (Strohl and Larkin, 1978; Güde *et al.*, 1981; Strohl *et al.*, 1981), which are able to oxidize sulfide only in the presence of organic compounds. Furthermore, also mixotrophic *Beggiatoa* strains were isolated (Pringsheim, 1967; Güde *et al.*, 1981) thus reflecting the diverse metabolisms present within the genus *Beggiatoa*.




Figure 1.7: Cell structures of *Beggiatoa* filaments. (A) Schematic representation of *Beggiatoa alba* strain B15LD indicating the location of sulfur globules [S] in the periplasm and poly-β-hydroxybutyrate [PHB] in the cytoplasm. (B) Transmission electron micrograph of a *Beggiatoa* sp. cross section. The cytoplasm of this large *Beggiatoa* filament is restricted to the edge of the cell and the interior mainly consists of a large central vacuole. (C) Transmission electron micrograph showing electron-dense inclusion bodies in the cytoplasm of *Beggiatoa* filaments probably consisting of polyphosphate [P]. (Images adapted and reproduced from Strohl *et al.*, 1982 [A]; Nelson *et al.*, 1989 [B]; de Albuquerque *et al.*, 2010 [C])

The investigated *Beggiatoa* sp. co-culture

The marine *Beggiatoa* sp. strain 35Flor investigated in this thesis was isolated in 2002 from a microbial community associated with scleractinian corals suffering from black band disease off the coast of Florida. This *Beggiatoa* sp. strain grows under chemolithoautotrophic conditions in an agar-stabilized oxygen-sulfide gradient medium gaining energy from the aerobic oxidation of sulfide. Both, a fixed carbon and a fixed nitrogen source are absent from the medium and nitrogen fixation in the investigated *Beggiatoa* sp. was determined earlier (Henze, 2005). Typical storage compounds of the genus *Beggiatoa*, such as sulfur, PHA and polyphosphate were found in the investigated filaments (Schwedt, unpublished data, Brock and Schulz-Vogt, 2011). A central vacuole is present (Kamp *et al.*, 2008; Brock and Schulz-Vogt, 2011), but the storage of nitrate could not be detected (Schwedt *et al.*, unpublished data).

The *Beggiatoa* sp. strain 35Flor is accompanied by only one type of organism (Bachmann, 2007), the *Pseudovibrio* sp. strain FO-BEG1. Unlike the *Beggiatoa* sp., the associated bacteria are able to grow in pure culture and could be isolated in artificial seawater medium. The investigated *Pseudovibrio* sp. is able to grow in pure artificial seawater medium under extreme nutrient-poor conditions (Bachmann, 2007) and thus belongs to the few so far cultured extremely oligotrophic organisms.

Bacterial growth under nutrient deficiency

The term ‘oligotroph’ was introduced by Weber (1907) to describe an organism growing under nutrient deficiency as opposed to that, bacteria growing under nutrient affluence are called ‘eutrophs’ (organisms living in nutrient-rich environments are sometimes also referred to as ‘copiotrophs’). Over time, several definitions of oligotrophy arose and today it is generally accepted that bacteria are referred to as oligotrophic when they are able to grow in medium containing less than 0.5 mg C L⁻¹ (e.g. Ishida *et al.*, 1989). When their growth is inhibited by high substrate concentrations, the bacteria are considered to be obligately oligotrophic, which is in contrast to facultatively oligotrophic bacteria, which are able to grow under both nutrient-poor and nutrient-rich conditions (Ishida *et al.*, 1989). Facultative oligotrophs are, therefore, successful in environments with changing nutrient conditions.

The open ocean, covering large parts of the earth's surface, is low in nutrients and contains less than 1 mg DOC in 1 L seawater (Schut *et al.*, 1997; Hansell *et al.*, 2009). Thus, it is denoted as an oligotrophic environment. 75% of the carbon consumed by the bacteria in the ocean can be composed of dissolved free amino acids (DFAA), dissolved combined amino acids (DCAA) and monosaccharides. The utilization of these substances can cover 5 to 93% of the carbon demand of the bacteria and 9 to 100% of the nitrogen demand (Fuhrman, 1987; Jørgensen, 1987; Stanley *et al.*, 1987; Keil and Kirchman, 1999; Cherrier and Bauer, 2004).

Attached and free-living marine bacteria

The particulate organic matter (POM) is an important part of the organic matter in the ocean. Particles larger than half a millimeter are so-called marine snow particles (Suzuki and Kato, 1953; Silver *et al.*, 1978). Besides the larger marine snow particles, there are also smaller microaggregates (Figure 1.8 A and B) and both consist of detrital organic and inorganic matter (Azam and Long, 2001), thereby representing hotspots of high nutrient concentration. The aggregates can be colonized by metazoans (e.g. Shanks and Edmondson, 1990; Kiørboe, 2000), protozoans (e. g. Silver *et al.*, 1978) and prokaryotes (e. g. Alldredge *et al.*, 1986; Smith *et al.*, 1992; Azam and Malfatti, 2007 and references therein), whereas only the latter was found on all types of aggregates studied so far. Extracellular hydrolytic enzymes produced by aggregate-associated bacteria can convert the POM of the sinking aggregates into cell biomass and non-sinking dissolved organic matter (DOM) (Smith *et al.*, 1992; Grossart *et al.*, 2007). While sinking down the particles leave behind a DOM plume that is composed mainly of carbon and nitrogen. The DOM plume is colonized by some of the attached bacteria but also by free-living bacteria from the surrounding water (Figure 1.8 C, Azam and Long, 2001; Kiørboe and Jackson, 2001).

Compared to the surrounding water, bacterial cell densities on aggregates are typically >100 times higher (e. g. Smith *et al.*, 1992; Turley and Mackie, 1994). Nevertheless, the particle-associated bacteria account only for <5% of the total bacterial numbers in seawater (e.g. Alldredge *et al.*, 1986; Alldredge and Gotschalk, 1990; Turley and Stutt, 2000) and contribute to only 3 to 12% of the total bacterial production (Alldredge *et al.*, 1986; Turley and Stutt, 2000). Even though the total activity is low, the per cell activity of the attached bacteria is higher compared to free-living bacteria, as demonstrated by higher incorporation rates and shorter doubling times (Alldredge *et al.*, 1986; Alldredge and Gotschalk, 1990; Smith *et al.*, 1992; Azam and Long, 2001; Kiørboe and Jackson, 2001). Furthermore, some

studies have shown that the free-living bacteria may either starve and not be active (Boylen and Ensign, 1970; Novitsky and Morita, 1976; 1977), while other studies show that they may be metabolically active (Azam and Hodson, 1977; Tabor and Neihof, 1982; Ishida *et al.*, 1989).




Figure 1.8: *In situ* photographs of (A) a marine snow aggregate in a pelagic environment and (B) microaggregates in a shallow environment (photos M. Lunau). (C) Scheme of a marine snow particle colonized by bacteria which excrete hydrolytic enzymes converting marine snow into DOM forming a plume behind the sinking aggregate that is also colonized by attached and free-living bacteria. The DOM consists mainly of carbon [C] and nitrogen [N]. (Images adapted and redrawn from Azam and Long, 2001 [C]; and reproduced from Simon *et al.*, 2002 [A and B])

The majority of the free-living bacteria in the open ocean is exposed to extremely low nutrient concentrations and many different survival strategies have evolved to cope with this nutrient limitation. These strategies include concentration-independent enzyme production (cells are considered to be prepared and have enzymes ready for substrates becoming available), depression of substrates (the use of one substrate is not repressed by another more efficient one) and the use of multiple substrates simultaneously (use different substrates at the same time, independent of their efficiency) (Egli, 2010 and references therein). Substrate tests on organisms grown under carbon limitation revealed that these cells can oxidize a much broader spectrum of organic compounds than cells that were pre-grown under carbon excess (Upton and Nedwell, 1989; Ihssen and Egli, 2005). The use of multiple carbon sources enables

growth on extremely low concentrations of each individual compound (Lendenmann *et al.*, 1996; Kovárová-Kovar and Egli, 1998) and is thus beneficial in an oligotrophic environment with a frequently changing supply of nutrients.

Cultivation of marine bacteria

So far, only about half of the known bacterial phyla have cultivable representatives (Hugenholtz, 2002), even though pure cultures are essential to study metabolic pathways of the different bacteria in detail. Possible reasons for the yet inability to cultivate many bacteria maybe unsuited growth conditions and could include a lack of nutrients or growth factors, inappropriate pH, pressure or temperature conditions or unsuitable levels of oxygen (reviewed in Vartoukian *et al.*, 2010). Furthermore, many of the used media contain very high amounts of nutrients, compared to most marine environments, and thus favor fast-growing bacteria rather than slow-growing ones. In turn, such conditions might even inhibit the growth of some oligotrophic bacteria (Ishida *et al.*, 1989; Koch, 1997; Connon and Giovannoni, 2002). Consequently, new strategies for the isolation of marine bacteria have to be developed to understand the different metabolic pathways of marine bacteria and their ecology and evolution (Grossart, 2010).

One approach to prevent overgrowth of slow-growing bacteria is the dilution-to-extinction method, that reduces the number of cells per sample until ideally solely single cells are left for cultivation (e. g. Button *et al.*, 1993; Connon and Giovannoni, 2002). Additionally, the use of low-nutrient natural seawater for isolation and *in vitro* simulation of the natural environment using diffusion chambers placed in natural seawater provoked isolation of new, so far uncultured bacteria (Connon and Giovannoni, 2002; Kaeberlein *et al.*, 2002; Rappe *et al.*, 2002; Zengler *et al.*, 2002; Bollmann *et al.*, 2007). However, the utilization of natural seawater always implies undefined conditions because merely a few percent of the highly diverse organic compounds in natural seawater is already characterized (Dittmar and Paeng, 2009). Hence, in order to study bacterial metabolism at the lower border of bacterial growth in detail and to identify the essential substances for growth, a defined artificial seawater medium is crucial. Those approaches so far reported to isolate and cultivate marine bacteria using artificial seawater contained either agar or vitamins, both of which represent a fixed carbon source, or were supplemented with at least 0.1 to 3 mg C L⁻¹ of organic substrates to support growth (Van der Kooij *et al.*, 1980; Ishida *et al.*, 1982; Schut *et al.*, 1993; Azam and Long, 2001; Vancanneyt *et al.*, 2001).

Aims of this study

This work was initiated by the question of how marine *Beggiatoa* spp. form mats and succeed in anoxic habitats. Until today, it was believed that only the presence of nitrate as alternative electron acceptor allows the population of anoxic environments by the large sulfide-oxidizing bacteria of the family *Beggiatoaceae*. Recently, I found that nitrate is not essential for the thriving of *Beggiatoa* filaments in anoxic parts. For these experiments, I used the marine *Beggiatoa* sp. 35Flor that is cultivated in gradient culture tubes. It was observed that filaments moved below the oxygen-sulfide interface without the presence of nitrate and aggregated in anoxic parts of the culture tube. Therefore, the aim of the first part of this thesis (Chapter 2) was to study this behavior and to reveal how the filaments can survive in the anoxic layers and why they leave the overlapping zone of oxygen and sulfide, where both electron acceptor and donor are present.

Already during my diploma thesis (Bachmann, 2007) I was able to show that the investigated *Beggiatoa* culture is not a pure culture. Instead, the *Beggiatoa* sp. 35Flor is in co-culture with a single accompanying organism, *Pseudovibrio* sp. FO-BEG1. Accordingly, the second objective of my PhD thesis (Chapter 3) was to examine whether the growth of the sulfide-oxidizer is dependent on the presence of the accompanying *Pseudovibrio* sp. and, if so, whether the *Pseudovibrio denitrificans* type strain (DSM number 17465) can also provoke growth of the *Beggiatoa* sp. 35Flor.

The accompanying *Pseudovibrio* sp. FO-BEG1 is able to grow in pure culture without the *Beggiatoa* sp. under extreme nutrient deficiency in artificial seawater medium (Bachmann, 2007). The physiology of the *Pseudovibrio* sp. should now be subject to a detailed physiological analysis. Despite omitting the addition of an energy source, DOC was detected in the range of $5 \mu\text{mol C L}^{-1}$ (0.06 mg C L^{-1}), which is 1 to 2 orders of magnitude below natural oligotrophic seawater (Schut *et al.*, 1997; Hansell *et al.*, 2009). This contamination could have potentially been used as an energy source. To address this question, the third objective of this thesis (Chapter 4) was to analyze the artificial medium used for cultivation, before and after growth of the *Pseudovibrio* strain, in order to find out which compounds were used by the bacteria. Eventually, other heterotrophic bacterial strains were isolated in the course of this thesis under nutrient limitation to estimate how common the ability among heterotrophic bacteria (associated with large sulfide-oxidizers) is to grow under nutrient limitation.

Chapter 2

Physiology and mat formation of a marine *Beggiatoa* culture

This second chapter of my PhD thesis deals with the physiology of the large, sulfide-oxidizing *Beggiatoa* sp. strain 35Flor. The focus is laid on mat formation processes and was motivated by a new observation that filaments migrate into deeper anoxic regions without the presence of nitrate (Figure 2). In the first part of this chapter, the physiology behind the observed migration event is discussed in detail in form of a manuscript. The second part of this chapter deals with the inducibility of this migration process by blue/green light and the influence of chemical substances on the mat. This part of the chapter is presented in form of a short communication.




Figure 2: Image of marine *Beggiatoa* cultivated under chemolithotrophic conditions without nitrate at a high sulfide flux ($43.1 \text{ mmol m}^{-2} \text{ d}^{-1}$) after two weeks. A subpopulation of filaments migrated downwards into deeper layers.

Contributions:

2.1 Sulfur respiration in a chemolithoautotrophic marine *Beggiatoa* strain

A. Schwedt, A.-C. Kreutzmann, L. Polerecky and H. N. Schulz-Vogt

The concept of the study was developed by me and H. N. Schulz-Vogt. All initial experiments were performed by me. The final experiments and data analysis were performed by me, A.-C. Kreutzmann and L. Polerecky. The manuscript was written together with A.-C. Kreutzmann and with the help of the other two co-authors.

2.2 Coordinated movement of *Beggiatoa* filaments in oxygen/sulfide gradients and the effect of blue/green light

H. N. Schulz-Vogt, T. Hohmann, A.-C. Kreutzmann, L. Polerecky and A. Schwedt

The concept of the study was developed by H. N. Schulz-Vogt and T. Hohmann. Major experiments were performed by H. N. Schulz-Vogt and T. Hohmann with the help of L. Polerecky during data analysis. I assisted during migration experiments and filming and performed c-di-GMP experiments.

2.1 Sulfur respiration in a marine chemolithoautotrophic *Beggiatoa* strain

Anne Schwedt^{1*}, Anne-Christin Kreutzmann¹, Lubos Polerecky¹, Heide N. Schulz-Vogt¹

¹Max Planck Institute for Marine Microbiology, Celsiusstr. 1, D-28359 Bremen, Germany

*Corresponding author

Manuscript submitted to *Frontiers in Microbiology* (22.09.2011), accepted with minor revisions (23.12.2011), and published online (09.01.2012). doi:10.3389/fmicb.2011.00276

Keywords: *Beggiatoa*; sulfur reduction; gradient cultivation; microelectrodes; migration

Abstract

The chemolithoautotrophic strain *Beggiatoa* sp. 35Flor shows an unusual migration behavior when cultivated in a gradient medium under high sulfide fluxes. As common for *Beggiatoa* spp., the filaments form a mat at the oxygen-sulfide interface. However, upon prolonged incubation, a subpopulation migrates actively downwards into the anoxic and sulfidic section of the medium, where the filaments become gradually depleted in their sulfur and polyhydroxyalkanoates (PHA) inclusions. This depletion is correlated with the production of hydrogen sulfide. The sulfur- and PHA-depleted filaments return to the oxygen-sulfide interface, where they switch back to depositing sulfur and PHA by aerobic sulfide oxidation. Based on these observations we conclude that internally stored elemental sulfur is respired at the expense of stored PHA under anoxic conditions. Until now, nitrate has always been assumed as the alternative electron acceptor in lithotrophic *Beggiatoa* spp. under anoxic conditions. As our medium and the filaments were free of oxidized nitrogen compounds we can exclude this metabolism. Furthermore, sulfur respiration with PHA under anoxic conditions has so far only been described for heterotrophic *Beggiatoa* spp., but our medium did not contain accessible organic carbon. Hence the PHA inclusions must originate from atmospheric CO₂ fixed by the filaments while at the oxygen-sulfide interface. We propose that the directed migration of filaments into the anoxic section of an oxygen-sulfide gradient system is a strategy to preserve cell integrity, which could be compromised by excessive sulfur deposition occurring in the presence of oxygen and high sulfide fluxes. The regulating mechanism of this migration is hitherto unknown.

Introduction

The genus *Beggiatoa* comprises large, filamentous bacteria that inhabit diverse sulfidic environments, such as sediments (Winogradsky, 1887; Jørgensen, 1977; Nelson and Castenholz, 1982; McHatton *et al.*, 1996), springs (Winogradsky, 1887; Macalady *et al.*, 2006) and activated sludge (Farquhar and Boyle, 1971). The motile filaments typically aggregate in a narrow overlapping zone of opposed oxygen and sulfide diffusion gradients where they form a sharply bounded mat (Faust and Wolfe, 1961; Nelson and Jannasch, 1983; Nelson *et al.*, 1986a). Within this mat, *Beggiatoa* spp. oxidize sulfide with oxygen and deplete both compounds (Nelson *et al.*, 1986a). This process is accompanied by deposition of elemental sulfur inside the filaments.

Several filamentous and non-filamentous members of the *Beggiatoaceae* (Salman *et al.*, 2011) are moreover capable of anaerobic sulfide oxidation with nitrate as an alternative electron acceptor (Fossing *et al.*, 1995; McHatton *et al.*, 1996; Schulz *et al.*, 1999). Dissimilatory nitrate reduction enables these organisms to colonize anoxic environments such as deeper layers in sediments, microbial mats or gradient cultures (Sweerts *et al.*, 1990; Mußmann *et al.*, 2003; Sayama *et al.*, 2005; Kamp *et al.*, 2006; Hinck *et al.*, 2007). Nitrate-based sulfide oxidation seems to have been of great importance for some members of the family *Beggiatoaceae*, as suggested by their ability to store nitrate within intracellular vacuoles at concentrations up to 10^4 fold higher than in the ambient water (Fossing *et al.*, 1995; McHatton *et al.*, 1996; Schulz *et al.*, 1999; Sayama, 2001; Mußmann *et al.*, 2003; Kalanetra *et al.*, 2004; Kalanetra *et al.*, 2005; Hinck *et al.*, 2007). However, also non-vacuolated strains were shown to use externally provided nitrate as a terminal electron acceptor (Sweerts *et al.*, 1990; Kamp *et al.*, 2006).

We cultivated the chemolithoautotrophic, marine strain *Beggiatoa* sp. 35Flor in an agar-stabilized oxygen-sulfide gradient medium. Upon prolonged incubation in the presence of medium to high sulfide fluxes, we observed an unusual migration behavior, where a subpopulation of filaments migrated downwards from the oxygen-sulfide interface. These filaments were able to survive although sulfide concentrations were high and terminal electron acceptors that are known to be utilized by *Beggiatoa* spp., i.e., oxygen and nitrate, were not detectable in medium nor filaments. In this study, we investigated the possibility of an alternative metabolism of *Beggiatoa* sp. 35Flor under anoxic, nitrate-free and sulfidic conditions, and discuss its possible ecological significance and link to the peculiar migration behavior.

Material and methods

Strain and cultivation

The strain *Beggiatoa* sp. 35Flor was originally enriched from a black band disease of scleractinian corals from the coast of Florida, and can so far only be cultivated in the presence of the *Pseudovibrio denitrificans* strain FO-BEG1 (Schwedt *et al.*, unpublished). Filaments of the strain 35Flor are about 6 µm wide, and the cells contain polyphosphate inclusions and a central vacuole filled with polyphosphate (Kamp *et al.*, 2008; Brock and Schulz-Vogt, 2011).

Cultivation was performed in tubes with an agar-based mineral gradient medium modified after Nelson *et al.* (1982) and Nelson and Jannasch (1983) using artificial seawater (Kamp *et al.*, 2008). The medium was composed of a sulfidic bottom agar plug (1.5% w/v agar) covered with a sulfide-free, semisolid top agar layer (0.25% w/v agar) of 5 cm height (Tables 2.1.1 and 2.1.2). The medium was prepared free of nitrate, nitrite and nitric oxide, as verified by measurements with an NO_x analyzer (CLD 66, Eco Physics, Rösrath, Germany). Gas exchange between headspace and the atmosphere was possible, and opposing gradients of oxygen and sulfide were allowed to form for one day before inoculation. The cultures were inoculated about 1 cm below the air-agar interface using 100 µL filament suspension from an established mat. Incubations were performed at room temperature in the dark.

During incubations, the distribution of filaments in the tube was measured simultaneously with the vertical profiles of H₂S and pH. Filaments from parallel culture tubes were subsampled and used for microscopic determination of their sulfur and PHA inclusions. Oxygen profiles were measured in parallel tubes.

Table 2.1.1 Solutions for the preparation of agar-stabilized gradient media.

Solution	Composition ^a
Artificial seawater	27.5 g NaCl, 5 g MgCl ₂ · 6H ₂ O, 4.1 g MgSO ₄ · 7H ₂ O, 0.66 g CaCl ₂ · 2H ₂ O, 1.02 g KCl
Mineral solution	555 mg K ₂ HPO ₄ , 28.72 mg Na ₂ MoO ₄ , 750 mg Na ₂ S ₂ O ₅ , 29 mg FeCl ₃ · 6H ₂ O
Trace element solution ^b	5.2 g EDTA, 1.5 g FeCl ₂ · 4H ₂ O, 70 mg ZnCl ₂ , 100 mg MnCl ₂ · 4H ₂ O, 62 mg H ₃ BO ₄ , 190 g CoCl ₂ · 6H ₂ O, 17 mg CuCl ₂ · 2H ₂ O, 24 mg NiCl ₂ · 6 H ₂ O, 36 mg Na ₂ MoO ₄ · 2H ₂ O
Vitamin solution ^c	0.1 g cyanocobalamine 0.1 g inositol 0.1 g biotin 0.1 g folic acid 1.0 g para-aminobenzoic acid 10 g nicotinic acid 10 g D-pantothenate 20 g thiamine

^a in 1 L distilled water

^b pH adjusted to 6.5

^c All vitamins were dissolved separately and then combined in a final stock solution (1 mL of each vitamin solution in a final volume of 100 mL distilled water) that was filtersterilized twice.

Table 2.1.2 Composition of gradient medium.

Medium	Component ^a	Composition
Bottom agar	A	100 mL artificial seawater, 2.9 g NaCl, 1 drop 1 mol L ⁻¹ KOH
	B	80 mL distilled water, 2.7 g agar ^b
	C	0.72 – 3.6 mL 1 mol L ⁻¹ Na ₂ S solution
Top agar	A	240 mL artificial seawater, 4.32 g NaCl
	B	96 mL distilled water, 0.9 g agar ^b
	C	24 mL mineral solution, 0.36 ml trace element solution, 7 drops 1 mol L ⁻¹ KOH
	D	0.72 mL 1 mol L ⁻¹ NaHCO ₃ solution
	E	150 µL vitamin solution

^a All components were sterilized separately before combination.

^b The agar was washed two times in distilled water before use.

Microsensor measurements

Microsensors for O₂ (OX-10 standard), H₂S (H2S-10), and pH (PH-10) were purchased from Unisense A/S (Aarhus, Denmark). The external reference for the pH electrode was manufactured and connected in-house. Calibration of the H₂S sensor was performed in anoxic, acidified artificial seawater (Table 2.1.1, pH<2) to which an anoxic Na₂S stock solution was added stepwise. The exact sulfide concentration of the Na₂S stock solution was determined by iodometric titration. Total sulfide (S_{tot}) profiles were calculated from measured H₂S and pH profiles using equation $S_{tot} = H_2S \times [1 + K_1 / H_3O^+]$, with pK₁ = 6.569 at 21°C and 39‰ salinity (Millero *et al.*, 1988). The oxygen sensor was two-point calibrated in a calibration chamber filled with artificial seawater. Signal readings were taken in water saturated with N₂ and ambient air. Oxygen concentrations at the respective salinity and temperature were calculated according to Weiss (1970). The pH electrode was calibrated using buffer solutions of pH 4.01, pH 7.00, and pH 9.21 (Mettler-Toledo, Giessen, Germany). All sensors were calibrated immediately before the measurement. In case of long time series measurements the sensor calibration was checked afterwards and a possible drift was corrected for.

Vertical profiling in 250 µm steps was performed with sensors mounted on a motorized linear positioner (VT-80, Pollux motor, Micos, Eschbach, Germany) controlled by a computer using a software for automated microsensor measurements (µ-Profiler, L. Polerecky, <http://www.microsen-wiki.net>). The sensors were aligned by manually adjusting their tips to the air-agar interface using a dissecting microscope (Stemi 2000-C, Zeiss, Jena, Germany).

Filament imaging

The distribution of sulfur-containing *Beggiatoa* sp. 35Flor filaments in the gradient cultures was monitored using time-lapse photography. An amber light-emitting diode (LXHL-NM98, Luxeon, Philips, San Jose, CA, USA) was positioned below the culture tube and switched on for one second when an image was taken with a cooled CCD camera (Sensicam, PCO, Kelheim, Germany). Illumination and image acquisition in 10 min intervals were controlled by a computer using a custom-written program (Look@Molli, B. Grunwald, <http://www.microsen-wiki.net>).

Intensities of the recorded images were horizontally averaged over an area with visible filaments (~5 mm wide, ~2 cm high), and the resulting vertical profiles were assembled into a 2D map with the x-axis representing incubation time and the y-axis corresponding to depth. Since the average image intensity was proportional to the density of sulfur globules, which were present exclusively inside filaments, vertical movement of sulfur-rich filaments was detected as a change in the shape of the vertical intensity profile. In contrast, an increase and decrease in the profile intensity that was not accompanied with the change in the profile shape indicated accumulation and depletion of sulfur inside the filaments, respectively. Because this method relied on light scattering from sulfur inclusions, it did not allow visualization of sulfur-free filaments.

Staining of internal PHA

Staining with Nile Red was used to visualize PHA inclusions in the filaments. A subsample from the culture tube (volume 90 μL) was incubated for 5 minutes with 10 μL of a Nile Red (Sigma-Aldrich, Steinheim, Germany) staining solution (25 mg L^{-1} in dimethyl sulfoxide). The filament suspension was transferred onto a poly-L-lysine (Sigma-Aldrich) coated microscope slide for immobilization of the filaments. Fluorescence of Nile Red was excited with a laser at 546 nm and emission was recorded above 590 nm (filter set 15, Zeiss, Jena, Germany) using an epifluorescence microscope (Axiophot equipped with AxioCam MRm, Zeiss, Jena, Germany).

Transfer experiment with sulfur-free filaments

To verify that sulfur-free filaments from the anoxic subpopulation of an aged culture (cultivated at high sulfide flux conditions) were alive, able to migrate back to the oxygen-sulfide interface and re-establish their sulfide-oxidizing metabolism, they were transferred

into the anoxic section of a fresh gradient medium (cultivated at low sulfide flux conditions). All cultivation media were prepared in plexiglass tubes (2×12 cm in size) with lateral holes (Brock and Schulz-Vogt, 2011). Fresh medium for inoculation with sulfur-free filaments was pre-incubated with the accompanying *Pseudovibrio* strain. This was done to ensure a sufficient cell density of *Pseudovibrio* sp. irrespective of the inoculum as the *Pseudovibrio* sp. is required for growth of *Beggiatoa* sp. 35Flor, but its abundance is negligible in the anoxic part of the gradient (Schwedt *et al.*, unpublished). Subsequently, sulfur-free filaments were removed laterally from the aged culture and injected laterally into the fresh medium at a depth of about 1 cm below the oxygen-sulfide interface. Lateral removal ensured that no sulfur-containing filaments from the oxygen-sulfide interface of the aged culture were transferred, whereas lateral injection prevented inoculation of the transferred sulfur-free filaments to the oxygen-sulfide interface of the fresh gradient media. The development of a mat at the oxygen-sulfide interface was inspected visually.

Results

Migration of *Beggiatoa* sp. 35Flor in gradient cultures

Beggiatoa sp. 35Flor filaments aggregated and formed a dense mat at the oxygen-sulfide interface within the gradient medium. In cultures with medium to high sulfide fluxes (Table 2.1.3 A) a subpopulation of filaments began a downward migration to the anoxic zone about 3–4 days after establishment of the mat. For medium sulfide fluxes, this migration resulted in a layer with a homogenous filament density extending up to 2–3 mm below the mat (Figure 2.1.1). In contrast, for high sulfide fluxes the migrating filaments were not homogeneously distributed, but progressively aggregated in a region distinctly separated from the mat at the oxygen-sulfide interface (Figure 2.1.1 and 2.1.2 B). Because the aggregation of filaments in the anoxic part increased the chance of detecting metabolic products all further experiments were conducted with cultures growing under a high sulfide flux.




Figure 2.1.1: Distribution of *Beggiatoa* sp. 35Flor filaments over depth in gradient cultures after 6 (open symbols) and 12 (closed symbols) days in the presence of different sulfide fluxes. The flux values represent theoretical maxima under the given cultivation conditions (Table 2.1.3 A).

Migration of filaments in cultures with a high sulfide flux followed a general pattern (Figure 2.1.2 E). During the initial 3–4 days of incubation, the mat at the oxygen-sulfide interface gradually formed. After about 6–7 days, the sulfur globule density in the mat decreased moderately, followed by a more pronounced decrease after 8–9 days. These decreases were correlated with two pronounced events of downward migration at days 5–6 and 7–8, respectively (arrows 1 and 2 in Figure 2.1.2 E). After reaching a depth of around 10 mm, the migrating filaments formed a layer of increased filament density. These filaments slowly disappeared from view due to a gradual loss of their internal sulfur granules. The disappearance of filaments was accompanied by a parallel increase in the sulfur globule density in the mat at the oxygen-sulfide interface (arrow 3 in Figure 2.1.2 E), suggesting that the filaments returned to this zone and switched back to sulfide oxidation, thereby depositing sulfur. This was confirmed by the transfer experiment, which showed that sulfur-free filaments transferred from the anoxic subpopulation of an aged culture into the anoxic section of a fresh gradient medium formed, within 12 days, a new mat of sulfur-containing filaments at the oxygen-sulfide interface.

Table 2.1.3 A Diffusive sulfide fluxes in gradient cultures from this study.

c(Na ₂ S) [mmol L ⁻¹] in bottom agar		Flux [mmol m ⁻² d ⁻¹]
4 (low flux)	theoretical maximum ^a	4.6
	measured (7 d)	8.0
	measured (13 d)	8.9
10 (medium flux)	theoretical maximum ^a	19.0
	measured (7 d)	20.1
	measured (13 d)	16.2
16 (high flux)	theoretical maximum ^a	40.1
	measured (7 d)	31.6
	measured (13 d)	25.3

^a The initial theoretical maximum of the sulfide flux in the gradient cultures, calculated using Fick's first law of diffusion ($J = -D \Delta c / \Delta x$). The diffusion coefficient D for HS⁻ was corrected for temperature (21°C) according to Jørgensen and Revsbech (1983), resulting in a value of $1.56 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}$. The concentration gradient was calculated from the height of the top agar ($\Delta x = 5 \text{ cm}$) and the initial sulfide concentration in the bottom agar, assuming that sulfide was depleted at the agar surface, i.e. $\Delta c = c(\text{Na}_2\text{S})$.

Table 2.1.3 B Diffusive sulfide fluxes in natural *Beggiatoa* spp. mats.

Sediment from	Reference	Flux [mmol m ⁻² d ⁻¹]
Lagoon	(Jørgensen and Revsbech, 1983)	38
Lagoon	(Mußmann <i>et al.</i> , 2003)	0.8
Arctic lagoon	(Jørgensen <i>et al.</i> , 2010)	34
Coast	(Preisler <i>et al.</i> , 2007)	4.3 ± 2
Harbor	(Fenchel and Bernard, 1995)	12-100
Deep sea mud volcano	(Lichtsclag <i>et al.</i> , 2010)	11.6
Deep sea mud volcano	(Girnth <i>et al.</i> , 2011)	40
Deep sea, gas hydrate-rich	(Sahling <i>et al.</i> , 2002)	63 ± 36

Sulfide production by filaments in the anoxic section

Throughout the incubation, sulfide oxidation in the mat at the oxygen-sulfide interface was confirmed by pronounced acidification and steep gradients of total sulfide (Figure 2.1.2 C and D). A small but detectable peak in the H₂S profile was observed at a depth of ~10 mm when the anoxic subpopulation was present (Figure 2.1.2 D). As pH varied only smoothly with depth in this region, the H₂S peak was not linked to pH variation, but indicated a true production of sulfide at and around this depth. This production was strongly spatially and

temporally correlated with the presence of the anoxic subpopulation (Figure 2.1.2 F), suggesting that it was linked to the metabolic activity of the filaments from this subpopulation.




Figure 2.1.2: Relationship between the migration of *Beggiatoa* sp. 35Flor filaments and the dynamics of O_2 , pH, H_2S and S_{tot} in the gradient culture tube. **(A-B)** Images of culture tubes showing the filament distribution after 2 and 11 days, respectively. **(C-D)** Examples of pH, H_2S and total sulfide profiles in the gradient culture incubated for 8 and 13 days. Shaded areas mark the oxic zone. **(E)** Average sulfur globule density as a function of time and depth, showing the dynamics of the filament distribution and their sulfur content. Arrows 1 and 2 indicate the onset of major downward migration events, arrow 3 indicates the onset of an increase in the filament density in the mat at the oxygen-sulfide interface. Although the timing of these events varied amongst experimental runs, the general pattern was reproducible. **(F)** H_2S excess as a function of time and depth, calculated by subtracting the measured H_2S profile from the background trend. The trend was derived from the H_2S concentrations measured above and below the peak (line indicated by arrow in panel D). Contour lines of the sulfur globule density from panel E are overlaid. Data shown in panels A, B, E and F are from the same culture tube, profiles in panels C and D are from a parallel culture tube.

Sulfur and PHA content in single filaments

Beggiatoa sp. strain 35Flor filaments accumulated elemental sulfur and PHA during growth at the oxic-anoxic interface. Sulfur inclusions were visible as dark, highly refractory globules in bright-field micrographs, while PHA inclusions appeared as strongly fluorescent globules in the images of Nile Red stained samples (Figure 2.1.3). When grown under high sulfide fluxes, most filaments from the mat at the oxygen-sulfide interface were densely filled with sulfur and PHA inclusions (Figure 2.1.3 B and 2.1.3 E). With increasing sulfide fluxes the amount of internal sulfur strongly increased (compare Figure 2.1.3A and 2.1.3 B), whereas PHA inclusions were equally abundant in all treatments (data not shown). In contrast, filaments from the anoxic subpopulation were heterogeneous with respect to their inclusion density; while some were densely filled with sulfur and PHA, others lacked both (Figure 2.1.3 D and 2.1.3 F). At high sulfide fluxes, filaments were frequently observed to burst, particularly in older cultures (Figure 2.1.3 C).




Figure 2.1.3: Bright-field and fluorescence micrographs illustrating the typical appearance of *Beggiatoa* sp. 35Flor filaments cultivated under different conditions. (A-B) Filaments from the mat at the oxygen-sulfide interface of 6 days old cultures growing under low (A) and high (B) sulfide flux conditions. (C-D) Filaments from cultures grown under high sulfide flux conditions, collected from the mat at the oxygen-sulfide interface after 27 days (C) and from the anoxic subpopulation after 12 days (D). (E-F) Nile Red-stained filaments from a 14 days old culture, collected from the mat at the oxygen-sulfide interface (E) and from the anoxic subpopulation (F). Bright fluorescence in panel E originates from PHA inclusions, whereas faded fluorescence in panel F is due to staining of the cell membrane.

Discussion

Sulfide production by members of the genus *Beggiatoa* is known from chemoheterotrophic strains that were cultivated in liquid medium and artificially exposed to short-term anoxic conditions (Schmidt *et al.*, 1987). Based on those experiments it was hypothesized that sulfur respiration may provide *Beggiatoa* spp. in gradient systems with energy for return from the anoxic zone to the oxygen-sulfide interface under changing environmental conditions. In this study, we cultivated the chemolithoautotrophic strain *Beggiatoa* sp. 35Flor in an oxygen-sulfide gradient medium, and we observed a directed migration of the filaments from the oxygen-sulfide interface into the anoxic and sulfidic zone where they reduced internal sulfur to sulfide. This suggested an alternative or additional function of sulfur respiration in *Beggiatoa* filaments.

We propose that the observed behavior is a survival strategy of *Beggiatoa* sp. 35Flor at prolonged incubation under high sulfide fluxes. Under this condition the filaments become densely filled with sulfur and were often observed to burst. By moving to the anoxic zone of the gradient system, the filaments can prevent further deposition of sulfur through aerobic sulfide oxidation and may even reduce the amount of storage compounds by sulfur respiration with PHA. We observed that filaments can migrate back to the oxygen-sulfide interface, where they resume aerobic sulfide oxidation and accumulate new sulfur globules.

Sulfur respiration for regulation of the amount of stored sulfur

The alteration between sulfide oxidation and sulfur reduction in spatially separated environments seems to allow *Beggiatoa* sp. 35Flor to control the amount of stored sulfur beyond the scope of enzymatic regulation. Sulfide is oxidized by *Beggiatoa* spp. in a two-step process via internally stored sulfur ($2 \text{H}_2\text{S} + \text{O}_2 \rightarrow 2 \text{S}^0 + 2 \text{H}_2\text{O}$) and further to sulfate ($2 \text{S}^0 + 3 \text{O}_2 + 2 \text{H}_2\text{O} \rightarrow 2 \text{SO}_4^{2-} + 4 \text{H}^+$). The regulation of these reactions is unknown in *Beggiatoa* spp., but the presence of internal sulfur globules demonstrates that the rates of the two reactions are not always well balanced. In principle, a balanced sulfur content can be achieved by either down-regulating sulfide oxidation or up-regulating sulfur oxidation. It is likely that sulfide oxidation is controlled kinetically and cannot be regulated by the cell, because both O_2 and H_2S are freely diffusing into the cytoplasm. This is supported by observations on two related genera *Thiomargarita* and *Marithioploca*, which both immediately increased their respiration rate upon addition of sulfide to the medium (Schulz and de Beer, 2002; Høglund *et al.*, 2009). Therefore, up-regulation of the sulfur oxidation

seems to be the more likely mechanism for balancing the internal sulfur content. However, at high sulfide fluxes bursting of *Beggiatoa* sp. 35Flor filaments densely filled with sulfur globules indicates that further up-regulation of sulfur oxidation was not possible, e.g. due to enzymatic rate limitation.

As an alternative to enzymatic regulation, the filaments may leave the overlapping zone of oxygen and sulfide in order to starve themselves of electron donor or acceptor, thereby interrupting sulfur deposition. A negative chemotactic response to oxygen (Møller *et al.*, 1985) presumably prevented the filaments from moving upwards into the oxic section of the gradient system. Instead, they migrated downwards into the anoxic and sulfidic section, where sulfide could no longer be oxidized to sulfur due to the lack of an electron acceptor. These filaments moved into the sulfidic zone, which is surprising, because elevated sulfide concentrations have previously been reported to be toxic for *Beggiatoa* spp. (Winogradsky, 1887; Keil, 1912; Nelson *et al.*, 1986a). However, all earlier studies were done under oxic conditions. Our study indicates that *Beggiatoa* can tolerate higher sulfide concentrations under anoxic conditions, whereas under oxic conditions high sulfide concentrations can cause cell death indirectly by inducing excessive sulfur accumulation.

Metabolism of *Beggiatoa* in the anoxic zone of gradient systems

The depletion of sulfur and polyhydroxyalkanoate inclusions together with the production of sulfide suggests that *Beggiatoa* sp. 35Flor reduced internal sulfur by oxidizing stored carbon in the anoxic part of the gradient system. It is not known which type of PHA was synthesized by *Beggiatoa* sp. 35Flor, but for the most frequent PHA, poly (3-hydroxybutyrate) (PHB), the reaction ($[C_4O_2H_6] + n \cdot 9 S^0 + n \cdot 6 H_2O \rightarrow n \cdot 4 CO_2 + n \cdot 9 H_2S$), which is pH-neutral, would be in agreement with the observed pH profiles. By reducing stored sulfur with stored PHA that derived from previously fixed CO₂, the filaments do not exploit an additional energy source in the anoxic environment. Instead, they use this process as the only possibility to empty storage space.

Presence of filamentous *Beggiatoaceae* in the anoxic section of oxygen-sulfide gradient systems has so far been shown in multiple laboratory and field studies (Jørgensen, 1977; Sweerts *et al.*, 1990; Mußmann *et al.*, 2003; Kamp *et al.*, 2006; Hinck *et al.*, 2007; Jørgensen *et al.*, 2010). However, in these systems either external or internal nitrate was present and could have been used for oxidizing sulfide in the anoxic zone of the sediment. Nitrate respiration could, however, be excluded in our experiments as NO_x compounds were absent from the medium and filaments.

The role of sulfur reduction by *Beggiatoa* spp. in the environment

The migration behavior and sulfur reduction by *Beggiatoa* filaments described in our study may occur and play the same role also in natural habitats. In the environment, filaments could respond to high sulfide fluxes either by moving laterally to an adjacent region with a lower sulfide flux or, if this is not possible, by migrating vertically to the sulfidic and anoxic sediment section below, where they respire sulfur (Figure 2.1.4). This is supported by the fact that sulfide fluxes in our culture tubes (Table 2.1.3 A) were in the range of fluxes measured in different natural *Beggiatoa* mats (Table 2.1.3 B), and that similar heterogeneity in internal sulfur content of *Beggiatoa* filaments was also observed for filaments collected from natural mats (Sassen *et al.*, 1993; Bernard and Fenchel, 1995). However, the conditions at which these phenomena occur will depend on the possible maximum oxidation rates of sulfide and ultimately sulfur, which likely define the tolerance of different *Beggiatoa* species towards high sulfide fluxes.




Figure 2.1.4: Proposed function of sulfur reduction as a survival strategy of *Beggiatoa* sp. 35Flor under high sulfide fluxes. In locations with high sulfide fluxes (right side) *Beggiatoa* filaments become excessively filled with sulfur (black dots inside the filaments), because the two oxidation rates of sulfide to sulfur (K_1) and sulfur to sulfate (K_2) are not well balanced ($K_1 > K_2$). To prevent bursting, filaments can move into a region with a lower sulfide flux (black arrow) where these two reactions may proceed in a balanced way. If this is not possible, filaments leave the oxygen-sulfide interface and move down into an anoxic region to reduce their internal sulfur deposits and thus prevent bursting (white arrow). They do so by using internally stored PHA as an electron donor to reduce S^0 to H_2S . After emptying the storage space, the filaments return to the oxygen-sulfide interface and continue with sulfide oxidation.

Acknowledgements

We thank Martina Meyer for technical support. The study was funded by the European Research Council (No 203364-ELNOX), the Max Planck Society and the German National Merit Foundation.

2.2 Coordinated movement of *Beggiatoa* filaments in oxygen-sulfide gradients and the effect of blue/green light

Heide N. Schulz-Vogt^{1*}, Tine Hohmann¹, Anne-Christin Kreutzmann¹, Lubos Polerecky¹,
Anne Schwedt¹

¹Max Planck Institute for Marine Microbiology, Celsiusstr. 1, D-28359 Bremen, Germany

*Corresponding author

Manuscript in preparation

Keywords: migration; blue/green light; *Beggiatoa*

Abstract

Filamentous sulfide-oxidizing bacteria of the genus *Beggiatoa* are gradient organisms. When grown autotrophically in opposing gradients of oxygen and sulfide, the filaments establish a thin and well-defined mat in the overlapping zone of oxygen and sulfide. We found that cyclic Adenosine-mono-phosphate (cAMP) and cyclic diguanylate (c-di-GMP) or blue light can modify or disturb this typical behavior.

Introduction

Large filamentous sulfur bacteria of the genus *Beggiatoa* form a thin mat in the overlapping zone of oxygen and sulfide (Nelson *et al.*, 1986a), where they oxidize the upwards diffusing sulfide with oxygen (e.g. Winogradsky, 1887; Keil, 1912; Jørgensen, 1977; Nelson and Castenholz, 1981). The formation of a thin mat between steep opposing gradients of oxygen and sulfide is thought to be the result of toxicity of both compounds for these bacteria (Winogradsky, 1887; Keil, 1912; Møller *et al.*, 1985). Furthermore, a migrational reaction was also observed when filaments were illuminated with light (Winogradsky, 1887; Nelson and Castenholz, 1982; Møller *et al.*, 1985), suggesting that light may also play a role in the migration of *Beggiatoa* during mat formation. In this study, we found that, in addition to light, chemical substances such as cyclic Adenosine-mono-phosphate (cAMP) or cyclic diguanylate (c-di-GMP) also influence the migrational behavior and mat formation by *Beggiatoa*. Our results suggest that light might directly lead to a coordinated migration of a subpopulation of filaments and that cAMP and c-di-GMP induce a dispersion of the mat.

Results and discussion

Beggiatoa filaments of strain 35Flor were cultivated in oxygen-sulfide gradient media (chapter 2.1) where they typically form a distinct mat in the overlaying zone of oxygen and sulfide. We found that several factors can modify or disturb the typical mat formation behavior of *Beggiatoa* filaments. The previously formed mat is disturbed and filaments distribute randomly throughout the gradient medium when cAMP and c-di-GMP are added to the culture (Figure 2.2.1). When grown under high sulfide fluxes ($> 40 \text{ mmol m}^{-2} \text{ d}^{-1}$), a part of the population from the mat in the oxygen-sulfide overlapping zone migrates down into the anoxic and sulfidic zone after an incubation time of 10–14 days. These filaments were shown to use sulfur as an electron acceptor for the oxidation of PHA, both stored internally during the time spent in the oxygen-sulfide overlapping zone (Chapter 2.1). The downward migration was proposed as the last resort strategy for the filaments to prevent cell disruption, which

would otherwise occur due to excessive accumulation of internal sulfur by sulfide oxidation. A very similar migration behavior was observed when the previously formed *Beggiatoa* mat was illuminated by light in the blue to green region (Fig. 2.2.2). When cyan light is applied to a four days old culture incubated at a high sulfide flux ($40 \text{ mmol m}^{-2} \text{ d}^{-1}$) a downward migration of a subpopulation of filaments can be observed immediately (Figure 2.2.2). In contrast to the migration in the dark, that occurs after 10-14 days, the migration can be induced with light even after incubation times of a few days.




Figure 2.2.1: *Beggiatoa* sp. 35Flor filament distribution at a low sulfide flux ($10.8 \text{ mmol m}^{-2} \text{ d}^{-1}$) is shown in A. The addition of $20 \mu\text{l}$ of c-di-GMP ($0.01 \mu\text{mol L}^{-1}$; B) or $10 \mu\text{l}$ cAMP (100 mM ; C) leads to the disruption of the mat starting after 2 hours (here shown after 8 hours). When Phosphate buffer or water was added there was no migrational reaction of the *Beggiatoa* filaments (data not shown).




Figure 2.2.2: Average sulfur globule density of *Beggiatoa* filaments at a high sulfide flux as a function of time and depth (calculation see Chapter 2.1). The culture was pre-incubated for 4 days in the dark where the mat at the oxygen sulfide interface established. The imaging started in the dark (black bar). After 4 hours the culture was illuminated with cyan light (cyan bar) and a subpopulation of filaments immediately migrated downwards.

The data presented here show that the known signalling molecules also have an effect on an established *Beggiatoa* mat and can lead to migrational reactions of the filaments. The mat is disturbed and the filaments distribute randomly in the culture tube. Therefore, it is possible that these substances might be used as signalling molecules among *Beggiatoa* filaments. In contrast to this, a directed migration known from cultures with a high sulfide flux (Chapter 2.1) can be induced by applying blue/green light to the culture, indicating that light might be involved in the migration of *Beggiatoa* filaments.

Acknowledgements

We thank Martina Meyer for technical support. The study was funded by the European Research Council (No 203364-ELNOX) and the Max Planck Society.

Chapter 3

Co-cultivation of a marine

Beggiatoa strain and *Pseudovibrio* sp.

In this chapter, the association between the large *Beggiatoa* sp. 35Flor and the accompanying *Pseudovibrio* sp. FO-BEG1 (Figure 3) is investigated. The first part of this chapter is a study on the co-culture of the two organisms written as a short communication. In order to study the association from the side of the *Pseudovibrio* sp, the genome of this bacterium was analyzed in detail. This second part of this chapter comprises an abstract about the genomic analysis and comparison of the isolated *Pseudovibrio* sp. and another *Pseudovibrio* sp. strain associated with a sponge. Both strains are able to grow in pure culture and both genomes reveal a versatile metabolism of the bacteria. During the genome analysis of *Pseudovibrio* sp. strain FO-BEG1 we found several genes encoding for more than 20 superoxide dismutases, catalases and peroxidases additionally to genes for the interaction with other cells. The manuscript to this second part of this chapter is added as an appendix to the thesis.




Figure 3: Differential interference contrast micrograph of a *Beggiatoa* sp. and *Pseudovibrio* sp. co-culture.

Contributions:

3.1 A chemolithoautotrophic *Beggiatoa* strain requiring the presence of a *Pseudovibrio* sp. for cultivation

A. Schwedt and H. N. Schulz-Vogt

The concept of this study was developed by both authors. I performed all experiments. The manuscript was written by me with the help of H. N. Schulz-Vogt.

3.2 The *Pseudovibrio* genus contains metabolically versatile and symbiotically interacting bacteria

V. Bondarev, M. Richter, J. Piel, S. Romano, A. Schwedt, and H. N. Schulz-Vogt

The study was initiated by V. Bondarev, H. N. Schulz-Vogt and me. Cultivation and DNA extraction for sequencing of the genome was performed by me. I coordinated the sequencing of the genome. Genome analysis was performed by V. Bondarev with assistance of M. Richter. J. Piel did NRPS and PKS cluster analysis. Cultivation experiments were performed by V. Bondarev. The manuscript was written by V. Bondarev with assistance of all co-authors.

3.1 A chemolithoautotrophic *Beggiatoa* strain requiring the presence of a *Pseudovibrio* sp. for cultivation

Anne Schwedt* and Heide N. Schulz-Vogt

Max Planck Institute for Marine Microbiology, Celsiusstr. 1, D-28359 Bremen, Germany

*Corresponding author

Manuscript in preparation

Keywords: co-cultivation; gradient medium; oxidative stress; catalase

Abstract

Numerous attempts to isolate larger chemolithoautotrophic members of the family *Beggiatoaceae* into pure culture have so far met little success. Here, we report the successful cultivation of a marine *Beggiatoa* sp. strain of 6 μm in diameter, which exclusively grows in the presence of a *Pseudovibrio* sp. strain.

Introduction

Beggiatoa spp. are colorless sulfur bacteria forming mats at the oxic-anoxic interface where they oxidize sulfide with oxygen (e. g. Winogradsky, 1887; Keil, 1912; Jørgensen, 1977; Nelson and Castenholz, 1981). Smaller chemolithoautotrophic strains (diameter $< 5 \mu\text{m}$) have been isolated into pure culture by using an agar based medium with opposing gradients of sulfide and oxygen (Nelson and Jannasch, 1983; Nelson *et al.*, 1986a). Besides these strains, only heterotrophic freshwater *Beggiatoa* spp. strains were obtained in pure culture (Strohl and Larkin, 1978). The reason for the general difficulty to isolate larger chemolithoautotrophic members of the family *Beggiatoaceae* in pure culture is still unknown. We obtained a marine *Beggiatoa* sp. strain of 6 μm in diameter into a stable culture, but did not succeed in growing it axenically. Apart from all attempts to clean the *Beggiatoa* sp. 35Flor a single type of organism, which was identified as *Pseudovibrio* sp. FO-BEG1 remained. Here, we present experiments indicating that the presence of this accompanying organism is required for the growth of the *Beggiatoa* sp. strain.

Results and discussion

Under higher sulfide fluxes ($> 40 \text{ mmol m}^{-2} \text{ d}^{-1}$) a subpopulation of the *Beggiatoa* filaments actively migrated into lower and anoxic parts of the culture tube (Figure 3.1) where they reduced the amount of internal sulfur by sulfur reduction (Chapter 2.2). As the accompanying *Pseudovibrio* sp. FO-BEG1 is predominantly found in the oxic part of the culture tube (Figure 3.1), these lower filaments were used to inoculate fresh culture tubes with only *Beggiatoa* filaments and relatively free of the accompanying *Pseudovibrio* sp. strain.

When *Beggiatoa* filaments from the mat at the oxic-anoxic interface were transferred into fresh medium a new mat developed in 4 out of 5 tubes. However, if filaments from the anoxic subpopulation with no or very few *Pseudovibrio* cells were transferred into fresh medium, only in 1 out of 7 tubes a mat developed. Examining this single mat-containing tube under the microscope, besides *Beggiatoa* filaments also *Pseudovibrio* cells were observed. When

filaments from the anoxic part were transferred into tubes containing already a *Pseudovibrio* population, most of the tubes were positive for growth of *Beggiatoa* sp. (5 positive tubes out of 7 tubes). Even another *Pseudovibrio* strain from the culture collection (*Pseudovibrio denitrificans* DSM number 17465) had a similar effect on growth (4 positive tubes out of 7). Therefore, we conclude that the growth of the *Beggiatoa* sp. depends on the presence of an accompanying bacterium of the genus *Pseudovibrio*.




Figure 3.1: Distribution of the accompanying *Pseudovibrio* sp. strain FO-BEG1 (red squares) in co-culture with the marine *Beggiatoa* sp. strain 35Flor (photograph, white filaments). The *Beggiatoa* filaments form a mat at the oxic-anoxic interface and a subpopulation migrates into the anoxic part of the tube. The accompanying *Pseudovibrio* strain is mainly present above the upper *Beggiatoa* mat in the oxic part of the culture tube.

There are a few indications about the possible reasons for a positive effect of the accompanying *Pseudovibrio* sp. on the growth of the large sulfide-oxidizer. *Pseudovibrio* cells populate mainly the oxic part of the culture tube, but *Beggiatoa* filaments also thrive well in the anoxic zone (Figure 3.1). Thus, it seems reasonable to assume that *Pseudovibrio* sp. is only required during the oxic metabolism of the *Beggiatoa* filaments. It may be suggested that *Pseudovibrio* sp. helps the *Beggiatoa* sp. in detoxifying oxygen radicals, as *Beggiatoa* typically do not possess the ability to produce the enzyme catalase (Larkin and Strohl, 1983). Moreover, earlier studies have shown a positive effect of catalase on the growth of *Beggiatoa*

spp. (Burton and Morita, 1964; Strohl and Larkin, 1978; Güde *et al.*, 1981; Nelson *et al.*, 1986b). In contrast, the accompanying *Pseudovibrio* strain is catalase-positive and the genome suggests further potential to detoxify oxygen radicals (Chapter 3.2). Therefore, we propose that the *Beggiatoa* sp. benefits from the oxygen protection system of the accompanying *Pseudovibrio* sp. strain and is consequently mandatory for the growth of the large sulfide-oxidizer.

Acknowledgements

This study was funded by the European Research Council (No 203364-ELNOX) and the Max Planck Society.

3.2 The *Pseudovibrio* genus contains metabolically versatile and symbiotically interacting bacteria

Vladimir Bondarev^{1*}, Michael Richter³, Jörn Piel², Stefano Romano, Anne Schwedt¹ and Heide N. Schulz-Vogt¹

¹Max Planck Institute for Marine Microbiology, Celsiusstr. 1, D-28359 Bremen, Germany

²Kekulé-Institut für Organische Chemie und Biochemie, Gerhard-Domagk Str. 1, D-53121 Bonn, Germany

³Ribocon GmbH, Fahrenheitsstr. 1, D-28359 Bremen, Germany

*Corresponding author

Manuscript in preparation, the whole manuscript is attached to this thesis in the appendix.

Keywords: Physiology; Genome; NRPS / PKS; colibactin; Secretion system; *Beggiatoa*

Abstract

The majority of strains belonging to the genus *Pseudovibrio* have been isolated from marine invertebrates like tunicates, corals and especially sponges, but the physiology of these bacteria is poorly understood. In this study, we analyze the genomes of two *Pseudovibrio* strains. One is a required symbiont of a cultivated *Beggiatoa* strain, a sulfide oxidizing, autotrophic bacterium. The other one was isolated from a sponge (Enticknap *et al.*, 2006). The data show that both strains are generalistic bacteria capable of importing and oxidizing a wide range of organic and inorganic compounds to meet their carbon, nitrogen, phosphorous and energy requirements under oxic and anoxic conditions. Several physiological traits encoded in the genome were verified in laboratory experiments with a pure culture of the *Pseudovibrio* strain originally associated with *Beggiatoa*. Besides the versatile metabolic abilities of both *Pseudovibrio* strains, our study reveals a number of open reading frames and gene clusters in the genomes that seem to be involved in symbiont-host interactions. *Pseudovibrio* has the genomic potential to attach to host cells, might be capable of interacting with the eukaryotic cell machinery, produce secondary metabolites and may supply the host with cofactors.

Chapter 4

Isolation and cultivation of *Pseudovibrio* sp. and other facultatively oligotrophic bacteria

The fourth chapter includes a detailed physiological investigation of the growth and substrate use of the isolated *Pseudovibrio* sp. FO-BEG1 under oligotrophic conditions. These results are presented in the first manuscript of this chapter. The investigated *Pseudovibrio* strain is on the one hand able to grow under extremely oligotrophic conditions and on the other hand it can grow on organic-rich media (Figure 4). Eventually, we were able to isolate further facultatively oligotrophic bacteria from Namibian sediments, which are highly populated by large sulfur bacteria. The isolation and cultivation of these bacteria belonging to different phylogenetic groups is presented in the second manuscript of this chapter.



Figure 4: Growth of *Pseudovibrio* sp. FO-BEG1 on organic-rich agar plates at day 2 forming colonies (left) and in pure artificial seawater (right).

Contributions:

4.1 Substrate use of *Pseudovibrio* sp. growing in extremely oligotrophic seawater

A. Schwedt, M. Seidel, T. Dittmar, M. Simon, V. Bondarev, S. Romano, G. Lavik, and H. N. Schulz-Vogt

I developed the concept of this study with assistance of H. N. Schulz-Vogt, M. Seidel and T. Dittmar. Furthermore, I performed the medium preparation and sampling with the help of M. Seidel. FT-ICR-MS measurements were performed together with M. Seidel. HPLC analysis was performed by the group of M. Simon. G. Lavik assisted during the measurement and analysis of the isotope-labelling experiments. V. Bondarev performed glucose and ammonia addition experiments and S. Romano the biolog experiment. I wrote the manuscript with assistance of the co-authors.

4.2 Facultatively oligotrophic bacteria isolated from the habitat of large sulfide-oxidizers

A. Schwedt, V. Bondarev, M. Seidel, T. Dittmar, H. N. Schulz-Vogt

This study was initiated by me with the help of H. N. Schulz-Vogt. I performed all isolations and cultivations. V. Bondarev assisted me with counting of the cells, M. Seidel and T. Dittmar measured DOC concentrations. The manuscript was written by me including comments of all co-authors.

4.1 Substrate use of *Pseudovibrio* sp. growing in extremely oligotrophic seawater

Anne Schwedt^{1*}, Michael Seidel^{1,2}, Thorsten Dittmar^{1,2}, Meinhard Simon², Vladimir Bondarev¹, Stefano Romano¹, Gaute Lavik¹, Heide N. Schulz-Vogt¹

¹Max Planck Institute for Marine Microbiology, Bremen, Germany

²Institute of Chemistry and Biology of the Marine Environment, Carl von Ossietzky University of Oldenburg, Germany

*Corresponding author

Manuscript in preparation

Keywords: low nutrient medium; detergents; amino acids; oligotrophy; *Pseudovibrio*

Abstract

Marine planktonic bacteria often live in habitats with extremely low concentrations of dissolved organic matter (DOM). To study the use of trace amounts of DOM by the facultatively oligotrophic bacterium *Pseudovibrio* sp. strain FO-BEG1, we investigated the composition of artificial and natural seawater before and after growth. For this, we determined dissolved organic carbon (DOC), total dissolved nitrogen (TDN), composition of DOM by electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry (ESI FT-ICR-MS) and amino acids by high performance liquid chromatography (HPLC). The DOC concentration of artificial seawater was 0.06 mg C L^{-1} ($5 \text{ } \mu\text{mol C L}^{-1}$), which was an order of magnitude below the concentration in the natural seawater (0.9 mg C L^{-1} or $75 \text{ } \mu\text{mol C L}^{-1}$). DOC concentrations did not decrease measurably during growth. Cell numbers increased from about 20 cells mL^{-1} to $20,000 \text{ cells mL}^{-1}$ in artificial and to $800,000 \text{ cells mL}^{-1}$ in natural seawater. No nitrogen fixation and minor CO_2 fixation ($< 1\%$) was observed. In both media, amino acids were not the major nutrient. Instead, we observed a decrease of compounds resembling detergents in artificial seawater, which also contained nitrogen. In natural seawater, we detected a decrease of two groups of compounds, one resembling amino- and thiosugars, which are also rich in nitrogen, and another group containing condensed hydrocarbons. The present study shows that heterotrophic bacteria can grow with even lower DOC concentrations than available in natural oligotrophic seawater and may use unexpected organic compounds to fuel their energy, carbon and nitrogen requirements.

Introduction

In open oceans, the amount of dissolved organic matter (DOM) is typically indicated as the concentration of dissolved organic carbon (DOC) and is generally below 1 mg C L^{-1} (Schut *et al.*, 1997; Hansell *et al.*, 2009). Consequently, marine bacteria are commonly exposed to very low concentrations of organic material. Marine DOM is an extremely diverse pool of different compounds, consisting of more than ten thousand types of molecules with different reactivities (Dittmar and Paeng, 2009). Previous studies demonstrated that most of the DOM in the ocean is inert to bacterial break-down and cannot be utilized by marine microorganisms (Bada and Lee, 1977; Ammerman *et al.*, 1984; Zweifel *et al.*, 1993). Measurements of the consumption of specific substrates revealed that a large fraction of the labile organic material that is used in the upper ocean can consist of dissolved free amino acids (DFAA), dissolved combined amino acids (DCAA) and monosaccharides. These substances can account for 5 to

100% of the bacterial carbon and nitrogen demand (Fuhrman, 1987; Jørgensen, 1987; Stanley *et al.*, 1987; Cowie and Hedges, 1992; Keil and Kirchman, 1999; Cherrier and Bauer, 2004). In many natural systems, amino acids are only present in nanomolar concentrations even though the flux of amino acids is comparably high due to a close coupling of their release and uptake by planktonic microorganisms (Fuhrman, 1987; Fuhrman, 1990; Keil and Kirchman, 1999). Still, amino acids and monosaccharides alone cannot fully explain the growth of the bacteria in all cases, where they represent only a small fraction of the C- and N-requirements (Fuhrman, 1987; Stanley *et al.*, 1987). The remaining demand must be covered by different DOM compounds, which might be present at only very low concentrations. Therefore, to specifically investigate the compounds consumed by the bacteria during growth under oligotrophic conditions, a method detecting a broad spectrum of different organic compounds with a low detection limit, such as electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry (ESI FT-ICR-MS), is necessary.

In contrast to natural systems where several processes supply nutrients simultaneously, a batch culture containing a single strain of bacteria is a closed system with defined amounts of nutrients. In such a system, it is possible to analyze compounds that are utilized during the growth of this strain. For the cultivation of marine bacteria, defined artificial seawater or natural seawater containing a diverse DOM pool can serve as a medium. In both seawater media, the risk of contamination especially with amino acids is a well known problem (Dittmar *et al.*, 2009). In the present study, we chose to investigate the substrate use of the marine *Pseudovibrio* sp. strain FO-BEG1, because it is capable to grow under oligotrophic conditions in extremely pure artificial seawater. In spite of special purification of the artificial seawater medium, a DOC concentration of 0.06 mg C L⁻¹ (5 μmol C L⁻¹) was measured, which is, nevertheless, low compared to the natural oligotrophic seawater with a DOC concentration of 0.9 mg C L⁻¹ (75 μmol C L⁻¹). We compared the substrate use in artificial seawater with natural oligotrophic seawater. In both media, DOC concentrations did not decrease measurably during growth of the bacteria. The composition of the DOM before and after growth in artificial and natural seawater was analyzed by high performance liquid chromatography (HPLC) measuring amino acids and by ESI FT-ICR-MS revealing changes on the level of individual intact molecules indicated by molecular formulas. The combination of these methods with isotope-labeling techniques to study incorporation of N₂ and CO₂ provides information of compounds used by the bacteria during growth under these extremely oligotrophic conditions.

Material and methods

Medium preparation and bacterial strain

The basal artificial oligotrophic seawater medium is composed of 30.3 g NaCl, 3.3 g $\text{MgCl}_2 \cdot 6 \text{H}_2\text{O}$, 2.8 g $\text{MgSO}_4 \cdot 2 \text{H}_2\text{O}$, 0.44 g $\text{CaCl}_2 \cdot 2 \text{H}_2\text{O}$, 0.7 g KCl, 50 μl 2 mol L^{-1} NaOH, 2 mL 1 mol L^{-1} NaHCO_3 , 1 mL 1 mmol L^{-1} K_2HPO_4 and 1 mL trace elements solution in 1 L MembraPure water (Optilab-Standard Water System, MembraPure, Bodenheim, Germany). The trace elements solution was composed of 2.1 mg L^{-1} $\text{FeSO}_4 \cdot 7 \text{H}_2\text{O}$, 13 mL L^{-1} HCl (25%), 60 mg L^{-1} H_3BO_3 , 100 mg L^{-1} $\text{MnCl}_2 \cdot 4 \text{H}_2\text{O}$, 190 mg L^{-1} $\text{CoCl}_2 \cdot 6 \text{H}_2\text{O}$, 24 mg L^{-1} $\text{NiCl}_2 \cdot 6 \text{H}_2\text{O}$, 2 mg L^{-1} $\text{CuCl}_2 \cdot 2 \text{H}_2\text{O}$, 144 mg L^{-1} $\text{ZnSO}_4 \cdot 7 \text{H}_2\text{O}$ and 36 mg L^{-1} $\text{Na}_2\text{MoO}_4 \cdot 2 \text{H}_2\text{O}$. All glassware, Teflon stoppers, screw caps and tubes were acid washed (0.1 mol L^{-1} HCl, Merck, Darmstadt, Germany) and rinsed with MembraPure water (Optilab-Standard Water System, MembraPure, Bodenheim, Germany). Glassware, aluminium foil and NaCl were combusted at 480°C for at least 3 hours. Serum bottles (Wheaton 125 mL serum bottles clear, Wheaton, Millville, NJ, USA) were used for cultivation, filled completely with medium and closed with Teflon stoppers. Before bottling of the medium, the artificial seawater was cleaned by solid phase extraction of dissolved organic matter (SPE-DOM, Dittmar *et al.*, 2008), i.e. organic contaminations. Incubations were performed without shaking at 28°C in the dark. For glucose and ammonium addition experiments the oligotrophic artificial seawater medium was not specifically purified and glucose and ammonium were added at a final concentration of 30 $\mu\text{mol L}^{-1}$. These cultures were incubated gently shaken (shaker GFL-3013, Gesellschaft für Labortechnik mbH, Burgwedel, Germany) at 28°C in the dark.

Natural surface seawater was collected in the South Pacific during Integrated Ocean Drilling (IODP) Expedition 329 from IODP Site U1368 (27.9°S ; 137.9°W) with a bucket and stored at 4°C. Before incubation, seawater was filtered through washed Acrodisc 25 mm syringe filters with a 0.2 μm GHP membrane (Pall Life Sciences, Ann Arbor, MI, USA) and filled into pre-combusted serum bottles that were closed with Teflon stoppers. Inoculated cultures were incubated without shaking at 28°C in the dark.

Media were inoculated with 100 μL of an oligotrophically pre-grown culture (about 2×10^4 cells mL^{-1}) of *Pseudovibrio* sp. FO-BEG1. Thus, the initial cell number after inoculation accounts to 4 to 20 cells mL^{-1}

Cell counts

Cells were stained with 1:5000 SYBR Green (SYBR Green I 10,000×, Sigma, Taufkirchen, Germany) for 20 minutes. The stained cells were filtered onto black filters (GTTP Isopore Membrane Filters 0.22 µm, Millipore, Schwalbach/Ts., Germany) using a Bio-Dot apparatus (Bio-Rad, München, Germany). Counting of cells was performed using a fluorescence microscope (Axiophot, Zeiss, Jena, Germany) at 450–490 nm excitation and 515–565 nm emission (filter set 10, Zeiss, Jena, Germany).

Sampling and solid phase extraction of dissolved organic matter (SPE-DOM)

About 900 mL medium (six serum bottles, 156 mL each) were pooled at four different time points (blank, t_0 , t_1 , t_2). The blank was taken directly after medium preparation without inoculation and inoculated samples were taken before (t_0), after 1 week (t_1) and after 3 weeks (t_2) of incubation. Samples were filtered into 1 L glass bottles using Acrodisc 25 mm syringe filters with a 0.2 µm GHP membrane (Pall Life Sciences, Ann Arbor, MI, USA) and acidified to pH 2 with 25% HCl. Samples for DOC measurements and amino acid analysis were taken after pooling the six serum bottles into a 1 L glass bottle. DOC samples were measured directly and samples for amino acid analysis were stored at -20°C . All samples were stored at 4°C until further analysis. From all four time points, DOM was extracted using the solid phase extraction of dissolved organic matter (SPE-DOM) method (Dittmar *et al.*, 2008). The extracts of dissolved organic matter were analyzed using ESI FT-ICR-MS (see below).

Measurement of dissolved organic carbon (DOC) and total dissolved nitrogen (TDN)

Dissolved organic carbon (DOC) and total dissolved nitrogen (TDN) were analyzed before, during and after growth of the bacteria using a Shimadzu TOC-VCPH total organic carbon analyzer (Shimadzu, Kyoto, Japan) equipped with a TNM-1 total nitrogen measuring unit and an ASI-V autosampler. To remove cells prior to analysis, samples were filtered through Acrodisc 25 mm syringe filters with a 0.2 µm GHP membrane (Pall Life Sciences, Ann Arbor, MI, USA). Within the instrument, samples were acidified with 1% v/v 2 mol L⁻¹ HCl and sparged with synthetic air for 2 minutes to remove inorganic carbon. Detection limits were 0.5 µmol L⁻¹ for DOC and TDN (0.006 mg C L⁻¹ and 0.007 mg N L⁻¹, respectively). Analytical errors based on the standard deviations for replicated measurements (at least three measurements per sample) were within 5% for DOC and TDN. Analytical precision and accuracy was tested in each run against deep Atlantic seawater reference material and low carbon water provided by the consensus reference materials program (D.A. Hansell,

University of Miami, Coral Gables, FL, USA). Procedural blanks, including the filtration step, were obtained with ultrapure water.

Amino acids analysis

The concentrations of dissolved free amino acids (DFAA) and total hydrolysable dissolved amino acids (THDAA) were analyzed using high performance liquid chromatography (HPLC) as described earlier (Lunau *et al.*, 2006). Derivatization was performed using an ortho-phthalaldehyde precolumn (Lindroth and Mopper, 1979). All samples were filtered through prewashed Acrodisc 25 mm syringe filters with a 0.2 μm GHP membrane (Pall Life Sciences, Ann Arbor, MI, USA). Prior to the analysis of THDAA, HCl was added to a final concentration of 1.7 mol L⁻¹ and incubated for 1 hour at 155°C under nitrogen atmosphere. The concentration of dissolved combined amino acids (DCAA) was calculated by subtracting the concentration of DFAA from the concentration of THDAA.

H¹³CO₃ and ¹⁵N₂ labeling experiments

For labeling experiments, oligotrophic artificial seawater medium was prepared as described above, however, DOM was not extracted prior to use and it was prepared under synthetic air atmosphere (20% O₂ in N₂; H₂O < 3 ppm-mol, C_nH_m < 0.1 ppm-mol, CO < 1 ppm-mol, CO₂ < 1 ppm-mol; Air Liquide, Krefeld, Germany). From each 156 mL serum bottle, which was completely filled with medium, 5 mL medium were exchanged with 5 mL ¹⁵N₂ gas and 300 μL of 1 mol L⁻¹ NaH¹³CO₃ were added. After inoculation with *Pseudovibrio* sp. FO-BEG1 (100 μL of a pre-culture containing about 2×10^4 cells mL⁻¹), samples were incubated for 12 hours, 3 and 5 days. Then the bacteria were filtered onto pre-combusted Glass Microfibre Filters (GF/F, Whatman, GE Healthcare, Dassel, Germany), frozen for 12 hours and dehydrated in an HCl desiccator. Filters were folded into tin cups and flash combusted (1050°C) within the mass spectrometer to release N₂ and CO₂. For analysis, we used an elemental analyzer (with autosampler) coupled to a Delta Plus Advantage mass spectrometer isotope-ratio monitoring mass spectrometer (ThermoFinnigan, Bremen, Germany). To calculate the amount of label in the cells the percentage of label added was taken into account.

Fourier transform ion cyclotron resonance mass spectrometry (FT-ICR-MS)

SPE-DOM extracts were diluted 1:1 with ultrapure water to yield DOC concentrations of 10 ppm for natural seawater and about 2 ppm for artificial seawater incubations, respectively. Diluted samples were analyzed on a solariX Fourier transform ion cyclotron resonance mass

spectrometer (FT-ICR-MS; Bruker Daltonik GmbH, Bremen, Germany) connected to a 15 Tesla superconducting magnet (Bruker Biospin, Wissembourg, France). Samples were infused at a rate of $2 \mu\text{L min}^{-1}$ into an electrospray source (ESI; Apollo II ion source, Bruker Daltonik GmbH, Bremen, Germany) with the capillary voltage set to 4 kV in negative mode and 4.5 kV in positive mode. Ions were accumulated in the hexapole for 0.3 seconds prior to transfer into the ICR cell. Data acquisition was done in broadband mode using 4 megaword data sets and a scanning range of 150–2000 Da. The instrument was externally calibrated with an in-house marine deep sea DOM standard (mass accuracy of less than 0.1 ppm). Before each sample set, blank checks with methanol/ultrapure water 1:1 were measured. For one mass spectrum, 500 scans were added and internally calibrated with the in-house marine deep sea DOM standard to yield a mass accuracy of less than 0.1 ppm for the internal calibrants (series of $\text{C}_x\text{H}_y\text{O}_z$ compounds from 281 to 621 Da). The molecular formulas were calculated in the mass range between 150 and 850 Da by applying the following restrictions: $^{12}\text{C}_{0-130}^{1}\text{H}_{0-200}^{16}\text{O}_{1-50}^{14}\text{N}_{0-4}^{32}\text{S}_{0-2}^{31}\text{P}_{0-2}$ (considering also ^{23}Na clusters in ESI positive mode). Assignment of molecular formulas was done with the Bruker software DataAnalysis 4.0 SP 4 using the criteria described by Koch *et al.* (2005). Molecular formulas were considered valid when the mass error was below 0.5 ppm. Only compounds with a signal-to-noise ratio of 3 and higher were used for further analysis. The achieved resolving power (full width half maximum) was on average 480,000 at m/z 400.

Relative peak heights were calculated by normalization to the ten most abundant peaks in each spectrum (triplicate measurements). These peak heights were then used to semi-quantitatively assess changes in DOM composition in natural oligotrophic seawater experiments. FT-ICR-MS is not a quantitative tool for DOM analysis due to the lack of standards and differences in ionization efficiencies for different compounds. However, this method yields reproducible results because the variability of relative peak heights is low under similar analytical conditions (Kido Soule *et al.*, 2010).

Sequential standard addition was applied to quantify the amount of polyethylene glycol (PEG) contamination in the artificial seawater medium using an authentic PEG standard (octaethylene glycol, $\text{C}_{16}\text{H}_{34}\text{O}_9$, $\geq 99\%$ oligomer purity, 370 g mol^{-1} , Sigma, Taufkirchen, Germany). PEG was added in concentrations of 1 to 3.6 nmol L^{-1} and FT-ICR mass spectra were recorded in ESI positive mode. To yield a linear response range in the MS, ion accumulation time was lowered to 0.1 seconds and 50 scans were added for one mass

spectrum. The amount of PEG in the samples was calculated from a linear calibration derived from peak heights in FT-ICR mass spectra. To investigate the suppression effect of the inorganic contamination on organic peaks during FT-ICR-MS, the in-house natural deep sea DOM standard was added in the range of 0.04 to 4 mg C L⁻¹ to a sample contaminated with the inorganic substances and measured in ESI negative mode.

Biolog experiment

A substrate respiration test was performed using a biolog GN2 plate (Hayward, CA, USA) with 95 different substrates. The bacteria were pre-grown in artificial seawater medium in 1 L bottles (Schott, Mainz, Germany). After 6 days, cells were concentrated by centrifugation for 2 hours at 11,000 × g and 15°C using a J-26XP Beckman centrifuge (Beckman Coulter GmbH, Krefeld, Germany). All centrifugation tubes were acid washed (0.1 mol L⁻¹ HCl) and rinsed with water. The obtained pellet was suspended in 20 mL of sterile saline solution (23.5 g L⁻¹ NaCl and 10.6 g L⁻¹ MgCl₂ · 6 H₂O). To verify the viability of the cells, 50 µL of the cell suspension was spread on organic-rich medium containing 2 g polypeptone, 0.5 g Bacto yeast extract (BD Diagnostics, Heidelberg, Germany), 30 g NaCl, 5 g MgCl₂ · 6 H₂O, 0.005 g CaCl₂ · 2 H₂O, 0.005 g Na₂MoO₄ · 7 H₂O, 0.004 g CuCl₂ · 2 H₂O, 0.006 g FeCl₃ · 6 H₂O, 15 g bactoagar in 1 L MembraPure water and the pH was adjusted to 8 with 1 mol L⁻¹ NaOH. The biolog plate was inoculated with 150 µL of the concentrated cells and incubated at 28°C in the dark in a humidity chamber to prevent excess evaporation. Activity of cells was checked visually each day for a total of 14 days.

Results

Growth under oligotrophic conditions

The investigated *Pseudovibrio* sp. strain FO-BEG1 was able to multiply in both artificial and natural seawater. The initial cell number was about 20 cells mL⁻¹ and in artificial seawater cell numbers increased to 2 × 10⁴ cells mL⁻¹ at time points t₁ and t₂ (one and three weeks after inoculation, respectively). Natural seawater medium contained about 8 × 10⁵ cell mL⁻¹ at both time points. In both media, the increase in cell numbers was detected during the first week of incubation (t₀ to t₁) and cell numbers did not change in the following two weeks (t₁ to t₂).

Dissolved organic carbon (DOC), total dissolved nitrogen (TDN) and dissolved free and combined amino acids (DFAA and DCAA)

In pure artificial seawater medium before bottling, the DOC contamination from salts was only $1 \mu\text{mol C L}^{-1}$. During the bottling of the medium and with the addition of NaHCO_3 , K_2HPO_4 , NaOH and trace elements, the concentration of DOC increased to about $4.5 \mu\text{mol C L}^{-1}$ (Table 4.1.1). The TDN was below limit of detection ($0.5 \mu\text{mol N L}^{-1}$) in the sterile artificial medium before and after bottling (Table 4.1.1). Inoculation did not introduce further measurable DOC or TDN to the medium (t_0). After one (t_1) and three weeks (t_2) of incubation, the concentrations of DOC and TDN did not change significantly in the medium (Table 4.1.1). In comparison to the artificial seawater medium, the DOC concentration in natural seawater was about 16 times higher ($71 \mu\text{mol L}^{-1}$; Table 4.1.1). However, no significant change in DOC concentration was detectable during growth, as DOC concentrations one and three weeks after inoculation were 78.5 and $75 \mu\text{mol C L}^{-1}$, respectively. A similar pattern was observed for TDN concentrations, which were around $12 \mu\text{mol N L}^{-1}$ at all time points (Table 4.1.1).

In contrast to DOC and TDN, the amount of DFAA decreased in the artificial seawater cultures within the first week of growth (from t_0 to t_1). This decrease represents a DFAA uptake of $0.06 \mu\text{mol C L}^{-1}$ and $0.02 \mu\text{mol N L}^{-1}$ (Table 4.1.1). However, the concentrations of DFAA remained constant until t_2 . The most abundant amino acids were serine, glycine and alanine, which together accounted for $0.05 \mu\text{mol C L}^{-1}$. The decrease in DFAA (from t_0 to t_1) was concurrently observed with an increase of DCAA. During the stationary growth phase (from t_1 to t_2), the DCAA concentrations decreased again (Table 4.1.1). Opposed to the artificial seawater incubations, the amino acids concentrations (THDAA) in natural seawater were very low and no change during incubation could be observed (Table 4.1.1).

Nitrogen and carbon dioxide fixation

The incorporation of labeled nitrogen ($^{15}\text{N}_2$) and carbon ($\text{NaH}^{13}\text{CO}_3$) into *Pseudovibrio* sp. FO-BEG1 growing in pure artificial seawater medium was examined using isotope-ratio monitoring mass spectrometry. Less than 1% of cellular carbon originated from carbon dioxide fixation under the oligotrophic growth conditions in artificial seawater. Nitrogen fixation was not detectable.

Table 4.1.1: Artificial and natural seawater medium: Amounts of dissolved organic carbon (DOC) and total dissolved nitrogen (TDN), dissolved free amino acids (DFAA), dissolved combined amino acids (DCAA) or total hydrolysable dissolved amino acids (THDAA) at the different time points (before bottling: only medium; blank: medium poured into serum bottles before inoculation; t_0 : directly after inoculation; t_1 : 1 week after inoculation; t_2 : 3 weeks after inoculation). (n.d. = not determined)

	Time point	DOC [$\mu\text{mol L}^{-1}$]	TDN [$\mu\text{mol L}^{-1}$]	C in	N in	C in	N in
				DFAA [$\mu\text{mol L}^{-1}$]	DFAA [$\mu\text{mol L}^{-1}$]	DCAA [$\mu\text{mol L}^{-1}$]	DCAA [$\mu\text{mol L}^{-1}$]
Artificial seawater	before bottling	1	< 0.5		n.d.		n.d.
	blank	4.5	< 0.5				
	t_0	4.5	< 0.5	0.08	0.03	0.05	0.02
	t_1	4.5	< 0.5	0.02	0.01	0.12	0.05
	t_2	4.5	< 0.5	0.02	0.01	0.06	0.02
				C in THDAA [$\mu\text{mol L}^{-1}$]		N in THDAA [$\mu\text{mol L}^{-1}$]	
Natural seawater	t_0	71	12		0.02		0.01
	t_1	78.5	13		0.02		0.01
	t_2	75	11		0.02		0.01

Composition of the dissolved organic matter

The analysis of extracted DOM from artificial and natural seawater using FT-ICR-MS was performed in the negative and positive electrospray ionization (ESI) mode. In ESI negative mode, 690 of 7347 peaks (10%) in artificial and 3685 of 14873 peaks (25%) in natural seawater were assigned with molecular formulas (Figure 4.1.2). However, in negative ESI mode inorganic contaminants (as indicated by a negative mass defect) suppressed ionization of organic molecules. To estimate this suppression effect, natural deep sea SPE-DOM was added to the artificial seawater extracts in the same, tenfold and hundredfold concentration (0.04, 0.4 and 4 mg C L⁻¹). The DOM signal was completely suppressed after addition of 0.04 mg C L⁻¹ (same range as the measured DOC contamination in the artificial seawater). In contrast, the typical distribution of natural DOM peaks in the mass spectrum was detected after addition of 0.4 and 4 mg C L⁻¹ natural DOM (ten- and hundredfold the DOC concentration in artificial seawater). As a consequence, FT-ICR-MS spectra of the artificial

seawater medium had to be analyzed in ESI positive mode, in which the inorganic contaminants were not detected.

The mass spectra obtained after positive ionization revealed two homologous series of high peaks with distances of 44.02567 Da ($C_2H_4O_1$) corresponding to a homologous series of polyethylene glycol (PEG) oligomers (Figure 4.1.1 A). The PEG peaks were present in the mass spectra of all sampled time points. Therefore, the mean intensity of five PEG peaks (m/z 305.157061, 349.183268, 393.209484, 437.235707, 481.261932) was used to normalize all other peaks in the mass spectra (Figure 4.1.1 A, arrows). The amount of PEG contamination was quantified by standard addition to the contaminated SPE-DOM blanks from artificial seawater using an authentic PEG standard (Figure 4.1.1 B and C). The total amount of PEG in the sample was then calculated to account for only 3 nmol $C L^{-1}$, which is three orders of magnitude below the measured concentration of the DOC contamination in the artificial seawater medium (5 $\mu\text{mol } C L^{-1}$). Thus, the PEG contamination was extremely low despite the very high peaks in the ESI FT-ICR mass spectra (Figure 4.1.1). A screening for potential PEG contaminations indicated NaOH as a likely source (data not shown).




Figure 4.1.1: Polyethylene glycol (PEG) contamination in artificial seawater: (A) FT-ICR mass spectra measured from the artificial seawater extracts (ESI positive mode) showing high peaks of PEG contamination with intervals of 44.02567 Da. Vertical arrows indicate the peaks used for normalization. (B) FT-ICR mass spectra after standard addition of PEG in the range of 0 to 3.6 nmol L^{-1} to the extracted DOM of the blank sample. (C) Linear calibration curve of PEG standard after addition to blank sample.




Figure 4.1.2: FT-ICR mass spectra of the negative electrospray ionization at the time points t_0 of (A) artificial seawater and (B) natural seawater. Asterisks indicate inorganic contaminations. Inserts show a zoom into the region of mass 325.0 to 325.2. In the negative mode inorganic contaminations suppressed the ionization of organic substances present in low amounts in the artificial seawater.

In van Krevelen diagrams, organic compounds are plotted based on their oxygen to carbon (O/C) and hydrogen to carbon (H/C) ratios. During the incubation in artificial seawater (from t_0 to t_2) the relative intensities of compounds with an $O/C \leq 0.4$ and $H/C \geq 1.5$ decreased strongly, while the relative intensity of only few compounds increased (Figure 4.1.3 A). After growth of the bacteria (from t_0 to t_2) in artificial seawater, many compounds with an $O/C \leq 0.4$ and $H/C \geq 1.5$ were not detected anymore. Most of the compounds that disappeared had molecular formulas containing nitrogen (Figure 4.1.3 B and C). No significant difference was found between the procedural blank of the artificial seawater and t_0 or between t_1 and t_2 (blank and t_0 before, t_1 and t_2 after growth).

Major changes in incubations with natural seawater during growth (from t_0 to t_2) were found in two clusters of compounds: (i) compounds with $O/C \leq 0.4$ and $H/C \leq 1.0$, and (ii) compounds with $O/C \geq 0.7$ and $HC \geq 1.25$ (Figure 4.1.3 D, blue). Compounds of cluster (i) with low O/C and H/C ratios contained sulfur but no nitrogen (Figures 4.1.3 E and F, yellow). Compounds of cluster (ii) with high O/C and H/C ratios contained mainly nitrogen and sulfur (Figures 4.1.3 E and F, red).




Figure 4.1.3: Van Krevelen diagrams (O/C = oxygen to carbon ratio and H/C = hydrogen to carbon ratio of the molecular formulas of single molecules detected) showing the change in DOM composition from t_0 to t_2 in artificial (**A-C**) and natural seawater (**D-F**). (**A+D**): Changes in normalized relative intensities are color-coded showing a decrease in blue, an increase in red and no significant change during incubation in grey. (**B+C, D+F**): Compounds containing no nitrogen and sulfur (purple), sulfur but no nitrogen (yellow), nitrogen but no sulfur (blue) and both nitrogen and sulfur (red) at t_0 (**B+E**) and t_2 (**C+F**). In artificial seawater the most notable decrease is found in the fraction of compounds containing high H/C and low O/C ratios and nitrogen but no sulfur. These compounds may correspond to detergents. In natural seawater a group of substances with high ratios of H/C and O/C (probably thio- and aminosugars) containing mostly nitrogen and sulfur decreased. In addition, compounds with low H/C and O/C ratios (probably condensed hydrocarbons) containing only sulfur were also observed to decrease.

Glucose and ammonium addition experiments

Cell numbers in artificial seawater medium without glucose and ammonium were about 3×10^5 cells mL⁻¹. The separate addition of glucose and ammonium did not lead to a strong increase in cell numbers, whereas the simultaneous addition of glucose and ammonium increased cell numbers to about 10^7 cells mL⁻¹ (Figure 4.1.4).




Figure 4.1.4: Growth curves of *Pseudovibrio* sp. strain FO-BEG1 incubated in artificial seawater without the addition of a fixed carbon or nitrogen source or with addition of ammonium or glucose. Addition of each substance alone did not lead to an increase in cell numbers. When both (ammonium and glucose) were added at the same time, cell numbers increased significantly.

Biolog experiment

Pseudovibrio sp. FO-BEG1 cells pre-grown in artificial seawater medium were used to inoculate a biolog GN2 microplate containing 95 different growth substrates. The bacteria showed activity after 8 to 14 days on the following substrates: D-raffinose, D-trehalose, turanose, D-gluconic acid, D-glucosaminic acid, D-glucuronic acid, acetic acid, glucoronamide, L-glutamic acid, glycyl-L-glutamic acid, L-serine and D-glucose-6-phosphate. No activity was observed on the remaining 84 substrates.

Discussion

In the present study, the heterotrophic *Pseudovibrio* sp. strain FO-BEG1 was grown either in artificial seawater medium with extremely low DOC concentrations or in natural seawater from an oligotrophic ocean (south Pacific, surface water) representing typical marine DOM. The bacterial growth was sustained by very low amounts of DOC in both incubation setups. However, growth in artificial seawater with only 0.06 mg DOC per liter seawater shows that bacterial growth can be sustained by even lower concentration as present in most natural oligotrophic seawaters, which typically contain less than 1 mg C L⁻¹ (Schut *et al.*, 1997; Hansell *et al.*, 2009).

Impurities in oligotrophic media

During the preparation of oligotrophic medium, different potential contamination risks emerge. In our case, an inorganic contamination was introduced to the artificial seawater after the incubation during the solid phase extraction of DOM (SPE-DOM). This led to ion suppression of the organic compounds in the FT-ICR mass spectra in ESI negative mode. Consequently, it was not possible to analyze the DOM composition in that mode and we chose the positive mode instead. This problem does not occur in natural seawater samples, because the concentration of DOC is higher. We could show that the organic substances present in the artificial medium are detectable if the overall background of DOM is 10 to 100 times higher. This observation indicates that a specific concentration of organic matter has to be present to be detected by ESI FT-ICR-MS if inorganic substances are introduced in high amounts during extraction or ionize extremely well.

In ESI positive mode, an organic contamination of the artificial seawater was observed. Two series of polyethylene glycol (PEG) oligomers were detected. As a potential source NaOH used for medium preparation was identified. Since PEGs ionize extremely well in ESI positive mode, the peaks are much higher compared to most other peaks in the sample. Using the standard addition method (described in material and methods), the total amount of PEG contaminations was estimated to be 1000fold less than the total DOC concentration in artificial seawater, and thus represented only a very small fraction of the entire DOC contamination. These impurities analyzed in the artificial seawater medium were not detected in the SPE-DOM of natural seawater samples and thus did not affect the FT-ICR-MS analysis of natural seawater incubations.

Growth and substrate use in artificial seawater

During growth in artificial seawater under extremely oligotrophic conditions, *Pseudovibrio* sp. strain FO-BEG1 multiplied from about 20 cells mL⁻¹ to 2 × 10⁴ cells mL⁻¹, even though the overall amount of DOC did not measurably decrease. Thus, the amount of compounds that were consumed was probably below detection limit (0.5 μmol C L⁻¹). Based on the increase in cell numbers when grown with glucose and ammonium (supplementary material Table S.4.1 and Figure S.4.1), it was calculated that about 1 to 3 μmol C L⁻¹ is needed for the observed cell numbers as carbon and energy source. Apparently, the uptake of amino acids alone could not sustain bacterial growth because the initial amino acid concentration (0.13 μmol L⁻¹) was already much lower than the required 1 to 3 μmol C L⁻¹. During the initial growth phase, we found a decrease in dissolved free amino acids (DFAA) concentration concurrently with an increase in dissolved combined amino acids (DCAA), whereas the overall amino acid concentration (DFAA + DCAA) did not decrease (Table 4.1.1). Within the growth phase, the DFAA might have been used as precursors, e.g. for exo-enzymes, which lead in turn to an increase in the DCAA concentration, but no overall decrease in the amino acid concentration. During stationary phase, we found a slight decrease of total amino acids. This may suggest that amino acids were rather used as substrate for maintaining non-growing cells.

The compositional analysis of DOM with ESI FT-ICR-MS revealed a decrease of nitrogen-containing compounds during the initial growth phase (Figure 4.1.3 B and C). Inferring from their molecular compositions (high H/C, low O/C, N-containing), these substances were most likely detergents, such as ampholytic amino oxides or betaines. Using this method we cannot quantify the amount of carbon corresponding to this decrease. In addition, a compound with a decreasing peak in the mass spectrum is not necessarily completely oxidized to CO₂. Thus, we cannot ultimately clarify whether the use of these substrates alone explains the observed growth. Nevertheless, the preferential decrease of nitrogen-containing compounds suggests that these substances at least served as nitrogen source. This agrees with the observation that N₂ fixation was not detectable.

Growth and substrate use in natural seawater

The overall concentration of amino acids in the natural seawater was already initially quite low and did not further decrease during the incubation. Thus, amino acids did not sustain growth or survival of cells. The compositional analysis of DOM showed a decrease of two

groups of compounds. One of them was likely containing carbohydrate-like compounds with nitrogen and sulfur. These compounds are most likely thiosugars and/or aminosugars and may therefore have also served as nitrogen source. In addition, we observed a decrease in relative intensities of compounds with low O/C and H/C ratios containing sulfur but no nitrogen. Compounds with low amounts of oxygen and hydrogen in comparison to carbon are typically condensed or aromatic hydrocarbons. Since it cannot be excluded that these compounds were lost abiotically, e.g. by absorption to the glassware, it needs to be further investigated if the bacteria indeed metabolized these complex molecules during growth. Notably, during growth of *Pseudovibrio* sp. strain FO-BEG1 in natural seawater, we observed a decrease of different groups of substances compared to artificial seawater and preferentially substances containing sulfur decreased.

The use of multiple substrates

The potential use of different substrates under oligotrophic conditions was shown by the biologic experiment. The bacteria were able to metabolize different types of organic compounds, such as sugars (e.g. D-raffinose and D-trehalose), amino acids (L-serine, glutamic acid), carboxylic acids (glucuronic acid, acetic acid) and amide (glucuronamide). These compound classes were also present as contaminations in the artificial seawater medium and decreased during growth. This is in agreement with studies on *E. coli* showing a broad potential to use different substrates after adaptation to carbon starvation (Ihssen and Egli, 2005).

Even though the substances decreasing during growth, as revealed by ESI FT-ICR-MS, tended to cluster into certain groups with a specific ratio of O/C and H/C, the overall pattern showed a broad simultaneous use of many substrates both in artificial and natural seawater. In contrast to the biologic experiment, this does not show the potential to use a substrate, but the actual decrease of compounds present in original concentrations. Using different compounds simultaneously may enable bacteria to grow on very low concentrations of each of the different substrates (Lendenmann *et al.*, 1996; Kovárová-Kovar and Egli, 1998), if a substrate does not repress enzymes for another less efficient one (reviewed in Egli, 2010). This strategy, together with the potential to use substrates, which are not present but may become available, enables bacteria to survive in habitats with a low and fluctuating supply of nutrients, as it is found in the oceans.

Conclusions

The data presented in this study demonstrate that contaminations can arise from many different sources (e.g. chemicals, bottles, humans or plastics), which in turn might support growth of bacteria under oligotrophic conditions. Moreover, even under extremely oligotrophic conditions, the isolated bacteria were not in a resting state, but showed a moderate growth, even though nitrogen, carbon and energy sources were limiting factors at the same time. The investigated *Pseudovibrio* sp. FO-BEG1 uses many different types of substrates under nutrient-poor conditions as demonstrated by the FT-ICR-MS data (Figure 4.1.3). In our case, amino acids were not the primary substrate for growth, but rather a complex mixture of organic compounds, preferably containing nitrogen. Furthermore, we were able to show that ESI FT-ICR-MS is a strong tool to investigate bacterial growth under low-nutrient conditions.

Acknowledgments

We thank S. Menger, M. Meyer, M. Friebe, K. Klapproth, B. Kürzel and T. Max for technical support and T. G. Ferdelman and W. Ziebis for providing us with natural seawater samples from the Integrated Ocean Drilling (IODP) Expedition 329. This study is funded by the European Research Council (No 203364-ELNOX) and the Max Planck Society.

Supplementary figure and table

Table S.4.1: Cell numbers of isolate *Pseudovibrio* strain FO-BEG1 derived from different amounts of carbon added to the medium. Carbon in form of glucose was added to a *Pseudovibrio* strain FO-BEG1 culture in 2 different concentrations. As a negative control a culture without addition of carbon was used. To ensure carbon-dependent growth, ammonium was added to the medium (with and without carbon addition) as nitrogen source.

Carbon added	Ammonium added	Cells mL ⁻¹ (day 7-8)
60 μmol L ⁻¹	10 mmol L ⁻¹	31 x 10 ⁵
180 μmol L ⁻¹	30 μmol L ⁻¹	72 x 10 ⁵
0 μmol L ⁻¹	10 mmol L ⁻¹	3 x 10 ⁵
0 μmol L ⁻¹	30 μmol L ⁻¹	10 x 10 ⁵




Figure S.4.1: Amount of carbon needed for specific cell number. From additions of glucose to the medium (Table S.4.1) the amount of carbon needed for growth of isolate FO-BEG1 was calculated by subtracting the cell numbers without carbon addition (as negative control) from the cell numbers with carbon addition and plotting amount of carbon versus cell numbers produced. The amount of carbon needed for 10⁵ cells mL⁻¹ was calculated to be 3 μmol C L⁻¹ and for 10⁴ cells 1 μmol C L⁻¹.

4.2 Facultatively oligotrophic bacteria isolated from the habitat of large sulfide-oxidizers

Anne Schwedt^{1*}, Vladimir Bondarev¹, Michael Seidel^{1,2}, Thorsten Dittmar^{1,2}, Heide N. Schulz-Vogt¹

¹Max Planck Institute for Marine Microbiology, Celsiusstr. 1, D-28359 Bremen, Germany

²Institute for Chemistry and Biology of the Marine Environment, Carl von Ossietzky University of Oldenburg, Germany

*Corresponding author

Manuscript in preparation

Keywords: cultivation; facultative oligotrophy; *Pseudovibrio*; *Marinobacter*

Abstract

Axenic cultivation of large chemolithoautotrophic bacteria belonging to the genus *Beggiatoa* succeed only rarely. Growth of the large sulfide-oxidizers often seems to be dependent on the presence of heterotrophic prokaryotes, similar to the often described associations of cyanobacteria and heterotrophs. Recently, we observed that the growth of the marine, chemolithoautotrophic *Beggiatoa* sp. strain 35Flor depends on the presence of the α -Proteobacterium *Pseudovibrio* sp. strain FO-BEG1. Furthermore, we found that this bacterium, besides heterotrophic growth on organic-rich medium, is capable of growth under extreme nutrient deficiency in artificial and natural seawater. This observation inspired us to investigate whether we could isolate other facultative oligotrophs from overlaying water of Namibian sediment, an environment known to contain a large number of different sulfide-oxidizers belonging to the family *Beggiatoaceae*. Indeed, we succeeded to obtain 14 new strains closely related to known marine bacteria, all of which were capable of growth under extreme nutrient deficiency. The potential of these isolates to support growth of the large sulfide-oxidizing bacteria can now be studied in culture-based experiments.

Introduction

Large bacteria, such as *Beggiatoa* spp. or filamentous cyanobacteria, often live together with heterotrophic prokaryotes and these associations seem to be the reason for the inability of axenic cultivation of the large bacteria (Burton and Morita, 1964; Cohen and Rosenberg, 1989; Palinska *et al.*, 1999; Morris *et al.*, 2008). Different reasons for these interactions have been proposed, such as the recycling of carbon dioxide or the reduction of the oxygen concentration (Kuentzel, 1969; Paerl and Pinckney, 1996). It is known that *Beggiatoa* spp. typically lack the gene for catalase (Larkin and Strohl, 1983), but since aerobic respiration produces reactive oxygen species (ROS, Tapley *et al.*, 1999) an efficient protection against such molecules is needed. Therefore, *Beggiatoa* spp. might depend on the enzyme catalase, which catalyzes the disproportionation of hydrogen peroxide to oxygen and water, or other protection systems of the associated heterotrophs against ROS. The positive effect of catalase and of accompanying heterotrophic bacteria on the growth of *Beggiatoa* filaments (Burton and Morita, 1964; Strohl and Larkin, 1978; Güde *et al.*, 1981; Nelson *et al.*, 1986b) and, furthermore, the accumulation of peroxides in cultures without catalase or accompanying bacteria was shown (Burton and Morita, 1964). Recently, we found that both *Pseudovibrio* sp. strain FO-BEG1 and *Pseudovibrio denitrificans* (DSM number 17465) support growth of the marine chemolithoautotrophic *Beggiatoa* sp. strain 35Flor (Chapter 3.1). Both bacterial

strains are heterotrophic organisms (Chapter 3.2, Shieh *et al.*, 2004). Additionally, we have shown that the newly isolated *Pseudovibrio* sp. strain FO-BEG1 can grow under extremely oligotrophic conditions and its substrate use in pure artificial and natural seawater was studied in detail (Chapter 4.1).

The aim of the present study was to isolate heterotrophic, facultatively oligotrophic bacteria from Namibian sediments, the habitat of large sulfide-oxidizers, to investigate how common facultative oligotrophy is among bacteria associated with large sulfide-oxidizers. Here, we report the successful isolation of 14 facultatively oligotrophic bacteria from water overlaying Namibian sediments using a method relying on the change from oligotrophic to eutrophic growth conditions, called the CANgrow–method (changing availability of nutrients growth–method). In contrast to earlier methods for the isolation of marine bacteria, the artificial oligotrophic medium used here, is defined and contains much lower nutrient concentrations. Three initial transfers strongly pre-select for bacteria, which can grow under extreme nutrient deficiency. Subsequently, three transfers on nutrient-rich agar plates select for facultatively oligotrophic bacteria and are used to obtain pure cultures. Finally, the ability of the isolates to grow oligotrophically is ensured by at least seven transfers in pure artificial seawater.

Material and methods

Samples

The new bacterial strains were isolated from oceanic bottom water overlaying Namibian sediments that harbor different large sulfur bacteria (sample acquisition described in Salman *et al.*, 2011). All samples were stored at 4°C. In addition to the new isolates, the *Pseudovibrio denitrificans* type strain (DSM number 17465) was purchased from the German culture collection DSMZ (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Braunschweig, Germany) and cultivated under oligotrophic conditions.

Growth media and cultivation conditions

For cultivation and isolation, two different media were used, an oligotrophic and a eutrophic medium. The liquid, oligotrophic medium was composed as described above (Chapter 4.1), but prepared under synthetic air atmosphere (20% O₂ in N₂; H₂O < 3 ppm-mol, C_nH_m < 0.1 ppm-mol, CO < 1 ppm-mol, CO₂ < 1 ppm-mol). Furthermore, bottles were filled only with 50 mL medium and the medium was not cleaned using solid phase extraction. This medium was used for all oligotrophic cultivation experiments. The solid, eutrophic medium was

composed as described above (Chapter 4.1, Methods section “Biolog experiment”). All incubations in oligotrophic and eutrophic media were performed without shaking at 28°C in the dark.

CANgrow–method (changing availability of nutrients growth–method)

For isolation, 50 mL of oligotrophic medium were inoculated with 100 µL seawater sample (from off shore Namibia). The cultures were transferred (100 µL enrichment in 50 mL fresh medium) at least three times in oligotrophic medium with incubation periods between the transfers of at least one week. Aliquots of the oligotrophic enrichments were then plated on eutrophic, solid medium and single colonies were transferred three times on eutrophic medium. Finally, at least another seven transfers (100 µL culture in 50 mL fresh medium) were performed in liquid, oligotrophic medium (Figure 4.2.1).

Sequencing of 16S rDNA genes and phylogenetic analysis

Eutrophically grown colonies were picked and directly transferred to a polymerase chain reaction (PCR) mix containing 1× PCR MasterMix (Promega, Mannheim, Germany) and 1 µmol L⁻¹ of each primer (GM3F and GM4R in Muyzer *et al.*, 1995). The PCR program applied was as follows: initial denaturation at 95°C for 5 minutes, 32 cycles of 95°C for 1 minute, 50°C for 30 seconds and 72°C for 90 seconds followed by a final elongation at 72°C for 7 minutes. PCR products were cloned using the TOPO TA Cloning[®] Kit for Sequencing (Invitrogen, Karlsruhe, Germany) according to manufacture’s instructions. Sequencing of the cloned inserts was performed using the Big Dye Cycle Sequencing Kit (Applied Biosystems, Carlsbad, CA, USA) and sequences were analyzed on an ABI Genetic Analyzer 3130x (Applied Biosystems, Carlsbad, CA, USA). Nearly full-length sequences were assembled with SeqMan (Lasergene software package, DNASTar, Madison, WI, USA) and deposited in the DDBJ/EMBL/GenBank databases under accession numbers FR716535 to FR716549. Phylogenetic analysis of the 16S rDNA sequences was performed using the ARB software package (Ludwig *et al.*, 2004) and release 102 of the SILVA SSURef database (Pruesse *et al.*, 2007). Tree reconstruction with maximum likelihood and neighbour joining methods was performed using 0, 30 and 50% positional conservatory filters that exclude highly variable regions. Finally, a consensus tree based on the different reconstruction methods was built. A total number of 102 nearly full-length sequences was used for initial calculation to stabilize tree topology. Displayed in the final tree (Figure 4.2.2) are the sequences of the 15 isolates grouped with their closest relatives.

Cell counts

Cell counts were performed as described in Chapter 4.1.

Measurement of dissolved organic carbon (DOC)


Dissolved organic carbon (DOC) was measured in the oligotrophic medium using a Shimadzu TOC-VCPH total organic carbon analyzer (Shimadzu, Kyoto, Japan). Acidification of samples was performed with 1% v/v 2 mol L⁻¹ HCl followed by sparging with synthetic air in order to remove inorganic carbon. The detection limit of the method was 5 μmol C L⁻¹ (0.06 mg C L⁻¹). The analytical accuracy was confirmed with reference material (deep Atlantic seawater) and low carbon water from the consensus reference materials program (D.A. Hansell, University of Miami, Coral Gables, FL, USA).

Results

Isolation of facultatively oligotrophic bacteria

Applying the CANgrow–method, which favors facultatively oligotrophic bacteria (Figure 4.2.1), we obtained 14 isolates of marine bacteria that were able to adapt from oligotrophic to eutrophic growth conditions and vice versa within 3 to 5 days. We were able to obtain pure cultures of these strains by transferring single colonies from organic-rich agar plates and could show that these colonies were able to grow oligotrophically by at least seven transfers in pure artificial seawater.

Figure 4.2.1: Comparison of strategies for the isolation of oligotrophic bacteria. Three different methods for the isolation of obligately or facultatively oligotrophic bacteria are compared with the newly developed CANgrow–method (Changing availability of nutrients growth–method).



Phylogeny of the isolates

Bacteria from three different phyla were isolated with the CANgrow–method. Phylogenetic relations are shown in a 16S rDNA sequence tree including the *Pseudovibrio* sp. strain FO-BEG1 showing > 99.5% sequence identity to the *Pseudovibrio denitrificans* type strain (Figure 4.2.2). Except for two isolates, which were closely related to *Arthrobacter* spp. (*Actinobacteria*) on 16S rDNA level (99.8 to 99.9% identity to closest related strain), all isolates were members of the *Proteobacteria*. The remaining 12 isolates were members of the *Gammaproteobacteria*, two of them grouping with *Kangiella* spp. (96.4% identity to closest related strain) and ten with *Marinobacter* spp. (98.7 to 100% identity to closest related strain).




Figure 4.2.2: Phylogenetic trees based on a total number of 102 nearly full-length sequences were calculated with maximum likelihood and neighbor joining methods using different positional conservatory filters. The displayed tree is an excerpt from the consensus tree that was inferred based on the different reconstruction approaches. The 14 new isolates and strain FO-BEG1 are grouped with the most closely related type strains. Isolated strains listed in one line feature an identical 16S rDNA sequence, whereas isolated strains listed directly one below the other are 99.6 to 99.9% identical in their 16S rDNA sequence. The isolates FO-NAM13, 14 were only able to grow for six transfers under oligotrophic conditions and are therefore marked grey in the tree.

Oligotrophic growth

The artificial seawater contained a DOC concentration of $0.18 \pm 0.06 \text{ mg C L}^{-1}$ ($15 \pm 5 \text{ } \mu\text{mol C L}^{-1}$). Growth curves in oligotrophic artificial seawater medium (Figure 4.2.3) were obtained for one isolate from each phylogenetic group (*Actinobacteria*, *Alpha-* and *Gammaproteobacteria*). We observed a clear increase in cell numbers starting from 4 to 20 cells mL^{-1} to a final density of 10^4 to 10^5 cells mL^{-1} . The proteobacterial isolates showed growth after 2 days and reached a stationary phase after 5 to 7 days of incubation. Both actinobacterial isolates were characterized by delayed growth that was detectable after 12 days. Here, the stationary phase was reached after about 20 days of incubation. Moreover, all isolates except for the two isolated *Kangiella* strains were able to grow after at least seven transfers in the oligotrophic seawater medium. The two isolates closely related to *Kangiella* spp. were not able to grow after more than six transfers under oligotrophic conditions. The isolates closely related to *Marinobacter* spp. reached the highest final cell numbers (Figure 4.2.3 D), whereas the actinobacterial isolates showed lowest final cell densities (Figure 4.2.3 C). Furthermore, we observed growth under oligotrophic conditions for the type strain *Pseudovibrio denitrificans* (DSM number 17465). The growth curve of this strain (Figure 4.2.3 B) showed the same pattern as the *Pseudovibrio* sp. FO-BEG1, which is currently growing in the 26th oligotrophic transfer in highly purified artificial seawater.




Figure 4.2.3: Oligotrophic growth curves of (A) strain FO-BEG1 (*Alphaproteobacteria*, related to *Pseudovibrio* spp.), (B) *Pseudovibrio denitrificans* type strain (*Alphaproteobacteria*), (C) isolate FO-NAM2 (*Actinobacteria*, related to *Arthrobacter* spp.) and (D) isolate FO-NAM6 (*Gammaproteobacteria*, related to *Marinobacter* spp.).

Discussion

Isolation of facultatively oligotrophic bacteria with the CANgrow–method

Each strategy that is applied for the isolation of bacteria selects for a specific physiology and metabolism. Most approaches used recently for the isolation of oligotrophic bacteria are based on the dilution to extinction method (Button *et al.*, 1993; Connon and Giovannoni, 2002) and thereby select for the most abundant microorganisms. In contrast, the CANgrow–method favors bacteria, which might not have been particularly abundant in the original inoculum, but can adapt fast to changes in nutrient availability. Previous studies have shown that many bacteria isolated under oligotrophic conditions can adapt to nutrient-rich media (Yanagita *et al.*, 1977; MacDonell and Hood, 1982; Carlucci *et al.*, 1986). Also, the single-cell encapsulation method (Zengler *et al.*, 2002) is based on oligotrophic growth followed by eutrophic growth conditions. This cultivation approach is similar to ours (Figure 4.2.1), but we used artificial seawater of very low DOC concentration rather than natural seawater. The measured DOC concentration of 0.18 mg C L^{-1} is two to five times lower than in natural seawater (Schut *et al.*, 1997; Hansell *et al.*, 2009). Nevertheless, we are certain to observe true growth under these extremely oligotrophic conditions, since we performed at least seven transfers in purified artificial seawater with each of the isolated strains and 26 transfers for *Pseudovibrio* sp. FO-BEG1, for which we also identified growth substrates under oligotrophic conditions (Chapter 4.1). The initial cell number after each transfer was 4 to 20 cells mL^{-1} . Thus, 9 to 15 divisions must have occurred between two consecutive transfers to account for a final cell number of 10^4 to 10^5 cells mL^{-1} as observed at the end of the growth phase. This accounts for 60 to 100 divisions during a total of 7 incubations. Therefore, we conclude that all isolates are viable under oligotrophic conditions by the definition of Button *et al.* (1993), who characterize organisms as viable after having performed 13 divisions which we observe already after 1 to 2 transfers.

Diverse phylogeny of facultatively oligotrophic bacteria

The isolated bacterial strains belong to different phylogenetic groups, namely *Alphaproteobacteria*, *Gammaproteobacteria* and *Actinobacteria*. Growth under oligotrophic conditions as observed for the isolates FO-NAM13 and FO-NAM14 (related to *Kangiella* spp.), has so far never been described for any member of the genus *Kangiella*. Long-term starvation and survival but not growth in the absence of external nutrients has previously been reported for *Arthrobacter* spp. and was proposed to be fueled by internally stored reserve material (Zevenhuizen, 1966). In our study, isolates FO-NAM1, FO-NAM2 (related to *Arthrobacter*

spp.) were transferred in oligotrophic medium more than seven times and active growth was always observed. Hence, we assume that the cells gained energy and produced biomass from an external source, since we determined growth and not only survival. Bacteria belonging to the genus *Marinobacter* are known to be diverse in physiology (e.g. Gauthier *et al.*, 1992; Huu *et al.*, 1999). Substrate uptake under low nutrient conditions was shown for *Marinobacter arcticus* (Button *et al.*, 2004), but oligotrophic growth was not studied in detail. In contrast, the isolates FO-NAM3 to FO-NAM12 (related to *Marinobacter* spp.) actively grow under nutrient deficiency.

Cell numbers of the isolated strains growing under oligotrophic conditions differed between the phylogenetic groups. The cell numbers of isolates related to *Marinobacter* spp. were higher than cell numbers of the other isolates, whereas the isolates related to *Kangiella* spp. did not grow for more than six transfers. This suggests that the different bacteria vary in their capabilities of adapting to oligotrophic growth conditions or that the present organic and inorganic material can be used differently among the isolates. This might be due to the presence of different metabolic pathways and a different number and type of high affinity transporters necessary for scavenging nutrients at such low concentrations. Taken together, our data support earlier findings (Yanagita *et al.*, 1977; MacDonell and Hood, 1982; Carlucci *et al.*, 1986) that the ability to switch between extreme nutrient deficiency and affluence of substrate is not unusual or restricted to a certain phylogenetic group, even if the level of adaptation might differ. Therefore, we propose that the ability to grow under extreme substrate limitation is much more widespread among known heterotrophic bacteria than currently recognized.

As expected, the heterotrophic bacteria isolated from the habitat of large sulfide-oxidizers are phylogenetically diverse. It was possible to isolate bacteria following a similar metabolic strategy – facultative oligotrophy. If these bacteria are associated with sulfide-oxidizers, if their metabolic activity is somehow related to the presence or absence of the lithotrophs or whether they can support growth of the large sulfur bacteria by scavenging ROS needs to be further investigated.

Acknowledgments

This study was funded by the European Research Council (No 203364-ELNOX) and the Max Planck Society. We thank A.-C. Kreutzmann for helpful remarks on the manuscript, V. Salman, M. Friebe, S. Menger and M. Meyer for technical support and the members of the Meteor cruise M76 (April – June 2008) for providing samples.

„The important thing in science is not so much to obtain new facts as to discover new ways of thinking about them.“

~Sir William Lawrence Bragg (1890-1971)

Chapter 5

Concluding remarks and outlook

The findings of this thesis contribute to different topics ranging from migration behavior of mat-forming sulfur bacteria via associations between different bacteria to the lower limits of bacterial growth. Nevertheless, all these different aspects are linked to each other. The growth of *Beggiatoa* sp. filaments, for which we discovered an unusual migration behavior (Chapter 2), depends on the presence of *Pseudovibrio* sp. (Chapter 3), which is capable of growth under extreme nutrient deficiency (Chapter 4). This chapter (Chapter 5) connects all these different aspects, including preliminary data, which are not presented in the previous chapters and considers the obtained results in a broader context. Furthermore, a detailed discussion of special proceedings and precautions during performed experiments is given. Finally, this chapter ends with an outlook for future research concerning the discussed topics.

Associations between small heterotrophic and larger bacteria

Axenic cultivation of large marine *Beggiatoa* spp. under chemolithoautotrophic conditions is difficult and only rarely successful (Nelson and Jannasch, 1983; Nelson *et al.*, 1986a). In this thesis, a successful and stable cultivation of the marine chemolithoautotrophic *Beggiatoa* sp. strain 35Flor (6 µm in diameter) is described. This *Beggiatoa* strain grows solely in the presence of an accompanying organism, the heterotrophic *Pseudovibrio* sp. strain FO-BEG1 (Chapters 2 and 3). We propose that the accompanying bacterium protects the *Beggiatoa* sp. from oxidative stress because we have shown that the genome of the accompanying organism possesses more than 20 genes for the enzymes catalase, superoxide dismutase and peroxidase (Chapter 3.2), whereas sulfide-oxidizing bacteria of the genus *Beggiatoa* typically do not possess the gene for the enzyme catalase (reviewed in Larkin and Strohl, 1983).

Similar to large sulfide-oxidizers, also large marine cyanobacteria can often be found associated with small heterotrophic bacteria and it is difficult to sustain axenic cultures (Palinska *et al.*, 1999; Morris *et al.*, 2008). The association of small heterotrophic bacteria

with larger bacteria, such as *Beggiatoa* spp. or cyanobacteria in microbial mats is a common observation (Cohen and Rosenberg, 1989; van Gemerden, 1993). The complex interactions between cyanobacteria and heterotrophic bacteria have been studied and different reasons for their interactions have been proposed. These reasons include recycling of carbon dioxide needed by the cyanobacteria (Kuentzel, 1969), production of growth factors and formation of anoxic microniches due to aerobic respiration (Paerl and Pinckney, 1996). Interestingly, for cyanobacteria of the genus *Prochlorococcus*, it was also proposed that the function of the heterotrophs is to scavenge reactive oxygen species (ROS) because the *Prochlorococcus* spp. themselves have no gene encoding for a catalase or peroxidase (Morris *et al.*, 2008). Furthermore, a mutant of the accompanying heterotroph lacking the gene for catalase was found to not support growth of the cyanobacteria, whereas the addition of catalase had a positive effect on their growth (Morris *et al.*, 2008). These observations concerning the association between cyanobacteria and accompanying heterotrophic bacteria point in the same direction as the observations presented here on the association between *Beggiatoa* sp. and *Pseudovibrio* sp. (Chapter 3). It suggests that the protection system of the large bacteria from ROS might be less efficient than the ones of smaller heterotrophic bacteria.

Cyanobacteria and *Beggiatoa* spp. are often found to share one habitat as for example in microbial mats (van Gemerden, 1993) or in biofilms associated with the black band disease of scleractinian corals (Richardson, 1996). Therefore, in these common habitats they might also live together with similar types of heterotrophic bacteria. The bacteria associated with cyanobacteria belong to a diverse range of phylogenetic groups, including *Actinobacteria*, *Bacteroidetes*, *Alpha-*, *Beta-*, *Gamma-* and *Deltaproteobacteria* (Salomon *et al.*, 2003; Kolmonen *et al.*, 2004; Hube *et al.*, 2009). The isolation of phylogenetically different bacteria from the habitat of large sulfide-oxidizers (Chapter 4.2) indicates that also the large sulfide-oxidizers are not restricted to the association with bacteria of one phylogenetic group. It seems more likely that the large bacteria depend on specialized functions performed by certain types of the associated bacteria rather than the presence of a specific phylogenetic group. Very recently, a metagenomic sequence analysis of bacterial communities associated with the green macroalgae *Ulva australis* revealed that not the phylogeny of the associated bacteria but the function of their genes was correlated with the associations (Burke *et al.*, 2011). Consequently, it was proposed that the functional genes rather than the 16S rDNA genes are more appropriate to investigate associations in microbial communities (Burke *et al.*, 2011).

Toxicity factors and migration behavior of *Beggiatoa* filaments

The growth of *Beggiatoa* spp. depends on oxygen and sulfide, but both substances can also be harmful to the bacteria if concentrations exceed a critical threshold (Winogradsky, 1887; Møller *et al.*, 1985). Close relatives of *Beggiatoa* spp. belonging to the genus *Thiomargarita* can be found in sulfidic sediments with sulfide concentrations of 100 to 800 $\mu\text{mol L}^{-1}$ (Schulz *et al.*, 1999) or even up to 22 mmol L^{-1} (Brüchert *et al.*, 2003). Moreover, these bacteria can also tolerate atmospheric oxygen levels while being exposed to lower sulfide concentrations (0 to 100 $\mu\text{mol L}^{-1}$) at a pH of 7.3 (Schulz *et al.*, 1999; Schulz and de Beer, 2002). However, *Thiomargarita* spp. cells are not as motile as *Beggiatoa* spp. and as a consequence they can not position themselves in the transition zone of oxygen and sulfide. Instead, the cells wait for resuspension of the sediment to get into contact with oxygenated, nitrate-containing seawater (Schulz *et al.*, 1999).

In contrast, *Beggiatoa* filaments are motile and built up mats between the opposing gradients of oxygen and sulfide (Winogradsky, 1887; Keil, 1912; Jørgensen, 1977; Nelson *et al.*, 1982; Nelson and Jannasch, 1983). Consequently, *Beggiatoa* filaments usually get into contact with oxygen and sulfide at the same time, since they consume the two gases, they lower the concentrations and steepen the gradients (Nelson *et al.*, 1986a). However, depending on the thickness of the *Beggiatoa* mat, some filaments might not be in direct contact with oxygen and sulfide at the same time (Nelson *et al.*, 1986b). Taken this into account, the results presented in this thesis (Chapter 2.1) suggest that sulfide is probably more harmful to the cells under oxic conditions because the aerobic sulfide oxidation leads to a decrease in pH (Winogradsky, 1887) and as a consequence more sulfide outside the cells is present in form of H_2S . In the presented experiments (Chapter 2.1) we measured a pH of 6.5 within the mat, leading to about 70% of the sulfide being present in form of H_2S (compared to only about 30% present as H_2S at pH 7.3). This uncharged, harmful gas can easily diffuse into the cells where it can act as a strong reductant and binds to iron in cytochromes, by this blocking the cellular respiration. Thus, the bacteria probably have to perform sulfide oxidation to detoxify the inside of the cells. At high sulfide fluxes the cells obviously accumulate large amounts of storage compounds during this process and might eventually burst. To prevent this, the bacteria have to dispose of the internal storage compounds. In the anoxic regions the pH is higher and sulfide predominantly occurs in form of HS^- which can not diffuse into the cells and might only get inside via ion channels or transport systems.

In this thesis (Chapter 2.1), a new strategy to cope with high sulfide fluxes is proposed. The bacteria actively migrate into anoxic regions with high sulfide concentrations. Here, the aerobic sulfide oxidation is stopped and with this also sulfur deposition. The bacteria can then reduce the intracellular sulfur with internal PHA to form sulfide. This strategy represents a novel explanation for the presence of *Beggiatoa* filaments in anaerobic habitats and shows that filaments actively migrate into anoxic, sulfidic regions. Previously, downwards migrations were typically observed in habitats containing nitrate and there the anaerobic sulfide oxidation with nitrate lowered the diffusion zone of sulfide and separated the oxygen and sulfide gradients over up to a few centimeters (Mußmann *et al.*, 2003; Sayama *et al.*, 2005; Hinck *et al.*, 2007; Preisler *et al.*, 2007).

Although *Beggiatoa* spp. also require oxygen for growth (unless an alternative electron acceptor is present) they show phobic reactions to higher (above 5% air saturation) oxygen concentrations (Winogradsky, 1887; Møller *et al.*, 1985). During aerobic sulfide oxidation the production of oxygen radicals and, moreover, chemiluminescence was shown to occur (Tapley *et al.*, 1999). As mentioned above, we propose that the *Beggiatoa* sp. 35Flor requires the accompanying organism to protect themselves against reactive oxygen species (Chapter 3.1). However, no correlation between the catalase activity of heterotrophic, associated cells and the beneficial effect of their presence on cyanobacteria (*Prochlorococcus* sp.) was found (Morris *et al.*, 2008). In fact, the heterotroph with the lowest catalase activity was the only strain able to support growth of all studied cyanobacteria. Consequently, the dependency of the large bacteria on small heterotrophic bacteria can not be completely explained by their possession of a catalase enzyme. During our studies on the co-culture of *Beggiatoa* sp. and *Pseudovibrio* sp., the possibility that nitric oxide (NO) might be involved in the reaction counteracting oxidative stress arose. In oxygen-sulfide gradient co-cultures with high sulfide concentrations, the NO signal in *Pseudovibrio* cells, stained with a copper-based fluorescent probe (CuFL, Lim *et al.*, 2006), was higher compared to cells exposed to a low sulfide flux (Figure 5.1). This increase in NO signal was also inducible when hydrogen peroxide (H₂O₂) was added to co-cultures with a low sulfide flux or to pure cultures of the isolated *Pseudovibrio* sp. (growing in seawater medium containing glucose and ammonia). Under both conditions, the enhanced signal was visible in the interior of the cells and not in the surrounding medium. The used dye is specific for NO and no interaction with H₂O₂ or other reactive oxygen and nitrogen species, such as HNO, NO₂⁻, NO₃⁻ and ONOO⁻, could be detected (Lim *et al.*, 2006).




Figure 5.1: Nitric oxide staining (copper-based fluorescent probe) of *Pseudovibrio* sp. FO-BEG1 cells in co-culture with *Beggiatoa* sp. 35Flor at high and low sulfide flux (12.6 and 50.4 mmol m⁻² d⁻¹, respectively) and in *Pseudovibrio* sp. FO-BEG1 pure cultures. White bars show measurements without the addition of hydrogen peroxide and grey bars show those with 0.35% H₂O₂ added. (preliminary data, imaging performed together with M. Beutler)

NO is known to protect against oxidative stress in other bacteria. In *Bacillus subtilis*, the addition of NO has been shown to lead to an increase in resistance against H₂O₂ by 100fold (Gusarov and Nudler, 2005). The enzymes of the group nitric oxide synthases (NOS) can produce NO and citrulline from L-arginine and oxygen and are typically known from eukaryotes (Alderton *et al.*, 2001). However, recently genes encoding for NOS-like enzymes were found also in different gram-positive and some gram-negative bacteria, and also the NOS-dependent NO production could be shown (Adak *et al.*, 2002a; Adak *et al.*, 2002b; Gusarov and Nudler, 2005; Agapie *et al.*, 2009; Schreiber *et al.*, 2011).

Interestingly, we found NOS-related domains (Filippovich, 2010) in the genome of the investigated *Pseudovibrio* sp. FO-BEG1 (Schwedt *et al.*, unpublished data). A small domain of yet unknown function is followed by an amine oxidase domain (known to oxidize L-amino acids) and a flavodoxin/nitric oxide synthase domain in the genome sequence. This opens up the possibility that NO production might be involved in the reaction to ROS in the investigated *Pseudovibrio* sp. strain FO-BEG1. The production of NO as a protective measure against oxidative stress in *Bacillus subtilis* cells leads to an enhanced catalase activity, which was shown by comparison of wild type to Δnos deletion mutants (Gusarov and Nudler, 2005). In that study, the bacteria were cultivated in a complex medium with yeast extract containing free amino acids. In the pure culture experiments presented here, the medium contained ammonia, as the sole nitrogen source, which could have potentially been used to produce NO.

In our co-culture experiments together with *Beggiatoa* sp., no fixed nitrogen source was present, but the investigated *Beggiatoa* sp. are able to fix N₂ (Henze, 2005). The genome of the accompanying organism does not contain any gene encoding for a nitrogenase enzyme (Schwedt *et al.*, unpublished data). The transfer of fixed nitrogen sources from cyanobacteria to associated bacteria is a common observation (Paerl, 1984). It may well be that also the *Beggiatoa* sp. transfer fixed nitrogen to the accompanying bacteria. Nevertheless, taking into consideration that dyes can unspecifically bind to different compounds and that the production of NO in our study was not proven by direct measurements, these considerations remain speculative and await further investigation (see outlook).

Apart from chemical substances, such as oxygen and sulfide, also light is known to be a potential cause of phobic migration reactions of *Beggiatoa* filaments (Winogradsky, 1887; Nelson and Castenholz, 1982; Møller *et al.*, 1985). The *Beggiatoa* sp. that we investigated showed an unusual reaction to blue/green light. Application of a blue or green light source (intensity: 67 $\mu\text{E m}^{-2} \text{s}^{-1}$) above or below the culture tube induced an immediate downwards movement of a subpopulation of filaments into the anoxic part of the culture tube when cultivated under a high sulfide flux (Chapter 2.2). The downward movement also occurred in the dark, but solely in cultures where a mat at the oxic-anoxic transition had already been established for about 10 days. The lower subpopulation was established 2 to 3 days after the downwards migration started (Chapter 2.1). Applying a blue or green light source in close proximity to the culture tube induced and also enhanced this filament movement and within a few hours the lower subpopulation was observed (Chapter 2.2). Because chemiluminescence occurs during chemical sulfide oxidation (Tapley *et al.*, 1999) at the transition between oxygen and sulfide, it can be speculated that light at low intensities might be involved in the migration and/or mat formation processes of *Beggiatoa* filaments, but this remains to be studied in the future (see outlook).

The lower limits of bacterial growth

In contrast to the large *Beggiatoa* filaments, most bacterial cells are not visible with the naked eye. However, bacterial growth can lead to macroscopically visible colonies on nutrient-rich agar plates or cause turbidity in nutrient-rich liquid media due to high cell densities, whereas under nutrient limitation cell densities will be low and the small bacterial cells might not cause turbidity. Therefore, the build-up of bacterial biomass in liquid low-nutrient media has to be quantified in a different way than optical turbidity measurements. Although, there might

be no measurable growth, the cells might still be metabolically active. This is due to the fact that there is a difference between biomass production and activity of bacterial cells. Although the consumption of substrate is essential for the cells to grow and build-up biomass, the opposite is not the case, as cells do not necessarily grow while utilizing substrate (del Giorgio and Cole, 1998). Therefore, the formation of biomass, called bacterial production (BP) has to be separated from the consumption of substrate, called bacterial respiration (BR). The bacterial growth efficiency (BGE) is defined as the quantity of bacterial biomass resulting from a certain amount of substrate respired (del Giorgio and Cole, 1998). The substrate is used by the bacteria for catabolic and anabolic purposes generating ATP (Adenosine-5'-triphosphate) and cell biomass, respectively. The BGE can be determined by isotope-labeling techniques (BR and BP), rates of protein or DNA synthesis (BP), cell counts (BP), oxygen consumption (BR), carbon dioxide production (BR) and changes in DOC and POC (BR and BP) (del Giorgio and Cole, 1998 and references therein).

Under oligotrophic conditions, the concentrations of substrates and cell numbers are extremely low. Therefore, most of these methods may not be sensitive enough and measured values are close to or below the limit of detection. Thus, new routines and methods have to be developed or methods have to be refined to be suitable for studying bacterial growth under extreme nutrient deficiency. Classical dissolved organic carbon (DOC) measurements, which are used as a parameter for the amount of dissolved organic matter (DOM) in environments with a low amount of nutrients, have a detection limit of about $0.5 \mu\text{mol C L}^{-1}$ ($0.006 \text{ mg C L}^{-1}$). However, the observed cell numbers in our medium (Chapter 4) are so extremely low that about $1 \mu\text{mol L}^{-1}$ of carbon would be enough to explain the observed growth. Consequently, the classical method might not provide sufficient sensitivity to detect potential small changes in DOC. The results presented in this thesis show that electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry (ESI FT-ICR-MS) provides sufficient resolution and can be used to study bacterial growth under oligotrophic conditions. Furthermore, this technique gives information on both potential fixed carbon and nitrogen sources of the bacteria. Combined with the measurement of amino acids by high performance liquid chromatography (HPLC) and isotope-labeling techniques, bacterial growth in oligotrophic seawater can be studied in detail. However, this technique does generally not give quantitative information and as a consequence respiration rates under oligotrophic conditions still remain unknown.

The risk of impurities

Even though no electron donor is added to an oligotrophic medium, there can still be impurities present that might support growth and lead to the formation of biomass and/or metabolic activity of the cells (Chapter 4). Studying bacterial growth under nutrient-poor conditions is challenging, since many different contaminations, such as macronutrients, trace elements or energy sources can occur. Accordingly, when studying the physiology of bacteria growing under oligotrophic conditions, it is crucial to keep the medium and equipment contamination-free for the mandatory substances for growth. Amino acids represent a serious contamination risk for seawater samples. Therefore, there are a number of precautions that have to be taken, such as working with gloves, combustion of glassware, avoidance of dust in the working room (working in laminar flow hood) and use of chemicals of highest available quality (Dittmar *et al.*, 2009). Also, during the measurements of quantity and composition of DOM typically all used equipment is pre-combusted if possible. All these precautions were applied during the preparation of the oligotrophic seawater medium and all further analyses presented in Chapter 4.1 of this thesis.

Furthermore, the medium might become contaminated with trace elements, which would probably not resemble a potential energy source but could be decisive for growth or no growth of microorganisms. To avoid the contamination with trace elements, equipment is usually washed with HNO_3^- (e.g. Fitzwater *et al.*, 1982), which, however, represents an easily utilizable nitrogen source. For the experiments presented in this thesis (Chapter 4), a fixed nitrogen source was considered as worse contamination than trace elements, and the equipment was not washed with HNO_3^- .

In spite of all precautions, it is still extremely difficult to prepare a seawater medium without any contaminations. However, if the contaminating substance is not used by the bacteria during growth, such as the PEG contamination in this study (Chapter 4.1), it might not disturb or influence further physiological analysis. The contamination can even be used to normalize other peaks in the mass spectra, if its peaks are high and consistent (as the PEG peaks in this study, Chapter 4.1).

Facultative oligotrophy

In addition to growth under oligotrophic conditions, the investigated *Pseudovibrio* sp. FO-BEG1 (Chapter 4) is capable of growing under eutrophic conditions as well and can switch within days from one trophic state to the other. Under oligotrophic conditions, the bacterial growth efficiency is typically very low and varies with the supply of nutrients (del Giorgio and Cole, 1998). Furthermore, the investigated bacteria are limited in carbon, nitrogen and energy at the same time (Chapter 4.1). Consequently, they are supposed to have high maintenance energy costs to maintain crucial transport systems and enzymes prepared (del Giorgio and Cole, 1998). With increasing nutrient supply, probably also the bacterial growth efficiency increases because the bacteria can exhibit higher growth rates and produce only enzymes necessary for the substrates available (reviewed in del Giorgio and Cole, 1998). Facultatively oligotrophic bacteria have to switch between oligotrophic and eutrophic growth conditions and as a consequence the cells have to switch also between both strategies for growth to adapt to the actual amount of substrates present.

The variation in nutrient availability is common in natural environments such as the open ocean. Bacteria attached to marine snow particles can exhibit very high growth rates (Alldredge *et al.*, 1986; Alldredge and Gotschalk, 1990; Smith *et al.*, 1992; Azam and Long, 2001; Kiørboe and Jackson, 2001), while free-living bacteria in the open ocean are restricted in nutrients and consequently their growth rates are low or they even starve (Boyle and Ensign, 1970; Novitsky and Morita, 1976; Azam and Hodson, 1977; Novitsky and Morita, 1977; Tabor and Neihof, 1982; Ishida *et al.*, 1989). Facultatively oligotrophic bacteria can adapt fast to nutrient affluence or deficiency, while they have to up- or down-regulate internal pathways of anabolism and catabolism. The investigated facultatively oligotrophic *Pseudovibrio* sp. FO-BEG1 is highly versatile with respect to its energy gain (Chapter 3.2 and Chapter 4.1). This feature makes the bacteria flexible and they are, moreover, capable to interact with other prokaryotes or possibly even eukaryotes (Chapter 3.2).

Already 30 years ago, it was suspected that two types of oligotrophic bacteria exist (Ishida and Kadota, 1981): 1. organisms, which disappear with increasing man-made eutrophication and 2. organisms, which can adapt fast to man-made eutrophication. In this thesis bacteria were not isolated from the oligotrophic open ocean, but from water directly overlaying marine sediments (Chapter 4.2). All of the isolated strains were able to adapt fast to the nutrient deficiency of the initial isolation medium. The observation that facultatively oligotrophic bac-

teria can also be isolated from non-oligotrophic water indicates that these bacteria might be more widespread and not limited to nutrient-poor environments. For example, they might live attached to marine snow particles (e. g. Alldredge *et al.*, 1986; Smith *et al.*, 1992; Azam and Malfatti, 2007 and references therein), which represent nutrient hotspots for heterotrophic bacteria, and possibly the bacteria sink down to the sediment with these particles. We assume that many facultatively oligotrophic bacteria have been overlooked so far because they were not searched for in non-oligotrophic environments. Likewise, many more already known heterotrophic bacteria may be capable of growing under much poorer nutrient conditions than currently assumed.

Conclusions

In conclusion, the results presented in this PhD thesis (summarized in Figure 5.2) show that filaments of *Beggiatoa* sp. strain 35Flor react to high sulfide fluxes by migration into anoxic regions, where they reduce the amount of internal storage compounds. This migration can be enhanced or induced by blue/green light for a yet unknown reason. We suggest that the accompanying *Pseudovibrio* sp. strain FO-BEG1 can detoxify reactive oxygen species (ROS), generally produced during sulfide oxidation (Tapley *et al.*, 1999), and might be responsible for the protection of *Beggiatoa* filaments, which are known to lack the gene for catalase (Larkin and Strohl, 1983). Possibly, production of nitric oxide (NO) by the *Pseudovibrio* sp. might also be involved in the protection against ROS. However, the origin of NO is unknown because no fixed nitrogen source was present in the medium of the co-culture and the heterotroph does not possess a gene encoding for nitrogenase. On the other hand, the *Beggiatoa* sp. can fix N_2 (Henze, 2005) and it might be that a fixed nitrogen source is transferred from the sulfide-oxidizer to the accompanying bacterium. The *Pseudovibrio* sp. examined in this thesis is a generalist able to gain energy in many different ways and can also grow under extremely nutrient-poor conditions. Furthermore, it possesses genes to interact with pro- and eukaryotes. We found that also other bacteria from the habitat of large sulfide-oxidizers can adapt fast from nutrient affluence to deficiency. Maybe these bacteria are able to also support growth of the large sulfide-oxidizers, but this needs to be investigated in the future.




Figure 5.2: Illustration of the results obtained in this PhD thesis concerning the migration behavior of *Beggiatoa* sp. and the co-occurring *Pseudovibrio* strain. Some results were supported by our data (black) while others are hypothesis and remain to be proven (blue). ROS = reactive oxygen species, NO = nitric oxide, N = nitrogen

Outlook

The results of my PhD thesis provide new insights into the migration behavior of large sulfur bacteria, associations between bacteria, and growth under extreme nutrient deficiency. Nevertheless, the obtained results also raise new questions that need to be addressed in the future.

We hypothesize that the *Beggiatoa* sp. filaments depend on the heterotrophic *Pseudovibrio* sp. because they do not themselves possess protective measure against oxidative stress (Chapter 3). However, direct evidence for this theory is still missing. To investigate the capability of other heterotrophic, catalase-positive bacteria or ROS scavenger to support growth of the *Beggiatoa* sp., filaments from the lower subpopulation should be transferred without (or with only few) accompanying bacteria into fresh media. If growth is positive in the presence of the other catalase-positive bacteria or ROS scavengers, the transfer of *Pseudovibrio* sp. from the inoculation source has to be excluded, which can be tested by Fluorescence *in situ* hybridization.

The preliminary data on the staining of nitric oxide opens up the possibility that NO might be involved in the reaction with ROS (Chapter 5). However, to proof this, the production of NO has to be verified by direct measurement of NO with a NO_x-analyzer or detection of NO by its molecular mass using mass spectrometry. The chemical reaction of NO with oxygen in water forming mainly nitrite (Ignarro *et al.*, 1993) might complicate the direct measurement of nitric oxide. The sole nitrogen source in the co-culture is N₂. Therefore, it can be speculated that the interaction between the two bacteria is not only one-sided because of the two partners only the *Beggiatoa* sp. are known to fix N₂ (Chapter 5). To investigate if a direct transfer of fixed nitrogen in the co-culture occurs, labeled nitrogen (¹⁵N₂) can be added and the label can be searched for in the *Pseudovibrio* sp. using mass spectrometry (e.g. nano secondary ion mass spectrometry).

We have shown that the *Pseudovibrio* sp. FO-BEG1 growing in pure culture under nutrient-poor conditions can also use typical substrates known from heterotrophic growth (Chapter 4.1). It still remains unclear if the observed growth is the same as typical heterotrophic growth only with less nutrients or if special regulating enzymes are needed. A proteomic comparison between bacteria grown under nutrient deficiency and grown under nutrient

affluence might provide more information on the growth and the regulation of enzymes under oligotrophic conditions.

The detailed analysis of substrates used during oligotrophic growth shows that the bacteria use multiple substrates at the same time under oligotrophic conditions (Chapter 4.1). However, only amino acids (which account only for a small fraction) were measured quantitatively and the overall respiration rate remains unknown. Therefore, the measurement of oxygen consumption (e.g. using microelectrode respiration chambers) under nutrient-poor conditions could allow quantification and calculation of the respiration rate under oligotrophic conditions. Furthermore, the fact that we have both metabolites (detected by ESI FT-ICR-MS) and genes (closed genome of *Pseudovibrio* sp. strain FO-BEG1) may allow us to connect metabolites with genes present in the investigated organism in the future. Maybe, this will also help predicting new metabolic pathways.

Eventually, several heterotrophic strains, which were able to grow under both nutrient deficiency and affluence, were isolated during our studies (Chapter 4.2). The possibility of heterotrophic bacteria to grow under nutrient-deficient conditions might be a more widespread feature. To test this hypothesis, common heterotrophic bacteria from bacterial strain collections can be ordered and their ability to grow under nutrient-poor conditions can be tested.

References

- Adak S, Aulak KS, Stuehr DJ. (2002a). Direct evidence for nitric oxide production by a nitric-oxide synthase-like protein from *Bacillus subtilis*. *Journal of Biological Chemistry* **277**: 16167–16171.
- Adak S, Bilwes AM, Panda K, Hosfield D, Aulak KS, McDonald JF, Tainer JA, Getzoff ED, Crane BR, Stuehr DJ. (2002b). Cloning, expression, and characterization of a nitric oxide synthase protein from *Deinococcus radiodurans*. *Proceedings of the National Academy of Sciences of the United States of America* **99**: 107–112.
- Agapie T, Suseno S, Woodward JJ, Stoll S, Britt RD, Marletta MA. (2009). NO formation by a catalytically self-sufficient bacterial nitric oxide synthase from *Sorangium cellulosum*. *Proceedings of the National Academy of Sciences of the United States of America* **106**: 16221–16226.
- Ahmad A, Barry JP, Nelson DC. (1999). Phylogenetic affinity of a wide, vacuolate, nitrate-accumulating *Beggiatoa* sp. from Monterey Canyon, California, with *Thioploca* spp. *Applied and Environmental Microbiology* **65**: 270–277.
- Ahmad A, Kalanetra KM, Nelson DC. (2006). Cultivated *Beggiatoa* spp. define the phylogenetic root of morphologically diverse, noncultured, vacuolate sulfur bacteria. *Canadian Journal of Microbiology* **52**: 591–598.
- Alderton WK, Cooper CE, Knowles RG. (2001). Nitric oxide synthases: Structure, function and inhibition. *Biochemical Journal* **357**: 593–615.
- Allredge AL, Cole JJ, Caron DA. (1986). Production of heterotrophic bacteria inhabiting macroscopic organic aggregates (marine snow) from surface waters. *Limnology and Oceanography* **31**: 68–78.
- Allredge AL, Gotschalk CC. (1990). The relative contribution of marine snow of different origins to biological processes in coastal waters. *Continental Shelf Research* **10**: 41–58.
- Ammerman JW, Fuhrman JA, Hagström Å, Azam F. (1984). Bacterioplankton growth in seawater: 1. Growth kinetics and cellular characteristics in seawater cultures. *Marine Ecology Progress Series* **18**: 31–39.
- Arrigo KR. (2005). Marine microorganisms and global nutrient cycles. *Nature* **437**: 349–355.
- Azam F, Hodson RE. (1977). Size distribution and activity of marine microheterotrophs. *Limnology and Oceanography* **22**: 492–501.
- Azam F, Long RA. (2001). Sea snow microcosms. *Nature* **414**: 495–498.
- Azam F, Malfatti F. (2007). Microbial structuring of marine ecosystems. *Nature Reviews Microbiology* **5**: 782–791.
- Bachmann A. (2007). Charakterisierung des dominanten Begleitorganismus eines marinen *Beggiatoa* Konsortiums. Diploma thesis, Leibniz University of Hannover, Germany.

- Bada JL, Lee C. (1977). Decomposition and alteration of organic compounds dissolved in seawater. *Marine Chemistry* **5**: 523–534.
- Barry JP, Greene HG, Orange DL, Baxter CH, Robison BH, Kochevar RE, Nybakken JW, Reed DL, McHugh CM. (1996). Biologic and geologic characteristics of cold seeps in Monterey bay, California. *Deep-Sea Research Part I-Oceanographic Research Papers* **43**: 1739–1762.
- Battley EH. (1995). An apparent anomaly in the calculation of ash-free dry weights for the determination of cellular yields. *Applied and Environmental Microbiology* **61**: 1655–1657.
- Bauer JE, Williams PM, Druffel ERM. (1992). ¹⁴C activity of dissolved organic carbon fractions in the north-central Pacific and Sargasso Sea. *Nature* **357**: 667–670.
- Bernard C, Fenchel T. (1995). Mats of colourless sulphur bacteria 2. Structure, composition of biota and successional patterns. *Marine Ecology – Progress Series* **128**: 171–179.
- Bolin B, Crutzen PJ, Vitousek PM, Woodmansee RG, Goldberg ED, Cook RB. (1983). Interactions of biogeochemical cycles. In: Bolin B, Cook RB (eds). *The major biogeochemical cycles and their interactions*: Ipswich, UK. pp 1–39.
- Bollmann A, Lewis K, Epstein SS. (2007). Incubation of environmental samples in a diffusion chamber increases the diversity of recovered isolates. *Applied and Environmental Microbiology* **73**: 6386–6390.
- Bondarev V. (2007). Physiologische und molekulare Untersuchungen an filamentösen Bakterien aus Süßwasser-Standorten. Diploma thesis, Leibniz University of Hannover, Germany.
- Boylen CW, Ensign JC. (1970). Intracellular substrates for endogenous metabolism during long-term starvation of rod and spherical cells of *Arthrobacter crystallopoietes*. *Journal of Bacteriology* **103**: 578–587.
- Brock J, Schulz-Vogt HN. (2011). Sulfide induces phosphate release from polyphosphate in cultures of a marine *Beggiatoa* strain. *ISME Journal* **5**: 497–506.
- Brüchert V, Jørgensen BB, Neumann K, Riechmann D, Schlösser M, Schulz H. (2003). Regulation of bacterial sulfate reduction and hydrogen sulfide fluxes in the central Namibian coastal upwelling zone. *Geochimica Et Cosmochimica Acta* **67**: 4505–4518.
- Burke C, Steinberg P, Rusch D, Kjelleberg S, Thomas T. (2011). Bacterial community assembly based on functional genes rather than species. *Proceedings of the National Academy of Sciences of the United States of America* **108**: 14289–14293.
- Burton SD, Morita RY. (1964). Effect of catalase and cultural conditions on growth of *Beggiatoa*. *Journal of Bacteriology* **88**: 1755–1761.
- Button DK, Schut F, Quang P, Martin R, Robertson BR. (1993). Viability and isolation of marine bacteria by dilution culture: Theory, procedures, and initial results. *Applied and Environmental Microbiology* **59**: 881–891.

- Button DK, Robertson B, Gustafson E, Zhao XM. (2004). Experimental and theoretical bases of specific affinity, a cytoarchitecture-based formulation of nutrient collection proposed to supercede the Michaels-Menten paradigm of microbial kinetics. *Applied and Environmental Microbiology* **70**: 5511–5521.
- Carlucci AF, Shimp SL, Craven DB. (1986). Growth characteristics of low-nutrient bacteria from the Northeast and Central Pacific Ocean. *Fems Microbiology Ecology* **38**: 1–10.
- Chameides WL, Perdue EM. (1997). *Biogeochemical cycles - A computer-interactive study of earth systems and global change*. Oxford University Press: New York.
- Cherrier J, Bauer JE, Druffel ERM. (1996). Utilization and turnover of labile dissolved organic matter by bacterial heterotrophs in eastern north Pacific surface waters. *Marine Ecology – Progress Series* **139**: 267–279.
- Cherrier J, Bauer JE. (2004). Bacterial utilization of transient plankton-derived dissolved organic carbon and nitrogen inputs in surface ocean waters. *Aquatic Microbial Ecology* **35**: 229–241.
- Cohen Y, Rosenberg E. (1989). *Microbial mats - Physiological ecology of benthic microbial communities*: Washington, D.C.
- Connon SA, Giovannoni SJ. (2002). High-throughput methods for culturing microorganisms in very-low-nutrient media yield diverse new marine isolates. *Applied and Environmental Microbiology* **68**: 3878–3885.
- Cowie GL, Hedges JJ. (1992). Sources and reactivities of amino acids in a coastal marine environment. *Limnology and Oceanography* **37**: 703–724.
- de Albuquerque JP, Keim CN, Lins U. (2010). Comparative analysis of *Beggiatoa* from hypersaline and marine environments. *Micron* **41**: 507–517.
- del Giorgio PA, Cole JJ. (1998). Bacterial growth efficiency in natural aquatic systems. *Annual Review of Ecology and Systematics* **29**: 503–541.
- Dittmar T, Koch B, Hertkorn N, Kattner G. (2008). A simple and efficient method for the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater. *Limnology and Oceanography: Methods* **6**: 230–235.
- Dittmar T, Cherrier J, Ludwichowski K-U. (2009). The analysis of amino acids in seawater. In: Wurl O (ed). *Practical guidelines for the analysis of seawater*.
- Dittmar T, Paeng J. (2009). A heat-induced molecular signature in marine dissolved organic matter. *Nature Geoscience* **2**: 175–179.
- Ducklow HW, Steinberg DK, Buesseler KO. (2001). Upper ocean carbon export and the biological pump. *Oceanography* **14**: 50–58.
- Egli T. (2010). How to live at very low substrate concentration. *Water Research* **44**: 4826–4837.

- Enticknap JJ, Kelly M, Peraud O, Hill RT. (2006). Characterization of a culturable alphaproteobacterial symbiont common to many marine sponges and evidence for vertical transmission via sponge larvae. *Applied and Environmental Microbiology* **72**: 3724–3732.
- Farquhar GJ, Boyle WC. (1971). Identification of filamentous microorganisms in activated sludge. *Journal Water Pollution Control Federation* **43**: 604–622.
- Faust L, Wolfe RS. (1961). Enrichment and cultivation of *Beggiatoa alba*. *Journal of Bacteriology* **81**: 99–106.
- Fenchel T, Bernard C. (1995). Mats of colourless sulphur bacteria 1. Major microbial processes. *Marine Ecology – Progress Series* **128**: 161–170.
- Fiadeiro ME. (1983). Physical-chemical processes in the open ocean. In: Bolin B, Cook RB (eds). *The major biogeochemical cycles and their interactions*: Ipswich, UK. pp 461–476.
- Filippovich SY. (2010). Bacterial NO synthases. *Biochemistry-Moscow* **75**: 1217–1224.
- Fitzwater SE, Knauer GA, Martin JH. (1982). Metal contamination and its effect on primary production measurements. *Limnology and Oceanography* **27**: 544–551.
- Fossing H, Gallardo VA, Jørgensen BB, Hüttel M, Nielsen LP, Schulz H, Canfield DE, Forster S, Glud RN, Gundersen JK, Küver J, Ramsing NB, Teske A, Thamdrup B, Ulloa O. (1995). Concentration and Transport of Nitrate by the Mat-Forming Sulfur Bacterium *Thioploca*. *Nature* **374**: 713–715.
- Francis CA, Beman JM, Kuypers MMM. (2007). New processes and players in the nitrogen cycle: the microbial ecology of anaerobic and archaeal ammonia oxidation. *Isme Journal* **1**: 19–27.
- Fuhrman JA. (1987). Close coupling between release and uptake of dissolved free amino acids in seawater studied by an isotope dilution approach. *Marine Ecology – Progress Series* **37**: 45–52.
- Fuhrman JA. (1990). Dissolved free amino acid cycling in an estuarine outflow plume. *Marine Ecology – Progress Series* **66**: 197–203.
- Gauthier MJ, Lafay B, Christen R, Fernandez L, Acquaviva M, Bonin P, Bertrand JC. (1992). *Marinobacter hydrocarbonoclasticus* gen. nov., sp. nov., a new, extremely halotolerant, hydrocarbon-degrading marine bacterium. *International Journal of Systematic Bacteriology* **42**: 568–576.
- Girnth AC, Grünke S, Lichtschlag A, Felden J, Knittel K, Wenzhofer F, de Beer D, Boetius A. (2011). A novel, mat-forming *Thiomargarita* population associated with a sulfidic fluid flow from a deep-sea mud volcano. *Environmental Microbiology* **13**: 495–505.
- Grossart HP, Tang KW, Kiørboe T, Ploug H. (2007). Comparison of cell-specific activity between free-living and attached bacteria using isolates and natural assemblages. *Fems Microbiology Letters* **266**: 194–200.

- Grossart HP. (2010). Ecological consequences of bacterioplankton lifestyles: changes in concepts are needed. *Environmental Microbiology Reports* **2**: 706–714.
- Güde H, Strohl WR, Larkin JM. (1981). Mixotrophic and heterotrophic growth of *Beggiatoa alba* in continuous culture. *Archives of Microbiology* **129**: 357–360.
- Gusarov I, Nudler E. (2005). NO-mediated cytoprotection: Instant adaptation to oxidative stress in bacteria. *Proceedings of the National Academy of Sciences of the United States of America* **102**: 13855–13860.
- Hansell DA, Carlson CA, Repeta DJ, Schlitzer R. (2009). Dissolved organic matter in the ocean a controversy stimulates new insights. *Oceanography* **22**: 202–211.
- Henze A-T. (2005). Stickstofffixierung bei *Beggiatoa*. Diploma thesis, University of Hannover, Germany.
- Hertkorn N, Benner R, Frommberger M, Schmitt-Kopplin P, Witt M, Kaiser K, Kettrup A, Hedges JJ. (2006). Characterization of a major refractory component of marine dissolved organic matter. *Geochimica Et Cosmochimica Acta* **70**: 2990–3010.
- Hinck S, Neu TR, Lavik G, Mussmann M, de Beer D, Jonkers HM. (2007). Physiological adaptation of a nitrate-storing *Beggiatoa* sp. to diel cycling in a phototrophic hypersaline mat. *Applied and Environmental Microbiology* **73**: 7013–7022.
- Hinze G. (1901). Über den Bau und der Zellen von *Beggiatoa mirabilis* Cohn. *Berichte der Deutschen Botanischen Gesellschaft*: 369–374.
- Høgslund S, Revsbech NP, Kuenen JG, Jørgensen BB, Gallardo VA, van de Vossenberg JV, Nielsen JL, Holmkvist L, Arning ET, Nielsen LP. (2009). Physiology and behaviour of marine *Thioploca*. *Isme Journal* **3**: 647–657.
- Hopkinson CS, Vallino JJ. (2005). Efficient export of carbon to the deep ocean through dissolved organic matter. *Nature* **433**: 142–145.
- Hube AE, Heyduck-Söllner B, Fischer U. (2009). Phylogenetic classification of heterotrophic bacteria associated with filamentous marine cyanobacteria in culture. *Systematic and Applied Microbiology* **32**: 256–265.
- Huettel M, Forster S, Klöser S, Fossing H. (1996). Vertical migration in the sediment-dwelling sulfur bacteria *Thioploca* spp. in overcoming diffusion limitations. *Applied and Environmental Microbiology* **62**: 1863–1872.
- Hugenholtz P. (2002). Exploring prokaryotic diversity in the genomic era. *Genome Biology* **3**: 1–8.
- Huu NB, Denner EBM, Ha DTC, Wanner G, Stan-Lotter H. (1999). *Marinobacter aquaeolei* sp. nov., a halophilic bacterium isolated from a Vietnamese oil-producing well. *International Journal of Systematic Bacteriology* **49**: 367–375.
- Ignarro LJ, Fukuto JM, Griscavage JM, Rogers NE, Byrns RE. (1993). Oxidation of nitric oxide in aqueous solution to nitrite but not nitrate: Comparison with enzymatically formed

nitric oxide from L-arginine. *Proceedings of the National Academy of Sciences of the United States of America* **90**: 8103–8107.

Ihssen J, Egli T. (2005). Global physiological analysis of carbon- and energy-limited growing *Escherichia coli* confirms a high degree of catabolic flexibility and preparedness for mixed substrate utilization. *Environmental Microbiology* **7**: 1568–1581.

Ishida Y, Kadota H. (1981). Growth patterns and substrate requirements of naturally occurring obligate oligotrophs. *Microbial Ecology* **7**: 123–130.

Ishida Y, Imai I, Miyagaki T, Kadota H. (1982). Growth and uptake kinetics of a facultatively oligotrophic bacterium at low nutrient concentrations. *Microbial Ecology* **8**: 23–32.

Ishida Y, Fukami K, Eguchi M, Yoshinaga I. (1989). Strategies for growth of oligotrophic bacteria in the pelagic environment. In: Hattori T, Ishida Y, Maruyama Y, Morita RY, Uchida A (eds). *Recent advances in microbial ecology*. Japan Scientific Society Press: Japan. pp 89–93.

Jiao N, Herndl GJ, Hansell DA, Benner R, Kattner G, Wilhelm SW, Kirchman DL, Weinbauer MG, Luo TW, Chen F, Azam F. (2010). Microbial production of recalcitrant dissolved organic matter: long-term carbon storage in the global ocean. *Nature Reviews Microbiology* **8**: 593–599.

Jørgensen BB. (1977). Distribution of colorless sulfur bacteria (*Beggiatoa* spp.) in a coastal marine sediment. *Marine Biology* **41**: 19–28.

Jørgensen BB. (1983). Processes at the sediment-water interface. In: Bolin B, Cook RB (eds). *The major biogeochemical cycles and their interactions*: Ipswich, UK. pp 477–509.

Jørgensen BB, Revsbech NP. (1983). Colorless sulfur bacteria, *Beggiatoa* spp. and *Thiovulum* spp. in O₂ and H₂S microgradients. *Applied and Environmental Microbiology* **45**: 1261–1270.

Jørgensen BB, Nelson DC. (2004). Sulfide oxidation in marine sediments: Geochemistry meets microbiology. In: Amend JP, Edwards KJ, Lyons TW (eds). *Sulfur biogeochemistry – Past and present*: Geological Society of America pp 63–81.

Jørgensen BB, Dunker R, Grünke S, Røy H. (2010). Filamentous sulfur bacteria, *Beggiatoa* spp., in arctic marine sediments (Svalbard, 79 degrees N). *Fems Microbiology Ecology* **73**: 500-513.

Jørgensen NOG. (1987). Free amino acids in lakes: Concentrations and assimilation rates in relation to phytoplankton and bacterial production. *Limnology and Oceanography* **32**: 97–111.

Kaeberlein T, Lewis K, Epstein SS. (2002). Isolating "uncultivable" microorganisms in pure culture in a simulated natural environment. *Science* **296**: 1127–1129.

Kalanetra KM, Huston SL, Nelson DC. (2004). Novel, attached, sulfur-oxidizing bacteria at shallow hydrothermal vents possess vacuoles not involved in respiratory nitrate accumulation. *Applied and Environmental Microbiology* **70**: 7487–7496.

- Kalanetra KM, Joye SB, Sunseri NR, Nelson DC. (2005). Novel vacuolate sulfur bacteria from the Gulf of Mexico reproduce by reductive division in three dimensions. *Environmental Microbiology* **7**: 1451–1460.
- Kamp A, Stief P, Schulz-Vogt HN. (2006). Anaerobic sulfide oxidation with nitrate by a freshwater *Beggiatoa* enrichment culture. *Applied and Environmental Microbiology* **72**: 4755–4760.
- Kamp A, Røy H, Schulz-Vogt HN. (2008). Video-supported analysis of *Beggiatoa* filament growth, breakage, and movement. *Microbial Ecology* **56**: 484–491.
- Keil F. (1912). Beiträge zur Physiologie der farblosen Schwefelbakterien. *Beiträge zur Biologie der Pflanzen* **11**: 335–372.
- Keil RG, Kirchman DL. (1999). Utilization of dissolved protein and amino acids in the northern Sargasso Sea. *Aquatic Microbial Ecology* **18**: 293–300.
- Kido Soule MC, Longnecker K, Giovannoni SJ, Kujawinski EB. (2010). Impact of instrument and experiment parameters on reproducibility of ultrahigh resolution ESI FT-ICR mass spectra of natural organic matter. *Organic Geochemistry* **41**: 725–733.
- Kiørboe T. (2000). Colonization of marine snow aggregates by invertebrate zooplankton: Abundance, scaling, and possible role. *Limnology and Oceanography* **45**: 479–484.
- Kiørboe T, Jackson GA. (2001). Marine snow, organic solute plumes, and optimal chemosensory behavior of bacteria. *Limnology and Oceanography* **46**: 1309–1318.
- Kirchman DL, Meon B, Ducklow HW, Carlson CA, Hansell DA, Steward GF. (2001). Glucose fluxes and concentrations of dissolved combined neutral sugars (polysaccharides) in the Ross Sea and Polar Front Zone, Antarctica. *Deep-Sea Research Part II* **48**: 4179–4197.
- Klas Z. (1937). Über den Formenkreis von *Beggiatoa mirabilis*. *Archives of Microbiology: Aus dem Ozeanographischen Institut Split und den Botanischen Institut Zagreb*: 312–320.
- Koch AL. (1997). Microbial physiology and ecology of slow growth. *Microbiology and Molecular Biology Reviews* **61**: 305–318.
- Koch BP, Witt MR, Engbrodt R, Dittmar T, Kattner G. (2005). Molecular formulae of marine and terrigenous dissolved organic matter detected by electrospray ionization Fourier transform ion cyclotron resonance mass spectrometry. *Geochimica Et Cosmochimica Acta* **69**: 3299–3308.
- Kolmonen E, Sivonen K, Rapala J, Haukka K. (2004). Diversity of cyanobacteria and heterotrophic bacteria in cyanobacterial blooms in Lake Joutikas, Finland. *Aquatic Microbial Ecology* **36**: 201–211.
- Kovárová-Kovar K, Egli T. (1998). Growth kinetics of suspended microbial cells: From single-substrate-controlled growth to mixed-substrate kinetics. *Microbiology and Molecular Biology Reviews* **62**: 646–666.

- Kuentzel LE. (1969). Bacteria, carbon dioxide, and algal blooms. *Journal Water Pollution Control Federation* **41**: 1737–1747.
- Larkin JM, Strohl WR. (1983). *Beggiatoa*, *Thiothrix*, and *Thioploca*. *Annual Review of Microbiology* **37**: 341–367.
- Larkin JM, Henk MC. (1996). Filamentous sulfide-oxidizing bacteria at hydrocarbon seeps of the Gulf of Mexico. *Microscopy Research & Technique* **33**: 23–31.
- Lendenmann U, Snozzi M, Egli T. (1996). Kinetics of the simultaneous utilization of sugar mixtures by *Escherichia coli* in continuous culture. *Applied and Environmental Microbiology* **62**: 1493–1499.
- Lichtsschlag A, Felden J, Brüchert V, Boetius A, de Beer D. (2010). Geochemical processes and chemosynthetic primary production in different thiotrophic mats of the Hakon Mosby Mud Volcano (Barents Sea). *Limnology and Oceanography* **55**: 931–949.
- Lim MH, Xu D, Lippard SJ. (2006). Visualization of nitric oxide in living cells by a copper-based fluorescent probe. *Nature Chemical Biology* **2**: 375–380.
- Lindroth P, Mopper K. (1979). High-performance liquid chromatographic determination of subpicomole amounts of amino acids by precolumn fluorescence derivatization with orthophthaldialdehyde. *Analytical Chemistry* **51**: 1667–1674.
- Ludwig W, Strunk O, Westram R, Richter L, Meier H, Yadhukumar, Buchner A, Lai T, Steppi S, Jobb G, Forster W, Brettske I, Gerber S, Ginhart AW, Gross O, Grumann S, Hermann S, Jost R, König A, Liss T, Lussmann R, May M, Nonhoff B, Reichel B, Strehlow R, Stamatakis A, Stuckmann N, Vilbig A, Lenke M, Ludwig T, Bode A, Schleifer KH. (2004). ARB: A software environment for sequence data. *Nucleic Acids Research* **32**: 1363–1371.
- Lunau M, Lemke A, Dellwig O, Simon M. (2006). Physical and biogeochemical controls of microaggregate dynamics in a tidally affected coastal ecosystem. *Limnology and Oceanography* **51**: 847–859.
- Macalady JL, Lyon EH, Koffman B, Albertson LK, Meyer K, Galdenzi S, Mariani S. (2006). Dominant microbial populations in limestone-corroding stream biofilms, Frasassi cave system, Italy. *Applied and Environmental Microbiology* **72**: 5596–5609.
- MacDonell MT, Hood MA. (1982). Isolation and characterization of ultramicrobacteria from a Gulf Coast Estuary. *Applied and Environmental Microbiology* **43**: 566–571.
- Maier S, Murray RGE. (1965). Fine Structure of *Thioploca Ingrida* and a comparison with *Beggiatoa*. *Canadian Journal of Microbiology* **11**: 645–655.
- McHatton SC, Barry JP, Jannasch HW, Nelson DC. (1996). High nitrate concentrations in vacuolate, autotrophic marine *Beggiatoa* spp. *Applied and Environmental Microbiology* **62**: 954–958.
- Millero FJ, Plese T, Fernandez M. (1988). The dissociation of hydrogen sulfide in seawater. *Limnology and Oceanography* **33**: 269–274.

- Møller MM, Nielsen LP, Jørgensen BB. (1985). Oxygen responses and mat formation by *Beggiatoa* spp. *Applied and Environmental Microbiology* **50**: 373–382.
- Morris JJ, Kirkegaard R, Szul MJ, Johnson ZI, Zinser ER. (2008). Facilitation of robust growth of *Prochlorococcus* colonies and dilute liquid cultures by "helper" heterotrophic bacteria. *Applied and Environmental Microbiology* **74**: 4530–4534.
- Mußmann M, Schulz HN, Strotmann B, Kjær T, Nielsen LP, Rosselló-Mora RA, Amann RI, Jørgensen BB. (2003). Phylogeny and distribution of nitrate-storing *Beggiatoa* spp. in coastal marine sediments. *Environmental Microbiology* **5**: 523–533.
- Muyzer G, Teske A, Wirsen CO, Jannasch HW. (1995). Phylogenetic relationships of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gradient gel-electrophoresis of 16S rDNA fragments. *Archives of Microbiology* **164**: 165–172.
- Nelson DC, Castenholz RW. (1981). Use of reduced sulfur-compounds by *Beggiatoa* sp. *Journal of Bacteriology* **147**: 140–154.
- Nelson DC, Castenholz RW. (1982). Light responses of *Beggiatoa*. *Archives of Microbiology* **131**: 146–155.
- Nelson DC, Waterbury JB, Jannasch HW. (1982). Nitrogen fixation and nitrate utilization by marine and fresh-water *Beggiatoa*. *Archives of Microbiology* **133**: 172–177.
- Nelson DC, Jannasch HW. (1983). Chemoautotrophic growth of a marine *Beggiatoa* in sulfide-gradient cultures. *Archives of Microbiology* **136**: 262–269.
- Nelson DC, Jørgensen BB, Revsbech NP. (1986a). Growth-pattern and yield of a chemoautotrophic *Beggiatoa* sp. in oxygen-sulfide microgradients. *Applied and Environmental Microbiology* **52**: 225–233.
- Nelson DC, Revsbech NP, Jørgensen BB. (1986b). Microoxic-anoxic niche of *Beggiatoa* spp. - microelectrode survey of marine and fresh water strains. *Applied and Environmental Microbiology* **52**: 161–168.
- Nelson DC, Wirsen CO, Jannasch HW. (1989). Characterization of large, autotrophic *Beggiatoa* spp. abundant at hydrothermal vents of the Guaymas Basin. *Applied and Environmental Microbiology* **55**: 2909–2917.
- Norland S, Fagerbakke KM, Heldal M. (1995). Light element analysis of individual bacteria by X-ray microanalysis. *Applied and Environmental Microbiology* **61**: 1357–1362.
- Novitsky JA, Morita RY. (1976). Morphological characterization of small cells resulting from nutrient starvation of a psychrophilic marine vibrio. *Applied and Environmental Microbiology* **32**: 617–622.
- Novitsky JA, Morita RY. (1977). Survival of a psychrophilic marine vibrio under long-term nutrient starvation. *Applied and Environmental Microbiology* **33**: 635–641.

- Paerl HW. (1984). Transfer of N₂ and CO₂ fixation products from *Anabaena oscillarioides* to associated bacteria during inorganic carbon sufficiency and deficiency. *Journal of Phycology* **20**: 600–608.
- Paerl HW, Pinckney JL. (1996). A mini-review of microbial consortia: Their roles in aquatic production and biogeochemical cycling. *Microbial Ecology* **31**: 225–247.
- Palinska KA, Becker C, Krumbein WE. (1999). Isolation and purification techniques for benthic marine cyanobacteria - biotechnology potential. *Bulletin de l'Institute océanographique, Monaco* **19**: 585–592.
- Preisler A, Lichtschlag A, Lavik G, Boetius A, Jørgensen BB, de Beer D. (2007). Biological and chemical sulfide oxidation in a *Beggiatoa* inhabited marine sediment. *ISME Journal* **1**: 341–353.
- Pringsheim EG. (1964). Heterotrophism and species concepts in *Beggiatoa*. *American Journal of Botany* **51**: 898–913.
- Pringsheim EG. (1967). Die Mixotrophie von *Beggiatoa*. *Archiv Für Mikrobiologie* **59**: 247–254.
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig WG, Peplies J, Glöckner FO. (2007). SILVA: A comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Research* **35**: 7188C7196.
- Rappe MS, Connon SA, Vergin KL, Giovannoni SJ. (2002). Cultivation of the ubiquitous SAR11 marine bacterioplankton clade. *Nature* **418**: 630–633.
- Richardson LL. (1996). Horizontal and vertical migration patterns of *Phormidium corallyticum* and *Beggiatoa* spp. associated with black-band disease of corals. *Microbial Ecology* **32**: 323–335.
- Sahling H, Rickert D, Lee RW, Linke P, Suess E. (2002). Macrofaunal community structure and sulfide flux at gas hydrate deposits from the Cascadia convergent margin, NE Pacific. *Marine Ecology – Progress Series* **231**: 121–138.
- Salman V, Amann R, Girnth A-C, Polerecky L, Bailey JV, Høgslund S, Jessen G, Pantoja S, Schulz-Vogt HN. (2011). A single-cell sequencing approach to the classification of large, vacuolated sulfur bacteria. *Systematic and Applied Microbiology* **34**: 243–259.
- Salomon PS, Janson S, Graneli E. (2003). Molecular identification of bacteria associated with filaments of *Nodularia spumigena* and their effect on the cyanobacterial growth. *Harmful Algae* **2**: 261–272.
- Sassen R, Roberts HH, Aharon P, Larkin J, Chinn EW, Carney R. (1993). Chemosynthetic bacterial mats at cold hydrocarbon seeps, Gulf of Mexico continental slope. *Organic Geochemistry* **20**: 77–89.
- Sayama M. (2001). Presence of nitrate-accumulating sulfur bacteria and their influence on nitrogen cycling in a shallow coastal marine sediment. *Applied and Environmental Microbiology* **67**: 3481–3487.

- Sayama M, Risgaard-Petersen N, Nielsen LP, Fossing H, Christensen PB. (2005). Impact of bacterial NO³⁻ transport on sediment biogeochemistry. *Applied and Environmental Microbiology* **71**: 7575–7577.
- Schlünz B, Schneider RR. (2000). Transport of terrestrial organic carbon to the oceans by rivers: re-estimating flux- and burial rates. *International Journal of Earth Sciences* **88**: 599–606.
- Schmidt TM, Arieli B, Cohen Y, Padan E, Strohl WR. (1987). Sulfur metabolism in *Beggiatoa alba*. *Journal of Bacteriology* **169**: 5466–5472.
- Schreiber F, Beutler M, Enning D, Lamprecht-Grandio M, Zafra O, Gonzalez-Pastor JE, de Beer D. (2011). The role of nitric-oxide-synthase-derived nitric oxide in multicellular traits of *Bacillus subtilis* 3610: biofilm formation, swarming, and dispersal. *Bmc Microbiology* **11**: 1–11.
- Schulz HN, Brinkhoff T, Ferdelman TG, Marine MH, Teske A, Jørgensen BB. (1999). Dense populations of a giant sulfur bacterium in Namibian shelf sediments. *Science* **284**: 493–495.
- Schulz HN, de Beer D. (2002). Uptake rates of oxygen and sulfide measured with individual *Thiomargarita namibiensis* cells by using microelectrodes. *Applied and Environmental Microbiology* **68**: 5746–5749.
- Schut F, Devries EJ, Gottschal JC, Robertson BR, Harder W, Prins RA, Button DK. (1993). Isolation of typical marine bacteria by dilution culture: Growth, maintenance, and characteristics of isolates under laboratory conditions. *Applied and Environmental Microbiology* **59**: 2150–2160.
- Schut F, Prins RA, Gottschal JC. (1997). Oligotrophy and pelagic marine bacteria: Facts and fiction. *Aquatic Microbial Ecology* **12**: 177–202.
- Shanks AL, Edmondson EW. (1990). The vertical flux of metazoans (holoplankton, meiofauna, and larval invertebrates) due to their association with marine snow. *Limnology and Oceanography* **35**: 455–463.
- Shieh WY, Lin YT, Jean WD. (2004). *Pseudovibrio denitrificans* gen. nov., sp. nov., a marine, facultatively anaerobic, fermentative bacterium capable of denitrification. *International Journal of Systematic and Evolutionary Microbiology* **54**: 2307–2312.
- Silver MW, Shanks AL, Trent JD. (1978). Marine snow - microplankton habitat and source of small-scale patchiness in pelagic populations. *Science* **201**: 371–373.
- Simon M, Grossart HP, Schweitzer B, Ploug H. (2002). Microbial ecology of organic aggregates in aquatic ecosystems. *Aquatic Microbial Ecology* **28**: 175–211.
- Smith DC, Simon M, Alldredge AL, Azam F. (1992). Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature* **359**: 139–142.
- Stanley SO, Boto KG, Alongi DM, Gillan FT. (1987). Composition and bacterial utilization of free amino acids in tropical mangrove sediments. *Marine Chemistry* **22**: 13–30.

- Strohl WR, Larkin JM. (1978). Enumeration, isolation, and characterization of *Beggiatoa* from freshwater sediments. *Applied and Environmental Microbiology* **36**: 755–770.
- Strohl WR, Cannon GC, Shively JM, Güde H, Hook LA, Lane CM, Larkin JM. (1981). Heterotrophic Carbon Metabolism by *Beggiatoa alba*. *Journal of Bacteriology* **148**: 572–583.
- Strohl WR, Howard KS, Larkin JM. (1982). Ultrastructure of *Beggiatoa alba* strain B15LD. *Journal of General Microbiology* **128**: 73–84.
- Strous M, Kuenen JG, Jetten MSM. (1999). Key physiology of anaerobic ammonium oxidation. *Applied and Environmental Microbiology* **65**: 3248–3250.
- Suess E. (1980). Particulate organic-carbon flux in the oceans - surface productivity and oxygen utilization. *Nature* **288**: 260–263.
- Suzuki N, Kato K. (1953). Studies on suspended materials *marnie snow* in the sea. Part I. sources of *marine snow*. *Bull Fac Fish Hokkaido Univ*: 132–137.
- Sweerts JPRA, De Beer D, Nielsen LP, Verdouw H, Van den Heuvel JC, Cohen Y, Cappenberg TE. (1990). Denitrification by sulphur oxidizing *Beggiatoa* spp. mats on freshwater sediments. *Nature* **344**: 762–763.
- Tabor PS, Neihof RA. (1982). Improved micro-auto-radiographic method to determine individual microorganisms active in substrate uptake in natural waters. *Applied and Environmental Microbiology* **44**: 945–953.
- Tapley DW, Buettner GR, Shick JM. (1999). Free radicals and chemiluminescence as products of the spontaneous oxidation of sulfide in seawater, and their biological implications. *Biological Bulletin* **196**: 52–56.
- Trevisan V. (1842). *Prospetto della Flora Euganea*. Padua: *Coi Tipi del Seminario*: 1–68.
- Turley CM, Mackie PJ. (1994). Biogeochemical significance of attached and free-living bacteria and the flux of particles in the NE Atlantic Ocean. *Marine Ecology – Progress Series* **115**: 191–203.
- Turley CM, Stutt ED. (2000). Depth-related cell-specific bacterial leucine incorporation rates on particles and its biogeochemical significance in the Northwest Mediterranean. *Limnology and Oceanography* **45**: 419–425.
- Upton AC, Nedwell DB. (1989). Nutritional flexibility of oligotrophic and copiotrophic antarctic bacteria with respect to organic substrates. *Fems Microbiology Ecology* **62**: 1–6.
- Van der Kooij D, Visser A, Hijnen WAM. (1980). Growth of *Aeromonas hydrophila* at low concentrations of substrates added to tap water. *Applied and Environmental Microbiology* **39**: 1198–1204.
- van Gemerden H. (1993). Microbial mats - a joint venture. *Marine Geology* **113**: 3–25.

- Vancanneyt M, Schut F, Snauwaert C, Goris J, Swings J, Gottschal JC. (2001). *Sphingomonas alaskensis* sp. nov., a dominant bacterium from a marine oligotrophic environment. *International Journal of Systematic and Evolutionary Microbiology* **51**: 73–80.
- Vartoukian SR, Palmer RM, Wade WG. (2010). Strategies for culture of 'unculturable' bacteria. *Fems Microbiology Letters* **309**: 1–7.
- Vaucher JP. (1803). Histoire des conferves d'eau douce, contenant leurs différents modes de reproduction, et la description de leurs principales espèces. *Paschoud, Geneva*: 1–285.
- Volkov II, Rozanov AG. (1983). The sulphur cycle in the oceans. Part 1. Reservoirs and Fluxes of sulphur. In: Ivanov MV, Freney JR (eds). *The global biogeochemical sulfur cycle*. Wiley: Chichester.
- Weber CA. (1907). Aufbau und Vegetation der Moore Norddeutschlands. *Botanische Jahrbuecher fuer Systematik, Pflanzengeschichte und Pflanzengeographie / Beiblatt*. pp 19–34.
- Weiss RF. (1970). Solubility of nitrogen, oxygen and argon in water and seawater. *Deep-Sea Research* **17**: 721–735.
- Winogradsky S. (1887). Ueber Schwefelbakterien. *Botanische Zeitung* **45**: 489–507, 513–523, 529–539, 545–559, 569–576, 585–594, 606–610.
- Yanagita T, Ichikawa T, Tsuji T, Kamata Y, Ito K, Sasaki M. (1977). Two trophic groups of bacteria, oligotrophs and eutrophs: Their distribution in fresh and sea water areas in the central northern Japan. *J Gen Appl Microbiol* **24**: 59–88.
- Zengler K, Toledo G, Rappe M, Elkins J, Mathur EJ, Short JM, Keller M. (2002). Cultivating the uncultured. *Proceedings of the National Academy of Sciences of the United States of America* **99**: 15681–15686.
- Zevenhuizen LPTM. (1966). Formation and function of glycogen-like polysaccharide of *Arthrobacter*. *Antonie Van Leeuwenhoek Journal of Microbiology and Serology* **32**: 356–372.
- Zweifel UL, Norrman B, Hagström Å. (1993). Consumption of dissolved organic carbon by marine bacteria and demand for inorganic nutrients. *Marine Ecology – Progress Series* **101**: 23–32.

List of abbreviations

AHL	N-acyl-homoserine lactone
ATP	adenosine-5'-triphosphate
BGE	bacterial growth efficiency
BP	bacterial production
BR	bacterial respiration
cAMP	cyclic adenosine monophosphate
DCAA	dissolved combined amino acids
DFAA	dissolved free amino acids
DOC	dissolved organic carbon
DOM	dissolved organic matter
DSM	Deutsche Sammlung von Mikroorganismen (German Collection of Microorganisms)
DSMZ	Deutsche Sammlung von Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell cultures)
ESI	electrospray ionization
FT-ICR-MS	Fourier transform ion cyclotron resonance mass spectrometry
HPLC	high performance liquid chromatography
LDOM	labile dissolved organic matter
NO	nitric oxide
NOS	nitric oxide synthases
NRPS	non ribosomal polypeptide synthases
PEG	polyethylene glycol
PHA	polyhydroxyalkanoates
PHB	poly- β -hydroxybutyric acid
PKS	polyketide synthases
POC	particulate organic carbon
POM	particulate organic matter
RDOM	recalcitrant dissolved organic matter
ROS	reactive oxygen species
THDAA	total hydrolysable dissolved amino acids

Appendix

The genus *Pseudovibrio* contains metabolically versatile and symbiotically interacting bacteria

Vladimir Bondarev^{1*}, Michael Richter¹, Jörn Piel², Stefano Romano¹, Anne Schwedt¹ and Heide N. Schulz-Vogt¹

¹Max Planck Institute for Marine Microbiology, Bremen, Germany

²Kekulé-Institut für Organische Chemie und Biochemie, Bonn, Germany

*Corresponding author

Manuscript in preparation

Keywords: genomics/functional genomics/comparative genomics; metabolism; microbe:microbe interactions; primary and secondary production; symbionts; microbe:higher organism interaction

Abstract

The majority of strains belonging to the genus *Pseudovibrio* have been isolated from marine invertebrates like tunicates, corals and especially sponges, but the physiology of these bacteria is poorly understood. In this study, we analyze the genomes of two *Pseudovibrio* strains. One is a required symbiont of a cultivated *Beggiatoa* strain, a sulfide oxidizing, autotrophic bacterium. The other one was isolated from a sponge (Enticknap *et al.*, 2006). The data show that both strains are generalistic bacteria capable of importing and oxidizing a wide range of organic and inorganic compounds to meet their carbon, nitrogen, phosphorous and energy requirements under oxic and anoxic conditions. Several physiological traits encoded in the genome were verified in laboratory experiments with a pure culture of the *Pseudovibrio* strain originally associated with *Beggiatoa*. Besides the versatile metabolic abilities of both *Pseudovibrio* strains, our study reveals a number of open reading frames and gene clusters in the genomes that seem to be involved in symbiont-host interactions. *Pseudovibrio* has the genomic potential to attach to host cells, might be capable of interacting with the eukaryotic cell machinery, produce secondary metabolites and may supply the host with cofactors.

Introduction

The first strain of the genus *Pseudovibrio* has been isolated from coastal seawater in 2004 and was described as *Pseudovibrio denitrificans* – a marine, heterotrophic, facultatively anaerobic bacterium capable of denitrification and fermentation (Shieh *et al.*, 2004). Two further type strains, *P. ascidiaceicola* (Fukunaga *et al.*, 2006) and *P. japonicus* (Hosoya and Yokota, 2007), were isolated from a tunicate and coastal seawater, respectively. Physiologically, these isolates were not notably different from *P. denitrificans*. Besides the three type strains, *Pseudovibrio* spp.-related bacteria have been found in various studies throughout the world either by 16S rRNA gene analysis or direct isolation methods (Hentschel *et al.*, 2001; Webster and Hill, 2001; Olson *et al.*, 2002; Thakur *et al.*, 2003; Thiel and Imhoff, 2003; Thoms *et al.*, 2003; Agogu e *et al.*, 2005; Lafi *et al.*, 2005; Enticknap *et al.*, 2006; Koren and Rosenberg, 2006; Sertan-de Guzman *et al.*, 2007; Muscholl-Silberhorn *et al.*, 2008; Riesenfeld *et al.*, 2008; Kennedy *et al.*, 2009; Rypien *et al.*, 2010; Santos *et al.*, 2010). Interestingly, besides *P. denitrificans*, *P. japonicus* and a *Pseudovibrio* spp.-related isolate from coastal, oligotrophic seawater (Agogu e *et al.*, 2005), all other strains belonging to this genus discovered until today have been found closely associated with marine invertebrates like tunicates, corals, and sponges. Especially Porifera seem to harbor *Pseudovibrio* populations, e.g., as the dominating species of the culturable bacterial community (Webster and Hill, 2001; Muscholl-Silberhorn

et al., 2008). Additionally, *Pseudovibrio* has been found in sponge larvae as the most abundant prokaryote, indicating vertical transmission of these bacteria in their hosts (Enticknap *et al.*, 2006). Such a consistent pattern of *Pseudovibrio* spp. associated with sponges suggests that they are symbionts of those metazoa (Webster and Hill, 2001; Enticknap *et al.*, 2006). Whether the nature of this symbiosis is mutualistic/ commensalistic or whether *Pseudovibrio* spp. rather represent pathogens/parasites is uncertain, but the fact that *Pseudovibrio* spp. have been isolated only from healthy sponges indicates that the bacteria do not harm the host and might be even required for its health (Webster and Hill, 2001). Another shared feature is the production of secondary metabolites by many of the cultured *Pseudovibrio* strains. For instance, heptylprodigiosin, a compound that exhibits antimicrobial activity, was isolated from a pure culture of *P. denitrificans* Z143-1 (Sertan-de Guzman *et al.*, 2007) and the production of additional bioactive compounds could be shown in several other studies (Hentschel *et al.*, 2001; Muscholl-Silberhorn *et al.*, 2008; Kennedy *et al.*, 2009; Santos *et al.*, 2010).

Despite the fact that members of the genus *Pseudovibrio* seem to be ubiquitous and important associates of marine invertebrates and are also found free-living, very little is known about the physiology and interactions with the host. In this study, we analyze the genomes of two *Pseudovibrio* strains. *Pseudovibrio* sp. FO-BEG1 has been isolated from an enrichment culture of a *Beggiatoa* strain, a filamentous, sulfide oxidizing bacterium (Brock and Schulz-Vogt, 2011; Chapters 2 and 3). Initially, this *Beggiatoa* strain was sampled from a coral suffering from the black band disease off the coast of Florida, which indicates that the strain *Pseudovibrio* FO-BEG1 could have been associated with the coral at the time of sampling – either in a commensalistic/mutualistic or pathogenic relationship – and is now available as an axenic culture in our lab. Intriguingly, strain FO-BEG1 is also maintained in a co-culture with a *Beggiatoa* sp., which seems to be unable to grow without *Pseudovibrio* and is therefore dependent on strain FO-BEG1. The second strain, *Pseudovibrio* sp. JE062, has been isolated in Florida from the sponge *Mycale laxissima* in the year 2006 and was described as a sponge symbiont by Enticknap *et al.* (2006). The analysis of these genomes gives us an insight into the physiological and symbiotic potential of both *Pseudovibrio* strains and reveals fascinating microorganisms that seem to be adapted to free-living and symbiotic life styles.

Materials and Methods

Growth conditions

For aerobic growth CM medium modified after Shieh *et al.* (2004) was used. After autoclaving, the medium was supplemented with K_2HPO_4 (1.15 mmol L^{-1}), glucose (10 mmol L^{-1} unless stated otherwise), 1 mL L^{-1} tungsten/selenium solution (Brysch *et al.*, 1987), 1 mL L^{-1} trace elements (Widdel and Pfennig, 1984), and 1 mL L^{-1} of four vitamin solutions prepared according to Aeckersberg *et al.* (1991). For measurement of SO_4^{2-} evolution during $\text{S}_2\text{O}_3^{2-}$ oxidation, 10 mmol L^{-1} $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5 \text{H}_2\text{O}$ and 5 mmol L^{-1} glucose were added and 2 g L^{-1} K_2SO_4 from the original recipe was replaced with 0.75 g L^{-1} KCl. To compare growth between a culture with and without $\text{Na}_2\text{S}_2\text{O}_3$, K_2SO_4 was not omitted from the medium and glucose and $\text{Na}_2\text{S}_2\text{O}_3 \cdot 5 \text{H}_2\text{O}$ were used in the same concentrations as described above. To investigate the growth with 4-hydroxybenzoic acid and benzoate, both compounds were added in a concentration of 2 mmol L^{-1} , respectively, without any other carbon source. Growth with phosphonoacetate (1 mmol L^{-1}) as phosphorus source was tested by adding this compound as the only phosphorus source and all vitamins were omitted from the medium. For fermentation and denitrification experiments under anoxic conditions, aged North Sea water was buffered with 50 mmol L^{-1} TRIS, supplemented with NH_4Cl (10 mmol L^{-1}) and the pH adjusted to 8. Preparation of the medium was performed according to Widdel and Bak (1992) in order to prepare the medium anoxically. Cooling was performed under N_2 atmosphere, except for experiments in which N_2 production was measured, in which Ar was used as the atmosphere instead. After autoclaving, the medium was supplemented with 10 mmol L^{-1} glucose, 1 mL L^{-1} tungsten/selenium solution, 1 mL L^{-1} trace elements and 1 mL L^{-1} of four vitamin solutions prepared as described above. NaNO_3 (10 mmol L^{-1}) was added for experiments investigating denitrification. To test CO oxidation, CM medium was prepared as described above, containing 400 $\mu\text{mol L}^{-1}$ glucose and supplied with 500 p.p.m. CO to the bottle headspace. For aerobic growth experiments, 250 mL Erlenmeyer flasks were filled with 100 mL medium. For anaerobic growth, 156 mL serum bottles (Wheaton, Millville, USA) were filled anoxically with 50 mL medium and closed with butyl rubber stoppers. For all experiments, 0.1% or 0.5% of a preculture grown aerobically in CM medium was used as inoculum. All growth experiments were performed with *Pseudovibrio* sp. FO-BEG1 in triplicates at 28°C in the dark with shaking at 110 rpm.

Chemical analyses

Bacterial growth was monitored as the optical density ($\text{OD}_{600 \text{ nm}}$) using an Eppendorf BioPhotometer (Eppendorf AG, Hamburg, Germany). SO_4^{2-} was measured with a Metrohm

761 Compact IC with conductivity detector (Metrohm AG, Herisau, Switzerland) equipped with a Metrosep A Supp 5-100 column with a carbonate eluent (3.2 mmol L⁻¹ Na₂CO₃/1 mmol L⁻¹ NaHCO₃ in deionised water) at a flow rate of 0.7 mL min⁻¹. Tetrathionate was measured according to Kamyshny (2009). Glucose and organic acids were determined using a HPLC system (Sykam GmbH) equipped with an anion neutral pre-column (4x20 mm; Sykam GmbH) and an Aminex HPX-87H separation column (300x7.8 mm; Biorad, Munich, Germany) at a temperature of 60 °C. The eluent consisted of 5 mM H₂SO₄ in HPLC-grade water with a flow rate of 0.6 mL min⁻¹. Quantification of glucose, succinate, lactate, formate, acetate, propionate and ethanol was performed with the 7515A RI detector (ERC, Riemerling, Germany); pyruvate was measured with the Sapphire UV-Vis detector at 210 nm (Ecom, Praha, Czech Republic). NO₃⁻ was quantified with a HPLC system (Sykam GmbH, Eresing, Germany) containing an anion neutral pre-column (4x20 mm; Sykam GmbH) and an IBJ A3 anion separation column (4x60 mm; Sykam GmbH) with a column temperature of 50 °C. The eluent consisted of 25 mmol L⁻¹ NaCl and 45% ethanol in deionised water with a flow rate of 1 mL min⁻¹. Detection of NO₃⁻ was conducted with Linear Uvis 200 (Thermo Fischer Scientific GmbH, Dreieich, Germany) at 220 nm. N₂ was measured as described by Zedelius *et al.* (2011). CO determination was conducted with a Shimadzu GC-8A (Shimadzu, Duisburg, Germany) gas chromatograph with a Molecular Sieve 5A column (80 to 100; 0.125 in. by 2 m; Restek, Bellefonte, USA) at a flow of 20 mL of synthetic air per minute at 40°C and an RGD2 reduction gas detector (Trace Analytical, Menlo Park, USA).

DNA extraction and sequencing

DNA was extracted from strain FO-BEG1 using the Fast DNA SPIN Kit for Soil (MP Biomedicals LLC, Illkirch, France), according to manufacturers' instructions. 454 sequencing was conducted by LGC Genomics GmbH with a 454 GS FLX System. The Newbler 2.0.00.22 software was used for sequence assembly and quality assessment. Overall, 522919 sequenced reads with an average length of 336.30 bp lead to a 29-fold sequence coverage. In order to close the gaps, a fosmid library with a 1.5-fold physical coverage was created and used for direct sequencing of the fosmid clones. For the residual gaps, 96 specific primers were designed and used for combinatorial PCR on DNA level, the products of which were sequenced via the Sanger method.

Gene prediction, annotation and data mining

Gene prediction was carried out by using the software Glimmer3 (Delcher *et al.*, 2007). Ribosomal RNA genes were detected by using the RNAmmer 1.2 software (Lagesen *et al.*, 2007) and transfer RNAs by tRNAscan-SE (Lowe and Eddy, 1997). Annotation was

performed by using the GenDB, version 2.2 system (Meyer *et al.*, 2003), supplemented by the tool JCoast, version 1.6 (Richter *et al.*, 2008). For each predicted open reading frame (ORF) observations have been collected from similarity searches against sequence databases NCBI-nr, Swiss-Prot, KEGG and genomesDB (Richter *et al.*, 2008) and for protein family databases from Pfam (Bateman *et al.*, 2004) and InterPro (Mulder *et al.*, 2005). SignalP has been used for signal peptide predictions (Bendtsen *et al.*, 2004) and TMHMM for transmembrane helix-analysis (Krogh *et al.*, 2001). Predicted protein coding sequences were automatically annotated by the in-house software MicHanThi (Quast, 2006). The MicHanThi software predicts gene functions based on similarity searches using the NCBI-nr (including Swiss-Prot) and InterPro database. The annotation of proteins highlighted within the scope of this study was subject of manual inspection. For all observations regarding putative protein functions, an e-value cutoff of 10^{-4} was considered.

Comparison of the shared gene content by reciprocal best matches (RBMs) and functional classification with Kyoto encyclopedia of genes and genomes (KEGG)

Determination of the shared gene content has been performed by a BLAST all versus all search between FO-BEG1 and JE062. Reciprocal best matches were counted by a BLAST result with an E value $<1e^{-5}$ each and a subject coverage of over 65%. For metabolic pathway identification, genes were searched for similarity against the KEGG database. A match was counted if the similarity search resulted in an expectation E value below $1e^{-5}$. All occurring KO (KEGG Orthology) numbers were mapped against KEGG pathway functional hierarchies and statistical analyzed.

Functional classification with cluster of orthologous groups (COG) and calculation of the Average nucleotide identity (ANI)

All predicted ORFs were also searched for similarity against the COG database (Tatusov *et al.*, 2003). A match was counted if the similarity search resulted in an E value below $1e^{-5}$. ANI between the whole-genome sequences of strain FO-BEG1 and the draft genome sequences of strain JE062 was determined by using the *in silico* DNA-DNA hybridization method of the JSpecies (Richter and Rosselló-Móra, 2009) software with default parameters.

Creation of circular genome maps and prediction of ABC and TRAP type transporters

Comparative circular genome maps of the RBMs shared between JE062 and FO-BEG1 have been drawn by using JCoast's plugin for CGView (Stothard and Wishart, 2005). Circular GC-plot and GC-skew representation has been drawn by using DNAPlotter (Carver *et al.*, 2009). As initial step for the identification of the ABC transporters, genes containing the Pfam domain ABC_tran (PF00005) were detected in the genome of strain FO-BEG1. For the

identification of the permease and the periplasmic binding protein, the close proximity of genes containing the ABC_tran domain was searched. Only ABC systems with at least one ABC_tran domain, one permease and one periplasmic binding protein were regarded as functioning ABC transporters and substrate specificity was predicted from the annotations of the subunits. In several cases, one subunit (e.g. the permease) was missing in close proximity of genes with the ABC_tran domain. In this case, a single permease gene located on any place in the genome with the same substrate specificity prediction but not belonging to any complete ABC system, was used to complement the transporter system. TRAP transporters were regarded as complete when the subunits DctM, DctQ and DctP were present in close proximity. When two subunits were identified in close proximity and the third was missing, the single subunit located on any place in the genome not belonging to any complete TRAP system was used to complement the transporter system. In the case of fusion of the DctQ and M subunits in one gene, only the DctP subunit was required to complete the transporter.

Accession numbers

The genome shotgun project of strain FO-BEG1 has been deposited at DDBJ/EMBL/GenBank under the accession number CP003147 for the chromosome and CP003148 for the plasmid. The draft genome sequence of strain JE062 has the DDBJ/EMBL/GenBank accession number ABXL00000000.

Results and Discussion

General genome characteristics

The genome size of strain FO-BEG1 is 5.9 Mbp, including a large plasmid of 0.4 Mbp (Figure S.1). The circular chromosome of 5.5 Mbp contains a large stretch of repeats at position 2,707,040. This area of unknown size could not be bridged with a direct sequencing approach despite the presence of this area on a fosmid, indicating strong secondary structures, and has been masked with the ambiguous nucleotide code 'N'. The G+C content is 52.5 mol% and is consistent with the known values of the described *Pseudovibrio* isolates (Shieh *et al.*, 2004; Fukunaga *et al.*, 2006; Hosoya and Yokota, 2007). Altogether, we found 5,478 ORFs, 398 of which were located on the plasmid, which correspond to about 87% of encoding DNA. Six complete rRNA operons and 69 tRNA encoding regions were annotated, indicating the capability of a quick response to changing conditions and fast growth when nutrients are available. The genome of strain JE062 has not been closed, but there are 19 contigs available with an overall size of 5.7 Mbp, 5,225 ORFs and 52.4 mol% GC content, which is almost

identical to the genome of strain FO-BEG1 (Figure S.1 A and B). It contains 72 tRNA genes and seven complete rRNA operons. Unfortunately, the repeat-rich area that could not be sequenced in the genome of strain FO-BEG1 shows an ambiguous sequence in strain JE062 as well, and could therefore not be used to close the gap in FO-BEG1.




Figure S.1. Comparative circular map of *Pseudovibrio* sp. FO-BEG1 chromosome (A) and plasmid (B). Most outer lane represents the reciprocal best match (RBM)-shared gene content between FO-BEG1 and JE062. Lane two and three represent all predicted open reading frames (ORFs) on the lagging (red) and leading (green) strands. The two inner lanes display the GC-plot and the GC-skew. The red arrow indicates the area of unknown size that could not be closed during sequencing. The bar chart (C) express the amino acid percentage identity of each RBM shared gene-content between FO-BEG1 and JE062. The blue bar is representing the FO-BEG1 chromosome and orange the corresponding plasmid.

Even though the genome of JE062 is not completely closed we assume that it also contains a plasmid with similar content, since most of the genes identified on the plasmid of FO-BEG1 were allocated in the genome of JE062 (Figure S.1 B). Table S.1 shows an overview of the genome characteristics of both strains as well as the assignment of the genes to COGs. The shared gene content between FO-BEG1 and the draft genome of JE062 comprises 84.4% (4,287 ORFs, Figure S.1 C). An ANI analysis conducted between strains FO-BEG1 and JE062 revealed a 94.5% ANI_b (87% genome alignment) and 95.4% ANI_m (86% genome

alignment) value. The values are in the range of the proposed species definition boundary (Richter and Rosselló-Móra, 2009) indicating a species level degree of similarity.

Table S.1. General genome features of *Pseudovibrio* sp. FO-BEG1 and JE062, including categorization of the genes into cluster of orthologous group (COG) categories.

Characteristics	FO-BEG1	JE062
Base pairs	5,916,782	5,726,521
G+C content (%)	52.5	52.4
No. of protein-coding genes	5,478	5,225
Percent coding	85.5	85
No. of rRNA operons	6	7
No. of tRNA genes	69	72
COG category		
[J] Translation, ribosomal structure and biogenesis	196	190
[K] Transcription	387	367
[L] Replication, recombination and repair	135	125
[D] Cell cycle control, cell division, chromosome partitioning	21	20
[T] Signal transduction mechanism	138	140
[M] Cell wall/membrane/envelope biogenesis	190	176
[N] Cell motility	153	149
[O] Posttranslational modification, protein turnover, chaperones	135	127
[C] Energy production and conversion	245	245
[G] Carbohydrate transport and metabolism	323	311
[E] Amino acid transport and metabolism	507	492
[F] Nucleotide transport and metabolism	99	92
[H] Coenzyme transport and metabolism	185	181
[I] Lipid transport and metabolism	148	142
[P] Inorganic ion transport and metabolism	291	287
[Q] Secondary metabolites biosynthesis, transport and catabolism	210	204
[R] General function prediction only	598	576
[S] Function unknown	281	272

Physiology

In both genomes we found a number of genes that indicate high metabolic variety of *Pseudovibrio* FO-BEG1 and JE062. Degradation of carbohydrates is most likely performed via the Entner-Doudoroff pathway, which is present in both genomes, due to absence of the phosphofructokinase (PFK), a key enzyme of the glycolysis (Emden-Meyerhoff-Parnas), which is a regularly encountered phenomenon within marine α -*Proteobacteria* (Fürch *et al.*, 2009; Tang *et al.*, 2009; Williams *et al.*, 2009). Besides the PFK, all other enzymes involved

in glycolysis can be identified in both genomes, including fructose-1,6-bisphosphatase I, the key enzyme for glyconeogenesis, indicating that the Emden-Meyerhoff-Parnas pathway can be used for anabolic purposes (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000). Genes encoding all enzymes of the citric acid cycle and pentose phosphate pathway are present. Additionally, both strains have the genetic potential to degrade aromatic compounds via the β -keto adipate pathway, which we demonstrated by growing *Pseudovibrio* sp. FO-BEG1 with 4-hydroxybenzoate as the only carbon and energy source under aerobic conditions (Figure S 2.1 A). Benzoate, however, was not degraded, indicating that either the uptake of benzoate is detained or the hydroxylation of the aromatic ring structure cannot be performed by *Pseudovibrio* FO-BEG1. Under anoxic conditions without nitrate, strain FO-BEG1 metabolized glucose in mixed acid type fermentation, as suggested by the present genes in both strains (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000), resulting in acidification of the medium and formation of mainly formate, lactate, acetate, and ethanol. Ethanol production during fermentation has not been described for any *Pseudovibrio* strain yet. Additionally, pyruvate, propionate, and succinate have been formed, but to a lesser extent (Figure S 2.2 A). Production of trace amounts of fumarate was detected, but could not be quantified. As expected, we found the complete set of genes essential for denitrification, including a membrane-bound (*nar*) and a periplasmic nitrate reductase (*nap*). In agreement, we observed a complete denitrification to N₂ in laboratory experiments with strain FO-BEG1 (**Error! Reference source not found.** and Figure S 2.2 C). For the type strain *P. denitrificans*, simultaneous denitrification and fermentation was described by Shieh *et al.* (2004) and could be confirmed in our experiments for strain FO-BEG1 with acetate, formate, lactate, and ethanol as the main fermentation products (Figure S 2.2 B). No genes for assimilatory nitrate reduction could be identified in the genome. A set of *sox* genes suggests that both bacteria can use reduced inorganic sulfur compounds as a source of energy to complement heterotrophy. We could show experimentally that the addition of thiosulfate to the medium enhances the aerobic growth of the *Pseudovibrio* sp. FO-BEG1 culture and sulfate is produced over the incubation period (Figure S 2.1 B and C). No tetrathionate could be measured as an intermediate (results not shown). Therefore, we propose that thiosulfate is oxidized completely to sulfate without any intermediates, as it is known for the typical Sox pathway in α -*Proteobacteria* (for review, see Ghosh and Dam, 2009). We identified genes encoding a small (*cutS*), medium (*cutM*) and large (*cutL*) subunit of the aerobic form II carbon monoxide dehydrogenase (CODH) with the accessory gene *coxG* in the operon (see

DDBJ/EMBL/GenBank accession numbers CP003147 and CP003148), indicating the capability of CO oxidation. However, uptake of CO could not be demonstrated under tested conditions (results not shown). Interestingly, our results confirm the hypothesis from a recent publication testing CO oxidation in bacteria containing type II CODH genes (Cunliffe, 2011), in which none of the isolates containing only the type II variant was capable of CO oxidation. Only bacteria containing the form I CODH have been shown to effectively oxidize CO, thereby questioning whether form II CODH is involved in the process of carbon monoxide oxidation, or if it has another primary function not known until now, as suggested by King and Weber (2007).

In both *Pseudovibrio* strains, we found genes for phosphonate import and degradation, which allows the bacteria to cleave the relatively stable C-P bonds of phosphonates (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000). Thereby, they can metabolize a less accessible phosphorous pool in times of phosphate limitation. We could demonstrate growth of *Pseudovibrio* sp. FO-BEG1 with phosphonoacetate as the only source of phosphorous (Figure S 2.3 A). Additionally, we could show adaptation of *Pseudovibrio* strain FO-BEG1 to oligotrophic conditions by culturing it with as little as 15 $\mu\text{mol C L}^{-1}$ (0.18 mg C L^{-1}) dissolved organic carbon in the medium (Chapter 4), which shows that *Pseudovibrio* FO-BEG1 is capable of growth under extreme nutrient depletion. The high metabolic variety of *Pseudovibrio* sp. FO-BEG1 and JE062 is also reflected in the analysis of encoded primary transporters. In the genome of strain FO-BEG1 we could identify 31 tripartite ATP-independent periplasmic (TRAP) type transporters (see DDBJ/EMBL/GenBank accession numbers CP003147 and CP003148) that are required for import of dicarboxylic acids like malate, succinate and fumarate, one of the highest numbers of TRAP type transporters reported in a genome of a marine prokaryote so far (Wagner-Döbler *et al.*, 2010). In strain JE062 we identified 27 TRAP transporters. Citric acid cycle intermediates seem therefore to be an important source of carbon and energy for the investigated *Pseudovibrio* strains. In addition, we reconstructed over 80 ATP-binding cassette (ABC) transporter systems with predicted substrate specificity from the genomic data of the strain FO-BEG1, including the plasmid, and over 70 ABC transporter systems for JE062 (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000). Sugars, oligopeptides and amino acids are the main substrates that are imported via the ABC systems. A large number of transporters for oligopeptides and amino acids in combination with over 85 genes encoding peptidases and proteases (over 75 genes in strain JE062, see

DDBJ/EMBL/GenBank accession number ABXL00000000) could help *Pseudovibrio* to hydrolyze complex particulate nutrients into oligopeptides and amino acids, which could serve as nutrition for both, the prokaryote and the host, as has been suggested by Siegl *et al.* (2011). Also iron seems to be an important trace element, for which we identified eight transporters including three siderophores and three transporters for hemin (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000).

Table S.2. Identified ATP-binding cassette (ABC) and tripartite ATP-independent periplasmic (TRAP) transporters in the genomes of both *Pseudovibrio* strains and their putative functions.

Transporter type and proposed function	Transporters identified	
	FO-BEG1	JE062
TRAP transporter for dicarboxylates	31	27
ABC Transporter for:		
Sugars	22	19
Oligopeptides	15	13
Amino acids	12	10
Putrescine/spermidine	5	5
Glycerol 3-phosphate	4	4
Glycine betaine/L-proline	3	3
Glycine betaine/carnitine/ choline	1	0
Taurine	1	1
Thiamine	1	1
Urea	1	1
Nopaline	1	1
Hemin	3	3
Enterobactin	1	1
Ferrichrome	1	1
Anguibactin	1	1
Iron	2	2
Manganese/zinc	2	2
Molybdenum	1	1
Cobalt	1	1
Sulfonate	2	2
Phosphate	1	1
Phosphonate	1	1

Vitamin synthesis

Growth of pro- and eukaryotes highly depends on their requirements for cofactors that the organism can or cannot synthesize on its own. Vitamins are important for many different enzymatic processes and the synthesis of some vitamins is mainly accomplished by bacteria, making the prokaryotes a necessary part of the eukaryotic diet or an important partner in symbiotic relationships. The genomes of *Pseudovibrio* sp. FO-BEG1 as well as JE062 contain genes encoding key enzymes of the biosynthesis pathways of biotin (H), thiamin (B₁), pyridoxin (B₆), cobalamin (B₁₂), riboflavin (B₂), folic acid (B₉) and lipoic acid (see

DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000). Independence of an external vitamin supply was confirmed during aerobic growth in the defined CM medium without the addition of any vitamins, which implies *de novo* synthesis of all required growth factors by strain FO-BEG1 under tested conditions (Figure S 2.3 B). *Pseudovibrio* spp. would therefore be beneficial companions for other prokaryotes or marine invertebrates, since the dependency on an external supply of those vitamins would be relieved.

Bioactive compounds

Symbiotic relationships between bacteria and marine invertebrates, especially sponges, are of special interest, because bacteria associated with sponges often produce novel bioactive compounds (Piel *et al.*, 2004; Taylor *et al.*, 2007; Fisch *et al.*, 2009). In the chromosome of *Pseudovibrio* FO-BEG1 we identified a genomic island of more than 50 kb containing among others a gene cluster of 20 genes predicted to be involved in secondary metabolite production (see DDBJ/EMBL/GenBank accession number CP003147). The cluster exhibits high sequence similarity to an architecturally almost identical hybrid nonribosomal peptide synthetase-polyketide synthase (NRPS-PKS) system previously reported from many pathogenic and commensal *Escherichia coli* strains (Figure S.2) (Nougayrède *et al.*, 2006). The *E. coli* metabolite, termed colibactin, remains structurally uncharacterized. However, transposon mutagenesis of the gene cluster suggested that colibactin is a pathogenicity determinant that induces DNA double strand breaks in eukaryotic host cells, eventually resulting in cell death. The only significant difference between the gene clusters in *Pseudovibrio* FO-BEG1 and *E. coli* is an additional set of genes in the former, encoding putative transporters and the presence of a different phosphopantetheinyl transferase gene variant likely involved in generating *holo*-proteins from *apo* forms of PKSs and NRPSs (Lambalot *et al.*, 1996). In addition, two *E. coli* genes are fused in the *Pseudovibrio* cluster. Despite these differences, the architecture strongly suggests that the product of the FO-BEG1 cluster is colibactin, providing new opportunities to unveil the identity of this elusive and biomedically relevant compound. Interestingly, we find this more than 50 kb NRPS/PKS fragment only in *Pseudovibrio* sp. FO-BEG1 but not in the genome of strain JE062, with flanking regions downstream and upstream of the inserted fragment highly conserved in synteny in strain JE062 (data not shown), indicating that it has been acquired via horizontal gene transfer. Additionally, the plasmid of strain FO-BEG1 contained an ORF encoding a type III PKS of a size of 7.4 kb, which could also be detected in strain JE062 (see DDBJ/EMBL/GenBank accession number CP003148).




Figure S.2. Nonribosomal peptide synthetase-polyketide synthase (NRPS-PKS) system in *Pseudovibrio* sp. FO-BEG1 and *Escherichia coli* strain IHE3034. White arrows represent the genes present in *Enterobacteriaceae* and strain FO-BEG1; black arrows represent the open reading frames (ORFs) present only in either *Enterobacteriaceae* or FO-BEG1 but presumably involved in the production of colibactin; the gray arrow shows a gene presumably not involved in the synthesis of colibactin. The symbol at ORF PSE_3331 represents a gene fusion of *E. coli* genes *clbG* and *clbH* in FO-BEG1; the symbol at PSE_3324-3321 represents gene insertion or deletion in strain FO-BEG1 or *E. coli* IHE3034, respectively.

DNA exchange and horizontal gene transfer

The genomes of both *Pseudovibrio* strains show a high metabolic variety. It is reasonable to assume that various genes were acquired via horizontal gene transfer from other microorganisms as is indicated e. g. by the presence of a 50 kb large NRPS-PKS island that can be found only in *Pseudovibrio* sp. FO-BEG1 but not in strain JE062, although both genomes are in general highly similar. In the genome of strain FO-BEG1 we identified a set of genes coding for a complete gene transfer agent (GTA) (in strain JE062 several genes were missing, see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000), a unit best described as a virus. It harbors small parts of the host DNA and capable of injecting it into appropriate cells, without having negative effects on the host cell (for reviews see Lang and Beatty, 2001; Lang and Beatty, 2007). By this process, *Pseudovibrio* could have taken up and dispersed DNA carried in virus-like particles, thereby gathering genes and establishing a diverse physiology for a symbiotic and a free-living lifestyle. Additionally, we found 14 integrase and 21 transposase elements in the genome of *Pseudovibrio* sp. FO-BEG1 (see DDBJ/EMBL/GenBank accession numbers CP003147 and CP003148), 9 of which are located adjacent to the hybrid NRPS-PKS gene cluster, which verifies acquisition of this genomic island via horizontal gene transfer.

Quorum sensing

We could identify 15 genes in strain FO-BEG1 and 14 in strain JE062 containing the LuxR domain, which represents the transcriptional regulator of the acetylated homoserine lactone (AHL) type, allowing the bacterium to detect AHL quorum sensing molecules and to initiate a response (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and

ABXL0000000). Intriguingly, we could not find any *luxI* genes, which code for AHL quorum sensing molecules. This observation leads us to the hypothesis that both *Pseudovibrio* strains do not communicate via AHL within their own species, but seem to use the LuxR as receptors to react to quorum sensing molecules produced by other species and initiate a respective answer. Such a scenario has been described before by Case *et al.* (2008) and was called ‘eavesdropping’. The response reaction could include the production of bioactive compounds to repel competing prokaryotes or to protect the host from pathogens or parasites. Alternatively, such LuxR-family ‘solos’ could participate in interkingdom signaling, as suggested by Subramoni and Venturi (2009), thereby facilitating prokaryote-host interactions of *Pseudovibrio* strains with marine invertebrates.

Growth with *Beggiatoa* sp. 35Flor

Pseudovibrio sp. FO-BEG1 is the single accompanying organism of the *Beggiatoa* strain 35Flor, which is growing in a chemolithoautotrophic sulfide-oxygen-gradient medium (Brock and Schulz-Vogt, 2011; Chapters 2 and 3). All attempts to grow *Beggiatoa* without *Pseudovibrio* failed and so far we could not identify the factors required by the *Beggiatoa* strain for autonomous growth. It is known, however, that *Beggiatoa* spp. do not possess catalases (Larkin and Strohl, 1983) and therefore are susceptible to reactive oxygen molecules originating from respiration. Addition of catalase to the medium is known to increase the viability of *Beggiatoa* sp. (Burton and Morita, 1964). We hence hypothesize that *Beggiatoa* sp. 35Flor depends on the radical protection system exhibited by *Pseudovibrio* sp. FO-BEG1 including genes coding for over 20 superoxide dismutases, catalases and peroxidases (see DDBJ/EMBL/GenBank accession numbers CP003147 and CP003148). The role of heterotrophic bacteria as scavenger of reactive oxygen species has also been described by Morris *et al.* (2008), which could establish robust growth of cyanobacteria after addition of ‘helper’ heterotrophs.

Secretion Systems

In the genomes of FO-BEG1 and JE062 we could identify two loci that encode type VI secretion systems (T6SS) as well as one type III secretion system (T3SS) including effector molecules, which indicates the capability of specific interactions with eukaryotes and the possibility of influencing their cell machinery.

The T6SS has been described as a major secretion system in the context of pathogenicity as a virulence factor in moribific bacteria (Mougous *et al.*, 2006; Pukatzki *et al.*, 2006) and a core of 13 highly conserved and essential subunits has been identified for this secretion system

(Boyer *et al.*, 2009). In both genomes of the *Pseudovibrio* strains, we found two gene clusters consisting of 12 (cluster I) and 20 (cluster II) genes that encode T6SSs. Cluster II contains the complete set of core subunits and therefore we assume that cluster II could, if expressed, produce a complete and functional type VI secretion system. In cluster I, two core genes are missing in the operon, *hcpI* and *vgrG*, which are main components of the injection apparatus with possible effector functions (Pukatzki *et al.*, 2009; Bönemann *et al.*, 2010). However, homologues of *hcpI* and *vgrG* could be identified in additional copies at other locations in the genomes of FO-BEG1 and JE062 (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000), which is a phenomenon regularly found in genomes containing T6SS (Pukatzki *et al.*, 2009). The possible role for type VI secretion systems in bacteria has not been completely elucidated so far, but several functions have been attributed to it already. Mainly, T6SS is described as a virulence factor of pathogenic bacteria delivering effector proteins into host cells (Filloux *et al.*, 2008). However, further studies reveal the involvement of T6SS in biofilm formation (Aschtgen *et al.*, 2008), quorum sensing (Weber *et al.*, 2009), interbacterial interactions (Hood *et al.*, 2010) and antipathogenesis (Chow and Mazmanian, 2010; Jani and Cotter, 2010). In conclusion, it can be assumed that the T6SS of both strains are functional since the genomes contain the main structural components of the type VI secretion system.

In addition to the T6SS, we identified a type III secretion system in the genomes of both *Pseudovibrio* strains, which is located in a genomic region encompassing around 35 ORFs with various highly conserved proteins known from T3S systems (Cornelis and Van Gijsegem, 2000) (Figure S.3 and see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000). Besides the secretion apparatus we also identified genes encoding homologues of three types of effector molecules in the genome of strain FO-BEG1 and two effector molecule types in strain JE062. Those effectors might be directly involved in the establishment of symbiosis between *Pseudovibrio* and its host. YpkA, IpgD (found in both genomes) and YopJ (only in strain FO-BEG1) are effector molecules that affect the cytoskeleton or the innate immune response of the host, respectively. YpkA is a serine/threonine kinase, which has negative effects on cytoskeletal dynamics due to its interaction with actin, thereby contributing to the resistance to phagocytosis (Cornelis, 2002). YpkA is present in three copies in both genomes. In Porifera, specialized amoeboid cells, the archaeocytes, resemble macrophages and eliminate non-self material via phagocytosis (Müller and Müller, 2003). *Pseudovibrio*, expressing and secreting the YpkA effector, could interfere

with this process, preventing archaeocytes from digesting *Pseudovibrio* cells. A similar effect could be induced by a homologue of IpgD found in both genomes, a virulence factor that is responsible for morphological changes of a host cell by increasing membrane detachment from the cytoskeleton (Niebuhr *et al.*, 2000; Niebuhr *et al.*, 2002).




Figure S.3. Operon coding for type III secretion system (T3SS) subunits and effector proteins. White arrows show annotated homologues of T3SS subunits including the gene name within the arrows; black arrows represent annotated effector homologues; dark gray arrows show annotated genes encoding proteins presumably not involved in T3SS; light gray arrows show hypothetical proteins with unknown function. The locus is indicated above and below some genes for orientation purposes.

In FO-BEG1 we additionally identified a homologue of the YopJ effector exhibiting a serine/threonine acetyltransferase function. By acetylation of serine and threonine residues of mitogen-activated protein (MAP) kinases it prevents phosphorylation of those molecules and therefore inhibits the innate immune response of the organism (Mukherjee *et al.*, 2006). Intriguingly, it has been shown that sponges possess a very efficient innate immune response system, using MAP kinases as the essential component of its response to bacterial endotoxin lipopolysaccharide (LPS) (Böhm *et al.*, 2001; Müller and Müller, 2003). This indicates that homologues of the acetyltransferase YopJ effector in *Pseudovibrio* could prevent phosphorylation of MAP kinases via acetylation, thereby playing a role in the inactivation of the immune answer of the host organism, allowing *Pseudovibrio* to avoid phagocytosis, as described by Bartsev *et al.* (2004) for a *Rhizobium* strain, and to remain in the host for establishment of a symbiosis. This hypothesis is further supported by the fact that a homologue of YopJ (NopJ) was shown to be an effector in symbiotic rhizobia (Deakin and Broughton, 2009) and Lackner *et al.* (2011) demonstrated that T3SS is involved in maintenance of a symbiosis between bacteria and fungi by enhancement of intracellular survival of the prokaryote within the host.

Adhesion

In both genomes we found homologues of genes coding for proteins responsible for adhesion to surfaces or other cells. These proteins, belonging to the group of amyloids, are extracellular

proteinaceous components and are known in Enterobacteriaceae as curli fibers. They are involved in adhesion to surfaces, cell aggregation, biofilm formation and mediate cell-cell adhesion and invasion of host cells (Barnhart and Chapman, 2006). The production of curli fibers in enteric bacteria is dependent on at least six proteins encoded by the operons *csgAB* and *csgDEFG* (*agf* in *Salmonella*) (Hammar *et al.*, 1995), the latter of which is required for assembly, stability and secretion of the amyloids (Hammar *et al.*, 1995). *csgAB* encodes the structural subunits of the curli fibers, both genes containing characteristic repeat motifs (Hammar *et al.*, 1996). A gene cluster in the genome of *Pseudovibrio* sp. FO-BEG1 resembles the curli formation operon in enteric bacteria (Figure S.4).




Figure S.4. Comparison of genes encoding amyloids in Enterobacteriaceae and the operon in *Pseudovibrio* sp. FO-BEG1. White arrows represent homologues of genes in enteric bacteria; gray arrows show genes present in Enterobacteriaceae only; black arrows show genes containing curli repeats, typical motifs of the amyloid structural subunits. The number within the black arrows shows the amount of curli repeats in the according gene.

Homologues of *csgF* and *csgG*, required for stabilization and secretion of the amyloids are found in direct proximity to three genes containing curlin associated repeats as typical structural components of the curli fibers. We hypothesize that the identified operon might code for amyloid structures comparable to curli fibers due to the existence of characteristic curlin repeat motifs and genes involved in the assembly and secretion of such structures, therefore allowing *Pseudovibrio* to attach to other cells or form biofilms or aggregates. Additionally, we identified 35 genes in strain FO-BEG1 and 37 in JE062 containing domains mediating prokaryote-eukaryote interactions, supporting the proposed role of *Pseudovibrio* as a symbiont with possibilities to attach and interact with the host organism (see DDBJ/EMBL/GenBank accession numbers CP003147, CP003148 and ABXL00000000).

Conclusions

In this study, we analyzed highly similar genomes of two *Pseudovibrio* strains that originate from the coast of Florida, the *Pseudovibrio* sp. FO-BEG1 sampled from a coral and maintained over 10 years in co-culture with *Beggiatoa* sp. and *Pseudovibrio* sp. JE062

sampled from a sponge in the same region (Enticknap *et al.*, 2006). The physiology of both strains is extremely versatile and the metabolic traits found in the genome could be partially verified in experiments with strain FO-BEG1. Here, we describe for the first time a *Pseudovibrio* strain that uses aromatic compounds as a carbon and electron source, oxidizes thiosulfate under aerobic conditions and uses phosphonates as a phosphorous source. Notably, strain FO-BEG1 grows under extreme nutrient limitation, which emphasizes its adaptation to life in the open ocean. The metabolic variety is confirmed by the numerous transporter systems that are encoded in the genome. Compared with other marine bacteria, like the prominent *Roseobacter* clade, which is known to be ubiquitous, multitudinous and physiologically versatile (Newton *et al.*, 2010), *Pseudovibrio* seems to be capable of a similarly generalistic life style, exploiting quite a number of sources for energy sources, nutrients and trace elements.

Aside from metabolic variety, the genomic data of both strains also confirm close associations with marine invertebrates and indicate several potential mechanisms for establishing and maintaining a symbiosis. The most striking discovery is the presence of effector homologues secreted by type III secretion systems, which could affect sponges by interacting with their immune response system (YopJ) or the cytoskeleton (YpkA, IpgD) and thereby have a drastic impact on the cell machinery of the host. Another fascinating discovery is the presence of the hybrid NRPS-PKS system in strain FO-BEG1, which has so far only been described for members of the Enterobacteriaceae family (Putze *et al.*, 2009), producing the bioactive compound colibactin with yet unknown *in-vivo* functions, but arresting eukaryotic cells in the G₂ phase, eventually leading to cell death (Nougayrède *et al.*, 2006). The presence of a gene cluster coding for a cytopathic compound in strain FO-BEG1 emphasizes the impact that *Pseudovibrio* cells might have on marine invertebrates. Intriguingly, strain FO-BEG1 seems to be a required partner in the *Beggiatoa* co-culture, indicating its important symbiotic role not only for marine invertebrates but also for prokaryotes. It is possible that *Pseudovibrio* has positive effects for certain bacteria under *in-vivo* conditions, e.g. by supplying vitamins or detoxifying metabolic intermediates or radical oxygen species.




Figure S.5. Schematic overview of the possible life styles and the physiologic capabilities derived from genetic information of both *Pseudovibrio* genomes. On the left hand side, physiologic abilities are depicted that could be used in free-living, oxic and anoxic conditions. On the right hand side, the attached or associated life style is illustrated. The host organism for the associated life style can be represented by a sponge, coral or tunicate. Biofilm formation, aggregation and attachment to host cells could be performed via e. g. amyloid-like structures. The proposed secretion systems could be involved in prokaryote-eukaryote interactions, influencing the cell machinery of the host. Additionally, *Pseudovibrio* could supply the host with cofactors like vitamins or synthesize secondary metabolites as a defense mechanism against other prokaryotes or the host.

The frequent identification and isolation of *Pseudovibrio* strains in many studies over the last years implies an important but rather unexplored role for this genus in marine habitats. According to the genomic and physiological data on *Pseudovibrio* spp., we propose a free-living and attached or associated life style model for this genus (Figure S.5). As a denitrifying heterotroph, *Pseudovibrio* has an obvious influence on the carbon and nitrogen cycles. Its ecological impact can now be extended to the sulfur and phosphorus cycles due to its ability to metabolize thiosulfate and phosphonates. Additionally, we hypothesize that, due to the predictions based on the genomic data, similar to *E. coli* in humans, *Pseudovibrio* is a commensalistic or even beneficial symbiont of marine invertebrates with a potential to become pathogenic.

Acknowledgments

We thank A. Meyerdierks, M. Mußmann, and R. Amann for comments on the manuscript. For technical support, we thank M. Meyer, S. Menger, C. Probian, and R. Appel. We also thank A. Kamyshny for tetrathionate measurements and K. Imhoff for sulfate determination. This study was funded by the European Research Council and the Max Planck Society.

References of appendix

- Aeckersberg F, Bak F, Widdel F. (1991). Anaerobic oxidation of saturated hydrocarbons to CO₂ by a new type of sulfate-reducing bacterium. *Arch Microbiol* **156**: 5–14.
- Agogué H, Casamayor EO, Bourrain M, Obernosterer I, Joux F, Herndl GJ *et al.* (2005). A survey on bacteria inhabiting the sea surface microlayer of coastal ecosystems. *FEMS Microbiol Ecol* **54**: 269–280.
- Aschtgen MS, Bernard CS, De Bentzmann S, Lloubès R, Cascales E. (2008). SciN is an outer membrane lipoprotein required for type VI secretion in enteroaggregative *Escherichia coli*. *J Bacteriol* **190**: 7523–7531.
- Barnhart MM, Chapman MR. (2006). Curli biogenesis and function. *Annu Rev Microbiol* **60**: 131–147.
- Bartsev AV, Deakin WJ, Boukli NM, McAlvin CB, Stacey G, Malnoe P *et al.* (2004). NopL, an effector protein of *Rhizobium* sp. NGR234, thwarts activation of plant defense reactions. *Plant Physiology* **134**: 871–879.
- Bateman A, Coin L, Durbin R, Finn RD, Hollich V, Griffiths-Jones S *et al.* (2004). The Pfam protein families database. *Nucleic Acids Res* **32**: D138–D141.
- Bendtsen JD, Nielsen H, von Heijne G, Brunak S. (2004). Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol* **340**: 783–795.
- Böhm M, Hentschel U, Friedrich AB, Fieseler L, Steffen R, Gamulin V *et al.* (2001). Molecular response of the sponge *Suberites domuncula* to bacterial infection. *Mar Biol* **139**: 1037–1045.
- Bönemann G, Pietrosiuk A, Mogk A. (2010). Tubules and donuts: a type VI secretion story. *Mol Microbiol* **76**: 815–821.
- Boyer F, Fichant G, Berthod J, Vandenbrouck Y, Attree I. (2009). Dissecting the bacterial type VI secretion system by a genome wide in silico analysis: what can be learned from available microbial genomic resources? *BMC Genomics* **10**: 104.
- Brock J, Schulz-Vogt HN. (2011). Sulfide induces phosphate release from polyphosphate in cultures of a marine *Beggiatoa* strain. *ISME J* **5**: 497–506.
- Brysch K, Schneider C, Fuchs G, Widdel F. (1987). Lithoautotrophic growth of sulfate-reducing bacteria, and description of *Desulfobacterium autotrophicum* gen. nov., sp. nov. *Arch Microbiol* **148**: 264–274.
- Burton SD, Morita RY. (1964). Effect of catalase and cultural conditions on growth of *Beggiatoa*. *J Bacteriol* **88**: 1755–1761.
- Carver T, Thomson N, Bleasby A, Berriman M, Parkhill J. (2009). DNAPlotter: circular and linear interactive genome visualization. *Bioinformatics* **25**: 119–120.
- Case RJ, Labbate M, Kjelleberg S. (2008). AHL-driven quorum-sensing circuits: their frequency and function among the Proteobacteria. *ISME J* **2**: 345–349.

- Chow J, Mazmanian SK. (2010). A pathobiont of the microbiota balances host colonization and intestinal inflammation. *Cell Host Microbe* **7**: 265–276.
- Cornelis GR. (2002). The *Yersinia* Ysc-Yop 'type III' weaponry. *Nat Rev Mol Cell Bio* **3**: 742–752.
- Cornelis GR, Van Gijsegem F. (2000). Assembly and function of type III secretory systems. *Annu Rev Microbiol* **54**: 735–774.
- Cunliffe M. (2011). Correlating carbon monoxide oxidation with *cox* genes in the abundant Marine Roseobacter Clade. *ISME J* **5**: 685–691.
- Deakin WJ, Broughton WJ. (2009). Symbiotic use of pathogenic strategies: rhizobial protein secretion systems. *Nat Rev Microbiol* **7**: 312–320.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. (2007). Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* **23**: 673–679.
- Enticknap JJ, Kelly M, Peraud O, Hill RT. (2006). Characterization of a culturable alphaproteobacterial symbiont common to many marine sponges and evidence for vertical transmission via sponge larvae. *Appl Environ Microbiol* **72**: 3724–3732.
- Filloux A, Hachani A, Bleves S. (2008). The bacterial type VI secretion machine: yet another player for protein transport across membranes. *Microbiol-SGM* **154**: 1570–1583.
- Fisch KM, Gurgui C, Heycke N, van der Sar SA, Anderson SA, Webb VL *et al.* (2009). Polyketide assembly lines of uncultivated sponge symbionts from structure-based gene targeting. *Nat Chem Biol* **5**: 494–501.
- Fukunaga Y, Kurahashi M, Tanaka K, Yanagi K, Yokota A, Harayama S. (2006). *Pseudovibrio ascidiaceicola* sp. nov., isolated from ascidians (sea squirts). *Int J Syst Evol Microbiol* **56**: 343–347.
- Fürch T, Preusse M, Tomasch J, Zech H, Wagner-Döbler I, Rabus R *et al.* (2009). Metabolic fluxes in the central carbon metabolism of *Dinoroseobacter shibae* and *Phaeobacter gallaeciensis*, two members of the marine *Roseobacter* clade. *BMC Microbiol* **9**: 209.
- Ghosh W, Dam B. (2009). Biochemistry and molecular biology of lithotrophic sulfur oxidation by taxonomically and ecologically diverse bacteria and archaea. *FEMS Microbiol Rev* **33**: 999–1043.
- Hammar M, Bian Z, Normark S. (1996). Nucleator-dependent intercellular assembly of adhesive curli organelles in *Escherichia coli*. *Proc Natl Acad Sci U S A* **93**: 6562–6566.
- Hammar M, Arnqvist A, Bian Z, Olsén A, Normark S. (1995). Expression of two *csg* operons is required for production of fibronectin- and congo red-binding curli polymers in *Escherichia coli* K-12. *Mol Microbiol* **18**: 661–670.
- Hentschel U, Schmid M, Wagner M, Fieseler L, Gernert C, Hacker J. (2001). Isolation and phylogenetic analysis of bacteria with antimicrobial activities from the mediterranean

- sponges *Aplysina aerophoba* and *Aplysina cavernicola*. *FEMS Microbiol Ecol* **35**: 305–312.
- Hood RD, Singh P, Hsu FS, Güvener T, Carl MA, Trinidad RRS *et al.* (2010). A type VI secretion system of *Pseudomonas aeruginosa* targets a toxin to bacteria. *Cell Host Microbe* **7**: 25–37.
- Hosoya S, Yokota A. (2007). *Pseudovibrio japonicus* sp. nov., isolated from coastal seawater in Japan. *Int J Syst Evol Microbiol* **57**: 1952–1955.
- Jani AJ, Cotter PA. (2010). Type VI secretion: not just for pathogenesis anymore. *Cell Host Microbe* **8**: 2–6.
- Kamysny A. (2009). Improved cyanolysis protocol for detection of zero-valent sulfur in natural aquatic systems. *Limnol Oceanogr-Meth* **7**: 442–448.
- Kennedy J, Baker P, Piper C, Cotter PD, Walsh M, Mooij MJ *et al.* (2009). Isolation and analysis of bacteria with antimicrobial activities from the marine sponge *Haliclona simulans* collected from Irish waters. *Mar Biotechnol* **11**: 384–396.
- King GM, Weber CF. (2007). Distribution, diversity and ecology of aerobic CO-oxidizing bacteria. *Nat Rev Microbiol* **5**: 107–118.
- Koren O, Rosenberg E. (2006). Bacteria associated with mucus and tissues of the coral *Oculina patagonica* in summer and winter. *Appl Environ Microbiol* **72**: 5254–5259.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**: 567–580.
- Lackner G, Moebius N, Hertweck C. (2011). Endofungal bacterium controls its host by an *hrp* type III secretion system. *ISME J* **5**: 252–261.
- Lafi FF, Garson MJ, Fuerst JA. (2005). Culturable bacterial symbionts isolated from two distinct sponge species (*Pseudoceratina clavata* and *Rhabdastrella globostellata*) from the Great Barrier Reef display similar phylogenetic diversity. *Microb Ecol* **50**: 213–220.
- Lagesen K, Hallin P, Rødland EA, Stærfeldt HH, Rognes T, Ussery DW. (2007). RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* **35**: 3100–3108.
- Lambalot RH, Gehring AM, Flugel RS, Zuber P, LaCelle M, Marahiel MA *et al.* (1996). A new enzyme superfamily - the phosphopantetheinyl transferases. *Chem Biol* **3**: 923–936.
- Lang AS, Beatty JT. (2001). The gene transfer agent of *Rhodobacter capsulatus* and "constitutive transduction" in prokaryotes. *Arch Microbiol* **175**: 241–249.
- Lang AS, Beatty JT. (2007). Importance of widespread gene transfer agent genes in α -*Proteobacteria*. *Trends Microbiol* **15**: 54–62.

- Larkin JM, Strohl WR. (1983). *Beggiatoa*, *Thiothrix*, and *Thioploca*. *Annu Rev Microbiol* **37**: 341–367.
- Lowe TM, Eddy SR. (1997). tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* **25**: 955–964.
- Meyer F, Goesmann A, McHardy AC, Bartels D, Bekel T, Clausen J *et al.* (2003). GenDB - an open source genome annotation system for prokaryote genomes. *Nucleic Acids Res* **31**: 2187–2195.
- Morris JJ, Kirkegaard R, Szul MJ, Johnson ZI, Zinser ER. (2008). Facilitation of robust growth of *Prochlorococcus* colonies and dilute liquid cultures by "helper" heterotrophic bacteria. *Appl Environ Microbiol* **74**: 4530–4534.
- Mougous JD, Cuff ME, Raunser S, Shen A, Zhou M, Gifford CA *et al.* (2006). A virulence locus of *Pseudomonas aeruginosa* encodes a protein secretion apparatus. *Science* **312**: 1526–1530.
- Mukherjee S, Keitany G, Li Y, Wang Y, Ball HL, Goldsmith EJ *et al.* (2006). *Yersinia* YopJ acetylates and inhibits kinase activation by blocking phosphorylation. *Science* **312**: 1211–1214.
- Mulder NJ, Apweiler R, Attwood TK, Bairoch A, Bateman A, Binns D *et al.* (2005). InterPro, progress and status in 2005. *Nucleic Acids Res* **33**: D201–D205.
- Müller WEG, Müller IM. (2003). Origin of the metazoan immune system: identification of the molecules and their functions in sponges. *Integr Comp Biol* **43**: 281–292.
- Muscholl-Silberhorn A, Thiel V, Imhoff JF. (2008). Abundance and bioactivity of cultured sponge-associated bacteria from the mediterranean sea. *Microb Ecol* **55**: 94–106.
- Newton RJ, Griffin LE, Bowles KM, Meile C, Gifford S, Givens CE *et al.* (2010). Genome characteristics of a generalist marine bacterial lineage. *ISME J* **4**: 784–798.
- Niebuhr K, Jouihri N, Allaoui A, Gounon P, Sansonetti PJ, Parsot C. (2000). IpgD, a protein secreted by the type III secretion machinery of *Shigella flexneri*, is chaperoned by IpgE and implicated in entry focus formation. *Mol Microbiol* **38**: 8–19.
- Niebuhr K, Giuriato S, Pedron T, Philpott DJ, Gaits F, Sable J *et al.* (2002). Conversion of PtdIns(4,5)P-2 into PtdIns(5)P by the *S.flexneri* effector IpgD reorganizes host cell morphology. *EMBO J* **21**: 5069–5078.
- Nougayrède JP, Homburg S, Taieb F, Boury M, Brzuszkiewicz E, Gottschalk G *et al.* (2006). *Escherichia coli* induces DNA double-strand breaks in eukaryotic cells. *Science* **313**: 848–851.
- Olson JB, Harmody DK, McCarthy PJ. (2002). α -Proteobacteria cultivated from marine sponges display branching rod morphology. *FEMS Microbiol Lett* **211**: 169–173.

- Piel J, Hui DQ, Wen GP, Butzke D, Platzer M, Fusetani N *et al.* (2004). Antitumor polyketide biosynthesis by an uncultivated bacterial symbiont of the marine sponge *Theonella swinhoei*. *Proc Natl Acad Sci U S A* **101**: 16222–16227.
- Pukatzki S, McAuley SB, Miyata ST. (2009). The type VI secretion system: translocation of effectors and effector-domains. *Curr Opin Microbiol* **12**: 11–17.
- Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC *et al.* (2006). Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc Natl Acad Sci U S A* **103**: 1528–1533.
- Putze J, Hennequin C, Nougayrède JP, Zhang WL, Homburg S, Karch H *et al.* (2009). Genetic structure and distribution of the colibactin genomic island among members of the family *Enterobacteriaceae*. *Infect Immun* **77**: 4696–4703.
- Quast C. (2006). MicHanThi - design and implementation of a system for the prediction of gene functions in genome annotation projects. Diploma thesis, Universität Bremen, Bremen.
- Richter M, Rosselló-Móra R. (2009). Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci U S A* **106**: 19126–19131.
- Richter M, Lombardot T, Kostadinov I, Kottmann R, Duhaime MB, Peplies J *et al.* (2008). JCoast - a biologist-centric software tool for data mining and comparison of prokaryotic (meta)genomes. *BMC Bioinformatics* **9**: 177.
- Riesenfeld CS, Murray AE, Baker BJ. (2008). Characterization of the microbial community and polyketide biosynthetic potential in the palmerolide-producing tunicate *Synoicum adareanum*. *J Nat Prod* **71**: 1812–1818.
- Rypien KL, Ward JR, Azam F. (2010). Antagonistic interactions among coral-associated bacteria. *Environ Microbiol* **12**: 28–39.
- Santos OCS, Pontes PVML, Santos JFM, Muricy G, Giambiagi-deMarval M, Laport MS. (2010). Isolation, characterization and phylogeny of sponge-associated bacteria with antimicrobial activities from Brazil. *Res Microbiol* **161**: 604–612.
- Sertan-de Guzman AA, Predicala RZ, Bernardo EB, Neilan BA, Elardo SP, Mangalindan GC *et al.* (2007). *Pseudovibrio denitrificans* strain Z143-1, a heptylprodigiosin-producing bacterium isolated from a Philippine tunicate. *FEMS Microbiol Lett* **277**: 188–196.
- Shieh WY, Lin YT, Jean WD. (2004). *Pseudovibrio denitrificans* gen. nov., sp. nov., a marine, facultatively anaerobic, fermentative bacterium capable of denitrification. *Int J Syst Evol Microbiol* **54**: 2307–2312.
- Siegl A, Kamke J, Hochmuth T, Piel J, Richter M, Liang CG *et al.* (2011). Single-cell genomics reveals the lifestyle of *Poribacteria*, a candidate phylum symbiotically associated with marine sponges. *ISME J* **5**: 61–70.
- Stothard P, Wishart DS. (2005). Circular genome visualization and exploration using CGView. *Bioinformatics* **21**: 537–539.

- Subramoni S, Venturi V. (2009). LuxR-family 'solos': bachelor sensors/regulators of signalling molecules. *Microbiol-SGM* **155**: 1377–1385.
- Tang KH, Feng XY, Tang YJJ, Blankenship RE. (2009). Carbohydrate metabolism and carbon fixation in *Roseobacter denitrificans* OCh114. *PLoS One* **4**: e7233.
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV *et al.* (2003). The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* **4**: 41.
- Taylor MW, Radax R, Steger D, Wagner M. (2007). Sponge-associated microorganisms: evolution, ecology, and biotechnological potential. *Microbiol Mol Biol Rev* **71**: 295–374.
- Thakur NL, Hentschel U, Krasko A, Pabel CT, Anil AC, Muller WEG. (2003). Antibacterial activity of the sponge *Suberites domuncula* and its primmorphs: potential basis for epibacterial chemical defense. *Aquat Microb Ecol* **31**: 77–83.
- Thiel V, Imhoff JF. (2003). Phylogenetic identification of bacteria with antimicrobial activities isolated from Mediterranean sponges. *Biomolecular Engineering* **20**: 421–423.
- Thoms C, Horn M, Wagner M, Hentschel U, Proksch P. (2003). Monitoring microbial diversity and natural product profiles of the sponge *Aplysina cavernicola* following transplantation. *Mar Biol* **142**: 685–692.
- Wagner-Döbler I, Ballhausen B, Berger M, Brinkhoff T, Buchholz I, Bunk B *et al.* (2010). The complete genome sequence of the algal symbiont *Dinoroseobacter shibae*: a hitchhiker's guide to life in the sea. *ISME J* **4**: 61–77.
- Weber B, Hasic M, Chen C, Wai SN, Milton DL. (2009). Type VI secretion modulates quorum sensing and stress response in *Vibrio anguillarum*. *Environ Microbiol* **11**: 3018–3028.
- Webster NS, Hill RT. (2001). The culturable microbial community of the Great Barrier Reef sponge *Rhopaloeides odorabile* is dominated by an α -proteobacterium. *Mar Biol* **138**: 843–851.
- Widdel F, Pfennig N. (1984). Dissimilatory sulfate- or sulfur-reducing bacteria. In: Krieg NR, Holt JG (eds). *Bergey's manual of systematic bacteriology*. Williams & Wilkins: Baltimore. pp 663–379.
- Widdel F, Bak F. (1992). Gram-negative mesophilic sulfate-reducing bacteria. In: Balows A, Trüper HG, Dworkin M, Harder W, Schleifer KH (eds). *The Prokaryotes*, 2nd edn. Springer: New York. pp 3352–3378.
- Williams TJ, Ertan H, Ting L, Cavicchioli R. (2009). Carbon and nitrogen substrate utilization in the marine bacterium *Sphingopyxis alaskensis* strain RB2256. *ISME J* **3**: 1036–1052.
- Zedelius J, Rabus R, Grundmann O, Werner I, Brodkorb D, Schreiber F *et al.* (2011). Alkane degradation under anoxic conditions by a nitrate-reducing bacterium with possible

involvement of the electron acceptor in substrate activation. *Environ Microbiol Rep* **3**: 125–135.

Supplementary figures of the appendix




Figure S 2.1. (A) Growth of *Pseudovibrio* strain FO-BEG1 with aromatic compounds as the sole carbon and energy source. (B) Growth and SO_4^{2-} evolution by *Pseudovibrio* sp. FO-BEG1 under oxic conditions with the addition of $10 \text{ mmol l}^{-1} \text{ Na}_2\text{S}_2\text{O}_3$ to the medium. Closed circles and crosses represent the SO_4^{2-} evolution and optical density in inoculated samples. Open circles and open crosses represent SO_4^{2-} evolution and optical density, respectively, in an uninoculated control to exclude chemical oxidation of $\text{Na}_2\text{S}_2\text{O}_3$. Initially, the medium did not contain any SO_4^{2-} in order to decrease the SO_4^{2-} background during measurements. (C) Growth of *Pseudovibrio* sp. FO-BEG1 with and without the addition of $10 \text{ mmol l}^{-1} \text{ Na}_2\text{S}_2\text{O}_3$ to the medium. The medium for this experiment contains $11.5 \text{ mmol l}^{-1} \text{ K}_2\text{SO}_4$ to ensure that the culture without $\text{Na}_2\text{S}_2\text{O}_3$ contains a sulfur source for growth. Error bars represent the standard deviation in biological triplicates.




Figure S 2.2. (A) Glucose consumption, growth and production of fermentation products by *Pseudovibrio* sp. FO-BEG1 grown under anoxic conditions without NO_3^- . (B) Glucose consumption, growth and production of fermentation products during simultaneous denitrification and fermentation by *Pseudovibrio* sp. FO-BEG1 grown under anoxic conditions with addition of $10 \text{ mmol l}^{-1} \text{NO}_3^-$. (C) Consumption of NO_3^- , evolution of gaseous nitrogen and growth during simultaneous denitrification and fermentation of *Pseudovibrio* sp. FO-BEG1 with $10 \text{ mmol l}^{-1} \text{NO}_3^-$. Error bars represent the standard deviation in biological triplicates.




Figure S 2.3. (A) Growth of *Pseudovibrio* sp. FO-BEG1 with 1 mmol L⁻¹ phosphonoacetate as the only phosphorus source and without the addition of any phosphorus to the medium. (B) Growth of *Pseudovibrio* sp. FO-BEG1 without the addition of vitamins to the medium. Error bars represent the standard deviation in biological triplicates.

Acknowledgements

I would like to express my deepest thanks to my “Doktormutter” Heide Schulz-Vogt. You have shown me the beauty of the large sulfide-oxidizers and you have always encouraged me to follow my ideas and to go into new directions. Thank you for providing me such an interesting topic and for all your support!

I would also like to thank Prof. Dr. Ulrich Fischer for writing the second review. Furthermore, I am thankful for the helpful discussion and remarks you made during my thesis committee meeting and all further discussions.

Thank you Thorsten Dittmar for being a member of my thesis and defense committee and also for all your help during the last years! I am very happy that I had the possibility to work with and learn from you and I am looking forward to perform the next experiments.

Moreover, I thank Rudi Amann for joining my thesis and defense committee and for all your helpful remarks during our discussions.

Anne-Christin Kreutzmann and Sascha Artelt, thank you for being members of my defense committee.

Dear Ecophysios (Jörg, Verena, Sandra, Stefano, Vladi, Martina, Susanne, Anne-Christin, Tine, Daniela, Sascha, Arthur), thank you all so much for the last years, I enjoyed working with you all! Especially I would like to thank Anne-Christin Kreutzmann, I hope we will write many more manuscripts together in the future! Martina Meyer and Susanne Menger, you are the best technicians I can imagine!

Michael Seidel, thank you for measuring so many samples with me and for cleaning up the lab without me late at night, when I already had to catch the last train back to Bremen. I also thank you for picking up the “Sorgentelefon” when I analyzed the FT-ICR-MS data! Furthermore, I would like to thank the whole Marine Geochemistry Group in Oldenburg for helping me (especially Matze, Katrin and Ina). I always enjoyed the time in Oldenburg!

Thanks a lot to Meinhard Simon and Birgit Kürzel for measuring the amino acid samples.

Acknowledgements

Thank you, Lubos Polerecky, for analysing all the data with me and for nice colourful figures.

Martin Beutler, thank you for taking many, many pictures with me at the CLSM.

Thanks to Michael Richter for his very spontaneous help with KEGG and his help during genome annotation.

Gaute Lavik and Thomas Max are thanked for their help during isotope-labelling experiments.

I thank Johannes Zedelius and Frank Schreiber for their help when I had “NO” questions.

I thank Bernd Stickford for his help during literature search. He was able to find also very, very old articles for me.

Many thanks to the whole Microbiology department at the MPI Bremen for the nice working atmosphere and all your help, representative I would like to mention Jens Harder and Prof. Widdel.

My dearest officemates: Jana, Julia and Verena. You were always there – celebrating when things worked out, discussing when something unexpected happened and encouraging me when experiments failed. Thank you so much!

Ines and Anna - lunch with you was always awesome ☺

I always enjoyed the “Mädelsabend” with my MPI girls (Anna, Ines, Frauke, Kirsten, Sandra, Ulli, Verena, Julia, Anne-Christin and Jana). I’ve never been alone with you at my side going through the ups and downs of the PhD life, thank you!

Meinen Freunden (Julia, Karo und ALLEN anderen) und meiner Familie (Andy, Mama, Papa, Lina, Bernd, Oma, Opa, Frank, Katha + XX und der GANZEN Sippe) möchte ich danken, dass sie mir immer zur Seite gestanden, mich unterstützt und wenn nötig auf andere Gedanken gebracht haben!

Andy – du bist der beste Mann den es gibt und dafür bin ich dir von ganzem Herzen dankbar!!!

Name: Anne Schwedt
Anschrift: Am Deich 75, 28199 Bremen

Bremen, 23.09.2011

ERKLÄRUNG

Hiermit erkläre ich, dass ich die Arbeit mit dem Titel:

„Physiology of a marine *Beggiatoa* strain and the accompanying organism *Pseudovibrio* sp. – a facultatively oligotrophic bacterium“

selbstständig verfasst und geschrieben habe und außer den angegebenen Quellen keine weiteren Hilfsmittel verwendet habe.

Ebenfalls erkläre ich hiermit eidesstattlich, dass es sich bei den von mir abgegebenen Arbeiten um 3 identische Exemplare handelt.

(Unterschrift)