





## REVIEW ARTICLE

# Current technical approaches to brain energy metabolism

L. Felipe Barros<sup>1</sup>  | Juan P. Bolaños<sup>2</sup> | Gilles Bonvento<sup>3</sup> |  
 Anne-Karine Bouzier-Sore<sup>4</sup> | Angus Brown<sup>5</sup> | Johannes Hirrlinger<sup>6,7</sup>  |  
 Sergey Kasparov<sup>8,9</sup>  | Frank Kirchhoff<sup>10</sup> | Anne N. Murphy<sup>11</sup> | Luc Pellerin<sup>12</sup> |  
 Michael B. Robinson<sup>13</sup>  | Bruno Weber<sup>14,15</sup>

<sup>1</sup>Centro de Estudios Científicos (CECs), Valdivia, 5110466, Chile

<sup>2</sup>Instituto de Biología Funcional y Genómica-CSIC, Universidad de Salamanca, CIBERFES, Salamanca 37007, Spain

<sup>3</sup>Commissariat à l'Energie Atomique et aux Energies Alternatives (CEA), Département de la Recherche Fondamentale (DRF), Institut de Biologie François Jacob, Molecular Imaging Research Center (MIRcen), CNRS UMR 9199, Université Paris-Sud, Université Paris-Saclay, Fontenay-aux-Roses, France

<sup>4</sup>Centre de Résonance Magnétique des Systèmes Biologiques UMR 5536, CNRS-Université Bordeaux 146 rue Léo-Saignat, Bordeaux, France

<sup>5</sup>School of Life Sciences, University of Nottingham, Nottingham, United Kingdom

<sup>6</sup>Carl Ludwig Institute of Physiology, University of Leipzig, Liebigstr. 27, D-04103 Leipzig, Germany

<sup>7</sup>Department of Neurogenetics, Max-Planck-Institute for Experimental Medicine, Hermann-Rein-Str. 3, Göttingen D-37075, Germany

<sup>8</sup>School of Physiology, Pharmacology and Neuroscience, University of Bristol, University Walk, BS8 1TD, United Kingdom

<sup>9</sup>Baltic Federal University, Kalinigrad, Russian Federation

<sup>10</sup>Molecular Physiology, Center for Integrative Physiology and Molecular Medicine, University of Saarland, Building 48, Homburg 66421, Germany

<sup>11</sup>Department of Pharmacology, University of California, San Diego, La Jolla, CA 92093

<sup>12</sup>Département de Physiologie, 7 rue du Bugnon, Lausanne CH1005, Switzerland

<sup>13</sup>Department of Pediatrics, and Department of Systems Pharmacology and Translational Therapeutics, Children's Hospital of Philadelphia Research Institute, University of Pennsylvania, Philadelphia, PA 19104, USA

<sup>14</sup>Institute of Pharmacology and Toxicology, University of Zurich, Zurich, Switzerland

<sup>15</sup>Neuroscience Center Zurich, Zurich, Switzerland

## Correspondence

L. Felipe Barros, Centro de Estudios Científicos (CECs), 5110466 Valdivia, Chile.

Email: fbarros@cecs.cl

## Funding information

Fondecyt, Grant Number: 1160317; MINECO, Grant Numbers: SAF2016-78114-R and RTC-2015-3237-1; CIBERFES, Grant Number: CB16/10/00282; SP3-People-MC-ITN program, Grant Number: 608381; EU BATCure, Grant Number: 666918; FEDER (European regional development fund); Investissement d'Avenir, Grant Number: ANR-11-INBS-0011; French State in the context of the "Investments for the future" Program IdEx and the LabEx TRAIL, Grant Numbers: ANR-10-IDEX and ANR-10-LABX-57; French-Swiss ANR-FNS, Grant Number: ANR-15-CE37-0012. University of Nottingham; BBSRC, Grant Numbers: BB/L019396/1 and BB/K009192/1; MRC, Grant Number: MR/L020661/1; Deutsche Forschungsgemeinschaft, Grant Numbers: DFG SPP 1757, SFB 894, and FOR 2289; European Commission, Grant Number: H2020-FETPROACT 732344; Neurofibres, Grant Number: H2020-MSCA-ITN-722053 EU-GliaPhD; US National Institutes of Health, Grant Number: R01NS087611; Teva Pharmaceuticals; Agilent Technologies. IdEx, Grant Number: ANR-10-IDEX-03-02; French-Swiss ANR-FNS, Grant number: 310030E-164271; National Institutes of Neurologic Disease and Stroke at the National Institutes of Health, Grant Number: R01 NS077773; University of Zurich and the Swiss National Science Foundation; Comisión Nacional de Investigación Científica y Tecnológica, Grant Number: PB 01; Fondo Nacional de Desarrollo Científico y Tecnológico, Grant Number: 1160317; Ministerio de Economía y Competitividad, Grant Number: RTC-2015-3237-1, SAF2016-78114-R; Agence Nationale de la Recherche, Grant Numbers: ANR-10-IDEX, ANR-10-IDEX-03-02, ANR-10-LABX-57, ANR-11-INBS-0011, and ANR-15-CE37-0012; Biotechnology and Biological Sciences Research Council, Grant Numbers: BB/L019396/1 and BB/K009192/1; Medical Research Council, Grant Number: MR/L020661/1.

**Abstract**

Neuroscience is a technology-driven discipline and brain energy metabolism is no exception. Once satisfied with mapping metabolic pathways at organ level, we are now looking to learn what it is exactly that metabolic enzymes and transporters do and when, where do they reside, how are they regulated, and how do they relate to the specific functions of neurons, glial cells, and their subcellular domains and organelles, in different areas of the brain. Moreover, we aim to quantify the fluxes of metabolites within and between cells. Energy metabolism is not just a necessity for proper cell function and viability but plays specific roles in higher brain functions such as memory processing and behavior, whose mechanisms need to be understood at all hierarchical levels, from isolated proteins to whole subjects, in both health and disease. To this aim, the field takes advantage of diverse disciplines including anatomy, histology, physiology, biochemistry, bioenergetics, cellular biology, molecular biology, developmental biology, neurology, and mathematical modeling. This article presents a well-referenced synopsis of the technical side of brain energy metabolism research. Detail and jargon are avoided whenever possible and emphasis is given to comparative strengths, limitations, and weaknesses, information that is often not available in regular articles.

**KEYWORDS**

*in vitro*, *in vivo*, organization level, spatio-temporal resolution

*“Each portion of matter may be conceived as like a garden full of plants and like a pond full of fishes. But each branch of every plant, each member of every animal, each drop of its liquid parts is also some such garden or pond.”*

Gottfried W. Leibniz, *Monadology* 66–69, 1714


**1 | INTRODUCTION**

Picture a student tantalized by one of our favorite questions: *How is energy production coupled to energy demand? What are the roles of neurons and glial cells in this coupling? Information processing controls energy metabolism, but can it also work the other way round? Is epilepsy a metabolic disease? Is energy handling at the core of neurodegeneration?* In search for guidance, she participates in the biennial International Conference on Brain Energy Metabolism (<http://cecs.cl/icbem/>). After listening to talks and reading posters, she realizes that many different techniques are being applied to many experimental models. Each expert uses a different jargon: reversal potential, State 3 respiration, transacceleration, cataplerosis, glutamate enrichment, hyperpolarized, pixel shift, blood oxygenation level dependent (BOLD), default mode network, and so on. One speaker shows single ion-channel recordings. The next one pyruvate carboxylation by mitochondria, and yet another links autofluorescence of mouse hippocampal slices to a genetic disease in humans. She leaves the meeting fascinated and bewildered. *Would the channel behave similarly in situ? Is mitochondrial metabolism sensitive to cytoplasmic context? How does invertebrate metabolism relate to human metabolism?*

Our purpose here is to offer some technological guidance to our prospective colleagues (and to ourselves). Length restrictions preclude covering the ever-expanding technical range, so we apologize for possible omissions. Techniques are described and their strengths are stated briefly. Special attention is allocated to limitations, problems, and ever elusive underlying assumptions. We are not focusing here on the merits of experimental models, which are discussed in the article that introduces this special issue. Before going to the specifics, a theoretical framework may help to explain why such diverse approaches are required and why none of them may claim preference over the others.

**2 | ENERGY METABOLISM IS ORGANIZED HIERARCHICALLY**


Energy enters the brain as glucose, and leaves it minutes later in the degraded forms of water, CO<sub>2</sub>, and heat. The incessant cascade of electrons is somehow harnessed to maintain the complex structure of the brain and to process information. The science of Brain Energy Metabolism aims to figure out how this is done. It is a relatively young branch of science, stemming from well-established crafts that bring their own concepts, experimental models and techniques, assumptions, and flaws: anatomy, histology, physiology, biochemistry, bioenergetics, cellular biology, molecular biology, development, neurology, mathematical modeling, and so on. This pleiotropy is necessary because energy metabolism spans multiple organizational levels, from nanometer-scale interactions between intermediate metabolites and enzymes to organ-wide modulation of metabolism by hormones and behavior (Figure 1). The existence of structural levels and discernible modules within levels is understood by thermodynamics as



**FIGURE 1** Hierarchical organization of brain energy metabolism. Structural levels result from binding forces becoming weaker as small structures accrete into larger ones but also have an evolutionary explanation (Simon, 1962). The hierarchical organization of brain metabolism explains the success of reductionist investigative approaches

the expected result of binding forces becoming weaker as small structures accrete into larger ones but also has an evolutionary explanation, as building in modular fashion dramatically reduces error (Simon, 1962). The hierarchical organization of living structures such as the brain makes them amenable to dissection into smaller parts that remain functional. Much fundamental knowledge has been acquired from the study of isolated brain mitochondria, but *which features are context-independent and which are not?* To what extent does behavior in culture or in slices represent that observed in vivo? A given transporter was described and characterized in muscle cells, *does it work the same way in astrocytes?* To what extent is it safe to extrapolate from mouse to human? There are no a priori answers to these key

questions but a rule of thumb may be applied: the shorter the range of the phenomenon, the more likely it is to be unaffected by context. For example, the 18 atoms of glutamate are joined by covalent bonds that withstand 198°C. Because covalent bonds are much stronger than the forces that keep organelles, cells and tissues together, we know for sure that the glutamate powder on the shelf is structurally and functionally identical to the glutamate released in your visual cortex as you read these words. Therefore, applying shelf glutamate to cells in culture is meaningful. Likewise, the ligand specificity of enzymes, transporters and channels is a robust property that may be fully characterized using reconstituted proteins. However, understanding their regulation demands an intact cellular environment.



**FIGURE 2** Technical domains in brain energy metabolism research. Different techniques perform over different domains, which are in turn defined by a combination of organizational level, experimental control and relevance to human physiology. For example, electrophysiology achieves excellent experimental control at small and intermediate hierarchies by *in vitro* recording from single channels, single cells or small cell assemblies, but it is not practical *in vivo* and cannot be used in humans. MRS, in contrast, may be applied to human subjects, but has limited experimental control and provides time- and space-averaged measurements

### 3 | PICK YOUR LEVEL, PICK YOUR TECHNIQUE

The ideal technique would cover the whole organizational range, offering precise control of experimental variables while providing physiologically relevant data that may be reproduced by other laboratories. It should also be affordable and accessible. Of course, there is hardly such technique and trade-offs become necessary. The illustration in Figure 2 shows how some popular approaches distribute over the organizational space. The structure of purified enzymes and transporters is studied by means of X-ray diffraction, while understanding their function may demand complex systems like vesicles. Oxidative phosphorylation is studied by respirometry of isolated mitochondria, where substrates and cofactors are manipulated at will and fast phenomena are followed in real time, or by respirometry of cell populations, where experimental control and temporal resolution are sacrificed for the benefit of subcellular context and physiological relevance. Cellular and organ level experiments could involve imaging of metabolites concentrations in brain cells *in vitro* and *in vivo* or optical control of signaling in cells of known phenotype. At the end of the spectrum are minimally

invasive techniques like magnetic resonance spectroscopy (MRS), positron emission tomography (PET) scanning, and functional magnetic resonance imaging (fMRI), the only methods that may be used in human subjects.

### 4 | THE STAGE: EXPRESSION, LOCATION AND INTERACTIONS OF ENZYMES AND TRANSPORTERS

Energy metabolism is distributed across the diverse cellular and subcellular compartments of brain tissue. Division of labor among compartments is evident from clear segregation of key enzyme activities, for example the almost exclusive astrocytic expression of glutamine synthase and pyruvate carboxylase (Martinez-Hernandez, Bell, & Norenberg, 1977; Schousboe, Svenneby, & Hertz, 1977; Yu, Drejer, Hertz, & Schousboe, 1983), which inform on the preferential role of these cells in glutamate recycling and anaplerosis. Differences between compartments have also emerged from the study of RNA transcripts in identified cell types acutely isolated from brain tissue by Fluorescence Activated Cell Sorting (Cahoy et al., 2008; Zhang et al., 2014). Recently, the possible introduction of



expression artifacts by the sorting procedure was minimized by the ingenious expedient of co-culturing neurons and glial cells from different species followed by RNA sequencing and in-silico separation of the transcripts (Hasel et al., 2017).

#### 4.1 | How to approach structure based on microscopy


Microscopy has been the technique of choice for investigating the subcellular localization of the various enzymes, transporters, and other proteins involved in brain metabolism. In order to make these targets visible under the microscope, different labeling techniques have been used. One group is based on the conversion by the target protein of a chemical substance that becomes a dye accumulating locally within the cell and that can be visualized by bright field microscopy (e.g., 1,4-dideoxy-1,4-imino-D-arabinitol hydrochloride (DAB) staining of cytochrome oxidase (Hevner & Wong-Riley, 1989; Wong-Riley, 1989). The other major group makes use of antibodies that can specifically recognize the protein of interest (Bergersen, Magistretti, & Pellerin, 2005; Chaudhry et al., 1995; Choeiri, Staines, & Messier, 2002; Lehre, Levy, Ottersen, Storm-Mathisen, & Danbolt, 1995; Martinez-Hernandez et al., 1977; Pierre, Magistretti, & Pellerin, 2002; Pierre et al., 2009; Vannucci, Maher, & Simpson, 1997). After applying a primary antibody specific for the target protein, a secondary antibody directed against the primary antibody and carrying a tag is used to visualize its localization. The nature of the tag can vary and will determine the type of microscopy that will be subsequently used to reveal the presence of the target protein. The tag can be an enzyme which, in the presence of the appropriate substrate, will convert it into a dye that accumulates locally and can be visualized with bright field microscopy (e.g., peroxidase/DAB). If the tag is fluorescent (e.g., FITC or Cy3), then either wide-field epifluorescence microscopy may be used, which is more suitable for analysis of tissue cultures and larger overviews, or single- and two-photon laser scanning microscopy, which can provide a three-dimensional (3D) diffraction-limited view of tissues. If the tag is a heavy metal atom (silver or gold), then we can use electron microscopy, which increases the spatial resolution power compared with conventional brightfield or fluorescence microscopy. The fluorescent peptide or protein can then be visualized by fluorescence microscopy (Figure 3a). Most reagents and antibodies are commercially available or can be obtained from the researcher who produced them. Two limitations of antibody detection, particularly for comparison purposes, are variable affinity and epitope-masking. As an alternative, the mRNA expression of a gene in a given cell type may be evaluated with genetic tools. The idea is to modify the gene of interest and place the sequence coding for a bright fluorescent protein of a particular color (e.g., eGFP, Venus, and tdTomato) as a reporter to be expressed under the control of the promoter of the target protein. One example of this type of approach is the expression of tdTomato under the control of the promoter of the lactate transporter MCT1 gene introduced as bacterial artificial chromosome, which revealed preferential MCT1 expression in oligodendrocytes, a finding with consequences for glial-to-neuron lactate shuttling and the pathophysiology of white matter neurodegeneration (Lee et al., 2012).

The main strengths of these various techniques and approaches are their capacity to distinguish the specific cell type within which a particular protein is found and then in which cellular compartment or subcellular domain the protein is localized. From this information, which may be obtained under normal, untreated conditions and also after stimulation or treatment, it is possible to glean information about the participation of a protein in a particular process. In the case of immunofluorescence combined with confocal microscopy, it is possible to determine whether two proteins colocalize, that is if they are located close to each other in the same subcellular compartment. The main limitations of these approaches are the specificity and availability of the visualizing reagents. It is essential to demonstrate the specificity of each antibody and sometimes good antibodies are unavailable. Resolution can also be an issue, depending on the degree of localization that is required. For the genetic techniques, discrepancies have been observed between the expression of the native protein and the reporter. Care should be taken to avoid over-expression of reporter proteins as this could cause abnormal expression patterns or toxic effects. Although ideal for structural/localization purposes, these methods are rather limited for the study of dynamic processes.

#### 4.2 | Protein-protein interactions, supramolecular complexes, and structural microdomains

Fick's law of diffusion states that the time required for a molecule to diffuse a specific distance increases as a function of distance squared. So, doubling the distance quadruples the time. Therefore, protein-protein interactions can reduce the distance between active sites of enzymes and increase the rate of flux of substrates through multistep reactions. In fact, there is good evidence that formation of supramolecular complexes within mitochondria increases the rate of oxidative phosphorylation (Boekema & Braun, 2007) and that proton antennae in neighboring proteins increase the speed of the astrocytic lactate transporter (Stridh et al., 2012). Formation of complexes also likely improves the specificity of some reactions within nanodomains of a cell. For example, interactions with a kinase or ubiquitin ligase might limit the number of proteins phosphorylated/ubiquitinated upon activation.

Astrocytes have fine processes that contact ~100,000 synapses (Bushong, Martone, Jones, & Ellisman, 2002; Figure 3a) and endfeet processes that ensheath arterioles and capillaries (Iadecola & Nedergaard, 2007). Therefore, much like neurons, astrocytes are likely polarized cells in which proteins are selectively sorted into these functionally distinct domains and assembled into multiprotein complexes. A few studies have identified proteins that interact with the  $\text{Na}^+/\text{K}^+$  ATPase or with glutamate transporters (Bauer et al., 2012; Genda et al., 2011; Rose et al., 2009). Because these proteins are enriched near synapses, this information may help us understand the functional properties of these nano-domains. Similarly aquaporin-4 is enriched in endfeet and forms complexes with other proteins (Nagelhus & Ottersen, 2013). Essentially nothing is known about the mechanisms involved in this sorting and there is still a lot to learn about the complexes that are formed in these nano-domains and their functional significance.



**FIGURE 3** Examples of recent data in brain energy metabolism. (a) Astrocytes in organotypic hippocampal slice culture were transfected with cDNA constructs encoding for hexokinase-EGFP and GLT1-mCherry using a gene-gun. (b) *In vivo* [<sup>14</sup>C]-2-deoxyglucose uptake, obtained as described in Boussicault et al. (2014). (c) <sup>1</sup>H-MRS data obtained in a rat, at rest or during whisker stimulation. The difference between the two conditions reveals an increase in tissue lactate. The BOLD fMRI response was elicited with the same paradigm. (d) Rat neurons in culture were exposed to 3-<sup>13</sup>C<sub>1</sub> glucose so that the labeled carbon is lost via the PDH reaction but retained by the anaplerotic reactions of pyruvate carboxylase and malic enzyme, from Divakaruni et al. (2017). (e) Perturbation of hippocampal fuel fluxes during a memory task (adapted from Newman et al. (2011)). (f) Lactate-mediated depolarization and firing of a neuron in the locus coeruleus after optogenetic stimulation of neighboring astrocytes (adapted from Tang et al. (2014)). (g) The graph shows the biphasic response to astrocytic lactate levels as detected with Laconic during local stimulation of the somatosensory cortex (from Sotelo-Hitschfeld et al. (2015)). The top image shows the expression of Laconic and Pyronic in neurons and astrocytes of the somatosensory cortex using viral vectors (adapted from Machler et al. (2016)). Bottom images were taken from a transgenic mouse expressing ATeam in neurons described in Trevisiol et al. (2017)

Many interactions are identified by a hypothesis-driven approach: two proteins are found to functionally interact and one tests for protein interactions (Robinson & Jackson, 2016). In many cases, antibodies are used to immunoprecipitate a protein of interest and Western blots are used to determine whether an interacting protein is found in the immunoprecipitate (Isono & Schwechheimer, 2010; Kaboord & Perr, 2008). With discovery-based strategies, novel interacting proteins can be identified, information which is used to generate and test hypotheses. Advancements in mass spectrometry have made it relatively easy to identify proteins in immunoprecipitates. It is also possible to fuse a domain from a protein of interest to glutathione S-transferase (GST) with cloning strategies. The subsequent cDNA is then used to over-express the protein in bacteria, which is then purified and incubated with solubilized tissues of interest. Complexes are then isolated using

glutathione-based affinity columns (Harper & Speicher, 2011). Variants of this strategy are used to test whether interactions between two proteins are direct and to map domains of proteins required for a given interaction. Finally, one can use yeast two-hybrid based strategies to identify novel interacting proteins. Here, a domain from a protein of interest is fused to part of a transcription factor. Then, the hybrid is used to screen a library of target sequences fused to the rest of the transcription factor and interactions are identified as transcriptional activity (Ito et al., 2001). Candidates can be tested for colocalization in tissues or cells of interest, for example by looking for covariance (Li et al., 2004). If one or both of the proteins are widely distributed, it is possible to determine whether the observed overlap occurs more frequently than that which would occur by chance using Monte-Carlo simulation (Genda et al., 2011).



There are several issues with any of these approaches. First, it is at least theoretically possible that some interactions form after solubilization. This post-solubilization aggregation is likely due to either electrostatic or hydrophobic interactions and will increase linearly with protein abundance. Sequential rinses with buffers containing different concentrations of salt can reduce these physiologically irrelevant interactions. Second, antibodies used to immunoprecipitate a protein may directly bind to and precipitate nonspecific targets. This can be partially addressed by performing “reverse immunoprecipitations” with an antibody against the putative interacting protein. Third, the GST and yeast two-hybrid approaches require guessing where potential interacting proteins might bind. Fourth, there are several different types of detergents, including nonionic, bile acid salts, ionic, and so on. One needs to empirically identify the detergents that best solubilize the protein of interest and reduce nonspecific binding of proteins in the lysate (Garavito & Ferguson-Miller, 2001; Le Maire, Champeil, & Moller, 2000; Phizicky & Fields, 1995; Seddon, Curnow, & Booth, 2004).

It is ultimately important to rigorously demonstrate the relevance of an interaction *in vivo*. Super-resolution microscopy has been improved to the point that the resolving power is in the 10–50 nm range. Although this resolution is not sufficient to demonstrate direct protein interactions, it can support the notion that two proteins exist in the same nanodomain. Mapping domains required for an interaction allow for generation of constructs or peptides that will function as dominant-negative inhibitors of the interaction. Of course, if multiple “interacting” proteins are identified, we must guess which proteins form direct interactions; this is a daunting task because there may be many proteins in a supramolecular complex. For example, the postsynaptic density has about 2,000 proteins (Focking et al., 2016). It is also possible to test for interactions by labeling two putatively interacting proteins with different fluorescent tags. One of the fluorophores is excited and energy is transferred to a second fluorophore by resonance. This technique is called intermolecular Förster resonance energy transfer (FRET) and only detects proteins that are within 10 nm of each other (Jares-Erijman & Jovin, 2003; Pietraszewska-Bogiel & Gadella, 2011; Sekar & Periasamy, 2003).

## 5 | THE PLAYERS: FUNCTION OF ENZYMES AND TRANSPORTERS

### 5.1 | Enzyme activity

Once the location and abundance of an enzyme have been established, the question of function is next to arise. To assess the specific activity of an enzyme time-dependent changes in the concentration of the substrate (or product) need to be determined, during the steady state, in tissue/cell homogenates under specified conditions (Lorsch, 2015), and at a wide range of substrate/ligand concentrations to obtain the main kinetic parameters: half-saturation or Michaelis constant ( $K_m$ ), half-inhibition constant ( $K_i$ ), maximal velocity constant ( $V_{max}$ ) and turnover number ( $k_{cat}$ ; Brooks et al., 2004; Purich, 1979; Purich, 1980; Purich, 1982). This is essential when predicting metabolic fluxes controlled by enzymes according to substrate concentrations (Herrero-Mendez et al.,

2009; Kaplan & Colowick, 1955), tissue damage (Adan, Kiraz, & Baran, 2016) or drug discovery (Widder & Green, 1985; Zhang, 2017).

The specific activity of enzymes is a very useful measurement, and is necessary for understanding the functional fraction. This is important for enzymes that, for instance, are highly susceptible to oxidation of key cysteine residues leading to inactivation, such as mitochondrial respiratory chain complexes (Bolanos, Peuchen, Heales, Land, & Clark, 1994; Bolanos, Heales, Land, & Clark, 1995). Subcellular fractionation followed by enzymatic analysis represents a good strategy for estimating specific activity within the natural environment (Murphy & Chan, 2014). The development of techniques for the use of in-gel activity assays for proteins in their native tertiary/quaternary structures, such as blue-native gel electrophoresis followed by NADH dehydrogenase activity for mitochondrial Complex I (Lopez-Fabuel et al., 2016; Murphy & Chan, 2014) may also represent progress towards the characterization of enzyme activities in their natural conformation.

The main drawbacks of enzymatic analyses are technical and of interpretation. Technically, the unfeasibility of detecting the substrate or the product forces that couple enzymatic reactions (Purich, 1982), although the product of which—usually NAD(P)H—can be easily determined. Whereas coupling often represents the only available approach for assessing the specific activity of an enzyme, it often generates parasitic reactions (Auld & Inglese, 2004) that mask, usually by overestimation, the actual enzyme activity. When the limitation is due to the low sensitivity of the method to detect the substrate/product, alternative approaches can be implemented to enhance sensitivity. Often, the use of fluorescent, luminescent, and radiometric detection methods can overcome the sensitivity problems (Dahlin, Baell, & Walters, 2004; Purich, 1980; Simeonov & Davis, 2004), although they require the use of more complex equipment and expensive probes. Kinetic parameters of enzymes responsible for post-translational changes in proteins are difficult to assess as they require more tedious analyses such as Western blotting followed by semi-quantitative estimation of the modified substrate (e.g., phosphorylation, ubiquitination, acetylation, and so on; Glickman, 2004), hence complicating the accurate knowledge of the kinetic parameters of the enzyme. Another limitation of this technique is the absence of the physiological environment of the enzyme when its activity is determined. Kinetic parameters of enzymes are determined by adding cofactors and substrates to the reaction mixture at concentrations that might be far from the physiological concentrations in the cell, especially if saturating concentrations are used for determining  $V_{max}$ . Furthermore, the specific cellular or subcellular localization of enzymes is not reflected in the reaction mixture, although efforts should be made to mimic the in-cell conditions (Dahlin et al., 2004). For enzymes that are naturally membrane bound, the use of detergents may help solubilize the protein for easier access of substrates and cofactors, but very often the detergent interferes with light, fluorescent or luminescent detection methods (Simeonov & Davis, 2004). When the detergent cannot be used, due to interferences, the insolubilized membranes cause light scattering, which may preclude spectrophotometric analyses (Brooks et al., 2004). Finally, the specific activity, which represents maximal activity, gives no information about the influence of other enzymes of the pathway on the metabolic flux (Almeida,

Moncada, & Bolanos, 2004). A well known example is hexokinase, which under physiological conditions works at 10% of its maximal activity, as it is tonically inhibited by its product glucose-6-phosphate (Wilson, 2003).

## 5.2 | Activity of transporters and channels: Tracers, fluorescence, and electrophysiology

The wide-spread availability of radioactive tracers at the dawn of the nuclear age led to detailed kinetic understanding of several membrane transporters well before their molecular identification. For practical reasons, the best characterized transporters were (and still are) those expressed in erythrocytes, in particular the facilitative hexose transporter GLUT1 (LeFevre, 1948; Widdas, 1954), later found to dominate the glucose permeability of endothelial cells, astrocytes, and oligodendrocytes. Radioactive indicator-dilution assays were developed for *in vivo* studies in animals that allowed the kinetic characterization of the transport mechanisms at the blood-brain barrier (Crone, 1965; Yudilevich & Sepulveda, 1976). In human, brain glucose transport parameters could be estimated by combining steady-state measurements of parenchymal glucose concentration obtained by MRS, microdialysis, and microelectrodes with mathematical modeling (Barros, Bittner, Loaiza, & Porras, 2007; Gruetter, Ugurbil, & Seaquist, 1998). Decades of radiotracer experiments in cultured cells and heterologous expression systems have shown that the selectivity and kinetic properties of transporters for energy metabolites such as glucose, lactate, pyruvate, glutamate, and glutamine are insensitive to cellular context. What is less well understood is their regulation, that is, if and how the intrinsic activity and subcellular localization of the transporters responds to the large changes in metabolic demand that accompany neural activity. Excellent in terms of signal-to-noise ratio, radiotracer methods offer limited spatial and temporal resolution, which precludes the study of single cells, complex tissues or rapid regulatory phenomena. In the case of  $H^+$ -coupled monocarboxylate transporters (MCTs), the carriers of lactate, pyruvate and ketone bodies, indirect single-cell characterization has been possible by means of intracellular pH electrodes or pH-sensitive dyes like 2',7'-bis-(2-carboxyethyl)-5-(6)-carboxyfluorescein. Single cell glucose transport determination is possible with the fluorescent glucose analogs 6- and 2-([N-(7-nitrobenz-2-oxa-1,3-diazol-4-yl)amino]-2-deoxy-D-glucose (6-NBDG and 2-NBDG; Speizer, Haugland, & Kutchai, 1985; Yoshioka et al., 1996). They are twice as big as glucose, and are therefore taken by GLUTs at a lower rate (Barros et al., 2009a), a property that can be taken advantage of to achieve real-time measurement of transport activity and thus detect fast regulatory changes (Loaiza, Porras, & Barros, 2003; Porras, Loaiza, & Barros, 2004; Porras, Ruminot, Loaiza, & Barros, 2008). Detected by confocal microscopy in brain tissue slices or by two-photon microscopy *in vivo*, they have served to estimate the contribution of neurons and astrocytes to the uptake of sugar (Barros et al., 2009b; Chuquet, Quilichini, Nimchinsky, & Buzsaki, 2010; Jakoby et al., 2014). At present, their performance has been surpassed by that of genetically-encoded fluorescent sensors (Section 6.3), which provide a stronger signal and a direct readout of the metabolite of interest, avoiding the need for

assumptions and controls about analog handling by transporters and enzymes.

A substantial fraction of the brain's ATP turnover is associated with the recovery of ion gradients dissipated by ion channels during synaptic activity (Harris, Jolivet, & Attwell., 2012). In addition, many metabolites possess a net charge and recent studies in glial cells have indicated that some of them may be transported by ion channels (Kara-giannis et al., 2016; Sotelo-Hitschfeld et al., 2015). This makes electrophysiology an important tool not only for the manipulation, but also for the understanding of brain energy metabolism. The electrical properties of brain cells are typically studied in acute tissue preparations and in cell cultures. In contrast to neurons, which are characterized by action potentials, glial cells have no swift electrical events taking place across their membranes. In addition, the plasma membranes of astrocytes and oligodendrocytes are extremely leaky due to high background activity of potassium channels. There are also no fast excitatory or inhibitory synapses on astrocytes or oligodendrocytes and therefore, stimulation of an afferent pathway commonly used for activation of neurons cannot be used to synchronously "fire" them. For these reasons, electrical recordings are less commonly used in studies of these cells. Recordings with extracellular electrodes can detect stimulus-evoked compound action potentials (CAPs; Stys, Ransom, & Waxman, 1991), where changes in the extracellular ion distribution due to glial transporters do play a role, but it is very hard to separate the glial component from the contributions of other tissue elements. From the metabolic point of view, the *in vitro* recording of CAPs offers the advantage that the tissue can be maintained for extended periods in a perfusion chamber with temperature, degree of oxygenation, and nutrient supply controlled by the experimenter. In a homogeneous tissue such as the optic nerve, the CAP represents the response of a common neural cell type, central white matter myelinated axons. However, in less homogeneous tissue, for example, the hippocampus, the recorded response is from heterogeneous cell types is recorded, therefore, the response may not be a true reflection of the contribution of individual neural cell types (Brown, Tekkok, & Ransom, 2003; Cater, Benham, & Sundstrom, 2001). *In vitro* recordings of acutely isolated tissue may be problematic with regards to the diffusion barrier created by large pieces of tissue, where inadequate diffusion of oxygen and glucose may lead to anoxic and/or hypoglycemic conditions at the center of the tissue (Tekkok, Brown, Westenbroek, Pellerin, & Ransom, 2005).

Recordings with sharp glass electrodes are possible for large glial cells (Deitmer, 1992). Intracellular recordings in patch-clamp mode seem to be more suitable for studies of astrocytes, which are usually much easier to patch than neurons in spite of their small size. Using the patch clamp approach, it is possible to perform an accurate pharmacological characterization of receptors expressed by astrocytes (Lalo et al., 2008; Palygin, Lalo, & Pankratov, 2011). However, such experiments work best when astrocytes are isolated, because it is not always easy to localize the site of action of the drugs when applied to tissues. Experiments *in vitro* allow tight control of the perfusion media including concentration of metabolites and drugs and the recordings are much more stable. However, astrocytes are very sensitive to stress, which is inevitable during preparation of slices and they easily convert to a





“reactive” phenotype that may have a somewhat different physiology to the naïve cells *in situ*. In addition, metabolic conditions in a slice never correctly mimic conditions *in vivo*. Most acute slice experiments are performed in solutions bubbled with 95% O<sub>2</sub>/5% CO<sub>2</sub> mix, resulting in hyper-oxygenation of the superficial cells. For some experiments, this is a problem as shown for example in studies of neuro-vascular coupling where the O<sub>2</sub> level was eventually reduced to 20% (Hall, Klein-Flugge, Howarth, & Attwell, 2012). However, using lower oxygen can easily lead to hypoxia in the center of a moderately thick slice (300 μm). In a similar vein, virtually all standard slice work is performed in solutions containing 10 mM glucose or more, whereas brain tissue glucose is ~1 mM (Barros et al., 2017). This overloads superficial astrocytes with glucose. Finally, cultured astrocytes are ideally suited for patch recordings but culturing affects the morphology, physiology, and metabolism of these cells, factors that can be ameliorated by coculturing with neurons (Hasel et al., 2017; Mamczur et al., 2015). In summary, patch clamp is perhaps the best electrophysiological approach for studying astrocytes, but caution is required when extrapolating from data obtained on isolated astrocytes. It should also be remembered that while recording with patch electrodes, we artificially impose on the cell our ideas about the intracellular concentration of ions and metabolites because pipette solutions rather quickly dialyze into the astrocyte. Therefore, our conclusions about the currents in these cells are only as accurate as our knowledge of their native intracellular ion composition.

### 5.3 | Optogenetic actuators: Manipulating metabolism

Optical control of electrical and signaling events in brain tissue *in situ* is in principle highly attractive because it seems to be noninvasive, fast, and can be targeted to individual cells and cellular populations. Although light-sensitive proteins that affect cellular functions in plants have been known of for a long time, a major shift in neuroscience took place around 2003–2006 when it was demonstrated that such proteins can be expressed in mammalian neurons and used to activate them (Bi et al., 2006; Boyden, Zhang, Bamberg, Nagel, & Deisseroth, 2005; Nagel et al., 2003). Channelrhodopsin-2 (ChR2), which was (and still is) the most popular member of this family behaves as a light-gated channel permeable to sodium, potassium, protons, and Ca<sup>2+</sup>. Manipulating sodium conductance presents an obvious way to depolarize neurons and trigger action potentials on demand. This is by far the most common use for this protein, which can be targeted to specific populations of cells using molecular and genetic approaches, and it will be very interesting to exploit it to investigate the role of specific neurons on local energy metabolism. ChR2 has also been used to activate astrocytes (Figure 3f; Gourine et al., 2010), where it triggered Ca<sup>2+</sup> elevations, although this required an application of several seconds of pulsing blue light (NB: constant blue light is a notorious source of artifacts). The Ca<sup>2+</sup> increases were largely dependent on thapsigargin-sensitive internal Ca<sup>2+</sup> stores (Figueiredo et al., 2014; Perea, Yang, Boyden, & Sur, 2014). ATP release was also documented from ChR2-stimulated astrocytes (Gourine et al., 2010). Many studies in the meantime have used ChR2 to activate astrocytes, although it should be

noted that this is not an ideal tool for controlling these cells. A significant Na<sup>+</sup> influx most likely affects multiple Na<sup>+</sup>-dependent transporters on astrocytes and the downstream effects become hard to disentangle. Prolonged ChR2-mediated activation leads to acidification of astrocytes due to a build-up of lactate (Tang et al., 2014). Since ChR2 does not mimic any obvious endogenous physiological mechanism operating in astrocytes, it is reasonable to explore alternatives. Attempts to use exotic approaches such as the light sensitive proton pump Arch have also been made (Beppu et al., 2014; Poskanzer & Yuste, 2016), but the actions of this protein on astrocytes are even less clear than those of ChR2. Light-activated G-protein coupled receptors (GPCR) have been used (Tang et al., 2014) with the assumption that via this route events can be evoked that are similar to the native signaling in the brain where astrocytes are mainly controlled via GPCRs. These constructs, however, need to be used with caution because, when expressed at high level, GPCRs have a tendency to signal without any external trigger. Therefore, overexpression, commonly used in many experiments, can be seen as a problem. An interesting new approach is to use a dimerization strategy for optical activation of signaling, for example, activities of kinases. For astrocyte research a construct has been introduced that triggers Ca<sup>2+</sup> entry via the ubiquitous Calcium Release-Activated Channels channels (Kyung et al., 2015). Even though this tool may not be fast enough for triggering really rapid events, it seems that for many cases it can be advantageous, because at least it is a clearly defined mechanism for increasing astrocytic Ca<sup>2+</sup>. To summarize, the use of optogenetics for the study of brain energy metabolism is just starting, with a promising future of ever more specific and powerful ways to manipulate individual cells.

## 6 | THE PLAY: METABOLITE CONCENTRATIONS AND FLUXES

### 6.1 | Respirometry

Neural activity is tightly coupled to local fuel oxidation. The oxygen consumption rate (OCR) is therefore highly valuable information when characterizing cellular bioenergetics or diagnosing diseases. After decades of productive use for research and teaching, the estimation of oxygen consumption with manometry (Warburg, 1928) was superseded by easier and more accurate, real-time determinations by using polarography (Clark, Wolf, Granger, & Taylor, 1953). While the traditional Clark-type electrode has delivered most of our current knowledge of mitochondrial bioenergetics, it has disadvantages (Zhang et al., 2012). Oxygen often leaks from the chamber through junctions and holes used for the administration of drugs and inhibitors; assembly of the equipment is tedious, time consuming and requires frequent calibration; the electrode is easily oxidized or damaged, and thus has a short lifespan; the need for continuous stirring to mix the cell or mitochondrial suspension usually harms the sample, in particular when analyzing isolated intact mitochondria; and since several of the inhibitors used to characterize the contribution of mitochondrial complexes to oxygen consumption are lipid soluble, the chamber needs to be thoroughly washed between samples. While the chamber size is often fixed, the internal volume can

be adjusted between 0.3 and 2 ml. However, the best results are often obtained with large sample volumes (usually, 1 ml), thus requiring large amounts of biological material for the analysis. This makes the system unsuitable for screenings. The appearance of multiwell-plate respirometers has fired up the field (e.g., Agilent technologies/Seahorse Bioscience, Presens, Santa Clara, CA). These devices are built to simultaneously determine OCR and extracellular acidification rate (ECAR) in 24- or 96-well plates of adherent rather than suspended cells. OCR is determined using an oxygen-sensing fluorophore instead of the traditional oxygen-consuming electrode, and ECAR, with a pH sensor. Although popular, the high cost of both the instrument and the single-use plates are a disadvantage. In addition, ECAR is often thought to reflect glycolysis rate because the release of lactate acidifies the extracellular medium. However, caution should be taken because the CO<sub>2</sub> released by the tricarboxylic acid activity significantly contributes to extracellular acidification (Mookerjee, Goncalves, Gerencser, Nicholls, & Brand, 2015; Mookerjee, Gerencser, Nicholls, & Brand, 2017). The Clark electrode and the fluorophore-based multi-well instruments are highly valuable tools for characterizing aerobic metabolism in biological samples. Choosing one or the other depends on sample type and on flexibility in the study design. For instance, the Clark-type electrode chambers are very useful when the sample size is not limiting, as it provides more accuracy and time resolution, but the method is time consuming, such that only a few samples can be assayed in a day. This approach is also suitable for tissue slices or isolated mitochondria in suspension, as well as intact or permeabilized cells in suspension. It should be noted that harvest of neurons from a culture dish for assay in suspension leaves behind the neuronal processes and the mitochondria within them. With both assay platforms, a large number of substrates, inhibitors or uncouplers can be used to better characterize the bioenergetics of the sample. Selective permeabilization of the plasma membrane allows control of substrate delivery to the mitochondria, and avoids the need for isolation of mitochondria from cultured cells (Divakaruni, Rogers, & Murphy, 2014). The multi-well apparatus, in contrast, is very useful for accurate screenings in intact adherent cells, permeabilized adherent cells, or isolated mitochondria and requires a smaller sample size (~1–3 × 10<sup>4</sup> cells/well or 1–3 micrograms of isolated mitochondria/well; Divakaruni et al., 2017; Rogers et al., 2011). This is particularly important when determining OCR in highly differentiated cells, such as neurons, given the low yield in these preparations. Note that multi-well respirometry is routinely employed in the absence of bicarbonate, a substrate of the anaplerotic enzyme pyruvate carboxylase and a regulator of astrocytic glycolysis (Ruminot et al., 2011). Multi-well respirometry is also constrained in terms of temporal resolution, with a typical rates integrated >2–5 min. One limitation common to both oxygen electrodes and multi-well methods is the need for homogenous cell populations, which is not ideal for the study of brain cells, as neurons and astrocytes foster their mutual functional and metabolic differentiation (Barres, Koroshetz, Chun, & Corey, 1990; Hasel et al., 2017; Mamczur et al., 2015). A limitation of the use of isolated brain mitochondria in either apparatus is the fact that the mitochondria arise from multiple different cells types, thus specific information on the function of neuronal versus astrocytic mitochondria is lost in this

approach. However, respirometry can be performed on preparations of synaptoneurosomes, which are resealed synaptic vesicles created upon homogenization of brain tissue that contain mitochondria from synaptic terminals (Choi, Gerencser, & Nicholls, 2009).

## 6.2 | Extracellular microdialysis and enzyme-based probes

Given that the interstitial fluid is the medium of exchange between blood, the cerebrospinal fluid and the intracellular compartments of neuron and glia (Ransom, 2009), the ability to measure interstitial metabolites is crucial. However, the space left between brain cells is only 20 nm wide, which makes probing its composition a major technical challenge. Dialysis probes have been used to measure metabolic substrates from awake, moving animals (Brown, Skamarauskas, Lister, Madjd, & Ray, 2011). Placement of the dialysis probe requires complex surgery from which the animal recovers. The probes can be permanently attached to the animals' skull with the sensing component extending into the brain parenchyma. The probe is guided through the brain parenchyma until a desired location is reached, which creates considerable localized damage and forms a third space that fills with the contents of damaged cells, and whose composition clearly does not represent the baseline interstitial fluid. However, this perturbation dissipates over time as local repair mechanisms tend to normalize the third space contents. The interstitial fluid is sampled by applying backpressure, such that a small volume is drawn up into the probe (5–10 μl) to be later analyzed by high-performance liquid chromatography (HPLC). Repeated samples can be withdrawn over extended periods of time (days to weeks) thereby facilitating longitudinal studies. Advantages of this technique include its use for extended periods and the accuracy of HPLC analysis. Some disadvantages are that it does not provide real-time information, it disrupts the tissue, has low spatial resolution. It is a hotly debated area whether the samples reflect the real environment in the interstitial space.

Real-time measurements of brain interstitial fluid can be obtained with enzyme-based microelectrodes. Probes can be as fine as 7 μm in diameter with the active tip between 100 and 500 μm in length. The greater the surface area of the tip, the greater the temporal resolution. Such probes are useful for recording from homogeneous brain structures such as isolated optic nerve (Yang et al., 2014b) and hippocampus (Figure 3e; Frenguelli, Llaudet, & Dale, 2003; Newman, Korol, & Gold, 2011). However, as the probe records the aggregate signal from the entire tip area, care should be taken in interpreting data acquired from brain areas with heterogeneous cell populations. Those probes currently available can sense a multitude of compounds of interest to metabolic studies, including ATP, adenosine, acetylcholine, glucose, lactate, D-serine, choline, and glutamate. The sensors can be calibrated before the experiment and are thus able to provide real-time estimates of concentrations, and they can be used for both *in vivo* and *in vitro* studies. A great advantage of modern commercially available sensors is that several probes may be used simultaneously to measure the concentrations of multiple metabolically relevant compounds, for example, glucose and lactate, ATP, and adenosine (Figure 3e; Frenguelli et al., 2007; Newman



et al., 2011). Microsensors can be used in parallel with other real-time techniques, for example, electrophysiology (Brown, Evans, Black, & Ransom, 2012; Yang et al., 2014b) and genetically-encoded intracellular nanosensors (Machler et al., 2016). The power of combining functional assays with the real-time measurement of metabolic substrates has proven to be very effective in determining key processes in both grey and white matter.

### 6.3 | Genetically-encoded fluorescent nanosensors for metabolites

Metabolite nanosensors are chimeric proteins that change their optical properties upon binding a ligand (Fehr, Lalonde, Lager, Wolff, & Frommer, 2003; Tsien, 2009). Their specificity is conferred by a recognition domain, such as a bacterial periplasmic protein or a transcription factor, and their fluorescence by one or two fluorescent proteins, such as GFP, mTFP, or Venus. Many of them are based on intramolecular FRET. Expressed in cellular compartments, the nanosensors are detected by microscopy: wide field, confocal, or two-photon. In brain cells, measurements have been reported for glucose, glutamate, ATP, ATP/ADP, NADH/NAD<sup>+</sup>, lactate, and pyruvate, reviewed in San Martín et al. (2014b). Most sensors are easily available through the non-profit repository Addgene. The main strength of these tools is resolution, which for high-end microscope setups is in the order of seconds and micrometers. Therefore, they provide a real-time estimation in single cells. Being genetically-encoded, they can be targeted to organelles and subcellular domains (Pandin, Greotti, Lefkimmiatis, & Pozzan, 2017). Their most common and robust use is to report metabolite fluctuations in before-and-after experiments, for example, when characterizing the effect of a physiological or pharmacological perturbation (Figure 3g). Using standard transfection or viral expression systems, sensor concentration is ~10  $\mu$ M (Miyawaki, Griesbeck, Heim, & Tsien, 1999), which is lower than the physiological concentration of most metabolites and therefore metabolite sequestration and slow dissociation kinetics are not usual concerns. An important exception is NADH, which is in the nanomolar range.

As they are proteins, nanosensors are potentially susceptible to their physical environment. Some of the early versions were sensitive to pH, which could be corrected by parallel pH measurements. Their main weakness is that they are difficult to calibrate. One important caveat, which is not always explicit, is that it is not safe to draw conclusions from differences in intensity between cells. Even for FRET-based and other ratiometric sensors, the readout is affected to some extent by expression level due to nonlinearities in the optical path. To compare cells, a calibration is therefore required. The simplest one-point calibration is achieved by forcing a reading at one end of the binding curve (i.e., zero or saturation) and then using the affinity constant and maximum fluorescence change determined *in vitro* to extract concentrations by interpolation (Barros, Baeza-Lehnert, Valdebenito, Ceballo, & Alegría, 2014). If both ends of the saturation curve are accessible, a more robust 2-point calibration protocol becomes feasible, where only affinity needs to be assumed as shown for the pyruvate sensor (San Martín et al., 2014a). Avoiding assumptions by recurring to a full *in situ*

calibration is possible (Bittner et al., 2010) but is seldom practical, as it demands harsh permeabilization and/or metabolic inhibition procedures that may perturb the sensor itself. When quantitative data are available, nanosensors may be used to measure absolute flux by means of inhibitor-stop protocols (Bittner et al., 2010; San Martín et al., 2013; San Martín et al., 2014a). As both calibration and quick transporter inhibition are needed, this kind of assay is currently restricted to cells in culture and in slices. Inhibitor-stop methods assume that there is no quick adaptation to the inhibition, which has to be evaluated on a case-to-case basis.

Nanosensing is an emerging technology and probes for other metabolites are surely on their way. One example is a sensor for NADP<sup>+</sup> (Cameron et al., 2016), which may help in tackling the cryptic pentose phosphate pathway (Bouzier-Sore & Bolanos, 2015). In parallel, existing sensors are being improved and extended in terms of brightness, affinity, dynamic range, and spectra (e.g., red-shifted for deeper tissue penetration by reducing scattering). One important objective at hand is the generation of extracellular sensors to approach the brain interstice in noninvasive manner.

### 6.4 | *In vivo* optical imaging: From autofluorescence to genetically-encoded sensors and transgenic animals

Brain energy metabolism has been the focus of studies since optical imaging methods began in the 1960s (Chance, Cohen, Jobsis, & Schoener, 1962), where changes in tissue UV autofluorescence observed upon altering respiration *in vivo* were interpreted as changes in the NAD<sup>+</sup>/NADH ratio (Mayevsky & Chance, 1975). This and its relative flavoprotein autofluorescence imaging have been widely applied (Kasischke, Vishwasrao, Fisher, Zipfel, & Webb, 2004; Reinert et al., 2011; Shibuki et al., 2003; Shuttleworth, 2010; Weber et al., 2004). In the early reports, the dynamics of hemoglobin oxygenation was discussed, a phenomenon that was later exploited by Grinvald and colleagues to map brain activity (Malonek & Grinvald, 1996), and which paved the way for the development of BOLD fMRI (Ogawa, Lee, Kay, & Tank, 1990). Oxy- and deoxyhemoglobin can be discerned optically due to their differential absorption spectrum. However, the intrinsic optical contrast mechanisms have only marginally helped to generate mechanistic insight into cell-specific metabolic processes, because the signals are difficult to discriminate from each other, and hard to quantify and interpret (see Box 1). After their successful use in cultured cells and brain tissue slices (Section 5.4), fluorescently labeled substrates were used to investigate energy metabolism in combination with two-photon microscopy (Denk, Strickler, & Webb, 1990; Zipfel, Williams, & Webb, 2003). Using the nonmetabolized fluorescent glucose analog 6-NBDG, an acute increase in glucose uptake was detected in astrocytes but not in neurons of the somatosensory cortex of stimulated rats (Chuquet et al., 2010). However, the optical properties of NBDGs are not favorable because of a very low two-photon cross-section. Alternative fluorescent glucose analogues were developed (Kovar, Volcheck, Sevick-Muraca, Simpson, & Olive, 2009; Tian et al., 2009) and used *in vivo* (Lundgaard et al., 2015), but the data remain difficult to interpret because some are hardly transported due to high molecular weights of

1.3 kDa (Kovar et al., 2009; Lundgaard et al., 2015). The advent of genetically encoded sensors for energy substrates (Section 6.3) promises to be a significant advance for the field of brain energy metabolism, as they offer cellular resolution and real-time measurements (Figure 3g; Lerchundi et al., 2015; Machler et al., 2016; Sotelo-Hitschfeld, Fernández-Moncada, & Barros, 2012). In combination with advanced viral expression vectors and hopefully transgenic mice (see below), these sensors are particularly well suited for *in vivo* research using two-photon microscopy. Quantitative measurements of partial pressure of oxygen ( $pO_2$ ) *in vivo* can be obtained with phosphorescence lifetime imaging microscopy, in which a phosphorescent dye produces a decaying light signal after a short illumination. This decay is shortened by the presence of oxygen and its half-life can be translated to  $pO_2$  in absolute units. New phosphorescence probes were recently developed that can be two-photon excited (Esipova, Rivera-Jacquez, Weber, Masunov, & Vinogradov, 2016; Esipova & Vinogradov, 2014), which greatly widens the range of *in vivo* applications (Lecoq et al., 2011; Lyons, Parpaleix, Roche, & Charpak, 2016; Parpaleix, Goulam, & Charpak, 2013; Sakadzic et al., 2010; Sakadzic et al., 2011). Until recently, high resolution imaging was only possible in anesthetized animals. A general concern for *in vivo* optical imaging and *in vivo* experimentation are the effects of anesthesia on metabolic processes. For this reason, an increasing number of laboratories are now using head-restrained awake rodents even with two-photon microscopy at single cell resolution.

#### 6.4.1 | Transgenic animals for metabolic imaging

The study of brain structure and function has been significantly facilitated by the availability of transgenic reporter animals, from nematodes to mammals, contributing, for example, to the discoveries of the neurogenic potential of embryonic glia (Malatesta et al., 2003) and microglial motility (Nimmerjahn, Kirchhoff, & Helmchen, 2005). More recently, genetically encoded sensors have been used as biosensors to visualize transmitter- and behavior-evoked  $Ca^{2+}$  signals (Akerboom et al., 2012; Paukert et al., 2014). In transgenic animals the sensors are expressed permanently at constant levels over generations, a distinct advantage over viral transduction, which requires injection of each animal, with the possibility of tissue injury and inflammation. In addition, transgenesis provides a higher degree of flexibility regarding cell-specific expression and size of the reporter construct. The generation of transgenic mammals is however time consuming, labor intensive, and expensive. Another important drawback is the scarcity of appropriate promoters to reach sufficient expression levels in specific cell types, but strategies have been developed to address these limitations (Hirrlinger et al., 2009; Madisen et al., 2015). To our knowledge, there are only two transgenic mouse models to measure metabolites in brain tissue at this time: one that ubiquitously expresses a nanosensor for pyruvate (Bulusu et al., 2017) and the other specific for neuronal ATP (Figure 3g; Trevisiol et al., 2017). The pyruvate sensor was used to reveal the spatiotemporal dynamics of tissue explant glycolytic activity during development and the ATP sensor allowed imaging, for the first time, of the frequency-dependent energy demand of optic nerve axons.

The age of metabolism studies using transgenic animals is just beginning. Novel genetic tools such as CRISPR/Cas9 (Cong et al., 2013; Wang et al., 2013) facilitate particularly the generation of mouse lines with loxP-flanked transcriptional “stop” sequences followed by nanosensors targeted to a single well-defined gene locus (e.g., at ROSA26). Combinatorial breeding of mice with expression of inducible and cell-specific Cre-DNA recombinase and ROSA-nanosensor mice will provide promising opportunities to explore the many facets of energy metabolism in the mammalian brain. Transgenic *C. elegans* and *D. melanogaster* lines have been generated that express an ATP sensor (Tsuyama et al., 2013). Recently, fruit fly lines selectively expressing pyruvate and glucose sensors in mushroom body neurons helped to show that energy metabolism can play an instructional role in long-term memory encoding (Plaçais et al., 2017), a role that appears to extend to behavior (Li-Byarlay, Rittschof, Massey, Pittendrigh, & Robinson, 2014). Another fruit fly study expressed a glucose sensor to characterize the delivery of glucose to brain cells (Volkenhoff, Hirrlinger, Kappel, Klämbt, & Schirmeier, 2017). Short life cycle, relatively inexpensive experimentation and the availability of powerful tools for genetic manipulation, renders worms and flies very attractive for comparative studies of brain energy metabolism using imaging probes.

#### 6.5 | 2-Deoxy-[ $^{14}C$ ]glucose and [ $^{18}F$ ]fluorodeoxyglucose methods for the measurement of local glucose utilization

2-Deoxy-D-glucose (2DG) is an analogue of glucose (a hydrogen atom replaces the hydroxyl group on the second carbon) and this single structural difference is responsible for the chemical properties that make 2DG so suitable for the measurement of the rates of glucose utilization in the brain *in vivo*. 2DG is transported in the cell and metabolized qualitatively like glucose but only through the first step of the glycolytic pathway to provide 2DG-6-phosphate (2DG-6P). Since 2DG-6P is not metabolized further to a significant extent, its rate of accumulation is proportional to the amount of glucose used by the cell. Using an appropriate mathematical model of enzyme kinetics and [ $^{14}C$ ]-labeled 2DG associated with autoradiography (a technique using X-ray films to visualize radioactive molecules with a spatial resolution of approx. 50  $\mu m$ ), the [ $^{14}C$ ]-2DG method provides quantitative metabolic maps (in  $\mu mol$  of glucose/g/min) of the brain in which regions of altered activity can be easily visualized (Sokoloff et al., 1977). This method can be used in different species, yields massive amounts of data and has provided unequivocal evidence that energy metabolism is linked to local functional activity. This method was adapted for use in humans with [ $^{18}F$ ]fluorodeoxyglucose (FDG) and a PET scanner (Reivich et al., 1979), which remains useful as a tool for mechanistic investigation of brain energy metabolism in animals (Zimmer et al., 2017).

Absolute quantification of glucose utilization is possible thanks to the use of  $^{14}C$ -2DG, but many laboratories lack authorization to work with long life radioactive isotopes. The operational equation based on the original model requires plasma glucose (glucose oxidase method) and [ $^{14}C$ ]-2DG concentrations (liquid scintillation counter) to be sampled frequently during a 45-min period, for which intravenous and



intra-arterial catheters are inserted. While this is feasible in large animals, this can represent a technical limitation in mice, the choice animal model in the neurosciences. Also, the method requires that the brain of the animal is processed for autoradiography soon after completion of the experiment. Therefore only one measurement is possible per animal. The brain is cut entirely into 20- $\mu\text{m}$ -thick sections with a cryostat and dried sections mounted on coverslips are left in contact with an X-ray film or phosphor screens. After exposure data can be read out with means of a specific image analysis system or phosphor-imager. The method cannot measure individual cells so that it is not possible to separate, for example, the contribution of neurons and glial cells. 2DG and to a larger extent FDG (high electronegativity of the fluorine-18 atom) do not have the same affinity as glucose for either glucose transporters or hexokinase. For quantitative analyses, these differences must be corrected for by the lumped constant, a parameter present in the operational equation used to determine the local rate of cerebral glucose utilization. While the operational equation has been thoroughly validated for its use in physiological conditions, it may not apply in specific pathophysiological conditions in which cardiovascular function or blood glucose levels are altered. FDG remains the most frequently used tracer in PET imaging, particularly for the detection and staging of tumors, in spite of the low spatial resolution of PET, the relative invasiveness of the method and the radiation dose due to the injection of [ $^{18}\text{F}$ ]-FDG. The use of fMRI-BOLD (see Box 1) now represents the gold standard to study the human brain at rest and during increased functional activity.

Even if the 2DG/FDG method cannot provide information on the fate of glucose downstream of the first hexokinase step, it is as yet the only one that provides quantitative measurement of local glucose

consumption in various structures of the brain. The recent development of computerized procedures for the automatic acquisition and 3D reconstruction of postmortem autoradiographic volume images now permits voxel-wise statistical methods to 3D-analyze the differences in glucose use in different groups of animals (without a priori identification of Region-of-Interest), as is done in humans (Figure 3b; Boussicault et al., 2014; Dubois et al., 2010).

## 6.6 | Magnetic resonance spectroscopy

Combined with MRI, MRS is a powerful technique that permits simultaneous visualization and quantification of dozens of metabolites contained in the brain area studied. The major advantage of MRS though is the possibility to obtain metabolic information in a noninvasive manner.

Nuclear magnetic resonance arises from the interaction between atomic nuclei that possess an intrinsic angular momentum, or spin, and a magnetic field. This interaction is a very weak one that confers a low sensitivity on this technique, depending on the nucleus considered, the signal being stronger for  $^1\text{H}$  and much weaker for  $^{13}\text{C}$ , for example. Most of the *in vivo* studies are thus  $^1\text{H}$ -MRS. Metabolites are analyzed in a voxel, or volume of interest, located on the MR image acquired just before MRS (Figure 3c). Accuracy is therefore strongly dependent on the immobility of the animal, which can be controlled by performing another MRI just after MRS. Anesthetic choice is also very important, especially for metabolic studies, since many drugs have an impact either on the vascular response or on metabolite contents (Horn & Klein, 2010; Tchaoussoff et al., 1991). Note that only metabolites present at a concentration higher than 1 mM may be detected by MRS.

### BOX 1 The BOLD-fMRI method: Relationship with energy metabolism

MRI is well known for its capacity to noninvasively provide high resolution images of brain structures. MRI can also be rendered sensitive to local magnetic field inhomogeneity. This principle has been applied to develop the most common modality of fMRI, where the so-called BOLD effect on magnetic field inhomogeneity is detected during brain activity. The BOLD effect relies on the local transient increase of the oxyhemoglobin-to-deoxyhemoglobin ratio during the haemodynamic response associated with neuronal activity (Ogawa et al., 1990). Since deoxyhemoglobin induces magnetic inhomogeneity, activated regions of the brain, with less deoxyhemoglobin, exhibit a stronger BOLD signal. BOLD-fMRI is either used with task-based paradigms, in which the mean of the baseline state is subtracted from the mean of the stimulated state to identify activated regions associated with the task, or it is used to study the brain at rest, in the absence of an explicit task (resting state BOLD-fMRI; Gusnard, Raichle, & Raichle, 2001). BOLD contrast fMRI has come to dominate the functional neuro-imaging field.

The BOLD-fMRI signal is not a straightforward measure of energy metabolism. The deoxyhemoglobin content within a voxel is determined by the balance between the supply of oxygenated blood, the rate of oxygen consumption and the fractional blood volume. This is because the supply of oxygen is not matched precisely with the demand that the BOLD-fMRI signals change in regions where the actual amount of oxygen remaining in blood vessels is modified during activation. Hence, the BOLD response is dependent upon hemodynamic (blood flow and volume) and metabolic (oxygen consumption) responses. Although the value of BOLD fMRI for functional brain mapping is undisputed, the more subtle questions of how to interpret and use BOLD response magnitude and changes to derive information about energy metabolism are still unresolved. A reduction in the BOLD signal can result from a reduced neuronal activation, abnormal neuron/astrocyte coupling, a reduced vascular response, or a combination of all these components of the neurovascular response.

Quantitative fMRI techniques have been developed to measure the hemodynamic and metabolic responses to modulations in brain activity (Pike, 2012). They include arterial spin labeling to measure variations in tissue perfusion in each pixel (which can help to disentangle confounding mechanisms), contrast agents to measure cerebral blood volume and calibrated fMRI using biophysical models of the BOLD signal to estimate changes in the cerebral metabolic rate of oxygen consumption ( $\text{CMRO}_2$ ). Calibrated fMRI may be used as a biomarker of brain disorders that could be applied for therapy monitoring.

Moreover, spatial and temporal resolutions are rather low, for voxels are in the  $\text{mm}^3$  range, thus preventing discrimination at the cellular level, and, to get enough signal-to-noise ratio, acquisitions are performed over minutes rather than seconds. This temporal resolution is even lower with  $^{13}\text{C}$ -MRS, which has a detection Level 4 orders of magnitude lower than that of  $^1\text{H}$ . Therefore, the use of  $^{13}\text{C}$ -labeled precursor is a prerequisite to follow brain metabolism.  $^{13}\text{C}$ -labeled metabolites are infused in the blood circulation of the animal, at a rather high concentration, which leads to hyperglycemia with the main  $^{13}\text{C}$ -labeled precursor used, that is  $^{13}\text{C}$ -glucose, a point that is often neglected but has an impact when studying brain metabolism. Moreover, this labeled substrate, infused in the blood circulation, is not only metabolized by the brain but also by other organs, such as the liver for example, which may scramble the position of the  $^{13}\text{C}$  in some metabolites that go back to the blood and then to the brain, and thus complicates the analyses at the cerebral level. However,  $^{13}\text{C}$ -MRS is a powerful technique since the fate of the  $^{13}\text{C}$  can be followed and the exact labeled carbon positions of each detected metabolite can be determined and quantified, allowing the pathway taken by the  $^{13}\text{C}$ -labeled precursor to be traced back (Rodrigues, Valette, & Bouzier-Sore, 2013; Rothman et al., 1999). Even if this technique does not reach cellular resolution, it can distinguish between a predominant neuronal or astrocytic metabolism. This is based on the principle that neurons and astrocytes differentially express some enzymes; for example, glutamine synthase and pyruvate carboxylase.

Several approaches have been developed to overcome the low sensitivity of MRS. First, since the signal is directly proportional to the field strength, companies develop magnets with increasing magnetic fields. Secondly, the  $^{13}\text{C}$  weak signal can be detected by  $^1\text{H}$ -MRS using proton-observed carbon-edited sequences. Finally, improved sensitivity for  $^{13}\text{C}$  can be attained with polarization transfer experiments, which partly transfer energy from electron to nucleus, increasing the signal by 10,000 times, but only for some tens of seconds. Even if MRS is not a very sensitive technique, this relative weakness is also its strength, enabling NMR to study living organisms without significantly disturbing them and thus yielding detailed metabolic information noninvasively, which allows longitudinal studies (de Graaf, 2013; Stagg & Rothman, 2014).

In parallel to *in vivo* MRS, *ex vivo* NMR spectroscopy is also often used to explore details of brain metabolism (Figure 3d). One of the biggest advantages of *ex vivo* NMR spectroscopy is achieving very high-resolution spectra and metabolic profiles. Spectra can be obtained either on perchloric acid extracts of cell cultures or tissues. Alternatively, with the use of special NMR probes such as HRMAS (high resolution magic angle spinning) it is now possible to achieve the same high spectral resolution on biopsies (without any biochemical treatment) as at obtained in liquid state NMR (extracts). The very recent development of the micro HRMAS probe (HR $\mu$ MAS) allows for high quality spectra on nanoliter-sized samples (Duong et al., 2016). An important issue when performing *ex vivo* analyses is to completely stop metabolism before acquiring the spectra. For cell cultures, the cellular layer is quickly rinsed with cold phosphate buffer sulfate (to avoid any contamination with medium metabolites)

and cells are further either frozen with liquid nitrogen or directly extracted with perchloric acid. For brain biopsies, the only technique to avoid any post-mortem metabolism is to use focused microwaves. In less than one second, brain temperature reaches 70–80°C, a temperature at which proteins are denatured and thus all enzymatic activities stop. The brain can be then taken out of the skull and a precise brain region can be either directly analyzed by HRMAS NMR spectroscopy or further processed with perchloric acid extraction of the metabolites for later liquid state NMR analyses.

The technical considerations described above are valid for all observed nuclei ( $^1\text{H}$ ,  $^{13}\text{C}$ ,  $^{31}\text{P}$ , etc). For  $^{13}\text{C}$ , the use of  $^{13}\text{C}$ -labeled precursors is also needed (see *in vivo* spectroscopy paragraph). However, natural abundance can also be observed with *ex vivo* NMR spectroscopic studies, compared to *in vivo* studies, since long time acquisition (several hours) can be performed. Finally, the high spectral resolution of *ex vivo* NMR spectroscopy permits analysis of what is called homonuclear spin coupling. When several  $^{13}\text{C}$  are located on the same molecule, the resonance of the observed carbon is split and multiplets will appear. The analysis of these homonuclear spin-coupling patterns represents a unique advantage over other metabolic monitoring techniques such as radioactivity (Rodrigues et al., 2013).

## 6.7 | Metabolomics, mass spectrometry and stable isotope tracing

### 6.7.1 | Metabolomics

Cataloguing the complement of metabolites in a biological sample can lend important insight, particularly in the search for disease biomarkers. Mass spectrometry and proton NMR are generally the methods of choice for metabolomics (Dumas & Davidovic, 2015). Proton NMR, a faster technique that can have a simpler sample preparation than mass spectrometry, measures metabolites in the low micromolar range, and has been used extensively in the characterization of CNS disorders (see Section 6.6 and Oz et al., 2014), whereas mass spectrometry can detect picomolar and even femtomolar quantities of metabolites with time-of-flight and quadrupole time-of-flight detection systems. However, mass spectrometry typically involves a more complex sample extraction, derivatization of the metabolites to facilitate detection, and separation techniques before injection of a sample into the mass spectrometer. In general, methanol:chloroform extraction allows separation of polar and nonpolar metabolites, and gas chromatography is used to further separate polar compounds, whereas liquid chromatography is used to separate nonpolar metabolites. Following separation, the samples are ionized, and are separated and/or further fragmented in the mass spectrometer that detects the species based upon their mass/charge ( $m/z$ ) ratio. Each metabolite generates its own pattern of peaks in the  $m/z$  spectrum, and the peak heights generally correlate with metabolite concentration. Huge quantities of data are generated by mass spectrometry, thus sophisticated systems biology data analysis tools are used to interpret the spectra, to detect differences between samples, and to define changes in patterns of related metabolites and metabolic pathways. Such approaches have been used to identify changes in the metabolome in a wide variety of



TABLE 1 Main strengths and limitations of some techniques of frequent application to the study of brain energy metabolism

	Technique	Strengths	Limitations
The Stage: Location	Immunohistochemistry	Tissue context Accessible Low cost	Reduced antibody availability Epitope-masking Mostly qualitative
	Immunoblotting	Semiquantitative Accessible	Reduced antibody availability Lacks cellular resolution
	Protein-Protein interaction assays (Immunoprecipitation, Two-Hybrid, FRET, etc)	Specificity Spatial resolution	High rate of false negatives High rate of false positives May miss indirect interactions in supramolecular complexes
The players: Functional properties	Enzyme activity	Precise functional characterization	Non-physiological physicochemical context
	Transporter activity Electrophysiology	Optimum manipulation of experimental parameters	Possible loss of cofactors and relevant interactions
	Optogenetic actuators	Cell type-specific manipulation	Non-physiological ion perturbations
The play: Concentrations and fluxes	Respirometry	High temporal resolution	Unavailable for adherent cells
	Clarke-type electrode		Requires cell populations
	Respirometry Multi-well plate	High throughput	Low temporal resolution Requires cell populations No HCO <sub>3</sub> <sup>-</sup> present
	Microdialysis	In vivo Available for many metabolites	Low spatial and temporal resolution Local tissue disruption
	Enzyme-based probes	In vivo	Low spatial resolution Local tissue disruption Available for few metabolites
	Fluorescent glucose analogs	Single cell resolution In vitro and in vivo Non-invasive	Variable kinetic properties
	Genetically-encoded sensors	High specificity Single cell resolution In vitro and in vivo Non-invasive	Calibration can be difficult Requires high expression levels Restricted to a small set of metabolites
	Mitochondrial autofluorescence	High temporal resolution	Small signal to noise Low spatial resolution Non-trivial interpretation
	Isotope-labelled glucose	Quantitative estimation of regional flux Human studies	Low temporal resolution Lacks cellular resolution
	Magnetic Resonance Spectroscopy	Multiple metabolites Non-invasive Human studies	Only abundant metabolites (<1mM) Low temporal resolution Hyperglycemia (13C-glucose)
Metabolomics	Multiple metabolites	Sample destruction Low throughput Intensive data analysis High cost	
Mathematical modeling	Tackles complexity Low cost	Key parameters are often unavailable Hidden assumptions	
BOLD-fMRI	Human studies	Unclear relationship with energy metabolism	

(BOLD = blood oxygenation level dependent; fMRI = functional magnetic resonance imaging; FRET = Förster resonance energy transfer).

neurodegenerative diseases in humans and in animal models (Dumas & Davidovic, 2015).

A particular challenge with metabolomics is to determine the meaning and significance of the multitude of findings with respect to cell and tissue health or to disease etiology. Relatively low throughput and intensive data analysis, as well as the high cost of state-of-the-art LC- and gas chromatography-mass spectrometers present additional challenges. In metabolomics, spatial and subcompartmental information is lost except for the regional information provided by NMR-based spectroscopy. However, mass spectroscopy imaging is beginning to address cellular, and even subcellular resolution of metabolites using fluorinated nanoparticles in nanostructure-initiator mass spectroscopy and related approaches (Kurczy et al., 2015). If a metabolite concentration increases, it is unclear whether the change arises from enhanced production, or decreased consumption of the intermediate, as metabolomics is generally based on data gathered from a single snapshot-in-time. Therefore, whether or not there are alterations in the flux of a metabolite through a metabolic pathway cannot readily be discerned. Computational flux balance can yield estimates of metabolic flux without labeling, but the approach is entirely reliant on existing knowledge of enzyme kinetics and pathways and thus has limitations (Aurich & Thiele, 2016).

### 6.7.2 | Stable isotope tracing

By labeling and following carbon through metabolic pathways, stable isotope tracing can address questions regarding the extent and pattern of flux through metabolic pathways (Buescher et al., 2015). Stable isotope tracers are nonradioactive and naturally occurring, and are therefore safe. The technique is typically destructive to the sample to enable mass spectrometry, but can be performed on cultured cells, tissues, and *in vivo*. NMR can instead be used for detection, with a significant sacrifice in sensitivity (Rodrigues et al., 2013).  $^{13}\text{C}$ -containing molecules like glucose and glutamine have been used extensively, as the  $^{13}\text{C}$  in the parent molecule is transferred to downstream metabolites that can be detected by mass spectrometry (Figure 3d). The extent to which metabolites become enriched with  $^{13}\text{C}$  combined with an analysis of the concentration of the metabolite is an indication of the flux through the metabolic pathway. The same, in principle, is true for  $^{15}\text{N}$ - and  $^3\text{H}$ -labeled substrates. For isotope tracing with cells in culture, a full complement of growth media is typically added with replacement of a fuel like glucose with  $^{13}\text{C}$ -labeled glucose. The system is incubated for a sufficient time to allow the sample to come to isotopic steady state, and then the media is removed and the sample is extracted and derivatized for analysis by mass spectrometry. The spectral peaks for each metabolite are then used to quantify its abundance as well as the mass isotopomer distribution (the fraction of a metabolite enriched with zero, 1, 2, 3, etc  $^{13}\text{C}$ -labeled carbons). From these data, as well as separate experiments using other metabolic tracers, the extent of carbon flux from different substrates into the pool of a particular metabolite and its precursors can be calculated (Kelleher & Nickol, 2015). The ability to resolve cell-type specific metabolism in brain tissue or co-cultures is somewhat limited, and is reliant on either pharmacologic agents or genetic manipulation of isoforms of metabolite transporters

or enzymes to control metabolism in particular cell types. Stable isotope tracing can determine the *relative* rates of flux of one labeled substrate versus another. Quantitative rates of flux can be determined by an extension of the technique termed *metabolic flux analysis* whereby the  $^{13}\text{C}$ -labeling pattern of intracellular metabolites is pinned to the rates of uptake and release of metabolites from a biological system. The data is applied to a computational network model of biochemical reactions that enables quantitative estimates of flux through specific pathways (Buescher et al., 2015). The approach is more labor- and data-intensive than stable isotope tracing, and is reliant on a robust computational model that accounts for all potential reaction pathways.

Finally, in a particularly powerful approach, stable isotope tracing with mass spectrometry can be coupled with real-time measurements of metabolic flux using respirometry (a technique described in detail in Section 6.1). The depth of insight provided by the combination of the two approaches is greater than either technique alone, and has been applied to the study of cultured cortical neurons (Divakaruni et al., 2017), the mouse retina (Du et al., 2016), and glioma metabolism and growth (Yang et al., 2014a).

## 6.8 | Mathematical modeling

Mathematical modeling aims to integrate the outcome of multiple experimental techniques in order to grasp complex behaviors that are otherwise impenetrable. It serves to identify variables and parameters that are critical and should therefore be given experimental attention, to then guide the design of experiments by a priori, inexpensive visualization of possible outcomes. At its best, it may prefigure novel phenomena. In the specific case of brain energy metabolism, models combine structural parameters of brain tissue with the microscopic properties of enzymes and transporters in view of dynamic readouts obtained by technologies like PET, MRS, fMRI, tissue electrodes, microdialysis, and genetically-encoded nanosensors. A key well-substantiated assumption of current models is that metabolism is compartmentalized, that is, that it occurs in a few well-mixed reservoirs (say neuronal and glial). Thus, chemical and transport reactions can be represented by a small set of ordinary differential equations that are solved by numerical simulation using standard computers. The outcome of a model is as good as what the model is fed, and so different groups have arrived at opposite conclusions: witness the astrocyte-to-neuron lactate shuttle controversy (Aubert & Costalat, 2007; Hyder et al., 2006; Jolivet, Coggan, Allaman, & Magistretti, 2015; Mangia, Simpson, Vannucci, & Carruthers, 2009;). While debate speaks of a lively discipline, assumptions about key parameters are often hidden, and so it has been hard for the noninitiated to judge the relative merits of competing models (Jolivet, Allaman, Pellerin, Magistretti, & Weber, 2010). In view of structural, functional, and likely metabolic heterogeneity of neurons and glial cells (DeFelipe et al., 2013; Freeman, 2015), future models will have to become more complex, as has recently occurred for neuronal ATP subcellular domains (Le Masson, Przedborski, & Abbott, 2014). The next generation of models will also have to deal with growing evidence for fast local regulation of enzymes, transporters and channels in response to neural activity (Barros et al., 2013). However, it is important to





understand that a more complex model (i.e., which include more equations and more parameters) is not necessarily better to adequately describe the different possible outcomes of a system. It all depends on the question(s) that need(s) to be addressed.

## CONCLUDING REMARKS

The diversity, range, and depth of the techniques applied to the study of brain energy metabolism are amazing, as is the complexity of the problem in hand. As we are now gaining access to the particular metabolic behavior of specific brain areas, cell types and even individual cells, techniques are becoming more and more sophisticated, each own having its own domain, strengths and limitations (Table 1). A challenge for the near future is to increase the crosstalk between technical viewpoints, perhaps through the active exchange of ideas and collaboration between dedicated laboratories.

## ACKNOWLEDGMENT

We thank Karen Everett for critical reading of the manuscript and Ray Swanson for helpful criticism. We would like to thank Anne-Sophie Herard, Dr. Joshua Jackson, and Dr. Ulrike Winkler for providing data and for helping to prepare Figure 3. LFB is partly funded by Fondecyt Grant 1160317. The Centro de Estudios Científicos (CECs) is supported by the Chilean Government through the Centers of Excellence Basal Financing Program of CONICYT. JPB is funded by MINECO (SAF2016-78114-R; RTC-2015-3237-1), CIBERFES (CB16/10/00282), E.U. SP3-People-MC-ITN programme (608381), EU BATCure grant (666918) and FEDER (European regional development fund). GB is affiliated to NeurATRIS: A Translational Research Infrastructure for Biotherapies in Neurosciences ("Investissement d'Avenir", ANR-11-INBS-0011). AKBS has received financial support from the French State in the context of the "Investments for the future" Programme IdEx and the LabEx TRAIL, reference ANR-10-IDEX and ANR-10-LABX-57, and a French-Swiss ANR-FNS grant reference ANR-15-CE37-0012. AMB was funded by the University of Nottingham. SK is funded by BBSRC: BB/L019396/1, BB/K009192/1 and MRC MR/L020661/1. FK was supported by grants from Deutsche Forschungsgemeinschaft DFG SPP 1757, SFB 894 and FOR 2289; European Commission H2020-FETPROACT 732344 Neurofibres and H2020-MSCA-ITN-722053 EU-GliaPhD. ANM is supported by US National Institutes of Health grant R01NS087611, Teva Pharmaceuticals, and Agilent Technologies. LP received an Excellence Chair from the Programme IdEx, reference ANR-10-IDEX-03-02 and obtained a French-Swiss ANR-FNS grant number 310030E-164271. MBR was supported by the National Institutes of Neurologic Disease and Stroke at the National Institutes of Health (R01 NS077773). BW is supported by the University of Zurich and the Swiss National Science Foundation and is a member of the Clinical Research Priority Program of the University of Zurich on Molecular Imaging.

## ORCID

L. Felipe Barros <http://orcid.org/0000-0002-6623-4833>

Sergey Kasparov <http://orcid.org/0000-0002-1824-1764>

Michael B. Robinson <http://orcid.org/0000-0003-0162-2218>

Johannes Hirrlinger <http://orcid.org/0000-0002-6327-0089>

## REFERENCES

- Adan, A., Kiraz, Y., & Baran, Y. (2016). Cell proliferation and cytotoxicity assays. *Current Pharmaceutical Biotechnology*, 17, 1213–1221.
- Akerboom, J., Chen, T. W., Wardill, T. J., Tian, L., Marvin, J. S., Mutlu, S., ... Looger, L. L. (2012). Optimization of a GCaMP calcium indicator for neural activity imaging. *Journal of Neuroscience*, 32, 13819–13840.
- Almeida, A., Moncada, S., & Bolanos, J. P. (2004). Nitric oxide switches on glycolysis through the AMP protein kinase and 6-phosphofructo-2-kinase pathway. *Nature Cell Biology*, 6, 45–51.
- Aubert, A., & Costalat, R. (2007). Compartmentalization of brain energy metabolism between glia and neurons: insights from mathematical modeling. *Glia*, 55, 1272–1279.
- Auld, D. S., & Inglese, J. (2004). Interferences with Luciferase Reporter Enzymes. <https://www.ncbi.nlm.nih.gov/books/NBK374281/>
- Aurich, M. K., & Thiele, I. (2016). Computational modeling of human metabolism and its application to systems biomedicine. *Methods in Molecular Biology*, 1386, 253–281.
- Barres, B. A., Koroshetz, W. J., Chun, L. L., & Corey, D. P. (1990). Ion channel expression by white matter glia: the type-1 astrocyte. *Neuron*, 5, 527–544.
- Barros, L. F., Baeza-Lehnert, F., Valdebenito, R., Ceballos, S., & Alegría, K. (2014). Fluorescent nanosensor based flux analysis: overview and the example of glucose. In: H. S. Waagepetersen, J. Hirrlinger, (Eds.). *Springer Protocols: Brain Energy Metabolism*. Berlin: Springer.
- Barros, L. F., Bittner, C. X., Loaiza, A., & Porras, O. H. (2007). A quantitative overview of glucose dynamics in the gliovascular unit. *Glia*, 55, 1222–1237.
- Barros, L. F., Bittner, C. X., Loaiza, A., Ruminot, I., Larenas, V., Moldenhauer, H., ... Alvarez, M. (2009a). Kinetic validation of 6-NBDG as a probe for the glucose transporter GLUT1 in astrocytes. *Journal of Neurochemistry*, 109(Suppl 1), 94–100.
- Barros, L. F., Courjaret, R., Jakoby, P., Loaiza, A., Lohr, C., & Deitmer, J. W. (2009b). Preferential transport and metabolism of glucose in Bergmann glia over Purkinje cells: a multiphoton study of cerebellar slices. *Glia*, 57, 962–970.
- Barros, L. F., San Martin, A., Sotelo-Hitschfeld, T., Lerchundi, R., Fernández-Moncada, I., Ruminot, I., ... Espinoza, D. (2013). Small is fast: Astrocytic glucose and lactate metabolism at cellular resolution. *Frontiers in Cell Neuroscience*, 7, 27.
- Barros, L. F., San, M. A., Ruminot, I., Sandoval, P. Y., Fernandez-Moncada, I., Baeza-Lehnert, F., ... Alegria, K. (2017). Near-critical GLUT1 and neurodegeneration. *Journal of Neuroscience Research*, 95, 2267–2274.
- Bauer, D. E., Jackson, J. G., Genda, E. N., Montoya, M. M., Yudkoff, M., & Robinson, M. B. (2012). The glutamate transporter, GLAST, participates in a macromolecular complex that supports glutamate metabolism. *Neurochemistry International*, 61, 566–574.
- Beppu, K., Sasaki, T., Tanaka, K. F., Yamanaka, A., Fukazawa, Y., Shigemoto, R., & Matsui, K. (2014). Optogenetic countering of glial acidosis suppresses glial glutamate release and ischemic brain damage. *Neuron*, 81, 314–320.
- Bergersen, L. H., Magistretti, P. J., & Pellerin, L. (2005). Selective postsynaptic co-localization of MCT2 with AMPA receptor GluR2/3

- subunits at excitatory synapses exhibiting AMPA receptor trafficking. *Cerebral Cortex*, 15, 361–370.
- Bi, A., Cui, J., Ma, Y. P., Olshevskaya, E., Pu, M., Dizhoor, A. M., & Pan, Z. H. (2006). Ectopic expression of a microbial-type rhodopsin restores visual responses in mice with photoreceptor degeneration. *Neuron*, 50, 23–33.
- Bittner, C. X., Loaiza, A., Ruminot, I., Larenas, V., Sotelo-Hitschfeld, T., Gutiérrez, R., ... Barros, L. F. (2010). High resolution measurement of the glycolytic rate. *Frontiers in Neuroenergetics*, 2, 1–11. <https://doi.org/10.3389/fnene.2010.00026>.
- Boekema, E. J., & Braun, H. P. (2007). Supramolecular structure of the mitochondrial oxidative phosphorylation system. *Journal of Biological Chemistry*, 282, 1–4.
- Bolanos, J. P., Heales, S. J., Land, J. M., & Clark, J. B. (1995). Effect of peroxynitrite on the mitochondrial respiratory chain: differential susceptibility of neurones and astrocytes in primary culture. *Journal of Neurochemistry*, 64, 1965–1972.
- Bolanos, J. P., Peuchen, S., Heales, S. J., Land, J. M., & Clark, J. B. (1994). Nitric oxide-mediated inhibition of the mitochondrial respiratory chain in cultured astrocytes. *Journal of Neurochemistry*, 63, 910–916.
- Boussicault, L., Herard, A. S., Calingasan, N., Petit, F., Malgorn, C., Merienne, N., ... Bonvento, G. (2014). Impaired brain energy metabolism in the BACHD mouse model of Huntington's disease: critical role of astrocyte-neuron interactions. *Journal of Cerebral Blood Flow and Metabolism* 34, 1500–1510.
- Bouzier-Sore, A. K., & Bolanos, J. P. (2015). Uncertainties in pentose-phosphate pathway flux assessment underestimate its contribution to neuronal glucose consumption: Relevance for neurodegeneration and aging. *Frontiers in Aging Neuroscience*, 7, 89.
- Boyden, E. S., Zhang, F., Bamberg, E., Nagel, G., & Deisseroth, K. (2005). Millisecond-timescale, genetically targeted optical control of neural activity. *Nature Neuroscience*, 8, 1263–1268.
- Brooks, H. B., Geeganage, S., Kahl, S. D., Montrose, C., Sittampalam, S., Smith, M. C., & Weidner, J. R. (2004). Basics of Enzymatic Assays for HTS. <https://www.ncbi.nlm.nih.gov/books/NBK92007/>
- Brown, A. M., Evans, R. D., Black, J., & Ransom, B. R. (2012). Schwann cell glycogen selectively supports myelinated axon function. *Annals Neurology*, 72, 406–418.
- Brown, A. M., Skamarasuskas, J., Lister, T., Madjd, A., & Ray, D. E. (2011). Differential susceptibility of astrocytic and neuronal function to 3-chloropropanediol in the rat inferior colliculus. *Journal of Neurochemistry*, 116, 996–1004.
- Brown, A. M., Tekkok, S. B., & Ransom, B. R. (2003). Glycogen regulation and functional role in mouse white matter. *Journal of Physiology*, 549, 501–512.
- Buescher, J. M., Antoniewicz, M. R., Boros, L. G., Burgess, S. C., Brunen-graber, H., Clish, C. B., ... Fendt, S. M. (2015). A roadmap for interpreting (13)C metabolite labeling patterns from cells. *Current Opinion in Biotechnology*, 34, 189–201.
- Bulusu, V., Prior, N., Snaebjornsson, M. T., Kuehne, A., Sonnen, K. F., Kress, J., ... Aulehla, A. (2017). Spatiotemporal analysis of a glycolytic activity gradient linked to mouse embryo mesoderm development. *Developmental Cell*, 40, 331–341.
- Bushong, E. A., Martone, M. E., Jones, Y. Z., & Ellisman, M. H. (2002). Protoplasmic astrocytes in CA1 stratum radiatum occupy separate anatomical domains. *Journal of Neuroscience*, 22, 183–192.
- Cahoy, J. D., Emery, B., Kaushal, A., Foo, L. C., Zamanian, J. L., Christopherson, K. S., ... Barres, B. A. (2008). A transcriptome database for astrocytes, neurons, and oligodendrocytes: A new resource for understanding brain development and function. *Journal of Neuroscience*, 28, 264–278.
- Cameron, W. D., Bui, C. V., Hutchinson, A., Loppnau, P., Graslund, S., & Rocheleau, J. V. (2016). Apollo-NADP(+): A spectrally tunable family of genetically encoded sensors for NADP(+). *Nature Methods*, 13, 352–358.
- Cater, H. L., Benham, C. D., & Sundstrom, L. E. (2001). Neuroprotective role of monocarboxylate transport during glucose deprivation in slice cultures of rat hippocampus. *Journal of Physiology*, 531, 459–466.
- Chance, B., Cohen, P., Jobsis, F., & Schoener, B. (1962). Intracellular oxidation-reduction states *in vivo*. *Science*, 137, 499–508.
- Chaudhry, F. A., Lehre, K. P., van Lookeren, C. M., Ottersen, O. P., Danbolt, N. C., & Storm-Mathisen, J. (1995). Glutamate transporters in glial plasma membranes: highly differentiated localizations revealed by quantitative ultrastructural immunocytochemistry. *Neuron*, 15, 711–720.
- Choeiri, C., Staines, W., & Messier, C. (2002). Immunohistochemical localization and quantification of glucose transporters in the mouse brain. *Neuroscience*, 111, 19–34.
- Choi, S. W., Gerencser, A. A., & Nicholls, D. G. (2009). Bioenergetic analysis of isolated cerebrocortical nerve terminals on a microgram scale: Spare respiratory capacity and stochastic mitochondrial failure. *Journal of Neurochemistry*, 109, 1179–1191.
- Chuquet, J., Quilichini, P., Nimchinsky, E. A., & Buzsaki, G. (2010). Pre-dominant enhancement of glucose uptake in astrocytes versus neurons during activation of the somatosensory cortex. *Journal of Neuroscience*, 30, 15298–15303.
- Clark, L. C., Jr., Wolf, R., Granger, D., & Taylor, Z. (1953). Continuous recording of blood oxygen tensions by polarography. *Journal of Applied Physiology*, 6, 189–193.
- Cong, L., Ran, F. A., Cox, D., Lin, S., Barretto, R., Habib, N., ... Zhang, F. (2013). Multiplex genome engineering using CRISPR/Cas systems. *Science*, 339, 819–823.
- Crone, C. (1965). Facilitated transfer of glucose from blood into brain tissue. *Journal of Physiology*, 181, 103–113.
- Dahlin, J. L., Baell, J., & Walters, M. A. (2004). Assay Interference by Chemical Reactivity. <https://www.ncbi.nlm.nih.gov/books/NBK326709/>
- de Graaf, R. A. (2013). *In Vivo NMR Spectroscopy: Principles and Techniques*. Chichester: John Wiley & Sons.
- DeFelipe, J., Lopez-Cruz, P. L., Benavides-Piccione, R., Bielza, C., Larranaga, P., Anderson, S., ... Ascoli, G. A. (2013). New insights into the classification and nomenclature of cortical GABAergic interneurons. *Nature Reviews Neuroscience*, 14, 202–216.
- Deitmer, J. W. (1992). Evidence for glial control of extracellular pH in the leech central nervous system. *Glia*, 5, 43–47.
- Denk, W., Strickler, J. H., & Webb, W. W. (1990). Two-photon laser scanning fluorescence microscopy. *Science*, 248, 73–76.
- Divakaruni, A. S., Rogers, G. W., & Murphy, A. N. (2014). Measuring mitochondrial function in permeabilized cells using the Seahorse XF analyzer or a Clark-type oxygen electrode. *Current Protocols in Toxicology*, 60, 25–16.
- Divakaruni, A. S., Wallace, M., Buren, C., Martyniuk, K., Andreyev, A. Y., Li, E., ... Murphy, A. N. (2017). Inhibition of the mitochondrial pyruvate carrier protects from excitotoxic neuronal death. *Journal of Cell Biology*, 216, 1091–1105.
- Du, J., Rountree, A., Cleghorn, W. M., Contreras, L., Lindsay, K. J., Sadi- lek, M., ... Hurley, J. B. (2016). Phototransduction influences metabolic flux and nucleotide metabolism in mouse retina. *Journal of Biological Chemistry*, 291, 4698–4710.
- Dubois, A., Herard, A. S., Delatour, B., Hantraye, P., Bonvento, G., Dhennain, M., & Delzescaux, T. (2010). Detection by voxel-wise statistical



- analysis of significant changes in regional cerebral glucose uptake in an APP/PS1 transgenic mouse model of Alzheimer's disease. *Neuroimage*, 51, 586–598.
- Dumas, M. E., & Davidovic, L. (2015). Metabolic profiling and phenotyping of central nervous system diseases: metabolites bring insights into brain dysfunctions. *Journal of Neuroimmune Pharmacology*, 10, 402–424.
- Duong, N. T., Endo, Y., Nemoto, T., Kato, H., Bouzier-Sore, A. K., Nishiyama, Y., & Wong, A. (2016). Evaluation of a high-resolution micro-sized magic angle spinning (HRIMAS) probe for NMR-based metabolomic studies of nanoliter samples. *Analytical Methods*, 216, 6815–6820.
- Esipova, T. V., Rivera-Jacquez, H. J., Weber, B., Masunov, A. E., & Vinogradov, S. A. (2016). Two-photon absorbing phosphorescent metalloporphyrins: Effects of pi-extension and peripheral substitution. *Journal of American Chemical Society*, 138, 15648–15662.
- Esipova, T. V., & Vinogradov, S. A. (2014). Synthesis of phosphorescent asymmetrically pi-extended porphyrins for two-photon applications. *Journal of Organic Chemistry*, 79, 8812–8825.
- Fehr, M., Lalonde, S., Lager, I., Wolff, M. W., & Frommer, W. B. (2003). *In vivo* imaging of the dynamics of glucose uptake in the cytosol of COS-7 cells by fluorescent nanosensors. *Journal of Biological Chemistry*, 278, 19127–19133.
- Figueiredo, M., Lane, S., Stout, R. F., Jr., Liu, B., Parpura, V., Teschemacher, A. G., & Kasparov, S. (2014). Comparative analysis of optogenetic actuators in cultured astrocytes. *Cell Calcium*, 56, 208–214.
- Focking, M., Dicker, P., Lopez, L. M., Hryniewiecka, M., Wynne, K., English, J. A., ... Cotter, D. R. (2016). Proteomic analysis of the postsynaptic density implicates synaptic function and energy pathways in bipolar disorder. *Translational Psychiatry*, 6, e959.
- Freeman, M. R. (2015). Drosophila central nervous system glia. *Cold Spring Harbor Perspectives Biology*, 7, a020552.
- Frenguelli, B. G., Llaudet, E., & Dale, N. (2003). High-resolution real-time recording with microelectrode biosensors reveals novel aspects of adenosine release during hypoxia in rat hippocampal slices. *Journal of Neurochemistry*, 86, 1506–1515.
- Frenguelli, B. G., Wigmore, G., Llaudet, E., & Dale, N. (2007). Temporal and mechanistic dissociation of ATP and adenosine release during ischaemia in the mammalian hippocampus. *Journal of Neurochemistry*, 101, 1400–1413.
- Garavito, R. M., & Ferguson-Miller, S. (2001). Detergents as tools in membrane biochemistry. *Journal of Biological Chemistry*, 276, 32403–32406.
- Genda, E. N., Jackson, J. G., Sheldon, A. L., Locke, S. F., Greco, T. M., O'donnell, J. C., ... Robinson, M. B. (2011). Co-compartmentalization of the astroglial glutamate transporter, GLT-1, with glycolytic enzymes and mitochondria. *Journal of Neuroscience*, 31, 18275–18288.
- Glickman, J. F. (2004). Assay Development for Protein Kinase Enzymes. <https://www.ncbi.nlm.nih.gov/books/NBK91991/>
- Gourine, A. V., Kasymov, V., Marina, N., Tang, F., Figueiredo, M. F., Lane, S., ... Kasparov, S. (2010). Astrocytes control breathing through pH-dependent release of ATP. *Science*, 329, 571–575.
- Gruetter, R., Ugurbil, K., & Seaquist, E. R. (1998). Steady-state cerebral glucose concentrations and transport in the human brain. *Journal of Neurochemistry*, 70, 397–408.
- Gusnard, D. A., Raichle, M. E., & Raichle, M. E. (2001). Searching for a baseline: Functional imaging and the resting human brain. *Nature Reviews in Neuroscience*, 2, 685–694.
- Hall, C. N., Klein-Flugge, M. C., Howarth, C., & Attwell, D. (2012). Oxidative phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information processing. *Journal of Neuroscience*, 32, 8940–8951.
- Harper, S., & Speicher, D. W. (2011). Purification of proteins fused to glutathione S-transferase. *Methods in Molecular Biology*, 681, 259–280.
- Harris, J. J., Jolivet, R., & Attwell, D. (2012). Synaptic energy use and supply. *Neuron*, 75, 762–777.
- Hasel, P., Dando, O., Jiwaji, Z., Baxter, P., Todd, A. C., Heron, S., ... Hardingham, G. E. (2017). Neurons and neuronal activity control gene expression in astrocytes to regulate their development and metabolism. *Nature Communications*, 8, 15132.
- Herrero-Mendez, A., Almeida, A., Fernandez, E., Maestre, C., Moncada, S., & Bolanos, J. P. (2009). The bioenergetic and antioxidant status of neurons is controlled by continuous degradation of a key glycolytic enzyme by APC/C-Cdh1. *Nature Cell Biology*, 11, 747–752.
- Hevner, R. F., & Wong-Riley, M. T. (1989). Brain cytochrome oxidase: Purification, antibody production, and immunohistochemical/histochemical correlations in the CNS. *Journal of Neuroscience*, 9, 3884–3898.
- Hirrlinger, J., Scheller, A., Hirrlinger, P. G., Kellert, B., Tang, W., Wehr, M. C., ... Kirchhoff, F. (2009). Split-cre complementation indicates coincident activity of different genes *in vivo*. *PLoS One*, 4, e4286.
- Horn, T., & Klein, J. (2010). Lactate levels in the brain are elevated upon exposure to volatile anesthetics: A microdialysis study. *Neurochemistry International*, 57, 940–947.
- Hyder, F., Patel, A. B., Gjedde, A., Rothman, D. L., Behar, K. L., & Shulman, R. G. (2006). Neuronal-glial glucose oxidation and glutamatergic-GABAergic function. *Journal of Cerebral Blood Flow and Metabolism*, 26, 865–877.
- Iadecola, C., & Nedergaard, M. (2007). Glial regulation of the cerebral microvasculature. *Nature Neuroscience*, 10, 1369–1376.
- Isono, E., & Schwachheimer, C. (2010). Co-immunoprecipitation and protein blots. *Methods in Molecular Biology*, 655, 377–387.
- Ito, T., Chiba, T., Ozawa, R., Yoshida, M., Hattori, M., & Sakaki, Y. (2001). A comprehensive two-hybrid analysis to explore the yeast protein interactome. *Proceedings of the National Academy of Science USA*, 98, 4569–4574.
- Jakoby, P., Schmidt, E., Ruminot, I., Gutierrez, R., Barros, L. F., & Deitmer, J. W. (2014). Higher transport and metabolism of glucose in astrocytes compared with neurons: A multiphoton study of hippocampal and cerebellar tissue slices. *Cerebral Cortex*, 24, 222–231.
- Jares-Erijman, E. A., & Jovin, T. M. (2003). FRET imaging. *Nature Biotechnology*, 21, 1387–1395.
- Jolivet, R., Allaman, I., Pellerin, L., Magistretti, P. J., & Weber, B. (2010). Comment on recent modeling studies of astrocyte-neuron metabolic interactions. *Journal of Cerebral Blood Flow and Metabolism*, 30, 1982–1986.
- Jolivet, R., Coggan, J. S., Allaman, I., & Magistretti, P. J. (2015). Multi-time-scale modeling of activity-dependent metabolic coupling in the neuron-glia-vasculature ensemble. *PLoS Computational Biology*, 11, e1004036.
- Kaboord, B., & Perr, M. (2008). Isolation of proteins and protein complexes by immunoprecipitation. *Methods in Molecular Biology*, 424, 349–364.
- Kaplan, N., & Colowick, N. (1955). *Preparation and Assay of Enzymes*. New York: Academic Press.
- Karagiannis, A., Sylantsev, S., Hadjihambi, A., Hosford, P. S., Kasparov, S., & Gourine, A. V. (2016). Hemichannel-mediated release of lactate. *Journal of Cerebral Blood Flow and Metabolism*, 36, 1202–1211.
- Kasischke, K. A., Vishwasrao, H. D., Fisher, P. J., Zipfel, W. R., & Webb, W. W. (2004). Neural activity triggers neuronal oxidative metabolism followed by astrocytic glycolysis. *Science*, 305, 99–103.

- Kelleher, J. K., & Nickol, G. B. (2015). Isotopomer spectral analysis: Utilizing nonlinear models in isotopic flux studies. *Methods in Enzymology*, 561, 303–330.
- Kovar, J. L., Volcheck, W., Sevick-Muraca, E., Simpson, M. A., & Olive, D. M. (2009). Characterization and performance of a near-infrared 2-deoxyglucose optical imaging agent for mouse cancer models. *Analytical Biochemistry*, 384, 254–262.
- Kurczmy, M. E., Zhu, Z. J., Ivanisevic, J., Schuyler, A. M., Lalwani, K., Santidrian, A. F., ... Siuzdak, G. (2015). Comprehensive bioimaging with fluorinated nanoparticles using breathable liquids. *Nature Communications*, 6, 5998.
- Kyung, T., Lee, S., Kim, J. E., Cho, T., Park, H., Jeong, Y. M., ... Heo, W. D. (2015). Optogenetic control of endogenous Ca(2+) channels in vivo. *Nature Biotechnology*, 33, 1092–1096.
- Lalo, U., Pankratov, Y., Wichert, S. P., Rossner, M. J., North, R. A., Kirchhoff, F., & Verkhratsky, A. (2008). P2X1 and P2X5 subunits form the functional P2X receptor in mouse cortical astrocytes. *Journal of Neuroscience*, 28, 5473–5480.
- Le Maire, M., Champeil, P., & Moller, J. V. (2000). Interaction of membrane proteins and lipids with solubilizing detergents. *Biochimica et Biophysica Acta*, 1508, 86–111.
- Le Masson, G., Przedborski, S., & Abbott, L. F. (2014). A computational model of motor neuron degeneration. *Neuron*, 83, 975–988.
- Lecoq, J., Parpaleix, A., Roussakis, E., Ducros, M., Houssen, Y. G., Vinogradov, S. A., & Charpak, S. (2011). Simultaneous two-photon imaging of oxygen and blood flow in deep cerebral vessels. *Nature Medicine*, 17, 893–898.
- Lee, Y., Morrison, B. M., Li, Y., Lengacher, S., Farah, M. H., Hoffman, P. N., ... Rothstein, J. D. (2012). Oligodendroglia metabolically support axons and contribute to neurodegeneration. *Nature*, 487, 443–448.
- Lefevre, P. G. (1948). Evidence of active transfer of certain non-electrolytes across the human red cell membrane. *Journal of General Physiology*, 31, 505–527.
- Lehre, K. P., Levy, L. M., Ottersen, O. P., Storm-Mathisen, J., & Danbolt, N. C. (1995). Differential expression of two glial glutamate transporters in the rat brain: Quantitative and immunocytochemical observations. *Journal of Neuroscience*, 15, 1835–1853.
- Lerchundi, R., Fernandez-Moncada, I., Contreras-Baeza, Y., Sotelo-Hitschfeld, T., Machler, P., Wyss, M. T., ... Barros, L. F. (2015). NH4+ triggers the release of astrocytic lactate via mitochondrial pyruvate shunting. *Proceedings of the National Academy of Science USA*, 112, 11090–11095.
- Li, Q., Lau, A., Morris, T. J., Guo, L., Fordyce, C. B., & Stanley, E. F. (2004). A syntaxin 1, Galpha(o), and N-type calcium channel complex at a presynaptic nerve terminal: Analysis by quantitative immunocolocalization. *Journal of Neuroscience*, 24, 4070–4081.
- Li-Byarlay, H., Rittschof, C. C., Massey, J. H., Pittendrigh, B. R., & Robinson, G. E. (2014). Socially responsive effects of brain oxidative metabolism on aggression. *Proceedings of the National Academy of Science USA*, 111, 12533–12537.
- Loaiza, A., Porras, O. H., & Barros, L. F. (2003). Glutamate triggers rapid glucose transport stimulation in astrocytes as evidenced by real-time confocal microscopy. *Journal of Neuroscience*, 23, 7337–7342.
- Lopez-Fabuel, I., Le, D. J., Logan, A., James, A. M., Bonvento, G., Murphy, M. P., ... Bolanos, J. P. (2016). Complex I assembly into supercomplexes determines differential mitochondrial ROS production in neurons and astrocytes. *Proceedings of the National Academy of Science USA*, 113, 13063–13068.
- Lorsch, J. R. (2015). Preface. Laboratory methods in enzymology: Protein part D. *Methods in Enzymology*, 559, xi.
- Lundgaard, I., Li, B., Xie, L., Kang, H., Sanggaard, S., Haswell, J. D., ... Nedergaard, M. (2015). Direct neuronal glucose uptake heralds activity-dependent increases in cerebral metabolism. *Nature Communications*, 6, 6807.
- Lyons, D. G., Parpaleix, A., Roche, M., & Charpak, S. (2016). Mapping oxygen concentration in the awake mouse brain. *Elife*, 5, e12024.
- Machler, P., Wyss, M. T., Elsayed, M., Stobart, J., Gutierrez, R., von Faber-Castell, A., ... Weber, B. (2016). In vivo evidence for a lactate gradient from astrocytes to neurons. *Cell Metabolism*, 23, 94–102.
- Madisen, L., Garner, A. R., Shimaoka, D., Chuong, A. S., Klapoetke, N. C., Li, L., ... Zeng, H. (2015). Transgenic mice for intersectional targeting of neural sensors and effectors with high specificity and performance. *Neuron*, 85, 942–958.
- Malatesta, P., Hack, M. A., Hartfuss, E., Kettenmann, H., Klinkert, W., Kirchhoff, F., & Gotz, M. (2003). Neuronal or glial progeny: Regional differences in radial glia fate. *Neuron*, 37, 751–764.
- Malonek, D., & Grinvald, A. (1996). Interactions between electrical activity and cortical microcirculation revealed by imaging spectroscopy: Implications for functional brain mapping. *Science*, 272, 551–554.
- Mamczur, P., Borsuk, B., Paszko, J., Sas, Z., Mozrzymas, J., Wisniewski, J. R., ... Rakus, D. (2015). Astrocyte-neuron crosstalk regulates the expression and subcellular localization of carbohydrate metabolism enzymes. *Glia*, 63, 328–340.
- Mangia, S., Simpson, I. A., Vannucci, S. J., & Carruthers, A. (2009). The in vivo neuron-to-astrocyte lactate shuttle in human brain: Evidence from modeling of measured lactate levels during visual stimulation. *Journal of Neurochemistry*, 109(Suppl 1), 55–62.
- Martinez-Hernandez, A., Bell, K. P., & Norenberg, M. D. (1977). Glutamine synthetase: Glial localization in brain. *Science*, 195, 1356–1358.
- Mayevsky, A., & Chance, B. (1975). Metabolic responses of the awake cerebral cortex to anoxia hypoxia spreading depression and epileptiform activity. *Brain Research*, 98, 149–165.
- Miyawaki, A., Griesbeck, O., Heim, R., & Tsien, R. Y. (1999). Dynamic and quantitative Ca2+ measurements using improved cameleons. *Proceedings of the National Academy of Science USA*, 96, 2135–2140.
- Mookerjee, S. A., Gerencser, A. A., Nicholls, D. G., & Brand, M. D. (2017). Quantifying intracellular rates of glycolytic and oxidative ATP production and consumption using extracellular flux measurements. *Journal of Biological Chemistry*, 292, 7189–7207.
- Mookerjee, S. A., Goncalves, R. L., Gerencser, A. A., Nicholls, D. G., & Brand, M. D. (2015). The contributions of respiration and glycolysis to extracellular acid production. *Biochimica et Biophysica Acta*, 1847, 171–181.
- Murphy, A. N., & Chan, D. (2014). *Mitochondrial Function*. 1 ed. New York: Academic Press.
- Nagel, G., Szellas, T., Huhn, W., Kateriya, S., Adeishvili, N., Berthold, P., ... Bamberg, E. (2003). Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. *Proceedings of the National Academy of Science USA*, 100, 13940–13945.
- Nagelhus, E. A., & Ottersen, O. P. (2013). Physiological roles of aquaporin-4 in brain. *Physiological Review*, 93, 1543–1562.
- Newman, L. A., Korol, D. L., & Gold, P. E. (2011). Lactate produced by glycogenolysis in astrocytes regulates memory processing. *PLoS One*, 6, e28427.
- Nimmerjahn, A., Kirchhoff, F., & Helmchen, F. (2005). Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science*, 308, 1314–1318.
- Ogawa, S., Lee, T. M., Kay, A. R., & Tank, D. W. (1990). Brain magnetic resonance imaging with contrast dependent on blood oxygenation. *Proceedings of the National Academy of Science USA*, 87, 9868–9872.



- Oz, G., Alger, J. R., Barker, P. B., Bartha, R., Bizzi, A., Boesch, C., ... Kauppinen, R. A. (2014). Clinical proton MR spectroscopy in central nervous system disorders. *Radiology*, *270*, 658–679.
- Palygin, O., Lalo, U., & Pankratov, Y. (2011). Distinct pharmacological and functional properties of NMDA receptors in mouse cortical astrocytes. *British Journal of Pharmacology*, *163*, 1755–1766.
- Parpaleix, A., Goulam, H. Y., & Charpak, S. (2013). Imaging local neuronal activity by monitoring PO(2) transients in capillaries. *Nature Medicine*, *19*, 241–246.
- Paukert, M., Agarwal, A., Cha, J., Doze, V. A., Kang, J. U., & Bergles, D. E. (2014). Norepinephrine controls astroglial responsiveness to local circuit activity. *Neuron*, *82*, 1263–1270.
- Pendin, D., Greotti, E., Lefkimiatis, K., & Pozzan, T. (2017). Exploring cells with targeted biosensors. *Journal of General Physiology*, *149*, 1–36.
- Perea, G., Yang, A., Boyden, E. S., & Sur, M. (2014). Optogenetic astrocyte activation modulates response selectivity of visual cortex neurons *in vivo*. *Nature Communications*, *5*, 3262.
- Phizicky, E. M., & Fields, S. (1995). Protein–protein interactions: Methods for detection and analysis. *Microbiology Review*, *59*, 94–123.
- Pierre, K., Chatton, J. Y., Parent, A., Repond, C., Gardoni, F., Di, L. M., & Pellerin, L. (2009). Linking supply to demand: the neuronal monocarboxylate transporter MCT2 and the alpha-amino-3-hydroxyl-5-methyl-4-isoxazole-propionic acid receptor GluR2/3 subunit are associated in a common trafficking process. *European Journal of Neuroscience*, *29*, 1951–1963.
- Pierre, K., Magistretti, P. J., & Pellerin, L. (2002). MCT2 is a major neuronal monocarboxylate transporter in the adult mouse brain. *Journal of Cerebral Blood Flow and Metabolism*, *22*, 586–595.
- Pietraszewska-Bogiel, A., & Gadella, T. W. (2011). FRET microscopy: From principle to routine technology in cell biology. *Journal of Microscopy*, *241*, 111–118.
- Pike, G. B. (2012). Quantitative functional MRI: Concepts, issues and future challenges. *Neuroimage*, *62*, 1234–1240.
- Plaças, P. Y., de Tredern, E., Scheunemann, L., Trannoy, S., Goguel, V., Han, K. A., ... Preat, T. (2017). Upregulated energy metabolism in the *Drosophila* mushroom body is the trigger for long-term memory. *Nature Communications*, *8*, 15510. <https://doi.org/10.1038/ncomms15510>.
- Porras, O. H., Loaiza, A., & Barros, L. F. (2004). Glutamate mediates acute glucose transport inhibition in hippocampal neurons. *Journal of Neuroscience*, *24*, 9669–9673.
- Porras, O. H., Ruminot, I., Loaiza, A., & Barros, L. F. (2008). Na(+)-Ca(2+) cosignaling in the stimulation of the glucose transporter GLUT1 in cultured astrocytes. *Glia*, *56*, 59–68.
- Poskanzer, K. E., & Yuste, R. (2016). Astrocytes regulate cortical state switching *in vivo*. *Proceedings of the National Academy of Science USA*, *113*, E2675–E2684.
- Purich, D. (1979). *Enzyme Kinetics and Mechanism, Part A: Initial Rate and Inhibitor Methods*. New York: Academic Press.
- Purich, D. (1980). *Enzyme Kinetics and Mechanism, Part B: Isotopic Probes and Complex Enzyme Systems*. New York: Academic Press.
- Purich, D. (1982). *Enzyme Kinetics and Mechanism, Part C: Intermediates, Stereochemistry, and Rate Studies*. New York: Academic Press.
- Ransom, B. (2009). The neuronal microenvironment. In: W. F. Boron, E. L. Boulpaep, (Eds.). *Medical Physiology* (pp. 289–309). Philadelphia: Saunders Elsevier.
- Reinert, K. C., Gao, W., Chen, G., Wang, X., Peng, Y. P., & Ebner, T. J. (2011). Cellular and metabolic origins of flavoprotein autofluorescence in the cerebellar cortex *in vivo*. *Cerebellum*, *10*, 585–599.
- Reivich, M., Kuhl, D., Wolf, A., Greenberg, J., Phelps, M., Ido, T., ... Sokoloff, L. (1979). The [18F]fluorodeoxyglucose method for the measurement of local cerebral glucose utilization in man. *Circ Res*, *44*, 127–137.
- Robinson, M. B., & Jackson, J. G. (2016). Astroglial glutamate transporters coordinate excitatory signaling and brain energetics. *Neurochemistry International*, *98*, 56–71.
- Rodrigues, T. B., Valette, J., & Bouzier-Sore, A. K. (2013). (13)C NMR spectroscopy applications to brain energy metabolism. *Frontiers in Neuroenergetics*, *5*, 9.
- Rogers, G. W., Brand, M. D., Petrosyan, S., Ashok, D., Elorza, A. A., Ferrick, D. A., & Murphy, A. N. (2011). High throughput microplate respiratory measurements using minimal quantities of isolated mitochondria. *PLoS One*, *6*, e21746.
- Rose, E. M., Koo, J. C., Antflick, J. E., Ahmed, S. M., Angers, S., & Hampson, D. R. (2009). Glutamate transporter coupling to Na, K-ATPase. *Journal of Neuroscience*, *29*, 8143–8155.
- Rothman, D. L., Sibson, N. R., Hyder, F., Shen, J., Behar, K. L., & Shulman, R. G. (1999). *In vivo* nuclear magnetic resonance spectroscopy studies of the relationship between the glutamate-glutamine neurotransmitter cycle and functional neuroenergetics. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, *354*, 1165–1177.
- Ruminot, I., Gutiérrez, R., Peña-Munzenmeyer, G., Añazco, C., Sotelo-Hitschfeld, T., Lerchundi, R., ... Barros, L. F. (2011). NBCe1 mediates the acute stimulation of astrocytic glycolysis by extracellular K<sup>+</sup>. *Journal of Neuroscience*, *31*, 14264–14271.
- Sakadzic, S., Roussakis, E., Yaseen, M. A., Mandeville, E. T., Srinivasan, V. J., Arai, K., ... Boas, D. A. (2010). Two-photon high-resolution measurement of partial pressure of oxygen in cerebral vasculature and tissue. *Nature Methods*, *7*, 755–759.
- Sakadzic, S., Roussakis, E., Yaseen, M. A., Mandeville, E. T., Srinivasan, V. J., Arai, K., Ruvinskaya, S., Wu, W., Devor, A., Lo, E. H., Vinogradov, S. A., & Boas, D. A. (2011). Cerebral blood oxygenation measurement based on oxygen-dependent quenching of phosphorescence. *Journal of Visualized Experiments*, *51*, 1694.
- San Martín, A., Ceballo, S., Baeza-Lehnert, F., Lerchundi, R., Valdebenito, R., Contreras-Baeza, Y., ... Barros, L. F. (2014a). Imaging mitochondrial flux in single cells with a FRET sensor for pyruvate. *PLoS ONE*, *9*, e85780.
- San Martín, A., Ceballo, S., Ruminot, I., Lerchundi, R., Frommer, W. B., & Barros, L. F. (2013). A genetically encoded FRET lactate sensor and its use to detect the Warburg effect in single cancer cells. *PLoS ONE*, *8*, e57712.
- San Martín, A., Sotelo-Hitschfeld, T., Lerchundi, R., Fernández-Moncada, I., Ceballo, S., Valdebenito, R., ... Barros, L. F. (2014b). Single-cell imaging tools for brain energy metabolism: A review. *Neurophotonics*, *1*, 011004.
- Schousboe, A., Svenneby, G., & Hertz, L. (1977). Uptake and metabolism of glutamate in astrocytes cultured from dissociated mouse brain hemispheres. *Journal of Neurochemistry*, *29*, 999–1005.
- Seddon, A. M., Curnow, P., & Booth, P. J. (2004). Membrane proteins, lipids and detergents: Not just a soap opera. *Biochimica et Biophysica Acta*, *1666*, 105–117.
- Sekar, R. B., & Periasamy, A. (2003). Fluorescence resonance energy transfer (FRET) microscopy imaging of live cell protein localizations. *Journal of Cell Biology*, *160*, 629–633.
- Shibuki, K., Hishida, R., Murakami, H., Kudoh, M., Kawaguchi, T., Watanabe, M., ... Tanaka, R. (2003). Dynamic imaging of somatosensory cortical activity in the rat visualized by flavoprotein autofluorescence. *Journal of Physiology*, *549*, 919–927.
- Shuttleworth, C. W. (2010). Use of NAD(P)H and flavoprotein autofluorescence transients to probe neuron and astrocyte responses to synaptic activation. *Neurochemistry International*, *56*, 379–386.

- Simeonov, A., & Davis, M. I. (2004). Interference with Fluorescence and Absorbance. <https://www.ncbi.nlm.nih.gov/books/NBK343429/>
- Simon, H. A. (1962). The architecture of complexity. *Proceedings of the American Philosophical Society*, 106, 467–482.
- Sokoloff, L., Reivich, M., Kennedy, C., Des Rosiers, M. H., Patlak, C. S., Pettigrew, K. D., ... Shinohara, M. (1977). The [<sup>14</sup>C]deoxyglucose method for the measurement of local cerebral glucose utilization: theory, procedure, and normal values in the conscious and anesthetized albino rat. *Journal of Neurochemistry*, 28, 897–916.
- Sotelo-Hitschfeld, T., Fernández-Moncada, I., & Barros, L. F. (2012). Acute feedback control of astrocytic glycolysis by lactate. *Glia*, 60, 674–680.
- Sotelo-Hitschfeld, T., Niemeyer, M. I., Machler, P., Ruminot, I., Lerchundi, R., Wyss, M. T., ... Barros, L. F. (2015). Channel-mediated lactate release by K<sup>+</sup>-stimulated astrocytes. *Journal of Neuroscience*, 35, 4168–4178.
- Speizer, L., Haugland, R., & Kutchai, H. (1985). Asymmetric transport of a fluorescent glucose analogue by human erythrocytes. *Biochimica et Biophysica Acta*, 815, 75–84.
- Stagg, A., & Rothman, D. (2014). *Magnetic Resonance Spectroscopy: Tools for Neuroscience Research and Emerging Clinical Applications*. New York: Academic Press.
- Stridh, M. H., Alt, M. D., Wittmann, S., Heidtmann, H., Aggarwal, M., Riederer, B., ... Becker, H. M. (2012). Lactate flux in astrocytes is enhanced by a non-catalytic action of carbonic anhydrase II. *Journal of Physiology*, 590, 2333–2351.
- Stys, P. K., Ransom, B. R., & Waxman, S. G. (1991). Compound action potential of nerve recorded by suction electrode: A theoretical and experimental analysis. *Brain Research*, 546, 18–32.
- Tang, F., Lane, S., Korsak, A., Paton, J. F. R., Gourine, A. V., Kasparov, S., & Teschemacher, A. (2014). Lactate-mediated glia-neuronal signalling in the mammalian brain. *Nature Communications*, 5,
- Tchoussouff, J., Meric, P., Borredon, J., Fedeli, O., Gillet, B., Berenger, G., ... Seylaz, J. (1991). Value of *in vivo* NMR spectroscopy in the study of cerebral metabolism under inhalation anesthesia. *Agressologie*, 32, 333–338.
- Tekkok, S. B., Brown, A. M., Westenbroek, R., Pellerin, L., & Ransom, B. R. (2005). Transfer of glycogen-derived lactate from astrocytes to axons via specific monocarboxylate transporters supports mouse optic nerve activity. *Journal of Neuroscience Research*, 81, 644–652.
- Tian, Y. S., Lee, H. Y., Lim, C. S., Park, J., Kim, H. M., Shin, Y. N., ... Cho, B. R. (2009). A two-photon tracer for glucose uptake. *Angewandte Chemie International Edition in English*, 48, 8027–8031.
- Trivisoli, A., Saab, A. S., Winkler, U., Marx, G., Imamura, H., Mobius, W., ... Hirrlinger, J. (2017). Monitoring ATP dynamics in electrically active white matter tracts. *Elife*, 6,
- Tsien, R. Y. (2009). Indicators based on fluorescence resonance energy transfer (FRET). *Cold Spring Harbor Protocols*, 2009, db.
- Tsuyama, T., Kishikawa, J., Han, Y. W., Harada, Y., Tsubouchi, A., Noji, H., ... Imamura, H. (2013). *In vivo* fluorescent adenosine 5'-triphosphate (ATP) imaging of *Drosophila melanogaster* and *Caenorhabditis elegans* by using a genetically encoded fluorescent ATP biosensor optimized for low temperatures. *Analytical Chemistry*, 2015, 7889–7896.
- Vannucci, S. J., Maher, F., & Simpson, I. A. (1997). Glucose transporter proteins in brain: Delivery of glucose to neurons and glia. *Glia*, 21, 2–21.
- Volkenhoff, A., Hirrlinger, J., Kappel, J. M., Klämbt, C., & Schirmeier, S. (2017). Live imaging using a FRET glucose sensor reveals glucose delivery to all cell types in the *Drosophila* brain. *Journal of Insect Physiology*, 18, S0022–1910(17)30082–30083. <https://doi.org/10.1016/j.jinsphys.2017.07.010>. [Epub ahead of print]
- Wilson, J. E. (2003). Isozymes of mammalian hexokinase: Structure, sub-cellular localization and metabolic function. *Journal of Experimental Biology*, 206, 2049–2057.
- Wang, H., Yang, H., Shivalila, C. S., Dawlaty, M. M., Cheng, A. W., Zhang, F., & Jaenisch, R. (2013). One-step generation of mice carrying mutations in multiple genes by CRISPR/Cas-mediated genome engineering. *Cell*, 153, 910–918.
- Warburg, O. (1928). The chemical constitution of the respiration ferment. *Science*, 68, 437–443.
- Weber, B., Burger, C., Wyss, M. T., von Schulthess, G. K., Scheffold, F., & Buck, A. (2004). Optical imaging of the spatiotemporal dynamics of cerebral blood flow and oxidative metabolism in the rat barrel cortex. *European Journal of Neuroscience*, 20, 2664–2670.
- Widdas, W. F. (1954). Facilitated transfer of hexoses across the human erythrocyte membrane. *Journal of Physiology*, 125, 163–180.
- Widder, K., & Green, R. (1985). *Drug and Enzyme Targeting, Part A*. New York: Academic Press.
- Wong-Riley, M. T. (1989). Cytochrome oxidase: An endogenous metabolic marker for neuronal activity. *Trends in Neuroscience*, 12, 94–101.
- Yang, C., Ko, B., Hensley, C. T., Jiang, L., Wasti, A. T., Kim, J., ... DeBerardinis, R. J. (2014a). Glutamine oxidation maintains the TCA cycle and cell survival during impaired mitochondrial pyruvate transport. *Molecular Cell*, 56, 414–424.
- Yang, X., Hamner, M. A., Brown, A. M., Evans, R. D., Ye, Z. C., Chen, S., & Ransom, B. R. (2014b). Novel hypoglycemic injury mechanism: N-methyl-D-aspartate receptor-mediated white matter damage. *Annals Neurology*, 75, 492–507.
- Yoshioka, K., Takahashi, H., Homma, T., Saito, M., Oh, K. B., Nemoto, Y., & Matsuoka, H. (1996). A novel fluorescent derivative of glucose applicable to the assessment of glucose uptake activity of *Escherichia coli*. *Biochimica et Biophysica Acta*, 1289, 5–9.
- Yu, A. C., Drejer, J., Hertz, L., & Schousboe, A. (1983). Pyruvate carboxylase activity in primary cultures of astrocytes and neurons. *Journal of Neurochemistry*, 41, 1484–1487.
- Yudilevich, D. L., & Sepulveda, F. V. (1976). The specificity of amino acid and sugar carriers in the capillaries of the dog brain studied *in vivo* by rapid indicator dilution. *Advances in Experimental Medicine and Biology*, 69, 77–87.
- Zhang, J., Nuebel, E., Wisidagama, D. R., Setoguchi, K., Hong, J. S., Van Horn, C. M., ... Teitell, M. A. (2012). Measuring energy metabolism in cultured cells, including human pluripotent stem cells and differentiated cells. *Nature Protocols*, 7, 1068–1085.
- Zhang, R. W. K. (2017). High performance enzyme kinetics of turnover, activation and inhibition for translational drug discovery. *Expert Opinion in Drug Discovery*, 12, 17–37.
- Zhang, Y., Chen, K., Sloan, S. A., Bennett, M. L., Scholze, A. R., O'keeffe, S., ... Wu, J. Q. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *Journal of Neuroscience*, 34, 11929–11947.
- Zimmer, E. R., Parent, M. J., Souza, D. G., Leuzy, A., Lecrux, C., Kim, H. I., ... Rosa-Neto, P. (2017). [<sup>18</sup>F]FDG PET signal is driven by astroglial glutamate transport. *Nature Neuroscience*, 20, 393–395.
- Zipfel, W. R., Williams, R. M., & Webb, W. W. (2003). Nonlinear magic: Multiphoton microscopy in the biosciences. *Nature Biotechnology*, 21, 1369–1377.

**How to cite this article:** Barros LF, Bolaños JP, Bonvento G, et al. Current technical approaches to brain energy metabolism. *Glia*. 2018;66:1138–1159. <https://doi.org/10.1002/glia.23248>