The anatomy of melancholia – focal abnormalities of cerebral blood flow in major depression

CHRISTOPHER J. BENCH,¹ KARL J. FRISTON, RICHARD G. BROWN,
LYNETTE C. SCOTT, RICHARD S. J. FRACKOWIAK AND RAYMOND J. DOLAN

From the Academic Department of Psychiatry, Royal Free Hospital and School of Medicine;
MRC Cyclotron Unit, Hammersmith Hospital; National Hospital for Neurology and Neurosurgery;
and MRC Human Movement and Balance Unit, London

SYNOPSIS Using positron emission tomography (PET) and ¹⁵Oxygen, regional cerebral blood flow (rCBF) was measured in 33 patients with primary depression, 10 of whom had an associated severe cognitive impairment, and 23 age-matched controls. PET scans from these groups were analysed on a pixel-by-pixel basis and significant differences between the groups were identified on Statistical Parametric Maps (SPMs). In the depressed group as a whole rCBF was decreased in the left anterior cingulate and the left dorsolateral prefrontal cortex ($P < 0.05$ Bonferroni-corrected for multiple comparisons). Comparing patients with and without depression-related cognitive impairment, in the impaired group there were significant decreases in rCBF in the left medial frontal gyrus and increased rCBF in the cerebellar vermis ($P < 0.05$ Bonferroni-corrected). Therefore an anatomical association has been described between the rCBF profiles associated with depressed mood and depression-related cognitive impairment. The pre-frontal and limbic areas identified in this study constitute a distributed anatomical network that may be functionally abnormal in major depressive disorder.

INTRODUCTION

Major depression is a common illness, with an annual incidence of 0.4–2.7 per 1000 (Nielsen, 1976; Helgason, 1977) and a lifetime prevalence of up to 6.7% (Robins *et al.* 1984). Increasing age and genetic loading are among the most potent risk factors for severe depression. Although presenting most commonly as mood disturbance, depression may also present under a number of other guises including psychomotor disturbance and cognitive impairment, so-called depressive pseudodementia. This latter presentation may lead to a misdiagnosis of primary degenerative dementia (Marsden & Harrison, 1972).

The mediating biological mechanisms in depression are unknown. A guiding assumption in research is that depression is associated with functional disturbance in one or more cerebral monoamine pathways. Evidence implicating

specific monoamine systems is indirect and based largely on examination of bodily fluids, such as CSF (van Praag *et al.* 1970), or inferred from abnormal neuroendocrine responses to specific probes (van Praag, 1982). Cell bodies of the monoamine systems are localized to the brainstem and send relatively diffuse projections to subcortical and cortical structures. Whether the proposed abnormalities are anatomically localized or globally distributed is not specified by the monoamine theories, although the limbic system is frequently invoked as a likely site of perturbed function. Consequently the precise neuroanatomical systems involved in depression and its associated impairments such as the cognitive impairment of depression have not been defined.

Developments in functional imaging techniques offer new possibilities for exploring brain behavioural relationships in the major psychiatric syndromes. In the adult brain neuronal activity is closely linked to regional blood flow (rCBF) and metabolism of oxygen and glucose. Positron Emission Tomography (PET)-

Address for correspondence: Dr Christopher J. Bench, MRC Cyclotron Unit, Hammersmith Hospital, DuCane Road, London W12 0HS.

derived measures of rCBF and metabolism are a sensitive and reliable index of local neuronal function (Raichle, 1987), and such measures may be used to explore the functional anatomy of specific psychiatric disorders. Using PET, regional decreases in glucose metabolic rates have been reported in bipolar and unipolar depressed patients with normalization on clinical recovery (Baxter *et al.* 1985). We report a PET study of regional cerebral blood flow in depressed subjects, combined with statistical parametric mapping (Friston & Frackowiak, 1991), which identifies dissociations between focal abnormalities in cerebral function due to mood disturbance on the one hand and associated cognitive impairment on the other.

METHOD

Patients and Controls

Patients with an upper age limit of 75 years were recruited from a regional psychiatric service and from a national referral centre. Potential participants were approached to ascertain willingness to co-operate with the study and ability to give informed consent. In those who consented, the Schedule for Affective Disorders and Schizophrenia (SADS) (Endicott & Spitzer, 1978) was administered. Patients who met Research Diagnostic Criteria (RDC) (Spitzer *et al.* 1977) for major depressive disorder were further evaluated, and excluded if there was a history of organic brain disease, drug or alcohol abuse or significant medical illness as assessed by clinical examination and haematological, biochemical and endocrinological parameters. Subjects were also excluded if they scored over 4 on the Hachinski ischaemic scale (Hachinski *et al.* 1975) or had evidence of focal abnormality on CT or MRI scans. Patients on psychotropic medication (19 of the 33) were entered into the study, as it was predicted that patterns of rCBF would reflect their depressed mental state rather than medication – a prediction which could be tested. The inclusion of medicated patients also allowed the examination of patients in whom drug washout was not ethical. Of the medicated patients nine were taking tricyclic antidepressants alone and one was taking a monoamine oxidase inhibitor. In the remaining nine patients medication additional to tricyclics included neuroleptics ($N = 5$), lithium ($N = 2$), carba-

mazepine ($N = 1$) and tryptophan ($N = 2$). Of the unmedicated patients, six were drug naïve, two had been drug-free for greater than 1 year and five had been drug-free for over 2 weeks. A single patient who had been taking antidepressants erratically was drug-free for 2 days. All patients had moderate to severe depression as rated on the 17-item Hamilton Rating Scale for Depression (Hamilton, 1960) and on the Montgomery and Åsberg Depression Rating Scale (Montgomery & Åsberg, 1979). Handedness was assessed according to Oldfield (Oldfield, 1971). Those subjects who satisfied the entry criteria received a comprehensive battery of neuropsychological tests, designed specifically to target cognitive functions thought to be impaired in depression. These data will be presented elsewhere. PET scans were performed with 3 days of neuropsychological assessment. This study was approved by the ethical committees of all referring hospitals and by the local committee of the Hammersmith Hospital, where the scans were performed. Permission to administer ^{15}O was obtained from the Administration of Radioactive Substances Advisory Committee (ARSAC). All subjects gave informed written consent. Demographic data are presented in Table 1. PET studies of 18 control subjects were taken from the library of normal scans available at the MRC Cyclotron Unit and supplemented with five subjects recruited prospectively by local advertisement. None of the controls had a history of psychiatric or significant medical illness, nor were they taking psychotropic medication (Table 1).

Table 1. Characteristics of the patients and controls

	Depressed	Controls
Number	33	23
Sex (M/F)	21/12	10/13
Age (yr mean $\pm$ S.D.)	56.8 ($\pm$ 12.8)	63.4 ($\pm$ 11.6)
Handedness (R/L)	30/3	22/1
Psychotropic medication ($\pm$)	19/14	—
HAM-D 17-item (mean $\pm$ S.D.)	25 ($\pm$ 4.1)	—
MADRS (mean $\pm$ S.D.)	30 ($\pm$ 5.5)	—
Unipolar/Bipolar	30/3	—

PET scanning

All patients and controls underwent steady-state measurement of cerebral blood flow using PET

and C^{15}O_2 inhalation. Studies were performed in a supine position in a room with lights and minimal noise other than the scanner's cooling system. For each scan a polystyrene head mould was used to ensure correct positioning and relative head immobilization. A cannula was inserted into a cannula in the Allen's test for collateral circulation of the skin with 1% Marcaine. Subjects were aligned in reference to a laser-beam system. The detectors were parallel to the (OM) line. They were asked to remain still during the study, but no other instructions were given. Measured attenuation correction scans were made using counts in blank and transmission scans using a ^{68}Ge ring source. Correction of the raw emission scans was performed into parametric images of whole-blood and plasma flow during the scans. Details of the reconstruction have been previously described (Spinks, 1980). After reconstruction pixel size was 2.05 mm^2 and the slice thickness was $8.5 \times 8.5 \times 7.0 \text{ mm}$. Studies were performed using a PET scanner (CTI PET System 931-08/12 CTI PET System, Knoxville, Tennessee, USA) whose performance has been described (Spinks, 1980). A retrospective control subject was scanned prior to recruitment of the prospective controls ($N = 5$) in a random order with the patient. The patient was acquired using the same equipment over a two-year period.

Image and statistical analysis

Image analysis was performed using SPARC Workstation (Sun Microsystems Inc., Sunnyvale, CA, USA). The data within the 10.5 cm scan were interpolated to the voxels approximately 2.05 mm^3 were checked for artefacts and motion. To enable intersubject comparison the scan data the images were transformed into standard stereotactic space. For this transformation was used the AC-PC line, which was defined from anatomical landmark

) and tryptophan ($N = 2$). Of the patients, six were drug naïve, drug-free for greater than 1 year and five had been taking antidepressants and were drug-free for 2 days. All had moderate to severe depression as assessed by the Hamilton Rating Scale for Depression (Hamilton, 1960) and on the Asberg Depression Rating Scale (Asberg & Asberg, 1979). Handedness was assessed according to Oldfield (Oldfield, 1971). Subjects who satisfied the entry criteria for a comprehensive battery of neuropsychological tests, designed specifically to assess cognitive functions thought to be impaired in depression. These data will be reported elsewhere. PET scans were performed as part of a neuropsychological assessment approved by the ethical committees of the participating hospitals and by the local Research Ethics Committee at Hammersmith Hospital, where the scans were performed. Permission to add subjects was obtained from the local Research Ethics Committee (ARSAC). All subjects gave written consent. Demographic data are given in Table 1. PET studies of 18 depressed patients were taken from the library of PET studies available at the MRC Cyclotron Unit, which was complemented with five subjects recruited by local advertisement. Controls had a history of psychiatric illness, nor were they taking any psychotropic medication (Table 1).

Characteristics of the patients and controls

	Depressed	Controls
Number	33	23
Age (mean $\pm$ S.D.)	21/12	10/13
Weight (kg)	56.8 ($\pm$ 12.8)	63.4 ($\pm$ 11.4)
Height (cm)	30/3	22/1
Education (years)	19/14	—
IQ (mean $\pm$ S.D.)	25 ($\pm$ 4.1)	—
Verbal IQ (mean $\pm$ S.D.)	30 ($\pm$ 5.5)	—
Non-verbal IQ (mean $\pm$ S.D.)	30/3	—

18 controls underwent steady-state PET studies of cerebral blood flow using PET

and $C^{15}O_2$ inhalation. Studies were performed in the supine position in a room with dimmed lights and minimal noise other than that of the scanner's cooling system. For each subject a custom-made polystyrene head mould was made which was used to ensure correct positioning in the scanner and relative head immobility. A 22 g Teflon cannula was inserted into a radial artery after Allen's test for collateral circulation and infiltration of the skin with 1% Marcain (Bupivacaine). Subjects were aligned in the scanner with reference to a laser-beam system so that the detectors were parallel to the orbito-meatal (OM) line. They were asked to close their eyes during the study, but no other instructions were given. Measured attenuation corrections of the emission scans were made with the ratio of counts in blank and transmission scans obtained using a ^{68}Ge ring source. After attenuation correction the raw emission data were transformed into parametric images of CBF using the whole-blood and plasma activities measured during the scans. Details of these methods have been previously described (Frackowiak *et al.* 1980). After reconstruction and filtering the pixel size was 2.05 mm^2 and the image resolution was $8.5 \times 8.5 \times 7.0 \text{ mm}$. Studies were performed using a PET scanner (CTI Knoxville model 931-08/12 CTI PET Systems, Knoxville, Tennessee, USA) whose physical performance has been described (Spinks *et al.* 1988). The retrospective control subjects ($N = 18$) were scanned prior to recruitment of the patients but the prospective controls ($N = 5$) were scanned in a random order with the patients. All scans were acquired using the same equipment and methods over a two-year period.

Image and statistical analysis

Image analysis was performed on a SUN 3/60 or SPARC Workstation (Sun Microsystems Europe Inc., Surrey, UK). The 15 original planes of data within the 10.5 cm field of view of the scanner were interpolated to 43 planes to render the voxels approximately cubic. The images were checked for artefacts and corrected for yaw and roll. To enable inter-subject averaging of scan data the images were then resized into a standard stereotactic space. The reference plane for this transformation was the intercommissural (AC-PC) line, which was identified directly from anatomical landmarks in the PET image

(Friston *et al.* 1989). The 43 planes of data were then resliced into the standard stereotactic space (Talairach & Tournoux, 1988). In this space each pixel represents $2 \times 2 \text{ mm}$ and the 26 new planes are 4 mm thick. The images were smoothed with a Gaussian filter to reduce the effect of individual variability in cortical gyral anatomy. Smoothing allows the constructive interference of equivalent sites of difference when images are averaged. This initial transformation of the data allowed subsequent analysis to be entirely data led and not subject to bias imposed by observer placement of regions of interest on the images.

Following stereotactic normalization the CBF images were adjusted for individual differences in global blood flow using an analysis of covariance (ANCOVA), as previously described (Friston *et al.* 1990). This procedure generated an adjusted mean blood flow map for each of the pooled groups (e.g. depressed and controls) and permitted an estimate of the error variance of the rCBF in each pixel. Group-by-group comparisons were then made on a pixel-by-pixel basis using the t statistic to compare adjusted (for whole-brain CBF) group means. The resulting statistical parametric maps (SPMs) of t values were displayed within the standard stereotactic space, plane by plane and as projections on to sagittal, coronal and transverse renderings of the brain.

In order to identify foci of significant difference throughout the brain volume without false positives, a threshold for significance was used to take into account the multiple simultaneous t tests performed. The (t)SPMs were therefore thresholded with a Bonferroni correction for repeated measurements (Friston *et al.* 1991). Although in practice the t tests were carried out on every pixel, the number of truly independent sites was far fewer, because neighbouring pixels are correlated due to averaging and smoothing. The correction applied corresponds to an expected false-positive rate of 1 per 20 planes.

RESULTS

Comparing the 33 depressed patients and 23 controls the (t)SPM identified two areas in the depressed group which showed significantly reduced rCBF. These areas were the left anterior

Table 2a. Co-ordinates of the pixels where the most significant differences in blood flow were identified. Co-ordinates refer to the stereotaxic atlas of Talairach and Tournoux. The CBF values are in units of ml/dl tissue/min, and have been adjusted for a global mean blood flow of 50 ml/dl/min. The Z score is a measure of the degree of significance of the difference and is the number of standard deviations from the mean t value in the (t) statistical map of the t value for the most significant pixel in the plane.

Location	Co-ordinates			Regional CBF		Z score
	x	y	z	Controls	Depressed	
(L) Anterior cingulate cortex	-14	24	36	57.4	54.5	3.46**
	-16	18	40	57.1	54.3	3.48**
(L) Dorsolateral prefrontal cortex	-38	34	20	50.0	47.4	3.74**

** $P < 0.05$ Bonferroni-corrected.

Table 2b. Blood flow in medicated and unmedicated patients at the regions where maximum difference was found in depressed patients compared with normal controls. The groups have been compared using an unpaired t test. The CBF values are in units of ml/dl tissue/min, and have been adjusted for a global mean blood flow of 50 ml/dl/min. Statistical maps at a threshold of $P < 0.01$ (non-corrected) did not detect any significant change at these co-ordinates when medicated and unmedicated patients were compared.

Location	Co-ordinates			Regional CBF		t	P
	x	y	z	Unmedicated	Medicated		
(L) Anterior cingulate cortex	-14	24	36	54.9	54.3	0.64	0.53
	-16	18	40	54.9	53.7	1.15	0.26
(L) Dorsolateral prefrontal cortex	-38	34	20	47.9	47.1	0.93	0.36

cingulate and the left dorsolateral prefrontal cortex respectively (Fig. 1). The CBF values for the pixels of maximum significance normalized to a global mean of 50 ml/dl/min are presented in Table 2. The first area identified was in the superior/anterior cingulate cortex (Brodmann's Area 32). This was centred at +36 mm with respect to the AC-PC line and extended from +32 mm to +40 mm. Results for the left dorsolateral prefrontal cortex (DLPFC) (BA 9,46) were of a similar order of significance and spread from +16 mm to +28 mm with respect to the AC-PC line. The most significant pixel was at +20 mm. The right DLPFC showed a strong trend to decreased blood flow (+24 to +28 mm), but did not survive Bonferroni correction. Two other regions in the depressed-patient sample also showed strong trends toward significant differences in rCBF ($P < 0.001$ non-Bonferroni-corrected). An increase in CBF was localized to the left posterior cingulate gyrus (BA 23,30). A decrease in CBF was localized to the left angular gyrus (BA 39) confluent with the

superior aspect of the superior temporal sulcus (Fig. 1).

To test the prediction that the effects of medication were not confounding the changes due to psychopathology, a second comparison was made between medicated ($N = 19$) and unmedicated ($N = 14$) depressives. This comparison was made at a lower threshold ($P < 0.01$, non-Bonferroni-corrected). No significant changes in rCBF were found at the regions previously identified. This finding does not allow the rejection of the null hypothesis that medication has no effect, but does indicate that medication effects cannot account for the profile of differences in rCBF between the depressed patients and controls. The rCBF values for medicated and unmedicated patients are presented in Table 2b.

A third comparison was made to examine specific changes within the depressed group associated with cognitive impairment. The patients were subdivided into three groups on the basis of their scores on the Mini-Mental

FIG. 1. Statistical parametric maps (SPM) showing significant differences in blood flow in the depressed group as a whole. The most significant differences were found in the cingulate cortex, superior/anterior cingulate cortex, and the left dorsolateral prefrontal cortex. The coronal slice is used with blue corresponding to the level of the transverse slice and the subjects' right is

FIG. 2. (t)SPM showing significant differences in blood flow in cognitively impaired depressed patients. White areas are t value

ices in blood flow were identified.
The CBF values are in units of
w of 50 ml/dl/min. The Z score
mber of standard deviations from
t significant pixel in the plane

al CBF

Depressed Z score

54.5	3.46**
54.3	3.48**
47.4	3.74**

regions where maximum difference
groups have been compared using
and have been adjusted for a global
 $P < 0.01$ (non-corrected) did not
and unmedicated patients were

CBF

Medicated t P

54.3	0.64	0.53
53.7	1.15	0.26
47.1	0.93	0.36

of the superior temporal sulcus

prediction that the effects of
re not confounding the changes
pathology, a second comparison
between medicated ($N = 19$) and
 $N = 14$) depressives. This com-
rade at a lower threshold ($P <$
ferroni-corrected). No significant
BF were found at the regions
ified. This finding does not allow
of the null hypothesis that med-
effect, but does indicate that
ects cannot account for the profile
in rCBF between the depressed
controls. The rCBF values for
unmedicated patients are presen-

b.
nparison was made to examine
ges within the depressed group
th cognitive impairment. The
subdivided into three groups on
their scores on the Mini-Mental

FIG. 1. Statistical parametric maps (t)SPM showing location of significant decreases in rCBF in the depressed group as a whole in comparison with the normal controls. White areas are t values at $P < 0.001$. The most significant decreases, in the left dorsolateral prefrontal cortex (DLPFC) and the anterior cingulate cortex, survive Bonferroni correction ($P < 0.05$). The group image is included for reference to show anatomical detail and is the average of the stereotactically normalized CBF data set from the 23 normal subjects. The choice of colour scale of the group image is arbitrary; in this case a rainbow scale is used with blue corresponding to lowest blood flow and white to highest. Numbers (e.g. +16) refer to level of the transverse slices in millimetres relative to the AC-PC line. Anterior is at the top of the image and the subjects' right is at the right. The grid corresponds to that used by Talairach & Tournoux (1988).

FIG. 2. (t)SPM showing location of significant increases (left) and decreases (right) in rCBF when the cognitively impaired depressed group is compared with non-cognitively impaired depressed patients. White areas are t values at $P < 0.05$ Bonferroni corrected.

Table 3. Characteristics of depressed patients with and without cognitive impairment as defined by their scores on the Mini-Mental State Examination

	Cognitively impaired depressed (MMSE $\leq$ 25)	Cognitively unimpaired depressed (MMSE $\geq$ 29)
Number	10	10
Age (yr mean $\pm$ S.D.)	60.9 ($\pm$ 7.8)	53.2 ($\pm$ 14.8)
HAM-D 17-item (mean $\pm$ S.D.)	25.7 ($\pm$ 4.2)	24.4 ($\pm$ 3.5)
Psychotropic medication ($\pm$)	8/2	4/6

State examination (MMSE). On this measure ten patients were unimpaired (score $\geq$ 29) and ten were impaired (score $\leq$ 25), while the remaining 13 patients achieved intermediate scores and were excluded from the subsequent analysis. These impaired patients also scored 85 or less on the CAMCOG subscale of the CAMDEX, which incorporates all of the items of the MMSE. The correlation between CAMCOG and MMSE scores within the depressed group was 0.93 ($P < 0.001$). The impaired and unimpaired groups were matched for age and severity of depression (Table 3). Comparing scans from the impaired and unimpaired groups the (t)SPM, Bonferroni-corrected, identified foci of significant change in the depressed cognitively impaired group. These consisted of decreases in rCBF in the left medial prefrontal cortex (BA 10) (0 mm to +12 mm) and increases in rCBF in the right aspect of the cerebellar vermis (-12 mm to -4 mm) (Fig. 2).

The co-ordinates of the most significant pixels and magnitude of changes are shown in Table 4.

A fourth and final comparison was made between the cognitively impaired depressed group ($N = 10$) and the controls ($N = 23$). As predicted, this analysis showed significant changes in rCBF characteristic of both the cognitive impairment of depression and major depression, i.e. decreases in the left medial frontal and dorsolateral prefrontal cortices and the anterior cingulate, and increases in the cerebellar vermis.

DISCUSSION

These findings indicate that functional changes in the left anterior cingulate and the DLPFC are associated with the syndrome of major depression, while additional changes in the medial prefrontal cortex are associated with the cognitive impairment of depression, so-called depressive pseudodementia. The findings offer direct *in vivo* evidence of limbic system involvement, specifically the anterior cingulate cortex, in the pathogenesis of depression. Using the Xenon-133 inhalation technique, reductions in rCBF have been described in selective frontal, central, superior temporal and anterior parietal regions (Sackeim *et al.* 1990). Previous PET studies of depressed patients have found decreased metabolism in the inferior frontal lobe (Buchsbbaum *et al.* 1984). More specific findings include lower hemispheric metabolic rates in bipolar patients (Baxter *et al.* 1985), relative caudate hypometabolism in unipolar patients

Table 4. Co-ordinates of the pixels where the most significant differences in blood flow were identified. Co-ordinates refer to the stereotaxic atlas of Talairach and Tournoux. The CBF values are in units of ml/dl tissue/min, and have been adjusted for a global mean blood flow of 50 ml/dl/min. The Z score is a measure of the degree of significance of the difference and is the number of standard deviations from the mean t value in the (t) statistical map of the t value for the most significant pixel in the plane

Location	Co-ordinates			Regional CBF		Z score
	x	y	z	Impaired	Unimpaired	
(L) Medial frontal cortex	-8	48	0	57.2	62.6	3.98**
	-6	52	4	55.1	60.2	4.55**
	-6	54	8	51.4	56.2	4.29**
	-6	54	12	50.1	54.1	3.56**
Cerebellar vermis	8	-66	-12	81.6	72.7	3.75**
	10	-60	-8	78.3	70.0	3.82**
	8	-48	-4	68.1	61.7	3.73**

** $P < 0.05$ Bonferroni-corrected.

(Schwartz *et al.* 1987) and impaired left DLPFC metabolism in both subgroups (Baxter *et al.* 1989).

The regions implicated by this study have, as individual deficits, little specificity for depression. In particular, functional abnormalities in the left DLPFC appear to be common to a number of major psychiatric syndromes (Weinberger *et al.* 1986). However, the pattern of deficits described in the present study may constitute a profile unique to the depressive syndrome. Although each regional deficit is associated with a symptomatic or neuropsychological component that is a part of the syndrome of depression, it can also be seen in other disorders. For example, impaired higher-order cognitive function and a left medial prefrontal deficit appear to be common to depressive pseudodementia and psychomotor poverty syndrome of schizophrenia (Liddle *et al.* 1990). Thus, functional deficits in this area may provide the basis for higher cognitive impairments seen in the major psychoses. The pattern of decreased rCBF in the medial prefrontal area is also distinct from the characteristic metabolic changes seen with cognitive impairment and a different neuropsychological profile in the more common neurodegenerative and vascular dementias (Frackowiak *et al.* 1981).

The lateralization of the significant decreases in rCBF to the left hemisphere is striking. Previous work on lateralization has tended to implicate the right hemisphere in the pathogenesis of depression (Flor-Henry, 1979). The findings in this study are consistent with some investigations of stroke patients which show that left-hemisphere lesions are associated with depression, and severity of depression correlates with the proximity of the lesion to the anterior pole of the left hemisphere (Robinson *et al.* 1983, 1984). This is a contentious area of research, and other studies have shown that the relationship with site of lesion is not upheld when the patient sample includes non-in-patients interviewed using standardized instruments (House *et al.* 1990; Sharpe *et al.* 1990).

Significant and non-significant changes in rCBF were noted in the cerebellum and posterior cingulate respectively. The pattern of significantly increased blood flow in the cerebellar vermis was unexpected. However, a number of converging lines of evidence implicate the cer-

ebellum in the regulation of emotion. The midline cerebellum (vermis and fastigial nucleus) has previously been demonstrated to be an integral part of a neural network for emotional expression with anatomical connections and functional relationships to the limbic system (Heath & Harper, 1974; Heath *et al.* 1978). Several case reports have described affective concomitants of vermal pathology (Heath *et al.* 1979; Hamilton *et al.* 1983). In a PET study of patients with anxiety disorder increased blood flow in the cerebellum was observed (Reiman *et al.* 1989).

Structural imaging was not performed routinely in the depressed patients, but 15 of the 33 had CT or MRI scans. No focal abnormalities were reported in these scans, although atrophic or involutional changes were reported in 3 of the 15. It is known that a proportion of depressed patients display CT evidence of structural brain change (Dolan *et al.* 1985, 1986). These abnormalities are generalized, unrelated to clinical features of depression and more pronounced in elderly subjects. Although the majority of studies have found these changes to be non-progressive, a recent report has raised the possibility of progressive loss of tissue in the temporal lobes with increasing duration of illness (Altshuler *et al.* 1991). Depressed patients with cognitive deficits may have ventricular brain ratios (VBRs) intermediate in size between those of depressed cognitively normal patients and patients with probable Alzheimer's disease (Pearlson *et al.* 1989). It seems unlikely that neuronal loss secondary to gross structural changes can account for the regional functional abnormalities we have described. Previous studies of Alzheimer's disease have shown that the majority of metabolic abnormalities identified by PET cannot be explained by cortical atrophy (Fazekas *et al.* 1989). It would require a systematic difference in brain structure, for example cortical thinning, within the depressed group to lead to the profile of abnormalities described in this study. Even if a component of the functional changes described is due to structural differences, either macroscopic or microscopic (decreases in synaptic or neuronal density), the regional specificity remains of great interest.

Psychotropic medication failed to account for the differences found, suggesting that the main

source of change in rCBF was the depressed mental state. The differences were mainly at the same (early) illness and treatment. It is possible that rCBF normalizes over a time course that parallels the therapeutic effects of antidepressant medication. Studies have shown that glucose metabolism in the left IFC normalizes with antidepressant medication in depression (Baxter *et al.* 1990). Follow-up scans in a particular phase will identify whether the changes in particular those specific to the depressed group, are reversible.

The limbic system, particularly the cingulate gyrus, has long been implicated in depression (Papez, 1937) (MacLean, 1952) of emotion. A classic formulation of the role of the limbic system, speculated that 'radiating processes from the gyri cinguli in the cerebral cortex would colouring to psychic processes'. The procedure of cingulotomy has reported success in both refractory and intractable pain. The procedure suggests that the emotional tone to lower-cerebral processes is modulated. Cingulotomy for intractable pain abolishes the sensation of pain and abolishes the associated unpleasantness (White, 1962). Among cerebellar cingulate gyrus shows the decrease in rCBF with age (1). This selective vulnerability of activity may provide a biological basis for the increased incidence of major depression in older age groups.

The role of the cingulate centre has recently been investigated (Pardo *et al.* 1990). The cingulate was inferior and posterior in the present study. That attentional impairment is a symptom in depressive illness, and these two centres demonstrate similar changes in these studies could implicate a common involvement of the anterior cingulate in depression. These data lend support to the view that the cingulate can be considered as a bridge between attention and emotion.

tion of emotion. The nucleus and fastigial nucleus) demonstrated to be an anatomical network for emotional connections and is related to the limbic system (Heath *et al.* 1978). We have described affective pathology (Heath *et al.* 1983). In a PET study of major depression increased blood flow was observed (Reiman *et al.*

was not performed routinely in depressed patients, but 15 of the 33 patients. No focal abnormalities were reported in 3 of the 15 scans, although atrophic changes were reported in 3 of the 15. A proportion of depressed patients showed evidence of structural brain changes (Pearlson *et al.* 1985, 1986). These abnormalities were localized, unrelated to clinical severity and more pronounced in older patients. Although the majority of studies suggest that depression is non-progressive, our study raised the possibility of progression of illness (Altshuler *et al.* 1989). We found that patients with cognitive impairment and higher ventricular brain ratios (VBRs) between those of depressed patients and patients with normal brain ratios (Pearlson *et al.* 1985) were more likely that neuronal loss was structural changes described. Previous studies have shown that the abnormalities identified by PET are explained by cortical atrophy (Pearlson *et al.* 1989). It would require a change in brain structure, for example, within the depressed state, to account for the profile of abnormalities identified in the present study. Even if a component of the changes described is due to atrophy, either macroscopic or microscopic, the specificity of the changes in synaptic or neuronal function remains of great interest. Medication failed to account for the changes, suggesting that the

source of change in rCBF was due to the depressed mental state. The depressed patients were mainly at the same (early) phase of their illness and treatment. It is conceivable that rCBF normalizes over a time scale which parallels the therapeutic effects of antidepressants. Studies have shown that reductions in glucose metabolism in the left DLPFC normalize with antidepressant medication and improvement of depression (Baxter *et al.* 1989; Martinot *et al.* 1990). Follow-up scans in the recovered phase will identify whether the changes observed, in particular those specific to the cognitively impaired group, are reversible, static or progressive.

The limbic system, particularly the cingulate gyrus, has long been implicated in either the expression (Papez, 1937) or modulation (MacLean, 1952) of emotion. Papez, in his classic formulation of the role of the limbic system, speculated that 'radiations of emotive processes from the gyri cinguli to other regions in the cerebral cortex would add emotional colouring to psychic processes'. The surgical procedure of cingulotomy has been used with reported success in both resistant depression and intractable pain. The efficacy of such procedures suggests that the cingulate imparts emotional tone to lower-order perceptions. Cingulotomy for intractable pain does not abolish the sensation of pain, but rather extinguishes the associated unpleasant effect (Foltz & White, 1962). Among cerebral structures the cingulate gyrus shows the greatest functional decrease in rCBF with age (Martin *et al.* 1991). This selective vulnerability of anterior cingulate activity may provide a biological basis for the increased incidence of major depression in older age groups.

The role of the cingulate as an attentional centre has recently been investigated using PET (Pardo *et al.* 1990). The cingulate area highlighted was inferior and posterior to the area identified in the present study. It is noteworthy that attentional impairments are an invariable symptom in depressive illness. The proximity of these two centres demonstrated in separate studies could implicate a more widespread involvement of the anterior cingulate in depression. These data lend support to the view that the cingulate can be considered an interface between attention and emotion (Powell & Hines,

1974). A non-significant increase in rCBF was noted in the left posterior cingulate (BA 23). The anterior and posterior cingulate have reciprocal anatomical connections, and the increase in rCBF supports a notion of functional as well as anatomical coupling (Baleydier & Mauguière, 1980).

The importance of the pre-frontal focal abnormalities described in the present study is highlighted by primate studies which indicate a pattern of reciprocal connections between the anterior cingulate, the dorsolateral pre-frontal cortices and the medial frontal cortices (Pandya *et al.* 1981). The pre-frontal areas are therefore sites of convergence for limbic inputs with highly processed associative information, and may serve the function of integration of thought and emotion (Mesulam, 1986). Such structures would, therefore, seem to be part of an extended neural network, whose functions include the regulation of mood and directed attention, which is disturbed in major depression.

C.J.B. and K.J.F. are supported by grants from the Wellcome Trust. We would like to thank the staff of the MRC Cyclotron Unit, in particular Claire Taylor and Graham Lewington, without whose assistance this study would not have been possible.

REFERENCES

- Altshuler, L. L., Conrad, A., Hauser, P., Li, X., Guze, B. H., Denikoff, K., Tourtellotte, W. & Post, R. (1991). Reduction of temporal lobe volume in bipolar disorder: a preliminary report of magnetic resonance imaging. *Archives of General Psychiatry* **48**, 482-483.
- Baleydier, C. & Mauguière, F. (1980). The duality of the cingulate gyrus in the monkey. *Brain* **103**, 525-554.
- Baxter, L. R., Phelps, M. E., Mazziotta, J. C., Schwartz, J. M., Gerner, R. H., Selin, C. E. & Sumida, R. M. (1985). Cerebral metabolic rates for glucose in mood disorders. Studies with positron emission tomography and fluorodeoxyglucose F18. *Archives of General Psychiatry* **42**, 441-447.
- Baxter, L. R., Schwartz, J. M., Phelps, M. E., Mazziotta, J. C., Guze, B. H., Selin, C. E., Gerner, R. H. & Sumida, R. M. (1989). Reduction of prefrontal cortex metabolism common to three types of depression. *Archives of General Psychiatry* **46**, 243-250.
- Buchsbaum, M. S., DeLisi, L. E., Holcomb, H. H., Cappelletti, J., King, A. C., Johnson, J., Hazlett, E., Dowling-Zimmerman, S., Post, R. M., Morihisa, J., Carpenter, W., Cohen, R., Pickar, D., Weinberger, D. R., Margolin, R. & Kessler, R. M. (1984). Anteroposterior gradients in cerebral glucose use in schizophrenia and affective disorders. *Archives of General Psychiatry* **41**, 1159-1166.
- Dolan, R. J., Calloway, S. P. & Mann, A. H. (1985). Cerebral ventricular size in depressed subjects. *Psychological Medicine* **15**, 873-878.
- Dolan, R. J., Calloway, S. P., Thacker, P. & Mann, A. H. (1986). The cerebral cortical appearance in depressed subjects. *Psychological Medicine* **16**, 775-779.

- Endicott, J. & Spitzer, R. L. (1978). A diagnostic interview. The Schedule for Affective Disorders and Schizophrenia. *Archives of General Psychiatry* **35**, 837-844.
- Fazekas, F., Alavi, A., Chawluk, J. B., Zimmerman, R. A., Hackney, D., Bilaniuk, L., Rosen, M., Alves, W. M., Hurtig, H. I., Jamieson, D. G., Kushner, M. J. & Reivich, M. (1989). Comparison of CT, MR and PET in Alzheimer's dementia and normal aging. *Journal of Nuclear Medicine* **30**, 1607-1615.
- Flor-Henry, P. (1979). On certain aspects of the localization of the cerebral systems regulating and determining emotion. *Biological Psychiatry* **14**, 677-698.
- Foltz, E. L. & White, L. E. Jr (1962). Pain 'relief' by frontal cingulotomy. *Journal of Neurosurgery* **19**, 89-100.
- Frackowiak, R. S. J., Lenzi, G. L., Jones, T. & Heather, J. D. (1980). Quantitative measurement of regional cerebral blood flow and oxygen metabolism in man using ^{15}O and positron emission tomography: theory, procedure, and normal values. *Journal of Computer Assisted Tomography* **4**, 727-736.
- Frackowiak, R. S. J., Pozzilli, C., Legg, N. J., Du Boulay, G. H., Marshall, J., Lenzi, G. L. & Jones, T. (1981). Regional cerebral oxygen supply and utilization in dementia. A clinical and physiological study with Oxygen-15 and positron tomography. *Brain* **104**, 753-778.
- Friston, K. J. & Frackowiak, R. S. J. (1991). Imaging functional anatomy. In *Brain Work and Mental Activity, Alfred Benzon Symposium 31* (ed. N. A. Lassen, D. H. Ingvar, M. E. Raichle and L. Friberg), pp. 267-277. Munksgaard: Copenhagen.
- Friston, K. J., Passingham, R. E., Nutt, J. G., Heather, J. D., Sawle, G. V. & Frackowiak, R. S. J. (1989). Localisation in PET images: direct fitting of the intercommissural (AC-PC) line. *Journal of Cerebral Blood Flow and Metabolism* **9**, 690-695.
- Friston, K. J., Frith, C. D., Liddle, P. F., Dolan, R. J., Lammertsma, A. A. & Frackowiak, R. S. J. (1990). The relationship between global and local changes in PET scans. *Journal of Cerebral Blood Flow and Metabolism* **10**, 458-466.
- Friston, K. J., Frith, C. D., Liddle, P. F. & Frackowiak, R. S. J. (1991). Comparing functional (PET) images: the assessment of significant change. *Journal of Cerebral Blood Flow and Metabolism* **11**, 690-699.
- Hachinski, V. C., Iliff, L. D., Zilhka, E., Du Boulay, G. H., McAllister, V. L., Marshall, J., Ross Russell, R. W. & Symon, L. (1975). Cerebral blood flow in dementia. *Archives of Neurology* **32**, 632-637.
- Hamilton, M. (1960). A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry* **23**, 56-62.
- Hamilton, N. G., Frick, R. B., Takahashi, T. & Hopping, M. W. (1983). Psychiatric symptoms and cerebellar pathology. *American Journal of Psychiatry* **140**, 1322-1326.
- Heath, R. G. & Harper, J. W. (1974). Ascending projections of the cerebellar fastigial nucleus to the hippocampus, amygdala, and other temporal lobe sites: evoked potential and histological studies in monkeys and cats. *Experimental Neurology* **45**, 268-287.
- Heath, R. G., Dempsey, C. W., Fontana, C. J. & Myers, W. A. (1978). Cerebellar stimulation: effects on septal region, hippocampus, and amygdala of cats and rats. *Biological Psychiatry* **13**, 501-529.
- Heath, R. G., Franklin, D. E. & Shraberg, D. (1979). Gross pathology of the cerebellum in patients diagnosed and treated as functional psychiatric disorders. *Journal of Nervous and Mental Disease* **167**, 585-591.
- Helgason, L. (1977). Psychiatric services and mental illness in Iceland. *Acta Psychiatrica Scandinavica Suppl.* **268**.
- House, A., Dennis, M., Warlow, C., Hawton, K. & Molyneux, A. (1990). Mood disorders after stroke and their relation to lesion location. A CT scan study. *Brain* **113**, 1113-1129.
- Liddle, P. F., Friston, K. J., Hirsch, S. R. & Frackowiak, R. S. J. (1990). Regional cerebral metabolic activity in chronic schizophrenia. *Schizophrenia Research* **3**, 23-24.
- MacLean, P. D. (1952). Some psychiatric implications of physiological studies on frontotemporal portion of limbic system (visceral brain). *Electroencephalography and Clinical Neurophysiology* **4**, 407-418.
- Marsden, C. D. & Harrison, M. J. G. (1972). Outcome of investigation of patients with presenile dementia. *British Medical Journal* **ii**, 249-252.
- Martin, A. J., Friston, K. J., Colebatch, J. G. & Frackowiak, R. S. J. (1991). Decreases in regional cerebral blood flow with normal aging. *Journal of Cerebral Blood Flow and Metabolism* **11**, 684-689.
- Martinot, J. L., Hardy, P., Feline, A., Hurte, J. D., Mazoyer, B., Attar-Levy, D., Pappata, S. & Syrota, A. (1990). Left prefrontal glucose hypometabolism in the depressed state: a confirmation. *American Journal of Psychiatry* **147**, 1313-1317.
- Mesulam, M.-M. (1986). Frontal cortex and behavior. *Annals of Neurology* **19**, 320-325.
- Montgomery, S. A. & Åsberg, M. (1979). A new depression scale designed to be sensitive to change. *British Journal of Psychiatry* **134**, 382-389.
- Nielsen, J. (1976). The Samsø project from 1957 to 1974. *Acta Psychiatrica Scandinavica* **54**, 198-222.
- Oldfield, R. C. (1971). The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* **9**, 97-113.
- Pandya, D. N., Van Hoesen, G. W. & Mesulam, M.-M. (1981). Efferent connections of the cingulate gyrus in the Rhesus monkey. *Experimental Brain Research* **42**, 319-330.
- Papez, J. W. (1937). A proposed mechanism of emotion. *Archives of Neurology and Psychiatry* **79**, 217-224.
- Pardo, J. V., Pardo, P. J., Janer, K. W. & Raichle, M. E. (1990). The anterior cingulate cortex mediates processing selection in the Stroop attentional conflict paradigm. *Proceedings of the National Academy of Sciences* **87**, 256-259.
- Pearlson, G. D., Rabins, P. V., Kim, W. S., Speedie, L. J., Moberg, P. J., Burns, A. & Bascom, M. J. (1989). Structural brain CT changes and cognitive deficits in elderly depressives with and without reversible dementia ('pseudodementia'). *Psychological Medicine* **19**, 573-584.
- Powell, E. & Hines, G. (1974). The limbic system: an interface. *Behavioral Biology* **12**, 149-164.
- Raichle, M. E. (1987). In *Handbook of Physiology, section 1, Nervous System, vol. 5, Higher Functions of the Brain* (ed. F. Plum), pp. 643-669. Oxford University Press: New York.
- Reiman, E. M., Raichle, M. E., Robins, E., Mintun, M., Fusselman, M. J., Fox, P. T., Price, J. L. & Hackman, K. A. (1989). Neuroanatomical correlates of a lactate-induced anxiety attack. *Archives of General Psychiatry* **46**, 493-500.
- Robins, L. N., Helzer, J. E., Weissman, M. M., Orvaschel, H., Gruenberger, E., Burke, J. D., Jr & Regier, D. A. (1984). Lifetime prevalence of specific psychiatric disorders in three sites. *Archives of General Psychiatry* **41**, 949-958.
- Robinson, R. G., Starr, L. B., Kubos, K. L. & Price, T. R. (1983). A two-year longitudinal study of post-stroke mood disorders: findings during the initial evaluation. *Stroke* **14**, 736-741.
- Robinson, R. G., Kubos, K. L., Starr, L. B., Rao, K. & Price, T. R. (1984). Mood disorders in stroke patients. Importance of location of lesion. *Brain* **107**, 81-91.
- Sackeim, H. A., Prohovnik, I., Moeller, J. R., Brown, R. P., Apter, S., Prudic, J., Devanand, D. P. & Mukherjee, S. (1990). Regional cerebral blood flow in mood disorders. I. Comparison of major depressives and normal controls at rest. *Archives of General Psychiatry* **47**, 60-70.
- Schwartz, J. M., Baxter, L. R., Mazziotta, J. C., Gerner, R. H. & Phelps, M. E. (1987). The differential diagnosis of depression. Relevance of Positron Emission Tomography studies of cerebral glucose metabolism to the bipolar-unipolar dichotomy. *Journal of the American Medical Association* **258**, 1368-1374.
- Sharpe, M., Hawton, K., House, A., Molyneux, A., Sandercock, P., Bamford, J. & Warlow, C. (1990). Mood disorders in long-term survivors of stroke: associations with brain lesion location and volume. *Psychological Medicine* **20**, 815-828.
- Spinks, T. J., Jones, T., Gilardi, M. C. & Heather, J. D. (1980). Physical performance of the latest generation of commercial positron scanner. *IEEE Transactions* **27**, 721-725.
- Spitzer, R. L., Endicott, J. & Robins, E. (1978). *Criteria for a Selected Group of F* Research Division, New York State Psychiatric Institute, New York.
- Talbot, J. & Tournoux, P. (1988). *The Human Brain*. Georg Thieme Verlag, Stuttgart.
- van Praag, H. M. (1982). The significance of the limbic system in the pathogenesis of affective disorders. *Archives of General Psychiatry* **39**, 1-11.

- temporal portion of limbic system (visceralography and Clinical Neurophysiology).
- son, M. J. G. (1972). Outcome of investigation in senile dementia. *British Medical Journal* **2**, 11-15.
- J., Colebatch, J. G. & Frackowiak, R. S. J. (1988). Regional cerebral blood flow with normal arterial blood flow and metabolism. *Journal of Cerebral Blood Flow and Metabolism* **8**, 684-689.
- l., Feline, A., Huret, J. D., Mazoyer, B., Le Bars, D. & Syrota, A. (1990). Left prefrontal activation in the depressed state: a confirmation. *Psychiatry* **53**, 1313-1317.
- Frontal cortex and behavior. *Annals of the New York Academy of Sciences* **557**, 1-10.
- berg, M. (1979). A new depression scale: a change. *British Journal of Psychiatry* **135**, 1-6.
- Samsø project from 1957 to 1974. *Acta Psychiatrica Scandinavica* **54**, 198-222.
- assessment and analysis of handedness. *Neuropsychologia* **9**, 97-113.
- sen, G. W. & Mesulam, M. M. (1981). The cingulate gyrus in the Rhesus monkey. *Neuroscience* **42**, 319-330.
- posed mechanism of emotion. *Archives of General Psychiatry* **79**, 217-224.
- Janer, K. W. & Raichle, M. E. (1990). The cingulate cortex mediates processing selection in the Stroop paradigm. *Proceedings of the National Academy of Sciences USA* **87**, 256-259.
- P. V., Kim, W. S., Speedie, L. J., Moberg, E. & Combs, M. J. (1989). Structural brain CT deficits in elderly depressives with and without dementia ('pseudodementia'). *Psychological Medicine* **19**, 1049-1054.
- (1974). The limbic system: an interface. *Handbook of Physiology, section 1, Nervous System, Functions of the Brain* (ed. F. Plum). University Press: New York.
- l., E., Robins, E., Mintun, M., Fusselman, M. J. & Hackman, K. A. (1989). Neurochemical changes in a lactate-induced anxiety attack. *Archives of General Psychiatry* **46**, 493-500.
- l., E., Weissman, M. M., Orvaschel, H., J. D., Jr & Regier, D. A. (1984). Lifetime prevalence of psychiatric disorders in three sites. *Archives of General Psychiatry* **41**, 949-958.
- B., Kubos, K. L. & Price, T. R. (1983). A study of post-stroke mood disorders: clinical evaluation. *Stroke* **14**, 736-741.
- K. L., Starr, L. B., Rao, K. & Price, T. R. (1983). Importance of location of stroke lesions. *Stroke* **14**, 91-95.
- l., I., Moeller, J. R., Brown, R. P., Apantaku, D. P. & Mukherjee, S. (1990). Regional mood disorders. I. Comparison of major depressive disorder with controls at rest. *Archives of General Psychiatry* **47**, 1049-1054.
- l., R., Mazziotta, J. C., Gerner, R. H. & (1988). The differential diagnosis of depression: emission Tomography studies of cerebral metabolism in the bipolar-unipolar dichotomy. *Journal of Affective Disorders* **25**, 1368-1374.
- house, A., Molyneux, A., Sandercock, P. A. G. (1990). Mood disorders in long-term survivors of stroke: associations with brain lesion location and outcome. *Stroke* **21**, 815-828.
- Gilardi, M. C. & Heather, J. D. (1988). The latest generation of computerized diagnosis of depressions. II. Hormonal variables. *Comprehensive Psychiatry* **23**, 216-226.
- van Praag, H. M., Korf, J. & Puite, J. (1970). 5-hydroxyindoleacetic acid levels in the cerebrospinal fluid of depressed patients treated with probenecid. *Nature* **225**, 1259-1260.
- Weinberger, D. R., Berman, K. F. & Zec, R. F. (1986). Physiologic dysfunction of dorsolateral prefrontal cortex in schizophrenia. I. Regional cerebral blood flow evidence. *Archives of General Psychiatry* **43**, 114-124.