

Supporting information for:

1-Butanol as a Solvent for Efficient Extraction of

Polar Compounds from Aqueous Medium:

Theoretical and Practical Aspects.

Gerhard König,^{*,†,‡} Manfred T. Reetz,^{†,¶} and Walter Thiel[†]

Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany, Laboratory for Biomolecular Simulation Research, Center for Integrative Proteomics Research, and Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, New Jersey 08854, United States, and Department of Chemistry, Philipps-University Marburg, 35032 Marburg, Germany

E-mail: gkoenig@mpi-muelheim.mpg.de

*To whom correspondence should be addressed

[†]Max-Planck-Institut für Kohlenforschung, 45470 Mülheim an der Ruhr, Germany

[‡]Laboratory for Biomolecular Simulation Research, Center for Integrative Proteomics Research, and Department of Chemistry and Chemical Biology, Rutgers University, Piscataway, New Jersey 08854, United States

[¶]Department of Chemistry, Philipps-University Marburg, 35032 Marburg, Germany

Figure S1. Example for the quantification of cis-cyclohexane-1,2-diol in the aqueous phase with HPLC. The abscissa denotes the retention time while the ordinate describes the refractive index change relative to the pure mobile phase.

