



Available online at www.sciencedirect.com

ScienceDirect

**NUCLEAR
PHYSICS B**

ELSEVIER

Nuclear Physics B 943 (2019) 114606

www.elsevier.com/locate/nuclphysb

Quantum entanglement in de Sitter space from stringy axion: An analysis using α vacua

Sayantan Choudhury ^{a,b,*}, Sudhakar Panda ^{c,d,e}

^a *Quantum Gravity and Unified Theory and Theoretical Cosmology Group, Max Planck Institute for Gravitational Physics (Albert Einstein Institute), Am Mühlenberg 1, 14476 Potsdam-Golm, Germany*

^b *Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India*

^c *Institute of Physics, Sachivalaya Marg, Bhubaneswar, Odisha 751005, India*

^d *National Institute of Science Education and Research, Jatni, Bhubaneswar, Odisha 752050, India*

^e *Homi Bhabha National Institute, Training School Complex, Anushakti Nagar, Mumbai 400085, India*

Received 14 September 2018; received in revised form 6 March 2019; accepted 24 March 2019

Available online 19 April 2019

Editor: Tommy Ohlsson

Abstract

In this work, we study the phenomena of quantum entanglement by computing de Sitter entanglement entropy from von Neumann measure. For this purpose we consider a bipartite quantum field theoretic set up for axion field, previously derived from **Type II B** string theory compactified to four dimensions. We consider the initial vacuum to be CPT invariant non-adiabatic α vacua state under **SO(1, 4)** isometry, which is characterised by a real one-parameter family. To implement this technique we use a S^2 which divide the de Sitter into two exterior and interior sub-regions. First, we derive the wave function of axion in an open chart for α vacua by applying Bogoliubov transformation on the solution for Bunch-Davies vacuum state. Further, we quantify the density matrix by tracing over the contribution from the exterior region. Using this result we derive entanglement entropy, Rényi entropy and explain the long-range quantum effects in primordial cosmological correlations. Our results for α vacua provides the necessary condition for generating non zero entanglement entropy in primordial cosmology.

© 2019 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

* Corresponding author.

E-mail addresses: sayantan@aei.mpg.de, sayantan.choudhury@aei.mpg.de, sayanphysicsisi@gmail.com (S. Choudhury), panda@iopb.res.in (S. Panda).

1. Introduction

It is well accepted fact that von Neumann entropy is a measure of quantum entanglement to quantify long range correlation in condensed matter physics [1–3] and cosmology [4–21]. In condensed matter physics entanglement entropy exactly mimics the role of an order parameter and the corresponding phase transition phenomena can be characterised by correlations at quantum level. Also, it is expected that, from this understanding of long range effects in quantum correlations, we can understand the underlying physics of the theory of multiverse, bubble nucleation etc. in de Sitter space [22]. As a consequence, we can observe a prompt response due to the local measurement in quantum physics, by violating causal structure of the space-time. In quantum theory such causality violation is known as Einstein-Podolsky-Rosen (EPR) paradox [23]. But in such type of local measurement causality remains unaffected as the required quantum information is not propagating. In this context Schwinger effect in de Sitter space [24,25] is one of the prominent examples of quantum entanglement. In Schwinger effect, particle pair creation takes place with a finite separation in de Sitter space-time in presence of a constant electric field [15] and the quantum states exhibit long range correlation.

To quantify entanglement entropy in the context of quantum field theory, one requires to have a bipartite system. In strong coupling regime of a such a theory we can compute entanglement entropy by using the principles of gauge gravity duality in the bulk [26–35]. Using this technique many issues have been addressed in the context of holographic entanglement entropy. Further, in ref. [10] the authors have constructed a completely different computational algorithm to quantify entanglement entropy using Bunch Davies initial state¹ in de Sitter space. Later, in ref. [15] this has been extended to the computation of entanglement entropy using α vacua initial state in de Sitter space by following the techniques presented in ref. [10]. Moreover, using Bunch Davies initial state in de Sitter space, we have computed entanglement entropy in a field theory where the effective action contains a linear source term [14]. We have shown that this result complements the necessary condition for the violation of Bell's inequality in primordial cosmology.

In quantum field theory in de Sitter background, one can construct a one parameter family of initial vacuum states which are CPT invariant under $\text{SO}(1, 4)$ isometry group [36,37]. These classes of states are characterised by a real parameter α , known as α vacua. For $\alpha = 0$ these states reduce to the usual Bunch Davies vacuum state. In a more technical terms, the α vacua can be treated as squeezed quantum states, which are created by an unitary operator acting on the Bunch Davies vacuum state. This leads to the generalisation of Wick's theorem in interacting quantum field theory, which allows us to describe any free quantum field theory Green's function computed using α vacua in terms of the products of the Green's functions computed using Bunch Davies vacuum [38]. See refs. [37,39,40] for more details on the quantum field theory of α vacua. For a specific α vacuum state as a quantum initial condition, it is possible to describe the long range correlations within the framework of quantum field theory. As a consequence, the non-local quantum phenomena can be associated with the long range effects, which is described by quantum entanglement of vacuum state as an initial condition. We note that till date no such experiment is available using which one can be able to test the local behaviour of quantum field theory in cosmological scale (Hubble scale). However, it is expected that in future it may be possible to test such prescriptions. Additionally, it is important to note that, propagators in free quantum field theory of de Sitter space computed in presence of adiabatic Bunch Davies vacuum

¹ We note that, Bunch Davies initial state is exact equivalent to the Euclidean or adiabatic vacuum state in quantum field theory.

state manifest Hadamard singularity which is consistent with the result obtained in the context of Minkowski flat space-time limit [41,42].

However, for interacting quantum field theory in the de Sitter space-time background, such singular propagators applicable for adiabatic Bunch Davies vacuum are dubious. Hence α vacuum state plays significant role, using which one can express the propagators in interaction picture. In the quantum field theory described by the α vacua state the real parameter α plays the role of super-selection number associated with a quantum state of a different bipartite Hilbert space. But it is still a debatable issue that whether the interaction picture of the quantum field theory with any arbitrary α vacua with any super-selection rule are consistent with the physical requirements of quantum mechanics or not [39]. In general, one can treat the α vacua as a family of quantum initial state, where we have quantum fluctuation around an excited state. Here it might be possible that the Hilbert space corresponding to excited state (for α vacua) and the adiabatic Bunch Davies vacuum coincides with each other. In such a situation it is perfectly consistent to describe quantum field theory of excited state in de Sitter space in terms of the adiabatic Bunch Davies vacuum in the ultraviolet regime.² As a result this specific identification allows us to write an effective field theory description in the ultraviolet regime. This implies that identifying the correct and more appropriate quantum α vacuum state is fine tuned. However, this fine tuning only allows us to describe the quantum field theory with any excited states compared to ground state described by the adiabatic Bunch Davies vacuum. Using this prescription apart from inflationary paradigm, one may be able to explain a lot of unexplored late time physical phenomena of nonstandard vacuum state. In this work, we further generalise the computational strategy of entanglement entropy for axion field using α vacua initial state in de Sitter space. This result will establish the generation of quantum entanglement entropy in early universe in a more generalised fashion. In this setup, while the possibility of EPR pair creation increases, it appears that the quantum long range correlation will increase naturally. As a consequence, the amount of entanglement entropy increases as the parameter α increases. In this report, we have investigated this possibility with a specific model of axion field theory previously derived from **Type IIB** string theory [43–46] setup. Here, we will demonstrate the Bell's inequality violation from nonzero entanglement entropy of axion field. This connection also will be helpful in future to provide a theoretical tool to compare various models of inflation [47–50]. We have also commented on Rényi entropy using the same setup which will finally give rise to a complete new interpretation to long range quantum correlation for the case of α vacua.

We note that for cosmology, it is crucial to know the observational constraints on the α (non-Bunch Davies) vacua from CMB maps [19,51,52]. It is expected that the (auto and cross) correlation functions of primordial fluctuations get modified significantly, which is an important information to understand the underlying new physics of α vacua. Also this will help us to discriminate between the physical outcomes of α vacua and the adiabatic Bunch Davies vacuum state.

Similarly, for the case of gravity, it is also important to understand the physical implications of the new physics originated from α vacua described in a specific curved gravitational background. Note that Einstein General Theory of relativity is a classical field theoretic description, which describes the interactions in astrophysical scales and constrained by galaxy rotation curves, dynamics of clusters etc. [39,53]. However, in the infrared regime of the gravity sector, we do not

² In the infrared regime, due to the nonremoval of divergences appearing from various interactions in quantum field theory, explaining the physics of excited states with the adiabatic vacuum is not a good approximation.

have observational probes to test the infrared correction to the classical field theory of gravity. From the theoretical perspective, if we describe the fluctuation in the metric in terms of spin 2 transverse, traceless degrees of freedom in de Sitter space-time, then using α vacua and the non local field redefinition in metric, one can express the scalar degrees of freedom also. This scheme needs to be developed in future.

This paper is organised as follows. In [section 2](#), we briefly review the basic set up using which we will compute the entanglement entropy and Rényi entropy using α vacua. In [section 3.1.1](#), we introduce the axion model from string theory. Then using this model we compute the expression for the wave function in a de Sitter hyperbolic open chart in presence of Bunch Davies vacuum in [Appendix A](#). Further using Bogoliubov transformation we express the solution in terms of new basis, called α vacua in [section 3.1.2](#). After that in [section 3.2](#), we construct the density matrix in presence of α vacua. Using this result further in [section 3.3](#), we derive the expression for the von Neumann entropy which is the measure of entanglement entropy in presence of α vacua. Next in [section 3.3](#), we compute Rényi entropy using the result of density matrix as derived in [section 3.2](#). Finally we conclude in [section 4](#) with some future prospects of the present work.

2. Basic setup: brief review

In this section we briefly review the computational method to derive entanglement entropy in de Sitter space following the work performed in ref. [10] and ref. [14]. We consider a time preserving space-like hypersurface S^2 for this purpose. As a result S^2 is divided into two sub regions, interior and exterior, which are characterised by $\text{RI} (\equiv \mathbf{L})$ and $\text{RII} (\equiv \mathbf{R})$. In terms of the Lorentzian signature an open chart in de Sitter space is described by three different subregions. Further [10,14]:

$$\mathbf{R} (\equiv \text{RII}) / \mathbf{L} (\equiv \text{RI}) : \begin{cases} \tau_E = \pi/2 - it_{\mathbf{R}}/\tau_E = -\pi/2 + it_{\mathbf{L}} \\ \rho_E = -ir_{\mathbf{R}/\mathbf{L}} \end{cases} \quad \begin{array}{l} \text{for } t_{\mathbf{R}/\mathbf{L}} \geq 0 \\ \text{for } r_{\mathbf{R}/\mathbf{L}} \geq 0. \end{array} \quad (2.1)$$

$$\mathbf{C} : \begin{cases} \tau_E = t_C \\ \rho_E = \pi/2 - ir_C \end{cases} \quad \begin{array}{l} \text{for } -\pi/2 \leq t_C \leq \pi/2 \\ \text{for } -\infty < r_C < \infty. \end{array} \quad (2.2)$$

Also in open chart the metric with Lorentzian signature can be written as [10,14]:

$$\mathbf{R} (\equiv \text{RII}) / \mathbf{L} = (\text{RI}) : \left\{ ds_{\mathbf{R}/\mathbf{L}}^2 = H^{-2} \left[-dt_{\mathbf{R}/\mathbf{L}}^2 + \sinh^2 t_{\mathbf{R}/\mathbf{L}} \left(dr_{\mathbf{R}/\mathbf{L}}^2 + \sinh^2 r_{\mathbf{R}/\mathbf{L}} d\Omega_2^2 \right) \right], \right. \quad (2.3)$$

$$\mathbf{C} : \left\{ ds_C^2 = H^{-2} \left[dt_C^2 + \cos^2 t_C \left(-dr_C^2 + \cosh^2 r_C d\Omega_2^2 \right) \right], \right. \quad (2.4)$$

where $H = \dot{a}/a$ is the Hubble parameter and $d\Omega_2^2$ represents angular part of the metric on S^2 .

Now let us assume that the total Hilbert space of the local quantum mechanical system is described by \mathcal{H} , which can be written using bipartite decomposition in a direct product space [54] as, $\mathcal{H} = \mathcal{H}_{\text{INT}} \otimes \mathcal{H}_{\text{EXT}}$. Here \mathcal{H}_{INT} and \mathcal{H}_{EXT} are the Hilbert space associated with interior and exterior region and describe the localised modes in RI and RII respectively. Consequently, one can construct the reduced density matrix for the internal RI region by tracing over the external RII region and the Von Neumann entropy measure, the entanglement entropy in de Sitter space can be expressed as:

$$\rho(\alpha) = \text{Tr}_{\mathbf{R}} |\alpha\rangle\langle\alpha| \implies S(\alpha) = -\text{Tr} [\rho(\alpha) \ln \rho(\alpha)]. \quad (2.5)$$

Here the vacuum state $|\alpha\rangle$ is the α vacuum. The reduced density matrix, which is a key ingredient for computing entanglement entropy, is obtained by tracing over the exterior (\mathbf{R}) region. Also it is

important to note that the total entanglement entropy can be expressed as a sum of UV divergent and finite contribution. In 3 + 1 D, the UV-divergent part of the entropy can be written as [10,14,15]:

$$S_{\text{UV-divergent}} = \mathbf{c}_1 \epsilon_{\text{UV}}^{-2} \mathcal{A}_{\text{ENT}} + \left[\mathbf{c}_2 + \left(\mathbf{c}_3 m^2 + \mathbf{c}_4 H^2 \right) \mathcal{A}_{\text{ENT}} \right] \ln(\epsilon_{\text{UV}} H), \quad (2.6)$$

$$S_{\text{UV-finite}} = \mathbf{c}_5 \mathcal{A}_{\text{ENT}} H^2 - \mathbf{c}_6 \ln \left(\sqrt{\mathcal{A}_{\text{ENT}}} H \right) + \text{finite terms}, \quad (2.7)$$

where ϵ_{UV} is the short distance lattice UV cut-off, \mathcal{A}_{ENT} is the proper area of the entangling region of S^2 and $\mathbf{c}_i \forall i = 1, 2, 3, 4$ are the coefficients. Here we restrict ourself only within the UV-finite part which contains the information of long range effects of quantum state. Here \mathbf{c}_6 quantify the long range effect. In general, \mathbf{c}_6 can be expressed as [10,14,15], $\mathbf{c}_6 \equiv S_{\text{intr}}$, where S_{intr} is the UV-finite relevant part which we quantify in later sections.

3. Quantum entanglement for axionic pair using α vacua

3.1. Wave function of axion in open chart

3.1.1. Model for axion effective potential

In this section our prime objective is to compute de Sitter entanglement entropy for axion field. Such axion field appears from RR sector of **Type IIB** string theory compactified on **CY³** in presence of **NS 5** brane. For details, see refs. [43–46,55]. Let us start with the following effective action for axion field:

$$\begin{aligned} S_{\text{axion}} &= \int d^4x \sqrt{-g} \left[-\frac{1}{2} (\partial\phi)^2 + \mu^3 \phi + \Lambda_G^4 \cos \left(\frac{\phi}{f_a} \right) \right] \\ &= \int d^4x \sqrt{-g} \left[-\frac{1}{2} (\partial\phi)^2 + \mu^3 \left[\phi + b f_a \cos \left(\frac{\phi}{f_a} \right) \right] \right], \end{aligned} \quad (3.1)$$

where μ^3 is the mass scale, f_a is the decay constant of axion and we introduce a parameter b , is defined as, $b = \frac{\Lambda_G^4}{\mu^3 f_a}$. Here Λ_G can be expressed as, $\Lambda_G = \sqrt{\frac{m_{\text{SUSY}} L^3}{\sqrt{\alpha'} g_s}} e^{-c S_{\text{inst}}}$, where S_{inst} is the instanton action, factor $c \sim \mathcal{O}(1)$, m_{SUSY} is SUSY breaking scale, α' represents Regge slope parameter, g_s characterises the string coupling constant and L^6 is the world volume factor. Here we restrict up to the linear term of the effective potential as given by $V(\phi) \approx \mu^3 \phi$, which can be interpreted as a massless source in the equation of motion. In the limit $\phi \ll f_a$, the total effective potential for axion can be approximated as, $V(\phi) \approx \mu^3 (b f_a + \phi) - \frac{m_{\text{axion}}^2}{2} \phi^2$, where we introduce the effective mass of the axion as, $m_{\text{axion}}^2 = \frac{\mu^3 b}{f_a} = \frac{\Lambda_G^4}{f_a^2}$.

3.1.2. Wave function for axion using α vacua

Here our prime objective is to derive results for α -vacua, which can be interpreted as a quantum state filled with particles defined by some hypothetical observer who initially belongs to the Bunch Davies vacuum state ($\alpha = 0$). Here the α vacua are invariant under **SO(1, 4)** isometry group of de Sitter space. Consequently we use the equivalent prescription followed in case of Bunch Davies vacuum by defining two subspaces in de Sitter space, **RI** and **RII** respectively. In general α -vacua is CPT invariant, which is parametrised by a single real positive parameter α which plays the role of super-selection quantum number. We use the results obtained for the

solution of the EOM where we expand the field in terms of creation and annihilation operators in Bunch Davies vacuum, and further using Bogoliubov transformation the mode functions for the α -vacua can be written as:

$$\Phi(r, t, \theta, \phi) = \int_0^\infty dp \sum_{\sigma=\pm 1} \sum_{l=0}^{\infty} \sum_{m=-l}^{+l} \left[d_{\sigma plm} \mathcal{E}_{\sigma plm}^{(\alpha)}(r, t, \theta, \phi) + d_{\sigma plm}^\dagger (\mathcal{E}_{\sigma plm}^{(\alpha)})^*(r, t, \theta, \phi) \right], \quad (3.2)$$

where the α -vacua state are defined as, $d_{\sigma plm} |\alpha\rangle = 0 \forall \sigma = (+1, -1); 0 < p < \infty; l = 0, \dots, \infty, m = -l, \dots, +l$. In this context, the α -vacua mode function $\mathcal{E}_{\sigma plm}^{(\alpha)}$ can be expressed in terms of Bunch Davies mode function $\mathcal{U}_{\sigma plm}(r, t, \theta, \phi)$ using Bogoliubov transformation as:

$$\mathcal{E}_{\sigma plm}^{(\alpha)} = \left[\cosh \alpha \mathcal{U}_{\sigma plm}(r, t, \theta, \phi) + \sinh \alpha \mathcal{U}_{\sigma plm}^*(r, t, \theta, \phi) \right]. \quad (3.3)$$

After substituting Eq (3.3) in Eq (3.2) we get the following expression for the wave function:

$$\begin{aligned} \Phi(r, t, \theta, \phi) = & \frac{H}{\sinh t} \int_0^\infty dp \sum_{l=0}^{\infty} \sum_{m=-l}^{+l} \sum_{\sigma=\pm 1} \left[d_{\sigma plm} \cosh \alpha \chi_{p,\sigma}(t) \right. \\ & \left. + d_{\sigma plm}^\dagger \sinh \alpha \chi_{p,\sigma}^*(t) \right] \mathcal{Y}_{plm}(r, \theta, \phi). \end{aligned} \quad (3.4)$$

Finally, the solution of the time dependent part of the wave function can be recast as:

$$\begin{aligned} \chi_{p,\sigma}(t) = & \sum_{q=\mathbf{R}, \mathbf{L}} \left\{ \frac{1}{\mathcal{N}_p} \left[\alpha_q^\sigma \mathcal{P}^q + \beta_q^\sigma \mathcal{P}^{q*} \right] \right. \\ & \left. + \sum_{n=0}^{\infty} \frac{1}{\mathcal{N}_{p_n} (p_n^2 - p^2)} \left[\bar{\alpha}_{q,n}^\sigma \bar{\mathcal{P}}^{q,n} + \bar{\beta}_{q,n}^\sigma \bar{\mathcal{P}}^{q*,n} \right] \right\}, \end{aligned} \quad (3.5)$$

where we use the following shorthand notation, $\bar{\mathcal{P}}^{q,n} = \mu^3 \sinh^2 t \int dt' \chi_{p_n, \sigma, q}^{(c)}(t') \mathcal{P}^{q,n}$. Additionally, here we use the shorthand notations $\mathcal{P}^q, \mathcal{P}^{*q}, \mathcal{P}^{q,n}, \mathcal{P}^{*q,n}$ for the Legendre polynomial, which is defined in ref. [14]. Also the coefficient functions $(\alpha_q^\sigma, \beta_q^\sigma)$ and $(\alpha_{q,n}^\sigma, \beta_{q,n}^\sigma)$, normalisation constants $\mathcal{N}_p, \mathcal{N}_{p_n}$ are explicitly mentioned in ref. [14].

For further computation α -vacua are defined in terms of Bunch Davies vacuum state as:

$$\begin{aligned} |\alpha\rangle = & \exp \left(\frac{1}{2} \tanh \alpha \sum_{\sigma=\pm 1} a_\sigma^\dagger a_\sigma \right) \\ & \times \exp \left(\frac{1}{2} \sum_{i,j=\mathbf{R}, \mathbf{L}} m_{ij} b_i^\dagger b_j^\dagger + \frac{1}{2} \sum_{i,j=\mathbf{R}, \mathbf{L}} \sum_{n=0}^{\infty} \bar{m}_{ij,n} \bar{b}_{i,n}^\dagger \bar{b}_{j,n}^\dagger \right) (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle). \end{aligned} \quad (3.6)$$

Further one can also write the \mathbf{R} and \mathbf{L} vacua as [14]:

$$|\mathbf{R}\rangle = |\mathbf{R}\rangle_{(c)} + \sum_{n=0}^{\infty} |\mathbf{R}\rangle_{(p),n}, \quad |\mathbf{L}\rangle = |\mathbf{L}\rangle_{(c)} + \sum_{n=0}^{\infty} |\mathbf{L}\rangle_{(p),n}, \quad (3.7)$$

with (c) and (p) representing the complementary and particular part respectively. Here the matrices m_{ij} and $\bar{m}_{ij,n}$ corresponding to complementary and particular part of the solution are

explicitly computed in ref. [14] for Bunch Davies vacuum. Also the creation and annihilation operators a_σ^\dagger and a_σ for the **R** and **L** vacuum are defined as [14]:

$$a_\sigma = \sum_{q=\mathbf{R},\mathbf{L}} \left\{ \left[\gamma_{q\sigma} b_q + \delta_{q\sigma}^* b_q^\dagger \right] + \sum_{n=0}^{\infty} \left[\bar{\gamma}_{q\sigma,n} \bar{b}_{q,n} + \bar{\delta}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger \right] \right\}, \quad (3.8)$$

$$a_\sigma^\dagger = \sum_{q=\mathbf{R},\mathbf{L}} \left\{ \left[\gamma_{q\sigma}^* b_q^\dagger + \delta_{q\sigma} b_q \right] + \sum_{n=0}^{\infty} \left[\bar{\gamma}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger + \bar{\delta}_{q\sigma,n} \bar{b}_{q,n} \right] \right\}, \quad (3.9)$$

with $\sigma = \pm 1$. Here it is important to note that, the coefficient matrices for the Bogoliubov transformation $\gamma_{q\sigma}$, $\delta_{q\sigma}$, $\bar{\gamma}_{q\sigma,n}$ and $\bar{\delta}_{q\sigma,n}$ helps us to write the a type of oscillators in terms of a new b type of oscillators. For more details see ref. [14] where all the symbols are explicitly defined. Here it is important to note that, the newly introduced b type of oscillators exactly satisfy the harmonic oscillator algebra, provided the oscillators corresponding to the solution of complementary and particular part of the time dependent solution of the wave function are not interacting with each other. This surely helps us to set up the rules for the operation of creation and annihilation operators of these oscillators in this context [14].

Below, we use the definition of α -vacuum state as given in Eq (3.6), which is very useful to compute entanglement entropy in de Sitter space. However, note that the technical steps for the computation of the entanglement entropy in de Sitter space from α -vacua are exactly similar to the steps followed for Bunch Davies vacuum. The difference will only appear when we use the creation and annihilation operators in the context of α -vacua, which can be written in terms of the creation and annihilation operators defined for **R** or **L** vacuum state as:

$$\begin{aligned} d_\sigma = & \sum_{q=\mathbf{R},\mathbf{L}} \left\{ \left[(\cosh \alpha \gamma_{q\sigma} - \sinh \alpha \delta_{q\sigma}) b_q + (\cosh \alpha \delta_{q\sigma}^* - \sinh \alpha \gamma_{q\sigma}^*) b_q^\dagger \right] \right. \\ & + \left[\left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n} \bar{b}_{q,n} - \sinh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n} \bar{b}_{q,n} \right) \right. \\ & \left. \left. + \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger \right) \right] \right\}, \end{aligned} \quad (3.10)$$

$$\begin{aligned} d_\sigma^\dagger = & \sum_{q=\mathbf{R},\mathbf{L}} \left\{ \left[(\cosh \alpha \gamma_{q\sigma}^* - \sinh \alpha \delta_{q\sigma}^*) b_q^\dagger + (\cosh \alpha \delta_{q\sigma} - \sinh \alpha \gamma_{q\sigma}) b_q \right] \right. \\ & + \left[\left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger \right) \right. \\ & \left. \left. + \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n} \bar{b}_{q,n} - \sinh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n} \bar{b}_{q,n} \right) \right] \right\}, \end{aligned} \quad (3.11)$$

where we use the definition of creation and annihilation operators in Bunch Davies vacuum as mentioned in Eq (3.9) and Eq (3.8). In this computation it is important to note that, under Bogoliubov transformation the original matrix $\gamma_{q\sigma}$, $\delta_{q\sigma}$, $\bar{\gamma}_{q\sigma,n}$ and $\bar{\delta}_{q\sigma,n}$ used for Bunch Davies vacuum are transformed in the context of α -vacua as:

$$\gamma_{q\sigma} \rightarrow (\cosh \alpha \gamma_{q\sigma} - \sinh \alpha \delta_{q\sigma}), \quad \delta_{q\sigma} \rightarrow (\cosh \alpha \delta_{q\sigma} - \sinh \alpha \gamma_{q\sigma}), \quad (3.12)$$

$$\bar{\gamma}_{q\sigma,n} \rightarrow (\cosh \alpha \bar{\gamma}_{q\sigma,n} - \sinh \alpha \bar{\delta}_{q\sigma,n}), \quad \bar{\delta}_{q\sigma,n} \rightarrow (\cosh \alpha \bar{\delta}_{q\sigma,n} - \sinh \alpha \bar{\gamma}_{q\sigma,n}).$$

Considering this fact, after Bogoliubov transformation α -vacua state can be written in terms of \mathbf{R} and \mathbf{L} vacua as:

$$|\alpha\rangle = \exp\left(\frac{1}{2} \sum_{i,j=\mathbf{R},\mathbf{L}} \tilde{m}_{ij} b_i^\dagger b_j^\dagger + \frac{1}{2} \sum_{i,j=\mathbf{R},\mathbf{L}} \sum_{n=0}^{\infty} \tilde{\bar{m}}_{ij,n} \bar{b}_{i,n}^\dagger \bar{b}_{j,n}^\dagger\right) (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle). \quad (3.13)$$

Here \tilde{m}_{ij} and $\tilde{\bar{m}}_{ij,n}$ represents the entries of the matrices corresponding to the complementary and particular solution in presence of α vacuum which we will compute in this paper.

Further one can write the annihilation of α vacuum in terms of the annihilations of the direct product state of \mathbf{R} and \mathbf{L} vacuum as:

$$d_\sigma |\alpha\rangle = \sum_{q=\mathbf{R},\mathbf{L}} \sum_{s=1}^4 \mathcal{J}_s^{(q)} = 0, \quad (3.14)$$

where neglecting contribution from the powers of creation operators, $\mathcal{J}_s^{(q)} \forall s = 1, 2, 3, 4, q = \mathbf{R}, \mathbf{L}$ are defined as:

$$\begin{aligned} \sum_{q=\mathbf{R},\mathbf{L}} \mathcal{J}_1^{(q)} &= \sum_{q=\mathbf{R},\mathbf{L}} (\cosh \alpha \gamma_{q\sigma} - \sinh \alpha \delta_{q\sigma}) b_q e^{\hat{\phi}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) \\ &\approx \sum_{i,j=\mathbf{R},\mathbf{L}} \tilde{m}_{ij} (\cosh \alpha \gamma_{j\sigma} - \sinh \alpha \delta_{j\sigma}) b_i^\dagger (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle), \end{aligned} \quad (3.15)$$

$$\begin{aligned} \sum_{q=\mathbf{R},\mathbf{L}} \mathcal{J}_2^{(q)} &= \sum_{q=\mathbf{R},\mathbf{L}} (\cosh \alpha \delta_{q\sigma}^* - \sinh \alpha \gamma_{q\sigma}^*) b_q^\dagger e^{\hat{\phi}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) \\ &\approx \sum_{q=\mathbf{R},\mathbf{L}} (\cosh \alpha \delta_{q\sigma}^* - \sinh \alpha \gamma_{q\sigma}^*) b_q^\dagger (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle), \end{aligned} \quad (3.16)$$

$$\begin{aligned} \sum_{q=\mathbf{R},\mathbf{L}} \mathcal{J}_3^{(q)} &= \sum_{q=\mathbf{R},\mathbf{L}} \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n} \bar{b}_{q,n} - \sinh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n} \bar{b}_{q,n} \right) e^{\hat{\phi}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) \\ &\approx \sum_{i,j=\mathbf{R},\mathbf{L}} \left(\cosh \alpha \sum_{n=0}^{\infty} \tilde{m}_{ij,n} \bar{\gamma}_{j\sigma,n} \bar{b}_{i,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \tilde{\bar{m}}_{ij,n} \bar{\delta}_{j\sigma,n} \bar{b}_{i,n}^\dagger \right) (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle), \end{aligned} \quad (3.17)$$

$$\begin{aligned} \sum_{q=\mathbf{R},\mathbf{L}} \mathcal{J}_4^{(q)} &= \sum_{q=\mathbf{R},\mathbf{L}} \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger \right) e^{\hat{\phi}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) \\ &\approx \sum_{q=\mathbf{R},\mathbf{L}} \sum_{n=0}^{\infty} \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{q\sigma,n}^* \bar{b}_{q,n}^\dagger \right) (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle). \end{aligned} \quad (3.18)$$

This directly implies that:

$$\begin{aligned} &[\tilde{m}_{ij} (\cosh \alpha \gamma_{j\sigma} - \sinh \alpha \delta_{j\sigma}) + (\cosh \alpha \delta_{i\sigma}^* - \sinh \alpha \gamma_{i\sigma}^*)] b_i^\dagger \\ &+ \left[\left(\cosh \alpha \sum_{n=0}^{\infty} \tilde{m}_{ij,n} \bar{\gamma}_{j\sigma,n} \bar{b}_{i,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \tilde{\bar{m}}_{ij,n} \bar{\delta}_{j\sigma,n} \bar{b}_{i,n}^\dagger \right) \right. \end{aligned}$$

$$+ \left(\cosh \alpha \sum_{n=0}^{\infty} \bar{\delta}_{i\sigma,n}^* \bar{b}_{i,n}^\dagger - \sinh \alpha \sum_{n=0}^{\infty} \bar{\gamma}_{i\sigma,n}^* \bar{b}_{i,n}^\dagger \right) = 0. \quad (3.19)$$

As we have already mentioned that the complementary and particular part of the solutions are completely independent of each other and hence vanish individually. Consequently, we get the following constraints in case of α vacuum:

$$[\tilde{m}_{ij} (\cosh \alpha \gamma_{j\sigma} - \sinh \alpha \delta_{j\sigma}) + (\cosh \alpha \delta_{i\sigma}^* - \sinh \alpha \gamma_{i\sigma}^*)] = 0, \quad (3.20)$$

$$[(\cosh \alpha \tilde{m}_{ij,n} \bar{\gamma}_{j\sigma,n} - \sinh \alpha \tilde{m}_{ij,n} \bar{\delta}_{j\sigma,n}) + (\cosh \alpha \bar{\delta}_{i\sigma,n}^* - \sinh \alpha \bar{\gamma}_{i\sigma,n}^*)] = 0 \forall n. \quad (3.21)$$

Further using Eq (3.20) and Eq (3.21), the matrices corresponding to the complementary and particular part of the solution can be expressed as:

$$\begin{aligned} \tilde{m}_{ij} &= -(\cosh \alpha \delta_{i\sigma}^* - \sinh \alpha \gamma_{i\sigma}^*) (\cosh \alpha \gamma - \sinh \alpha \delta)_{\sigma j}^{-1} \\ &= -\frac{\Gamma(v + \frac{1}{2} - ip)}{\Gamma(v + \frac{1}{2} + ip)} \frac{2 \mathcal{D}_{ij}^{(v)}}{e^{2\pi p} (\cosh \alpha - \sinh \alpha e^{-2\pi p})^2 + e^{2i\pi v} (\cosh \alpha + \sinh \alpha e^{-2i\pi v})^2}, \end{aligned} \quad (3.22)$$

$$\begin{aligned} \tilde{m}_{ij,n} &= -(\cosh \alpha \bar{\delta}_{i\sigma,n}^* - \sinh \alpha \bar{\gamma}_{i\sigma,n}^*) (\cosh \alpha \bar{\gamma} - \sinh \alpha \bar{\delta})_{\sigma j,n}^{-1} \\ &= -\frac{\Gamma(v + \frac{1}{2} - ip_n)}{\Gamma(v + \frac{1}{2} + ip_n)} \\ &\times \frac{2 \mathcal{D}_{ij}^{(v,n)}}{e^{2\pi p_n} (\cosh \alpha - \sinh \alpha e^{-2\pi p_n})^2 + e^{2i\pi v} (\cosh \alpha + \sinh \alpha e^{-2i\pi v})^2} \\ &\forall (i, j) = \mathbf{R}, \mathbf{L}. \end{aligned} \quad (3.23)$$

Here we define the \mathcal{D} matrices as:

$$\mathcal{D}_{ij}^{(v)} = \begin{pmatrix} \mathcal{D}_{\mathbf{RR}}^{(v)} & \mathcal{D}_{\mathbf{RL}}^{(v)} \\ \mathcal{D}_{\mathbf{LR}}^{(v)} & \mathcal{D}_{\mathbf{LL}}^{(v)} \end{pmatrix}, \quad \mathcal{D}_{ij}^{(v,n)} = \begin{pmatrix} \mathcal{D}_{\mathbf{RR}}^{(v,n)} & \mathcal{D}_{\mathbf{RL}}^{(v,n)} \\ \mathcal{D}_{\mathbf{LR}}^{(v,n)} & \mathcal{D}_{\mathbf{LL}}^{(v,n)} \end{pmatrix}, \quad (3.24)$$

and the corresponding entries of the \mathcal{D} matrices are given by:

$$\mathcal{D}_{\mathbf{RR}}^{(v)} = \mathcal{D}_{\mathbf{LL}}^{(v)} = (\cosh^2 \alpha e^{i\pi v} + \sinh^2 \alpha e^{-i\pi v}) \cos \pi v - \sinh 2\alpha \sinh^2 \pi p, \quad (3.25)$$

$$\mathcal{D}_{\mathbf{RL}}^{(v)} = \mathcal{D}_{\mathbf{LR}}^{(v)} = i (\cosh^2 \alpha e^{i\pi v} + \sinh^2 \alpha e^{-i\pi v} + \sinh 2\alpha \cos \pi v) \sinh \pi p, \quad (3.26)$$

$$\mathcal{D}_{\mathbf{RR}}^{(v,n)} = \mathcal{D}_{\mathbf{LL}}^{(v,n)} = (\cosh^2 \alpha e^{i\pi v} + \sinh^2 \alpha e^{-i\pi v}) \cos \pi v - \sinh 2\alpha \sinh^2 \pi p_n, \quad (3.27)$$

$$\mathcal{D}_{\mathbf{RL}}^{(v,n)} = \mathcal{D}_{\mathbf{LR}}^{(v,n)} = i (\cosh^2 \alpha e^{i\pi v} + \sinh^2 \alpha e^{-i\pi v} + \sinh 2\alpha \cos \pi v) \sinh \pi p_n. \quad (3.28)$$

Before further discussion here we point out few important features from the obtained results:

- We see that for the complementary and particular part of the solution

$$\begin{aligned} \tilde{m}_{\mathbf{RR}} &= \tilde{m}_{\mathbf{LL}} \\ &= -\frac{\Gamma(v + \frac{1}{2} - ip)}{\Gamma(v + \frac{1}{2} + ip)} \end{aligned}$$

$$\times \frac{2 \left[(\cosh^2 \alpha e^{i\pi\nu} + \sinh^2 \alpha e^{-i\pi\nu}) \cos \pi\nu - \sinh 2\alpha \sinh^2 \pi p \right]}{e^{2\pi p} (\cosh \alpha - \sinh \alpha e^{-2\pi p})^2 + e^{2i\pi\nu} (\cosh \alpha + \sinh \alpha e^{-2i\pi\nu})^2}, \quad (3.29)$$

$$\begin{aligned} \tilde{m}_{\mathbf{RR},n} &= \tilde{m}_{\mathbf{LL},n} \\ &= -\frac{\Gamma(\nu + \frac{1}{2} - ip_n)}{\Gamma(\nu + \frac{1}{2} + ip_n)} \\ &\times \frac{2 \left[(\cosh^2 \alpha e^{i\pi\nu} + \sinh^2 \alpha e^{-i\pi\nu}) \cos \pi\nu - \sinh 2\alpha \sinh^2 \pi p_n \right]}{e^{2\pi p_n} (\cosh \alpha - \sinh \alpha e^{-2\pi p_n})^2 + e^{2i\pi\nu} (\cosh \alpha + \sinh \alpha e^{-2i\pi\nu})^2}, \end{aligned} \quad (3.30)$$

which is non vanishing for $0 < \nu \leq 3/2$ and $\nu > 3/2$. For $\nu = 3/2$ we get the non vanishing result using α -vacuum and this result is significantly different from the result obtained for Bunch Davies vacuum state.

Finally to implement numerical analysis we use the following approximated expressions for the entries of the coefficient matrices as given by³:

$$\begin{aligned} \tilde{m}_{\mathbf{RR}} &= e^{i\theta} \frac{\sqrt{2} e^{-p\pi} \cos \pi\nu}{\sqrt{\cosh 2\pi p + \cos 2\pi\nu}} \\ &\times \frac{\left[(\cosh^2 \alpha + \sinh^2 \alpha e^{-2i\pi\nu}) - \sinh 2\alpha \sinh^2 \pi p e^{-i\pi\nu} \sec \pi\nu \right]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p+i\nu)})}, \end{aligned} \quad (3.31)$$

$$\begin{aligned} \tilde{m}_{\mathbf{RR},n} &= e^{i\theta} \frac{\sqrt{2} e^{-p_n\pi} \cos \pi\nu}{\sqrt{\cosh 2\pi p_n + \cos 2\pi\nu}} \\ &\times \frac{\left[(\cosh^2 \alpha + \sinh^2 \alpha e^{-2i\pi\nu}) - \sinh 2\alpha \sinh^2 \pi p_n e^{-i\pi\nu} \sec \pi\nu \right]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p_n+i\nu)})}. \end{aligned} \quad (3.32)$$

- We see that for the complementary and particular part of the solution:

$$\begin{aligned} \tilde{m}_{\mathbf{RL}} &= \tilde{m}_{\mathbf{LR}} \\ &= -\frac{\Gamma(\nu + \frac{1}{2} - ip)}{\Gamma(\nu + \frac{1}{2} + ip)} \\ &\times \frac{2 i \left[(\cosh^2 \alpha e^{i\pi\nu} + \sinh^2 \alpha e^{-i\pi\nu} + \sinh 2\alpha \cos \pi\nu) \sinh \pi p \right]}{e^{2\pi p} (\cosh \alpha - \sinh \alpha e^{-2\pi p})^2 + e^{2i\pi\nu} (\cosh \alpha + \sinh \alpha e^{-2i\pi\nu})^2}, \end{aligned} \quad (3.33)$$

$$\begin{aligned} \tilde{m}_{\mathbf{RL},n} &= \tilde{m}_{\mathbf{LR},n} \\ &= -\frac{\Gamma(\nu + \frac{1}{2} - ip)}{\Gamma(\nu + \frac{1}{2} + ip)} \\ &\times \frac{2 i \left[(\cosh^2 \alpha e^{i\pi\nu} + \sinh^2 \alpha e^{-i\pi\nu} + \sinh 2\alpha \cos \pi\nu) \sinh \pi p_n \right]}{e^{2\pi p_n} (\cosh \alpha - \sinh \alpha e^{-2\pi p_n})^2 + e^{2i\pi\nu} (\cosh \alpha + \sinh \alpha e^{-2i\pi\nu})^2}. \end{aligned} \quad (3.34)$$

³ For rest of the analysis we absorb this overall phase factor $e^{i\theta}$.

Additionally, the non vanishing entries of the off diagonal components of the coefficient matrix for both of the cases in presence of α -vacuum indicates the existence of quantum entanglement in the present computation, which we will explicitly show that finally give rise to a non vanishing entanglement entropy.

Finally to interpret the result numerically we use the following approximated expressions for the entries of the coefficient matrices as given by:

$$\tilde{m}_{\mathbf{RL}} = e^{i(\theta+\frac{\pi}{2})} \frac{\sqrt{2} e^{-p\pi} \sinh \pi p}{\sqrt{\cosh 2\pi p + \cos 2\pi v}} \times \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{-2i\pi v} + \sinh 2\alpha \cos \pi v e^{-i\pi v}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p+iv)})}, \quad (3.35)$$

$$\tilde{m}_{\mathbf{RL},n} = e^{i(\theta+\frac{\pi}{2})} \frac{\sqrt{2} e^{-p_n\pi} \sinh \pi p_n}{\sqrt{\cosh 2\pi p_n + \cos 2\pi v}} \times \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{-2i\pi v} + \sinh 2\alpha \cos \pi v e^{-i\pi v}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p_n+iv)})}. \quad (3.36)$$

To find a suitable basis first of all we trace over all possible contributions from **R** and **L** region. To implement this we need to perform another Bogoliubov transformation introducing new sets of operators as given by:

$$\begin{aligned} \tilde{c}_{\mathbf{R}} &= \tilde{u} b_{\mathbf{R}} + \tilde{v} b_{\mathbf{R}}^\dagger, \quad \tilde{c}_{\mathbf{L}} = \tilde{u} b_{\mathbf{L}} + \tilde{v} b_{\mathbf{L}}^\dagger, \\ \tilde{c}_{\mathbf{R},n} &= \tilde{U}_n b_{\mathbf{R},n} + \tilde{V}_n b_{\mathbf{R},n}^\dagger, \quad \tilde{c}_{\mathbf{L},n} = \tilde{U}_n b_{\mathbf{L},n} + \tilde{V}_n b_{\mathbf{L},n}^\dagger, \end{aligned} \quad (3.37)$$

where following conditions are satisfied:

$$|\tilde{u}|^2 - |\tilde{v}|^2 = 1, \quad |\tilde{u}|^2 - |\tilde{v}|^2 = 1, \quad |\tilde{U}_n|^2 - |\tilde{V}_n|^2 = 1, \quad |\tilde{U}_n|^2 - |\tilde{V}_n|^2 = 1. \quad (3.38)$$

Using these new sets of operators one can write the α -vacuum state in terms of new basis represented by the direct product of **R'** and **L'** vacuum state as:

$$|\alpha\rangle = e^{\hat{Q}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) = (\tilde{\mathcal{N}}_p^{(\alpha)})^{-1} e^{\hat{Q}} (|\mathbf{R}'\rangle \otimes |\mathbf{L}'\rangle)^{(\alpha)}, \quad (3.39)$$

where we introduce a new composite operator \hat{Q} which is defined in the new transformed basis as:

$$\hat{Q} = \gamma_p^{(\alpha)} \tilde{c}_{\mathbf{R}}^\dagger \tilde{c}_{\mathbf{L}}^\dagger + \sum_{n=0}^{\infty} \Gamma_{p,n}^{(\alpha)} \tilde{C}_{\mathbf{R},n}^\dagger \tilde{C}_{\mathbf{L},n}^\dagger, \quad (3.40)$$

where $\gamma_p^{(\alpha)}$ and $\Gamma_{p,n}^{(\alpha)}$ are defined corresponding to the complementary and particular solution, which we will explicitly compute further for α vacuum. Additionally, it is important to note that the overall normalisation factor $\tilde{\mathcal{N}}_p^{(\alpha)}$ is defined as:

$$\tilde{\mathcal{N}}_p^{(\alpha)} = \left| e^{\hat{Q}} (|\mathbf{R}'\rangle \otimes |\mathbf{L}'\rangle)^{(\alpha)} \right| \approx \left[1 - \left(|\gamma_p^{(\alpha)}|^2 + \sum_{n=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^2 \right) \right]^{-1/2}, \quad (3.41)$$

which reduces to the result obtained for Bunch Davies vacuum in ref. [14] for $\alpha = 0$. In this calculation due to the second Bogoliubov transformation the direct product of the **R** and **L** vacuum state is connected to the direct product of the new **R'** and **L'** vacuum state as:

$$(|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle) \rightarrow (|\mathbf{R}'\rangle \otimes |\mathbf{L}'\rangle)^{(\alpha)} = \tilde{\mathcal{N}}_p^{(\alpha)} e^{-\hat{Q}} e^{\hat{O}} (|\mathbf{R}\rangle \otimes |\mathbf{L}\rangle). \quad (3.42)$$

Let us now mention the commutation relations of the creation and annihilation operators corresponding to the new sets of oscillators describing the \mathbf{R}' and \mathbf{L}' vacuum state as:

$$\begin{aligned} [\tilde{c}_i, \tilde{c}_j^\dagger] &= \delta_{ij}, \quad [\tilde{c}_i, \tilde{c}_j] = 0 = [\tilde{c}_i^\dagger, \tilde{c}_j^\dagger], \\ [\tilde{C}_{i,n}, \tilde{C}_{j,m}^\dagger] &= \delta_{ij} \delta_{nm}, \quad [\tilde{C}_{i,n}, \tilde{C}_{j,m}] = 0 = [\tilde{C}_{i,m}^\dagger, \tilde{C}_{j,m}^\dagger]. \end{aligned} \quad (3.43)$$

Here, for α vacuum, the oscillator algebra is exactly same as that obtained for Bunch Davies vacuum. However for α vacuum the structure of these operators are completely different and also they are acting in a different Hilbert space \mathcal{H}_α , which is characterised by one parameter α . Here it is important to note that for $\alpha = 0$ these oscillators will act on Bunch Davies vacuum state where the corresponding Hilbert space, \mathcal{H}_{BD} is the subclass of \mathcal{H}_α .

The action of creation and annihilation operators defined on the α vacuum state are appended below:

$$\begin{aligned} \tilde{c}_{\mathbf{R}}|\alpha\rangle &= \gamma_p^{(\alpha)} \tilde{c}_{\mathbf{L}}^\dagger|\alpha\rangle, \quad \tilde{c}_{\mathbf{L}}|\alpha\rangle = \gamma_p^{(\alpha)} \tilde{c}_{\mathbf{R}}^\dagger|\alpha\rangle, \\ \tilde{C}_{\mathbf{R},n}|\alpha\rangle &= \Gamma_{p,n}^{(\alpha)} \tilde{C}_{\mathbf{L},n}^\dagger|\alpha\rangle, \quad \tilde{C}_{\mathbf{L},n}|\alpha\rangle = \Gamma_{p,n}^{(\alpha)} \tilde{C}_{\mathbf{R},n}^\dagger|\alpha\rangle. \end{aligned} \quad (3.44)$$

Further, one can express the new c type annihilation operators in terms of the old b type annihilation operators as:

$$\tilde{c}_J = b_I \tilde{\mathcal{G}}_J^I = b_I \begin{pmatrix} \tilde{U}_q & \tilde{V}_q^* \\ \tilde{V}_q & \tilde{U}_q^* \end{pmatrix}, \quad \tilde{C}_{J(n)} = \bar{b}_{J(n)} (\tilde{\mathcal{G}}_{(n)})_J^I = \bar{b}_{J(n)} \begin{pmatrix} \tilde{\bar{U}}_{q,n} & \tilde{\bar{V}}_{\sigma,q,n}^* \\ \tilde{\bar{V}}_{q,n} & \tilde{\bar{U}}_{q,n}^* \end{pmatrix}. \quad (3.45)$$

Here the entries of the matrices for α vacuum are given by, $\tilde{U}_q \equiv \mathbf{diag}(\tilde{u}, \tilde{\bar{u}})$, $\tilde{V}_q \equiv \mathbf{diag}(\tilde{v}, \tilde{\bar{v}})$, $\tilde{\bar{U}}_{q,n} \equiv \mathbf{diag}(\tilde{U}_n, \tilde{\bar{U}}_n)$, $\tilde{\bar{V}}_{q,n} \equiv \mathbf{diag}(\tilde{V}_n, \tilde{\bar{V}}_n)$. Further using Eq (3.37), in Eq (3.44), we get the following sets of homogeneous equations:

For complementary solution :

$$\begin{aligned} \tilde{m}_{\mathbf{RR}} \tilde{u} + \tilde{v} - \gamma_p^{(\alpha)} \tilde{m}_{\mathbf{RL}} \tilde{\bar{v}}^* &= 0, \quad \tilde{m}_{\mathbf{RR}} \tilde{\bar{u}} + \tilde{\bar{v}} - \gamma_p^{(\alpha)} \tilde{m}_{\mathbf{RL}} \tilde{v}^* &= 0, \\ (3.46) \end{aligned}$$

$$\begin{aligned} \tilde{m}_{\mathbf{RL}} \tilde{u} - \gamma_p^{(\alpha)} \tilde{\bar{u}}^* - \gamma_p^{(\alpha)} \tilde{m}_{\mathbf{RR}} \tilde{\bar{v}}^* &= 0, \quad \tilde{m}_{\mathbf{RL}} \tilde{\bar{u}} - \gamma_p^{(\alpha)} \tilde{u}^* - \gamma_p^{(\alpha)} \tilde{m}_{\mathbf{RR}} \tilde{v}^* &= 0, \\ (3.47) \end{aligned}$$

For particular solution :

$$\begin{aligned} \tilde{m}_{\mathbf{RR},n} \tilde{U}_n + \tilde{V}_n - \Gamma_{p,n}^{(\alpha)} \tilde{m}_{\mathbf{RL},n} \tilde{\bar{V}}_n^* &= 0, \quad \tilde{m}_{\mathbf{RR},n} \tilde{\bar{U}}_n + \tilde{\bar{V}}_n - \Gamma_{p,n}^{(\alpha)} \tilde{m}_{\mathbf{RL},n} \tilde{V}_n^* &= 0, \\ (3.48) \end{aligned}$$

$$\begin{aligned} \tilde{m}_{\mathbf{RL},n} \tilde{U}_n - \Gamma_{p,n}^{(\alpha)} \tilde{\bar{U}}_n^* - \Gamma_{p,n}^{(\alpha)} \tilde{m}_{\mathbf{RR},n} \tilde{\bar{V}}_n^* &= 0, \\ \tilde{m}_{\mathbf{RL},n} \tilde{\bar{U}}_n - \Gamma_{p,n}^{(\alpha)} \tilde{U}_n^* - \Gamma_{p,n}^{(\alpha)} \tilde{m}_{\mathbf{RR},n} \tilde{V}_n^* &= 0. \end{aligned} \quad (3.49)$$

Now with α vacuum, it is not sufficient to use $\tilde{v}^* = \tilde{\bar{v}}$, $\tilde{u}^* = \tilde{\bar{u}}$ for particular part and also $\tilde{V}_n^* = \tilde{\bar{V}}_n$, $\tilde{U}_n^* = \tilde{\bar{U}}_n$ for the complementary part. In this case, the system of four equations, each for

complementary and particular part will not be reduced to two sets of simplified equations. This is an outcome of the fact that in case of α vacuum, the entries of the coefficient matrices \tilde{m}_{ij} and $\tilde{m}_{ij,n}$ are complex in nature. On the other hand, they are either real or imaginary for Bunch Davies vacuum state. To solve these equations for $\gamma_p^{(\alpha)}$ and $\Gamma_{p,n}^{(\alpha)}$, we also need to use the normalisation conditions, $|\tilde{u}|^2 - |\tilde{v}|^2 = 1$ and $|\tilde{U}_n|^2 - |\tilde{V}_n|^2 = 1$.

Finally, the non trivial solutions obtained from these systems of equations can be expressed as:

$$\begin{aligned}\gamma_p^{(\alpha)} &= \frac{1}{\sqrt{2}|\tilde{m}_{\text{RL}}|} \left[\left(1 + |\tilde{m}_{\text{RL}}|^4 + |\tilde{m}_{\text{RR}}|^4 - 2|\tilde{m}_{\text{RR}}|^2 - \tilde{m}_{\text{RR}}^2 (\tilde{m}_{\text{RL}}^*)^2 - \tilde{m}_{\text{RL}}^2 (\tilde{m}_{\text{RR}}^*)^2 \right) \right. \\ &\quad \pm \left\{ \left(-1 - |\tilde{m}_{\text{RL}}|^4 - |\tilde{m}_{\text{RR}}|^4 + 2|\tilde{m}_{\text{RR}}|^2 + \tilde{m}_{\text{RR}}^2 (\tilde{m}_{\text{RL}}^*)^2 + \tilde{m}_{\text{RL}}^2 (\tilde{m}_{\text{RR}}^*)^2 \right)^2 \right. \\ &\quad \left. \left. - 4|\tilde{m}_{\text{RL}}|^4 \right\}^{\frac{1}{2}} \right],\end{aligned}\quad (3.50)$$

$$\begin{aligned}\Gamma_{p,n}^{(\alpha)} &= \frac{1}{\sqrt{2}|\tilde{m}_{\text{RL},n}|} \left[\left(1 + |\tilde{m}_{\text{RL},n}|^4 + |\tilde{m}_{\text{RR},n}|^4 - 2|\tilde{m}_{\text{RR},n}|^2 - \tilde{m}_{\text{RR},n}^2 (\tilde{m}_{\text{RL},n}^*)^2 \right. \right. \\ &\quad \left. \left. - \tilde{m}_{\text{RL},n}^2 (\tilde{m}_{\text{RR},n}^*)^2 \right) \pm \left\{ \left(-1 - |\tilde{m}_{\text{RL},n}|^4 - |\tilde{m}_{\text{RR},n}|^4 + 2|\tilde{m}_{\text{RR},n}|^2 \right. \right. \\ &\quad \left. \left. + \tilde{m}_{\text{RR},n}^2 (\tilde{m}_{\text{RL},n}^*)^2 + \tilde{m}_{\text{RL},n}^2 (\tilde{m}_{\text{RR},n}^*)^2 \right)^2 - 4|\tilde{m}_{\text{RL},n}|^4 \right\}^{\frac{1}{2}} \right],\end{aligned}\quad (3.51)$$

where the components $\tilde{m}_{\text{RR}} = \tilde{m}_{\text{LL}}$, $\tilde{m}_{\text{RL}} = \tilde{m}_{\text{LR}}$ and $\tilde{m}_{\text{RR},n} = \tilde{m}_{\text{LL},n}$, $\tilde{m}_{\text{RL},n} = \tilde{m}_{\text{LR},n}$ are defined in Eqn (3.29), Eqn (3.30) and Eqn (3.33), Eqn (3.34) respectively. Note that, in both the solutions for $\gamma_p^{(\alpha)}$ and $\Gamma_{p,n}^{(\alpha)}$ we absorb the overall phase factor.

After further simplification we get the following expressions for the non trivial solutions for arbitrary ν can be written as:

$$\begin{aligned}\gamma_p^{(\alpha)} &\approx i \frac{\sqrt{2}}{\sqrt{\cosh 2\pi p + \cos 2\pi\nu} \pm \sqrt{\cosh 2\pi p + \cos 2\pi\nu + 2}} \\ &\quad \times \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{2i\pi\nu} + \sinh 2\alpha \cos \pi\nu e^{i\pi\nu}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p-i\nu)})},\end{aligned}\quad (3.52)$$

$$\begin{aligned}\Gamma_{p,n}^{(\alpha)} &\approx i \frac{\sqrt{2}}{\sqrt{\cosh 2\pi p_n + \cos 2\pi\nu} \pm \sqrt{\cosh 2\pi p_n + \cos 2\pi\nu + 2}} \\ &\quad \times \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{2i\pi\nu} + \sinh 2\alpha \cos \pi\nu e^{i\pi\nu}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p_n-i\nu)})}.\end{aligned}\quad (3.53)$$

3.2. Construction of density matrix using α vacua

In this subsection our prime objective is construct the density matrix using the α vacuum state which is expressed in terms of another set of annihilation and creation operators in the Bogoliubov transformed frame. Here the Bunch Davies vacuum state can be expressed as a product of the quantum state for each oscillator in the new frame after Bogoliubov transformation. Each oscillator is labelled by the quantum numbers p, l and m in this calculation. After tracing over

the right part of the Hilbert space we get the following expression for the density matrix and the left part of the Hilbert space can be written as, $(\rho_{\mathbf{L}}(\alpha))_{p,l,m} = \text{Tr}_{\mathbf{R}}|\alpha\rangle\langle\alpha|$, where the α vacuum state can be written in terms of \tilde{c} type of oscillators as:

$$|\alpha\rangle \approx \left[1 - \left(|\gamma_p^{(\alpha)}|^2 + \sum_{n=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^2 \right) \right]^{1/2} \\ \times \exp \left[\gamma_p^{(\alpha)} \tilde{c}_{\mathbf{R}}^\dagger \tilde{c}_{\mathbf{L}}^\dagger + \sum_{n=0}^{\infty} \Gamma_{p,n}^{(\alpha)} \tilde{C}_{\mathbf{R},n}^\dagger \tilde{C}_{\mathbf{L},n}^\dagger \right] (|\mathbf{R}'\rangle \otimes |\mathbf{L}'\rangle)^{(\alpha)}, \quad (3.54)$$

which is already derived in the earlier section. Further using Eq (3.54), we find the following simplified expression for the density matrix for the left part of the Hilbert space for α vacuum as:

$$(\rho_{\mathbf{L}}(\alpha))_{p,l,m} = \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \sum_{k=0}^{\infty} |\gamma_p^{(\alpha)}|^{2k} |\widetilde{k; p, l, m}\rangle \langle \widetilde{k; p, l, m}| \\ + (f_p^{(\alpha)})^2 \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^{2r} |\widetilde{n, r; p, l, m}\rangle \langle \widetilde{n, r; p, l, m}|, \quad (3.55)$$

where $\gamma_p^{(\alpha)}$ and $\Gamma_{p,n}^{(\alpha)}$ are derived in the earlier section. Also we define the α parameter dependent source normalisation factor $f_p^{(\alpha)}$ as, $f_p^{(\alpha)} = \left(\sum_{n=0}^{\infty} (1 - |\Gamma_{p,n}^{(\alpha)}|^2)^{-1} \right)^{-1}$. In this computation the states $|\widetilde{k; p, l, m}\rangle$ and $|\widetilde{n, r; p, l, m}\rangle$ are defined in terms of the quantum state $|\mathbf{L}'\rangle$ as:

$$|\widetilde{k; p, l, m}\rangle = \frac{1}{\sqrt{n!}} (\tilde{c}_{\mathbf{L}}^\dagger)^k |\mathbf{L}'\rangle, \quad |\widetilde{n, r; p, l, m}\rangle = \frac{1}{\sqrt{r!}} (\tilde{C}_{\mathbf{L},n}^\dagger)^r |\mathbf{L}'\rangle. \quad (3.56)$$

Here we note that:

1. For α vacuum density matrix is diagonal for a given set of the **SO(1, 3)** quantum numbers p, l, m and additionally depends on the parameter α explicitly. This leads to the total density matrix to take the following simplified form as:

$$\rho_{\mathbf{L}}(\alpha) = \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \mathbf{diag} \left(1, |\gamma_p^{(\alpha)}|^2, |\gamma_p^{(\alpha)}|^4, |\gamma_p^{(\alpha)}|^6 \dots \right) \\ + (f_p^{(\alpha)})^2 \sum_{n=0}^{\infty} \mathbf{diag} \left(1, |\Gamma_{p,n}^{(\alpha)}|^2, |\Gamma_{p,n}^{(\alpha)}|^4, |\Gamma_{p,n}^{(\alpha)}|^6 \dots \right). \quad (3.57)$$

2. To find out an acceptable normalisation of the total density matrix in presence of α vacuum state, we use the following limiting results:

$$\sum_{k=0}^{\infty} |\gamma_p^{(\alpha)}|^{2k} = \lim_{k \rightarrow \infty} \frac{1 - |\gamma_p^{(\alpha)}|^{2k}}{1 - |\gamma_p^{(\alpha)}|^2} \xrightarrow{|\gamma_p^{(\alpha)}| < 1 \forall \alpha} \frac{1}{1 - |\gamma_p^{(\alpha)}|^2}, \quad (3.58)$$

$$\sum_{n=0}^{\infty} \sum_{r=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^{2r} = \sum_{n=0}^{\infty} \lim_{r \rightarrow \infty} \frac{1 - |\Gamma_{p,n}^{(\alpha)}|^{2r}}{1 - |\Gamma_{p,n}^{(\alpha)}|^2} \xrightarrow{|\Gamma_{p,n}^{(\alpha)}| < 1 \forall n, \alpha} \sum_{n=0}^{\infty} \frac{1}{1 - |\Gamma_{p,n}^{(\alpha)}|^2} = (f_p^{(\alpha)})^{-1}. \quad (3.59)$$

Consequently using these results for α vacuum we get:

$$\begin{aligned} \text{Tr} \left[\left(1 - |\gamma_p^{(\alpha)}|^2 \right) \mathbf{diag} \left(1, |\gamma_p^{(\alpha)}|^2, |\gamma_p^{(\alpha)}|^4, |\gamma_p^{(\alpha)}|^6 \dots \right) \right] \\ = \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \sum_{k=0}^{\infty} |\gamma_p^{(\alpha)}|^{2k} = 1, \end{aligned} \quad (3.60)$$

$$\begin{aligned} \text{Tr} \left[(f_p^{(\alpha)})^2 \sum_{n=0}^{\infty} \mathbf{diag} \left(1, |\Gamma_{p,n}^{(\alpha)}|^2, |\Gamma_{p,n}^{(\alpha)}|^4, |\Gamma_{p,n}^{(\alpha)}|^6 \dots \right) \right] \\ = (f_p^{(\alpha)})^2 \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^{2r} = f_p^{(\alpha)}. \end{aligned} \quad (3.61)$$

Consequently the normalisation condition of this total density matrix in presence of α vacuum state is given by, $\text{Tr} \rho_{\mathbf{L}}(\alpha) = 1 + f_p^{(\alpha)}$. This result is consistent with the ref. [10] where $f_p^{(\alpha)} = 0 \forall \alpha$ and also ref. [14] where $\alpha = 0$ and $f_p^{(0)} = f_p$. But for simplicity it is better to maintain always $\text{Tr} \rho_{\mathbf{L}}(\alpha) = 1$ and to get this result for α vacuum the total density matrix can be redefined by changing the normalisation constant as:

$$\begin{aligned} (\rho_{\mathbf{L}}(\alpha))_{p,l,m} &= \frac{\left(1 - |\gamma_p^{(\alpha)}|^2 \right)}{1 + f_p^{(\alpha)}} \sum_{k=0}^{\infty} |\gamma_p^{(\alpha)}|^{2k} |k; \widetilde{p, l, m}\rangle \langle k; \widetilde{p, l, m}| \\ &\quad + \frac{(f_p^{(\alpha)})^2}{1 + f_p^{(\alpha)}} \sum_{n,r=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^{2r} |n, r; \widetilde{p, l, m}\rangle \langle n, r; \widetilde{p, l, m}|. \end{aligned} \quad (3.62)$$

In this context equivalent convention for normalisation factors can also be chosen such that it always satisfies $\text{Tr} \rho_{\mathbf{L}}(\alpha) = 1$, even in the presence of source contribution.⁴

3. For each set of values of the **SO(1, 3)** quantum numbers p, l, m , the density matrix yields $(\rho_{\mathbf{L}})_{p,l,m}$ and so that the total density matrix can be expressed as a product of all such possible contributions:

$$\rho_{\mathbf{L}}(\alpha) = \prod_{p=0}^{\infty} \prod_{l=0}^{p-1} \prod_{m=-l}^{+l} (\rho_{\mathbf{L}}(\alpha))_{p,l,m}. \quad (3.64)$$

This also indicates that in such a situation entanglement is absent among all states which carries non identical **SO(1, 3)** quantum numbers p, l, m .

4. Finally, the total density matrix can be written in terms of entanglement modular Hamiltonian of the axionic pair as, $\rho_{\mathbf{L}}(\alpha) = e^{-\beta \mathcal{H}_{\text{ENT}}}$, where at finite temperature T_{dS} of de Sitter

⁴ Here one can choose the following equivalent ansatz for total density matrix in presence of α vacuum as:

$$\begin{aligned} (\rho_{\mathbf{L}}(\alpha))_{p,l,m} &= \left[\frac{1}{1 - |\gamma_p^{(\alpha)}|^2} + f_p^{(\alpha)} \right]^{-1} \left[\sum_{k=0}^{\infty} |\gamma_p^{(\alpha)}|^{2k} |k; \widetilde{p, l, m}\rangle \langle k; \widetilde{p, l, m}| \right. \\ &\quad \left. + (f_p^{(\alpha)})^2 \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^{2r} |n, r; \widetilde{p, l, m}\rangle \langle n, r; \widetilde{p, l, m}| \right]. \end{aligned} \quad (3.63)$$

space $\beta = 2\pi/T_{\text{dS}}$. If we further assume that the dynamical Hamiltonian in de Sitter space is represented by entangled Hamiltonian then for a given principal quantum number p the Hamiltonian for axion can be expressed as:

$$\mathcal{H}_p(\alpha) = \left[E_p^{(\alpha)} \tilde{c}_p^\dagger \tilde{c}_p + \sum_{n=0}^{\infty} \mathcal{E}_{p,n}^{(\alpha)} \tilde{C}_{p,n}^\dagger \tilde{C}_{p,n} \right]. \quad (3.65)$$

Acting this Hamiltonian on the α vacuum state we find:

$$\begin{aligned} \mathcal{H}_p(\alpha)|\alpha\rangle &\approx \left[1 - \left(|\gamma_p^{(\alpha)}|^2 + \sum_{n=0}^{\infty} |\Gamma_{p,n}^{(\alpha)}|^2 \right) \right]^{1/2} \left[E_p^{(\alpha)} \tilde{c}_p^\dagger \tilde{c}_p + \sum_{n=0}^{\infty} \mathcal{E}_{p,n}^{(\alpha)} \tilde{C}_{p,n}^\dagger \tilde{C}_{p,n} \right] \\ &\times \exp \left[\gamma_p^{(\alpha)} \tilde{c}_{\mathbf{R}}^\dagger \tilde{c}_{\mathbf{L}}^\dagger + \sum_{m=0}^{\infty} \Gamma_{p,m}^{(\alpha)} \tilde{C}_{\mathbf{R},m}^\dagger \tilde{C}_{\mathbf{L},m}^\dagger \right] (|\mathbf{R}'\rangle \otimes |\mathbf{L}'\rangle)^{(\alpha)} = E_{\mathbf{T},p}^{(\alpha)}|\alpha\rangle, \end{aligned} \quad (3.66)$$

where the total energy spectrum of this system for α vacuum can be written as:

$$\begin{aligned} E_{\mathbf{T},p}^{(\alpha)} &= E_p^{(\alpha)} + \sum_{n=0}^{\infty} \mathcal{E}_{p,n}^{(\alpha)} \quad \forall \alpha , \\ \text{with } E_p^{(\alpha)} &= -\frac{1}{2\pi} \ln(|\gamma_p^{(\alpha)}|^2), \quad \mathcal{E}_{p,n}^{(\alpha)} = -\frac{1}{2\pi} \ln \left(\frac{1}{1 - |\Gamma_{p,n}^{(\alpha)}|^2} \right) \end{aligned} \quad (3.67)$$

where spectrum for the complementary and particular part for α vacuum state is defined by $E_p^{(\alpha)}$ and $\mathcal{E}_{p,n}^{(\alpha)}$.

Now if we consider any arbitrary mass parameter v and any arbitrary value of the parameter α , the **SO(1, 3)** principal quantum number p dependent spectrum can be expressed as:

$$\begin{aligned} E_p^{(\alpha)} &= -\frac{1}{2\pi} \ln \left\{ \frac{1}{2|\tilde{m}_{\mathbf{RL}}|^2} \left[\left(1 + |\tilde{m}_{\mathbf{RL}}|^4 + |\tilde{m}_{\mathbf{RR}}|^4 \right. \right. \right. \\ &\quad \left. \left. \left. - 2|\tilde{m}_{\mathbf{RR}}|^2 - \tilde{m}_{\mathbf{RR}}^2 (\tilde{m}_{\mathbf{RL}}^*)^2 - \tilde{m}_{\mathbf{RL}}^2 (\tilde{m}_{\mathbf{RR}}^*)^2 \right) \pm \left\{ \left(-1 - |\tilde{m}_{\mathbf{RL}}|^4 - |\tilde{m}_{\mathbf{RR}}|^4 \right. \right. \right. \\ &\quad \left. \left. \left. + 2|\tilde{m}_{\mathbf{RR}}|^2 + \tilde{m}_{\mathbf{RR}}^2 (\tilde{m}_{\mathbf{RL}}^*)^2 + \tilde{m}_{\mathbf{RL}}^2 (\tilde{m}_{\mathbf{RR}}^*)^2 \right)^2 - 4|\tilde{m}_{\mathbf{RL}}|^4 \right\}^{\frac{1}{2}} \right] \right\}, \end{aligned} \quad (3.68)$$

$$\begin{aligned} \mathcal{E}_{p,n}^{(\alpha)} &= \frac{1}{2\pi} \ln \left\{ 1 - \frac{1}{2|\tilde{m}_{\mathbf{RL},n}|^2} \left[\left(1 + |\tilde{m}_{\mathbf{RL},n}|^4 + |\tilde{m}_{\mathbf{RR},n}|^4 \right. \right. \right. \\ &\quad \left. \left. \left. - 2|\tilde{m}_{\mathbf{RR},n}|^2 - \tilde{m}_{\mathbf{RR},n}^2 (\tilde{m}_{\mathbf{RL},n}^*)^2 - \tilde{m}_{\mathbf{RL},n}^2 (\tilde{m}_{\mathbf{RR},n}^*)^2 \right) \pm \left\{ \left(-1 - |\tilde{m}_{\mathbf{RL},n}|^4 \right. \right. \right. \\ &\quad \left. \left. \left. - |\tilde{m}_{\mathbf{RR},n}|^4 + 2|\tilde{m}_{\mathbf{RR},n}|^2 + \tilde{m}_{\mathbf{RR},n}^2 (\tilde{m}_{\mathbf{RL},n}^*)^2 + \tilde{m}_{\mathbf{RL},n}^2 (\tilde{m}_{\mathbf{RR},n}^*)^2 \right)^2 \right. \right. \\ &\quad \left. \left. \left. - 4|\tilde{m}_{\mathbf{RL},n}|^4 \right\}^{\frac{1}{2}} \right] \right\}. \end{aligned} \quad (3.69)$$

Here the components $\tilde{m}_{\mathbf{RR}} = \tilde{m}_{\mathbf{LL}}$, $\tilde{m}_{\mathbf{RL}} = \tilde{m}_{\mathbf{LR}}$ and $\tilde{m}_{\mathbf{RR},n} = \tilde{m}_{\mathbf{LL},n}$, $\tilde{m}_{\mathbf{RL},n} = \tilde{m}_{\mathbf{LR},n}$ are defined in Eqn (3.29), Eqn (3.30) and Eqn (3.33), Eqn (3.34) respectively.

Further using Eq (3.50) and Eq (3.51), we get the following simplified expressions:

$$\begin{aligned} E_p^{(\alpha)} &\approx -\frac{1}{2\pi} \ln \left[\frac{2}{(\sqrt{\cosh 2\pi p + \cos 2\pi v} \pm \sqrt{\cosh 2\pi p + \cos 2\pi v + 2})^2} \right. \\ &\quad \times \left. \left| \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{2i\pi v} + \sinh 2\alpha \cos \pi v e^{i\pi v}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p-i)v})} \right|^2 \right], \\ \mathcal{E}_{p,n}^{(\alpha)} &\approx \frac{1}{2\pi} \ln \left[1 - \frac{2}{(\sqrt{\cosh 2\pi p_n + \cos 2\pi v} \pm \sqrt{\cosh 2\pi p_n + \cos 2\pi v + 2})^2} \right. \\ &\quad \times \left. \left| \frac{[\cosh^2 \alpha + \sinh^2 \alpha e^{2i\pi v} + \sinh 2\alpha \cos \pi v e^{i\pi v}]}{(\cosh^2 \alpha + \sinh^2 \alpha e^{-2\pi(p_n-i)v})} \right|^2 \right]. \end{aligned} \quad (3.70)$$

These results imply that for arbitrary parameter v and α the entangled Hamiltonian (\mathcal{H}_{ENT}) and the Hamiltonian for axion (\mathcal{H}_p) _{$\mathbf{R} \times \mathbf{H}^3$} are significantly different, compared to the result obtained in absence of linear source term in case of Bunch Davies vacuum.

3.3. Computation of entanglement entropy using α vacua

In this subsection our prime objective is to derive the expression for entanglement entropy in de Sitter space in presence of α vacuum state. In general the entanglement entropy with arbitrary α can be expressed as:

$$S(p, v, \alpha) = -\mathbf{Tr} [\rho_{\mathbf{L}}(p, \alpha) \ln \rho_{\mathbf{L}}(p, \alpha)], \quad (3.71)$$

where the parameter v and the corresponding α vacuum state are defined in the earlier section. In this context the expression for entanglement entropy for a given $\mathbf{SO}(1, 3)$ principal quantum number p can be expressed as⁵:

$$\begin{aligned} S(p, v, \alpha) &= - \left(1 + \frac{f_p^{(\alpha)}}{1 + f_p^{(\alpha)}} \right) \left[\ln \left(1 - |\gamma_p^{(\alpha)}|^2 \right) + \frac{|\gamma_p^{(\alpha)}|^2}{\left(1 - |\gamma_p^{(\alpha)}|^2 \right)} \ln \left(|\gamma_p^{(\alpha)}|^2 \right) \right] \\ &\quad - \left(1 - f_p^{(\alpha)} \right) \ln \left(1 + f_p^{(\alpha)} \right). \end{aligned} \quad (3.73)$$

Then the final expression for the entanglement entropy in de Sitter space can be expressed as a sum over all possible quantum states which carries $\mathbf{SO}(1, 3)$ principal quantum number p . Consequently, the final expression for the entanglement entropy in de Sitter space is given by the following expression:

$$S(v, \alpha) = \sum_{\text{States}} \sum_{p=0}^{\infty} S(p, v, \alpha) \rightarrow V_{\mathbf{H}^3} \int_{p=0}^{\infty} dp \mathcal{D}_3(p) S(p, v, \alpha) = \mathbf{c}_6(\alpha, v) V_{\mathbf{H}^3} / V_{\mathbf{H}^3}^{\text{REG}}, \quad (3.74)$$

⁵ If we follow the equivalent ansatz of density matrix as mentioned in Eq (3.63), the expression for entanglement entropy for a given $\mathbf{SO}(1, 3)$ principal quantum number p can be expressed as:

$$\begin{aligned} S(p, v, \alpha) &= -\ln a_p^{(\alpha)} - a_p^{(\alpha)} \frac{|\gamma_p^{(\alpha)}|^2}{\left(1 - |\gamma_p^{(\alpha)}|^2 \right)^2} \ln \left(|\gamma_p^{(\alpha)}|^2 \right) \left(1 + f_p^{(\alpha)} \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \right) \\ &\quad - a_p^{(\alpha)} f_p^{(\alpha)} \ln \left(1 + f_p^{(\alpha)} \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \right) - a_p^{(\alpha)} (f_p^{(\alpha)})^2 \left(1 - |\gamma_p^{(\alpha)}|^2 \right) \ln \left(1 + f_p^{(\alpha)} \right). \end{aligned} \quad (3.72)$$

For our computation we will not use this ansatz any further.

where $\mathcal{D}_3(p) = p^2/2\pi^2$ characterise the density of states for radial functions on the Hyperboloid \mathbf{H}^3 . Additionally, it is important to note that the volume of the hyperboloid \mathbf{H}^3 is denoted by the overall factor $V_{\mathbf{H}^3}$. Here the regularised volume of the hyperboloid \mathbf{H}^3 for $r \leq L_c$ can be written as:

$$\begin{aligned} V_{\mathbf{H}^3} &= V_{\mathbf{S}^2} \int_{r=0}^{L_c} dr \sinh^2 r \xrightarrow{\text{large } L_c} \frac{\pi}{2} \left[e^{2L_c} - 4L_c \right] \\ &= \left[\mathbf{A}_{\text{ENT}} - \pi \ln \mathbf{A}_{\text{ENT}} + \pi \ln \left(\frac{\pi}{2} \right) \right] = V_{\mathbf{H}^3}^{\text{REG}} \left[\frac{1}{4\eta^2} + \ln \eta \right], \end{aligned} \quad (3.75)$$

where \mathbf{A}_{ENT} is the entangling area and we use $V_{\mathbf{S}^2} = 4\pi$. Here the cutoff L_c can be written as, $L_c \sim -\ln \eta$. In this context we define regularised volume of the hyperboloid \mathbf{H}^3 as, $V_{\mathbf{H}^3}^{\text{REG}} = V_{\mathbf{S}^3}/2 = 2\pi$.

In $3+1$ D, for the case of α vacuum, long range quantum correlation is measured by $\mathbf{c}_6(\alpha, v)$, which is defined as:

$$\mathbf{c}_6(\alpha, v) \equiv S_{\text{intr}}(\alpha, v) = \left[\left(1 + \frac{f_p^{(\alpha)}}{1 + f_p^{(\alpha)}} \right) \mathcal{I}^{(\alpha)} + \left(1 - f_p^{(\alpha)} \right) \ln \left(1 + f_p^{(\alpha)} \right) \mathcal{V} \right], \quad (3.76)$$

where the integrals $\mathcal{I}^{(\alpha)}$ and \mathcal{V} can be written in $3+1$ dimensional space-time as:

$$\begin{aligned} \mathcal{I}^{(\alpha)} &= -\frac{1}{\pi} \int_{p=0}^{\infty} dp \, p^2 \left[\ln \left(1 - |\gamma_p^{(\alpha)}|^2 \right) + \frac{|\gamma_p^{(\alpha)}|^2}{\left(1 - |\gamma_p^{(\alpha)}|^2 \right)} \ln \left(|\gamma_p^{(\alpha)}|^2 \right) \right], \\ \mathcal{V} &= -\frac{1}{\pi} \int_{p=0}^{\infty} dp \, p^2. \end{aligned} \quad (3.77)$$

Here it is important to mention that:

- The integral \mathcal{V} is divergent. To make it finite, we need to regularise it by introducing a change in variable by using $x = 2\pi p$ and by introducing a UV cut off Λ_C leading to:

$$\begin{aligned} \mathcal{V} &= -\frac{1}{8\pi^4} \int_{x=0}^{\Lambda_C} dx \, x^2 = -\frac{\Lambda_C^3}{24\pi^4} = -\frac{L^3}{3\pi} = -\frac{1}{4\pi^2} V_{\mathbf{S}^2} \\ \text{where } V_{\mathbf{S}^2} &= \frac{4}{3}\pi L^3 = \frac{1}{6\pi^2} \Lambda_C^3. \end{aligned} \quad (3.78)$$

The magnitude of this integral represents the finite volume of the configuration space in which we are computing the entanglement entropy from Von Neumann measure. In principle, this volume can be infinite, but after fixing the cut-off, this integral actually proportional to a finite volume of a sphere of radius Λ_C . Actually, the cut-off of the parameter x fix the highest accessible value of the characteristic length scale p , which mimics the role of some sort of principal quantum number as appearing in hydrogen atom problem. Only the difference is, here p is restricted within the window, $0 < p < \Lambda_C/2\pi$ and it takes all possible values within this window (i.e. continuous). After fixing the cut-off, the length of the system under consideration can be expressed as, $L = \int_0^{\Lambda_C/2\pi} dp = \Lambda_C/2\pi$. Now it is obvious that

if the cut-off $\Lambda_C \rightarrow \infty$, the entanglement entropy is diverging, which physically implies that the effect of long range quantum correlation is very large at very large length scale, $L = \int_0^\infty dp \rightarrow \infty$ i.e. at late time scales in de Sitter space.

- Further we analyse the integral $\mathcal{I}^{(\alpha)}$ using both of the solutions obtained for arbitrary v and $v = 3/2$. Following the previous argument, we also put a cut-off Λ_C to perform the integral on the rescaled **SO(1, 3)** quantum number $x = 2\pi p$ and after performing the integral we can study the behaviour of both of the results. First of all we start with the following integral with “ \pm ” signature, as given by:

$$\mathcal{I}^{(\alpha)} = -\frac{1}{8\pi^4} \int_{x=0}^{\Lambda_C} dx x^2 \left[\ln(1 - 2G_\pm(x, v, \alpha)) + \frac{2G_\pm(x, v, \alpha)}{(1 - 2G_\pm(x, v, \alpha))} \ln(2G_\pm(x, v, \alpha)) \right], \quad (3.79)$$

where $G_\pm(x, v, \alpha)$ for any arbitrary value of the parameter α is defined as:

$$G_\pm(x, v, \alpha) = \frac{1}{4|\tilde{m}_{\mathbf{RL}}|^2} \left[\left(1 + |\tilde{m}_{\mathbf{RL}}|^4 + |\tilde{m}_{\mathbf{RR}}|^4 - 2|\tilde{m}_{\mathbf{RR}}|^2 - \tilde{m}_{\mathbf{RR}}^2 (\tilde{m}_{\mathbf{RL}}^*)^2 - \tilde{m}_{\mathbf{RL}}^2 (\tilde{m}_{\mathbf{RR}}^*)^2 \right) \pm \left\{ \left(-1 - |\tilde{m}_{\mathbf{RL}}|^4 - |\tilde{m}_{\mathbf{RR}}|^4 + 2|\tilde{m}_{\mathbf{RR}}|^2 + \tilde{m}_{\mathbf{RR}}^2 (\tilde{m}_{\mathbf{RL}}^*)^2 + \tilde{m}_{\mathbf{RL}}^2 (\tilde{m}_{\mathbf{RR}}^*)^2 \right)^2 - 4|\tilde{m}_{\mathbf{RL}}|^4 \right\}^{\frac{1}{2}} \right], \quad (3.80)$$

where the components $\tilde{m}_{\mathbf{RR}} = \tilde{m}_{\mathbf{LL}}$ and $\tilde{m}_{\mathbf{RL}} = \tilde{m}_{\mathbf{LR}}$ are redefined in terms of the new variable $x = 2\pi p$.

Here small axion mass ($v^2 > 0$) limiting situations are considered in $v = 1/2$ conformal mass as well in $v = 3/2$ case in presence of an additional arbitrary parameter α . Additionally, we consider large axion mass ($v^2 < 0$ where $v \rightarrow -i|v|$) limiting situation. In this large axion mass limiting situation we consider the window of **SO(1, 3)** principal quantum number is $0 < p < |v|$. Consequently, the entries of the coefficient matrix \tilde{m} can be approximated as:

$$\tilde{m}_{\mathbf{RR}} = -\sqrt{\frac{\cosh(|v| - p)}{\cosh(|v| + p)}} \frac{2 \left[\cosh 2\alpha \cosh^2 \pi|v| - \sinh 2\alpha \sinh^2 \pi p + \frac{1}{2} \sinh 2\pi|v| \right]}{(e^{2\pi p} + e^{2\pi|v|}) \cosh^2 \alpha + (e^{2\pi p} + e^{2\pi|v|}) \sinh^2 \alpha}, \quad (3.81)$$

$$\tilde{m}_{\mathbf{RL}} = -\sqrt{\frac{\cosh(|v| - p)}{\cosh(|v| + p)}} \frac{2i \left[(\cosh 2\alpha + \sinh 2\alpha) \cosh \pi|v| + \sinh \pi|v| \right]}{(e^{2\pi p} + e^{2\pi|v|}) \cosh^2 \alpha + (e^{2\pi p} + e^{2\pi|v|}) \sinh^2 \alpha}. \quad (3.82)$$

This implies that for α vacuum if we consider the large axion mass ($v^2 < 0$ where $v \rightarrow -i|v|$) limiting situation we get always real value for $\tilde{m}_{\mathbf{RR}}$ and imaginary value for $\tilde{m}_{\mathbf{RL}}$. Consequently one can easily reduce the four sets of Eqn. (3.46) and Eqn. (3.47) into two sets of equations as exactly we have done in ref. [14] for Bunch Davies vacuum. In this large axion mass ($v^2 < 0$ where $v \rightarrow -i|v|$) limiting situation the two solutions for the $\gamma_p^{(\alpha)}$ for α vacuum are given by:

$$\gamma_p^{(\alpha)} \approx \frac{1}{2|\tilde{m}_{\mathbf{RL}}|} \left[\left(1 + |\tilde{m}_{\mathbf{RL}}|^2 - \tilde{m}_{\mathbf{RR}}^2 \right) \pm \sqrt{\left(1 + |\tilde{m}_{\mathbf{RL}}|^2 - \tilde{m}_{\mathbf{RR}}^2 \right)^2 - 4|\tilde{m}_{\mathbf{RL}}|^2} \right]. \quad (3.83)$$

Small mass limiting situations are explicitly appearing in $v = 1/2$ and $v = 3/2$ case. For our study here we consider large mass limiting situation which is important to study the physical outcomes. In this situation we divide the total window of p into two regions, as given by $0 < p < |v|$ and $|v| < p < \Lambda_C$. Here in these regions of interests the two solutions for $\gamma_p^{(\alpha)}$ in presence of α vacuum can be approximately written as:

$$|\gamma_p^{(\alpha)}| \approx \begin{cases} \frac{e^{-\pi|v|}(1 + \tan \alpha)}{e^{-\pi p}(1 + \tan \alpha)(1 + \tan \alpha e^{2\pi|v|})} & \text{for } 0 < p < |v| \\ \frac{e^{-\pi p}(1 + \tan \alpha)(1 + \tan \alpha e^{2\pi|v|})}{(1 + \tan^2 \alpha e^{-2\pi p})} & \text{for } |v| < p < \Lambda_C/2\pi, \end{cases} \quad (3.84)$$

and

$$|\gamma_p^{(\alpha)}| = \begin{cases} \frac{e^{\pi|v|}(1 + \tan \alpha)}{e^{\pi p}(1 + \tan \alpha)(1 + \tan \alpha e^{2\pi|v|})} & \text{for } 0 < p < |v| \\ \frac{e^{\pi p}(1 + \tan \alpha)(1 + \tan \alpha e^{2\pi|v|})}{(1 + \tan^2 \alpha e^{-2\pi p})} & \text{for } |v| < p < \Lambda_C/2\pi. \end{cases} \quad (3.85)$$

As a result, for large mass limiting range the α parameter dependent regularised integral $\mathcal{I}_1^{(\alpha)}$ for the first solution for $|\gamma_p^{(\alpha)}|$ can be written as:

$$\mathcal{I}_1^{(\alpha)} = \begin{cases} -\frac{\mathcal{A}(v)}{8\pi^4} \left[\ln(1 - e^{-2\pi v}(1 + \tan \alpha)^2) + \frac{(2 \ln(1 + \tan \alpha) - 2\pi v) e^{-2\pi v}(1 + \tan \alpha)^2}{(1 - e^{-2\pi v}(1 + \tan \alpha)^2)} \right] \\ \text{for } 0 < x < 2\pi|v| \\ -\frac{\mathcal{B}(v, \alpha, \Lambda_C)}{8\pi^4} \quad \text{for } 2\pi|v| < x < \Lambda_C, \end{cases} \quad (3.86)$$

and for the second solution for $|\gamma_p^{(\alpha)}|$ we get:

$$\mathcal{I}_1^{(\alpha)} = \begin{cases} -\frac{\mathcal{A}(v)}{8\pi^4} \left[\ln(1 - e^{2\pi v}(1 + \tan \alpha)^2) + \frac{(2 \ln(1 + \tan \alpha) + 2\pi v) e^{2\pi v}(1 + \tan \alpha)^2}{(1 - e^{2\pi v}(1 + \tan \alpha)^2)} \right] \\ \text{for } 0 < x < 2\pi|v| \\ -\frac{\mathcal{C}(v, \alpha, \Lambda_C)}{8\pi^4} \quad \text{for } 2\pi|v| < x < \Lambda_C. \end{cases} \quad (3.87)$$

In Eq. (3.86) and Eq (3.87) coefficients $\mathcal{A}(v)$, $\mathcal{B}(v, \alpha, \Lambda_C)$ and $\mathcal{C}(v, \alpha, \Lambda_C)$ are defined by the following expressions:

$$\mathcal{A}(v) = \int_{x=0}^{2\pi v} dx x^2 = \frac{8\pi^3}{3} v^3, \quad (3.88)$$

$$\begin{aligned} \mathcal{B}(v, \alpha, \Lambda_C) \approx & \int_{x=2\pi v}^{\Lambda_C} dx x^2 \left[\ln \left(1 - \frac{e^{-x}(1 + \tan \alpha)^2(1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2} \right) \right. \\ & \left. + \frac{\frac{e^{-x}(1 + \tan \alpha)^2(1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2}}{\left(1 - \frac{e^{-x}(1 + \tan \alpha)^2(1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2} \right)} \left\{ 2 \ln((1 + \tan \alpha)(1 + \tan \alpha e^{2\pi|v|})) \right. \right. \\ & \left. \left. - 2 \ln(1 - \tan^2 \alpha e^{-x}) - x \right\} \right], \end{aligned} \quad (3.89)$$

$$\begin{aligned} \mathcal{C}(v, \alpha, \Lambda_C) = & \int_{x=2\pi v}^{\Lambda_C} dx x^2 \left[\ln \left(1 - \frac{e^x (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2} \right) \right. \\ & + \frac{\frac{e^x (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2}}{\left(1 - \frac{e^x (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi|v|})^2}{(1 - \tan^2 \alpha e^{-x})^2} \right)} \left\{ 2 \ln((1 + \tan \alpha) (1 + \tan \alpha e^{2\pi|v|})) \right. \\ & \left. \left. - 2 \ln(1 - \tan^2 \alpha e^{-x}) + x \right\} \right]. \end{aligned} \quad (3.90)$$

Further within the window $0 < x < 2\pi|v|$ we take the large mass limit $|v| \gg 1$ in the first solution for $|\gamma_p^{(\alpha)}|$ in presence of α vacuum:

$$\lim_{|v| \gg 1} \mathcal{I}_1^{(\alpha)} \approx \frac{2v^4}{3} e^{-2\pi v} (1 + \tan \alpha)^2 \left\{ 1 - \frac{1}{\pi v} \ln(1 + \tan \alpha) \right\} \left[1 + (1 + \tan \alpha)^2 \mathcal{O}(v^{-1}) \right]. \quad (3.91)$$

Similarly the integral \mathcal{V} can be written as:

$$\mathcal{V} = \begin{cases} -\frac{1}{8\pi^4} \int_{x=0}^{2\pi|v|} dx x^2 = -\frac{|v|^3}{3\pi} & \text{for } 0 < x < 2\pi|v| \\ -\frac{1}{8\pi^4} \int_{x=2\pi|v|}^{\Lambda_C} dx x^2 = -\frac{1}{24\pi^4} (\Lambda_C^3 - 8\pi^3 |v|^3), & \text{for } 2\pi|v| < x < \Lambda_C. \end{cases} \quad (3.92)$$

Consequently in the large mass limiting situation ($0 < x < 2\pi|v|$) we get the following expression for the entanglement entropy:


$$\begin{aligned} \lim_{|v| \gg 1} \mathbf{c}_6(\alpha, v) \approx & \left(1 + \frac{f_p^{(\alpha)}}{1 + f_p^{(\alpha)}} \right) \frac{2v^4}{3} e^{-2\pi v} (1 + \tan \alpha)^2 \\ & \times \left\{ 1 - \frac{1}{\pi v} \ln(1 + \tan \alpha) \right\} \left[1 + (1 + \tan \alpha)^2 \mathcal{O}(v^{-1}) \right] \\ & - \left(1 - f_p^{(\alpha)} \right) \ln(1 + f_p^{(\alpha)}) \frac{v^3}{3\pi}. \end{aligned} \quad (3.93)$$

Further in absence of the source contribution in the large mass limit the long range quantum correlation can be expressed as:

$$\begin{aligned} \lim_{|v| \gg 1, f_p \rightarrow 0} \mathbf{c}_6(\alpha, v) \approx & \frac{2v^4}{3} e^{-2\pi v} (1 + \tan \alpha)^2 \left\{ 1 - \frac{1}{\pi v} \ln(1 + \tan \alpha) \right\} \\ & \times \left[1 + (1 + \tan \alpha)^2 \mathcal{O}(v^{-1}) \right]. \end{aligned} \quad (3.94)$$

For the second solution of $|\gamma_p^{(\alpha)}|$ in presence of α vacuum, we get:

$$\lim_{|v| \gg 1} \mathcal{I}_1^{(\alpha)} = -\frac{v^3}{3\pi} \left[\ln(1 - e^{2\pi v} (1 + \tan \alpha)^2) \right]$$



(a) Normalized entanglement entropy vs ν^2 in 3 + 1 D de Sitter space in presence of α vacuum and in absence Sitter space in presence of α vacuum and axionic source axionic source ($f_p^{(\alpha)} = 0$). (b) Normalized entanglement entropy vs ν^2 in 3 + 1 D de Sitter space in presence of α vacuum and in absence Sitter space in presence of α vacuum and axionic source ($f_p^{(\alpha)} = 10^{-7}$).

Fig. 1. Normalised entanglement entropy $S_{intr}/S_{v=1/2}$ vs mass parameter ν^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for '+' branch of solution of α vacuum i.e. $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. In both the situations we have normalised with conformal $v = 1/2$ result in presence of α vacuum. (For interpretation of the colours in the figure(s), the reader is referred to the web version of this article.)

$$+ \frac{(2 \ln(1 + \tan \alpha) + 2\pi v) e^{2\pi v} (1 + \tan \alpha)^2}{(1 - e^{2\pi v} (1 + \tan \alpha)^2)} \Bigg]. \quad (3.95)$$

Consequently in the large mass limiting situation ($0 < x < 2\pi|\nu|$) we get the following expression for the entanglement entropy:

$$\begin{aligned} \lim_{|\nu| \gg 1} \mathbf{c}_6(\alpha, \nu) \approx & - \left(1 + \frac{f_p^{(\alpha)}}{1 + f_p^{(\alpha)}} \right) \frac{\nu^3}{3\pi} \left[\ln \left(1 - e^{2\pi v} (1 + \tan \alpha)^2 \right) \right. \\ & + \frac{(2 \ln(1 + \tan \alpha) + 2\pi v) e^{2\pi v} (1 + \tan \alpha)^2}{(1 - e^{2\pi v} (1 + \tan \alpha)^2)} \Bigg] \\ & - \left(1 - f_p^{(\alpha)} \right) \ln \left(1 + f_p^{(\alpha)} \right) \frac{\nu^3}{3\pi}. \end{aligned} \quad (3.96)$$

Further in absence of the source contribution in the large mass limit the long range quantum correlation can be expressed as:

$$\begin{aligned} \lim_{|\nu| \gg 1, f_p \rightarrow 0} \mathbf{c}_6(\alpha, \nu) \approx & - \frac{\nu^3}{3\pi} \left[\ln \left(1 - e^{2\pi v} (1 + \tan \alpha)^2 \right) \right. \\ & + \frac{(2 \ln(1 + \tan \alpha) + 2\pi v) e^{2\pi v} (1 + \tan \alpha)^2}{(1 - e^{2\pi v} (1 + \tan \alpha)^2)} \Bigg]. \end{aligned} \quad (3.97)$$

In Fig. 1(a) and Fig. 1(b), we have demonstrated the behaviour of entanglement entropy in $D = 4$ de Sitter space in absence ($f_p^{(\alpha)} = 0$) and in presence ($f_p^{(\alpha)} = 10^{-7}$) of axionic source with respect to the mass parameter ν^2 . In both the cases we have normalised the entanglement entropy with the result obtained from conformal mass parameter $v = 1/2$ in presence of α vacuum. In Fig. 1(a), it is clearly observed that in absence of axionic source in the large mass regime (where $\nu^2 < 0$) the normalised entanglement entropy $S_{intr}(\alpha)/S_{v=1/2}(\alpha)$ asymptotically approaches towards zero. In the large mass regime the measure of long range

correlation (or more precisely the entanglement entropy) in presence of α vacuum for axion can be expressed for $\gamma_p^{(\alpha)} = e^{-\pi|m_{axion}/H|} (1 + \tan \alpha)$ as:

$$\begin{aligned} \mathbf{c}_6 \left(\alpha, |v| \approx \frac{m_{axion}}{H} \right) &= S_{\text{intr}} \left(\alpha, |v| \approx \frac{m_{axion}}{H} \right) \\ &\approx \frac{2}{3} \left(\frac{m_{axion}}{H} \right)^4 e^{-\frac{2\pi m_{axion}}{H}} (1 + \tan \alpha)^2 \left\{ 1 - \frac{H}{\pi m_{axion}} \ln (1 + \tan \alpha) \right\} \\ &\quad \times \left[1 + (1 + \tan \alpha)^2 \mathcal{O} \left(\frac{H}{m_{axion}} \right) \right]. \end{aligned} \quad (3.98)$$

If we further use Eq (3.98) then it is clearly observed that in presence of α vacuum one is able to get considerably large entanglement compared to the result obtained for Bunch Davies vacuum ($\alpha = 0$) for large mass regime ($v^2 < 0$). To demonstrate this clearly we have depicted the numerical values of the entanglement entropy for $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green) and $\alpha = 0.3$ (violet). Now from the Fig. 1(a) it is observed that in $v^2 > 0$ region $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ reaches its maximum value at $\alpha = 0.1$ (green) and $\alpha = 0.3$ (violet) with $v = 0$ (or $m_{axion} = 3H/2$), as given by, $(S_{\text{intr}}(0.1)/S_{v=1/2}(0.1))_{\max, v=0} \sim 1.2$ and $(S_{\text{intr}}(0.3)/S_{v=1/2}(0.3))_{\max, v=0} \sim 2.1$. On the other hand, at $\alpha = 0.03$ (blue) and $\alpha = 0$ and $\alpha = 0.3$ (red) with $v = 1/2$ (or $m_{axion} = \sqrt{2}H$) the maximum value of $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ is given by, $(S_{\text{intr}}(0.03)/S_{v=1/2}(0.03))_{\max, v=1/2} \sim (S_{\text{intr}}(0)/S_{v=1/2}(0))_{\max, v=1/2} \sim 1$. Further if we consider the interval $3/2 < v < 5/2$ then $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ show one oscillation with different amplitude for all values of the parameter α . After that it reaches its maximum value for $\alpha = 0$ and $\alpha = 0.03$, as given by, $(S_{\text{intr}}(0.03)/S_{v=1/2}(0.03))_{\max, 3/2 < v < 5/2} \sim (S_{\text{intr}}(0)/S_{v=1/2}(0))_{\max, 3/2 < v < 5/2} \sim 1$. On the other hand, $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ reaches its minimum value for $\alpha = 0.1$ and $\alpha = 0.3$, as given by, $(S_{\text{intr}}(0.1)/S_{v=1/2}(0.1))_{\min, 3/2 < v < 5/2} \sim 1$ and $(S_{\text{intr}}(0.3)/S_{v=1/2}(0.3))_{\min, 3/2 < v < 5/2} \sim 1$. Similarly in the interval $5/2 < v < 7/2$ we can observe the same feature for the same values of α with larger period of oscillation. In Fig. 1(b), the significant role of axionic source term is explicitly shown. In both $v^2 < 0$ and $v^2 > 0$ regime the behaviour of $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ is exactly same as depicted in Fig. 1(a). But in presence of axionic source term the amount of $S_{\text{intr}}(\alpha)/S_{v=1/2}(\alpha)$ increase for $\alpha = 0, \alpha = 0.03$ and decrease for $\alpha = 0.1, \alpha = 0.3$ compared to Fig. 1(a). Also it is important to note that, the amplitude of the maximum and minimum of the oscillations change in presence of axionic source term.

On the other hand, for $\gamma_p = e^{\pi|m_{axion}/H|} (1 + \tan \alpha)$ the entanglement entropy for axion in the large mass limiting range is given by the following expression:

$$\begin{aligned} \mathbf{c}_6 \left(\alpha, |v| \approx \frac{m_{axion}}{H} \right) &= S_{\text{intr}} \left(\alpha, |v| \approx \frac{m_{axion}}{H} \right) \\ &\approx -\frac{1}{3\pi} \left(\frac{m_{axion}}{H} \right)^3 \left[\ln \left(1 - e^{\frac{2\pi m_{axion}}{H}} (1 + \tan \alpha)^2 \right) \right. \\ &\quad \left. + \frac{\left(2 \ln (1 + \tan \alpha) + \frac{2\pi m_{axion}}{H} \right) e^{\frac{2\pi m_{axion}}{H}} (1 + \tan \alpha)^2}{\left(1 - e^{\frac{2\pi m_{axion}}{H}} (1 + \tan \alpha)^2 \right)} \right]. \end{aligned} \quad (3.99)$$

Next, in Fig. 2, we have depicted the behaviour of entanglement entropy $S_{\text{intr}}(\alpha)$ with respect to the parameter α in absence ($f_p^{(\alpha)} = 0$) and presence ($f_p^{(\alpha)} = 10^{-7}$) of axionic source




Fig. 2. Entanglement entropy $S_{intr}(\alpha)$ vs parameter α in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we fix the value of the parameter v^2 at different positive and negative values including zero.

for the mass parameter $v^2 < 0$ and $v^2 > 0$ respectively. In Fig. 2(a) and Fig. 2(b) it is observed that a crossover takes place for $v^2 = 1/4, 9/4, 25/4$ (green), $v^2 = 1/16, 9/16, 25/16$ (blue) and $v^2 = 0$ (red) with small values of the parameter α . We also observe that for $v^2 = 1/4, 9/4, 25/4$ (green) entanglement entropy decreases with increasing value of the parameter α . On the other hand, for $v^2 = 1/16, 9/16, 25/16$ (blue) and $v^2 = 0$ (red) entanglement entropy increases with increasing value of the parameter α . Additionally, we observe that, in presence of axionic source the entanglement entropy is significantly larger compared to the result obtained in absence of source contribution. In Fig. 2(c) and Fig. 2(d) it is observed that no crossover takes place for $v^2 = -1/2$ (green), $v^2 = -1/4, -9/4, -25/4$ (blue) and $v^2 = -1/16, -9/16, -25/16$ (red) with all values of the parameter α . Also it is important to note that, for all values of $v^2 < 0$ entanglement entropy increases with increasing value of the parameter α .

3.4. Computation of Rényi entropy using α vacua

In this context one can further use the density matrix to compute the Rényi entropy for α vacuum, which is defined as:

$$S_q(p, v, \alpha) = \frac{1}{1-q} \ln \text{Tr} [\rho_{\mathbf{L}}(p, \alpha)]^q. \quad \text{with } q > 0. \quad (3.100)$$

The obtained solution for α vacuum with a given **SO(1, 3)** principal quantum number p can be written as:

$$\begin{aligned} S_q(p, v, \alpha) = & \frac{1}{1-q} \left[q \ln \left(\frac{1 - |\gamma_p^{(\alpha)}|^2}{1 + f_p^{(\alpha)}} \right) - \ln \left(1 - |\gamma_p^{(\alpha)}|^{2q} \right) \right] \\ & + \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{(1 - |\gamma_p^{(\alpha)}|^2)^{-k}}{(1 - |\gamma_p^{(\alpha)}|^{-2k})} \right], \end{aligned} \quad (3.101)$$

using which the interesting part of the Rényi entropy in de Sitter space for α vacuum can be written as:

$$S_{q,\text{intr}}(\alpha, v) = \frac{1}{\pi} \int_{p=0}^{\infty} dp p^2 S_q(p, v, \alpha). \quad (3.102)$$

Now to study the properties of the derived result we check the following physical limiting situations as given by:

- If we take the limit $q \rightarrow 1$ limit then from the Rényi entropy in α vacuum we get, $\lim_{q \rightarrow 1} S_q(p, v, \alpha) \neq S(p, v, \alpha)$, which shows that in presence of axionic source, the entanglement entropy and Rényi entropy are not same in the limit $q \rightarrow 1$. Now if we take further $f_p \rightarrow 0$ then entanglement entropy and Rényi entropy both are same.
- Further if we take the limit $q \rightarrow \infty$ limit then from the Rényi entropy in α vacuum we get:

$$\lim_{q \rightarrow \infty} S_q(p, v, \alpha) = -\ln[\rho_{\mathbf{L}}]_{\max} \approx \ln \left(\frac{1 + f_p^{(\alpha)}}{1 - |\gamma_p^{(\alpha)}|^2} \right), \quad (3.103)$$

which directly implies the largest eigenvalue of density matrix. Now if we take further $f_p \rightarrow 0$ in Eqn (3.103) then entanglement entropy and Rényi entropy both are same.

Further substituting the expression for entanglement entropy $S(p, v, \alpha)$ computed in presence of axion for α vacuum and integrating over all possible **SO(1, 3)** principal quantum number, lying within the window $0 < p < \infty$, we get:

$$S_{q,\text{intr}}(\alpha, v) = \left[\mathcal{M}_{1,q} + \ln \left(1 + f_p^{(\alpha)} \right) \mathcal{M}_{2,q}^{(\alpha)} + \mathcal{M}_{3,q}^{(\alpha)} \right], \quad (3.104)$$

where the integrals $\mathcal{M}_{1,q}$, $\mathcal{M}_{2,q}^{(\alpha)}$ and $\mathcal{M}_{3,q}^{(\alpha)}$ can be written as:

$$\mathcal{M}_{1,q} = \frac{1}{\pi} \int_{p=0}^{\infty} dp p^2 \left[\frac{q}{1-q} \ln \left(1 - |\gamma_p^{(\alpha)}|^2 \right) - \frac{1}{1-q} \ln \left(1 - |\gamma_p^{(\alpha)}|^{2q} \right) \right], \quad (3.105)$$

$$\mathcal{M}_{2,q}^{(\alpha)} = -\frac{1}{\pi} \frac{q}{1-q} \int_{p=0}^{\infty} dp p^2, \quad (3.106)$$

$$\mathcal{M}_{3,q}^{(\alpha)} = \frac{1}{\pi} \frac{1}{1-q} \int_{p=0}^{\infty} dp p^2 \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{(1 - |\gamma_p^{(\alpha)}|^2)^{-k}}{(1 - |\gamma_p^{(\alpha)}|^{-2k})} \right]. \quad (3.107)$$

Here it is important to note that:

- Here the integral $\mathcal{M}_{2,q}^{(\alpha)}$ diverges. Further introducing a change in variable to $x = 2\pi p$ along with a cut-off Λ_C the regularised version of this integral can be written as:

$$\mathcal{M}_{2,q}^{(\alpha)} = -\frac{1}{8\pi^4} \frac{q}{1-q} \int_{x=0}^{\Lambda_C} dx x^2 = -\frac{\Lambda_C^3}{24\pi^4} \frac{q}{1-q}. \quad (3.108)$$

- On the other hand for arbitrary v and α we get:

$$\begin{aligned} \mathcal{M}_{1,q}^{(\alpha)} &= \frac{1}{8\pi^4} \int_{x=0}^{\Lambda_C} dx x^2 \left[\frac{q}{1-q} \ln(1 - 2G_{\pm}(x, v, \alpha)) \right. \\ &\quad \left. - \frac{1}{1-q} \ln(1 - (2G_{\pm}(x, v, \alpha))^q) \right], \end{aligned} \quad (3.109)$$

$$\mathcal{M}_{3,q}^{(\alpha)} = \frac{1}{8\pi^4} \frac{1}{1-q} \int_{x=0}^{\Lambda_C} dx x^2 \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{(1 - 2G_{\pm}(x, v, \alpha))^{-k}}{(1 - (2G_{\pm}(x, v, \alpha))^{-k})} \right], \quad (3.110)$$

where $G_{\pm}(x, v, \alpha)$ is defined in Eqn (3.80). We consider large axion mass ($v^2 < 0$ where $v \rightarrow -i|v|$) limiting situation which is important to study the physics from this case. In this large axion mass limiting situation we consider the window of **SO(1, 3)** principal quantum number is $0 < p < |v|$.

As a result, the regularised integral $\mathcal{M}_{1,q}^{(\alpha)}$ and $\mathcal{M}_{3,q}^{(\alpha)}$ for the first solution for $|\gamma_p^{(\alpha)}|$ in presence of α vacuum can be expressed as:

$$\mathcal{M}_{1,q}^{(\alpha)} = \begin{cases} \frac{\mathcal{A}(v)}{8\pi^4} \left[\frac{q}{1-q} \ln(1 - e^{-2\pi v} (1 + \tan \alpha)^2) \frac{1}{1-q} \ln(1 - e^{-2\pi v q} (1 + \tan \alpha)^{2q}) \right] \\ \text{for } 0 < x < 2\pi|v| \\ \frac{\mathcal{D}(v, \alpha, \Lambda_C, q)}{8\pi^4} \text{ for } 2\pi|v| < x < \Lambda_C. \end{cases} \quad (3.111)$$

$$\mathcal{M}_{3,q}^{(\alpha)} = \begin{cases} \frac{\mathcal{A}(v)}{8\pi^4} \frac{1}{1-q} \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{(1 - e^{-2\pi v} (1 + \tan \alpha)^2)^{-k}}{(1 - e^{2\pi v k} (1 + \tan \alpha)^{-2k})} \right] \\ \text{for } 0 < x < 2\pi|v| \\ \frac{1}{8\pi^4} \frac{1}{1-q} \int_{x=2\pi v}^{\Lambda_C} dx x^2 \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{\left(1 - \frac{e^{-x} (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi v})^2}{(1 + \tan^2 \alpha e^{-x})^2} \right)^{-k}}{\left(1 - \frac{e^{xk} (1 + \tan \alpha)^{-2k} (1 + \tan \alpha e^{2\pi v})^{-2k}}{(1 + \tan^2 \alpha e^{-x})^{-2k}} \right)} \right] \\ \text{for } 2\pi|v| < x < \Lambda_C, \end{cases} \quad (3.112)$$

and for the second solution for $|\gamma_p^{(\alpha)}|$ in presence of α vacuum we get:

$$\mathcal{M}_{1,q}^{(\alpha)} = \begin{cases} \frac{\mathcal{A}(\nu)}{8\pi^4} \left[\frac{q}{1-q} \ln \left(1 - e^{2\pi\nu} (1 + \tan \alpha)^2 \right) \right. \\ \left. - \frac{1}{1-q} \ln \left(1 - e^{2\pi\nu q} (1 + \tan \alpha)^{2q} \right) \right] & \text{for } 0 < x < 2\pi|\nu| \\ \frac{\mathcal{W}(\nu, \alpha, \Lambda_C, q)}{8\pi^4} & \text{for } 2\pi|\nu| < x < \Lambda_C. \end{cases} \quad (3.113)$$

$$\mathcal{M}_{3,q}^{(\alpha)} = \begin{cases} \frac{\mathcal{A}(\nu)}{8\pi^4} \frac{1}{1-q} \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{(1 - e^{2\pi\nu} (1 + \tan \alpha)^2)^{-k}}{(1 - e^{-2\pi\nu k} (1 + \tan \alpha)^{-2k})} \right] \\ \text{for } 0 < x < 2\pi|\nu| \\ \frac{1}{8\pi^4} \frac{1}{1-q} \int_{x=2\pi\nu}^{\Lambda_C} dx x^2 \ln \left[1 + \sum_{k=1}^q {}^q \mathbf{C}_k (f_p^{(\alpha)})^k \frac{\left(1 - \frac{e^x (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi\nu})^2}{(1 + \tan^2 \alpha e^{-x})^2} \right)^{-k}}{\left(1 - \frac{e^{-xk} (1 + \tan \alpha)^{-2k} (1 + \tan \alpha e^{2\pi\nu})^{-2k}}{(1 + \tan^2 \alpha e^{-x})^{-2k}} \right)} \right] \\ \text{for } 2\pi|\nu| < x < \Lambda_C. \end{cases} \quad (3.114)$$

Here the coefficient function $\mathcal{A}(\nu)$ is defined in Eq (3.89) and other α parameter dependent functions $\mathcal{D}(\nu, \alpha, \Lambda_C)$ and $\mathcal{W}(\nu, \alpha, \Lambda_C)$ are defined as:

$$\begin{aligned} \mathcal{D}(\nu, \alpha, \Lambda_C, q) = & \int_{x=2\pi\nu}^{\Lambda_C} dx x^2 \left[\frac{q}{1-q} \ln \left(1 - \frac{e^{-x} (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi\nu})^2}{(1 + \tan^2 \alpha e^{-x})^2} \right) \right. \\ & \left. - \frac{1}{1-q} \ln \left(1 - \frac{e^{-xq} (1 + \tan \alpha)^{2q} (1 + \tan \alpha e^{2\pi\nu})^{2q}}{(1 + \tan^2 \alpha e^{-x})^{2q}} \right) \right], \end{aligned} \quad (3.115)$$

$$\begin{aligned} \mathcal{W}(\nu, \alpha, \Lambda_C, q) = & \int_{x=2\pi\nu}^{\Lambda_C} dx x^2 \left[\frac{q}{1-q} \ln \left(1 - \frac{e^x (1 + \tan \alpha)^2 (1 + \tan \alpha e^{2\pi\nu})^2}{(1 + \tan^2 \alpha e^{-x})^2} \right) \right. \\ & \left. - \frac{1}{1-q} \ln \left(1 - \frac{e^{xq} (1 + \tan \alpha)^{2q} (1 + \tan \alpha e^{2\pi\nu})^{2q}}{(1 + \tan^2 \alpha e^{-x})^{2q}} \right) \right]. \end{aligned} \quad (3.116)$$

Further, using the results obtained from the first solution for $|\gamma_p^{(\alpha)}|$, within the range $0 < x < 2\pi|\nu|$ with $\nu^2 < 0$, we take $q \rightarrow 1$ limit. This gives the following simplified expression for the integral $\mathcal{M}_{1,q}^{(\alpha)}$:

$$\lim_{q \rightarrow 1} \mathcal{M}_{1,q}^{(\alpha)} = \frac{\nu^3}{3} \left[\frac{2(1 + \tan \alpha)^2 \left\{ \nu - \frac{1}{\pi} \ln(1 + \tan \alpha) \right\}}{(e^{2\pi\nu} - (1 + \tan \alpha)^2)} - \frac{\ln(1 - e^{-2\pi\nu} (1 + \tan \alpha)^2)}{\pi} \right]. \quad (3.117)$$

Now further using $|\nu| \gg 1$ approximation in Eq (3.117) we get:

$$\lim_{|\nu| >> 1, q \rightarrow 1} \mathcal{M}_{1,q}^{(\alpha)} = \frac{2\nu^4}{3} e^{-2\pi\nu} (1 + \tan\alpha)^2 \left\{ 1 - \frac{1}{\pi\nu} \ln(1 + \tan\alpha) \right\} \\ \times \left[1 + (1 + \tan\alpha)^2 \mathcal{O}(\nu^{-1}) \right]. \quad (3.118)$$

In this context further if we take the source less limit $f_p^{(\alpha)} \rightarrow 0$ then the integral $\mathcal{M}_{3,q}^{(\alpha)}$ vanishes:

$$\lim_{q \rightarrow 1, |\nu| >> 1, f_p \rightarrow 0} \mathcal{M}_{3,q}^{(\alpha)} = 0. \quad (3.119)$$

As a result in the large mass limiting situation with $q \rightarrow 1$ the long range correlation can be expressed in terms of Rényi entropy as:

$$\lim_{q \rightarrow 1, |\nu| >> 1, f_p^{(\alpha)} \rightarrow 0} S_{q,\text{intr}}(\alpha) \approx \frac{2\nu^4}{3} e^{-2\pi\nu} (1 + \tan\alpha)^2 \left\{ 1 - \frac{1}{\pi\nu} \ln(1 + \tan\alpha) \right\} \\ \times \left[1 + (1 + \tan\alpha)^2 \mathcal{O}(\nu^{-1}) \right] \\ = S_{\text{intr}}(\alpha) = \lim_{|\nu| >> 1, f_p^{(\alpha)} \rightarrow 0} \mathbf{c}_6(\alpha, \nu). \quad (3.120)$$

Similarly using the results obtained from the second solution for $|\gamma_p^{(\alpha)}|$, within the range $0 < x < 2\pi|\nu|$ with $\nu^2 < 0$, we take $q \rightarrow 1$ limit. This gives the following simplified expression for the integral $\mathcal{M}_{1,q}^{(\alpha)}$:


$$\lim_{q \rightarrow 1} \mathcal{M}_{1,q}^{(\alpha)} = \frac{\nu^3}{3} \left[\frac{2e^{2\pi\nu} (1 + \tan\alpha)^2 \{\nu + \ln(1 + \tan\alpha)\}}{e^{2\pi\nu} \tan^2\alpha + 2e^{2\pi\nu} \tan\alpha + e^{2\pi\nu} - 1} \right. \\ \left. - \frac{\ln(1 - e^{2\pi\nu} (1 + \tan\alpha)^2)}{\pi} \right], \quad (3.121)$$

$$\lim_{q \rightarrow 1, |\nu| >> 1, f_p \rightarrow 0} \mathcal{M}_{3,q}^{(\alpha)} = 0. \quad (3.122)$$

As a result in the large mass limiting situation with $q \rightarrow 1$ the long range correlation can be expressed in terms of Rényi entropy as:


$$\lim_{q \rightarrow 1, |\nu| >> 1, f_p^{(\alpha)} \rightarrow 0} S_{q,\text{intr}}(\alpha) \approx \frac{\nu^3}{3} \left[\frac{2e^{2\pi\nu} (1 + \tan\alpha)^2 \{\nu + \ln(1 + \tan\alpha)\}}{e^{2\pi\nu} \tan^2\alpha + 2e^{2\pi\nu} \tan\alpha + e^{2\pi\nu} - 1} \right. \\ \left. - \frac{\ln(1 - e^{2\pi\nu} (1 + \tan\alpha)^2)}{\pi} \right] \\ = S_{\text{intr}}(\alpha) = \lim_{|\nu| >> 1, f_p^{(\alpha)} \rightarrow 0} \mathbf{c}_6(\alpha, \nu). \quad (3.123)$$

In Fig. 3(a), Fig. 4(a), Fig. 5(a), Fig. 6(a), Fig. 7(a), we have demonstrated the behaviour of Rényi entropy for $q = 0.9$, $q = 0.7$, $q = 0.5$, $q = 0.3$ and $q = 0.1$ with respect to the mass parameter ν^2 . Here we did the computation in $D = 4$ de Sitter space in absence ($f_p^{(\alpha)} = 0$) of axionic source. Similarly in Fig. 3(b), Fig. 4(b), Fig. 5(b), Fig. 6(b), Fig. 7(b), we have demonstrated the behaviour of Rényi entropy for $q = 0.9$, $q = 0.7$, $q = 0.5$, $q = 0.3$ and $q = 0.1$ with respect to the mass parameter ν^2 . Additionally, the largest eigenvalue of the



(a) Normalized Rényi entropy vs v^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 0$). (b) Normalized Rényi entropy vs v^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 10^{-7}$).

Fig. 3. Normalised Rényi entropy $S_{q,intr}(\alpha)/S_{q,v=1/2}(\alpha)$ vs mass parameter v^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.9$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (violet) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.



(a) Normalized Rényi entropy vs v^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 0$). (b) Normalized Rényi entropy vs v^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 10^{-7}$).

Fig. 4. Normalised Rényi entropy $S_{q,intr}(\alpha)/S_{q,v=1/2}(\alpha)$ vs mass parameter v^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.7$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (violet) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.

density matrix ($q \rightarrow \infty$) in absence and presence of axionic source are plotted in Fig. 8(a) and Fig. 8(b). Here we did the computation in $D = 4$ de Sitter space in presence ($f_p^{(\alpha)} = 10^{-7}$) of axionic source. In this both the cases we also have normalised the Rényi entropy with the result obtained from conformal mass parameter $v = 1/2$ in presence of α vacuum. For a given value of the parameter q we have shown the plots for $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green) and $\alpha = 0.3$ (violet) in both the cases. Here we observe the following features:

- For $q = 0.9$ case in absence of the axionic source (see Fig. 3(a)) in the large mass parameter range ($v^2 < 0$) normalised Rényi entropy asymptotically approaches towards zero value. On the other hand in the small mass parameter range ($v^2 > 0$) it shows oscillations




Fig. 5. Normalised Rényi entropy $S_{q,intr}(\alpha)/S_{q,v=1/2}(\alpha)$ vs mass parameter ν^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.5$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (magenta) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.






Fig. 6. Normalised Rényi entropy $S_{q,intr}(\alpha)/S_{q,v=1/2}(\alpha)$ vs mass parameter ν^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.3$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (magenta) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.

in a periodic fashion. Here the amplitude of the oscillation is larger for $\alpha = 0.3$ compared to the other values of α . Also it is important to note that, at $v = 1/2$, $v = 3/2$ and $v = 5/2$ we get extrema for the oscillation. Further in presence of the axionic source (see Fig. 3(b)) in the large mass parameter range ($\nu^2 < 0$) normalised Rényi entropy rapidly approaches to zero value for all values of the parameter α considered in this paper. Also in the small mass parameter range ($\nu^2 > 0$) the amplitude of the oscillation is significantly large for $\alpha = 0.3$. Also it is observed that for $\nu^2 > 0$ the long range correlation is larger in presence of the axionic source. But for $\nu^2 < 0$ the long range correlation is rapidly decaying with $f_p^{(\alpha)} = 10^{-7}$ and asymptotically decaying with $f_p^{(\alpha)} = 0$ for all values of α .



(a) Normalized Rényi entropy vs ν^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 0$).
(b) Normalized Rényi entropy vs ν^2 in 3 + 1 D de Sitter space without axionic source ($f_p^{(\alpha)} = 10^{-7}$).

Fig. 7. Normalised Rényi entropy $S_{q,\text{intr}}(\alpha)/S_{q,v=1/2}(\alpha)$ vs mass parameter ν^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.1$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (violet) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.



(a) Normalized Rényi entropy vs ν^2 in 3 + 1 D de Sitter space without axionic source ($f_p = 0$).
(b) Normalised Rényi entropy vs ν^2 in de Sitter space with axionic source ($f_p = 10^{-7}$).

Fig. 8. Normalised Rényi entropy $S_{q \rightarrow \infty, \text{intr}}(\alpha)/S_{q \rightarrow \infty, v=1/2}(\alpha)$ vs mass parameter ν^2 in 3 + 1 D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) and in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q \rightarrow \infty$ and $\alpha = 0$ (red), $\alpha = 0.03$ (blue), $\alpha = 0.1$ (green), $\alpha = 0.3$ (violet) with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$, which quantifies largest eigenvalue of density matrix. Here we set the cut-off $\Lambda_C = 300$ for numerical computation.

- For other values of the parameter q i.e. $q = 0.7$, $q = 0.5$, $q = 0.3$ and $q = 0.1$ cases in absence of the axionic source (see Fig. 4(a), Fig. 5(a), Fig. 6(a) and Fig. 7(a)) in the large mass parameter range ($\nu^2 < 0$) normalised Rényi entropy asymptotically approaches towards zero value. On the other hand in the small mass parameter range ($\nu^2 > 0$) it shows oscillations in a periodic fashion. Here the amplitude of the oscillation is larger for $\alpha = 0.3$ compared to the other values of α . Also it is important to note that, at $v = 1/2$, $v = 3/2$ and $v = 5/2$ we get extrema for the oscillation. Further in presence of the axionic source (see Fig. 4(b), Fig. 5(b), Fig. 6(b) and Fig. 7(b)) one can observe the exact behaviour as observed without any source contribution. It also implies that for all $q < 0.9$ the normalised Rényi entropy is insensitive to the source contribution.




Fig. 9. Rényi entropy $S_{q,intr}(\alpha)$ vs parameter α plot in $3 + 1$ D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) for $q = 0.1$, $q = 0.3$, $q = 0.5$, $q = 0.7$, $q = 0.9$ with ‘+’ branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$.

- For $q \rightarrow \infty$ case in absence (see Fig. 8(a)) and presence of the axionic source (see Fig. 8(b)) variation of normalised Rényi entropy with v^2 for all values of the parameter α is similar. It is important to note that the amplitudes of the oscillations in $v^2 > 0$ region and the saturation value in $v^2 < 0$ region is larger in presence of axionic source.

Next, in Fig. 9(a), Fig. 9(b), Fig. 9(c), Fig. 9(d) and Fig. 10(a), Fig. 10(b), Fig. 10(c), Fig. 10(d), we have depicted the behaviour of Rényi entropy with respect to the parameter α in absence ($f_p^{(\alpha)} = 0$) and presence ($f_p^{(\alpha)} = 10^{-7}$) of axionic source for the mass parameter $v^2 > 0$. In all figures it is observed that a crossover takes place for $v^2 = 1/4, 9/4, 25/4$ (green), $v^2 = 1/16, 9/16, 25/16$ (blue) and $v^2 = 0$ (red) with small values of the parameter α . We also observe that for $v^2 = 1/4, 9/4, 25/4$ (green) Rényi entropy decreases with increasing value of the parameter α . On the other hand, for $v^2 = 1/16, 9/16, 25/16$ (blue) and $v^2 = 0$ (red) Rényi entropy increases with increasing value of the parameter α . Additionally, in presence of axionic source the Rényi entropy is slightly larger compared to the result obtained in absence of source contribution. Also it is observed that no crossover takes place for $v^2 = -1/2$ (green), $v^2 = -1/4, -9/4, -25/4$ (blue) and $v^2 = -1/16, -9/16, -25/16$ (red) with all values of the parameter α . Also it is important to note that, for all values of $v^2 < 0$ Rényi entropy increases with increasing value of the parameter α (see Figs. 11 and 12). Further in Fig. 13(a), Fig. 13(b)), Fig. 13(c), Fig. 13(d), we have shown the variation of Rényi entropy with respect to the parameter q in absence and presence of axionic source for $\alpha = 0$ (purple), $\alpha = 0.03$ (orange), $\alpha = 0.1$ (cyan) and $\alpha = 0.3$ (brown) respec-




Fig. 10. Rényi entropy $S_{q,\text{intr}}(\alpha)$ vs parameter α plot in $3 + 1$ D de Sitter space in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.1, q = 0.3, q = 0.5, q = 0.7, q = 0.9$ with ‘+’ branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$.

tively. It is observed that for small values of the parameter q the value of the Rényi entropy for a given value of α always increase. On the other hand for small values of the parameter q Rényi entropy saturates to a finite small value.

4. Summary

To summarise, in this paper, we have addressed the following issues:

- First we have presented the computation of entanglement entropy in de Sitter space in presence of axion with a linear source contribution in the effective potential as originating from **Type IIB** string theory. To demonstrate this we have derived the axion wave function in an open chart.
- Next using the α vacuum state we have expressed the wave function in terms of creation and annihilation operators. Further applying Bogoliubov transformation on α vacuum state we have constructed the expression for reduced density matrix.
- Further, using reduced density matrix we have derived the entanglement entropy, which is consistent with ref. [10] if we set $\alpha = 0$. In the $\nu^2 < 0$ range we have derived analytical result for the entanglement entropy. Finally, we have used numerical approximations to estimate entanglement entropy with any value of ν^2 .




Fig. 11. Rényi entropy $S_{q=0,n,intr}(\alpha)$ vs parameter α plot in $3 + 1$ D de Sitter space in absence of axionic source ($f_p^{(\alpha)} = 0$) for $q = 0.1, q = 0.3, q = 0.5, q = 0.7, q = 0.9$ with ‘+’ branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$.

- We have also computed the Rényi entropy in presence of axion source. In absence of the source this result is consistent with ref. [10] in $q \rightarrow 1$ limit. Here in $\nu^2 < 0$ region we have provided the analytical expression for the Rényi entropy. We have also used numerical techniques to study the behaviour of Rényi entropy and largest eigenvalue of the density matrix with any value of ν^2 .
- Our result provides the necessary condition to generate non vanishing entanglement in primordial cosmology due to axion.

The future directions of this paper are appended below:

- Using the derived expression for density matrix for generalised α vacua, one can further compute any n point long range quantum correlation to study the implications in the context of primordial cosmology. It is expected that this result will surely help to understand the connection between the Bell’s inequality violation, quantum entanglement and primordial non-Gaussianity.
- Till now we have studied the necessary condition for generating non zero entanglement entropy in primordial cosmology. In this connection one can further compute quantum discord, entanglement negativity etc., which play a significant role to quantify long range quantum correlations without necessarily involving quantum entanglement.




Fig. 12. Rényi entropy $S_{q,\text{intr}}(\alpha)$ vs parameter α plot in 3 + 1 D de Sitter space in presence of axionic source ($f_p^{(\alpha)} = 10^{-7}$) for $q = 0.1, q = 0.3, q = 0.5, q = 0.7, q = 0.9$ with '+' branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$.

Acknowledgements

SC would like to thank Quantum Gravity and Unified Theory and Theoretical Cosmology Group, Max Planck Institute for Gravitational Physics, Albert Einstein Institute and Inter University Center for Astronomy and Astrophysics, Pune for providing the Post-Doctoral Research Fellowship. SP acknowledges the J. C. Bose National Fellowship for support of his research. Last but not the least, we would all like to acknowledge our debt to the people of India for their generous and steady support for research in natural sciences.

Appendix A. Wave function for axion using Bunch Davies vacuum

Further using Eqn (3.1) the field equation of motion for the axion can be written as [14]:

$$\left[(a(t))^{-3} \partial_t \left(a^3(t) \partial_t \right) - (Ha(t))^{-2} \hat{\mathbf{L}}_{\mathbf{H}^3}^2 + m_{axion}^2 \right] \phi = \mu^3, \quad (\text{A.1})$$

where the scale factor $a(t)$ in de Sitter open chart is given by, $a(t) = \sinh t / H$. Here the Laplacian operator $\hat{\mathbf{L}}_{\mathbf{H}^3}^2$ in \mathbf{H}^3 satisfies the following eigenvalue equation [56]:

$$\begin{aligned}\hat{\mathbf{L}}_{\mathbf{H}^3}^2 \mathcal{Y}_{plm}(r, \theta, \phi) &= \frac{1}{\sinh^2 r} \left[\partial_r \left(\sinh^2 r \partial_r \right) + \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\phi^2 \right] \mathcal{Y}_{plm}(r, \theta, \phi) \\ &= -(1 + p^2) \mathcal{Y}_{plm}(r, \theta, \phi),\end{aligned}\quad (\text{A.2})$$




Fig. 13. Rényi entropy $S_{q,intr}(\alpha)$ vs q plot in $3 + 1$ D de Sitter space in absence and presence of axionic source for $\alpha = 0$, $\alpha = 0.03$, $\alpha = 0.1$ and $\alpha = 0.3$ with ‘+’ branch of solution of $|\gamma_p^{(\alpha)}|$ and $|\Gamma_{p,n}^{(\alpha)}|$.

where $\mathcal{Y}_{plm}(r, \theta, \phi)$ represents orthonormal eigenfunctions which can be written as:

$$\mathcal{Y}_{plm}(r, \theta, \phi) = \frac{\Gamma(ip + l + 1)}{\Gamma(ip + 1)} \frac{p}{\sqrt{\sinh r}} \mathcal{P}_{(ip - \frac{1}{2})}^{-l} (\cosh r) Y_{lm}(\theta, \phi). \quad (\text{A.3})$$

The total solution of the equations of motion can be written as:

$$\Phi(t, r, \theta, \phi) = \int_0^\infty dp \sum_{\sigma=\pm 1} \sum_{l=0}^\infty \sum_{m=-l}^{+l} \left[a_{\sigma plm} \mathcal{U}_{\sigma plm}(t, r, \theta, \phi) + a_{\sigma plm}^\dagger \mathcal{U}_{\sigma plm}^*(t, r, \theta, \phi) \right], \quad (\text{A.4})$$

where $\mathcal{U}_{\sigma plm}(t, r, \theta, \phi)$ forms complete basis of mode function, $\mathcal{U}_{\sigma plm}(t, r, \theta, \phi) = \frac{H}{\sinh t} \chi_{p,\sigma}(t) \mathcal{Y}_{plm}(r, \theta, \phi)$. Here $\chi_{p,\sigma}(t)$ forms a complete set of positive frequency function. Also this can be written as a sum of complementary ($\chi_{p,\sigma}^{(c)}(t)$) and particular integral ($\chi_{p,\sigma}^{(p)}(t)$) part, as given by $\chi_{p,\sigma}(t) = \chi_{p,\sigma}^{(c)}(t) + \chi_{p,\sigma}^{(p)}(t)$. Explicitly the complementary and particular integral part can be expressed as [14]:

$$\chi_{p,\sigma}^{(c)}(t) = \chi_{-p,\sigma}^{(c)}(t) = \frac{1}{2 \sinh \pi p} \left[\frac{(e^{\pi p} - i\sigma e^{-i\pi\nu})}{\Gamma(\nu + \frac{1}{2} + ip)} \mathcal{P}_{(\nu - \frac{1}{2})}^{ip}(\cosh t_{\mathbf{R}/\mathbf{L}}) \right. \\ \left. - \frac{(e^{-\pi p} - i\sigma e^{-i\pi\nu})}{\Gamma(\nu + \frac{1}{2} - ip)} \mathcal{P}_{(\nu - \frac{1}{2})}^{-ip}(\cosh t_{\mathbf{R}/\mathbf{L}}) \right], \quad (\text{A.5})$$

$$\chi_{p,\sigma}^{(p)}(t) = \mu^3 \sinh^2 t \sum_{n=0}^{\infty} \frac{1}{(p^2 - p_n^2)} \chi_{p_n,\sigma}^{(c)}(t) \int dt' \chi_{p_n,\sigma}^{(c)}(t'), \quad (\text{A.6})$$

where the parameter ν is defined as, $\nu = \sqrt{\frac{9}{4} - \frac{m_{axion}^2}{H^2}} = \sqrt{\frac{9}{4} - \frac{\mu^3 b}{f_a H^2}} = \sqrt{\frac{9}{4} - \frac{\Lambda_G^4}{f_a^2 H^2}}$.

References

- [1] L. Amico, R. Fazio, A. Osterloh, V. Vedral, Entanglement in many-body systems, Rev. Mod. Phys. 80 (2008) 517, arXiv:quant-ph/0703044.
- [2] R. Horodecki, P. Horodecki, M. Horodecki, K. Horodecki, Quantum entanglement, Rev. Mod. Phys. 81 (2009) 865, arXiv:quant-ph/0702225.
- [3] N. Laflamme, Quantum entanglement in condensed matter systems, Phys. Rep. 646 (2016) 1, arXiv:1512.03388 [cond-mat.str-el].
- [4] E. Martin-Martinez, N.C. Menicucci, Cosmological quantum entanglement, Class. Quantum Gravity 29 (2012) 224003, arXiv:1204.4918 [gr-qc].
- [5] Y. Nambu, Entanglement of quantum fluctuations in the inflationary universe, Phys. Rev. D 78 (2008) 044023, arXiv:0805.1471 [gr-qc].
- [6] D. Campo, R. Parentani, Quantum correlations in inflationary spectra and violation of bell inequalities, Braz. J. Phys. 35 (2005) 1074, arXiv:astro-ph/0510445.
- [7] Y. Nambu, Y. Ohsumi, Classical and quantum correlations of scalar field in the inflationary universe, Phys. Rev. D 84 (2011) 044028, arXiv:1105.5212 [gr-qc].
- [8] G.L. Ver Steeg, N.C. Menicucci, Entangling power of an expanding universe, Phys. Rev. D 79 (2009) 044027, arXiv:0711.3066 [quant-ph].
- [9] D. Mazur, J.S. Heyl, Characterizing entanglement entropy produced by nonlinear scalar interactions during inflation, Phys. Rev. D 80 (2009) 023523, arXiv:0810.0521 [gr-qc].
- [10] J. Maldacena, G.L. Pimentel, Entanglement entropy in de Sitter space, J. High Energy Phys. 1302 (2013) 038, arXiv:1210.7244 [hep-th].
- [11] J. Maldacena, A model with cosmological Bell inequalities, Fortschr. Phys. 64 (2016) 10, arXiv:1508.01082 [hep-th].
- [12] S. Choudhury, S. Panda, R. Singh, Bell violation in the sky, Eur. Phys. J. C 77 (2) (2017) 60, arXiv:1607.00237 [hep-th].
- [13] S. Choudhury, S. Panda, R. Singh, Bell violation in primordial cosmology, Universe 3 (1) (2017) 13, arXiv:1612.09445 [hep-th].
- [14] S. Choudhury, S. Panda, Entangled de Sitter from stringy axionic bell pair I: an analysis using Bunch Davies vacuum, arXiv:1708.02265 [hep-th];
S. Choudhury, S. Panda, Spectrum of cosmological correlation from vacuum fluctuation of Stringy Axion in entangled De Sitter space, arXiv:1809.02905 [hep-th].
- [15] S. Kanno, J. Murugan, J.P. Shock, J. Soda, Entanglement entropy of α -vacua in de Sitter space, J. High Energy Phys. 1407 (2014) 072, arXiv:1404.6815 [hep-th].
- [16] S. Kanno, J. Soda, Infinite violation of Bell inequalities in inflation, arXiv:1705.06199 [hep-th].
- [17] S. Kanno, J.P. Shock, J. Soda, Quantum discord in de Sitter space, Phys. Rev. D 94 (12) (2016) 125014, arXiv:1608.02853 [hep-th].
- [18] S. Kanno, J.P. Shock, J. Soda, Entanglement negativity in the multiverse, J. Cosmol. Astropart. Phys. 1503 (03) (2015) 015, arXiv:1412.2838 [hep-th].
- [19] S. Kanno, Impact of quantum entanglement on spectrum of cosmological fluctuations, J. Cosmol. Astropart. Phys. 1407 (2014) 029, arXiv:1405.7793 [hep-th].

- [20] W. Fischler, S. Kundu, J.F. Pedraza, Entanglement and out-of-equilibrium dynamics in holographic models of de Sitter QFTs, *J. High Energy Phys.* 1407 (2014) 021, arXiv:1311.5519 [hep-th].
- [21] W. Fischler, P.H. Nguyen, J.F. Pedraza, W. Tangarife, Fluctuation and dissipation in de Sitter space, *J. High Energy Phys.* 1408 (2014) 028, arXiv:1404.0347 [hep-th].
- [22] S.R. Coleman, F. De Luccia, Gravitational effects on and of vacuum decay, *Phys. Rev. D* 21 (1980) 3305.
- [23] J.S. Bell, On the Einstein-Podolsky-Rosen paradox, *Physics* 1 (1964) 195.
- [24] M.B. Fröb, J. Garriga, S. Kanno, M. Sasaki, J. Soda, T. Tanaka, A. Vilenkin, Schwinger effect in de Sitter space, *J. Cosmol. Astropart. Phys.* 1404 (2014) 009, arXiv:1401.4137 [hep-th].
- [25] W. Fischler, P.H. Nguyen, J.F. Pedraza, W. Tangarife, Holographic Schwinger effect in de Sitter space, *Phys. Rev. D* 91 (8) (2015) 086015, arXiv:1411.1787 [hep-th].
- [26] S. Ryu, T. Takayanagi, Holographic derivation of entanglement entropy from AdS/CFT, *Phys. Rev. Lett.* 96 (2006) 181602, arXiv:hep-th/0603001.
- [27] S. Ryu, T. Takayanagi, Aspects of holographic entanglement entropy, *J. High Energy Phys.* 0608 (2006) 045, arXiv: hep-th/0605073.
- [28] T. Nishioka, S. Ryu, T. Takayanagi, Holographic entanglement entropy: an overview, *J. Phys. A* 42 (2009) 504008, arXiv:0905.0932 [hep-th].
- [29] M. Rangamani, T. Takayanagi, Holographic entanglement entropy, *Lect. Notes Phys.* 931 (2017), arXiv:1609.01287 [hep-th].
- [30] V.E. Hubeny, M. Rangamani, T. Takayanagi, A covariant holographic entanglement entropy proposal, *J. High Energy Phys.* 0707 (2007) 062, arXiv:0705.0016 [hep-th].
- [31] X. Dong, Holographic entanglement entropy for general higher derivative gravity, *J. High Energy Phys.* 1401 (2014) 044, arXiv:1310.5713 [hep-th].
- [32] J. Camps, Generalized entropy and higher derivative gravity, *J. High Energy Phys.* 1403 (2014) 070, arXiv:1310.6659 [hep-th].
- [33] S. Banerjee, A. Bhattacharyya, A. Kaviraj, K. Sen, A. Sinha, Constraining gravity using entanglement in AdS/CFT, *J. High Energy Phys.* 1405 (2014) 029, arXiv:1401.5089 [hep-th].
- [34] A. Bhattacharyya, M. Sharma, On entanglement entropy functionals in higher derivative gravity theories, *J. High Energy Phys.* 1410 (2014) 130, arXiv:1405.3511 [hep-th].
- [35] S.S. Pal, S. Panda, Entanglement temperature with Gauss–Bonnet term, *Nucl. Phys. B* 898 (2015) 401, arXiv: 1507.06488 [hep-th].
- [36] E. Mottola, Particle creation in de Sitter space, *Phys. Rev. D* 31 (1985) 754.
- [37] B. Allen, Vacuum states in de Sitter space, *Phys. Rev. D* 32 (1985) 3136.
- [38] K. Goldstein, D.A. Lowe, A note on alpha vacua and interacting field theory in de Sitter space, *Nucl. Phys. B* 669 (2003) 325, arXiv:hep-th/0302050.
- [39] J. de Boer, V. Jejjala, D. Minic, Alpha-states in de Sitter space, *Phys. Rev. D* 71 (2005) 044013, arXiv:hep-th/ 0406217.
- [40] R. Brunetti, K. Fredenhagen, S. Hollands, A remark on alpha vacua for quantum field theories on de Sitter space, *J. High Energy Phys.* 0505 (2005) 063, arXiv:hep-th/0503022.
- [41] B.S. Kay, R.M. Wald, Theorems on the uniqueness and thermal properties of stationary, nonsingular, quasifree states on space-times with a bifurcate killing horizon, *Phys. Rep.* 207 (1991) 49.
- [42] R.M. Wald, Quantum Field Theory in Curved Space-Time and Black Hole Thermodynamics, Univ. Pr., Chicago, USA, 1994, 205 pp.
- [43] L. McAllister, E. Silverstein, A. Westphal, Gravity waves and linear inflation from axion monodromy, *Phys. Rev. D* 82 (2010) 046003, arXiv:0808.0706 [hep-th].
- [44] E. Silverstein, A. Westphal, Monodromy in the CMB: gravity waves and string inflation, *Phys. Rev. D* 78 (2008) 106003, arXiv:0803.3085 [hep-th].
- [45] L. McAllister, E. Silverstein, A. Westphal, T. Wrane, The powers of monodromy, *J. High Energy Phys.* 1409 (2014) 123, arXiv:1405.3652 [hep-th].
- [46] S. Panda, Y. Sumitomo, S.P. Trivedi, Axions as quintessence in string theory, *Phys. Rev. D* 83 (2011) 083506, arXiv:1011.5877 [hep-th].
- [47] S. Choudhury, S. Panda, COSMOS- e' -GTachyon from string theory, *Eur. Phys. J. C* 76 (5) (2016) 278, arXiv: 1511.05734 [hep-th];
S. Choudhury, COSMOS- e' - soft higgsotic attractors, *Eur. Phys. J. C* 77 (7) (2017) 469, arXiv:1703.01750 [hep-th];
S. Choudhury, S. Pal, Primordial non-Gaussian features from DBI Galileon inflation, *Eur. Phys. J. C* 75 (6) (2015) 241, arXiv:1210.4478 [hep-th];
S. Choudhury, S. Pal, DBI Galileon inflation in background SUGRA, *Nucl. Phys. B* 874 (2013) 85, arXiv:1208.4433 [hep-th];

- S. Choudhury, S. Pal, Fourth level MSSM inflation from new flat directions, *J. Cosmol. Astropart. Phys.* 1204 (2012) 018, arXiv:1111.3441 [hep-ph];
 S. Choudhury, S. Pal, Brane inflation in background supergravity, *Phys. Rev. D* 85 (2012) 043529, arXiv:1102.4206 [hep-th];
 S. Choudhury, Can effective field theory of inflation generate large tensor-to-scalar ratio within Randall–Sundrum single braneworld?, *Nucl. Phys. B* 894 (2015) 29, arXiv:1406.7618 [hep-th];
 S. Choudhury, B.K. Pal, B. Basu, P. Bandyopadhyay, Quantum gravity effect in torsion driven inflation and CP violation, *J. High Energy Phys.* 1510 (2015) 194, arXiv:1409.6036 [hep-th];
 S. Choudhury, Reconstructing inflationary paradigm within effective field theory framework, *Phys. Dark Universe* 11 (2016) 16, arXiv:1508.00269 [astro-ph.CO];
 S. Choudhury, A. Mazumdar, An accurate bound on tensor-to-scalar ratio and the scale of inflation, *Nucl. Phys. B* 882 (2014) 386, arXiv:1306.4496 [hep-ph];
 S. Choudhury, A. Mazumdar, Primordial blackholes and gravitational waves for an inflection-point model of inflation, *Phys. Lett. B* 733 (2014) 270, arXiv:1307.5119 [astro-ph.CO];
 S. Choudhury, A. Mazumdar, Reconstructing inflationary potential from BICEP2 and running of tensor modes, arXiv:1403.5549 [hep-th];
 S. Choudhury, A. Mazumdar, E. Pukartas, Constraining $\mathcal{N} = 1$ supergravity inflationary framework with non-minimal Kähler operators, *J. High Energy Phys.* 1404 (2014) 077, arXiv:1402.1227 [hep-th];
 S. Choudhury, Constraining $\mathcal{N} = 1$ supergravity inflation with non-minimal Kähler operators using δN formalism, *J. High Energy Phys.* 1404 (2014) 105, arXiv:1402.1251 [hep-th];
 S. Choudhury, A. Mazumdar, S. Pal, Low & high scale MSSM inflation, gravitational waves and constraints from Planck, *J. Cosmol. Astropart. Phys.* 1307 (2013) 041, arXiv:1305.6398 [hep-ph].
- [48] J. Maharana, S. Mukherji, S. Panda, Notes on axion, inflation and graceful exit in stringy cosmology, *Mod. Phys. Lett. A* 12 (1997) 447, arXiv:hep-th/9701115;
 A. Mazumdar, S. Panda, A. Perez-Lorenzana, Assisted inflation via tachyon condensation, *Nucl. Phys. B* 614 (2001) 101, arXiv:hep-ph/0107058;
 D. Choudhury, D. Ghoshal, D.P. Jatkar, S. Panda, Hybrid inflation and brane - anti-brane system, *J. Cosmol. Astropart. Phys.* 0307 (2003) 009, arXiv:hep-th/0305104;
 D. Choudhury, D. Ghoshal, D.P. Jatkar, S. Panda, On the cosmological relevance of the tachyon, *Phys. Lett. B* 544 (2002) 231, arXiv:hep-th/0204204;
 P. Chingangbam, S. Panda, A. Deshamukhya, Non-minimally coupled tachyonic inflation in warped string background, *J. High Energy Phys.* 0502 (2005) 052, arXiv:hep-th/0411210;
 A. Deshamukhya, S. Panda, Warm tachyonic inflation in warped background, *Int. J. Mod. Phys. D* 18 (2009) 2093, arXiv:0901.0471 [hep-th];
 P. Vargas Moniz, S. Panda, J. Ward, Higher order corrections to heterotic M-theory inflation, *Class. Quantum Gravity* 26 (2009) 245003, arXiv:0907.0711 [astro-ph.CO];
 A. Ali, A. Deshamukhya, S. Panda, M. Sami, Inflation with improved D3-brane potential and the fine tunings associated with the model, *Eur. Phys. J. C* 71 (2011) 1672, arXiv:1010.1407 [hep-th];
 A. Bhattacharjee, A. Deshamukhya, S. Panda, A note on low energy effective theory of chromo-natural inflation in the light of BICEP2 results, *Mod. Phys. Lett. A* 30 (11) (2015) 1550040, arXiv:1406.5858 [astro-ph.CO];
 S. Panda, M. Sami, S. Tsujikawa, Inflation and dark energy arising from geometrical tachyons, *Phys. Rev. D* 73 (2006) 023515, arXiv:hep-th/0510112;
 S. Panda, M. Sami, S. Tsujikawa, J. Ward, Inflation from D3-brane motion in the background of D5-branes, *Phys. Rev. D* 73 (2006) 083512, arXiv:hep-th/0601037;
 S. Panda, M. Sami, S. Tsujikawa, Prospects of inflation in delicate D-brane cosmology, *Phys. Rev. D* 76 (2007) 103512, arXiv:0707.2848 [hep-th].
- [49] D. Baumann, TASI lectures on inflation 2009, arXiv:0907.5424 [hep-th];
 D. Baumann, A. Dymarsky, I.R. Klebanov, L. McAllister, Towards an explicit model of D-brane inflation, *J. Cosmol. Astropart. Phys.* 0801 (2008) 024, arXiv:0706.0360 [hep-th];
 D. Baumann, L. McAllister, Advances in inflation in string theory, *Annu. Rev. Nucl. Part. Sci.* 59 (2009) 67, arXiv:0901.0265 [hep-th];
 V. Assassi, D. Baumann, D. Green, Symmetries and loops in inflation, *J. High Energy Phys.* 1302 (2013) 151, arXiv:1210.7792 [hep-th];
 D. Baumann, L. McAllister, Inflation and string theory, arXiv:1404.2601 [hep-th];
 D. Baumann, A. Dymarsky, S. Kachru, I.R. Klebanov, L. McAllister, Holographic systematics of D-brane inflation, *J. High Energy Phys.* 0903 (2009) 093, arXiv:0808.2811 [hep-th];

- H.V. Peiris, D. Baumann, B. Friedman, A. Cooray, Phenomenology of D-brane inflation with general speed of sound, *Phys. Rev. D* 76 (2007) 103517, arXiv:0706.1240 [astro-ph].
- [50] N. Agarwal, R. Bean, L. McAllister, G. Xu, Universality in D-brane inflation, *J. Cosmol. Astropart. Phys.* 1109 (2011) 002, arXiv:1103.2775 [astro-ph.CO];
R. Flauger, L. McAllister, E. Pajer, A. Westphal, G. Xu, Oscillations in the CMB from axion monodromy inflation, *J. Cosmol. Astropart. Phys.* 1006 (2010) 009, arXiv:0907.2916 [hep-th].
- [51] S. Kanno, A note on initial state entanglement in inflationary cosmology, *Europhys. Lett.* 111 (6) (2015) 60007, arXiv:1507.04877 [hep-th].
- [52] F.V. Dimitrakopoulos, L. Kabir, B. Mosk, M. Parikh, J.P. van der Schaar, Vacua and correlators in hyperbolic de Sitter space, *J. High Energy Phys.* 1506 (2015) 095, arXiv:1502.00113 [hep-th].
- [53] A. Aguirre, C.P. Burgess, A. Friedland, D. Nolte, Astrophysical constraints on modifying gravity at large distances, *Class. Quantum Gravity* 18 (2001) R223, arXiv:hep-ph/0105083.
- [54] C.G. Callan Jr., F. Wilczek, On geometric entropy, *Phys. Lett. B* 333 (1994) 55, arXiv:hep-th/9401072.
- [55] P. Svrcek, E. Witten, Axions in string theory, *J. High Energy Phys.* 0606 (2006) 051, arXiv:hep-th/0605206.
- [56] M. Sasaki, T. Tanaka, K. Yamamoto, Euclidean vacuum mode functions for a scalar field on open de Sitter space, *Phys. Rev. D* 51 (1995) 2979, arXiv:gr-qc/9412025.