

## REVIEW ARTICLE

# Bacterial–fungal interactions: ecology, mechanisms and challenges

Aurélie Deveau<sup>1,\*</sup>, Gregory Bonito<sup>2</sup>, Jessie Uehling<sup>3,4</sup>, Mathieu Paoletti<sup>5</sup>, Matthias Becker<sup>6</sup>, Saskia Bindschedler<sup>7</sup>, Stéphane Hacquard<sup>9</sup>, Vincent Hervé<sup>7,8,†</sup>, Jessy Labbé<sup>10,11</sup>, Olga A. Lastovetsky<sup>12,‡</sup>, Sophie Mieszkina<sup>1</sup>, Larry J. Millet<sup>13</sup>, Balázs Vajna<sup>14</sup>, Pilar Junier<sup>7</sup>, Paola Bonfante<sup>15</sup>, Bastiaan P. Krom<sup>16</sup>, Stefan Olsson<sup>17</sup>, Jan Dirk van Elsas<sup>18</sup> and Lukas Y. Wick<sup>19</sup>

<sup>1</sup>Université de Lorraine, INRA, UMR IAM, 54280 Champenoux, France, <sup>2</sup>Department of Plant Soil and Microbial Sciences, Michigan State University, East Lansing, MI 48824, USA, <sup>3</sup>Biology Department, Duke University, Box 90338, Durham, NC 27705, USA, <sup>4</sup>Plant and Microbial Biology, University of California, Berkeley, CA 94703, USA, <sup>5</sup>Institut de Biologie et Génétique Cellulaire, UMR 5095 CNRS et Université de Bordeaux, 1 rue Camille Saint-Saëns, 33077 Bordeaux cedex, France, <sup>6</sup>IGZ, Leibniz-Institute of Vegetable and Ornamental Crops, 14979 Großbeeren, Germany, <sup>7</sup>Laboratory of Microbiology, Institute of Biology, University of Neuchâtel, CH-2000 Neuchâtel, Switzerland, <sup>8</sup>Laboratory of Biogeosciences, Institute of Earth Surface Dynamics, University of Lausanne, CH-1015 Lausanne, Switzerland, <sup>9</sup>Department of Plant Microbe Interactions, Max Planck Institute for Plant Breeding Research, 50829 Cologne, Germany, <sup>10</sup>Biosciences Division, Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA, <sup>11</sup>Department of Biochemistry and Cellular and Molecular Biology, University of Tennessee, Knoxville, TN 37996, USA, <sup>12</sup>Graduate Field of Microbiology, Cornell University, Ithaca, NY 14853, USA, <sup>13</sup>Joint Institute for Biological Science, University of Tennessee, and the Biosciences Division of Oak Ridge National Laboratory, Oak Ridge, TN 37831, USA, <sup>14</sup>Department of Microbiology, Eötvös Loránd University, Pázmány Péter sétány 1/C, 1117 Budapest, Hungary, <sup>15</sup>Department of Life Science and Systems Biology, University of Torino, 10125 Torino, Italy, <sup>16</sup>Department of Preventive Dentistry, Academic Centre for Dentistry, G. Mahlerlaan 3004, 1081 LA, Amsterdam, The Netherlands, <sup>17</sup>State Key Laboratory of Ecological Pest Control for Fujian and Taiwan Crops, Fujian Agriculture and Forestry University (FAFU), Fuzhou 350002, China, <sup>18</sup>Microbial Ecology group, GELIFES, University of Groningen, 9747 Groningen, The Netherlands and <sup>19</sup>Helmholtz Centre for Environmental Research—UFZ, Department of Environmental Microbiology, Permoserstraße 15, 04318 Leipzig, Germany

\*Corresponding author: Université de Lorraine, INRA, UMR IAM, 54280 Champenoux, France. Tel: +33-3-83-39-40-88; Fax: +33-3-83-39-40-69; E-mail: aurelie.deveau@inra.fr

†Present address: Department of Biogeochemistry, Max Planck Institute for Terrestrial Microbiology, 35043 Marburg, Germany

‡Present address: School of Biology and Environmental Science, University College Dublin, Dublin, Ireland.

One sentence summary: Bacterial–fungal interactions are cornerstones of numerous processes that impact ecosystem functions, animal and plant physiology, and industrial activities.

Editor: Gerhard Braus

†Vincent Hervé, <http://orcid.org/0000-0002-3495-561X>

## ABSTRACT

Fungi and bacteria are found living together in a wide variety of environments. Their interactions are significant drivers of many ecosystem functions and are important for the health of plants and animals. A large number of fungal and bacterial families engage in complex interactions that lead to critical behavioural shifts of the microorganisms ranging from mutualism to antagonism. The importance of bacterial–fungal interactions (BFI) in environmental science, medicine and biotechnology has led to the emergence of a dynamic and multidisciplinary research field that combines highly diverse approaches including molecular biology, genomics, geochemistry, chemical and microbial ecology, biophysics and ecological modelling. In this review, we discuss recent advances that underscore the roles of BFI across relevant habitats and ecosystems. A particular focus is placed on the understanding of BFI within complex microbial communities and in regard of the metaorganism concept. We also discuss recent discoveries that clarify the (molecular) mechanisms involved in bacterial–fungal relationships, and the contribution of new technologies to decipher generic principles of BFI in terms of physical associations and molecular dialogues. Finally, we discuss future directions for research in order to stimulate synergy within the BFI research area and to resolve outstanding questions.

**Keywords:** bacterial–fungal interactions; metaorganisms; microbiome; mechanism; microbial logistics

## INTRODUCTION

Bacteria and fungi often share microhabitats where they assemble into dynamic co-evolving communities. Such bacterial–fungal communities have been described to exist in nearly all ecosystems and include microbial species from a wide diversity of fungal and bacterial families (Peleg, Hogan and Mylonakis 2010; Scherlach, Graupner and Hertweck 2013). Interactions between fungi and bacteria play a key role in the functioning of numerous ecosystems: they are cornerstone members of communities driving biochemical cycles, and contribute to both the health and diseases of plants and animals (Fig. 1). Moreover, bacteria and fungi have been exploited by humans for centuries to manufacture food products, antibiotics and secondary metabolites for pharmacology and biotechnological applications (Frey-Klett et al. 2011). As a consequence, by-products of bacterial–fungal interactions (BFI) have been harnessed to improve many human activities in agriculture, horticulture, forestry, environmental protection, food processing, biotechnology and medical applications.

BFI intrinsically modulate the behaviour of either or both of the interacting partners. Such modulation cannot be easily predicted based on our knowledge of the biology of the isolated microorganisms grown in pure cultures. Different levels and degrees of specificity of BFI have been reported. On one end of the spectrum, co-occurrence patterns of bacteria and fungi result from intimate biophysical and metabolic interactions during which bacterial and fungal partners interdependently develop and co-evolve. On the other end, co-occurrence may not be representative of any causal relationships, being the result of stochastic ‘mixing’ within the microbial community. Depending on the degree of interaction, the molecular dialogue between the partners may be very simple, highly refined or absent. Depending on the species involved in BFI, interactions can be highly specific, like the intimate interaction between endofungal bacteria and early emerging fungi (Bonfante and Desirò 2017), or they can involve a broad spectrum of species. For instance, the opportunistic human pathogens *Candida albicans* and *Pseudomonas aeruginosa* frequently interact with each other, but also with

numerous additional bacteria and fungi, respectively (Leclair and Hogan 2010). Such multipartner interactions can occur within a single environment—such as in the oral plaque (Janus, Willems and Krom 2016), in soil (Warmink, Nazir and van Elsas 2009), in a single food product (Kastman et al. 2016) or across multiple environments. Opportunistic microorganisms such as the aforementioned *P. aeruginosa* colonise a wide variety of environments including human tissues, plant root systems and soils, in which they engage in different interactions with local fungal species (Walker et al. 2004). Whatever the environment considered, BFI can produce a diverse range of interactions—from antagonism to mutualism—that influence the biology and ecology of the fungal and bacterial partners at different levels, i.e. with respect to growth, reproduction, transport/movement, nutrition, stress resistance and pathogenicity. The outcomes of these interactions are the combined results of the physical associations (biofilm, free cells, intracellular), the molecular dialogue between the organisms (direct or indirect), and the environmental conditions and/or the host activity (Fig. 1).

Within the past decade, a range of multidisciplinary studies on diverse BFI, which integrate tools from molecular biology, genomics, chemical and microbial ecology, biophysics and ecological modelling, have emerged. The more than 300 studies dealing with BFI, as published within the last five years across divergent fields of research (e.g. medicine, agriculture, environment science, biotechnology and food processing), have culminated in a better understanding of interaction mechanisms and consequences of BFI. Striking mechanistic generalities have emerged that extend beyond known BFI despite the intricacies inherent to each system analysed, as first outlined by Frey-Klett et al. (2011) and Scherlach, Graupner and Hertweck (2013). Such generalist patterns mirror the remarkable similarities shared between plants and animals recently documented in microbiota-assisted host nutrition (Hacquard et al. 2015). Here, we review the main findings obtained with respect to BFI in different fields in the past years, including the latest advances with respect to the roles and mechanisms involved, as well as the emerging opportunities and applications to biotechnology and ecology.



**Figure 1.** Relevance, applications and drivers of BFI. Biotic and abiotic environments filter microbial communities selecting for specific consortia, and conversely interactions between microorganisms influence the biotic and abiotic environment. Bacterial-fungal phenotypes emerge from physical and molecular interactions between members of the microbial community and are highly dependent on time and scale.

## BFI WITHIN COMPLEX NETWORKS OF INTERACTIONS: FROM FUNGAL MICROBIOMES TO METAORGANISMS AND HOLOBIONTS

The exponential development of molecular tools aimed at describing the diversity of microorganisms in many biomes and environments on Earth has brought to light the huge diversity of microorganisms and potential interactions between them (Thompson et al. 2017). As a consequence, the traditional concept of BFI as bipartite bacterial-fungal or bacterial-fungal-host interactions is now shifting towards BFI as complex networks of multiple interacting organisms. In these networks, there may be different levels of complexity depending on the environment and the scale of analysis. The networks can be envisioned at different levels depending on the habitats considered: ranging from networks restricted to microorganisms on abiotic matrices and surfaces such as soils, wood, hydrothermal vents, water pipes and medical catheters (Hervé et al. 2014; Lindsay and Hogan 2014; Urich et al. 2014; Douterelo et al. 2016; de Menezes, Richardson and Thrall 2017) to networks involving higher organisms, in which BFI occur within the microbiomes of hosts such as lichens (Grube et al. 2015), corals (Moree et al. 2013), nematodes (Wang et al. 2014), insects (Aylward et al. 2014), batrachians (Longo and Zamudio 2017) or mammals (Hacquard et al. 2015; Hoyt et al. 2015). In this regard, the interacting microorganisms together may be conceptually regarded as one metaorganism (Olsson, Bonfante and Pawlowska 2017, see box 1).

## The fungal microbiome

The hyphosphere (box 1) provides microhabitats that are colonised by specific bacterial communities (Frey-Klett et al. 2011). In a seminal paper, some bacterial associates of soil fungi were called bacterial 'fungiphiles' (Warmink and van Elsas 2009, box 1). The diversity of these communities can range from a few to several hundreds of species, depending on the fungus and the organ considered (Grube et al. 2015; Wolfe and Dutton 2015; Schulz-Bohm et al. 2016; Ghodsali et al. 2017). Filamentous fungi can produce differentiated tissues (e.g. mycelium, fruiting bodies, spores, mycorrhizae) that are colonised by distinguished sets of microbiomes (Zagriadskaia et al. 2013; Deveau et al. 2016; El-Jurdi and Ghannoum 2017). While some bacteria, such as *Burkholderia* spp., can colonise a large set of fungal species given their abilities to utilise fungal-derived metabolites and overcome fungal defence mechanisms (Haq et al. 2014; Stopnisek et al. 2016; Jung et al. 2018), others may have a more specific and intimate relationship with their fungal hosts (Warmink, Nazir and van Elsas 2009). Similar to plant and animal microbiomes, which are known to contribute to the 'extended phenotype' of their hosts, it is likely that fungal microbiomes also contribute to the biology of their fungal hosts. Indeed, treatments with antibiotics that suppress or alter fungal-associated bacterial communities impaired mycelial growth, secondary metabolite production and/or reproduction (Vahdatzadeh, Deveau and Splivallo 2015; Schulz-Bohm et al. 2016; Mondo et al. 2017; Uehling et al. 2017).

**Box 1. Definitions of terms and concepts**

**Bacterial fungiphile.** Bacterial strain that preferentially associates with fungi and for which the hyphosphere is the main habitat (Warmink and van Elsland 2009).

**Hub microorganism.** Highly interconnected species that drives community responses through microbe-microbe interactions.

**Hyphosphere.** The microhabitat surrounding hyphal cells.

**Keystone species.** Species on which relies the functioning of the community.

**Metaorganism & holobiont.** The metaorganism concept has been defined as “a community of interacting biological entities that is indicated by a metagenome” (Bosch and McFall-Ngai 2011). It thus is a dynamic entity in dependency of the boundaries set by the researchers. Similarly, the holobiont concept is often used in reference to microbiomes associated with hosts, being both parts of this association subjected to evolutionary selection (Bordenstein and Theis 2015). When it comes to microbial interactions, it is often difficult to determine who is the “host”, as is the case in, for example, the fungal-algal-bacterial holobiont (Aschenbrenner et al. 2016). Here, we redefine a holobiont as a “unit of biological organization composed of several distinct genomes, that, in principle, influence the genomic evolution of each other”. This definition is “host neutral”, and avoids the inclusion of temporary opportunistic assemblages, while at the same time focusing on the evolutionary importance of the holobiont concept (Bordenstein and Theis 2015).

**Microbial logistics.** Effective provision of microbes, matter and energy for microbial ecosystem functioning and targeted substrate turnover (Fester 2014).

**-omics.** The suffix “omics” designates an approach that permits to study given molecules in their globality within a sample: metabolomics for metabolites, proteomics for proteins, volatilomics for volatiles, genomics for genes and transcriptomics for transcripts. Moreover, adding the prefix ‘meta’ implies that the omics method will, within technical limits, measure all genes, transcripts, proteins, metabolites and volatiles in a given sample containing more than one organism.

Understanding of the fungal microbiome is an important challenge in food processing and production, as microbiomes are often involved in fermentation of alcoholic beverages (e.g. wine and beers), dairy products (e.g. cheese, sourdough) and other fermented foods (for review, see Wolfe and Dutton 2015), as well as cultivation of edible mushrooms (Bánfi et al. 2015; Murat 2015).

### The endofungal microbiome

Bacteria that live inside fungal cells (i.e. endofungal bacteria or endobacteria) have first been described in the seminal work by Barbara Mosse (Mosse 1970). They were originally considered as biological curiosities; however, numerous emerging studies have demonstrated their omnipresence in fungi, as well as their clear effects on fungal biology (Bonfante and Desirò 2017). To date, endobacteria have been reported in fungi with diverse lifestyles and of broad taxonomic origins, including endophytic Ascomycetes (Hoffman and Arnold 2010; Arendt et al. 2016; Shaffer et al. 2016), symbiotic, pathogenic and endophytic Basidiomycetes (Bertaux et al. 2003; Ruiz-Herrera et al. 2015;

Glaeser et al. 2016) as well as saprotrophic and symbiotic fungi in the Mucoromycota (Partida-Martínez 2017; Uehling et al. 2017; Desirò et al. 2018). The best-studied fungal endobacteria belong to the family Burkholderiaceae, and are associated with early-diverging lineages of terrestrial fungi within the Mucoromycota (Bonfante and Desirò 2017; Uehling et al. 2017). These associations appear to be specific, and have presumably tightly co-evolved over millions of years (Mondo et al. 2012; Desirò et al. 2015; Uehling et al. 2017). This has resulted in host dependency and significant genome reductions for the bacterial endosymbionts (Ghignone et al. 2012; Uehling et al. 2017). Endobacteria can have profound effects on fungal host biology, including aspects of host reproduction (Partida-Martínez et al. 2007; Mondo et al. 2017), growth (Shaffer et al. 2017; Uehling et al. 2017; Desirò et al. 2018), energy dynamics (Salvioli et al. 2016; Vannini et al. 2016), primary metabolism (Lastovetsky et al. 2016; Salvioli et al. 2016; Vannini et al. 2016; Li et al. 2017; Uehling et al. 2017) and secondary metabolism (Rohm et al. 2010; Hoffman et al. 2013).

Several examples of fungi in the Mucoromycota and their endobacteria offer lessons in fungal endobacterial biology. First, the association between *Paraburkholderia rhizoxinica* (formerly *Burkholderia rhizoxinica*) and *Rhizopus microsporus* is mutualistic, whereby the bacterium provides its host with a toxin, which facilitates fungal pathogenicity on rice. Remarkably, the vertically transmitted endobacteria impact fungal reproduction, as their removal abolishes asexual sporulation and significantly reduces mating (Mondo et al. 2017). A recent study leveraged this endobacterial control over fungal mating into identifying reproductive genes in the Mucoromycota, a group of fungi that is notoriously recalcitrant to genetic approaches, as well as reconstructing key reproductive pathways across the fungal kingdom (Mondo et al. 2017). Moreover, studying the pre-symbiotic interaction between *R. microsporus* and *Paraburkholderia* revealed that the fungus undergoes specific lipid metabolic changes in order to accommodate endobacteria, which, when perturbed, shift the interaction from mutualistic into antagonistic (Lastovetsky et al. 2016). Clearly, the *Rhizopus-Paraburkholderia* system is a token of the key role that bacteria can play in modulating the basic biology of their host fungi.

A second example of a well-studied endobacterial-fungal system is the association between members of the arbuscular mycorrhizal fungal family Gigasporaceae, Glomeromycotina) and *Candidatus Glomeribacter gigasporarum* (CaGg, Burkholderiaceae). These bacteria are vertically transmitted between the fungal generations (Bianciotto et al. 2004) and have a strong effect on the pre-symbiotic phase of the fungus. In the pre-symbiotic phase, they raise the fungal bioenergetic capacity, increase ATP production and elicit reactive oxygen detoxification mechanisms (Salvioli et al. 2016). Recent work discovered a new aspect of the endobacterial biology, in that a toxin-antitoxin system was active (Salvioli Di Fossalunga et al. 2017), as well as the whole operon for vitamin B<sub>12</sub> production (Ghignone et al. 2012). This indicates potential metabolic assistance by the endobacterium, not only for the fungal host, but also for the plant mycorrhizal partner. Interestingly, sharing of B-vitamins was also described for the lichen *Lobaria pulmonaria*, where lichen-associated bacteria have been hypothesised to support photosynthesis by provision of vitamin B<sub>12</sub> (Grube et al. 2015).

A third fungal endosymbiont example is the endobacterium *Mycovoidus cysteinexigens*, an endosymbiont of the saprotrophic fungus *Mortierella elongata* (Mortierellomycotina) (Uehling et al. 2017). Despite the close phylogenetic affiliation to CaGg, its impact on host fungal growth is strikingly different. CaGg promotes the growth of its fungal host, while *M. cysteinexigens* decreases

fungus growth (Uehling et al. 2017) suggesting that these later endobacteria utilise host fungal metabolic products. This behaviour likely reflects their ancient divergence from CaMg and co-evolution with the fungal host.

Interestingly, many Mucoromycota, including Glomeromycotina, Endogone and Mortierellamycotina, can host endosymbionts belonging to the Mollicutes group of bacteria (Naumann, Schüssler and Bonfante 2010; Desirò et al. 2014, 2015, 2018); some have been identified as novel types named 'Candidatus Moeniiplasma glomeromycotinum' (CaMg) (Naito et al. 2017). The effect of these Mollicutes endobacteria on their fungal hosts is still being unraveled. Molecular evolution analyses of Glomeromycotina indicate that CaMg is a parasite of the fungus. Experimental work with Mortierellamycotina indicates that the Mollicutes-related endosymbionts are conditional parasites (Toomer et al. 2015; Desirò et al. 2018). Remarkably, genome sequencing of selected CaMg revealed evidence of horizontal gene transfer events, in particular of fungal genes involved in post-translational modification (Naito, Morton and Pawlowska 2015; Torres-Cortés et al. 2015). Ongoing studies now aim at determining the function of CaMg; in particular, the striking observation of the presence of multiple lineages of endobacteria within a single fungal host calls for further scrutiny (Desirò et al. 2014).

In contrast to the endobacteria of Mucoromycota, endofungal bacteria reported in the Ascomycota and Basidiomycota appear to be more transient in nature, yet they can also influence host phenotype and fitness (Hoffman and Arnold 2010; Spraker et al. 2016). Such transient bacterial–fungal associations may be ecologically important in local habitat-associated adaptation, in which the fungal hosts may serve as environmental reservoirs or refuges for the bacteria (Spraker et al. 2016).

The mechanisms by which endofungal bacteria colonise their hosts have been deciphered for only a few examples. *Paraburkholderia rhizoxinica* actively secretes chitinolytic enzymes by means of the type II secretion system, to penetrate the hyphae of the *R. microsporus* host (Moebius et al. 2014). In contrast, *Ralstonia solanacearum* requires the production of the lipopeptide ralsomycin to invade the chlamydospores of its fungal hosts (Spraker et al. 2016). Once inside the mycelium, some bacteria, much like mitochondria, are able to move through dolipore septa (Bertaux et al. 2005). Some can also be vertically transmitted between generations through fungal spores (Spraker et al. 2016), with the bacterial type III secretion system possibly playing a role (Lackner, Moebius and Hertweck 2011). Further studies are necessary to decipher how widespread these mechanisms are among the endofungal bacteria. Interestingly, in some cases endosymbionts influence fungal host biology and the ability of the fungus to interact with its own host through beneficial (Hoffman et al. 2013; Vannini et al. 2016; Guo et al. 2017) or detrimental (Lackner and Hertweck 2011) associations, giving rise to multilevel interkingdom interactions.

Bacterial DNA is often detected in fungal genome-sequencing projects, opening the question of whether endobacteria are more common in fungi than previously thought. Such 'contaminating' DNA could belong to external bacteria or to endobacteria (either transient or stable). With improvements in genome-sequencing technology, it has become possible to assemble entire bacterial genomes from a fungal–bacterial DNA preparation (Uehling et al. 2017). Researchers are urged to keep an open mind to the possibility of endobacterial associates in their fungi before discarding these 'contaminating' bacterial reads from their projects. Although presence of bacterial reads in fungal sequencing projects is indicative of potential endobacterial symbionts, the presence and taxonomic identity

of endofungal bacteria should still be demonstrated by other evidence, such as provided by transmission electron microscopy and fluorescence in situ hybridisation (FISH).

### BFI in complex microbial communities, metaorganisms and holobionts

Despite the increasing number of in-depth analyses of microbial communities in multiple systems, studies that consider fungi and bacteria together are still limited in number. Clearly, Next Generation Sequencing (NGS) offers unprecedented opportunities for obtaining a broad view of potential BFI across habitats (reviewed in de Menezes, Richardson and Thrall 2017), yet it only permits co-occurrence inferences that may not represent true interactions. Network inference can help to identify those microbes that potentially interact. In a recent study, co-occurrence analyses between bacterial and fungal OTUs across 266 soil samples revealed a significant association between bacteria belonging to the genus *Burkholderia* and a wide range of soil fungi (Stopnisek et al. 2016). This ubiquitous association, together with co-cultivation experiments under laboratory conditions, suggests that specific soil bacteria have evolved strategies to utilise fungal-secreted metabolites and overcome fungal defence mechanisms (Stopnisek et al. 2016). Interactions involving hub microorganisms or keystone species (box 1) can be then further investigated at the molecular level. Agler and co-workers thus identified the yeast *Dioszegia* as a fungal hub of the phyllosphere microbiome of *Arabidopsis thaliana*, as well as its bacterial interactants (Agler et al. 2016). This methodology has already identified BF networks and the drivers that govern community assembly in leaf litter (Purahong et al. 2016), soils (de Menezes et al. 2014; Ma et al. 2016; Stopnisek et al. 2016), floral nectar (Álvarez-Pérez and Herrera 2013), plants (Bell et al. 2014; Agler et al. 2016) and human microbiomes (Mukherjee et al. 2014; Trosvik and de Muinck 2015). All these studies revealed non-random associations between fungi and bacteria and an over-representation of positive associations compared to negative ones. Such positive associations are likely to reflect commonalities of habitats between the microorganisms and potential positive interactions. However, they can also be the result of the common colonisation of a habitat via the same selective or dispersal agent, as in the case of some microorganisms in flowers that are transported by bees (Álvarez-Pérez and Herrera 2013). Networks can vary from a few dozens of microorganisms (as in the oral microbiome) to over 50,000 (in soil) (Mukherjee et al. 2014; Ma et al. 2016). Linking network-inferred prediction with functional analyses will represent an important step forward to decipher the potential link between BFI and ecosystem functioning (Ma et al. 2016; Purahong et al. 2016). For instance, the co-occurrence of the lignocellulose decomposer fungi *Clitocybe* and *Mycena* spp. with potential N<sub>2</sub>-fixing bacterial taxa was correlated with nitrogen (N) deposition in the soil during the decay of leaves, indicating that some bacteria may contribute to the N nutrition of fungi while fungi make C available for bacteria (Purahong et al. 2016).

An ecological balance within the microbiome and between the microbiome and the host (host–microbiota homeostasis) has been hypothesised to be fundamental to maintaining the health of both animal and plant hosts (Krom and Oskam 2014; Hacquard et al. 2017). Understanding how microbiomes shift between healthy symbiosis and unhealthy dysbiosis, and how BFI are involved in such process, is therefore of rising interest in many research fields. For example, BFI can be a factor that

modulates human disease if the ecological balance between the partners shifts. This is illustrated by the recurrent interactions between fungi and bacteria in infections of burn wounds, denture stomatitis, lungs of cystic fibrosis and immunocompromised patients, as well as in (recurrent) bowel disease, or related to the use of invasive medical devices (Dhamgaye, Qu and Peleg 2016; Förster et al. 2016). Consequently, BFI in such associations may impact the virulence of both partners of the interactome. In plants, the critical role of the microbiota for suppression of plant pathogens has been extensively reported (e.g. Santhanam et al. 2015; Ritpitakphong et al. 2016; Expósito et al. 2017). Similar to what has been described for human diseases, several plant infections are often associated with dysbiosis and the loss of diversity in the microbiome (Santhanam et al. 2015; Koskella, Hall and Metcalf 2017). The mechanisms leading to bacterial–fungal homeostasis in plant tissues remain unclear, but likely involve a combination of host-dependent and host-independent mechanisms, such as metabolic and nutritional interdependencies among microbes, secretion of antimicrobials and production of protective barriers (Wei et al. 2015; Mousa et al. 2016).

Such complex interactions are starting to be taken into account when designing new strategies to improve the growth and health of crops (Panke-Buisse et al. 2015; Poudel et al. 2016), or treating dysbiosis in animals and plants using microbiome-based strategies (Fraune et al. 2015; Santhanam et al. 2015; Adam et al. 2016). For instance, there is growing awareness that we now need to consider potential synergisms between BF pathogenic communities in order to analyse and treat diseases (Lamichane and Venturi 2015), with an emphasis on the interactions between microorganisms in the context of pathogenesis (Lopes, Azevedo and Pereira 2014). In addition, positive BFI effects on human health might allow to use fungi and/or bacteria as probiotics. Microbiome-based analyses are also used to improve food processes such as cheese or wine making (Pinto et al. 2014; Dugat-Bony et al. 2015; Liu et al. 2017), and could be applied to many other systems including energy production and bioremediation.

### The emerging importance of Archaea among microbiomes

Besides bacteria, Archaea are now also recognised as important members of Earth's biosphere in terms of their contribution to ecosystem functioning (Moissl-Eichinger et al. 2017). They play key roles in global carbon and nitrogen cycles, for instance in methanogenesis, anaerobic methane oxidation (methanotrophy) and ammonia oxidation. Interestingly, Archaea are found in niches where BFI occur, such as decaying wood (Rinta-Kanto et al. 2016), the mycorrhizosphere (Bomberg and Timonen 2009), rhizosphere (Thion et al. 2016), soil (Ma et al. 2016), rumen (Kumar et al. 2015) and human gut (Hoffmann et al. 2013). However, to date, only few studies have investigated the bacterial–archaeal (Raymann et al. 2017), archaeal–fungal (Hoffmann et al. 2013; Kumar et al. 2015) and fungal–bacterial–archaeal (Ma et al. 2016) interactions or co-occurrences. Altogether, this suggests that Archaea should be integrated into the metaorganism concept, especially since they are known to be involved in different microbial interactions including syntrophy (Morris et al. 2013).

## MECHANISMS OF INTERACTION

A suite of molecular mechanisms may underlie BFI in different systems relying on a combination of physical and chemical interactions, as outlined in Frey-Klett et al. (2011) and


illustrated in Table S1, Supporting Information. Such mechanisms were conceptually divided into four classes, i.e. (i) antibiotic involving metabolite exchange, (ii) signalling and chemotaxis involving metabolite sensing and conversion, (iii) physicochemical changes following adhesion and (iv) protein secretion. Clearly, the above division in four mechanistic types allows for overlap, as it is likely that in all four cases signalling, signal perception and modulation of gene expression in either or both of the partner organisms play a crucial role. Hence, we present a strong focus on the ways by which BFI depend on signal (or metabolite) exchange. We also focus specifically on recent advances on physical interactions during BFI and the particular importance of 'microbial logistics' in BFI.

### Signalling and recognition during BFI

Whether and to what extent fungi and bacteria have the ability to perceive and recognise other microorganisms is a question that animates the BFI field for years. Transcriptomic analyses of several BFI have demonstrated that both fungi and bacteria react to the presence of the partner microorganism and respond differentially depending on the interacting partners (e.g. Mela et al. 2011; Sztajer et al. 2014; Gkarmiri et al. 2015; Haq et al. 2017; Tomada et al. 2017). Several cues may be used by the microorganisms for mutual detection, and most are based on small signalling molecules (Scherlach and Hertweck 2017) (Fig. 2). The underlying modes of action vary as well as specificity; from highly specific signals which are solely perceived as a direct sign of the presence of the interacting partner, to compounds that interfere with signalling pathways in the interacting partner and induce a specific response. This second class of compounds is the most reported one in the literature so far.

One example is quorum sensing (QS). QS has long been considered to constitute a means by which bacteria sense and communicate their population density to coordinate their activities. Recently, QS was shown to be also involved in fungal processes such as morphogenesis, germination, apoptosis, pathogenicity and biofilm development (reviewed in Wongsuk, Pumeesat and Luplertlop 2016). Furthermore, both bacterial and fungal QS molecules were shown to play significant roles in cross-kingdom signalling (Cugini et al. 2007; Stanley et al. 2014; Sztajer et al. 2014; Dixon and Hall 2015). Indeed, certain bacteria react to fungal QS molecules (e.g. farnesol, tyrosol, phenylethanol, tryptophol; Wongsuk, Pumeesat and Luplertlop 2016), and, conversely, fungi may react to bacterium-secreted compounds (e.g. quinolone signals, homoserine lactones; Dixon and Hall 2015; Fourie et al. 2016). Such interkingdom signalling is likely to be a common mechanism of communication between microbes in mixed fungal–bacterial biofilms in which these molecules are abundantly produced (Trejo-Hernández et al. 2014; Dixon and Hall 2015; Fourie et al. 2016). This intricate dialogue has been particularly well studied in *C. albicans*–*P. aeruginosa*/S. *gordonii*/S. *aureus* interactions (Lindsay and Hogan 2014). Signalling may be involved in a broader number of habitats since QS may also intervene in bacterial endofungal symbioses (Kai et al. 2012).

Other soluble compounds released by fungi are also sensed by bacteria. Examples are organic acids, sugars, polyols and even toxins. These compounds induce bacterial chemotaxis towards the hyphae of fungi that excrete them. Among these compounds, oxalic acid is of peculiar interest since it induces chemotaxis in the soil bacterium *Collimonas* without being consumed (Rudnick, van Veen and de Boer 2015; Haq 2016). This is in contrast to most other compounds (e.g. glycerol) that are later used as a source of nutrients by fungal-associated bacteria




**Figure 2.** Short and long distance signalling in BFI. Diverse small molecules, either soluble or volatile, are perceived as a cue for the presence of the fungal/bacterial interactant during BFI. These molecules affect positively or negatively the fungal/bacterial partner, but also sometimes the hosts of BFI or additional organisms.

(Boersma et al. 2009; Haq et al. 2016). Oxalic acid would therefore serve as a sole probe of the presence of fungi in the present case.

The importance of volatile organic compounds (VOCs) in BFI signalling has long been overlooked. However, reports involving 'long-distance' signalling during BFI through VOCs originating from bacteria (Briard, Heddergott and Latgé 2016; Jones et al. 2017), fungi (Schmidt 2015), or synergistically from both partners (Spraker et al. 2014; Vahdatzadeh, Deveau and Splivallo 2015; Schmidt et al. 2017; Uehling et al. 2017) have recently accumulated. VOCs encompass a broad range of small compounds that easily diffuse through water- and gas-filled pores or tissues (reviewed in Effmert et al. 2012; Schmidt et al. 2015). In addition to their well-described fungistatic and bacteriostatic activities (Cordero et al. 2014; Cernava et al. 2015), VOCs such as terpenes or dimethyl sulphide stimulate microbial activities during BFI. For instance, the VOCs produced by *P. aeruginosa* stimulate the growth of the opportunistic pathogen *Aspergillus fumigatus*, favouring invasion of lung parenchyma by the fungus (Briard, Heddergott and Latgé 2016). Conversely, the plant-pathogenic fungus *Fusarium culmorum* produces terpenes that induce motility in the bacterium *Serratia plymuthica* (Schmidt et al. 2017). Interestingly, VOC production is highly influenced by nutrient availability (Hacquard 2017), and it has been proposed that microorganisms sense changes in their environments via shifts in VOC blends, adapting their behaviour accordingly (Garbeva et al. 2014). Intriguingly, some VOCs, such as the terpene sordorifen, are produced by both fungi and bacteria. This has led to the

hypothesis that VOCs may serve as a *lingua franca* between microorganisms (Schmidt et al. 2017). Elucidating VOC perception mechanisms in both fungi and bacteria may answer the question whether a shared language is used by bacteria and fungi during their interactions. To date, volatile receptors have not been identified in either fungi or bacteria and the effects of VOCs on cell membrane depolarisation-based signalling during BFI remain to be measured.

Lastly, fungi may also recognise bacteria during BFI using receptors similar to plant and animal immune receptors that detect microbe-associated molecular patterns (MAMPs). Transcriptomic data have recently revealed that fungi react to similar MAMPs as plants and animals (Ipcho et al. 2016), and a recent survey of fungal genomes has uncovered a repertoire of putative Nod-like immune receptors or NLRs (Dyrka et al. 2014; Uehling, Deveau and Paoletti 2017). Some NLRs could directly recognise the presence of these MAMPs in the environment. Noteworthy is the fact that a subset of NLRs has the ability to rapidly generate new binding specificities through recombination of tandem repeat sequences (Dyrka et al. 2014; Uehling, Deveau and Paoletti 2017) that could favour fast adaptation to new ligands. A fungal lectin that binds bacterial lipopolysaccharide was also found to be upregulated during the interaction of the fungus *Laccaria bicolor* with different soil bacteria (Deveau et al. 2014; Wohlschlager et al. 2014). Whether these different receptors trigger immunity-like responses or are used to detect more generic BFI still needs to be determined.



**Figure 3.** Microbial logistics in BFI. Fungal hyphae can efficiently colonise heterogeneous environmental habitats, create new microhabitats and thereby enable a variety of emerging ecosystem processes and services that can be beneficial or detrimental to bacteria. The mycosphere functions depend on the environment and the microorganisms involved and are highly susceptible to disturbance.

### Mycelia as networks for bacterial transport

The spatial structure of the microbial habitat has been recognised to be crucial for its ecology as it drives the composition and activity of microbiomes (Andersson et al. 2014; Tecon and Or 2017). Spatial aspects also drive BFI (Harms, Schlosser and Wick 2011) and their better understanding will assist their use in microbial resource management. Similar to logistics of human resources and goods, microbial logistics (box 1) are essential for the functioning of microbial systems (Fig. 3). Given that mycelia vastly extend in soils (up to  $10^2$  m g<sup>-1</sup>,  $10^3$  m g<sup>-1</sup> and  $10^4$  m g<sup>-1</sup> length in arable, pasture and forest soils, respectively) (Ritz and Young 2004; Joergensen and Wichern 2018), mycelia can be considered to constitute ideal transport paths and scaffolds for bacteria. The fractal structure of mycelia enables fungi to effectively exploit the three-dimensional space and to easily adapt to environmental disturbances. Fungi also cope well with heterogeneous distribution of nutrients (Boswell et al. 2007). A relevant feature of microbial logistics related to mycelial growth is the translocation of compounds between 'feeder' hyphae growing in optimal environments to hyphal expansion/exploration of more unfavourable areas (i.e. resource transport, Fig. 3). Likewise, fungi recycle and re-allocate their hyphal biomass from substrate-depleted regions to the benefit of exploratory colonisation of new habitats (Fricker et al. 2017). Hydrophobic cell wall proteins (hydrophobins) further enable hyphae to cross air-water interfaces and access heterogeneously

distributed nutrients in vadose environments. Important for BFI ecology is the observation that hyphae serve as dispersal vectors for motile bacteria ('fungal highways', Kohlmeier et al. 2005; see <https://www.youtube.com/watch?v=AnsYh6511lc> for a time lapse movie). In soil, fungal hyphae may thereby preferentially invade the larger pores that are most likely air-filled under typical field conditions (Falconer et al. 2012) and hence allow for bacterial dispersal at vadose conditions. This enables random and directed (e.g. chemotactic) access to new habitats and nutrients (Furuno et al. 2010). For instance, experiments and model simulations showed that mycelia-based bacterial dispersal stimulates contaminant biodegradation in situations where chemicals and/or bacteria are heterogeneously distributed and where the active movement of bacteria to pollutant reservoirs is limited by physical barriers (e.g. air-filled pores) (Banitz et al. 2011; Tecon and Or 2016; Worrich et al. 2016). The hyphosphere is also an ideal hotspot for the foraging of bacterial prey populations (Otto et al. 2016; Otto, Harms and Wick 2017) and for horizontal gene transfer, including those for antibiotic resistance, by facilitating dispersal and preferential contact of bacteria in the hyphosphere (Zhang et al. 2014; Berthold et al. 2016; Nazir et al. 2017). Mycelia-facilitated bacterial dispersal may likewise promote new niche colonisation (Warmink and van Elsas 2009; Martin et al. 2012; Simon et al. 2017) and contribute to bacterial food spoilage (Lee et al. 2014), or the co-invasion of tissues during pathogenesis (Schlecht et al. 2015; Jung et al. 2018). It may be a critical issue in the medical field, as recent studies have revealed

the existence of a variety of diverse mycobiomes related to human niches (Kalan et al. 2016; El-Jurdi and Ghannoum 2017).

### Is it all about food acquisition? New aspects of nutrient-based BFI

It has long been known that many BFI, whether antagonistic or synergistic, rely on competition or cooperation for the acquisition of nutrients, both organic and inorganic ones (Fig. 3). Competition for nutrients has led to the development of a large chemical arsenal in both fungi and bacteria over the millions of years of interaction. Antimicrobial peptides (e.g. cospin—Essig et al. 2014), biosurfactants (e.g. surfactin, nunnamycin—Raaijmakers et al. 2010; Hennessy et al. 2017), phenol and quinone derivatives (e.g. penicillin, atromentin—Kong, Schnepfer and Mathee 2010; Reen et al. 2016; Tauber et al. 2016), pyrrol nitrin (Costa et al. 2009), phenazines (e.g. pyocyanin—Morales et al. 2010), QS inhibitors (Scopel et al. 2013; de Carvalho et al. 2016) to name a few, are all microbial compounds that are naturally involved in BFI (Table 1). These compounds act through a wide variety of mechanisms that include cell membrane disruption, inhibition of cell wall biosynthesis and primary metabolism, formation of reactive oxygen species against a fungus, starvation or disruption by a fungus of bacterial QS signalling (Table 1). The production of these compounds varies depending on the organisms and on environmental conditions, as exemplified by the interaction between *P. aeruginosa* and *A. fumigatus* or *C. albicans* (Lindsay and Hogan 2014; Ferreira et al. 2015). In response, defensive mechanisms (e.g. active efflux of antibiotics or degrading enzymes) have also been developed by target microorganisms to protect themselves (Künzler 2015). The development of protection mechanisms against toxins can also lead to cooperative behaviours between toxic fungi and bacteria as in the case of the plant pathogens *B. glumae* and *F. graminearum* (Jung et al. 2018). Chemical warfare in BFI can be exploited to search for new drugs and antibiotics (Reen et al. 2016). A number of novel compounds, e.g. glionitrin A or new members of enacyloxin family, has been uncovered through BFI analyses in the past years (Park et al. 2009; Ross et al. 2014; Tyc et al. 2014; Barkal et al. 2016). High-throughput screening of BFI has been developed to uncover cryptic or new secondary metabolites (Tyc et al. 2014; Navarri et al. 2016). Antibiotics are probably the most commonly sought-after compounds; however, other compounds such as QS inhibitors could also prove to be valuable (Scopel et al. 2013; de Carvalho et al. 2016). Given the huge unexplored metabolome space, there is great potential for the discovery of novel therapeutic approaches or methods to limit food spoilage (Debbab et al. 2010; Navarri et al. 2016). For instance, lactic acid bacteria, via the production of organic acids, hydroxyl fatty acids, hydrogen peroxide or reuterin, may restrict food spoilage (Gänzle 2015). Conversely, some compounds may be detrimental, as exemplified by the production of rhizoxin and rhizonin toxins through BFI in soybean fermentations, which can cause hepatic lesions when ingested (Rohm et al. 2010).

Less recognised is maybe the importance of BFI in food webs and nutrient cycling. Numbers of bacteria have the ability to degrade fungal cell walls, and bacteria are likely to have an important role in fungal biomass decomposition (Brabcová et al. 2016; Lladó, López-mondéjar and Baldrian 2017). Lysis of fungal cells by bacteria also stimulates biogeochemical processes such as carbon flow within the mycorrhizosphere (Ballhausen and de Boer 2016) or cellulose degradation from plant biomass as exemplified by the activities of the forest soil

bacterium *Clostridium phytofermentans* (Tolonen et al. 2015). Plant biomass degradation often involves the action of both bacteria and fungi (Žifčáková et al. 2017). In the case of fungus-growing termites of the order Macrotermitinae, it has been demonstrated that the gut microbiomes of both fungal ectosymbiont (*Termytomycetes*) and termite (workers) participate in plant biomass decomposition by providing a full set of complementary carbohydrate-active enzymes (Poulsen et al. 2014). In addition to decomposition, a large array of rhizosphere bacteria can directly consume fungal exudates, and so fungal hyphae may be an important source of nutrients in this habitat as well as in soil (Rudnick, van Veen and de Boer 2015). Some bacteria can kill and consume living fungi (i.e. mycophagy, Fig. 3). *Collimonas fungivorans* is the best-described ‘mycophagous’ bacterium so far, whereas other bacteria such as *S. marcescens* can also live off living fungi (Rudnick, van Veen and de Boer 2015; Ballhausen and de Boer 2016; Hover et al. 2016). While *Collimonas* relies on the production of secondary metabolites and chitinases to destabilise and degrade fungal cell walls (Mela et al. 2012), the killer activity of *S. marcescens* is independent of chitinase production and relies instead on the ability to form biofilms on the hyphae (Hover et al. 2016). The abilities to produce antifungal compounds are phylogenetically conserved in collimonads, suggesting the existence of co-evolution processes in this nutrient-based BFI (Ballhausen and de Boer 2016). Fungi may also be able to take advantage of their bacterial partner to improve their nutrition. Pion et al. (2013) demonstrated that the fungus *Morchella crassipes* is able to exploit bacterial biomass through a sophisticated mechanism coined bacterial farming, in which the fungus first feeds the bacterium *P. putida* and then harvests this self-created C source.

By contrast, mycelia of fungi (*F. oxysporum* and *Lyophyllum* sp. strain Karsten) and oomycetes (*Pythium ultimum*) may enable bacterial activity by nutrient and water transfer from the hyphae to the bacterial cells exposed to oligotrophic habitats (Worrich et al. 2017) or favour microbial activity in dry soils (Guhr et al. 2015) (Fig. 3). Mycelia have also been found to mobilise entrapped polycyclic aromatic hydrocarbons (PAHs) via vesicle-bound cytoplasmic transport (‘hyphal pipelines’, Furuno et al. 2012) and to render them available to degrader bacteria (Fester et al. 2014; Schamfuß et al. 2013). Altogether, we see an emerging picture of fungi promoting ecosystem functioning in heterogeneous habitats by transporting resources from high nutrient level and water activity areas to nutrient-poor and dry areas.

### BFI mediated habitat modification

Bacteria and fungi can indirectly interact by modifying their environment in ways that positively or negatively affect their partners (i.e. niche modulation, Fig. 3). pH has been frequently reported as an important factor involved in tinkering with BFI (Frey-Klett et al. 2011). Fungi sense and actively modulate the pH in their surroundings (Nazir et al. 2010; Bignell 2012; Braunsdorf, Mailänder-Sánchez and Schaller 2016). For instance, *Lyophyllum* sp. strain Karsten growing through soil was shown to raise the soil pH from levels below pH 5.0 to just above this threshold for survival of the pH sensitive *Variovorax paradoxus* and other fungal-associated bacterial strains (Nazir et al. 2010). Also, *C. albicans* has been shown to influence the pH of the phagolysosome to increase its chances of survival in phagocytic cells of the immune system (Vylkova and Lorenz 2014; Vylkova 2017). In addition, in combination with *Streptococcus mutans*, a cariogenic acid producing oral bacterium, *C. albicans* actively raises the environmental pH (Willems et al. 2016). Increasing the pH from acid

Table 1. Illustration of the diversity of antimicrobial compounds and of their targets produced by bacteria and fungi during their interactions.

Class of compounds	Compound	Producer	Target	Natural environment	Mechanisms of action	References
Antimicrobial aldehyde	Reuterin	B: <i>Lactobacillus reuteri</i>	Multiples: bacteria, fungi and more	Food	Oxidative stress	Ganzel 2015
Antimicrobial peptide	Copsin	F: <i>Coprinosia cinerea</i>	B: Gram + and Gram –	Dung	Interference with cell wall synthesis	Essig et al. 2014
Antimicrobial peptide	Micasin	F: <i>Microsporium canis</i>	B: Gram + and Gram –	Human skin	Interference with protein folding	Zhu et al. 2012
Antimicrobial peptide	Dipeptide cis-cyclo (Leucyl-Tyrosyl)	F: <i>Penicillium</i> sp.	B: <i>Staphylococcus aureus</i>	Sponge associated	QS inhibition	Scopel et al. 2013
Lipopeptide	Viscosin	B: <i>Pseudomonas</i> sp.	Multiples: bacteria, fungi and more	Multiple, mainly soils and plants but also human gut, deepsea...	Membrane permeabilisation	Raaijmakers et al. 2010
Organo-halogenic metabolite	Pyrrol nitrin	B: Some Gram negative bacteria	F: broad spectrum	Soil	Inhibition of electron transport	Costa et al. 2009
Phenazines	Pyocyanine	B: <i>Pseudomonas aeruginosa</i>	F: <i>Candida albicans</i> , <i>Aspergillus fumigatus</i>	Mammals, plants	Redox and iron homeostasis	Briard, Heddergott and Latgé 2016; Morales et al. 2013
QS molecule	Quinolones	B: <i>Pseudomonas aeruginosa</i>	F: <i>Aspergillus fumigatus</i> , <i>Candida albicans</i>	Mammals, plants	Biofilm disruption	Reen et al. 2016
QS molecule	Farnesol	F: <i>Candida albicans</i>	B: <i>Pseudomonas aeruginosa</i>	Human	Interference with PQS signalling	Cugini et al. 2007
Siderophore	Pyoverdine	Multiples : bacteria and fungi	Multiples: bacteria, fungi and more	Multiple	Iron scavenging	Deveau et al. 2016
VOC	Terpenes	F: <i>Fusarium culmorum</i>	B: <i>Serratia</i> and <i>Collimonas</i>	Soil	Unknown—effect on chemotaxis	Schmidt et al. 2017
VOC	Blends of volatiles	B: <i>Collimonas</i> sp.	F: soil fungi	Soil	Unknown—effects on morphology and sporulation	Garbeva et al. 2014

towards a more neutral value directly stimulates overall bacterial growth and metabolism, as low pH commonly inhibits the growth of most bacteria.

Recent studies also identified oxygen level as an important BFI modulator, particularly for *C. albicans*—bacteria interactions. Early reports indicated that biofilms of *C. albicans* provide an anoxic environment (Bonhomme et al. 2011). This was later confirmed by co-culturing *C. albicans* with a variety of strict anaerobic bacterial species (Fox et al. 2014). In the oral cavity, rapid respiration by *C. albicans* and several other *Candida* species creates an anaerobic niche by reducing the level of dissolved oxygen (Lambooij et al. 2017). This favours anaerobic bacteria and antagonises aerobic ones, thereby directly influencing the composition of the microbiome (Janus, Willems and Krom 2016; Janus et al. 2017). Notwithstanding the afore described anaerobism, aerobic respiration is facilitated by the structure of *Candida* biofilms, and inhibition of respiration (e.g. by bacterial metabolites such as phenazines) inhibits biofilm formation by the fungus (Morales et al. 2013). Conversely, ethanol production by *C. albicans* stimulates phenazine production by *P. aeruginosa* and biofilm formation by the bacteria through a feedback loop, which theoretically increases virulence of both microorganisms (Chen et al. 2014). In light of the diversity of other fungi commonly found in the oral cavity, this oxygen-mediated effect may play an important role in more BFI in this habitat as well as many other human-associated niches.

BFI can also occur indirectly, via host behaviour modulations. For instance, bacteria and fungi induce different innate immune defences in the nematode *Caenorhabditis elegans* (Pukkila-Worley, Ausubel and Mylonakis 2011). By this means, co-infection by bacteria and fungi can alter the outcome of the disease and favour or reduce the development of the pathogens (Arvanitis and Mylonakis 2015). *Candida albicans* and *S. aureus* resulted in increased end-organ damage in murine peritonitis and higher mortality compared with single-pathogen infection. This was mediated by higher levels of circulating inflammatory cytokines (Peters and Noverr 2013). Interplays between bacteria, fungi and the innate ‘immune systems’ are also expected in plants (Hacquard et al. 2017).

### Use of -omics to obtain an integrated view of BFIs

The molecular dialogue that occurs during BFI usually relies on intricate and multiple cell responses as highlighted in Table S1, Supporting Information. ‘Omics’ approaches are ideally suited to address such dialogues and -omics tools can be used to analyse BFI from ‘simple’ *in vitro* dual interactions to complex natural multispecies interactions (box 1). The past years have seen a multitude of applications of -omics to BFI (e.g. Mela et al. 2012; Deveau et al. 2014; Phelan et al. 2014; Benoit et al. 2015; Gkarmiri et al. 2015; Lamacchia et al. 2016; Li et al. 2017; Haq et al. 2017; Schmidt et al. 2017; Uehling et al. 2017; Jung et al. 2018). As an overriding theme, responses of partner organisms were commonly found, yet the magnitudes of the responses varied greatly, probably reflecting dependency on the types of interactions, their context, and the technology used. Interestingly, most studies demonstrated regulation of primary metabolisms including nutrient transporters, stress response, cell wall remodelling and secondary metabolite production during BFI. Noteworthy is the fact that genes/proteins with unknown functions, or showing restricted phylogenetic distribution, often represent a significant part of genes regulated in BFI. Emerging studies have been made on tripartite interactions between fungi, bacteria and a host,

shedding light on the complex cross-talks occurring (Kurth et al. 2015; Vannini et al. 2016).

Complex microbial communities, being most realistic, should be examined using the combination of such analyses. The following questions emerge as relevant: ‘Who is there?’, ‘What are they capable of?’, ‘Who is actively doing what?’ and ‘What are the factors that modify the output of the interaction?’ Combining metagenomics and metaproteomics analyses, Grube deciphered the multifaceted roles of the bacteriome of the lichen *L. pulmonaria* (Grube et al. 2015). In this fungus-alga-bacteria symbiosis, more than 800 bacterial species contributed to the nutrient supply of the lichen, helped its resistance against fungal pathogens and abiotic stress and provided essential hormones and vitamins. Similarly, by using a combination of multi-omics approaches and soil biological techniques, Nuccio et al. demonstrated, for the first time, that the AMF *Glomus hoi* in *Plantago lanceolata*, significantly modified 10% of the bacterial community in decomposing litter (Nuccio et al. 2013). Moreover, the AMF was shown to affect the physicochemical environment in the decomposing litter by preferentially exporting N, for which it appeared to acquire N primarily in the inorganic form. This implied that the export of N from litter is one mechanism by which AMF alter the composition of the bacterial community and decomposition processes in soil.

In addition, to pinpoint functional activities within microbiomes, metaomics approaches help in determining the active players in natural conditions in microbiomes of cheese, soil or the human gut (Huttenhower et al. 2012; Dugat-Bony et al. 2015; Perazzolli et al. 2016; Ghodsavali et al. 2017). Identifying keystone members of such microbiomes and their responses to perturbations is a current challenge of microbial ecology. To allow such studies, synthetic communities may be designed that reproduce patterns of community formation and dynamics of natural systems as well as their functional outputs. So far, much progress has been achieved in fermented food ecosystems (Wolfe and Dutton 2015). In surface-ripened cheese, key functions involved in cheese maturation process such as carbohydrate, lipid and protein metabolisms were highlighted using synthetic bacterial and fungal communities (Dugat-Bonny 2015). The consumption of lactate produced by *Lactobacillus lactis*, by the fungi *Debaryomyces hansenii* and *Geotrichum candidum* was evidenced by a high level of lactate dehydrogenase transcripts (Dugat-Bonny 2015). Moreover, the dominance of *Staphylococcus equorum* in cheese was maintained due to the presence of the fungus *Scopulariopsis* sp. via a molecular mechanism based on the iron utilisation pathways such as a homolog of the *S. aureus* staphyloferrin B siderophore operon pathway (Kastman et al. 2016).

One of the future challenges will be to take into account the spatial and temporal scales of BFI in the analyses. Even though fungi and bacteria co-colonise the same habitat, they do not have the same lifestyle in terms of colonisation area. This is particularly true for soils, in which bacterial habitats may be reduced to a soil particle of a mm<sup>3</sup> or specific zones in a biofilm on a root, while the hyphae of the fungus with which they locally interact forage across centimetres to meters and also interact with other plants, wood debris, microorganisms and microfauna. We here argue that there are fundamental differences in the way that bacteria and fungi respond to biotic and abiotic cues. For example, plant-associated bacterial communities show more resistance to perturbations such as in land use and pH modification than plant-associated fungal communities, while the fungal communities are more resistant to drought than bacterial communities (Uroz et al. 2016). Moreover, the interactive populations tend to be spatiotemporally heterogeneous, so each part

of the interacting system may be (slightly) different from each other, at different points in time during development. In addition, BFI tend to be dynamic (Young and Crawford 2004; Hennessy, Stougaard and Olsson 2017). Current methodologies used to analyse bacterial and fungal microbiomes do not allow one to take into account such complex spatiotemporal organisation. However, their use in combination with microscopy, FISH and analytic techniques such as Raman spectroscopy, Imaging Mass Spectrometry or nanoSIMS may help to overcome this limitation (Behrens et al. 2008; Kaltenpoth, Strupat and Svatoš 2016; Wang et al. 2016).

## FUTURE PERSPECTIVES OF BFI RESEARCH

As highlighted in this review, important progress has been made in the understanding of BFI in model microorganisms, as well as in the description of complex microbial communities involving BFI. Within the last two decades, it has become clear that BFI are crucial to the functions in both natural and anthropogenic ecosystems, including human health. At the ecosystem scale, BFI present all types of outcomes, from positive to negative. As a result, on the one hand they represent a great potential to be harnessed, for instance in sustainable agriculture. On the other hand, the recognition of BFI with negative properties, for instance in human health, could lead to improved therapeutics. However, there is still an important gap between studies performed in laboratory conditions and the 'in vivo' reality that impedes our ability to extrapolate generic principles of BFI at the (eco)system scale. The rapid technological advances in methodological fields related to the study of microorganisms may help in reaching such goal. The manipulation of host-associated microbiomes using either synthetic microbial communities, dilution of natural communities, CRISPR-Cas9, *Agrobacterium*-mediated and other transgenesis tools or antibiotic manipulation of microbial communities and/or germ-free hosts combined with modelling will help to identify the driving factors of BFI and of their interactions with their hosts and/or environment. Moreover, although the number of researchers integrating BFI into their studies is expanding, the field needs to become more interdisciplinary. As a result, we expect that both the methodological aspect and the interdisciplinary contribution will bring new development in the BFI research field.

Finally, BFI could also have a broader impact in science if they are used as model systems to analyse complex interactions. Indeed, apart from being an object of study, the BFI holobiont also provides an interesting and relatively simple model for the study of eukaryote–bacterial interactions. One advantage is the fact that many fungi are haploid, easy to transform (Michielse et al. 2005), and may be grown both in the absence or presence of bacterial partners. In this way, the BFI holobiont can become a model system for the assessment of evolutionarily conserved molecular interactions between eukaryotic cells and bacteria. A key characteristic of eukaryotic metaorganisms/holobionts is the modulated recognition of bacterial symbionts by the hosts' innate immune systems, welcoming mutualists and resisting pathogens (Artis 2008; Zamioudis and Pieterse 2012). It has long been proposed that fungi, similar to plants and animals, possess an innate immune system (Paoletti and Saupe 2009; Salvioli et al. 2016), and have receptors for recognising bacteria (Dyrka et al. 2014; Uehling, Deveau and Paoletti 2017) and indeed do so with fast transcriptomic responses (Ipcho et al. 2016). This opens up an exciting avenue of research into the conservation of innate immune systems across

phylogenetically distant eukaryotes. BFI also serve as useful models for the study of evolutionary theory. For example, how do symbiotic bacteria–eukaryote interactions remain stable under different environmental conditions and over time (Olsson, Bonfante and Pawłowska 2017). Fungal mycelia have recently been proposed to be a driving factor of the evolution of bacterial diversity by enabling preferential contact of spatially distinct bacteria and acting as focal point for horizontal gene transfer (Zhang et al. 2014; Berthold et al. 2016). Thus, BFI may serve as models to study other eukaryotes–prokaryotes interactions, in an analogous way to how fruit flies or worms are used as models to study processes occurring in human cells (Olsson, Bonfante and Pawłowska 2017). Fungal and bacterial model systems have the advantages of being fast-growing, and easy to manipulate and track genetically. Based on these premises, BFI research should expand rapidly, not only to better understand the fundamental processes involved in BFI across research fields, and commercial and industrial settings, but also to take advantage of the fantastic properties of BFI to exploit them as model systems.

## SUPPLEMENTARY DATA

Supplementary data are available at [FEMSRE](https://femsre.oup.com/femsre/article-abstract/42/3/335/4875924) online.

## ACKNOWLEDGEMENTS

This cooperative work was initiated following the workshop on Bacterial–Fungal Interactions at the 2016 European Conference on Fungal Genetics (ECFG) and the authors would like to acknowledge the ECFG to have supported the organisation of such networking workshop. We thank Guillaume Cailleau for his critical help in figure designing. We also thank the anonymous reviewers for their comments and suggestions which greatly increased the quality of this manuscript.

## FUNDING

AD and SM are supported by the French National Research Agency through the Laboratory of Excellence ARBRE (ANR-12-LABXARBRE-01) and the INRA Metaprogram MEM. BPK is supported by a grant from the University of Amsterdam for research into the focal point 'Oral Infections and Inflammation'. GB was supported through the US National Science Foundation (NSF) DEB 1737898 and Michigan State University AgBioResearch NIFA project MICL02416. The work of LYW was co-funded by the Deutsche Forschungsgemeinschaft (DFG) CRC 1076 'AquaDiva' and the Helmholtz Centre for Environmental Research. LM, JL and JU were supported by the US Department of Energy, Office of Science, Biological and Environmental Research as part of the Plant-Microbe Interfaces Scientific Focus Area (<http://pmi.ornl.gov>). Oak Ridge National Laboratory is managed by UT-Battelle, LLC, for the US Department of Energy under contract DE-AC05-00OR22725. SB and PJ supported by several grants from the Swiss National Science Foundation (SNSF) and the Commission for Technology and Innovation (CTI) of the Swiss confederation. VH was supported by the Swiss National Science Foundation through Grant FN CR3212-149853/1. VB was supported by the European Union and the State of Hungary, co-financed by the European Social Fund in the framework of TÁMOP 4.2.4. A/1-11-1-2012-0001 'National Excellence Program' (A1-MZPD-12-0166). SO acknowledge FAFU for research support. PB was supported by the University of Torino (60% Project). SH is

supported by the Max Planck Society and the European Research Council Starting Grant (MICRORULES).

**Conflict of interest.** None declared.

## REFERENCES

- Adam E, Groenenboom AE, Kurm V et al. Controlling the microbiome: microhabitat adjustments for successful biocontrol strategies in soil and human gut. *Front Microbiol* 2016;7:1079.
- Agler MT, Ruhe J, Kroll S et al. Microbial hub taxa link host and abiotic factors to plant microbiome variation. *PLoS Biol* 2016;14:e1002352.
- Álvarez-Pérez S, Herrera CM. Composition, richness and nonrandom assembly of culturable bacterial-microfungal communities in floral nectar of Mediterranean plants. *FEMS Microbiol Ecol* 2013;83:685–99.
- Andersson MGI, Berga M, Lindström E et al. The spatial structure of bacterial communities is influenced by historical environmental conditions. *Ecology* 2014;95:1134–40.
- Arendt KR, Hockett KL, Araldi-Brondolo SJ et al. Isolation of endohyphal bacteria from foliar ascomycota and in vitro establishment of their symbiotic associations. *Appl Environ Microb* 2016;82:2943–9.
- Artis D. Epithelial-cell recognition of commensal bacteria and maintenance of immune homeostasis in the gut. *Nat Rev Immunol* 2008;8:411–20.
- Arvanitis M, Mylonakis E. Fungal-bacterial interactions and their relevance in health. *Cell Microbiol* 2015;17:1442–6.
- Aschenbrenner IA, Cernava T, Berg G et al. Understanding microbial multi-species symbioses. *Front Microbiol* 2016;7:180.
- Aylward FO, Suen G, Biedermann PHW et al. Convergent bacterial microbiotas in the fungal agricultural systems of insects. *mBio* 2014;5:e02077.
- Ballhausen MB, de Boer W. The sapro-rhizosphere: carbon flow from saprotrophic fungi into fungus-feeding bacteria. *Soil Biol Biochem* 2016;102:14–7.
- Bánfi R, Pohner Z, Kovács J et al. Characterisation of the large-scale production process of oyster mushroom (*Pleurotus ostreatus*) with the analysis of succession and spatial heterogeneity of lignocellulolytic enzyme activities. *Fungal Biol* 2015;119:1354–63.
- Banitz T, Fetzter I, Johst K et al. Assessing biodegradation benefits from dispersal networks. *Ecol Model* 2011;222:2552–60.
- Barkal LJ, Theberge AB, Guo C-J et al. Microbial metabolomics in open microscale platforms. *Nat Commun* 2016;7:e10610.
- Behrens S, Lösekann T, Pett-Ridge J et al. Linking microbial phylogeny to metabolic activity at the single-cell level by using enhanced element labeling-catalyzed reporter deposition fluorescence in situ hybridization (EL-FISH) and NanoSIMS. *Appl Environ Microb* 2008;74:3143–50.
- Bell TH, El-Din Hassan S, Lauron-Moreau A et al. Linkage between bacterial and fungal rhizosphere communities in hydrocarbon-contaminated soils is related to plant phylogeny. *ISME J* 2014;8:331–43.
- Benoit I, van den Esker MH, Patyshakuliyeva A et al. *Bacillus subtilis* attachment to *Aspergillus niger* hyphae results in mutually altered metabolism. *Environ Microbiol* 2015;17:2099–113.
- Bertaux J, Schmid M, Hutzler P et al. Occurrence and distribution of endobacteria in the plant-associated mycelium of the ectomycorrhizal fungus *Laccaria bicolor* S238N. *Environ Microbiol* 2005;7:1786–95.
- Bertaux J, Schmid M, Prevost-Boure NC et al. In situ identification of intracellular bacteria related to *Paenibacillus* spp. in the mycelium of the ectomycorrhizal fungus *Laccaria bicolor* S238N. *Appl Environ Microb* 2003;69:4243–8.
- Berthold T, Centler F, Hübschmann T et al. Mycelia as a focal point for horizontal gene transfer among soil bacteria. *Sci Rep* 2016;6:e36390.
- Bianciotto V, Genre A, Jargeat P et al. Vertical transmission of endobacteria in the arbuscular mycorrhizal fungus *Gigaspora margarita* through generation of vegetative spores. *Appl Environ Microb* 2004;70:3600–8.
- Bignell E. The molecular basis of pH sensing, signaling, and homeostasis in Fungi. *Advances in Applied Microbiology*. 2012;79:1–18.
- Boersma FG, Warmink JA, Andreote FA et al. Selection of Sphingomonadaceae at the base of *Laccaria proxima* and *Russula exalbicans* fruiting bodies. *Appl Environ Microb* 2009;75:1979–89.
- Bomberg M, Timonen S. Effect of tree species and mycorrhizal colonization on the archaeal population of boreal forest rhizospheres. *Appl Environ Microb* 2009;75:308–15.
- Bonfante P, Desirò A. Who lives in a fungus? The diversity, origins and functions of fungal endobacteria living in *Mucoromycota*. *ISME J* 2017;11:1727–35.
- Bonhomme J, Chauvel M, Goyard S et al. Contribution of the glycolytic flux and hypoxia adaptation to efficient biofilm formation by *Candida albicans*. *Mol Microbiol* 2011;80:995–1013.
- Bordenstein SR, Theis KR. Host biology in light of the microbiome: ten principles of holobionts and hologenomes. *PLoS Biol* 2015;13:e1002226.
- Bosch TCG, McFall-Ngai MJ. Metaorganisms as the new frontier. *Zoology* 2011;114:185–90.
- Boswell GP, Jacobs H, Ritz K et al. The development of fungal networks in complex environments. *Bull Math Biol* 2007;69:605–34.
- Brabcová V, Nováková M, Davidová A et al. Dead fungal mycelium in forest soil represents a decomposition hotspot and a habitat for a specific microbial community. *New Phytol* 2016;210:1369–81.
- Braunsdorf C, Mailänder-Sánchez D, Schaller M. Fungal sensing of host environment. *Cell Microbiol* 2016;18:1188–200.
- Briard B, Heddergott C, Latgé J-P. Volatile compounds emitted by *Pseudomonas aeruginosa* stimulate growth of the fungal pathogen *Aspergillus fumigatus*. *mBio* 2016;7:e00219.
- Cernava T, Aschenbrenner IA, Grube M et al. A novel assay for the detection of bioactive volatiles evaluated by screening of lichen-associated bacteria. *Front Microbiol* 2015;6:e398.
- Chen AI, Dolben EF, Okegbe C et al. *Candida albicans* ethanol stimulates *Pseudomonas aeruginosa* WspR-controlled biofilm formation as part of a cyclic relationship involving phenazines. *PLoS Pathog* 2014;10:e1004480.
- Cordero P, Príncipe A, Jofré E et al. Inhibition of the phytopathogenic fungus *Fusarium proliferatum* by volatile compounds produced by *Pseudomonas*. *Arch Microbiol* 2014;196:803–9.
- Costa R, Van Aarle IM, Mendes R et al. Genomics of pyrrolnitrin biosynthetic loci: Evidence for conservation and whole-operon mobility within Gram-negative bacteria. *Environ Microbiol* 2009;11:159–75.
- Cugini C, Calfee MW, Farrow JM, 3rd et al. Farnesol, a common sesquiterpene, inhibits PQS production in *Pseudomonas aeruginosa*. *Mol Microbiol* 2007;65:896–906.
- de Carvalho MP, Gulotta G, do Amaral MW et al. Coprinus-lactone protects the edible mushroom *Coprinus comatus*

- against biofilm infections by blocking both quorum-sensing and MurA. *Environ Microbiol* 2016;**18**:4254–64.
- de Menezes AB, Prendergast-Miller MT, Richardson AE et al. Network analysis reveals that bacteria and fungi form modules that correlate independently with soil parameters. *Environ Microbiol* 2014;**17**:2677–89.
- de Menezes AB, Richardson AE, Thrall PH. Linking fungal-bacterial co-occurrences to soil ecosystem function. *Curr Opin Microbiol* 2017;**37**:135–41.
- Debbab A, Aly AH, Lin WH et al. Bioactive compounds from marine bacteria and fungi. *Microb Biotechnol* 2010;**3**:544–63.
- Desirò A, Faccio A, Kaech A et al. Endogone, one of the oldest plant-associated fungi, host unique Mollicutes-related endobacteria. *New Phytol* 2015;**205**:1464–72.
- Desirò A, Hao Z, Liber JA et al. Mycoplasma-related endobacteria within Mortierellomycotina fungi: diversity, distribution and functional insights into their lifestyle. *ISME J* 2018;**1**.
- Desirò A, Salvioli A, Ngonkeu EL et al. Detection of a novel intracellular microbiome hosted in arbuscular mycorrhizal fungi. *ISME J* 2014;**8**:257–70.
- Deveau A, Antony-Babu S, Le Tacon F et al. Temporal changes of bacterial communities in the Tuber melanosporum ectomycorrhizosphere during ascocarp development. *Mycorrhiza* 2016;**26**:389–99.
- Deveau A, Barret M, Diedhiou AG et al. Pairwise transcriptomic analysis of the interactions between the ectomycorrhizal fungus *Laccaria bicolor* S238N and three beneficial, neutral and antagonistic soil bacteria. *Microb Ecol* 2014;**69**:145–59.
- Dhamgaye S, Qu Y, Peleg AY. Polymicrobial infections involving clinically relevant Gram-negative bacteria and fungi. *Cell Microbiol* 2016;**18**:1716–22.
- Dixon EF, Hall RA. Noisy neighbourhoods: quorum sensing in fungal-polymicrobial infections. *Cell Microbiol* 2015;**17**:1431–41.
- Douterelo I, Jackson M, Solomon C et al. Microbial analysis of in situ biofilm formation in drinking water distribution systems: implications for monitoring and control of drinking water quality. *Appl Microbiol Biot* 2016;**100**:3301–11.
- Dugat-Bony E, Straub C, Teissandier A et al. Overview of a surface-ripened cheese community functioning by meta-omics analyses. *PLoS One* 2015;**10**:e0124360.
- Dyrka W, Lamacchia M, Durrens P et al. Diversity and variability of NOD-Like receptors in Fungi. *Genome Biol Evol* 2014;**6**:3137–58.
- Effmert U, Kalderás J, Warnke R et al. Volatile mediated interactions between bacteria and fungi in the soil. *J Chem Ecol* 2012;**38**:665–703.
- El-Jurdi N, Ghannoum MA. The mycobiome: impact on health and disease states. *Microbiol Spectr* 2017;**5**. DOI: 10.1128/microbiolspec.FUNK-0045-2016.
- Essig A, Hofmann D, Münch D et al. Copsin, a novel peptide-based fungal antibiotic interfering with the peptidoglycan synthesis. *J Biol Chem* 2014;**289**:34953–64.
- Expósito RG, De Bruijn I, Postma J et al. Current insights into the role of rhizosphere bacteria in disease suppressive soils. *Front Microbiol*, 2017. DOI: 10.3389/fmicb.2017.02529.
- Falconer RE, Houston AN, Otten W et al. Emergent behavior of soil fungal dynamics. *Soil Sci* 2012;**17**:111–9.
- Ferreira JAG, Penner JC, Moss RB et al. Inhibition of *Aspergillus fumigatus* and its biofilm by *Pseudomonas aeruginosa* is dependent on the source, phenotype and growth conditions of the bacterium. *PLoS One* 2015;**10**:e0134692.
- Fester T, Giebler J, Wick LY et al. Plant-microbe interactions as drivers of ecosystem functions relevant for the biodegradation of organic contaminants. *Curr Opin Biotechnol* 2014;**27**:168–75.
- Förster TM, Mogavero S, Dräger A et al. Enemies and brothers in arms: *Candida albicans* and gram-positive bacteria. *Cell Microbiol* 2016;**18**:1709–15.
- Fourie R, Ells R, Swart CW et al. *Candida albicans* and *Pseudomonas aeruginosa* interaction, with focus on the role of eicosanoids. *Front Physiol* 2016;**7**:64.
- Fox EP, Cowley ES, Nobile CJ et al. Anaerobic bacteria grow within *Candida albicans* biofilms and induce biofilm formation in suspension cultures. *Curr Biol* 2014;**24**:2411–6.
- Fraune S, Anton-Erxleben F, Augustin R et al. Bacteria-bacteria interactions within the microbiota of the ancestral metazoan Hydra contribute to fungal resistance. *ISME J* 2015;**9**:1543–56.
- Frey-Klett P, Burlinson P, Deveau A et al. Bacterial-Fungal interactions: hyphens between agricultural, clinical, environmental, and food microbiologists. *Microbiol Mol Biol Rev* 2011;**75**:583–609.
- Fricker MD, Heaton LLM, Jones NS et al. The mycelium as a network. *Microbiol Spectr* 2017;**5**. DOI: doi:10.1128/microbiolspec.FUNK-0033-2017.
- Furuno S, Pazolt K, Rabe C et al. Fungal mycelia allow chemotactic dispersal of polycyclic aromatic hydrocarbon-degrading bacteria in water-unsaturated systems. *Env Microbiol* 2010;**12**:1391–8.
- Furuno S, Foss S, Wild E et al. Mycelia promote active transport and spatial dispersion of polycyclic aromatic hydrocarbons. *Environ Sci Technol* 2012;**46**:5463–70.
- Gänzle MG. Lactic metabolism revisited: metabolism of lactic acid bacteria in food fermentations and food spoilage. *Curr Opin Food Sci* 2015;**2**:106–17.
- Garbeva P, Hordijk C, Gerards S et al. Volatiles produced by the mycophagous soil bacterium *Collimonas*. *FEMS Microbiol Ecol* 2014;**87**:639–49.
- Ghignone S, Salvioli A, Anca I et al. The genome of the obligate endobacterium of an AM fungus reveals an inter-phylum network of nutritional interactions. *ISME J* 2012;**6**:136–45.
- Ghodsavali B, Svenningsen NB, Hao X et al. A novel baiting microcosm approach used to identify the bacterial community associated with *Penicillium bilaii* hyphae in soil. *PLoS One* 2017;**12**. DOI: 10.1371/journal.pone.0187116.
- Gkarmiri K, Finlay RD, Alström S et al. Transcriptomic changes in the plant pathogenic fungus *Rhizoctonia solani* AG-3 in response to the antagonistic bacteria *Serratia proteamaculans* and *Serratia plymuthica*. *BMC Genomics* 2015;**16**:630.
- Glaeser SP, Imani J, Alabid I et al. Non-pathogenic *Rhizobium radiobacter* F4 deploys plant beneficial activity independent of its host *Piriformospora indica*. *ISME J* 2016;**10**:871–84.
- Grube M, Cernava T, Soh J et al. Exploring functional contexts of symbiotic sustain within lichen-associated bacteria by comparative omics. *ISME J* 2015;**9**:412–24.
- Guhr A, Borken W, Spohn M et al. Redistribution of soil water by a saprotrophic fungus enhances carbon mineralization. *P Natl Acad Sci USA* 2015;**112**:14647–51.
- Guo H, Glaeser SP, Alabid I et al. The abundance of endofungal bacterium *Rhizobium radiobacter* (syn. *Agrobacterium tumefaciens*) increases in its fungal host *Piriformospora indica* during the tripartite sebacinal symbiosis with higher plants. *Front Microbiol* 2017;**8**:1–13.
- Hacquard S. Commentary: microbial small talk: volatiles in fungal-bacterial interactions. *Front Microbiol* 2017b;**6**. DOI: 10.3389/fmicb.2015.01495.

- Hacquard S, Garrido-Oter R, González A et al. Microbiota and host nutrition across plant and animal kingdoms. *Cell Host Microbe* 2015;17:603–16.
- Hacquard S, Spaepen S, Garrido-Oter R et al. Interplay between innate immunity and the plant microbiota. *Annu Rev Phytopathol* 2017;55:565–89.
- Haq IU. Interactions of *Burkholderia terrae* with soil fungi: mechanisms, gene expression patterns and ecological behavior of *Burkholderia terrae* BS001 during its interaction with *Lyophyllum* sp. strain Karsten and *Trichoderma asperellum* 302 in soil. Ph.D. Thesis. University of Groningen 2016; 199.
- Haq IU, Calixto da ROR, Yang P et al. Chemotaxis and adherence to fungal surfaces are key components of the behavioral response of *Burkholderia terrae* BS001 to two selected soil fungi. *FEMS Microbiol Ecol* 2016. DOI:10.1093/femsec/fiw164;92.
- Haq IU, Dini-Andreote F, van Elsas JD. Transcriptional responses of the bacterium *Burkholderia terrae* BS001 to the fungal host *Lyophyllum* sp. strain Karsten under soil-mimicking conditions. *Microb Ecol* 2017;73:236–52.
- Haq IU, Zhang M, Yang P et al. The interactions of bacteria with fungi in soil: emerging concepts. *Adv Appl Microbiol* 2014;89:185–215.
- Harms H, Schlosser D, Wick LY. Untapped potential: exploiting fungi in bioremediation of hazardous chemicals. *Nat Rev Microbiol* 2011;9:177–92.
- Hennessy RC, Phippen CBW, Nielsen KF et al. Biosynthesis of the antimicrobial cyclic lipopeptides nunamycin and nunapeptin by *Pseudomonas fluorescens* strain In5 is regulated by the LuxR-type transcriptional regulator NunF. *Microbiologyopen* 2017;e516:1–14.
- Hennessy RC, Stougaard P, Olsson S. A Microplate reader-based system for visualizing transcriptional activity during in vivo microbial interactions in space and time. *Sci Rep* 2017;7: 281.
- Hervé V, Le Roux X, Uroz S et al. Diversity and structure of bacterial communities associated with *Phanerochaete chrysosporium* during wood decay. *Environ Microbiol* 2014;16:2238–52.
- Hoffman MT, Arnold AE. Diverse bacteria inhabit living hyphae of phylogenetically diverse fungal endophytes. *Appl Environ Microb* 2010;76:4063–75.
- Hoffman MT, Gunatilaka MK, Wijeratne K et al. Endohyphal bacterium enhances production of indole-3-acetic acid by a foliar fungal endophyte. *PLoS One* 2013;8:e73132.
- Hoffmann C, Dollive S, Grunberg S et al. Archaea and Fungi of the human gut microbiome: correlations with diet and bacterial residents. *PLoS One* 2013;8. DOI: 10.1371/journal.pone.0066019.
- Hover T, Maya T, Ron S et al. Mechanisms of bacterial (*Serratia marcescens*) attachment to, migration along, and killing of fungal hyphae. *Appl Environ Microb* 2016;82:2585–94.
- Hoyt JR, Cheng TL, Langwig KE et al. Bacteria isolated from bats inhibit the growth of *Pseudogymnoascus destructans*, the causative agent of white-nose syndrome. *PLoS One* 2015;10:e0121329.
- Huttenhower C, Gevers D, Knight R et al. Structure, function and diversity of the healthy human microbiome. *Nature* 2012;486:207–14.
- Ipcho S, Sundelin T, Erbs G et al. Fungal innate immunity induced by bacterial microbe-associated molecular patterns (MAMPs). *G3* 2016;6:1585–95.
- Janus MM, Crielgaard W, Volgenant CMC et al. *Candida albicans* alters the bacterial microbiome of early in vitro oral biofilms. *J Oral Microbiol* 2017;9:1270613.
- Janus MM, Willems HME, Krom BP. *Candida albicans* in multi-species oral communities; a keystone commensal? *Adv Exp Med Biol* 2016;931:13–20.
- Joergensen RG, Wichern F. Alive and kicking: Why dormant soil microorganisms matter. *Soil Biol Biochem* 2018;116:419–30.
- Jones SE, Ho L, Rees CA et al. *Streptomyces* exploration is triggered by fungal interactions and volatile signals. *Elife* 2017;6:4268–73.
- Jung B, Park J, Kim N et al. Cooperative interactions between seed-borne bacterial and air-borne fungal pathogens on rice. *Nat Commun* 2018;9:31.
- Kai K, Furuyabu K, Tani A et al. Production of the quorum-sensing molecules N-acylhomoserine lactones by endobacteria associated with *Mortierella alpina* A-178. *ChemBioChem* 2012;13:1776–84.
- Kalan L, Loesche M, Hodgkinson BP et al. Redefining the chronic-wound microbiome: fungal communities are prevalent, dynamic, and associated with delayed healing. *mBio* 2016;7:1–12.
- Kaltenpoth M, Strupat K, Svatoš A. Linking metabolite production to taxonomic identity in environmental samples by (MA)LDI-FISH. *ISME J* 2016;10:527–31.
- Kastman EK, Kamelamel N, Norville JW et al. Biotic interactions shape the ecological distributions of *Staphylococcus* species. *mBio* 2016;7:e01157–16.
- Kohlmeier S, Smits THM, Ford RM et al. Taking the fungal highway: mobilization of pollutant-degrading bacteria by fungi. *Environ Sci Technol* 2005;15:4640–6.
- Kong K-F, Schnepel L, Mathee K. Beta-lactam antibiotics: from antibiosis to resistance and bacteriology. *APMIS* 2010;118:1–36.
- Koskella B, Hall L, Metcalf C. The microbiome beyond the horizon of ecological and evolutionary theory. *Nat Ecol Evol* 2017;1:1606–15.
- Krom BP, Oskam J. Microbial biofilms and wound healing: an ecological hypothesis. *Phlebology* 2014;29:168–73.
- Kumar S, Indugu N, Vecchiarelli B et al. Associative patterns among anaerobic fungi, methanogenic archaea, and bacterial communities in response to changes in diet and age in the rumen of dairy cows. *Front Microbiol* 2015;6:1–10.
- Künzler M. Hitting the sweet spot: glycans as targets of fungal defense effector proteins. *Molecules* 2015;20:8144–67.
- Kurth F, Feldhahn L, Bönn M et al. Large scale transcriptome analysis reveals interplay between development of forest trees and a beneficial mycorrhiza helper bacterium. *BMC Genomics* 2015;16:658.
- Lackner G, Hertweck C. Impact of endofungal bacteria on infection biology, food safety, and drug development. *PLoS Pathog* 2011;7:e1002096.
- Lackner G, Moebius N, Hertweck C. Endofungal bacterium controls its host by an hrp type III secretion system. *ISME J* 2011;5:252–61.
- Lamacchia M, Dyrka W, Breton A et al. Overlapping *Podospora anserina* transcriptional responses to bacterial and fungal non self indicate a multilayered innate immune response. *Front Microbiol* 2016;7. DOI: 10.3389/fmicb.2016.00471.
- Lambooij JM, Hoogenkamp MA, Brandt BW et al. Fungal mitochondrial oxygen consumption induces the growth of strict anaerobic bacteria. *Fungal Genet Biol* 2017;109:1–6.
- Lamichhane JR, Venturi V. Synergisms between microbial pathogens in plant disease complexes: a growing trend. *Front Plant Sci* 2015;6. DOI: 10.3389/fpls.2015.00385.
- Lastovetsky OA, Gaspar ML, Mondo SJ et al. Lipid metabolic changes in an early divergent fungus govern the

- establishment of a mutualistic symbiosis with endobacteria. *P Natl Acad Sci USA* 2016;**113**:15102–7.
- Leclair LW, Hogan DA. Mixed bacterial-fungal infections in the CF respiratory tract. *Med Mycol* 2010;**48**:S125–32.
- Lee K, Kobayashi N, Watanabe M et al. Spread and change in stress resistance of Shiga toxin-producing *Escherichia coli* O157 on fungal colonies. *Microb Biotechnol* 2014;**7**:621–9.
- Li Z, Yao Q, Dearth SP et al. Integrated proteomics and metabolomics suggests symbiotic metabolism and multimodal regulation in a fungal-endobacterial system. *Environ Microbiol* 2017;**19**:1041–53.
- Lindsay AK, Hogan DA. *Candida albicans*: molecular interactions with *Pseudomonas aeruginosa* and *Staphylococcus aureus*. *Fungal Biol Rev* 2014;**28**:85–96.
- Liu Y, Rousseaux S, Tourdot-Maréchal R et al. Wine microbiome, a dynamic world of microbial interactions. *Crit Rev Food Sci Nutr* 2017;**57**:856–73.
- Lladó S, López-mondéjar R, Baldrian P. Forest soil bacteria: diversity, involvement in ecosystem processes, and response to global change. *Microbiol Mol Biol Rev* 2017;**81**:1–27.
- Longo A V, Zamudio KR. Environmental fluctuations and host skin bacteria shift survival advantage between frogs and their fungal pathogen. *ISME J* 2017;**11**:349–61.
- Lopes SP, Azevedo NF, Pereira MO. Microbiome in cystic fibrosis: shaping polymicrobial interactions for advances in antibiotic therapy. *Crit Rev Microbiol* 2014;**78**:281–13.
- Ma B, Wang H, Dsouza M et al. Geographic patterns of co-occurrence network topological features for soil microbiota at continental scale in eastern China. *ISME J* 2016;**10**:1891–901.
- Martin G, Guggiari M, Bravo D et al. Fungi, bacteria and soil pH: the oxalate-carbonate pathway as a model for metabolic interaction. *Environ Microbiol* 2012;**14**:2960–70.
- Mela F, Fritsche K, de Boer W et al. Dual transcriptional profiling of a bacterial/fungal confrontation: *Collimonas fungivorans* versus *Aspergillus niger*. *ISME J* 2011;**5**:1494–504.
- Mela F, Fritsche K, de Boer W et al. Comparative genomics of bacteria from the genus *Collimonas*: linking (dis)similarities in gene content to phenotypic variation and conservation. *Environ Microbiol Rep* 2012;**4**:424–32.
- Michielse CB, Hooykaas PJJ, van den Hondel CA et al. *Agrobacterium*-mediated transformation as a tool for functional genomics in fungi. *Curr Genet* 2005;**48**:1–17.
- Moebius N, Üzümlü Z, Dijksterhuis J et al. Active invasion of bacteria into living fungal cells. *Elife* 2014;**3**. DOI: 10.7554/eLife.03007.
- Moissl-Eichinger C, Pausen M, Taffner J et al. Archaea are interactive components of complex microbiomes. *Trends Microbiol* 2017. DOI: 10.1016/j.tim.2017.07.004.
- Mondo SJ, Lastovetsky OA, Gaspar ML et al. Bacterial endosymbionts influence host sexuality and reveal reproductive genes of early divergent fungi. *Nat Commun* 2017;**8**: 1843.
- Mondo SJ, Toomer KH, Morton JB et al. Evolutionary stability in a 400-million-year-old heritable facultative mutualism. *Evolution* 2012;**66**:2564–76.
- Morales DK, Grahl N, Okegbe C et al. Control of *Candida albicans* metabolism and biofilm formation by *Pseudomonas aeruginosa* phenazines. *mBio* 2013;**4**:e00526–12.
- Morales DK, Jacobs NJ, Rajamani S et al. Antifungal mechanisms by which a novel *Pseudomonas aeruginosa* phenazine toxin kills *Candida albicans* in biofilms. *Mol Microbiol* 2010;**78**:1379–92.
- Moree WJ, Yang JY, Zhao X et al. Imaging mass spectrometry of a coral microbe interaction with fungi. *J Chem Ecol* 2013;**39**:1045–54.
- Morris BEL, Henneberger R, Huber H et al. Microbial syntrophy: interaction for the common good. *FEMS Microbiol Rev* 2013;**37**:384–406.
- Mosse B. Honey-coloured, sessile *Endogone* spores: II. Changes in fine structure during spore development. *Archiv Mikrobiol* 1970;**74**:129–45.
- Mousa WK, Shearer C, Limay-Rios V et al. Root-hair endophyte stacking in finger millet creates a physicochemical barrier to trap the fungal pathogen *Fusarium graminearum*. *Nat Microbiol* 2016;**1**:16167.
- Mukherjee PK, Chandra J, Retuerto M et al. Oral mycobiome analysis of HIV-infected patients: identification of *Pichia* as an antagonist of opportunistic fungi. *PLoS Pathog* 2014;**10**. DOI: 10.1371/journal.ppat.1003996.
- Murat C. Forty years of inoculating seedlings with truffle fungi: past and future perspectives. *Mycorrhiza* 2015;**25**: 77–81.
- Naito M, Desirò A, Gonzalez JB et al. “*Candidatus Moenioplasma glomeromycetorum*”, an endobacterium of arbuscular mycorrhizal fungi. *Int J Syst Evol Microbiol* 2017. DOI: 10.1099/ijsem.0.001785.
- Naito M, Morton JB, Pawlowska TE. Minimal genomes of mycoplasma-related endobacteria are plastic and contain host-derived genes for sustained life within Glomeromycota. *P Natl Acad Sci USA* 2015;**112**:7791–6.
- Naumann M, Schüssler A, Bonfante P. The obligate endobacteria of arbuscular mycorrhizal fungi are ancient heritable components related to the Mollicutes. *ISME J* 2010;**4**: 862–71.
- Navarri M, Jégou C, Meslet-Cladière L et al. Deep Subseafloor Fungi as an untapped reservoir of amphipathic antimicrobial compounds. *Mar Drugs* 2016;**14**:50.
- Nazir R, Boersma FGH, Warmink JA et al. *Lyophyllum* sp. strain Karsten alleviates pH pressure in acid soil and enhances the survival of *Variovorax paradoxus* HB44 and other bacteria in the mycosphere. *Soil Biol Biochem* 2010;**42**:2146–52.
- Nazir R, Shen J-P, Wang J-T et al. Fungal networks serve as novel ecological routes for enrichment and dissemination of antibiotic resistance genes as exhibited by microcosm experiments. *Sci Rep* 2017;**7**:15457.
- Nuccio EE, Hodge A, Pett-Ridge J et al. An arbuscular mycorrhizal fungus significantly modifies the soil bacterial community and nitrogen cycling during litter decomposition. *Environ Microbiol* 2013;**15**:1870–81.
- Olsson S, Bonfante P, Pawlowska TE. Ecology and evolution of Fungal-Bacterial interactions. In: Dighton J, White JF (eds). *The Fungal Community: Its Organization and Role in the Ecosystem*, 4th edn. Boca Raton: CRC Press, 2017.
- Otto S, Bruni EP, Harms H et al. Catch me if you can: dispersal and foraging of *Bdellovibrio bacteriovorus* 109J along mycelia. *ISME J* 2017 **11**:386–93.
- Otto S, Harms H, Wick LY. Effects of predation and dispersal on bacterial abundance and contaminant biodegradation. *FEMS Microbiol Ecol* 2017. DOI: 10.1093/femsec/fiw241:93.
- Panke-Buisse K, Poole AC, Goodrich JK et al. Selection on soil microbiomes reveals reproducible impacts on plant function. *ISME J* 2015;**9**:980–9.
- Paoletti M, Saupe SJ. Fungal incompatibility: evolutionary origin in pathogen defense? *Bioessays* 2009;**31**:1201–10.
- Park HB, Kwon HC, Lee C et al. Glionitrin A, an antibiotic? antitumor metabolite derived from competitive interaction between abandoned mine microbes. *J Nat Prod* 2009;**72**:248–52.
- Partida-Martínez LP. The fungal holobiont: evidence from early diverging fungi. *Environ Microbiol* 2017;**19**:2919–23.

- Partida-Martinez LP, Monajembashi S, Greulich K-O et al. Endosymbiont-dependent host reproduction maintains bacterial-fungal mutualism. *Curr Biol* 2007;17:773–7.
- Peleg AY, Hogan DA, Mylonakis E. Medically important bacterial-fungal interactions. *Nat Rev Microbiol* 2010;8:340–9.
- Perazzolli M, Herrero N, Sterck L et al. Transcriptomic responses of a simplified soil microcosm to a plant pathogen and its biocontrol agent reveal a complex reaction to harsh habitat. *BMC Genomics* 2016;17:838.
- Peters BM, Noverr MC. *Candida albicans*-*Staphylococcus aureus* polymicrobial peritonitis modulates host innate immunity. *Infect Immun* 2013;81:2178–89.
- Phelan VV, Moree WJ, Aguilar J et al. Impact of a transposon insertion in *phzF2* on the specialized metabolite production and interkingdom interactions of *Pseudomonas aeruginosa*. *J Bacteriol* 2014;196:1683–93.
- Pinto C, Pinho D, Sousa S et al. Unravelling the diversity of grapevine microbiome. *PLoS One* 2014;9:e85622.
- Pion M, Spangenberg JE, Simon A et al. Bacterial farming by the fungus *Morchella crassipes*. *P Roy Soc B-Biol Sci* 2013;280:20132242.
- Poudel R, Jumpponen A, Schlatter DC et al. Microbiome networks: a systems framework for identifying candidate microbial assemblages for disease management. *Phytopathology* 2016;106:1083–96.
- Poulsen M, Hu H, Li C et al. Complementary symbiont contributions to plant decomposition in a fungus-farming termite. *P Natl Acad Sci USA* 2014;111:14500–5.
- Pukkila-Worley R, Ausubel FM, Mylonakis E. *Candida albicans* infection of *Caenorhabditis elegans* induces antifungal immune defenses. *PLoS Pathog* 2011;7. DOI: 10.1371/journal.ppat.1002074.
- Purahong W, Wubet T, Lentendu G et al. Life in leaf litter: novel insights into community dynamics of bacteria and fungi during litter decomposition. *Mol Ecol* 2016;25:4059–74.
- Raaijmakers JM, De Bruijn I, Nybroe O et al. Natural functions of lipopeptides from *Bacillus* and *Pseudomonas*: more than surfactants and antibiotics. *FEMS Microbiol Rev* 2010;34:1037–62.
- Raymann K, Moeller AH, Goodman AL et al. Unexplored archaeal diversity in the great ape gut microbiome. *mSphere* 2017;2:e00026–17.
- Reen FJ, Phelan JP, Woods DF et al. Harnessing bacterial signals for suppression of biofilm formation in the nosocomial fungal pathogen *Aspergillus fumigatus*. *Front Microbiol* 2016;7:2074.
- Rinta-Kanto JM, Sinkko H, Rajala T et al. Natural decay process affects the abundance and community structure of bacteria and archaea in *Picea abies* logs. *FEMS Microbiol Ecol* 2016; DOI: 10.1093/femsec/fiw087:92.
- Ritpitakphong U, Falquet L, Vimoltust A et al. The microbiome of the leaf surface of *Arabidopsis* protects against a fungal pathogen. *New Phytol* 2016;210:1033–43.
- Ritz K, Young IM. Interactions between soil structure and fungi. *Mycologist* 2004;18:52–9.
- Rohm B, Scherlach K, Möbius N et al. Toxin production by bacterial endosymbionts of a *Rhizopus microsporus* strain used for tempe/sufu processing. *Int J Food Microbiol* 2010;136:368–71.
- Ross C, Opel V, Scherlach K et al. Biosynthesis of antifungal and antibacterial polyketides by *Burkholderia gladioli* in coculture with *Rhizopus microsporus*. *Mycoses* 2014;57:48–55.
- Rudnick MB, van Veen JA, de Boer W. Oxalic acid: a signal molecule for fungus-feeding bacteria of the genus *Colimonas*? *Environ Microbiol Rep* 2015;7:709–14.
- Ruiz-Herrera J, León-Ramírez C, Vera-Núñez A et al. A novel intracellular nitrogen-fixing symbiosis made by *Ustilago maydis* and *Bacillus* spp. *New Phytol* 2015;207:769–77.
- Salvioli A, Ghignone S, Novero M et al. Symbiosis with an endobacterium increases the fitness of a mycorrhizal fungus, raising its bioenergetic potential. *ISME J* 2016;10:130–44.
- Salvioli Di Fossalunga A, Lipuma J, Venice F et al. The endobacterium of an arbuscular mycorrhizal fungus modulates the expression of its toxin-antitoxin systems during the life cycle of its host. *ISME J* 2017;11:2394–8.
- Santhanam R, Luu VT, Weinhold A et al. Native root-associated bacteria rescue a plant from a sudden-wilt disease that emerged during continuous cropping. *P Natl Acad Sci USA* 2015;112:E5013–20.
- Schamfuß S, Neu TR, Roelof Van Der Meer J et al. Impact of mycelia on the accessibility of fluorene to PAH-degrading bacteria. *Environ Sci Technol* 2013;47:6908–15.
- Scherlach K, Graupner K, Hertweck C. Molecular bacteria-fungi interactions: effects on environment, food, and medicine. *Annu Rev Microbiol* 2013;67:375–97.
- Scherlach K, Hertweck C. Mediators of mutualistic microbe-microbe interactions. *Nat Prod Rep* 2017, DOI: 10.1039/C7NP00035A.
- Schlecht LM, Peters BM, Krom BP et al. Systemic *Staphylococcus aureus* infection mediated by *Candida albicans* hyphal invasion of mucosal tissue. *Microbiology* 2015;161:168–81.
- Schmidt R, de Jager V, Zühlke D et al. Fungal volatile compounds induce production of the secondary metabolite Sodorifen in *Serratia plymuthica* PRI-2C. *Sci Rep* 2017;7:862.
- Schmidt R, Etalo DW, de Jager V et al. Microbial small talk: volatiles in fungal-bacterial interactions. *Front Microbiol* 2015;6:1495.
- Schulz-Bohm K, Tyc O, de Boer W et al. Fungus-associated bacteriome in charge of their host behavior. *Fungal Genet Biol* 2016. DOI: 10.1016/j.fgb.2016.07.011.
- Scopel M, Abraham W-R, Henriques AT et al. Dipeptide cyclo(Leucyl-Tyrosyl) produced by sponge associated *Penicillium* sp. F37 inhibits biofilm formation of the pathogenic *Staphylococcus epidermidis*. *Bioorg Med Chem Lett* 2013;23:624–6.
- Shaffer JP, Sarmiento C, Zalamea P-C et al. Diversity, Specificity, and phylogenetic relationships of endohyphal bacteria in fungi that inhabit tropical eeds and leaves. *Front Ecol Evol* 2016;4:116.
- Shaffer JP, U'Ren JM, Gallery RE et al. An endohyphal bacterium (*Chitinophaga*, *Bacteroidetes*) alters carbon source use by *Fusarium keratoplasticum* (*F. solani* species complex, *Nectriaceae*). *Front Microbiol* 2017;8. DOI: 10.3389/fmicb.2017.00350.
- Simon A, Hervé V, Al-Dourobi A et al. An in situ inventory of fungi and their associated migrating bacteria in forest soils using fungal highway columns. *FEMS Microbiol Ecol* 2017. DOI: 10.1093/femsec/fiw217:93.
- Spraker JE, Jewell K, Roze L V et al. A volatile relationship: profiling an inter-kingdom dialogue between two plant pathogens, *Ralstonia solanacearum* and *Aspergillus flavus*. *J Chem Ecol* 2014;40:502–13.
- Spraker JE, Sanchez LM, Lowe TM et al. *Ralstonia solanacearum* lipopeptide induces chlamydospore development in fungi and facilitates bacterial entry into fungal tissues. *ISME J* 2016;10:2317–30.
- Stanley CE, Stöckli M, van Swaay D et al. Probing bacterial-fungal interactions at the single cell level. *Integr Biol* 2014;6:935–45.

- Stopnisek N, Zühlke D, Carlier A et al. Molecular mechanisms underlying the close association between soil *Burkholderia* and fungi. *ISME J* 2016;**10**:253–64.
- Sztajer H, Szafranski SP, Tomasch J et al. Cross-feeding and interkingdom communication in dual-species biofilms of *Streptococcus mutans* and *Candida albicans*. *ISME J* 2014;**8**:2256–71.
- Tauber JP, Schroeckh V, Shelest E et al. Bacteria induce pigment formation in the basidiomycete *Serpula lacrymans*. *Environ Microbiol* 2016;**18**:5218–27.
- Tecon R, Or D. Bacterial flagellar motility on hydrated rough surfaces controlled by aqueous film thickness and connectedness. *Sci Rep* 2016;**6**:1–11.
- Tecon R, Or D. Biophysical processes supporting the diversity of microbial life in soil. *FEMS Microbiol Rev* 2017;**41**:599–623.
- Thion CE, Poirel JD, Cornulier T et al. Plant nitrogen-use strategy as a driver of rhizosphere archaeal and bacterial ammonia oxidiser abundance. *FEMS Microbiol Ecol* 2016;**92**:1–11.
- Thompson LR, Sanders JG, McDonald D et al. A communal catalogue reveals Earth's multiscale microbial diversity. *Nature* 2017;**551**:457–63.
- Tolonen AC, Cerisy T, El-Sayyed H et al. Fungal lysis by a soil bacterium fermenting cellulose. *Environ Microbiol* 2015;**17**:2618–27.
- Tomada S, Sonogo P, Moretto M et al. Dual RNA-Seq of *Lyso bacter capsici* AZ78 – *Phytophthora infestans* interaction shows the implementation of attack strategies by the bacterium and unsuccessful oomycete defense responses. *Environ Microbiol* 2017;**19**:4113–25.
- Toomer KH, Chen X, Naito M et al. Molecular evolution patterns reveal life history features of mycoplasma-related endobacteria associated with arbuscular mycorrhizal fungi. *Mol Ecol* 2015;**24**:3485–500.
- Torres-Cortés G, Ghignone S, Bonfante P et al. Mosaic genome of endobacteria in arbuscular mycorrhizal fungi: Transkingdom gene transfer in an ancient mycoplasma-fungus association. *P Natl Acad Sci USA* 2015;**112**:7785–90.
- Trejo-Hernández A, Andrade-Domínguez A, Hernández M et al. Interspecies competition triggers virulence and mutability in *Candida albicans*-*Pseudomonas aeruginosa* mixed biofilms. *ISME J* 2014;**8**:1974–88.
- Trosvik P, de Muinck EJ. Ecology of bacteria in the human gastrointestinal tract—identification of keystone and foundation taxa. *Microbiome* 2015;**3**:44.
- Tyc O, van den Berg M, Gerards S et al. Impact of interspecific interactions on antimicrobial activity among soil bacteria. *Front Microbiol* 2014;**5**. DOI: 10.3389/fmicb.2014.00567.
- Uehling J, Deveau A, Paoletti M. Innate immune response and NLR based non-self recognition in Fungi, Plants and Animals. *Pearls PLoS Pathog* 2017. DOI:10.1371/journal.ppat.1006578
- Uehling J, Gryganskyi A, Hameed K et al. Comparative genomics of *Mortierella elongata* and its bacterial endosymbiont *Mycoavidus cysteinexigens*. *Environ Microbiol* 2017. DOI: 10.1111/1462-2920.13669.
- Urich T, Lanzén A, Stokke R et al. Microbial community structure and functioning in marine sediments associated with diffuse hydrothermal venting assessed by integrated metabolomics. *Environ Microbiol* 2014;**16**:2699–710.
- Uroz S, Buée M, Deveau A et al. Ecology of the forest microbiome: highlights of temperate and boreal ecosystems. *Soil Biol Biochem* 2016;**103**:471–88.
- Vahdatzadeh M, Deveau A, Splivallo R. The role of the microbiome of truffles in aroma formation: a meta-analysis approach. *Appl Environ Microb* 2015;**81**:6946–52.
- Vannini C, Carpentieri A, Salvioli A et al. An interdomain network: the endobacterium of a mycorrhizal fungus promotes antioxidative responses in both fungal and plant hosts. *New Phytol* 2016;**211**:265–75.
- Vylkova S. Environmental pH modulation by pathogenic fungi as a strategy to conquer the host. *PLoS Pathog* 2017;**13**:1–6.
- Vylkova S, Lorenz MC. Modulation of phagosomal pH by *Candida albicans* promotes hyphal morphogenesis and requires Stp2p, a regulator of amino acid transport. *PLoS Pathog* 2014;**10**. DOI: 10.1371/journal.ppat.1003995.
- Walker TS, Bais HP, Déziel E et al. *Pseudomonas aeruginosa*-plant root interactions. Pathogenicity, biofilm formation, and root exudation. *Plant Physiol* 2004;**134**:320–31.
- Wang X, Li G-H, Zou C-G et al. Bacteria can mobilize nematode-trapping fungi to kill nematodes. *Nat Commun* 2014;**5**:5776.
- Wang Y, Huang WE, Cui L et al. Single cell stable isotope probing in microbiology using Raman microspectroscopy. *Curr Opin Biotechnol* 2016;**41**:34–42.
- Warmink JA, Nazir R, van Elsas JD. Universal and species-specific bacterial “fungiphiles” in the mycospheres of different basidiomycetous fungi. *Environ Microbiol* 2009;**11**:300–12.
- Warmink JA, van Elsas JD. Migratory response of soil bacteria to *Lyophyllum* sp. strain Karsten in soil microcosms. *Appl Environ Microb* 2009;**75**:2820–30.
- Wei Z, Yang T, Friman VP et al. Trophic network architecture of root-associated bacterial communities determines pathogen invasion and plant health. *Nat Commun* 2015;**6**:1–9.
- Willems HM, Kos K, Jabra-Rizk MA et al. *Candida albicans* in oral biofilms could prevent caries. *Pathog Dis* 2016;**74**:1–6.
- Wohlschlager T, Butschli A, Grassi P et al. Methylated glycans as conserved targets of animal and fungal innate defense. *P Natl Acad Sci USA* 2014;**111**:2787–96.
- Wolfe BE, Dutton RJ. Fermented foods as experimentally tractable microbial ecosystems. *Cell* 2015;**161**:49–55.
- Wongsuk T, Pumeesat P, Luplertlop N. Fungal quorum sensing molecules: role in fungal morphogenesis and pathogenicity. *J Basic Microbiol* 2016;**56**:440–7.
- Worrich A, König S, Miltner A et al. Mycelium-like networks increase bacterial dispersal, growth, and biodegradation in a model ecosystem at various water potentials. *Appl Environ Microb* 2016;**82**:2902–8.
- Worrich A, Stryhanyuk H, Musat N et al. Mycelium-mediated transfer of water and nutrients stimulates bacterial activity in dry and oligotrophic environments. *Nat Commun* 2017. DOI:10.1038/ncomms15472.
- Young IM, Crawford JW. Interactions and self-organization in the soil-microbe complex. *Science* 2004;**304**:1634–7.
- Zagladskaja YA, Lysak L V., Sidorova II et al. Bacterial complexes of the fruiting bodies and hyphosphere of certain basidiomycetes. *Biol Bull* 2013;**40**:358–64.
- Zamioudis C, Pieterse CMJ. Modulation of host immunity by beneficial microbes. *Mol Plant Microbe In* 2012;**25**:139–50.
- Zhang M, de Pereira Silva MC, De Mares Maryam C et al. The mycosphere constitutes an arena for horizontal gene transfer with strong evolutionary implications for bacterial-fungal interactions. 2014. DOI: 10.1111/1574-6941.12350.
- Žižňáková L, Větrovský T, Lombard V et al. Feed in summer, rest in winter: microbial carbon utilization in forest topsoil. *Microbiome* 2017;**5**:122.
- Zhu S, Gao B, Harvey PJ et al. Dermatophytic defensin with anti-infective potential. *Proc Natl Acad Sci* 2012;**109**:8495–500.