
Letter to Glyco-Forum

Ten years of *CAZypedia*: a living encyclopedia of carbohydrate-active enzymes

The *CAZypedia* Consortium*

*A list of contributors at the time of publication is provided in the Acknowledgments. All past and future *CAZypedia* Editors and Authors are invited to cite this article in reference to their invaluable contributions to this community resource.

Dedication: CAZypedia is dedicated to Emeritus Professor Bruce Stone (1928–2008†), whose enthusiasm to create a comprehensive encyclopedia of carbohydrate-active enzymes was essential to the genesis of this resource.

Abstract

CAZypedia was initiated in 2007 to create a comprehensive, living encyclopedia of the carbohydrate-active enzymes (CAZymes) and associated carbohydrate-binding modules involved in the synthesis, modification and degradation of complex carbohydrates. *CAZypedia* is closely connected with the actively curated CAZy database, which provides a sequence-based foundation for the biochemical, mechanistic and structural characterization of these diverse proteins. Now celebrating its 10th anniversary online, *CAZypedia* is a successful example of dynamic, community-driven and expert-based biocuration. *CAZypedia* is an open-access resource available at URL <http://www.cazypedia.org>.

Key words: biocuration, bioinformatics, carbohydrate-active enzymes, glycoscience, glycobiology

Background

The Carbohydrate-Active Enzymes (CAZymes) classification groups catalytic and substrate-binding modules of proteins responsible for the assembly and breakdown of complex carbohydrates into sequence-based families. Since the original definition of 35 glycoside hydrolase (GH) families in 1991 (Henrissat 1991), the CAZy database (available at URL <http://www.cazy.org/>) continues to grow and currently (October 2017) encompasses 105 glycosyltransferase (GT) families, 145 GH families, 27 polysaccharide lyase (PL) families, 16 carbohydrate esterase (CE) families, 13 auxiliary activity (AA) families and 81 carbohydrate-binding module (CBM) families (Levasseur et al. 2013; Lombard et al. 2014). As a result of vigorous biocuration [as defined by Bourne and McEntyre (2006)] and tireless technical development in response to an ever-increasing rate of gene sequencing, the CAZy database has become the de facto framework that unites protein sequence, biochemical and structural data among the tremendous diversity of CAZymes in nature [see (Davies and Sinnott 2008) for an accessible primer and review].

The CAZy database is arranged in a conventional format, with individual family pages consisting of tables of protein names, GenBank and/or UniProt sequence accession codes, EC numbers (when activity has been experimentally defined) and Protein Data Bank accession codes (when a structure has been solved). Each family

page contains a compact header that summarizes key information on substrate specificity, catalytic mechanism, three-dimensional protein fold and carbohydrate ligand complexes. Additionally, individual genome pages provide a convenient census of all CAZyme families in individual organisms (Lombard et al. 2014). In keeping with its primary function to list individual family members, family pages in the CAZy database are efficiently minimalistic. *CAZypedia* arose from the idea that a more detailed and directly accessible summary of the key research on individual CAZy families would be of significant value to glycoscience researchers, particularly highlighting the primacy of key research discoveries in a family, and supporting the activities of all scientists interested in CAZymes.

Genesis

CAZypedia's roots can be traced to renowned polysaccharide biochemist Professor Bruce Stone (1928–2008†; Whelan 2009) who proposed the idea of a comprehensive encyclopedia of the CAZymes. Bruce initially raised this idea informally at the 23rd International Carbohydrate Symposium (ICS; Whistler, Canada; July 2006) among a select group of glycoscientists, including Harry Brumer, Anthony Clarke, Gideon Davies, Harry Gilbert, Bernard Henrissat, Antoni Planas, Birte Svensson, David Vocadlo, Spencer Williams, Stephen

Withers and others. Bruce's original vision was to produce a traditional printed book or series, comprising chapters written by specific experts on individual families. It was recognized early on that the sheer number of families at that time (>100 GH families alone), combined with rapid advancements in the field, would make the timely completion of a printed work with lasting value a Sisyphean task.

Further ad hoc discussions about the best way to bring Bruce's vision to fruition continued through subsequent months, culminating at a second, larger group discussion at the seventh Carbohydrate Bioengineering Meeting (CBM7; Braunschweig, Germany; April 2007). Among those in attendance were (again) Bruce Stone, Harry Brumer, Anthony Clarke, Harry Gilbert, Antoni Planas and Birte Svensson, as well as Vincent Bulone, Marco Moracci, Warren Wakarchuk, Tony Warren, Lisa Willis and others. Here, there was general agreement that only an online, internet-based format would have sufficient flexibility and immediacy to match the rapid advances being made in CAZymology. Inspired by the growing impact of *Wikipedia* as a community-based publishing model of encyclopedic information, the idea to use a *wiki* (see definition at URL <https://en.wikipedia.org/wiki/Wiki>) approach to develop an online "Encyclopedia of Carbohydrate-Active Enzymes" was adopted. Hence, *CAZypedia* was born in May 2007 when Harry Brumer, then of the Kungliga Tekniska Högskolan in Stockholm, established *CAZypedia* using the MediaWiki software (freely available at URL <https://www.mediawiki.org/>).

Content

Content creation for *CAZypedia* was focussed initially on the GH families, due to a particularly long and rich history of biochemical and structural characterization of these enzymes (Sinnott 1990; Davies and Henrissat 1995). An original set of pages covering families GH1, GH2, GH10 and GH11 by Stephen Withers, together with GH27 and GH36 by Harry Brumer, were produced and refined with editorial input from Bernard Henrissat through the summer of 2007. In this process, a streamlined page format was devised (Figure 1), comprising individual sections on "Substrate specificities", "Kinetics and mechanism", "Catalytic residues" and "Three-dimensional structures", which present a concise summary of common features of each family. A "Family Firsts" section provides a brief, itemized list of references to seminal publications that define the key mechanistic and structural features of the family: the first reaction stereochemistry determination, catalytic residue identification and three-dimensional structure solution. An overarching goal in page design was to provide a rapid entry into the key primary literature on each family (which is not directly available in the CAZy database), through an abbreviated and consistent format. *CAZypedia* pages may be beneficially embellished with figures, although this is optional.

As part of an explicit design intent, *CAZypedia* pages do not necessarily strive to provide comprehensive reviews of all the available literature on individual families, although it should be noted that there is formally no prescribed page length. The reasons for this are largely practical. Initially, pages can be composed rapidly by focussing on the key defining literature. Compilation of a comprehensive corpus of the published work on a family, which is in many cases extensive when all individual biochemical characterization studies are considered, is therefore not required. This focus also helps to future-proof pages in a rapidly evolving field: first achievements will always remain historically significant, regardless of the number of subsequent publications on a family. For the same reason, pages explicitly avoid enumeration of time-sensitive data, such

as the number of sequences or structures for individual families, which can otherwise be gleaned from the continually updated CAZy database (individual *CAZypedia* and CAZy database pages are cross-linked for this purpose). Thus, *CAZypedia* pages are designed to be perpetually accurate, regardless of the frequency of future updates from page authors. The appellation "Curator Approved" is given to each newly minted family page once all sections contain a basic coverage of the seminal literature (see also "Technical aspects" section below).

From the initial seed of six GH families, *CAZypedia* has grown to include over 100 individual Curator Approved GH family pages, produced by a similar number of expert contributors from the CAZyme/glycoscience community. Indeed, July 2014 marked a watershed in *CAZypedia*'s history, with the completion of the GH12 page by Gerlind Sulzenbacher as the 100th Curator Approved GH page. Pages on other groups of CAZymes (i.e., Glycosyltransferases (Coutinho et al. 2003), Polysaccharide Lyases (Lombard et al. 2010) and Auxiliary Activity redox enzymes (Levasseur et al. 2013)) and non-catalytic CBMs (Boraston et al. 2004) continue to be incorporated through growing community engagement. Notable *CAZypedia* firsts include the completion of the GT42 page by Warren Wakarchuk in April 2010, the PL2 page by Wade Abbott in September 2013, the AA9 lytic polysaccharide mono-oxygenase page by Paul Harris in September 2013, and the CBM32 page by Elizabeth Ficko-Blean and Alisdair Boraston in May 2013. *CAZypedia*'s History page (available at URL <https://www.cazypedia.org/index.php/CAZypedia:History>, accessed via the About *CAZypedia* menu.) serves as a repository for these and future major milestones, while the News page (available at URL <https://www.cazypedia.org/index.php/News> and via *CAZypedia*'s Main Page.) covers recent Curator Approved pages and other newsworthy items.

In recognition of the complex nature of carbohydrate chemistry and CAZymes, *CAZypedia* also incorporates a Lexicon that provides a definition of key terms, explanation of specialist nomenclature and tutorial reviews of concepts that are relevant to individual family pages. The Lexicon provides a touchstone for new readers to support their understanding and interpretation of individual families, and is hyperlinked within the text of family pages. The Lexicon and category pages for each major CAZyme class are conveniently accessed under the Content menu, prominently displayed on the left side of all *CAZypedia* pages (Figure 1).

At its 10th anniversary online, *CAZypedia* currently comprises 106 GH, 10 CBM, 6 PL, 2 AA, 2 GT and 22 Lexicon pages with Curator Approved status. The MediaWiki software upon which *CAZypedia* relies tracks usage statistics, which are available through the Special Pages menu item. These statistics reveal over 12 million total page views, and over one hundred thousand views for several of the most popular GH and Lexicon pages. More conservative estimates of activity provided by Google Analytics indicate that *CAZypedia* access has increased to thousands of international users per week since data recording on that utility began in the autumn of 2009 (Figure 2). Regardless of the absolute values, these data highlight the sustained and growing value of *CAZypedia* to specialists and non-specialists alike.

Editorial framework

During the birth of *CAZypedia*, there was significant concern about the potential pitfalls of applying directly the *Wikipedia* model, which allows author anonymity and lacks formal editorial oversight, to the publication of a rigorous scientific encyclopedia. Thus,

CAZypedia
CARBOHYDRATE-ACTIVE
ENZYMES

Page | Discussion | View | View source | History | Search | Go | Search

New to the CAZy classification? Read this first.

GLYCOSIDE HYDROLASE FAMILY 66

This page has been approved by the Responsible Curator as essentially complete. CAZypedia is a living document, so further improvement of this page is still possible. If you would like to suggest an addition or correction, please contact the page's Responsible Curator directly by e-mail, or using this form.

- Author: Ryoichiro Suzuki
- Responsible Curator: Zui Fujimoto

Contents [\[hide\]](#)

- Substrate specificities
- Kinetics and Mechanism
- Catalytic Residues
- Three-dimensional structures
- Family Firsts
- References

Substrate specificities

Glycoside hydrolases of family GH66 include *endo*-acting dextranases (Dex; EC 3.2.1.11 [\[d\]](#)) and cyclotransferases (CITase; EC 2.4.1.48 [\[d\]](#)). Family GH66 enzymes are classified into the following three types: Type I Dexs, Type II Dexs with low CITase activity, and Type III CITases [\[1, 2\]](#).

Dex enzymes hydrolyze α -1,6-linkages of dextran and produce isomaltosaccharides (IGs) of varying length. Dex enzymes from oral streptococci have been studied since the 1970s [\[3, 4, 5\]](#). Dexs are classified into families GH49 and GH66. CITases catalyze intramolecular transglycosylation to produce cyclotransferase products (CIs; cyclodextrins) with degree of polymerization of 7-17 [\[6\]](#). CITases produce CIs from IG4 and larger IGs [\[7\]](#). CITase from *Bacillus* sp. T-3040 (CITase-T3040) produced CI-8 predominantly from dextran 40, whereas the major product of CITase from *Paenibacillus* sp. 598K (CITase-598K) was CI-7 [\[7, 8\]](#). CITases contain a CITase-specific insertion (about 90 residues) inside the catalytic domain. The insertion region is a family 35 carbohydrate-binding module (CBM35) domain [\[1\]](#). Some Dexs displaying strong dextranolytic activity with low cyclization activity have been discovered [\[1, 2\]](#).

Kinetics and Mechanism

GH66 enzymes are *retaining* enzymes, as first shown by structural analysis of cyclic dextrins formed by transglycosylation from α -1,6-glucan by *Bacillus* sp. T-3040 CITase-T3040 [\[9\]](#). This has been supported by subsequent structural [\[10\]](#) and chemical rescue studies [\[1\]](#). GH66 enzymes appear to operate through a classical Koshland retaining mechanism. The K_{cat} and K_M values of Dex from *Bacteroides thetaiotaomicron* VP1-5482 (BDex) toward dextran T2000 were determined to be 86.7 s^{-1} and 0.029 mM , respectively [\[2\]](#). Both CITase-T3040 and CITase-598K showed the same K_M value for dextran 40 (0.18 mM) [\[7\]](#). The K_{cat} values of CITase-T3040 and CITase-598K against dextran 40 were 3.2 s^{-1} and 5.8 s^{-1} , respectively [\[7\]](#). Dexs from family GH49 are inverting enzymes.

Catalytic Residues

Catalytic residues of several GH66 enzymes have been identified by mutational and structural studies [\[1, 7, 10, 11\]](#). The catalytic nucleophile is aspartic acid and the general acid/base is glutamic acid. Asp385 and Glu453 are nucleophile and acid/base catalyst, respectively. In Dex from *Streptococcus mutans* SmDex [\[10, 11\]](#), Asp340 and Glu412 in Dex from *Paenibacillus* sp. (PsDex) [\[1\]](#), Asp270 and Glu342 in CITase-T3040 [\[7, 12\]](#), and Asp269 and Glu341 in CITase-598K [\[7\]](#).

Three-dimensional structures

Crystal structures of a truncated mutant of *Streptococcus mutans* SmDex (lacking the N-terminal 99 and C-terminal 118 residues) have been reported as the first three-dimensional structure of a GH66 enzyme [\[10\]](#). Three structures, ligand free (PDB ID [3vnn](#) [\[d\]](#)), in complex with IG3 (PDB ID [3vno](#) [\[d\]](#)), and in complex with 4',5'-epoxypentyl α -D-glucopyranoside (PDB ID [3vnp](#) [\[d\]](#)), have been solved [\[10\]](#). The catalytic domain of SmDex is a β (α) β -barrel fold, accompanied by N-terminal immunoglobulin-like β -sandwich fold and C-terminal β -sandwich structure containing two Greek key motifs. These three domains are the common structural components in GH66 enzymes.

A structure for a GH66 CITase-T3040 (PDB ID [3vnn](#) [\[d\]](#), [3vno](#) [\[d\]](#)) has been reported [\[12\]](#). CITase-T3040 has a similar domain arrangement to that of SmDex, but a CBM35 domain is inserted into the catalytic module, which assists substrate uptake and production of the dominant cytoctylisomaltose (CI-8).

Family Firsts

First stereochemistry determination
Bacillus sp. T-3040 CITase-T3040 by structural analysis of transglycosylation products using $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ spectroscopy [\[9\]](#).

First catalytic nucleophile identification
Streptococcus mutans SmDex and *Paenibacillus* sp. PsDex by structural study [\[10\]](#) and chemical rescue approach [\[1\]](#), respectively.

First general acid/base residue identification
SmDex and PsDex by structural study [\[10\]](#) and chemical rescue approach [\[1\]](#), respectively.

First 3-D structure
Truncated mutant of SmDex [\[10\]](#).

References

- Kim YM, Kiso Y, Muraki T, Kang MS, Nakai H, Saburi W, Lang W, Kang HK, Okuyama M, Mori H, Suzuki R, Funane K, Suzuki N, Momma M, Fujimoto Z, Oguma T, Kobayashi M, Kim D, and Kimura A. Novel dextranase catalyzing cyclotransferase formation and identification of catalytic amino acids and their functions using chemical rescue approach. *J Biol Chem*. 2012 Jun 8;287(24):19927-35. DOI:10.1074/jbc.M111.339036 | PubMed ID:22461618 | PubMed [Xin2012A]
- Kim YM, Yamamoto E, Kang MS, Nakai H, Saburi W, Okuyama M, Mori H, Funane K, Momma M, Fujimoto Z, Kobayashi M, Kim D, and Kimura A. *Bacteroides thetaiotaomicron* VP1-5482 glycoside hydrolase family 66 homolog catalyzes dextranolytic and cyclization reactions. *FEBS J*. 2012 Sep;279(17):3185-91. DOI:10.1111/j.1742-4658.2012.08698.x | PubMed ID:22776355 | PubMed [Xin2012B]
- Staat RH and Schachtel CF. Evolution of dextranase production by the cariogenic bacterium *Streptococcus mutans*. *Infect Immun*. 1974 Feb;9(2):467-9. PubMed ID:4816468 | PubMed [Staat1974]
- Hamada S, Mizuno J, Murayama Y, Ooshima Y, and Masuda N. Effect of dextranase on the extracellular polysaccharide synthesis of *Streptococcus mutans*; chemical and scanning electron microscopy studies. *Infect Immun*. 1975 Dec;12(6):1415-25. PubMed ID:1205620 | PubMed [Hamada1975]
- Ellis DW and Miller CH. Extracellular dextran hydrolase from *Streptococcus mutans* strain 6715. *J Dent Res*. 1977 Jan;56(1):57-69. DOI:10.1177/00220345770560011301 | PubMed ID:141177 | PubMed [Ellis1977]
- Funane K, Terasawa K, Mizuno Y, Ono H, Gibu S, Tokashiki T, Kawabata Y, Kim YM, Kimura A, and Kobayashi M. Isolation of *Bacillus* and *Paenibacillus* bacterial strains that produce large molecules of cyclic isomaltosaccharides. *Biosci Biotechnol Biochem*. 2008 Dec;72(12):3277-80. PubMed ID:19060390 | PubMed [Funane2008]
- Suzuki R, Terasawa K, Kimura K, Fujimoto Z, Momma M, Kobayashi M, Kimura A, and Funane K. Biochemical characterization of a novel cyclotransferase from *Paenibacillus* sp. 598K. *Biochim Biophys Acta*. 2011 Mar;1814(3):428-34. DOI:10.1016/j.bbapap.2010.12.009 | PubMed ID:21193067 | PubMed [Funane2011]
- Funane K, Kawabata Y, Suzuki R, Kim YM, Kang HK, Suzuki N, Kimura A, and Kobayashi M. Deletion analysis of regions at the C-terminal part of cyclotransferase from *Bacillus circulans* T-3040. *Biochim Biophys Acta*. 2011 Mar;1814(3):428-34. DOI:10.1016/j.bbapap.2010.12.009 | PubMed ID:21193067 | PubMed [Funane2011]
- Oguma T, Horuchi T, and Kobayashi M. Novel Cyclic Dextrins, Cyclotransferase Products, from *Bacillus* sp. T-3040 Culture. *Biosci Biotechnol Biochem*. 1993 S7(7):1225-1227. DOI:10.1271/bbb.57.1225d [Oguma1993]
- Suzuki N, Kim YM, Fujimoto Z, Momma M, Okuyama M, Mori H, Funane K, and Kimura A. Structural elucidation of dextran degradation mechanism by streptococcus mutans dextranase belonging to glycoside hydrolase family 66. *J Biol Chem*. 2012 Jun 8;287(24):19916-26. DOI:10.1074/jbc.M112.340444 | PubMed ID:22337884 | PubMed [Suzuki2012]
- Igarashi T, Morisaki H, Yamamoto A, and Goto N. An essential amino acid residue for catalytic activity of the dextranase of *Streptococcus mutans*. *Oral Microbiol Immunol*. 2002 Jun;17(3):193-6. PubMed ID:12020973 | PubMed [Igarashi2002]
- Suzuki N, Fujimoto Z, Kim YM, Momma M, Nishine N, Suzuki R, Suzuki S, Kitamura S, Kobayashi M, Kimura A, and Funane K. Structural elucidation of the cyclization mechanism of α -1,6-glucan by *Bacillus circulans* T-3040 cyclotransferase. *J Biol Chem*. 2014 Apr 25;289(17):12040-51. DOI:10.1074/jbc.M114.547992 | PubMed ID:24616103 | PubMed [Suzuki2014]

All Medline abstracts: PubMed | PubMed

Categories: Curator approved | Glycoside Hydrolase Families

This page was last modified on 6 September 2017, at 17:33.
This page has been accessed 43,646 times.
© 2007-2017 The Authors and Curators of CAZypedia | Open Access | How to cite CAZypedia articles
About CAZypedia | Disclaimers

CAZypedia | |

Fig. 1. Layout of a typical CAZyme family page in CAZypedia.

although CAZypedia adopts many of the general principles and rules of Wikipedia, CAZypedia draws on best-practice authoring and editing principles of peer-reviewed, wiki-based encyclopedia such as *Citizendium* (available at URL <http://en.citizendium.org/>) and *Scholarpedia* (available at URL <http://www.scholarpedia.org/>). CAZypedia strives to be a dynamic, community-based resource, which at the same time balances the need for careful content curation. A full description of CAZypedia's editorial policies is available on the About page (see URL <https://www.cazypedia.org/index.php/>

CAZypedia:About, accessed via Introduction to CAZypedia under the About CAZypedia menu); however, a few points deserve special comment.

The editorial organization of CAZypedia is designed with a minimum of bureaucratic and administrative overhead, because it is entirely volunteer-based and has no direct funding support. CAZypedia generally adopts Wikipedia's Simplified Ruleset (available at URL http://en.wikipedia.org/wiki/Wikipedia:Simplified_Ruleset), particularly the concepts of using a *neutral point-of-view*, writing *verifiable text*, including




Fig. 2. CAZypedia usage statistics from Google Analytics. Access tracking with this service was initiated in August 2009. Sharp dips correspond to December holidays and broad troughs correspond to summer in the northern hemisphere.

only peer-reviewed information (no original research), being *civil and well-behaved*, and *not infringing copyright*. As a culmination of these principles, CAZypedia reports on—but does not engage in critique of—the published literature, and supports all statements of fact with primary citations. Not least, *Wikipedia's* extensive “What Wikipedia is not” page (available at URL http://en.wikipedia.org/wiki/Wikipedia:What_Wikipedia_is_not) can be translated to “What CAZypedia is not” essentially point-by-point.

Following the *Citizenium* model, transparency is achieved through the use of contributors' real names in CAZypedia. Additionally, individual biographical pages enable readers to evaluate directly each contributor's expertise in the field. To maintain editorial quality control, every Family and Lexicon page in CAZypedia is overseen by a *Responsible Curator*, who is primarily responsible for overall content. Responsible Curators are selected by a panel of *Senior Curators* based on established expertise and a willingness to participate in the active maintenance of specific pages. In turn, Responsible Curators are tasked with recruiting and managing *Authors* to participate in content creation; Responsible Curators may also contribute directly to composing page content.

In the spirit of a community-driven resource, individuals are encouraged to self-nominate to become Responsible Curators or Authors. In general, individuals at any career stage are welcomed to participate as Authors, including keen undergraduates, postgraduate students and post-doctoral scientists. Indeed, the current list of contributors (see below) includes many junior scientists (or scientists who were at least junior at the time of their first contribution). Ultimately, the quality of entries in CAZypedia, like *Wikipedia*, relies upon the keen eye of readers at-large to identify errors and omissions. All users who spot such oversights are encouraged to contact the Responsible Curator for that page, so that a correction can be made.

CAZypedia is an open-access publication, i.e., it is freely available online for anyone to read, study and otherwise use for scholarly pursuits. However, the Authors and Curators of CAZypedia assert their copyright for the sole purpose of preventing outright duplication and uncontrolled modification of the content, which could undermine the

expert-based nature of this resource. Although we strongly advocate that readers should cite the primary research literature directly, individual CAZypedia pages may also be cited when practical, analogous to a book chapter or review article. Citation details are provided in the footer and via the Tools menu on each page (Figure 1).

Technical aspects

Wiki-wiki

As introduced above, CAZypedia runs on MediaWiki, the free, open source PHP software originally developed for *Wikipedia*. This choice was based on the demonstrated robustness and scalability of MediaWiki, as well as the availability of diverse software extensions to add functionality. As *Wikipedia* is unlikely to disappear anytime soon, so too is MediaWiki's active community of developers likely to persist well into the future, thereby ensuring continued maintenance of the software running CAZypedia. A full technical and functional description of MediaWiki is beyond the scope of this Letter; interested readers should visit MediaWiki.org for more details.

For the content contributor and user, the most important practical aspect of the use of MediaWiki is that CAZypedia is a *wiki*: edits are displayed instantaneously when saved and do not require approval before appearing online. This enables dynamic development of page content driven by individual Authors. In the initial stages of development, pages are clearly marked as “Under Construction”, with a warning that content is under revision and may be subject to major changes. Once vetted by the Responsible Curator, a page may be upgraded to “Curator Approved” status to indicate that it is factually accurate and essentially complete. However, “completeness” is not absolute: as a wiki, CAZypedia is a living document, so further development of page content is forever possible.

Creating content for CAZypedia is relatively intuitive. Once a new Author has been provided with a login, page editing can be conducted within a modern web browser using a simplified markup language. A boilerplate pre-populates the page with the major template features, and Authors can view the code of other pages to get ideas of ways to insert features like hyperlinks, references and figures. A “Getting Started Guide”, along with concise pages that provide help with editing, references and adding images provide guidance to assist the novice. Here, too, the use of MediaWiki as software platform is a considerable benefit, due to vast extant help resources on editing. Finally, assistance is always at hand from CAZypedia Curators, who are able to activity monitor edits via the global “Recent Changes” and individual “History” pages.

BiblioPlus

MediaWiki functionality can be enhanced through extensions, and CAZypedia utilizes several, including those for user administration, defining page boilerplate content and integrating Google Analytics. Among these, BiblioPlus (freely available at URL <https://www.mediawiki.org/wiki/Extension:BiblioPlus>) deserves special mention as the MediaWiki extension that drives bibliographic referencing. BiblioPlus is the result of a significant effort by CAZypedia contributor Karen Eddy to correct compatibility issues arising in the original Biblio extension by Martin Jambon and others (see URL <https://www.mediawiki.org/wiki/Extension:Biblio>).

Like its predecessor, BiblioPlus performs automated retrieval and formatting of citations from PubMed and the ISBN databases in MediaWiki pages. Similar to other reference formatting software, BiblioPlus automatically numbers in-text citations and generates a

reference section, which is included at the bottom of a page. Notably, the reference section contains hyperlinks to original sources, specifically PubMed or the ISBNdb, HubMed and DOI hyperlinks. BiblioPlus was specifically re-coded to utilize the modern NCBI Entrez Programming Utilities (E-utilities) interface (Anonymous 2010). A full description of features and usage instructions is available on the BiblioPlus Mediawiki extension page (URL <https://www.mediawiki.org/wiki/Extension:BiblioPlus>). It should be noted that BiblioPlus is freely available and will work together with any modern MediaWiki implementation, so that it may be broadly deployed in any wiki, scientific or otherwise.

The next 10 years: CAZyedia needs you!

The continued success of *CAZyedia* will remain entirely dependent on the diligence and commitment of experts and keen junior scientists to voluntarily contribute to the maintenance and growth of this reference work. The job of building *CAZyedia* is by no means complete, and as a living encyclopedia, it never will be—especially as research continues to reveal new CAZyme families, tertiary structures and mechanistic details (e.g., Campos et al. 2016; Venditto et al. 2016; Abe et al. 2017; Munoz-Munoz, Cartmell, Terrapon, Basle 2017; Munoz-Munoz, Cartmell, Terrapon, Henrissat 2017; Ndeh et al. 2017). Currently, many pages remain to be written and existing pages would benefit from regular updates as new data come to hand, which requires expert volunteers willing to assume the responsibility for page creation and maintenance.

Thus, the *CAZyedia* Consortium openly invites all interested glycoscientists, regardless of career stage (including keen undergraduate and postgraduate students, post-doctoral researchers, industrial scientists and professors) to peruse the “Unassigned Pages” lists for each CAZyme class and see if they might be able to help. The growth of *CAZyedia* will depend exclusively on the generous and selfless contributions of the existing and new generations of CAZyepedians. We invite you to join us! Contact information is available at URL <http://www.cazyedia.org/>.

Acknowledgments

We thank Stephen MacDonald and Vince Tingey (Michael Smith Laboratories, University of British Columbia), and Eric Björkqvall (School of Biotechnology, Kungliga Tekniska Högskolan), for invaluable IT support. Dr. Karen Eddy (Brumer group, MSL, UBC) is thanked for developing the BiblioPlus extension.

CAZyedia is the result of many hours of effort by the following group of current contributors (A continually updated list is available at URL <http://www.cazyedia.org/index.php/Category:Contributors>, accessed via the “About *CAZyedia*” menu.): Wade Abbott, Agriculture and Agri-Food Canada, Canada; Orly Alber, Weizmann Institute of Science, Israel; Ed Bayer, Weizmann Institute of Science, Israel; Jean-Guy Berrin, Institut National de la Recherche Agronomique, France; Alisdair Boraston, University of Victoria, Canada; Harry Brumer, University of British Columbia, Canada; Ryszard Brzezinski, Université de Sherbrooke, Canada; Anthony Clarke, University of Guelph, Canada; Beatrice Cobucci-Ponzano, National Research Council of Italy, Italy; Darrell Cockburn, Penn State University, United States of America; Pedro Coutinho, Aix Marseille Université, France; Mirjam Czejek, Centre National de la Recherche Scientifique, France; Bareket Dassa, Weizmann Institute of Science, Israel; Gideon John Davies, University of York, United Kingdom; Vincent Eijsink, Norwegian University of Life Sciences, Norway; Jens Eklöf, University of British Columbia, Canada; Alfons Felice, Universität für Bodenkultur, Austria; Elizabeth Ficko-Blean, Centre National de la Recherche Scientifique, France; Geoff Fincher, University of Adelaide, Australia; Thierry Fontaine, Institut Pasteur, France; Zui Fujimoto, National Agriculture and Food Research Organisation, Japan; Kiyotaka Fujita, Kagoshima University, Japan; Shinya Fushinobu, University of Tokyo,

Japan; Harry Gilbert, Newcastle University, United Kingdom; Tracey Gloster, University of St. Andrews, United Kingdom; Ethan Goddard-Borger, Walter and Eliza Hall Institute of Medical Research, Australia; Ian Greig, Simon Fraser University, Canada; Jan-Hendrik Hehemann, Max Planck Institute for Marine Microbiology, Germany; Glyn Hemsworth, University of Leeds, United Kingdom; Bernard Henrissat, Centre National de la Recherche Scientifique, France; Masafumi Hidaka, University of Tokyo, Japan; Ramon Hurtado-Guerrero, University of Zaragoza, Spain; Kiyohiko Igarashi, University of Tokyo, Japan; Takuya Ishida, University of Tokyo, Japan; Stefan Janecek, Slovak Academy of Sciences, Slovakia; Seino Jongkees, University of Tokyo, Japan; Nathalie Juge, Quadram Institute, United Kingdom; Satoshi Kaneko, University of the Ryukyus, Japan; Takane Katayama, Ishikawa Prefectural University, Japan; Motomitsu Kitaoka, National Agriculture and Food Research Organisation, Japan; Naotake Konno, Utsunomiya University, Japan; Daniel Kracher, Universität für Bodenkultur, Austria; Anna Kulminskaya, Petersburg Nuclear Physics Institute, Russia; Alicia Lammerts van Bueren, University of Groningen, Netherlands; Sine Larsen, University of Copenhagen, Denmark; Junho Lee, University of British Columbia, Canada; Markus Linder, Aalto University, Finland; Leila LoLeggio, University of Copenhagen, Denmark; Roland Ludwig, Universität für Bodenkultur, Austria; Ana Luis, University of Lisbon, Portugal; Mirko Maksimainen, University of Oulu, Finland; Brian Mark, University of Manitoba, Canada; Richard McLean, University of Lethbridge, Canada; Gurvan Michel, Centre National de la Recherche Scientifique, France; Cedric Montanier, Institut National de la Recherche Agronomique, France; Marco Moracci, National Research Council of Italy, Italy; Haruhide Mori, Hokkaido University, Japan; Hiroyuki Nakai, Niigata University, Japan; Wim Nerinckx, Ghent University, Belgium; Takayuki Ohnuma, Kinki University, Japan; Richard Pickersgill, Queen Mary University of London, United Kingdom; Kathleen Piens, Sveriges Lantbruksuniversitet, Sweden; Tirso Pons, National Centre for Biotechnology, Spain; Etienne Rebuffet, Centre National de la Recherche Scientifique, France; Peter Reilly, Iowa State University, United States of America; Magali Remaud-Simeon, Institut National des Sciences Appliquées, France; Brian Rempel, University of British Columbia, Canada; Kyle Robinson, University of British Columbia, Canada; David Rose, University of Waterloo, Canada; Juha Rouvinen, University of Eastern Finland, Finland; Wataru Saburi, Hokkaido University, Japan; Yuichi Sakamoto, Iwate Biotechnology Research Center, Japan; Mats Sandgren, Sveriges Lantbruksuniversitet, Sweden; Fathima Shaikh, University of British Columbia, Canada; Yuval Shoham, Technion, Israel; Franz St. John, United States Department of Agriculture, United States of America; Jerry Stahlberg, Sveriges Lantbruksuniversitet, Sweden; Michael Suits, Wilfrid Laurier University, Canada; Gerlind Sulzenbacher, Centre National de la Recherche Scientifique, France; Tomomi Sumida, RIKEN, Japan; Ryuichiro Suzuki, Akita Prefectural University, Japan; Birte Svensson, Danmarks Tekniske Universitet, Denmark; Toki Taira, University of the Ryukyus, Japan; Ed Taylor, University of Lincoln, United Kingdom; Takashi Tonozuka, Tokyo University of Agriculture and Technology, Japan; Breeanna Urbanowicz, University of Georgia, United States of America; Gustav Vaaje-Kolstad, Norwegian University of Life Sciences, Norway; Wim Van den Ende, Katholieke Universiteit Leuven, Belgium; Annabelle Varrot, Centre National de la Recherche Scientifique, France; Maxime Versluys, Katholieke Universiteit Leuven, Belgium; Florence Vincent, Centre National de la Recherche Scientifique, France; David Voadlo, Simon Fraser University, Canada; Warren Wakarchuk, Ryerson University, Canada; Tom Wenekes, Universiteit Utrecht, Netherlands; Rohan Williams, University of Melbourne, Australia; Spencer Williams, University of Melbourne, Australia; David Wilson†, Cornell University, United States of America; Stephen Withers, University of British Columbia, Canada; Katsuro Yaoi, National Institute of Advanced Industrial Science and Technology, Japan; Vivian Yip, University of British Columbia, Canada; Ran Zhang, University of British Columbia, Canada.

This Letter was composed by H. Brumer (brumer@msl.ubc.ca) and S.J. Williams (sjwill@unimelb.edu.au), with input from B. Svensson, B. Henrissat, G.J. Davies, H.J. Gilbert, A.J. Clarke, W.W. Wakarchuk, D.W. Abbott, D.J. Voadlo, E. Ficko-Blean, A.B. Boraston, A. Planas, and S. Fushinobu.

Page charges for this article were supported by a Natural Sciences and Engineering Research Council of Canada (NSERC) Discovery Grant to H. Brumer.

References

- Abe K, Nakajima M, Yamashita T, Matsunaga H, Kamisuki S, Nihira T, Takahashi Y, Sugimoto N, Miyanaga A, Nakai H et al. 2017. Biochemical and structural analyses of a bacterial endo-beta-1,2-glucanase reveal a new glycoside hydrolase family. *J Biol Chem.* 292:7487–7506.
- Anonymous. 2010. *Entrez Programming Utilities Help* National Center for Biotechnology Information: Bethesda, MD, USA. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK25501/>.
- Boraston AB, Bolam DN, Gilbert HJ, Davies GJ. 2004. Carbohydrate-binding modules: Fine-tuning polysaccharide recognition. *Biochem J.* 382:769–781.
- Bourne PE, McEntyre J. 2006. Biocurators: Contributors to the world of science. *PLoS Comput Biol.* 2:1185–1185.
- Campos BM, Liberato MV, Alvarez TM, Zanthorlin LM, Ematsu GC, Barud H, Polikarpov I, Ruller R, Gilbert HJ, Zeri ACD et al. 2016. A novel carbohydrate-binding module from sugar cane soil metagenome featuring unique structural and carbohydrate affinity properties. *J Biol Chem.* 291:23734–23743.
- Coutinho PM, Deleury E, Davies GJ, Henrissat B. 2003. An evolving hierarchical family classification for glycosyltransferases. *J Mol Biol.* 328:307–317.
- Davies G, Henrissat B. 1995. Structures and mechanisms of glycosyl hydrolases. *Structure.* 3:853–859.
- Davies GJ, Sinnott ML. 2008. Sorting the diverse. *Biochemist.* 30:26–32.
- Henrissat B. 1991. A classification of glycosyl hydrolases based on amino acid sequence similarities. *Biochem J.* 280:309–316.
- Levasseur A, Drula E, Lombard V, Coutinho PM, Henrissat B. 2013. Expansion of the enzymatic repertoire of the CAZy database to integrate auxiliary redox enzymes. *Biotechnol Biofuels.* 6:14.
- Lombard V, Bernard T, Rancurel C, Brumer H, Coutinho PM, Henrissat B. 2010. A hierarchical classification of polysaccharide lyases for glycogenomics. *Biochem J.* 432:437–444.
- Lombard V, Ramulu HG, Drula E, Coutinho PM, Henrissat B. 2014. The carbohydrate-active enzymes database (CAZy) in 2013. *Nucleic Acids Res.* 42:D490–D495.
- Munoz-Munoz J, Cartmell A, Terrapon N, Basle A, Henrissat B, Gilbert HJ. 2017. An evolutionarily distinct family of polysaccharide lyases removes rhamnose capping of complex arabinogalactan proteins. *J Biol Chem.* 292:13271–13283.
- Munoz-Munoz J, Cartmell A, Terrapon N, Henrissat B, Gilbert HJ. 2017. Unusual active site location and catalytic apparatus in a glycoside hydrolase family. *Proc Natl Acad Sci U S A.* 114:4936–4941.
- Ndeh D, Rogowski A, Cartmell A, Luis AS, Basle A, Gray J, Venditto I, Briggs J, Zhang XY, Labourel A et al. 2017. Complex pectin metabolism by gut bacteria reveals novel catalytic functions. *Nature.* 544:65–70.
- Sinnott ML. 1990. Catalytic mechanisms of enzymatic glycosyl transfer. *Chem Rev.* 90:1171–1202.
- Venditto I, Luis AS, Rydahl M, Schuckel J, Fernandes VO, Vidal-Melgosa S, Bule P, Goyal A, Pires VMR, Dourado CG et al. 2016. Complexity of the *Ruminococcus flavefaciens* cellulosome reflects an expansion in glycan recognition. *Proc Natl Acad Sci U S A.* 113:7136–7141.
- Whelan W. 2009. Obituary: Bruce A. Stone. *IUBMB Life.* 61:84–84.