




Max-Planck-Institut für Meteorologie

REPORT No. 213



MULTI - PATTERN FINGERPRINT METHOD FOR DETECTION AND ATTRIBUTION OF CLIMATE CHANGE

by
Klaus Hasselmann

HAMBURG, August 1996

Multi-pattern fingerprint method for detection and
attribution of climate change

K. Hasselmann

August 23, 1996

ISSN 0937-1060

AUTHORS:

Klaus Hasselmann

Max-Planck-Institut
für Meteorologie

MAX-PLANCK-INSTITUT
FÜR METEOROLOGIE
BUNDESSTRASSE 55
D - 20146 HAMBURG
GERMANY

Tel.: +49-(0)40-4 11 73-0
Telefax: +49-(0)40-4 11 73-298
E-Mail: <name> @ dkrz.de

Report No. 1-150

Please order the reference-list from the MPI for Meteorology, Hamburg

-
- Report No. 151**
January 1995
Changes in the Winter Precipitation in Romania and its Relation to the Large Scale Circulation
Aristita Busuioc, Hans von Storch
* Tellus, 1996 (in press)
- Report No. 152**
January 1995
Climate Variability and Change
Hans von Storch, Klaus Hasselmann
* G. Hempel (Ed.): New Trends in Ocean and Polar Sciences, 33-58, 1996
- Report No. 153**
February 1995
Regional climate changes as simulated in time-slice experiments
Ulrich Cubasch, Jürgen Waszkewitz, Gabi C. Hegerl, Jan Perlwitz
- Report No. 154**
February 1995
Water Isotope Modeling in The Asian Monsoon Region
Georg Hoffmann, Martin Heimann
* Quaternary International, 1995 (submitted)
- Report No. 155**
February 1995
A comparison of satellite observations and model simulations of column integrated moisture and upper tropospheric humidities
C.-T. Chen, Erich Roeckner, Brian J. Soden
* Journal of Climate, 1996 (in press)
- Report No. 156**
February 1995
The implementation of the semi-implicit scheme in cell-integrated semi-Lagrangian models
Bennert Machenhauer, Markus Oik
- Report No. 157**
February 1995
**The 19th century discussion of climate variability and climate chance:
Analogies for the present debate ?**
Nico Stehr, Hans von Storch, Moritz Flügel
* World Resources Review, 7, 589-604, 1996
- Report No. 158**
February 1995
Recent Northern Winter Climate Trends due to Ozone Changes and Increased Greenhouse Gas Forcing ?
Hans-F. Graf, Judith Perlwitz, Ingo Kirchner, Ingrid Schult
* Contributions to Atmospheric Physics 68, 233-248, 1995
- Report No. 159**
March 1995
The role of water vapor and convection during the Central Equatorial Pacific Experiment (CEPEX) from observations and model simulations
Ulrike Lohmann, Erich Roeckner, William D. Collins, Andrew J. Heymsfield, Greg McFarquhar, Timothy P. Barnett
* Journal of Geophysical Research, 100 (D12), 26.229-26.245, 1995
- Report No. 160**
March 1995
Sodar for precipitation measurements
Shixuan Pang, Hartmut Graßl
- Report No. 161**
May 1995
The Reduction of Complex Dynamical Systems Using Principal Interaction Patterns
Frank Kwasniok
* Physica D 92, 26-60, 1996
- Report No. 162**
May 1995
The Water Vapour Continuum and its representation in ECHAM4
Marco Giorgetta, Martin Wild
- Report No. 163**
May 1995
Modeling bio-geophysical feedback in the Sahel
Martin Claussen
- Report No. 164**
May 1995
**Stratospheric climate and variability from a general circulation model and observations.
Part II: Results for March-May, June-August and September-November.**
Elisa Manzini, Lennart Bengtsson

-
- Report No. 165**
May 1995
Tropical Intraseasonal Oscillation Appearing in Operational Analyses and in a Family of General Circulation Models
Silvio Gualdi, Hans von Storch, Antonio Navarra
- Report No. 166**
June 1995
Validation of the Earth radiation budget as simulated by the Max Planck Institute for Meteorology general circulation model ECHAM4 using satellite observations of the Earth Radiation Budget Experiment (ERBE)
C.-T. Chen, Erich Roeckner
* Journal of Geophysical Research, 101 (D2), 4.269-4.287, 1996
- Report No. 167**
June 1995
Statistically Optimal Approaches to Detecting Anthropogenic Climate Change
Gabriele C. Hegerl, Gerald R. North
* Journal of Climate, 1995 (in press) (note section)
- Report No. 168**
July 1995
Detection of Anthropogenic Climate Change using a Fingerprint Method
Klaus Hasselmann, Lennart Bengtsson, Ulrich Cubasch, Gabriele C. Hegerl, Henning Rodhe, Erich Roeckner, Hans von Storch, Reinhard Voss, Jürgen Waszkewitz
- Report No. 169**
July 1995
Techniques for Asynchronous and Periodically-synchronous Coupling of Atmosphere and Ocean Models
Part I: General Strategy and Application to the Cyclo-stationary Case
Robert Sausen, Reinhard Voss
* Climate Dynamics (accepted)
- Report No. 170**
July 1995
On the Benefit of Sea Level Assimilation in the Tropical Pacific
Martin Fischer, Mojib Latif, Moritz Flügel, Ming Ji
* Monthly Weather Review, 1996 (in press)
- Report No. 171**
July 1995
Interactions between ocean circulation and the biological pumps in the global warming
Ernst Maier-Reimer, Uwe Mikolajewicz, Arne Winguth
- Report No. 172**
October 1995
The metron model : elements of a unified deterministic theory of fields and particles
Klaus Hasselmann
* Physics Essays (accepted)
- Report No. 173**
September 1995
AGCM experiments on the Younger Dryas climate
Hans Renssen, Michael Lautenschlager, Lennart Bengtsson, Uwe Schulzweida
- Report No. 174**
September 1995
Techniques for Asynchronous and Periodically-synchronous Coupling of Atmosphere and Ocean Models
Part II: Impact of Variability
Reinhard Voss, Robert Sausen
* Climate Dynamics (submitted)
- Report No. 175**
September 1995
Greenhouse Warming, Decadal Variability, or El Niño? An Attempt to Understand the Anomalous 1990s
Mojib Latif, Richard Kleeman, Christian Eckert
* Journal of Climate (submitted)
- Report No. 176**
September 1995
High resolution GCM simulations over Europe: Surface processes
Martin Wild, Lydia Dümenil, Jan-Peter Schulz
- Report No. 177**
September 1995
Mean circulation and internal variability in an ocean primitive equation model
Sybren Drijfhout, Christoph Heinze, Mojib Latif, Ernst Maier-Reimer
* Journal of Physical Oceanography (in press)

-
- Report No. 178**
November 1995
ENSO variability and atmospheric response in a global coupled atmosphere-ocean GCM
Erich Roeckner, Josef M. Oberhuber, Andreas Bacher, Michael Christoph, Ingo Kirchner
* Climate Dynamics, 1996 (in press)
- Report No. 179**
November 1995
Introduction of a prognostic cloud ice scheme in the ECHAM general circulation model: Impact on climate and climate sensitivity
Ulrike Lohmann, Erich Roeckner
* Climate Dynamics, 1996 (in press)
- Report No. 180**
December 1995
The Chemistry of the polluted Atmosphere over Europe: Simulations and Sensitivity Studies with a regional Chemistry-Transport-Model
Bärbel Langmann, Hans-F. Graf
* Atmospheric Environment, 1996 (submitted)
- Report No. 181**
December 1995
Evaluation of Satellite-derived Latent Heat Fluxes
Jörg Schulz, Jens Meywerk, Stefan Ewald, Peter Schlüssel
* Journal of Climate, 1996 (submitted)
- Report No. 182**
December 1995
Improvements of cloud particle sizing with a 2D-Grey probe
Andreas Reuter, Stephan Bakan
- Report No. 183**
January 1996
Estimates of Climate Change in Southern Europe using Different Downscaling Techniques
Ulrich Cubasch, Hans von Storch, Jürgen Waszkewitz, Eduardo Zorita
* Climate Research, 1996 (in press)
- Report No. 184**
January 1996
The coupled GCM ECHO-2 Part I: The Tropical Pacific
Helmut Frey, Mojib Latif, Timothy Stockdale
* Monthly Weather Review's special issue on „Coupled Ocean-Atmosphere Models“, 1996 (in press)
- Report No. 185**
January 1996
A Multi-Limit Formulation for the Equilibrium Depth of a Stably Stratified Boundary Layer
Sergej Zilitinkevich, Dmitrii V. Mironov
* Boundary-Layer Meteorology, 1996 (submitted)
- Report No. 186**
January 1996
Monitoring the Temperature of the Troposphere by Means of a General Circulation Model
Martin Stendel, Lennart Bengtsson
* Journal of Geophysical Research, 1996 (submitted)
- Report No. 187**
January 1996
A Mechanism of Decadal Climate Variability
Mojib Latif, Anselm Groetzner, Matthias Münnich, Ernst Maier-Reimer, Stephan Venzke, Timothy P. Barnett
* NATO winter school on decadal variability, Les Houches, 1995
- Report No. 188**
February 1996
The Stability of the Thermohaline Circulation in a Coupled Ocean-Atmosphere General Circulation Model
Andreas Schiller, Uwe Mikolajewicz, Reinhard Voss
* Climate Dynamics, 1996 (submitted)
- Report No. 189**
February 1996
A meltwater induced collapse of the 'conveyor belt' thermohaline circulation and its influence on the distribution of $\Delta^{14}\text{C}$ and $\delta^{18}\text{O}$ in the oceans
Uwe Mikolajewicz

-
- Report No. 190**
February 1996
Modelling the Snow Cover for Climate Studies
Part I: Long-Term Integrations under Different Climatic Conditions Using a Multi-Layered Snow-Cover Model
Part II: The Sensitivity to Internal Snow Parameters and Interface Processes
Bettina Loth, Hans F. Graf
- Report No. 191**
February 1996
Validation of present-day regional climate simulations over Europe: nested LAM and variable resolution global model simulations with observed or mixed layer ocean boundary conditions
Bennert Machenhauer, Martin Windelband, Michael Botzet, Richard G. Jones, Michel Déqué
- Report No. 192**
March 1996
SECOND EDITION
June 1996
Optimization of CO₂ emissions using coupled integral climate response and simplified cost models. A sensitivity study
Klaus Hasselmann, Susanne Hasselmann, Ralf Giering, Victor Ocaña, Hans von Storch
- Report No. 193**
May 1996
Cloud simulations with the Max-Planck-Institute for Meteorology general circulation model ECHAM4 and comparison with observations
C.-T. Chen, Erich Roeckner
* Journal of Geophysical Research, 1996 (submitted)
- Report Nr. 194**
May 1996
Decadal climate variability over the North Pacific and North America: Dynamics and Predictability
Mojib Latif, Timothy P. Barnett
* Journal of Climate, 1996 (in press)
- Report Nr. 195**
May 1996
INSIDE SCIENCE
A Preliminary Investigation of the Case of Global Warming
Dennis Bray, Hans von Storch
- Report Nr. 196**
May 1996
El Hermanito: El Niño's Overlooked Little Brother in the Atlantic
Mojib Latif, Anselm Groetzner, Helmut Frey
- Report Nr. 197**
May 1996
Transient tracers in a global OGCM - source functions and simulated distributions
Christoph Heinze, Ernst Maier-Reimer, Peter Schlosser
- Report Nr. 198**
May 1996
A 1260-year control integration with the coupled ECHAM1/LSG general circulation model
Jin-Song von Storch, Viatcheslav Kharin, Ulrich Cubasch, Gabriele C. Hegerl, Dierk Schriever, Hans von Storch, Eduardo Zorita
- Report Nr. 199**
July 1996
ENSO dynamics and seasonal cycle in the tropical Pacific as simulated by the ECHAM4 / OPYC3 coupled general circulation model
Andreas Bacher, Josef M. Oberhuber, Erich Roeckner
* Climate Dynamics, 1996 (submitted)
- Report Nr. 200**
July 1996
Improved representation of surface and atmospheric radiation budgets in the ECHAM4 General Circulation Model
Martin Wild, Atsumu Ohmura, Hans Gilgen, Erich Roeckner, Marco Giorgetta
- Report Nr. 201**
July 1996
Interpretation of Interbasin exchange in an Isopycnal Ocean Model
Xue-Hong Zhang, Josef M. Oberhuber, Andreas Bacher, Erich Roeckner
* Climate Dynamics, 1996 (submitted)

-
- Report Nr. 202**
July 1996
Predictability of a Stochastically Forced Hybrid Coupled Model of El Niño
Christian Eckert, Mojib Latif
*Journal of Climate, 1996 (submitted)
- Report Nr. 203**
July 1996
On the sensitivity of an ocean general circulation model to glacial boundary conditions
Arne M.E. Winguth, Ernst Maier-Reimer, Uwe Mikolajewicz, Jean-Claude Duplessy
* Paleoceanography, 1996 (submitted)
- Report Nr. 204**
July 1996
Large-Eddy-Simulation of Contrails
Andreas Chlond
* Journal of the Atmospheric Sciences, 1996 (submitted)
- Report Nr. 205**
July 1996
Volcanic Sulfur Emissions: Estimates of Source Strength and its Contribution to the Global Sulfate Distribution
Hans-F. Graf, Johann Feichter, Bärbel Langmann
* Journal of Geophysical Research, 1996 (submitted)
- Report Nr. 206**
July 1996
Simulation with an O-AGCM of the influence of variations of the solar constant on the global climate
Ulrich Cubasch, Gabriele C. Hegerl, Reinhard Voss, Jürgen Waszkewitz, Thomas J. Crowley
* Geophysical Research Letters, 1996 (submitted)
- Report Nr. 207**
July 1996
On multi-fingerprint detection and attribution of greenhouse gas- and aerosol forced climate change
Gabriele C. Hegerl, Klaus Hasselmann, Ulrich Cubasch, John F.B. Mitchell, Erich Roeckner, Reinhard Voss, Jürgen Waszkewitz
- Report Nr. 208**
August 1996
A decadal climate cycle in the North Atlantic Ocean as simulated by the ECHO coupled GCM
Anselm Grötzner, Mojib Latif, Timothy P. Barnett
- Report Nr. 209**
August 1996
Multi-actor optimization of greenhouse gas emission paths using coupled integral climate response and economic models
Klaus Hasselmann, Susanne Hasselmann
- Report Nr. 210**
August 1996
Space Refractive Tomography of the Atmosphere: Modeling of Direct and Inverse Problems
M. E. Gorbunov, S. V. Sokolovsky, Lennart Bengtsson
- Report Nr. 211**
August 1996
Advanced Algorithms of Inversion of GPS/MET Satellite Data and their Application to Reconstruction of Temperature and Humidity
M. E. Gorbunov, A. S. Gurvich, Lennart Bengtsson
- Report Nr. 212**
August 1996
Recipes for Adjoint Code Construction
Ralf Giering, Thomas Kaminski
* ACM Transactions on Mathematical Software, 1996 (submitted)
- Report No. 213**
August 1996
Multi-pattern fingerprint method for detection and attribution of climate change
Klaus Hasselmann

ISSN 0937-1060

ABSTRACT

The multi-variate optimal fingerprint method for the detection of an externally forced climate change signal in the presence of natural internal variability is extended to the attribution problem. To determine whether a climate change signal which has been detected in observed climate data can be attributed to a particular climate forcing mechanism, or combination of mechanisms, the predicted space-time dependent climate change signal patterns for the candidate climate forcings must be specified. In addition to the signal patterns, the method requires input information on the space-time dependent covariance matrices of the natural climate variability and the predicted signal pattern errors. The detection and attribution problem is treated as a sequence of individual consistency tests applied to all candidate forcing mechanisms, as well as to the null hypothesis that no climate change has taken place, within the phase space spanned by the predicted climate change patterns. As output the method yields a significance level for the detection of a climate change signal in the observed data and individual confidence levels for the consistency of the retrieved climate change signal with each of the forcing mechanisms. A statistically significant climate change signal is regarded as consistent with a given forcing mechanism if the statistical confidence level exceeds a given critical value, but is attributed to that forcing only if all other climate change mechanisms are rejected at that confidence level. The analysis is carried out using tensor notation, with a metric given by the natural-variability covariance matrix. This clarifies the relation between the covariant signal patterns and their contravariant fingerprint counterparts. The signal patterns define the vector space in which the climate trajectories are analyzed, while the fingerprints are needed to project the climate trajectories onto this space.

1 Introduction

There is mounting evidence that the global warming due to increasing atmospheric greenhouse gas concentrations predicted by state-of-the-art coupled ocean-atmosphere global circulation models (CGCMs) is beginning to emerge from the background noise of natural climate variability (cf. summary in IPCC Second Assessment Report, Santer *et al*, 1995b). However, much of the evidence is still qualitative or circumstantial. There have been relatively few attempts to assign a quantitative measure to the probability that a climate change signal distinct from natural climate variability can be detected in observed climate data.

A basic obstacle for quantitative signal-to-noise analyses is that they require information on the space-time structure of both the predicted climate signal and the climate variability. While the predicted signal properties can be inferred from model computations, the estimation of the required space-time covariance structure of natural climate variability from model simulations and observations is more difficult. Thus although general multi-variate theories for the optimal detection of a space-time dependent climate change signal in the presence of natural climate variability noise have now been developed (Hasselmann, 1979, Bell, 1982, 1986, Hasselmann, 1993, referred to in the following as H, North *et al*, 1995, North and Kim, 1995),

gives a summary of the results and presents some conclusions.

2 The detection problem

We review in this section briefly the multi-fingerprint method of multi-variate climate change detection, following the approach of H for the general space-time dependent problem, but returning – for better illustration of the interrelationship between fingerprint and signal patterns – to the co- and contra-variant tensor notation of Hasselmann’s (1979) earlier analysis of the spatial signal-to-noise problem (see also Thacker, 1995).

Terminology

We shall use the term climate change in the following to denote the response of the climate system to external forcing, as opposed to natural internal climate variability generated by interactions within the climate system. According to this terminology, climate variations due to volcanic activity or variations in the solar constant are classed as (natural) climate change, rather than as climate variability. An alternative terminology refers to these variations also as natural variability, climate variability being regarded as a superposition of externally forced and internally generated components, the term climate change being reserved for anthropogenic climate modifications only. However, for the detection and attribution problem our definitions will be found to be more convenient. Thus climate change in our terminology can be of either natural or of anthropogenic origin, while climate variability is always natural. The definition loses precision if interactions between climate change and internal natural climate variability are considered, but in our applications we shall regard the climate state to first order simply as a linear superposition of climate change and climate variability.

The present definitions are more consistent than alternative earlier attempts to distinguish between climate change and climate variability on the basis of time scales, or in terms of climate change ‘events’ as opposed to ‘continuous’ climate fluctuations. In practice, the time scales of internal climate variability and externally forced climate change overlap, so that for a given finite time scale it is not possible to distinguish between ‘events’ and ‘continuous fluctuations’. Indeed, the impossibility of distinguishing between externally generated climate change and internal climate variability on the basis of time scale considerations alone is the essence of the detection and attribution problem.

We consider a vector time series $\phi_a(t)$ of climate data, which we assume can be represented as a superposition

$$\phi_a = \phi_a^s + \tilde{\phi}_a \quad (1)$$

of a climate change signal ϕ_a^s and a natural-variability component $\tilde{\phi}_a$. The index a refers to different types of climate data, e.g. temperature or precipitation, and to the location or averaging region of the data. The data set can represent either observed data or synthetic data from a model simulation. The climate vector ϕ_a need not represent a dynamically complete description of the climate state. In fact,

the covariance matrix C_{ij} . Thus the operations of index raising and lowering are defined by

$$\begin{aligned} X^{\dots i \dots} &= C^{ij} X^{\dots j \dots} \\ X^{\dots j \dots} &= C_{ij} X^{\dots i \dots} \end{aligned} \quad (6)$$

The definition of the climate trajectory vector as a covariant vector is arbitrary in the present context. The role of co- and contravariant variables can be interchanged. We adopt here the original assignments of Hasselmann (1979).

For each trajectory ψ there exists a constant probability surface $\rho^2(\tilde{\psi}) = C^{ij} \tilde{\psi}_i \tilde{\psi}_j = \text{const} = C^{ij} \psi_i \psi_j$ which contains the vector ψ . We consider then the integral

$$\bar{P}_\rho = \int_{\tilde{\rho}^2 > \rho^2} p(\tilde{\psi}) d\tilde{\psi}_1 \cdots d\tilde{\psi}_n \quad (7)$$

of the n -dimensional probability density over the region $\tilde{\rho}^2(\tilde{\psi}) > \rho^2(\psi)$ outside the surface $\rho^2(\tilde{\psi}) = \rho^2(\psi) = \text{const}$. If \bar{P}_ρ is small, 5%, say, the null hypothesis that ψ represents a realization of the natural variability ensemble is said to be rejected with a risk of \bar{P}_ρ . Conversely, a climate change signal is said to have been detected in the data at a significance level of $P_\rho = (1 - \bar{P}_\rho)$ (95%).

Reduction of the detection space

In practice, this straightforward statistical detection test can be applied successfully only if the vector dimension n of the climate state trajectory is small. Unfortunately, the situation is normally just the reverse: the discretization of a set of time series of gridded climate data will normally yield a vector ψ of very high dimension. The problem or many dimensions is that even a relatively large climate change signal ψ^s relative to the noise component in some given but unknown direction in phase space cannot be detected in the presence of noise distributed over a large number of other components. For successful detection and attribution, the dimension of the detection space must be strongly reduced – ideally to a single dimension by specifying the direction of the anticipated climate change signal, or to a small number of climate change patterns if more than one candidate forcing mechanism is considered.

The impact of the number of dimensions on the detection power can best be demonstrated by transforming to ortho-normal variables

$$\psi'_i = T_i^j \psi_j \quad (8)$$

$$\psi'^i = \hat{T}^i_j \psi^j, \quad (9)$$

where \hat{T}^i_j denotes the transposed inverse of the transformation matrix T_i^j ,

$$T_i^j \hat{T}^i_k = \delta_k^j. \quad (10)$$

In the ortho-normal system, the covariance matrix and its inverse are transformed to the unit co- and contra-variant matrices I_{ij} and I^{ij} , respectively,

$$C'_{ij} = \langle \tilde{\psi}'_i \tilde{\psi}'_j \rangle = T_i^k T_j^l C_{kl} = I_{ij} \quad (11)$$

$$C'^{ij} = \langle \tilde{\psi}'^i \tilde{\psi}'^j \rangle = \hat{T}^i_k \hat{T}^j_l C^{kl} = I^{ij}. \quad (12)$$

The optimal fingerprint

In the ortho-normal coordinate system, it is self-evident from the isotropic symmetry of the problem that if the signal lies in the direction of the first coordinate, the univariate detection test should also be carried out with respect to the first coordinate. How does this result transform to a signal ψ_i^s oriented in some given guess-pattern direction g_i in an arbitrary coordinate system? To estimate the amplitude of the signal from the observed data ψ in the general case we write

$$\psi_i = d g_i + \psi_i^r \quad (18)$$

where the coefficient d (the *detection variable*) is determined by the scalar multiplication of the observed data with a suitably defined fingerprint f^i ,

$$d = f^i \psi_i, \quad (19)$$

and ψ_i^r is a residual which we wish to minimize.

It is common practice in many applications to determine the coefficient d by minimizing the mean square error $\sum_i \langle (\psi_i^r)^2 \rangle$. However, in the present case this is not appropriate. Firstly, the mean square error is not invariant with respect to linear transformations to other variables. Secondly, our goal for the purpose of detection is to not to maximize the explained variance in a particular reference system, but rather to maximize the squared signal-to-noise ratio $d^2 / \langle \tilde{d}^2 \rangle$ for an arbitrary reference system, where $\tilde{d} = f^i \tilde{\psi}_i$ is the detection variable determined by the natural climate variability in the absence of a climate change signal. Since the signal-to-noise ratio is independent of the scaling of d , for detection applications we need to determine only the direction of the fingerprint. It was shown in H, and is shown again trivially below (see also Hasselmann, 1979), that the maximization of the signal-to-noise ratio yields the fingerprint

$$f^i = C^{ij} g_j \equiv g^i. \quad (20)$$

where the signal pattern and fingerprint can be normalized, without loss of generality, such that

$$C^{ij} g_i g_j = 1, \quad \text{or} \quad (21)$$

$$C_{ij} f^i f^j = f^i g_i = 1. \quad (22)$$

Thus the optimal fingerprint represents the contravariant counterpart of the covariant guess pattern. (We nevertheless use different symbols for the fingerprint and signal rather than distinguishing the two only by the position of the index to emphasize the basic difference in the role of the two patterns. In the detection literature this distinction is sometimes overseen.)

In the present co- and contravariant notation the result (20)-(22) follows immediately from the argument indicated above that in the special case of an ortho-normal reference system, $C'^{ij} = I^{ij} =$ unit matrix, the fingerprint and signal pattern must have the same directions for reasons of isotropic symmetry:

$$f'^i = I^{ij} g'_j = C'^{ij} g'_j, \quad (23)$$

of the guess patterns (applying the summation convention also to the indices ν of the p guess patterns), the condition that the quadratic form $\rho^{r^2} = \rho^2(\psi^r)$, cf. eq.(4), for the residual is minimized (maximizing also the multi-variate signal-to-noise ratio for the coefficient vector $\mathbf{d} = (d^\nu)$) yields as determining equations for the coefficients d^ν of the retrieved climate change signal the set of p linear equations

$$D_{\nu\mu}d^\mu = f_\nu^i\psi_i \quad (\nu = 1, \dots, p), \quad (29)$$

where

$$f_\nu^i = C^{ij}g_{\nu j} (= g_\nu^i) \quad (30)$$

denotes the fingerprint of the ν 'th guess pattern, in analogy with the definition (20) in the single pattern case, and

$$D_{\nu\mu} = f_\nu^i g_{\mu i} = C^{ij} g_{\nu i} g_{\mu j}. \quad (31)$$

The solution can be expressed in a concise form by introducing the operations of index raising and lowering also for Greek guess-pattern indices, using as metric the matrix $D_{\nu\mu}$ defined by the scalar products of the signal patterns. Introducing the covariant multi-pattern detection coefficients, given, in analogy with the definition for the scalar single-pattern detection coefficient d , eq.(19), by

$$d_\nu = f_\nu^i\psi_i, \quad (32)$$

the contravariant detection coefficients may be expressed as

$$d^\nu = D^{\nu\mu}f_\mu^i\psi_i = f^{\nu i}\psi_i, \quad (33)$$

where $D^{\nu\mu}$ denotes the inverse of $D_{\nu\mu}$,

$$D^{\nu\mu}D_{\mu\lambda} = \delta_\lambda^\nu. \quad (34)$$

It follows from eq.(33) that $D^{\nu\mu}$ represents the covariance matrix of the natural variability components \tilde{d}^ν of the contravariant detection coefficients,

$$D^{\nu\mu} = \langle \tilde{d}^\nu \tilde{d}^\mu \rangle = f^{\nu i} f^{\mu j} \langle \psi_i \psi_j \rangle = f^{\nu i} g_i^\mu, \quad (35)$$

while

$$D_{\nu\mu} = \langle \tilde{d}_\nu \tilde{d}_\mu \rangle = f_\nu^i f_\mu^j \langle \psi_i \psi_j \rangle = f_\nu^i g_{\mu i} \quad (36)$$

represents the corresponding covariance matrix of the natural variability contribution of the covariant detection coefficients.

Depending on the context, the multi-pattern detection problem is seen to lead to a detection vector which can appear either in a co- or a contravariant form with respect to the metric $D_{\nu\mu}$. We shall refer to the contravariant detection coefficients d^ν , which appear in the original representation (28) of the climate trajectory in terms of the signal patterns, as *pattern amplitudes*. The covariant detection coefficients d_ν , defined by the straightforward generalization, eq.(32), of the expression (19) for the scalar detection variable, will be termed simply the *detection variables*. The detection variables are the variables which arise naturally in the multivariate

attributed to internal natural climate variability. For the attribution problem we need to consider now further hypotheses regarding the cause of a detected climate change. We assume there exist generally several candidate mechanisms $\nu = 1, \dots, p$, each of which is characterized by a predicted climate change signal. In contrast to the detection problem, where we needed to know only the normalized directions \mathbf{g}_ν of the signal patterns, we specify now also the predicted amplitudes a^ν of the signals.

To decide whether the climate change signal $\psi_{(\nu)}^o$ inferred from observations is consistent with a given signal $\psi_{(\nu)}^m$ predicted from a model simulation, we must assign to each predicted climate change signal an error covariance matrix – in analogy with the natural variability covariance matrix required for the detection test. We assume again that the error distributions are Gaussian. The consistency of the retrieved climate change signal with the predicted signal is then tested by comparing the difference between the two signals with the differences which could be expected from the estimated signal errors. We shall be concerned only with the distinction between different signals in the space spanned by the p predicted signal patterns. Thus we need consider only the projection of the signal pattern errors in this signal pattern space.

We assume that the p predicted signal patterns are linearly independent and therefore do indeed span a p -dimensional space. However, we can allow also additional forcing mechanisms which generate climate change signals lying in this space (for example, by explicitly considering linear combinations of the p basic forcing mechanisms, such as a combined greenhouse gas and aerosol forcing, cf. Hegerl *et al.*, 1996b). If the pattern amplitudes of such linearly combined climate change signals are prescribed, the attribution (or consistency) tests can be applied in the same way to these signals as to the p base signals. Formally, one needs only to replace one of the original base signals by the linear combination selected for the consistency test (note that the signal patterns \mathbf{g}_ν are assumed to be normalized by eq.(21), but are not necessarily orthogonal).

The consistency test described in the following is carried out for each forcing mechanism separately. The outcome can be that one, none, or some sub-set of the forcings is consistent with the inferred climate change. If the observations are found to be consistent with exactly one forcing mechanism, and the null hypothesis that the retrieved climate change signal is consistent with natural climate variability is rejected, the retrieved climate change is attributed to that mechanism.

3.1 Consistency and attribution tests

Having retrieved the observed climate change signal

$$\psi^o = d^\mu \mathbf{g}_\mu, \quad (43)$$

with pattern amplitudes d^μ given by the solutions of eqs. (29), we investigate now for each proposed forcing mechanism ν whether the retrieved signal is consistent with the predicted climate change signal

$$\psi^m = a^{(\nu)} \mathbf{g}_{(\nu)} \quad (44)$$

For the consistency test we apply the same approach as in the detection test. The null hypothesis is replaced now by the consistency hypothesis, and the retrieved pattern amplitude vector by the difference amplitude vector. Apart from this change in terminology, the concepts are identical to those introduced for the detection test. For any given amplitude difference vector $\epsilon_{(\nu)}$ there exists a surface $\rho_\epsilon^2 = \text{const}$ which contains the vector. We consider then the integral

$$\bar{P}_{\rho_\epsilon} = \int_{\tilde{\rho}_\epsilon^2 > \rho_\epsilon^2} p_\epsilon(\tilde{\epsilon}_{(\nu)}) d\tilde{\epsilon}_{(\nu)}^1 \cdots d\tilde{\epsilon}_{(\nu)}^p \quad (51)$$

of the p -dimensional probability density p_ϵ over the region $\tilde{\rho}_\epsilon^2(\tilde{\epsilon}_{(\nu)}) > \rho_\epsilon^2(\epsilon_{(\nu)})$ outside the surface $\tilde{\rho}_\epsilon^2(\tilde{\epsilon}_{(\nu)}) = \rho_\epsilon^2(\epsilon_{(\nu)}) = \text{const}$.

If \bar{P}_{ρ_ϵ} is small, 5%, say, the hypothesis that the retrieved climate change signal is consistent with the forcing mechanism ν is said to be rejected with a risk of \bar{P}_{ρ_ϵ} , or at a significance level of $P_{\rho_\epsilon} = (1 - \bar{P}_{\rho_\epsilon})$ (95%).

We note that a positive outcome of the statistical detection test (i.e. the rejection of the null hypothesis) is formally analagous to a negative outcome of the consistency test (i.e. the rejection of the consistency hypothesis). A positive outcome of the consistency test should therefore be expressed formally in the double negative form that the retrieved climate change signal is not inconsistent with the proposed forcing mechanism at a given significance level P . However, if the chosen significance level P is high, 95%, say, this statement is rather weak (a high significance level is normally chosen to yield a strong statement for the converse case that the attribution test is rejected). To avoid the cumbersome double negative wording, while at the same time conveying more accurately the statistical significance of a positive outcome of a consistency test, we shall replace the statement that ‘a retrieved climate change signal is *not inconsistent* with a given forcing mechanism at a *significance level* of P (95%)’ by the simpler positive statement that ‘a climate change signal is *consistent* with the forcing mechanism within the P (95%)- *confidence region*’ (in analogy with the terminology of power spectral analysis) or ‘at a *confidence level* of \bar{P} (5%)’. Note that the stringency of the consistency test increases with decreasing P or increasing \bar{P} . For $P \rightarrow 0$, the confidence region contracts to zero, requiring zero error between the retrieved and predicted pattern amplitudes for a positive outcome of the consistency test, while the confidence level \bar{P} for a consistent signal increases to 100%. For the acceptance of a consistency test as positive, it will generally be advisable to select a consistency confidence level somewhat higher than 5%, of the order of 10% - 20%. Still higher confidence levels, however, incur the risk of erroneously rejecting valid attributions.

As outcome of the combined multi-pattern detection/attribution exercise we can then assign a statistical *significance level*, defined by eq.(42), for the detection of a climate change signal within the space spanned by the p predicted signal patterns; and a *consistency confidence level* for each proposed climate change mechanism ν , defined, in analogy with the risk associated with the null hypothesis, by eq.(51).

The result of the test will consist generally of one of the following combinations (cf. Figure 1):

1. A statistically significant climate change signal a consisting of a superposition of predicted climate change signals is detected in the observed data at a given

5. The retrieved climate change signal e is not statistically significant and the retrieved climate change signal is not consistent with any of the predicted signals.

We note that the attribution of a detected climate change signal to a particular forcing mechanism is successful only in the first of these possible outcomes.

One can consider various modifications of the test procedure outlined above. Rather than determining the retrieved climate change signal in the p -dimensional space of all proposed signal patterns, the detection and attribution test can be carried out as a single-pattern analysis separately for each individual mechanism (yielding the same set of possible test outcomes). This has the advantage of enhancing the probability of detection of any given forcing signal. However, it provides less discrimination between competing mechanisms when the signal patterns are not orthogonal. The signal pattern a of Figure 1, for example, fails the consistency test for the forcing mechanism 2 in the full signal pattern space, but would pass an individual pattern consistency test for this process (as is apparent from a visual projection of the retrieved signal vector onto the direction of the signal pattern 2). Thus in contrast to the two-pattern analysis, a unique attribution is no longer achieved in this case using individual single pattern consistency tests (see also the similar example discussed in Hegerl *et al*, 1996b)).

Another modification is suggested if one of the predicted signals is consistent with a zero amplitude with acceptable probability, and the detection/attribution test also returns a small amplitude for that signal. One can then repeat the test leaving out that forcing mechanism, in the expectation that the significance and confidence levels for the detection and attribution of the other signals are thereby enhanced.

We note, however, that in our formulation of the attribution problem we have not considered the possibility that a proposed forcing mechanism, once introduced, simply does not exist. A proposed mechanism can only be rejected as not consistent statistically with the observations, or the retrieved signal, although consistent statistically with the predicted signal, can be so small that it is nevertheless not distinguishable statistically from zero.

To establish an optimal trade-off between a high detection significance level (requiring a small number of patterns) and the ability to discriminate between different competing climate forcing mechanisms (requiring a larger number of patterns), one can apply also a series of detection/attribution tests at different levels, each successive level involving an increase in the number of patterns. A similar optimal trade-off between statistical significance and the number of predictors has been applied in the construction of a hierarchy of statistical linear prediction models from a finite data set, cf. Barnett and Hasselmann (1979).

3.2 Maximum likelihood estimate of the climate change signal

If a detected climate change signal has been successfully attributed to a particular forcing mechanism ν , one may ask whether the climate change signal retrieved from the observations is necessarily the best estimate of the climate change signal. The retrieved signal is determined by projection of the observed climate trajectory onto

climate change signal is always trivially consistent in this limit to the proposed forcing mechanism, since the retrieved climate change signal will always lie within the very large error bounds of the prediction.

The opposite limit of large $\hat{E}_{\mu\lambda}^{(\nu)}$ compared with $D_{\mu\lambda}$, i.e. very accurately determined differences between the predicted and retrieved pattern coefficients relative to the statistical errors in the retrieved optimal-detection pattern coefficients, formally yields the solution

$$d_{(\nu)}^{\mu} = \delta_{(\nu)}^{\mu} a^{(\nu)}, \quad (54)$$

i.e. the maximum likelihood signal is identical to the predicted signal. However, this limit is unaccessable, since the errors in the differences between the predicted and retrieved pattern amplitudes are always larger, according to eq.(47), than the statistical errors in the retrieved optimal-detection pattern coefficients. The largest values of $\hat{E}_{\mu\lambda}^{(\nu)}$ are obtained when the model errors $M_{(\nu)}^{\mu\lambda}$ vanish, so that eq.(47) yields $E_{(\nu)}^{\mu\lambda} = D^{\mu\lambda}$. In this case eq.(53) reduces to

$$2D_{\mu\lambda}d_{(\nu)}^{\lambda} = d_{\mu} + \hat{D}_{\mu(\nu)}a^{(\nu)}. \quad (55)$$

Multiplication from the left with $D^{\sigma\mu}$ yields the solution

$$d_{(\nu)}^{\sigma} = \frac{1}{2} \left(d^{\sigma} + \delta_{(\nu)}^{\sigma} a^{(\nu)} \right), \quad (56)$$

i.e. the maximum likelihood solution is given by the mean of the predicted and original retrieved solution.

In practice, neither limiting case will apply, and the maximum likelihood solution will lie somewhere between the original retrieved climate change signal and the limiting, maximally modified solution (56) (cf. Figure 1, signal vector ml).

4 Summary and conclusions

The general multi-pattern optimal fingerprint method for the detection of a space-time dependent climate change signal in the presence of natural climate variability can be readily extended to the problem of attribution. A co- and contra-variant tensor notation, based on a metric given by the space-time dependent covariance matrix C_{ij} of the natural climate variability, simplifies the analysis considerably. The optimal fingerprint patterns f_{ν}^i for detection are identified as the contravariant counterparts of the covariant signal patterns, $f_{\nu}^i = C^{ij}g_{\nu j} = g_{\nu}^i$. For the multi-pattern problem it is useful to introduce a second metric $D_{\nu\mu}$, defined by the scalar products $D_{\nu\mu} = g_{\nu i}g_{\mu j}C^{ij}$, in the p -dimensional space of signal patterns $g_{\nu i}$. The covariant detection variables $d_{\nu} = f_{\nu}^i\psi_i$ represent then the simplest set of coefficients for establishing the detection significance level, while the contravariant coefficients $d^{\nu} = D^{\nu\mu}d_{\mu}$, which require the inversion of the metric $D_{\nu\mu}$, define the amplitudes of the signal patterns g_{ν} estimated from the observed data. The matrices $D_{\nu\mu}$ and $D^{\nu\mu}$ represent also the covariance matrices of the natural variability of the co- and contravariant detection coefficients, respectively.

References

- [1] Barnett, T.P. and K.Hasselmann (1979) Techniques of linear prediction, with application to oceanic and atmospheric fields in the tropical Pacific, *Rev.Geophys.*, *17*, 949-968.
- [2] Bell, T.L. (1982) Optimal weighting of data to detect climate change: Application to the carbon dioxide problem, *J.Geophys.Res.* *87*, 11161-11170.
- [3] Bell, T.P. (1986) Theory of optimal weighting of data to detect climate change, *J.Atmos.Sci.*, *43*, 1694-1710.
- [4] Cubasch, U., B.D.Santer, A.Hellbach, G.Hegerl, H.Höck, E.Maier-Reimer, U.Mikolajewicz, A.Stössel and R.Voss, (1994), Monte Carlo climate change forecasts with a global coupled ocean-atmosphere model, *Climate Dynamics*, *10*, 1- 19.
- [5] Hasselmann, K. (1979) On the signal-to-noise problem in atmospheric response studies, in *Meteorology of Tropical Oceans*, edited by D.B.Shaw, pp. 251-259, Roy.Met.Soc., London.
- [6] Hasselmann, K. (1993) Optimal fingerprints for the detection of time-dependent climate change, *J.Climate*, *6*, 1957-1969
- [7] Hasselmann, K. L.Bengtsson, U.Cubasch, G.C.Hegerl, H.Rodhe, E.Roeckner, H.v.Storch, R.Voss, J.Waskewitz (1995) Detection of anthropogenic climate change using a fingerprint method, *Modern Dynamical Meteorology*, Proc. Wiin-Nielsson Symposium, ECMWF Report, 203-221.
- [8] Hegerl, G.C., H.v.Storch, K.Hasselmann, B.D.Santer, U.Cubasch, and P.D.Jones (1996a) Detecting greenhouse gas induced climate change with an optimal fingerprint method, *J.Climate* (in press)
- [9] Hegerl, G.C., K.Hasselmann, U.Cubasch, J.F.B.Mitchell, E.Roeckner, R.Voss, J.Waskewitz, On multi-fingerprint detection and attribution of greenhouse gas and aerosol forced climate change (1996b), *Climate Dynamics*, (submitted)
- [10] North, G.R., K.Y.Kim, S.P.Shen and J.W.Hardin, (1995) Detection of forced climate signals, Part I: Filter theory, *J.Climate*, *8*, 401-408
- [11] North, G.R. and K.Y.Kim (1995) Detection of forced climate signals, Part II: Simulation results, *J.Climate*, *8*, 409-417
- [12] Pennell, W.T., T.P.Barnett, K.Hasselmann, W.R.Holland, T.R.Karl, G.R.North, M.C.MacCracken, M.E.Moss, G.Pearman, E.M.Rasmusson, B.D.Santer, W.K.Smith, H.v.Storch, P.Switzer and F.Zwiers (1993) The detection of anthropogenic climate change, *Proc.4'th Symp.Global Change Studies*, Anaheim (Cal.), Jan. 17-19, 1993, Amer.Met.Soc., 21-28.
- [13] Santer, B.D., U.Mikolajewicz, W.Brüggemann, U.Cubasch, K.Hasselmann, H.Höck, E.Maier-Reimer and T.M.Wigley (1995a) Ocean variability