Coupled monoubiquitylation of the co–E3 ligase DCNL1 by Ariadne-RBR E3 ubiquitin ligases promotes cullin-RING ligase complex remodeling

Ian R. Kelsall¹, Yosua A. Kristariyanto¹, Axel Knebel¹, Nicola T. Wood¹, Yogesh Kulathu¹, Arno F. Alpi^{1,#,*}

¹ MRC Protein Phosphorylation and Ubiquitylation Unit, School of Life Sciences, University of Dundee, DD1 5EH, UK.

Running title: DCNL1 monoubiquitylation by Ariadne E3 ligases

[#] Present address: Department of Molecular Machines and Signaling, Max Planck Institute of Biochemistry, Martinsried, Germany.

To whom correspondence should be addressed: Arno F. Alpi: Department of Molecular Machines and Signaling, Max Planck Institute of Biochemistry, Martinsried, Germany; *aalpi@biochem.mpg.de;* Tel.(0049-89-85782480

Keywords: Post-translational modifications, Ubiquitin, Ubiquitylation, Ubiquitin ligases, ARIH1, ARIH2, Neddylation, DCNL1, Cell signaling, Coupled monoubiquitylation

SUPPORTING INFORMATION

Figure S1

Figure S2





Figure S1: In vitro monoubiquitylation and purification of DCNL1. (A) Flow chart summerizing the sequential purification steps of *in vitro* monoubiquitylated DCNL1 as described in Experimental procedures (left), and Coomassie stained SDS PAGE showing reaction products of the monoubiquitylation reaction, fraction after imidazole and HA-peptide elution (right). (B) Coomassie stained SDS-PAGE showing recombinant purified DCNL1 variants.



Figure S2 Identification of ubiquitlated lysin residues on DCNL1 (A) Reconstitution of DNCL1 monoubiquitylation with purified recombinant TRIAD1, neddylated CUL5–RBX2 (N8-CUL5–RBX2), and UBCH7. Products of complete and drop-out (-) reactions were separated on SDS-PAGE and detected by coomassie stain. Boxed band indicating mono-ubiquitylated DCNL1 was excised and subjected to MS analysis. (B) Table and cartoon summarizing identified Lys-diGG sites including published lysine sites.