



Draft Genome Sequence of Actinobacterial Strain *Kineosporia* sp. R_H_3, a Neutrophilic Iron-Depositing Bacterium

 Burga Braun,^a Sven Künzel,^b Josephin Schröder,^a Ulrich Szewzyk^a

^aTechnische Universität Berlin, Berlin, Germany

^bMax Planck Institute for Evolutionary Biology, Plön, Germany

ABSTRACT The draft genome sequence of a neutrophilic iron-depositing actinobacterial strain, *Kineosporia* sp. R_H_3, is reported here. Detailed analysis of the genome can elucidate the role of specific cytochromes for Fe oxidation and how this organism might receive energy from Fe oxidation. To date, this is the second publicly available genome sequence of a *Kineosporia* strain.

Kineosporia sp. strain R_H_3 is a small Gram-positive bacterium that formed very small dark-brown colonies with a diameter of less than 1 mm on agar plates containing iron and manganese. BLAST searches of a 1,504-bp sequence of the 16S RNA gene using the Nucleotide Collection Nr/Nt database (1) and EzBioCloud (2) revealed five of the seven described *Kineosporia* strains as the closest relatives, namely, strains NBRC 16234^T, JCM 9954^T, VN05A0415^T, JCM 9957^T, and YIM 65293^T, each with an identity of 96%.

Strain R_H_3 was isolated from ochrous formations of a 320-m-deep dewatering well of the open cast mining area in Hambach, Germany (Rhenish lignite-mining area). For isolation, diluted ochrous samples were spread on H₂O medium [2 g liter⁻¹ MnCO₃ hydrate, 0.2 g liter⁻¹ Fe(NH₄)₂(SO₄)₂, 2 ml vitamin solution (3), 2 ml trace element solution SL 9 (4), 15 g agar, and 1 liter deionized water] and incubated at room temperature until dark-brown colonies developed. The iron deposition ability of the strain was verified according to Braun et al. (5). Determination of the physiological characteristics using Biolog GN2 microplates (I&L Biosystems GmbH, Königswinter, Germany) showed that strain R_H_3 was able to metabolize only Tween 40 and Tween 80. pH-dependent growth rates were determined in R2A medium (DSMZ 830), and optimal growth occurred at pH 6. On agar plates, strain R_H_3 was able to grow on H₂O medium and on LSM2 (6), CY (DSMZ 67), and R2A and NA (DSMZ 1) media.

Extraction of genomic DNA was done as previously described (7), and a paired-end library was prepared according to the Illumina Nextera XT DNA library prep kit protocol. Genome sequencing was performed on an Illumina NextSeq 500 sequencer using NextSeq mid output kit v2 (300-cycle) chemistry, which generated 19,900,890 raw reads. Demultiplexing was done with bcl2fastq v2.18.0.12, and quality filtering of raw reads was performed using Trimmomatic v0.36 (8), resulting in 13,319,870 filtered reads. Subsequently, reads were checked for ambiguous base calls and low complexity employing the DUST algorithm (9) and filtered accordingly with an R script in Microsoft R Open v3.3.2 (<http://www.r-project.org/>), followed by preassembly with SPAdes v3.10.0 (10) using default k-mer lengths up to 99 bp. Scaffolds of ≥ 500 bp of this preassembly were subject to extension and second-round scaffolding with SSPACE standard v3.0 (11). Scaffolds of $\geq 2,500$ bp were assigned to genome bins by MetaBAT v0.32.4 (12), and functional annotation of the draft genomes was performed with Prokka v1.12b (13). The draft genome included 270 scaffolds with an N_{50} assembly quality of 39,507 and an L_{50} value of 53. The shortest scaffold was 2,548 bp, and the longest scaffold was 206,123 bp. The total size of the scaffolds was 7,260,683 bp, with a GC content of 74%.

Received 18 March 2018 **Accepted** 20 March 2018 **Published** 12 April 2018

Citation Braun B, Künzel S, Schröder J, Szewzyk U. 2018. Draft genome sequence of actinobacterial strain *Kineosporia* sp. R_H_3, a neutrophilic iron-depositing bacterium. *Genome Announc* 6:e00335-18. <https://doi.org/10.1128/genomeA.00335-18>.

Copyright © 2018 Braun et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Burga Braun, Burga.braun@tu-berlin.de.

Annotation resulted in 6,590 coding regions for 6,678 genes, 514 signal peptides, no clustered regularly interspaced short palindromic repeat (CRISPR) repeat units, 3 rRNAs (5S, 16S, and 23S), 68 tRNAs, 1 transfer-messenger RNA (tmRNA), and 16 miscellaneous RNAs.

Accession number(s). This whole-genome shotgun project has been deposited at DDBJ/ENA/GenBank under the accession number [MWLL00000000](https://doi.org/10.1093/genomeA.00897-16). The version described in this paper is version MWLL01000000.

ACKNOWLEDGMENTS

Bioinformatic consulting was provided by omics2view.consulting GbR, Kiel (Germany).

This work was supported by the Bundesministerium für Bildung und Forschung (BMBF) (grant O2WT1184).

REFERENCES

- Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. 1997. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 25:3389–3402. <https://doi.org/10.1093/nar/25.17.3389>.
- Yoon S, Ha S, Kwon S, Lim J, Kim Y, Seo H, Chun J. 2017. Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67: 1613–1617. <https://doi.org/10.1099/ijsem.0.001755>.
- Schlegel HG, Zaborosch C. 1992. Allgemeine mikrobiologie Thieme, Stuttgart, Germany.
- Tschech A, Pfennig N. 1984. Growth yield increase linked to caffeate reduction in *Acetobacterium woodii*. *Arch Microbiol* 137:163–167. <https://doi.org/10.1007/BF00414460>.
- Braun B, Künzel S, Schröder J, Szewzyk U. 2017. Draft genome sequence of strain R_RK_3, an iron-depositing isolate of the genus *Rhodomicrobium*, isolated from a dewatering well of an opencast mine. *Genome Announc* 5:e00864-17. <https://doi.org/10.1128/genomeA.00864-17>.
- Mulder EG, Van Veen W. 1963. Investigations on the *Sphaerotilus-Leptothrix* group. *Antonie Van Leeuwenhoek* 29:121–153. <https://doi.org/10.1007/BF02046045>.
- Braun B, Szewzyk U. 2016. Complete genome sequence of “*Candidatus Viadribacter manganicus*” isolated from a German floodplain area. *Genome Announc* 4:e00897-16. <https://doi.org/10.1128/genomeA.00897-16>.
- Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* 30:2114–2120. <https://doi.org/10.1093/bioinformatics/btu170>.
- Morgulis A, Gertz EM, Schäffer AA, Agarwala R. 2006. A fast and symmetric DUST implementation to mask low-complexity DNA sequences. *J Comput Biol* 13:1028–1040. <https://doi.org/10.1089/cmb.2006.13.1028>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham SON, Pribelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MAXA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Boetzer M, Henkel CV, Jansen HJ, Butler D, Pirovano W. 2011. Scaffolding pre-assembled contigs using SSPACE. *Bioinformatics* 27: 578–579. <https://doi.org/10.1093/bioinformatics/btq683>.
- Kang DD, Froula J, Egan R. 2014. A robust statistical framework for reconstructing genomes from metagenomic data. *bioRxiv*. <https://doi.org/10.1101/011460>.
- Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30:2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>.