

# FREE POISSON FIELDS AND THEIR AUTOMORPHISMS

Leonid Makar-Limanov<sup>1</sup> and Ualbai Umirbaev<sup>2</sup>

ABSTRACT. Let  $k$  be an arbitrary field of characteristic 0. We prove that the group of automorphisms of a free Poisson field  $P(x, y)$  in two variables  $x, y$  over  $k$  is isomorphic to the Cremona group  $\text{Cr}_2(k)$ . We also prove that the universal enveloping algebra  $P(x_1, \dots, x_n)^e$  of a free Poisson field  $P(x_1, \dots, x_n)$  is a free ideal ring and give a characterization of the Poisson dependence of two elements of  $P(x_1, \dots, x_n)$  via universal derivatives.

**Mathematics Subject Classification (2010):** Primary 17B63, 17B40; Secondary 17A36, 16W20.

**Key words:** Poisson algebras, automorphisms, universal enveloping algebras, left dependence.

## 1. INTRODUCTION

It is well known [3, 7, 8, 9] that the automorphisms of polynomial algebras and free associative algebras in two variables are tame. It is also known [23, 31] that polynomial algebras and free associative algebras in three variables in the case of characteristic zero have wild automorphisms. It was recently proved [11] that the automorphisms of free Poisson algebras in two variables are tame in characteristic zero. Note that the Nagata automorphism [16, 23] gives an example of a wild automorphism of a free Poisson algebra in three variables.

The famous Max Noether Theorem on the Cremona groups (see [15, 20]) says that the Cremona group  $\text{Cr}_2(k)$  of birational automorphisms of the projective plane  $\mathbb{P}_k^2$  over an algebraically closed field  $k$  is generated by the standard quadratic transformation and the projective transformations. The Cremona group  $\text{Cr}_2(k)$  is isomorphic to the group of automorphisms of the field of rational functions  $k(x, y)$  over  $k$ . There are some generalizations of Noether's theorem to the case in which the ground field  $k$  is not algebraically closed (see, for example [14]). Defining relations of the Cremona group  $\text{Cr}_2(k)$  are described in [4] (see also [5]).

Let  $P\langle x_1, \dots, x_n \rangle$  be a free Poisson algebra over a field  $k$  freely generated by  $x_1, \dots, x_n$  [21]. Denote by  $P(x_1, \dots, x_n)$  the field of fractions of the commutative and associative

---

<sup>1</sup>Supported by an NSA grant H98230-09-1-0008, by an NSF grant DMS-0904713, and a Fulbright fellowship awarded by the United States-Israel Educational Foundation; The Weizmann Institute of Science, Rehovot, Israel, University of Michigan, Ann Arbor, and Wayne State University, Detroit, MI 48202, USA, e-mail: [lml@math.wayne.edu](mailto:lml@math.wayne.edu)

<sup>2</sup>Supported by an NSF grant DMS-0904713 and by a grant of Kazakhstan; Eurasian National University, Astana, Kazakhstan and Wayne State University, Detroit, MI 48202, USA, e-mail: [umirbaev@math.wayne.edu](mailto:umirbaev@math.wayne.edu)

algebra  $P\langle x_1, \dots, x_n \rangle$ . The Poisson bracket  $\{\cdot, \cdot\}$  on  $P\langle x_1, \dots, x_n \rangle$  can be uniquely extended to a Poisson bracket on  $P(x_1, \dots, x_n)$ . The field  $P(x_1, \dots, x_n)$  with this Poisson bracket is called the free Poisson field in the variables  $x_1, \dots, x_n$  over  $k$  [10].

The main goal of this paper is to prove that the group of automorphisms of the free Poisson field  $P(x, y)$  over a field  $k$  of characteristic 0 is isomorphic to the Cremona group  $Cr_2(k)$ . Notice that there are several different proofs [11, 13, 10, 30] of the tameness of automorphisms of the free Poisson algebra  $P\langle x, y \rangle$  in characteristic zero. In [30] the technic of universal derivations or Fox derivatives is used. We adopt this method to the case of free fields.

For any Poisson algebra  $P$  denote by  $P^e$  its universal enveloping algebra. The main property of the universal enveloping algebra  $P^e$  is that the notion of a Poisson  $P$ -module (see, for example [17]) is equivalent to the notion of a left module over the associative algebra  $P^e$  (see, for example [6]). The universal enveloping algebras of Poisson algebras were first studied in [17]. A linear basis of the universal enveloping algebra is constructed in [18] for Poisson polynomial algebras. The structure of the universal enveloping algebras of free Poisson algebras were studied in [30].

It is well known that the universal enveloping algebras of free Lie algebras are free associative algebras. P. Cohn [2] proved that every left ideal of a free associative algebra is a free left module, i.e., a free associative algebra is a free ideal ring (also fir or FI-ring [2]). In fact, it follows from this theorem [29] that subalgebras of free Lie algebras are free [24, 32] and automorphisms of finitely generated free Lie algebras are tame [1]. Unfortunately the universal enveloping algebras of free Poisson algebras are not free ideal rings [30]. In this paper we prove that the universal enveloping algebras of free Poisson fields are free ideal rings.

Recall that any two elements of a free Poisson field are Poisson dependent if and only if they are algebraically dependent [10]. In [30] a characterization of Poisson dependence of two elements of free Poisson algebras via universal derivations is given. We extend this result to free Poisson fields.

This paper is organized as follows. In Section 2 we recall the definition of the universal enveloping algebras of Poisson algebras and construction of free Poisson algebras and free Poisson fields. In Section 3 we describe the structure of the universal enveloping algebras of free Poisson fields and prove that they are free ideal rings. Section 4 is devoted to the study of Poisson dependence of two elements. Developed technics are used to describe automorphisms of free Poisson fields in two variables.

## 2. DEFINITIONS AND NOTATIONS

Recall that a vector space  $P$  over  $k$  endowed with two bilinear operations  $x \cdot y$  (a multiplication) and  $\{x, y\}$  (a Poisson bracket) is called a *Poisson algebra* if  $P$  is a commutative associative algebra under  $x \cdot y$ ,  $P$  is a Lie algebra under  $\{x, y\}$ , and  $P$  satisfies the following identity (the Leibniz identity):

$$\{x, y \cdot z\} = \{x, y\} \cdot z + y \cdot \{x, z\}.$$

Let  $P$  be a Poisson algebra over  $k$ . A vector space  $V$  over  $k$  is called a *Poisson module* over  $P$  (or *Poisson  $P$ -module*) if there are two bilinear maps

$$P \times V \longrightarrow V, ((x, v) \mapsto x \cdot v), \quad P \times V \longrightarrow V, ((x, v) \mapsto \{x, v\}),$$

such that the relations

$$\begin{aligned} (x \cdot y) \cdot v &= x \cdot (y \cdot v), \\ \{\{x, y\}, v\} &= \{x, \{y, v\}\} - \{y, \{x, v\}\}, \\ \{x \cdot y, v\} &= y \cdot \{x, v\} + x \cdot \{y, v\}, \\ \{x, y\} \cdot v &= \{x, y \cdot v\} - y \cdot \{x, v\} \end{aligned}$$

hold for all  $x, y \in P$  and  $v \in V$ . A Poisson module  $V$  is called unitary if  $1 \cdot v = v$  for all  $v \in V$ .

The universal enveloping algebra  $P^e$  of  $P$  is an associative algebra for which the category of Poisson modules over  $P$  and the category of (left) modules over  $P^e$  are equivalent. It can be given by generators and defining relations. Let  $m_P = \{m_a | a \in P\}$  and  $h_P = \{h_a | a \in P\}$  be two copies of the vector space  $P$  endowed with two linear isomorphisms  $m : P \longrightarrow m_P$  ( $a \mapsto m_a$ ) and  $h : P \longrightarrow h_P$  ( $a \mapsto h_a$ ). Then  $P^e$  is an associative algebra over  $k$ , with an identity 1, generated by two linear spaces  $m_P$  and  $h_P$  and defined by the relations

$$\begin{aligned} m_{xy} &= m_x m_y, \\ h_{\{x, y\}} &= h_x h_y - h_y h_x, \\ h_{xy} &= m_y h_x + m_x h_y, \\ m_{\{x, y\}} &= h_x m_y - m_y h_x \end{aligned}$$

for all  $x, y \in P$  [30]. Additionally we put  $m_1 = 1$  since we consider only unitary Poisson modules.

If  $V$  is an arbitrary Poisson  $P$ -module, then  $V$  becomes a left  $P^e$ -module under the actions

$$m_x v = x \cdot v, \quad h_x v = \{x, v\},$$

for all  $x \in P$  and  $v \in V$ . Conversely, if  $V$  is a left  $P^e$ -module then the same formulae turn  $V$  to a Poisson  $P$ -module.

The first example of a Poisson  $P$ -module is  $V = P$  under the actions  $x \cdot v$  and  $\{x, v\}$  which provides us with a representation  $\rho$  of  $P^e$  by homomorphisms of  $V$ . It is clear that  $\rho(m_x)(v)$  is  $xv$  and hence we have an imbedding

$$m : P \longrightarrow P^e \quad (x \mapsto m_x).$$

Therefore we can identify  $m_x$  with  $x$ . After this identification the defining relations of  $P_e$  are

$$\begin{aligned} (1) \quad h_{\{x, y\}} &= h_x h_y - h_y h_x, \\ (2) \quad h_{xy} &= y h_x + x h_y, \\ (3) \quad \{x, y\} &= h_x y - y h_x, \end{aligned}$$

for all  $x, y \in P$ . It follows from (3) that

$$(4) \quad h_x y - y h_x = x h_y - h_y x.$$

for all  $x, y \in P$ . From (2) we obtain that  $h_1 = 0$  and if  $x \in P$  is invertible then

$$(5) \quad h_{x^{-1}} = -x^{-2} h_x.$$

A word of caution,  $\rho$  is not necessarily an exact representation. A question for which  $P$   $\rho$  is exact seems to be an interesting problem.

Let  $g$  be a Lie algebra with a linear basis  $e_1, e_2, \dots, e_k, \dots$ . The Poisson symmetric algebra  $PS(g)$  of  $g$  is the usual polynomial algebra  $k[e_1, e_2, \dots, e_k, \dots]$  endowed with the Poisson bracket defined by

$$\{e_i, e_j\} = [e_i, e_j]$$

for all  $i, j$ , where  $[x, y]$  is the multiplication in the Lie algebra  $g$ .

Denote by  $P\langle x_1, x_2, \dots, x_n \rangle$  the free Poisson algebra over  $k$  in the variables  $x_1, x_2, \dots, x_n$ . From now on let  $g = \text{Lie}\langle x_1, x_2, \dots, x_n \rangle$  be the free Lie algebra with free (Lie) generators  $x_1, x_2, \dots, x_n$ . It is well-known (see, for example [21]) that the Poisson symmetric algebra  $PS(g)$  is the free Poisson algebra  $P\langle x_1, x_2, \dots, x_n \rangle$  in the variables  $x_1, x_2, \dots, x_n$ .

Let us choose a multihomogeneous linear basis

$$x_1, x_2, \dots, x_n, [x_1, x_2], \dots, [x_1, x_n], \dots, [x_{n-1}, x_n], [[x_1, x_2], x_3], \dots$$

of a free Lie algebra  $g$  and denote the elements of this basis by

$$(6) \quad e_1, e_2, \dots, e_m, \dots$$

The algebra  $P\langle x_1, x_2, \dots, x_n \rangle$  coincides with the polynomial algebra on the elements (6). Consequently, the set of all words of the form

$$(7) \quad u = e^\alpha = e_1^{i_1} e_2^{i_2} \dots e_m^{i_m},$$

where  $0 \leq i_k, 1 \leq k \leq m$ , and  $m \geq 0$ , forms a linear basis of  $P$ .

By  $\text{deg}$  we denote the Poisson degree function on  $P\langle x_1, x_2, \dots, x_n \rangle$  as the polynomial algebra on the elements (6), i.e.  $\text{deg}(e_i)$  is the Lie degree of this element. Note that

$$\text{deg}\{f, g\} = \text{deg } f + \text{deg } g$$

if  $f$  and  $g$  are homogeneous and  $\{f, g\} \neq 0$ . By  $\text{deg}_{x_i}$  we denote the degree function on  $P$  with respect to  $x_i$ . i.e.  $\text{deg}_{x_i}(e_j)$  is the number of appearances of  $x_i$  in the Lie monomial  $e_j$ . If  $f$  is homogeneous with respect to each  $\text{deg}_{x_i}$ , where  $1 \leq i \leq n$ , then  $f$  is called multihomogeneous. The basis (7) is multihomogeneous since so is (6).

Denote by  $P(x_1, x_2, \dots, x_n)$  the field of fractions of the polynomial algebra  $k[e_1, \dots, e_n, \dots]$  in the variables (6). The Poisson bracket  $\{\cdot, \cdot\}$  on  $k[e_1, \dots, e_n, \dots] = P\langle x_1, x_2, \dots, x_n \rangle$  can be uniquely extended to a Poisson bracket on  $P(x_1, x_2, \dots, x_n)$  and

$$\left\{ \frac{a}{b}, \frac{c}{d} \right\} = \frac{\{a, c\}bd - \{a, d\}bc - \{b, c\}ad + \{b, d\}ac}{b^2 d^2}$$

for all  $a, b, c, d \in P\langle x_1, x_2, \dots, x_n \rangle$  with  $bd \neq 0$ .

The field  $P(x_1, x_2, \dots, x_n)$  with this Poisson bracket is called the *free Poisson field* over  $k$  in the variables  $x_1, x_2, \dots, x_n$  [10].

### 3. ENVELOPING ALGEBRAS OF FREE POISSON FIELDS

**Proposition 1.** *Let  $Q = P(x_1, x_2, \dots, x_n)$  be the free Poisson field over  $k$  in the variables  $x_1, x_2, \dots, x_n$  and let  $Q^e$  be its universal enveloping algebra. Then the following statements are true:*

- (i) *The subalgebra  $A$  (with identity) of  $Q^e$  generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$  is a free associative algebra freely generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ ;*
- (ii) *The left vector space  $Q^e$  over  $Q$  is isomorphic to the left vector space  $Q \otimes_k A$  over  $Q$ .*

*Proof.* Let  $M = P(x_1, x_2, \dots, x_n, y)$  be the free Poisson field in the variables  $x_1, x_2, \dots, x_n, y$ . Then  $M$  is a Poisson  $Q$ -module under the actions  $x \cdot v$  and  $\{x, v\}$  and correspondingly a  $Q^e$  module. Denote by  $N$  the submodule of  $M$  generated by  $y$ .

Consider the elements

$$(8) \quad h_{x_{i_1}} h_{x_{i_2}} \dots h_{x_{i_k}}(y) = \{x_{i_1}, \{x_{i_2}, \dots, \{x_{i_k}, y\} \dots\}\}$$

of  $N$ . They are a subset of the multihomogeneous liner basis of  $Lie\langle x_1, x_2, \dots, x_n, y \rangle$  and are linearly independent. Consequently, the elements of the form

$$(9) \quad h_{x_{i_1}} h_{x_{i_2}} \dots h_{x_{i_k}}$$

are linearly independent in  $Q^e$  and  $A$  is a free associative algebra freely generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ .

The relations (3) allow to express every element of  $Q^e$  as a linear combination  $\sum_i q_i w_i$ , where  $q_i \in Q$  and  $w_i$  are different elements of the form (9). Then  $\sum_i q_i w_i(y) \neq 0$ . Indeed,  $M$  is a polynomial ring and  $q_i$  are polynomials in variables which are algebraically independent with the variables (8) since all these variables are different elements of the liner basis of  $Lie\langle x_1, x_2, \dots, x_n, y \rangle$ . This means that (ii) is true.  $\square$

**Corollary 1.** *Every nonzero element  $u$  of the universal enveloping algebra  $Q^e$  can be uniquely written in the form*

$$(10) \quad u = \sum_{i=1}^k q_i w_i,$$

where  $0 \neq q_i \in Q$  for all  $i$  and  $w_1, w_2, \dots, w_k$  are different elements of the form (9).

Let  $u$  be an element of  $Q^e$  written in the form (10). Put  $\text{hdeg } u = \max_{i=1}^k \text{hdeg } w_i$  where  $\text{hdeg}$  is the homogeneous degree on the free algebra  $A$  and  $\text{hdeg } 0 = -\infty$ . We say that  $u$  is homogeneous with respect to  $\text{hdeg}$  if  $\text{hdeg } w_1 = \text{hdeg } w_2 = \dots = \text{hdeg } w_k$ .

Put  $h_{x_i} < h_{x_j}$  if  $i < j$ . Let  $u, v$  be two elements of the form (9). Then put  $u < v$  if  $\text{hdeg } u < \text{hdeg } v$  or  $\text{hdeg } u = \text{hdeg } v$  and  $u$  precedes  $v$  in the lexicographical order.

Let  $u$  be an element of the form (10). We may assume that  $w_1 < w_2 < \dots < w_k$ . Then  $w_k$  is called the *leading monomial* of  $u$  and  $q_k$  is called the *leading coefficient* of  $u$ . We write  $w_k = \text{ldm}(u)$  and  $q_k = \text{ldc}(u)$ . The *leading term* of  $u$  is defined to be  $\text{ldt}(u) = \text{ldc}(u)\text{ldm}(u)$ .

**Lemma 1.** *If  $u$  and  $v$  are arbitrary nonzero elements of  $Q^e$  then*

$$\text{ldc}(uv) = \text{ldc}(u)\text{ldc}(v) \quad \text{and} \quad \text{ldm}(uv) = \text{ldm}(u)\text{ldm}(v).$$

*Proof.* Note that if  $u$  and  $v$  are two elements of the form (10) then to put the product  $uv$  into the form (10) again we need to use only the relations of type (3) which imply that  $h_{x_i}$  and  $y \in P$  commute modulo terms of smaller degrees in the variables  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ . Consequently, we can put  $uv$  into the form (10) with the leading monomial  $\text{ldm}(u)\text{ldm}(v)$  and the leading coefficient  $\text{ldc}(u)\text{ldc}(v)$ .  $\square$

It follows directly from Lemma 1 that

$$\text{hdeg } uv = \text{hdeg } u + \text{hdeg } v$$

for every  $u$  and  $v$  from  $Q^e$ , i.e.,  $\text{hdeg}$  is a degree function on  $Q^e$ .

Denote by  $U_i$  the subset of all elements  $u$  of  $Q^e$  with  $\text{hdeg } u \leq i$ . Then

$$Q = U_0 \subset U_1 \subset U_2 \subset \dots \subset U_k \subset \dots,$$

is a filtration of  $Q^e$ , i.e.,  $U_i U_j \subseteq U_{i+j}$  for all  $i, j \geq 0$  and  $\cup_{i \geq 0} U_i = Q^e$ . Put

$$\text{gr } Q^e = \text{gr } U_0 \oplus \text{gr } U_1 \oplus \text{gr } U_2 \oplus \dots \oplus \text{gr } U_k \oplus \dots,$$

where  $\text{gr } U_0 = Q$  and  $\text{gr } U_i = U_i/U_{i-1}$  for all  $i \geq 1$ . Denote by  $\varphi_i : U_i \rightarrow \text{gr } U_i$  the natural projection for every  $i \geq 1$  and define

$$(11) \quad \varphi = \{\varphi_i\}_{i \geq 0} : Q^e \rightarrow \text{gr } Q^e$$

by  $\varphi(u) = \varphi_i(u)$  if  $u \in U_i \setminus U_{i-1}$  for every  $i \geq 1$  and  $\varphi(u) = u$  if  $u \in Q$ .

The multiplication on  $Q^e$  induces a multiplication on  $\text{gr } Q^e$  and the graded vector space  $\text{gr } Q^e$  becomes an algebra.

Recall that  $A$  is the free associative subalgebra of  $Q^e$  generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$  (Proposition 1). Let  $B = Q \otimes_k A$  be the tensor product of associative algebras over  $k$ . Then  $B$  is a free associative algebra over  $Q$  freely generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ .

**Proposition 2.** *The graded algebra  $\text{gr } Q^e$  is isomorphic to  $B = Q \otimes_k A$ .*

*Proof.* By (3),  $Q$  is in the center of the algebra  $\text{gr } Q^e$  and  $\text{gr } Q^e$  is generated by  $\varphi(h_{x_1}), \varphi(h_{x_2}), \dots, \varphi(h_{x_n})$  as an algebra over  $Q$ . Note that  $B = Q \otimes_k A$  is a free associative algebra over  $Q$ . Hence there is a  $Q$ -algebra homomorphism  $\psi : B \rightarrow \text{gr } Q^e$  such that  $\psi(h_{x_i}) = \varphi(h_{x_i})$  for all  $i$ .

Let  $T_s$  be the space of  $\text{hdeg}$  homogeneous elements of  $Q^e$  of degree  $s \geq 0$  and  $B_s$  be the space of homogeneous elements of degree  $s$  of  $B$ . There is an obvious isomorphism between the  $Q$ -linear spaces  $T_s$  and  $B_s$  established by Corollary 1. Note that  $U_s = U_{s-1} + T_s$ ,  $U_s/U_{s-1} \simeq T_s \simeq B_s$ , and  $\psi|_{B_s} : B_s \rightarrow \text{gr } U_s$  is an isomorphism of  $Q$ -modules. Consequently,  $\text{Ker}(\psi) = 0$  and  $\psi$  is an isomorphism of algebras.  $\square$

Therefore  $\text{gr } Q^e$  is a free associative algebra over  $Q$  and  $\text{gr } Q^e$  is a free ideal ring [2], i.e., every left (right) ideal of  $\text{gr } Q^e$  is a free left (right)  $\text{gr } Q^e$ -module of unique rank. From now on we will identify  $\text{gr } Q^e$  with  $B = Q \otimes_k A$  by means of the isomorphism from Proposition 2. Obviously, in this identification the elements  $\varphi(h_{x_1}), \dots, \varphi(h_{x_n}) \in \text{gr } Q^e$  correspond to  $h_{x_1}, \dots, h_{x_n} \in B$ . Denote by  $\text{Deg}$  a  $Q$ -degree function on  $B$  such that  $\text{Deg}(h_{x_i}) = 1$  for all  $i$ . Notice that for any  $u \in Q^e$  we have  $\text{hdeg}(u) = \text{Deg}(\varphi(u))$ .

**Theorem 1.** *Let  $Q = P(x_1, x_2, \dots, x_n)$  be the free Poisson field over  $k$  in the variables  $x_1, x_2, \dots, x_n$  and  $Q^e$  be its universal enveloping algebra. Then  $Q^e$  satisfies the weak algorithm for hdeg and is a free ideal ring.*

*Proof.* We consider only the left dependence in  $Q^e$  since the right dependence can be treated similarly. It is sufficient to prove that  $Q^e$  satisfies the weak algorithm for hdeg [2], i.e., for any finite set of left  $Q^e$ -dependent elements  $s_1, s_2, \dots, s_k$  of  $Q^e$  with  $\text{hdeg}(s_1) \leq \text{hdeg}(s_2) \leq \dots \leq \text{hdeg}(s_k)$  there exist  $i, 1 \leq i \leq k$  and  $v_1, \dots, v_{i-1} \in Q^e$  such that  $\text{hdeg}(s_i - v_1 s_1 - \dots - v_{i-1} s_{i-1}) < \text{hdeg}(s_i)$  and  $\text{hdeg}(v_j) + \text{hdeg}(s_j) \leq \text{hdeg}(s_i)$  for all  $j, 1 \leq j \leq i-1$ .

Suppose that

$$\sum_{r=1}^k u_r s_r = 0.$$

Put  $m = \max\{\text{hdeg}(u_i) + \text{hdeg}(s_i) | 1 \leq i \leq k\}$ . Let  $j_1, \dots, j_r$  be the set of indexes  $i$  with  $\text{hdeg}(u_i) + \text{hdeg}(s_i) = m$ . Then the last equality induces

$$\varphi(u_{i_1})\varphi(s_{i_1}) + \varphi(u_{i_2})\varphi(s_{i_2}) + \dots + \varphi(u_{i_r})\varphi(s_{i_r}) = 0$$

in the free associative algebra  $B$  over  $Q$ . This is a nontrivial left dependence of the homogeneous elements  $\varphi(s_{i_1}), \varphi(s_{i_2}), \dots, \varphi(s_{i_r})$  in  $B$ . Since the free associative algebra  $B$  over the field  $Q$  satisfies the weak algorithm for Deg [2] it follows that there exist  $t$  and  $v_{i_1}, \dots, v_{i_{t-1}} \in Q^e$  such that

$$\varphi(s_{i_t}) = \varphi(v_{i_1})\varphi(s_{i_1}) + \dots + \varphi(v_{i_{t-1}})\varphi(s_{i_{t-1}})$$

and  $\text{Deg}(\varphi(v_{i_j})) + \text{Deg}(\varphi(s_{i_j})) = \text{Deg}(\varphi(s_{i_t}))$  for all  $j, 1 \leq j \leq t-1$ . Then

$$\text{hdeg}(s_{i_t} - v_{i_1} s_{i_1} - \dots - v_{i_{t-1}} s_{i_{t-1}}) < \text{hdeg}(s_{i_t})$$

and  $\text{hdeg}(v_{i_j}) + \text{hdeg}(s_{i_j}) \leq \text{hdeg}(s_{i_t})$  for all  $j, 1 \leq j \leq t-1$ .  $\square$

The proof of this theorem is constructive. Hence the standard algorithms (see, for example [25, 28, 29]) give the next result.

**Corollary 2.** (i) *The left ideal membership problem for  $Q^e$  is algorithmically decidable;*  
(ii) *The left dependence of a finite system of elements of  $Q^e$  is algorithmically recognizable.*

#### 4. POISSON DEPENDENCE OF TWO ELEMENTS

Let  $P$  be an arbitrary Poisson algebra over  $k$ . Any finite set of elements  $p_1, p_2, \dots, p_k$  of  $P$  is called *Poisson dependent* if there exists a nonzero element  $f$  of the free Poisson algebra  $P\langle x_1, x_2, \dots, x_k \rangle$  such that  $f(p_1, p_2, \dots, p_k) = 0$ . Otherwise  $p_1, p_2, \dots, p_k$  are called *Poisson free* or *Poisson independent*. If  $p_1, p_2, \dots, p_k$  are Poisson free then the Poisson subalgebra of  $P$  generated by these elements is a free Poisson algebra in these variables. Similarly we can define Lie dependence and associative dependence [25, 26, 28, 29].

Denote by  $\Omega_Q$  the left ideal of  $Q^e$  generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ . By Proposition 1,

$$\Omega_Q = Q^e h_{x_1} \oplus Q^e h_{x_2} \oplus \dots \oplus Q^e h_{x_n},$$

i.e.,  $\Omega_Q$  is a free left  $Q^e$ -module. Note that

$$Q^e = Q \oplus \Omega_Q.$$

Consider

$$h : Q \rightarrow \Omega_Q$$

such that  $h(q) = h_q$  for all  $q \in Q$ . It follows from (1) and (2) that  $h$  is a derivation of the Poisson algebra  $Q$  with coefficients in the Poisson  $Q$ -module  $\Omega_Q$ . It is proved in [30] that  $h$  is the universal derivation of the Poisson algebra  $Q$  and  $\Omega_Q$  is the universal differential module of  $Q$ . It means (see, for example [28, 29]) that for any Poisson  $Q$ -module  $V$  and for any derivation  $d : Q \rightarrow V$  there exists a unique homomorphism  $\tau : \Omega_Q \rightarrow V$  of Poisson  $Q$ -modules such that  $d = \tau h$ .

The proof of the next lemma is standard [28, 29, 30].

**Lemma 2.** *Let  $f_1, f_2, \dots, f_k$  be arbitrary elements of the free Poisson field  $Q$ . If  $f_1, f_2, \dots, f_k$  are Poisson dependent then  $h_{f_1}, h_{f_2}, \dots, h_{f_k}$  are left dependent over  $Q^e$ .*

*Proof.* We repeat the proof given in [30] for a free Poisson algebra. Let  $f = f(z_1, z_2, \dots, z_k)$  be a nonzero element of  $T = P\langle z_1, z_2, \dots, z_k \rangle$  such that  $f(f_1, f_2, \dots, f_k) = 0$  with the minimal degree possible. Then  $h_f$  can be written in  $\Omega_T$  as

$$h_f = u_1 h_{z_1} + u_2 h_{z_2} + \dots + u_k h_{z_k},$$

and

$$0 = h_{f(f_1, f_2, \dots, f_k)} = u'_1 h_{f_1} + u'_2 h_{f_2} + \dots + u'_k h_{f_k},$$

where  $u'_i = u_i(f_1, f_2, \dots, f_k)$  for all  $i$ .

Since  $f \in T \setminus k$  and hence  $f$  is not in the Poisson center of  $T$  it is clear that  $h_f$  is not identically zero. Therefore we may assume that  $u_1 = u_1(z_1, z_2, \dots, z_k) \neq 0$ . If  $u'_1 \neq 0$  then the last equation gives a nontrivial dependence of  $h_{f_1}, h_{f_2}, \dots, h_{f_k}$ . Suppose that  $u'_1 = 0$ . Note that  $u_1 = t + w$ , where  $t \in T$  and  $w \in \Omega_T$ , since  $U(T) = T \oplus \Omega_T$ . Obviously,  $t(f_1, f_2, \dots, f_k) \in Q$  and it follows from (1)–(2) that  $w(f_1, f_2, \dots, f_k) \in \Omega_Q$ . Then,  $t(f_1, f_2, \dots, f_k) = 0$  and  $w(f_1, f_2, \dots, f_k) = 0$  since  $0 = u'_1 = t(f_1, f_2, \dots, f_k) + w(f_1, f_2, \dots, f_k) \in Q \oplus \Omega_Q$ . For any monomial  $M \in T$  we have  $h_M = \sum_i t_i w_i$  where  $t_i \in T$ ,  $w_i$  belong to a free associative algebra generated by  $h_{z_1}, h_{z_2}, \dots, h_{z_k}$  and  $\deg t_i + \text{hdeg } w_i = \deg M$ . Therefore  $\deg t \leq \deg f - 1$  and  $t = 0$  since the degree of  $f$  was minimal possible. Hence  $w \neq 0$  and  $w' = w(f_1, f_2, \dots, f_k)$  provides a "smaller" dependence between  $h_{f_1}, h_{f_2}, \dots, h_{f_k}$  and after a finite number of steps we get a nontrivial dependence between  $h_{f_1}, h_{f_2}, \dots, h_{f_k}$  over  $Q^e$ .  $\square$

For every  $i \geq 1$  denote by  $\partial_{e_i}$  the usual partial derivation of the field of rational functions  $Q$  in the variables (6). The next easy lemma is useful.

**Lemma 3.** (i) *For every  $q \in Q$  we have*

$$h_q = \sum_{i \geq 1} \partial_{e_i}(q) h_{e_i};$$

(ii) *For every  $l = l(x_1, x_2, \dots, x_n) \in g = \text{Lie}\langle x_1, x_2, \dots, x_n \rangle$  we have*

$$h_l = l(h_{x_1}, h_{x_2}, \dots, h_{x_n}) \in L = \text{Lie}\langle h_{x_1}, h_{x_2}, \dots, h_{x_n} \rangle.$$



*Proof.* As mentioned above, by (1) and (2), the mapping  $h$  is a derivation with respect to both operations on  $Q$ .  $\square$

**Corollary 3.** *For every  $q \in Q$  we have  $h_q \in QL$  and  $\text{hdeg}(h_q) \leq 1$  if and only if  $q \in k(x_1, x_2, \dots, x_n)$ .*

We need the next lemma.

**Lemma 4.** [25, 27, 28] *Let  $H$  be a free Lie algebra over a field  $k$  and  $U(H)$  be its universal enveloping algebra. Then arbitrary elements  $f_1, \dots, f_k$  of  $H$  are Lie dependent if and only if they are left dependent over  $U(H)$ .*

Recall that  $A$  is a free associative algebra freely generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ . Hence  $L$  is the Lie subalgebra of the Lie algebra  $A^{(-)}$ . Notice that  $B = Q \otimes_k A$  is a free associative algebra over  $Q$  and  $Q \otimes_k L$  is a free Lie algebra over  $Q$  freely generated by  $h_{x_1}, h_{x_2}, \dots, h_{x_n}$ . Furthermore  $B$  is the universal enveloping algebra of the Lie algebra  $Q \otimes_k L$  over  $Q$ .

**Theorem 2.** *Let  $Q = P(x_1, x_2, \dots, x_n)$  be the Poisson free field in the variables  $x_1, x_2, \dots, x_n$  over a field  $k$  of characteristic zero and  $f$  and  $g$  be arbitrary elements of  $Q$ . Then the following conditions are equivalent:*

- (i)  $f$  and  $g$  are Poisson dependent;
- (ii)  $h_f$  and  $h_g$  are left dependent over  $Q^e$ ;
- (iii)  $f$  and  $g$  are polynomially dependent, i.e., they are algebraically dependent in the commutative algebra  $Q$  over  $k$ ;
- (iv) there exists  $a \in Q$  such that  $f, g \in k(a)$ ;
- (v)  $\{f, g\} = 0$  in  $Q$ .

*Proof.* The implication (iii)  $\rightarrow$  (iv) follows from the generalized Lüroth Theorem [19]. The implications (iv)  $\rightarrow$  (v) and (v)  $\rightarrow$  (i) are obvious. By Lemma 2, (i) implies (ii). To finish the proof it is sufficient to show that (ii) implies (iii).

Notice that  $h_f = 0$  if and only if  $f \in k$  by Lemma 3. Suppose that  $h_f$  and  $h_g$  are both nonzero and

$$uh_f + vh_g = 0, \quad (u, v) \neq (0, 0), \quad u, v \in Q^e.$$

It follows that

$$\text{hdeg}(u) + \text{hdeg}(h_f) = \text{hdeg}(v) + \text{hdeg}(h_g)$$

by Lemma 1 and the mapping  $\varphi$  defined in (11) gives

$$\varphi(u)\varphi(h_f) + \varphi(v)\varphi(h_g) = 0, \quad (\varphi(u), \varphi(v)) \neq (0, 0)$$

in  $B = Q \otimes_k A$ . So,  $\varphi(h_f)$  and  $\varphi(h_g)$  are left dependent over  $B$ .

By Lemma 3,  $h_f, h_g \in QL$ . Consequently,  $\varphi(h_f), \varphi(h_g) \in Q \otimes_k L$ . Thus  $\varphi(h_f)$  and  $\varphi(h_g)$  are elements of the free Lie algebra  $Q \otimes_k L$  over  $Q$  which are left dependent over  $B$ . By Lemma 4,  $\varphi(h_f)$  and  $\varphi(h_g)$  are Lie dependent over  $Q$ . It is well-known (see, for example [24, 32]) that in this case  $\varphi(h_f)$  and  $\varphi(h_g)$  are linearly dependent over  $Q$ , i.e.,  $\varphi(h_f) = \lambda\varphi(h_g)$ . Consequently,  $\text{hdeg}(h_f) = \text{hdeg}(h_g) > \text{hdeg}(h_f - \lambda h_g)$ .

Suppose that  $h_f - \lambda h_g \neq 0$ . Then  $h_f - \lambda h_g$  and  $h_g$  are left  $B$ -dependent again since so are  $h_f$  and  $h_g$ . Notice that  $h_f - \lambda h_g \in QL$ . Consequently, repeating the same considerations as above we get  $\text{hdeg}(h_f - \lambda h_g) = \text{hdeg}(h_g)$ , which is a contradiction.

Therefore  $h_f = \lambda h_g$ . Then,

$$\sum_{i \geq 1} (\partial_{e_i}(f) - \lambda \partial_{e_i}(g)) h_{e_i} = 0$$

and hence

$$\partial_{e_i}(f) - \lambda \partial_{e_i}(g) = 0$$

for all  $i \geq 1$ . It is well-known (see, for example [22]) that in this case  $f$  and  $g$  are algebraically dependent.  $\square$

Notice that the equivalence of conditions (i) and (iii) is proved in [10] and the equivalence of conditions (iii) and (v) for free Poisson algebras is proved in [12]. All statements of this theorem for free Poisson algebras are established in [30].

**Theorem 3.** *Every automorphism of the free Poisson field  $P(x, y)$  in two variables  $x, y$  over a field  $k$  of characteristic 0 is an automorphism of the field of rational functions  $k(x, y)$ , i.e.,*

$$\text{Aut } P(x, y) \cong \text{Aut } k(x, y) = \text{Cr}_2(k).$$

*Proof.* Let  $\psi$  be an automorphism of  $Q = P(x, y)$ . Put  $\psi(x) = f$  and  $\psi(y) = g$ . We will show that  $f, g \in k(x, y)$ . By Corollary 3, it is sufficient to prove that  $\text{hdeg}(h_f) = \text{hdeg}(h_g) = 1$ .

Suppose that  $\text{hdeg}(h_f) + \text{hdeg}(h_g) \geq 3$ . Note that  $\Omega_Q = Q^e h_f + Q^e h_g = Q^e h_x + Q^e h_y$ . Consequently,  $Q^e h_f + Q^e h_g$  contains two elements  $h_x$  and  $h_y$  of  $h$ -degree 1. It follows that  $\varphi(h_f)$  and  $\varphi(h_g)$  are left dependent in the free associative algebra  $Q \otimes_k \text{Asso}\langle h_x, h_y \rangle$ .

As in the proof of Theorem 2,  $\text{hdeg}(h_f) = \text{hdeg}(h_g)$  and there exist  $0 \neq \lambda \in Q$  such that  $\text{hdeg}(h_f - \lambda h_g) < \text{hdeg}(h_f)$ . Put  $T = h_f - \lambda h_g$ . Then  $\Omega_Q = Q^e T + Q^e h_g$ ,  $\text{hdeg}(h_f) + \text{hdeg}(T) \geq 3$ , and  $T \in Q \otimes_k \text{Lie}\langle h_x, h_y \rangle$ . Hence  $\text{hdeg}(T) = \text{hdeg}(h_g)$ , which is impossible. Therefore  $\text{hdeg}(h_f) = \text{hdeg}(h_g) = 1$  and  $f, g \in k(x, y)$ . Similarly  $\psi^{-1}(x), \psi^{-1}(y) \in k(x, y)$  and the restriction of  $\psi$  on  $k(x, y)$  is an automorphism.

The Theorem is proved since every automorphism of  $k(x, y)$  can be uniquely extended to an automorphism of  $P(x, y)$ .  $\square$

## Acknowledgments

We are grateful to Max-Planck Institute für Mathematik for hospitality and excellent working conditions, where part of this work has been done.

## REFERENCES

- [1] Cohn, P.M., Subalgebras of free associative algebras. Proc. London Math. Soc., 56 (1964), 618–632.
- [2] Cohn, P.M., Free ideal rings and localization in general rings. New Mathematical Monographs, 3. Cambridge University Press, Cambridge, 2006.
- [3] A. J. Czerniakiewicz, Automorphisms of a free associative algebra of rank 2, I, II, Trans. Amer. Math. Soc., 160 (1971), 393–401; 171 (1972), 309–315.
- [4] Gizatullin, M.Kh., Defining relations for the Cremona group of the plane. Izv. Akad. Nauk SSSR Ser. Mat. 46 (1982), no. 5, 909–970; English translation: in Math. USSR Izvestiya 21 (1983), 211–268.
- [5] Iskovskikh, V.A., Proof of a theorem on relations in two dimensional Cremona group. Uspekhi Mat. Nauk 40 (1985), no. 5, 255–256.

- [6] Jacobson, N., Structure and Representations of Jordan Algebras. American Mathematical Society, Providence, R.I. 1968.
- [7] Jung, H.W.E., Über ganze birationale Transformationen der Ebene. *J. reine angew. Math.*, 184 (1942), 161–174.
- [8] van der Kulk, W., On polynomial rings in two variables. *Nieuw Archief voor Wiskunde*, (3)1 (1953), 33–41.
- [9] L. Makar-Limanov, The automorphisms of the free algebra with two generators, *Funktional. Anal. i Prilozhen.* 4(1970), no.3, 107-108; English translation: in *Functional Anal. Appl.* 4 (1970), 262–263.
- [10] Makar-Limanov, L., Shestakov, I., Polynomial and Poisson dependence in free Poisson algebras and free Poisson fields. *Journal of Algebra* 349 (2012), 372–379.
- [11] Makar-Limanov, L., Turusbekova, U., Umirbaev, U., Automorphisms and derivations of free Poisson algebras in two variables. *Journal of Algebra* 322 (2009), 3318–3330.
- [12] Makar-Limanov, L., Umirbaev, U., Centralizers in free Poisson algebras. *Proc. Amer. Math. Soc.* 135 (2007), no. 7, 1969–1975.
- [13] Makar-Limanov, L., Umirbaev, U., The Freiheitssatz for Poisson algebras. *J. Algebra* 328 (2011), 495–503.
- [14] Manin, Yu.I., Rational surfaces over perfect fields. II. (Russian) *Mat. Sb. (N.S.)* 72 (1967), 161–192.
- [15] Nagata, M., On rational surfaces. II. Irreducible curves of arithmetic genus 0 or 1. *Mem. Coll. Sci. Univ. Kyoto Ser. A Math.* 33 (1960/1961), 271–293.
- [16] Nagata, M., On the automorphism group of  $k[x, y]$ . *Lect. in Math.*, Kyoto Univ., Kinokuniya, Tokio, 1972.
- [17] Oh, S.-Q., Poisson enveloping algebras. *Comm. Algebra* 27 (1999), no. 5, 2181–2186.
- [18] Oh, S.-Q., Park, C.-G., Shin, Y.-Y., A Poincare-Birkhoff-Witt theorem for Poisson enveloping algebras. *Comm. Algebra* 30 (2002), no. 10, 4867–4887.
- [19] Samuel, P., Some remarks on Lüroth’s theorem. *Mem. Coll. Sci. Univ. Kyoto. Ser. A. Math.* 27, (1953), 223–224.
- [20] Shafarevich, I.R., Averbukh, B.G., Vainberg, Yu.R., Zhizhchenko, A.B., Manin, Ju.I., Moishezon, B.G., Tjurina, G.N., Tjurin, A.N., Algebraic surfaces. (Russian) *Trudy Mat. Inst. Steklov.* 75 (1965), 1-215.
- [21] Shestakov, I.P., Quantization of Poisson superalgebras and speciality of Jordan Poisson superalgebras. *Algebra and Logic*, 32(1993), no. 5, 309–317.
- [22] Shestakov, I.P., Umirbaev, U.U., Poisson brackets and two generated subalgebras of rings of polynomials. *Journal of the American Mathematical Society*, 17 (2004), 181–196.
- [23] Shestakov, I.P., Umirbaev, U.U., Tame and wild automorphisms of rings of polynomials in three variables. *Journal of the American Mathematical Society*, 17 (2004), 197–227.
- [24] Shirshov, A.I., Subalgebras of free Lie algebras. *Mat. Sbornik*, 33(75) (1953), 441–452.
- [25] Umirbaev, U.U., On the approximation of free Lie algebras with respect to entry. *Monoids, rings and algebras. Tartu: Tartuskij Universitet, Tartu Uelik. Toim., Mat.-Meh.-Alaseid Toeid*, 878 (1990), 147–152.
- [26] Umirbaev, U.U., Algorithmic problems in associative algebras. *Algebra and Logic*, 32 (1993), 244–255.
- [27] Umirbaev, U.U., Partial derivations and endomorphisms of some relatively free Lie algebras. *Sib. Math. J.* 34 (1993), No. 6, 1161–1170.
- [28] Umirbaev, U.U., On Schreier varieties of algebras. *Algebra Logic* 33 (1994), No. 3, 180–193.
- [29] Umirbaev, U.U., Universal derivations and subalgebras of free algebras. In *Proc. 3rd Internat. Conf. Algebra, Krasnoyarsk, Russia, Walter de Gruyter, Berlin, 1996*, 255–271.
- [30] Umirbaev, U.U., Universal enveloping algebras and universal derivations of Poisson algebras. *Journal of Algebra* 354 (2012), 77–94.
- [31] U. U. Umirbaev, The Anick automorphism of free associative algebras, *J. Reine Angew. Math.* 605 (2007), 165–178.
- [32] Witt, E., Die Unterringe der freien Lieschen Ringe. *Math. Z.* 64 (1956), 195–216.