

Motivic Cohomology Spectral Sequence and Steenrod Operations

Serge Yagunov

To Victoria

ABSTRACT

For a prime number p , it is shown that differentials d_n in the motivic cohomology spectral sequence with p -local coefficients vanish unless $p-1$ divides $n-1$. We obtain an explicit formula for the first non-trivial differential d_p , expressing it in terms of motivic Steenrod p -power operations and Bockstein maps. For this end, we compute the algebra of operations of weight $p-1$ with p -local coefficients. Finally, we construct examples of varieties, having non-trivial differentials d_p in their motivic spectral sequences.

1. Introduction

The motivic cohomology spectral sequence (MCSS) is an algebro-geometric analogue of the Atiyah–Hirzebruch spectral sequence in topology. Its second term consists of motivic cohomology groups and the sequence converges to algebraic K -theory.

The spectral sequence was initially constructed for fields by Bloch and Lichtenbaum. Unfortunately, their arguments contained a gap and the construction can now be found only in unpublished preprint [BL95]. Later, different constructions were built by Grayson [Gr95] and Friedlander–Suslin [FS02]. These two constructions not only globalized the MCSS to the whole category of smooth varieties, but also showed that it is supplied with multiplicative structure. The equivalence of two approaches was established in [Su03].

Voevodsky [Vo02a, Vo02b] observed that the slice-filtration of the motivic Eilenberg–Mac Lane spectrum leads (modulo some conjectures) to another model of MCSS. This approach was developed by Levine and he has also shown the equivalence of all three constructions [Le08]. These steps made it possible to extend the MCSS to the category of Voevodsky’s spaces. More historical issues can be found in Weibel’s “K-Book” [We, VI.4.4] on his web-page.

The behavior of differentials in the MCSS is quite similar to the topological case. Being taken with rational coefficients the sequence collapses at its E_2 -term (see [GSo99]). On the other hand, its structure with integer coefficients becomes too tangled, because of the interrelation of different p -prime effects involved. The purpose of the current paper is to investigate the case of $\mathbf{Z}_{(p)}$ -coefficients that allows to “distill” the p -prime effects. In this case one gets non-trivial differentials of rather high degree and that makes their computation an interesting quest.

Differentials in the Atiyah–Hirzebruch spectral sequence were computed by Buchstaber long time ago [Bu69]. In the current paper we establish the parallel result for the MCSS. Philosoph-

ically, our approach is quite similar to Buchstaber's one, but the technique is certainly rather different.

The strategy of the proof is the following. Firstly, we show, using Adams operations, that the first non-trivial differential may appear only in E_p -term (Proposition 3.1). Then, computing the motivic Steenrod algebra in the corresponding degree, it is possible to show that the differential in question is a scalar multiple of some concrete cohomological operation. Finally, to check that the scalar in question is not zero, we construct examples of varieties such that the differentials d_p in their motivic spectral sequences are non-trivial (Proposition 6.2, Example 6.3).

The significant part of our results becomes trivial in the case $p = 2$. So, this case is systematically avoided in the paper. However, we give an evidence (see Example 6.1) of non-triviality of the differential d_2 in the MCSS with $\mathbf{Z}_{(2)}$ -coefficients.

Let us, finally, mention that the scalar appearing in our result is actually a unit in the field \mathbf{Z}/p and, therefore, plays a negligible role in the spectral sequence structure. So that, our theorem gives a full control over all differentials up to d_{2p-2} . To compute the differential d_{2p-1} and other possibly non-zero differentials $d_{k(p-1)+1}$ we need a good description of secondary (and higher) cohomological operations. As far as this description is currently not available, this makes studying higher cohomological operations in motivic cohomology an interesting topic.

The computation of the p -local Steenrod algebra is based on Voevodsky's result on the structure of the motivic Steenrod algebra with finite coefficients. Originally, the statement was proven only for fields of characteristic zero, but recent work [HKØ13] extends Voevodsky's construction to fields of characteristic mutually prime to p .

1.1 Notation remarks.

We fix a prime number p and denote by $\mathbf{Z}_{(p)}$ the localization of the ring of integers at the prime ideal (p) . We also denote by \mathbf{Z}/p^∞ the p -cyclotomic group, i.e. $\varinjlim \mathbf{Z}/p^m \mathbf{Z}$. Unless it is specified, we always assume that $p > 2$.

We always assume the field k to be perfect and $(\text{Char } k, p) = 1$. Here and below by $\text{Char } k$ we denote the characteristic exponent of k . We denote by Sm/k the category of smooth separated schemes of finite type (smooth varieties) over a field k . We also denote by \mathbf{Spc} the category of pointed Nisnevich sheaves over Sm/k (pointed Voevodsky spaces) and by \mathbf{Sp} the homotopy category of \mathbf{A}^1 -spectra. The reader is referred to [Vo98, Sect. 2] for the constructions of the categories as well as for the description of a closed model category structure on \mathbf{Spc} . We denote by $H^{*,*}(-)$ the motivic cohomology [MVW06, 3.4] (cf. also Sect. 4) and by $K_*(-)$ Quillen's K -groups [Qu73, §7]. We often call the first index of motivic cohomology groups *degree* and the second index *weight*.

- $\mathbf{pt} := \text{Spec } k$;
- $X_+ := X \sqcup \mathbf{pt}$;
- \mathbf{A}^n (resp. \mathbf{P}^n) denotes affine (resp. projective) space of dimension n in Sm/k ;
- $T := \mathbf{A}^1/(\mathbf{A}^1 - \{0\})$ is the Tate object.
- We denote T -suspension functor by $T \wedge -$. The natural morphism $X \rightarrow T \wedge X$ induces an isomorphism in motivic cohomology. For consistency we call the inverse map $\tilde{H}^{*,*}(-) \xrightarrow{\cong} \tilde{H}^{*,*}(T \wedge -)$ the T -suspension isomorphism and denote in by Σ_T .
- $\sigma_T := \Sigma_T(1) \in \tilde{H}^{2,1}(T)$ is often called the Tate element.
- $H^{*,*} := \tilde{H}^{*,*}(\text{Spec } k, \mathbf{Z}/p)$ (see 4.2)

We often denote by $[X, Y, Z]$ the Bockstein homomorphism $H^{*,*}(-, Z) \rightarrow H^{**+1,*}(-, X)$ corresponding to the short exact sequence $0 \rightarrow X \rightarrow Y \rightarrow Z \rightarrow 0$ of abelian groups.

Finally, we summarize here some vanishing results, which we will use below.

Statement 1.1 *For $X \in Sm/k$, one has $H^{p,q}(X) = 0$ if:*

- (i) $p > 2q$;
- (ii) $p > q + \dim X$;
- (iii) $q < 0$;
- (iv) $q = 0$ and $p \neq 0$;

Proof. See [MVW06]: Theorem 19.3 for (i), Theorem 3.5 for (ii), Corollary 4.2 for (iii,iv). □

2. Main Result and Outline of the Proof

As was shown in [FS02], for any $X \in Sm/k$ there exists the Motivic Cohomology Spectral Sequence:

$$E_2^{i,j} = H^{i-j,-j}(X) \Rightarrow K_{-i-j}(X), \tag{2.1}$$

starting from the motivic cohomology groups $H^{*,*}(X)$ and converging to the algebraic K -groups of the variety X . The differentials in this spectral sequence are: $d_n: E_n^{i,j} \rightarrow E_n^{i+n,j-n+1}$ ($n \geq 2$).

Theorem 2.1 *Let p be an odd prime and k be a perfect field of characteristic l such that either $l = 0$, or $(l, p) = 1$. For a variety $X \in Sm/k$ the motivic cohomology spectral sequence*

$$E_2^{i,j} = H^{i-j,-j}(X, \mathbf{Z}_{(p)}) \Rightarrow K_{-i-j}(X, \mathbf{Z}_{(p)})$$

has zero differentials d_n for $p-1 \nmid n-1$. The differential d_p coincides with the bistable operation $\mathfrak{B}\alpha P^1 r$, where r denotes the coefficient reduction corresponding to the residue map $\mathbf{Z}_{(p)} \rightarrow \mathbf{Z}/p$, the operation P^1 is the first \mathbf{Z}/p motivic Steenrod power, α denotes multiplication of coefficients by an element of \mathbf{Z}/p^\times , and $\mathfrak{B} = [\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}, \mathbf{Z}/p]$ is the Bockstein map. Moreover, for any l satisfying the theorem conditions, one can find a field F of characteristic l and a variety $X \in Sm/F$ such that the differential d_p in the corresponding MCSS is non-trivial.

In the next section we prove, following the strategy of Buchstaber, the first statement of the theorem. (Let us also mention that a similar technique was also used by Merkurjev [Me10] to analyze the structure of the Brown–Gersten–Quillen spectral sequence.) Then, in Section 4 the differential d_p is interpreted as a bistable motivic cohomology operation of bidegree $(2p-1, p-1)$ i.e., as an element of the corresponding motivic Steenrod algebra, which is computed in Section 5. Finally, in Section 6 we construct examples of varieties for which differentials d_p in MCSS are non-trivial that completes the proof of the Theorem.

3. Differentials and Adams operations

The purpose of the current section is to prove the following proposition.

Proposition 3.1 *$d_n = 0$ for $p-1 \nmid n-1$.*

Proof. As was shown in [GSo99], for every integer k such that $\frac{1}{k} \in \mathbf{Z}_{(p)}$ the Adams operation ψ_k on $K_*(X, \mathbf{Z}_{(p)})$ can be represented as an operation acting on the whole motivic cohomology spectral sequence. Moreover, the action of this operation on the E_2 -term is given by the equality: $\psi_k(\alpha) = k^{-q}\alpha$ for $\alpha \in H^{*,q}(X)$. Therefore, all topological arguments proposed by Buchstaber [Bu69] work in this case as well. Since Adams operations commute with differentials, for every integer $n > 1$, we get:

$$d_n \psi_k = \psi_k d_n : H^{*,*}(X) \rightarrow H^{*+2n-1, *+n-1}(X).$$

Hence, one has: $(k^{n-1} - 1)d_n = 0$. Let us now define the number $M(i)$ as the greatest common divisor of the following sequence:

$$M(i) := \text{g.c.d.}\{k^N(k^i - 1)\}_{k>1}, \tag{3.1}$$

where $N \gg i$. One can easily verify that the numbers $M(i)$ are well-defined. The integers $M(i)$ are sometime called Kervaire–Milnor–Adams numbers, probably, after the paper [KM60]. Their values are presented in the lemma below. Obviously, $M(n-1)d_n = 0$. Since for $p-1 \nmid n-1$, we have: $p \nmid M(n-1)$, the differentials of these degrees vanish. \square

Lemma 3.2 *For a prime p and a positive integer n denote by $\nu_p(n)$ the greatest dividing p -exponent¹ of n . The sequence of Kervaire–Milnor–Adams numbers is determined as follows. For $i \geq 1$ and a prime number p , one has: $M(2i-1) = 2$ and*

$$\nu_p(M(2i)) = \begin{cases} 1 + \nu_p(4i) & \text{for } (p-1) \mid 2i \\ 0 & \text{else.} \end{cases}$$

Proof. See [Ad65]. \square

Corollary 3.3 *The motivic spectral sequence with \mathbf{Q} -coefficients degenerates at E_2 -term.*

Proof. Any differential vanishes after multiplication by an invertible number. \square

Corollary 3.4 *For $p > 2$, one has: $pd_p = 0$ in the MCSS with $\mathbf{Z}_{(p)}$ -coefficients..*

Proof. Since, by Lemma 3.2, one has: $\nu_p M(p-1) = 1$, the corollary follows. \square

Remark 3.5 *It is interesting to mention that the sequence $M(2i)$*

$$24, 240, 504, 480, 65520, \dots$$

can be identified with denominators of terms of sequences $\frac{1}{2}\zeta(1-2i)$ or $\frac{B_{2i}}{4i}$.

4. Differentials as cohomology operations

Let us give a brief explanation of the construction of motivic Eilenberg–Mac Lane spaces, following, almost literally, the exposition of [Vo03].

For a variety $X \in Sm/k$ consider the presheaf $\mathbf{Z}_{tr}(X)$ of abelian groups on the category Sm/k , which takes a variety U to the free abelian group, generated by all cycles on $X \times U$, which are finite and equidimensional over U . For an abelian group A we set $A_{tr} := A \otimes \mathbf{Z}_{tr}$ and define presheaves of abelian groups:

$$K_{n,A}^{pre} : U \mapsto A_{tr}(\mathbf{A}^n)(U)/A_{tr}(\mathbf{A}^n - \{0\})(U). \tag{4.1}$$

¹For example, for any positive integer n , one has: $n = 2^{\nu_2(n)} 3^{\nu_3(n)} 5^{\nu_5(n)} \dots$

On the Nisnevitch site $(Sm/k)_{\text{Nis}}$ one can sheafify $K_{n,A}^{pre}$. Applying to the resulting sheaves the functor which forgets the abelian group structure, one obtains the family of pointed sheaves of sets $K_n(A)$ that play the role of Eilenberg–Mac Lane spaces in the category **Spc**.

Alternatively, one can start from the presheaf $K_{n,\mathbf{Z}}^{pre}$ and obtain a complex $\mathbf{Z}(n)$ of sheaves of abelian groups on $(Sm/k)_{\text{Nis}}$ (see the construction in [VSF00, Ch. 5]). For any $i, j \in \mathbf{Z}$, a smooth scheme X , and an abelian group A one defines motivic cohomology groups as hypercohomology groups $H^{i,j}(X, A) := \mathbf{H}^i(X_{\text{Nis}}, A(j))$, where $A(j) = A \otimes \mathbf{Z}(j)$. Let $K(i, j, A)$ be a simplicial abelian group sheaf corresponding to the complex $A(j)[i]$. Applying again the forgetful functor one gets the simplicial sheaf of sets that determines an object (also denoted by $K(i, j, A)$) of the motivic homotopy category of spaces $\mathcal{H}o_{\mathbf{A}^1}$. The sheaves $K(i, j, A)$ are \mathbf{A}^1 -local [De09, sect. 2.2–2.4] and for any smooth scheme X , one has: $H^{i,j}(X, A) = \text{Hom}_{\mathcal{H}o_{\mathbf{A}^1}}(X_+, K(i, j, A))$. For any pointed simplicial sheaf F_\bullet on $(Sm/k)_{\text{Nis}}$ one can take the following definition of reduced motivic cohomology:

$$\tilde{H}^{i,j}(F_\bullet, A) = \text{Hom}_{\mathcal{H}o_{\mathbf{A}^1}}(F_\bullet, K(i, j, A)). \quad (4.2)$$

It is shown in *loc. cit.* that there exists a weak equivalence between $K_n(A)$ and $K(2n, n, A)$, so the two constructions of Eilenberg–Mac Lane spaces agree. This extends the definition of motivic cohomology groups to the whole category of spaces.

We shall also need a notion of a cohomological operation.

Definition 4.1 *A collection $\{\varphi\}_{p,q}$ of natural transformations of functors on **Spc***

$$\varphi_{p,q}: \tilde{H}^{p,q}(-, A) \rightarrow \tilde{H}^{p+i, q+j}(-, B),$$

where A and B are abelian groups and the index (p, q) runs through $\mathbf{Z} \times \mathbf{Z}$ is called an (unstable) cohomological A - B -operation of degree i and weight j .

Let us recall that in the category **Spc** there are two circles and hence two different suspension functors. Among all the cohomological operations there are special ones that commute with both suspension isomorphisms. These operations are called *bistable*, and Voevodsky showed, using a simple trick [Vo03, Prop 2.6], that there exists a bijection between bistable operations and operations that *a priori* commute only with the T -suspension. (Recall that here T is the Tate object.) We will call operations of the latter type *stable*.

Notation. *We denote the set of all stable cohomological A - B -operations of degree i and weight j by $\mathcal{OP}^{i,j}(A, B)$. We always implicitly assume that all considered operations have nonnegative degree and weight. Since, by [Vo03, Cor. 2.10], stable operations are additive, this set has a natural structure of an abelian group, induced by addition in cohomology.*

If A (resp. B) has a ring structure, the set $\mathcal{OP}^{*,*}(A, B)$ also has a natural structure of a bigraded left (resp. right) $H^{*,*}$ -module.

It is reasonable to expect that natural transformations of motivic cohomology functors can be classified by cohomology groups of motivic Eilenberg–Mac Lane spaces.

For every Eilenberg–Mac Lane space $K_n(A)$ one can choose a universal element

$$\iota_n \in \tilde{H}^{2n, n}(K_n(A)),$$

corresponding to the identity morphism of the space $K(2n, n, A)$. Applying the T -suspension isomorphism map $\Sigma_T: \tilde{H}^{*,*}(-) \rightarrow \tilde{H}^{*+2, *+1}(T \wedge -)$ to the element ι_n , one obtains the element $\Sigma_T \iota_n \in \tilde{H}^{2n+2, n+1}(T \wedge K_n(A))$, corresponding to some homotopy class $\alpha_n \in [T \wedge K_n(A), K_{n+1}(A)]$.

This class coincides with the homotopy class of the n -th structure morphism of the Eilenberg–Mac Lane spectrum $\mathbf{H}(A)$.

Finally, using the collection of classes $\{\alpha_\bullet\}$, one can construct an inverse system of the groups $\tilde{H}^{i+2n,j+n}(K_n(A), B)$ as shown in the diagram below.

$$\begin{array}{ccc}
 \vdots & & \\
 \tilde{H}^{i+2n+2,j+n+1}(K_{n+1}(A), B) & \xrightarrow{\alpha_n^*} & \tilde{H}^{i+2n+2,j+n+1}(T \wedge K_n(A), B) \\
 \downarrow & \nearrow \Sigma_T & \\
 \tilde{H}^{i+2n,j+n}(K_n(A), B) & & \\
 \vdots & &
 \end{array} \tag{4.3}$$

A natural modification of [Vo03, Prop. 2.7.] shows that

$$\mathcal{O}\mathcal{P}^{i,j}(A, B) = \lim_{\leftarrow n} \tilde{H}^{i+2n,j+n}(K_n(A), B). \tag{4.4}$$

We will see that the module $\mathcal{O}\mathcal{P}^{*,*}(\mathbf{Z}/p, \mathbf{Z}/p)$ is naturally isomorphic to the motivic Steenrod algebra by Voevodsky (see the discussion on page 8).

Proposition 4.2 *Consider the motivic cohomology spectral sequence $(E_*^{*,*}, d_*)$. Let us fix an integer $n > 1$ and assume that for every $1 < i < n$ and any variety $X \in Sm/k$ the differentials $d_i: H^{*,*}(X) \rightarrow H^{*+2i-1,*+i-1}(X)$ are trivial. Then, the differential d_n can be identified with a stable cohomological operation of bidegree $(2n-1, n-1)$ up to multiplication by ± 1 .*

Proof. Since all the previous differentials vanish, the differential d_n actually acts on the E_2 -term of the spectral sequence. To prove the stability, one has to check the commutativity of the following diagram:

$$\begin{array}{ccc}
 H^{i,j}(X) & \xrightarrow{d_n} & H^{i+2n-1,j+n-1}(X) \\
 \Sigma_T \downarrow & \pm 1 & \downarrow \Sigma_T \\
 \tilde{H}^{i+2,j+1}(T \wedge X_+) & \xrightarrow{d_n} & \tilde{H}^{i+2n+1,j+n}(T \wedge X_+).
 \end{array} \tag{4.5}$$

Though the space $T \wedge X_+$ does not belong to Sm/k , its cohomology is a direct summand of the cohomology of the scheme $\mathbf{P}^1 \times X$ due to the existence of the retraction $\text{Spec}(k) \rightarrow \mathbf{P}^1 \leftarrow T$. Actually, the space $T \wedge X_+$ happens to be \mathbf{A}^1 -homotopically equivalent to $(\mathbf{P}^1, \infty) \wedge X$ that allows us to apply differentials to its cohomology groups.

The motivic cohomology groups of $T \wedge X_+$ are the $(2, 1)$ -shifted cohomology groups of X and the isomorphism Σ_T is delivered by multiplication with the image of the Tate element σ_T . The MCSS is functorial and has a canonical multiplicative structure that is compatible with multiplication in motivic cohomology (see [FS02, § 14]). Hence, its differentials satisfy the Leibnitz rule and one has: $d_n(\sigma_T \wedge x) = d_n(\sigma_T) \wedge x \pm \sigma_T \wedge d_n(x)$. Now, to prove the commutativity of 4.5 up to the sign, it suffices to verify that $d_n(\sigma_T) = 0$. This element should lie in the cohomology group of the variety \mathbf{P}^1 of bidegree $(2n+1, n)$ that vanishes, since $2n+1 > 2n$ (see 1.1.i). So, the commutativity result follows for dimension reasons.

In order to complete the proposition proof we only need to extend the differential to the whole category of spaces. It can be done, using Levine’s [Le08] identification between MCSS and the spectral sequence built by the slice-filtration. Due to the functoriality of the spectral sequence construction, the differential d_n becomes a motivic cohomological operation of bidegree

$(2n-1, n-1)$. It is not hard to show that the arguments above are also applicable to the category \mathbf{Spc} and prove the stability of the operation. \square

5. Some calculations in Steenrod modules

In this section we are going to perform some computations with cohomology of motivic Eilenberg–Mac Lane spaces and spectra, and we need some preliminary results and notation. We denote by $\mathfrak{K}_n: \mathcal{A}b \rightarrow \mathbf{Spc}$ (resp. $\mathfrak{K}: \mathcal{A}b \rightarrow \mathbf{Sp}$) the functor sending an abelian group A to the Eilenberg–Mac Lane space $K_n(A)$ (resp. Eilenberg–Mac Lane spectrum $\mathbf{H}(A)$).

Proposition 5.1 *For every $n > 0$ the functor \mathfrak{K}_n preserves*

- (i) *limits;*
- (ii) *filtered colimits.*

Proof. The functor \mathfrak{K}_n can be considered as the following chain of functors:

$$\mathcal{A}b \rightarrow (\text{Presheaves of } \mathcal{A}b) \rightarrow (\text{Presheaves of } \mathbf{Sets}) \rightarrow (\text{Nisnevich Sheaves}).$$

Since the groups $\mathbf{Z}_{tr}(X)(U)$ are free abelian groups, one can easily check that the first functor preserves limits and filtered colimits.

Limits and colimits of presheaves are computed objectwise. The forgetful functor $\mathcal{A}b \rightarrow \mathbf{Sets}$ preserves limits, because it has a left adjoint functor sending every set X to the free abelian group $\mathbf{Z}[X]$ and also preserves filtered colimits (see, for example, [Ar62, Sect 1.1]).

Finally, it is well known that the sheafification functor preserves arbitrary limits and colimits. \square

For a field k we call an abelian group k -admissible if it has a $\mathbf{Z}[\frac{1}{l}]$ -module structure for $l = \text{Char } k$.

Proposition 5.2 *Let k be a perfect field. Then, the functor \mathfrak{K} sends every short exact sequence of k -admissible groups to a distinguished triangle in the category \mathbf{Sp} .*

Proof. The following result was established by Østvær and Röndigs [RØ08, Thm. 1] for fields of characteristic zero and by Hoyois, Kelly, and Østvær [HKØ13, Thm. 5.8] for perfect fields of positive characteristic. Let k be a perfect field and R a ring such that $\text{Char } k$ is invertible in R . Then, Voevodsky’s big category of motives $\text{DM}(Sm/k, R)$ is equivalent to the homotopy category $\mathbf{H}(R)\text{-mod}$ of modules over the Eilenberg–Mac Lane spectrum $\mathbf{H}(R)$. The equivalence preserves the monoidal and triangulated structures.

Now, it is not hard to check that the short exact sequence of k -admissible abelian groups leads to a distinguished triangle of motives in $\text{DM}(Sm/k, R)$. Since the category of $\mathbf{H}(R)$ -modules is a triangulated subcategory of \mathbf{Sp} , this proves the proposition. \square

Remark 5.3 *For the short exact sequence $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ of abelian groups the morphism $\mathfrak{K}(C) \rightarrow \mathfrak{K}(A)[1]$ in the corresponding distinguished triangle of T -spectra induces Bockstein map $\beta = [A, B, C]$ in motivic cohomology. In particular, this implies functoriality of Bockstein maps with respect to morphisms of short exact sequences.*

All the relations below, involving Bockstein maps are obvious consequences of this remark.

Statement 5.4 *Let k be a perfect field of characteristic exponent mutually prime to p . Then, the groups $H^{*,*}(\mathbf{H}(\mathbf{Z}/p), \mathbf{Z}/p)$ and $\mathcal{O}\mathcal{P}^{*,*}(\mathbf{Z}/p, \mathbf{Z}/p)$ are naturally isomorphic.*

Proof. It is exactly the statement of [HKØ13, Thm. 3.2] and [Vo10, Cor. 2.71] that the \lim^1 groups in the short exact sequences:

$$0 \rightarrow \lim_{\leftarrow n} {}^1\tilde{H}^{i+2n-1, j+n}(K_n(\mathbf{Z}/p), \mathbf{Z}/p) \rightarrow H^{i, j}(\mathbf{H}(\mathbf{Z}/p), \mathbf{Z}/p) \rightarrow \lim_{\leftarrow n} \tilde{H}^{i+2n, j+n}(K_n(\mathbf{Z}/p), \mathbf{Z}/p) \rightarrow 0 \quad (5.1)$$

vanish (conf. also [HKØ13, Cor. 3.3]). We can identify the right-hand term with the group $\mathcal{O}\mathcal{P}^{i, j}(\mathbf{Z}/p, \mathbf{Z}/p)$ using construction (4.4). \square

Our current aim is to compute the module of stable operations from cohomology with \mathbf{Z}/p^∞ -coefficients. We start with Voevodsky's computation of the motivic Steenrod algebra.

The module $\mathcal{O}\mathcal{P}^{*,*}(\mathbf{Z}/p, \mathbf{Z}/p)$ has a natural bigraded algebra structure given by composites of operations. Consider its bigraded subalgebra, generated by Steenrod power operations P^i (of bidegrees $(2i(p-1), i(p-1))$) for $i > 0$, Bockstein homomorphism $\beta = [\mathbf{Z}/p, \mathbf{Z}/p^2, \mathbf{Z}/p]$ (of bidegree $(1, 0)$) and operations of the form $x \mapsto ax$ for $a \in H^{*,*}$. This subalgebra is called *motivic Steenrod algebra* $\mathcal{A}^{*,*}(k, \mathbf{Z}/p)$ in [Vo03, Sect. 11, Lemma 9.5].

Let us also consider sequences $I = (\varepsilon_0, s_1, \varepsilon_1, s_2, \dots, s_k, \varepsilon_k)$ of non-negative integers and such that one has: $\varepsilon_i \in \{0, 1\}$ and $s_i \geq ps_{i+1} + \varepsilon_i$ for every index i . These sequences are called *admissible*. To every admissible sequence I one can correspond the operation $P^I = \beta^{\varepsilon_0} P^{s_1} \beta^{\varepsilon_1} \dots P^{s_k} \beta^{\varepsilon_k}$. (Here we assume $\beta^0 = P^0 = \text{id}$.) These operations are called *admissible monomials*. There is a natural graded module map from the free graded left $H^{*,*}$ -module generated by all admissible monomials to $\mathcal{A}^{*,*}(k, \mathbf{Z}/p)$.

It is proven in *loc.cit* Lemma 11.1 that the latter homomorphism of $H^{*,*}$ -modules is an epimorphism and in *loc.cit* Corollary 11.5 that the admissible monomials are linearly independent with respect to the left $H^{*,*}$ -module structure.

Moreover, Voevodsky showed [Vo10, Thm. 3.49] that over a field k of characteristic 0 there is a natural isomorphism of graded left $H^{*,*}$ -modules between $\mathcal{O}\mathcal{P}^{*,*}(\mathbf{Z}/p, \mathbf{Z}/p)$ and $\mathcal{A}^{*,*}(k, \mathbf{Z}/p)$.

In the sequel we are mostly dealing with the operations of weight $p-1$ and degree $> p$, so we will often omit the second (weight) index in the notation for operation and cohomology groups and implicitly assume that the first (degree) index is greater than p .

Up to the end of this section we will omit, for brevity, mentioning \mathbf{Z}/p -coefficients and write $\mathcal{O}\mathcal{P}^*(-)$ for $\mathcal{O}\mathcal{P}^{*, p-1}(-, \mathbf{Z}/p)$. We will also write $H^*(A, B)$ for $H^{*, p-1}(\mathbf{H}(A), B)$.

The arguments above immediately imply that $\mathcal{O}\mathcal{P}^*(\mathbf{Z}/p)$ (we assume that $* > p$) is a free \mathbf{Z}/p -module with the set of generators $\{P^1, \beta P^1, P^1\beta, \beta P^1\beta\}$.

Remark 5.5 *Voevodsky's Theorems [Vo10, Thm. 3.49, Prop. 3.55] mentioned in the previous discussion were originally proven only for base fields of characteristic 0. However, recently Hoyois, Kelly, and Østrvær [HKØ13] could eliminate this annoying restriction and extend the result to the case of a perfect field k such that $(\text{Char } k, p) = 1$.*

Remark 5.6 *Using Voevodsky's computation of the motivic Steenrod algebra, it is possible to extend the results of this section to operations of weights $\leq p^2 - p$, provided that $p > 3$. We leave all the details to the reader.*

Now we will explicitly compute weight $p-1$ cohomology groups of the T -spectra $\mathbf{H}(\mathbf{Z}/p^m)$ with integral and finite coefficients.

Proposition 5.7 *For $m > 0$ there are natural isomorphisms $H^*(\mathbf{Z}/p^m) \cong \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^m)$ and $H^*(\mathbf{Z}/p^m, \mathbf{Z}) \cong \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^m, \mathbf{Z})$. The groups $H^*(\mathbf{Z}/p^m)$ and $H^*(\mathbf{Z}/p^m, \mathbf{Z})$ are the free graded \mathbf{Z}/p -modules with the following generators in the corresponding degrees:*

| | $2p - 2$ | $2p - 1$ | $2p$ |
|-----------------------------------|-------------|--------------------------------|----------------------------------|
| $H^*(\mathbf{Z}/p^m)$ | $P^1 r_m$ | $P^1 \beta_m, \beta_1 P^1 r_m$ | $\beta_1 P^1 \beta_m$ |
| $H^*(\mathbf{Z}/p^m, \mathbf{Z})$ | \emptyset | $\beta_{\mathbf{Z}} P^1 r_m$ | $\beta_{\mathbf{Z}} P^1 \beta_m$ |

Here r_m is induced by the coefficient reduction $\mathbf{Z}/p^m \rightarrow \mathbf{Z}/p$, $\beta_m = [\mathbf{Z}/p, \mathbf{Z}/p^{m+1}, \mathbf{Z}/p^m]$, and $\beta_{\mathbf{Z}} = [\mathbf{Z}, \mathbf{Z}, \mathbf{Z}/p]$.

Proof. We start with the case of \mathbf{Z}/p -coefficients. Setting $m = 1$, we get just Voevodsky's result cited above. Since in this case the higher inverse limits vanish, cohomology of spectra coincide with groups of operations. We now assume that $p > 3$. The case $p = 3$, which is similar, but requires a bit more calculations, is left to the reader. Consider the short exact sequence:

$$0 \rightarrow \mathbf{Z}/p \rightarrow \mathbf{Z}/p^{m+1} \xrightarrow{r} \mathbf{Z}/p^m \rightarrow 0$$

and assume the groups $H^*(\mathbf{Z}/p^m)$ satisfy the theorem conclusions. By Theorem 5.2, one has a distinguished triangle of spectra:

$$\mathbf{H}(\mathbf{Z}/p) \rightarrow \mathbf{H}(\mathbf{Z}/p^{m+1}) \rightarrow \mathbf{H}(\mathbf{Z}/p^m). \quad (5.2)$$

Consider the following fragment of the corresponding cohomology long exact sequence:

$$\begin{array}{ccccccc}
 0 & \longrightarrow & H^{2p-2}(\mathbf{Z}/p^m) & \xrightarrow{r^*} & H^{2p-2}(\mathbf{Z}/p^{m+1}) & \longrightarrow & H^{2p-2}(\mathbf{Z}/p) \\
 & & & & \downarrow \beta_m^* & & \\
 & & H^{2p-1}(\mathbf{Z}/p^m) & \xrightarrow{r^*} & H^{2p-1}(\mathbf{Z}/p^{m+1}) & \longrightarrow & H^{2p-1}(\mathbf{Z}/p) \\
 & & & & \downarrow & & \\
 & & H^{2p}(\mathbf{Z}/p^m) & \longrightarrow & H^{2p}(\mathbf{Z}/p^{m+1}) & \longrightarrow & H^{2p}(\mathbf{Z}/p) \longrightarrow 0
 \end{array}$$

Since the map β_m^* delivers an isomorphism between the group $H^{2p-2}(\mathbf{Z}/p)$ and the direct summand of $H^{2p-1}(\mathbf{Z}/p^m)$ generated by the operation $P^1 \beta_m$, one gets the isomorphism $H^{2p-2}(\mathbf{Z}/p^m) \cong H^{2p-2}(\mathbf{Z}/p^{m+1})$, which sends the generator $P^1 r_m$ to $r^*(P^1 r_m) = P^1 r_m r = P^1 r_{m+1}$.

In the same way, one can see that the direct summand of $H^{2p-1}(\mathbf{Z}/p^m)$ generated by the operation $\beta_1 P^1 r_m$ maps onto the direct summand of $H^{2p-1}(\mathbf{Z}/p^{m+1})$ with the generator $r^*(\beta_1 P^1 r_m) = \beta_1 P^1 r_m r = \beta_1 P^1 r_{m+1}$. The map $\bar{\beta}_{m+1} = [\mathbf{Z}/p^{m+1}, \mathbf{Z}/p^{m+2}, \mathbf{Z}/p]$ sends the group generated by $\beta_1 P^1 r_{m+1}$ to the group $H^{2p}(\mathbf{Z}/p)$ in such a way that the composite $\bar{\beta}_{m+1}^* r^* = \bar{\beta}_m^*$ makes an isomorphism between the direct summand of $H^{2p-1}(\mathbf{Z}/p^m)$ generated by the operation $\beta_1 P^1 r_m$ and the group $H^{2p}(\mathbf{Z}/p)$. Hence, the group $H^{2p-1}(\mathbf{Z}/p^{m+1})$ splits into two direct \mathbf{Z}/p -summands.

The rest of the exact sequence can be treated in a similar way. One can also immediately check that $H^i(\mathbf{Z}/p^{m+1}) = 0$ for $i > 2p$ and $p < i < 2p - 2$.

It is also easy to show that the natural epimorphism $H^*(\mathbf{Z}/p^{m+1}) \twoheadrightarrow \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^{m+1})$ is a monomorphism. For example, the image of $P^1 r_{m+1}$ is a non-zero element in the group $\mathcal{O}\mathcal{P}^{2p-2}(\mathbf{Z}/p^{m+1})$, since we know that the operation $(P^1 r_{m+1})\bar{\beta}_{m+1} = P^1(r_{m+1}\bar{\beta}_{m+1}) = P^1 \beta_1$ is non-trivial in $\mathcal{O}\mathcal{P}^{2p-1}(\mathbf{Z}/p)$.

As a result we conclude that $H^*(\mathbf{Z}/p^{m+1}) \cong \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^{m+1})$ and

$$\lim_{\leftarrow n} {}^1\tilde{H}^{*+2n-1, *+n}(K_n(\mathbf{Z}/p^{m+1}), \mathbf{Z}/p) = 0.$$

The case of finite coefficients now follows by induction.

In order to proceed with the case of integral operations, we need the following simple lemma.

Lemma 5.8 *Let*

$$A \xrightarrow{\varphi} B \xrightarrow{\chi} B \xrightarrow{\psi} C$$

be an exact sequence of groups such that the composite $\psi\varphi: A \rightarrow C$ is an isomorphism. Then, $B \cong A \oplus Q$, for such a group Q that the restricted map $\chi: Q \rightarrow Q$ is an automorphism.

Proof. The map $(\psi\varphi)^{-1}\psi$ (resp. $\varphi(\psi\varphi)^{-1}$) splits the exact sequence on the left (resp. right). Therefore, the group A is a direct summand of B . Denoting $B/\varphi A$ by Q , one can easily see that the four-term exact sequence splits into isomorphisms $A \xrightarrow{\varphi} B/Q$, $Q \xrightarrow{\chi} Q$, and $B/Q \xrightarrow{\psi} C$. \square

End of the proof of Proposition 5.7. Assuming now, that the theorem holds for the groups $H^*(\mathbf{Z}/p^m)$ we derive the integral case. Consider the fragment of the coefficient long exact sequence:

$$H^{2p-2}(\mathbf{Z}/p^m) \xrightarrow{\beta_{\mathbf{Z}}} H^{2p-1}(\mathbf{Z}/p^m, \mathbf{Z}) \xrightarrow{p} H^{2p-1}(\mathbf{Z}/p^m, \mathbf{Z}) \xrightarrow{r} C,$$

where $C = \text{Ker } \beta_{\mathbf{Z}}$ is the direct summand of $H^{2p-1}(\mathbf{Z}/p^m)$ generated by the element $\beta_1 P^1 r_m$ and $\beta_{\mathbf{Z}} = [\mathbf{Z}, \mathbf{Z}, \mathbf{Z}/p]$. As one can easily verify, the map $\beta_1 = r\beta_{\mathbf{Z}}$ provides an isomorphism between $H^{2p-2}(\mathbf{Z}/p^m)$ and C . Therefore, the conditions of the above lemma are satisfied. Let us also mention that all the groups $H^*(\mathbf{Z}/p^m, \mathbf{Z})$ are p -groups. Together with the lemma above, this gives us an isomorphism: $H^{2p-2}(\mathbf{Z}/p^m) \xrightarrow{\beta_{\mathbf{Z}}} H^{2p-1}(\mathbf{Z}/p^m, \mathbf{Z})$. The case of degree $2p$ can be verified in the same way. Similarly, we can also check that $H^*(\mathbf{Z}/p^m, \mathbf{Z}) = 0$ for $p < * < 2p - 1$ and $* > 2p$. \square

Thus, we reproved one classical Cartan's result [Ca54] in the motivic context.

The group inclusions $i_m: \mathbf{Z}/p^m \hookrightarrow \mathbf{Z}/p^{m+1}$ induce morphisms of spectra: $\mathbf{H}(\mathbf{Z}/p^m) \rightarrow \mathbf{H}(\mathbf{Z}/p^{m+1})$. Passing to cohomology, one obtains the inverse system of groups (with arbitrary coefficients):

$$H^*(\mathbf{Z}/p) \xleftarrow{i_1^*} H^*(\mathbf{Z}/p^2) \xleftarrow{i_2^*} \dots$$

Corollary 5.9

$$\lim_{\leftarrow m} H^l(\mathbf{Z}/p^m, \mathbf{Z}/p) = \begin{cases} \mathbf{Z}/p & \text{for } l = 2p - 1, 2p \\ 0 & \text{otherwise.} \end{cases}$$

and $\lim_{\leftarrow m} H^l(\mathbf{Z}/p^m, \mathbf{Z}) = \mathbf{Z}/p$ for $l = 2p$ and 0 otherwise.

Proof. We consider the case of the \mathbf{Z}/p -coefficients. The integral case is similar and left to the reader. Applying the map $i_m^*: H^*(\mathbf{Z}/p^{m+1}) \rightarrow H^*(\mathbf{Z}/p^m)$ to the generators, one has:

$$i_m^*(P^1 r_{m+1}) = 0, \quad i_m^*(P^1 \beta_{m+1}) = P^1 \beta_m, \quad i_m^*(\beta_1 P^1 r_{m+1}) = 0, \quad \text{and} \quad i_m^*(\beta_1 P^1 \beta_{m+1}) = \beta_1 P^1 \beta_m.$$

Therefore, $\text{Im}(i_m^*) \subseteq H^*(\mathbf{Z}/p^m)\beta_m$. Hence, only the elements of the form

$$\{X\beta_1 \leftarrow X\beta_2 \leftarrow \dots\}$$

“survive” in the projective limit. The corollary follows immediately. \square

To complete the computation of p -cyclotomic operations we need a lemma.

Lemma 5.10 *Let $X_1 \xrightarrow{\varphi_1} X_2 \xrightarrow{\varphi_2} \dots$ be a sequence of abelian groups. Then, for an abelian group W , one has:*

$$\lim_{\leftarrow i} \mathcal{O}\mathcal{P}^{*,*}(X_i, W) \cong \mathcal{O}\mathcal{P}^{*,*}(\lim_{\rightarrow i} X_i, W).$$

Proof. The system $\{X_i, \varphi_i\}$ induces the projective system of groups:

$$\mathcal{O}\mathcal{P}^{*,*}(X_1, W) \xleftarrow{\varphi_1^\#} \mathcal{O}\mathcal{P}^{*,*}(X_2, W) \xleftarrow{\varphi_2^\#} \dots$$

Let $\alpha \in \lim_{\leftarrow} \mathcal{O}\mathcal{P}^{*,*}(X_i, W)$. In other words, one has a system of operations $\{\alpha_i \in \mathcal{O}\mathcal{P}^{*,*}(X_i, W)\}$ such that $\alpha_i = \varphi_i^\#(\alpha_{i+1})$.

Let us also consider an element $y \in H^{*,*}(-, \lim_{\rightarrow} X_i)$. Since the homology functor on the category of complexes of abelian groups commutes with direct limits, it implies

$$H^{*,*}(-, \lim_{\rightarrow} X_i) \cong \lim_{\rightarrow} H^{*,*}(-, X_i).$$

Hence, the element y determines a set of elements $\{y_j \in H^{*,*}(-, X_j)\}_{j \gg 0}$ such that $\varphi_j^\#(y_j) = y_{j+1}$. We construct $\check{\alpha} \in \mathcal{O}\mathcal{P}^{*,*}(\lim_{\rightarrow} X_i, W)$, setting $\check{\alpha}(y) := \alpha_N(y_N)$ for $N \gg 0$. Since

$$\alpha_{N+1}(y_{N+1}) = \alpha_{N+1}(\varphi_N^\#(y_N)) = (\varphi_N^\# \alpha_{N+1})(y_N) = \alpha_N(y_N),$$

the operation $\check{\alpha}$ is well-defined.

In order to construct the map in the opposite direction, let us start with an operation $\gamma \in \mathcal{O}\mathcal{P}^{*,*}(\lim_{\rightarrow} X_j, W)$ and construct for every index j the operation $\hat{\gamma}_j \in \mathcal{O}\mathcal{P}^{*,*}(X_j, W)$ given by the through map

$$H^{*,*}(-, X_j) \rightarrow H^{*,*}(-, \lim_{\rightarrow} X_j) \xrightarrow{\gamma} H^{*,*}(-, W),$$

where the first arrow is canonical and the second is given by the operation γ . These operations fit together to make an element of the projective system and, therefore, the operation $\hat{\gamma} \in \lim_{\leftarrow j} \mathcal{O}\mathcal{P}^{*,*}(X_j, W)$. One can easily verify that given constructions are mutually inverse. \square

Corollary 5.11 *The natural map $H^*(\mathbf{Z}/p^\infty, G) \rightarrow \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^\infty, G)$ is an isomorphism for $G = \mathbf{Z}/p$ or \mathbf{Z} .*

Proof. We have already seen above that $H^*(\mathbf{Z}/p^m, G) \cong \mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^m, G)$. These groups and their maps were explicitly computed in Proposition 5.7 and Corollary 5.9. The computation also implies that $\lim_{\leftarrow m} {}^1H^{*,p-1}(\mathbf{Z}/p^m, G) = 0$. The desired result now follows from the short exact sequence:

$$0 \rightarrow \lim_{\leftarrow m} {}^1H^{*-1,p-1}(\mathbf{Z}/p^m, G) \rightarrow H^{*,p-1}(\mathbf{Z}/p^\infty, G) \rightarrow \lim_{\leftarrow m} H^{*,p-1}(\mathbf{Z}/p^m, G) \rightarrow 0.$$

\square

Corollary 5.12 *If $G = \mathbf{Z}/p$, the \mathbf{Z}/p -module $\mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^\infty, G)$ has two generators $P^1\beta_\infty, \beta_1 P^1\beta_\infty$, lying in degrees $2p - 1, 2p$, correspondingly. If $G = \mathbf{Z}$, it is generated by the element $\beta_{\mathbf{Z}} P^1\beta_\infty$. Here $\beta_\infty = [\mathbf{Z}/p, \mathbf{Z}/p^\infty, \mathbf{Z}/p^\infty]$.*

Proof. Corollaries 5.9 and 5.11 give us an explicit description of the generators of the module $\mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^\infty, G)$. Since it follows from Proposition 5.1.ii that there is a natural identification $\beta_\infty = \lim \beta_m$, this completes the proof. \square

Let us now return back from p -cyclotomic coefficients to p -local. Further, we will also need some auxiliary results about rational operations, which are presented in the appendix.

Proposition 5.13 *The Bockstein homomorphism $B = [\mathbf{Z}_{(p)}, \mathbf{Q}, \mathbf{Z}/p^\infty]$ induces an isomorphism of \mathbf{Z}/p -modules: $\mathcal{O}\mathcal{P}^*(\mathbf{Z}_{(p)}) \cong \mathcal{O}\mathcal{P}^{*+1}(\mathbf{Z}/p^\infty)$. So that, the group $\mathcal{O}\mathcal{P}^l(\mathbf{Z}_{(p)})$ is \mathbf{Z}/p in degrees $l = 2p - 2, 2p - 1$ and trivial otherwise. One can take operations $P^1r, \beta_1 P^1r \in \mathcal{O}\mathcal{P}^*(\mathbf{Z}_{(p)})$ as generators in the corresponding degrees. Here $r: \mathbf{Z}_{(p)} \rightarrow \mathbf{Z}/p$ is the coefficient reduction map.*

Proof. Let us recall that by (4.4) one has: $\mathcal{O}\mathcal{P}^{i,j}(A, B) = \lim_{\leftarrow n} \tilde{H}^{i+2n, j+n}(K_n(A), B)$. Consider the following commutative square:

$$\begin{array}{ccc} H^{*+1, p-1}(\mathbf{Z}/p^\infty) & \xrightarrow{\cong} & \mathcal{O}\mathcal{P}^{*+1}(\mathbf{Z}/p^\infty) \\ \uparrow B & & \uparrow \\ H^{*, p-1}(\mathbf{Z}_{(p)}) & \twoheadrightarrow & \lim_{\leftarrow m} \tilde{H}^{*+2m, *+m}(K_m(\mathbf{Z}_{(p)})), \end{array}$$

where the vertical arrows are induced by the Bockstein homomorphism B and both horizontal arrows are epimorphisms from (5.1). The top horizontal arrow is an isomorphism by Lemma 5.11.

Taking the short exact sequence of abelian groups $0 \rightarrow \mathbf{Z}_{(p)} \rightarrow \mathbf{Q} \rightarrow \mathbf{Z}/p^\infty \rightarrow 0$ and applying Proposition 5.2, one gets a distinguished triangle

$$\mathbf{H}(\mathbf{Z}_{(p)}) \rightarrow \mathbf{H}(\mathbf{Q}) \rightarrow \mathbf{H}(\mathbf{Z}/p^\infty) \quad (5.3)$$

of spectra. Using the triangle and lemma A.1, one shows that the map B in the diagram is also an isomorphism. Hence, all maps in the diagram are isomorphisms. This proves the isomorphism $\mathcal{O}\mathcal{P}^*(\mathbf{Z}_{(p)}) \stackrel{B}{\cong} \mathcal{O}\mathcal{P}^{*+1}(\mathbf{Z}/p^\infty)$.

Finally, the equality $rB = \beta_\infty = [\mathbf{Z}/p, \mathbf{Z}/p^\infty, \mathbf{Z}/p^\infty]$ together with the description of the groups $\mathcal{O}\mathcal{P}^*(\mathbf{Z}/p^\infty, \mathbf{Z}/p)$ given above, supplies us with the desired set of generators. This proves the proposition. \square

Our current purpose is to compute the group $\mathcal{O}\mathcal{P}^*(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)})$.

Proposition 5.14

$$H^l(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) = \begin{cases} \mathbf{Z}/p & \text{for } l = 2p - 1 \\ 0 & \text{for } l \geq 2p. \end{cases}$$

Proof. We show, first, that $H^l(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)})$ is a p -group for $l \geq 2p - 1$. Using distinguished triangle (5.3) and the universal coefficient formula, one can write the exact sequence:

$$H^l(\mathbf{Q}, \mathbf{Z}_{(p)}) \longrightarrow H^l(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \longrightarrow H^{l+1}(\mathbf{Z}/p^\infty, \mathbf{Z}) \otimes \mathbf{Z}_{(p)}.$$

By Corollary 5.12 we already know that $H^{l+1}(\mathbf{Z}/p^\infty, \mathbf{Z})$ is either \mathbf{Z}/p for $l = 2p - 1$, or 0. So, it suffices to show that $H^l(\mathbf{Q}, \mathbf{Z}_{(p)})$ is a p -group. By A.2, one has: $0 = H^l(\mathbf{Q}, \mathbf{Q}) = H^l(\mathbf{Q}, \mathbf{Z}_{(p)}) \otimes_{\mathbf{Z}_{(p)}} \mathbf{Q}$

and this group is the p -localization of $H^l(\mathbf{Q}, \mathbf{Z}_{(p)})$. So, the statement follows.

Now, consider the fragment of the coefficient long exact sequence:

$$H^{2p-2}(\mathbf{Z}_{(p)}) \xrightarrow{\mathfrak{B}} H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \xrightarrow{p} H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \xrightarrow{r} H^{2p-1}(\mathbf{Z}_{(p)}),$$

where $\mathfrak{B} = [\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}, \mathbf{Z}/p]$. As we already know from the computation above (Proposition 5.13), both the groups with finite coefficients are isomorphic to \mathbf{Z}/p and the isomorphism between them can be performed by the map $\beta_1: H^{2p-2}(\mathbf{Z}_{(p)}) \rightarrow H^{2p-1}(\mathbf{Z}_{(p)})$. One can easily verify the relation $\beta_1 = \mathfrak{B}r$. From Lemma 5.8 and the fact that $H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)})$ is a p -group, we have:

$H^{2p-2}(\mathbf{Z}_{(p)}) \xrightarrow{\mathfrak{B}} H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)})$. The same arguments can be used to show that $H^l(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) = 0$ for $l \geq 2p$. One just should mention, in addition, that the map $r: H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \rightarrow H^{2p-1}(\mathbf{Z}_{(p)})$ in the sequence above is an isomorphism. \square

Corollary 5.15 *The Bockstein homomorphism \mathfrak{B} induces the isomorphism:*

$$\mathcal{O}\mathcal{P}^{2p-2}(\mathbf{Z}_{(p)}) \xrightarrow{\mathfrak{B}} \mathcal{O}\mathcal{P}^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}).$$

Proof. In the comutative diagram

$$\begin{array}{ccc} H^{2p-2}(\mathbf{Z}_{(p)}) & \longrightarrow & H^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \\ \downarrow & & \downarrow \\ \mathcal{O}\mathcal{P}^{2p-2}(\mathbf{Z}_{(p)}) & \xrightarrow{\mathfrak{B}} & \mathcal{O}\mathcal{P}^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}) \end{array}$$

the top arrow is an isomorphism by Proposition 5.14. The right vertical arrow is an epimorphism by (5.1). So, the map \mathfrak{B} is an epimorphism. Since $\mathcal{O}\mathcal{P}^{2p-2}(\mathbf{Z}_{(p)}) \cong \mathbf{Z}/p$ and, as it follows from the next section results, the group $\mathcal{O}\mathcal{P}^{2p-1}(\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)})$ is non-trivial, the statement follows. \square

Summarizing the results of 5.13, 5.14, and 5.15, we obtain the following

Theorem 5.16 *There are no non-trivial stable cohomological operations of weight $p - 1$ and degree greater than $2p - 1$. Every non-trivial bistable operation*

$$H^{*,*}(-, \mathbf{Z}_{(p)}) \rightarrow H^{*+2p-1, *+p-1}(-, \mathbf{Z}_{(p)})$$

of degree $2p - 1$ in motivic cohomology coincides, after multiplication by a unit of \mathbf{Z}/p , with the operation $\mathfrak{B}P^1r$, where P^1 denotes the first \mathbf{Z}/p motivic Steenrod power, $\mathfrak{B} = [\mathbf{Z}_{(p)}, \mathbf{Z}_{(p)}, \mathbf{Z}/p]$, and r is the corresponding coefficient reduction operation.

6. $d_p \neq 0$

The purpose of the current section is to construct for every prime p a smooth variety having the property that the p -th differential d_p is non-zero. Although in the previous discussion we systematically avoided the case of $p = 2$, in this section we decided to give slightly more general statements for completeness. So, let p just be a prime number. All coefficient rings are by default $\mathbf{Z}_{(p)}$. Abusing the notation, we omit mentioning coefficients unless it is absolutely necessary.

Below we give two examples, demonstrating non-triviality of the differential d_2 for $p = 2$ and d_p for odd primes.

Example 6.1 Consider the motivic cohomology spectral sequence for the variety $\text{Spec } \mathbf{Q}$. One can check that the Milnor symbol $\{-1, -1, -1, -1\} \in K_4^M(\mathbf{Q})$ is non-trivial of order 2. The group $K_4^M(\mathbf{Q}) = \mathbf{Z}/2$ is canonically isomorphic to $E_2^{0,-4}$. On the other hand, the spectral sequence converges in the degree $i + j = -4$ to $K_4(\mathbf{Q})$ and the map $K_4^M(\mathbf{Q}) \rightarrow K_4(\mathbf{Q})$ should pass through the stable homotopy group of the sphere spectrum π_S^4 . The latter group is trivial, therefore, one gets from the short exact sequence $E_2^{-2,-3} \xrightarrow{d_2} E_2^{0,-4} \rightarrow E_\infty^{0,-4}$ that the differential $d_2: H^{1,3}(\text{Spec } \mathbf{Q}) \rightarrow E_2^{0,-4} = K_4^M(\mathbf{Q})$ is non-zero. This is, certainly, true with $\mathbf{Z}_{(2)}$ coefficients as well.

More detailed explanation can be found in [We, VI.4.3, Ex. IV.1.12, Ex. III.7.2].

Proposition 6.2 Let us assume that for an odd prime number p and a variety $G \in \text{Sm}/k$ the following conditions are satisfied:

- (i) $K_0(G, \mathbf{Z}_{(p)}) = \mathbf{Z}_{(p)} \cdot \mathbf{1}$, where the class $\mathbf{1}$ lies in codimension 0;
- (ii) $CH^{p+1}(G, \mathbf{Z}_{(p)}) \neq 0$.

Then, the differential $d_p: E_p^{1,-2} \rightarrow E_p^{p+1,-p-1}$ in the motivic spectral sequence

$$E_2^{i,j} = H^{i-j,-j}(G, \mathbf{Z}_{(p)}) \Rightarrow K_{-i-j}(G, \mathbf{Z}_{(p)})$$

is non-trivial.

Proof. Since motivic cohomology groups coincide with higher Chow groups, the term

$$E_2^{p+1,-p-1} = CH^{p+1}(G, 0) = CH^{p+1}(G) \neq 0$$

by (ii). By Proposition 3.1, one has: $E_2 = E_p$. On the other hand, $E_\infty^{p+1,-p-1} = 0$, since, by (i) the whole group $K_0(G)$ is concentrated in the term $E_\infty^{0,0} = \mathbf{Z}_{(p)}$. Again, by Proposition 3.1 and the triviality of groups $E_2^{i,j} = H^{i-j,-j}(X)$ for $j > 0$ or $i + j > 0$ (see 1.1.(i,iii)), one also has: $E_{p+1}^{p+1,-p-1} = 0$. (For the case $p = 3$ we also need to use triviality of the group $H^{-1,0}(X)$ (see 1.1.iv.) Hence, there should be a non-trivial differential, that “kills” the term $E_p^{p+1,-p-1}$ and the only possibility is that $0 \neq d_p: E_p^{1,-2} \rightarrow E_p^{p+1,-p-1}$. \square

Example 6.3 Consider a non-split central simple algebra \mathcal{D} of degree p over k . Set $G = SL_{1,\mathcal{D}}$ to be the norm variety, the subvariety of \mathcal{D} , given by the equation $\text{Nrd } x = 1$, where Nrd denotes the reduced norm (see [GSz06, Sect. 2.6]). This gives us an example of a variety with $d_p \neq 0$.

Now we are almost ready to complete the proof of Theorem 2.1.

Let us fix an odd prime number p and a base-field characteristic l such that either $l = 0$, or $(l, p) = 1$. We also introduce a field F of characteristic l , setting

$$F = \begin{cases} \mathbf{Q} & \text{for } l = 0 \\ \mathbf{F}_l(t) & \text{for } l > 0. \end{cases}$$

Here $\mathbf{F}_l(t)$ is the field of rational functions over the prime finite field \mathbf{F}_l . The global class field theory tells us that in all the cases the Brauer group $\text{Br}(F)$ has many non-trivial p -torsion elements, so one can construct non-split central simple algebras over F and for any characteristic l we obtain examples of fields and varieties over them with non-trivial differentials d_p .

Due to Levine’s result mentioned before [Le08], the MCSS coincides with the slice spectral sequence restricted to the category of smooth varieties. In particular, this means that the differentials in MCSS can be considered as stable cohomological operations on the category \mathbf{Spc}

of Voevodsky's spaces. Let us now consider the case $k = \mathbf{F}_l$ for a prime number l . A non-split central simple algebra over $\mathbf{F}_l(t)$ can be seen as a motivic space over k . As we have shown in the previous paragraph, the corresponding slice spectral sequence has non-trivial differential d_p . This implies that the coefficient α in the Theorem's 2.1 relation $d_p = \mathfrak{B}\alpha P^1 r$ is non-zero over k . This argument shows that $\alpha \neq 0$ for every prime field, because we already know the conclusion is true for the case of characteristic 0. Using the functoriality of MCSS and cohomological operations, one can show that the same statement holds for an arbitrary field. This completes the proof of Theorem 2.1. \square

It is left to show that the variety G from Example 6.3 satisfies the assumptions of Proposition 6.2. The first one is checked in [Su91, Thm. 6.1]. The rest of the paper is devoted to prove the second one.

Below we denote by $X = SB(\mathcal{D})$ the Severi–Brauer variety, corresponding to the algebra \mathcal{D} (see [GSz06, Ch. 5]). This is a twisted form of the projective space \mathbf{P}^{p-1} . So, one has: $\dim X = p - 1$. Let us also mention that since G is a twisted form of SL_p , one has $\dim G = p^2 - 1$.

Proposition 6.4 *For the variety $G = SL_{1,\mathcal{D}}$ introduced above, one has: $CH^{p+1}(G) \neq 0$.*

Proof. Setting, as above, $X = SB(\mathcal{D})$, for the projection map $G \times X \rightarrow G$ consider a filtration of the base by codimension of points and write down the corresponding spectral sequence (see Rost[Ro96, Sect. 8]):

$$E_1^{st}(n) = \coprod_{g \in G^{(s)}} H^t(X_{F(g)}, \mathcal{K}_{n-s}) \Rightarrow H^{s+t}(G \times X, \mathcal{K}_n), \quad (6.1)$$

where $X_{F(g)} = X \times \text{Spec } F(g)$ is a fiber over the generic point g of codimension s . This spectral sequence is a natural generalization of the Brown–Gersten–Quillen (BGQ) spectral sequence (cohomology groups here are \mathcal{K} -cohomology).

For convenience, we have included a diagram below of the case $n = p + 1$, which is the most important case for us. For brevity we have used the following notation:

$$\coprod_{g \in G^{(s)}} H^t(X_{F(g)}, \mathcal{K}_u) =: R_{t,u}^s.$$

The non-zero part of the E_1 -term is concentrated in the strip given by the inequalities: $0 \leq t \leq p - 1$ and $s + t \leq n$. Let us denote the spectral sequence $E_*^{*,*}(p + 1)$ by $E_*^{*,*}$. So that, $E_1^{s,t} = R_{t,p+1-s}^s$.

$$\begin{array}{cccccc}
 R_{p-1,p+1}^0 & R_{p-1,p}^1 & R_{p-1,p-1}^2 & 0 & 0 & \\
 \vdots & \vdots & & \ddots & 0 & 0 \\
 R_{1,p+1}^0 & R_{1,p}^1 & \cdots & R_{1,2}^{p-1} & R_{1,1}^p & 0 \\
 R_{0,p+1}^0 & R_{0,p}^1 & \cdots & & R_{0,1}^p & R_{0,0}^{p+1}
 \end{array}$$

Let the following statements hold:

- (i) $E_2^{p+1,0} = CH^{p+1}(G)$;
- (ii) The boundary map $H^p(G \times X, \mathcal{K}_{p+1}) \xrightarrow{\varphi} E_p^{1,p-1}$ is not an epimorphism.

Then the proposition follows easily. Actually, just consider a fragment of the boundary short exact sequence:

$$H^p(G \times X, \mathcal{K}_{p+1}) \xrightarrow{\varphi} E_p^{1,p-1} \xrightarrow{d_2^p} E_p^{p+1,0}.$$

By (ii) φ is not an epimorphism, and so one has: $E_p^{p+1,0} \neq 0$. But by (i) and for dimension reasons there exists an epimorphism $CH^{p+1}(G) = E_2^{p+1,0} \twoheadrightarrow E_p^{p+1,0}$ that proves the desired result.

The rest of the paper is devoted to the proof of the auxiliary statements: (i) is established in Lemma 6.5 right below, (ii) is proven in Proposition 6.10. \square

Lemma 6.5 *In the spectral sequence considered in Proposition 6.4 above, one has: $E_2^{p+1,0} = CH^{p+1}(G)$.*

Proof. One has: $E_2^{p+1,0} = R_{0,0}^{p+1}/R_{0,1}^p$. Decoding the notation, we get:

$$E_2^{p+1,0} = \text{Coker} \left(\prod_{g \in G^{(p)}} F(g)^* \rightarrow \prod_{g \in G^{(p+1)}} \mathbf{Z} \right) = CH^{p+1}(G) \quad (6.2)$$

that completes the proof. The same is, certainly, true with $\mathbf{Z}_{(p)}$ coefficients. \square

In order to check statement (ii), we should perform some computation with the term $E_p^{1,p-1}$. The following lemma simplifies our life showing that we actually work with the term $E_2^{1,p-1}$.

Lemma 6.6 *In the spectral sequence in Proposition 6.4, one has: $E_2^{1,p-1} = E_p^{1,p-1}$.*

Proof. By the next lemma, one has: $E_2^{p+1-t,t} = 0$ for $1 \leq t \leq p-1$. So, for dimension reasons, the only non-trivial differential with domain $E_*^{1,p-1}$ is d_p . \square

Lemma 6.7 *The differential maps $d_1^t: R_{t,t+1}^{p-t} \rightarrow R_{t,t}^{p-t+1}$ are epimorphisms, provided that $1 \leq t \leq p-1$. In other words, in these cases $E_2^{p+1-t,t} = 0$.*

Proof. We have to prove that the maps

$$\prod_{g \in G^{(p-t)}} H^t(X_{F(g)}, \mathcal{K}_{t+1}) \rightarrow \prod_{g \in G^{(p+1-t)}} H^t(X_{F(g)}, \mathcal{K}_t)$$

are epimorphisms. The inner groups $H^t(X_{F(g)}, \mathcal{K}_{t+m})$ can be computed using the Brown–Gersten–Quillen spectral sequence. Writing down Gersten resolutions for different values of t one gets natural maps of the resolutions, induced by embeddings of points of different codimensions. This implies natural maps of BGQ spectral sequences and, finally, natural maps of \mathcal{K} -cohomology groups

$$\cdots \rightarrow H^t(X_{F(g)}, \mathcal{K}_{t+m}) \rightarrow H^{t+1}(X_{F(g)}, \mathcal{K}_{t+1+m}) \rightarrow \cdots$$

By Statement 6.8, these maps are isomorphisms for $m = 0, 1$ and $1 \leq t \leq p-1$. By functoriality of the construction this implies that

$$E_2^{p+1-t,t} = R_{t,t}^{p+1-t}/R_{t,t+1}^{p-t} \simeq R_{p-1,p-1}^{p+1-t}/R_{p-1,p}^{p-t} = E_2^{p+1-t,p-1}(2p-t).$$

The rest follows from Lemma 6.9 below. \square

In the proof of the previous proposition we used a result of Merkurjev and Suslin, which we reproduce here.

Statement 6.8 ([MS82, Cor. 8.7.2]) *Let \bar{k} be the algebraic closure of k . For a Severi–Brauer variety X of dimension $p - 1$, set $\bar{X} = X \times \text{Spec } \bar{k}$. Then*

$$H^i(X, \mathcal{K}_i) = CH^i(X) = p\mathbf{Z}_{(p)} \subset \mathbf{Z}_{(p)} = CH^i(\bar{X}) \quad (6.3)$$

and

$$H^i(X, \mathcal{K}_{i+1}) = \text{Nrd } \mathcal{D}^* \subset \bar{k}^* = H^i(\bar{X}, \mathcal{K}_{i+1}), \quad (6.4)$$

provided that $1 \leq i \leq p - 1$. (Here Nrd denotes the group of the reduced norms.)

Lemma 6.9 *For $n > p$, one has: $E_2^{n-p+1, p-1}(n) = 0$.*

Proof. Consider now $G \times X$ as a group-variety over X . By Suslin’s computations [Su91, Thm. 4.2], $H^*(G \times X, \mathcal{K}_*)$ becomes a module over $H^*(X, \mathcal{K}_*)$ generated by Chern classes c_j for $j \geq 1$, where $c_j \in H^j(G \times X, \mathcal{K}_{j+1})$. In particular, this implies that $CH^i(G \times X) = 0$ for $i > p - 1$. Therefore, the spectral sequence converges to zero in the n -th diagonal. In particular, $E_\infty^{n-p+1, p-1}(n) = 0$. For dimension reasons there are no differentials affecting the term $E_2^{n-p+1, p-1}(n)$. So one has: $E_2^{n-p+1, p-1}(n) = E_\infty^{n-p+1, p-1}(n) = 0$. \square

Proposition 6.10 *The map $\varphi: H^p(G \times X, \mathcal{K}_{p+1}) \rightarrow E_p^{1, p-1}$ has non-trivial cokernel.*

Proof. Let us mention, first, that by the previous lemma, one has: $E_p^{1, p-1} = E_2^{1, p-1}$. Consider the base-change commutative diagram corresponding to the morphism $\text{Spec } \bar{k} \rightarrow \text{Spec } k$, where \bar{k} is the algebraic closure of k . Later we denote $E_2^{1, p-1}$ by V and the corresponding group $E_2^{1, p-1}$ over \bar{k} by \bar{V} .

$$\begin{array}{ccc} H^p(G \times X, \mathcal{K}_{p+1}) & \xrightarrow{\varphi} & V \\ \chi \downarrow & & \psi \downarrow \\ H^p(\bar{G} \times \bar{X}, \mathcal{K}_{p+1}) & \xrightarrow{\bar{\varphi}} & \bar{V} \end{array} \quad (6.5)$$

The desired statement can be derived easily from the following three claims:

- (i) $\text{Im } \chi$ is divisible by p ;
- (ii) $\psi: V \rightarrow \bar{V}$ is an epimorphism;
- (iii) $\bar{V} = \mathbf{Z}_{(p)}$.

Assume that φ is an epimorphism. Since ψ is also an epimorphism, we can chose an element $x \in H^p(G \times X, \mathcal{K}_{p+1})$ such that $\psi\varphi(x) = 1$. Then, by (i), $1 = \bar{\varphi}\chi(x)$ is p -divisible. This gives a contradiction. We prove (i) in Lemma 6.11 and (ii) in Proposition 6.13 below. Finally, (iii) appears in the proof of 6.13 as an indirect result. \square

Lemma 6.11 *For the base-change morphism $\chi: H^p(G \times X, \mathcal{K}_{p+1}) \rightarrow H^p(\bar{G} \times \bar{X}, \mathcal{K}_{p+1})$ the image of χ is divisible by p .*

Proof. This follows from the above mentioned (see the proof of Lemma 6.9) decomposition

$$H^p(G \times X, \mathcal{K}_{p+1}) = \coprod_{i>0} c_i CH^{p-i}(X) \quad (6.6)$$

and the fact that the map $CH^i(X) \rightarrow CH^i(\bar{X})$ is a multiplication by p due to Statement 6.8. \square

Before we can prove the last proposition, we need to construct one map. For this end, let us reproduce here one important definition (see [Pa91, 3.1] for details).

Definition 6.12 *For a quasi-compact locally Noetherian scheme Y , let A be a sheaf of algebras on Y locally isomorphic in the étale topology to the sheaf of split algebras $M_n(\mathcal{O}_Y)$. In other words, A is an Azumaya algebra on Y .*

Consider the category of sheaves of left A -modules and denote by $\mathcal{P}(Y; A)$ its full subcategory, whose objects are locally free coherent \mathcal{O}_Y -modules. We set $K_(Y; A) := K_*(\mathcal{P}(Y; A))$, where the functor on the right-hand side is obtained by application of Quillen's Q -construction [Qu73].*

We will also write, for brevity, $H^(G, \mathcal{K}_*; A)$ for $H^*(G, \mathcal{K}_*(-; A))$.*

Currently, we are going to construct a natural epimorphism $\tilde{\rho}: V \rightarrow H^1(G, \mathcal{K}_2; \mathfrak{D})$, where $\mathfrak{D} := \mathcal{D}^{\otimes(p-1)}$ and $V = E_2^{1,p-1}$ (see Proposition 6.10).

First, consider the BGQ spectral sequence converging to the K -groups of the Severi–Brauer variety X . Since $(p-1)!$ is invertible in the coefficient ring, this spectral sequence has no non-trivial differentials affecting the two highest diagonals. Moreover, if the base field is algebraically closed, all the differentials in the spectral sequence vanish (see [MS82, 8.6.2]). Again, by the invertibility of $(p-1)!$ the topological filtration on the K -groups coincides with γ -filtration. The latter filtration is generated by the image of the corresponding γ -operation.

E_∞ -term of the BGQ consists of consequent factor-filtration groups of $K(X)$. Taking into account the triviality of differentials, mentioned in the previous paragraph, there exist boundary maps:

$$H^{p-1}(X, \mathcal{K}_{p-1+m}) \rightarrow K_m(X)^{(p-1)}, \quad (6.7)$$

where $m = 0, 1, 2$ and we have the smallest non-trivial filtration group on the right-hand side. These maps are isomorphisms for $m = 0, 1$. Provided that the base field is algebraically closed, they are isomorphisms also for $m = 2$.

By Quillen's computation of K -groups of Severi–Brauer varieties [Qu73], one has isomorphisms: $K_m(X)^{(p-1)} \simeq K_m(\mathfrak{D})$. So that, we obtain the maps: $H^{p-1}(X_g, \mathcal{K}_{p-1+m}) \xrightarrow{\rho_m} K_m(F(g); \mathfrak{D})$ for $m = 0, 1, 2$, which are isomorphisms for $m = 0, 1$ and isomorphism for $m = 2$ provided that the base field is algebraically closed. As a result, one gets the map of complexes ρ_* :

$$\begin{array}{ccccc} R_{p-1,p+1}^0 & \longrightarrow & R_{p-1,p}^1 & \longrightarrow & R_{p-1,p-1}^2 \\ \downarrow \rho_2 & & \downarrow \rho_1 & & \downarrow \rho_0 \\ K_2(F(G); \mathfrak{D}) & \longrightarrow & \coprod_{g \in G^{(1)}} K_1(F(g); \mathfrak{D}) & \longrightarrow & \coprod_{g \in G^{(2)}} K_0(F(g); \mathfrak{D}), \end{array} \quad (6.8)$$

inducing the epimorphism map $\tilde{\rho}$ on the middle-term homology groups. The latter map becomes an isomorphism after passing to the algebraic closure. The middle-term homology groups in the upper and bottom lines can be identified with V and $H^1(G, \mathcal{K}_2; \mathfrak{D})$, correspondingly, that gives us the desired epimorphism $\tilde{\rho}$.

Proposition 6.13 *Let V and \bar{V} be as before. Then, the map $\psi: V \rightarrow \bar{V}$ is an epimorphism.*

Proof. Let us consider the base-change diagram corresponding to the morphism $\text{Spec } \bar{k} \rightarrow \text{Spec } k$:

$$\begin{array}{ccc} V & \xrightarrow{\psi} & \bar{V} \\ \tilde{\rho} \downarrow & & \parallel \\ H^1(G, \mathcal{K}_2; \mathfrak{D}) & \xrightarrow{\omega} & H^1(\bar{G}, \mathcal{K}_2; \bar{\mathfrak{D}}). \end{array} \quad (6.9)$$

Observe now, that $\bar{G} = SL_n(\bar{k})$ and $H^1(\bar{G}, \mathcal{K}_2; \bar{\mathcal{D}}) = H^1(SL_n, \mathcal{K}_2) = \mathbf{Z}_{(p)}$ with a natural choice of a generator, given by the first Chern class (see [Su91, Thm. 2.7]). This gives us the following commutative diagram:

$$\begin{array}{ccc}
 V & \xrightarrow{\psi} & \bar{V} \\
 \bar{\rho} \downarrow & & \parallel \\
 H^1(G, \mathcal{K}_2; \mathcal{D}) & \xrightarrow{\omega} & H^1(SL_n, \mathcal{K}_2) \\
 \uparrow c_1 & & \uparrow \bar{c}_1 \\
 K_1(G; \mathcal{D}) & \xrightarrow{f} & K_1(SL_n)
 \end{array} \tag{6.10}$$

Consider the universal element $\alpha \in K_1(G; \mathcal{D})$ determined as in [Su91, Sect. 4]. It is constructed in such a way that its image $f(\alpha)$ in $K_1(SL_n)$ is the universal matrix element. Then, due to [Su91, Thm. 2.7], $\bar{c}_1 f(\alpha) = 1$. Hence, the map ω is an epimorphism and so is ψ . \square

Appendix A. Something about the groups of rational operations

In this short appendix we give two statements concerning cohomology groups of the spectrum $\mathbf{H}(\mathbf{Q})$, which we need in the paper.

Statement A.1 *All motivic cohomology groups of the spectrum $\mathbf{H}(\mathbf{Q})$ with \mathbf{Z}/p -coefficients vanish.*

Proposition A.2 *For integers $n, \varepsilon > 0$, one has: $H^{2n+\varepsilon, n}(\mathbf{H}(\mathbf{Q}), \mathbf{Q}) = 0$.*

Proof. We want to compute the group $H^{2n+\varepsilon, n}(\mathbf{H}(\mathbf{Q}), \mathbf{Q}) = [\mathbf{H}(\mathbf{Q}), \mathbf{H}(\mathbf{Q})[2n + \varepsilon](n)]$. Since the spectrum $\mathbf{H}(\mathbf{Q})$ is a direct summand of the spectrum $\mathbf{BGL}_{\mathbf{Q}}$, it suffices to show that $[\mathbf{BGL}_{\mathbf{Q}}, \mathbf{BGL}_{\mathbf{Q}}[2n + \varepsilon](n)] = 0$. Using the Bott periodicity and [Ri10, Cor. 5.3.1], we have:

$$[\mathbf{BGL}_{\mathbf{Q}}, \mathbf{BGL}_{\mathbf{Q}}[2n + \varepsilon](n)] = [\mathbf{BGL}_{\mathbf{Q}}, \mathbf{BGL}_{\mathbf{Q}}[\varepsilon]] = \varprojlim (K_{-\varepsilon}(k))^{\Omega}.$$

(Here we use the notation of [Ri10].) Since the group $K_{-\varepsilon}(k)$ is the algebraic K -group of the base field k and, obviously, vanishes for $\varepsilon > 0$, then so does $\varprojlim (K_{-\varepsilon}(k))^{\Omega}$. \square

ACKNOWLEDGEMENTS

I would like to thank Andrei Suslin and Ivan Panin for their interest in the topic and multiple fruitful discussions during my work. I am deeply grateful to Alexander Merkurjev, whose help with the construction of a variety with $d_p \neq 0$ can not be overestimated. I am also in debt to Charles Weibel, who gave me, besides many valuable remarks, a reference to Example 6.1. I thank Alexander Vishik and Marc Levine for their invitations to visit the University of Nottingham and Universität Duisburg–Essen and thank both the universities for their hospitality. Finally, I'm very grateful to anonymous referee for his/her questions and remarks, which pointed my attention to some gaps in the preliminary version of the paper.

The significant part of this paper was written during my visit to IHES (Bures-sur-Yvette). I cordially thank the Institute for its hospitality and excellent work conditions. Writing of the revised version was partially supported by research grant 14-11-00456 from Russian Science Foundation. The author is grateful to the foundation for its support.

REFERENCES

- Ad65 Adams, J.F. *On the groups $J(X)$ -II*, Topology Vol. 3, 1965, pp. 137-171.
- Ar62 Artin, M. *Grothendieck Topologies*. Lecture Notes, Harvard University Math. Dept., Cambridge, Mass., 1962.
- BL95 Bloch, S. and Lichtenbaum, S. *A Spectral Sequence for Motivic Cohomology*. www.math.uiuc.edu/K-theory/0062, (1995) (preprint).
- Bu69 (Buchstaber, V.M.) Бухштабер, В.М. *Модули дифференциалов спектральной последовательности Атья–Хирцебруха*. Матем. сб. 78 (120), 2, (1969), 307–320.
English translation: Buchstaber, V.M. *Modules of differentials of the Atiyah–Hirzebruch spectral sequence*. Math. USSR Sbornik Vol.7, No.2, (1969), pp. 299–313.
- Ca54 Cartan, H. *Algebres d’Eilenberg–Mac Lane et homotopie*, Seminaire H. Cartan, Ecole Norm. Sup., vol. 7, No. 1, 1954–5.
- De09 Deligne, P. *Voevodsky’s Lectures on motivic cohomology 2000/2001*. Algebraic Topology. The Abel Symposium 2007., Springer Verlag, 2009., pp. 355–409.
- FS02 Friedlander, E. and Suslin, A. *The spectral sequence relating algebraic K-theory to motivic cohomology*. Ann. Sci. Ecole Norm. Sup. (4) 35 (2002), no. 6, 773–875.
- GSo99 Gillet, H. and Soulé, C., *Filtrations on higher algebraic K-theory*. Proc. Sympos. Pure Math., 67, 89–148, (1999).
- GSz06 Gille, P. and Szamuely, T., *Central simple algebras and Galois cohomology*. Cambridge Studies in Advanced Mathematics, Vol. 101, Cambridge University Press, Cambridge, 2006. 343p.
- Gr95 Grayson, D. *Weight filtrations via commuting automorphisms*. K-Theory 9 (1995), no. 2, 139–172.
- HKØ13 Hoyois, M., Kelly, S., and Østvær, P.A. *The motivic Steenrod algebra in positive characteristic*. arXiv.org. 1305.5690v2, 2013
- KM60 Kervaire, M. and Milnor, J. *Bernoulli numbers, homotopy groups, and a theorem of Rohlin*. 1960 Proc. Internat. Congress Math. 1958 pp. 454–458 Cambridge Univ. Press, N.-Y.
- Le08 Levine, M. *The homotopy coniveau tower*. J. Topology 1 (2008), pp. 217–267.
- MVW06 Mazza, C., Voevodsky, V., and Weibel, C. *Lecture notes on motivic cohomology*. Clay Mathematics Monographs, 2, AMS, Providence, RI; Clay Math. Inst., Cambridge, MA, 2006, 216p.
- Me10 Merkurjev, A. *Adams operations and the Brown-Gersten-Quillen spectral sequence*. Quadratic forms, linear algebraic groups, and cohomology, Dev. Math., Vol. 18, pp. 305–313, 2010
- MS82 Merkurjev, A. and Suslin, A. *K-cohomology of Severi-Brauer varieties and the norm residue homomorphism*. Izv. Akad. Nauk SSSR Ser. Mat. 46 (1982), no. 5, 1011–1046.
- Pa91 Panin, I. A., *On algebraic K-theory of generalized flag fiber bundles and some of their twisted forms*. Algebraic K-theory, Adv. Soviet Math., 4 Providence, AMS, 1991 21–46.
- Qu73 Quillen, D. *Higher Algebraic K-theory I*. Lecture Notes in Math., 341, Springer-Verlag, 1973, 85-147.
- Ri10 Riou, J. *Algebraic K-theory, A^1 -homotopy and Riemann–Roch theorems*. Journal of Topology 3 (2010), 229–264.
- Ro96 Rost, M. *Chow groups with coefficients*. Doc. Math. 1 (1996), No. 16, 319–393 (electronic).
- RØ08 Röndigs, O., and Østvær, P.A. *Modules over motivic cohomology*. Adv. Math. 219 (2008), 689–727.
- Su91 Suslin, A. *K-theory and K-cohomology of certain group varieties*. Algebraic K-theory, 53–74, Adv. Soviet Math., 4, Amer. Math. Soc., Providence, RI, 1991.
- Su03 Suslin, A. *On the Grayson spectral sequence*. Proc. Steklov Inst. Math. (2003), no. 2 (241), 202–237.
- Vo98 Voevodsky, V. A^1 -homotopy theory. Proc. of the ICM, Vol.1, Berlin, 1998, Documenta Mathematica, 1998, Extra Vol. I, 579–604 (electronic).
- Vo02a Voevodsky, V. *Open problems in the motivic stable homotopy theory I*. Motives, Polylogarithms and Hodge Theory I, 2002, 3–34.

MOTIVIC SPECTRAL SEQUENCE AND STEENROD OPERATIONS

- Vo02b Voevodsky, V. *A possible new approach to the motivic spectral sequence for algebraic K-theory.* Contemp. Math. vol. 293 (2002), 371–379.
- Vo03 Voevodsky, V. *Reduced power operations in motivic cohomology.* Publ. Math. Inst. Hautes Etudes Sci. No. 98 (2003), 1–57.
- Vo10 Voevodsky, V. *Motivic Eilenberg-Mac Lane spaces.* Publ. Math. Inst. Hautes Etudes Sci. No. 112 (2010), 1–99.
- VSF00 Voevodsky, V., Suslin, A., and Friedlander, E. *Cycles, transfers, and motivic homology theories.* Annals of Mathematics Studies, 143, Princeton University Press, Princeton, NJ, 2000
- We Weibel, Ch. www.math.rutgers.edu/~weibel/Kbook.html

Serge Yagunov yagunov@gmail.com

Steklov Mathematical Institute (St.Petersburg), Fontanka, 27, St.Petersburg, 191023, RUSSIA
and Max-Planck-Institut für Mathematik, Vivatsgasse, 7, 53111, Bonn, GERMANY