2

On Direct Integration for Mirror Curves of Genus Two and an Almost Meromorphic Siegel Modular Form

Albrecht Klemm, Maximilian Poretschkin, Thorsten Schimannek, and Martin Westerholt-Raum

Abstract

This work considers aspects of almost holomorphic and meromorphic Siegel modular forms from the perspective of physics and mathematics. The first part is concerned with (refined) topological string theory and the direct integration of the holomorphic anomaly equations. Here, a central object to compute higher genus amplitudes, which serve as the generating functions of various enumerative invariants, is provided by the so-called propagator. We derive a universal expression for the propagator for geometries that have mirror curves of genus two which is given by the derivative of the logarithm of Igusa's cusp form χ_{10} . In addition, we illustrate our findings by solving the refined topological string on the resolutions of the three toric orbifolds $\mathbb{C}^3/\mathbb{Z}_3$, $\mathbb{C}^3/\mathbb{Z}_6$.

In the second part, we give explicit expressions for lowering and raising operators on Siegel modular forms, and define almost holomorphic Siegel modular forms based on them. Extending the theory of Fourier-Jacobi expansions to almost holomorphic Siegel modular forms and building up on recent work by Pitale, Saha, and Schmidt, we can show that there is no analogue of the almost holomorphic elliptic Eisenstein series \tilde{E}_2 . In the case of genus 2, we provide an almost meromorphic substitute for it. This, in particular, leads us to a generalization of Ramanujan's differential equation for \tilde{E}_2 .

The two parts are intertwined by the observation that the meromorphic analogue of \tilde{E}_2 coincides with the physical propagator. In addition, the generalized Ramanujan identities match precisely the physical consistency conditions that need to be imposed on the propagator.

Contents

Introduction and Summary

1	introduction and Summary	2
2	Calculating Refined BPS Invariants	6
	2.1 The refined A-model	6
	2.2 The refined B-model	9
	2.3 The refined holomorphic anomaly equations	13
	2.4 Background independence of the topological string partition function and modularity	15
2	Invariants of Piomann surfaces and Their Medular Properties	16
3	invariants of Memanin surfaces and Their Modular Properties	10
3	3.1 Invariants of genus one curves	10 16
3	3.1 Invariants of genus one curves	16 16
3	3.1 Invariants of genus one curves 3.2 Invariants of genus two curves Direct Integration Algorithms based on Normal Forms of Mirror Curves	16 16 16
3	3.1 Invariants of genus one curves	16 16 16 18 18

5	Examples	22					
	5.1 Solving the Topological String on local \mathbb{P}^2	23					
	5.2 Solving the Topological String on $\mathbb{C}^3/\mathbb{Z}_5$	24					
	5.3 Solving the Topological String on $\mathbb{C}^3/\mathbb{Z}_6$	30					
6	Almost Holomorphic Siegel Modular Forms	34					
	6.1 Preliminaries	34					
	6.2 Differential operators for Siegel modular forms	35					
	6.3 Almost holomorphic functions	41					
	6.4 Definition of almost holomorphic Siegel modular forms	41					
7	Fourier-Jacobi expansions	42					
	7.1 Preliminaries	42					
	7.2 Differential operators for genus 1 Jacobi forms	44					
	7.3 Almost holomorphic functions (genus 1)	44					
	7.4 Almost holomorphic Jacobi forms (genus 1)	45					
	7.5 Covariant operators from vector valued weights (genus 1)	45					
		45					
8	Classification of Almost Holomorphic Siegel Modular Forms	46					
	8.1 The non-commutative algebra of Siegel modular forms	47					
	8.2 Hyper-algebras	47					
	8.3 The differential graded hyper-algebra of almost holomorphic slegel modular forms	48					
		43					
9	A meromorphic replacement for E_2	50					
	9.1 An almost meromorphic Siegel modular form	50					
	9.2 Fourier coefficients of f_{RS}	52					
Re	eferences	53					
Α	The Ambiguities for $\mathbb{C}^3/\mathbb{Z}_5$	56					
В	The Ambiguities for $\mathbb{C}^3/\mathbb{Z}_6$	58					
C	Siegel modular forms of genus two	58					
D	Refined GV invariants for $\mathbb{C}^3/\mathbb{Z}_5$	61					
Е	Refined GV invariants for $\mathbb{C}^3/\mathbb{Z}_6$	64					
F	Fourier coefficients of Siegel modular forms 65						

1 Introduction and Summary

The topological string theory on non-compact Calabi-Yau geometries M provides exact moduli dependent amplitudes in the N = 2 low energy effective action of type II string theory on these backgrounds. Expanded in flat coordinates for the moduli, these exact amplitudes are generating functions of related symplectic invariants, namely the Gromov-Witten-, Donaldson-Thomas- or Pandharipande-Thomas invariants. From the physics point of view the most natural invariants are the actual numbers $N_{j_L,j_R}^{\beta} \in \mathbb{N}$ of refined BPS states with a given representation (j_L, j_R) of the 5d little group and the charge $\beta \in H_2(M, \mathbb{Z})$. Non-compact Calabi-Yau geometries exhibit typically an $U(1)_{\mathscr{R}}$ isometry, which is most obvious in the toric cases. This $U(1)_{\mathscr{R}}$ symmetry allows a motivic decomposition of the Pandharipande-Thomas moduli space of stable pairs and a mathematical definition of the N_{j_L,j_R}^{β} [CKK14].

Topological string theory on non-compact Calabi-Yau spaces exhibits mirror symmetry and is related by string-gauge theory duality to three dimensional Chern-Simons theory [Wit95] and matrix models [AKMV04]. This makes A-model localization techniques [KZ01], B-model approaches, see e.g. [HKR08], and large N-duality methods available for its solutions. The large N Chern-Simons theory leads to the topological vertex [AKMV05]. The Chern-Simons theory large N expansion can be seen as mirror dual to a matrix model large N expansion [AKMV04]. The Ward identities of the matrix models leads to an iterative solution of the topological string in the genus [Mar08],[BKMP09]. Recently possible non-perturbative completions of the topological string were studied using a free Fermi-gas approach [MP12] and quantum mechanical systems [GHM14]. This led to a profound connection of topological string theory to operator analysis [KM15].

The full moduli dependence of the closed amplitudes $F^{(g)}$ is most efficiently studied in the Bmodel using modularity and the holomorphic anomaly equation of BCOV [BCOV94]. For topological strings the $F^{(g)}$ can be compared at special points in the moduli space with the A-model localization computation e.g. of Gromov-Witten invariants. A first check of mirror symmetry at higher genus was performed in [KZ01] for the non-compact toric $\mathcal{O}(-3) \rightarrow \mathbb{P}^2$ Calabi-Yau space. The BCOV equations can be generalized to the refined holomorphic anomaly equations (2.6) [HK12][KW11]¹, which can be used to compute the refined amplitudes $F^{(n,g)}$. The corresponding check of refined mirror symmetry for topological strings was done in [CKK14] on the same geometry $\mathcal{O}(-3) \rightarrow \mathbb{P}^2$.

The B-model of a wide class of local Calabi-Yau geometries is defined by a mirror geometry, whose data are a family of Riemann surfaces $\mathcal{C}_{\mathfrak{g}}$ of genus \mathfrak{g} and a meromorphic differential λ . These data ($\mathcal{C}_{\mathfrak{g}}, \lambda$) define also N = 2 Seiberg-Witten theories, where $\mathcal{C}_{\mathfrak{g}}$ is known as Seiberg-Witten curve and λ as Seiberg-Witten differential [SW94], and matrix models [Mar08],[BKMP09] with algebraic spectral curve $\mathcal{C}_{\mathfrak{g}}$. In this case λ defines the filling fractions. So far the B-model approach using modularity properties has been developed for $\mathfrak{g} = 1$ starting with [HK07]. Here we will extend the formalism to $\mathfrak{g} = 2$. This extension applies to the calculation of closed amplitudes in all the settings mentioned above.

The occurrence of elliptic quasi-modular- and almost holomorphic modular forms [Shi86] in the solution of the holomorphic anomaly equation was noticed in [HK07]. In particular the identification of the propagator

$$S^{tt} = c_{\tau}^{t} \frac{1}{2\pi i} \frac{\partial}{\partial \tau} F^{(0,1)} = \frac{1}{2\pi i} \frac{\partial}{\partial \tau} c_{\tau}^{t} \left(\log(y) + 4\log(|\eta||f_{1}|) \right) = \frac{c_{\tau}^{t}}{12} \widetilde{E}_{2} + G_{2}$$
(1.1)

with the almost-holomorphic Eisenstein series $\tilde{E}_2 = E_2 - \frac{3}{\pi y}$ up to a meromorphic ambiguous piece G_2 was found ². This was in the context of N = 2 Seiberg Witten theories of rank one on \mathbb{R}^4 , which do emerge in the field theory limit of Type II theories on non-compact Calabi-Yau spaces, whose mirror geometry is a family of elliptic curves $\mathscr{C}_{\mathfrak{g}=1}$. Here η is the Dedekind eta-function, $\tau \in \mathbb{H}^{(\mathfrak{g}=1)}$ is in the Siegel (Poincaré) upper half space (plane) and we denote³ $\tau = x + iy$. c_{τ}^t is a constant whose geometrical significance is explained following eq. (1.3). Further $f_1(z)$, a rational function of the absolute modular invariant z^4 of the family \mathscr{C}_1 , is the holomorphic ambiguity at genus one. G_2 is a congruent subgroup and often a meromorphic weight 2 form of Γ does not exits, which fixes $f_1 = 1$. In any case we may set $f_1 = 1$ and hence G_2 to zero in the definition of the propagator. This merely modifies the form of the higher genus holomorphic or modular ambiguity. It was argued in [HK07] that the latter can be fixed by regularity and the *conifold gap* for the unrefined theory. In [HKR08] this fact has been more carefully established.

¹The one of [KW11] is correct after a shift.

²It is also implicit in the invariant N = 4 sector of the $\mathbb{Z}_3 \otimes \mathbb{Z}_3$ orbifold example in [BCOV94].

³Due to a conflict with other parameters we use in the physical part also explicitly $\operatorname{Re}(\tau)$ and $\operatorname{Im}(\tau)$ for x and y.

⁴It is traditionally called $u = \frac{1}{2} \text{Tr} \Phi^2$ for Seiberg Witten curves.

Historically almost holomorphic (elliptic) modular forms were defined by Shimura [Shi86], who called them "nearly holomorphic". The occurrence of the regularized quasi modular Eisenstein series E_2 in topological string theory can be traced back to the world-sheet contact terms in the derivation of the holomorphic anomaly equation [BCOV94]. Related occurrences of E_2 due to space contact terms in Seiberg-Witten theory on compact target spaces, where one integrates over the phase space, were noted in [MW97] and in 2d QCD on the torus in [Dij95]. The occurrence of E_2 in the latter case was related systematically to almost holomorphic forms in [KZ95], where the isomorphism of the ring of quasi modular and almost holomorphic modular forms, the closures of these rings under the Serre- and Maass derivative respectively and the relation to weak Jacobiforms⁵ was explained. Quasi modular forms have Fourier expansions. Fourier coefficients attached to quasi modular forms are not only counting refined BPS states, but they have been connected to a vast variety of combinatorial and geometric quantities – see, for example, [BO00] and papers citing it. In many cases no recursive holomorphic anomaly is known. E.g. for the 2d QCD example it was argued in [Rud94] that such a recursion does not exist.

The general structure on almost holomorphic modular forms, which is particularly useful when holomorphic anomaly equations govern the theory, is as follows: The non-holomorphic Maass derivative acts as raising operator R on weight k modular forms as $\partial_{\tau} - \frac{ki}{2}y^{-1}$. In addition, there is the lowering operator $L = y^2 \partial_{\overline{\tau}}$, which acts trivially on quasi modular forms, but non-trivially on almost holomorphic ones. The commutation relation $[L, R] = \frac{-k}{4}$ is crucial when proving that \widetilde{E}_2 is essentially the only proper quasi modular form that does not arise from differentiating other ones.

This structure allows for the *direct integration* of the holomorphic anomaly equations (2.6) and an effective way to determine the $F^{(g,n)}$ for mirror curves $\mathscr{C}_{g=1}$ of genus one [ABK08][GKMW07]. The covariant derivative in the holomorphic anomaly equation (2.6) becomes the Maass derivative after changing from the flat coordinate *t* to the τ coordinate using $\frac{dt}{d\tau} = C_{ttt}^{-1}$. The fact that the Maass derivative closes on the generators of the ring of holomorphic forms by the Ramanujan identities, e.g. if the modular group of the family is $\Gamma_0 = PSL(2, \mathbb{Z})$ they read

$$\frac{1}{2\pi i}\partial_{\tau} \widetilde{E}_2 = \frac{1}{12} (\widetilde{E}_2^2 - E_4), \quad \frac{1}{2\pi i}\partial_{\tau} E_4 = \frac{1}{3} (E_2 E_4 - E_6), \quad \frac{1}{2\pi i}\partial_{\tau} E_6 = \frac{1}{2} (E_2 E_6 - E_4^2), \quad (1.2)$$

implies that the r.h.s. of (2.6) is a polynomial in \tilde{E}_2 , with coefficients that are rational functions in the absolute modular invariant z. The uniqueness of the an-holomorphic \tilde{E}_2 implies further that the an-holomorphic derivative on the l.h.s of the holomorphic anomaly equation (2.6) can be replaced by a derivative by \tilde{E}_2 . Hence the recursive equations (2.6) can be solved by integration w.r.t. to \tilde{E}_2 up to an integration constant. This constant in \tilde{E}_2 is another rational function in z called the holomorphic or modular ambiguity. In the refined case it can also be completly determined from regularity requirements and a refined gap condition [KW11] [HK12][HKPK13]. Note that $F^{(g)}$ or $F^{(g,n)}$ are meromorphic functions and in fact it is convenient to define a meromorphic almost holomorphic generator (4.5).

The purpose of this paper is to extend this analysis from families of elliptic mirror curves $\mathscr{C}_{g=1}$ to families of mirror curves of genus two $\mathscr{C}_{g=2}$. A key observation is that in the generalization of (1.1) the Dedekind η -function is replaced by the Igusa cusp form ⁶ χ_{10} of weight 10, so that the propagators becomes

$$S^{ij} = C^{i}_{p} C^{j}_{q} \frac{1}{2\pi i} \frac{\partial}{\partial \tau_{pq}} F^{(0,1)}_{mod} = \frac{1}{2\pi i} \frac{1}{10} \left(\partial_{\tau_{pq}} \log(\chi_{10}) + \frac{5}{i} (\mathrm{Im}\tau)^{-1}_{pq} \right) C^{i}_{p} C^{j}_{q}.$$
(1.3)

Here we set in $F_{mod}^{(0,1)}$ the holomorphic ambiguity $f_1 = 1$. Of course this corresponds just to choice of the ambiguity in the propagator. In this universal relation the indices *i*, *j* refer to the flat coor-

⁵Based on [KMW12] quite spectacular all genus results have been obtained in [HKK15] for compact elliptical fibered Calabi-Yau 3 folds using this relation to weak Jacobi-forms.

⁶There are two classical normalizations of the weight 10 genus two cusp form. The Maass lift $\phi_{10}^{(2)}$ used in section nine is related to it by $4\chi_{10} = \phi_{10}^{(2)}$.

dinates as in (1.1) and the relative coefficients of the terms on the r.h.s. are fixed by the weight. The matrix C_p^i is the intersection matrix of the homology basis of a genus two surface, which is in general different from the identity matrix. In fact, for toric geometries, the non-standard intersection form C_p^i is determined by the intersection pairing in the even-dimensional homology of the A-model geometry due to mirror symmetry. In addition, we show that the physical consistency conditions on the propagator coincide with generalized Ramanujan identities which are derived in the mathematical half of this paper. In the genus 2 case one has due to the Riemann Roch theorem a normal form with a corresponding theory of classifying invariants, the Igusa invariants, which replace the j-function of the elliptic case. The Fourier expansion of these invariants allows to extract the period matrix τ given the algebraic normal form of the Riemann surface, without any further calculation.

Even though Shimura studied nearly holomorphic modular forms in full generality the theory of quasi Siegel modular forms has not yet been developed. We fill this gap in this paper, and reveal several interesting, possibly unexpected phenomena. The theory of Siegel differential operators is well-known, but it has not yet been made explicit. Section 6 contains expressions for raising (6.13) and lowering (6.6) operators, as well as their commutation relations (Lemma 6.11 and 6.12).

Fourier Jacobi expansions are important tools, when investigating Siegel modular forms. The theory of Jacobi forms which arise from vector valued Siegel modular forms was initiated recently in [IK11], and we extend it to the case of genus 2 almost holomorphic Siegel modular forms. Furthermore, we set up a framework for Jacobi forms that arise from vector valued Siegel modular forms of arbitrary genus. We do not characterize them as precisely as in the genus 2 case; rather this will be the theme of a sequel to this paper.

In Section 8, we classify almost holomorphic Siegel modular forms of genus 2, and find that every proper quasi Siegel modular form arises from differentiating other ones. We thus complete the independent classification by Pitale, Saha, and Schmidt [PSS15], which exhausted all almost holomorphic modular forms except for those of weight det³sym^l. As opposed to the case of elliptic modular forms constant modular forms do not occur in the image of the lowering operator. This insight naturally leads us to Section 9, in which we study the genus 2 case in detail. We introduce almost meromorphic Siegel modular forms with determined singular locus, which comprise the inverse of Igusa's χ_{10} [Igu62]. All of a sudden, we observe behavior very similar to the genus 1 case: There is an almost meromorphic Siegel modular forms cannot be obtained as a derivative of any meromorphic Siegel modular forms cannot be obtained as a derivative of any meromorphic Siegel modular forms cannot be obtained as a derivative of any meromorphic Siegel modular forms cannot be obtained as a derivative of any meromorphic Siegel modular forms (2). In Proposition 9.3 we provide an analogue of the Ramanujan differential equation for $S^{(2)}$.

Increasing the genus of the mirror curve is e.g. very natural for the mirror curves of resolved orbifolds for the A-model geometry. Consider e.g. the $\mathbb{C}^3/\mathbb{Z}_3$ orbifold. Resolving the singularity leads to $\mathcal{O}(-3) \to \mathbb{P}^2$ geometry. The Kähler parameter of the \mathbb{P}^2 is mapped to the one modulus of the genus one mirror geometry. If one considers the $\mathbb{C}^3/\mathbb{Z}_5$ orbifold the mirror will have $\mathfrak{g} = 2$. More precisely, it is a special two parameter family of genus two curves, where the two parameters correspond to the two blow up divisors of the $\mathbb{C}^3/\mathbb{Z}_5$ blowup geometry. These blow ups have toric descriptions and the genus g of the mirror curve is given by the interior points of the 2d toric diagram, which represents the trace of the non-compact 3d cone that represents M. It is not directly related to the number of (Kähler) parameters n_k , which is given by the total number of points in the 2d toric diagram minus 3. The inner points correspond to dynamical fields of the 4d gauge theory, while the rest of n_k parameters correspond to mass or non-dynamical parameters. The latter are given by the non-vanishing residua of the meromorphic form λ . Their number is not directly related to \mathfrak{q} , even though by the Riemann Roch theorem the latter determines an upper bound on the number of non dynamical parameters. An easy example, where such non-dynamical fields arise in the orbifold construction with a genus two mirror curve is the $\mathbb{C}^3/\mathbb{Z}_6$ orbifold. Our main computational contribution is to solve the $\mathbb{C}^3/\mathbb{Z}_5$ and the $\mathbb{C}^3/\mathbb{Z}_6$ cases to n+g=3 and n+g=1 respectively, checking the general formalism in great detail.

As reviewed in [BGHZ08], Siegel modular forms can be constructed from products of θ functions

$$\theta[\epsilon] = \sum_{m \in \mathbb{Z}^g} e^{2\pi i \left[(m + \epsilon'/2)^t \tau (m + \epsilon'/2) + (m + \epsilon'/2) \epsilon''/2 \right]},\tag{1.4}$$

where $\epsilon = \begin{pmatrix} \epsilon' \\ \epsilon'' \end{pmatrix}$ with $\epsilon', \epsilon'' \in \{0, 1\}^{\mathfrak{g}}$ encoding the spin structure. The simplest way is to raise the product of the $2^{\mathfrak{g}-1}(2^{\mathfrak{g}-1}+1)\theta$ -functions with *even spin structure* to a power so that no phase transformation occurs. E.g. for $\mathfrak{g} = 1$: $2^{\mathfrak{g}}\eta^{24} = \left(\theta \begin{bmatrix} 0 \\ 0 \end{bmatrix} \theta \begin{bmatrix} 0 \\ 1 \end{bmatrix} \theta \begin{bmatrix} 1 \\ 0 \end{bmatrix}\right)^{\mathfrak{g}}$, for $\mathfrak{g} = 2$: $2^{14}\chi_{10} = -\prod_{\epsilon \text{ even }} \theta[\epsilon]^2$, while for $\mathfrak{g} = 3$ the product of the 36 even theta functions defines a Siegel cusp form of weight 18 under $\operatorname{Sp}_6(\mathbb{Z})$. It is therefore natural to speculate that (1.1,1.3) get replaced, at least for genus 3, using the corresponding canonical cusp form and adapting the coefficients to the corresponding weight.

Acknowledgments It is a pleasure to thank Mirjam Cvetic, Hans Jockers, Ameya Pitale, and Don Zagier for discussions. AK thanks for support by KL 2271/1-1 and NSF DMS-11-01089. The work of MP has been supported by a scholarship of the Deutsche Telekom Stiftung and by the grant DE-SC0007901. He also thanks the BCTP for hospitality. TS is supported by a scholarship by the graduate school BCGS. MWR thanks the Max Planck Institute for Mathematics for their hospitality.

2 Calculating Refined BPS Invariants

In this section we provide some necessary background material. We start with a review of the notion of refined BPS numbers in paragraph 2.1. Afterwards, we present the mirror symmetry for toric A-model geometries. We move on with a review of (refined) B-model computational techniques in 2.2. In particular, we rigorously discuss how the non-symplectic basis of the mirror curve modifies the well-known relations that apply to the case of a standard homology basis. Finally, we end by pointing out the quasi- respectively almost modular structure of the generating functions of the (un-)refined BPS invariants in paragraph 2.4.

§2.1 The refined A-model. In this subsection we provide some necessary background material for the discussion of refined BPS invariants. These can be meaningfully defined within five-dimensional theories with eight supercharges and an additional $U(1)_{\mathscr{R}}$ symmetry. Such theories are naturally engineered by M-theory compactifications on toric Calabi-Yau threefolds *X*. Physically, these refined BPS invariants count particles arising from M2-branes that wrap holomorphic curves within *X*. Besides the homology class $\beta \in H_2(X, \mathbb{Z})$, these particles are in addition classified by their spins (j_L, j_R) under the five-dimensional little group $SU(2)_L \times SU(2)_R$. The multiplicities N_{j_L, j_R}^{β} are called the refined BPS invariants. These are counted by a five-dimensional index which reads

$$Z_{BPS}(\epsilon_1, \epsilon_2) = \operatorname{Tr}(-1)^{2(J_L + J_R)} \exp\left[-\left((\epsilon_1 - \epsilon_2)J_L^3 + (\epsilon_1 + \epsilon_2)J_R^3 + (\epsilon_1 + \epsilon_2)J_{\mathscr{R}}^3 + \beta H\right)\right].$$
(2.1)

Here we have denoted the respective Cartan generators by J_* . It is crucial to note that the invariance under deformations follows from the twist of the generator J_R by $J_{\mathscr{R}}$. It turns out to be convenient to re-organize the refined BPS invariants in the so-called refined free energy that takes the form

$$\mathscr{F}(\epsilon_1,\epsilon_2,t) = \sum_{\substack{j_L,j_R=0\\k=1}}^{\infty} (-1)^{2(j_L+j_R)} \frac{N_{j_L,j_R}^{\beta}}{k} \frac{\sum_{m_L=-j_L}^{j_L} q_L^{km_L}}{2\sinh\left(\frac{k\epsilon_1}{2}\right)} \frac{\sum_{m_R=-j_R}^{j_R} q_R^{km_R}}{2\sinh\left(\frac{k\epsilon_2}{2}\right)} Q^{k\beta}, \qquad Q^{\beta} = e^{-\beta \cdot t}.$$
(2.2)

Here t_i denote the Kähler moduli and $q_{L/R} = e^{\epsilon_1 \pm \epsilon_2}$. There is a second set of invariants, denoted by n_{g_L,g_R}^{β} which are related to the refined BPS numbers by a change of spin basis given as

$$I_{*}^{n} = \left(2 \left[0\right]_{*} + \left[\frac{1}{2}\right]_{*}\right)^{\otimes n}, \qquad * \in \{L, R\},$$
(2.3)

which is easily recognized as the spin content of the Lefshetz decomposition of an *n*-torus. In terms of this new basis, the n_{g_L,g_R}^{β} are explicitly given as

$$\sum_{g_L,g_R} n_{g_L,g_R}^{\beta} I_L^{g_L} \otimes I_R^{g_R} = \sum_{j_L,j_R} N_{j_L j_R}^{\beta} \left[\frac{j_L}{2} \right] \otimes \left[\frac{j_R}{2} \right].$$
(2.4)

While both sets of invariants take values in the integers, the refined BPS invariants N_{j_L,j_R}^{β} are in addition non-negative. It is also useful to rewrite the refined free energy as

$$\mathscr{F}(\epsilon_1,\epsilon_2,t) = \sum_{n,g=0}^{+\infty} (\epsilon_1 + \epsilon_2)^{2n} (\epsilon_1 \epsilon_2)^{g-1} \mathscr{F}^{(n,g)}(t).$$
(2.5)

One recognizes the genus expansion of the unrefined topological string in the limit $\epsilon_1 = -\epsilon_2$. The $\mathscr{F}^{(n,g)}(t)$ can be recursively determined using the refined holomorphic anomaly equations

$$\bar{\partial}_{\bar{i}}F^{(n,g)} = \frac{1}{2}\bar{C}_{\bar{i}}^{jk} \left(D_j D_k F^{(n,g-1)} + \sum_{m,h}' D_j F^{(m,h)} D_k F^{(n-m,g-h)} \right), \quad n+g>1,$$
(2.6)

which are discussed in more detail in section 2.3 and taking the holomorphic limit

$$\mathscr{F}^{(n,g)}(t) = \lim_{\bar{t} \to i\infty} F^{(n,g)}(t,\bar{t}).$$

$$(2.7)$$

Mathematically, the formulation of refined BPS invariants is based on the notion of stable pairs. These are defined as a pure sheaf \mathscr{F} of complex dimension one with

$$\operatorname{ch}_2(\mathscr{F}) = \beta, \qquad \chi(\mathscr{F}) = n,$$
(2.8)

where $\beta \in H_2(X, \mathbb{Z})$ gets identified with the D2-brane charge and *n* determines the number of D0branes. In addition, one requires a section $s \in H^0(\mathcal{F})$ that generates \mathcal{F} outside a finite set of points *Q*. All this information can be organized within an exact sequence

$$0 \longrightarrow I_C \longrightarrow \mathscr{O}_X \xrightarrow{s} \mathscr{F} \longrightarrow Q \longrightarrow 0,$$

where I_C denotes the annihilation ideal of a curve *C* representing β . The moduli space of stable pairs with charges β , *n* is denoted $P_n(X, \beta)$ and carries a symmetric and perfect obstruction theory. This implies the existence of a virtual fundamental class $[P_n(X, \beta)]^{\text{vir}}$ which can be integrated to a number, the Pandharipande Thomas invariant

$$P_{n,\beta} = \int_{P_n(X,\beta)} \left[P_n(X,\beta) \right]^{\text{vir}}.$$
(2.9)

These are organized in the following generating function

$$Z_{PT} = \sum_{n,\beta} P_{n,\beta} q^n Q^{\beta}.$$
(2.10)

The latter is conjectured, and in the toric case proven, to coincide with the generating function of disconnected Gromov-Witten invariants

$$Z_{GW} = \exp\left(\mathscr{F}_{GW}(\lambda, Q)\right), \qquad \mathscr{F}_{GW}(\lambda, Q) = \sum_{\beta \neq 0} \sum_{g} \lambda^{2g-2} Q^{\beta}, \tag{2.11}$$

provided one identifies $q = -e^{i\lambda}$.

Only recently the notion of refinement on the level of PT-invariants has been established in [CKK14]. It requires an \mathbb{C}^* -action on $P_n(X,\beta)$ with finitely many isolated fixed points p and gives

rise to a virtual Bialynicki-Birula cell decomposition. More precisely, for any fixed point p one considers the decomposition of the tangent space into eigenspaces of positive and negative characters χ of the \mathbb{C}^* -action

$$T_p X = \bigoplus_{\chi \in X(\mathbb{C}^*)} T_p^{\chi} X = \bigoplus_{\substack{\chi > 0 \\ T_p^+ X}} T_p^{\chi} X \oplus \bigoplus_{\substack{\chi < 0 \\ T_p^- X}} T_p^{\chi} X,$$
(2.12)

and defines $d_p^{\pm} = \dim \left(T_p^{\pm} X \right)$. This construction allows to define the virtual motive

$$\left[P_n(X,\beta)\right]^{\text{vir}} = \sum_{p \in P_n(X,\beta)^{\mathbb{C}^*}} \left(-\mathbb{L}^{-\frac{1}{2}}\right)^{d_p^+ - d_p^-},$$
(2.13)

where \mathbb{L} denotes the absolute motive of \mathbb{C} . Interpreting the right-hand side as a direct sum of irreducible SU(2) characters, one ends up with the following decomposition into refined PT invariants

$$[P_{1-p_a}(X,\beta)]^{\nu ir} = \sum_{j_R} (-1)^{2j_R} N^{\beta}_{p_a/2,j_R} [j_R]$$
(2.14)

where $[j_R]$ denotes the representations associated to the respective characters. In particular the refined PT partition function

$$Z_{PT}^{ref} = \prod_{\beta, j_L, j_R} \prod_{m_{L/R} = -j_{L/R}}^{j_{L/R}} \prod_{m=1}^{\infty} \prod_{j=0}^{m-1} \left(1 - \mathbb{L}^{-m/2 + 1/2 + j - m_R} (-q)^{m-2m_L} Q^{\beta} \right)^{(-1)^{2(j_L + j_R)} N_{j_L, j_R}^{\beta}},$$
(2.15)

coincides with the refined string partition function

$$Z^{ref} = \exp\left[\mathscr{F}(\epsilon_1, \epsilon_2, t)\right],\tag{2.16}$$

where $\mathscr{F}(\epsilon_1, \epsilon_2, t)$ denotes the refined free energy defined in (2.2).

§2.1.1 Local A-model geometries. A toric (i.e. in particular non-compact) Calabi-Yau three-fold M is specified as the quotient of some open subset of \mathbb{C}^{k+3} by a group $G = (\mathbb{C}^*)^k \times C$, where the last factor denotes a possible discrete part

$$M = \left(\mathbb{C}^{k+3} - Z \right) / G.$$
 (2.17)

The continuous part of the group action is specified by *k* charge vectors $Q^{\alpha} \in \mathbb{Z}^{k+3}$ satisfying the Calabi-Yau condition

$$\sum_{i=1}^{k+3} Q_i^{\alpha} = 0, \qquad (2.18)$$

that act on the open subset as

$$x_i \mapsto \mu_{\alpha}^{Q_i^a} x_i \,. \tag{2.19}$$

Finally, Z denotes the Stanley-Reisner ideal which is the fix point set of the group action (2.19) and needs to be substracted in order to make M being well-defined as a variety. All this information is encoded in the so-called toric diagram.

$$\begin{pmatrix}
a_{1} \quad b_{1} \quad 1 & Q_{1}^{1} & \dots & Q_{1}^{n} \leftarrow D_{1} \\
\vdots & & \vdots & & \\
a_{k} \quad b_{k} \quad 1 & Q_{k}^{1} & \dots & Q_{k}^{n} \leftarrow D_{k} \\
& & \uparrow & \uparrow & \\
& & & C^{1} & C^{n} &
\end{pmatrix}$$
(2.20)

Here $D_k = \{x_k = 0\}$ denotes the divisor associated to the coordinate x_k corresponding to the ray $e_k = (a_k \ b_k \ 1)$ within the toric diagram. The C^{α} denote a basis of the Mori cone and constitute a

distinguished basis of $H_2(X, \mathbb{Z})$. In particular, the intersection number between the divisor D_i and the curve C^{α} is given by

$$Q_i^{\alpha} = C^{\alpha} \cdot D_i \,. \tag{2.21}$$

From a physics point of view, the toric variety (2.17) has an interpretation as the vacuum field configuration of a two-dimensional Sigma-model. Here the coordinates x_i get identified as the vacuum expectation values of the scalars of the chiral multiplets having charge vectors Q_i under the gauge group $U(1)^k$. The vacuum field configuration arises from the D-term constraints

$$D^{\alpha} = \sum_{i=1}^{k+3} Q_i^{\alpha} |x_i|^2 = r^{\alpha}.$$
 (2.22)

The inequivalent vacua are obtained by dividing out the group $U(1)^k$. In addition, $r^{\alpha} = \int_{C^{\alpha}} J + iB$ correspond to the complexified Kähler parameters, where *J* denotes the Kähler class and *B* is the Kalb Ramond field.

§2.2 The refined B-model.

§2.2.1 Local B-model geometries. The local B-model geometry can also be constructed in terms of the toric charge vectors using the Hori-Vafa method. For this one introduces two \mathbb{C} -valued coordinates w^+ , w^- as well as the homogeneous coordinates $x_i := e^{y_i}$ which are subject to the rescaling

$$x_i \mapsto \lambda x_i, \qquad \lambda \in \mathbb{C}^*,$$
 (2.23)

and are further constrained by

$$(-1)^{Q_0^{\alpha}} \prod_{i=1}^{k+3} x_i^{Q_i^{\alpha}} = z_{\alpha} \,. \tag{2.24}$$

The local mirror geometry is then defined by

$$w^+ w^- = H = \sum_{i=1}^{k+3} x_i, \qquad (2.25)$$

where the constraints (2.24) as well as the rescaling relation (2.23) can be used to eliminate all but two (x, y) of the homogeneous coordinates such that the geometry takes the form of a conic bundle over a family of Riemann surfaces $H(x, y, z_{\alpha})$ parametrized by the complex structure moduli z_{α} . The local B-model geometry (2.25) naturally inherits a holomorphic three-form from the ambient space which is given as

$$\Omega = \frac{dHdxdy}{Hxy}.$$
(2.26)

After integrating out the non-compact directions this gets reduced to a meromorphic one-form defined on H

$$\lambda = \frac{\log(y)dx}{x}.$$
(2.27)

As the Riemann surfaces arising in this way are generically non-compact, one does not expect to be able to find a normalizable symplectic basis in general. Instead the most standard form of a basis A^i , B_j of $H_1(H,\mathbb{Z})$ is subject to the intersections

$$A^{i} \cap A^{j} = 0, \qquad B_{i} \cap B_{j} = 0, \qquad A^{i} \cap B_{j} = n_{j}^{i}, \quad n_{j}^{i} \in \mathbb{Z}.$$
 (2.28)

In fact, these intersections can be determined from toric diagram of the A-model geometry, see also the discussion in section 2.2.3. For the following discussion we always make use of a normalizable basis and devote the section 2.2.2 to analyse the modifications in the case of a non-normalizable basis.

One refers to the respective period integrals of λ along this basis as the A- and B-periods respectively, denoting them by t_i and t_D^i respectively. These periods are annihilated by a set of linear differential operators, called the Picard-Fuchs equations

$$D_{\alpha} = \prod_{Q_i^{\alpha} > 0} \partial_{x_i}^{Q_i^{\alpha}} - \prod_{Q_i^{\alpha} < 0} \partial_{x_i}^{-Q_i^{\alpha}}.$$
 (2.29)

The solutions are constructed by the Frobenius method. Starting with the fundamental period [CKYZ99]

$$w_0(\vec{z},\vec{\rho}) = \sum_{\vec{n}} \frac{1}{\Gamma(Q_i^{\alpha}(n^{\alpha} + \rho^{\alpha}) + 1)} ((-1)^{Q_0^{\alpha}} z^{\alpha})^{n^{\alpha} + \rho^{\alpha}}, \qquad (2.30)$$

one obtains a constant solution, as well as a number of linear logarithmic solutions.

$$X^{0} = w_{0}(\vec{z}, 0) = 1, \qquad t_{i} = \frac{1}{2\pi i} \frac{\partial}{\partial \rho^{i}} w_{0}(\vec{z}, \vec{\rho})|_{\rho=0}.$$
(2.31)

Second order derivatives with respect to $\vec{\rho}$ are not solutions to (2.29), but suitable linear combinations ⁷ of them constitute the B-periods t_D^i . The number of these solutions is combinatorically determined by the toric diagram as well. In fact, any inner point gives rise to a single as well as a doubly logarithmic solution (that are dual to each other), while the other points give rise to further linear logarithmic solutions. The first class of solutions corresponds to true complex structure moduli of the Riemann surface, while the second class of solutions is identified with additional residues m_k of the meromorphic one-form that correspond to non-normalizable directions in the complex structure moduli space and therefore are merely parameters of the geometry rather than real moduli. As the residues of the meromorphic one-form give rise to masses of fundamental matter in the Seiberg-Witten theory engineered by this Riemann surface, these parameters are also referred to as mass parameters. From a homological point of view, these moduli correspond to curves that have no compact dual within the geometry, compare also the discussion in 2.2.3. Yet another interpretation is given as isomonodromic deformations. In fact, the periods may undergo non-trivial monodromies when encircling singularities in the complex structure moduli space. As these monodromies have to respect the symplectic form, they have to constitute a subgroup $\Gamma \subset$ $\operatorname{Sp}_{2\mathfrak{g}}(\mathbb{Z})$. This action takes explicitly the form

$$\tilde{t}_i = a_i^j t_j + b_{ij} t_D^j + E_i^k m_k$$
(2.32)

$$\tilde{t}_{D}^{i} = c^{ij}t_{j} + c^{i}_{j}t_{D}^{j} + F^{ik}m_{k}$$
(2.33)

$$\widetilde{m}_k = m_k \qquad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{Sp}_{2\mathfrak{g}}(\mathbb{Z}).$$
(2.34)

The pairs of true periods and their duals are related to the prepotential $F^{(0,0)}$ as

$$t_D^i = \frac{\partial F^{(0,0)}}{\partial t_i}.$$
(2.35)

The period matrix of the mirror geometry $\tau = \{\tau_{ij}\}_{(i,j=1,\dots,g)}$ is related to the prepotential as

$$\tau_{ij} = -\frac{\partial^2}{\partial t_k \partial t_l} F^{(0,0)} \,. \tag{2.36}$$

⁷Clearly, one may also add logarithmic solutions of order zero and one to the double-logarithmic solutions. The latter do not affect the instanton part of the prepotential, but only its classical contribution. Nevertheless, they are crucial in order to obtain a set of solutions that forms an integral basis for the monodromies. See [Hos06] for a conjecture how this basis is obtained. Recently, the choice of basis has been related to the $\hat{\Gamma}$ -classes and the hemisphere partition function, see [CBKL14] for an application to fourfolds and more detailed references.

The third derivatives of the prepotential $F^{(0,0)}$

$$C_{ijk} = \partial_{t_i} \partial_{t_i} \partial_{t_k} F^{(0,0)}, \qquad (2.37)$$

are called the Yukawa couplings. Finally there is the Kähler potential

$$K = \frac{1}{2i} \left(t_i \bar{t}_D^i - \bar{t}_i t_D^i \right).$$
 (2.38)

Note that this form only holds true in the local case, in contrast to the compact case where the right hand side of (2.38) gets dressed by a logarithm. From this one deduces the Kähler metric

$$g_{t_i\bar{t}_j} = \frac{\partial^2 K}{\partial t_i \partial \bar{t}_j} = \mathrm{Im}\tau.$$
(2.39)

§2.2.2 Modifications for non-compact Riemann surfaces. In this subsection we discuss the modifications of the results obtained in the previous section for a canonically normalized basis of the Riemann surface.

We start with a basis (A_i, B^j) of A- and B-cycles of a Riemann surface of genus g. Their intersection matrix defines the symplectic form

$$J_{\mathfrak{g}} = \begin{pmatrix} & \mathbb{1}_{\mathfrak{g}} \\ -\mathbb{1}_{\mathfrak{g}} & \end{pmatrix}, \qquad (2.40)$$

and with respect to this one defines the pair of dual cycles

$$t_i = \int_{A_i} \lambda, \qquad t_D^j = \int_{B^j} \lambda, \tag{2.41}$$

where λ is the meromorphic one-form that is inherited from the Calabi-Yau three-form. As already discussed, t_i and t_D^j provide an integral basis for the monodromy group and are related via

$$t_D^i = \frac{\partial F^{(0,0)}}{\partial t_i} \,. \tag{2.42}$$

In contrast, if there is no normalizable basis, i.e. the A- and B- cycles intersect as

$$J_{\mathfrak{g}}^{\text{non-norm.}} = \begin{pmatrix} C \\ -C \end{pmatrix}, \qquad (2.43)$$

the dual periods are given as non-trivial linear combinations of derivatives of the prepotential [Hos06]

$$t_D^i = C_j^i \frac{\partial F^{(0,0)}}{\partial t_j}, \qquad (2.44)$$

and it is the pair (t_i, t_D^j) that provides an integral basis for the monodromy group.

In order to derive the modification of the period matrix, we note the following behavior under the symplectic transformation

$$\begin{pmatrix} A \\ B \end{pmatrix} \mapsto \begin{pmatrix} C & 0 \\ 0 & C^{-1} \end{pmatrix} \begin{pmatrix} A \\ B \end{pmatrix} = \begin{pmatrix} CA \\ C^{-1}B \end{pmatrix}, \qquad \tau \mapsto C\tau C \quad \text{for} \quad \begin{pmatrix} C & 0 \\ 0 & C^{-1} \end{pmatrix} \in \operatorname{Sp}_{2\mathfrak{g}}(\mathbb{Z}).$$
 (2.45)

Instead of a change from (A, B) to $(CA, C^{-1}B)$, we can interpret (2.45) also as transforming into the basis $(A, C^{-1}B)$ while changing $J_{\mathfrak{g}}$ into $J_{\mathfrak{g}}^{\text{non-norm.}}$ at the same time. I.e. (2.45) changes the canonical symplectic form into a non-canonical form which is characterized by the intersection matrix *C*. Accordingly, the period matrix is connected to the prepotential as

$$\tau_{ij} = -C_i^k C_j^l \frac{\partial^2 F^{(0,0)}}{\partial t_k \partial t_l}.$$
(2.46)

§2.2.3 Mirror symmetry and the mirror map. Mirror symmetry relates the B-model on a three-fold *X* to the A-model on the mirror three-fold *Y*. Mathematically, it is a conjectured equivalence between the bounded derived category of coherent sheaves on *X* and the the derived Fukaya category DFuk(Y, J) with the Kähler form *J* viewed as a symplectic form. This equivalence gives in particular rise to a symplectomorphism

$$mir: K^{c}(X) \longrightarrow H_{3}(Y, \mathbb{Z}).$$

$$(2.47)$$

Here, $K^c(X)$ denotes the Grothendieck or K-theory group of coherent sheaves on *X*. As our main interest is in non-compact three-folds, we have to demand that the latter are compactly supported. As a consequence of the Grothendieck Riemann Roch theorem the Chern character map gives rise to a further symplectomorphism

$$K^{c}(X) \otimes \mathbb{Q} \cong A^{c}_{\text{even}}(X) \otimes \mathbb{Q}, \qquad (2.48)$$

that identifies the pairing on $K^{c}(X)$

$$\chi(R,S) = \int \operatorname{ch}(R^*)\operatorname{ch}(S)\operatorname{td}(X), \qquad (2.49)$$

with the intersection pairing on $A^c_{\text{even}}(X)$. The latter denotes the compactly supported Chow group which coincides with the homology group in case of crepant resolutions of \mathbb{C}^3/G where *G* is an abelian subgroup of SL(3, \mathbb{Z}) which is the concern of the present publication. For this class of geometries the generators of as well as the pairing in the compact K-theory group can be calculated group-theoretically using the McKay correspondence⁸.

Roughly speaking, *mir* identifies A-cycles in the mirror geometry with curves in *X* and B-cycles with divisors such that the natural intersection pairing between curves and divisors on the left side gets identified with that of A- and B-cycles on the right side. To be more precise, we have the following identification, as compactly summarized in table 1. For every generator C^{α} of the Mori cone there is a dual element of the Kähler cone J_k . Also we have a basis of $H_4(X, \mathbb{Z})$ given by divisors D_k . To all these elements we associate the following sheaves [Hos06; OFS02]

$$C_{\alpha} \longrightarrow \mathscr{O}_{C_{\alpha}}(-J_{\alpha}) := \mathscr{O}_{C} \otimes \mathscr{O}(-J_{\alpha}), \tag{2.50}$$

$$D_k \longrightarrow \mathcal{O}_{D_k}(\mathscr{L}_k) := \mathcal{O}_D \otimes \mathscr{L}_k, \quad \mathscr{L}_k = \mathcal{O}_X(-a_{k,1}J_1 \dots - a_{k,l}J_l).$$
(2.51)

Here, $\mathcal{O}_{C_{\alpha}}$ and \mathcal{O}_{D_k} are supported⁹ on C_{α} and D_k . The coefficients in (2.51) are chosen such that

$$\chi\left(\mathcal{O}_{D_i}(\mathcal{L}_j), \mathcal{O}_{D_k}(\mathcal{L}_k)\right) = 0.$$
(2.52)

Finally there is the D0-brane charge which is supported on a skyscraper sheaf \mathcal{O}_p and corresponds to the trivial solution of the Picard-Fuchs equations. By construction, the only non-trivial intersection between the K-theory generators is given by

$$C_{\alpha\beta} = \chi \left(\mathscr{O}_{C_{\alpha}}(-J_{\alpha}), \mathscr{O}_{D_{\beta}}(\mathscr{L}_{\beta}) \right), \qquad (2.53)$$

which corresponds to the intersection between divisors and curves and can directly be read off from (2.20). We end by noting that the matrix *C* defined by (2.53) is in general not regular. This can be traced back to the fact that there are curves which have no compact dual in $H_4(X, \mathbb{Q})$. These are precisely given by the mass parameters discussed in section 2.25.

⁸Strictly speaking, this is only guaranteed for the resolution given by *G*-Hilb.

⁹One way to construct these sheaves is to consider a coherent sheaf S_V on a compact sub-manifold $\iota : V \hookrightarrow X$ and to compute the compact Chern character $ch^c(S) := ch(\iota_*S_V)$ by the Grothendieck Riemann Roch theorem $\iota_*(ch(S_V)td(V)) = ch(\iota_*S_V)td(X)$.

Solution of the PF eqn.	Charge	$H_{\text{even}}(X,\mathbb{Z})$	$K^{c}(X)$	comp. dual
1	D0	$H_0(X,\mathbb{Z})$	\mathcal{O}_p	no
transcendental single log.	D2	$H_2(X,\mathbb{Z})$	$\mathcal{O}_{C^{\alpha}}(-J_{\alpha})$	yes
rational single log.	D2	$H_2(X,\mathbb{Z})$	$\mathcal{O}_{C^{\alpha}}(-J_{\alpha})$	no
double log.	D4	$H_4(X,\mathbb{Z})$	$\mathscr{O}_{D_j}(\mathscr{L}_j)$	yes

Table 1: The elements of the compact K-theory group together with their associated D-brane charges and mirror maps.

In the second part of this subsection we aim to discuss a a second - obviously not unrelated - notion of mirror map which provides an identification of the Kähler moduli in the A-model on Y in terms of the complex structure moduli of the B-model on X. Concretely, one identifies for every normalizable complex structure parameter z_i the logarithmic solution given as

$$t_i = \frac{1}{2\pi i} \log(z_i) + \Sigma(z_j),$$
 (2.54)

where $\Sigma(z_j)$ denotes a power series with the Kähler modulus t_i . The relation with the homological mirror map is obtained by noting that the complex structure moduli are parameterized by $H^{(0,1)}(X, T^{(1,0)}X) \cong H^{(2,1)}(X)$ and the Kähler moduli by $H^{(1,1)}(X)$ and passing to the respective Poincaré dual homology. The transcendental mirror map is finally given by inverting (2.54) in favor of the monodromy invariant variables $Q_i = e^{2\pi i t_i}$. In contrast, moduli corresponding to nonnormalizable directions have rational mirror maps.

§2.3 The refined holomorphic anomaly equations. Refined holomorphic anomaly equations have been proposed in [HK12; HKPK13]. They read

$$\bar{\partial}_{\bar{i}}F^{(n,g)} = \frac{1}{2}\bar{C}_{\bar{i}}^{jk} \left(D_j D_k F^{(n,g-1)} + \sum_{m,h}' D_j F^{(m,h)} D_k F^{(n-m,g-h)} \right), \quad n+g>1.$$
(2.55)

Here the prime signals the omission of (m, h) = (0, 0) and (n, g). The covariant derivatives are with regard to the bundle $T^{(1,0)}X$ and its symmetric powers. More precisely, for a correlation function of k operators they take the form

$$D_t = \partial_t - k\Gamma_t, \qquad (2.56)$$

where Γ denotes the Christoffel symbol of the Kähler metric (2.39). In general, the free energies at genus *g* transform as a section of a line bundle \mathscr{L}^{2g-2} that gives an additional contribution to the connection in (2.56). However, in the local limit, \mathscr{L} becomes trivial and (2.56) is exact.

While the unrefined holomorphic anomaly equations can be rigorously derived, a worldsheet interpretation of the refined holomorphic anomaly equations is so far missing. However, it was conjectured in [HKPK13] that the second deformation parameter corresponds to the insertion of *n* times an operator \mathcal{O} into the worldsheet of genus *g*. Accordingly, the refined free energies take now the form

$$F^{(n,g)} = \int_{\bar{\mathcal{M}}_g} \langle \mathcal{O}^n \prod_{k=1}^{3g-3} \beta^k \bar{\beta}^k \rangle_g dm \wedge d\bar{m}.$$
(2.57)

Thus, the second sum in (2.55) gets an interpretation as the distribution of the *n* points among the degenerated components of the Riemann surface. In particular, it was conjectured that

$$F^{(n+1,0)} = \langle \phi^{(0)}(0)\phi^{(0)}(1)\phi^{(0)}(\infty)\mathcal{O}^n \rangle_{g=0}.$$
(2.58)

This implies in particular that $F^{(1,0)}$ is purely holomorphic such that its most general ansatz reads

$$F^{(1,0)} = \frac{1}{24} \log \left(\Delta \prod_{i} u_{i}^{a_{i}} \prod_{j} m_{j}^{b_{j}} \right),$$
(2.59)

where Δ denotes the discriminant locus of the mirror curve and the constants a_i, b_j are fixed by requiring regularity at infinity and the knowledge of a few BPS numbers. In contrast, $F^{(0,1)}$ corresponds to the unrefined free energy and obeys a holomorphic anomaly equation as well. The most general ansatz for $F^{(0,1)}$ is given in the local limit by

$$F^{(0,1)} = \frac{1}{2} \log \left(\Delta^a \prod_i u_i^{a_i} m_j^{b_j} |g_{i\bar{j}}^{-1}| \right).$$
(2.60)

Here $g_{z_i \bar{z}_j} = \partial_{z_i} \bar{\partial}_{\bar{z}_j} K$ refers to the Kähler metric (2.39) transformed into the coordinates z_i . This can be re-expressed as

$$g_{z_i\bar{z}_j} = \left(\frac{\partial^2 K}{\partial t_k \partial \bar{t}_{\bar{l}}}\right) \frac{\partial t_k}{\partial z_i} \frac{\partial \bar{t}_{\bar{l}}}{\partial \bar{z}_{\bar{j}}} = (\operatorname{Im} \tau_{kl}) G_i^k \bar{G}_{\bar{j}}^{\bar{l}}.$$
(2.61)

The quantity

$$G_i^j = \frac{\partial t_i}{\partial z_j},\tag{2.62}$$

is often referred to as the topological metric in the local and holomorphic limit. Clearly, the factor $|\text{Im}\tau_{ij}|$ is not holomorphic. Accordingly, $F^{(0,1)}$ can be split as

$$F^{(0,1)} = \frac{1}{2} \log(|\mathrm{Im}\tau_{ij}|^{-1}) + \frac{1}{2} \log(|(\bar{G}^{-1})_{\bar{i}}^{\bar{j}}|) + \frac{1}{2} \log\left(\Delta^{a} \prod_{i} u_{i}^{a_{i}} m_{j}^{b_{j}}|(G^{-1})_{i}^{j}|\right)$$
$$= \frac{1}{2} \log(|\mathrm{Im}\tau_{ij}|^{-1}) + \frac{1}{2} \log(|(\bar{G}^{-1})_{\bar{i}}^{\bar{j}}|) + \mathscr{F}^{(0,1)}.$$
(2.63)

As discussed in detail in section 4 for genus one and two, $F^{(0,1)}$ is invariant under modular transformations, but not holomorphic. In contrast, $F^{(0,1)}$ alone is not modular invariant.

Having discussed the antagonistic behavior under modularity respectively holomorphicity of $F^{(0,1)}$, we remark that the ambiguity can be fixed with the help of a few known BPS numbers, analogously to the case of $F^{(1,0)}$. Alternatively, they can also be determined from the behavior of $\mathscr{F}^{(0,1)}$ near singularities, e.g.

$$\lim_{z_i \to 0} \mathscr{F}^{(0,1)} = -\frac{1}{24} \sum_{j=1}^{h^{2,1}} t_j \int_X c_2 J_j.$$
(2.64)

Besides the prepotential $F^{(0,0)}$, the knowledge of the free energies at genus one is sufficient to determine the higher genus free energies $F^{(n,g)}$ using the refined holomorphic anomaly equations up to an integration constant, called the ambiguity. The direct integration method proceeds by calculating the so-called propagators S^{ij} . They can be used to re-express the anti-holomorphic derivative of $F^{(n,g)}$ as

$$\bar{\partial}_{\bar{i}}F^{(n,g)} = C_{\bar{i}}^{jk}\frac{\partial F^{(n,g)}}{\partial S^{jk}},$$
(2.65)

which implies that $F^{(n,g)}$ is a polynomial in S^{ij} of degree 3(g + n) - 3. The propagators are determined from a - in the multi-moduli case overdetermined - set of equations that reads for local Calabi-Yau manifolds as follows

$$D_i S^{kl} = -C_{imn} S^{km} S^{lm} + f_i^{kl}, (2.66)$$

$$\Gamma_{ij}^k = -C_{ijl}S^{kl} + \tilde{f}_i^{kl}, \qquad (2.67)$$

$$\partial_i F^{(0,1)} = \frac{1}{2} C_{ijk} S^{jk} + A_i.$$
(2.68)

In these equations the ambiguities f_i^{kl} , \tilde{f}_i^{kl} , A_i can be solved from the following ansätze

$$\begin{aligned} f_i^{kl} &= \frac{h(z_i)}{\prod_i z_i^{m_i} \Delta^p}, \\ A_i &= \partial_i \big(\tilde{a}_j \log \Delta_j + \tilde{b}_j \log z_j \big). \end{aligned}$$

The first formula applies to both, f_i^{kl} and \tilde{f}_i^{kl} and p is either zero or one while $h(z_i)$ is a polynomial.

Finally, the integration constant in (2.55) can be fixed by matching the refined free energies with a very particular behaviour of the refined free energies at the conifold locus known as the gap condition. It arises from integrating out a particle becoming massless by performing a Schwinger Loop computation, where the coordinate *t* denotes the flat coordinate normal to the singularity

$$\begin{aligned} \mathscr{F}(s,g_{s},t) &= \int_{0}^{\infty} \frac{ds}{s} \frac{\exp(-st)}{4\sinh(s\epsilon_{1}/2)\sinh(s\epsilon_{2}/2)} + \mathscr{O}(t^{0}) \end{aligned} \tag{2.69} \\ &= \left[-\frac{1}{12} + \frac{1}{24}(\epsilon_{1} + \epsilon_{2})^{2}(\epsilon_{1}\epsilon_{2})^{-1} \right] \log(t) \\ &+ \frac{1}{\epsilon_{1}\epsilon_{2}} \sum_{g=0}^{\infty} \frac{(2g-3)!}{t^{2g-2}} \sum_{m=0}^{g} \hat{B}_{2g} \hat{B}_{2g-2m} \epsilon_{1}^{2g-2m} \epsilon_{2}^{2m} + \dots \\ &= \left[-\frac{1}{12} + \frac{1}{24}sg_{s}^{-2} \right] \log(t) + \left[-\frac{1}{240}g_{s}^{2} + \frac{7}{1440}s - \frac{7}{5760}s^{2}g_{s}^{-2} \right] \frac{1}{t^{2}} \\ &+ \left[\frac{1}{1008}g_{s}^{4} - \frac{41}{20160}sg_{s}^{2} + \frac{31}{26880}s^{2} - \frac{31}{161280}s^{3}g_{s}^{-2} \right] \frac{1}{t^{4}} + \mathscr{O}(t^{0}) \end{aligned}$$

+ contributions to 2(g+n) - 2 > 4. (2.70)

Here, $\hat{B}_m = (\frac{1}{2^{m-1}} - 1) \frac{B_m}{m!}$ with B_m denoting the Bernoulli numbers.

§2.4 Background independence of the topological string partition function and modularity. Besides the (refined) holomorphic anomaly equations, there is a second way to compute the higher genus free energies which (so far) only applies to the unrefined case. This method is based on an analysis of the monodromy group Γ that strongly constrains the form of the free energies. Namely it requires them to transform as almost/quasi modular forms of Γ , as explained in the following. In this case the topological string partition function is interpreted as a state $|Z\rangle$ in the Hilbert space obtained by quantizing $H^3(M, \mathbb{Z})$ where g_s plays the role of \hbar . While $|Z\rangle$ as an abstract state is invariant under the monodromy action, this does not neccessarily have to hold for its wavefunction (i.e. the projection of $|Z\rangle$ onto a particular set of coordinates). There are two important choices of coordinates. The first set $\{z_i, \bar{z}_i\}$ refers to picking a complex structure (holomorphic polarization), while the second set $\{x_i, p_i\}$ refers to a background symplectic structure (real polarization).

The results of this analysis, performed in [ABK08] are as follows:

1. In the holomorphic polarization the free energies are almost modular forms of the monodromy group Γ (i.e. modular invariant but not holomorphic) having an expansion

$$\widetilde{F}^{(g)} = F_0^{(g)} + \sum_{i=1}^{3g-3} h_g^i(\tau) (\mathrm{Im}\tau)^{-i}$$

with $h^i(\tau)$ being holomorphic functions. From a physics point of view these are determined by the holomorphic anomaly equation up to an integration constant.

- 2. In the real polarization the free energies are quasi modular forms of the monodromy group Γ (i.e. holomorphic but not modular invariant) $F^{(g)}$ and in fact $F^{(g)} \equiv F_0^{(g)}$.
- 3. The holomorphic anomaly equation can be derived from transforming from the real polarization to the complex polarization.
- 4. The propagator can be shown to be a modular form of weight two

$$\widetilde{E}^{IJ}(\tau, \overline{\tau}) = E^{IJ}(\tau) + \left((\mathrm{Im}\tau)^{-1} \right)^{IJ},$$

transforming as

$$\widetilde{E}^{IJ}(\tau,\bar{\tau}) \mapsto (c\tau+d)^{I}_{K}(c\tau+d)^{J}_{I}\widetilde{E}^{KL}(\tilde{\tau},\bar{\tilde{\tau}}), \qquad (2.71)$$

under

$$\tau \mapsto \gamma \tau = (a\tau + b)(c\tau + d)^{-1}, \qquad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{Sp}_{2\mathfrak{g}}(\mathbb{Z}).$$
 (2.72)

Here *E* takes values in $\mathbb{C}^{\mathfrak{g} \times \mathfrak{g}}$. In fact, any free energy $\widetilde{F}^{(g)}$ in holomorphic polarization can be written as a polynomial in $\widetilde{E}^{IJ}(\tau, \overline{\tau})$ where the coefficients are given by meromophic modular forms.

3 Invariants of Riemann surfaces and Their Modular Properties

This section is devoted to discuss the invariants of elliptic and genus two curves respectively. We mainly focus on the Igusa invariants that classify the latter. In particular, their Fourier expansion is used in the sections 4 and 5 in order to extract the period matrix of the mirror curve.

§3.1 Invariants of genus one curves. Any genus one curve can be represented in Weierstrass normal form

$$y^{2} = 4x^{3} - g_{2}(u, m_{i})x - g_{3}(u, m_{i}).$$
(3.1)

Here *u* denotes the true modulus of the curve that corresponds to the complex structure modulus τ , while the parameters m_i denote possible isomonodromic deformations as discussed in section 2.2.1. The coefficients g_2, g_3 are not true invariants, but enjoy a rescaling symmetry

$$g_2 \mapsto r^4 g_2, \qquad g_3 \mapsto r^6 g_3.$$
 (3.2)

In particular there is an *r* such that

$$E_4 = 12r^4g_2, \qquad E_6 = 216r^6g_3, \qquad \Delta_{\rm mod} = r^{12}\Delta_{\rm dis}.$$
 (3.3)

Here, we denote the discriminant locus respectively the modular discriminant by

$$\Delta_{\text{mod}} = \frac{1}{1728} \left(E_4^3(\tau) - E_6^2(\tau) \right), \qquad \Delta_{\text{dis}} = g_2^3(u, m_i) - 27g_3^2(u, m_i). \tag{3.4}$$

In contrast to g_2 , g_3 , the associated *j*-function

$$j = \frac{g_2^3(u, m_i)}{\Delta(u, m_i)} = \frac{E_4^3(\tau)}{E_4^3(\tau) - E_6^2(\tau)} = \frac{1}{q} + 744 + \dots, \qquad q = e^{2\pi i \tau},$$
(3.5)

is a true invariant.

§3.2 Invariants of genus two curves. Any Riemann surface *R* of genus g = 2 can be represented as hyperelliptic curve

$$y^{2} = v_{0}x^{6} - v_{1}x^{5} + v_{2}x^{4} - v_{3}x^{3} + v_{4}x^{2} - v_{5}x + v_{6},$$

$$= \prod_{i=1}^{6} (x - \lambda_{i}) \quad v_{i}, \lambda_{i} \in \mathbb{C}.$$
 (3.6)

To see this one notices that a hyperelliptic curve is a double cover of \mathbb{P}^1 which is equivalent to a function of degree two on *R*. The existence of the latter is guaranteed by the Riemann Roch theorem.

Two such hyperelliptic curves are conformally equivalent and define therefore the same Riemann surface if their branch points λ_i differ by a fractional linear transformation. Thus the moduli space of the hyperelliptic curves is locally parametrized by the λ_i and has dimension

$$\dim \mathcal{M}_g^{\text{hyp}} = 2g + 2 - \dim \text{PSL}_2(\mathbb{Z}) = 2g - 1.$$
(3.7)

Note that this coincides for g = 2 with dimension of the moduli space of Riemann surfaces

$$\dim \mathcal{M}_g = 3g - 3. \tag{3.8}$$

§3.2.1 The Igusa invariants. In this subsection we discuss the Igusa invariants which provide a natural generalization of g_2 , g_3 which occur in the Weierstrass normal form and the absolute *j*-invariant.

Given a genus two curve in the form (3.6), the Igusa invariants are defined as

$$A(v_i) = v_0^2 \sum_{15} (12)^2 (34)^2 (56)^2, \qquad (3.9)$$

$$B(v_i) = v_0^4 \sum_{10} (12)^2 (23)^2 (31)^2 (45)^2 (56)^2 (64)^2, \qquad (3.10)$$

$$C(v_i) = v_0^6 \sum_{60} (12)^2 (23)^2 (31)^2 (45)^2 (56)^2 (64)^2 (14)^2 (25)^2 (36)^2, \qquad (3.11)$$

$$D(v_i) = v_0^{10} \prod_{i < j} (\lambda_i - \lambda_j).$$
(3.12)

In these expressions we have summed over all permutations $\sigma \in \mathfrak{S}_6$ and used the abbreviation $(ij) = (\lambda_{\sigma(i)} - \lambda_{\sigma(j)})$. The invariants (3.9), (3.10), (3.11), (3.12) for a sextic of the form (3.6) are given by

$$\begin{split} &A = 6v_3^2 - 16v_2v_4 + 40v_1v_5 - 240v_0v_6, \\ &B = 48v_6v_2^3 + 4v_4^2v_2^2 - 12v_3v_5v_2^2 + 300v_0v_5^2v_2 + 4v_1v_4v_5v_2 - 180v_1v_3v_6v_2 - 504v_0v_4v_6v_2 + 48v_0v_4^3 \\ &- 12v_1v_3v_4^2 - 80v_1^2v_5^2 + 1620v_0^2v_6^2 + 36v_1v_3^2v_5 - 180v_0v_3v_4v_5 + 324v_0v_3^2v_6 + 300v_1^2v_4v_6 \\ &- 540v_0v_1v_5v_6, \\ &C = -36v_5^2v_2^4 - 160v_4v_6v_2^4 - 24v_4^3v_2^3 - 96v_0v_6^2v_2^3 + 76v_3v_4v_5v_2^3 + 60v_3^2v_6v_2^3 + 616v_1v_5v_6v_2^3 \\ &+ 8v_3^2v_4^2v_2^2 + 26v_1v_3v_5v_2^2 - 640v_0v_4v_5^2v_2^2 - 900v_1^2v_6^2v_2^2 - 24v_3^3v_5v_2^2 + 28v_1v_4^2v_5v_2^2 \\ &+ 424v_0v_4^2v_6v_2^2 + 492v_1v_3v_4v_6v_2^2 - 876v_0v_3v_5v_6v_2^2 - 160v_0v_4^4v_2 + 76v_1v_3v_4^3v_2 \\ &+ 1600v_0v_1v_5^3v_2 + 330v_0v_3^2v_5^2v_2 + 64v_1^2v_4v_5^2v_2 + 3060v_0v_1v_3v_6^2v_2 + 20664v_0^2v_4v_6^2v_2 \\ &+ 492v_0v_3v_4^2v_5v_2 - 238v_1v_3^2v_4v_5v_2 - 198v_1v_3^3v_6v_2 - 640v_1^2v_4^2v_6v_2 - 18600v_0^2v_5^2v_6v_2 \\ &- 468v_0v_3^2v_4v_6v_2 - 1860v_1^2v_3v_5v_6v_2 + 3472v_0v_1v_4v_5v_6v_2 - 36v_1^2v_4^4 + 60v_0v_3^2v_4^3 - 320v_1^3v_5^3 \\ &+ 2250v_0^2v_3v_5^3 - 119880v_0^3v_6^3 - 24v_1v_3^3v_4^2 + 176v_1^2v_3^2v_5^2 - 900v_0^2v_4^2v_5^2 - 1860v_0v_1v_3v_4v_5^2 \\ &- 10044v_0^2v_3^2v_6^2 + 2250v_1^3v_3v_6 - 18600v_0v_1^2v_4v_6^2 + 59940v_0^2v_1v_5v_6^2 + 72v_1v_4^3v_5 + 616v_0v_1v_4^3v_5 \\ &+ 330v_1^2v_3^2v_4v_6 + 1818v_0v_1v_3^2v_5v_6 + 1600v_1^3v_4v_5v_6 + 3060v_0^2v_3v_4v_5v_6, \\ D = v_0^2\Delta. \end{split}$$

The invariants A, B, C, D are the genus two analogues of g_2 and g_3 in the Weierstrass form. I.e. the rescaled coordinates

$$r^{2}A, r^{4}B, r^{6}C, r^{10}D, r \in \mathbb{C},$$
 (3.14)

define the same curve. In particular, one can find a proportionality constant such that one has ¹⁰

$$A = -24r^2 \frac{\chi_{12}}{\chi_{10}}, \qquad B = 4r^4 E_4, \qquad C' = \frac{1}{2}(AB - 3C) = 4r^6 E_6, \quad D = -2^{14}r^{10}\chi_{10}. \tag{3.15}$$

Altogether these invariants define a point in $\mathbb{P}^{1,2,3,5}$ and the complement of the divisor D = 0 describes the moduli space of genus two curves. However, if three roots in (3.6) coincide, the genus two curve degenerates into a product of elliptic curves. At the same time, the three invariants B, C, D vanish simultaneously such that the two-dimensional subspace parametrizing the two factors gets mapped onto the point [1:0:0:0]. One can show [Igu62; Igu67] that the following absolute invariants

$$x_1 = 2^4 3^2 \frac{B}{A^2}, \qquad x_2 = -2^6 3^3 \frac{AB - 3C}{A^3}, \qquad x_3 = 2 \cdot 3^5 \frac{D}{A^5},$$
 (3.16)

¹⁰The Maass lift $\phi_{10}^{(2)}$ used in section 9 is related to Igusa's cusp form by $4\chi_{10} = \phi_{10}^{(2)}$

serve as good uniformizing parameters around [1:0:0:0]. In particular, by introducing the following quantities

$$y_1 = \frac{x_1^3}{x_3}, \qquad y_2 = \frac{x_2^2}{x_3}, \qquad y_3 = \frac{x_1^2 x_2}{x_3}.$$
 (3.17)

one obtains the relations between the invariants of the degenerated genus two curve and its respective elliptic factors as

$$y_1 = j(\tau_1)j(\tau_2), \qquad y_2 = (j(\tau_1) - 2^6 3^3)(j(\tau_2) - 2^6 3^3).$$
 (3.18)

In this work we will eventually be interested to use the Fourier expansion of (3.16) in order to extract the complex structure parameters of a given genus two curve. We review some basic properties of the ring of Siegel modular forms and the computation of their Fourier expansions in the appendix C.

4 Direct Integration Algorithms based on Normal Forms of Mirror Curves

The methods presented in section 2.3 are perfectly sufficient in order to compute the (refined) free energies up to arbitrary worldsheet genus. However, for mirror curves of genus one and two, the fact that these possess a normal form allows to directly extract many quantities from the toric diagram. The key observation is that all quantities of the associated topological field theory can be expressed in terms of invariants of the curves and their period matrices. The latter are obtained by inverting the Fourier expansion of the associated absolute invariants which are constituted by the j-function respectively the Igusa invariants. In contrast, there is no way to circumvent the direct integration procedure to obtain the higher genus invariants in general. However there is one universal object, the propagator, which in fact can be written down once the period matrices are known. It is given by the second Eisenstein series E_2 in the case of genus one and by the derivative of $\log(\chi_{10})$ in the case of genus two.

Mathematically, for the case of genus one this is reflected in the fact that it is precisely the second Eisenstein series whose adjunction turns the ring of modular forms into the ring of almost respectively quasi modular forms. We show in section 9 that a similar statement can be made for genus two. The complete modular structure which is ensured to hold at least in the unrefined case is encoded in the monodromy group of the individual mirror curves and needs to be determined case by case.

In the following two subsections we discuss the algorithmic integration procedures for genus one and two separately.

§4.1 Direct Integration for genus one mirror curves. We summarize the integration procedure in figure 1 and discuss in addition some salient points of the algorithm.

§4.1.1 Obtaining the periods. The relation between the A-period *t* of the meromorphic differential and the period of the mirror curve at a cusp point in the moduli space is given by

$$\frac{dt}{du} = \sqrt{\frac{E_4(\tau)g_3(u,m_i)}{E_6(\tau)g_2(u,m_i)}} \stackrel{!}{=} \frac{1}{2\pi i} \int_{\mu} \frac{dx}{y}.$$
(4.1)

It is obvious that this expression is not invariant under the rescalings (3.2). This ambiguity is fixed by demanding that it coincides with the explicit integral over the vanishing cycle μ which implies the leading behaviour $t = \frac{1}{2\pi i} \log(u) + \dots$ It is trivial, but also important to note that (4.1) is also the holomorphic limit of the topological metric and is to be identified¹¹ with the proportionality constant *r* in (3.3).

¹¹Up to a numerical factor of 18.

§4.1.2 Obtaining the propagator. It is also possible to derive a closed expression for the propagator. One notices from (2.60) and (2.68) that one can choose A_t in such a way that one obtains 12

$$\frac{1}{2}C_{ttt}\mathscr{S}^{tt} = -\frac{1}{24}\partial_t \log\left(\Delta_{\mathrm{mod}}\right).$$
(4.2)

From this one easily obtains

$$\mathscr{S}^{tt} = \frac{c_0}{12} E_2.$$
 (4.3)

We denote by c_0 the intersection number of the A-cycle with the B-cycle. Here we made use of (2.46) and the well-known relation

$$E_2(\tau) = \frac{1}{2\pi i} \frac{d}{d\tau} \log(\Delta_{\text{mod}}).$$
(4.4)

After tensor transforming to the *u*-coordinates, one obtains

$$\mathscr{S}^{uu} = \frac{c_0}{12} \frac{g_3(u, m_i)}{g_2(u, m_i)} \frac{E_2(\tau) E_4(\tau)}{E_6(\tau)}.$$
(4.5)

In this derivation we have used the generic form (2.63) of $\mathscr{F}^{(0,1)}$. Taking into account the modular complement in equation (2.61), the right hand side of (4.2) picks up an additional summand

$$-\frac{1}{2}\log(\mathrm{Im}\tau), \qquad (4.6)$$

which precisely cancels the modular factor of the discriminant¹³. The relation (4.3) gets accordingly replaced by

$$S^{tt} = \frac{c_0}{12} \left(E_2 - \frac{3}{\pi \mathrm{Im}\tau} \right).$$
(4.7)

In addition, one has to check in general that the relations (2.66) and (2.67) are satisfied as well. Here we note that for genus two curves the relation (2.66) becomes just the Ramanujan identity

$$\partial_{\tau} E_2 = \frac{1}{12} \left(E_2^2 - E_4 \right). \tag{4.8}$$

This follows from (4.3) and (2.46). In particular, one finds that

$$f_t^{tt} = \frac{c_0^2}{144} C_{ttt} E_4(q) \,. \tag{4.9}$$

After transforming to *u*-coordinates one obtains

$$f_z^{zz} = \frac{c_0^2}{12} g_2 C_{zzz}.$$
(4.10)

 $^{^{12}}$ In the following, we use ${\cal S}$ to denote the holomorphic part of the propagator.

¹³Actually this is not completely true. Under a modular transformation $\tau \mapsto \frac{a\tau+b}{c\tau+d}$, $\frac{1}{\mathrm{Im}\tau}$ transforms into $\frac{|c\tau+d|^2}{\mathrm{Im}\tau}$. But (2.61) also contains a term $\bar{\partial}_{\bar{u}}\bar{t}$ which is not taken into account in (4.2). Altogether these three contributions are modular invariant.



Figure 1: This diagram shows the algorithm based on genus one mirror curves to compute the free energies via direct integration. Boxes show quantities that can derived purely from the mirror curve while clouds denote information that is needed in addition.



Figure 2: This diagram shows the algorithm based on genus two mirror curves to compute the free energies via direct integration. Boxes show quantities that can derived purely from the mirror curve while clouds denote information that is needed in addition. In contrast to the genus one case, also the knowledge of the A-periods is required.

§4.2 Direct integration for genus two mirror curves.

§4.2.1 Obtaining the periods and the topological metric. Unfortunately there seems to be no known generalization of (4.1) in order to obtain the A-periods of the meromorphic differential directly. Therefore these have to be computed by making use of the Picard-Fuchs equations in the case of genus two. However we conjecture that (4.1), interpreted as the topological metric, generalizes as follows to the case of genus two

$$|G_{ij}| = \text{Det}(\partial_{u_i} t_j) = \sqrt{\frac{E_6(\tau)B(u_i, m_j)}{E_4(\tau)C'(u_i, m_j)}}.$$
(4.11)

Clearly, also this expression is not invariant under rescalings and we further conjecture that this freedom can be fixed by requiring a leading term of $\prod_{i} \frac{1}{u_i}$. We have successfully checked this con-

jecture for the examples of the resolutions of $\mathbb{C}^3/\mathbb{Z}_5$ and $\mathbb{C}^3/\mathbb{Z}_6$, see also the discussion in sections 5.2 and 5.3. Finally we note that (4.11) provides up to a numerical factor the proportionality constant *r* in (3.14).

§4.2.2 Obtaining the propagator. The same derivation as above applies to the case of genus two. Analogously to the previous discussion, one chooses the ambiguity in (2.68), such that one obtains

$$\frac{1}{2}C_{ijk}S^{jk} = -\frac{1}{20}\partial_i \log(\chi_{10}) - \frac{1}{2}\partial_i \log(\det \mathrm{Im}\tau) .$$
(4.12)

Here, we abbreviate t_i by i and have used (3.14).

In this case one finds the following equality for the propagator

$$S^{ij} = \frac{1}{2\pi i} \frac{1}{10} \left(\partial_{\tau_{pq}} \log(\chi_{10}) + \frac{5}{i} (\mathrm{Im}\tau)_{pq}^{-1} \right) C_i^p C_j^q.$$
(4.13)

As for the well-known case of genus one, it is interesting to wonder whether the ambiguity (2.66) is also fixed by a generalized Ramanujan identity. In fact, we derive such an identity for the almost meromorphic Siegel modular form $\partial_{\tau} \log(\chi_{10})$ in section 9, see (9.3) and (9.5) for an explicit realization.

We recall the constraint (2.66) for the propagator which reads in the flat coordinates t_i

$$\partial_i \mathscr{S}^{kl} = -C_{imn} \mathscr{S}^{km} \mathscr{S}^{lm} + f_i^{kl}.$$
(4.14)

Again, we can make use of the relation (2.46) to re-write this as

$$(R_{\text{sym}^2}\mathscr{S})^{rs,mn} = t(\mathscr{S} \otimes \mathscr{S})^{rs,mn} + \frac{\partial t^i}{\partial \tau_{rs}} \left(C^{-1}\right)_k^m \left(C^{-1}\right)_l^m f_i^{kl}.$$
(4.15)

By abuse of notation, \mathscr{S}^{mn} denotes here only $\partial_{\tau_{mn}} \log(\chi_{10})$ and the definition of the operators R_{sym^2} and $t(S \otimes S)$ is provided in section 6.1. Comparing this expression to (9.3), it follows that the ambiguities f_i^{mn} are universally determined and can be expressed in terms of the meromorphic Siegel modular form f_{RS} of weight sym² \otimes sym² as

$$f_i^{mn} = -C_{irs} C_k^r C_l^s C_o^m C_p^n (f_{\rm RS})^{kl,op}.$$
(4.16)

5 | Examples

In this section we exemplify the theory which has been developed in sections 2, 3 and 4 by solving the topological string on three different toric geometries. The first one is given by local \mathbb{P}^2 . Besides being the standard example of a geometry with genus one mirror curve it also serves as an interesting limiting case of the resolution of $\mathbb{C}^3/\mathbb{Z}_5$. The latter geometry has a genus two mirror curve with two real moduli and no mass parameter deformations. Finally, we consider an example with a mass parameter which is provided by the resolution of $\mathbb{C}^3/\mathbb{Z}_6$. In all calculations we find our formalism to be perfectly confirmed.



§5.1 Solving the Topological String on local \mathbb{P}^2 .

§5.1.1 The geometry and its mirror. First we will solve the refined topological string on the anticanonical bundle $\mathbb{K}_{\mathbb{P}^2} = \mathcal{O}(-3) \to \mathbb{P}^2$. The solution to this problem is well known and will serve as a reference for the latter cases. Here the mirror curve is of genus one and all physical quantities can be expressed through modular or quasi-modular forms. We read off from the toric diagram that the A-cycle intersects the B-cycle -3 times. In addition, we find that the coordinates

$$X_0 = uxyw, \qquad X_1 = x^2y, \qquad X_2 = wy^2, \qquad X_3 = w^2x,$$
 (5.2)

automatically fulfill the constraint

$$z = \frac{X_1 X_2 X_3}{X_0^3}, \qquad u^{-3} = z, \tag{5.3}$$

that can be read off from the toric diagram. Accordingly, the mirror curve $\Sigma(z)$ is given by

$$x^{2}y + w^{2}y + wx^{2} + uxyw = 0.$$
 (5.4)

Here *z* denotes the complex structure modulus that takes values in the punctured sphere $\mathcal{M}(\Sigma(z)) = \mathbb{P}^1 \setminus \{z = 0, z = -\frac{1}{27}, \frac{1}{z} = 0\}$. Using birational transformations as e.g. provided by Nagell's algorithm this can be brought into Weierstrass form

$$y^{2} = 4x^{3} - \frac{1}{12}(1 + 24z)x - \frac{1}{216}(1 + 36z + 216z^{2}).$$
 (5.5)

As a next step, one inverts the *j*-function in order to evaluate τ as a function of *z* and obtains

$$q(z) = -z^3 + 45z^4 - 1512z^5 + 45672z^6 + \mathcal{O}(z^7), \qquad q = e^{2\pi i \tau}.$$
(5.6)

We can also use the knowledge about q in order to evaluate the A-period according to (4.1) as

$$t = \log(z) - 6z + 45z^2 - 560z^3 + \frac{17325}{2}z^4 + \mathcal{O}(z^5).$$
(5.7)

As a consistency check one notices that the Picard-Fuchs operator

$$\mathcal{D} = \Theta^2 + z(3\Theta + 2)(3\Theta + 1)3\Theta = \mathcal{L}\Theta, \qquad \Theta = z\partial_z, \qquad (5.8)$$

annihilates the A-period, as expected. Having computed the period we can exploit (2.36) to determine the prepotential

$$F^{(0,0)} = -\frac{1}{18}X_A^3 + 3Q - \frac{45}{8}Q^2 + \frac{244}{9}Q^3 - \frac{12333}{64}Q^4 + \frac{211878}{125}Q^5 + \mathcal{O}(Q^6).$$
(5.9)

Here, we have introduced the monodromy invariant variable $Q = e^t$. Also, we note that the correctly normalized B-period is given as

$$t_D = -\frac{1}{6}\log(z)^2 + \frac{1}{3}X_A\log(z) - 3z + \frac{141}{4}z^2 + \frac{1486}{3}z^3 - \frac{129805}{16}z^4 + \mathcal{O}(z^5).$$
(5.10)

Finally, we find that the Yukawa couplings are given as

$$C_{zzz} = -\frac{1}{3} \frac{1}{z^3 (1+27z)} \,. \tag{5.11}$$

The genus one free energies are given by

$$\mathscr{F}^{(0,1)} = -\frac{1}{12}\log(z^7(1+27z)) - \frac{1}{2}\log(\frac{\partial X_A}{\partial z}), \qquad (5.12)$$

and

$$F^{(1,0)} = \frac{1}{24} \log(\frac{1+27z}{z}).$$
(5.13)

§5.1.2 Direct integration. In this subsection we discuss the direct integration procedure applied to the example of local \mathbb{P}^2 . First of all, one notices that

$$z_c = 1 + 27z, (5.14)$$

is a good coordinate to perform computations around the conifold locus. Indeed, using (4.1) one directly obtains the A-period around the conifold locus as

$$t_c = \alpha \left(z_c + \frac{11}{18} z_c^2 + \frac{109}{243} z_c^3 + \frac{9389}{26244} z_c^4 + \frac{88351}{295245} z_c^5 + \mathcal{O}(z_c^6) \right)$$
(5.15)

Here, the proportionality constant α parametrizes the matching of the conifold period with the mass of D2-branes wrapping the vanishing cycle. In practice, it gets fixed by the gap condition (2.69). As a next step, we use the results of section 4.1 in order to determine the propagator,

$$S^{zz} = \frac{3}{4}z^2 + 9z^3 - 54z^4 + 756z^5 - 13284z^6 + 260496z^7 - 5451624z^8 + 119116656z^9 + \mathcal{O}\left(z^{10}\right).$$
(5.16)

Now the holomorphic anomaly equation (2.55) can be solved recursively and the holomorphic ambiguity at each step can be fixed by implementing the gap condition (2.70).

§5.2 Solving the Topological String on $\mathbb{C}^3/\mathbb{Z}_5$.

§5.2.1 The geometry and its mirror. We now apply the direct integration procedure to the topological string on a mirror-pair where the A-model geometry is the resolution of the orbifold $\mathbb{C}^3/\mathbb{Z}_5$. The toric data is given in (5.17) and one can see that the Kaehler-structure is parametrised by two moduli, both corresponding to normalisable directions in the moduli-space.

$$\begin{pmatrix} 0 & 0 & 1 & | & -3 & 1 & | & x_0 \\ 1 & 0 & 1 & | & 1 & -2 & | & x_1 \\ 2 & 0 & 1 & | & 0 & 1 & | & x_2 \\ 0 & 1 & 1 & | & 1 & 0 & | & x_3 \\ -1 & -1 & 1 & | & 1 & 0 & | & x_4 \\ & & & \uparrow & \uparrow & \\ & & & & \mathbb{P}_b^1 & \mathbb{P}_f^1 & | & \end{pmatrix}$$
(5.17)

By investigating the scaling relations, one easily sees that the divisor obtained by setting $x_0 = 0$ is given by a \mathbb{P}^2 while one obtains a \mathbb{F}_3 from $x_1 = 0$. In addition there are two generators of the Mori cone. They correspond to the hyperplane class of \mathbb{P}^2 , which can also be identified with the base \mathbb{P}^1_b of \mathbb{F}_3 while the curve \mathbb{P}^1_f gets identified with the fiber of the latter. There are no mass parameters present and the mirror curve is now given by a Riemann surface of genus two. From the two generators of the Mori cone we construct the mirror curve and read off the moduli

$$z_1 = \frac{X_1 X_3 X_4}{X_0^3}, \qquad z_2 = \frac{X_0 X_2}{X_1^2}.$$
 (5.18)



Figure 3: Toric diagram of the A-model geometry, which is the resolution of $\mathbb{C}^3/\mathbb{Z}_5$.

The constraints can be easily solved using the following choice of coordinates

$$X_0 = u x y z, \quad X_1 = v x^2 y, \quad X_2 = \frac{x^3 y}{z}, \quad X_3 = y^2 z, \quad X_4 = x z^2,$$
 (5.19)

where the relation between u, v and z_1 , z_2 is given as

$$u = z_1^{-\frac{2}{5}} z_2^{-\frac{1}{5}}, \qquad v = z_1^{-\frac{1}{5}} z_2^{-\frac{3}{5}}.$$
 (5.20)

Note that the coordinates u, v are the natural coordinates from a diagrammatic point of view and signal two normalizable complex structure moduli. In contrast, z_1 , z_2 are the physical complex structure moduli that appear in the mirror map. Having clarified the choice of coordinates, the mirror curve takes the form

$$y^{2} = -4x^{5} + z_{1}^{-\frac{4}{5}} z_{2}^{-\frac{2}{5}} x^{4} + 2z_{1}^{-\frac{3}{5}} z_{2}^{-\frac{4}{5}} x^{3} + (1 + 2z_{2}) z_{1}^{-\frac{2}{5}} z_{2}^{-\frac{6}{5}} x^{2} + 2z_{1}^{-\frac{1}{5}} z_{2}^{-\frac{3}{5}} x + 1.$$
(5.21)

The discriminant locus¹⁴ of the curve is given by

$$\Delta = 1 + 27z_1 + 3125z_1^2 z_2^3 + 4z_2^2 (4 + 125z_1) - z_2 (8 + 225z_1).$$
(5.22)

In addition, one finds three Picard Fuchs operators

$$\begin{aligned} \mathscr{L}_{1} &= (\Theta_{1} - 2\Theta_{2})\Theta_{1}^{2} - z_{1} \left[(\Theta_{2} - 3\Theta_{1})^{3} - 3(\Theta_{2} - 3\Theta_{1})^{2} + 2\Theta_{2} - 6\Theta_{1} \right], \\ \mathscr{L}_{2} &= (\Theta_{2} - 3\Theta_{1})\Theta_{2} - z_{2} \left[(2\Theta_{2} - \Theta_{1})^{2} + 2\Theta_{2} - \Theta_{1} \right], \\ \mathscr{L}_{3} &= \Theta_{2}\Theta_{1}^{2} + z_{1}z_{2}(\Theta_{2} - 3\Theta_{1})(2\Theta_{2} - \Theta_{1})(\Theta_{2} - 3\Theta_{1} - 1). \end{aligned}$$
(5.23)

Here we have used the common abbreviation for logarithmic derivatives $\Theta_i = z_i \frac{d}{dz_i}$. Note that linear dependent relations of the generators in the toric diagram can lead to linear independent Picard-Fuchs operators which is what happens here. In particular, the third differential operator is needed in order to exclude a triple logarithmic solution which is allowed by the first two operators. If one does not take into account the third operator, the solutions to the first two operators take the form $(1, t^i, \partial_i F, 2F - t^i \partial_i F)$ which is familiar from the compact case but not expected for the non-compact case.

After having excluded the triple-logarithmic solution, one then finds five solutions that are anni-

 $^{^{14}}$ We refer to the discriminant locus as the product of non-trivial irreducible components of the discriminant. The discriminant is up to a numerical factor given by the Igusa invariant *D* in (5.33).

hilated by the Picard-Fuchs operators \mathcal{L}_i ,

$$\begin{aligned}
\sigma_{0} &= 1 \\
\sigma_{1} &= \log(z_{1}) - 6z_{1} - z_{2} + 45z_{1}^{2} - \frac{3}{2}z_{2}^{2} - 560z_{1}^{3} - 18z_{1}^{2}z_{2} - \frac{10}{3}z_{2}^{3} + \mathcal{O}(z^{4}) \\
\sigma_{2} &= \log(z_{2}) + 2z_{1} + 2z_{2} - 15z_{1}^{2} + 3z_{2}^{2} + \frac{560}{3}z_{1}^{3} + 6z_{1}^{2}z_{2} + \frac{20}{3}z_{2}^{3} + \mathcal{O}(z^{4}) \\
\sigma_{3} &= 3z_{1} - z_{2}^{2} - z_{1}z_{2} - \frac{141}{4}z_{1}^{2} - z_{2}\log(z_{2}) - \frac{3}{2}z_{2}^{2}\log(z_{2}) - \frac{3}{10}\log(z_{2})^{2} \\
&- \frac{1}{5}\log(z_{1})\log(z_{2}) + 2z_{1}\log(z_{1}) - 15z_{1}^{2}\log(z_{1}) - \frac{1}{5}\log(z_{1})^{2} + \mathcal{O}(z^{3}), \\
\sigma_{4} &= -2z_{2} - 7z_{2}^{2} - z_{2}\log(z_{2}) - \frac{3}{2}z_{2}^{2}\log(z_{1}) - \frac{3}{5}\log(z_{1})\log(z_{2}) - \frac{1}{10}\log(z_{1})^{2} \\
&- 3z_{2}\log(z_{2}) - \frac{9}{2}z_{2}^{2}\log(z_{2}) - \frac{9}{10}\log(z_{2})^{2} + \mathcal{O}(z^{3}),
\end{aligned}$$
(5.24)

and a basis for the periods takes the form [Hos06]

$$\Pi = \begin{pmatrix} \tilde{t}_1^D \\ \tilde{t}_2^D \\ t_1 \\ t_2 \\ 1 \end{pmatrix} = \begin{pmatrix} -3 & 1 & \lambda_{11} & \lambda_{12} & \rho_1 \\ 1 & -2 & \lambda_{21} & \lambda_{22} & \rho_2 \\ 0 & 0 & 1 & 0 & \rho_3 \\ 0 & 0 & 0 & 1 & \rho_4 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \partial_{t_1} F^{(0,0)} \\ \partial_{t_2} F^{(0,0)} \\ t_1 \\ t_2 \\ 1 \end{pmatrix},$$
(5.25)

with

$$\left(\partial_{t_1} F^{(0,0)}, \partial_{t_2} F^{(0,0)}, t_1, t_2\right) = \left(\frac{1}{(2\pi i)^2}\sigma_3, \frac{1}{(2\pi i)^2}\sigma_4, \frac{1}{2\pi i}\sigma_1, \frac{1}{2\pi i}\sigma_2\right).$$
(5.26)

Here, the λ_{kl} as well as ρ_m parameterize the ambiguity to add logarithmic solutions of order zero and one to the double logarithmic solutions and do not influence the non-perturbative part of the prepotential but they are crucial in order to obtain an integral basis. However, as we do not refer to such a basis in the following we do not fix these ambiguities. By inverting the A-periods σ_1, σ_2 (5.24), we readily obtain the mirror map as

$$z_1(Q_1, Q_2) = Q_1 + 6Q_1^2 + Q_1Q_2 + 9Q_1^3 + 10Q_1^2Q_2 + \mathcal{O}(Q^4), \qquad (5.27)$$

$$z_2(Q_1, Q_2) = Q_2 - 2Q_1Q_2 - 2Q_2^2 + 5Q_1^2Q_2 + 6Q_1Q_2^2 + 3Q_2^3 + \mathcal{O}(Q^4), \qquad (5.28)$$

where we have introduced the monodromy invariant variables $Q_i = e^{\sigma_i}$.

§5.2.2 Extracting the complex structure moduli from the mirror. As a next step we extract the period matrix of the mirror curve that takes the explicit form

$$\tau = \begin{pmatrix} \tau_{11} & \frac{1}{2}\tau_{12} \\ \frac{1}{2}\tau_{12} & \tau_{22} \end{pmatrix}.$$
 (5.29)

In contrast to the torus, where the number of normalizable complex structure moduli of the mirror curve matches the moduli of a generic elliptic curve, one finds in the case at hand that the family of mirror curves parametrized by z_1, z_2 only defines a hypersurface within the Siegel fundamental domain $\mathscr{F}_2 = \Gamma_2 \setminus \mathbb{H}^{(2)}$. Using the formulae (3.13), one obtains for the mirror curve (5.21)

$$A = -8z_1^{\frac{2}{5}}z_2^{\frac{4}{5}}(-1+z_2(4+40z_1)), \qquad (5.30)$$

$$B = 4z_1^{\frac{4}{5}} z_2^{\frac{8}{5}} \left(1 + 24z_1 + 2400z_1^2 z_2^3 - 8z_2 \left(1 + 25z_1\right) + z_2^2 \left(16 + 440z_1 - 80z_1^2\right)\right),$$
(5.31)

$$C = -8z_1^{\frac{5}{5}}z_2^{\frac{15}{5}} \left(-1 - 20z_1 + 72z_1^2 + 8z_1^2z_2^4 (1009 + 10900z_1) + 4z_2 \left(3 + 75z_1 + 92z_1^2\right) - 4z_2^2 \left(12 + 365z_1 + 1652z_1^2\right) + 16z_2^3 \left(4 + 145z_1 + 948z_1^2 + 320z_1^3\right)\right),$$
(5.32)

$$D = 4096z_1^6 z_2^9 \left(1 + 27z_1 + 3125z_1^2 z_2^3 + 4z_2^2 \left(4 + 125z_1 \right) - z_2 \left(8 + 225z_1 \right) \right).$$
(5.33)

Computing the absolute invariants (3.16) and comparing them to their Fourier expansions (C.18) whose computation is presented in appendix C one obtains

$$q_{1}(z_{1}, z_{2}) = -z_{1}^{3} + z_{1}^{3}z_{2} + 45z_{1}^{4} - z_{1}^{3}z_{2}^{2} - 60z_{1}^{4}z_{2} - 1512z_{1}^{5} + \mathcal{O}(z^{6}),$$

$$q_{2}(z_{1}, z_{2}) = z_{1}^{2}z_{2}^{5} + 16z_{1}^{2}z_{2}^{6} - 5z_{1}^{3}z_{2}^{5} + \mathcal{O}(z^{9}),$$

$$r(z_{1}, z_{2}) = -z_{1} - 3z_{1}z_{2} + 15z_{1}^{2} - 13z_{1}z_{2}^{2} + 20z_{1}^{2}z_{2} - 279z_{1}^{3} + \mathcal{O}(z^{4}).$$
(5.34)

Here, we have denoted $q_1 = e^{2\pi i \tau_{11}}$, $q_2 = e^{2\pi i \tau_{22}}$ and $r = e^{2\pi i \tau_{12}}$. In addition, we note that in the limit of restricting to the local \mathbb{P}^2 model, corresponding to $z_2 \rightarrow 0$ only q_1 survives and gets indeed identified with (5.6). It is also satisfying to check that x_1 and x_2 reduce to the Weierstrass coefficients g_2, g_3 in (3.1). Using the knowledge of the τ -matrix as well as of the periods, we can determine the prepotential. For this purpose we make use of (2.46) noting that

$$C = \begin{pmatrix} -3 & 1\\ 1 & -2 \end{pmatrix},\tag{5.35}$$

and find that

$$F^{(0,0)} = -\frac{1}{15}T_1^3 - \frac{1}{10}T_1^2T_2 - \frac{3}{10}T_1T_2^2 - \frac{3}{10}T_2^3 + 3Q_1 - 2Q_2 - \frac{45}{8}Q_1^2 + 4Q_1Q_2 - \frac{1}{4}Q_2^2 + \frac{244}{9}Q_1^3 - 10Q_1^2Q_2 + 3Q_1Q_2^2 - \frac{2}{27}Q_2^3 + \mathcal{O}(Q^4).$$
(5.36)

In particular, denoting $K_{ij} = \partial_{t_i^A} \partial_{t_i^A} F^{(0,0)}$, one finds the explicit relations¹⁵

$$\tau_{11} = -4K_{11} + 4K_{12} - K_{22}, \qquad (5.37)$$

$$\tau_{22} = -K_{11} + 6K_{12} - 9K_{22}, \qquad (5.38)$$

$$\tau_{12} = -2K_{11} + 7K_{12} - 3K_{22}. \tag{5.39}$$

In addition, once the prepotential is known, we can directly write down the Yukawa couplings which are found to be

$$C_{z_1 z_1 z_1} = -\frac{2 + 9z_1 - 16z_2 - 95z_1 z_2 + 32z_2^2 + 300z_1 z_2^2}{5z_1^3 \Delta},$$
(5.40)

$$C_{z_1 z_1 z_2} = -\frac{1 + 27z_1 - 8z_2 - 210z_1 z_2 + 16z_2^2 + 400z_1 z_2^2}{5z_1^2 z_2 \Delta},$$
(5.41)

$$C_{z_1 z_2 z_2} = -\frac{3 + 81z_1 - 14z_2 - 405z_1 z_2 + 8z_2^2 + 325z_1 z_2^2}{5z_1 z_2^2 \Delta},$$
(5.42)

$$C_{z_2 z_2 z_2} = -\frac{9 + 243 z_1 - 17 z_2 - 540 z_1 z_2 + 4 z_2^2 + 225 z_1 z_2^2}{5 z_2^3 \Delta}.$$
 (5.43)

The knowledge of the A-periods enables us to check the conjecture (4.11) and we find in this case that

$$|G_{ij}| = \sqrt{\frac{E_6(\tau)B(z_i, m_j)}{z_1^{\frac{8}{5}} z_2^{\frac{6}{5}} E_4(\tau)C'(z_i, m_j)}},$$
(5.44)

where we have fixed the rescaling ambiguity accordingly. Finally, with the help of a few known BPS numbers the unkowns in (2.59) and (2.60) can be determined and one finds the refined free energies at genus one

$$F^{(1,0)} = \frac{1}{24} \log\left(\Delta z_1^{-2} z_2^{-3}\right), \qquad \mathscr{F}^{(0,1)} = -\frac{1}{12} \log\left(\Delta z_1^{\frac{38}{5}} z_2^{\frac{39}{5}}\right) - \frac{1}{2} \log\left(|G_{ij}|\right),$$

¹⁵Clearly, the prepotential can also be integrated by determing the B-periods first. From this perspective, this is merely a check of (2.46).

$$F^{(1,0)} = -\frac{1}{12}T_1 - \frac{1}{8}T_2 + \frac{7}{8}Q_1 - \frac{1}{6}Q_2 - \frac{129}{16}Q_1^2 + \frac{5}{6}Q_1Q_2 - \frac{1}{12}Q_2^2 + \frac{589}{6}Q_1^3 - \frac{65}{6}Q_1^2Q_2 + \frac{7}{8}Q_1Q_2^2 - \frac{1}{18}Q_2^3 + \mathcal{O}(Q^4), \qquad (5.45)$$

$$\mathscr{F}^{(0,1)} = -\frac{2}{15}T_1 - \frac{3}{20}T_2 + \frac{1}{4}Q_1 - \frac{1}{6}Q_2 - \frac{3}{8}Q_1^2 + \frac{1}{3}Q_1Q_2 - \frac{1}{12}Q_2^2 -\frac{23}{3}Q_1^3 - \frac{5}{6}Q_1^2Q_2 + \frac{1}{4}Q_1Q_2^2 - \frac{1}{18}Q_2^3 + \mathcal{O}(Q^4).$$
(5.46)

After this discussion, we are prepared to proceed to the direct integration which provides us with the higher genus free invariants.

§5.2.3 Direct Integration. In order to determine the higher genus free energies, we first have to calculate the propagator. Recall that the propagator is in the genus two case given by a matrix

$$\mathscr{S} = \begin{pmatrix} \mathscr{S}^{11}(z_1, z_2) & \mathscr{S}^{12}(z_1, z_2) \\ \mathscr{S}^{21}(z_1, z_2) & \mathscr{S}^{22}(z_1, z_2) \end{pmatrix}.$$
 (5.47)

Instead of using equations (2.66), (2.67) and (2.68), we follow the procedure in section 4.2. Using the relation (5.39) one finds that the following ansatz for the components of (5.47) solves (2.68),

$$\mathscr{S}^{t_1 t_1} = \frac{1}{10} \left(9 \frac{\partial}{\partial \tau_{11}} \log(\chi_{10}) + 3 \frac{\partial}{\partial \tau_{12}} \log(\chi_{10}) + \frac{\partial}{\partial \tau_{22}} \log(\chi_{10}) \right), \tag{5.48}$$

$$\mathscr{S}^{t_1 t_2} = \frac{1}{10} \left(-3 \frac{\partial}{\partial \tau_{11}} \log(\chi_{10}) - \frac{7}{2} \frac{\partial}{\partial \tau_{12}} \log(\chi_{10}) - 2 \frac{\partial}{\partial \tau_{22}} \log(\chi_{10}) \right), \tag{5.49}$$

$$\mathscr{S}^{t_2 t_2} = \frac{1}{10} \left(\frac{\partial}{\partial \tau_{11}} \log(\chi_{10}) + 2 \frac{\partial}{\partial \tau_{12}} \log(\chi_{10}) + 4 \frac{\partial}{\partial \tau_{22}} \log(\chi_{10}) \right).$$
(5.50)

This expression can be directly written down without any further computation, once the toric diagram of the A-model geometry is known and the τ -matrix has been extracted from the mirror curve by inverting the Fourier expansion of the Igusa invariants as performed in the previous section. After tensor transforming to the coordinates z_i these read explicitly

$$\mathscr{S}^{z_1 z_1} = \frac{7}{10} z_1^2 + 9 z_1^3 - \frac{3}{10} z_1^2 z_2 - 54 z_1^4 - 6 z_1^3 z_2 + \mathcal{O}(z^5), \qquad (5.51)$$

$$\mathscr{S}^{z_1 z_2} = -\frac{3}{20} z_1 z_2 - 3 z_1^2 z_2 + \frac{3}{5} z_1 z_2^2 + 18 z_1^3 z_2 + 7 z_1^2 z_2^2 + \mathcal{O}(z^5), \qquad (5.52)$$

$$\mathcal{S}^{z_2 z_2} = \frac{3}{10} z_2^2 + z_1 z_2^2 - \frac{6}{5} z_2^3 - 6 z_1^2 z_2^2 - 4 z_1 z_2^3 + \mathcal{O}(z^5).$$
(5.53)

In addition, we have checked that these propagators obey the over-determined system of equations provided by (2.67). Even more importantly, the ambiguities take precisely the form¹⁶ (4.16). Let us emphasize once more that this reflects the fact that the ambiguity of the propagator is completely fixed by the generalization of the Ramanujan identities for genus two as presented in (9).

In order to fix the holomorphic ambiguities $f^{(g_L,g_R)}$ that arise after integration of the holomorphic anomaly equation (2.55), we have to impose the gap condition at the conifold. To do this we have to choose a rational point on the discriminant locus in the moduli space and repeat the previous analysis. Throughout the following discussion, we will work with the point

$$(z_1, z_2) = (3, -2/9). \tag{5.54}$$

¹⁶For the ambitious reader who wants to re-check this calculation we note that our internal conventions for $f_{\rm RS}^{mn,kl}$ differ by a factor of -1/2 for the off-diagonal elements on each side.

The logarithmic singularities in the periods at the conifold are of the form $log(\Delta)$. However since we will need to transform objects from the large radius point to the conifold it is desirable that the new set of variables $z_i^c(z)$ can easily be inverted. The most convenient ansatz is a linear transformation

$$z_i^c = b_i \left(z_1 - z_1^{c,0} \right) + c_i \left(z_2 - z_2^{c,0} \right).$$
(5.55)

For the expansion of the singular periods to be well behaved in these coordinates we choose z_o^c orthogonal and z_p^c parallel to the conifold locus, i.e. $\nabla_z z_o^c \sim \nabla_z \Delta$ and $\nabla_z z_p^c \cdot \nabla_z \Delta = 0$. Then one can expand the logarithm as

$$\log(\Delta) = \log(z_{o}^{c} + \mathcal{O}\left[(z^{c})^{2}\right]) = \log(z_{o}^{c}) + \log\left(1 + \frac{\mathcal{O}\left[(z^{c})^{2}\right]}{z_{o}^{c}}\right)$$
$$= \log(z_{o}^{c}) + \sum_{n}^{\infty} \frac{(-1)^{n+1}}{n} \left(\frac{\mathcal{O}\left[(z^{c})^{2}\right]}{z_{o}^{c}}\right)^{n}.$$
(5.56)

and observes that a period $\omega = \pi \log(\Delta) + f$ has the structure

$$\omega(z_o^c, z_p^c) = \pi(z_o^c, z_p^c) \log(z_o^c) + \pi(z_o^c, z_p^c) \sum_n^\infty \frac{(-1)^{n+1}}{n} \left(\frac{\mathscr{O}\left[(z^c)^2\right]}{z_o^c}\right)^n + f(z_o^c, z_p^c).$$
(5.57)

Using a linear ansatz we find the variables

$$z_{o}^{c} = -9 + z_{1} - 27z_{2},$$

$$z_{p}^{c} = -\frac{727}{9} + 27z_{1} + z_{2},$$
(5.58)

at the point (3, -2/9) which are orthogonal and tangential to the discriminant locus. After transforming the Picard-Fuchs operators into these new variables one obtains again five periods,

$$\begin{split} t_c^0 &= 1, \end{split} \tag{5.59} \\ t_c^{A,1} &= z_o^c - \frac{271476409}{2127869700} \left(z_o^c \right)^2 + \frac{116172}{16120225} z_o^c z_p^c - \frac{2187}{5861900} \left(z_p^c \right)^2 + \mathcal{O}(z^3), \\ t_c^{A,2} &= z_p^c - \frac{8420610821}{590880856630} z_o^c z_p^c - \frac{2408185701}{295440428315} \left(z_p^c \right)^2 + \mathcal{O}(z^3), \\ t_c^{B,1} &= \frac{271062557874}{35748291826115} z_o^c z_p^c - \frac{6636689883}{12999378845860} \left(z_p^c \right)^2 + z_o^c \log(z_o^c) - \frac{271476409}{2127869700} \left(z_o^c \right)^2 \log(z_o^c) \\ &+ \frac{116172}{16120225} z_o^c z_p^c \log(z_o^c) - \frac{2187}{5861900} \left(z_p^c \right)^2 \log(z_o^c) + \mathcal{O}(z^3), \\ t_c^{B,2} &= \left(z_o^c \right)^2 - \frac{1359666z_o^c z_p^c}{809425831} - \frac{793881}{809425831} \left(z_p^c \right)^2 + \mathcal{O}(z^3). \end{split}$$

Note that these are not analytical continuations of the periods (5.24), (5.25) obtained at the large radius point. As there are besides the trivial solution three more solutions that take the form of a power series and only one logarithmic solution, a comment how the splitting into A- and B- periods is performed is in order. One identifies the power series in front of the logarithm $t_c^{B,1}$ as the period $t_c^{A,1}$. $t_c^{A,2}$ is choosen so that it starts with z_p^c and we found a possible addition of $X_c^{B,2}$ to cancel out in all physical quantities. Also note that $X_c^{B,1}$ contains terms with negative powers in z_o^c at higher order. Finally, we find the following mirror map

$$z_{o}^{c} = T_{1}^{c} + \frac{271476409}{2127869700} (T_{1}^{c})^{2} - \frac{116172}{16120225} T_{1}^{c} T_{2}^{c} + \frac{2187}{5861900} (T_{2}^{c})^{2} + \mathcal{O}(T^{3}),$$

$$z_{p}^{c} = T_{2}^{c} + \frac{8420610821}{590880856630} T_{1}^{c} T_{2}^{c} + \frac{2408185701}{295440428315} (T_{2}^{c})^{2} + \mathcal{O}(T^{3}).$$
(5.61)

At the large radius point the ambiguities A_i , f_i^{mn} , \tilde{f}_{ij}^m and Yukawa couplings C_{ijk} are rational expressions in the moduli z_i and can be tensor transformed into the conifold coordinates z^c . Again the topological metric can be calculated from the A-periods via

$$G_{ij}^c = \frac{\partial X_c^{A,i}}{\partial z_i^c},\tag{5.62}$$

and solving the over-constrained system of equations (2.66), (2.67) and (2.68) yields the propagator at the conifold. We have integrated the refined holomorphic anomaly equation (2.55) and implemented the gap-condition (2.69) up to genus three. The first two orders of the refined free energies at genus two are given by

$$\mathcal{F}^{(2,0)} = \frac{7}{46080} + \frac{47}{1280}z_1 - \frac{1}{240}z_2 - \frac{753}{640}z_1^2 - \frac{13}{3840}z_1z_2 - \frac{1}{60}z_2^2 + \mathcal{O}(z^3),$$

$$\mathcal{F}^{(1,1)} = -\frac{7}{11520} - \frac{73}{200}z_1 + \frac{19}{600}z_2 + \frac{2949}{200}z_1^2 + \frac{31}{300}z_1z_2 + \frac{19}{150}z_2^2 + \mathcal{O}(z^3),$$

$$\mathcal{F}^{(0,2)} = \frac{1}{1920} + \frac{109}{640}z_1 - \frac{11}{1200}z_2 - \frac{23331}{3200}z_1^2 - \frac{233}{9600}z_1z_2 - \frac{11}{300}z_2^2 + \mathcal{O}(z^3),$$

(5.63)

and the leading terms for genus three are

$$\mathcal{F}^{(3,0)} = \frac{31}{77414400} - \frac{3229}{1152000} z_1 + \frac{83}{2304000} z_2 + \mathcal{O}(z^3),$$

$$\mathcal{F}^{(2,1)} = -\frac{31}{12902400} + \frac{346669}{10240000} z_1 - \frac{1023}{1280000} z_2 + \mathcal{O}(z^3),$$

$$\mathcal{F}^{(1,2)} = \frac{41}{9676800} - \frac{104421}{1280000} z_1 + \frac{13277}{2880000} z_2 + \mathcal{O}(z^3),$$

$$\mathcal{F}^{(0,3)} = -\frac{1}{483840} + \frac{26454323}{460800000} z_1 - \frac{27563}{4800000} z_2 + \mathcal{O}(z^3).$$
(5.64)

The refined GV invariants are listed in Appendix D. In addition we find the following non-vanishing refined BPS invariants

$$N_{0,1}^{(1,0)} = N_{0,0}^{(1,1)} = N_{0,1}^{(1,1)} = N_{0,1}^{(1,2)} = 1.$$
(5.65)

§5.3 Solving the Topological String on $\mathbb{C}^3/\mathbb{Z}_6$. In the previous section we discussed local $\mathbb{C}^3/\mathbb{Z}_5$ which is the simplest geometry with mirror curve of genus two as it only has two true moduli. Now we present the discussion of the resolution¹⁷ of the orbifold $\mathbb{C}^3/\mathbb{Z}_6$ which provides an example with three moduli out of which one is just a deformation (mass) parameter.



Figure 4: The A-model geometry of $\mathbb{C}^3/\mathbb{Z}_6$.

 $^{^{17}}$ Note that there are two ways to make \mathbb{Z}_6 act on \mathbb{C}^3 . The second way leads to a geometry whose mirror curve has genus one.

The toric diagram is shown in figure 4 and the corresponding data is listed in (5.66).

We note that the orbifold is resolved by two Hirzebruch surfaces, \mathbb{F}_2 and \mathbb{F}_4 which correspond to the divisors $\{x_0 = 0\}$ and $\{x_1 = 0\}$ respectively. In addition there is the non-compact divisor $\{x_4 = 0\}$ that has the topology of $\mathbb{C} \times \mathbb{P}^1$ which resolves the singularity $\mathbb{C}^2/\mathbb{Z}_2$ corresponding to the point on the edge of the toric diagram.

By analogous considerations to those for the case $\mathbb{C}^3/\mathbb{Z}_5$ one finds the mirror curve by the Hori-Vafa method to be given as

$$y^{2} = \left(z_{3}z_{1}^{2} - \frac{z_{1}^{2}}{4}\right)x^{6} - \frac{z_{1}}{2}x^{5} + \left(\frac{z_{1}}{2} + \frac{1}{4}\right)x^{4} - \left(\frac{z_{1}z_{2}}{2} + \frac{1}{2}\right)x^{3} - \left(\frac{z_{2}}{2} + \frac{1}{4}\right)x^{2} - \frac{z_{2}}{2}x - \frac{z_{2}^{2}}{4}.$$
 (5.67)

Note that the variables z_i are already those which correspond to the three Mori vectors of the toric Diagram which need to be contrasted to the two real moduli and the mass parameters appearing in the toric diagram. The discriminant locus of the mirror curve is found to consist of two irreducible components which are given as

$$\begin{split} \Delta_{1} = &1 - 8z_{1} + 16z_{1}^{2} - 64z_{1}^{2}z_{3} - 8z_{2} + 68z_{1}z_{2} - 144z_{1}^{2}z_{2} + 576z_{1}^{2}z_{2}z_{3} + 16z_{2}^{2} - 144z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} \\ &- 1512z_{1}^{2}z_{2}^{2}z_{3} + 216z_{1}^{3}z_{2}^{2} - 864z_{1}^{3}z_{2}^{2}z_{3} + 216z_{1}^{2}z_{3}^{2} + 864z_{1}^{2}z_{2}^{3}z_{3} - 972z_{1}^{3}z_{3}^{3} + 3888z_{1}^{3}z_{2}^{3}z_{3} \\ &+ 729z_{1}^{4}z_{2}^{4} - 5832z_{1}^{4}z_{2}^{4}z_{3} + 11664z_{1}^{4}z_{2}^{4}z_{3}^{2}, \end{split}$$

$$(5.68)$$

Here Δ_2 corresponds to a locus with enhanced *SU*(2) gauge symmetry where the theory stays regular, compare also the discussion in [KM96; KMP96].

The three Picard Fuchs operators that correspond to the Mori vectors of the toric diagram

$$\begin{aligned} \mathscr{L}_{1} &= \Theta_{1}(\Theta_{1} - 2\Theta_{2} - 2\Theta_{3}) + 4\Theta_{2}\Theta_{3} - z_{1}(2\Theta_{1} - \Theta_{2} + 1)(2\Theta_{1} - \Theta_{2}), \\ \mathscr{L}_{2} &= \Theta_{2}(\Theta_{2} - 2\Theta_{1}) + z_{2}(2\Theta_{2} - \Theta_{1} + 1)(2\Theta_{2} - \Theta_{1}), \\ \mathscr{L}_{3} &= \Theta_{3}^{2} + z_{3}(2\Theta_{3} - \Theta_{1} + 1)(2\Theta_{3} - \Theta_{1}), \end{aligned}$$
(5.69)

need also in this case to be supplemented by a fourth operator corresponding to the sum of the three mori cone vectors

$$\mathscr{L}_{4} = \Theta_{2}\Theta_{3}^{2} + z_{1}z_{2}z_{3}(\Theta_{1} - 2\Theta_{2})(\Theta_{1} - 2\Theta_{3})(2\Theta_{1} - \Theta_{2}).$$
(5.70)

The latter discards one double-logarithmic and a triple-logarithmic solution of the three operators (5.69). In addition, we observe again that the system of solutions of (5.69) takes the form (1, t^i , $\partial_i F$, $2F - t^i \partial_i F$). Having remarked this we present the solution to the complete system that includes (5.70),

$$\begin{aligned} \sigma_{0} &= 1, \\ \sigma_{1} &= \log(z_{1}) + 2z_{1} + 3z_{1}^{2} - z_{3} - \frac{3}{2}z_{3}^{2} - z_{2} - \frac{3}{2}z_{2}^{2} + \mathcal{O}(z^{3}), \\ \sigma_{2} &= \log(z_{2}) - z_{1} - \frac{3}{2}z_{1}^{2} + 2z_{2} + 3z_{2}^{2} + \mathcal{O}(z^{3}), \\ \sigma_{3} &= \log(z_{3}) + 2z_{3} + 3z_{3}^{2} + \frac{20}{3}z_{3}^{3} + \frac{35}{2}z_{3}^{4} + \frac{252}{5}z_{3}^{5} + \mathcal{O}(z^{6}), \\ \sigma_{4} &= -2z_{1} - 2z_{1}\log(z_{1}) - z_{1}\log(z_{3}) + \frac{2}{3}z_{3}\log(z_{3}) - \frac{2}{3}\log(z_{1})\log(z_{3}) - 2z_{2}\log(z_{2}) \\ &- \frac{2}{3}\log(z_{1})\log(z_{1}) - \frac{1}{3}\log(z_{3})\log(z_{2}) - \frac{2}{3}\log(z_{2})^{2} + \mathcal{O}(z^{2}), \\ \sigma_{5} &= -2z_{2} - 2z_{2}\log(z_{1}) - \frac{1}{3}\log(u)^{2} + \frac{1}{3}z_{3}\log(z_{3}) - z_{2}\log(z_{3}) - \frac{1}{3}\log(z_{1})\log(z_{3}) - 4z_{2}\log(z_{2}) \\ &- \frac{4}{3}\log(z_{1})\log(z_{2}) - \frac{2}{3}\log(z_{3})\log(z_{2}) - \frac{4}{3}\log(z_{2})^{2} + \mathcal{O}(z^{2}), \end{aligned}$$
(5.71)

with

$$\left(\partial_1 F^{(0,0)}, \partial_2 F^{(0,0)}, t_1, t_2, t_3\right) = \left(\frac{1}{(2\pi i)^2}\sigma_4, \frac{1}{(2\pi i)^2}\sigma_5, \frac{1}{2\pi i}\sigma_1, \frac{1}{2\pi i}\sigma_2, \frac{1}{2\pi i}\sigma_3\right).$$
(5.72)

Using the monodromy invariant variables $Q_1 = e^{\sigma_1}$, $Q_2 = e^{\sigma_2}$ and $Q_3 = e^{\sigma_3}$ the mirror map is given by

$$z_{1}(Q_{1}, Q_{2}, Q_{3}) = Q_{1} - 2Q_{1}^{2} + 3Q_{1}^{3} + Q_{1}Q_{3} - 4Q_{1}^{2}Q_{3} + Q_{1}Q_{2} - 3Q_{1}^{2}Q_{2} + Q_{1}Q_{2}Q_{3} + \mathcal{O}(Q^{4}),$$

$$z_{2}(Q_{1}, Q_{2}, Q_{3}) = Q_{2} + Q_{1}Q_{2} + Q_{1}Q_{2}Q_{3} - 2Q_{2}^{2} - 3Q_{1}Q_{2}^{2} + 3Q_{2}^{3} + \mathcal{O}(Q^{4}),$$

$$z_{3}(Q_{1}, Q_{2}, Q_{3}) = Q_{3} - 2Q_{3}^{2} + 3Q_{3}^{3} - 4Q_{3}^{4} + 5Q_{3}^{5} - 6Q_{3}^{6} + 7Q_{3}^{7} + \mathcal{O}(Q^{8}),$$

(5.73)

where $z_3(Q_1, Q_2, Q_3)$ can be expressed as

$$z_3(Q_3) = \frac{Q_3}{(1+Q_3)^2}.$$
(5.74)

Integrating $\partial_1 F^{(0,0)}$ and $\partial_2 F^{(0,0)}$ we obtain the prepotential

$$F^{(0,0)} = -2Q_1 - 2Q_2 - \frac{2}{9}T_1^3 - \frac{1}{3}T_1^2T_3 - \frac{1}{3}T_1^2T_2 - \frac{1}{3}T_1T_2T_3 - \frac{2}{3}T_1T_2^2 - \frac{1}{3}T_2^2T_3 - \frac{4}{9}T_2^3 + \mathcal{O}(Q^2).$$
(5.75)

We list the corresponding instanton numbers in appendix E. Having computed the prepotential, we can determine the Yukawa couplings

$$C_{111} = -\frac{1}{3z_1^3 \Delta_1} (4 - 8z_1 + 16z_1^2 - 64z_1^2 z_3 - 32z_2 + 80z_1 z_2 - 168z_1^2 z_2 + 672z_1^2 z_2 z_3 + 64z_2^2 - 192z_1 z_2^2 + 540z_1^2 z_2^2 - 2160z_1^2 z_2^2 z_3 - 432z_1^2 z_2^3 + 1728z_1^2 z_2^3 z_3),$$

$$C_{112} = -\frac{1}{3z_1^2 z_2 \Delta_1} (2 - 16z_1 + 32z_1^2 - 128z_1^2 z_3 - 16z_2 + 124z_1 z_2 - 264z_1^2 z_2 + 1056z_1^2 z_2 z_3 + 32z_2^2 - 240z_1 z_2^2 + 594z_1^2 z_2^2 - 2376z_1^2 z_2^2 z_3 - 216z_1^2 z_2^3 + 864z_1^2 z_2^3 z_3),$$

$$C_{122} = -\frac{1}{3z_1^2 z_2 \Delta_1} (4 - 32z_1 + 64z_1^2 - 256z_1^2 z_3 - 20z_2 + 176z_1 z_2 - 384z_1^2 z_2 + 1536z_1^2 z_2 z_3 + 16z_2^2 - 192z_1 z_2^2 + 540z_1^2 z_2^2 - 2160z_1^2 z_2^2 z_3 - 108z_1^2 z_2^3 + 432z_1^2 z_2^3 z_3),$$

$$C_{222} = -\frac{1}{3z_3^3 \Delta_1} (8 - 64z_1 + 128z_1^2 - 512z_1^2 z_3 - 22z_2 + 208z_1 z_2 - 480z_1^2 z_2 + 1920z_1^2 z_2 z_3 + 8z_2^2 - 132z_1 z_2^2 + 432z_1^2 z_2^2 - 1728z_1^2 z_2^2 z_3 - 54z_1^2 z_2^3 + 216z_1^2 z_2^3 z_3).$$

$$(5.76)$$

We also use the knowledge of the prepotential in order to successfully check the relation (2.46). Here, the two derivatives have to be taken with respect to the two true moduli t_1 and t_2 . The toric data (5.66) provides us with the following intersection matrix

$$C = \begin{pmatrix} -2 & 1\\ 1 & -2 \end{pmatrix}.$$
 (5.77)

We obtain

$$\begin{aligned} q_1(z_1, z_2, z_3) &= z_1^4 z_2^6 z_3^2 + 4 z_1^5 z_2^6 z_3^2 + 16 z_1^6 z_2^6 z_3^2 + 16 z_1^4 z_2^7 z_3^2 + 56 z_1^5 z_2^7 z_3^2 + 160 z_1^4 z_2^8 z_3^2 + \mathcal{O}(z^{15}), \\ q_2(z_1, z_2, z_3) &= z_1^4 z_3^2 + 16 z_1^5 z_3^2 + 160 z_1^6 z_3^2 - 2 z_1^4 z_2 z_3^2 - 40 z_1^5 z_2 z_3^2 + z_1^4 z_2^2 z_3^2 + \mathcal{O}(z^9), \\ r(z_1, z_2, z_3) &= z_1^2 z_3 + 8 z_1^3 z_3 + 48 z_1^4 z_3 + 2 z_1^2 z_2 z_3 + 6 z_1^3 z_2 z_3 + 9 z_1^2 z_2^2 z_3 + \mathcal{O}(z^6), \end{aligned}$$
(5.78)

and plugging these expressions into the respective Fourier expansions of the absolute Igusa invariants (3.16) we find perfect agreement with the rational expressions obtained from the mirror curve (5.67). In addition, we also verify our conjecture for the expression of the determinant of the topological metric (4.11)

$$|G_{ij}| = \sqrt{\frac{E_6(\tau)B(z_i)}{16z_1^2 z_2^2 E_4(\tau)C'(z_i)}},$$
(5.79)

where the indices *i*, *j* are again only running over true moduli t_1 , t_2 .

As a next step, we use the known GV invariants for local \mathbb{F}^2 [Ali09; HKP13] to fix the ambiguities for the free energies at genus one

$$F^{(1,0)} = \frac{1}{24} \log \left(\Delta_1 z_1^{-4} z_2^{-4} z_3^{-2} \right), \qquad \mathscr{F}^{(0,1)} = -\frac{1}{12} \log \left(\Delta_1 z_1^8 z_2^8 z_3 \right) - \frac{1}{2} \log \left(|G_{ij}| \right).$$

The corresponding refined GV invariants can be found in Appendix E. Finally using the logarithmic derivatives of χ_{10} we obtain the propagator from (4.13) as

$$\mathscr{S}_{11} = \frac{3z_1^2}{10} - \frac{6z_1^3}{5} - 8z_1^4 z_3 - 32z_1^5 z_3 - \frac{3}{10}z_1^2 z_2 + \frac{9}{5}z_1^3 z_2 + 16z_1^4 z_2 z_3 + \mathcal{O}(z^7),$$

$$\mathscr{S}_{12} = -\frac{3}{20}z_1 z_2 + \frac{3}{5}z_1^2 z_2 + 4z_1^3 z_2 z_3 + 16z_1^4 z_2 z_3 + \frac{3}{5}z_1 z_2^2 - \frac{9}{4}z_1^2 z_2^2 - 14z_1^3 z_2^2 z_3 + \mathcal{O}(z^7),$$

$$\mathscr{S}_{22} = \frac{3z_2^2}{10} - \frac{3}{10}z_1 z_2^2 - 2z_1^2 z_2^2 z_3 - 8z_1^3 z_2^2 z_3 - \frac{6z_2^3}{5} + \frac{9}{5}z_1 z_2^3 + 10z_1^2 z_2^3 z_3 + \mathcal{O}(z^7).$$
(5.80)

We have successfully checked that this propagator obeys the over-determined system of equations (2.66) and (2.67).

6 | Almost Holomorphic Siegel Modular Forms

The propagator that was described in (4.13) is an almost meromorphic Siegel modular form. To subsume it under a satisfying structure theory, we first develop a theory of almost holomorphic Siegel modular forms. Shimura studied them very early on [Shi86] and called them nearly holomorphic. Subsequent work by Zagier [Zag94] and Kaneko-Zagier [KZ98] employed "almost holomorphic" as terminology, and defined quasi modular form as the "constant part" of almost holomorphic ones.

From a geometric and representation theoretic point of view, it is clear how to define almost holomorphic Siegel modular forms. We discuss this in Remark 6.14. A classical, explicit description, however, is not available in the literature, and we start the mathematical part of this work by filling that gap. In fact, not even all covariant differential operators are explicitly known. They are the theme of Section 6.2. In Section 6.3, we provide a definition of almost holomorphic functions. It serves as a foundation for Section 6.4, which contains the definition and basic properties of almost holomorphic Siegel modular forms.

§6.1 Preliminaries. We revisit some basic aspects of the theory of Siegel modular forms. The reader can find a good introduction, which goes more into detail, in [Fre83] or [BGHZ08]. Siegel's foundational work [Sie51] provides one of the most general contexts in which Siegel modular forms can be understood.

The upper half spaces. We write $\mathbb{H}^{(n)} = \{\tau \in \operatorname{Mat}_{n}^{t}(\mathbb{C}) : \mathfrak{Im}(\tau) \text{ is positive definite} \}$ for the degree *n* Siegel upper half space. The real and imaginary parts of τ , which are matrices of size $n \times n$, are written $x = \mathfrak{Re}(\tau), y = \mathfrak{Im}(\tau)$, throughout. Coordinates of τ , *x*, and *y* are denoted respectively by τ_{ij} , x_{ij} , and y_{ij} . We use the ambiguous notation $y_{ij}^{-1} = (y^{-1})_{ij}$ for the *ij*-th entry of y^{-1} (as opposed to the inverse of γ_{ii}).

The symplectic group. The Siegel upper half space carries an action of the symplectic group. Let $J_n = \begin{pmatrix} 0 & \mathbb{I}_n \\ -\mathbb{I}_n & 0 \end{pmatrix}$ be the standard symplectic form, where \mathbb{I}_n is the $n \times n$ identity matrix. The symplectic group is its stabilizer

$$\mathbf{G}^{(n)} = \operatorname{Sp}_n(\mathbb{R}) = \{ g \in \operatorname{Mat}_{2n}(\mathbb{R}) : {}^{\mathrm{t}}gJ_ng = J_n \}.$$
(6.1)

Elements of $G^{(n)}$ are written as $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $a, b, c, d \in Mat_n(\mathbb{R})$. The action of $G^{(n)}$ on $\mathbb{H}^{(n)}$ is then defined by

$$g\tau = (a\tau + b)(c\tau + d)^{-1}.$$
 (6.2)

We fix special elements inv = $\begin{pmatrix} 0 & -\mathbb{I}_n \\ \mathbb{I}_n & 0 \end{pmatrix} \in G^{(n)}$ and trans $(b) = \begin{pmatrix} \mathbb{I}_n & b \\ 0 & \mathbb{I}_n \end{pmatrix} \in G^{(n)}$ for $b \in \operatorname{Mat}_n^t(\mathbb{R})$, where $\operatorname{Mat}_n^t(\mathbb{R})$ denotes the space of $n \times n$ symmetric matrices with real entries. Note that inv and trans(b), for *b* running trough $\operatorname{Mat}_n^t(\mathbb{R})$, generate $G^{(n)}$.

The integral points of $G^{(n)}$ are denoted by $\Gamma^{(n)} = \operatorname{Sp}_n(\mathbb{Z})$. In analogy to the previous observation, $\Gamma^{(n)}$ is generated by inv and all trans(*b*) with $b \in \operatorname{Mat}_n^t(\mathbb{Z})$.

Slash actions. Given a representation σ of $GL_n(\mathbb{C})$, we set

$$(f|_{\sigma}g)(\tau) = \sigma(c\tau + d)^{-1}f(g\tau)$$

for $f \in C^{\infty}(\mathbb{H}^{(n)} \to V(\sigma))$ and $g \in G^{(n)}$. We call $|_{\sigma}$ the slash action associated to the weight σ , or less strictly, the weight σ slash action.

If we do not state otherwise, σ is assumed to be finite dimensional and complex. In the special case of $\sigma = \det^k$, we write $f|_k g$ for $f|_{\det^k} g$.

Weights. Write std for the standard representation of $GL_n(\mathbb{C})$, defined by $(g, v) \mapsto gv$. Let std^{\vee} be its contragredient, given by $(g, v) \mapsto {}^tg^{-1}v$. Bases for them will be denoted by \mathfrak{e}_i and \mathfrak{e}_i^{\vee} with $1 \le i \le n$. We identify the symmetric square sym^2 of std and its contragredient $sym^{\vee 2}$ with the representations $(g, m) \mapsto gm{}^tg$ and $(g, m) \mapsto {}^tg^{-1}mg^{-1}$ on $Mat_n^t(\mathbb{C})$. Bases in this case, are \mathfrak{e}_{ij} and \mathfrak{e}_{ij}^{\vee} with $1 \le i, j \le n$.

Since every representation of $SL_n(\mathbb{C})$ is a subrepresentation of std^l for some $0 \le l \in \mathbb{Z}$ (cf. Theorem 5.5.11 of [GW09]), we find that every representation of $GL_n(\mathbb{C})$ can be embedded into $std^{l_1} \otimes std^{\vee l_2}$ for some $0 \le l_1, l_2 \in \mathbb{Z}$. We will often write tensor products as $\sigma\sigma' = \sigma \otimes \sigma'$, if σ and σ' are representations of the same group. The outer tensor product is denoted by \boxtimes .

The space of homomorphisms between two representations is written $\operatorname{Hom}(\sigma, \sigma')$. If σ and σ' are irreducible, then by Schur's Lemma either $\operatorname{Hom}(\sigma, \sigma') \cong \mathbb{C}$ or $\operatorname{Hom}(\sigma, \sigma') = 0$. Do mind the difference between $\operatorname{Hom}(\sigma, \sigma')$ and the generally large space of vector space homomorphisms $\operatorname{Hom}(V(\sigma), V(\sigma'))$. To simplify notation, if we are given a $\operatorname{Hom}(V(\sigma), V(\sigma'))$ -valued function m on $\mathbb{H}^{(n)}$ and a function f taking values in $V(\sigma)$ then we write m f for $\tau \mapsto m(\tau) f(\tau)$.

Since we often swap components of tensor product, we fix notation once and for all. We let

$$t_{ij}:\sigma_1 \otimes \cdots \otimes \sigma_a \longrightarrow \sigma_1 \otimes \cdots \otimes \sigma_{i-1} \otimes \sigma_j \otimes \sigma_{i+1} \otimes \cdots \otimes \sigma_{j-1} \otimes \sigma_i \otimes \sigma_{j+1} \otimes \cdots \otimes \sigma_a \tag{6.3}$$

be the canonical isomorphism, if, say, $1 \le i < j \le a$. Now fix distinct integers i_1, \ldots, i_b and j_1, \ldots, j_c between 1 and *a*. The average of components is denoted by

$$\mathbf{t}_{(i_1,\dots,i_b)(j_1,\dots,j_c)} = \frac{1}{bc} \sum_{k=1}^b \sum_{l=1}^c \mathbf{t}_{i_k j_l}.$$
(6.4)

In what follows, we will few sym² and sym^{$\vee 2$} as subrepresentations of std² and std^{$\vee 2$}, if they originate in the application of lowering or raising operators.

§6.2 Differential operators for Siegel modular forms. A differential operator D from $C^{\infty}(\mathbb{H}^{(n)} \to V(\sigma))$ to $C^{\infty}(\mathbb{H}^{(n)} \to V(\sigma'))$ is called covariant if it intertwines the associated slash actions:

$$D(f|_{\sigma}g) = (Df)|_{\sigma'}g$$

for all $f \in C^{\infty}(\mathbb{H}^{(n)} \to V(\sigma))$ and all $g \in G^{(n)}$. The space of such differential operators is denoted by $\mathbb{D}(\sigma, \sigma')$.

Differential operators for Siegel modular forms in the case of $\sigma = \det^k$ were found by Maass [Maa79] and by Shimura [Shi73]. For our purpose we need a basic understanding of the analytic properties of covariant differential operators for any (holomorphic) weight.

We proceed in three steps to compute them. We first give the lowering operators (in $\mathbb{D}(\sigma, \operatorname{sym}^{\vee 2} \otimes \sigma)$) for arbitrary holomorphic weight. This amounts to a direct verification. Second, we deduce from this the raising operators (in $\mathbb{D}(\sigma, \operatorname{sym}^2 \otimes \sigma)$) in case that σ is the standard representation or its contragredient. As a third step, we deduce all properties of covariant differential operators that we need from the previous computations and the fact that the standard representation generates the representation ring of $\operatorname{SL}_n(\mathbb{C})$.

Notation for differentials. Given $1 \le i, j \le n$, we set

$$\partial_{ij} = \partial_{\tau,ij} = \partial_{\tau_{ij}} = \frac{d}{d\tau_{ij}}, \quad \partial_{\overline{ij}} = \partial_{\overline{\tau},ij} = \partial_{\overline{\tau}_{ij}} = \frac{d}{d\overline{\tau_{ij}}}.$$
 (6.5)

Differentials with respect to *y* are analogously denoted by $\partial_{y,ij}$.

Following both Maass and Shimura, we define the following matrices of differential operators.

$$\partial_{\tau} = \left(\frac{1+\delta_{ij}}{2}\partial_{\tau,ij}\right)_{i,j}$$
 and $\partial_{\overline{\tau}} = \left(\frac{1+\delta_{ij}}{2}\partial_{\overline{\tau},ij}\right)_{i,j}$

where δ_{ij} is the Dirac symbol.

Lowering operators. For holomorphic weight σ , we define the lowering operator

$$\mathbf{L} = \mathbf{L}^{(n)} = \mathbf{L}^{(n)}_{\sigma} : f \mapsto y^{\mathsf{t}} (y\partial_{\overline{\tau}}) \otimes f.$$
(6.6)

Proposition 6.1. Let σ be a complex representation of $GL_n(\mathbb{C})$. Then $L_{\sigma} \in \mathbb{D}(\sigma, \operatorname{sym}^{\vee 2} \otimes \sigma)$.

Remark 6.2. Note that in the genus 1 case, sym^{$\vee 2$} is the same as det⁻². Thus (6.6) reproduces the classical Maass raising operator (see p. 177 of [Maa71]), which is $L_k = y^2 \partial_{\overline{\tau}}$ up to normalizing, multiplicative scalars.

Lemma 6.3. We have

$$y \circ \operatorname{inv} = \overline{\tau}^{-1} y \tau^{-1} = \tau^{-1} y \overline{\tau}^{-1}.$$

Proof. Since $y = \frac{1}{2i}(\tau - \overline{\tau})$, we have $y \circ inv = \frac{1}{2i}(-\tau^{-1} + \overline{\tau}^{-1}) = \tau^{-1}y\overline{\tau}^{-1}$.

Proof of Proposition 6.1. Since σ is, by assumption, holomorphic, application of the cocycle associated with σ commutes with application of $\partial_{\overline{\tau}}$. We can therefore reduce our considerations to $f \in C^{\infty}(\mathbb{H}^{(n)})$ and the trivial weight $\sigma = \mathbb{1}$. This, in particular, simplifies notation: Instead of $y^{t}(y\partial_{\overline{\tau}}) \otimes f$ we can write $y(\partial_{\overline{\tau}} f)y$.

It is clear that L intertwines trans(*b*) \in G^(*n*) for all *b* \in Mat^t_{*n*}(\mathbb{R}). Compatibility of L with the action of inv thus implies the statement. Using $y \circ inv = \overline{\tau}^{-1}y\tau^{-1} = \tau^{-1}y\overline{\tau}^{-1}$ (see Lemma 6.3) and $(\partial_{\overline{\tau}}f) \circ inv = \overline{\tau}(\partial_{\overline{\tau}}(f \circ inv))\overline{\tau}$, verified in Equation (6.7) below, we find that

$$y(\partial_{\overline{\tau}}f)y|_{\mathrm{sym}^{\vee 2}}\operatorname{inv} = \tau \Big(\tau^{-1}y\overline{\tau}^{-1}\overline{\tau}\big(\partial_{\overline{\tau}}(f\circ\operatorname{inv})\big)\overline{\tau}\overline{\tau}^{-1}y\tau^{-1}\Big)\tau = y\big(\partial_{\overline{\tau}}(f\circ\operatorname{inv})\big)y.$$

This proves the proposition after we have checked

$$(\partial_{\tau} f) \circ \operatorname{inv} = \tau (\partial_{\tau} (f \circ \operatorname{inv})) \tau.$$
(6.7)

Consider

$$\left(\partial_{\tau}(f\circ\mathrm{inv})\right)(\tau) = \partial_{\overline{\tau}}f\left(-\tau^{-1}\right) = \sum_{i,j} \mathfrak{e}_{ij} \frac{1+\delta_{ij}}{2}\overline{\partial}_{ij}f\left(-\tau^{-1}\right) = \sum_{i,j,k,l} \mathfrak{e}_{ij}\left(\frac{1+\delta_{kl}}{2}\overline{\partial}_{kl}f\right)\left(-\tau^{-1}\right) \frac{1+\delta_{ij}}{2}\overline{\partial}_{ij}\left(-\overline{\tau}^{-1}\right).$$

The inner differential $\partial_{\overline{\tau}}(-\overline{\tau}^{-1})$ can be computed using the product rule, giving

$$0 = \partial_{\tau} \otimes (\tau \tau^{-1}) = (\partial_{\tau} \otimes \tau) (\mathbb{1}_n \otimes \tau^{-1}) + (\mathbb{1}_n \otimes \tau) (\partial_{\tau} \otimes \tau^{-1}),$$

which yields

$$\partial_{\tau} \otimes \tau^{-1} = - \big(\mathbb{1}_n \otimes \tau^{-1} \big) \Big(\sum_{i,j} \mathfrak{e}_{ij} \otimes \frac{1}{2} (\mathfrak{e}_{ij} + \mathfrak{e}_{ji}) \Big) \big(\mathbb{1}_n \otimes \tau^{-1} \big).$$

Symmetry of τ allows us to shift the occurrence of τ^{-1} from the second to the first tensor component:

$$\partial_{\tau} \otimes \tau^{-1} = - \big(\tau^{-1} \otimes \mathbb{I}_n \big) \big(\sum_{i,j} \mathfrak{e}_{ij} \otimes \frac{1}{2} (\mathfrak{e}_{ij} + \mathfrak{e}_{ji}) \big) \big(\tau^{-1} \otimes \mathbb{I}_n \big).$$

This establishes the transformation behavior in (6.7).

The raising operator for the standard representation. To deduce raising operators from lowering operators, we will use the intertwining operators

$$C_{\text{std}}f = yf$$
 and $C_{\text{sym}^2}f = yfy.$ (6.8)

Lemma 6.4. The operator C_{std} is covariant from std to std^V, and C_{sym^2} is covariant from sym² to sym^{V2}.

Proof. Covariance of Cstd follows from

$$y\overline{f}|_{\mathrm{std}^{\vee}}g = {}^{\mathrm{t}}(c\tau+d) {}^{\mathrm{t}}(c\tau+d)^{-1}y(c\overline{\tau}+d)^{-1}\overline{f(g\tau)} = y\overline{(c\tau+d)^{-1}f(g\tau)}$$

Covariance of C_{sym^2} can be established by a similar computation.

Remark 6.5. The inverses of the above operators are given explicitly by

$$C_{\text{std}^{\vee}} f = C_{\text{std}}^{-1} f = y^{-1} \overline{f} \text{ and } C_{\text{sym}^{\vee 2}} f = C_{\text{sym}^2}^{-1} f = y^{-1} \overline{f} y^{-1}.$$
 (6.9)

We set

$$R_{\rm std} = C_{\rm sym^2}^{-1} \otimes C_{\rm std}^{-1} \circ L_{\rm std^{\vee}} \circ C_{\rm std} \quad \text{and} \quad R_{\rm std^{\vee}} = C_{\rm sym^2}^{-1} \otimes C_{\rm std} \circ L_{\rm std} \circ C_{\rm std}^{-1}.$$
(6.10)

From the covariance properties of the defining operators, we deduce that

$$R_{\rm std} \in \mathbb{D}\left({\rm std}, {\rm sym}^{\vee 2} \otimes {\rm std}\right) \quad \text{and} \quad R_{\rm std^{\vee}} \in \mathbb{D}\left({\rm std}^{\vee}, {\rm sym}^{\vee 2} \otimes {\rm std}^{\vee}\right). \tag{6.11}$$

Recall that we denote the canonical basis of V(std) and $V(\text{sym}^2)$ by e_i $(1 \le i \le n)$ and e_{ij} $(1 \le i, j \le n)$, respectively. In a formal way, we write $t_{(12)3} : V(\text{sym}^2 \otimes \text{std}) \to V(\text{sym}^2 \otimes \text{std})$ for the map that descends from

$$\frac{1}{2}(\mathfrak{t}_{13} + \mathfrak{t}_{23}): V(\mathsf{std} \otimes \mathsf{std} \otimes \mathsf{std}) \longrightarrow V(\mathsf{std} \otimes \mathsf{std} \otimes \mathsf{std})$$

$$\mathfrak{e}_l \otimes \mathfrak{e}_m \otimes \mathfrak{e}_n \longmapsto \frac{1}{2}(\mathfrak{e}_n \otimes \mathfrak{e}_m \otimes \mathfrak{e}_l + \mathfrak{e}_l \otimes \mathfrak{e}_n \otimes \mathfrak{e}_m).$$
(6.12)

Proposition 6.6. For $f \in C^{\infty}(\mathbb{H}^{(n)} \to V(\text{std}))$, we have

$$\mathbf{R}_{\mathrm{std}} f = \partial_{\tau} \otimes f - \frac{i}{2} \mathbf{t}_{(12)3} (y^{-1} \otimes f), \tag{6.13}$$

and for $f \in C^{\infty}(\mathbb{H}^{(n)} \to V(\mathrm{std}^{\vee}))$, we have

$$\mathbf{R}_{\mathrm{std}^{\vee}} f = \partial_{\tau} \otimes f + \frac{i}{2} \mathbf{t}_{(12)3} \big(\mathbb{1}_n \otimes y^{-1} f \big).$$
(6.14)

Proof. Denote coordinates of f corresponding to the e_i by f_i . To compute R_{std} , we start by computing $L_{std^{\vee}} \circ C_{std} f$, which equals

$$\sum_{i,j,k,l} \mathfrak{e}_{ij} y_{ik} y_{jl} \frac{1+\delta_{kl}}{2} \overline{\partial}_{kl} \sum_{m,n} \mathfrak{e}_m y_{mn} \overline{f_n} = \sum_{i,j,m} \mathfrak{e}_{ij} \mathfrak{e}_m \Big(\sum_{k,l,n} y_{ik} y_{jl} y_{mn} \frac{1+\delta_{kl}}{2} \overline{\partial}_{kl} \overline{f_n} + \frac{i}{4} \sum_n \big(y_{im} y_{jn} + y_{in} y_{jm} \big) \overline{f_n} \Big).$$

Recall that we write y_{ij}^{-1} for the ij-th entry of y^{-1} . Now, $C_{sym^{\vee 2}} \otimes C_{std^{\vee}}$ maps the previous expression to

$$\sum_{i',j',m'} \mathfrak{e}_{i'j'} \mathfrak{e}_{m'} y_{ii'}^{-1} y_{jj'}^{-1} y_{mm'}^{-1} \sum_{i,j,m} \left(\sum_{k,l,n} y_{ik} y_{jl} y_{mn} \frac{1+\delta_{kl}}{2} \partial_{kl} f_n - \frac{i}{2} \sum_n \frac{1}{2} (y_{im} y_{jn} + y_{in} y_{jm}) f_n \right)$$

= $\sum_{k,l,n} \mathfrak{e}_{kl} \mathfrak{e}_n \frac{1+\delta_{kl}}{2} \partial_{kl} f_n - \frac{i}{4} \sum_{m,n,m'} \mathfrak{e}_{mn} \mathfrak{e}_{m'} (y_{mm'}^{-1} f_n + y_{nm'}^{-1} f_m).$

The expression for $R_{std^{\vee}}$ can be derived in an analogous way.

Differential operators for tensor products of weights. To extend our considerations to arbitrary representations, we have to study how raising operators combine when taking the tensor product of two weights. For a representation σ , the order 1 raising operator R_{σ} is of the form

$$\mathbf{R}_{\sigma} f = \partial_{\tau} f + \mathbf{K}_{\sigma} f, \quad \text{for} \quad \mathbf{K}_{\sigma} \in \mathbf{C}^{\infty} \big(\mathbb{H}^{(n)} \to \operatorname{Hom} \big(V(\sigma), V(\operatorname{sym}^{2} \otimes \sigma) \big) \big).$$

Given two representations σ and σ' , we write ι_{σ} for the inclusion

$$\iota_{\sigma} : \operatorname{Hom}\left(V(\sigma), V(\operatorname{sym}^{2} \otimes \sigma)\right) \longrightarrow \operatorname{Hom}\left(V(\sigma \otimes \sigma'), V(\operatorname{sym}^{2} \otimes \sigma \otimes \sigma')\right)$$
$$\phi \longmapsto \phi \otimes \operatorname{id.}$$

Similarly, $\iota_{\sigma'}$ is defined as an inclusion of Hom $(V(\sigma'), V(\text{sym}^2 \otimes \sigma'))$ into the same codomain.

Lemma 6.7. Using above notation, we have

$$\mathbf{K}_{\sigma \otimes \sigma'} = \iota_{\sigma} (\mathbf{K}_{\sigma}) + \iota_{\sigma'} (\mathbf{K}_{\sigma'}).$$

Proof. A computation gives for $K_{\sigma \otimes \sigma'}(f|_{\sigma \sigma'}g)$ the expression

$$\left(\partial_{\tau} \otimes \sigma(c\tau+d)^{-1}\right) \otimes \sigma'(c\tau+d)^{-1} f(g\tau) + t_{12} \left(\sigma(c\tau+d)^{-1} \otimes \left(\partial_{\tau} \otimes \sigma'(c\tau+d)^{-1}\right)\right) f(g\tau)$$

+ $K_{\sigma \otimes \sigma'} \left(\sigma(c\tau+d)^{-1} \otimes \sigma'(c\tau+d)^{-1} f(g\tau)\right).$

The transposition t_{12} , defined in (6.3), switches the first and second component of $\sigma \otimes \text{sym}^2 \otimes \sigma'$. By definition, we have

$$\left(\partial_{\tau} \otimes \sigma (c\tau + d)^{-1}\right) f(g\tau) + \mathcal{K}_{\sigma} \left(\sigma (c\tau + d)^{-1} f(g\tau)\right) = \left(\mathcal{K}_{\sigma} f\right)|_{\operatorname{sym}^{2} \otimes \sigma} g,$$

implying that

$$\left(\partial_{\tau} \otimes \sigma(c\tau+d)^{-1}\right) \otimes \sigma'(c\tau+d)^{-1} f(g\tau) + \iota_{\sigma}(\mathbf{K}_{\sigma}) \left(\sigma(c\tau+d)^{-1} \otimes \sigma'(c\tau+d)^{-1} f(g\tau)\right)$$

equals $(\iota_{\sigma}(K_{\sigma}) f)|_{\text{sym}^2 \otimes \sigma \otimes \sigma'} g$. The analogue equation holds for $K_{\sigma'}$ and its image under $\iota_{\sigma'}$. After plugging this into the first equality and comparing terms, this implies the lemma.

Let $a, b, c, i_1, ..., i_b$, and $j_1, ..., j_c$ be as in (6.4). The raising operators for powers of std^V involve the following modification of $t_{(i_1,...,i_b)(j_1,...,j_c)}$:

$$\mathbf{t}_{(i_1,\dots,i_b)(j_1,\dots,j_c)}^{\mathbf{y}} = \frac{1}{bc} \sum_{k=1}^{b} \sum_{l=1}^{c} \mathbf{t}_{i_k j_l} \circ \mu_{j_l}^{\mathbf{y}}, \tag{6.15}$$

where μ_{j}^{γ} is multiplication with y^{-1} in the *j*-th component.

Corollary 6.8. Explicit expressions for the corvariant differential operators

$$R_{std^{l}} \in \mathbb{D}(std^{l}, sym^{\vee 2} \otimes std^{l}), \quad R_{std^{\vee l}} \in \mathbb{D}(std^{\vee l}, sym^{\vee 2} \otimes std^{\vee l}), \quad and$$

$$R_{std^{l_{1}}std^{\vee l_{2}}} \in \mathbb{D}(std^{l_{1}}std^{\vee l_{2}}, sym^{\vee 2} \otimes std^{l_{1}}std^{\vee l_{2}})$$
(6.16)

are

$$R_{\text{std}^{l}} f = \partial_{\tau} \otimes f - \frac{il}{2} t_{(12)(3\cdots 2+l)}(y^{-1} \otimes f), \qquad R_{\text{std}^{\vee l}} f = \partial_{\tau} \otimes f + \frac{il}{2} t_{(12)(3\cdots 2+l)}^{\vee}(\mathbb{1}_{n} \otimes f), \quad and$$

$$R_{\text{std}^{l_{1}} \text{std}^{\vee l_{2}}} = \partial_{\tau} \otimes f - \frac{il_{1}}{2} t_{(12)(3\cdots 2+l_{1})}(y^{-1} \otimes f) + \frac{il_{2}}{2} t_{(12)(3+l_{1}\cdots 2+l_{1}+l_{2})}^{\vee}(\mathbb{1}_{n} \otimes f). \tag{6.17}$$

The Lie algebra of Sp_{*n*}. To describe $\mathbb{D}(\sigma, \operatorname{sym}^2 \sigma)$ and $\mathbb{D}(\sigma, \operatorname{sym}^{\vee 2} \sigma)$, we employ the Lie theoretic description of differential operators given by Helgason [Hel59; Hel92] and, in context of discrete series representations, by Schmid [Sch76].

Let \mathfrak{sp}_n denote the complexified Lie algebra of $G^{(n)} = \operatorname{Sp}_n(\mathbb{R})$. Fix

$$\mathfrak{t}_n = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} : a = -{}^{\mathrm{t}}a \in \mathrm{Mat}_n(\mathbb{C}), \ b = {}^{\mathrm{t}}b \in \mathrm{Mat}_n(\mathbb{C}) \right\},\tag{6.18}$$

which is the complexified Lie algebra of the maximal compact subgroup

$$\mathbf{K}_n = \left\{ \begin{pmatrix} a & b \\ -b & a \end{pmatrix} : a, b \in \mathrm{Mat}_n(\mathbb{R}), ai + b \in \mathbf{U}_n(\mathbb{R}) \right\} \subseteq \mathrm{Sp}_n(\mathbb{R})$$

Denote the center of \mathfrak{k}_n by $\mathfrak{k}_{n,\mathbb{C}}$ or, suppressing the genus, $\mathfrak{k}_{\mathbb{C}}$. To a representation σ of $GL_n(\mathbb{C})$ we can attach a representation of \mathfrak{k}_n by taking the differential of its restriction to K_n . Since K_n is connected, we can and will refer to representations of \mathfrak{k}_n by corresponding representations of $GL_n(\mathbb{C})$.

As a \mathfrak{k}_n module, we have the decomposition

$$\mathfrak{sp}_n \cong \mathfrak{k}_n \oplus \mathfrak{p}_n^+ \oplus \mathfrak{p}_n^-, \tag{6.19}$$

where \mathfrak{p}_n^{\pm} is the differential of the K_n representations sym² and sym^{$\vee 2$}. A suitable generator of $\mathfrak{k}_{\mathbb{C}}$ acts on them by ± 2 . We will throughout refer to eigenvalues of this generator as $\mathfrak{k}_{\mathbb{C}}$ -eigenvalues.

We denote invariants of a Lie algebra representation V of \mathfrak{k}_n by $\mathrm{H}^0(\mathfrak{k}_n, V)$. Fix two representation σ and σ' of $\mathrm{GL}_n(\mathbb{C})$. Write $\mathbb{D}_o(\sigma, \sigma')$ for the space of order o differential operators that are covariant from $|_{\sigma}$ to $|_{\sigma'}$. In Theorem 10 of [Hel59], Helgason described them by the following vector space isomorphism:

$$\mathbb{D}_{o}(\sigma,\sigma') \cong \mathrm{H}^{0}(\mathfrak{k}_{n}, \operatorname{sym}^{o}(\mathfrak{p}_{n}^{+} \oplus \mathfrak{p}_{n}^{-}) \otimes \sigma^{\vee} \otimes \sigma').$$
(6.20)

We reiterate Helgson's remark that this isomorphism is not compatible with composition of differential operators.

Helgason's result allows us to restrict the order of covariant differential operators.

Proposition 6.9. *Fix irreducible representations* σ *and* σ' *of* $GL_n(\mathbb{C})$ *.*

- (*i*) If the $\mathfrak{k}_{\mathbb{C}}$ -eigenvalues of σ and σ' agree, then $\mathbb{D}_1(\sigma, \sigma') = \mathbb{D}_0(\sigma, \sigma')$.
- (ii) If the $\mathfrak{k}_{\mathbb{C}}$ -eigenvalues of σ and σ' differ by ± 2 , then $\mathbb{D}_2(\sigma, \sigma') = \mathbb{D}_1(\sigma, \sigma')$.
- (*iii*) We have $\mathbb{D}_0(\sigma, \sigma') \cong \operatorname{Hom}(\sigma, \sigma')$.
- (iv) The space $\mathbb{D}_1(\sigma, \sigma')$ is spanned by operators $\pi_L \circ L_\sigma$ and $\pi_R \circ R_{\sigma'}$, where $\pi_L \in \text{Hom}(\text{sym}^2\sigma, \sigma')$ and $\pi_R \in \text{Hom}(\text{sym}^{\vee 2}\sigma, \sigma')$, respectively.

Proof. Case (iii) is a direct consequence of (6.20). To prove case (i) and (ii), it suffices to compute $\mathfrak{t}_{\mathbb{C}}$ -eigenvalues in sym^{*o*}($\mathfrak{p}_n^+ \oplus \mathfrak{p}_n^-$) $\otimes V(\sigma)^{\vee} \otimes \sigma^{\vee} \otimes \sigma'$ for $o \in \{1, 2\}$.

The $\mathfrak{k}_{\mathbb{C}}$ -eigenvalues of $\mathbb{1} \cong \operatorname{sym}^{0}(\mathfrak{p}_{n}^{+} \oplus \mathfrak{p}_{n}^{-})$ and $\mathfrak{p}_{n}^{+} \otimes \mathfrak{p}_{n}^{-} \subset \operatorname{sym}^{2}(\mathfrak{p}_{n}^{+} \oplus \mathfrak{p}_{n}^{-})$ are 0. On \mathfrak{p}_{n}^{+} and \mathfrak{p}_{n}^{-} they are ± 2 , by definition, and on $\operatorname{sym}^{2}\mathfrak{p}_{n}^{+}$ and $\operatorname{sym}^{2}\mathfrak{p}_{n}^{-}$ they equal ± 4 . In case (i) the $\mathfrak{k}_{\mathbb{C}}$ -eigenvalue of $\sigma^{\vee} \otimes \sigma'$ equals $-a + a \pm 2 = \pm 2$ for some $a \in \mathbb{C}$, and in case (ii) it equals 0. Combining these values we deduce all cases but (iv).

To establish the remaining statement (iv), note that the case of lowering operators is implied by the isomorphism

$$\mathrm{H}^{0}(\mathfrak{k}_{n}, \operatorname{sym}^{\vee 2} \otimes \sigma^{\vee} \otimes \sigma') \cong \mathrm{Hom}_{\mathfrak{k}_{n}}(\operatorname{sym}^{2} \otimes \sigma, \sigma').$$

The case of raising operators can be dealt with analogously.

Commutation relations. By abuse of notation, we denote two projections by the same symbol:

$$\pi_{\rm sym^2} : \rm sym^2 \otimes \rm sym^{\vee 2} \otimes \sigma, \, \rm sym^{\vee 2} \otimes \rm sym^2 \otimes \sigma \longrightarrow \sigma.$$
(6.21)

To concisely state commutation relations of differential operators, we suppress the composition symbol: $LR_{\sigma} = L \circ R_{\sigma}$. Powers of operators will be understood as repeated composition: $L^{d} = L \circ \cdots \circ L$ for $0 \le d \in \mathbb{Z}$.

Lemma 6.10. We have

$$\pi_{\operatorname{sym}^2} \operatorname{LR}_{\operatorname{sym}^{\vee 2} \otimes \sigma} \operatorname{L}_{\sigma} - \operatorname{L} \pi_{\operatorname{sym}^2} \operatorname{R}_{\operatorname{sym}^{\vee 2} \otimes \sigma} \operatorname{L}_{\sigma} \in \mathbb{D}_1(\sigma, \operatorname{sym}^2 \otimes \sigma).$$

In particular, it can be expressed as a linear combination of constituents of L_{σ} .

Proof. It is straightforward to see that the given expression is a differential operator of order at most 2, since symbols of differential operators commute. We may assume that σ is irreducible. Denote its $\mathfrak{k}_{\mathbb{C}}$ eigenvalue by *a*. Further, observe that the $\mathfrak{k}_{\mathbb{C}}$ eigenvalues of sym^{$\vee 2$} \otimes sym^{$\vee 2$} \otimes sym^{$\vee 2$} $\otimes \sigma$ equals a-1. By Helgason's theory as stated in Proposition 6.9, the left hand side is an order 1 differential operator, which is a linear combination of the constituents of L_{σ} by the same proposition.

Lemma 6.11. We have

$$L_{sym^2\sigma}R_{\sigma} - t_{12}R_{sym^{\vee 2}\sigma}L_{\sigma} \in Hom(\sigma, sym^2sym^{\vee 2}\sigma).$$

Proof. The left hand side is a covariant differential operator of order at most 1 that preserves $\mathfrak{k}_{\mathbb{C}}$ eigenvalues. Therefore, we can deduce the statement from Proposition 6.9.

Lemma 6.12. Let σ be a representation with $\mathfrak{k}_{\mathbb{C}}$ eigenvalue a. Then

$$\pi_{\operatorname{sym}^2}\left(\mathcal{L}_{\operatorname{sym}^2\sigma}\operatorname{R}_{\sigma} - \operatorname{t}_{12}\operatorname{R}_{\operatorname{sym}^{\vee 2}\sigma}\operatorname{L}_{\sigma}\right) = \frac{-(1+n)\,a}{8}.$$

Proof. By Lemma 6.11, the left hand side lies in Hom(σ , σ). We may assume that σ is irreducible, so that Hom(σ , σ) $\cong \mathbb{C}$. To compute the precise value, note that in the following expression for LR – t₁₂RL

$$(y^{t}(y\partial_{\overline{\tau}}))(\partial_{\tau} + K_{\sigma}) - t_{12}((\partial_{\tau} + K_{sym^{\vee 2}\sigma})(y^{t}(y\partial_{\overline{\tau}})))$$

order 0 terms can only arise from $y^{t}(y\partial_{\overline{\tau}})K_{\sigma}$. Using Lemma 6.7, we reduce ourselves to the cases $\sigma = \text{std}$ and $\sigma = \text{std}^{\vee}$. We focus on the former one, and leave the latter one to the reader. For any $1 \le o \le n$, we have

$$y^{\mathfrak{t}}(y\partial_{\overline{\tau}}) \otimes \frac{-i}{2} \mathfrak{t}_{(12)3}(y^{-1} \otimes \mathfrak{e}_{o}) = \frac{-i}{2} \sum_{i,j,k,l} \mathfrak{e}_{ij} y_{ik} y_{jl} \frac{1+\delta_{kl}}{2} \overline{\partial}_{kl} \sum_{m,n,o} \frac{1}{2} \big(\mathfrak{e}_{mo} \mathfrak{e}_{n} + \mathfrak{e}_{on} \mathfrak{e}_{m} \big) y_{mn}^{-1}$$
$$= \frac{1}{4} \sum_{i,j,k,l,m,n} \mathfrak{e}_{ij} y_{ik} y_{jl} \frac{1}{2} \big(\mathfrak{e}_{mo} \mathfrak{e}_{n} + \mathfrak{e}_{on} \mathfrak{e}_{m} \big) \frac{-1}{2} \big(y_{km}^{-1} y_{ln}^{-1} + y_{kn}^{-1} y_{lm}^{-1} \big).$$

We execute multiplication of *y* and y^{-1} twice

$$\frac{-1}{16}\sum_{i,j,m,n}\mathfrak{e}_{ij}\left(\mathfrak{e}_{mo}\mathfrak{e}_{n}+\mathfrak{e}_{on}\mathfrak{e}_{m}\right)\left(\delta_{im}\delta_{jn}+\delta_{in}\delta_{jm}\right)=\frac{-1}{16}\sum_{i,j}\mathfrak{e}_{ij}\left(\mathfrak{e}_{io}\mathfrak{e}_{j}+\mathfrak{e}_{oj}\mathfrak{e}_{i}+\mathfrak{e}_{jo}\mathfrak{e}_{i}+\mathfrak{e}_{oi}\mathfrak{e}_{j}\right).$$

Applying π_{svm^2} , we obtain

$$\frac{-1}{16}\sum_{i,j} \left(\delta_{jo} + \delta_{io} + \delta_{ij} \delta_{jo} \mathfrak{e}_o + \delta_{io} \delta_{ij} \right) \mathfrak{e}_o = \frac{-(1+n)}{8} \mathfrak{e}_o.$$

§6.3 Almost holomorphic functions. We throughout write $\mathcal{O}(\mathbb{H}^{(n)})$ for the space of holomorphic functions on $\mathbb{H}^{(n)}$.

Definition 6.13. A smooth function $f : \mathbb{H}^{(n)} \to V(\sigma)$ is almost holomorphic of depth *d* if $\mathbb{L}^{d+1} f = 0$.

Remark 6.14. This definition is motivated by the theory of holomorphic discrete series, whose behavior under differential operators is well-known [Sch76]. Very recently, a work that focuses on this perspective has appear as a preprint: [PSS15].

The following are basic properties of almost holomorphic functions on $\mathbb{H}^{(n)}$ which are straight forward to prove.

Proposition 6.15. Suppose that $f : \mathbb{H}^{(n)} \to V(\sigma)$ and $g : \mathbb{H}^{(n)} \to V(\sigma')$ are almost holomorphic functions of depth d and d'. Then the tensor product $f \otimes g$ is almost holomorphic of depth d + d'.

Proof. We abbreviate d'' = d + d'. It suffices to inspect the equation $L^{d''+1}f \otimes g = \sum_{t=0}^{d''+1} t_t ((L^t f) \otimes (L^{d''+1-t}g))$ with suitable permutations of tensor components $t_t : (sym^2)^t \sigma(sym^2)^{d''+1-t}\sigma' \to (sym^2)^{d''+1}\sigma\sigma'$.

Lemma 6.16. If f is almost holomorphic of depth d, then π_{sym^2} LRL $f - L\pi_{sym^2}$ RL f has depth d - 1.

Proof. This is a consequence of Lemma 6.10, seen that f, which has depth d, vanishes after applying d + 1 times arbitrary constituents of L.

Analytic properties. Almost holomorphic functions in the setting of elliptic modular forms can be easily recognized as a polynomial in y^{-1} whose coefficients are holomorphic functions. Their total degree coincides with the depth.

Theorem 6.17. A function $f \in C^{\infty}(\mathbb{H} \to V(\sigma))$ is almost holomorphic of depth d if and only if it is a polynomial in the entries of y^{-1} of degree at most d and with holomorphic coefficients.

Proof. Note that Ly^{-1} is constant. After embedding σ into $(sym^2 sym^{\vee 2})^d \otimes \sigma$, it follows that every polynomial in the entries of y^{-1} of at most degree *d* is indeed almost holomorphic of depth *d*.

We proceed as in the case of elliptic modular forms to establish the converse. Note that it suffices to treat irreducible σ . Since the lowering operator is the same for representations σ and det^k σ for any $k \in \mathbb{Z}$, we may always modify σ correspondingly. In particular, we can assume that the $\mathfrak{k}_{\mathbb{C}}$ -eigenvalue of σ is arbitrarily large. Then Lemma 6.12, shows that the space of almost holomorphic functions of depth d is contained in the span of depth d - 1 functions and their image under $\pi_{\text{sym}^2} R_{\text{sym}^2 \sigma}$. Since the case d = 0 is clear, the result follows by induction.

Theorem 6.17 tells us that an almost holomorphic function f is a polynomial in the entries of y^{-1} . For $t \in Mat_n^t(\frac{1}{2}\mathbb{Z})$ with integral diagonal entries, set $y^{-t} = \exp(\operatorname{trace}(\operatorname{tlog}(y^{-1})))$. Then f can be written as

$$f(\tau) = \sum_{t} f_t(\tau) y^{-t}$$
(6.22)

with holomorphic f_t , finitely many of which are non-zero. We call f_t the *t*-th part of *f*, and f_0 is called the constant part of *f*.

§6.4 Definition of almost holomorphic Siegel modular forms. Seen that one goal of this work is to settle basic tools for the treatment of almost holomorphic Siegel modular forms, we give a slightly more general definition, including vector valued modular Siegel modular forms with respect to representations of $\Gamma^{(n)} = \text{Sp}_n(\mathbb{Z})$, than the one that is necessary to prove Theorem 8.5. Note also that statements in this paper can be extended to representations of the metaplectic double cover $\text{Mp}_{2n}(\mathbb{Z})$ of $\text{Sp}_n(\mathbb{Z})$ without difficulties, since they depend only on analytic properties of corresponding modular forms.

Definition 6.18. Let ρ be a finite dimensional, complex representation of $\Gamma^{(n)}$ and σ a finite dimensional, complex representation of $\operatorname{GL}_n(\mathbb{C})$. An almost holomorphic function $f : \mathbb{H}^{(n)} \to V(\sigma) \otimes V(\rho)$ of depth *d* that satisfies

$$f|_{\sigma,\rho} \gamma = f$$
 for all $\gamma \in \Gamma^{(n)}$,

and, if n = 1, is bounded by some power of y^{-1} as $y \to \infty$, is called an almost holomorphic Siegel modular form of genus n, depth d, weight σ , and type ρ .

We write $M^{(n)}(\sigma^{[d]} \boxtimes \rho)$ for the space of such functions; notation which is inspired by the fact that almost holomorphic Siegel modular forms correspond to automorphic forms whose "*K*-type at the infinite place", related to σ , is not the highest possible, but *d* steps away from it. Proposition 6.15 implies that

$$\mathbf{M}^{(n)}\big(\sigma^{[d]} \boxtimes \rho\big) \otimes \mathbf{M}^{(n)}\big(\sigma'^{[d']} \boxtimes \rho'\big) \subseteq \mathbf{M}^{(n)}\big((\sigma\sigma')^{[d+d']} \boxtimes \rho\rho'\big).$$
(6.23)

An almost holomorphic Siegel modular form f has Fourier expansion

$$f(\tau) = \sum_{t \in \operatorname{Mat}_{n}^{t}(\mathbb{Q})} c(f; t, y) \exp\left(2\pi i \operatorname{trace}(t\tau)\right)$$
(6.24)

where every Fourier coefficient c(f; t, y) is a polynomial in the entries of y^{-1} with coefficients in \mathbb{C} .

7 | Fourier-Jacobi expansions

Fourier-Jacobi expansions of Siegel modular forms have developed to a major tool to study Siegel modular forms. They have been abstractly been considered in [EZ85]. Roughly, a Siegel modular form f can be written as

$$f(\tau) = \sum_{m \in \mathbb{Z}} \phi_m(\tau_1, z) \exp(2\pi i \, m \tau_2)$$

where we use, here and throughout, the decomposition $\tau = \begin{pmatrix} \tau_1 & \tau_2 \\ z & \tau_2 \end{pmatrix}$ with $\tau_1 \in \mathbb{H}^{(n-1)}$, $\tau_2 \in \mathbb{H}^{(1)}$, and $z \in \mathbb{C}^{n-1}$. Each ϕ_m is a Jacobi form, which can be related to Siegel modular forms of genus n-1. The theory of Fourier-Jacobi expansions has recently culminate in [BWR14], where the last named author and his coauthor prove that such formal expansions converge, if they exhibit the symmetry relations of Fourier coefficients that hold for Fourier expansions of Siegel modular forms.

In this section, we present the foundations of Fourier Jacobi expansions of almost holomorphic Siegel modular forms. In particular, we attach almost holomorphic Fourier-Jacobi coefficients to almost holomorphic Siegel modular forms and show that they determine them uniquely. In the case of genus n = 2, we characterize the spaces of almost holomorphic Jacobi forms using the approach in [IK11]. Note that it is possible to extend this to arbitrary genera, but the focus of this paper lies on n = 2.

§7.1 Preliminaries. Jacobi forms were defined in [EZ85] and generalized in [Zie89]. Here, we revisit basic notation.

The Jacobi upper half space. Write $\mathbb{H}^{(n)J} = \mathbb{H}^{(n)} \times \mathbb{C}^n = \{(\tau, z) : \tau \in \mathbb{H}^{(n)}, z \in \mathbb{C}^n\}$ for the Jacobi upper half space of genus *n* (and cogenus 1). The real and imaginary part of τ and *z* are denoted by $\tau = x + iy$ and z = u + iv. If n = 1, then we suppress the genus, writing \mathbb{H}^J instead of $\mathbb{H}^{(n)J}$.

The Heisenberg group. Fix the Heisenberg group

$$H(\mathbb{R}^n) = \left\{ h = (\lambda, \mu, \kappa) : \lambda, \mu \in \mathbb{R}^n, \kappa \in \operatorname{Mat}_n(\mathbb{R}) \right\}$$

with addition

$$(\lambda_1, \mu_1, \kappa_1) + (\lambda_2, \mu_2, \kappa_2) = (\lambda_1 + \lambda_2, \mu_1 + \mu_2, \kappa_1 + \kappa_2 + \lambda_1^{t} \mu_2 - \mu_1^{t} \lambda_2).$$

It carries a right action of $G^{(n)}$ defined by the natural action on row vectors of length 2*n*:

$$(\lambda, \mu, \kappa) \begin{pmatrix} a & b \\ c & d \end{pmatrix} = (\lambda a + \mu c, \lambda b + \mu d, \kappa).$$

The Jacobi group. The real Jacobi group $G^{(n)J} = G^{(n)J}(\mathbb{R})$ is the non-trivial extension $0 \to H(\mathbb{R}^n) \to G^{(n)J} \to G^{(n)} \to 1$ of $G^{(n)}$ by $H(\mathbb{R}^n)$. Multiplication in $G^{(n)J}$ is defined by

$$(g_1, h_1)(g_2, h_2) = (g_1g_2, h_1g_2 + h_1).$$

We often consider elements of $H(\mathbb{R}^n)$ as elements of $G^{(n)J}$ via the inclusion $H(\mathbb{R}^n) \subset G^{(n)J}$. Shorthand notation for elements of $G^{(n)}$ considered as elements of $G^{(n)J}$ is provided by the following section to $G^{(n)J} \rightarrow G^{(n)}$:

$$g \in \mathbf{G}^{(n)} \longmapsto (g, (0, 0, 0)) \in \mathbf{G}^{(n)\mathbf{J}}$$

If n = 1, we drop the superscript (*n*), writing G^J instead of $G^{(1)J}$. Recall the shorthand notation a, b, c, d for the entries of $g \in G^{(n)}$. Similarly, the entries of $h \in H(\mathbb{R}^n)$ will be denoted by λ , μ , and κ without further mentioning it.

We have an action of $G^{(n)J}$ on $\mathbb{H}^{(n)J}$:

$$g^{\rm J}(\tau,z) = \left(g,(\lambda,\mu,\kappa)\right)(\tau,z) = \left((a\tau+b)(c\tau+d)^{-1},(z+\lambda\tau+\mu)(c\tau+d)^{-1}\right).$$
(7.1)

Remark 7.1. The connection between Jacobi forms and Siegel modular forms, that we are going to exhibit, is based on the embedding

$$G_{\text{sym}}^{(n)J} \hookrightarrow G^{(n)}, \quad \left(\begin{pmatrix} a & b \\ c & d \end{pmatrix}, (\lambda, \mu, \kappa) \right) \longmapsto \begin{pmatrix} a & 0 & b & a^{t}\mu - b^{t}\lambda \\ \lambda & 1 & \mu & \kappa \\ c & 0 & d & c^{t}\mu - d^{t}\lambda \\ 0 & 0 & 0 & 1 \end{pmatrix},$$
(7.2)

where $G_{\text{sym}}^{(n)J}$ is the subgroup of $G^{(n)J}$ such that $\kappa + \mu^{t}\lambda$ is symmetric. Details can be found in [Zie89].

The (discrete) Jacobi group $\Gamma^{(n)J} \subset G^{(n)J}$ is defined as

$$\Gamma^{(n)\mathsf{J}} = \big\{ \gamma^{\mathsf{J}} = \big(\gamma, (\lambda, \mu, 0)\big) : g \in \Gamma^{(n)}, \, \lambda, \mu \in \mathbb{Z}^n \big\}.$$

Slash actions. Let σ be a finite dimensional, complex representation of the affine group

$$\operatorname{Aff}_{n}(\mathbb{C}) = \operatorname{GL}_{n}(\mathbb{C}) \ltimes \mathbb{C}^{n}.$$
(7.3)

We attach to it a slash action on $C^{\infty}(\mathbb{H}^{(n)J} \to V(\sigma))$ by

$$(\phi|_{\sigma,m}^{J} g^{J})(\tau,z) = \sigma (c\tau + d, c^{t}z + c^{t}\mu - d^{t}\lambda)^{-1}$$

$$+ \exp \left(2\pi i m \left(-(z + \lambda\tau + \mu)c(c\tau + d)^{-1} t(z + \lambda\tau + \mu) + 2\lambda^{t}z + \lambda\tau^{t}\lambda + \operatorname{trace}(\kappa)\right)\right) \phi (g^{J}(\tau,z)).$$

$$(7.4)$$

This is called the Jacobi slash action of weight σ and index m. If $\sigma = \det^k$ for some $k \in \mathbb{Z}$, then we write $|_{k,m}^J$ for the corresponding slash action.

Weights. Via the embeddings

$$\operatorname{GL}_n(\mathbb{C}) \longrightarrow \operatorname{Aff}_n(\mathbb{C}), \qquad g \longmapsto (g,0), \text{ and}$$

 $\operatorname{Aff}_n(\mathbb{C}) \longrightarrow \operatorname{GL}_{n+1}(\mathbb{C}), \quad (g,h) \longmapsto \begin{pmatrix} g & h \\ 0 & 1 \end{pmatrix}$

representations of $\operatorname{GL}_n(\mathbb{C})$ and $\operatorname{GL}_{n+1}(\mathbb{C})$ give rise to $\operatorname{Aff}_n(\mathbb{C})$ -representations. We ambiguously use std^{*j*} and sym^{*j*} for (symmetric) powers of the standard representation of $\operatorname{GL}_n(\mathbb{C})$ for any *n*. When considering them as representations of $\operatorname{Aff}_n(\mathbb{C})$, we will add *n* or *n* + 1 as a subscript to distinguish them: std^{*j*}_{*n*} and std^{*j*}_{*n*+1} are then the representations of $\operatorname{Aff}_n(\mathbb{C})$ that arise from the first and second inclusion, respectively.

§7.2 Differential operators for genus 1 **Jacobi forms.** The first, physical, part of this paper only made use of genus 2 Siegel modular form. We have so far formulated the theory of Jacobi forms in the case of arbitrary genus *n*. For the rest of this section, we will focus on the case of genus 1 Jacobi forms (which arise from genus 2 Siegel modular forms). The main reason is that the explicit theory of differential operators has not yet been fully developed for all genera. It will be the theme of a separate paper.

We revisit differential operators, following closely the exposition in [CR10], but use different notation for the raising and lowering operators. To define differential operators, we use the following notation:

$$\begin{aligned} \partial_{\tau} &:= \frac{\partial}{\partial \tau} = \frac{1}{2} \Big(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y} \Big), \\ \partial_{\overline{\tau}} &:= \frac{\partial}{\partial \overline{\tau}} = \frac{1}{2} \Big(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y} \Big), \\ \partial_{\overline{z}} &:= \frac{\partial}{\partial \overline{z}} = \frac{1}{2} \Big(\frac{\partial}{\partial u} - i \frac{\partial}{\partial v} \Big), \end{aligned}$$

The heat operator, normalized as in [IK11], is

$$\mathbb{L}_m = \frac{4m}{2\pi i} \partial_\tau - \frac{1}{4\pi^2} \partial_z^2.$$
(7.5)

The raising operators and lowering operators with respect to Jacobi slash actions $|_{k,m}^{J}$ are given by

$$\begin{aligned} \mathbf{R}_{k,m}^{\mathrm{J}} &:= 2i \left(\partial_{\tau} + \frac{\nu}{y} \partial_{z} + 2\pi i m \frac{\nu^{2}}{y^{2}} \right) + \frac{k}{y}, \qquad \qquad \mathbf{L}_{k,m}^{\mathrm{J}} &:= -2i y \left(y \partial_{\overline{\tau}} + \nu \partial_{\overline{z}} \right), \\ \mathbf{R}_{k,m}^{\mathrm{JH}} &:= i \partial_{z} - 4\pi m \frac{\nu}{y}, \qquad \qquad \qquad \mathbf{L}_{k,m}^{\mathrm{JH}} :- i y \partial_{\overline{z}}. \end{aligned}$$

Bernd and Schmidt described their covariance properties in [BS98]. For $g \in G^J$ and $\phi \in C^{\infty}(\mathbb{H}^J)$, we have

$$L_{k,m}^{J}(\phi|_{k,m}g) = (L_{k,m}^{J}\phi)|_{k-2,m}g, \qquad R_{k,m}^{J}(\phi|_{k,m}g) = (R_{k,m}^{J}\phi)|_{k+2,m}g L_{k,m}^{JH}(\phi|_{k,m}g) = (L_{k,m}^{JH}\phi)|_{k-1,m}g, \qquad R_{k,m}^{JH}(\phi|_{k,m}g) = (R_{k,m}^{JH}\phi)|_{k+1,m}g.$$

As usual, we suppress the subscripts *k* and *m*. Note, however, that $(\mathbf{R}^{J})^{d} = \mathbf{R}_{k+2d-2,m}^{J} \cdots \mathbf{R}_{k,m}^{J}$ and $(\mathbf{R}^{JH})^{d^{H}} = \mathbf{R}_{k+d^{H}-1,m}^{JH} \cdots \mathbf{R}_{k,m}^{JH}$. The commutation rules for the raising and lowering operator are:

$$[L^{J}, L^{JH}] = [R^{J}, R^{JH}] = 0, \quad [L^{J}, R^{J}] = -k, \quad [L^{JH}, R^{JH}] = im, \quad [L^{J}, R^{JH}] = -L^{JH}, \quad [L^{JH}, R^{J}] = R^{JH}.$$
(7.6)

§7.3 Almost holomorphic functions (genus 1). Let $\mathcal{O}(\mathbb{H}^J)$ be the space of holomorphic functions on \mathbb{H}^J .

Definition 7.2. A smooth function $\phi : \mathbb{H}^{J} \to \mathbb{C}$ is almost holmorphic of depth (d, d^{H}) with $0 \le d, d^{H} \in \mathbb{Z}$ if $(L^{J})^{d} \phi = (L^{JH})^{d^{H}} \phi = 0$.

The analogues of Proposition 6.15 and Theorem 6.17 are as follows.

Proposition 7.3. Suppose that $\phi : \mathbb{H}^{J} \to \mathbb{C}$ and $\psi : \mathbb{H}^{J} \to \mathbb{C}$ are almost holomorphic functions of depth (d, d^{H}) and (d', d'^{H}) . Then the product fg is almost holomorphic of depth $(d + d', d^{H} + d'^{H})$.

Theorem 7.4. A function $\phi \in C^{\infty}(\mathbb{H}^{J})$ is almost holomorphic of depth (d, d^{H}) if and only if it is a polynomial in y^{-1} and $v y^{-1}$ of degree at most d and d^{H} , respectively, with holomorphic coefficients.

Proof. This follows from the expressions for L^J and L^{JH} , using induction. Indeed, up to non-zero multiplicative constants, $L^J y^{-a} (vy^{-1})^b$ equals $ay^{-(a-1)} (vy^{-1})^b$, and $L^{JH} y^{-a} (vy^{-1})^b$ is $by^{-a} (vy^{-1})^{b-1}$.

§7.4 Almost holomorphic Jacobi forms (genus 1). We define almost holomorphic Jacobi forms only for weight $\sigma = \det^k$, since in the general case, we have not studied the corresponding lowering operators.

Definition 7.5. An almost holomorphic function $\phi \in C^{\infty}(\mathbb{H}^{J})$ is called an almost holomorphic Jacobi form of depth (d, d^{H}) , weight k, and index m if it satisfies

- (i) $\phi|_{k,m}^{J} \gamma = \phi$ for all $\gamma \in \Gamma^{J}$.
- (ii) For all $\alpha, \beta \in \mathbb{Q}$, $\phi(\tau, \alpha\tau + \beta)$ is bounded by some power of *y* as $\tau \to i\infty$.

We write $J_{k,m}^{[d,d^H]}$ for the space of such function. The following proposition generalizes Lemma 6.3 in [BRR12]. It says that there is no analogue of the exceptional almost holomorphic elliptic modular form E_2 .

Proposition 7.6. For every $k, m \in \mathbb{Z}$ and every pair of positive integers d and d^{H} , we have

$$\mathbf{J}_{k,m}^{[d,d^{\mathrm{H}}]} = \bigoplus_{t=0}^{d} \bigoplus_{t^{\mathrm{H}}=0}^{d^{\mathrm{H}}} \left(\mathbf{R}^{\mathrm{J}}\right)^{t} \left(\mathbf{R}^{\mathrm{J}\mathrm{H}}\right)^{t^{\mathrm{H}}} \mathbf{J}_{k,m}$$

Proof. It suffices to know that there is no Jacobi form of integral weight and index that is either constant in τ or z. Then the statement follows along the lines of the proof of Lemma 6.3 in [BRR12].

§7.5 Covariant operators from vector valued weights (genus 1). Recall that we focus on the case of genus n = 1. In this section, we reinterpret a result by Ibukiyama and Kyomura [IK11] in terms of covariant operators

$$C^{J}_{\det^{k} \operatorname{sym}_{2}^{l}} : C^{\infty} \big(\mathbb{H}^{J} \to \det^{k} \operatorname{sym}_{2}^{l} \big) \longrightarrow C^{\infty} \big(\mathbb{H}^{J} \to \bigoplus_{j=0}^{l} V(\det^{k+j}) \big), \quad \phi \longmapsto \big(\iota_{0}(\phi), \dots, \iota_{l}(\phi) \big), \tag{7.7}$$

where ι_i is defined below.

Fix a basis f_j $(0 \le j \le l)$ for $V(\det^k \operatorname{sym}_2^l)$, where $\begin{pmatrix} a & 0 \\ 0 & 1 \end{pmatrix} \in G^J$ acts on f_j by multiplication with a^j . Coordinates of a function $\phi : \mathbb{H}^J \to V(\operatorname{sym}_2^l)$ with respect to this basis are denoted by $\langle \phi, f_i \rangle$. Ibukiyama and Kyomura defined the following maps ι_i on $C^{\infty}(\mathbb{H}^J \to V(\text{sym}_2^l))$ on page 795 of [IK11]:

$$\iota_{j}(\phi) = \sum_{t=0}^{\lfloor j/2 \rfloor} \sum_{j'=0}^{j-2t} \binom{l-j'}{j-2t-j'} \binom{l-j+2t}{2t} \frac{(2k+2j-2t-5)!!(2t-1)!!}{(-2m)^{j-j'}(2k+2j-5)!!} \partial_{z}^{j-2t-j'} \mathbb{L}_{m}^{t} \langle \phi, \mathfrak{f}_{j'} \rangle,$$
(7.8)

where $n!! = (2n)!/(2^n n!)$. They prove in Theorem 2.1 that

$$\iota_{\nu}(\phi|_{\det^{k}\operatorname{sym}_{\nu}^{l},m}g^{J}) = \iota_{\nu}(\phi)|_{k+\nu,m}g^{J}$$

for all $g^{J} \in G^{J}$. This, in particular, implies covariance of $C_{det^{k}sym^{l}}$.

§7.6 Fourier Jacobi coefficients. In this section, we briefly return to the case of general genus *n*. We attach Fourier Jacobi coefficients to every almost holomorphic Siegel modular form.

Decompose τ as

$$\begin{pmatrix} \tau_1 & \tau_2 \\ z & \tau_2 \end{pmatrix}$$
, $\tau_1 \in \mathbb{H}^{(n-1)}$, $\tau_2 \in \mathbb{H}$, and $z \in \mathbb{C}^{n-1}$

Write y_1 and y_2 for the imaginary part of τ_1 and τ_2 . The ambiguous notation y_1^{-1} will be used to denote $(y^{-1})_1$. Every almost holomorphic Siegel modular form has a weak Fourier Jacobi expansion

$$f(\tau) = \sum_{m \in \mathbb{Q}} \widetilde{\phi}_m(\tau_1, z, y_2) \exp(2\pi i \, m \tau_2).$$
(7.9)

We define Fourier Jacobi coefficients of f by

$$\phi_m(\tau_1, z) = \lim_{y_2 \to \infty} \widetilde{\phi}_m(\tau_1, z, y_2), \tag{7.10}$$

which is well-defined, since $y^{-1} \rightarrow \begin{pmatrix} y_1^{-1} & 0 \\ 0 & 0 \end{pmatrix}$ as $y_2 \rightarrow \infty$. If *f* has trivial type and weight σ , then $m \in \mathbb{Z}$ if $\widetilde{\phi}_m \neq 0$, and for every y_2 , $\widetilde{\phi}_m(\tau_1, z, y_2)$ is invariant under the $|_{\sigma,m}^J$ action of $\Gamma^{(n-1)J}$.

Remark 7.7. Observe that $\det y^{-1} \to 0$ as $y_2 \to \infty$. In particular, it is possible to find subexpressions of, for example, $\mathbb{R}^n f$ for a Siegel modular form f, whose term of highest depth vanishes when passing to the Fourier Jacobi expansion.

8 Classification of Almost Holomorphic Siegel Modular Forms

The graded ring of elliptic modular forms is defined as

$$\mathbf{M}(\bullet) = \mathbf{M}^{(1)}(\bullet) = \bigoplus_{0 \le k \in \mathbb{Z}} \mathbf{M}^{(1)} \big(\det^k \boxtimes \mathbb{1} \big).$$

A minimal set of (algebraically independent) generators is given by the Eisenstein series E_4 and E_6 , where

$$E_k(\tau) = 1 - \frac{2k}{B_k} \sum_{m=1}^{\infty} \sigma_{k-1}(m) \exp(2\pi i \, m\tau) \qquad k \ge 4.$$

When extending this to a differential graded algebra with derivative R acting as $R_k = \partial_\tau - \frac{ki}{2}y^{-2}$ on the *k*-th graded piece, we obtain

$$\mathbf{R}^{\infty}\mathbf{M}(\bullet) = \bigoplus_{d=0}^{\infty} \mathbf{R}^{d} \mathbf{M}(\bullet) \subset \bigcup_{0 \le d \in \mathbb{Z}} \bigoplus_{0 \le k \in \mathbb{Z}} \mathbf{M}^{(1)} \big(\det^{k[d]} \boxtimes \mathbb{1} \big).$$

This carries an action of the lowering operator L, which is not surjective. Instead, we have

$$\operatorname{coker}_{R^{\infty}M(\bullet)} L = R^{\infty}M(\bullet) / L(R^{\infty}M(\bullet)) = \operatorname{span} 1.$$
(8.1)

The (modular) weight 2 Eisenstein series is

$$E_2(\tau) = \frac{-3}{\pi} y^{-1} + 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) \exp(2\pi i \, n\tau) \in \mathcal{M}^{(1)}(\det^{2[1]}).$$
(8.2)

While E_2 is not in the range of R, the Ramanujan relation

$$\frac{1}{2\pi i} \mathbf{R} E_2 = \frac{1}{2\pi i} \mathbf{R}_{\det^2} E_2 = \frac{1}{12} \left(E_2^2 - E_4 \right)$$
(8.3)

implies that

$$\widehat{\mathbf{M}}^{(1)}(\bullet) = \bigcup_{0 \le d} \bigoplus_{0 \le k \in \mathbb{Z}} \mathbf{M}^{(1)} \left(\det^{k[d]} \right) = \bigoplus_{d=0}^{\infty} \mathbf{R}^d \left(\mathbb{C}[E_4, E_6] + E_2 \mathbb{C}[E_4, E_6] \right) = \mathbb{C} \left[E_2, E_4, E_6 \right]$$

is a differential algebra. Since $LE_2^d = \frac{3d}{2\pi i}E_2^{d-1}$ we have an exact sequence

$$0 \longrightarrow M^{(1)}(\bullet) \longrightarrow \widehat{M}^{(1)}(\bullet) \xrightarrow{L} \widehat{M}^{(1)}(\bullet) \longrightarrow 0.$$
(8.4)

The goal of this section is to show that the situation is more complicated for higher genera. We focus on the case n = 2, and find that

$$\operatorname{coker}_{\widehat{M}^{(2)}} L = \widehat{M}^{(2)} / L \widehat{M}^{(2)} = \operatorname{span} 1.$$

In particular, we prove that there is no analogue of E_2 in genus 2. In Section 9, we will then provide a replacement for it.

§8.1 The non-commutative algebra of Siegel modular forms. It is common to consider the graded ring of Siegel modular forms, summing over all spaces of scalar valued Siegel modular forms like so

$$\bigoplus_{k\in\mathbb{Z}} \mathbf{M}^{(n)}(\det^k). \tag{8.5}$$

Note that the sum runs over all isomorphism classes of complex 1-dimensional representations of $\operatorname{GL}_n(\mathbb{C})$. The above is, indeed, a ring: The tensor product of two Siegel modular forms of weight k and k' is a modular form of weight $\det^k \otimes \det^{k'}$, which is isomorphic to $\det^{k+k'}$. Further, it is possible to consistently choose bases of $V(\det^k)$ for all k, which turn all the isomorphisms $\det^k \otimes \det^{k'} \cong \det^{k+k'}$ into isomorphisms of framed representations. One simply chooses the unit vector $\mathfrak{e}^k \in V(\det^k)$ as a basis, and uses isomorphisms $\mathfrak{e}^k \otimes \mathfrak{e}^{k'} \mapsto \mathfrak{e}^{k+k'}$.

What seems like excess of precision in the case of scalar valued Siegel modular forms, becomes important in the vector valued setting. Given two representations σ and σ' of $GL_n(\mathbb{C})$ the tensor product $\sigma \otimes \sigma'$ in general decomposes into many irreducibles with multiplicities possibly larger than 1. Also, it is not trivial to consistently choose bases of all irreducibles that behave well with respect to tensor products. A similar problem arises from types of Siegel modular forms.

There are two approaches to accommodate this problem, neither of which seems to have appeared in the literature. It is possible to sum over tensor powers of the standard representation. One would consider

$$\bigoplus_{0 \le l \in \mathbb{Z}} \mathbf{M}^{(n)}(\mathrm{std}^l) \tag{8.6}$$

By a result of Weissauer [Wei83], every irreducible representation σ for which $M^{(n)}(\sigma \boxtimes \rho)$ is not trivial for some ρ is contained in some tensor power of std^{*l*}, so that (8.6) contains all Siegel modular forms of level 1. More specifically, det^{*k*} \subseteq std^{*nk*}. In particular, the non-commutative graded ring (8.6) of Siegel modular forms contains the classical one (8.5).

The type of Siegel modular forms, if $n \ge 2$, can be expressed in terms of Weyl representations $\rho_{\mathscr{L}}$ attached to (positive definite and integral) lattices \mathscr{L} . ONe can thus appy a similar construction, to obtain a non-commutative ring that contains all Siegel modular forms of arbitrary weights and types. We arbitrary will not go into details, though.

The second approach, which seems most suitable to study the analogue of (8.1), is to sacrifice the algebra structure in favor of a grading with respect to irreducible representations of $GL_n(\mathbb{C})$ and $\Gamma^{(n)}$. The direct sum

$$\mathbf{M}^{(n)}(\bullet \boxtimes \bullet) = \bigoplus \mathbf{M}^{(n)}(\sigma \boxtimes \rho), \tag{8.7}$$

where σ and ρ run through fixed representatives of isomorphism classes of representations, is a commutative hyper-algebra.

§8.2 Hyper-algebras. A hyper-group is group whose multiplication is multi-valued [Wal37]. We take the idea of multi-valued operators to define hyper-algebras, which we could not find in the literature.

Let SubVec(*A*) be the set of all finite rank submodules of a vector space *A*.

Definition 8.1. A triple $(A, +, \cdot)$ with (A, +) a *K*-vector space and a binary operator

$$\cdot : A \times A \longrightarrow \text{SubVec}(A)$$

is called a hyper-algebra (with strong identity) if

- (i) there is a unique element $e \in A$ such that $e \cdot r = r \cdot e = \operatorname{span} r$ for all $r \in A$, and for every $a \in A$ there is a' such that $e \in aa', a'a$.
- (ii) for $k \in K \setminus \{0\}$ and $a, a' \in A$, we have aa' = (ka)a' = a(ka').

- (iii) for $a, a', a'' \in A$, we have $a(a' + a'') \subseteq aa' + aa''$.
- (iv) for $a, a', a'' \in A$, we have a(a'a'') = (aa')a'', where the product is linearly extended to SubVec(A) × SubVec(A).

Further, we say that

- (1) *A* is commutative if for $a, a' \in A$, we have aa' = a'a.
- (2) *A* is *G*-graded for a hyper-group *G* if $A = \bigoplus_{g} A_g$ and $A_g A_{g'} \subseteq \sum_{h \in gg'} A_h$.
- (3) A is a differential hyper-algebra, if there is a K-linear map

$$\partial: A \longrightarrow \operatorname{SubVec}(A)$$

satisfying the Leibniz rule $\partial(aa') \subseteq (\partial a)a' + a(\partial a')$ for all $a, a' \in A$. If *A* is graded by *G*, then we say that ∂ has grading $g \in G$, if $\partial A_{g'} \subseteq \sum_{h \in gg'} A_h$ for all $g' \in G$.

Remark 8.2. The notion of hyper-groups makes it obvious to call the structure that we have defined a hyper-algebra. But the reader should be warned that this is not standard. In fact, it has nothing in common with hyper-algebras associated with groups in, for example, [Sul78].

§8.3 The differential graded hyper-algebra of almost holomorphic Siegel modular forms. Recall that isomorphism classes of irreducible, finite-dimensional, complex representations of $GL_n(\mathbb{C})$ form a hyper-group $\operatorname{Rep}(GL_n(\mathbb{C}))$ with multiplication given by the decomposition of tensor products. Typically, this is formulated in terms of fusion algebras (see [Eho95] for one instance) or Grothendieck rings, but Definition 8.1 suggests that it is more convenient to work with hyper-groups. Similarly, irreducible, finite-dimensional representations of $\Gamma^{(n)}$ with finite index kernel, yield a hyper-group $\operatorname{Rep}(\Gamma^{(n)})$. By fixing representatives once and for all, we consider elements of $\operatorname{Rep}(\operatorname{GL}_n(\mathbb{C}))$ and $\operatorname{Rep}(\Gamma^{(n)})$ as actual representations.

Set

$$\widehat{\mathbf{M}}^{(n)} = \bigcup_{0 \le d \in \mathbb{Z}} \bigoplus_{\substack{\sigma \in \operatorname{Rep}(\operatorname{GL}_n(\mathbb{C}))\\\rho \in \operatorname{Rep}(\Gamma^{(n)})}} \mathbf{M}^{(n)} \big(\sigma^{[d]} \boxtimes \rho \big).$$
(8.8)

For $f \in M^{(n)}(\sigma^{[d]} \boxtimes \rho)$ and $g \in M^{(n)}(\sigma'^{[d']} \boxtimes \rho')$, set

$$f \otimes g = \sup_{\substack{\kappa_{\sigma}: \sigma \otimes \sigma' \to \sigma_{\rm irr} \\ \kappa_{\rho}: \rho \otimes \rho' \to \rho_{\rm irr}}} (\kappa_{\sigma} \boxtimes \kappa_{\rho}) (f \otimes g), \tag{8.9}$$

where the span runs through all homomorphism representations with irreducible σ_{irr} and ρ_{irr} . The \mathbb{C} -module $\widehat{M}^{(n)}$ carries a graded commutative hyper- \mathbb{C} -algebra structure when equipped with multiplication as in (8.9).

The raising operators $\mathbb{R}_{\text{std}^l}$ can be extended to $\widehat{\mathbb{M}}^{(n)}$. Given σ choose an embedding $\iota : \sigma \hookrightarrow \text{std}^l$ for some $0 \le l \in \mathbb{Z}$. Then

$$Rf = \sup_{\kappa: \operatorname{sym}^2 \otimes \operatorname{std}^l \to \sigma_{\operatorname{irr}}} \kappa(R_{\operatorname{std}^l} f).$$
(8.10)

When equipped with the differential R, then $\widehat{M}^{(n)}$ becomes a differential graded hyper-algebra, and R has grading sym². Similarly, we obtain an action of L on $\widehat{M}^{(n)}$.

§8.4 Exception Siegel modular forms. We finish the classification of almost holomorphic Siegel modular forms of genus 2 that was initiated by Pitale, Saha, and Schmidt in [PSS15]. The goal of this section is to prove the following theorem.

Theorem 8.3. *Given* σ *,* ρ *, and* $0 \le d$ *, we have*

$$\mathbf{M}^{(2)}\big(\sigma^{[d]} \boxtimes \rho\big) = \sum_{t=0}^{d} \big(\pi_{\mathrm{sym}^2} \mathbf{R}\big)^t \mathbf{M}^{(n)}\big((\mathrm{sym}^{\vee 2})^t \sigma \boxtimes \rho\big).$$

That is, Siegel modular forms generate $\widehat{M}^{(2)}$ as a differential hyper-algebra.

Proof. This was almost proved in [PSS15], except for the cases $\sigma = \det^3 \operatorname{sym}^l$ for some $0 \le l \in \mathbb{Z}$. As noted there, it suffices to show that dim $M^{(2)}(\operatorname{detsym}^l \boxtimes \rho) = 0$ for all l and all ρ to rule out the remaining cases.

Suppose that $f \in M^{(2)}(\text{detsym}^l \boxtimes \rho)$ is a non-zero Siegel modular form. We consider its Fourier Jacobi expansion

$$f(\tau) = \sum_{0 \le m \in \mathbb{Z}} \phi_m(\tau_1, z) e(m\tau_2).$$

Applying the operator $C^{J}_{detsym_{2}^{l}}$ and the map $\langle \cdot, \mathfrak{f}_{l} \rangle$ studied in Section 7.7, we find that

 $\langle \phi_m(\tau_1, z), \mathfrak{f}_l \rangle$

is a classical Jacobi form of weight 1 and index *m*. By results of Ibukiyama and Skoruppa it vanishes [IS07]. Applying the next Lemma 8.4 yields the result.

Lemma 8.4. Let $f \in M^{(n)}(\sigma \boxtimes \rho)$ and $v \in \sigma^{\vee}$. If $v \circ f = 0$ then $(GL_n(\mathbb{C}) v) \circ f = 0$.

Proof. This follows from invariance of f under the action of the Levi factor. For $u \in GL_n(\mathbb{Z})$, we have

$$(uv) f(\tau) = (uv) uf({}^{t}u\tau u) = v f({}^{t}u^{-1}\tau u^{-1}) = 0.$$

The uv for $u \in GL_n(\mathbb{Z})$ span $GL_n(\mathbb{C})$ $v \subseteq V(\sigma^{\vee})$, and this implies the lemma.

The classification by Pitale, Saha, and Schmidt, which appeared shortly before the present manuscript was finished, shows that there is no modular level 1 Siegel modular form of weight sym² if n = 2. We extend this to all genera by applying a variant of the argument in the proof of Theorem 8.3.

Proposition 8.5. *If* $n \ge 2$ *, then for any* ρ *, we have*

$$\mathbf{M}^{(n)}(\mathbf{sym}^{2\,[1]}\boxtimes \mathbb{1}) = \mathbf{M}^{(n)}(\mathbf{sym}^{2}\boxtimes \mathbb{1}).$$

Proof. We can and will assume that ρ is irreducible. We claim that every $f \in M^{(n)}(\text{sym}^{2[1]})$ is of the form $f(\tau) = g(\tau) + cy^{-1}$ for some constant $c \in \mathbb{C}$ and holomorphic g. Indeed, assume that $f(\tau) = g(\tau) + h(y^{-1})$ for a holomorphic function g and a linear function $h \in \text{sym}^2 \otimes \text{sym}^{\vee 2}$ without constant term. A direct computation shows that the image of f under L equals $-t_{12}h$. It is a modular form of weight $\text{sym}^{\vee 2} \text{sym}^2$, which by Lemma 8.7 is supported on the span of $\sum_{ij} \mathfrak{e}_{ij}^{\vee} \frac{1}{2}(\mathfrak{e}_{ij} + \mathfrak{e}_{ji})$. This proves the claim.

It suffices to show that there is no $f^{(n)} \in M^{(n)}(\text{sym}^{2[1]})$ of the form $f^{(n)}(\tau) = g^{(n)}(\tau) + y^{-1}$. The 0-th Fourier Jacobi coefficient of $f^{(n)}$, defined in (7.10), is a Siegel modular form of genus n - 1. Since dim $M^{(n-1)}(\text{sym}^2) = 0$, we find that it equals $f^{(n-1)}$. We are therefore reduced to the case n = 2.

Consider the case n = 2. We employ the Fourier Jacobi expansion. As a next step we show that vanishing of the 0-th Fourier Jacobi coefficient of $f^{(2)}$ contradicts its existence. Assuming that $f^{(2)}$

exists, let *m* be minimal subject to the condition that the *m*-th Fourier Jacobi coefficient, say ϕ_m of $f^{(2)}$ is non-zero. Combining (7.10) and (7.7), we see that $C_{\text{sym}^2}^{\text{J}} \phi_m$ has Fourier expansion starting at *m*:

$$\sum_{n\leq n;r}c(n,r)\exp\left(2\pi i(n\tau_1+rz)\right).$$

Its components are elements of $J_{0,m}^{[0,0]}$, $J_{1,m}^{[0,0]}$, and $J_{2,m}^{[1,0]}$. Using the same argument as in [BWR14], these correspond via the maps $\mathcal{D}_{2\nu}$ in (9) of [EZ85] to almost modular forms modular forms of weight at most 2 + 2m, whose Fourier expansions starts at m. If $m \ge 1$, they are zero by the theory of almost holomorphic elliptic modular forms. This implies that $\phi_0 \ne 0$, if $f^{(2)}$ exists.

It suffices to show that the 0-th Fourier Jacobi coefficient of $f^{(2)}$ would have to vanish, contradicting its existence. The Fourier Jacobi expansions of $f^{(2)}$ starts like

$$f^{(2)}(\tau) = \begin{pmatrix} \phi_{0,3}(\tau_1, z) & \phi_{0,2}(\tau_1, z) \\ \phi_{0,2}(\tau_1, z) & \phi_{0,1}(\tau_1, z) \end{pmatrix} + y^{-1} + \begin{pmatrix} \phi_{1,3}(\tau_1, z) & \phi_{1,2}(\tau_1, z) \\ \phi_{1,2}(\tau_1, z) & \phi_{1,1}(\tau_1, z) \end{pmatrix} e(\tau_2) + O(e(2\tau_2)),$$

where $\tau = \begin{pmatrix} \tau_1 & z \\ z & \tau_2 \end{pmatrix}$ as in Section 7. The covariant operator $C_{sym^2}^J$ in (7.7) yields Jacobi forms in

$$J_{0,0} = \{0\}, J_{0,1} = \{0\}, J_{0,2}^{[1,0]} = \operatorname{span}(E_2), J_{1,0} = \{0\}, J_{1,1} = \{0\}, J_{1,2}^{[1,0]} = \{0\}.$$

This implies that

$$\phi_{0,1} = c, \quad \phi_{0,2} = 0, \quad \phi_{0,3}(\tau_1, z) + y_1^{-1} = \frac{-\pi}{3} E_2(\tau_1), \quad \phi_{1,1} = 0, \quad \phi_{1,2} = 0, \quad \phi_{1,3} = 0$$

for some constant $c \in \mathbb{C}$. The symmetry relations $c(f; t, y) = {}^{t}uc(f; {}^{t}utu, y)u$ for $u \in GL_2(\mathbb{Z})$, when applied to $t = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}$ and $u = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ contradicts the given equality for $\phi_{0,3}$. This establishes the statement.

Remark 8.6. In the case $n \ge 4$, one could prove Proposition 8.5 by combining [Wei87] and [Wei86].

Lemma 8.7. Suppose that σ and ρ are irreducible. If the $\mathfrak{k}_{\mathbb{C}}$ -eigenvalue of σ is negative, or if it is zero and $\sigma \neq 1$ or $\rho \neq 1$, then dim $M^{(n)}(\sigma \boxtimes \rho) = 0$.

Proof. This is a consequence of Theorem 2 in [Wei83]. Indeed, for negative $\mathfrak{k}_{\mathbb{C}}$ -eigenvalue, we obtain the statement directly. If it vanishes, than the corank of σ in the sense defined in [Wei83] equals n, so that σ is a power of det, implying $\sigma = 1$. Then $\rho = 1$ is a consequence of the fact ρ is irreducibe, and hence its invariant subspaces are all trivial.

9 A meromorphic replacement for E_2

Throughout this section, we focus on the case n = 2. Physical considerations naturally gave rise to the logarithmic derivative $\partial_{\tau} \log(\phi_{10}^{(2)})$ of $\phi_{10}^{(2)} = 4\chi_{10}$ for Igusa's $\chi_{10} \in M^{(2)}(\det^{10})$, defined on page 195 of [Igu62]. The goal of this section is to describe $\partial_{\tau} \log(\chi_{10})$ as a meromorphic replacement of the constant part of the weight 2 Eisenstein series E_2 in the genus 1 case. In particular, we will find a ring-like structure equipped with the (vector valued) differential ∂_{τ} acting on it.

§9.1 An almost meromorphic Siegel modular form. In (4.13), we find the following definition:

$$S(\tau) = S^{(2)}(\tau) = \frac{1}{10}\partial_{\tau}\log(\phi_{10}^{(2)}) - \frac{i}{2}y^{-1},$$
(9.1)

where ∂_{τ} is the vector valued derivative defined in Section 6.2. We can conveniently express S as

$$S(\tau) = \frac{1}{10}\phi_{10}^{(2)-1} \operatorname{R}_{\det^{10}}\phi_{10}^{(2)} = \frac{1}{10}\phi_{10}^{(2)-1} \left(\partial_{\tau} - 5iy^{-1}\right)\phi_{10}^{(2)},\tag{9.2}$$

from which modularity of S becomes immediately clear.

In analogy with Definition 6.13, we define almost meromorphic functions for complex representations σ of $GL_n(\mathbb{C})$, subsuming *S*.

Definition 9.1. Let $D \subset \mathbb{H}^{(n)}$ be a smooth, locally closed submanifold of codimension 1. A smooth function $f : \mathbb{H}^{(n)} \setminus D \to V(\sigma)$ is almost meromorphic of depth d if $L^{d+1} f = 0$.

It is immediately clear that Theorem 6.17 generalizes. That is, almost meromorphic functions are polynomials in the entries of y^{-1} whose coefficients are meromorphic functions. The constant term of this polynomial is called its constant part.

Let us summarize immediate consequences of Equation (9.2).

Proposition 9.2. The function *S* defined in (9.1) is an almost meromorphic function, whose singularities lie on the submanifolds z = 0 and its $\Gamma^{(2)}$ translates. For any $\gamma \in \Gamma^{(2)}$ we have $S|_{sym^2} \gamma = S$.

The Ramanujan relation (8.3) is generalized by the next proposition. Before stating it, recall the averaged transposition $t = t_{(12)(34)}$ defined in (6.4). When applying it to sym²sym², as below, it has the form

$$\mathbf{t}(\mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{11}) = \mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{11}, \mathbf{t}(\mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{12}) = \frac{1}{2}\mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{12} + \frac{1}{2}\mathbf{\hat{e}}_{12}\mathbf{\hat{e}}_{11}, \mathbf{t}(\mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{22}) = \frac{1}{4}\mathbf{\tilde{e}}_{12}\mathbf{\tilde{e}}_{12},$$
$$\mathbf{t}(\mathbf{\hat{e}}_{12}\mathbf{\tilde{e}}_{12}) = \mathbf{\hat{e}}_{11}\mathbf{\hat{e}}_{22} + \mathbf{\hat{e}}_{22}\mathbf{\hat{e}}_{11} + \frac{1}{2}\mathbf{\tilde{e}}_{12}\mathbf{\tilde{e}}_{12}, \quad \text{where} \quad \mathbf{\tilde{e}}_{ij} = \frac{\mathbf{\hat{e}}_{ij} + \mathbf{\hat{e}}_{ji}}{1 + \delta_{ij}}.$$

Proposition 9.3. We have

$$R_{svm^2} S = t(S \otimes S) + f_{RS}(\tau), \qquad (9.3)$$

where $f_{\rm RS}$ is a meromorphic Siegel modular forms of weight sym² \otimes sym², which has a pole of order 2 at z = 0; that is, $\phi_{10}^{(2)} f_{\rm RS}$ is holomorphic.

Proof. We prove this by first comparing non-constant parts and then checking Fourier coefficients. To summarize, we have

$$100 S \otimes S = \phi_{10}^{(2)-2} \partial_{\tau} \phi_{10}^{(2)} \otimes \partial_{\tau} \phi_{10}^{(2)} - 5i\phi_{10}^{(2)-1} \partial_{\tau} \phi_{10}^{(2)} \otimes y^{-1} - 5i\phi_{10}^{(2)-1} y^{-1} \otimes \partial_{\tau} \phi_{10}^{(2)} - 25 y^{-1} \otimes y^{-1},$$

$$10 R_{\text{sym}^{2}} S = -\phi_{10}^{(2)-2} \left(\partial_{\tau} \phi_{10}^{(2)} \otimes \partial_{\tau} \phi_{10}^{(2)}\right) + \phi_{10}^{(2)-1} \left(\partial_{\tau} \otimes \partial_{\tau}\right) \phi_{10}^{(2)} - 5i\frac{-i}{2} \partial_{y} y^{-1} - i\phi_{10}^{(2)-1} t_{(12)(34)} \left(y^{-1} \otimes \partial_{\tau} \phi_{10}^{(2)}\right) - 5 t_{(12)(34)} \left(y^{-1} \otimes y^{-1}\right).$$

Set $\Delta = \sum_{ij} \mathfrak{e}_{ij} \frac{1}{2} (\mathfrak{e}_{ij} + \mathfrak{e}_{ji})$. We have $\partial_y y^{-1} = -(y^{-1} \otimes \mathbb{I}_2) \Delta(y^{-1} \otimes \mathbb{I}_2)$, and a straightforward computation reveals that $\mathfrak{t}_{(12)(34)}(y^{-1} \otimes y^{-1}) = (y^{-1} \otimes \mathbb{I}_2) \Delta(y^{-1} \otimes \mathbb{I}_2)$. It is obvious from the definition of t that $\mathfrak{t}_{(12)(34)}(y^{-1} \otimes \partial_\tau \phi_{10}^{(2)}) = \mathfrak{t}_{(12)(34)}(\partial_\tau \phi_{10}^{(2)} \otimes y^{-1})$. Taking these equalities, we find that $\operatorname{R} S - \mathfrak{t}(S \otimes S)$ is meromorphic.

To finish the proof consider the holomorphic modular forms

$$100\phi_{10}^{(2)\,2} \left(\mathsf{R}_{\text{sym}^2} \, S - \mathsf{t}(S \otimes S) \right) = -10 \left(\partial_\tau \phi_{10}^{(2)} \otimes \partial_\tau \phi_{10}^{(2)} \right) + 10 \, \phi_{10}^{(2)} \left(\partial_\tau \otimes \partial_\tau \right) \phi_{10}^{(2)} - \mathsf{t}_{(12)(34)} \left(\partial_\tau \phi_{10}^{(2)} \otimes \partial_\tau \phi_{10}^{(2)} \right). \tag{9.4}$$

We compute its expansion around z = 0. Let $\Delta_{12} = \Delta(\tau_1)\Delta(\tau_2)$ be the product of the modular discriminant depending on τ_1 and τ_2 . Then $\frac{100}{(2\pi i)^4}\phi_{10}^{(2)2}(R_{sym^2}S - t(S \otimes S))$ has expansion

$$-10(\tilde{\mathfrak{e}}_{12}z\Delta_{12}+\mathcal{O}(z^3))(\tilde{\mathfrak{e}}_{12}z\Delta_{12}+\mathcal{O}(z^3))-t((\tilde{\mathfrak{e}}_{12}z\Delta_{12}+\mathcal{O}(z^3))(\tilde{\mathfrak{e}}_{12}z\Delta_{12}+\mathcal{O}(z^3)))$$

+10($z^2\Delta_{12}+\mathcal{O}(z^4)$)($\mathfrak{e}_{11}\tilde{\mathfrak{e}}_{12}z\partial_{\tau_1}\Delta_{12}+\mathfrak{e}_{22}\tilde{\mathfrak{e}}_{12}z\partial_{\tau_2}\Delta_{12}+\tilde{\mathfrak{e}}_{12}\mathfrak{e}_{11}z\partial_{\tau_1}\Delta_{12}$
+ $\tilde{\mathfrak{e}}_{12}\mathfrak{e}_{22}z\partial_{\tau_2}\Delta_{12}+\tilde{\mathfrak{e}}_{12}\tilde{\mathfrak{e}}_{12}\frac{1}{2}\Delta_{12}+\mathcal{O}(z^3)$).

The leading term is therefore

$$z^{2}\Delta_{12}^{2}\left(\frac{-11}{2}\tilde{\mathfrak{e}}_{12}\tilde{\mathfrak{e}}_{12}-\frac{1}{4}\mathfrak{e}_{11}\mathfrak{e}_{22}-\frac{1}{4}\mathfrak{e}_{22}\mathfrak{e}_{11}\right)+10z^{3}\Delta_{12}\left(\partial_{\tau_{1}}\Delta_{12}(\mathfrak{e}_{11}\tilde{\mathfrak{e}}_{12}+\tilde{\mathfrak{e}}_{12}\mathfrak{e}_{11})+\partial_{\tau_{2}}\Delta_{12}(\mathfrak{e}_{22}\tilde{\mathfrak{e}}_{12}+\tilde{\mathfrak{e}}_{12}\mathfrak{e}_{22}),$$

which, in particular, implies that $\phi_{10}^{(2)} f$ is holomorphic.

§9.2 Fourier coefficients of f_{RS} . Our goal is to compute Fourier coefficients of f_{RS} defined in (9.3). We use an Ansatz to find an expression for $\phi_{10}^{(2)} f_{RS}$ in terms of usual generators. This expression is displayed in Equation (9.5). We obtain it by first decomposing det¹⁰ sym² sym² explicitly, finding a basis for the corresponding spaces of Siegel modular forms, and then comparing coefficients with (9.4). Multiplication with $\phi_{10}^{(2)-1}$ then yields Fourier coefficients of $f_{\rm RS}$ (in a prescribed Weyl chamber).

In this Section we will frequently refer to Sage [Ste+14] scripts. They can all be found on the last named author's homepage. Note that several of them make us of the implementation of Jacobi forms [Rau12] that is currently only available at Sage-Trac, i.e., as a branch u/mraum/ticket/16448 at Sage-Git.

We will use polynomials p in two variables v_1 and v_2 with the action $(gp)(v_1v_2) = p((v_1v_2)g)$ as a basis for symmetric power representations of $SL_2(\mathbb{C})$. An isomorphism to our previous basis \mathfrak{e}_{11} , $\mathfrak{e}_{12} + \mathfrak{e}_{21}$, \mathfrak{e}_{22} of sym² is given by

$$\mathfrak{e}_{11} \mapsto v_1^2, \quad \mathfrak{e}_{12} + \mathfrak{e}_{21} \mapsto 2v_1v_2, \quad \mathfrak{e}_{22} \mapsto v_2^2.$$

A decomposition of sym² \otimes sym². By the Clebsch-Gordan rules we know that sym² \otimes sym² \cong sym⁰ \oplus sym² \oplus sym⁴ as a representation of SL₂(\mathbb{C}). Let us find an explicit basis in terms of v_1 and v_2 . To simplify notation, we will denote by u_1 , u_2 and v_1 , v_2 the variables of the polynomials in the first and second tensor component. Those in the image will be denoted by w_1 and w_2 .

The trivial representation, one checks, is spanned by

$$u_1^2 v_2^2 - 2u_1 u_2 v_1 v_2 + u_2^2 v_1^2$$
.

Bases for sym² and sym⁴, respectively, are given by

$$w_1^2 = u_1^2 v_1 v_2 - u_1 u_2 v_1^2$$
, $w_1 w_2 = \frac{1}{2} (u_1^2 v_2^2 - u_2^2 v_1^2)$, $w_2^2 = u_1 u_2 v_2^2 - u_2^2 v_1 v_2$,

and

$$w_1^4 = u_1^2 v_1^2, \quad w_1^3 w_2 = \frac{1}{2} (u_1^2 v_1 v_2 + u_1 u_2 v_1^2), \quad w_1^2 w_2^2 = \frac{1}{6} (u_2^2 v_1^2 + 4 u_1 u_2 v_1 v_2 + u_1^2 v_2^2), w_1 w_2^3 = \frac{1}{2} (u_2^2 v_1 v_2 + u_1 u_2 v_2^2), \quad w_2^4 = u_2^2 v_2^2.$$

Table of basic Fourier coefficients. In Table 13, we give the first few Fourier coefficients of classical scalar valued Siegel modular forms that have been taken from [LMF15]; Or equivalently computed by the Sage script basic_fourier_expansions.sage. On the left of Table 13, we give names of Siegel modular forms, which are all explained in below. On the right, we give their Fourier coefficients according to the scheme dictated by the first 9 lines.

The space M⁽²⁾(det¹²) has dimension 3. A natural basis consists of the Siegel Eisenstein series $E_{12}^{(2)}$, the Klingen Eisenstein series $E_{\Delta}^{(2)}$ attached to the unique weight 12 cusp form Δ in genus 1, and the Maass lift $\phi_{12}^{(2)}$ of the unique Jacobi cusp form ϕ_{12} of weight 12 and index 1. The space $M^{(2)}(det^{10}sym^4)$ is one dimensional. A generator was found by Ibukiyama in [Ibu12] using $E_4^{(2)}$ and $E_6^{(2)}$ as input data. The Ibukiyama-Rankin-Cohen bracket is

$$\{f, g\}_{\operatorname{sym}^4} \in \operatorname{M}^{(2)}(\operatorname{det}^{k+k'}\operatorname{sym}^4) \text{ for } f \in \operatorname{M}^{(2)}(\operatorname{det}^k) \text{ and } g \in \operatorname{M}^{(2)}(\operatorname{det}^{k'}).$$

With respect to our choice of basis w_1^4, \ldots, w_2^4 , it equals

$$\begin{split} & \frac{1}{2}k'(k'+1)g\left(\partial_{11}^2f \,w_1^4 + 2\partial_{11}\partial_{12}f \,w_1^3 \,w_2 + \left(\partial_{12}^2f + 2\partial_{11}\partial_{22}f\right)w_1^2 w_2^2 + 2\partial_{12}\partial_{22}f \,w_1 w_2^3 + \partial_{22}^2f \,w_2^4\right) \\ & -(k+1)(k'+1)\left((\partial_{11}f)(\partial_{11}g) \,w_1^4 + \left((\partial_{11}f)(\partial_{12}g) + (\partial_{12}f)(\partial_{11}g)\right)w_1^3 w_2 \\ & + \left((\partial_{11}f)(\partial_{22}g) + (\partial_{12}f)(\partial_{12}g) + (\partial_{22}f)(\partial_{11}g)\right)w_1^2 w_2^2 \\ & + \left((\partial_{22}f)(\partial_{12}g) + (\partial_{12}f)(\partial_{22}g)\right)w_1 w_2^3 + (\partial_{22}f)(\partial_{22}g) \,w_1^4\right) \\ & + \frac{1}{2}k(k+1)f\left(\partial_{11}^22 \,w_1^4 + 2\partial_{11}\partial_{12}g \,w_1^3 \,w_2 + \left(\partial_{12}^2g + 2\partial_{11}\partial_{22}g\right)w_1^2 w_2^2 + 2\partial_{12}\partial_{22}g \,w_1 w_2^3 + \partial_{22}^2g \,w_2^4\right). \end{split}$$

We do not reproduce the Fourier coefficients of $\{E_4^{(2)}, E_6^{(2)}\}_{sym^4}$, but refer the reader to the Sage script mentioned above.

An explicit expression for $\phi_{10}^{(2)} f_{RS}$. Using the Sage script fRS_phi2_10_fourier_expansion.sage, we compute an explicit representation of $\phi_{10}^{(2)} f_{RS}$. We obtain

$$\phi_{10}^{(2)} f_{\rm RS} = \frac{-1}{72576000} \left\{ E_4^{(2)}, E_6^{(2)} \right\}_{\rm sym^4} + \frac{1}{400} \phi_{12}^{(2)}, \tag{9.5}$$

where the embedding of det^{10} sym⁴ and det^{12} into det^{10} sym² sym² is as above. From this expression, we obtain the Fourier coefficients in Table 14.

Fourier expansion of $\phi_{10}^{(2)-1}$. The Fourier expansion of $\phi_{10}^{(2)-1}$ is not unique, but depends on a Weyl chamber as explained in [Bor98]. The zero locus of $\phi_{10}^{(2)}$ is $\Gamma^{(2)} \{\tau : z = 0\}$, and a Weyl chamber is a connected component of its complement in $\mathbb{H}^{(2)}$. We consider the one that contains $i \begin{pmatrix} 1 & t^2 \\ t^2 & t \end{pmatrix}$ for all sufficiently small, positive *t*. In this Weyl chamber, $\phi_{10}^{(2)-1}$ has Fourier expansion

$$q_1\zeta q_2 \prod_{(n,r,m)>0} \left(1-q_1^n\zeta^r q_2^m\right)^{c(nm-r^2)},$$

where $q_1 = \exp(2\pi i \tau_1)$, $\zeta = \exp(2\pi i z)$, $q_2 = \exp(2\pi i \tau_2)$, and

$$(n, r, m) > 0 \Leftrightarrow (n > 0) \lor (n = 0 \land m > 0) \lor (n = m = 0 \land r > 0).$$

The c(n) are the Fourier coefficients of the weak Jacobi form of weight 0 and index 1 that is uniquely determined by c(-1) = 2. Its Fourier expansion is computed to the necessary extend in the Sage script phi2_10_inv_fourier_expansion.sage.

To compute the Fourier expansion of $\phi_{10}^{(2)-1}$ correctly the bound on (n, r, m) that were found in [GKR13] have to be taken into account. This, eventually yields the Fourier coefficients, some of which are displayed in Table 15. The reader is also referred to the file phi2_10_inv_fe.sobj. This file is computed by phi2_10_inv_fourier_expansion.sage and contains a Sage-readable dictionary of Fourier coefficients

chi10inv_fe.sobj:
$$(n, r, m) \mapsto c(\phi_{10}^{(2)-1}, \begin{pmatrix} n & r/2 \\ r/2 & m \end{pmatrix})$$

for many more (*n*, *r*, *m*).

Fourier coefficients of f_{RS} . Computing Fourier coefficients of f_{RS} in the Weyl chamber that was previously defined, we simply have to multiply $\phi_{10}^{(2)} f_{RS}$ by $\phi_{10}^{(2)-1}$. This is performed in the Sage script fRS_fourier_expansion.sage. The resulting Fourier coefficients are displayed in Table 16. The previously mentioned Sage script will produce a file fRS_fe.sobj, which contains a dictionary of Fourier coefficients

fRS_fe.sobj:
$$(n, r, m) \mapsto c(f_{\text{RS}}, \binom{n r/2}{r/2 m})$$
.

References

[ABK08]	M. Aganagic, V. Bouchard, and A. Klemm, "Topological strings and (almost) modular forms", <i>Comm. Math. Phys.</i> , vol. 277, no. 3, 2008.
[AKMV04]	M. Aganagic, A. Klemm, M. Mariño, and C. Vafa, "Matrix model as a mirror of Chern-Simons theory", <i>J. High Energy Phys.</i> , no. 2, 2004, ISSN: 1126-6708. DOI: 10.1088/1126-6708/2004/02/010.
[AKMV05]	—, "The topological vertex", <i>Comm. Math. Phys.</i> , vol. 254, no. 2, 2005, ISSN: 0010-3616. DOI: 10.1007/s00220-004-1162-z.
[Ali09]	M. Alim, "Mirror symmetry, toric branes and topological string amplitudes as polynomials", 2009.

[BCOV94]	M. Bershadsky, S. Cecotti, H. Ooguri, and C. Vafa, "Kodaira-Spencer theory of gravity and exact results for quantum string amplitudes", <i>Commun.Math.Phys.</i> , vol. 165, 1994. DOI: 10.1007/BF02099774. arXiv: hep-th/9309140 [hep-th].
[BGHZ08]	J. H. Bruinier, G. van der Geer, G. Harder, and D. B. Zagier, <i>The 1-2-3 of modular forms</i> , K. Ranestad, Ed., ser. Universitext. Springer-Verlag, Berlin, 2008, Lectures from the Summer School on Modular Forms and their Applications held in Nordfjordeid, June 2004.
[BKMP09]	V. Bouchard, A. Klemm, M. Mariño, and S. Pasquetti, "Remodeling the B-model", <i>Comm. Math. Phys.</i> , vol. 287, no. 1, 2009, ISSN: 0010-3616. DOI: 10.1007/s00220-008-0620-4.
[BO00]	S. Bloch and A. Okounkov, "The character of the infinite wedge representation", <i>Adv. Math.</i> , vol. 149, no. 1, 2000.
[Bor98]	R. E. Borcherds, "Automorphic forms with singularities on grassmannians", <i>Invent. Math.</i> , vol. 132, no. 3, 1998.
[BRR12]	K. Bringmann, M. Raum, and O. K. Richter, <i>Harmonic Maass-Jacobi forms with singularities and a theta-like decomposition</i> , arXiv:1207.5600, 2012.
[BS98]	R. Berndt and R. Schmidt, <i>Elements of the representation theory of the Jacobi group</i> , ser. Progress in Mathematics. Basel: Birkhäuser Verlag, 1998, vol. 163.
[BWR14]	J. H. Bruinier and M. Westerholt-Raum, <i>Kudla's modularity conjecture and formal fourier-jacobi series</i> , arXiv:1409.4996, 2014.
[CBKL14]	N. Cabo Bizet, A. Klemm, and D. V. Lopes, "Landscaping with fluxes and the E8 Yukawa Point in F-theory", 2014. arXiv: 1404.7645 [hep-th].
[CKK14]	J. Choi, S. Katz, and A. Klemm, "The refined BPS index from stable pair invariants", <i>Comm. Math. Phys.</i> , vol. 328, no. 3, 2014.
[CKYZ99]	T. Chiang, A. Klemm, ST. Yau, and E. Zaslow, "Local mirror symmetry: Calculations and interpretations", <i>Adv.Theor.Math.Phys.</i> , vol. 3, 1999. arXiv: hep-th/9903053 [hep-th].
[CR10]	C. Conley and M. Raum, <i>Harmonic Maaß–Jacobi forms of degree</i> 1 <i>with higher rank indices</i> , arXiv:1012.2897, 2010.
[Dij95]	R. Dijkgraaf, "Mirror symmetry and elliptic curves", <i>The moduli space of curves (Texel Island, 1994)</i> , ser. Progr. Math. Vol. 129, Birkhäuser Boston, Boston, MA, 1995.
[Eho95]	W. Eholzer, "On the classification of modular fusion algebras", <i>Comm. Math. Phys.</i> , vol. 172, no. 3, 1995.
[EZ85]	M. Eichler and D. B. Zagier, <i>The theory of Jacobi forms</i> , ser. Progress in Mathematics. Boston, MA: Birkhäuser Boston Inc., 1985, vol. 55.
[Fre83]	E. Freitag, <i>Siegelsche Modulfunktionen</i> , ser. Grundlehren der Mathematischen Wissenschaften. Berlin: Springer-Verlag, 1983, vol. 254.
[GHM14]	A. Grassi, Y. Hatsuda, and M. Marino, "Topological strings from quantum mechanics", 2014. arXiv: 1410.3382 [hep-th].
[GKMW07]	T. W. Grimm, A. Klemm, M. Mariño, and M. Weiss, "Direct integration of the topological string", <i>J. High Energy Phys.</i> , no. 8, 2007, ISSN: 1126-6708. DOI: 10.1088/1126-6708/2007/08/058.
[GKR13]	D. Gehre, J. Kreuzer, and M. Raum, "Computing Borcherds products", <i>LMS J. Comput. Math.</i> , vol. 16, 2013.
[GW09]	R. Goodman and N. R. Wallach, <i>Symmetry, representations, and invariants,</i> ser. Graduate Texts in Mathematics. Dordrecht: Springer, 2009, vol. 255.
[Hel59]	S. Helgason, "Differential operators on homogenous spaces", Acta Math., vol. 102, 1959.
[Hel92]	—, "Some results on invariant differential operators on symmetric spaces", <i>Amer. J. Math.</i> , vol. 114, no. 4, 1992.
[HK07]	Mx. Huang and A. Klemm, "Holomorphic Anomaly in Gauge Theories and Matrix Models", <i>JHEP</i> , vol. 0709, 2007. DOI: 10.1088/1126-6708/2007/09/054. arXiv: hep-th/0605195 [hep-th].
[HK12]	—, "Direct integration for general Ω backgrounds", <i>Adv. Theor. Math. Phys.</i> , vol. 16, no. 3, 2012.
[HKK15]	Mx. Huang, S. Katz, and A. Klemm, "Topological String on elliptic CY 3-folds and the ring of Jacobi forms", 2015. arXiv: 1501.04891 [hep-th].
[HKP13]	MX. Huang, A. Klemm, and M. Poretschkin, "Refined stable pair invariants for E-, M- and [<i>p</i> , <i>q</i>]-strings", <i>JHEP</i> , vol. 1311, 2013. DOI: 10.1007/JHEP11(2013)112. arXiv: 1308.0619 [hep-th].
[HKPK13]	Mx. Huang, AK. Kashani-Poor, and A. Klemm, "The Ω deformed B-model for rigid $N = 2$ theories", <i>Ann. Henri Poincaré</i> , vol. 14, no. 3, 2013.
[HKR08]	B. Haghighat, A. Klemm, and M. Rauch, "Integrability of the holomorphic anomaly equations", <i>J. High Energy Phys.</i> , no. 10, 2008.

[Hos06]	S. Hosono, "Central charges, symplectic forms, and hypergeometric series in local mirror symmetry", <i>Mirror symmetry. V</i> , ser. AMS/IP Stud. Adv. Math. Vol. 38, Amer. Math. Soc., Providence, RI, 2006.
[Ibu12]	T. Ibukiyama, "Lifting conjectures from vector valued Siegel modular forms of degree two", <i>Comment. Math. Univ. St. Pauli</i> , vol. 61, no. 1, 2012.
[Igu62]	J. Igusa, "On Siegel modular forms of genus two", Amer. J. Math., vol. 84, 1962.
[Igu67]	Ji. Igusa, "Modular forms and projective invariants", Amer. J. Math., vol. 89, 1967.
[IK11]	T. Ibukiyama and R. Kyomura, "A generalization of vector valued Jacobi forms", <i>Osaka J. Math.</i> , vol. 48, no. 3, 2011.
[IS07]	T. Ibukiyama and NP. Skoruppa, "A vanishing theorem for Siegel modular forms of weight one", <i>Abh. Math. Sem. Univ. Hamburg</i> , vol. 77, 2007.
[KM15]	R. Kashaev and M. Marino, "Operators from mirror curves and the quantum dilogarithm", 2015. arXiv: 1501.01014 [hep-th].
[KM96]	A. Klemm and P. Mayr, "Strong coupling singularities and nonAbelian gauge symmetries in N=2 string theory", <i>Nucl.Phys.</i> , vol. B469, 1996. DOI: 10.1016/0550-3213(96)00108-3. arXiv: hep-th/9601014 [hep-th].
[KMP96]	S. H. Katz, D. R. Morrison, and M. R. Plesser, "Enhanced gauge symmetry in type II string theory", <i>Nucl.Phys.</i> , vol. B477, 1996. DOI: 10.1016/0550-3213(96)00331-8. arXiv: hep-th/9601108 [hep-th].
[KMW12]	A. Klemm, J. Manschot, and T. Wotschke, "Quantum geometry of elliptic Calabi-Yau manifolds", <i>Commun. Number Theory Phys.</i> , vol. 6, no. 4, 2012, ISSN: 1931-4523. DOI: 10.4310/CNTP.2012.v6.n4.a5.
[KW11]	D. Krefl and J. Walcher, "Extended Holomorphic Anomaly in Gauge Theory", <i>Lett.Math.Phys.</i> , vol. 95, 2011. DOI: 10.1007/s11005-010-0432-2. arXiv: 1007.0263 [hep-th].
[KZ01]	A. Klemm and E. Zaslow, "Local mirror symmetry at higher genus", <i>Winter School on Mirror Symmetry, Vector Bundles and Lagrangian Submanifolds (Cambridge, MA, 1999),</i> ser. AMS/IP Stud. Adv. Math. Vol. 23, Amer. Math. Soc., Providence, RI, 2001.
[KZ95]	M. Kaneko and D. B. Zagier, "A generalized Jacobi theta function and quasimodular forms", <i>The moduli space of curves (Texel Island, 1994)</i> , ser. Progr. Math. Vol. 129, Birkhäuser Boston, Boston, MA, 1995.
[KZ98]	——, "Supersingular <i>j</i> -invariants, hypergeometric series, and Atkin's orthogonal polynomials", <i>Computational perspectives on number theory (Chicago, IL, 1995)</i> , ser. AMS/IP Stud. Adv. Math. Vol. 7, Amer. Math. Soc., Providence, RI, 1998.
[Maa71]	H. Maass, <i>Siegel's modular forms and Dirichlet series</i> , ser. Lecture Notes in Mathematics, Vol. 216. Berlin: Springer-Verlag, 1971.
[Maa79]	——, "Über eine Spezialschar von Modulformen zweiten Grades", <i>Invent. Math.</i> , vol. 52, no. 1, 1979.
[Mar08]	M. Mariño, "Open string amplitudes and large order behavior in topological string theory", J. High Energy Phys., no. 3, 2008, ISSN: 1126-6708. DOI: 10.1088/1126-6708/2008/03/060.
[MP12]	M. Marino and P. Putrov, "Abjm theory as a fermi gas", <i>J.Stat.Mech.</i> , vol. 1203, 2012. DOI: 10.1088/1742-5468/2012/03/P03001. arXiv: 1110.4066 [hep-th].
[MW97]	G. W. Moore and E. Witten, "Integration over the u plane in Donaldson theory", <i>Adv.Theor.Math.Phys.</i> , vol. 1, 1997. arXiv: hep-th/9709193 [hep-th].
[OFS02]	X. de la Ossa, B. Florea, and H. Skarke, "D-branes on noncompact Calabi-Yau manifolds: <i>K</i> -theory and monodromy", <i>Nuclear Phys. B</i> , vol. 644, no. 1-2, 2002.
[PSS15]	A. Pitale, A. Saha, and R. Schmidt, <i>Lowest weight modules of</i> $Sp_4(\mathbb{R})$ <i>and nearly holomorphic Siegel modular forms</i> , arXiv:1501.00524, 2015.
[Rau12]	M. Raum, <i>Computing Genus 1 Jacobi Forms</i> , Accepted for publication in Math. Comp. arXiv:1212.1834, 2012.
[Rud94]	R. E. Rudd, "The String partition function for QCD on the torus", 1994. arXiv: hep-th/9407176 [hep-th].
[Sch76]	W. Schmid, " L^2 -cohomology and the discrete series", Ann. of Math. (2), vol. 103, no. 2, 1976.
[Shi73]	G. Shimura, "On modular forms of half integral weight", Ann. of Math. (2), vol. 97, 1973.
[Shi86]	——, "On a class of nearly holomorphic automorphic forms", <i>Ann. of Math. (2)</i> , vol. 123, no. 2, 1986.
[Sie51]	C. L. Siegel, <i>Die Modulgruppe in einer einfachen involutorischen Algebra</i> , Festschr. Akad. Wiss. Göttingen, MathPhys. Kl, 1951.
[Ste+14]	W. Stein <i>et al.</i> , <i>Sage Mathematics Software (Version 6.4)</i> , http://www.sagemath.org, The Sage Development Team, 2014.

(A.1)

(A.2)

(A.3) (A.4)

(A.5)

(A.6)

(A.7)

(A.8)

(A.9)

(A.10)

(A.11)

(A.12)

(A.13)

[Sul78]	J. B. Sullivan, "Representations of the hyperalgebra of an algebraic group", <i>Amer. J. Math.</i> , vol. 100, no. 3, 1978.
[SW94]	N. Seiberg and E. Witten, "Electric - magnetic duality, monopole condensation, and confinement in N=2 supersymmetric Yang-Mills theory", <i>Nucl.Phys.</i> , vol. B426, 1994. DOI: 10.1016/0550-3213(94)90124-4. arXiv: hep-th/9407087 [hep-th].
[Wal37]	H. S. Wall, "Hypergroups", Amer. J. Math., vol. 59, no. 1, 1937.
[Wei83]	R. Weissauer, "Vektorwertige Siegelsche Modulformen kleinen Gewichtes", <i>J. Reine Angew. Math.</i> , vol. 343, 1983.
[Wei86]	——, "Über die Konstruktion von Spitzenformen mit Hilfe von Thetareihen", <i>Abh. Math. Sem. Univ. Hamburg</i> , vol. 56, 1986.
[Wei87]	——, "Divisors of the Siegel modular variety", <i>Number theory (New York, 1984–1985)</i> , ser. Lecture Notes in Math. Vol. 1240, Springer, Berlin, 1987.
[Wit95]	E. Witten, "Chern-Simons gauge theory as a string theory", <i>The Floer memorial volume</i> , ser. Progr. Math. Vol. 133, Birkhäuser, Basel, 1995.
[Zag94]	D. Zagier, "Modular forms and differential operators", <i>Proc. Indian Acad. Sci. Math. Sci.</i> , vol. 104, no. 1, 1994, K. G. Ramanathan memorial issue.
[Zie89]	C. D. Ziegler, "Jacobi forms of higher degree", Abh. Math. Sem. Univ. Hamburg, vol. 59, 1989.
[LMF15]	T. LMFDB Collaboration, <i>The L-functions and Modular Forms Database</i> , http://www.lmfdb.org, 2015.

ŀ

A The Ambiguities for $\mathbb{C}^3/\mathbb{Z}_5$
$A_1 = \frac{1}{30z_2\Delta} \Big(500z_1z_2^2 + 16z_2^2 - 450z_1z_2 - 16z_2 + 81z_1 + 3 \Big)$
$A_2 = \frac{1}{30z_1\Delta} \Big(500z_1z_2^2 + 32z_2^2 - 225z_1z_2 - 16z_2 + 27z_1 + 2 \Big)$
$f_1^{11} = -\frac{1}{400\Delta} \left(-100000z_1^2 z_2^5 + 150000z_1^2 z_2^4 - 16800z_1 z_2^4 - 576z_2^4 - 1000z_1^2 z_2^3 + 30160z_1 z_2^3 + 1008z_2^3 - 3600z_1^2 z_2^2 - 11190z_1 z_2^2 - 396z_2^2 + 1215z_1 z_2 + 45z_2 \right) = -\frac{1}{400\Delta} \left(-$
$f_1^{12} = -\frac{1}{400\Delta} \Big(50000z_1^3 z_2^4 + 25000z_1^3 z_2^3 + 6900z_1^2 z_2^3 + 288z_1 z_2^3 - 12000z_1^3 z_2^2 + 3695z_1^2 z_2^2 + 96z_1 z_2^2 + 10800z_1^3 z_2 - 2555z_1^2 z_2 - 102z_1 z_2 + 405z_1^2 + 15z_1 \Big) \Big) \Big(-100z_1^3 z_1^2 + 100z_1^3 z_2^2 + 100z_1^3 z_2^2 + 100z_1^3 z_2^2 + 100z_1^3 z_2^2 + 10z_1^3 z_$
$f_1^{22} = -\frac{1}{400z_2\Delta} \Big(-25000z_2^4z_1^4 + 81000z_2^2z_1^4 - 32400z_2z_1^4 - 2700z_2^3z_1^3 + 6265z_2^2z_1^3 - 2460z_2z_1^3 + 135z_1^3 - 144z_2^3z_1^2 + 152z_2^2z_1^2 - 49z_2z_1^2 + 5z_1^2 \Big) \Big) = -\frac{1}{400z_2\Delta} \Big(-25000z_2^4z_1^4 + 81000z_2^2z_1^4 - 32400z_2z_1^4 + 2625z_2^2z_1^3 - 2460z_2z_1^3 + 135z_1^3 - 144z_2^3z_1^2 + 152z_2^2z_1^2 - 49z_2z_1^2 + 5z_1^2 \Big) \Big) = -\frac{1}{400z_2\Delta} \Big(-25000z_2^4z_1^4 + 81000z_2^2z_1^4 - 32400z_2z_1^4 - 2700z_2^3z_1^3 + 6265z_2^2z_1^3 - 2460z_2z_1^3 + 135z_1^3 - 144z_2^3z_1^2 + 152z_2^2z_1^2 - 49z_2z_1^2 + 5z_1^2 \Big) \Big)$
$f_2^{11} = -\frac{1}{400z_1\Delta} \Big(-200000z_1^2 z_2^6 + 50000z_1^2 z_2^5 - 31600z_1 z_2^5 - 1152z_2^5 + 3000z_1^2 z_2^4 + 21320z_1 z_2^4 + 864z_2^4 - 1200z_1^2 z_2^3 - 4815z_1 z_2^3 - 216z_2^3 + 365z_1 z_2^2 + 18z_2^2 \Big) \Big) \Big(-20000z_1^2 z_2^6 + 50000z_1^2 z_2^5 - 1152z_2^5 + 3000z_1^2 z_2^4 + 21320z_1 z_2^4 + 864z_2^4 - 1200z_1^2 z_2^3 - 216z_2^3 + 365z_1 z_2^2 + 18z_2^2 \Big) \Big) \Big) \Big) \Big(-20000z_1^2 z_2^6 + 50000z_1^2 z_2^5 - 31600z_1 z_2^5 - 1152z_2^5 + 3000z_1^2 z_2^4 + 21320z_1 z_2^4 + 864z_2^4 - 1200z_1^2 z_2^3 - 4815z_1 z_2^3 - 216z_2^3 + 365z_1 z_2^2 + 18z_2^2 \Big) $
$f_2^{12} = -\frac{1}{400\Delta} \Big(10000z_1^2 z_2^5 - 12500z_1^2 z_2^4 + 12800z_1 z_2^4 + 576z_2^4 - 14000z_1^2 z_2^3 - 7360z_1 z_2^3 - 432z_2^3 + 3600z_1^2 z_2^2 + 1220z_1 z_2^2 + 108z_2^2 - 45z_1 z_2 - 9z_2 \Big) \Big) \Big(-\frac{1}{400\Delta} \Big) \Big(-\frac{1}{400\Delta} \Big) \Big) \Big(-\frac{1}{400\Delta} \Big) \Big) \Big(-\frac{1}{400\Delta} \Big) \Big) \Big) \Big(-\frac{1}{400\Delta} \Big) $
$f_2^{22} = -\frac{1}{400\Delta} \Big(-50000z_1^3 z_2^4 + 125000z_1^3 z_2^3 - 4900z_1^2 z_2^3 - 288z_1 z_2^3 + 57000z_1^3 z_2^2 + 31405z_1^2 z_2^2 + 1216z_1 z_2^2 - 10800z_1^3 z_2 - 12985z_1^2 z_2 - 554z_1 z_2 + 1485z_1^2 + 67z_1 \Big) - 5000z_1^3 z_2^3 - 5000z_1^3 z_2^3 - 5000z_1^3 z_2^3 + 57000z_1^3 z_2^3 + 5700z_1^3 z_2^3 + 5700z_1^3 z_2^3 + 57000z_1^3 z_2^3 + 5700z_1^3 z_2^3 + 57000z_1^3 z_2^3 + 5700z_1^3 z_2^3$
$\bar{f}_{11}^1 = \frac{1}{20z_2\Delta} \Big(-105000z_1^2 z_2^3 - 100z_1 z_2^3 - 4500z_1^2 z_2^2 - 16535z_1 z_2^2 - 528z_2^2 + 6957z_1 z_2 + 248z_2 - 783z_1 - 29 \Big) - 528z_1^2 - 528z_2^2 - 528z_2^2 + 6957z_1 z_2 - 528z_2^2 - 528z_2^2$
$\bar{f}_{11}^2 = \frac{1}{20z_2^2\Delta} \Big(15000z_2^3 z_1^3 + 13500z_2^2 z_1^3 + 50z_2^3 z_1^2 + 2855z_2^2 z_1^2 - 216z_2 z_1^2 - 81z_1^2 + 84z_2^2 z_1 - 9z_2 z_1 - 3z_1 \Big)$
$\bar{f}_{12}^1 = \frac{1}{20z_1\Delta} \Big(-10000z_1^2 z_2^3 - 200z_1 z_2^3 - 1500z_1^2 z_2^2 - 1470z_1 z_2^2 - 48z_2^2 + 664z_1 z_2 + 24z_2 - 81z_1 - 3 \Big)$
$\bar{f}_{12}^2 = \frac{1}{20z_2\Delta} \Big(-7500z_1^2 z_2^3 + 100z_1 z_2^3 + 4500z_1^2 z_2^2 - 465z_1 z_2^2 - 16z_2^2 + 243z_1 z_2 + 8z_2 - 27z_1 - 1 \Big)$
$\bar{f}_{22}^{1} = \frac{1}{20z_{1}\Delta} \Big(5000z_{1}z_{2}^{4} - 400z_{2}^{4} - 500z_{1}z_{2}^{3} + 760z_{2}^{3} - 297z_{2}^{2} + 33z_{2} \Big)$
$\bar{f}_{22}^2 = \frac{1}{20z_1\Delta} \Big(-90000z_1^2 z_2^3 + 200z_1 z_2^3 + 1500z_1^2 z_2^2 - 13530z_1 z_2^2 - 400z_2^2 + 6086z_1 z_2 + 200z_2 - 729z_1 - 25 \Big)$

Any other combination of indices follows by symmetry. We find the following holomorphic ambiguities:

$f^{(2,0)} = -\frac{1}{\Delta^2} \left(\frac{31}{9216} - \frac{151}{2880} z_1 + \frac{49}{160} z_1^2 - \frac{143}{180} z_1^3 + \frac{139}{180} z_1^4 + \frac{51}{512} z_2 - \frac{2431}{1536} z_1 z_2 + \frac{10943}{152} z_1^2 z_2 - \frac{2557}{96} z_1^3 z_2 + \frac{2185}{72} z_1^4 z_2 + \frac{13851}{5120} z_2^2 - \frac{23553}{512} z_1 z_2^2 + \frac{309015}{1024} z_1^2 z_2^2 - \frac{23553}{1024} z_1^2 z_2^2 + \frac{13851}{1024} z_1^2 $	
$-\frac{4593875}{4608}z_1^3z_2^2+\frac{453125}{288}z_1^4z_2^2-\frac{179375}{288}z_1^5z_2^2-\frac{264375}{512}z_1^3z_2^3+\frac{1571875}{1536}z_1^4z_2^3+\frac{671875}{128}z_1^5z_2^3+\frac{212890625}{3072}z_1^6z_2^4\right)$	(A.14)
$f^{(1,1)} = -\frac{1}{\Delta^2} \left(-\frac{331}{57600} + \frac{257}{3600} z_1 - \frac{61}{200} z_1^2 + \frac{109}{225} z_1^3 - \frac{7}{45} z_1^4 - \frac{9}{640} z_2 - \frac{549}{640} z_1 z_2 + \frac{3493}{288} z_1^2 z_2 - \frac{1205}{24} z_1^3 z_2 + \frac{1175}{18} z_1^4 z_2 - \frac{2511}{1280} z_2^2 + \frac{2673}{128} z_1 z_2^2 - \frac{16035}{256} z_1^2 z_2^2 + \frac{310625}{1152} z_1^2 z_2 + \frac{1175}{128} z_1^2 z_2 - \frac{1175}{1280} z_1^2 z_2 + \frac{1175}{1280} z_2 + \frac{1175}{1280} z_2 + \frac{1175}{1$	$z_1^3 z_2^2$
$-\frac{113375}{72}z_1^4z_2^2+\frac{190625}{72}z_1^5z_2^2+\frac{406875}{128}z_1^3z_2^3-\frac{2615625}{128}z_1^4z_2^3+\frac{3015625}{96}z_1^5z_2^3+\frac{1953125}{256}z_1^6z_2^4\bigg)$	(A.15)
$f^{(0,2)} = -\frac{1}{\Delta^2} \left(\frac{167}{14400} - \frac{121}{720} z_1 + \frac{271}{300} z_1^2 - \frac{479}{225} z_1^3 + \frac{416}{225} z_1^4 + \frac{1419}{3200} z_2 - \frac{60691}{9600} z_1 z_2 + \frac{13353}{400} z_1^2 z_2 - \frac{27973}{360} z_1^3 z_2 + \frac{610}{9} z_1^4 z_2 + \frac{3807}{640} z_2^2 - \frac{28197}{320} z_1 z_2^2 + \frac{63135}{128} z_1^2 z_2^2 - \frac{28197}{320} z_1 z_2 + \frac{12353}{320} z_1^2 z_2 - \frac{28197}{360} z_1^2 z_2 - \frac{28197}{360} z_1^2 z_2 - \frac{28197}{360} z_1^2 z_2 - \frac{28197}{360} z_1^2 z_2 - \frac{28197}{320} z_1 z_2 + \frac{28197}{320} z_1^2 z_2 - \frac{28197}{320} z_1^$	
$-\frac{1513675}{1152}z_1^3z_2^2+\frac{119125}{72}z_1^4z_2^2-\frac{16375}{24}z_1^5z_2^2-\frac{38625}{64}z_1^3z_2^3+\frac{610625}{192}z_1^4z_2^3-\frac{828125}{288}z_1^5z_2^3+\frac{5078125}{384}z_1^6z_2^4\right)$	(A.16)

A. Klemm, M. Poretschkin, T. Schimannek, M. Westerholt-Raum

 $f^{(3,0)} = \frac{1}{\Delta^4} \left(-\frac{45893}{3222432000} + \frac{76957}{145152000} z_1 - \frac{5771}{907200} z_1^2 + \frac{82289}{1814400} z_1^3 - \frac{193997}{907200} z_1^4 + \frac{391327}{567000} z_1^5 - \frac{105361}{70875} z_1^6 + \frac{27334}{14175} z_1^7 - \frac{2294}{2025} z_1^8 + \frac{4703}{4300800} z_2 - \frac{2092763}{46448640} z_1 z_2 + \frac{14325557}{19353600} z_1^2 z_2 - \frac{30369853}{4838400} z_1^3 z_2 + \frac{110692871}{3628800} z_1^4 z_2 - \frac{27287759}{130200} z_1^5 z_2 + \frac{13058167}{76600} z_1^6 z_2 - \frac{2544593}{11340} z_1^7 z_2 + \frac{3407}{211} z_1^8 z_2 - \frac{1088733}{1638400} z_2^2 + \frac{74510981}{3440640} z_1 z_2^2 + \frac{3117875}{3446471} z_1^9 z_2^2 - \frac{281433165299}{19353600} z_1^4 z_2^2 + \frac{25441364153}{483840} z_1^5 z_2^2 - \frac{963971939}{48384} z_1^5 z_2^2 - \frac{963971939}{48384} z_1^6 z_2^2 + \frac{270034265}{1512} z_1^7 z_2^2 - \frac{148713575}{1008} z_1^8 z_2^2 + \frac{3117875}{54} z_1^9 z_2^2 + \frac{2716225221}{249120} z_1^3 z_2^3 + \frac{2928229314891}{2989728} z_1^4 z_2^3 - \frac{200845661185}{165888} z_1^5 z_2^3 + \frac{306851519075}{96768} z_1^6 z_2^3 - \frac{269782983125}{48384} z_1^7 z_2^3 + \frac{293629314891}{483840} z_1^2 z_2^4 + \frac{685564539978}{48384} z_1^2 z_2^4 + \frac{685564539978}{48384} z_1^2 z_2^4 + \frac{685564539728}{48384} z_1^2 z_2^4 + \frac{685564539728}{483840} z_1^2 z_2^4 + \frac{2984309687875}{48384} z_1^7 z_2^3 + \frac{2992829314891}{2989728} z_1^2 z_2^3 + \frac{200845661185}{165888} z_1^5 z_2^3 + \frac{306851519075}{96768} z_1^6 z_2^3 - \frac{269782983125}{48384} z_1^7 z_2^3 + \frac{2992829314891}{2989728} z_1^2 z_2^4 + \frac{685564539372}{485752} z_1^3 z_2^4 + \frac{879922720925}{688128} z_1^4 z_2^4 + \frac{2984309687875}{4128768} z_1^5 z_2^4 + \frac{2984309687875}{48384} z_1^7 z_2^4 + \frac{2984309687875}{78767} z_1^7 z_2^5 - \frac{307776826953125}{78767} z_1^7 z_2^5 - \frac{7648528515625}{72576} z_1^1 z_2^4 + \frac{949714875}{6182} z_1^2 z_2^4 + \frac{2984309687875}{7168} z_1^7 z_2^5 - \frac{307776826953125}{38772} z_1^8 z_2^8 + \frac{5600248046875}{48384} z_1^9 z_2^5 - \frac{15414913125}{7168} z_1^4 z_2^5 + \frac{1470126953125}{7168} z_1^4 z_2^5 - \frac{15414913125}{7168} z_1^4 z_2^5 + \frac{16567179677}{7168} z_1^8 z_2^6 - \frac{7648528515625}{72576}$

 $f^{(2,1)} = \frac{1}{\Delta 4} \left(\frac{625649}{387072000} - \frac{961951}{241920000} z_1 + \frac{255053}{6048000} z_1^2 - \frac{772259}{3024000} z_1^3 + \frac{1507181}{1512000} z_1^4 - \frac{2560261}{945000} z_1^5 + \frac{624748}{118125} z_1^6 - \frac{161746}{12625} z_1^7 + \frac{2902}{675} z_1^8 - \frac{590831}{43008000} z_2 + \frac{217453951}{387072000} z_1 z_2 - \frac{300910403}{32256000} z_1^2 z_2 + \frac{73848433}{386000} z_1^3 z_2 - \frac{2595602309}{6048000} z_1^4 z_2 + \frac{692727399}{504000} z_1^5 z_2 - \frac{2289351}{8400} z_1^6 z_2 - \frac{60540149}{18900} z_1^7 z_2 - \frac{189563}{105} z_1^8 z_2 - \frac{14828469}{4096000} z_2^2 - \frac{159168133}{14336000} z_1^2 z_2 + \frac{12528897251}{179200} z_1^4 z_2^2 - \frac{442311668733}{1612800} z_1^5 z_2^2 + \frac{8998561657}{13440} z_1^2 z_2 - \frac{4141160927}{40022} z_1^7 z_2^2 + \frac{115010095}{126} z_1^8 z_2^2 - \frac{13846175}{36} z_1^9 z_2^2 - \frac{13846175}{36} z_1^9 z_2^2 - \frac{13846175}{322} z_1^9 z_2^2 - \frac{13846175}{322} z_1^9 z_2^2 - \frac{11384717}{1228000} z_1^3 z_2^3 + \frac{17273342717}{1228000} z_1^3 z_2^3 - \frac{11697769909939}{16428000} z_1^4 z_2^3 + \frac{296490468347}{10592} z_1^5 z_2^3 - \frac{221032737295}{32256} z_1^6 z_2^3 + \frac{261061606225}{3064} z_1^7 z_2^3 - \frac{232881795375}{3024} z_1^2 z_2^3 + \frac{17273342717}{1433600} z_1^4 z_2^4 - \frac{3753997083}{1433600} z_1^2 z_2^4 - \frac{572678109}{143360} z_1^2 z_2^4 - \frac{572678109}{143360} z_1^2 z_2^4 - \frac{5523242725}{32256} z_1^6 z_2^3 + \frac{26046168627}{258048} z_1^7 z_2^3 + \frac{26494058347}{143360} z_1^2 z_2^4 - \frac{572678109}{143360} z_1^2 z_2^4 - \frac{5523242725}{14336} z_1^3 z_2^5 + \frac{260236067875}{258048} z_1^4 z_2^5 - \frac{260236067875}{14336} z_1^4 z_2^5 - \frac{260236067875}{14336} z_1^4 z_2^5 - \frac{260236067875}{14336} z_1^2 z_2^5 - \frac{272775390625}{16390} z_1^2 z_2^4 - \frac{5523242725}{14336} z_1^3 z_2^5 + \frac{260236067875}{14336} z_1^4 z_2^5 + \frac{27277539062}{14336} z_1^2 z_2^5 - \frac{3761371484375}{14336} z_1^2 z_2^5 - \frac{280236067875}{14335} z_1^2 z_2^5 + \frac{260236067875}{14336} z_1^2 z_2^5 - \frac{28023606787$

 $f^{(1,2)} = \frac{1}{\Delta^4} \left(-\frac{127}{193536} + \frac{107687}{672000} z_1 - \frac{2554393}{15120000} z_1^2 + \frac{772703}{756000} z_1^2 - \frac{254039}{63000} z_1^2 + \frac{522729}{756000} z_1^2 - \frac{2709046}{118125} z_1^6 + \frac{1213208}{39375} z_1^7 - \frac{66784}{3375} z_1^8 + \frac{1117523}{80640000} z_2 - \frac{92939087}{96768000} z_1 z_2 + \frac{1962311}{100000} z_1^2 z_2 - \frac{1150991209}{6048000} z_1^3 z_2 + \frac{792346361}{756000} z_1^4 z_2 - \frac{6475451077}{1890000} z_1^5 z_2 + \frac{3122609}{4725} z_1^6 z_2 - \frac{33107323}{4725} z_1^7 z_2 + \frac{3047144}{945} z_1^8 z_2 - \frac{1053221}{204800} z_1^2 z_2^2 + \frac{542284461}{16128000} z_1^2 z_2^2 + \frac{542284621}{16128000} z_1^4 z_2 - \frac{6475451077}{1512000} z_1^4 z_2^2 + \frac{105139716709}{302400} z_1^2 z_2^2 - \frac{27852246421}{302400} z_1^2 z_2^2 + \frac{236359459}{1512} z_1^7 z_2^2 - \frac{20906555}{189} z_1^8 z_2^2 + \frac{18496600}{27} z_1^9 z_2^2 + \frac{24849600}{27} z_1^9 z_2^2 + \frac{24849610}{27} z_1^2 z_2^2 + \frac{24849610}{1027} z_1^2 z_2^2 + \frac{24849600}{12096} z_1^4 z_2^2 + \frac{104053359459}{1512} z_1^7 z_2^2 - \frac{209026555}{189} z_1^8 z_2^2 + \frac{18496600}{27} z_1^9 z_2^2 + \frac{24849600}{1206} z_1^2 z_2^2 + \frac{24849600}{12096} z_1^2$

 $f^{(0,3)} = \frac{1}{\Delta 4} \left(\frac{64675591}{5866800000} - \frac{54856777}{1814400000} z_1 + \frac{65194601}{181440000} z_1^2 - \frac{55731167}{22680000} z_1^3 + \frac{242045929}{22680000} z_1^4 - \frac{216143287}{7087500} z_1^5 + \frac{200947999}{3543750} z_1^6 - \frac{22452548}{3543750} z_1^6 - \frac{22452548}{3543750} z_1^7 + \frac{1654522}{5625} z_1^8 + \frac{40894657}{645120000} z_2 + \frac{8658850111}{5866800000} z_1 z_2 + \frac{697315817}{48384000} z_1^2 z_2 - \frac{594293389}{8064000} z_1^3 z_2 + \frac{122562197}{567000} z_1^4 z_2 - \frac{975682873}{2520000} z_1^5 z_2 + \frac{54401369}{105000} z_1^6 z_2 - \frac{187652267}{283500} z_1^7 z_2 + \frac{2428696}{4725} z_1^8 z_2 + \frac{84392949}{20480000} z_2^2 + \frac{24515809627}{215040000} z_1 z_2 + \frac{598212759449}{430080000} z_1^2 z_2^2 - \frac{19527458020663}{1935380000} z_1^3 z_2^2 + \frac{1494655943567}{393580000} z_1^4 z_2^2 - \frac{1392743766707}{8064000} z_1^5 z_2^2 + \frac{25836031797}{604800} z_1^6 z_2^2 - \frac{944837177}{1344} z_1^7 z_2^2 + \frac{149465943567}{1935380000} z_1^2 z_2^2 + \frac{789111}{128000} z_1^3 z_2^3 + \frac{1595793}{512000} z_1 z_2^3 + \frac{2572634392021}{8040000} z_1^3 z_2^3 + \frac{165039233141773}{202243200} z_1^4 z_2^3 - \frac{908991091601}{331776} z_1^5 z_2^3 + \frac{157768}{4032} z_1^2 z_2^3 - \frac{2572634392021}{1832000} z_1^3 z_2^3 + \frac{165039233141773}{202243200} z_1^4 z_2^3 - \frac{90891091601}{331776} z_1^5 z_2^3 + \frac{157768}{4032} z_1^2 z_2^3 + \frac{15197463}{3584000} z_1^2 z_2^3 - \frac{2572634392021}{1832000} z_1^3 z_2^3 + \frac{165039233141773}{232432000} z_1^4 z_2^3 - \frac{90891091601}{331776} z_1^5 z_2^3 + \frac{151768}{4032} z_1^2 z_2^3 + \frac{15197463}{3584000} z_1^2 z_2^3 + \frac{15197463}{13832000} z_1^2 z_2^3 + \frac{15197463}{13832000} z_1^2 z_2^3 + \frac{15197453}{232432000} z_1^2 z_2^3 + \frac{2533321117}{1792} z_1^2 z_2^4 - \frac{2039769849}{1792} z_1^3 z_2^4 + \frac{10569277770}{12504} z_1^2 z_2^3 + \frac{1046099778775}{264} z_1^2 z_2^3 - \frac{28840191740875}{129024} z_1^2 z_2^2 + \frac{10461879}{193566} z_1^2 z_2^3 + \frac{10569277776}{12504} z_1^2 z_2^3 + \frac{105696777775}{25777} z_1^2 z_2^3 + \frac{105696777775}{2527777} z_1^2 z_2^3 + \frac{105696777775}{25277777} z_1^2 z_2^3 + \frac{104609778777}{2567} z_1^$

B | The Ambiguities for $\mathbb{C}^3/\mathbb{Z}_6$

$A_{1} = \frac{1}{15z_{1}\Delta_{1}} \left(2 - 12z_{1} + 16z_{1}^{2} - 64z_{3}z_{1}^{2} - 16z_{2} + 102z_{1}z_{2} - 144z_{1}^{2}z_{2} + 576z_{3}z_{1}^{2}z_{2} + 32z_{2}^{2} - 216z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} - 1512z_{3}z_{1}^{2}z_{2}^{2} + 108z_{1}^{3}z_{2}^{2} - 432z_{3}z_{1}^{3}z_{2}^{2} + 216z_{1}^{2}z_{2}^{3} - 216z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} - 1512z_{3}z_{1}^{2}z_{2}^{2} + 108z_{1}^{3}z_{2}^{2} - 432z_{3}z_{1}^{3}z_{2}^{2} + 216z_{1}^{2}z_{2}^{3} - 216z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} - 1512z_{3}z_{1}^{2}z_{2}^{2} + 108z_{1}^{3}z_{2}^{2} - 432z_{3}z_{1}^{3}z_{2}^{2} + 216z_{1}^{2}z_{2}^{3} - 216z_{1}z_{2}^{3}z_{2}^{2} - 216z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} - 1512z_{3}z_{1}^{2}z_{2}^{2} + 108z_{1}^{3}z_{2}^{2} - 432z_{3}z_{1}^{3}z_{2}^{2} + 216z_{1}^{2}z_{2}^{3} - 216z_{1}z_{2}^{3}z_{2}^{3} - 216z_{1}z_{2}^{3} - 216z_{1}z_{2}^{3}$	
$+864z_3z_1^2z_2^3 - 486z_1^3z_2^3 + 1944z_3z_1^3z_2^3 \Big)$	(B.1)
$A_{2} = \frac{1}{15z_{2}\Delta_{1}} \left(2 - 16z_{1} + 32z_{1}^{2} - 128z_{3}z_{1}^{2} - 12z_{2} + 102z_{1}z_{2} - 216z_{1}^{2}z_{2} + 864z_{3}z_{1}^{2}z_{2} + 16z_{2}^{2} - 144z_{1}z_{2}^{2} + 270z_{1}^{2}z_{2}^{2} - 1512z_{3}z_{1}^{2}z_{2}^{2} + 216z_{1}^{3}z_{2}^{2} - 864z_{3}z_{1}^{3}z_{2}^{2} + 108z_{1}^{2}z_{2}^{3} \right)$	
$+432z_3z_1^2z_2^3 - 486z_1^3z_2^3 + 1944z_3z_1^3z_2^3$	(B.2)
$f_1^{11} = \frac{1}{100\Delta_1} \left(-9z_1 + 72z_1^2 - 144z_1^3 + 320z_3z_1^3 + 81z_1z_2 - 684z_1^2z_2 + 1440z_1^3z_2 - 3200z_3z_1^3z_2 - 216z_1z_2^2 + 1908z_1^2z_2^2 - 3726z_1^3z_2^2 + 10428z_3z_1^3z_2^2 - 1944z_1^4z_2^2 + 6720z_3z_1^4z_2 + 10428z_3z_1^3z_2 - 1042z_1^3z_2 + 10428z_3z_1^3z_2 - 1042z_1^3z_2 + 10428z_3z_1^3z_2 - 1042z_1^3z_2 + 1042z_1^3z_2 - 1042z_1^3z_2 + 1042z_1^3z_2 - 104z_1^3z_2 - 104z_1^3z_2 - 104z_1^3z_2 - 104z_1^3z_2 - 104z_1^3z_2 - 104z_$	2
- -4800z_3z_1^5z_2^2 + 19200z_3^2z_1^5z_2^2 + 144z_1z_2^3 - 1296z_1^2z_2^3 + 486z_1^3z_2^3 - 12492z_3z_1^3z_2^3 + 10692z_1^4z_2^3 - 36960z_3z_1^4z_2^3 + 28800z_3z_1^5z_2^3 - 115200z_3^2z_1^5z_2^3 + 1944z_1^3z_2^4 + 7728z_3z_1^3z_2^3 + 10692z_1^4z_2^3 - 36960z_3z_1^4z_2^3 + 28800z_3z_1^5z_2^3 - 115200z_3^2z_1^5z_2^3 + 1944z_1^3z_2^4 + 7728z_3z_1^3z_2^3 + 10692z_1^4z_2^3 - 36960z_3z_1^4z_2^3 + 28800z_3z_1^5z_2^3 - 115200z_3^2z_1^5z_2^3 + 1944z_1^3z_2^4 + 7728z_3z_1^3z_2^3 + 10692z_1^4z_2^3 - 36960z_3z_1^4z_2^3 + 28800z_3z_1^5z_2^3 - 115200z_3^2z_1^5z_2^3 + 1944z_1^3z_2^4 + 7728z_3z_1^3z_2^3 + 10692z_1^4z_2^3 - 1260z_1^3z_1^3 - 1260z_1^3 - 1260z_1^3 - 1260z_1^3 - 1260z_1^3 - 1260z_1^3 - 1260z_1^3	$z_1^3 z_2^4$
$-8748z_1^4z_2^4+28656z_3z_1^4z_2^4-6561z_1^5z_2^4+6804z_3z_1^5z_2^4+77760z_3^2z_1^5z_2^4+6561z_1^5z_2^5-52164z_3z_1^5z_2^5+103680z_3^2z_1^5z_2^5\Big)$	(B.3)
$f_1^{12} = \frac{1}{200\Delta_1} \left(9z_2 - 72z_1z_2 + 144z_1^2z_2 - 320z_3z_1^2z_2 - 108z_2^2 + 900z_1z_2^2 - 1872z_1^2z_2^2 + 4832z_3z_1^2z_2^2 + 432z_3^2 - 3744z_1z_3^2 + 7614z_1^2z_2^3 - 25164z_3z_1^2z_2^3 + 1944z_1^3z_2^3 - 6720z_3z_1^2z_2 + 4832z_3z_1^2z_2^2 + 432z_3^2 - 3744z_1z_3^2 + 7614z_1^2z_2^3 - 25164z_3z_1^2z_2^3 + 1944z_1^3z_2^3 - 6720z_3z_1^2z_2 + 4832z_3z_1^2z_2^2 + 432z_3^2 - 3744z_1z_2^3 - 3744z_1z_2^3 - 25164z_3z_1^2z_2^3 + 1944z_1^3z_2^3 - 6720z_3z_1^2z_2 + 4832z_3z_1^2z_2^2 + 432z_3^2 - 3744z_1z_2^3 - 3744z_1z_2^3 - 25164z_3z_1^2z_2^3 + 1944z_1^3z_2^3 - 6720z_3z_1^2z_2 + 4832z_3z_1^2z_2^2 + 432z_3^2 - 3744z_1z_2^3 -$	z_{2}^{3}
$+4800z_3z_1^4z_2^3 - 19200z_3^2z_1^4z_2^3 - 576z_2^4 + 5184z_1z_2^4 - 7776z_1^2z_2^4 + 51552z_3z_1^2z_2^4 - 16524z_1^3z_2^4 + 57912z_3z_1^3z_2^4 - 26640z_3z_1^4z_2^4 + 106560z_3^2z_1^4z_2^4 - 7776z_1^2z_2^5 - 30912z_3z_1^2z_2^4 + 57912z_3z_1^3z_2^4 - 57912z_3^5 -$	$z_1^2 z_2^5$
$+ 34992z_1^3z_2^5 - 124128z_3z_1^3z_2^5 + 6561z_1^4z_2^5 - 16524z_3z_1^4z_2^5 - 38880z_3^2z_1^4z_2^5 - 26244z_1^4z_2^6 + 208656z_3z_1^4z_2^6 - 414720z_3^2z_1^4z_2^6 + 2286z_3^2z_1^4z_2^6 + 2286z_3^2z_1^2z_2^6 + 2286z_3^2z_2^6 + 2286z_3^2 $	(B.4)
$f_1^{22} = \frac{1}{100z_1\Delta_1} \left(-9z_2^2 + 72z_1z_2^2 - 144z_1^2z_2^2 + 512z_3z_1^2z_2^2 + 108z_3^2 - 900z_1z_2^3 + 1872z_1^2z_2^3 - 6800z_3z_1^2z_2^3 - 432z_2^4 + 3744z_1z_2^4 - 7614z_1^2z_2^4 + 31836z_3z_1^2z_2^4 - 1944z_1^3z_2^4 + 7512z_1^2z_2^3 - 7614z_1^2z_2^4 + 7512z_1^2z_2^4 - 7612z_1^2z_2^4 - 7614z_1^2z_2^4 - 7612z_1^2z_2^4 - 7612z_1^2z_$	$z_3 z_1^3 z_2^4$
$-1200z_3z_1^4z_2^4 + 4800z_3^2z_1^4z_2^4 + 576z_2^5 - 5184z_1z_2^5 + 7776z_1^2z_2^5 - 59040z_3z_1^2z_2^5 + 16524z_1^3z_2^5 - 63456z_3z_1^3z_2^5 + 6120z_3z_1^4z_2^5 - 24480z_3^2z_1^4z_2^5 + 7776z_1^2z_2^6 + 30912z_3z_1^2z_2^5 + 16524z_1^3z_2^5 - 63456z_3z_1^3z_2^5 + 6120z_3z_1^4z_2^5 - 24480z_3^2z_1^4z_2^5 + 7776z_1^2z_2^6 + 30912z_3z_1^2z_2^5 + 16524z_1^3z_2^5 - 63456z_3z_1^3z_2^5 + 6120z_3z_1^4z_2^5 - 24480z_3^2z_1^4z_2^5 + 7776z_1^2z_2^6 + 30912z_3z_1^2z_2^5 + 16524z_1^3z_2^5 - 63456z_3z_1^3z_2^5 + 6120z_3z_1^4z_2^5 - 24480z_3^2z_1^4z_2^5 + 7776z_1^2z_2^6 + 30912z_3z_1^2z_2^5 + 16524z_1^3z_2^5 - 63456z_3z_1^3z_2^5 + 6120z_3z_1^4z_2^5 - 24480z_3^2z_1^4z_2^5 + 7776z_1^2z_2^6 + 30912z_3z_1^2z_2^5 + 7776z_1^2z_2^6 + 77776z_1^2z_2^6 + 7776z_1^2z$.6 2
$-34992z_1^3z_2^6 + 133632z_3z_1^3z_2^6 - 6561z_1^4z_2^6 + 45684z_3z_1^4z_2^6 - 77760z_3^2z_1^4z_2^6 + 26244z_1^4z_2^7 - 208656z_3z_1^4z_2^7 + 414720z_3^2z_1^4z_2^7 + 26244z_1^4z_2^6 + 2624z_1^4z_2^6 + 2624z_1^4z_1^6 + 2624z_1^4z_1^6 + 2624z_1^4z_1^6 + 2624z_1^4z_1^6 + 2624z_1^4z_1^6 + 2624z_1^6 + 2$	(B.5)
$f_2^{11} = \frac{1}{200\Delta_1} \left(9z_1^2 - 72z_1^3 + 144z_1^4 - 320z_3z_1^4 - 72z_1^2z_2 + 612z_1^3z_2 - 1296z_1^4z_2 + 1872z_3z_1^4z_2 + 9600z_3z_1^5z_2 - 19200z_3z_1^6z_2 + 76800z_3^2z_1^6z_2 + 144z_1^2z_2^2 - 1296z_1^3z_2^2 + 2430z_1^4z_2 + 1872z_3z_1^4z_2 + 9600z_3z_1^5z_2 - 19200z_3z_1^6z_2 + 76800z_3^2z_1^6z_2 + 144z_1^2z_2^2 - 1296z_1^3z_2^2 + 2430z_1^4z_2 + 1872z_3z_1^4z_2 + 9600z_3z_1^5z_2 - 19200z_3z_1^6z_2 + 76800z_3^2z_1^6z_2 + 144z_1^2z_2^2 - 1296z_1^3z_2^2 + 2430z_1^4z_2 + 1872z_3z_1^4z_2 + 9600z_3z_1^5z_2 - 19200z_3z_1^6z_2 + 76800z_3^2z_1^6z_2 + 144z_1^2z_2^2 - 1296z_1^3z_2^2 + 2430z_1^4z_2 + 1872z_3z_1^4z_2 + 1872z_3z_1^4z_2 + 1872z_3z_1^4z_2 + 1872z_3z_1^4z_2 + 1872z_3z_1^5z_2 - 19200z_3z_1^5z_2 + 76800z_3^2z_1^6z_2 + 144z_1^2z_2^2 - 1296z_1^3z_2^2 + 2430z_1^4z_2 + 1872z_3z_1^4z_2 + 1872z_3z_2 + 1872z_3z_2 + 1872z_3z_2 + 1872z_3z_2 + 1872z_3z_2 + 1872z$	z_{2}^{2}
$-5796z_3z_1^4z_2^2 + 1944z_1^5z_2^2 - 35520z_3z_1^5z_2^2 + 72000z_3z_1^6z_2^2 - 288000z_3^2z_1^6z_2^2 + 1944z_1^4z_2^3 + 7728z_3z_1^4z_2^3 - 8748z_1^5z_2^3 + 28872z_3z_1^5z_2^3 - 38880z_3z_1^6z_2^3 + 155520z_3^2z_1^6z_2^3 + 155520z_3^2z_1^2z_2^3 + 155520z_3^2z_1^2z_2^$	
$+6561z_1^6z_2^4 - 52164z_3z_1^6z_2^4 + 103680z_3^2z_1^6z_2^4 \Big)$	(B.6)
$f_2^{12} = \frac{1}{100\Delta_1} \left(-9z_1z_2 + 72z_1^2z_2 - 144z_1^3z_2 + 512z_3z_1^3z_2 + 72z_1z_2^2 - 612z_1^2z_2^2 + 1296z_1^3z_2^2 - 4332z_3z_1^3z_2^2 - 2400z_3z_1^4z_2^2 + 4800z_3z_1^5z_2^2 - 19200z_3^2z_1^5z_2^2 - 144z_1z_2^3 + 1296z_1^3z_2^2 - 612z_1^2z_2^2 + 1296z_1^3z_2^2 - 4332z_3z_1^3z_2^2 - 2400z_3z_1^4z_2^2 + 4800z_3z_1^5z_2^2 - 19200z_3^2z_1^5z_2^2 - 144z_1z_2^3 + 1296z_1^3z_2^2 - 612z_1^2z_2^2 + 1296z_1^3z_2^2 - 4332z_3z_1^3z_2^2 - 2400z_3z_1^4z_2^2 + 4800z_3z_1^5z_2^2 - 19200z_3^2z_1^5z_2^2 - 144z_1z_2^3 + 1296z_1^3z_2^2 - 612z_1^2z_2^2 - 612z$	³ ₂₂
$-2430z_1^3z_2^3+11628z_3z_1^3z_2^3-1944z_1^4z_2^3+13632z_3z_1^4z_2^3-15840z_3z_1^5z_2^3+63360z_3^2z_1^5z_2^3-1944z_1^3z_2^4-7728z_3z_1^3z_2^4+8748z_1^4z_2^4-31248z_3z_1^4z_2^4+4860z_3z_1^5z_2^4-2860z_3z_1^5z_2^3+63360z_3z_1^5z_2^5+6326z_3z_1^5+6z_2^5$	
$-19440z_3^2 z_1^5 z_2^4 - 6561z_1^5 z_2^5 + 52164z_3 z_1^5 z_2^5 - 103680z_3^2 z_1^5 z_2^5 \Big)$	(B.7)
$f_2^{22} = -\frac{1}{200\Delta_1} \left(-27z_2 + 216z_1z_2 - 432z_1^2z_2 + 1728z_3z_1^2z_2 + 252z_2^2 - 2124z_1z_2^2 + 4464z_1^2z_2^2 - 17792z_3z_1^2z_2^2 - 720z_2^3 + 6336z_1z_2^3 - 12474z_1^2z_2^3 + 60684z_3z_1^2z_2^3 - 5832z_1^3z_2^3 $	
$+25728z_3z_1^3z_2^3-4800z_3z_1^4z_2^3+19200z_3^2z_1^4z_2^3+576z_2^4-5184z_1z_2^4+3888z_1^2z_2^4-75744z_3z_1^2z_2^4+34020z_1^3z_2^4-140856z_3z_1^3z_2^4+13680z_3z_1^4z_2^4-54720z_3^2z_1^4z_2^4+7776z_1^4z_2^4+2888z_1^2z_2^4-75744z_3z_1^2z_2^4+34020z_1^3z_2^4-140856z_3z_1^3z_2^4+13680z_3z_1^4z_2^4-54720z_3^2z_1^4z_2^4+7776z_1^4z_2^4+2888z_1^2z_2^4-75744z_3z_1^2z_2^4+34020z_1^3z_2^4-140856z_3z_1^3z_2^4+13680z_3z_1^4z_2^4-54720z_3^2z_1^4z_2^4+7776z_2^4+2888z_1^2z_2^4-75744z_3z_1^2z_2^4+34020z_1^3z_2^4-140856z_3z_1^3z_2^4+13680z_3z_1^4z_2^4-54720z_3^2z_1^4z_2^4+7776z_2^4+2888z_1^2z_2^4+7776z_2^4+2882z_1^2z_2^4+7776z_2^4+2880z_1^2z_2^4+7776z_2^4+77776z_2^4+77776z_2^4+77776z_2^4+777777777777777777777777777777777777$	$z_1^2 z_2^5$
$+ 30912z_3z_1^2z_2^5 - 34992z_1^3z_2^5 + 134496z_3z_1^3z_2^5 - 19683z_1^4z_2^5 + 156492z_3z_1^4z_2^5 - 311040z_3^2z_1^4z_2^5 + 26244z_1^4z_2^6 - 208656z_3z_1^4z_2^6 + 414720z_3^2z_1^4z_2^6 - 208656z_3z_1^4z_2^6 + 208656z_3z_1^6z_2^6 + 208656z_3z_1^6 + 208656z_3z_2^6 + 208656z_3z_1^6z_2^6 + 208656z_3z_1^6z$	(B.8)
-1] (-1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1 -1	2 2
$f_{11}^{1} = \frac{1}{10z_{1}\Delta_{1}} \left(-13 + 112z_{1} - 240z_{1}^{2} + 960z_{3}z_{1}^{2} + 104z_{2} - 949z_{1}z_{2} + 2128z_{1}^{2}z_{2} - 8672z_{3}z_{1}^{2}z_{2} + 80z_{1}^{2}z_{2} - 320z_{3}z_{1}^{3}z_{2} - 208z_{2}^{2} + 2008z_{1}z_{2}^{2} - 3911z_{1}^{2}z_{2}^{2} + 22932z_{3}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{2}^{2} + 2008z_{1}^{2}z_{2} - 300z_{1}^{2}z_{1}^{2}z_{1}^{2}z_{2}^{2} + 2008z_{1}^{2}z_{2}^{2} - 3911z_{1}^{2}z_{2}^{2} + 22932z_{3}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{2}^{2} + 2008z_{1}^{2}z_{1}^{2}z_{2}^{2} + 2008z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2}z_{2}^{2} + 2008z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{2}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z_{1}^{2} - 390z_{1}^{2}z$	$\int z_1^3 z_2^2$
$+15600z_3z_1^3z_2^2 - 16z_1z_3^2 - 3252z_1^2z_3^2 - 13584z_3z_1^2z_3^2 + 15849z_1^3z_3^2 - 63396z_3z_1^3z_2^3 + 540z_1^4z_3^2 - 4320z_3z_1^4z_3^2 + 8640z_3^2z_1^4z_3^2 - 108z_1^3z_2^4 + 432z_3z_1^3z_2^4 - 12960z_1^4z_2^4 + 432z_3z_1^3z_3^4 - 12960z_1^4z_2^4 + 12960z_1^4z_2^4 + 1292z_3z_1^3z_3^4 - 12960z_1^4z_2^4 + 1292z_3^2z_3^4 - 1292z_3^2z_3^2 - 1292z_3^2z_3^2 - 1292z_3^2 - 1292z_$	
$+103680z_3z_1^2z_2^2 - 207360z_3^2z_1^2z_2^2$	(B.9)
$f_{11}^{z} = \frac{1}{5z_{1}\Delta_{1}} \left[-2z_{2} + 8z_{1}z_{2} - 32z_{3}z_{1}z_{2} + 17z_{2}^{z} - 67z_{1}z_{2}^{z} + 308z_{3}z_{1}z_{2}^{z} - 20z_{1}^{z}z_{2}^{z} + 40z_{2}^{z} + 128z_{1}z_{2}^{z} - 912z_{3}z_{1}z_{2}^{z} + 264z_{1}^{z}z_{2}^{z} - 1056z_{3}z_{1}^{z}z_{2}^{z} + 48z_{1}z_{2}^{z} + 48z_{$	
$+768z_3z_1z_2^4 - 783z_1^2z_2^4 + 3132z_3z_1^2z_2^4 - 135z_1^3z_2^4 + 1080z_3z_1^3z_2^4 - 2160z_3^2z_1^3z_2^4 + 108z_1^2z_2^5 - 432z_3z_1^2z_2^5 + 810z_1^3z_2^5 - 6480z_3z_1^3z_2^5 + 12960z_3^2z_1^3z_2^5 \right)$	(B.10)
$\bar{f}_{12}^{1} = \frac{1}{5\Delta_1} \left(3z_1 - 32z_1^2 - 32z_3z_1^2 + 80z_1^3 - 320z_3z_1^3 - 11z_1z_2 + 160z_1^2z_2 + 240z_3z_1^2z_2 - 480z_1^3z_2 + 1920z_3z_1^3z_2 - 4z_1z_2^2 - 102z_1^2z_2^2 - 552z_3z_1^2z_2^2 + 459z_1^3z_2^2 - 1836z_3z_1^3z_2^2 - 1826z_3z_1^3z_2^2 - 182$	
$+540z_1^4z_2^2 - 4320z_3z_1^4z_2^2 + 8640z_3^2z_1^4z_2^2 - 27z_1^3z_2^3 + 108z_3z_1^3z_2^3 - 810z_1^4z_2^3 + 6480z_3z_1^4z_2^3 - 12960z_3^2z_1^4z_2^3 - 12960z_3^2z_1^2z_2^3 - 12960z_3^2z_2^3 - 12960z_3^2 - 12960z_3^2 - 12960z_3^2 - 12960z_3^3 - 12960z_3^3 - 12960z_3^3 - 12960z_3^3 - 12960z_3^3 - 12960z_3^3 - 12960z_3$	(B.11)
$\tilde{f}_{12}^2 = \frac{1}{10z_1\Delta_1} \left(-3 + 24z_1 - 48z_1^2 + 192z_3z_1^2 + 24z_2 - 207z_1z_2 + 464z_1^2z_2 - 1696z_3z_1^2z_2 - 80z_1^3z_2 + 320z_3z_1^3z_2 - 48z_2^2 + 440z_1z_2^2 - 949z_1^2z_2^2 + 4284z_3z_1^2z_2^2 - 204z_1^3z_2^2 + 816z_1^2z_2 - 80z_1^3z_1^2 + 816z_1^2z_1^2 - 80z_1^3z_1^2 - 80z_1^3z_1^3 - 80z_1^3z_1^3 - 80z_1^3z_1^3 - 80z_1^3z_1^3 - 80z_1^3z_1^3 - 80z_1^3z_1^$	$z_3 z_1^3 z_2^2$
$+16z_{1}z_{2}^{3}-636z_{1}^{2}z_{2}^{3}-1968z_{3}z_{1}^{2}z_{2}^{3}+2619z_{1}^{3}z_{2}^{3}-10476z_{3}z_{1}^{3}z_{2}^{3}-540z_{1}^{4}z_{2}^{3}+4320z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+108z_{1}^{3}z_{2}^{4}-432z_{3}z_{1}^{3}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+108z_{1}^{3}z_{2}^{4}-432z_{3}z_{1}^{3}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+108z_{1}^{3}z_{2}^{4}-432z_{3}z_{1}^{3}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+108z_{1}^{3}z_{2}^{4}-432z_{3}z_{1}^{3}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+108z_{1}^{3}z_{2}^{4}-432z_{3}z_{1}^{3}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{3}z_{1}^{4}z_{2}^{4}-1620z_{1}^{4}z_{2}^{4}+12960z_{1}^{4}z_{2}^{4}+12$	
$-25920z_3^2 z_1^4 z_2^4$	(B.12)
$\tilde{f}_{22}^{1} = \frac{1}{10z_{2}\Delta_{1}} \left(-13z_{1} + 124z_{1}^{2} - 368z_{1}^{3} + 832z_{3}z_{1}^{3} + 320z_{1}^{4} - 1280z_{3}z_{1}^{4} + 52z_{1}z_{2} - 521z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{2} + 1552z_{1}^{3}z_{2} - 4128z_{3}z_{1}^{3}z_{2} - 1200z_{1}^{4}z_{2} + 4800z_{3}z_{1}^{4}z_{2} - 4z_{1}^{2}z_{2}^{2} + 609z_{1}^{3}z_{2}^{2} + 2292z_{1}^{2}z_{1} + 220z_{1}^{2}z_{2} + 220z_{1}$	$z_3 z_1^3 z_2^2$
$-3024z_1^4z_2^2 + 12096z_3z_1^4z_2^2 + 2160z_1^5z_2^2 - 17280z_3z_1^5z_2^2 + 34560z_3^2z_1^5z_2^2 - 27z_1^4z_2^3 + 108z_3z_1^4z_2^3 + 1620z_1^5z_2^3 - 12960z_3z_1^5z_2^3 + 25920z_3^2z_1^5z_2^3 + 25920z_3^2z_1^2z_2^3 + 25920z_3^2z_1^2 + 25920z_3^2z_1^2 + 25920z_3^2 + 25920z_3^2 + 25920z_3^2 + 259$	(B.13)
$\tilde{f}_{22}^2 = \frac{1}{5z_2\Delta_1} \left(-8 + 64z_1 - 128z_1^2 + 512z_3z_1^2 + 51840z_3z_1^4 + 68z_2 - 581z_1z_2 + 1256z_1^2z_2 - 4864z_3z_1^2z_2 - 80z_1^3z_2 + 320z_3z_1^3z_2 - 144z_2^2 + 1307z_1z_2^2 - 2590z_1^2z_2^2 + 13368z_3z_1^2 + 512z_3z_1^2 + 512z_3z_1^2$	z_2^2
$-1464z_{1}^{3}z_{2}^{2}+5856z_{3}z_{1}^{3}z_{2}^{2}+4z_{1}z_{2}^{3}-1950z_{1}^{2}z_{2}^{3}-7656z_{3}z_{1}^{2}z_{2}^{3}+8775z_{1}^{3}z_{2}^{3}-35100z_{3}z_{1}^{3}z_{2}^{3}-540z_{1}^{4}z_{2}^{3}+4320z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+27z_{1}^{3}z_{2}^{4}-108z_{3}z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{3}-540z_{1}^{4}z_{2}^{3}+4320z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+27z_{1}^{3}z_{2}^{4}-108z_{3}z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{3}-540z_{1}^{4}z_{2}^{3}+4320z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+27z_{1}^{3}z_{2}^{4}-108z_{3}z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{3}-540z_{1}^{4}z_{2}^{3}+4320z_{3}z_{1}^{4}z_{2}^{3}-8640z_{3}^{2}z_{1}^{4}z_{2}^{3}+27z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+875z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+872z_{1}^{3}z_{2}^{4}-8640z_{1}^{3}z_{2}^{4}+822z_{1}^{3}z_{2}^{4}+822z_{1}^{3}z_{2}^{4}+822z_{1}^{3}z_{1}^{3}z_{2}^{4}-6480z_{1}^{4}z_{2}^{3}+872z_{1}^{3}z_{2}^{4}+822z_{1}^{3}z_{2}^{4}+822z_{1}^{3}z_{2}^{4}+82z_{1}^{3}+82z_{1}^{3}z_{2}^{4}+82z_{1}^{3}z_{2}^{4}+82z_{1}^{3}+82z_{1}^{3}+82z_{1}^{3}+82z_{1}^{3}+82z_{1}^{3}+82z_{1}^{3}+8z_{1}^{3}+82z_{1}^{3}+8z_{1}^{3}+82z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{1}^{3}+8z_{$	4 2)
	(B.14)

C | Siegel modular forms of genus two

In this appendix we introduce the basics of Siegel modular forms that are needed. Although many statements and definitions hold for arbitrary genus, we restrict ourselves to genus two. The Siegel upper halfplane is denoted by

$$\mathbb{H}^{(2)} = \left\{ \tau \in \mathbb{C}^{2 \times 2} | \tau^T = \tau, \quad \operatorname{Im} \tau \ge 0 \right\},$$
(C.1)

on which the homogenous modular group $\Gamma_2 = \text{Sp}(4, \mathbb{Z})$ operates by

$$\tau \mapsto (a\tau + b)(c\tau + d)^{-1}, \qquad \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \operatorname{Sp}(4, \mathbb{Z}).$$
 (C.2)

The quotient $\mathbb{H}^{(2)}$ by this action is called the Siegel fundamental domain $\mathscr{F}_2 = \Gamma_2 \setminus \mathbb{H}^{(2)}$. A Siegel modular form of weight *w* is a holomorphic function $f : \mathbb{H}^{(2)} \longrightarrow \mathbb{C}$, such that for all $\tau \in \mathbb{H}^{(2)}$ and $\gamma \in \Gamma_2$

$$f(\gamma\tau) = \det(c\tau + d)^{w} f(\tau), \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix},$$
(C.3)

holds. The space of all Siegel modular forms of weight w is denoted by $M_w(\mathbb{H}^{(2)})$. As $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ is contained in Γ_2 , any Siegel modular form f admits a Fourier expansion which reads

$$f = \sum_{T} a(T) \exp(2\pi i \operatorname{Tr}(T\tau)).$$
(C.4)

Here the summation is over all half integer matrices $T \in \frac{1}{2}\mathbb{Z}^{2\times 2}$ which have integer diagonal elements. Next we introduce the Siegel operator Φ , which is a map $M_w(\mathbb{H}^{(2)}) \longrightarrow M_w(\mathbb{H}^{(1)})$ and is defined by

$$\Phi f = \lim_{t \to \infty} f \begin{pmatrix} \tau & 0 \\ 0 & it \end{pmatrix}, \quad \tau \in \mathbb{H}, \ t \in \mathbb{R}.$$
(C.5)

The elements of ker Φ are called cusp forms. For $w \ge 4$ we define the Eisenstein series by

$$E_{w}(\tau) = \sum_{C,D} \det(c\tau + d)^{-w}.$$
 (C.6)

The summation is over all inequivalent bottom rows $\begin{pmatrix} C & D \end{pmatrix}$ of elements of Γ_2 . A classical theorem by Igusa states that the space of Siegel modular forms of genus two has a representation as

$$M = \mathbb{C}[E_4, E_6, \chi_{10}, \chi_{12}, \chi_{35}] / \{\chi_{35}^2 = R\},$$
(C.7)

Here E_4 and E_6 denote the Eisenstein series of degree four and six, while the cusp forms are given as follows

$$\chi_{10} = -\frac{43867}{2^{12}3^5 5^2 7 \cdot 53} (E_4 E_6 - E_{10}), \qquad (C.8)$$

$$\chi_{12} = -\frac{131 \cdot 593}{2^{13}3^7 5^3 7^2 337} \left(3^2 7^2 E_4^3 + 2 \cdot 5^3 E_6^2 - 691 E_{12} \right). \tag{C.9}$$

Note that χ_{10} as defined by Igusa differs from the Maas lift $\phi_{10}^{(2)}$ considered in section 9 by a factor of four. *R* denotes a polynomial in E_4 , E_6 , χ_{10} , χ_{12} whose explicit form can be found in [Igu67].

Fourier Expansion of Eisenstein series of genus two. In this subsection we discuss how to compute the Fourier coefficients of Eisenstein series. We start by recalling some mathematics terminology.

• Let *d* be a square-free integer and consider the field extension $K = \mathbb{Q}[\sqrt{d}]$ of the rational numbers. The discriminant of *K* is given as

$$\Delta_{K} = \begin{cases} d & \text{if } d \equiv 1 \mod 4 \\ 4d & \text{if } d \equiv 2, 3 \mod 4 \end{cases}$$
(C.10)

• The Möbius function $\mu : \mathbb{N} \longrightarrow \{-1, 0, 1\}$ is defined as follows

 $\mu(n) = \begin{cases} 1 \text{ if } n \text{ is a square-free, positive integer with an even number of prime factors} \\ -1 \text{ if } n \text{ is a square-free, positive integer with an odd number of prime factors} \\ 0 \text{ if } n \text{ has a squared prime factor} \end{cases}$

(C.11)

• The divisor function $\sigma_k(n)$ with $k \in \mathbb{C}$ is defined as

$$\sigma_k(n) = \sum_{d|n} d^k \tag{C.12}$$

- A Dirichlet character is a function $\chi : \mathbb{Z} \longrightarrow \mathbb{C}$ with the following properties
 - 1. There is a $k \in \mathbb{Z}$, such that $\chi(n) = \chi(n+k)$ for all n

2.
$$\chi(n) \begin{cases} = 0 \text{ if } \gcd(n, k) > 1 \\ \neq 0 \text{ if } \gcd(n, k) = 1 \end{cases}$$

$$(\neq 0 \text{ if } gcu(n, k))$$

3. $\chi(mn) = \chi(m)\chi(n)$

In particular, a Dirichlet character χ is a group homomorphism $(\mathbb{Z}/(k\mathbb{Z}))^* \longrightarrow \mathbb{C}^*$. Vice versa, any character of the unit group of $\mathbb{Z}/(k\mathbb{Z})$ extends to a Dirichlet character by setting $\chi(n) = 0$ for $n \notin (\mathbb{Z}/(k\mathbb{Z}))^*$

• A Dirichlet L-function associated to a Dirichlet character is given by

$$L(s,\chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}, \quad \text{Re}(s) > 1$$
 (C.13)

• Given a prime *p*, the Kroneckersymbol $\left(\frac{a}{p}\right)$ is for $a \in \mathbb{Z}$ defined as

$$\left(\frac{a}{p}\right) = \begin{cases} 1, \text{ if } a \text{ is a quadratic rest modulo } n \\ -1, \text{ if } a \text{ is not a quadratic rest modulo } n \\ 0, \text{ if } a \equiv 0 \text{ modulo } n \end{cases}$$
(C.14)

For general $n \in \mathbb{N}$ with prime factorization $n = p_1^{\nu_1} \dots p_k^{\nu_k}$ one puts $\left(\frac{a}{p_1}\right) = \left(\frac{a}{p_1}\right)^{\nu_1} \dots \left(\frac{a}{p_k}\right)^{\nu_k}$. Note that $\left(\frac{\cdot}{n}\right)$ is a Dirichlet character modulo n.

Theorem. (E. g. [EZ85]) Let E_w be an Eisenstein series of weight w, $T = \begin{pmatrix} a & b/2 \\ b/2 & c \end{pmatrix} \in \frac{1}{2}\mathbb{Z}^{2\times 2}$ be positive semi-definite. Denote $D = b^2 - 4ac \le 0$ and let D_0 be the discriminant of $\mathbb{Q}(\sqrt{D})$. Then the Fourier coefficient a(T) is one, if a = b = c = 0 and

$$\frac{-2w}{B_w} \sum_{d|gcd(a,b,c)} d^{w-1} \alpha(D/d^2)$$
(C.15)

otherwise. Here B_k denotes the kth Bernoulli number, α is defined by $\alpha(0) = 1$ and

$$\alpha(D) = \frac{1}{\zeta(3-2w)} C(w-1, D).$$
(C.16)

Here the Cohen function C is defined by

$$C(s-1,D) = L_{D_0}(2-s)\sum_{d|f} \mu(d) \left(\frac{D_0}{d}\right) d^{s-2} \sigma_{2s-3}(f/d), \qquad D = D_0 f^2.$$
(C.17)

In this expression, ζ denotes the Dedekind zet a-function and L_{D_0} is the Dirichlet L-series associated to the character $(\frac{\cdot}{D_0})$.

In the following we list the explicit Fourier expansions of the above generators

$$\begin{split} E_4(q_1,q_2,r) &= 1+240q_1+240q_2+2160q_1^2+30240q_1q_2+2160q_2^2+240\frac{q_1q_2}{r^2}+13440\frac{q_1q_2}{r}\\ &+13440q_1q_2r+240q_1q_2r^2+6720q_1^3+181440q_1^2+181440q_1q_2^2+6720q_2^3\\ &+30240\frac{q_1^2q_2}{r^2}+30240\frac{q_1q_2^2}{r^2}+138240\frac{q_1^2q_2}{r}+138240\frac{q_1q_2}{r}+138240q_1q_2r^2\\ &+138240q_1q_2^2r+30240q_1^2q_2r^2+302240q_1q_2r^2\\ &E_6(q_1,q_2,r) &= 1-504q_1-504q_2-16632q_1^2+166320q_1q_2-16632q_2^2-504\frac{q_1q_2}{r^2}+44352\frac{q_1q_2}{r}\\ &+44352q_1q_2r-504q_1q_2r^2-122976q_1^3+3792096q_1^2q_2+3792096q_1q_2^2-122976q_2^3\\ &+166320\frac{q_1^2q_2}{r^2}+166320q_1^2q_2r^2+166320q_1q_2r^2\\ &+2128896q_1q_2^2r+166320q_1^2q_2r^2+166320q_1q_2r^2\\ &+2128896q_1q_2^2r+166320q_1^2q_2r^2+66320q_1q_2^2r^2\\ &\chi_{10}(q_1,q_2,r) &= \frac{1}{2}q_1q_2-9q_1^2q_2-9q_1q_2^2+\frac{q_1q_2}{q_1q_2}+\frac{q_1q_2}{q_1q_2}+\frac{q_1q_2}{q_1q_2}+\frac{q_1q_2}{q_1q_2}+\frac{q_1q_2}{r^2}-\frac{q_1q_2}{q_1r}+\frac{4q_1q_2}{r}-\frac{22}{q_1q_2}-\frac{1}{4}q_1q_2r\\ &+4q_1^2q_2r+4q_1q_2^2r+\frac{1}{2}q_1^2q_2r^2+\frac{5}{6}q_1^2q_2r^2+\frac{5}{6}q_1q_2r^2\\ &\chi_{12}(q_1,q_2,r) &= \frac{5}{6}q_1q_2-11q_1^2q_2-11q_1q_2^2+\frac{5}{6}q_1^2q_2r+\frac{5}{6}q_1^2q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2^2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{22}{3}q_1^2q_2r-\frac{22}{3}q_1q_2r-\frac{22}{3}q_1q_2r+\frac{5}{6}q_1^2q_2r^2+\frac{5}{6}q_1^2q_2r^2+\frac{5}{6}q_1q_2^2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{22}{3}q_1^2q_2r-\frac{22}{3}q_1q_2r+\frac{5}{2}q_1q_2r+\frac{5}{6}q_1^2q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2^2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{22}{3}q_1^2q_2r-\frac{22}{3}q_1q_2r-\frac{22}{3}q_1q_2r+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{22}{3}q_1^2q_2r-\frac{22}{3}q_1q_2r-\frac{22}{3}q_1q_2r+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{22}{3}q_1q_2r-\frac{22}{3}q_1q_2r-\frac{22}{3}q_1q_2r-\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{23}{3}q_1q_2r-\frac{23}{3}q_1q_2r-\frac{23}{3}q_1q_2r-\frac{23}{2}q_1q_2r-\frac{5}{2}q_1q_2r-\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2+\frac{5}{6}q_1q_2r^2\\ &+\frac{1}{12}q_1q_2r-\frac{23}{q_1q_2}+\frac{4}{3}q_1q_2r-\frac{23}{2}q_1q_2r-\frac{5}{2}q_1q_2r-\frac{5}{2}q_1q_2r-\frac{5}{6}q_1q_2r-\frac$$

D | Refined GV invariants for $\mathbb{C}^3/\mathbb{Z}_5$

	d_1	0	1	2	3	4	5	6	7	8	9	10
d_2												
0			3	-6	27	-192	1695	-17064	188454	-2228160	27748899	-360012150
1		-2	4	-10	64	-572	6076	-71740	909760	-12146622	168604540	-2412582616
2		0	3	-12	91	-980	12259	-166720	2394779	-35737460	548460000	-8599208436
3		0	5	-12	108	-1332	18912	-289440	4632120	-76306398	1282295808	-21860004816
4		0	7	-24	150	-1808	26983	-443394	7665776	-136440800	2471539911	-45269668626
5		0	9	-56	294	-2982	42005	-689520	12254816	-227540162	4331108122	-83626566000
6		0	11	-140	675	-5992	76608	-1192644	20764870	-386343036	7482057534	-148451081248
7		0	13	-324	1738	-13550	158814	-2322056	38750866	-703362386	13488597425	-268229722780
8		0	15	-686	4732	-33552	359898	-4954570	79050699	-1387505216	25992283043	-509660731360
9		0	17	-1328	12960	-88746	874588	-11327904	172924796	-2932945300	53475853968	-1026989105240
10		0	19	-2394	34357	-245520	2245125	-27363700	399648535	-6552913216	116272411761	-2183695179370

Table 2: GV invariants at genus (0,0)

		d_1	0	1	2	3	4	5	6	7	8	9	10
6	l_2												
	0			0	0	-10	231	-4452	80948	-1438086	25301295	-443384578	7760515332
	1		0	0	0	-18	576	-13968	305244	-6329628	127275876	-2508961104	48786866820
	2		0	0	0	-24	896	-25636	650852	-15418734	349139480	-7658224250	164057415432
	3		0	0	0	-28	1152	-37032	1056780	-27964428	701652588	-16919331450	395589996732
	4		0	0	0	-30	1407	-48966	1515448	-43561508	1185905652	-30938142568	779767395056
	5		0	0	9	-66	2061	-68908	2174157	-65084016	1863846681	-51392001108	1371429906120
	6		0	0	68	-280	4500	-119124	3489856	-102704154	2969225052	-83868926462	2312642950872
1	7		0	0	300	-1410	13413	-261576	6617379	-181806634	5100476481	-142921977728	3965155214627
	8		0	0	988	-6760	48183	-695664	14702120	-365286402	9681953781	-262834540958	7179092104476
	9		0	0	2698	-29360	187770	-2131298	37329793	-822940764	20261551070	-525109967206	13926533807541
1	10		0	0	6444	-113186	751019	-7150716	105558998	-2048434992	46290047925	-1133487473126	28930878789904

Table 3: GV invariants at genus (1,0)

	d_1	0	1	2	3	4	5	6	7	8	9	10
d_2												
0			-4	35	-386	5161	-74368	1117672	-17319898	274571953	-4428464750	72399258543
1		1	-4	45	-750	13174	-235148	4227874	-76326692	1381543835	-25051915148	454880537144
2		0	-4	46	-900	19554	-420472	8861756	-183661746	3754800426	-75910228250	1520587166090
3		0	-20	46	-944	23394	-578872	13923300	-325336476	7413730499	-165426093894	3626709524196
4		0	-56	164	-1370	29417	-750734	19452681	-494871808	12281325148	-297499294404	7050658845633
5		0	-120	643	-3602	51118	-1121972	28252291	-735181136	19077844392	-487791469432	12245596115880
6		0	-220	2522	-11456	121392	-2132580	47798426	-1186598986	30575571450	-794192203582	20520959813816
7		0	-364	8526	-41314	340762	-4920912	95935665	-2184901598	53716745464	-1367832764572	35294081485451
8		0	-560	24835	-154752	1078545	-12985696	220762885	-4561068642	105019097003	-2565242151798	64643620048364
9		0	-816	63278	-566320	3665611	-37876670	564821327	-10560913916	226102977980	-5244562425062	127513019275063
10		0 ·	-1140	145113	-1945582	12930429	-118326784	1568446692	-26605876980	528222136235	-11577646359018	269921674605741

Table 4: GV invariants at genus (0,1)

	dı	0	1	2	3	4	5	6	7	8	9	10
da		-	-	-		•	0	0	•	0	0	10
đ			0	0	0	-102	5430	-194022	5784837	-155322234	3894455457	-93050366010
1		0	0	0	0	-216	15012	-659088	23286132	-722619936	20558391372	-549482611284
2		0	0	0	0	-312	25402	-1299948	52772850	-1855049556	59041090140	-1746828786552
3		0	0	0	0	-390	34912	-1994490	90371068	-3526024576	123699660448	-4006116067574
4		0	0	0	0	-450	43803	-2714134	133676408	-5667402346	215531532522	-7541030566234
5		0	0	0	0	-564	56340	-3636050	188364003	-8450098622	341039465436	-12674180528160
6		0	0	-12	33	-1302	89949	-5397344	277712616	-12685944558	527916004781	-20372727166960
7		0	0	-116	471	-5616	205699	-9811006	463444751	-20541312540	851408769126	-33210201574872
8		0	0	-628	4498	-31902	647792	-22397190	906948972	-37292448660	1490395858323	-57261471814196
9		0	0	-2488	32985	-194232	2563379	-62839000	2080200874	-76550883550	2874492150421	-106583214770442
10		0	0	-8036	190910	-1170468	11563235	-209199272	5533285795	-177007441804	6108659068296	-215066543042220

Table 5: GV invariants at genus (2,0)

	d	1^{0}	1	2	3	4	5	6	7	8	9	10
d2												
0			0	0	165	-7448	239030	-6574236	165498745	-3934550232	89865837855	-1992688534060
1		0	0	0	249	-16032	663816	-22351692	665886174	-18286464756	473803348152	-11751841988652
2		0	0	0	284	-21788	1085422	-43131672	1486268819	-46417626916	1348910501365	-37100685165104
3		0	0	0	294	-24852	1407460	-63647560	2476239454	-86461707148	2782756496277	-84053632394048
4		0	0	0	295	-27592	1688645	-83386604	3553493834	-135672322024	4756388163712	-155776855706696
5		0	0	-120	865	-40020	2221550	-110664544	4919143810	-198534576552	7396590575826	-257815753553060
6		0	0	-1484	5370	-101208	3849720	-169488448	7298405957	-296633814860	11342997342789	-409933588199632
7		0	0	-9632	39827	-381244	9203294	-321797124	12472772327	-484935911712	18313654586964	-665820277600268
8		0	0	-43732	268944	-1812560	28546696	-754793300	25028162865	-894682148776	32327818731481	-1151304488157956
9		0	0	-157232	1579910	-9323548	107088059	-2132204912	58508571346	-1866608513100	63045566722361	-2156999365698592
10		0	0	-477892	7945885	-48404096	447651280	-6998081720	157304121266	-4377953683428	135534530407883	-4387648213279700

Table 6: GV invariants at genus (1,1)

Г	(d ₁ 0	1	2	3	4	5	6	7	8	9	10
1	l_2											
	0		1	-56	1710	-45478	1108429	-25510644	565560106	-12208303534	258259176252	-5377587417300
	1	0	1	-62	2823	-101210	3123131	-87398508	2285563966	-56886814500	1363692509267	-31741367003420
	2	0	1	-62	3083	-135046	5051631	-167324604	5070030051	-143679833838	3866332989620	-99855731010794
	3	0	21	-62	3117	-150012	6401972	-242529558	8329863808	-264680950084	7904947970683	-224565445050932
	4	0	126	-370	5015	-187166	7888757	-317686030	11857695191	-411277565110	13383285674817	-412600220588284
	5	0	462	-2332	17188	-364870	12034886	-449182560	16816027403	-605858293698	20790005285269	-679898745656230
	6	0	1287	-14442	70155	-1022430	25116729	-783594714	26879295676	-941310618136	32492448213165	-1089619106809056
	7	0	3003	-72430	338223	-3391610	65165109	-1690999402	51032170076	-1653086126192	54861619097005	-1817288123911204
	8	0	6188	-293502	1718392	-13023266	195187516	-4260040764	112884832140	-3323304791332	103316703343033	-3287506711857818
	9	0	11628	-991856	8413737	-54852288	656650202	-12066214776	281352005882	-7507672419570	216602339119797	-6526511388596102
11	10	0	20349	-2908410	37685481	-242056852	2402813660	-37503651456	771051279956	-18655847273504	498243332814881	-14133822653295814

Table 7: GV invariants at genus (0,2)

_												
	d_1	0	1	2	3	4	5	6	7	8	9	10
d_2												
0			0	0	0	15	-3672	290853	-15363990	649358826	-23769907110	786400843911
1		0	0	0	0	28	-9038	895004	-56764044	2800998468	-117271264390	4368276685200
2		0	0	0	0	39	-14318	1647535	-120373298	6755811733	-317790113114	13156914112074
3		0	0	0	0	48	-18996	2412872	-196006904	12206895204	-633769608952	28779314566504
4		0	0	0	0	55	-23102	3156548	-277709398	18773083610	-1056971776506	51911426998599
5		0	0	0	0	60	-27784	4008354	-372807278	26742919550	-1601413604094	83702954155532
6		0	0	0	0	140	-40296	5522988	-516972356	38066404131	-2363089642518	128764263234726
7		0	0	15	-56	1090	-94614	9521920	-811965094	58217079594	-3617744837466	200176789418018
8		0	0	176	-1538	11205	-358426	22162654	-1538663154	100876901423	-6029485692286	329118801193692
9		0	0	1130	-21898	117201	-1881714	68569525	-3571773320	202306292870	-11205365856114	587561193934288
10		0	0	5232	-201110	1135775	-11788422	270108872	-10102578488	470971719089	-23362479231458	1149433429910848

Table 8: GV invariants at genus (3,0)

Γ		d_1	0 1	2	3	4	5	6	7	8	9	10
Γ	d_2											
1	õ		0	0	0	3646	-332500	18307188	-784000382	28746178815	-947455553070	28883283581120
1	1		0 0	0	0	6736	-817658	56229716	-2890271136	123724696980	-4664622899914	160128325772898
1	2		0 0	0	0	8591	-1240914	100812817	-6020174066	294558138571	-12516025972770	478543796677390
	3		0 0	0	0	9595	-1541556	141315405	-9507560084	520483728764	-24542285301304	1033022927270955
	4		0 0	0	0	10060	-1762130	176472063	-13005635874	779144061743	-40074259816370	1832137941830936
	5		0 0	0	0	11942	-2095360	218310699	-16998820726	1083695909005	-59481598311246	2903077344820940
	6		0 0	286	-704	31327	-3252496	304703564	-23423999308	1522934680590	-86512345468570	4402967053374246
	7		0 0	4101	-14832	173458	-7932290	542315279	-37210408886	2328821117728	-131676086618278	6787311660887958
1	8		0 0	30706	-201910	1335968	-29094066	1282419545	-71414914086	4056835940421	-219450368091250	11121462357918974
1	9		0 0	160494	-2029890	10923049	-143171810	3973169453	-167074649216	8171560102165	-408274358839182	19828985332878797
L	10		0 0	660490	-15457318	86650077	-819422430	15457087268	-475302472716	19096999811219	-851813505320610	38743456874318870

Table 9: GV invariants at genus (2,1)

	$d_1 = 0$	1	2	3	4	5	6	7	8	9	10
d_2											
0		0	0	-792	74640	-4150308	177499683	-6464352018	211097449143	-6368926838778	180940547430291
1	0	0	0	-1044	141232	-10305512	547583663	-23880243270	909350539113	-31362711810712	1002969680533521
2	0	0	0	-1100	173984	-15317486	968113645	-49239570774	2148320040446	-83640362553570	2982529054781422
3	0	0	0	-1106	185762	-18382200	1322846349	-76293814476	3741031288415	-162136840377466	6379219588997229
4	0	0	0	-1106	200166	-20872804	1625640316	-102612370488	5515121845123	-261276623451526	11187678438452042
5	0	0	462	-4208	308575	-27242952	2075005637	-134967532118	7642082210213	-384898656237490	17575381337394925
6	0	0	9757	-34468	914095	-50492366	3204053300	-195554839650	10983999127237	-564354092549442	26674496909199694
7	0	0	95115	-363608	4096989	-134471610	6429657554	-338709281142	17714458393384	-885389528147694	41780584233048109
8	0	0	602395	-3442542	24297636	-473005618	16339619628	-709931460036	33124639634829	-1550574847245638	70711217658486539
9	0	0	2872686	-27531174	159003934	-2069571994	50880368591	-1767456288994	71569396226022	-3060796330213490	131835980366899491
10	0	0	11166111	-182163160	1053629099	-10307057780	187616361586	-5139288077960	176685724431350	-6773022816685924	271069267170147567

Table 10: GV invariants at genus (1,2)

	d_1	0	1	2	3	4	5	6	7	8	9	10
d_2												
σ			0	36	-3552	199129	-8477200	305202621	-9831079874	292451827680	-8193427901442	219053227327973
1		0	0	37	-5132	391895	-21456544	951816717	-36559985102	1265167969027	-40459733831784	1216486396522567
2		0	0	37	-5332	482505	-31872458	1680167027	-75235065398	2982734988243	-107682958186546	3610552800188045
3	l l	0	-8	37	-5342	512892	-37960032	2277288722	-115697208032	5159487585801	-207532502054646	7683566995242455
4	i i	0	-120	403	-9694	650547	-45492860	2845195737	-155997880696	7584685164224	-332849815705118	13405043007333180
5	l l	0	-792	4032	-44232	1456988	-72718424	4003499489	-214623624454	10718340308348	-493969134826666	21087866983228178
6		0	-3432	40032	-227316	4852264	-169805898	7373281571	-346895883190	16401686036909	-749687050494070	32594365559442838
7		0	-11440	303420	-1430960	18827088	-497163382	17348476209	-692265842390	29328965878560	-1259736657853762	53323814192441639
8		0	-31824	1731540	-9791434	86102063	-1681294978	48005169030	-1640618797164	61576392616975	-2420210604499130	96594746682174718
9		0	-77520	7822110	-64355702	441639078	-6453703762	150166863631	-4417828314310	147334819728793	-5270128198705030	195620751168796920
10		0	-170544	29481458	-378136780	2411472333	-27268753568	519116905858	-13161855037508	390992623491806	-12736589773157528	438155527249781100

Table 11: GV invariants at genus (0,3)

$\textbf{E} \mid \textbf{Refined GV} \text{ invariants for } \mathbb{C}^3/\mathbb{Z}_6$

							GV invari	ants a	at gei	nus (0	,0):						
0 1 2 3 4 5 6 7 2 0 1 2 3 4 5 6 6 7 4 0 1 2 3 4 5 6 7 4 0 1 2 3 4 5 6 7 7 7	$\begin{array}{c} 0 \\ -2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ $	$\begin{array}{c} 1 \\ 0 \\ -2 \\ -4 \\ -6 \\ -8 \\ -10 \\ -12 \\ -14 \\ 1 \\ 0 \\ 0 \\ -6 \\ -12 \\ -18 \\ -24 \\ -30 \\ 0 \\ -6 \\ -10 \\ -36 \\ 1 \\ 0 \\ -6 \\ -10 \\ -20 \\ -30 \\ -40 \\ -50 \\ \end{array}$	$\begin{array}{c} 2\\ 0\\ 0\\ -32\\ -310\\ -388\\ -644\\ -2\\ 0\\ 0\\ -288\\ -644\\ -2\\ 0\\ 0\\ -1280\\ -3204\\ -3204\\ -3204\\ -3204\\ -3204\\ -3206\\ -296\\ -1280\\ -3204\\ -3204\\ -3206\\ -2016\\ -5580\\ \end{array}$	$\begin{array}{c} 3\\ 0\\ 0\\ -8\\ -8\\ -110\\ -756\\ -3356\\ -3356\\ 0\\ 0\\ 0\\ -3350\\ -22176\\ -3350\\ -22176\\ -3350\\ -22576\\ -6368\\ -6368\\ -42400 \end{array}$	4 0 0 0 -288 -3556 4 0 0 0 0 -22176 -22176 -22176 -22176 -30 -2016 -42400	$\begin{array}{c} 5\\ 0\\ 0\\ 0\\ 0\\ -12\\ -644\\ 0\\ 0\\ -30\\ 0\\ 0\\ -304\\ -304\\ -304\\ -304\\ -5580\\ \end{array}$	$\begin{array}{c} 6 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	$\begin{array}{c} 7 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	1 0 1 2 3 4 5 6 7 3 4 5 6 7 5 0 1 2 3 4 5 6 7 5 0 1 2 3 4 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 6 7 5 7 5 6 7 5 7 5 6 7 5 7 5 7 5 6 7 5 6 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 5 6 7 7 7 7 5 7 7 7 7 7 7 7 7	$\begin{array}{c} 0 \\ -2 \\ -2 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ $	$\begin{array}{c} 1 \\ 0 \\ -2 \\ -6 \\ -10 \\ -14 \\ -18 \\ -22 \\ -26 \\ 1 \\ 0 \\ 0 \\ -4 \\ -12 \\ -20 \\ -28 \\ -36 \\ -44 \\ -1 \\ -28 \\ -36 \\ -44 \\ -1 \\ -38 \\ -30 \\ -42 \\ -54 \\ \end{array}$	$\begin{array}{c} 2\\ 0\\ 0\\ -10\\ -70\\ -270\\ -770\\ -1820\\ 0\\ 0\\ -12\\ -110\\ -518\\ -1710\\ -4510\\ -518\\ -1710\\ -4510\\ 0\\ 0\\ 0\\ -14\\ -126\\ -630\\ -2254\\ -6426\end{array}$	$\begin{array}{c} 3\\ 0\\ 0\\ -14\\ -270\\ -2204\\ -11544\\ 3\\ 0\\ 0\\ 0\\ -518\\ -5292\\ -3300\\ 0\\ 0\\ -518\\ -5292\\ -3300\\ 0\\ 0\\ 0\\ -630\\ -7308\\ -50652\\ \end{array}$	$\begin{array}{c} 4\\ 0\\ 0\\ 0\\ 0\\ -18\\ -770\\ -11544\\ \hline 4\\ 0\\ 0\\ 0\\ 0\\ -28\\ -1710\\ -3300\\ -28\\ -1710\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ -330\\ -2254\\ -50652\\ \end{array}$	$\begin{array}{c} 5\\ 0\\ 0\\ 0\\ -36\\ -36\\ -36\\ -4510\\ 0\\ 0\\ -36\\ -4510\\ 0\\ 0\\ -36\\ -42\\ -6426\\ -42\\ -6426\\ \end{array}$	$\begin{array}{c} 6 & 7 \\ \hline 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 &$	
							GV invari	ants a	nt ger	nus (0	,1):						_
0 0 1 2 3 4 5 6 7 2 0 1 2 3 4 5 6 7 7 4 1 2 3 4 5 6 7 7 4 5 6 7 7 7 1 2 3 4 5 6 7 7 7 7 1 2 3 4 5 6 7 7 7 7 7 7 7 7 7 7 7 7 7	0 0	$ \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	2 0 0 0 0 0 0 0 0	$\begin{array}{c} 3\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\$	4 0		6 0	$\begin{array}{c} 7 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	$\begin{array}{c} 1 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 7 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 7 \\ 1 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 7 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1$		$ \begin{array}{c} 1 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0$	$\begin{array}{c} 2\\ 2\\ 0\\ 0\\ 0\\ 16\\ 144\\ 704\\ 2496\\ 2496\\ 2\\ 2\\ 0\\ 0\\ 0\\ 248\\ 35368\\ 2\\ 2\\ 2\\ 248\\ 5368\\ 2\\ 0\\ 0\\ 0\\ 0\\ 24\\ 288\\ 1736\\ 7272 \end{array}$	$\begin{array}{c} 3\\ 0\\ 0\\ 0\\ 0\\ 144\\ 2608\\ 22672\\ 3\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\$	$\begin{array}{c} 4\\ -4\\ 0\\ 0\\ 0\\ 0\\ -2672\\ -4\\ -22672\\ -4\\ -0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0\\ 0$	$-\frac{5}{0}$ 0 0 0 2496 $-\frac{5}{5}$ 0 0 0 0 0 0 0 0	$\begin{array}{c} 6 & 7 \\ \hline 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 &$	
							GV invari	ants a	nt ger	nus (1	,0):						
0 0 1 2 3 4 5 6 7 2 0 1 2 3 4 5 6 7 4 0 1 2 3 3 4 5 6 7 7 6 7 7	0 1 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{array}{c} 1 \\ 0 \\ 1 \\ 10 \\ 35 \\ 84 \\ 165 \\ 286 \\ 455 \\ \hline 1 \\ 0 \\ 0 \\ 11 \\ 46 \\ 129 \\ 284 \\ 535 \\ 906 \\ 129 \\ 284 \\ 535 \\ 906 \\ 1 \\ 0 \\ 0 \\ 35 \\ 45 \\ 130 \\ 295 \\ 580 \\ 1025 \\ \end{array}$	2 0 0 35 359 1987 7620 23414 2 0 0 0 461 7310 5380 0 0 461 7311 5380 0 0 0 0 45 731 6036 31835 126302	3 0 0 0 1987 20554 134882 3 0 0 0 0 5380 78341 652272 652272 652272 0 0 0 0 0 0 0 0 0 0 0 0 0 0 130 0 6036 103132 1000100	4 0 0 0 0 165 7620 134882 4 0 0 0 0 0 0 0 652272 652272 652274 0 0 <	5 0 0 286 23414 5 0 0 0 0 0 5355 90234 90234 90234 90234 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{array}{c} 6 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	$\begin{array}{c} 7 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \\$	$\begin{array}{c} 1 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 3 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 5 \\ 0 \\ 1 \\ 2 \\ 3 \\ 4 \\ 5 \\ 6 \\ 7 \\ 1 \\ 0 \\ 1 \\ 2 \\ 3 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1 \\ 1$	0 1 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{array}{c} 1 \\ 0 \\ 1 \\ 11 \\ 45 \\ 119 \\ 249 \\ 451 \\ 741 \\ \hline 0 \\ 0 \\ 0 \\ 10 \\ 46 \\ 130 \\ 294 \\ 130 \\ 294 \\ 570 \\ 990 \\ \hline 1 \\ 0 \\ 0 \\ 0 \\ 84 \\ 119 \\ 129 \\ 295 \\ 581 \\ 1035 \\ \end{array}$	2 0 0 45 635 78110 2 0 0 0 461 7511 2 9 0 0 0 461 75891 29847 113047 2 0 0 1199 1023 6627 33551	3 0 0 19 19 51555 384852 3 0 0 0 0 0 3 3 3 0 0 0 0 0 0 0 0 0 0	4 0 0 0 249 17545 384852 4 0 0 0 0 0 0 0 0 0 0 0 0 0	5 0 0 0 0 58110 58110 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0	$\begin{array}{c} 6 & 7 \\ \hline 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 &$	

Table 12: Refined GV invariants for $\mathbb{C}^3/\mathbb{Z}_6$. Degree d_1 goes from left to right, d_3 from top to bottom and d_2 is indexed in the upper-left corner in each table.

(<i>n</i> , <i>r</i> , <i>m</i>)	$\begin{array}{c} (0,0,0)\\ (1,1,1)\\ (1,1,3)\\ (2,2,2)\\ (2,2,3)\\ (2,2,4)\\ (3,3,3)\\ (3,3,4)\\ (4,4,4) \end{array}$	$\begin{array}{c} (0,0,1) \\ (1,0,1) \\ (1,0,3) \\ (2,1,2) \\ (2,1,3) \\ (2,1,4) \\ (3,2,3) \\ (3,2,4) \\ (4,3,4) \end{array}$	$\begin{array}{c} (0,0,2)\\ (1,1,2)\\ (1,1,4)\\ (2,0,2)\\ (2,0,3)\\ (2,0,4)\\ (3,1,3)\\ (3,1,4)\\ (4,2,4) \end{array}$	(0, 0, 3) (1, 0, 2) (1, 0, 4) (3, 0, 3) (3, 0, 4) (4, 1, 4)	(0,0,4) (4,0,4)
E ₄ ⁽²⁾	1 13440 362880 604800 1814400 5114880 3642240 10644480 20818560	240 30240 497280 9967680 2903040 5986780 5987520 13426560 24192000	2160 138240 967680 1239840 2782080 7439040 6531840 17418240 35804160	6720 181440 997920 8467200 15980160 34974720	17520
$E_{6}^{(2)}$	1 44352 15422400 24881472 234311616 1126185984 883802304 4864527360 12809611584	-504 166320 23462208 65995776 453458488 1724405760 1945345248 8158449600 22751511552	-16632 2128896 65995776 90644400 530228160 2066692320 2818924416 11304437760 34911765504	-122976 3792096 85322160 3327730560 12013404480 42077629440	-532728 46585733040
X10	0 1 99 240 -1800 4352 15399 -6864 135424	0 -2 -272 -240 2736 -6816 -19008 -26928 -22000	0 -16 -240 32 -1464 -576 27270 44064 65280	0 36 1056 -43920 12544 -36432	0-279040
E ⁽²⁾ 12	53678953 22266840960 18728449326561920 46765376055216000 9975044773603728000 341726484448551121920 232922965682969976960 11078677991333226193920 98074242084224505939840	5089790160 456798756960 46719773564929920 486707206711864320 43296648289119912960 994500257327275345920 1387314320747141666880 3929644083457709619840 409369370741219790720000	10428980037840 162868282536960 486707206711864320 958912407409188960 67657429630096225920 1388669119242478032960 3553058504433847368960 78584973005595527976960 1021197866857079343344640	901646147263680 661522702800960 957976883554934880 4778384234988952224000 97978466594063423571840 1703669378274682309370880	21358556207286480 2010986989911829611208800
$E^{(2)}_{\Delta}$	0 158976 44644753152 -3910809600 -210642882048 -10065668456448 2356270891776 18898645911552 -245453236125696	12096 2146176 111108644352 -25315338240 -097120692224 -28301296693248 8269812764928 108975837336704 -584737669785600	-290304 377726976 1155817838592 -54993613824 -1622158990848 -39925363636224 24778417466880 198828131659776 -1744357612732416	3048192 1558123776 2276348012928 31936386585600 245736461090304 -2891211375900672	-17805312 -3257305227141120
φ ⁽²⁾ 12	0 1 1275 2784 13080 -64768 48303 -806520 3392512	0 10 736 -8040 -14136 389520 389520 38016 938400 2311640	0 -88 -8040 17600 -54120 -232320 -236410 -256410 -1227600 -5917440	0 -132 -2880 1073520 -2309120 6141960	0

F | Fourier coefficients of Siegel modular forms

Table 13: Fourier coefficients $c(f, \binom{n}{r/2} m^{r/2})$ of some genus 2 Siegel modular forms in [LMF15].

(<i>n</i> , <i>r</i> , <i>m</i>)	$u_1^2 v_1^2$	$u_1 u_2 v_1^2$	$u_2^2 v_1^2$	$u_1^2 v_1 v_2$	$u_1 u_2 v_1 v_2$	$u_2^2 v_1 v_2$	$u_1^2 v_2^2$	$u_1 u_2 v_2^2$	$u_2^2 v_2^2$
(1, 1, 1)	$-\frac{1}{100}$	$-\frac{1}{100}$	$-\frac{1}{400}$	$-\frac{1}{100}$	$-\frac{1}{40}$	$-\frac{1}{100}$	$-\frac{1}{400}$	$-\frac{1}{100}$	$-\frac{1}{100}$
(1,0,1)	$\frac{1}{50}$	0	$-\frac{1}{200}$	0	$-\frac{17}{100}$	0	$-\frac{1}{200}$	0	$\frac{1}{50}$
(1,1,2)	$\frac{4}{25}$	$\frac{4}{25}$	3	$\frac{4}{25}$	<u>39</u> 25	$\frac{19}{25}$	3	19 25	$-\frac{28}{25}$
(1,0,2)	$-\frac{9}{25}$	0	100	0	$\frac{117}{50}$	0	100	0	<u>63</u> 25
(1, 1, 3)	$-\frac{\overline{99}}{100}$	$-\frac{99}{100}$	$-\frac{243}{400}$	$-\frac{99}{100}$	$-\frac{4311}{200}$	$-\frac{1089}{100}$	$-\frac{243}{400}$	$-\frac{1089}{100}$	$\frac{\overline{693}}{100}$
(1, 0, 3)	$\frac{68}{25}$	0	$-\frac{18}{25}$	0	$-\frac{348}{-25}$	0	$-\frac{18}{25}$	0	$-\frac{576}{-25}$
(1, 1, 4)	$\frac{12}{5}$	$\frac{12}{5}$	$\frac{33}{10}$	$\frac{12}{5}$	<u>669</u> 5	69	$\frac{33}{10}$	69	$\frac{192}{5}$
(1, 0, 4)	$-\frac{264}{25}$	0	$\frac{84}{25}$	0	$\frac{1416}{25}$	0	$\frac{84}{25}$	0	$-\frac{264}{25}$
(2, 2, 2)	$\frac{84}{5}$	<u>84</u> 5	$\frac{384}{25}$	<u>84</u> 5	<u>492</u> 25	<u>84</u> 5	$\frac{384}{25}$	<u>84</u> 5	$\frac{84}{5}$
(2, 1, 2)	$-\frac{84}{5}$	$-\frac{903}{25}$	$-\frac{2541}{50}$	$-\frac{903}{25}$	$-\frac{2067}{25}$	$-\frac{903}{25}$	$-\frac{2541}{50}$	$-\frac{903}{25}$	$-\frac{84}{5}$
(2, 0, 2)	<u>56</u> 25	0	$\frac{344}{5}$	0	<u>56</u> 5	0	<u>344</u> 5	0	25
(2, 2, 3)	-126	-126	$\frac{3573}{50}$	-126	$\frac{2241}{252}$	$\frac{4482}{25}$	$\frac{3573}{50}$	$\frac{4482}{25}$	2466 25
(2, 1, 3)	4788 25	25	<u>2331</u> 50	25	25	$\frac{324}{25}$	<u>2331</u> 50_0	324 25	$-\frac{10116}{25}$
(2, 0, 3)	$-\frac{2562}{25}$	0	$-\frac{1179}{10}$	0	1701 50704	0	$-\frac{1179}{10}$	0	18738
(2, 2, 4)	<u>7616</u> 25	$\frac{7616}{25}$	$-\frac{13248}{25}$	$\frac{7616}{25}$	$-\frac{28704}{25}$	$-\frac{31408}{25}$	$-\frac{13248}{25}$	$-\frac{31408}{25}$	$\frac{14272}{25}$
(2, 1, 4)	$-\frac{11928}{25}$	$-\frac{35994}{25}$	25010	$-\frac{35994}{25}$	$-\frac{21282}{25}$	24966	25010	24966	25070
(2,0,4)	$-\frac{1008}{25}$	0	$-\frac{42912}{25}$	0	$-\frac{84528}{25}$	0	$-\frac{42912}{5025}$	0	$-\frac{68976}{25}$
(3, 3, 3)	$\frac{3969}{100}$	100	$\frac{36241}{400}$	100	$-\frac{32427}{200}$	<u>3969</u> 100	$\frac{36241}{400}$	$\frac{3969}{100}$	$\frac{3969}{100}$
(3, 2, 3)	25 25 115587	- 3304 79891	$-\frac{10032}{12563}$	- 3304 72891	- <u>80784</u> 51597	- 3304 72581	$-\frac{10032}{17563}$	- 3304 72881	25 25 115587
(3, 1, 3)	- <u>115567</u> 50	$-\frac{72501}{50}$	$-\frac{12003}{40}$	$-\frac{72561}{50}$	20	$-\frac{72561}{50}$	$-\frac{12003}{40}$	$-\frac{72501}{50}$	- <u>115587</u> 50
(3,0,3)	500772	0	<u>50101</u>	0	_ <u>55000</u>	0	<u>50101</u>	0	27456
(3, 3, 4)	200772 125136	278784	70794	<u>200112</u> 278784	25 68724	<u>132557</u> 6336	70794	<u>132557</u> 6336	25308
(3,2,4)	125150 19656	71786	25 72009	71786	- 00724 25 748386	$\frac{0330}{37314}$	72009	2514	- <u>25</u> 176256
(3, 1, 4)	$\frac{13030}{25}$	25	25 32112	25	44736	25	25 32112	25	- 1702.50 488384
(3,0,4)	3136	0 3136	210464	0 3136	430336	0 3136	210464	0 3136	25
(4,4,4)	$-\frac{3130}{25}$	$-\frac{5130}{125041}$	25 60217	$-\frac{5130}{122941}$	- 203307	$-\frac{3130}{125041}$	25 60217	$-\frac{5130}{125941}$	$-\frac{5130}{25}$
(4,3,4)	3520 42816	102192	- <u>55536</u>	102192	_ <u>233807</u> 221664	102192	$-\frac{55217}{10}$	102192	3520 42816
(4,2,4)	145728	1227897	230571	1227897	921789	1227897	230571	1227897	145728
(4, 1, 4)	25	25	254656	- 1227037	- 5	- 1227037	254656	- 1227037	25
(4, 0, 4)	- 134344	0	5	0	-34816	0	5	0	- 134344

Table 14: Fourier coefficients $c(\phi_{10}^{(2)} f_{\text{RS}}, \binom{n}{r/2} \binom{r/2}{m})$ of $\phi_{10}^{(2)} f_{\text{RS}}$ with notation as in Section 9.2.

(<i>n</i> , <i>r</i> , <i>m</i>)	$\begin{array}{c} (-1,1,-1) \\ (1,1,-1) \\ (-1,0,0) \\ (1,0,0) \\ (-1,-1,1) \\ (1,-1,1) \\ (-1,-2,2) \\ (1,-2,2) \end{array}$	$\begin{array}{c} (-1,2,-1) \\ (1,2,-1) \\ (-1,1,0) \\ (1,1,0) \\ (-1,0,1) \\ (1,0,1) \\ (-1,-1,2) \\ (1,-1,2) \end{array}$	$\begin{array}{c} (0,0,-1) \\ (2,-2,-1) \\ (-1,2,0) \\ (1,2,0) \\ (-1,1,1) \\ (1,1,1) \\ (-1,0,2) \\ (1,0,2) \end{array}$	$\begin{array}{c} (0,1,-1) \\ (2,-1,-1) \\ (0,-1,0) \\ (2,-2,0) \\ (-1,2,1) \\ (1,2,1) \\ (-1,1,2) \\ (1,1,2) \end{array}$	$\begin{array}{c} (0,2,-1)\\ (2,0,-1)\\ (0,1,0)\\ (2,-1,0)\\ (0,-2,1)\\ (2,-2,1)\\ (-1,2,2)\\ (1,2,2)\end{array}$	$\begin{array}{c} (1,-1,-1)\\ (2,1,-1)\\ (0,2,0)\\ (2,0,0)\\ (0,-1,1)\\ (2,-1,1)\\ (0,-2,2)\\ (2,-2,2)\end{array}$	$\begin{array}{c} (1,0,-1)\\ (2,2,-1)\\ (1,-2,0)\\ (2,1,0)\\ (0,0,1)\\ (2,0,1)\\ (0,-1,2)\\ (2,-1,2)\end{array}$	(1, -1, 0) (2, 2, 0) (0, 1, 1) (2, 1, 1) (0, 0, 2) (2, 0, 2)	(0, 2, 1)(2, 2, 1)(0, 1, 2)(2, 1, 2)	(1, -2, 1) (0, 2, 2) (2, 2, 2)
	13272-6483253534-50064	2 648 24 8376 48 -50064 72 561576	2 4 48 15600 327 130329 648 -1127472	24 72 24 1152 648 209304 3272 1598376	$\begin{array}{c} 48\\ 648\\ 600\\ 8376\\ 48\\ -50064\\ 6404\\ 2023536\end{array}$	3 3272 1152 -12800 600 561576 1152 -3859456	48 6404 48 85176 -648 -1127472 8376 18458000	600 154752 8376 1598376 -12800 -32861184	15600 2023536 85176 28698000	-648 154752 16620544

Table 15: Fourier coefficients $c(\phi_{10}^{(2)-1}, \binom{n}{r/2} m)$ of $\phi_{10}^{(2)-1}$ in the Weyl chamber described in Section 9.2.

(n r m)	$u^2 u^2$	11 11 11 11 12 11 ²	$u^2 u^2$	11 ² 111 110	111 110 111 110	11 ² 111 110	² ² ²	111 110 v ²	$u^2 u^2$
(n, r, m)	^u 1 ^v 1	1	^u 2 ^v 1	1	1	1	^u 1 ^v 2	1	^u 2 ^v 2
(0, 0, 0)	$-\frac{1}{100}$	$\frac{100}{1}$	$-\frac{1}{400}$	$\frac{100}{1}$	- 10	$\frac{100}{1}$	$-\frac{1}{400}$	$\frac{100}{1}$	$-\frac{1}{100}$
(0, 1, 0) (0, 2, 0)	0	$\frac{50}{1}$	$\frac{-100}{1}$	$\frac{50}{1}$	$\frac{-50}{11}$	$\frac{50}{1}$	$\frac{-100}{-1}$	$\frac{50}{1}$	0
(1, -1, 0)	$-\frac{4}{25}$	$\frac{50}{4}$	50 1 100	$\frac{50}{4}$	$-\frac{25}{70}$	$\frac{50}{-\frac{1}{50}}$	50 1	$\frac{50}{-\frac{1}{50}}$	0
(1,0,0)	$-\frac{8}{5}$	$-\frac{23}{25}$	100 1 50	$-\frac{23}{25}$	$\frac{11}{25}^{50}$	0	$\frac{1}{50}$	0	0
(1, 1, 0)	$-\frac{4}{25}$	$-\frac{16}{25}$	$\frac{31}{100}$	$-\frac{16}{25}$	$-\frac{7}{50}$	$\frac{1}{50}$	$\frac{31}{100}$	$\frac{1}{50}$	0
(1, 2, 0)	0	$-\frac{12}{25}$	0	$-\frac{12}{25}$	0	0	0	0	0
(2, -2, 0)	$-\frac{2}{5}$	<u>2</u> 5 ₇	0	5	$-\frac{9}{25}$	$-\frac{1}{50}$	0	$-\frac{1}{50}$	0
(2, -1, 0)	$-\frac{30}{25}$ 372	10 18	100	10 18	$-\frac{5}{50}$	- 50	<u>100</u>	- 50	0
(2, 0, 0) (2, 1, 0)	38	$\frac{-125}{107}$	59	$\frac{-128}{187}$	25 3	1	<u>59</u>	1	0
(2, 2, 0)	$-\frac{25}{5}$	$-\frac{50}{25}$	100 0	$-\frac{50}{25}$	$-\frac{50}{25}$	50 1 50	100 0	$\frac{50}{50}$	0
(0, -1, 1)	0	$-\frac{23}{50}$	$\frac{1}{100}$	$-\frac{23}{50}$	$-\frac{29}{50}$	$\frac{4}{25}$	$\frac{1}{100}$	$\frac{4}{25}$	$-\frac{4}{25}$
(0, 0, 1)	0	0	$\frac{1}{50}$	0	$\frac{11}{25}$	$-\frac{6}{25}$	$\frac{1}{50}$	$-\frac{6}{25}$	$-\frac{8}{5}$
(0, 1, 1)	0	$\frac{1}{50}$	$\frac{1}{100}$	$\frac{1}{50}$	$-\frac{7}{50}$	$-\frac{16}{25}$	$\frac{1}{100}$	$-\frac{16}{25}$	$-\frac{4}{25}$
(0, 2, 1)	0 8	0 86	0 91	0 86	0 173	$-\frac{12}{25}$	0 91	$-\frac{12}{25}$	0 8
(1, -2, 1) (1, -1, 1)	<u>-</u> 5 256	$\frac{25}{256}$	$\frac{50}{314}$	$\frac{25}{256}$	2825	$\frac{25}{256}$	$\frac{50}{314}$	$\frac{25}{256}$	2 <u>56</u>
(1, 0, 1)	$-\frac{432}{35}$	25 0	$\frac{25}{561}$	25 0	_ <u>6</u>	25 0	$\frac{25}{561}$	25 0	$\frac{25}{432}$
(1,1,1)	$\frac{256}{25}^{3}$	256	$\frac{314}{25}^{25}$	256	$\frac{28}{5}$	256	$\frac{314}{25}^{3}$	256	256 25
(1, 2, 1)	-85	$-\frac{86}{25}$	$-\frac{291}{50}$	$-\frac{86}{25}$	$-\frac{173}{25}$	$-\frac{86}{25}$	$-\frac{91}{50}$	$-\frac{86}{25}$	-85
(2, -2, 1)	$-\frac{2016}{25}$	$\frac{2016}{25}$	$-\frac{993}{257}$	$\frac{2016}{25}$	$-\frac{1758}{25}$	$\frac{864}{25}$	$-\frac{993}{25}$	$\frac{864}{25}$	$-\frac{432}{25}$
(2, -1, 1)	25 16448	$-\frac{10145}{50}$	$\frac{20407}{100}$ 1658	$-\frac{10143}{50}$	50 508	$-\frac{2052}{25}$	$\frac{20407}{100}$ 1658	- 2052	25 3232
(2, 0, 1) (2, 1, 1)	$\frac{25}{10278}$	<u>10143</u>	20487	<u>10143</u>	3903	<u>2052</u>	20487	2052	2052
(2, 1, 1)	$-\frac{2516}{2516}$	$-\frac{2016}{25}$	$-\frac{100}{993}$	$-\frac{2016}{25}$	$-\frac{50}{1758}$	$-\frac{25}{35}$	$-\frac{100}{993}$	$-\frac{25}{364}$	$-\frac{25}{432}$
(0, -2, 2)	0	$-\frac{1}{50}$	0	$-\frac{1}{50}$	$-\frac{9}{25}^{3}$	2 5	0	$\frac{2}{5}^{23}$	$-\frac{2}{5}$
(0, -1, 2)	0	$-\frac{1}{50}$	$\frac{3}{100}$	$-\frac{1}{50}$	$-\frac{3}{50}$	$\frac{7}{10}$	$\frac{3}{100}$	$\frac{7}{10}$	$-\frac{38}{25}$
(0,0,2)	0	0	<u>50</u>	0	$\frac{33}{25}_{3}$	$-\frac{16}{25}$	<u>50</u>	$-\frac{10}{25}$	$-\frac{372}{25}$
(0, 1, 2) (0, 2, 2)	0	50	100	<u>50</u> 1	- 50	$-\frac{101}{50}$ - 46	100	$-\frac{101}{50}$ - 46	$-\frac{30}{25}$
(0, 2, 2) (1, -2, 2)	$-\frac{432}{2}$	$\frac{50}{864}$	_ <u>993</u>	$\frac{50}{864}$	$\frac{25}{1758}$	$\frac{25}{2016}$	_ <u>993</u>	$\frac{25}{2016}$	$\frac{5}{2016}$
(1, -1, 2)	$\frac{2052}{25}$	$\frac{25}{2052}$	20487 100	$-\frac{25}{2052}$	3903 50	$-\frac{25}{10143}$	20487 100	$-\frac{25}{10143}$	$\frac{10278}{25}$
(1,0,2)	$-\frac{3232}{25}$	0	$-\frac{1658}{5}$	0	$-\frac{508}{25}$	0	$-\frac{1658}{5}$	0	$-\frac{16448}{25}$
(1, 1, 2)	2052 252	2052 25	$\frac{20487}{100}$	2052 25	<u>3903</u> 50	$\frac{10143}{50}$	$\frac{20487}{100}$	$\frac{10143}{5016}$	$\frac{10278}{2516}$
(1,2,2)	$-\frac{432}{25}$	$-\frac{994}{25}$ 79104	- ⁹⁹⁵ 25 78936	$-\frac{904}{25}$ 79104	$-\frac{1730}{25}$ 15888	$-\frac{2016}{25}$ 79104	- 993 25 78936	$-\frac{2016}{25}$ 79104	$-\frac{2016}{25}$
(2, -2, 2) (2, -1, 2)	<u>235482</u>	25 234789	$\frac{941511}{941511}$	25 234789	<u>30081</u>	25 234789	$\frac{25}{941511}$	25 234789	235482
(2, -1, 2) (2, 0, 2)	$\frac{25}{332568}$	- 50	$100 \\ 332979$	- 50	$\frac{10}{6978}$	- 50	$100 \\ 332979$	- <u>50</u>	$\frac{25}{332568}$
(2, 1, 2)	$\frac{25}{235482}$	<u>234789</u> 50	$\frac{941511}{100}$	<u>234789</u> 50	$\frac{30081}{10}$	<u>234789</u> 50	$\frac{941511}{100}$	<u>234789</u> 50	235482 235482
(2,2,2)	- <u>79104</u> 25	$-\frac{79104}{25}$	$-\frac{78936}{25}$	$-\frac{79104}{25}$	$-\frac{15888}{5}$	$-\frac{79104}{25}$	$-\frac{78936}{25}$	$-\frac{79104}{25}$	_ <u>79104</u> 25

Table 16: Fourier coefficients $c(f_{RS}, \binom{n r/2}{m})$ of f_{RS} in the Weyl chamber described in Section 9.2.

Bethe Center for Theoretical Physics and Hausdorff Center for Mathematics Universität Bonn D-53115 Bonn, Germany E-mail: aklemm@th.physik.uni-bonn.de

Rittenhouse Laboratory University of Pennsylvania 19102 Philadelphia, USA E-mail: mporet@sas.upenn.edu

Bethe Center for Theoretical Physics Universität Bonn D-53115 Bonn, Germany E-mail: schimann@th.physik.uni-bonn.de

Max Planck Institute for Mathematics, Vivatsgasse 7, D-53111, Bonn, Germany E-mail: martin@raum-brothers.eu Homepage: http://raum-brothers.eu/martin