

## On the topology of $\mathcal{H}(2)$

Duc-Manh Nguyen

**Abstract.** The space  $\mathcal{H}(2)$  consists of pairs  $(M, \omega)$ , where  $M$  is a Riemann surface of genus two, and  $\omega$  is a holomorphic 1-form which has only one zero of order two. There exists a natural action of  $\mathbb{C}^*$  on  $\mathcal{H}(2)$  by multiplication to the holomorphic 1-form. In this paper, we single out a proper subgroup  $\Gamma$  of  $\mathrm{Sp}(4, \mathbb{Z})$  generated by three elements, and show that the space  $\mathcal{H}(2)/\mathbb{C}^*$  can be identified with the quotient  $\Gamma \backslash \mathcal{J}_2$ , where  $\mathcal{J}_2$  is the Jacobian locus in the Siegel upper half space  $\mathfrak{S}_2$ . A direct consequence of this result is that  $[\mathrm{Sp}(4, \mathbb{Z}) : \Gamma] = 6$ . The group  $\Gamma$  can also be interpreted as the image of the fundamental group of  $\mathcal{H}(2)/\mathbb{C}^*$  in the symplectic group  $\mathrm{Sp}(4, \mathbb{Z})$ .

**Mathematics Subject Classification (2010).** 57S30, 22E40.

**Keywords.** Riemann surface, translation surface.

### 1. Introduction

In this paper we are concerned with translation surfaces in the stratum  $\mathcal{H}(2)$ . Each element of  $\mathcal{H}(2)$  can be either considered as a translation surface having only one singularity of angle  $6\pi$  together with a parallel line field, or a pair  $(M, \omega)$ , where  $M$  is a Riemann surface of genus two, and  $\omega$  is a holomorphic 1-form having a single zero of order two on  $M$ . Using the latter viewpoint, we see that  $\mathbb{C}^*$  acts naturally on  $\mathcal{H}(2)$  by multiplication to the holomorphic 1-form. Note that, if  $\omega$  has only one zero on  $M$ , then this zero must be a Weierstrass point of  $M$ . Therefore, the quotient  $\mathcal{H}(2)/\mathbb{C}^*$  consists of pairs  $(M, W)$ , where  $M$  is a Riemann surface of genus two, and  $W$  is a Weierstrass point of  $M$ , two pairs  $(M_1, W_1)$  and  $(M_2, W_2)$  are identified if there exists a conformal homeomorphism  $\phi: M_1 \rightarrow M_2$  such that  $\phi(W_1) = W_2$ .

The space  $\mathcal{H}(2)$  is well known to be a complex orbifold of dimension 4. For any pair  $(M, \omega)$ , let  $\gamma_1, \dots, \gamma_4$  be a basis of the group  $H_1(M, \mathbb{Z})$ . Then the *period mapping*

$$(M, \omega) \mapsto \left( \int_{\gamma_1} \omega, \dots, \int_{\gamma_4} \omega \right) \in \mathbb{C}^4$$

gives a local chart for  $\mathcal{H}(2)$  in a neighborhood of  $(M, \omega)$  (see [2], [7], [14]). Consequently, we see that  $\mathcal{H}(2)/\mathbb{C}^*$  can be endowed with a complex projective orbifold structure.

In Section 3, we introduce a construction of translations surface in  $\mathcal{H}(2)$  from triples of parallelograms by a unique gluing model. This construction gives rise to the notion of *parallelogram decomposition* of surfaces in  $\mathcal{H}(2)$ . Actually, given a translation surface  $(M, \omega)$  in  $\mathcal{H}(2)$ , there exist infinitely many parallelogram decompositions of  $(M, \omega)$ . From a fixed parallelogram decomposition, one can obtain other ones by applying some elementary moves, which are called  $T$ ,  $S$  and  $R$ , these moves are realized by some homeomorphisms of the surface  $M$  whose actions on the group  $H_1(M, \mathbb{Z})$  (in an appropriate basis) are given by the following matrices respectively:

$$T = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \end{pmatrix},$$

and

$$R = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Let  $\Gamma$  be the group generated by the matrices  $T$ ,  $S$  and  $R$ , then  $\Gamma$  is a proper subgroup of  $\text{Sp}(4, \mathbb{Z})$ . The main result of this paper is the following:

**Theorem 1.1.** *There exists a homeomorphism from  $\mathcal{H}(2)/\mathbb{C}^*$  to the quotient  $\Gamma \backslash \mathcal{J}_2$ , where  $\mathcal{J}_2$  is the Jacobian locus of Riemann surfaces of genus two in the Siegel upper half space  $\mathcal{S}_2$ . As a consequence, we have  $[\text{Sp}(4, \mathbb{Z}) : \Gamma] = 6$ .*

Since every Riemann surface of genus two is a two-sheeted branched cover of the sphere  $\mathbb{C}\mathbb{P}^1$ , there exists a natural identification between  $\mathcal{H}(2)/\mathbb{C}^*$  and the moduli space of pairs  $(\lambda_0, \underline{\lambda})$ , where  $\lambda_0 \in \mathbb{C}\mathbb{P}^1$ , and  $\underline{\lambda}$  is a subset of cardinal five of  $\mathbb{C}\mathbb{P}^1 \setminus \{\lambda_0\}$  up to action of  $\text{Aut}(\mathbb{C}\mathbb{P}^1)$ .

Let  $\text{Mod}_{0,6}$  denote the mapping class group of the sphere with six punctures, and  $\text{Mod}_{0,5}^*$ , the subgroup of index 6 in  $\text{Mod}_{0,6}$  that fixes one of the punctures. The space  $\mathcal{H}(2)/\mathbb{C}^*$  can be identified with the quotient  $\mathcal{T}_{0,6}/\text{Mod}_{0,5}^*$ , where  $\mathcal{T}_{0,6}$  is the Teichmüller space of the sphere with six punctures. The group  $\Gamma$  can be then considered as the image of  $\text{Mod}_{0,5}^*$  in  $\text{Sp}(4, \mathbb{Z})$ . Note that we have an isomorphism between  $\text{Mod}_{0,5}^*$  and  $B_5/Z(B_5)$ , where  $B_5$  is the braid group of the closed disk with five punctures, and  $Z(B_5)$  is the center of  $B_5$ .

It is well known that the Jacobian locus  $\mathcal{J}_2$  is the quotient of  $\mathcal{T}_{0,6}$  by the Torelli group  $\mathcal{I}_2$ . Thus we have the following commutative diagram

$$\begin{array}{ccc}
 \mathcal{T}_{0,6} & \xrightarrow{\quad} & \mathcal{I}_2 \cong \mathcal{T}_{0,6}/\mathcal{I}_2 \\
 \downarrow & \swarrow \text{---} & \\
 \mathcal{H}(2)/\mathbb{C}^* \cong \mathcal{T}_{0,6}/\text{Mod}_{0,5}^* & & 
 \end{array}$$

Since the action of  $-\text{Id}$  on  $\mathfrak{S}_2$  is trivial, a direct consequence of Theorem 1.1 is the following

**Corollary 1.2.** *We have the following exact sequence*

$$1 \rightarrow \mathcal{I}_2 \rightarrow \text{Mod}_{0,5}^* \cong B_5/Z(B_5) \rightarrow \Gamma/\{\pm\text{Id}\} \rightarrow 1. \tag{1}$$

We refer to [11] for a general discussion on symplectic and unitary representations of braid groups.

The paper is organized as follows: in Section 2, we recall the basic properties of the Theta functions and give a brief explanation of how these functions allow us to compute the branched points of hyperelliptic coverings. In the following section, we introduce the notion of *parallelogram decomposition* for surfaces in  $\mathcal{H}(2)$  and the three elementary moves on those decompositions. We then define the group  $\Gamma$  as the group generated by the homology action of the elementary moves. Note that there are parallelogram decompositions for which some elementary moves can not be carried out. This means that the set of parallelogram decompositions is not the right place to study the action of the group generated by the elementary moves. To fix this problem, in Section 4, we introduce the notion of *admissible decomposition*, which generalizes the one of parallelogram decomposition. For every admissible decomposition, all the elementary moves can be carried out. The key ingredient of the proof of Theorem 1.1 is the fact that the symplectic homology bases associated to two admissible decompositions are always related by an element of the group  $\Gamma$ . This fact is the content of Theorem 5.1, which is proven in Section 5. The proof Theorem 1.1 is then given in Section 6. In the Appendices, we give the proof of the fact that every surface in  $\mathcal{H}(2)$  admits parallelogram decompositions, and we give an explicit family of  $\Gamma$ -right cosets in  $\text{Sp}(4, \mathbb{Z})$ .

**Acknowledgements.** The author is grateful to Université Paris Sud, Max Planck Institute for Mathematics, and Institute for Mathematical Sciences, NUS Singapore. He warmly thanks François Labourie for the advices and stimulating discussions. He thanks the referee for his patience and the helpful comments.

## 2. Siegel upper half space and Theta functions

In this section, we recall the definition and some basic properties of the theta functions. Our main references are [13] and [4].

**2.1. Siegel upper half space and Jacobian locus.** For any integer  $g \geq 1$ , the Siegel upper half space  $\mathfrak{S}_g$  is the space of complex symmetric matrices  $Z$  in  $\mathfrak{M}(g, \mathbb{C})$  such that  $\text{Im}(Z)$  is positive definite. This space is the quotient of the real symplectic group  $\text{Sp}(2g, \mathbb{R})$  by the compact subgroup  $U(g)$ . The integral symplectic group  $\text{Sp}(2g, \mathbb{Z})$  is a lattice in  $\text{Sp}(2g, \mathbb{R})$  which acts properly discontinuously on  $\mathfrak{S}_g$ . The quotient  $\mathcal{A}_g = \text{Sp}(2g, \mathbb{Z}) \backslash \mathfrak{S}_g$  is the moduli space of principally polarized Abelian varieties of dimension  $g$ . In this paper, by the real symplectic group  $\text{Sp}(2g, \mathbb{R})$  we mean the group of real  $2g \times 2g$  matrices preserving the symplectic form

$$\begin{pmatrix} J & & 0 \\ & \ddots & \\ 0 & & J \end{pmatrix}, \quad \text{where } J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}.$$

Let  $M$  be a Riemann surface of genus  $g$  and let  $\{a_1, b_1, \dots, a_g, b_g\}$  be a symplectic basis of  $H_1(M, \mathbb{Z})$ , that is,

$$\langle a_i, a_j \rangle = \langle b_i, b_j \rangle = 0 \quad \text{and} \quad \langle a_i, b_j \rangle = \delta_{ij},$$

where  $\langle \cdot, \cdot \rangle$  is the intersection form of  $H_1(M, \mathbb{Z})$ . There exist  $g$  holomorphic 1-forms  $(\phi_1, \dots, \phi_g)$  on  $M$  uniquely determined by the following condition:

$$\int_{a_j} \phi_i = \delta_{ij}.$$

The matrix  $\Pi = (\pi_{ij})_{i,j=1,\dots,g}$ , where  $\pi_{ij} = \int_{b_j} \phi_i$ , belongs to  $\mathfrak{S}_g$ , and we then have a mapping from the set of pairs  $(M, \{a_1, b_1, \dots, a_g, b_g\})$  into  $\mathfrak{S}_g$ . The image of this mapping is called the *Jacobian locus* and is denoted by  $\mathcal{J}_g$ .

For the case  $g = 2$ , it is well known that the complement of  $\mathcal{J}_2$  in  $\mathfrak{S}_2$  is a union of countably many copies of  $\mathfrak{S}_1 \times \mathfrak{S}_1$ , where  $\mathfrak{S}_1$  is the upper half plane  $\mathfrak{S}_1 = \{z \in \mathbb{C} : \text{Im}z > 0\}$  (see [6], [12]).

**2.2. Theta function.** Fix an integer  $g \geq 1$  and let  $\mathfrak{S}_g$  be the Siegel upper half space of genus  $g$ . The Riemann's *theta function* is a complex value function defined on  $\mathbb{C}^g \times \mathfrak{S}_g$  by the following formula

$$\theta(z, \sigma) = \sum_{N \in \mathbb{Z}^g} \exp\left(2\pi i \left(\frac{1}{2} {}^t N \sigma N + {}^t N z\right)\right).$$

The function  $\theta$  is holomorphic on  $\mathbb{C}^g \times \mathfrak{S}_g$ . We also consider functions defined on  $\mathbb{C}^g \times \mathfrak{S}_g$  by

$$\theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z, \sigma) = \sum_{N \in \mathbb{Z}^g} \exp \left\{ 2\pi i \left[ \frac{1}{2} {}^t(N + \frac{\epsilon}{2}) \sigma (N + \frac{\epsilon}{2}) + {}^t(N + \frac{\epsilon}{2})(z + \frac{\epsilon'}{2}) \right] \right\}$$

where  $\epsilon, \epsilon'$  are integer vectors. These functions are called first order theta functions with integer characteristic  $\left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right]$ .

**Proposition 2.1.** *The first order theta function with integer characteristic  $\left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right]$  has the following properties:*

- i)  $\theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z + e^{(k)}, \sigma) = \exp 2\pi i \left[ \frac{\epsilon_k}{2} \right] \theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z, \sigma),$
- ii)  $\theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z + \sigma^{(k)}, \sigma) = \exp 2\pi i \left[ -z_k - \frac{\sigma_{kk}}{2} - \frac{\epsilon'_k}{2} \right] \theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z, \sigma),$
- iii)  $\theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (-z, \sigma) = \exp 2\pi i \left[ \frac{{}^t \epsilon \epsilon'}{2} \right] \theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z, \sigma),$
- iv)  $\theta \left[ \begin{matrix} \epsilon + 2v \\ \epsilon' + 2v' \end{matrix} \right] (z, \sigma) = \exp 2\pi i \left[ \frac{{}^t \epsilon v'}{2} \right] \theta \left[ \begin{matrix} \epsilon \\ \epsilon' \end{matrix} \right] (z, \sigma),$  with  $v, v' \in \mathbb{Z}^g,$

where  $e^{(k)}$  and  $\sigma^{(k)}$  are the  $k$ -th column of the matrices  $\text{Id}_g$  and  $\sigma$  respectively.

By the Torelli theorem, we know that a closed Riemann surface  $M$  is uniquely determined by its Jacobian variety  $J(M)$ , or equivalently, by the period matrix associated to a symplectic basis of  $H_1(M, \mathbb{Z})$ . If  $M$  is hyperelliptic, then we can get more information from the period matrix by using theta functions. We have (cf. [4], VII.4)

**Theorem 2.2.** *The branched points of the two sheeted representation of a hyperelliptic Riemann surface are holomorphic functions of the period matrix. Furthermore, the hyperelliptic surface is completely determined by its period matrix.*

To illustrate the ideas of the proof, we will indicate below the method to compute some of the branched points; details of the calculations can be found in [4], VII.1 and VII.4.

Assume that  $M$  is the two-sheeted branched covering of  $\mathbb{C}\mathbb{P}^1$  ramified above  $\lambda_1, \dots, \lambda_{2g+2}$ . Let  $s_1, \dots, s_{g+1}$  be  $g+1$  simple arcs in  $\mathbb{C}\mathbb{P}^1$  such that the endpoints of  $s_i$  are  $\lambda_{2i-1}, \lambda_{2i}$ , for  $k = 1, \dots, g+1$ , and  $s_i \cap s_j = \emptyset$ , for  $i \neq j$ . We can consider  $M$  as the Riemann surface obtained by gluing two copies of  $\mathbb{C}\mathbb{P}^1$  slit along  $s_1, \dots, s_{g+1}$ . Let  $z$  be the meromorphic function on  $M$  realizing the two-sheeted branched cover from  $M$  to  $\mathbb{C}\mathbb{P}^1$ . Let  $P_i$  denote  $z^{-1}(\lambda_i)$ ,  $i = 1, \dots, 2g+2$ , then  $\{P_1, \dots, P_{2g+2}\}$  is the set of Weierstrass points of  $M$ .

Using  $\text{PSL}(2, \mathbb{C})$ , we may assume that  $\lambda_1 = 0, \lambda_2 = 1$  and  $\lambda_{2g+2} = \infty$ . The surface  $M$  is then the curve defined by the equation

$$w^2 = z(z - 1) \prod_{i=3}^{2g-1} (z - \lambda_i).$$

The function  $z$  is then characterized, up to a non zero multiplicative constant, by the property that it has a double zero at  $P_1$ , a double pole at  $P_{2g+2}$ , and it is holomorphic and non zero elsewhere. First, we specify a symplectic basis  $\{a_1, b_1, \dots, a_g, b_g\}$  of  $H_1(M, \mathbb{Z})$  as follows:

- $b_k = z^{-1}(s_k), k = 1, \dots, g + 1$ . By construction,  $b_k$  is a simple closed curve containing  $P_{2k-1}$  and  $P_{2k}$ , and  $b_k$  is preserved by the hyperelliptic involution.
- Let  $\alpha_k, k = 1, \dots, g$ , be  $g$  simple closed curves pairwise disjoint in  $\mathbb{C}\mathbb{P}^1$  satisfying
  - $\alpha_k$  intersects transversely  $s_k$  and  $s_{g+1}$ ,
  - $\text{Card}\{\alpha_k \cap s_k\} = \text{Card}\{\alpha_k \cap s_{g+1}\} = 1$ ,
  - $\alpha_k \cap s_j = \emptyset$  if  $j \notin \{k, g + 1\}$ ,
  - $\alpha_k \cap \{\lambda_1, \dots, \lambda_{2g+2}\} = \emptyset$ .

Let  $a_k$  be a connected component of  $z^{-1}(\alpha_k)$ . Note that  $a_k$  and its image under the hyperelliptic involution are disjoint.

It follows from the construction that the family  $\{a_1, b_1, \dots, a_g, b_g\}$  is a symplectic basis of  $H_1(M, \mathbb{Z})$ . Let  $\{\zeta_1, \dots, \zeta_g\}$  be the basis of  $\mathcal{H}^1(M)$ , the space of holomorphic 1-forms on  $M$ , dual to  $\{a_1, \dots, a_g\}$ , that is

$$\int_{a_j} \zeta_i = \delta_{ij}.$$

Let  $\Pi = (\pi_{ij})_{i,j=1,\dots,g}$ , with  $\pi_{ij} = \int_{b_j} \zeta_i$ , be the period matrix associated to  $\{a_1, b_1, \dots, a_g, b_g\}$ . By definition, the Jacobian variety  $J(M)$  of  $M$  is the quotient  $\mathbb{C}^g/\Lambda$ , where  $\Lambda$  is the lattice in  $\mathbb{C}^g$  which is generated by the column vectors of the  $g \times 2g$  matrix  $(\text{Id}_g, \Pi)$ . We denote by  $e^{(k)}$  and  $\pi^{(k)}, k = 1, \dots, g$ , the  $k$ -th columns of  $\text{Id}_g$  and  $\Pi$  respectively. Let  $\varphi: M \rightarrow J(M)$  be the map defined by

$$\varphi(P) = \left( \int_{P_1}^P \zeta_1, \dots, \int_{P_1}^P \zeta_g \right) \in \mathbb{C}/\Lambda,$$

where the integrals are taken along any path joining  $P_1$  to  $P$ . From the construction of the basis  $\{a_1, b_1, \dots, a_g, b_g\}$ , one can explicitly compute  $\varphi(P_i), i = 1, \dots, 2g + 2$ , as functions of  $(e^{(1)}, \dots, e^{(g)}, \pi^{(1)}, \dots, \pi^{(g)})$ . Namely, we have (see Figure 1)




Figure 1. Symplectic basis on a hyperelliptic Riemann surface.

- $\varphi(P_1) = 0,$
- $\varphi(P_2) = \frac{1}{2}\pi^{(1)},$
- $\varphi(P_3) = \frac{1}{2}(\pi^{(1)} + e^{(1)} + e^{(2)}),$
- ...
- $\varphi(P_{2k+1}) = \frac{1}{2}(\pi^{(1)} + \dots + \pi^{(k)} + e^{(1)} + \dots + e^{(k+1)}),$
- $\varphi(P_{2k+2}) = \frac{1}{2}(\pi^{(1)} + \dots + \pi^{(k+1)} + e^{(1)} + \dots + e^{(k+1)}),$
- ...
- $\varphi(P_{2g+1}) = \frac{1}{2}(\pi^{(1)} + \dots + \pi^{(g)} + e^{(1)}),$
- $\varphi(P_{2g+2}) = \frac{1}{2}e^{(1)}.$

Since  $\Pi \in \mathfrak{S}_g$ , we can now consider the first order theta functions with integer characteristic  $\theta \left[ \begin{smallmatrix} \epsilon' \\ \epsilon' \end{smallmatrix} \right] (z, \Pi)$ . From Proposition 2.1, we see that for any  $\nu \in \Lambda$ ,  $\theta \left[ \begin{smallmatrix} \epsilon' \\ \epsilon' \end{smallmatrix} \right] (z + \nu, \Pi)$  differs from  $\theta \left[ \begin{smallmatrix} \epsilon' \\ \epsilon' \end{smallmatrix} \right] (z, \Pi)$  by a multiplicative factor. It turns out that the multiplicative behavior of the theta functions is such that

$$f(P) := \frac{\theta^2 \left[ \begin{smallmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \end{smallmatrix} \right] (\varphi(P), \Pi)}{\theta^2 \left[ \begin{smallmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \end{smallmatrix} \right] (\varphi(P), \Pi)}$$

is a meromorphic function on  $M$  with divisor  $P_1^2 P_{2g+2}^{-2}$ . Hence

$$f = cz \quad \text{where } c \in \mathbb{C}^*.$$

The constant  $c$  can be evaluated at  $P_2$  since we have  $f(P_2) = cz(P_2) = c$ . Using the fact that  $\varphi(P_2) = \frac{1}{2}\pi^{(1)}$ , we have

$$c = f(P_2) = \frac{\theta^2 \left[ \begin{smallmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \end{smallmatrix} \right] (\frac{1}{2}\pi^{(1)}, \Pi)}{\theta^2 \left[ \begin{smallmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \end{smallmatrix} \right] (\frac{1}{2}\pi^{(1)}, \Pi)}.$$

It follows that

$$z(P) = \frac{\theta^2 \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \end{bmatrix} (\frac{1}{2}\pi^{(1)}, \Pi) \theta^2 \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \end{bmatrix} (\varphi(P), \Pi)}{\theta^2 \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 1 & 1 & 0 & \dots & 0 \end{bmatrix} (\frac{1}{2}\pi^{(1)}, \Pi) \theta^2 \begin{bmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & 1 & 0 & \dots & 0 \end{bmatrix} (\varphi(P), \Pi)}. \tag{2}$$

By setting  $P = P_j$ , for  $j = 3, \dots, 2g + 1$ , we get a formula for  $\lambda_j$ . Note that this formula is only useful for  $j = 3, 4, 6, \dots, 2g$ , for other values of  $j$ , both the numerator and denominator vanish. By replacing  $P_1$  by another Weierstrass point in the definition of  $\varphi$ , we can get similar formulae for the  $\lambda_j$  which can not be computed directly from (2).

### 3. The group $\Gamma$

In this section, we define the group  $\Gamma$  and prove some of its properties.

**3.1. Construction of surfaces in  $\mathcal{H}(2)$  by gluing parallelograms.** Any parallelogram in  $\mathbb{R}^2$  is determined by a pair of vectors in  $\mathbb{R}^2$ . In this section, we will represent any parallelogram by a pair of complex numbers  $(z_1, z_2)$  such that  $\text{Im}(\bar{z}_1 z_2) > 0$ ,  $\text{Im}(\bar{z}_1 z_2)$  is actually the area of the parallelogram. Note that by this convention, the pairs  $(z_1, z_2), (z_2, -z_1), (-z_1, -z_2), (-z_2, z_1)$  represent the same parallelogram.

Let  $\mathcal{P}^+$  denote the set  $\{(z_1, z_2) \in \mathbb{C}^2 : \text{Im} > 0(\bar{z}_1 z_2) > 0\}$ . Given 4 complex numbers  $(z_1, \dots, z_4)$  such that  $(z_1, z_2), (z_2, z_3), (z_3, z_4)$  all belong to  $\mathcal{P}^+$ , let  $A, B, C$  denote the parallelograms determined by the pairs  $(z_1, z_2), (z_2, z_3), (z_3, z_4)$  respectively. We can construct a translation surface in  $\mathcal{H}(2)$  from  $A, B, C$  as follows:

- Glue two sides of  $A$  corresponding to  $z_1$  together.
- Glue two sides of  $A$  corresponding to  $z_2$  to two sides of  $B$  also corresponding to  $z_2$ .
- Glue two sides of  $B$  corresponding to  $z_3$  to two sides of  $C$  also corresponding to  $z_3$ .
- Glue two sides of  $C$  corresponding to  $z_4$  together.

It is easy to check that the surface  $M$  obtained from this construction is of genus 2, equipped with a flat metric structure with a single cone singularity, which arises from the identification of all the vertices of  $A, B, C$ , we denote this point by  $W$ . Since all the gluings are realized by translations,  $M$  is a translation surface. We also get naturally a holomorphic 1-form on  $M$ , considered as a Riemann surface, defined as follows: since translations of  $\mathbb{R}^2$  preserve the holomorphic 1-form  $dz$ , the restrictions of  $dz$  into the parallelograms  $A, B, C$  are compatible with the gluings and give rise to a holomorphic 1-form  $\omega$  on  $M$  with only one zero at  $W$ , which is necessarily of order two. Clearly,  $(M, \omega) \in \mathcal{H}(2)$ .






Figure 2. Construction of surfaces in  $\mathcal{H}(2)$  from three parallelograms.

We know that  $M$  is a hyperelliptic surface, it is easy to visualize the hyperelliptic involution of  $M$  from its construction by gluing  $A, B, C$ . For each of the parallelograms  $A, B, C$ , consider the reflection through its center, one can easily check that these reflections agree with the gluing on the boundary of  $A, B, C$ . Thus, we have a conformal automorphism  $\tau$  of  $M$ . One can check that  $\tau^2 = \text{Id}$ , and the action of  $\tau$  on  $H_1(M, \mathbb{Z})$  is given by  $-\text{Id}$ , therefore  $\tau$  must be the hyperelliptic involution of  $M$ . We can also determine without difficulty the 6 fixed points of  $\tau$ , which are the Weierstrass points of  $M$ : two of which are contained in  $A$ , two in  $C$ , one in the interior of  $B$ , and the last one is  $W$ .

**3.2. Parallelogram decompositions and the group  $\Gamma$ .** Recall that, on translation surfaces  $(M, \omega)$ , a *saddle connection* is a geodesic segment (with respect to the flat metric) which joins singularity to singularity, the endpoints of a saddle connection may coincide. A *cylinder*  $C$  is a subset of  $M$  which is isometric to  $\mathbb{R} \times ]0, h[ / \mathbb{Z}$ , where the action of  $\mathbb{Z}$  is generated by  $(x, y) \mapsto (x + \ell, y)$  with  $\ell > 0$  and maximal with respect to this property (that is,  $C$  can not be embedded into a larger subset  $C'$  isometric to  $\mathbb{R} \times ]0, h'[ / \mathbb{Z}$ , with  $h' > h$ ). In other words,  $C$  is the union of all simple closed geodesics in the same free homotopy class. The construction of translation surfaces in  $\mathcal{H}(2)$  by gluing parallelograms with the model presented above suggests the following

**Definition 3.1.** Let  $(M, \omega)$  be a pair in  $\mathcal{H}(2)$ . A *parallelogram decomposition* of  $(M, \omega)$  is a family of six oriented saddle connections  $\{a, b_1, b_2, c_1, c_2, d\}$  verifying the following conditions:

- The intersection of any pair of saddle connections in this family is the unique zero of  $\omega$ .
- $b_1 \cup b_2$  (resp.  $c_1 \cup c_2$ ) is the boundary of a cylinder which contains  $a$  (resp.  $d$ ).
- The complement of  $a \cup b_1 \cup b_2 \cup c_1 \cup c_2 \cup d$  has three components, each of which is isometric to an open parallelogram in  $\mathbb{R}^2$ .

- The orientations of the saddle connections in this family are chosen such that  $\langle a, b_1 \rangle = \langle b_1, c_1 \rangle = \langle c_1, d \rangle = 1$ ,  $a$  goes from  $b_1$  to  $b_2$  and  $d$  goes from  $c_2$  to  $c_1$  (see Figure 3).




Figure 3. Parallelogram decomposition of surfaces in  $\mathcal{H}(2)$ .

If  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  is a parallelogram decomposition of  $(M, \omega)$  then  $(a, b_1, c_1, d)$  is not a symplectic basis of  $H_1(M, \mathbb{Z})$ . Let  $b$  (resp.  $c$ ) be a simple closed curve in the cylinder bounded by  $b_1$  and  $b_2$  (resp. by  $c_1$  and  $c_2$ ). Let  $e$  be a simple closed curve in the free homotopy class of the closed curve  $d * (-b_1)$ , that is  $e = d - b_1$  in  $H_1(M, \mathbb{Z})$ , then  $(a, b, c, e)$  is a symplectic basis of  $H_1(M, \mathbb{Z})$ , we will call it the *symplectic basis associated to  $\mathcal{D}$* .

Given a surface  $(M, \omega)$  in  $\mathcal{H}(2)$  which is obtained from three parallelograms  $A = (z_1, z_2), B = (z_2, z_3), C = (z_3, z_4)$  as in the previous section, let  $\tilde{A}, \tilde{B}, \tilde{C}$  be the subsets of  $M$  which correspond to  $A, B, C$  respectively. By construction,  $\tilde{A}$  and  $\tilde{C}$  are two cylinders, while  $\tilde{B}$  is an embedded parallelogram. Let  $a$  (resp.  $d$ ) denote the saddle connection in  $\tilde{A}$  (resp.  $\tilde{C}$ ) which corresponds to the  $z_1$  sides of  $A$  (resp.  $z_4$  sides of  $C$ ). Let  $b_1, b_2$  (resp.  $c_1, c_2$ ) denote the boundary components of  $\tilde{A}$  (resp.  $\tilde{C}$ ). We choose the orientations of  $a, b_1, b_2, c_1, c_2, d$  in such a way that

$$\int_a \omega = z_1, \quad \int_{b_1} \omega = \int_{b_2} \omega = z_2, \quad \int_{c_1} \omega = \int_{c_2} \omega = z_3, \quad \int_d \omega = z_4.$$

We also choose the numbering of  $(b_1, b_2)$  (resp.  $(c_1, c_2)$ ) such that the orientation of  $a$  goes from  $b_1$  to  $b_2$  and the orientation of  $d$  goes from  $c_2$  to  $c_1$ . By definition  $\{a, b_1, b_2, c_1, c_2, d\}$  is a parallelogram decomposition of  $(M, \omega)$ .

A surface  $(M, \omega)$  in  $\mathcal{H}(2)$  always admits a parallelogram decompositions (see Appendix A). The following operations allow us to get other decompositions from a particular one  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$ .

1. *The T move.* let  $a'$  be the saddle connection which is obtained from  $a$  by a Dehn twist in  $\tilde{A}$ , then  $\mathcal{D}' = \{a', b_1, b_2, c_1, c_2, d\}$  is another parallelogram decomposition of  $(M, \omega)$  (see Figure 4). Here, we consider both “left” and “right” Dehn twists.




Figure 4. The  $T$  move.

2. *The  $S$  move.* If  $\{a, b_1, b_2, c_1, c_2, d\}$  is a parallelogram decomposition of  $(M, \omega)$  then  $\mathcal{D}' = \{d, -c_2, -c_1, b_1, b_2, -a\}$  is also a parallelogram decomposition of  $(M, \omega)$ , the minus sign designates the same saddle connection with the inverse orientation (see Figure 5).




Figure 5. The  $S$  move.

3. *The  $R$  move.* Cut  $M$  along the saddle connections  $b_1, b_2$  and  $d$ , we then obtain two annuli, one of which is  $\tilde{A}$ , we denote the other by  $\tilde{A}'$ . Let  $c'_1, c'_2$  denote the images of  $c_1, c_2$  respectively under a Dehn twist in  $\tilde{A}'$ . Assume that  $c'_1, c'_2$  can be made into saddle connections in  $\tilde{A}'$ , then  $\{a, b_1, b_2, c'_1, c'_2, d\}$  is a parallelogram decomposition of  $(M, \omega)$  (see Figure 6).




Figure 6. The  $R$  move.

Let  $\mathcal{D}'$  be another parallelogram decomposition of  $(M, \omega)$  which is obtained from  $\mathcal{D}$  by one the moves presented above. Let  $(a, b, c, e)$  (resp.  $(a', b', c', e')$ ) be the symplectic basis of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  (resp.  $\mathcal{D}'$ ). The following lemma follows directly from the definitions.

**Lemma 3.2.** a) If  $\mathcal{D}'$  is obtained from  $\mathcal{D}$  by a  $T$  move then

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \begin{pmatrix} 1 & \pm 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

b) If  $\mathcal{D}'$  is obtained from  $\mathcal{D}$  by an  $S$  move then

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

c) If  $\mathcal{D}'$  is obtained from  $\mathcal{D}$  by an  $R$  move then

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & \pm 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

We denote by  $T, S, R$  the matrices of basis change corresponding to the moves  $T, S, R$  respectively, to fix ideas, we can take

$$T = \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad S = \begin{pmatrix} 0 & 1 & 0 & 1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ -1 & 0 & 1 & 0 \end{pmatrix}$$

and

$$R = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

We denote by  $\Gamma$  the subgroup of  $\mathrm{Sp}(4, \mathbb{Z})$  generated by those matrices.

### 3.3. Properties of $\Gamma$

**Lemma 3.3.** We have

(i)  $S^2 = -\mathrm{Id}_4$ .

- (ii)  $\begin{pmatrix} \text{Id}_2 & 0 \\ 0 & \text{SL}(2, \mathbb{Z}) \end{pmatrix} \subset \Gamma$ .
- (iii)  $\Gamma \subsetneq \text{Sp}(4, \mathbb{Z})$ .
- (iv)  $\Gamma$  is not a normal subgroup of  $\text{Sp}(4, \mathbb{Z})$ .

*Proof.* (i) follows from direct calculation.

(ii) We have

$$S^{-1}TS = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & -1 & 1 \end{pmatrix}.$$

Since  $\text{SL}(2, \mathbb{Z})$  is generated by

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad \text{and} \quad \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix},$$

we deduce that  $\begin{pmatrix} \text{Id}_2 & 0 \\ 0 & \text{SL}(2, \mathbb{Z}) \end{pmatrix}$  is contained in  $\Gamma$ .

(iii) The group  $\text{Sp}(4, \mathbb{Z})$  acts transitively on  $(\mathbb{Z}/2\mathbb{Z})^4 \setminus \{0\}$ , but  $\Gamma$  has two orbits:  $\mathcal{O}_1$  containing  $\mathbf{e}_1 = (1, 0, 0, 0)$  and  $\mathcal{O}_2$  containing  $\mathbf{e}_2 = (0, 1, 0, 0)$ . As a matter of fact, we have

$$\mathcal{O}_1 = \left\{ \begin{array}{lll} (1, 0, 0, 0) & (1, 1, 0, 0) & \\ (0, 1, 1, 0) & (0, 1, 0, 1) & (0, 1, 1, 1) \end{array} \right\}$$

and

$$\mathcal{O}_2 = \left\{ \begin{array}{lll} (0, 1, 0, 0) & & \\ (1, 0, 1, 0) & (1, 0, 0, 1) & (1, 0, 1, 1) \\ (1, 1, 1, 0) & (1, 1, 0, 1) & (1, 1, 1, 1) \\ (0, 0, 1, 0) & (0, 0, 0, 1) & (0, 0, 1, 1) \end{array} \right\}.$$

Here, we consider the action of  $\text{Sp}(4, \mathbb{Z})$  and  $\Gamma$  on  $(\mathbb{Z}/2\mathbb{Z})^4$  by right multiplication.

(iv) Let

$$T' = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Remark that  $T'$  does not belong to  $\Gamma$  since it sends  $\mathbf{e}_2$  to an element in  $\mathcal{O}_1$ . We have

$$T'^{-1}TT' = \begin{pmatrix} 2 & 1 & 0 & 0 \\ -1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}.$$

Observe that  $T'^{-1}TT'$  sends  $\mathbf{e}_1$  to  $\mathbf{e}_2$ , thus it does not belong to  $\Gamma$ . We can then conclude that  $\Gamma$  is not a normal subgroup of  $\text{Sp}(4, \mathbb{Z})$ . □

**Lemma 3.4.** *Set*

$$U = \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix}.$$

Then the integral symplectic group  $\mathrm{Sp}(4, \mathbb{Z})$  is generated by  $T$ ,  $S$ ,  $R$  and  $U$ .

*Proof.* See Appendix B. □

#### 4. Admissible decomposition

**4.1. Definition.** We have seen that for any surface  $(M, \omega)$  in  $\mathcal{H}(2)$ , given a parallelogram decomposition, other decompositions of  $(M, \omega)$  can be obtained by the elementary moves  $T$ ,  $S$  and  $R$ . However, while  $T$  and  $S$  are always realizable, the  $R$  move is not, it is only realizable when  $d * (-b_1) * c_2$  is homotopic to a saddle connection. In this section, we enlarge the set of decompositions so that the three elementary moves are always realizable. This leads us to the notion of *admissible decomposition* with respect to the pair  $(M, W)$ . The definition of admissible decomposition is inspired from parallelogram decomposition and relies on the action of the hyperelliptic involution of  $M$ .

Throughout this section,  $M$  is a fixed closed Riemann surface of genus two,  $W$  is a Weierstrass point and  $\tau$  is the hyperelliptic involution of  $M$ . For any closed curve  $\gamma$  with basepoint  $W$ , we denote by  $[\gamma]$  the homotopy class of  $\gamma$  in  $\pi_1(M, W)$ .

**Definition 4.1** (Admissible decomposition). Let  $\{a, b_1, b_2, c_1, c_2, d\}$  be six oriented simple closed curves containing  $W$ . We say that  $\{a, b_1, b_2, c_1, c_2, d\}$  is an *admissible decomposition* for the pair  $(M, W)$  if

- the intersection of any pair of curves in this family is  $\{W\}$ ,
- $\tau(a) = -a, \tau(d) = -d$ ,
- $\tau(b_1) = -b_2, \tau(c_1) = -c_2$ ,
- $a \setminus \{W\}$  is contained in an open annulus  $A_b$  bounded by  $b_1$  and  $b_2$ ,
- $d \setminus \{W\}$  is contained in an open annulus  $A_c$  bounded by  $c_1$  and  $c_2$ ,
- $M \setminus (\bar{A}_b \cup \bar{A}_c)$  is homeomorphic to an open disk.

The orientations of  $\{a, b_1, b_2, c_1, c_2, d\}$  are chosen such that

- $\langle a, b_1 \rangle = \langle a, b_2 \rangle = \langle c_1, d \rangle = \langle c_2, d \rangle = 1$ , where  $\langle \cdot, \cdot \rangle$  is the intersection form of  $H_1(M, \mathbb{Z})$ ,
- in the annulus  $A_b$ , the orientation of  $a$  goes from  $b_1$  to  $b_2$ ,
- in the annulus  $A_c$ , the orientation of  $d$  goes from  $c_2$  to  $c_1$ ,

- the boundary of the disk  $M \setminus (\bar{A}_b \cup \bar{A}_c)$  with the induced orientation is the concatenation  $b_1 * c_1 * (-b_2) * (-c_2)$ , which implies in particular that  $[b_1][c_1][b_2]^{-1}[c_2]^{-1} = 1$  in  $\pi_1(M, W)$ .

**Example.** If  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  is a parallelogram decomposition for a pair  $(M, \omega)$ , where  $\omega$  is a holomorphic 1-form with double zero at  $W$ , then  $\mathcal{D}$  is an admissible decomposition for the pair  $(M, W)$ . Note that on a fixed translation surface in  $\mathcal{H}(2)$ , there are admissible decompositions which can not be realized as parallelogram decompositions.

**4.2. Projection to the sphere and associated homology basis.** Let  $\rho: M \rightarrow \mathbb{CP}^1$  be the two-sheeted branched covering from  $M$  onto  $\mathbb{CP}^1$  ramified at the Weierstrass points of  $M$ . By definition, we have  $\rho(P) = \rho(P')$  if and only if  $P' \in \{P, \tau(P)\}$ . Let  $P_0, \dots, P_5$  denote the images of the Weierstrass points of  $M$  by  $\rho$ , with  $P_0 = \rho(W)$ . Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  be an admissible decomposition for the pair  $(M, W)$ . The projections of the curves in  $\mathcal{D}$  satisfy (see Figure 7).

- $\rho(a) = \bar{a}$  is a simple arc joining  $P_0$  to another point in  $\{P_1, \dots, P_5\}$ . Without loss of generality, we can assume that the endpoints of  $\bar{a}$  are  $P_0$  and  $P_1$ .
- $\rho(b_1) = \rho(b_2) = \bar{b}$  is a simple closed curve with basepoint  $P_0$ ,  $\bar{b}$  is the boundary of an open disc  $D_1$  which contains  $\bar{a} \setminus \{P_0\}$  and two points of the set  $\{P_1, \dots, P_5\}$ . We can assume that  $D_1 \cap \{P_1, \dots, P_5\} = \{P_1, P_2\}$ .
- $\rho(d) = \bar{d}$  is a simple arc disjoint from  $D_1$ , joining  $P_0$  to a point in  $\{P_3, \dots, P_5\}$ . We can assume that the endpoints of  $\bar{d}$  are  $P_0$  and  $P_3$ .
- $\rho(c_1) = \rho(c_2) = \bar{c}$  is a simple closed curve with basepoint  $P_0$  disjoint from  $D_1$ ,  $\bar{c}$  is the boundary of an open disc  $D_2$  which contains  $\bar{d} \setminus \{P_0\}$  and two points in the set  $\{P_3, \dots, P_5\}$ . We can assume that  $D_2 \cap \{P_1, \dots, P_5\} = \{P_3, P_4\}$ .




Figure 7. Projection of an admissible decomposition to  $\mathbb{CP}^1$ .

**Remark.** The orientations of the curves in  $\mathcal{D}$  do not determine the orientations of the curves  $\{\bar{a}, \bar{b}, \bar{c}, \bar{d}\}$ .

Let  $\bar{b}_*$  be a simple arc contained in  $D_1$  which joins  $P_1$  to  $P_2$  such that  $\bar{b}_* \cap \bar{a} = \{P_1\}$ . Observe that  $b = \rho^{-1}(\bar{b}_*)$  is a simple closed curve in  $M$ , freely homotopic to  $b_1$  and  $b_2$ , and we have  $\tau(b) = -b$ . Similarly, let  $\bar{c}_*$  be a simple arc in  $D_2$  joining  $P_3$  to  $P_4$  such that  $\bar{c}_* \cap \bar{d} = \{P_3\}$ , then  $c = \rho^{-1}(\bar{c}_*)$  is a simple closed curve freely homotopic to  $c_1, c_2$  such that  $\tau(c) = -c$ .

Now, let  $\bar{e}$  be a simple arc in  $\mathbb{C}\mathbb{P}^1$  which joins  $P_5$  to  $P_4$  such that  $\bar{e} \cap (\bar{b} \cup \bar{d}) = \emptyset$  (see Figure 7), then  $e = \rho^{-1}(\bar{e})$  is a simple closed curve such that  $\tau(e) = -e$ . Note that  $\bar{e}$  is unique up to homotopy with fixed endpoints in  $\mathbb{C}\mathbb{P}^1 \setminus (\bar{b} \cup \bar{d})$ .

Recall that  $a, d$  are already oriented, and we have an orientation for  $b$  which is induced by the orientation of  $b_1$  and  $b_2$ . Choose the orientation of  $e$  such that  $\langle c, e \rangle = 1$ , then  $(a, b, d, e)$  is a symplectic basis of  $H_1(M, \mathbb{Z})$ . We will call  $(a, b, c, e)$  the basis associated to the decomposition  $\mathcal{D}$ . It is easy to see that this basis is also the one described in Section 2. Note also that if  $\mathcal{D}$  is a parallelogram decomposition, then the two definitions of associated symplectic basis of  $H_1(M, \mathbb{Z})$  agree (see Figure 8).




Figure 8. Symplectic homology basis associated to an admissible decomposition.

**4.3. Elementary moves.** Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  be an admissible decomposition for the pair  $(M, W)$ . Let  $\bar{a}, \bar{b}, \bar{c}, \bar{d}, \bar{e}, D_1, D_2$  be as above.

- *The T move.* Let  $\bar{a}'$  be a simple arc joining  $P_0$  to  $P_2$  such that  $(\bar{a}' \setminus \{P_0\}) \subset D_1$  and  $\bar{a}' \cap \bar{a} = \{P_0\}$ . The preimage  $a'$  of  $\bar{a}'$  in  $M$  is a simple closed curves contained in the annulus bounded by  $b_1, b_2$  which satisfies  $\tau(a') = -a'$ . By choosing an appropriate orientation for  $a'$ , we see that the family  $\mathcal{D}' = \{a', b_1, b_2, c_1, c_2, d\}$  is an admissible decomposition for  $(M, W)$ . The symplectic homology basis associated to  $\mathcal{D}'$  is  $(a', b, c, e)$  and we have

$$\begin{pmatrix} a' \\ b \\ c \\ e \end{pmatrix} = \begin{pmatrix} 1 & \pm 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ d \end{pmatrix}.$$

We call this transformation the  $T$  move (see Figure 9).






Figure 9.  $T$  move for an admissible decomposition.

- *The  $S$  move.* We define the  $S$  move for admissible decompositions in the same way as for parallelogram decompositions, that is, an  $S$  move transforms  $\mathcal{D}$  into the family  $\mathcal{D}' = \{d, -c_2, -c_1, b_1, b_2, -a\}$ . To see what happens to the curves in  $\mathbb{C}\mathbb{P}^1$ , let  $\bar{e}'$  be a simple arc in  $\mathbb{C}\mathbb{P}^1$  joining  $P_5$  to  $P_2$  such that  $\bar{e}' \cap (\bar{a} \cup \bar{c}) = \emptyset$ , then the symplectic homology basis associated to  $\mathcal{D}'$  is  $(d, -c, b, e')$  where  $e' = \rho^{-1}(\bar{e}')$ . It is easy to check that the symplectic homology bases associated to  $\mathcal{D}$  and  $\mathcal{D}'$  are related by the matrix  $S$ .




Figure 10.  $S$  move for an admissible decomposition.

- *The  $R$  move.* Let  $\bar{c}'$  be a simple closed curve in  $\mathbb{C}\mathbb{P}^1$  with basepoint  $P_0$  which is disjoint from  $D_1$  and bounds an open disc  $D'_2$  such that (see Figure 11)

- $D'_2 \cap \{P_1, \dots, P_5\} = \{P_3, P_5\}$ ,
- $(\bar{d} \setminus \{P_0\}) \subset D'_2$ .

The preimage of  $\bar{c}'$  in  $M$  is the union of two simple closed curves  $c'_1, c'_2$  with basepoint  $W$ . By choosing an appropriate orientation of  $c'_1, c'_2$ , we see that the family  $\mathcal{D}' = \{a, b_1, b_2, c'_1, c'_2, d\}$  is an admissible decomposition. Let  $\bar{c}'_*$  be a simple arc in  $D'_2$  joining  $P_3$  to  $P_5$  and let  $c'$  be the preimage of  $\bar{c}'_*$  in  $M$ . Then  $(a, b, c', e)$  is the symplectic homology basis associated to  $\mathcal{D}'$ . One can

easily check that

$$\begin{pmatrix} a \\ b \\ c' \\ e \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & \pm 1 \\ 0 & 0 & 0 & 1 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$




Figure 11.  $R$  move for an admissible decomposition.

The elementary moves  $T$ ,  $S$ ,  $R$  defined above transform an admissible decomposition for the pair  $(M, W)$  into another admissible decomposition for the same pair. Let us now introduce an elementary move which transforms an admissible decomposition for the pair  $(M, W)$  into an admissible decomposition for another pair  $(M, W')$ .

Let  $\bar{a}'$  be a simple arc in  $\mathbb{C}\mathbb{P}^1$  joining  $P_5$  to  $P_4$  such that  $\bar{a}' \cap (\bar{b} \cup \bar{d}) = \emptyset$ . Let  $\bar{d}'$  be a simple arc joining  $P_5$  to  $P_2$  such that  $\bar{d}' \cap (\bar{a} \cup \bar{c}) = \emptyset$ . Let  $\bar{b}'$  and  $\bar{c}'$  be two simple closed curves with basepoint  $P_5$  which are respectively the boundary of the open disks  $D'_1$  and  $D'_2$  satisfying (see Figure 12)

- $D'_1 \cap D'_2 = \emptyset, D'_i \cap D_i = \emptyset, i = 1, 2,$
- $D'_1 \cap \{P_0, \dots, P_4\} = \{P_3, P_4\}, D'_2 \cap \{P_0, \dots, P_4\} = \{P_1, P_2\},$
- $\text{int}(\bar{a}') \subset D'_1, \text{int}(\bar{d}') \subset D'_2.$

Then the preimages of  $\bar{a}', \bar{b}', \bar{c}', \bar{d}'$  is an admissible decomposition  $\mathcal{D}'$  for the pair  $(M, \rho^{-1}(P_5))$ . Actually, the family  $(\bar{a}', \bar{b}', \bar{c}', \bar{d}')$  gives rise to two admissible decompositions, each of which is determined by the orientation of  $a' = \rho^{-1}(\bar{a}')$ . The symplectic homology bases associated those decompositions are related by  $-\text{Id}$ . Let  $(a', b', c', e')$  be the symplectic homology basis associated to  $\mathcal{D}'$ , we have

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \pm \begin{pmatrix} 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix} = U^{\pm 1} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

We will call the transformation from  $\mathcal{D}$  into  $\mathcal{D}'$  the  $U$  move.

From the definitions, the following lemma is clear.




Figure 12.  $U$  move for admissible decompositions.

**Lemma 4.2.** *Let  $\mathcal{D}$  be an admissible decomposition for the pair  $(M, W)$  and  $\gamma$  an element of the group  $\Gamma$ . Then there exists an admissible decomposition  $\mathcal{D}'$  for the same pair such that the symplectic homology bases associated to  $\mathcal{D}$  and  $\mathcal{D}'$  are related by  $\gamma$ .*

From Lemma 3.4, we know that the family  $\{T, S, R, U\}$  generates the group  $\text{Sp}(4, \mathbb{Z})$ , therefore we have

**Lemma 4.3.** *Let  $\mathcal{D}$  be an admissible decomposition for the pair  $(M, W)$  and  $A$  an element of  $\text{Sp}(4, \mathbb{Z})$ . Then there exist a Weierstrass point  $W'$  of  $M$  and an admissible decomposition  $\mathcal{D}'$  for the pair  $(M, W')$  such that the symplectic homology bases associated to  $\mathcal{D}$  and  $\mathcal{D}'$  are related by  $A$ .*

### 5. Symplectic homology bases associated to admissible decompositions and the group $\Gamma$

Let  $(M, W)$  be an element of the space  $\mathcal{H}(2)/\mathbb{C}^*$ . Our aim in this section is to prove the following theorem, which is the key ingredient of the proof of Theorem 1.1.

**Theorem 5.1.** *Let  $\mathcal{D}$  and  $\mathcal{D}'$  be two admissible decompositions for the pair  $(M, W)$  with the associated symplectic homology bases  $(a, b, c, e)$  and  $(a', b', c', e')$  respectively. Then there exists an element  $\gamma$  in  $\Gamma$  such that*

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \gamma \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

In what follows, we denote by  $\tau$  the hyperelliptic involution of  $M$ , by  $\rho$  the two-sheeted branched covering from  $M$  onto  $\mathbb{C}\mathbb{P}^1$ , and by  $P_0, \dots, P_5$  the images of the Weierstrass points of  $M$  by  $\rho$ , where  $P_0 = \rho(W)$ . We also equip  $M$  with the

hyperbolic metric in the conformal class of the Riemann surface structure. Note that  $\tau$  is now an isometry of  $M$ . Recall that, for any closed curve  $\alpha$  in  $M$  which contains  $W$ , we denote by  $[\alpha]$  the homotopy class of  $\alpha$  in  $\pi_1(M, W)$ .

**5.1. Admissible decompositions with common subfamily.** Let us first prove the following

**Proposition 5.2.** *Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  and  $\mathcal{D}' = \{a', b'_1, b'_2, c'_1, c'_2, d'\}$  be two admissible decompositions for the pair  $(M, W)$ . Assume that  $\bar{b} = \rho(b_1)$  and  $\bar{b}' = \rho(b'_1)$  are homotopic in  $\pi_1(\mathbb{C}\mathbb{P}^1 \setminus \{P_1, \dots, P_5\}, P_0)$ . Then there exists an element  $\gamma \in \Gamma$  such that the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\mathcal{D}'$  are related by  $\gamma$ .*

*Proof.* Since  $\bar{b}$  and  $\bar{b}'$  are homotopic in  $\pi_1(\mathbb{C}\mathbb{P}^1 \setminus \{P_1, \dots, P_5\}, P_0)$ , there exists a homeomorphism  $\Phi$  of  $\mathbb{C}\mathbb{P}^1$  isotopic to the identity relative to  $\{P_0, \dots, P_5\}$  such that  $\Phi(\bar{b}) = \bar{b}'$  (see [1], Theorem A.4, or [3], Lemma 2.9). The homotopy from the identity of  $\mathbb{C}\mathbb{P}^1$  to  $\Phi$  can be lifted to a homotopy of  $M$  which is identity on the set of Weierstrass points of  $M$ . Therefore, we can assume that  $b_1 \cup b'_1 = b'_1 \cup b'_2$  as subsets of  $M$ . A priori, the orientations of  $(b_1, b_2)$  and  $(b'_1, b'_2)$  may not be the same, but since  $-\text{Id} \in \Gamma$ , we can assume that they have the same orientation which means that  $b_1 = b'_1$  in  $H_1(M, \mathbb{Z})$ . Note that the orientation of  $(b_1, b_2)$  determines the orientation of  $a$  by the condition  $\langle a, b_1 \rangle = 1$ , and consequently, we get a unique numbering of the pair  $(b_1, b_2)$  by the condition that  $a$  goes from  $b_1$  to  $b_2$ .

By definition,  $b_1 \cup b_2$  is the boundary of an open annulus  $A_b$  which contains both  $a \setminus \{W\}$  and  $a' \setminus \{W\}$ , therefore there exists an integer  $n$  such that  $a'$  is homotopic to the image of  $a$  by  $n$  Dehn twists in  $A_b$ . Thus, by applying the  $T$  move  $n$  times, we can assume that  $a' = a$  as subsets of  $M$ . Let  $(a, b, c, e)$  and  $(a', b', c', e')$  be the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\mathcal{D}'$  respectively. It follows that we have the following equality in  $H_1(M, \mathbb{Z})$

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \begin{pmatrix} \text{Id}_2 & 0 \\ X & Y \end{pmatrix} \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix},$$

with  $X, Y \in \mathbf{M}_{2 \times 2}(\mathbb{Z})$ . Since  $\begin{pmatrix} \text{Id}_2 & 0 \\ X & Y \end{pmatrix}$  belongs to  $\text{Sp}(4, \mathbb{Z})$ , simple computations show that we must have  $X = 0$  and  $Y \in \text{SL}(2, \mathbb{Z})$ . Since the group  $\begin{pmatrix} \text{Id}_2 & 0 \\ 0 & \text{SL}(2, \mathbb{Z}) \end{pmatrix}$  is contained in  $\Gamma$  (cf. Lemma 3.3), the proposition follows.  $\square$

**5.2. Standard decomposition.** Let us now prove the following

**Lemma 5.3.** *Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  be an admissible decomposition of the pair  $(M, W)$ . Let  $b_0$  and  $c_0$  be the simple closed geodesic in the free homotopy class of  $b_1$  and  $c_1$  respectively, then we have*

- a)  $b_0 \cap c_0 = \emptyset$ ,
- b)  $\tau(b_0) = -b_0, \tau(c_0) = -c_0$ ,
- c)  $W \notin b_0 \cup c_0$ .

*Proof.* a) Let  $b$  (resp.  $c$ ) be a simple closed curve freely homotopic to  $b_1$  (resp. to  $c_1$ ) which is contained in the annulus bounded by  $b_1$  and  $b_2$  (resp.  $c_1$  and  $c_2$ ). By construction  $b$  and  $c$  are freely homotopic to  $b_0$  and  $c_0$ , respectively, and  $b \cap c = \emptyset$ . It is well known that  $\text{Card}\{b_0 \cap c_0\}$  is the intersection number  $\iota(b, c)$  of the free homotopy classes of  $b$  and  $c$ , thus we have that  $\text{Card}\{b_0 \cap c_0\} = 0$ .

b) By definition, we see that  $b_1$  is freely homotopic to  $b_2$  and  $\tau(b_1) = -b_2$ , therefore  $\tau(b_1)$  is freely homotopic to  $-b_1$ . Now, since  $b_0$  is the unique simple closed geodesic in the free homotopy class of  $b_1$ ,  $-b_0$  is then the unique simple closed geodesic in the free homotopy class of  $-b_1$ . Since  $\tau$  is an isometry,  $\tau(b_0)$  must be simple closed geodesic in the free homotopy class of  $\tau(b_1)$ , hence  $\tau(b_0)$  is freely homotopic to  $-b_1$ , therefore we must have  $\tau(b_0) = -b_0$ .

c) Suppose that  $W \in b_0$ . Since  $b_0$  is freely homotopic to  $b_1$ , there exists  $[h] \in \pi_1(M, W)$  such that  $[b_0] = [h][b_1][h]^{-1}$ . Note that we have  $[\tau(b_0)] = [b_0]^{-1}$ , therefore

$$[h][b_1]^{-1}[h]^{-1} = [\tau(h)][\tau(b_1)][\tau(h)]^{-1} = [\tau(h)][a]^{-1}[b_1]^{-1}[a][\tau(h)]^{-1}.$$

It follows that

$$[a][\tau(h)]^{-1}[h][b_1]^{-1} = [b_1]^{-1}[a][\tau(h)]^{-1}[h]. \tag{3}$$

We deduce that  $[b_1]^{-1}$  and  $[a][\tau(h)]^{-1}[h]$  commute. But  $[b_1]$  is a simple closed non-separating curve, therefore we have

$$[a][\tau(h)]^{-1}[h] = [b_1]^n \quad \text{with } n \in \mathbb{Z}. \tag{4}$$

Recall that  $\tau$  acts by  $-\text{Id}$  on  $H_1(M, \mathbb{Z})$ , thus (4) implies the following equality in  $H_1(M, \mathbb{Z})$

$$nb_1 - a = 2h. \tag{5}$$

It follows that  $\langle a, b_1 \rangle = 0 \pmod{2}$ , but by construction we know that  $\langle a, b_1 \rangle = 1$  and we get a contradiction. We can then conclude that  $b_0$  does not contain  $W$ . The same arguments apply to  $c_0$  and the lemma follows.  $\square$

**Remark.** If a simple closed curve  $g$  satisfies  $\tau(g) = -g$ , then  $g$  contains exactly two fixed points of  $\tau$ , which are Weierstrass points of  $M$ .

Let  $g$  be a simple closed geodesic in  $M$ . We will say that  $g$  has Property ( $\mathcal{P}$ ), if it satisfies

$$\tau(g) = -g \quad \text{for } W \notin g.$$

By Lemma 5.3, we know that each admissible decomposition provides us with a pair of disjoint simple closed geodesics in  $M$  satisfying Property ( $\mathcal{P}$ ). Conversely,

let  $(g_1, g_2)$  be such a pair of simple closed geodesics, we will construct an admissible decomposition of  $(M, W)$  associated to  $(g_1, g_2)$  as follows:

- Cut open  $M$  along  $g_1$  and  $g_2$ , we obtain a 4-holed sphere  $N$  which is equipped with a hyperbolic metric with geodesic boundary. Let  $g_i^+, g_i^-$  denote the boundary components of  $N$  corresponding to  $g_i, i = 1, 2$ . The curves  $g_i^\pm$  inherit the orientation of  $g_i$ . We choose the notations so that the orientation of  $g_i^+$  agrees with the one induced by the orientation of  $N$ . Note that the hyperelliptic involution  $\tau$  of  $M$  induces an isometric involution of  $N$  which interchanges  $g_i^+$  and  $g_i^-$ . By a slight abuse of notation, we will also denote by  $\tau$  the involution of  $N$ .

Let  $s_i^+$  be a path of minimal length in  $N$  from  $W$  to  $g_i^+, i = 1, 2$ . Note that by definition, we have  $s_1^+ \cap s_2^+ = \{W\}$ . Let  $s_i^-$  denote the image of  $s_i^+$  under  $\tau$ . Since  $\tau$  is an isometry,  $s_i^-$  is a path realizing the distance from  $W$  to  $g_i^-$  in  $N$ . In particular, it follows that  $s_i^+ \cap s_j^- = \{W\}$ . Note also that, since the action  $\tau$  on the tangent space at  $W$  is  $-\text{Id}$ , the union  $s_i$  of  $s_i^+$  and  $s_i^-$  is a simple geodesic segment.

Let us denote by  $\tilde{s}_i^+, \tilde{s}_i^-, \tilde{s}_i$  the geodesic segments in  $M$  corresponding to  $s_i^+, s_i^-$  and  $s_i$ , respectively. Let  $P_i^+, P_i^-$  denote the endpoints of  $\tilde{s}_i^+$  and  $\tilde{s}_i^-$  in  $g_i$ , respectively (it may happen that  $P_i^+ = P_i^-$ ), and let  $r_i$  be a simple arc in  $g_i$  which joins  $P_i^+$  to  $P_i^-$ . It follows that  $h_i = r_i \cup \tilde{s}_i$  is a simple closed curve in  $M$  containing  $W$ .

- We choose the orientations of  $h_1$  and  $h_2$  such that  $\langle h_1, g_1 \rangle = 1$  and  $\langle h_2, g_2 \rangle = 1$ . Observe that the induced orientation of  $\tilde{s}_1$  is from  $P_1^-$  to  $P_1^+$ , and the orientation of  $\tilde{s}_2$  is from  $P_2^+$  to  $P_2^-$ . There exist simple closed curves  $\tilde{b}$  and  $\tilde{c}$  containing  $W$  which satisfy the following:

- $\tilde{b}$  and  $\tilde{c}$  are disjoint from  $g_1 \sqcup g_2$ ,
- $\tilde{b}$  and  $g_1$  bound an annulus  $C_1^+$  which contains  $\tilde{s}_1^+$ ,
- $\tilde{c}$  and  $g_2$  bound an annulus  $C_2^+$  which contains  $\tilde{s}_2^+$ ,
- $\tilde{b} \cap \tilde{c} = \{W\}$ ,
- $\tilde{b} \cap \tau(\tilde{b}) = \tilde{c} \cap \tau(\tilde{c}) = \{W\}$ ,
- $\tilde{b} \cap \tau(\tilde{c}) = \tilde{c} \cap \tau(\tilde{b}) = \{W\}$ .

For  $\tilde{b}$  we can take a curve in a neighborhood of  $s_1^+ \cup g_1^+$  in  $N$ , and for  $\tilde{c}$  we can take a curve in the neighborhood of  $s_2^+ \cup g_2^+$  in  $N$ . By definition, we have

$$[\tilde{b}] = [\tilde{s}_1^+ * g_1 * (-\tilde{s}_1^+)] \quad \text{and} \quad [\tilde{c}] = [(-\tilde{s}_2^+) * g_2 * \tilde{s}_2^+] \quad \text{in } \pi_1(M, W).$$

Since  $\tau(g_i) = -g_i$ , the curves  $\tau(\tilde{b})$  and  $g_1$  (resp.  $\tau(\tilde{c})$  and  $g_2$ ) also bound an annulus denoted by  $C_1^-$  (resp.  $C_2^-$ ) which contains  $\tilde{s}_1^-$  (resp.  $\tilde{s}_2^-$ ). The union

$C_i$  of  $C_i^+$  and  $C_i^-$  is an annulus which contains  $g_i$  as a core curve. Following the orientation of  $g_2$ , we have two cases:

1.  $[\tilde{b}][\tilde{c}][\tau(\tilde{b})][\tau(\tilde{c})] = 1$  in  $\pi_1(M, W)$  (see Figure 13). In this case, we take  $a = h_1, b_1 = \tilde{b}, b_2 = -\tau(\tilde{b}), c_1 = \tilde{c}, c_2 = -\tau(\tilde{c}), d = h_2$ , then  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  is an admissible decomposition for  $(M, W)$ .




Figure 13. Admissible decomposition associated to a pair of geodesics satisfying  $(\mathcal{P})$ : Case 1.

2.  $[\tilde{b}][\tau(\tilde{c})][\tau(\tilde{b})][\tilde{c}] = 1$  in  $\pi_1(M, W)$ . Consider the two-sheeted ramified covering  $\rho: M \rightarrow \mathbb{C}\mathbb{P}^1$ . The images of  $g_1$  and  $g_2$  are two simple arcs in  $\mathbb{C}\mathbb{P}^1$  whose endpoints are images of the branched points of  $\rho$ . Recall that  $\rho$  has six branched points, one of which is  $W$ , four are contained in  $g_1$  and  $g_2$ . We can assume that the last one is mapped to  $\infty$ . Let  $\tilde{u}_2$  be a simple arc in  $\mathbb{C}\mathbb{P}^1$  as shown on the right of Figure 14. Then  $\tilde{u}_2 = \rho^{-1}(\tilde{u}_2)$  is a simple arc which joins  $P_2^+$  to  $P_2^-$  and passes through  $W$ . We denote by  $\tilde{u}_2^+$  the subarc of  $\tilde{u}_2$  from  $P_2^+$  to  $W$ . Observe  $\tilde{u}_2^- = \tau(\tilde{u}_2^+)$  is the subarc from  $P_2^-$  to  $W$ . One can easily check that

- $(-\tilde{s}_2^+) * \tilde{u}_2^+$  is homotopic to  $\tilde{s}_1^+ * \tilde{g}_1^+ * (-\tilde{s}_1^+)$  in  $\pi_1(M, W)$ ,
- $\tilde{u}_2^+ \cap \tau(\tilde{u}_2^-) = \{W\}$ ,
- $\tilde{u}_2 \cap C_1 = \{W\}$ .

Consequently, there exists a simple closed curve  $\tilde{c}'$  in  $M$  containing  $W$  such that

- $\tilde{c}'$  and  $g_2$  bound an annulus  $C_2'^+$  which contains  $\tilde{u}_2^+$ ,
- $\tau(\tilde{c}')$  and  $g_2$  bound an annulus  $C_2'^-$  which contains  $\tilde{u}_2^-$ ,
- $\tilde{c}' \cap \tilde{b} = \tilde{c}' \cap \tau(\tilde{b}) = \{W\}$ ,
- $\tilde{c}' \cap \tau(\tilde{c}') = \{W\}$ .




Figure 14. Admissible decomposition associated to a pair of geodesics satisfying  $(\mathcal{P})$ : Case 2.

We have

$$\begin{aligned}
 [\tilde{c}'] &= [(-\tilde{u}_2^+) * g_2 * \tilde{u}_2^+], \\
 &= [(-\tilde{u}_2^+) * \tilde{s}_2^+][(-\tilde{s}_2^+) * g_2 * \tilde{s}_2^+][(-\tilde{s}_2^+) * u_2^+], \\
 &= [\tilde{b}]^{-1}[\tilde{c}][\tilde{b}].
 \end{aligned}$$

Therefore,  $[\tilde{b}][\tilde{c}'][\tau(\tilde{b})][\tau(\tilde{c}')] = 1$  in  $\pi_1(M, W)$ . Let  $h'_2$  denote the simple closed curve  $r_2 \cup \tilde{u}_2^+ \cup \tilde{u}_2^-$ , we choose the orientation of  $h'_2$  in such a way that  $\langle g_2, h'_2 \rangle = 1$ . It follows that, if we take  $a = h_1, b_1 = \tilde{b}, b_2 = -\tau(\tilde{b}), c_1 = \tilde{c}', c_2 = -\tau(\tilde{c}')$  and  $d = h'_2$ , then  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  is an admissible decomposition for  $(M, W)$  (see Figure 14).

In both cases, we will call  $\mathcal{D}$  a standard decomposition associated to the pair of oriented geodesics  $(g_1, g_2)$ .

From Lemma 5.3, we know that if  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  is an admissible decomposition for the pair  $(M, W)$ , and  $b_0$  (resp.  $c_0$ ) is the simple closed geodesic in the free homotopy class of  $b_1$  (resp.  $c_1$ ), then  $b_0$  and  $c_0$  satisfy Property  $(\mathcal{P})$ . Hence, we can consider the standard decompositions associated to  $(b_0, c_0)$ . The following proposition tells us that the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to the two decompositions are related by an element of the group  $\Gamma$ .

**Proposition 5.4.** *Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  be an admissible decomposition for the pair  $(M, W)$ . Let  $b_0, c_0$  be the simple closed geodesics in the free homotopy classes of  $b_1$  and  $c_1$  respectively. Let  $\hat{\mathcal{D}} = \{\hat{a}, \hat{b}_1, \hat{b}_2, \hat{c}_1, \hat{c}_2, \hat{d}\}$  be the standard*



decomposition associated to the pair  $(b_0, c_0)$ , then the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\hat{\mathcal{D}}$  are related by an element in  $\Gamma$ .

*Proof.* Since  $b_1$  and  $\hat{b}_1$  are freely homotopic, there exists a closed curve  $h$  with basepoint  $W$  such that  $[\hat{b}_1] = [h]^{-1}[b_1][h]$  in  $\pi_1(M, W)$ . We will show that  $h \in \mathbb{Z}a \oplus \mathbb{Z}b_1 \oplus \mathbb{Z}c_1$  in  $H_1(M, \mathbb{Z})$ . By definition, we have

$$\begin{aligned} [\tau(\hat{b}_1)] &= [\hat{a}]^{-1}[\hat{b}_1]^{-1}[\hat{a}] \\ \implies [\tau(h)]^{-1}[\tau(b_1)][\tau(h)] &= [\hat{a}]^{-1}[h]^{-1}[b_1]^{-1}[h][\hat{a}] \\ \implies [\tau(h)]^{-1}[a]^{-1}[b_1]^{-1}[a][\tau(h)] &= [\hat{a}]^{-1}[h]^{-1}[b_1]^{-1}[h][\hat{a}] \\ \implies [b_1]^{-1}([a][\tau(h)][\hat{a}]^{-1}[h]^{-1}) &= ([a][\tau(h)][\hat{a}]^{-1}[h]^{-1})[b_1]^{-1}. \end{aligned}$$

It follows that  $[b_1]$  and  $[a][\tau(h)][\hat{a}]^{-1}[h]^{-1}$  commute. Since  $b_1$  is a simple closed curve, there exists  $k \in \mathbb{Z}$  such that

$$[a][\tau(h)][\hat{a}]^{-1}[h]^{-1} = [b_1]^k.$$

Therefore, in  $H_1(M, \mathbb{Z})$ , we have  $\hat{a} = a - kb_1 - 2h$ . We know that  $\langle \hat{a}, \hat{c}_1 \rangle = \langle \hat{a}, c_1 \rangle = 0$  and  $\langle a, c_1 \rangle = \langle b_1, c_1 \rangle = 0$ , hence,  $\langle h, c_1 \rangle = 0$ , which implies

$$h \in c_1^\perp = \mathbb{Z}a \oplus \mathbb{Z}b_1 \oplus \mathbb{Z}c_1.$$

Let  $(a, b, c, e)$  and  $(\hat{a}, \hat{b}, \hat{c}, \hat{e})$  be the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\hat{\mathcal{D}}$  respectively. We know that there exists  $\gamma \in \text{Sp}(4, \mathbb{Z})$  such that

$$\begin{pmatrix} \hat{a} \\ \hat{b} \\ \hat{c} \\ \hat{e} \end{pmatrix} = \gamma \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

Since in  $H_1(M, \mathbb{Z})$  we have  $\hat{b} = b$ ,  $\hat{c} = c$  and  $\hat{a} = a + kb + 2h \in \mathbb{Z}a \oplus \mathbb{Z}b \oplus \mathbb{Z}c$ , it follows that  $\gamma$  is of the form

$$\gamma = \begin{pmatrix} x & y & z & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ x' & y' & z' & t' \end{pmatrix}.$$

Now,  $\langle \hat{a}, \hat{b} \rangle = 1$  implies  $x = 1$ , and since  $z$  is the  $c$ -coordinate of  $2h$  in  $\mathbb{Z}a \oplus \mathbb{Z}b \oplus \mathbb{Z}c$ , we have

$$\hat{a} = a + mb + 2\ell c \quad \text{with } m, \ell \in \mathbb{Z}.$$

It follows that

$$\begin{aligned} \langle \hat{b}, \hat{e} \rangle = 0 &\implies x' = 0, \\ \langle \hat{c}, \hat{e} \rangle = 1 &\implies t' = 1, \\ \langle \hat{a}, \hat{e} \rangle = 0 &\implies y' = -z = -2\ell. \end{aligned}$$

We deduce that

$$\gamma = \begin{pmatrix} 1 & m & 2\ell & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2\ell & n & 1 \end{pmatrix},$$

with  $\ell, m, n$  in  $\mathbb{Z}$ . The proposition follows from Lemma 5.5 here below. □

**Lemma 5.5.** *For any integers  $\ell, m, n$ , the matrix*

$$\gamma = \begin{pmatrix} 1 & m & 2\ell & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2\ell & n & 1 \end{pmatrix}$$

*belongs to the group  $\Gamma$ .*

*Proof.* We have shown that the group  $\begin{pmatrix} \text{Id}_2 & 0 \\ 0 & \text{SL}(2, \mathbb{Z}) \end{pmatrix}$  is included in  $\Gamma$ . Thus

$$X = S \cdot \begin{pmatrix} \text{Id}_2 & 0 \\ 0 & -\text{Id}_2 \end{pmatrix} \cdot S \cdot \begin{pmatrix} \text{Id}_2 & 0 \\ 0 & -\text{Id}_2 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 2 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -2 & 0 & 1 \end{pmatrix} \in \Gamma.$$

Set

$$Y = S^{-1}T^{-1}S = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 1 \end{pmatrix}.$$

Now, straight computations show that  $T, X$  and  $Y$  commute, and  $\gamma = T^m X^\ell Y^n$ . □

**Corollary 5.6.** *Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  and  $\mathcal{D}' = \{a', b'_1, b'_2, c'_1, c'_2, d'\}$  be two admissible decompositions of the pair  $(M, W)$  such that  $b_1$  is freely homotopic to  $b'_1$  and  $c_1$  is freely homotopic to  $\pm c'_1$ . Then the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\mathcal{D}'$  are related by an element in  $\Gamma$ .*

*Proof.* Let  $b_0, c_0, c'_0$  be the simple closed geodesics in the free homotopy classes of  $b_1, c_1$  and  $c'_1$  respectively. Let

$$\hat{\mathcal{D}} = \{\hat{a}, \hat{b}_1, \hat{b}_2, \hat{c}_1, \hat{c}_2, \hat{d}\}$$

and

$$\hat{\mathcal{D}}' = \{\hat{a}', \hat{b}'_1, \hat{b}'_2, \hat{c}'_1, \hat{c}'_2, \hat{d}'\}$$

be the standard decompositions associated to the pairs  $(b_0, c_0)$  and  $(b_0, c'_0)$  respectively. From Proposition 5.4, the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\hat{\mathcal{D}}$  (resp.  $\mathcal{D}'$  and  $\hat{\mathcal{D}}'$ ) are related by an element of  $\Gamma$ .

By assumption, we have  $c_0 = \pm c'_0$ . From the construction of standard decompositions, we can assume that  $\hat{b}_1 = \hat{b}'_1$ ,  $\hat{b}_2 = \hat{b}'_2$ , and the corollary follows from Proposition 5.2. □

**5.3. Reducing the number of intersections.** By Proposition 5.4, we can now restrict ourselves into the case of standard decompositions. Our strategy for the proof of Theorem 5.1 is to find a sequence  $\{(g_1^j, g_2^j)\}_{0 \leq j \leq k}$  of pairs of disjoint simple closed geodesics in  $M$  satisfying  $(\mathcal{P})$  such that  $(g_1^0, g_2^0)$  is the pair associated to  $\mathcal{D}$ ,  $(g_1^k, g_2^k)$  is the pair associated to  $\mathcal{D}'$ , and for any  $0 < j \leq k$ , up to a renumbering, we have  $g_1^{j-1} = g_1^j$  and  $\text{Card}\{g_2^{j-1} \cap g_2^j\} = 1$ . We start by proving the following key lemma

**Lemma 5.7.** *Let  $g, g_1, g_2$  be three simple closed geodesics of  $M$  verifying Property  $(\mathcal{P})$ . Assume that*

$$g_1 \cap g_2 = \emptyset \quad \text{and} \quad \text{Card}\{g \cap (g_1 \cup g_2)\} = n > 1.$$

*Then there exists a simple closed geodesic  $g_3$  verifying Property  $(\mathcal{P})$  such that  $\text{Card}\{g_3 \cap (g_1 \cup g_2)\} = 1$ . Also, if  $g_3 \cap g_1 = \emptyset$ , then  $\text{Card}\{g \cap (g_1 \cup g_3)\} < \text{Card}\{g \cap (g_1 \cup g_2)\}$ . Moreover, for  $i = 1, 2$ , if  $g \cap g_i = \emptyset$ , then we can find  $g_3$  such that  $g_3 \cap g_i = \emptyset$ .*

*Proof.* We know that each of the curves  $g_1, g_2$  contains two Weierstrass points. Let  $W' \neq W$  be the other Weierstrass point of  $M$  which is not contained in  $g_1 \cup g_2$ . We have two possibilities:

*Case 1:*  $W' \in g$ . Let  $s$  be the segment of  $g$  which contains  $W'$  with endpoints in  $g \cap (g_1 \cup g_2)$ . We denote by  $Q_1, Q_2$  the two endpoints of  $s$  and choose the orientation of  $s$  to be from  $Q_1$  to  $Q_2$ . Since  $\tau(g) = -g$  and  $\tau(W') = W'$ , we deduce that  $\tau(s) = -s$  and  $Q_1, Q_2$  are interchanged by  $\tau$ . It follows that  $Q_1$  and  $Q_2$  are both contained in either  $g_1$  or  $g_2$ . Without loss of generality, we can assume that  $Q_1, Q_2$  are contained in  $g_2$ .

We know that  $\tau$  preserves the orientation of  $M$ , since  $\tau$  reverses the orientation of  $s$  and  $g_2$ , we deduce that  $s$  meets both sides of  $g_2$ . Let  $r$  be one of the two subsegments of  $g_2$  with endpoints  $Q_1, Q_2$ , and let  $W''$  be the Weierstrass point which is contained in  $r$ . Note that  $Q_1$  and  $Q_2$  must be distinct, otherwise  $s = g$  and  $\text{Card}\{g \cap (g_1 \cup g_2)\} = 1$ , which is discarded by the hypothesis.

Consider the simple closed curve  $g'_3$  which is composed by  $s$  and  $r$ . Note that we have  $\langle g'_3, g_2 \rangle = \pm 1$ , therefore  $g'_3 \neq 0$  in  $H_1(M, \mathbb{Z})$  and, in particular,  $g'_3$  is an essential, non-separating curve. By construction, we see that  $\tau(g'_3) = -g'_3$ . We

can move  $g'_3$  slightly in its free homotopy class so that the following conditions are satisfied:

- $g'_3 \cap s = \{W'\}$ ,
- $g'_3 \cap g_2 = \{W''\}$ ,
- $\tau(g'_3) = -g'_3$ .

By construction, we have

- $g'_3 \cap g_1 = \emptyset$ .
- $\text{Card}\{g'_3 \cap g\} = \text{Card}\{\text{int}(r) \cap g\} + 1 \leq \text{Card}\{g_2 \cap g\} - 2 + 1 = \text{Card}\{g_2 \cap g\} - 1$ .




Figure 15. Case  $W' \in g_3$ .

Let  $g_3$  be the simple closed geodesic in the free homotopy class of  $g'_3$ . Since  $\tau(g'_3) = -g'_3$ , we have  $\tau(g_3) = -g_3$ , as  $\tau(g_3)$  is the simple closed geodesic in the free homotopy class of  $\tau(g'_3)$ . It follows from Lemma 5.8 below that  $g_3$  does not contain  $W$ , therefore we can conclude that  $g_3$  verifies Property ( $\mathcal{P}$ ).

Let us now show that  $g_3$  satisfies the conditions in the conclusion of the lemma. Let  $\iota$  denote the geometric intersection number between free homotopy classes of simple closed curves in  $M$ . Recall that  $\iota(\alpha, \beta) = \text{Card}\{\alpha_0, \beta_0\}$ , where  $\alpha_0$  and  $\beta_0$  are the simple closed geodesics in the free homotopy classes of  $\alpha$  and  $\beta$  respectively (see [3], [5]). We have

- $\text{Card}\{g_3 \cap g_1\} = \iota(g'_3, g_1) = 0$ ,
- $\text{Card}\{g_3 \cap g_2\} = \iota(g'_3, g_2) \leq \text{Card}\{g'_3 \cap g_2\} = 1$ ,
- $\text{Card}\{g_3 \cap g\} = \iota(g'_3, g) \leq \text{Card}\{g'_3 \cap g\} < \text{Card}\{g_2 \cap g\}$ .

By construction, we have  $\langle g_3, g_2 \rangle = \langle g'_3, g_2 \rangle = \pm 1$ , therefore  $g_3 \cap g_2 \neq \emptyset$ . We deduce that  $\text{Card}\{g_3 \cap g_2\} = 1$ , and the lemma is proven for this case.

*Case 2:*  $W' \notin g$ . Cutting  $M$  along the curves  $g_1, g_2$ , we then get a 4-holed sphere  $N$  which is equipped with a hyperbolic metric with geodesic boundary. Let  $\hat{g}$  denote the union of geodesic arcs with endpoints in  $\partial N$  corresponding to sub-segments of  $g$  with endpoints in  $g_1 \cup g_2$ . Let  $\hat{s}_1$  be a geodesic arc realizing the distance  $\mathbf{d}_N(W', \hat{g})$ . Note that  $\hat{s}_1$  does not meet  $\partial N$ . The involution  $\tau$  of  $M$  induces an involution on  $N$ , which will be denoted by  $\tau_N$ . Let  $\hat{s}_2$  denote the image of  $\hat{s}_1$  by  $\tau_N$ . Note that  $\hat{s}_2$  is also a geodesic arc realizing the distance  $\mathbf{d}_N(W', \hat{g})$ .

From the fact that both  $\hat{s}_1, \hat{s}_2$  realize the distance in  $N$  from  $W'$  to  $\hat{g}$ , we derive that  $\hat{s}_1 \cap \hat{s}_2 = \{W'\}$ , since if any other common point exists, it must be the other common endpoint in  $\hat{g}$  of both segments, hence it is fixed by  $\tau_N$ . But  $\tau_N$  has only two fixed points in  $N$  corresponding to  $W$  and  $W'$ , and by assumption  $W, W' \notin \hat{g}$ , therefore we get a contradiction. Since  $\tau$  acts like  $-\text{Id}$  on the tangent plane at  $W'$ ,  $\hat{s} = \hat{s}_1 \cup \hat{s}_2$  is in fact a geodesic arc.

Let  $s$  be the geodesic arcs in  $M$  corresponding to  $\hat{s}$ , and let  $Q_i, i = 1, 2$ , denote the endpoint of  $s$ . By construction, we have  $\text{int}(s) \cap (g_1 \cup g_2) = \emptyset, W \notin s$  and  $\tau(s) = -s$ . Let  $R_1$  be an endpoint of the subarc of  $g$  (with endpoints in  $g_1 \cap g_2$ ) that contains  $Q_1$ , and let  $r_1$  denote the oriented subarc from  $Q_1$  to  $R_1$ . We denote by  $R_2$  and  $r_2$  the images of  $R_1$  and  $r_1$  by  $\tau$  respectively. The curve  $c = (-r_1)*(-s_1)*s_2*r_2$  is then a simple arc joining  $R_1$  to  $R_2$  verifying  $\tau(c) = -c$ .

Since  $\tau(R_1) = R_2$ , it follows that either  $\{R_1, R_2\} \subset g_1$ , or  $\{R_1, R_2\} \subset g_2$ . Without loss of generality, we can assume that  $R_1, R_2$  are contained in  $g_2$ , then the same argument as in Case 1 shows that  $c$  meets both sides of  $g_2$ . Here we have two issues

- $R_1 = R_2$ : in this case  $c$  is actually a simple closed curve which satisfies
  - (a)  $\tau(c) = -c$ ,
  - (b)  $c \cap g_1 = \emptyset$ ,
  - (c)  $\text{Card}\{c \cap g_2\} = 1$ .

We can then find in the neighborhood of  $c$  a simple closed curve  $c'$ , freely homotopic to  $c$ , satisfying (a), (b), (c) such that  $c' \cap g = \{R_1\}$ . Let  $g'_3$  be the image of  $c'$  by the Dehn twist about  $g_2$ , observe that  $g'_3$  satisfies (a), (b), (c), and  $\text{Card}\{g \cap g'_3\} = \text{Card}\{g \cap g_2\} - 1$  (see Figure 16). We denote by  $g_3$  the simple closed geodesic in the free homotopy class of  $g'_3$ .

- $R_1 \neq R_2$ : let  $d$  be one of the two simple arcs in  $g_2$  with endpoints  $R_1, R_2$ , then  $c \cup d$  is a simple closed curve invariant under  $\tau$ . We can choose  $d$  in such a way that there exists in the neighborhood of  $c \cup d$  a simple closed curve  $g'_3$  which satisfies (a), (b), (c), and  $\text{Card}\{g \cap g'_3\} \leq \text{Card}\{g \cap g_2\} - 2$ . We then take  $g_3$  to be the closed geodesic in the free homotopy class of  $g'_3$ .

In both cases, using the same arguments as in Case 1, we see that  $g_3$  verifies the required properties. It is also clear from the arguments above that, if in addition, we have  $g \cap g_1 = \emptyset$ , then  $g_3 \cap g_1 = \emptyset$ . The proof of the lemma is now complete.  $\square$




Figure 16. Case  $W' \notin g$  and  $R_1 = R_2$ .

**Lemma 5.8.** *Let  $c$  be a simple closed curve in  $M$  such that  $\tau(c) = -c$  and  $W \notin c$ . Let  $c_0$  be the simple closed geodesic in the free homotopy class of  $c$ . Then we also have  $\tau(c_0) = -c_0$  and  $W \notin c_0$ .*

*Proof.* Since  $\tau(c) = -c$ , the image of  $c$  by the ramified covering  $\rho: M \rightarrow \mathbb{C}\mathbb{P}^1$  is a simple arc  $\bar{c}$  with endpoints in  $\{P_1, \dots, P_5\}$  such that  $\text{int}(\bar{c}) \cap \{P_0, \dots, P_5\} = \emptyset$ . Assume that  $P_1, P_2$  are the endpoints of  $\bar{c}$ . Let  $\tilde{c}$  be a simple closed curve in  $\mathbb{C}\mathbb{P}^1$  passing through  $P_0 = \rho(W)$ , which bounds a disc  $D$  containing  $\bar{c}$  such that  $\text{int}(D) \cap \{P_0, \dots, P_5\} = \{P_1, P_2\}$ . Let  $\tilde{d}$  be a simple arc in  $D$  which joins  $P_0$  to an endpoint of  $\bar{c}$ . We have

- $\rho^{-1}(\tilde{c})$  is the union of two simple closed curve  $c_1, c_2$ , freely homotopic to  $c$ , such that  $c_1 \cap c_2 = \{W\}$ ,
- $\rho^{-1}(\tilde{d})$  is a simple closed curve  $d$  passing through  $W$  such that  $\tau(d) = -d$ ,
- $\rho^{-1}(\text{int}(D))$  is an open annulus bounded by  $c_1, c_2$  which contains  $d \setminus \{W\}$ .

Therefore, in the group  $\pi_1(M, W)$ , we have  $[\tau(c_1)] = [c_2]^{-1} = [d][c_1]^{-1}[d]^{-1}$ ,  $[\tau(d)] = [d]^{-1}$ . Clearly, in the homology level, we have  $\langle c, d \rangle = \pm 1$ . The lemma then follows from the same arguments as in Lemma 5.3 b) and c).  $\square$

Using Lemma 5.7, we can now prove

**Proposition 5.9.** *Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  and  $\mathcal{D}' = \{a', b'_1, b'_2, c'_1, c'_2, d'\}$  be two admissible decompositions for  $(M, W)$ . As usual, let  $(a, b, c, e)$  and  $(a', b', c', e')$  be the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\mathcal{D}'$  respectively. Assume that  $b_1$  is freely homotopic to  $b'_1$ , then there exists  $\gamma \in \Gamma$  such that*

$$\begin{pmatrix} a' \\ b' \\ c' \\ e' \end{pmatrix} = \gamma \cdot \begin{pmatrix} a \\ b \\ c \\ e \end{pmatrix}.$$

*Proof.* Let  $b_0, c_0, c'_0$  be the simple closed geodesics in the free homotopy classes of  $b_1, c_1, c'_1$  respectively. According to Lemma 5.5, we know that  $b_0, c_0, c'_0$  verify Property ( $\mathcal{P}$ ) and  $b_0 \cap c_0 = b_0 \cap c'_0 = \emptyset$ .

If  $c'_0 = \pm c_0$ , the proposition follows from Corollary 5.6. Hence, we only need to consider the case where  $c'_0 \neq c_0$  as subsets of  $M$ . In this case, since each of the curves  $b_0, c_0, c'_0$  contains exactly two Weierstrass points of  $M$  and  $W \notin b_0 \cup c_0 \cup c'_0$ , we deduce that  $c'_0 \cap c_0 \neq \emptyset$ . Let  $n$  be the number of intersections of  $c_0$  and  $c'_0$ . The proposition will be proved by induction.

*Case  $n = 1$ .* Note that in this case, the intersection of  $c_0$  and  $c'_0$  must be a Weierstrass point. We will show that there exist two admissible decompositions  $\{\hat{a}, \hat{b}_1, \hat{b}_2, \hat{c}_1, \hat{c}_2, \hat{d}\}$  and  $\{\hat{a}', \hat{b}'_1, \hat{b}'_2, \hat{c}'_1, \hat{c}'_2, \hat{d}'\}$  such that  $\hat{b}_i = \hat{b}'_i$  and  $\hat{b}_i, \hat{c}, \hat{c}'$  are freely homotopic to  $b_0, c_0$  and  $c'_0$ , respectively. We can then use Proposition 5.2 to conclude.

Observe that  $\bar{b}_0 = \rho(b_0), \bar{c}_0 = \rho(c_0), \bar{c}'_0 = \rho(c'_0)$  are three simple arcs on  $\mathbb{C}\mathbb{P}^1$ , which satisfy

- $P_0 \notin (\bar{b}_0 \cup \bar{c}_0 \cup \bar{c}'_0)$ .
- $\bar{b}_0 \cap (\bar{c}_0 \cup \bar{c}'_0) = \emptyset$ .
- $\bar{c}_0$  and  $\bar{c}'_0$  have a common endpoint and  $\text{Card}\{\bar{c}_0 \cap \bar{c}'_0\} = 1$ .

We choose the numbering of  $\{P_1, \dots, P_5\}$  such that  $P_1$  and  $P_2$  are the endpoints of  $\bar{b}_0$ ,  $P_3$  and  $P_4$  are the endpoints of  $\bar{c}_0$  and  $P_4$  and  $P_5$  are the endpoints of  $\bar{c}'_0$ . Let  $\tilde{b}$  be a simple closed curve in  $\mathbb{C}\mathbb{P}^1$ , passing through  $P_0$ , which separates  $\bar{b}_0$  from  $\bar{c}_0 \cup \bar{c}'_0$  (see Figure 17). The pre-image of  $\tilde{b}$  in  $M$  is the union of two simple closed curves passing through  $W$ , denoted by  $\hat{b}_1$  and  $\hat{b}_2$ , which bound an open annulus containing  $b_0$ . Let  $\tilde{c}$  (resp.  $\tilde{c}'$ ) be a simple closed curve in  $\mathbb{C}\mathbb{P}^1$  passing through  $P_0$  surrounding  $\bar{c}_0$  (resp.  $\bar{c}'_0$ ) as shown in Figure 17. Like  $\rho^{-1}(\tilde{b}), \rho^{-1}(\tilde{c})$  (resp.  $\rho^{-1}(\tilde{c}')$ ) is the union of two simple closed curves  $\hat{c}_1, \hat{c}_2$  (resp.  $\hat{c}'_1, \hat{c}'_2$ ) which bound an embedded open annulus in  $M$  containing  $c_0$  (resp.  $c'_0$ ). We denote by  $D_1, D_2$  and  $D'_2$  the open disks in  $\mathbb{C}\mathbb{P}^1$  bounded by  $\tilde{b}, \tilde{c}$  and  $\tilde{c}'$  which contain  $\bar{b}_0, \bar{c}_0$  and  $\bar{c}'_0$ , respectively.




Figure 17. Case  $b_0 = b'_0, \text{Card}\{c_0 \cap c'_0\} = 1$ .

We equip  $\hat{b}_1$  and  $\hat{b}_2$  with the orientation induced by  $b_0$  (via free homotopy). The numbering is chosen such that the orientation of  $\hat{b}_2$  is also the one induced by the orientation of the annulus  $\rho^{-1}(D_1)$ . Observe that the family of curves  $\{\hat{b}_1, \hat{b}_2, \hat{c}_1, \hat{c}_2\}$  (resp.  $\{\hat{b}_1, \hat{b}_2, \hat{c}'_1, \hat{c}'_2\}$ ) cuts  $M$  into two open annulus and one open quadrilateral, which will be denoted by  $Q$  (resp.  $Q'$ ). We will choose the numbering and the orientation of  $\hat{c}_1, \hat{c}_2$  (resp.  $\hat{c}'_1, \hat{c}'_2$ ) such that  $\hat{b}_1 * \hat{c}_1 * (-\hat{b}_2) * (-\hat{c}_2)$  (resp.  $\hat{b}_1 * \hat{c}'_1 * (-\hat{b}_2) * (-\hat{c}'_2)$ ) is the boundary of  $Q$  (resp.  $Q'$ ) with the induced orientation. Remark that it can happen that  $\hat{c}_i$  belongs to the free homotopy class of  $-c_0$  and  $\hat{c}'_i$  belongs to the free homotopy class of  $-c'_0$ .

We can then add to these families some simple closed curves to obtain two admissible decompositions for  $(M, W)$  as follows: let  $\tilde{a}$  be a simple arc in  $D_1$  joining  $P_0$  to an endpoint of  $\tilde{b}_0$ , and let  $\tilde{d}$  (resp.  $\tilde{d}'$ ) denote a simple arc in  $D_2$  (resp. in  $D'_2$ ) joining  $P_0$  to an endpoint of  $\tilde{c}_0$  (resp. of  $\tilde{c}'_0$ ). Set  $\hat{a} = \rho^{-1}(\tilde{a})$ ,  $\hat{d} = \rho^{-1}(\tilde{d})$  and  $\hat{d}' = \rho^{-1}(\tilde{d}')$ . By choosing appropriate orientations for  $\hat{a}, \hat{d}$ , and  $\hat{d}'$ , we see that  $\hat{\mathcal{D}} = \{\hat{a}, \hat{b}_1, \hat{b}_2, \hat{c}_1, \hat{c}_2, \hat{d}\}$  and  $\hat{\mathcal{D}}' = \{\hat{a}, \hat{b}_1, \hat{b}_2, \hat{c}'_1, \hat{c}'_2, \hat{d}'\}$  are two admissible decompositions for  $(M, W)$ .

Clearly, by construction, we have  $\hat{c}_i$  (resp.  $\hat{c}'_i$ ) is freely homotopic to  $\pm c_0$  (resp. to  $\pm c'_0$ ). Therefore, by Corollary 5.6, the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\hat{\mathcal{D}}$  are related by an element of  $\Gamma$ . Similarly, the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}'$  and  $\hat{\mathcal{D}}'$  are also related by an element of  $\Gamma$ . Now, by Proposition 5.2, we know that the symplectic bases associated to  $\hat{\mathcal{D}}$  and  $\hat{\mathcal{D}}'$  are related by an element of  $\Gamma$ . Hence, the proposition is proven for this case.

*Case  $n > 1$ :* By Lemma 5.7, there exists a simple closed geodesic  $c''_0$  verifying Property ( $\mathcal{P}$ ) such that

- $\text{Card}\{c''_0 \cap b_0\} = 0,$
- $\text{Card}\{c''_0 \cap c_0\} = 1,$
- $\text{Card}\{c'_0 \cap c''_0\} < \text{Card}\{c'_0 \cap c_0\} = n.$

Let  $\mathcal{D}'' = \{a'', b''_1, b''_2, c''_1, c''_2, d''\}$  be the standard decomposition associated to the pair  $(b_0, c''_0)$ . The arguments in Case  $n = 1$  show that the symplectic bases associated to  $\mathcal{D}''$  and  $\mathcal{D}$  are related by an element of  $\Gamma$ . Now, since  $\text{Card}\{c'_0 \cap c''_0\} < n$ , the induction hypothesis implies that the symplectic bases associated to  $\mathcal{D}''$  and  $\mathcal{D}'$  are also related by an element of  $\Gamma$ , and the proposition follows.  $\square$

**5.4. Proof of Theorem 5.1.** We can now give the proof of Theorem 5.1. Let  $b_0, c_0$  be the simple closed geodesics in the free homotopy classes of  $b_1$  and  $c_1$  respectively. Note that the roles of  $b_1$  and  $c_1$  are interchanged by the  $S$  move, and the orientation of  $b_1$  is irrelevant because  $-\text{Id} = S^2 \in \Gamma$ . Let  $b'_0$  denote the simple closed geodesic in the free homotopy class of  $b'_1$ . If  $b'_0 = \pm b_0$ , or  $b'_0 = \pm c_0$ , then Proposition 5.9 allows us to conclude immediately. Assume that  $b'_0 \neq \pm b_0$  and  $b'_0 \neq \pm c_0$ . Let  $n$  be the number of intersections of  $b'_0$  and  $b_0 \cup c_0$ . We first remark that  $n > 0$ ,



since each of the curves  $b_0, c_0, b'_0$  contains two Weierstrass points, and by definition  $W \notin (b_0 \cup c_0 \cup b'_0)$ , but there are only six Weierstrass points on  $M$ . We proceed by induction.

*Case  $n = 1$ :* in this case, we can suppose that  $b'_0 \cap b_0 = \emptyset$  and  $\text{Card}\{b'_0 \cap c_0\} = 1$ . Let  $\mathcal{D}''$  be the standard decomposition associated to the pair  $(b_0, b'_0)$ . By Proposition 5.9, we know that the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and to  $\mathcal{D}''$  are related by an element of  $\Gamma$ , and the symplectic bases associated to  $\mathcal{D}'$  and  $\mathcal{D}''$  are also related by an element of  $\Gamma$ , Theorem 5.1 is then proven for this case.

*Case  $n > 1$ :* by Lemma 5.7, there exists a simple closed geodesic  $g$  verifying Property ( $\mathcal{P}$ ) such that

- $\text{Card}\{g \cap (b_0 \cup c_0)\} = 1$ ,
- $\text{Card}\{b'_0 \cap g\} < \text{Card}\{b'_0 \cap c_0\}$  if  $g \cap b_0 = \emptyset$ , and  $\text{Card}\{b'_0 \cap g\} < \text{Card}\{b'_0 \cap b_0\}$  if  $g \cap c_0 = \emptyset$ .

Without loss of generality, we can assume that  $g \cap b_0 = \emptyset$ . Let  $\mathcal{D}''$  be the standard decomposition associated to the pair of geodesics  $(b_0, g)$ . From Proposition 5.9, we know that the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}$  and  $\mathcal{D}''$  are related by an element of  $\Gamma$ . Since  $\text{Card}\{b'_0 \cap (b_0 \cup g)\} < \text{Card}\{b'_0 \cap (b_0 \cup c_0)\}$ , by the induction hypothesis, the symplectic bases of  $H_1(M, \mathbb{Z})$  associated to  $\mathcal{D}''$  and  $\mathcal{D}'$  are related by an element  $\Gamma$ . Theorem 5.1 is then proven.  $\square$

## 6. Proof of Theorem 1.1

**6.1. The map  $\Xi$ .** Let  $\mathcal{M}$  denote the quotient  $\mathcal{H}(2)/\mathbb{C}^*$ . We define the map  $\Xi$  from  $\mathcal{M}$  to  $\Gamma \backslash \mathcal{S}_2$  as follows: given a pair  $(M, W)$  in  $\mathcal{M}$ , we associate to  $(M, W)$  the  $\Gamma$ -orbit of the period matrix of the symplectic homology basis associated to an admissible decomposition for  $(M, W)$ . It follows from Theorem 5.1 that the map  $\Xi$  is well defined. We will show that  $\Xi$  is a homeomorphism from  $\mathcal{H}(2)/\mathbb{C}^*$  and  $\Gamma \backslash \mathcal{F}_2$ , which implies Theorem 1.1.

**6.2. Injectivity of  $\Xi$ .** Let  $(M, W)$  and  $(M', W')$  be two pairs in  $\mathcal{M}$ . Assume that  $M$  and  $M'$  are defined by the equations  $w^2 = \prod_{i=0}^5 (z - \lambda_i)$  and  $w^2 = \prod_{i=0}^5 (z - \lambda'_i)$  so that  $W$  and  $W'$  correspond to  $\lambda_0$  and  $\lambda'_0$  respectively. Let  $\Pi$  (resp.  $\Pi'$ ) be the period matrix of the symplectic basis associated to an admissible decomposition  $\mathcal{D}$  (resp.  $\mathcal{D}'$ ) for the pair  $(M, W)$  (reps.  $(M', W')$ ). Assume that there exists an element  $\gamma$  of  $\Gamma$  such that  $\tilde{\Pi}' = \gamma \cdot \Pi$ . By Lemma 4.2, we know that there exists an admissible decomposition  $\tilde{\mathcal{D}}$  for the pair  $(M, W)$  such that the symplectic homology bases of  $M$  associated to  $\tilde{\mathcal{D}}$  and  $\mathcal{D}$  are related by  $\gamma$ . It follows that the period matrix of the basis associated to  $\tilde{\mathcal{D}}$  is equal to  $\Pi'$ .

Using an element of  $\mathrm{PSL}(2, \mathbb{C})$  we can assume that  $\lambda_1 = \lambda'_1 = 0, \lambda_2 = \lambda'_2 = 1$ , and  $\lambda_0 = \lambda'_0 = \infty$ . Then from Theorem 2.2, we see that the values of  $\lambda_i$  and  $\lambda'_i$  ( $i = 3, 4, 5$ ) can be computed by the same theta functions, with the same period matrix. Therefore we have  $\lambda_i = \lambda'_i, i = 3, 4, 5$ , and it follows that there exists a conformal homeomorphism  $\phi: M \rightarrow M'$  such that  $\phi(W) = W'$ .

**6.3. Surjectivity of  $\Xi$ .** Let  $\tilde{\Pi}$  be a matrix in  $\mathcal{J}_2$ , we will show that there exists a pair  $(M, W)$  in  $\mathcal{M}$  such that  $\Xi((M, W)) = \Gamma \cdot \tilde{\Pi}$ . Since  $\tilde{\Pi} \in \mathcal{J}_2$ , there exists a Riemann surface of genus two  $M$  and a symplectic homology basis whose period matrix is  $\tilde{\Pi}$ . Let  $W_0$  be a Weierstrass point of  $M$ , and let  $\mathcal{D}$  be an admissible decomposition for the pair  $(M, W_0)$ . Let  $\Pi$  be the period matrix of the symplectic homology basis of  $M$  associated to  $\mathcal{D}$ . By definition, there exists an element  $A \in \mathrm{Sp}(4, \mathbb{Z})$  such that  $\tilde{\Pi} = A \cdot \Pi$ . According to Lemma 4.3, there exists an admissible decomposition  $\tilde{\mathcal{D}}$  for a pair  $(M, W)$ , where  $W$  is also a Weierstrass point of  $M$ , such that the symplectic homology bases associated to  $\mathcal{D}$  and  $\tilde{\mathcal{D}}$  is related by  $A$ . Consequently, the period matrix of the basis associated to  $\tilde{\mathcal{D}}$  is  $\tilde{\Pi}$  and by definition  $\Gamma \cdot \tilde{\Pi} = \Xi((M, W))$ .

**6.4. Continuity of  $\Xi$ .** To prove the continuity of  $\Xi$  we will consider the inverse map  $\Xi^{-1}: \Gamma \backslash \mathcal{J}_2 \rightarrow \mathcal{M}$ . Let  $\Pi$  be a matrix in  $\mathcal{J}_2$ , then  $\Pi$  is the period matrix of the symplectic homology basis associated to an admissible decomposition for a pair  $(M, W)$  in  $\mathcal{M}$ . There exist complex numbers  $\{\lambda_0, \lambda_1, \dots, \lambda_5\}$  such that  $M$  is the surface defined by the equation  $w^2 = \prod_{i=0}^5 (z - \lambda_i)$ . We can assume that  $W$  is the Weierstrass point corresponding to  $\lambda_0$ . A neighborhood of  $\Pi$  in  $\mathcal{S}_2$  consists of period matrices of the same symplectic homology basis on Riemann surfaces close to  $M$ .

Using  $\mathrm{PSL}(2, \mathbb{C})$ , we can assume that  $\lambda_0 = 0$ , it follows that  $\omega = \frac{zdz}{w}$  is a holomorphic 1-form with double zero at  $W$ . Let  $\mathcal{D} = \{a, b_1, b_2, c_1, c_2, d\}$  be an admissible decomposition for the pair  $(M, W)$ . Let  $(a, b, c, e)$  be the symplectic homology basis associated to  $\mathcal{D}$ , then the map

$$\Phi: \mathcal{U} \rightarrow \mathbb{C}^4, \quad (M, \omega) \mapsto \left( \int_a \omega, \int_b \omega, \int_c \omega, \int_e \omega \right),$$

is a local chart for  $\mathcal{H}(2)$  in the neighborhood  $\mathcal{U}$  of  $(M, \omega)$ . Let  $\rho: M \rightarrow \mathbb{CP}^1$  be the two-sheeted branched cover from  $M$  onto  $\mathbb{CP}^1$ . Recall that by construction,  $\rho(a) = \bar{a}, \rho(b) = \bar{b}_*, \rho(c) = \bar{c}_*, \rho(e) = \bar{e}$  are simple arcs in  $\mathbb{CP}^1$  with endpoints in  $\{\lambda_0, \dots, \lambda_5\}$ . We have

$$\int_a \omega = 2 \int_{\bar{a}} \frac{zdz}{w}, \quad \int_b \omega = 2 \int_{\bar{b}_*} \frac{zdz}{w}, \quad \int_c \omega = 2 \int_{\bar{c}_*} \frac{zdz}{w}, \quad \int_e \omega = 2 \int_{\bar{e}} \frac{zdz}{w}.$$

Clearly, the integrals of  $\frac{zdz}{w}$  along  $\bar{a}, \bar{b}_*, \bar{c}_*, \bar{e}$  depend continuously on  $(\lambda_1, \dots, \lambda_5)$ . Since  $\lambda_i$  can be computed from  $\Pi$  by some theta functions, we get a continuous map  $\Psi$  from a neighborhood of  $\Pi$  in  $\mathcal{J}_2$  into  $\mathbb{C}^4$ . Now, in a neighborhood of  $\Pi$ , the map

$\Xi^{-1}$  is the composition of  $\Psi$  and the natural projection  $\mathbb{C}^4 \setminus \{0\} \rightarrow \mathbb{C}\mathbb{P}^3$ . It follows immediately that  $\Xi^{-1}$  is continuous. Since  $\dim_{\mathbb{C}} \mathcal{J}_2 = \dim_{\mathbb{C}} \mathcal{M} = 3$ , we conclude that  $\Xi$  is a homeomorphism.

**6.5.  $[\mathrm{Sp}(4, \mathbb{Z}) : \Gamma] = 6$ .** We have a natural projection from  $\mathcal{M}$  onto the moduli space of closed Riemann surface of genus two  $\mathfrak{M}_2$ , which is homeomorphic to  $\mathrm{Sp}(4, \mathbb{Z}) \backslash \mathcal{J}_2$ , by associating to any element  $(M, W)$  of  $\mathcal{M}$  the point  $M$  in  $\mathfrak{M}_2$ . Every Riemann surface of genus two has six Weierstrass points, and the group of automorphisms of a generic one contains exactly two elements, the identity and the hyperelliptic involution, both fix all the Weierstrass points. Therefore, the pre-image of a generic point in  $\mathfrak{M}_2$  contains exactly six points. Note that  $-\mathrm{Id}_4 \in \mathrm{Sp}(4, \mathbb{Z})$  acts trivially on  $\mathfrak{S}_2$  and the action of  $\mathrm{Sp}(4, \mathbb{Z}) / \{\pm \mathrm{Id}_4\}$  on  $\mathfrak{S}_2$  is effective. Since  $\{\pm \mathrm{Id}_4\} \subset \Gamma$ , we derive  $[\mathrm{Sp}(4, \mathbb{Z}) : \Gamma] = 6$ . The proof Theorem 1.1 is now complete.  $\square$

## Appendices

### A. Existence of parallelogram decompositions

Recall that on a translation surface  $(M, \omega)$ , a *saddle connection* is a geodesic segment whose endpoints are singularities of the flat metric (the endpoints need not to be distinct). A *cylinder* is a subset of  $M$  which is isometric to  $\mathbb{R} \times ]0, h[ / \mathbb{Z}$  and maximal with respect to this property, where the action of  $\mathbb{Z}$  is generated by  $(x, y) \mapsto (x + \ell, y)$ . The boundary of a cylinder  $C$  is the set  $\bar{C} \setminus C$ , this boundary is a union of saddle connections. We say that  $C$  is a *simple cylinder* if the boundary of  $C$  is the union of only two saddle connections.

**Proposition A.1** (Existence of parallelogram decompositions). *For any translation surface  $(M, \omega)$  in  $\mathcal{H}(2)$ , there always exists a parallelogram decomposition on  $M$ .*

*Proof.* We first prove that there always exists a simple cylinder in  $M$ . By a well known theorem of Masur (see [8], or [9]), there always exists a closed geodesic  $\gamma$  in  $M$ . Let  $C_\gamma$  be the (open) cylinder consisting of closed geodesics freely homotopic to  $\gamma$ . Since  $M$  is of genus two, and has only one singular points, we have three cases:

- a)  $C_\gamma$  is a simple cylinder.
- b)  $M \setminus \bar{C}_\gamma$  is a simple cylinder.
- c)  $\bar{C}_\gamma = M$ .

If *a*) or *b*) occurs then we are done. If we are in case *c*), then the pair  $(M, \omega)$  is obtained from a single parallelogram as shown in Figure 18. Therefore, in this situation, one can easily find a simple cylinder in another direction.




Figure 18. Surface admitting a decomposition in one cylinder.

**Remark.** This argument has been used in the proof of Theorem 7.1 in [10].

Let  $C_1$  be a simple cylinder in  $M$ . Let  $b_1, b_2$  denote the two saddle connections which bound  $C_1$ . Cutting off  $C_1$  from  $M$  along  $b_1$  and  $b_2$ , we get a torus minus two open disks whose boundaries meet at one point, which is the unique singular point  $p$  of  $M$ . Splitting the point  $p$  into two points, we obtain then a flat torus with an open disk removed whose boundary is the union of two parallel segments of same length. Gluing these two segments together, we obtain a flat torus  $M_1$  with a marked simple geodesic arc  $b$ . Denote by  $p_1, p_2$  the endpoints of  $b$  in  $M_1$ . Let us show that there always exist two parallel closed geodesic  $c_1, c_2$  in  $M_1$  such that  $c_1 \cap b = \{p_1\}$ , and  $c_2 \cap b = \{p_2\}$ .

Choose a direction  $\theta \in [0; 2\pi]$  which is not parallel to  $b$  and consider the geodesic flow  $\psi_\theta$  of  $M_1$  in this direction. Let  $t_0$  be the smallest value of  $t$  such that  $t_0 > 0$  and  $\psi_\theta^{t_0}(b) \cap b \neq \emptyset$ . The value  $t_0$  must be finite because otherwise the area of stripe  $S_t = \bigcup_{0 \leq s \leq t} \psi_\theta^s(b)$  tends to infinity as  $t \rightarrow \infty$ , which is impossible.

Since  $\psi_\theta^{t_0}(b)$  and  $b$  are parallel and have the same length,  $\psi_\theta^{t_0}(b)$  contains at least one endpoint of  $b$ . Without loss of generality, we can assume that  $\psi_\theta^{t_0}(b)$  contains  $p_1$ . Remark that the stripe  $S_{t_0} = \bigcup_{0 \leq s \leq t_0} \psi_\theta^s(a)$  is the image of a parallelogram  $P$  under an isometric immersion  $\varphi: P \rightarrow M$  such that the restriction of  $\varphi$  to  $\text{int}(P)$  is injective, and  $b$  and  $\psi_\theta^{t_0}(b)$  are the images of two opposite sides of  $P$ . By assumption,  $\varphi^{-1}(p_1)$  consists of two points, which belong to two opposite sides of  $P$ . It follows that the image of the segment in  $P$  joining these two points by  $\varphi$  is a closed geodesic  $c_1$  on  $M_1$  which satisfies  $c_1 \cap b = \{p_1\}$ . Consequently, the closed geodesic  $c_2$  passing through  $p_2$  and parallel to  $c_1$  also verifies  $c_2 \cap b = \{p_2\}$ .

By construction, we can identify  $M_1 \setminus b$  with the complement of  $\bar{C}_1$  in  $M$ . By this identification, the closed geodesics  $c_1$  and  $c_2$  correspond to two saddle connections which bound an open cylinder  $C_2$  disjoint from  $C_1$ , and the complement of  $\bar{C}_1 \cup \bar{C}_2$  is an open disk isometric to an open parallelogram in  $\mathbb{R}^2$ .

Let  $a$  (resp.  $d$ ) be a saddle connection contained in  $\bar{C}_1$  (resp.  $\bar{C}_2$ ) which intersects all the closed geodesics parallel to  $b_1$  and  $b_2$  (resp. parallel to  $c_1$  and  $c_2$ ). One can easily check that the family  $\{a, b_1, b_2, c_1, c_2, d\}$ , with appropriate orientations, is a parallelogram decomposition of  $M$ . □

It follows from this proposition that, on any surface  $(M, \omega)$  in  $\mathcal{H}(2)$ , there exist infinitely many parallelogram decompositions, since if we have one, we can get infinitely many others by using elementary moves  $T, S, R$ . It is also possible to show that any two parallelogram decompositions are related by elementary moves.

### B. Proof of Lemma 3.4

For any  $g \geq 1$ , let  $\sigma$  be the permutation of  $\{1, 2, \dots, 2g\}$  that transpose  $2i$  and  $2i - 1$ , for  $i = 1, \dots, g$ . The *elementary symplectic matrices* are the matrices

$$E_{ij} = \begin{cases} \text{Id}_{2g} + e_{ij} & \text{if } i = \sigma(j); \\ \text{Id}_{2g} + e_{ij} - (-1)^{i+j} e_{\sigma(j)\sigma(i)} & \text{otherwise,} \end{cases}$$

where  $i \neq j$ , and  $e_{ij}$  is the matrix whose the  $(i, j)$ -th entry is 1 and all other entries are 0. It is a classical fact that  $\text{Sp}(2g, \mathbb{Z})$  is generated by elementary symplectic matrices ([3], Chapter 7). For the case  $g = 2$ , we have

$$\begin{aligned} \bullet E_{12} = E'_{21} &= \begin{pmatrix} 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & E_{34} = E'_{43} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \\ \bullet E_{13} = E^{-1}_{42} &= \begin{pmatrix} 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}, & E_{31} = E^{-1}_{24} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & -1 \\ 1 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \\ \bullet E_{14} = E_{32} &= \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, & E_{41} = E_{23} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 1 & 0 & 0 & 1 \end{pmatrix}. \end{aligned}$$

All we need is to verify that  $E_{ij}$  ( $i \neq j$ ) is contained in the group  $\Gamma'$  generated by  $\{T, R, S, U\}$ . It is clear that  $E_{12}, E_{34}, E_{43}$  belong to  $\Gamma \subset \Gamma'$ . We have

- $E_{21} = U^{-1}T^{-1}U \in \Gamma'$ .
- $E_{13} = SU \in \Gamma'$ .

Since  $\Gamma$  contains

$$\begin{pmatrix} \text{Id}_2 & 0 \\ 0 & \text{SL}(2, \mathbb{Z}) \end{pmatrix},$$

we have

$$S_1 = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix} \in \Gamma,$$

therefore

$$S_2 = U^{-1}S_1U = \begin{pmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \in \Gamma'.$$

It follows that

- $E_{31} = (S_1S_2)E_{13}^{-1}(S_1S_2)^{-1} \in \Gamma'$ ,
- $E_{14} = S_1E_{13}S_1^{-1} \in \Gamma'$ ,
- $E_{41} = S_2E_{13}S_2^{-1} \in \Gamma'$ .

The lemma is then proven. □

### C. A family of $\Gamma$ right cosets in $\text{Sp}(4, \mathbb{Z})$

In this section, we give a partition of the group  $\text{Sp}(4, \mathbb{Z})$  into  $\Gamma$  right cosets. Recall that  $\Gamma$  is not a normal subgroup of  $\text{Sp}(4, \mathbb{Z})$  (see Lemma 3.3), and the group  $\text{Sp}(4, \mathbb{Z})$  is generated by the matrices  $T, S, R$  and  $U$  (see Lemma 3.4). Set

$$\mathcal{F} = \{\Gamma, U \cdot \Gamma, RU \cdot \Gamma, SRU \cdot \Gamma, URU \cdot \Gamma, USRU \cdot \Gamma\}.$$

By Lemma 3.3, we know that the action of  $\Gamma$  on  $(\mathbb{Z}/2\mathbb{Z})^4 \setminus \{0\}$  has two orbits  $\mathcal{O}_1$  and  $\mathcal{O}_2$ , therefore we have a simple criterion to show that an element of  $\text{Sp}(4, \mathbb{Z})$  does *not* belong to  $\Gamma$ . Consequently, it is easy to verify that the elements in the family  $\mathcal{F}$  are all distinct.

We will also determine explicitly the action of  $T^{\pm 1}, R^{\pm 1}, S^{\pm 1}, U^{\pm 1}$  on  $\mathcal{F}$  by multiplication from the left. Note that, since  $S^{-1} = -S$  (resp.  $U^{-1} = -U$ ) and  $-\text{Id}_4 \in \Gamma$ , the actions of  $S$  and  $S^{-1}$  (resp.  $U$  and  $U^{-1}$ ) are identical. Details of the calculations are lengthy and uninteresting, hence will be omitted. The final result is resumed in the following table.

	$\Gamma$	$U \cdot \Gamma$	$RU \cdot \Gamma$	$SRU \cdot \Gamma$	$URU \cdot \Gamma$	$USRU \cdot \Gamma$
$T$	$\Gamma$	$U \cdot \Gamma$	$RU \cdot \Gamma$	$URU \cdot \Gamma$	$SRU \cdot \Gamma$	$USRU \cdot \Gamma$
$R$	$\Gamma$	$RU \cdot \Gamma$	$U \cdot \Gamma$	$SRU \cdot \Gamma$	$URU \cdot \Gamma$	$USRU \cdot \Gamma$
$S$	$\Gamma$	$U \cdot \Gamma$	$SRU \cdot \Gamma$	$RU \cdot \Gamma$	$USRU \cdot \Gamma$	$URU \cdot \Gamma$
$U$	$U \cdot \Gamma$	$\Gamma$	$URU \cdot \Gamma$	$USRU \cdot \Gamma$	$RU \cdot \Gamma$	$SRU \cdot \Gamma$
$T^{-1}$	$\Gamma$	$U \cdot \Gamma$	$RU \cdot \Gamma$	$URU \cdot \Gamma$	$SRU \cdot \Gamma$	$USRU \cdot \Gamma$
$R^{-1}$	$\Gamma$	$RU \cdot \Gamma$	$U \cdot \Gamma$	$SRU \cdot \Gamma$	$URU \cdot \Gamma$	$USRU \cdot \Gamma$

## References

- [1] P. Buser, *Geometry and spectra of compact Riemann surfaces*. Progr. Math. 106, Birkhäuser, Boston, MA, 1992. [Zbl 0770.53001](#) [MR 1183224](#)
- [2] A. Eskin and A. Okounkov, Asymptotics of numbers of branched coverings of a torus and volumes of moduli spaces of holomorphic differentials. *Invent. Math.* **145** (2001), no. 1, 59–104. [Zbl 1019.32014](#) [MR 1839286](#)
- [3] B. Farb and D. Margalit, *A primer on mapping class groups*. Princeton Math. Ser. 49, Princeton University Press, Princeton, NJ, 2012. [Zbl 1245.57002](#) [MR 2850125](#)
- [4] H. Farkas and I. Kra, *Riemann surfaces*. Second edition, Grad. Texts in Math. 71, Springer-Verlag, New York 1992. [Zbl 0764.30001](#) [MR 1139765](#)
- [5] F. Gardiner, *Teichmüller theory and quadratic differentials*. Pure Appl. Math. (N. Y.), John Wiley & Sons, New York 1987. [Zbl 0629.30002](#) [MR 0903027](#)
- [6] R. Hain, Finiteness and Torelli spaces. In *Problems on mapping class groups and related topics*, B. Farb, ed., Proc. Symp. Pure Math. 74, Amer. Math. Soc., Providence, RI, 2006. [Zbl 1222.14014](#) [MR 2264130](#)
- [7] P. Hubert, H. Masur, T. A. Schmidt, and A. Zorich, Problems on billiards, flat surfaces and translation surfaces. In *Problems on mapping class groups and related topics*, B. Farb, ed., Proc. Symp. Pure Math. 74, Amer. Math. Soc., Providence, RI, 2006. [Zbl 05124686](#) [MR 2264543](#)
- [8] H. Masur, Closed trajectories for quadratic differentials with an application to billiards. *Duke Math. J.* **53**(1986), no. 2, 307–314. [Zbl 0616.30044](#) [MR 0850537](#)
- [9] H. Masur and S. Tabachnikov, Rational billiards and flat structures. In *Handbook of dynamical systems*, B. Hasselblatt and A. Katok, eds., Vol. 1A, North-Holland, Amsterdam, 2002, 1015–1089. [Zbl 1057.37034](#) [MR 1928530](#)
- [10] C. T. McMullen, Dynamics of  $SL_2(\mathbb{R})$  over moduli space in genus two. *Ann. of Math.* (2) **165** (2007), no. 2, 397–456. [Zbl 1131.14027](#) [MR 2299738](#)
- [11] C. T. McMullen, Braid groups and Hodge theory. *Math. Ann.* **355** (2013), no. 3, 893–946. [Zbl 06149478](#) [MR 3020148](#)
- [12] G. Mess, The Torelli groups for genus 2 and 3 surfaces. *Topology* **31** (1992), no. 4, 775–790. [Zbl 0772.57025](#) [MR 1191379](#)
- [13] H. Rauch and H. Farkas, *Theta functions with applications to Riemann surfaces*. The Williams & Wilkins Company, Baltimore, Md., 1974. [Zbl 0292.30015](#) [MR 0352108](#)
- [14] A. Zorich, Flat surfaces. In *number theory, physics, and geometry. I*, On random matrices, zeta functions and dynamical systems, Les Houches 2003, Springer-Verlag, Berlin 2006, 437–583. [Zbl 1129.32012](#) [MR 2261104](#)

Received December 7, 2009; revised October 24, 2012

D.-M. Nguyen, Max Planck Institut für Mathematik, Vivatsgasse 7, 53111 Bonn, Germany;  
current address: IMB Bordeaux, Université de Bordeaux 351, cours de la Libération,  
33405 Talence Cedex, France

E-mail: [duc-manh.nguyen@math.u-bordeaux1.fr](mailto:duc-manh.nguyen@math.u-bordeaux1.fr)