

VARIETIES WITH VANISHING HOLOMORPHIC EULER CHARACTERISTIC

JUNGKAI ALFRED CHEN, OLIVIER DEBARRE, AND ZHI JIANG

ABSTRACT. We study smooth complex projective varieties X of maximal Albanese dimension and of general type satisfying $\chi(X, \mathcal{O}_X) = 0$. We prove that the Albanese variety of X has at least three simple factors. Examples were constructed by Ein and Lazarsfeld, and we prove that in dimension 3, these examples are (up to abelian étale covers) the only ones. By results of Ueno, another source of examples is provided by varieties X of maximal Albanese dimension and of general type satisfying $h^0(X, K_X) = 1$. Examples were constructed by Chen and Hacon, and again, we prove that in dimension 3, these examples are (up to abelian étale covers) the only ones. We also formulate a conjecture on the general structure of these varieties in all dimensions.

1. INTRODUCTION

A smooth complex projective variety X is said to have *maximal Albanese dimension* if its Albanese mapping $X \rightarrow \text{Alb}(X)$ is generically finite (onto its image).

Green and Lazarsfeld showed in [GL] that such a variety satisfies $\chi(X, \omega_X) \geq 0$. Ein and Lazarsfeld later constructed in [EL] a smooth projective threefold X of maximal Albanese dimension and of general type with $\chi(X, \omega_X) = 0$ (see Examples 4.1 and 4.2).

We are interested here in describing the structure of varieties X of maximal Albanese dimension (and of general type) with $\chi(X, \omega_X) = 0$. This class of varieties is stable by modifications, étale covers, and products with any other variety of maximal Albanese dimension (and of general type). More generally, if X is a smooth projective variety of maximal Albanese dimension with a fibration whose general fiber F satisfies $\chi(F, \omega_F) = 0$, then $\chi(X, \omega_X) = 0$ ([HP], Proposition 2.5).

So we study smooth projective varieties X of general type with $\chi(X, \omega_X) = 0$ and a generically finite morphism $X \rightarrow A$ to an abelian variety. In §3, we prove a general structure theorem (Theorem 3.1) which implies among other things that A has at least three simple factors. Examples where A is the product of any three given non-zero factors can be constructed following Ein and Lazarsfeld, and we speculate that their construction should (more or less) describe all cases where A has three simple factors but, although we prove several results in §4 in this direction (Propositions 4.3, 4.5, and 4.7) and arrive at the rather rigid picture (5), we are only able to get a complete description when X has dimension

2010 *Mathematics Subject Classification.* 14J10, 14J30, 14F17, 14E05.

Key words and phrases. Vanishing theorems, generic vanishing, cohomological loci, varieties of general type, Albanese dimension, Albanese variety, Euler characteristic, isotrivial fibrations.

O. Debarre is part of the project VSHMOD-2009 ANR-09- BLAN-0104-01.

3: we prove that *a smooth projective threefold X of maximal Albanese dimension and of general type satisfies $\chi(X, \omega_X) = 0$ if and only if it has an abelian étale cover which is an Ein-Lazarsfeld threefold* (Theorem 5.1).

Another source of examples is provided by varieties X of maximal Albanese dimension and $h^0(X, \omega_X) = 1$: it follows from work of Ueno ([U]) that they satisfy $\chi(X, \omega_X) = 0$. Chen and Hacon constructed examples of general type (see Example 4.2). We gather some properties of these varieties in §6. However, this class of examples is not stable under étale covers and does not lend itself well to our methods of study, *except in dimension 3*, where the precise Theorem 5.1 allows us to give *a complete description of all smooth projective threefolds X of maximal Albanese dimension and of general type, such that $P_1(X) = 1$: they are all modifications of abelian étale covers of Chen-Hacon threefolds* (Theorem 6.3).

In §7, we propose a conjecture on the possible general structure of smooth projective varieties X of maximal Albanese dimension and of general type satisfying $\chi(X, \omega_X) = 0$. It seems difficult to give a complete classification, but based on the examples that we know, we conjecture that, after taking modifications and étale covers, there should exist a non-trivial fibration $X \rightarrow Y$ which is either isotrivial, or whose general fiber F satisfies $\chi(F, \omega_F) = 0$. For the converse, one does have $\chi(X, \omega_X) = 0$ in the second case by [HP], Proposition 2.5, but not necessarily in the first case, of course. Both cases do happen (Example 7.2).

We work over the field of complex numbers.

Acknowledgements. The first-named author is partially supported by NCTS and the National Science Council of Taiwan. This work started during the second-named author's visit to Taipei under the support of the bilateral Franco-Taiwanese Project Orchid and continued during the first-named author's visit to Institut Henri Poincaré in Paris and the third-named author's stay at the Max Planck Institut for Mathematics in Bonn. The authors are grateful for the support they received on these occasions.

2. NOTATION AND PRELIMINARIES

For any smooth projective variety X , we set $\widehat{X} = \text{Pic}^0(X)$. For $\xi \in \widehat{X}$, we will denote by P_ξ an algebraically trivial line bundle on X that represents ξ .

Following standard terminology, we will say that a morphism $f : X \rightarrow A$ to an abelian variety A is *minimal* if the induced group morphism $\widehat{f} : \widehat{A} \rightarrow \widehat{X}$ is injective. Equivalently, $f(X)$ generates A as an algebraic group and f factors through no non-trivial abelian étale covers of A . The Albanese mapping a_X has this property. Any $f : X \rightarrow A$ factors as $f : X \xrightarrow{f'} A' \rightarrow A$, where A' is an abelian variety and f' is minimal.

An *algebraic fibration* (or simply a fibration) is a surjective morphism between normal projective varieties, with connected fibers.

In the rest of this section, X will be a smooth projective variety, of dimension n , with a *generically finite* morphism $f : X \rightarrow A$ to an abelian variety A . In particular, X has maximal Albanese dimension.

2.1. **Cohomological loci.** For each integer i , we define the cohomological loci

$$\begin{aligned} V_i(\omega_X, f) &= \{\xi \in \widehat{A} \mid H^i(X, \omega_X \otimes f^*P_\xi) \neq 0\} \\ &= \{\xi \in \widehat{A} \mid H^{n-i}(X, f^*P_{-\xi}) \neq 0\}. \end{aligned}$$

If Y is a smooth projective variety and $\varepsilon : Y \dashrightarrow X$ is birational, we have $R^j\varepsilon_*\omega_Y = 0$ for $j > 0$ and $\varepsilon_*\omega_Y \simeq \omega_X$ ([K2], Theorem 2.1), hence $\chi(X, \omega_X) = \chi(Y, \omega_Y)$ and $V_i(\omega_X, f) = V_i(\omega_Y, f \circ \varepsilon)$ for all i . In particular, these loci do not change when X is replaced with Y .

2.1.1. Since $R^j f_*\omega_X = 0$ for $j > 0$, we have for all i

$$V_i(\omega_X, f) = V_i(f_*\omega_X) := \{\xi \in \widehat{A} \mid H^i(A, f_*\omega_X \otimes P_\xi) \neq 0\}.$$

2.1.2. Each irreducible component of $V_i(\omega_X, f)$ is an abelian subvariety of \widehat{A} of codimension $\geq i$ ([EL], Remark 1.6 and Theorem 1.2) translated by a torsion point ([Si]).

2.1.3. There is a chain of inclusions ([EL], Lemma 1.8)

$$\text{Ker}(\widehat{f}) = V_n(\omega_X, f) \subseteq V_{n-1}(\omega_X, f) \subseteq \cdots \subseteq V_0(\omega_X, f) \subseteq \widehat{A},$$

and $\text{codim}(V_n(\omega_X, f)) \geq n$.

2.1.4. If $V_0(\omega_X, f)$ has a component of codimension i , this component is contained in (hence is an irreducible component of) $V_i(\omega_X, f)$ ([EL], (1.10)), so that we have $i \leq n$ and $f(X)$ is fibered by i -dimensional abelian subvarieties of A ([EL], Theorem 3).

2.1.5. For $\xi \in \widehat{A}$ general, $\chi(X, \omega_X) = h^0(X, \omega_X \otimes f^*P_\xi) \geq 0$ (use 2.1.2) and

$$\begin{aligned} \chi(X, \omega_X) = 0 &\iff V_0(\omega_X, f) \neq \widehat{A} \\ &\iff V_0(\omega_X, f) \text{ has a component of codimension } i \\ &\qquad\qquad\qquad \text{for some } i \in \{1, \dots, n-1\} \\ &\implies V_i(\omega_X, f) \text{ has a component of codimension } i \\ &\qquad\qquad\qquad \text{for some } i \in \{1, \dots, n-1\}, \end{aligned}$$

where the last implication is not always an equivalence.

2.1.6. The variety X is of general type if and only if $V_0(\omega_X, f)$ generates \widehat{A} ([CH1], Theorem 2.3).

2.1.7. If $V_0(\omega_X, f)$ is finite, Ein and Lazarsfeld proved that X is birational to an abelian variety ([CH1], Theorem 1.3.2). In particular, if f is moreover minimal, it is a birational isomorphism.

2.2. Composing f with a generically finite morphism. If Y is smooth projective and $h : Y \twoheadrightarrow X$ is surjective and generically finite, the trace map $h_*\omega_Y \rightarrow \omega_X$ splits the natural inclusion $\omega_X \rightarrow h_*\omega_Y$, hence $H^i(X, \omega_X \otimes f^*P_\xi)$ injects into $H^i(X, h_*\omega_Y \otimes f^*P_\xi)$ for all $\xi \in \widehat{A}$. Thus, using also 2.1.1, we obtain

$$V_i(\omega_X, f) \subseteq V_i(h_*\omega_Y, f) = V_i(\omega_Y, f \circ h).$$

Moreover,

$$(1) \quad \chi(Y, \omega_Y) \geq \chi(X, \omega_X).$$

When h is étale, we have

$$\chi(Y, \omega_Y) = \deg(h)\chi(X, \omega_X).$$

Finally, when h is obtained from an isogeny $\eta : B \twoheadrightarrow A$ as in the cartesian diagram

$$\begin{array}{ccc} Y = X \times_A B & \xrightarrow{g} & B \\ \downarrow h & \square & \downarrow \eta \\ X & \xrightarrow{f} & A \end{array}$$

we have $V_i(\omega_Y, g) = \widehat{\eta}(V_i(\omega_X, f))$. Combining this with 2.1.2, we see that after making a suitable étale base change, we can always make all the components of $V_i(\omega_X, f)$ pass through 0.

3. COMPONENTS OF V_i OF CODIMENSION i

Let X be a smooth projective variety with a generically finite morphism $f : X \rightarrow A$ to an abelian variety. If $\chi(X, \omega_X) = 0$, it follows from 2.1.5 that $V_i(\omega_X, f)$ has a component of codimension i for some $i \in \{1, \dots, n-1\}$. We prove a structure theorem under this weaker assumption.

Theorem 3.1. *Let X be a smooth projective variety of dimension n , let A be an abelian variety, and let $f : X \rightarrow A$ be a minimal generically finite morphism. Assume that for some $i \in \{0, \dots, n\}$, the locus $V_i(\omega_X, f)$ has a component V of codimension i in \widehat{A} . Let B be the abelian variety \widehat{V} , let $K := \text{Ker}(A \twoheadrightarrow B)^0$, and assume $f(X) + K = f(X)$. For a suitable modification X' of an abelian étale cover of X , the Stein factorization of the morphism $X' \rightarrow A \twoheadrightarrow B$ induces a surjective morphism $X' \twoheadrightarrow Y$ where Y is smooth of dimension $n-i$, of general type, with $\chi(Y, \omega_Y) > 0$.*

Remark 3.2. The condition $f(X) + K = f(X)$ holds:

- when f is surjective;
- when V is also a component of $V_0(\omega_X, f)$ ([EL], proof of Theorem 3); this applies in particular when $\chi(X, \omega_X) = 0$ (2.1.5).

Proof of Theorem 3.1. By (2.1.2) and §2.2, we may assume, after isogeny, that $A = B \times K$ and $V = \widehat{B}$. Let $p : A \twoheadrightarrow B$ be the projection. Considering the Stein factorization of

$\pi = p \circ f : X \rightarrow B$, and replacing X by a suitable modification, we may assume that π factors as

$$\pi : X \xrightarrow{g} Y \xrightarrow{h} B,$$

where Y is smooth, h is generically finite, and g is surjective with connected fibers. Since $f(X) + K = f(X)$, the image of π has dimension $\dim(X) - \dim(K)$, hence general fibers of g have dimension $\dim(K) = i$.

We then have $R^i g_* \omega_X \simeq \omega_Y$ ([K2], Proposition 7.6). Moreover, the sheaves $R^k g_* \omega_X$ on Y satisfy the generic vanishing theorem ([HP], Theorem 2.2), hence

$$V_j(R^k g_* \omega_X, h) \neq \widehat{B} \quad \text{for all } j > 0 \text{ and all } k.$$

For all

$$\xi \in \widehat{B} - \bigcup_{j>0, k} V_j(R^k g_* \omega_X, h),$$

we have

$$H^j(Y, R^k g_* \omega_X \otimes h^* P_\xi) = 0 \quad \text{for all } j > 0 \text{ and all } k.$$

Hence, by the Leray spectral sequence, we obtain

$$h^i(X, \omega_X \otimes f^* P_\xi) = h^i(X, \omega_X \otimes \pi^* P_\xi) = h^0(Y, R^i g_* \omega_X \otimes h^* P_\xi) = h^0(Y, \omega_Y \otimes h^* P_\xi)$$

and these numbers are non-zero because $\widehat{B} = V \subseteq V_i(\omega_X, f)$. In particular, $V_0(\omega_Y, h) = \widehat{B}$. By 2.1.6 and 2.1.5, Y is of general type and $\chi(Y, \omega_Y) > 0$. This completes the proof. \square

We prove a partial converse to Theorem 3.1: assume that there is a generically finite morphism $f : X \rightarrow A$ and a quotient abelian variety $A \twoheadrightarrow B$ such that $f(X) + K = f(X)$, where $K := \text{Ker}(A \twoheadrightarrow B)^0$, and denote by $X \dashrightarrow Y \rightarrow B$ a modification of the Stein factorization of $X \rightarrow A \twoheadrightarrow B$, where Y is smooth of dimension $n - i$ (we set $i := \dim(K)$).

Proposition 3.3. *In this situation, if Y is not birational to an abelian variety, $V_j(\omega_X, f)$ has a component of codimension j for some $j \in \{i, \dots, n - 1\}$.*

Proof. Replacing X with a modification of an étale cover (which is allowed by §2.1 and §2.2), we may assume that we have a factorization

$$f : X \xrightarrow{(g,k)} Y \times K \xrightarrow{h \times \text{Id}_K} B \times K,$$

where (g, k) is surjective and $h : Y \rightarrow B$ is generically finite of degree > 1 . We obtain, as in the proof of Theorem 3.1, for ξ general in \widehat{B} ,

$$(2) \quad h^i(X, \omega_X \otimes f^* P_\xi) = h^0(Y, R^i g_* \omega_X \otimes h^* P_\xi) = h^0(Y, \omega_Y \otimes h^* P_\xi).$$

If $\chi(Y, \omega_Y) > 0$, we have $V_0(\omega_Y, h) = \widehat{B}$, the number on the right-hand-side of (2) is non-zero for all ξ , hence $V_i(\omega_X, f)$ contains the i -codimensional abelian subvariety \widehat{B} of \widehat{A} .

If $\chi(Y, \omega_Y) = 0$, since Y is not birational to an abelian variety, $V_0(\omega_Y, h)$ has (by 2.1.7 and 2.1.5) a component of codimension $l \in \{1, \dots, n - i - 1\}$ in \widehat{B} . Thus, by Remark 3.2, we can apply Theorem 3.1 to $h : Y \rightarrow B$: after taking an étale cover and a modification, h factors through a morphism $Y \rightarrow Z \times C$, where C is an abelian variety of dimension l ,

$\chi(Z, \omega_Z) > 0$, and $\dim(Z) = n - i - l$. We are therefore reduced to the first case and we conclude again that $V_{i+l}(\omega_X, f)$ contains an $(i + l)$ -codimensional component. \square

Remark 3.4. Under the hypotheses of Theorem 3.1 and the assumption $A = B \times K$ made in its proof, we obtain a surjective morphism $k : X \xrightarrow{f} B \times K \xrightarrow{p_2} K$ and, from its Stein factorization, morphisms

$$k : X \xrightarrow{l} Z \xrightarrow{m} K,$$

where Z is smooth of dimension i , m is generically finite, and l has connected (generically $(n - i)$ -dimensional) fibers. We have again $R^{n-i}l_*\omega_X \simeq \omega_Z$ and, for ξ general in \widehat{K} ,

$$h^{n-i}(X, \omega_X \otimes f^*P_\xi) = h^0(Z, \omega_Z \otimes m^*P_\xi).$$

Then,

- a) either $V_0(\omega_Z, m) \subsetneq \widehat{K}$ and $\chi(Z, \omega_Z) = 0$;
- b) or $V_0(\omega_Z, m) = \widehat{K}$ and $\chi(Z, \omega_Z) > 0$, in which case \widehat{K} is contained in (hence is a component of) $V_{n-i}(\omega_X, f)$. There is a surjective generically finite map $X \rightarrow Y \times Z$, hence $\chi(X, \omega_X) \geq \chi(Y, \omega_Y)\chi(Z, \omega_Z) > 0$ (this also follows from Corollary 3.5.a) below).

Finally, if F is a general fiber of $l : X \rightarrow Z$, there is a surjective generically finite map $F \rightarrow Y$, hence $\chi(F, \omega_F) > 0$ (see (1)).

We now deduce some consequences of Theorem 3.1 on the possible components of $V_0(\omega_X, f)$ and the number of simple factors of the abelian variety A .

Corollary 3.5. *Let X be a smooth projective variety with $\chi(X, \omega_X) = 0$ and a generically finite morphism $f : X \rightarrow A$ to an abelian variety.*

- a) *The locus $V_0(\omega_X, f)$ does not have complementary components.¹*
- b) *If X is in addition of general type, A has at least three simple factors.*

Proof. If $V_0(\omega_X, f)$ has complementary components V_1, \dots, V_r , with duals B_1, \dots, B_r , the image $f(X)$ is stable by translation by $\prod_{j \neq i} B_j$ for each i (Remark 3.2), hence f is surjective if $r \geq 2$. We obtain from Theorem 3.1, after passing to an étale cover and a modification of X , a generically finite surjective map $X \rightarrow Y_1 \times \dots \times Y_r$, with $\chi(Y_i, \omega_{Y_i}) > 0$ for all i . Since $\chi(X, \omega_X) \geq \prod_i \chi(Y_i, \omega_{Y_i})$ (by (1)), this is absurd. This proves a). Item b) then follows from 2.1.6. \square

Proposition 3.6. *Let X be a smooth projective variety of general type, with $\chi(X, \omega_X) = 0$ and a generically finite morphism $f : X \rightarrow A$ to an abelian variety. Then $V_0(\omega_X, f)$ has no 1-dimensional components.*

Proof. Assume $V_0(\omega_X, f)$ has a one-dimensional component and write it as $\tau_1 + \widehat{B}_1$ for some torsion point $\tau_1 \in \widehat{A}$ and some quotient elliptic curve $A \rightarrow B_1$. By [J], Proposition 1.7, and since $V_0(\omega_X, f)$ generates \widehat{A} (2.1.6), \widehat{B}_1 cannot be maximal for inclusion: more precisely,

¹By that, we mean components such that the sum morphism induces an isogeny from their product onto \widehat{A} .

there must exist two maximal components (in the sense of [J], Definition 1.6) $\tau_2 + \widehat{B}_2$ and $\tau_3 + \widehat{B}_3$ of $V_0(\omega_X, f)$ with $\widehat{B}_1 \subsetneq \widehat{B}_j \subsetneq \widehat{A}$ for each $j \in \{2, 3\}$ and $\widehat{B}_2 \neq \widehat{B}_3$. There are corresponding factorizations $A \twoheadrightarrow B_j \twoheadrightarrow B_1$.

As in the proof of Theorem 3.1, after passing to an étale cover and a modification of X , we may assume $\tau_1 = \tau_2 = \tau_3 = 0$ and that we have

- Stein factorizations

$$X \xrightarrow{g_i} Y_i \xrightarrow{h_i} B_i,$$

where h_1, h_2 , and h_3 are generically finite, Y_1, Y_2 , and Y_3 are smooth, Y_1 is a curve of genus ≥ 2 , and $\chi(Y_2, \omega_{Y_2})$ and $\chi(Y_3, \omega_{Y_3})$ are both positive (Theorem 3.1),

- a commutative diagram

$$\begin{array}{ccccc}
 & & Y_2 & \xrightarrow{h_2} & B_2 \\
 & g_2 \nearrow & & \searrow h_{21} & \\
 X & \xrightarrow{g_1} & Y_1 & & \\
 & g_3 \searrow & & \nearrow h_{31} & \\
 & & Y_3 & \xrightarrow{h_3} & B_3.
 \end{array}$$

We may further assume that the induced morphism $X \rightarrow Y_2 \times_{Y_1} Y_3$ factors as:

$$\begin{array}{ccccccc}
 & & & & Y_2 & \xrightarrow{h_2} & B_2 \\
 & & & & \nearrow & \searrow h_{21} & \\
 X & \longrightarrow & Y & \xrightarrow{\varepsilon} & Y_2 \times_{Y_1} Y_3 & \longrightarrow & Y_1, \\
 & & \nearrow q & & \nearrow & \searrow h_{31} & \\
 & & & & Y_3 & \xrightarrow{h_3} & B_3
 \end{array}$$

where ε is a resolution of singularities.

Now take $\xi_2 \in \widehat{B}_2$ and $\xi_3 \in \widehat{B}_3$. By [M], Lemma 4.10.(ii),² there is an inclusion

$$q_*(\omega_{Y/Y_1} \otimes \varepsilon^*(h_2^*P_{\xi_2} \otimes h_3^*P_{\xi_3})) \subseteq h_{21*}(\omega_{Y_2/Y_1} \otimes h_2^*P_{\xi_2}) \otimes h_{31*}(\omega_{Y_3/Y_1} \otimes h_3^*P_{\xi_3}),$$

of locally free sheaves of the same rank on the curve Y_1 . Moreover, we saw during the proof of Theorem 3.1 that for $j \in \{2, 3\}$, we have

$$0 \neq h^0(Y_j, \omega_{Y_j} \otimes h_j^*P_{\xi_j}) = h^0(Y_1, h_{j1*}(\omega_{Y_j} \otimes h_j^*P_{\xi_j})).$$

It follows that the sheaf $h_{j1*}(\omega_{Y_j} \otimes h_j^*P_{\xi_j})$ is non-zero, hence so is the sheaf $h_{j1*}(\omega_{Y_j/Y_1} \otimes h_j^*P_{\xi_j})$. All in all, we have obtained that the locally free sheaf $q_*(\omega_{Y/Y_1} \otimes \varepsilon^*(h_2^*P_{\xi_2} \otimes h_3^*P_{\xi_3}))$ is non-zero.

Assume now that ξ_2 and ξ_3 are torsion. By [V], Corollary 3.6,³ this vector bundle is nef, hence has non-negative degree. Since Y_1 is a curve of genus ≥ 2 , the Riemann-Roch

²This is stated in [M] for $\xi_2 = \xi_3 = 0$, but the same proof works in general.

³This is stated there for $\xi_2 = \xi_3 = 0$, but the general case follows by the étale covering trick.

theorem then implies

$$0 \neq h^0(Y_1, q_*(\omega_Y \otimes \varepsilon^*(h_2^*P_{\xi_2} \otimes h_3^*P_{\xi_3}))) = h^0(Y, \omega_Y \otimes \varepsilon^*(h_2^*P_{\xi_2} \otimes h_3^*P_{\xi_3})).$$

Finally, note that both X and Y have maximal Albanese dimensions. This implies that $\omega_{X/Y}$ is effective, hence $h^0(X, \omega_X \otimes f^*(P_{\xi_2} \otimes P_{\xi_3}))$ is also non-zero. It follows that $\xi_2 + \xi_3$ is in $V_0(\omega_X, f)$, which therefore contains $\widehat{B}_2 + \widehat{B}_3$. This contradicts the fact that \widehat{B}_2 is maximal. \square

4. CASE WHEN A HAS THREE SIMPLE FACTORS

Ein and Lazarsfeld constructed an example of a smooth projective threefold X of maximal Albanese dimension and of general type with $\chi(X, \omega_X) = 0$, whose Albanese variety is the product of three elliptic curves. After presenting their construction (and a variant due to Chen and Hacon), we prove some general results when $\text{Alb}(X)$ has three simple factors. In the next section, we will show that the Ein-Lazarsfeld example is essentially the only one in dimension 3 (Theorem 5.1).

Example 4.1 ([EL], Example 1.13). Let E_1, E_2 , and E_3 be elliptic curves and let $\rho_j : C_j \rightarrow E_j$ be double coverings, where C_j is a smooth curve of genus ≥ 2 and $\rho_{j*}\omega_{C_j} \simeq \mathcal{O}_{E_j} \oplus \delta_j$. Denote by ι_j the corresponding involution of C_j . Let $A = E_1 \times E_2 \times E_3$, and consider the quotient Z of $C_1 \times C_2 \times C_3$ by the involution $\iota_1 \times \iota_2 \times \iota_3$ and the tower of Galois covers:

$$C_1 \times C_2 \times C_3 \xrightarrow{g} Z \xrightarrow{f} A$$

of degrees 2 and 4 respectively. Observe that Z has rational singularities and is minimal of general type. Let $\varepsilon : X \rightarrow Z$ be any desingularization. The Albanese map of X is $a_X = f \circ \varepsilon$ and

$$a_{X*}\omega_X \simeq \mathcal{O}_A \oplus (L_1 \otimes L_2) \oplus (L_3 \otimes L_1) \oplus (L_2 \otimes L_3),$$

where L_j is the inverse image of δ_j by the projection $A \rightarrow E_j$, hence

$$(3) \quad V_0(\omega_X, a_X) = V_1(\omega_X, a_X) = (\widehat{E}_1 \times \widehat{E}_2 \times \{0\}) \cup (\widehat{E}_1 \times \{0\} \times \widehat{E}_3) \cup (\{0\} \times \widehat{E}_2 \times \widehat{E}_3),$$

whereas $V_2(\omega_X, a_X) = V_3(\omega_X, a_X) = \{0\}$.

This provides three-dimensional examples. Obviously, the same construction works starting from double coverings $\rho_j : X_j \rightarrow A_j$ of abelian varieties with smooth ample branch loci and provides examples in all dimensions ≥ 3 . One can also extend it to any *odd* number $2r + 1$ of factors and get examples where the Albanese mapping is birationally a $(\mathbf{Z}/2\mathbf{Z})^{2r}$ -covering.

Example 4.2 ([CH2], §4, Example). A variant of the construction above was given by Chen and Hacon. Keeping the same notation, choose points $\xi_j \in \widehat{E}_j$ of order 2 and consider the induced double étale covers $C'_j \rightarrow C_j$, with associated involution σ_j , and $E'_j \rightarrow E_j$. The involution ι_j on C_j pulls back to an involution ι'_j on C'_j (with quotient E'_j). Let Z' be the quotient of $C'_1 \times C'_2 \times C'_3$ by the group of automorphisms generated by $\text{id}_1 \times \sigma_2 \times \iota'_3$, $\iota'_1 \times \text{id}_2 \times \sigma_3$, $\sigma_1 \times \iota'_2 \times \text{id}_3$, and $\sigma_1 \times \sigma_2 \times \sigma_3$, and let $\varepsilon' : X' \rightarrow Z'$ be a desingularization. There is a morphism $f' : X' \rightarrow A$ of degree 4, the Albanese map of X' is $a_{X'} = f' \circ \varepsilon'$, and

$$(4) \quad a_{X'*}\omega_{\widetilde{X}'} \simeq \mathcal{O}_A \oplus (L_1 \otimes L_2^\xi \otimes P_{\xi_3}) \oplus (L_1^\xi \otimes P_{\xi_2} \otimes L_3) \oplus (P_{\xi_1} \otimes L_2 \otimes L_3^\xi),$$

where $L_j^\xi = L_j \otimes P_{\xi_j}$. In particular, $P_1(X') = 1$, and

$$V_0(\omega_{X'}, a_{X'}) = V_1(\omega_{X'}, a_{X'}) = \{0\} \cup (\widehat{E}_1 \times \widehat{E}_2 \times \{\xi_3\}) \cup (\widehat{E}_1 \times \{\xi_2\} \times \widehat{E}_3) \cup (\{\xi_1\} \times \widehat{E}_2 \times \widehat{E}_3).$$

Of course, the étale cover $E'_1 \times E'_2 \times E'_3 \rightarrow E_1 \times E_2 \times E_3$ pulls back to an étale cover $X'' \rightarrow X'$, where X'' is an Ein-Lazarsfeld threefold.

Again, this construction still works starting from double coverings of abelian varieties with smooth ample branch loci, providing examples in all dimensions ≥ 3 , and for any *odd* number $2r + 1$ of factors, providing examples where the Albanese mapping is birationally a $(\mathbf{Z}/2\mathbf{Z})^{2r}$ -covering.

Proposition 4.3. *Let X be a smooth projective variety of general type with $\chi(X, \omega_X) = 0$ and a generically finite morphism $f : X \rightarrow A$ to an abelian variety A with exactly three simple factors A_1, A_2, A_3 .*

- a) *The map f is surjective.*
- b) *After passing to an abelian étale cover, we may assume $A = A_1 \times A_2 \times A_3$ and that $\widehat{A}_1 \times \widehat{A}_2 \times \{0\}$, $\widehat{A}_1 \times \{0\} \times \widehat{A}_3$, and $\{0\} \times \widehat{A}_2 \times \widehat{A}_3$ are irreducible components of $V_0(\omega_X, f)$.*

Proof. We begin with the proof of item b). As in the proof of Theorem 3.1, we may assume, after passing to an abelian étale cover, that A is $A_1 \times A_2 \times A_3$ and, by Corollary 3.5.a) and 2.1.6, that $\widehat{A}_1 \times \widehat{A}_2 \times \{0\}$ and $\widehat{A}_1 \times \{0\} \times \widehat{A}_3$ are irreducible components of $V_0(\omega_X, f)$.

Assume that the projection $V_0(\omega_X, f) \rightarrow \widehat{A}_2 \times \widehat{A}_3$ is not surjective. For ξ_2 and ξ_3 general torsion points in \widehat{A}_2 and \widehat{A}_3 respectively, we then have

$$(\xi_2 + \xi_3 + \widehat{A}_1) \cap V_0(\omega_X, f) = \emptyset.$$

Consider the morphism $f_1 = p_1 \circ f : X \rightarrow A_1$ and the sheaf $\mathcal{E} = f_{1*}(\omega_X \otimes f_2^* P_{\xi_2} \otimes f_3^* P_{\xi_3})$ on A_1 . By [HP], Theorem 2.2, the cohomological loci of \mathcal{E} satisfy the chain of inclusions 2.1.3. On the other hand, for all $\xi \in \widehat{A}_1$, we have $H^0(A_1, \mathcal{E} \otimes P_\xi) = 0$, hence $V_0(\mathcal{E}) = \emptyset$. It follows that for all $\xi \in \widehat{A}_1$ and all $i \geq 0$, we have $H^i(A_1, \mathcal{E} \otimes P_\xi) = 0$. This implies $\mathcal{E} = 0$ by Fourier-Mukai duality ([Mu]). But the rank of \mathcal{E} is at least $\chi(F_1, \omega_{F_1})$, where F_1 is a component of a general fiber of f_1 ([HP], Corollary 2.3) and this is impossible: F_1 is of general type and generically finite over $A_2 \times A_3$, hence $\chi(F_1, \omega_{F_1}) > 0$ (Corollary 3.5.b)).

The projection $V_0(\omega_X, f) \rightarrow \widehat{A}_2 \times \widehat{A}_3$ is therefore surjective. Since A_1 is simple, this implies that, after passing to a (split) étale cover of A , there are morphisms $u_2 : \widehat{A}_2 \rightarrow \widehat{A}_1$ and $u_3 : \widehat{A}_3 \rightarrow \widehat{A}_1$ such that

$$\{u_2(\xi_2) + u_3(\xi_3) + \xi_2 + \xi_3 \mid \xi_2 \in \widehat{A}_2, \xi_3 \in \widehat{A}_3\}$$

is a component of $V_0(\omega_X, f)$. Composing this cover with the automorphism $(a_1, a_2, a_3) \mapsto (a_1, a_2 - \widehat{u}_2(a_1), a_3 - \widehat{u}_3(a_1))$ of A , we obtain b).

Item a) then follows from the fact that $f(X)$ is stable by translation by each A_i (Remark 3.2) hence is equal to A . \square

Remark 4.4. As shown by considering the product with a curve of genus ≥ 2 of any variety X of general type and maximal Albanese dimension with $\chi(X, \omega_X) = 0$, the conclusion of Proposition 4.3.a) does not hold in general as soon as A has at least four simple factors.

Proposition 4.5. *Let X be a smooth projective variety of general type of dimension n with $\chi(X, \omega_X) = 0$ and a generically finite morphism $X \rightarrow A$ to an abelian variety A with exactly three simple factors. We have:*

- a) $q(X) = n$;
- b) the general fiber F of any non-constant fibration $X \rightarrow Y$ satisfies $\chi(F, \omega_F) > 0$;
- c) any morphism from X to a curve of genus ≥ 2 is constant;
- d) $V_{n-1}(\omega_X, a_X) = \{0\}$.

Proof. Let us prove b) first. The fiber F is of general type and is generically finite but not surjective over A , hence $\chi(F, \omega_F) > 0$ by Proposition 4.3.a). The other items then follow from the lemma below. \square

Lemma 4.6. *Let X be a smooth projective variety of dimension n , of maximal Albanese dimension, of general type, with $\chi(X, \omega_X) = 0$. Assume that the general fiber F of any non-constant fibration $X \rightarrow Y$ satisfies $\chi(F, \omega_F) > 0$. Then,*

- a) the Albanese mapping a_X is surjective ($q(X) = n$);
- b) any morphism from X to a curve of genus ≥ 2 is constant;
- c) $V_{n-1}(\omega_X, a_X) = \{0\}$.

Proof. Item a) follows from [CH2], Theorem 4.2, and item b) from [HP], Theorem 2.4. Let us prove c).

If $V_{n-1}(\omega_X, a_X) \neq \{0\}$ is non-empty, it contains a torsion point by 2.1.2, which defines a connected étale cover $\pi : \tilde{X} \rightarrow X$ such that $q(\tilde{X}) > q(X) = n$. By [CH2], Theorem 4.2, again, there exists a non-constant fibration $\tilde{X} \rightarrow Y$ with general fiber F of maximal Albanese dimension, of general type, with $\chi(F, \omega_F) = 0$, such that $a_{\tilde{X}}(F)$ is a translate of a fixed abelian subvariety \tilde{K} of $\text{Alb}(\tilde{X})$ and $\dim(\tilde{K}) = \dim(F) < \dim(X)$. We consider the image K of \tilde{K} by the induced map $\text{Alb}(\pi) : \text{Alb}(\tilde{X}) \rightarrow \text{Alb}(X)$ and the commutative diagram:

$$\begin{array}{ccccc}
 F & \xrightarrow{\pi|_F} & \pi(F) & \xrightarrow{a_{\tilde{X}}|_{\pi(F)}} & K + x_F \\
 \downarrow & & \downarrow & & \downarrow \\
 \tilde{X} & \xrightarrow{\pi} & X & \xrightarrow{a_X} & \text{Alb}(X) \\
 f \downarrow & & & \searrow h & \downarrow \\
 Y & & & & \text{Alb}(X)/K.
 \end{array}$$

The map $a_{\tilde{X}}|_{\pi(F)}$ is generically finite, hence $\dim(K) = \dim \pi(F) = \dim(F)$, and

$$\dim(\text{Alb}(X)/K) = n - \dim(K) = n - \dim(F).$$

It follows that $\pi(F)$ is a general fiber of the Stein factorization of h . Since $\chi(\pi(F), \omega_{\pi(F)}) = 0$, this contradicts our hypothesis on X . \square

Assume now that we are in the situation of Proposition 4.3.b) and consider, as in Remark 3.4, the maps $f_i := p_i \circ f : X \rightarrow A_i$. Since $\chi(X, \omega_X) = 0$, we are in case a) of that remark, hence, with the notation therefrom, $V_0(\omega_Z, m)$ must be finite, because A_i is simple. If f is minimal, so is f_i and it follows from 2.1.7 that Z is birational to A_i , hence f_i is a fibration. A general fiber F_i satisfies $\chi(F_i, \omega_{F_i}) > 0$ by Proposition 4.5.b).

Proposition 4.7. *Let X be a smooth projective variety of general type with $\chi(X, \omega_X) = 0$ and a minimal generically finite morphism $f : X \rightarrow A$ to an abelian variety A product of three simple factors A_1, A_2 , and A_3 .*

Let $\{i, j, k\} = \{1, 2, 3\}$. For ξ_j and ξ_k general torsion points in \widehat{A}_j and \widehat{A}_k respectively, the sheaf $f_{i}(\omega_X \otimes f_j^* P_{\xi_j} \otimes f_k^* P_{\xi_k})$ on A_i is locally free, homogeneous, of positive rank $\chi(F_i, \omega_{F_i})$.*

Proof. We follow the proof of [CH2], Corollary 2.3. As in the proof of Proposition 4.3, since ξ_j and ξ_k are torsion, the cohomological loci of $\mathcal{E} := f_{i*}(\omega_X \otimes f_j^* P_{\xi_j} \otimes f_k^* P_{\xi_k})$ satisfy the chain of inclusions 2.1.3. On the other hand, since ξ_j and ξ_k are general and A_i is simple, the intersection

$$(\xi_j + \xi_k + \widehat{A}_i) \cap V_0(\omega_X, f)$$

is finite by 2.1.2, hence so is $V_0(\mathcal{E})$. It follows that all $V_l(\mathcal{E})$ are finite, hence \mathcal{E} is locally free and homogeneous ([Mu], Example 3.2). Its rank is $h^0(F_i, \omega_{F_i} \otimes f_j^* P_{\xi_j} \otimes f_k^* P_{\xi_k}) = \chi(F_i, \omega_{F_i})$. \square

Remark 4.8. Recall that a homogeneous vector bundle is a direct sum of twists of unipotent vector bundles (successive extensions of trivial line bundles) by algebraically trivial line bundles which, in our case, are torsion by Simpson's theorem (or rather its extension [HP], Theorem 2.2.b)).

When A_i is an elliptic curve, the sheaf of Proposition 4.7 is actually a direct sum of torsion line bundles (this is explained at the bottom of page 362 of [K3] when $\xi_j = \xi_k = 0$ and holds in general by the étale covering trick).

Finally, from the proof of Theorem 3.1, we have (after replacing X with a suitable modification), for each $\{i, j, k\} = \{1, 2, 3\}$, Stein factorizations

$$p_{jk} \circ f : X \xrightarrow{g_i} S_i \xrightarrow{(h_{ij}, h_{ik})} A_j \times A_k,$$

where S_i is smooth of general type with $\chi(S_i, \omega_{S_i}) > 0$ (this follows also from Corollary 3.5.b)). Since f_j has connected fibers, so does $h_{ij} : S_i \rightarrow A_j$.

All in all, we have for each $\{i, j, k\} = \{1, 2, 3\}$ a commutative diagram:

$$(5) \quad \begin{array}{ccccc} & & X & & \\ & \nearrow^{f_j} & \downarrow^{f_k} & \nwarrow_{g_j} & \\ & S_i & & S_j & \\ & \nwarrow_{h_{ij}} & \searrow_{h_{ik}} & \swarrow_{h_{jk}} & \searrow_{h_{ji}} \\ A_j & & A_k & & A_i \end{array}$$

where all the morphisms are fibrations.

Question 4.9. Are the h_{ij} isotrivial? Are the f_i isotrivial? Is X rationally dominated by a product $X_1 \times X_2 \times X_3$, where X_i dominates and is generically finite over A_i ? We are inclined to think that the answers to all these questions should be affirmative, but we were only able to go further in the case where the A_i are all elliptic curves.

5. THE 3-DIMENSIONAL CASE

We now come to our main result, which completely describes all smooth projective threefolds X of maximal Albanese dimension and of general type, with $\chi(X, \omega_X) = 0$.

Theorem 5.1. *Let X be a smooth projective threefold of maximal Albanese dimension and of general type, with $\chi(X, \omega_X) = 0$.*

There exist elliptic curves E_1, E_2 , and E_3 , double coverings $C_j \rightarrow E_j$ with associated involutions ι_j , and a commutative diagram

$$\begin{array}{ccc} & (C_1 \times C_2 \times C_3)/\iota_1 \times \iota_2 \times \iota_3 & \\ \varepsilon \nearrow & & \searrow \\ \tilde{X} & \xrightarrow{a_{\tilde{X}}} & E_1 \times E_2 \times E_3 \\ \downarrow & \square & \downarrow \eta \\ X & \xrightarrow{a_X} & \text{Alb}(X), \end{array}$$

where η is an isogeny and ε a desingularization.

In other words, up to abelian étale covers, the Ein-Lazarsfeld examples (Example 4.1) are the only ones (in dimension 3)! Note also that a_X is not finite, but that it is finite on the canonical model of X .

Corollary 5.2. *Under the hypotheses of the theorem, the Albanese mapping of X is birationally a $(\mathbf{Z}/2\mathbf{Z})^2$ -covering.*

Proof. With the notation of Example 4.1, we set $A := E_1 \times E_2 \times E_3$ and $\tilde{X}_i := \text{Spec}(\mathcal{O}_A \oplus L_i^\vee)$, so that f factors through the double coverings $\tilde{f}_i := \tilde{X}_i \rightarrow A$. Since the action of $\text{Ker}(\eta)$ on A by translations lifts to \tilde{X} , it leaves

$$a_{\tilde{X}*} \mathcal{O}_{\tilde{X}} \simeq \mathcal{O}_A \oplus (L_1^\vee \otimes L_2^\vee) \oplus (L_3^\vee \otimes L_1^\vee) \oplus (L_2^\vee \otimes L_3^\vee)$$

invariant, hence also each L_i^\vee . It follows that this action lifts to each \tilde{X}_i , hence \tilde{f}_i descends to a double covering $X_i \rightarrow \text{Alb}(X)$ through which a_X factors. \square

Proof of the theorem. By Proposition 4.5.a), a_X is surjective and $q(X) = 3$ (see also [CH2], Corollary 4.3).

Moreover, by Corollary 3.5.b), $\text{Alb}(X)$ is isogeneous to the product of three elliptic curves and, after passing to étale covers, we may assume that a_X can be written as

$$a_X : X \xrightarrow{(f_1, f_2, f_3)} E_1 \times E_2 \times E_3,$$

where each $f_i : X \rightarrow E_i$ is a fibration, and that $\widehat{E}_1 \times \widehat{E}_2 \times \{0\}$, $\widehat{E}_1 \times \{0\} \times \widehat{E}_3$, and $\{0\} \times \widehat{E}_2 \times \widehat{E}_3$ are irreducible components of $V_0(\omega_X, a_X)$ (Proposition 4.3.b)).

Let $\{i, j, k\} = \{1, 2, 3\}$. As in §4, we have a commutative diagram (5), where each S_i is a smooth minimal surface of general type and $A_i = E_i$. The proof of the theorem is very long, so we will divide it in several steps. The general scheme of proof goes as follows:

- In the diagram (5), the fibrations $h_{ij} : S_i \rightarrow E_j$ are all isotrivial (Step 1); we let C_{ij} be a (constant) general fiber.
- There exist finite groups G_i acting on C_{ij} such that $C_{ij}/G_i \simeq E_j$ and $C_{ik}/G_i \simeq E_k$, the surface S_i is birational to $(C_{ij} \times C_{ik})/G_i$, and h_{ij} and h_{ik} are the two projections (Step 2).
- At this point, it is quite easy to show that X dominates a threefold Y which is dominated by a product of 3 curves (Step 3).
- Taking an étale cover of X , we may assume that all the irreducible components of $V_0(\omega_X, a_X)$ pass through 0. We then show (Step 4) that $V_0(\omega_X, a_X)$ has the same form as the corresponding locus of an Ein-Lazarsfeld threefold (see (3)), from which we deduce that X is birationally isomorphic to Y (Step 5) hence is also dominated by a product of 3 curves.
- Using the fact that $V_0(\omega_X, a_X)$ has no “extra” components, we finish the proof by showing that the groups G_i all have order 2 (Step 6).

Step 1. *The fibrations $h_{ij} : S_i \rightarrow E_j$ are all isotrivial.*

We will denote by C_{ij} a general (constant) fiber of h_{ij} .

Proof. By the semi-stable reduction theorem ([KKMS], Chapter II), there exist a finite cover $h : C \rightarrow E_j$, where C is a smooth curve and commutative diagrams (for each $\alpha \in \{i, k\}$)

$$\begin{array}{ccccc} S'_\alpha & \xrightarrow{\varepsilon_\alpha} & C \times_{E_j} S_\alpha & \longrightarrow & S_\alpha \\ & \searrow h_\alpha & \downarrow & & \downarrow h_{\alpha j} \\ & & C & \xrightarrow{h} & E_j, \end{array}$$

where ε_α is a modification and the fibers of h_α are all reduced connected curves, with non-singular components crossing transversally.

We also make a modification $\tau : X' \rightarrow C \times_{E_j} X$ such that there exists a commutative diagram of morphisms between smooth varieties:

$$\begin{array}{ccccc}
 & & X' & & \\
 & \nearrow^{f'_k} & & \nwarrow_{f'_i} & \\
 & S'_i & & S'_k & \\
 & \nwarrow_{h'_{ik}} & & \swarrow_{h'_{ki}} & \\
 E_k & & C & & E_i.
 \end{array}$$

f'_k (top left), f'_i (top right), g'_i (middle left), g'_k (middle right), f' (vertical center), h_i (bottom left), h_k (bottom right), h'_{ik} (diagonal bottom left), h'_{ki} (diagonal bottom right).

Let $\xi_\alpha \in \widehat{E}_\alpha$. By [V], Lemma 3.1,⁴ we have an inclusion

$$(6) \quad f'_*(\omega_{X'/C} \otimes f_i^* P_{\xi_i} \otimes f_k^* P_{\xi_k}) \subseteq h^* f_{j*}(\omega_X \otimes f_i^* P_{\xi_i} \otimes f_k^* P_{\xi_k})$$

of locally free sheaves on C . Since h_α is flat with irreducible general fibers, $S'_i \times_C S'_k$ is irreducible, and we have a surjective morphism

$$g'_{ik} : X' \xrightarrow{(g'_i, g'_k)} S'_i \times_C S'_k.$$

Moreover, h_i and h_k are semistable, hence by [AK], Proposition 6.4, $S'_i \times_C S'_k$ has only rational Gorenstein singularities. After further modification of X' , we may assume that g'_{ik} factors through a desingularization of $S'_i \times_C S'_k$:

$$g'_{ik} : X' \rightarrow Y'_{ik} \xrightarrow{\varepsilon} S'_i \times_C S'_k.$$

By definition of rational Gorenstein singularities, we have $\varepsilon_* \omega_{Y'_{ik}} = \omega_{S'_i \times_C S'_k}$. Since $\omega_{X'/Y'_{ik}}$ is effective, we obtain an inclusion

$$p_i^*(\omega_{S'_i/C} \otimes h'_{ik*} P_{\xi_k}) \otimes p_k^*(\omega_{S'_k/C} \otimes h'_{ki*} P_{\xi_i}) \subseteq g'_{ik*}(\omega_{X'/C} \otimes f_i^* P_{\xi_i} \otimes f_k^* P_{\xi_k})$$

of sheaves on $S'_i \times_C S'_k$. Pushing forward these sheaves to C , we obtain

$$(7) \quad \begin{aligned} h_{i*}(\omega_{S'_i/C} \otimes h'_{ik*} P_{\xi_k}) \otimes h_{k*}(\omega_{S'_k/C} \otimes h'_{ki*} P_{\xi_i}) &\subseteq f'_*(\omega_{X'/C} \otimes f_i^* P_{\xi_i} \otimes f_k^* P_{\xi_k}) \\ &\subseteq h^* f_{j*}(\omega_X \otimes f_i^* P_{\xi_i} \otimes f_k^* P_{\xi_k}), \end{aligned}$$

where the second inclusion comes from (6). Let $\{\alpha, \beta\} = \{i, k\}$. Both sheaves $h_{\alpha*}(\omega_{S'_\alpha/C} \otimes h'_{\alpha\beta*} P_{\xi_\beta})$ are nef ([V], Corollary 3.6). On the other hand, for ξ_i and ξ_k general and torsion, the sheaf in (7) has degree 0 by Proposition 4.7, hence

$$\deg(h_{\alpha*}(\omega_{S'_\alpha/C} \otimes h'_{\alpha\beta*} P_{\xi_\beta})) = 0.$$

By [K1], Corollary 10.15, both sheaves $R^1 h_{\alpha*}(\omega_{S'_\alpha/C} \otimes h'_{\alpha\beta*} P_{\xi_\beta})$ are torsion-free and generically 0, hence 0.⁵ On the other hand, we have $R^1 h_{\alpha*} \omega_{S'_\alpha/C} = \mathcal{O}_C$ ([K2], Proposition 7.6) hence, by the Grothendieck-Riemann-Roch theorem,

$$[\text{ch}(h_{\alpha*}(\omega_{S'_\alpha/C})) - \text{ch}(\mathcal{O}_C)] \text{Td}(C) = \text{ch}(h_{\alpha*}(\omega_{S'_\alpha/C} \otimes h'_{\alpha\beta*} P_{\xi_\beta})) \text{Td}(C)$$

⁴This is proved there for $\xi_i = \xi_k = 0$, but the same proof works in general.

⁵Note that up to this point, the proof works in the more general situation where a_X is surjective and $\text{Alb}(X)$ has a 1-dimensional simple factor E_1 .

in the ring of cycles modulo numerical equivalence on C . This implies $\deg(h_{\alpha*}\omega_{S'_\alpha/C}) = 0$, and h_α is locally trivial (see e.g., [BPV], Theorem III.17.3). Hence $h_{\alpha j}$ is isotrivial (for each $\alpha \in \{i, k\}$). \square

Step 2. *There exist finite groups G_i acting on C_{ij} such that $C_{ij}/G_i \simeq E_j$ and $C_{ik}/G_i \simeq E_k$, the surface S_i is birational to the quotient $(C_{ij} \times C_{ik})/G_i$ for the diagonal action of G_i , and h_{ij} and h_{ik} are identified with the two projections.*

This is a consequence of the following (probably classical) result.

Lemma 5.3. *Let S be a smooth projective surface with an isotrivial fibration $h_1 : S \twoheadrightarrow \Gamma_1$ onto an irrational curve with (constant) irrational general fiber F_1 .*

a) *There exist a smooth curve F_2 and a finite group H acting faithfully on F_1 and F_2 such that Γ_1 is isomorphic to F_2/H , the surface S is birationally isomorphic to the diagonal quotient $(F_1 \times F_2)/H$, and h_1 is the composition $S \xrightarrow{\sim} (F_1 \times F_2)/H \twoheadrightarrow F_2/H \simeq \Gamma_1$. Let h_2 be the composition $S \xrightarrow{\sim} (F_1 \times F_2)/H \twoheadrightarrow F_1/H$.*

b) *Assume S is of general type. Any isotrivial fibration $h : S \twoheadrightarrow \Gamma$ onto an irrational curve Γ is either h_1 or h_2 followed by an isomorphism between F_1/H or F_2/H with Γ .*

Proof. Item a) is well-known and can be found in [S]. Let us prove b). Since Γ is irrational, h induces an isotrivial fibration $h' : (F_1 \times F_2)/H \twoheadrightarrow \Gamma$. Let D_2 be a general (constant irrational) fiber of h' . The quotient map $\pi : F_1 \times F_2 \twoheadrightarrow (F_1 \times F_2)/H$ is étale outside a finite set. Hence the Stein factorization g of $h' \circ \pi$ in the diagram

$$\begin{array}{ccccc} F_1 \times F_2 & \xrightarrow{\pi} & (F_1 \times F_2)/H & \xrightarrow{h'} & \Gamma \\ & \searrow g & & \nearrow & \\ & & D_1 & & \end{array}$$

is also isotrivial, with general fiber D'_2 a (fixed) étale cover of D_2 . By a), there is a base change $D'_1 \twoheadrightarrow D_1$ and a surjective morphism $t = (t_1, t_2) : D'_1 \times D'_2 \twoheadrightarrow F_1 \times F_2$. Since S is of general type, F_1 and F_2 are each of genus ≥ 2 , hence each t_i must factor through one of the projections $p_j : D'_1 \times D'_2 \twoheadrightarrow D'_j$.

If h factors through neither h_1 nor h_2 , the curve D'_2 dominates both F_1 and F_2 , hence t_1 and t_2 cannot factor through p_1 . Thus they must factor through p_2 , which contradicts the fact that t is surjective. \square

Let now Y_j be a resolution of singularities of the irreducible threefold $S_i \times_{E_j} S_k$ and let Y be a resolution of singularities of the component of $Y_1 \times_{E_2 \times E_3} S_1$ that dominates both Y_1

and Y_3 . After modification of X , we obtain a diagram

$$(8) \quad \begin{array}{ccccccc} X & \xrightarrow{g} & Y & & & & \\ & & \swarrow & & \searrow & & \\ & & Y_1 & & Y_3 & & \\ & \swarrow g_{13} & & \swarrow g_{12} & & \swarrow g_{32} & \searrow g_{31} \\ S_3 & & \square & & \square & & S_1 \\ & \swarrow h_{32} & & \swarrow h_{31} & & \swarrow h_{21} & \searrow h_{23} \\ E_2 & & E_1 & & E_3 & & E_2 \end{array}$$

where the squares are birationally cartesian and the isotrivial morphisms $h_{ij} : S_i \rightarrow E_j$ fit into diagrams

$$\begin{array}{ccc} C_{ik} \times C_{ij} & \xrightarrow{p_2} & C_{ij} \\ \downarrow /G_i & & \downarrow /G_i \\ C_{ik} \hookrightarrow S_i & \xrightarrow{h_{ij}} & E_j. \end{array}$$

Step 3. *The threefold Y is dominated by a product of three curves.*

The dominant maps $C_{31} \times C_{32} \dashrightarrow S_3$ and $C_{21} \times C_{23} \dashrightarrow S_2$ induce a factorization

$$((\rho_{31}, \rho_{21}), \rho_{32}, \rho_{23}) : (C_{31} \times_{E_1} C_{21}) \times C_{32} \times C_{23} \dashrightarrow S_3 \times_{E_1} S_2 \twoheadrightarrow E_1 \times E_2 \times E_3.$$

The (Stein factorization of the) morphism $Y_1 \rightarrow E_2 \times E_3$ is therefore isotrivial (its fibers are dominated by the curve $C_{31} \times_{E_1} C_{21}$). Thus, Y is dominated by the product

$$(9) \quad (C_{31} \times_{E_1} C_{21}) \times (C_{12} \times_{E_2} C_{32}) \times (C_{23} \times_{E_3} C_{13})$$

of three (possibly reducible) curves.

Going back to the proof of Theorem 5.1, after passing to an étale cover, we may and will assume, from now on, that the following holds (§2.2):

$$(10) \quad \text{All the irreducible components of } V_0(\omega_X, a_X) \text{ pass through } 0.$$

One checks, using §2.1, §2.2, and Proposition 4.5.d), that if we want X to be birationally covered by a product $\Gamma_1 \times \Gamma_2 \times \Gamma_3$, with morphisms $\Gamma_i \rightarrow E_i$, as in the conclusion of the theorem, we must have the following.

Step 4. *We have*

$$V_0(\omega_X, a_X) = (\widehat{E}_1 \times \widehat{E}_2 \times \{0\}) \cup (\widehat{E}_1 \times \{0\} \times \widehat{E}_3) \cup (\{0\} \times \widehat{E}_2 \times \widehat{E}_3).$$

We already know that $V_0(\omega_X, a_X)$ contains the right-hand-side (Proposition 4.3.b)) and we must prove that it has no other components.

Proof. Assume $V_0(\omega_X, a_X)$ has another component \widehat{T} . It has dimension 2 (Corollary 3.5.a)) and, after possibly permuting the indices, we may assume the neutral component \widehat{E}'_1 of $\widehat{T} \cap (\widehat{E}_1 \times \widehat{E}_2 \times \{0\})$ is neither $\widehat{E}_1 \times \{0\} \times \{0\}$ nor $\{0\} \times \widehat{E}_2 \times \{0\}$. This yields an elliptic curve E'_1 which is a quotient of $E_1 \times E_2$ which does not factor through either projection. As

we saw right before Proposition 4.7, the induced map $f_4 : X \rightarrow E'_1$ is a fibration. It factors as

$$f_4 : X \twoheadrightarrow S_3 \xrightarrow{h_{34}} E'_1$$

where, by Step 1, h_{34} is isotrivial. By Lemma 5.3.b), h_{34} must factor through one of the projections $h_{31} : S_3 \rightarrow E_1$ or $h_{32} : S_3 \rightarrow E_2$ so we reach a contradiction. \square

Step 5. *The morphism $g : X \rightarrow Y$ is birational.*

Proof. Consider, in the diagram (8), the generically finite morphism $v_1 : X \rightarrow Y_1$ and the three fibrations $f'_\alpha : Y_1 \rightarrow E_\alpha$, for $\alpha \in \{1, 2, 3\}$. Since X , Y_1 , and S_3 are all of maximal Albanese dimensions, ω_{X/Y_1} and ω_{Y_1/S_3} are effective, hence

$$h^0(X, \omega_X \otimes a_X^* P_\xi) \geq h^0(Y_1, \omega_{Y_1} \otimes (f'_1, f'_2)^* P_\xi) \geq h^0(S_3, \omega_{S_3} \otimes (h_{31}, h_{32})^* P_\xi)$$

for all $\xi \in \widehat{E}_1 \times \widehat{E}_2$. Moreover, for ξ non-zero, we have by Proposition 4.5.d)

$$h^2(X, \omega_X \otimes a_X^* P_\xi) = h^3(X, \omega_X \otimes a_X^* P_\xi) = 0$$

hence, since $\chi(X, \mathcal{O}_X) = 0$,

$$h^0(X, \omega_X \otimes a_X^* P_\xi) = h^1(X, \omega_X \otimes a_X^* P_\xi).$$

Finally, for ξ general in $\widehat{E}_1 \times \widehat{E}_2$, we have, as in the proof of Theorem 3.1, since $g_3 : X \rightarrow S_3$ has connected fibers,

$$h^1(X, \omega_X \otimes a_X^* P_\xi) = h^0(S_3, \omega_{S_3} \otimes (h_{31}, h_{32})^* P_\xi).$$

Therefore, for $\xi \in \widehat{E}_1 \times \widehat{E}_2$ general, we obtain

$$h^0(X, \omega_X \otimes a_X^* P_\xi) = h^0(Y_1, \omega_{Y_1} \otimes (f'_1, f'_2)^* P_\xi).$$

The induced morphism $Y \rightarrow E_1 \times E_2 \times E_3$ is the Albanese mapping of Y . Since in any event, we always have

$$h^0(X, \omega_X \otimes a_X^* P_\xi) \geq h^0(Y, \omega_Y \otimes a_Y^* P_\xi) \geq h^0(Y_1, \omega_{Y_1} \otimes (f'_1, f'_2)^* P_\xi)$$

for all $\xi \in \widehat{E}_1 \times \widehat{E}_2$, we obtain

$$(11) \quad h^0(X, \omega_X \otimes a_X^* P_\xi) = h^0(Y, \omega_Y \otimes a_Y^* P_\xi)$$

for ξ general in $\widehat{E}_1 \times \widehat{E}_2$, hence also, by Step 3, for ξ general in $V_0(\omega_X, a_X)$. But for $\xi \notin V_0(\omega_X, a_X)$, both sides of (11) vanish. By Lemma 5.4 below, we conclude that g is a birational morphism. \square

The following lemma (used in the proof above) is in the spirit of [HP], Theorem 3.1.

Lemma 5.4. *Let $X \xrightarrow{g} Y \xrightarrow{f} A$ be generically finite morphisms between smooth projective threefolds, where A is an abelian threefold, such that f and $f \circ g$ are both minimal. Assume that X is of general type with $\chi(X, \omega_X) = 0$ and that there exists an open subset $U \subseteq \widehat{A}$ with $\text{codim}_{\widehat{A}}(\widehat{A} - U) \geq 2$ such that*

$$h^0(X, \omega_X \otimes g^* f^* P_\xi) = h^0(Y, \omega_Y \otimes f^* P_\xi)$$

for all $\xi \in U$. Then g is birational.

Proof. By §2.2, we can write $g_*\omega_X \simeq \omega_Y \oplus \mathcal{E}$, and we need to show that the sheaf \mathcal{E} is zero. Since \mathcal{E} is torsion-free and f is generically finite, it is sufficient to prove $f_*\mathcal{E} = 0$.

As we saw at the beginning of §5, we have $q(X) = 3$, hence $f \circ g$ is the Albanese mapping of X . By Proposition 4.5.d), for each $i \in \{2, 3\}$, we then have $\{0\} = V_i(\omega_X, f \circ g) = V_i(g_*\omega_X, f)$, hence $V_i(f_*\mathcal{E}) \subseteq \{0\}$.

Since $q(X) = 3$, we also have $q(Y) = 3$, hence $h^i(Y, g_*\omega_X) = h^i(Y, \omega_Y)$. It follows that $V_i(f_*\mathcal{E})$ is empty.

The assumption $\chi(X, \omega_X) = 0$ implies $\chi(Y, \omega_Y) = 0$ by (1). Thus,

$$V_0(f_*\mathcal{E}) = V_1(f_*\mathcal{E}) \subseteq \widehat{A} - U.$$

Since $\text{codim}_{\widehat{A}}(\widehat{A} - U) > 1$, the sheaf $f_*\mathcal{E}$ is therefore M-regular in the sense of [PP], Definition 2.1 (see also Remark 2.3), hence continuously globally generated ([PP], Definition 2.10 and Proposition 2.13). Since $H^0(A, f_*\mathcal{E} \otimes P_\xi) = 0$ for all $\xi \in U$, we obtain $f_*\mathcal{E} = 0$. \square

Let us summarize what we know. Let $\{i, j, k\} = \{1, 2, 3\}$. The curve C_{ij} is the (constant) general fiber of the isotrivial fibration $S_i \rightarrow E_k$; it is acted on by a group G_i and $C_{ij}/G_i \simeq E_j$ (Step 2). The fibration $g_i : X \rightarrow S_i$ is also isotrivial; as we saw in Step 3, its general fiber C_i is dominated by the curve $C_{ji} \times_{E_i} C_{ki}$ but also maps onto C_{ji} and C_{ki} . Finally, a general fiber F_k of the isotrivial fibration $f_k : X \rightarrow E_k$ is an isotrivial fibration over C_{ij} with (constant) general fiber C_i . The situation is summarized in the following diagram:

$$\begin{array}{ccccc}
 & C_{ij} & \xrightarrow{\text{fiber}} & S_i & \\
 \text{fiber } C_i \nearrow & & & \nearrow g_i & \searrow h_{ik} \\
 F_k \hookrightarrow & \xrightarrow{\text{fiber}} & X & \xrightarrow{f_k} & E_k \\
 \text{fiber } C_j \searrow & & & \searrow g_j & \nearrow h_{jk} \\
 & C_{ji} & \xrightarrow{\text{fiber}} & S_j &
 \end{array}$$

By Lemma 5.3, there exists a finite group H_k acting faithfully on C_i and C_j such that $C_{ij} \simeq C_j/H_k$, $C_{ji} \simeq C_i/H_k$, and F_k is isomorphic to the diagonal quotient $(C_i \times C_j)/H_k$. Moreover, the maps to C_{ij} and C_{ji} are the natural projections. So we have diagrams

$$(12) \quad \begin{array}{ccc}
 & C_{ji} & \\
 /H_k \nearrow & & \searrow /G_j \\
 C_i & & E_i \\
 /H_j \searrow & & \nearrow /G_k \\
 & C_{ki} &
 \end{array}$$

Let D_1 be the Galois closure of C_1 over E_1 and set $G = \text{Gal}(D_1/E_1)$. Let $\{j, k\} = \{2, 3\}$. There is a normal subgroup $N_j \triangleleft G$ such that $G_j = G/N_j$ and G acts on C_{jk} via this quotient. By Step 2, the surface S_j is birationally isomorphic to $(C_{j1} \times C_{jk})/G_j$, hence to $(D_1 \times C_{jk})/G$. Therefore, the modification Y_1 of $S_2 \times_{E_1} S_3$ (see (8)) is birationally isomorphic to $(D_1 \times C_{23} \times C_{32})/G$.

Step 6. *The group G is isomorphic to $\mathbf{Z}/2\mathbf{Z}$.*

We begin with a lemma which is probably well-known. We denote by $\text{Irr}(G)$ the set of isomorphism classes of irreducible representations of G .

Lemma 5.5. *Let E be an elliptic curve and let $\pi : D \rightarrow E$ be a Galois cover with group G . We can write*

$$\pi_* \mathcal{O}_D = \bigoplus_{\chi \in \text{Irr}(G)} \bigoplus_i \mathcal{V}_{\chi,i},$$

where each vector bundle $\mathcal{V}_{\chi,i}$ is semistable and G -invariant, and the representation of G on the general fiber of each $\mathcal{V}_{\chi,i}$ is a direct sum of χ . Moreover, for each $\chi \neq 1$, the dual vector bundle $\mathcal{V}_{\chi,i}^\vee$ is either ample or a direct sum of non-zero torsion line bundles.

Proof. The groups G acts on $\pi_* \mathcal{O}_D$. Identifying each representation χ with its character, we consider the endomorphism $\sum_{g \in G} \chi(g)g$ of $\pi_* \mathcal{O}_D$ and we denote by \mathcal{V}_χ^\vee its image. We then have

$$(13) \quad \pi_* \mathcal{O}_D = \bigoplus_\chi \mathcal{V}_\chi,$$

where the general fiber of \mathcal{V}_χ is a (nonzero) direct sum of χ as a G -module.

The Harder-Narasimhan filtration

$$0 = \mathcal{V}_\chi^\ell \subseteq \mathcal{V}_\chi^{\ell-1} \subseteq \dots \subseteq \mathcal{V}_\chi^0 = \mathcal{V}_\chi$$

is preserved by the G -action. Serre duality shows that since we are on an elliptic curve, all the corresponding extensions are trivial, hence \mathcal{V}_χ is the direct sum of the G -invariant semistable bundles $\mathcal{V}_{\chi,i} = \mathcal{V}_\chi^i / \mathcal{V}_\chi^{i+1}$, for $0 \leq i < \ell$.

As a direct summand of $\pi_* \omega_D = (\pi_* \mathcal{O}_D)^\vee$, each vector bundle $\mathcal{V}_{\chi,i}^\vee$ is nef ([V], Corollary 3.6). Moreover, it is ample if it has positive degree. Consider the maximal degree-0 subsheaf \mathcal{F} of $\pi_* \mathcal{O}_D$, i.e., the direct sum of all $\mathcal{V}_{\chi,i}$ that have degree 0. By [KP], Lemma 3.2 and 3.4, \mathcal{F} is a G -invariant subalgebra and induces an étale cover of E , hence is a direct sum of torsion line bundles. \square

Let us continue with the Galois cover $\pi : D \rightarrow E$ with group G as in the lemma and assume moreover that for each $j \in \{2, 3\}$, we have a Galois cover $\pi_j : D_j \rightarrow E_j$ with Galois group $G_j = G/N_j$, where $g(D_j) \geq 2$ and E_j is an elliptic curve.

Then G acts on $D_1 \times D_2 \times D_3$ diagonally. Let Z be the quotient, let $\varepsilon : Y \rightarrow Z$ be a resolution, and consider

$$t : Y \xrightarrow{\varepsilon} Z \rightarrow E_1 \times E_2 \times E_3.$$

Lemma 5.6. *Assume*

$$V_0(\omega_Y, t) \subseteq (\widehat{E}_1 \times \widehat{E}_2 \times \{0\}) \cup (\widehat{E}_1 \times \{0\} \times \widehat{E}_3) \cup (\{0\} \times \widehat{E}_2 \times \widehat{E}_3)$$

and $V_2(\omega_Y, t) \cup V_3(\omega_Y, t) \subseteq \{0\}$. Then $N_2 = N_3$ and $G_2 \simeq G_3 \simeq \mathbf{Z}/2\mathbf{Z}$.

Proof. We decompose $\pi_* \mathcal{O}_D$ as in (13) and write similarly

$$\pi_{j*} \mathcal{O}_{D_j} = \bigoplus_{\mu \in \text{Irr}(G_j)} \bigoplus_i \mathcal{V}_{\mu,i}^j.$$

Since quotient singularities are rational, we have as in Example 4.1

$$t_*\omega_Y \simeq (q_*\mathcal{O}_Z)^\vee \simeq ((\pi_*\mathcal{O}_D)^\vee \boxtimes ((\pi_{2*}\mathcal{O}_{D_2})^\vee \boxtimes ((\pi_{3*}\mathcal{O}_{D_3})^\vee)^G.$$

Let μ be a non-trivial element of $\text{Irr}(G_2)$. Since G_2 is a quotient of G , the representation μ and its complex conjugate $\bar{\mu}$ are also in $\text{Irr}(G)$. Then, the vector bundle

$$\mathcal{G} := (\mathcal{V}_{\bar{\mu},1}^\vee \boxtimes \mathcal{V}_{\mu,1}^{2\vee} \boxtimes \mathcal{O}_{E_3})^G$$

on $E_1 \times E_2 \times E_3$ is a non-zero direct summand of both $\mathcal{V}_{\bar{\mu},1}^\vee \boxtimes \mathcal{V}_{\mu,1}^{2\vee} \boxtimes \mathcal{O}_{E_3}$ and $t_*\omega_Y$.

Assume that $\mathcal{V}_{\mu,1}^2$ has degree 0, hence is a direct sum of non-trivial torsion line bundles.

- If $\deg(\mathcal{V}_{\bar{\mu},1}^\vee) = 0$, the sheaf \mathcal{G} is a direct sum of non-trivial torsion line bundles, which is impossible since $V_3(\mathcal{G}) \subseteq V_3(\omega_Y, t) = \{0\}$.
- If $\mathcal{V}_{\bar{\mu},1}^\vee$ is ample, we can write

$$\mathcal{G} = \bigoplus_k (\mathcal{G}_k \boxtimes P_{\xi_k} \boxtimes \mathcal{O}_{E_3}),$$

where \mathcal{G}_k is a direct summand of $\mathcal{V}_{\bar{\mu},1}^\vee$, hence ample, and the ξ_k are non-zero torsion points in \widehat{E}_2 . This is again impossible, because $V_2(\mathcal{G}) \subseteq V_2(\omega_Y, t) = \{0\}$.

Therefore, $\mathcal{V}_{\mu,1}^{j\vee}$ is ample for all μ non-trivial in $\text{Irr}(G_j)$.

If $\text{Card}(G_2) > 2$, or if $N_2 \neq N_3$, we may take non-trivial $\chi \in \text{Irr}(G)$, $\mu \in \text{Irr}(G_2)$, and $\nu \in \text{Irr}(G_3)$ such that χ is a subrepresentation of $\mu \otimes \nu$. The vector bundle

$$\mathcal{H} := (\mathcal{V}_{\bar{\chi},1}^\vee \boxtimes \mathcal{V}_{\mu,1}^{2\vee} \boxtimes \mathcal{V}_{\nu,1}^{3\vee})^G$$

is then non-zero and a direct summand of $t_*\omega_Y$ (and $\mathcal{V}_{\mu,1}^{2\vee}$ and $\mathcal{V}_{\nu,1}^{3\vee}$ are ample).

If $\mathcal{V}_{\bar{\chi},1}^\vee$ is ample, since \mathcal{H} is a direct summand of $\mathcal{V}_{\bar{\chi},1}^\vee \boxtimes \mathcal{V}_{\mu,1}^{2\vee} \boxtimes \mathcal{V}_{\nu,1}^{3\vee}$, we have $V_m(\mathcal{H}) = \emptyset$ for all $m \in \{1, 2, 3\}$. Hence $h^0(E_1 \times E_2 \times E_3, \mathcal{H} \otimes P_\xi)$ is a non-zero constant for all $\xi \in \widehat{E}_1 \times \widehat{E}_2 \times \widehat{E}_3$ and $V_0(\mathcal{H}) = \widehat{E}_1 \times \widehat{E}_2 \times \widehat{E}_3$, which contradicts our assumptions.

If $\mathcal{V}_{\bar{\chi},1}^\vee$ is a direct sum of non-trivial torsion line bundles, we may write

$$\mathcal{H} = \bigoplus_k (P_{\xi_k} \boxtimes \mathcal{H}_k),$$

where the ξ_k are non-zero torsion points in \widehat{E}_1 and \mathcal{H}_k is a direct summand of $\mathcal{V}_{\mu,1}^{2\vee} \boxtimes \mathcal{V}_{\nu,1}^{3\vee}$. Then $V_0(\mathcal{H})$, hence also $V_0(\omega_Y, t)$, contains $\{-\xi_1\} \times \widehat{E}_2 \times \widehat{E}_3$, which contradicts our assumptions. \square

We now apply this second lemma to the Galois covers $\pi : D_1 \twoheadrightarrow E_1$, $\pi_2 : C_{23} \twoheadrightarrow E_2$, and $\pi_3 : C_{32} \twoheadrightarrow E_3$. The variety Y of the lemma is the variety Y_1 of the proof, and since $V_0(\omega_{Y_1}, t) \subseteq V_0(\omega_X, a_X)$ (see §2.2), the hypotheses of the lemma are satisfied (Step 4).

We obtain $N_2 = N_3$, hence the coverings $C_{ji} \twoheadrightarrow E_i$ and $C_{ki} \twoheadrightarrow E_i$ are the same (see (12)), and also $G/N_j \simeq \mathbf{Z}/2\mathbf{Z}$, so that they are double covers. Denote them by $C'_i \twoheadrightarrow E_i$. By the proof of Step 3 (see (9)), X is birational to $(C'_1 \times C'_2 \times C'_3)/(\mathbf{Z}/2\mathbf{Z})$. Since the latter

variety contains no rational curves, there is a birational *morphism* from X to it. This finishes the proof of Theorem 5.1. \square

6. VARIETIES WITH $P_1 = 1$

It follows from [U] and 2.1.5 that varieties X of maximal Albanese dimension and $P_1(X) = 1$ satisfy $\chi(X, \omega_X) = 0$. We presented in Example 4.2 a construction of Chen and Hacon of such a variety which is in addition of general type. We gather here some properties of these varieties (most of them taken from [U]).

Proposition 6.1. *Let X be a smooth projective variety of maximal Albanese dimension n , with $P_1(X) = 1$.*

a) *We have an isomorphism*

$$a_X^* : \bigwedge^{\bullet} H^0(A, \Omega_A) \simeq H^0(X, \Omega_X^{\bullet}).$$

In particular, $h^j(X, \mathcal{O}_X) = \binom{n}{j}$ for all j , hence $\chi(X, \omega_X) = 0$, and the Albanese mapping $a_X : X \rightarrow \text{Alb}(X)$ is surjective.

b) *The point 0 is isolated in $V_0(\omega_X, a_X)$.*

Proof. Replacing X with a modification, we may assume that there is a factorization $a_X : X \twoheadrightarrow Z \rightarrow \text{Alb}(X)$, where Z is a desingularization of $a_X(X)$, so that $P_1(Z) \leq P_1(X) = 1$. It follows from [U] (or [M], Corollary (3.5)) that $a_X(X)$ is a translate of an abelian subvariety of $\text{Alb}(X)$, hence a_X is surjective. Item a) then follows from another result of Ueno ([U], or [M], Corollary (3.4)).

By §2.2, we can write $a_{X*}\omega_X \simeq \omega_A \oplus \mathcal{E} \simeq \mathcal{O}_A \oplus \mathcal{E}$. The sheaf \mathcal{E} then satisfies $V_i(\mathcal{E}) - \{0\} = V_i(\omega_X, a_X) - \{0\}$ for all i . Since $1 = P_1(X) = 1 + h^0(A, \mathcal{E})$, the point 0 is not in the closed set $V_0(\mathcal{E})$, hence is isolated in $V_0(\omega_X, a_X)$. This proves b). \square

Remark 6.2. Regarding item b), to be more precise, a smooth projective variety X of maximal Albanese dimension satisfies $P_1(X) = 1$ *if and only if* 0 is isolated in $V_0(\omega_X, a_X)$.

Theorem 6.3. *Let X be a smooth projective threefold of maximal Albanese dimension and of general type. If $P_1(X) = 1$, the variety X is a modification of an abelian étale cover of a Chen-Hacon threefold.*

It is then very easy to describe all smooth projective threefolds X of maximal Albanese dimension and of general type, with $P_1(X) = 1$. Start from a Chen-Hacon threefold Y as in Example 4.2, with Albanese mapping $a_Y : Y \twoheadrightarrow E_1 \times E_2 \times E_3$. It satisfies

$$V_0(\omega_Y, a_Y) = \{0\} \cup (\widehat{E}_1 \times \widehat{E}_2 \times \{\xi_3\}) \cup (\widehat{E}_1 \times \{\xi_2\} \times \widehat{E}_3) \cup (\{\xi_1\} \times \widehat{E}_2 \times \widehat{E}_3),$$

where each $\xi_j \in \widehat{E}_j$ has order 2. Take an isogeny $A \twoheadrightarrow E_1 \times E_2 \times E_3$ corresponding to a (finite) subgroup of $\widehat{E}_1 \times \widehat{E}_2 \times \widehat{E}_3$ which contains none of the points ξ_1, ξ_2, ξ_3 . Finally, take for X a modification of $Y \times_{E_1 \times E_2 \times E_3} A$.

Proof of the theorem. Replacing a_X with its Stein factorization, we will assume that X is normal and a_X is finite. By Theorem 5.1, there exist elliptic curves E_1, E_2 , and E_3 , double coverings $\rho_i : C_i \rightarrow E_i$, with involution ι_i , and a commutative diagram

$$\begin{array}{ccc}
C_1 \times C_2 \times C_3 & & \\
\downarrow & \searrow^{(\rho_1, \rho_2, \rho_3)} & \\
\tilde{X} & \xrightarrow{a_{\tilde{X}}} & E_1 \times E_2 \times E_3 \\
\downarrow & \square & \downarrow \eta \\
X & \xrightarrow{a_X} & \text{Alb}(X),
\end{array}$$

where η is an isogeny (the variety \tilde{X} is the variety Z of Example 4.1) and both a_X and $a_{\tilde{X}}$ are $(\mathbf{Z}/2\mathbf{Z})^2$ -Galois coverings. In particular, X has rational singularities.

We denote by K the (finite) kernel of η and by K_i the image of the projection $K \rightarrow E_i$, so that K is a subgroup of $\tilde{K} := K_1 \times K_2 \times K_3$. The elliptic curve $F_i := E_i/K_i$ embeds in $\text{Alb}(X)$; let $\pi_i : \text{Alb}(X) \rightarrow F_i$ be the quotient. The natural morphism $h_i : X \rightarrow F_i$ is an isotrivial fibration and we denote by D_i its general (constant) fiber. Now we consider the square restricted to fibers:

$$\begin{array}{ccc}
C_i & \xrightarrow[2:1]{\rho_i} & E_i \\
\text{étale} \downarrow & & \downarrow \lambda_i \text{ étale} \\
D_i & \xrightarrow[4:1]{t_i} & F_i,
\end{array}$$

where t_i is a $(\mathbf{Z}/2\mathbf{Z})^2$ -cover. Since $D_i \times_{F_i} E_i$ is disconnected and factors as $C_i \sqcup C_i \rightarrow C_i \xrightarrow{\rho_i} E_i$, there is a non-zero 2-torsion point $\xi_i \in \tilde{F}_i$ such that $\xi_i \in \text{Ker}(t_i^*) \cap \text{Ker}(\lambda_i^*)$, the morphism t_i factors as $D_i \xrightarrow{s_i} D'_i \rightarrow F_i$, where s_i is a double étale cover, and $C_i \simeq D'_i \times_{F_i} E_i$. It follows that the group K_i acts on C_i , and the involution ι_i and the K_i -action commute. Therefore, \tilde{K} acts on $C_1 \times C_2 \times C_3$ and this action commutes with the involution $(\iota_1, \iota_2, \iota_3)$. It follows that \tilde{K} acts on \tilde{X} and the Albanese mapping $a_{\tilde{X}}$ is \tilde{K} -equivariant. Set $Y := \tilde{X}/\tilde{K}$.

Lemma 6.4. *The quotient morphism $\tilde{X} \rightarrow Y$ is étale and factors as $\tilde{X} \twoheadrightarrow X \rightarrow Y$. The variety Y has maximal Albanese dimension, is of general type, $P_1(Y) = 1$, and its Albanese variety is $F_1 \times F_2 \times F_3$.*

Proof. We have a cartesian diagram

$$(14) \quad \begin{array}{ccc}
\tilde{X} & \xrightarrow{a_{\tilde{X}}} & E_1 \times E_2 \times E_3 \\
\downarrow / \tilde{K} & \square & \downarrow \lambda / \tilde{K} \\
Y & \xrightarrow{a_Y} & F_1 \times F_2 \times F_3.
\end{array}$$

Since the rightmost quotient morphism is étale, so is the leftmost quotient morphism. Since the top morphism is the Albanese mapping of \tilde{X} , the bottom morphism is the Albanese

mapping of Y . Furthermore, since K is a subgroup of \tilde{K} , the leftmost quotient morphism factors as $\tilde{X} \twoheadrightarrow X \twoheadrightarrow Y$. Therefore, $P_1(Y) = 1$. \square

We now claim that Y is a Chen-Hacon threefold. By Theorem 5.1 (or the diagram (14)), a_Y is a $(\mathbf{Z}/2\mathbf{Z})^2$ -Galois covering and by [P], we can write

$$a_{Y*}\mathcal{O}_Y = \bigoplus_{\chi \in (\mathbf{Z}/2\mathbf{Z})^{2*}} L_\chi^\vee = \mathcal{O}_{\text{Alb}(Y)} \oplus L_{\chi_1}^\vee \oplus L_{\chi_2}^\vee \oplus L_{\chi_3}^\vee,$$

or equivalently, since Y has rational singularities,

$$(15) \quad a_{Y*}\omega_Y = \mathcal{O}_{F_1 \times F_2 \times F_3} \oplus L_{\chi_1} \oplus L_{\chi_2} \oplus L_{\chi_3},$$

where L_{χ_1} , L_{χ_2} , and L_{χ_3} are line bundles on $F_1 \times F_2 \times F_3$. Moreover, by [P], Theorem 2.1, we have the following ‘‘building data’’: there are effective divisors D_1 , D_2 , and D_3 on $F_1 \times F_2 \times F_3$ satisfying:

$$L_{\chi_i} + L_{\chi_j} \sim_{\text{lin}} L_{\chi_k} + D_k \quad \text{and} \quad L_{\chi_i}^2 \sim_{\text{lin}} D_j + D_k$$

for any $\{i, j, k\} = \{1, 2, 3\}$. These data pull back to the analogous building data on \tilde{X} , hence λ^*D_i is the pull-back on $E_1 \times E_2 \times E_3$ of the branch divisor $\Delta_i \sim_{\text{lin}} 2\delta_i$ of ρ_i . It follows that there exists an ample line bundle δ'_i on F_i which pulls back to δ_i on E_i and such that D_i is also the pull-back on $F_1 \times F_2 \times F_3$ of a divisor $\Delta'_i \sim_{\text{lin}} 2\delta'_i$ on F_i . Let L'_i be the pull-back on $F_1 \times F_2 \times F_3$ of δ'_i . Because of the relations $L_{\chi_i}^2 \sim_{\text{lin}} D_j + D_k$, we can write

$$L_{\chi_i} \simeq P_{\xi_i} \otimes (L'_j \otimes P_{\xi_{i,j}}) \otimes (L'_k \otimes P_{\xi_{i,k}}),$$

where $\xi_i \in \widehat{E}_i$, $\xi_{i,j} \in \widehat{E}_j$, and $\xi_{i,k} \in \widehat{E}_k$ are 2-torsion points. From (15) and the fact that $P_1(Y) = 1$, we deduce $H^0(E_i, L_{\chi_i}) = 0$, hence each ξ_i has order 2 and is in the kernel of $\widehat{\lambda}_i$. From the relations $L_{\chi_i} + L_{\chi_j} \sim_{\text{lin}} L_{\chi_k} + D_k$, we deduce

$$\xi_{i,k} + \xi_{j,k} = \xi_k \quad \text{and} \quad \xi_{i,j} + \xi_j = \xi_{k,j}.$$

Since $\lambda_1^*\xi_1 = 0$, we may always change L'_1 to $L'_1 \otimes P_{\xi_1}$, so we may assume $\xi_{3,1} = 0$ and similarly, $\xi_{1,2} = 0$ and $\xi_{2,3} = 0$. The $\mathcal{O}_{F_1 \times F_2 \times F_3}$ algebra $a_{Y*}\mathcal{O}_Y$ is then the algebra associated to a Chen-Hacon threefold (see (4)). We conclude that Y is a Chen-Hacon threefold. \square

7. A CONJECTURE

As mentioned in the introduction, we end this article with a conjecture on the possible general structure of smooth projective varieties X of maximal Albanese dimension, of general type, with $\chi(X, \omega_X) = 0$.

Conjecture. *Let X be a smooth projective variety of maximal Albanese dimension, of general type, with $\chi(X, \omega_X) = 0$. Then there exist a smooth projective variety X' , a morphism $X' \twoheadrightarrow X$ which is a composition of modifications and abelian étale covers, and a fibration $g: X' \twoheadrightarrow Y$ with general fiber F , such that $0 < \dim(Y) < \dim(X)$ and*

- a) *either g is isotrivial;*
- b) *or $\chi(F, \omega_F) = 0$.*

Remarks 7.1. 1) Conversely, in the situation b) above, $\chi(X, \omega_X) = 0$ ([HP], Proposition 2.5). Moreover, $\text{Alb}(X)$ has at least 4 simple factors by Corollary 3.5.b) and Proposition 4.5.b). Of course, in case a), without further constraints, one might have $\chi(X, \omega_X) > 0$, but we were unable to find necessary and sufficient conditions on the isotrivial fibration g (assuming X does not fall into case b)) to ensure $\chi(X, \omega_X) = 0$.

3) If we are *not* in case b), it follows from Lemma 4.6 that if $X' \rightarrow X$ is any composition of modifications and abelian étale covers, we have $q(X') = \dim(X)$ and any morphism from X' to a curve of genus ≥ 2 is constant.

The Ein-Lazarsfeld example (Example 4.1) falls into case a) of the conjecture, and not into case b) by Remark 7.1.1) above. We present an example that falls into case b), but not into case a). It is basically a non-isotrivial fibration whose general fibers are Ein-Lazarsfeld threefolds.

Example 7.2. Consider a smooth projective curve C of genus ≥ 2 , elliptic curves $E_1, E_2,$ and E_3 , and smooth double coverings $S_j \rightarrow C \times E_j$ ramified along ample divisors. Denote by ι_j the corresponding involution of S_j . We may moreover assume that the fibrations $f_j : S_j \rightarrow C$ are all semistable and not isotrivial.

The fourfold $T := S_1 \times_C S_2 \times_C S_3$ has only rational Gorenstein singularities, and so does its quotient Z by the involution $\iota_1 \times \iota_2 \times \iota_3$. Let $\varepsilon : X \rightarrow Z$ be a desingularization. We have a diagram

$$\begin{array}{ccccccc}
 & & & & & & T \\
 & & & & & & \downarrow g \quad 2:1 \\
 & & & & & & Z \xleftarrow{\varepsilon} X \\
 & & & & & & \downarrow f \quad 4:1 \\
 S_1 & \leftarrow & S_2 & \leftarrow & S_3 & \leftarrow & C \times E_1 \times E_2 \times E_3 \\
 \downarrow 2:1 & & \downarrow 2:1 & & \downarrow 2:1 & & \downarrow \\
 C \times E_1 & & C \times E_2 & & C \times E_3 & & C \\
 & & & & & & \downarrow \\
 & & & & & & C
 \end{array}$$

The variety X is of general type and has maximal Albanese dimension because $C \times E_1 \times E_2 \times E_3$ does. A general fiber of the fibration $X \rightarrow C$ is one of the examples constructed in Example 4.1, hence $\chi(X, \omega_X) = 0$ by [HP], Proposition 2.5, and X falls into case b) of the conjecture. One can prove that it does not fall into case a).

REFERENCES

- [AK] Abramovich, D., Karu, K., Weak semistable reduction in characteristic 0, *Invent. Math.* **139** (2000), 241–273.
- [BPV] Barth, W., Peters, C., Van de Ven, A., *Compact complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete **4** (1984), Springer Verlag, Berlin-Heidelberg-New York.
- [CH1] Chen, J.A., Hacon, C.D., Pluricanonical maps of varieties of maximal Albanese dimension, *Math. Ann.* **320** (2001), 367–380.
- [CH2] Chen, J.A., Hacon, C.D., On the irregularity of the image of the Iitaka fibration, *Comm. Algebra* **32** (2004), 203–215.

- [EL] Ein, L., Lazarsfeld, R., Singularities of theta divisors and the birational geometry of irregular varieties, *J. Amer. Math. Soc.* **10** (1997), 243–258.
- [GL] Green, M., Lazarsfeld, R., Deformation theory, generic vanishing theorems, and some conjectures of Enriques, Catanese and Beauville, *Invent. Math.* **90** (1987), 389–407.
- [HP] Hacon, C.D., Pardini, R., Birational characterization of products of curves of genus 2, *Math. Research Letters* **12** (2005), 129–140.
- [J] Jiang, Z., Varieties with $q(X) = \dim(X)$ and $P_2(X) = 2$, eprint [arXiv:1010.4709v1](https://arxiv.org/abs/1010.4709v1), *Ann. Scuola Norm. Sup. Pisa*, to appear.
- [KP] Kebekus, S., Peternell, T., A refinement of Stein factorization and deformations of surjective morphisms, *Asian J. Math.* **12** (2008), 365–389.
- [KKMS] Kempf, G., Knudsen, F., Mumford, D., Saint-Donat, B., *Toroidal Embeddings I*, Springer Lecture Notes in Mathematics **339**, 1973.
- [K1] Kollár, J., *Shafarevich Maps and Automorphic Forms*, Princeton University Press, 1995.
- [K2] Kollár, J., Higher direct images of dualizing sheaves. I, *Ann. of Math.* **123** (1986), 11–42.
- [K3] Kollár, J., Subadditivity of the Kodaira dimension: fibers of general type, in *Algebraic geometry, Sendai, 1985*, 361–398, Adv. Stud. Pure Math. **10**, North-Holland, Amsterdam, 1987.
- [M] Mori, S., Classification of higher-dimensional varieties, in *Algebraic geometry, Bowdoin, 1985* (Brunswick, Maine, 1985), 269–331, Proc. Sympos. Pure Math. **46**, Part 1, Amer. Math. Soc., Providence, RI, 1987.
- [Mu] Mukai, S., Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves, *Nagoya Math. J.* **81** (1981), 153–175.
- [P] Pardini, R., Abelian covers of algebraic varieties, *J. reine angew. Math.* **417** (1991), 191–213.
- [PP] Pareschi, G., Popa, M., Regularity on abelian varieties I, *J. Amer. Math. Soc.* **16** (2003), 285–302.
- [S] Serrano, F., Isotrivial fibred surfaces, *Ann. Mat. Pura Appl. (4)* **171** (1996), 63–81.
- [Si] Simpson, C., Subspaces of moduli spaces of rank one local systems, *Ann. Sci. École Norm. Sup.* **26** (1993), 361–401.
- [U] Uneo, K., *Classification theory of algebraic varieties and compact complex spaces*, notes written in collaboration with P. Cherenack, Springer Lecture Notes in Mathematics **439**, Berlin-New York, 1975.
- [V] Viehweg, E., *Positivity of direct image sheaves and applications to families of higher dimensional manifolds*, ICTP Lecture Notes **6** (2001), 249–284.

TAIPEI INSTITUTE FOR MATHEMATICAL SCIENCES, NATIONAL CENTER FOR THEORETICAL SCIENCES, TAIPEI OFFICE, AND DEPARTMENT OF MATHEMATICS, 1 SEC. 4, ROOSEVELT RD. TAIPEI 106, TAIWAN

E-mail address: jkchen@math.ntu.edu.tw

DÉPARTEMENT MATHÉMATIQUES ET APPLICATIONS, UMR CNRS 8553, ÉCOLE NORMALE SUPÉRIEURE, 45 RUE D’ULM, 75230 PARIS CEDEX 05, FRANCE

E-mail address: olivier.debarre@ens.fr

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, 53111 BONN, GERMANY

E-mail address: flipz@mpim-bonn.mpg.de