

QUATERNIONIC MODULAR FORMS OF ANY WEIGHT

RICCARDO BRASCA

ABSTRACT. In this work we construct an eigencurve for p -adic modular forms attached to an indefinite quaternion algebra over \mathbb{Q} . Our theory includes the definition, both as rules on test objects and sections of line bundle, of p -adic modular forms, convergent and overconvergent, of any p -adic weight. We prove that our modular forms can be put in analytic families over the weight space and we introduce the Hecke operators U and T_l , that can also be put in families. We show that the U -operator acts compactly on the space of overconvergent modular forms. We finally construct the eigencurve, a rigid analytic variety whose points correspond to systems of overconvergent eigenforms of finite slope with respect to the U -operator.

INTRODUCTION

Let D be an indefinite quaternion algebra over \mathbb{Q} . Under certain hypothesis, the Jacquet-Langlands correspondence gives a map from the space of modular forms for D to the space of modular forms for GL_2/\mathbb{Q} . In [Kas99], Kassaei developed the theory of p -adic modular forms (where p is a prime that does not divide δ , the discriminant of D), of integral weight, attached to D . The Jacquet-Langlands correspondence works also for p -adic modular forms, of level coprime with p . Since nowadays the theory of p -adic modular forms for GL_2/\mathbb{Q} is well developed, it is natural to seek for a similar theory also for modular forms relative to D . One of the most important result in the elliptic case is the existence of the eigencurve: a rigid space whose points correspond to systems of overconvergent eigenforms (of any p -adic weight) of finite slope with respect to the U_p -operator. The final goal of this paper is the construction of the eigencurve for modular forms attached to D . We are going to use the general machinery developed by Buzzard in [Buz07]. We therefore need a weight space \mathcal{W} and a notion of analytic family of modular forms over \mathcal{W} . Our weights are continuous characters $\mathbb{Z}_p^* \rightarrow K^*$, where K is a finite extension of \mathbb{Q}_p , and we take as \mathcal{W} the rigid analytic curve such that $\mathcal{W}(K) = \mathrm{Hom}_{\mathrm{cont}}(\mathbb{Z}_p^*, K^*)$. We define the space of modular forms, both convergent and overconvergent, with respect to D , level N coprime with p , with coefficients in K (where K is big enough), and weight any $\chi \in \mathcal{W}(K)$. We then show that this modular forms can be put in families. We have the Hecke operators T_l , for l a prime not dividing δN , and the U -operator, that acts completely continuously on the space of overconvergent modular forms. The main result of the paper (Theorem 6.5) is the following

Theorem. *There is a rigid space $\mathcal{C} \subseteq \mathcal{W} \times \mathbb{A}_K^{1,\mathrm{rig}}$, called the eigencurve, whose L -points, where L is a finite extension of K , correspond naturally to systems of*

Date: June 26, 2012.

2010 Mathematics Subject Classification. Primary: 11F85; Secondary: 14G35, 11G18.

Key words and phrases. p -adic modular forms, quaternionic modular forms, modular forms of non-integral weight.

eigenvalues of overconvergent modular forms of finite slope with respect to the U-operator, defined over L . Let π_i be the projection to the i -th factor. If $x \in \mathcal{C}(L)$, let \mathcal{M}_x be the set of overconvergent modular forms corresponding to x . Then the elements of \mathcal{M}_x have weight $\pi_1(x) \in \mathcal{W}(L)$ and the U-operator acts on \mathcal{M}_x with eigenvalue $\pi_2(x)^{-1}$. We have that π_1 is, locally on \mathcal{W} and on \mathcal{C} , finite and surjective.

In [Bra12], we have developed a theory of p -adic modular forms of any weight attached to a quaternion algebra over a totally real field F different from \mathbb{Q} , following the approach of [AIS11]. With this paper we want to fill the gap left by the case $F = \mathbb{Q}$, using the same techniques. The main technical results of [Bra12] work also in the case $F = \mathbb{Q}$, but there are some differences, that we explain the following description of the paper.

In Section 1, we describe the basic objects of our work, following [DT94]. We fix a prime $p > 3$ and an indefinite quaternion algebra D over \mathbb{Q} with discriminant δ . We assume $p \nmid \delta$. Let N be an integer with $N > 4$ and coprime with p . We fix a suitable compact open subgroup $K(N) \subseteq D^* \otimes_{\mathbb{Q}} \mathbb{A}^f$, where \mathbb{A}^f is the ring of finite adele of \mathbb{Q} . There is a smooth and proper PEL Shimura curve $\mathcal{M}(N)$ defined over \mathbb{Z}_p that parametrizes abelian surfaces (more precisely ‘false elliptic curves’) with N -level structure. We have also the curves $\mathcal{M}(Np^r)$ for any $r \geq 0$, analogue to the usual compactified modular curve of level $\Gamma_1(Np^r)$. Using the moduli problem solved by $\mathcal{M}(Np^r)$, we define the space of classical modular forms with respect to D , level Np^r , coefficients in K (a finite extension of \mathbb{Q}_p), and weight $k \in \mathbb{Z}$. We empathize the approach of considering a modular form as a ‘rule defined on test objects’ rather than a global section of a line bundle. This is a big difference with [Bra12], and we believe that it makes the theory more transparent (though we need to know that our modular forms actually are sections of some line bundle to construct the eigencurve). We take the rigid analytic viewpoint from the very beginning, so we will consider only modular forms over K .

In Section 2 we define p -adic modular forms of integral weights and we recall the theory of the canonical subgroup, following [Kas99] and [Bra12]. In Section 3 we study the (Cartier dual of the) canonical subgroup in detail and we introduce the map $d \log$, that will be crucial in the sequel. We obtain some of the results of [AIS11], but our approach needs much less calculations and it is completely explicit. In particular, we prove that, over $\mathcal{M}(Np)$, the dual of the canonical subgroup admits a canonical point, and we study the image of this point under the map $d \log$. We do this using a suitable Lubin-Tate group for \mathbb{Z}_p instead of the formal multiplicative group. This idea comes from [Bra12], where it was needed to take into account the action of a finite extension of \mathbb{Z}_p on various objects. In our situation, this is not necessary, but it is in any case very useful, as we explain comparing our approach with the one of [AIS11]. In Section 4, we explain the technical results about the map $d \log$ that we will need.

In Section 5, we recall the basic property of the weight space \mathcal{W} . Let K be a finite extension of \mathbb{Q}_p and let $\chi: \mathbb{Z}_p^* \rightarrow K^*$ be a continuous character. If K contains all the p^r -th roots of unity, where r depends on χ , we define the space of p -adic modular forms with coefficients in K and weight χ , of any level. As mentioned above, our definition is quite simple and it is based on test objects. This makes the definition and the study of the diamond operators very clear. If $\chi(t) = t^k$, then, as usual, we obtain the space of modular forms of integral weight k . We then show that our modular forms are sections of a line bundle on a certain tubular neighborhood

of the ordinary locus of the (rigid analytic) Shimura curve. We finally show that our modular forms can be put in analytic families over the weight space. This paves the way to the construction of the eigencurve via Buzzard’s machinery.

In Section 6 we finally construct the eigencurve. It remains to define the Hecke operators U and T_l , where l is a prime not dividing δN . Using test objects this is not difficult, but we need the approach via line bundles to prove that they are continuous and that U is completely continuous on the space of overconvergent modular forms. All the assumptions of [Buz07] are verified, so we obtain Theorem 6.5.

Acknowledgments. I would like to thank Fabrizio Andreatta and Marco Seveso for their encouragements in writing this work. This paper was carried out when the author was a guest researcher at the Max Planck Institute for Mathematics. I would like to thank that institution for the hospitality and the excellent work conditions provided.

1. SHIMURA CURVES AND CLASSICAL MODULAR FORMS

Let $p > 3$ be a rational prime, fixed from now on. Let $v(\cdot)$ be the valuation of \mathbb{Q}_p normalized in such a way that $v(p) = 1$. We will write $|\cdot|$ for the absolute value of \mathbb{Q}_p such that $|x| = p^{-v(x)}$ for all $x \in \mathbb{Q}_p$. We extend both $v(\cdot)$ and $|\cdot|$ to the whole \mathbb{C}_p , that is the completion of a fixed algebraic closure of \mathbb{Q}_p . We will write $[\cdot]: \mathbb{F}_p^* \rightarrow \mathbb{Z}_p$ for the Teichmüller character, and we set $[0] := 0$. All the schemes we are going to consider will be defined over \mathbb{Z}_p . When there is no possibility of confusion, we will denote base change with a subscript.

In this section we recall the theory of classical modular forms attached to a quaternion algebra, see [DT94] and [Kas99] for details. Let D be a quaternion algebra over \mathbb{Q} that is indefinite and with discriminant δ that satisfies $p \nmid \delta$. We assume that D is a division algebra and we fix \mathcal{O}_D , a maximal order of D . We fix identifications $D \otimes_{\mathbb{Q}} \mathbb{R} = M_2(\mathbb{R})$ and $\mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Z}_l = M_2(\mathbb{Z}_l)$ for all primes $l \nmid \delta$. Let $x \mapsto x'$ be the canonical involution of D . We choose $t \in \mathcal{O}_D$ such that $t^2 = -\delta$ and we define another involution on D by $x \mapsto x^* := t^{-1}x't$. We write \mathbb{A}^f for the ring of *finite* adèle of \mathbb{Q} . Let G/\mathbb{Q} be the reductive algebraic group such that its R -points, for any \mathbb{Q} -algebra R , are $G(R) = (D \otimes_{\mathbb{Q}} R)^*$. We have that $G(\mathbb{A}^f)$ is the restricted tensor product of the D_l^* ’s, where l is a prime and $D_l := D \otimes_{\mathbb{Q}} \mathbb{Q}_l$. Let \mathcal{O}_{D_l} be $\mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Z}_l$ and let K be a compact open subgroup of $\prod_l \mathcal{O}_{D_l}^*$. We will assume that K is ‘small enough’, in particular the following conditions will suffice

- $(N_K, \delta) = 1$, where N_K (the *level* of K) is the smallest positive integer N such that K contains all the elements of $\prod_l \mathcal{O}_{D_l}$ which are congruent to 1 modulo N ;
- $K \subseteq V_1(N)$ for some integer $N \geq 4$ such that $(N, \delta) = 1$, where $V_1(N)$ is the subgroup of $\prod_l \mathcal{O}_{D_l}^*$ given by elements which are congruent to $\begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix}$ modulo N ;
- $\det(K) = \widehat{\mathbb{Z}}^*$, where $\det: G(\mathbb{A}^f) \rightarrow \mathbb{A}^f$ is the reduced norm.

In particular we can take $K = V_1(N)$ with $N > 4$.

Let T be any scheme over \mathbb{Z}_p . By a *false elliptic curve* over T we mean an abelian surface $\mathcal{A} \rightarrow T$ together with an embedding $i: \mathcal{O}_D \hookrightarrow \text{End}_T(\mathcal{A})$. If $(N, \delta) = 1$, by a *N -level structure* on a false elliptic curve (\mathcal{A}, i) over T we mean an isomorphism

of left \mathcal{O}_D -modules $\alpha: \mathcal{A}[N] \xrightarrow{\sim} (\mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Z}/N\mathbb{Z})_T$. If K is as above, a K -level structure on (\mathcal{A}, i) is a N_K -level structure defined up to right K -equivalence.

We now fix $K(N) \subseteq G(\mathbb{A}^f)$ of level N that satisfies the above assumptions and we assume that $(p, N) = 1$. It can be proved that, on any false elliptic curve (\mathcal{A}, i) over T , there is a unique principal polarization whose corresponding Rosati involution on $\text{End}(\mathcal{A}_x)$ (where x is any geometric point of T) restricts to $*$ on \mathcal{O}_D . In particular, any false elliptic curve is canonically principally polarized. The functor \mathbb{Z}_p -schemes $\rightarrow \mathbf{set}$ that sends T to the set of isomorphism classes of false elliptic curves with K -level structure is representable by a geometrically connected scheme $\mathcal{M}(N)$ defined over \mathbb{Z}_p . We have that $\mathcal{M}(N) \rightarrow \text{Spec}(\mathbb{Z}_p)$ is smooth, proper, and of relative dimension 1. The universal object will be denoted $\mathcal{A}(N) \rightarrow \mathcal{M}(N)$.

Let \mathcal{C} be a pseudo-abelian category and let X be an object of \mathcal{C} with an action of $\mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Z}_p$. Choosing a non-trivial idempotent $e \in \mathcal{O}_D \otimes_{\mathbb{Z}} \mathbb{Z}_p \cong M_2(\mathcal{O}_{\mathcal{P}})$ such that $e^* = e$, we obtain a decomposition

$$X = X^1 \oplus X^2,$$

where each X^i has an action of \mathbb{Z}_p . Note that X^1 and X^2 are isomorphic. This notation will be used throughout the paper.

Let $n \geq 0$ be an integer and let $K(N, p^n)$ be the subset of $K(N)$ given by the elements whose component at p is upper triangular modulo p^n . There is a regular scheme $\mathcal{M}(N, p^n)$ that represents the functor \mathbb{Z}_p -schemes $\rightarrow \mathbf{set}$ that sends T to the set of isomorphism classes of quadruples $(\mathcal{A}, i, \alpha, \mathcal{C})$ where (\mathcal{A}, i) is a false elliptic curve over T , α is a K -level structure on (\mathcal{A}, i) and \mathcal{C} is a finite and flat subgroup of order p^n of $\mathcal{A}[p^n]^1$. We have that $\mathcal{M}(N, p^n) \rightarrow \text{Spec}(\mathbb{Z}_p)$ is proper and of relative dimension 1, generically smooth but not smooth. Indeed, the special fiber is a normal crossing divisor with two components, each one isomorphic to $\mathcal{M}(N)_{\mathbb{F}_p}$. The two components intersect at the supersingular points (see below). The universal object will be denoted $\mathcal{A}(N, p^n) \rightarrow \mathcal{M}(N, p^n)$.

Remark 1.1. Any \mathcal{C} as in the above moduli problem gives \mathcal{D} , a finite and flat subgroup of order p^n of $\mathcal{A}[p^n]^2$, and we have that $\mathcal{C} \oplus \mathcal{D}$ is a finite and flat subgroup of $\mathcal{A}[p^n]$ of order p^{2n} , stable under the action of \mathcal{O}_D . All such subgroups of $\mathcal{A}[p^n]$ arise in this way. The canonical principal polarization of \mathcal{A} induces isomorphism $\mathcal{A}[p^n]^1 \cong (\mathcal{A}[p^n]^2)^\vee$ (Cartier dual). This isomorphism takes $\mathcal{C} \hookrightarrow \mathcal{A}[p^n]^1$ to $(\mathcal{A}[p^n]^2/\mathcal{D})^\vee \hookrightarrow \mathcal{D}^\vee$.

Let $K(Np^n)$ be the subset of $K(N, p^n)$ given by the elements whose component at p is congruent to $\begin{pmatrix} * & * \\ 0 & 1 \end{pmatrix}$ modulo p^n . There is a regular scheme $\mathcal{M}(Np^n)$ that represents the functor \mathbb{Z}_p -schemes $\rightarrow \mathbf{set}$ that sends T to the set of isomorphism classes of quadruples $(\mathcal{A}, i, \alpha, P)$ where (\mathcal{A}, i, α) is as above and P is a point of exact order p^n of $\mathcal{A}[p^n]^1$. We have that $\mathcal{M}(Np^n) \rightarrow \text{Spec}(\mathbb{Z}_p)$ is proper, generically smooth, and of relative dimension 1. The universal object will be denoted $\mathcal{A}(Np^n) \rightarrow \mathcal{M}(Np^n)$.

The curves $\mathcal{M}(N)$, $\mathcal{M}(N, p^n)$, and $\mathcal{M}(Np^n)$ are the Shimura curves we are interested in. There are several morphisms between them, given by obvious natural transformations of functors. Let K be any one of $K(N)$, $K(N, p^n)$ or $K(Np^n)$ and let \mathcal{M} be the corresponding curve. If $e: \mathcal{M} \rightarrow \mathcal{A}$ is the zero section

of the universal object, we define

$$\underline{\omega} := \underline{\omega}_K := \left(e^* \Omega_{\mathcal{A}/\mathcal{M}}^1 \right)^1,$$

that is a locally free sheaf of rank 1 on \mathcal{M} . Given $\text{Spec}(R) \rightarrow \mathcal{M}$, where R is a \mathbb{Z}_p -algebra, the pullback of $\underline{\omega}$ to $\text{Spec}(R)$ will be denoted $\underline{\omega}_R$ or $\underline{\omega}_{\mathcal{A}/R}$, where \mathcal{A} is the pullback of the universal object to $\text{Spec}(R)$.

Let V be a finite extension of \mathbb{Z}_p , with field of fraction K (we will make several assumptions about V during the paper). From now on we will usually work over V or over K . To simplify the notation, we will omit the subscripts V and K . We will also be interested in formal schemes and we will use the following convention: algebraic objects defined over V will be denoted using Italic letter, like $\mathcal{M}(N)$. The completion along the subscheme defined by $p = 0$ will be denoted using the corresponding Gothic letter, like $\mathfrak{M}(N)$.

Definition 1.2. Let $r \geq 0$ be an integer. A (classical) test object of level $K(Np^r)$ is a couple $((\mathcal{A}/R, i, \alpha, P), \omega)$, where:

- R is a V -algebra;
- $(\mathcal{A}/R, i, \alpha, P)$ is an object of the moduli problem of level $K(Np^r)$ (we will usually omit P from the notation if $r = 0$), with \mathcal{A} defined over R and such that $\underline{\omega}_R$ is free;
- ω is a basis of $\underline{\omega}_R$.

Definition 1.3. Let k be an integer. A (classical) *modular form with respect to D , level $K(Np^r)$ and weight k , with coefficients in K* , is a rule that assigns to every test object $T = ((\mathcal{A}/R, i, \alpha, P), \omega)$, where T is of level $K(Np^r)$, an element $f(T) \in R_K$ such that

- $f(T)$ depends only on the isomorphism class of T ;
- if $\varphi: R \rightarrow R'$ is a morphism of V -algebras and we denote with T' the base change of T to R' , we have $f(T') = \varphi(f(T))$;
- for any $\lambda \in R^*$, we have

$$f(((\mathcal{A}/R, i, \alpha, P), \lambda\omega)) = \lambda^{-k} f(T).$$

We write $S^D(K, K(Np^r), k)$ for the space of such modular forms.

Proposition 1.4. *There is an invertible sheaf $\Omega^k(Np^r)$ on $\mathfrak{M}(Np^r)^{\text{rig}}$ such that*

$$S^D(K, K(Np^r), k) = H^0(\mathfrak{M}(Np^r)^{\text{rig}}, \Omega^k(Np^r)).$$

Proof. We take for $\Omega^k(Np^r)$ the rigidification of formal completion of $\underline{\omega}^{\otimes k}$. \square

Remark 1.5. The above proposition is trivial, and we can state a much stronger result (namely that we have an invertible sheaf over $\mathcal{M}(Np^r)$ that is a model of $\Omega^k(Np^r)$), but later on we will consider p -adic modular forms of any weight, and the above result has a direct analogue, while the stronger version has not.

Definition 1.3 is given only for modular forms over K , since this is the case we will be mainly interested in, but using Proposition 1.4, we have the notion of classical modular form defined over any \mathbb{Z}_p -algebra, as usual.

2. p -ADIC MODULAR FORMS OF INTEGRAL WEIGHT

We now analyze more closely what happens at level $K(N)$. There is a Kodaira-Spencer isomorphism $\underline{\omega}^{\otimes 2} \cong \Omega_{\mathcal{M}(N)/\mathbb{Z}_p}^1$ and we have $\deg(\underline{\omega}) = g - 1$, where g is the genus of any geometric fiber of $\mathcal{M}(N)$. Let (\mathcal{A}, i, α) be an object of the moduli problem, with \mathcal{A} defined over a V -algebra R . We have that $\mathcal{A}[p^\infty]^1$ is a p -divisible group of dimension 1 and height 2. Let \mathfrak{A} be the p -adic completion of \mathcal{A} and let $\widehat{\mathfrak{A}}$ be the completion of \mathfrak{A} along the zero section. Then $\widehat{\mathfrak{A}}^1$ is a formal group of dimension 1. If $\widehat{\mathfrak{A}}^1$ is a Barsotti-Tate group, its height is either 1 or 2. We say that (\mathcal{A}, i, α) , or simply \mathcal{A} , is *ordinary* if $\widehat{\mathfrak{A}}^1$ has height 1. Otherwise we say that \mathcal{A} is *supersingular*. We have that \mathcal{A} is ordinary (supersingular) if and only if $\mathcal{A}_{\overline{\mathbb{F}}_p}$ is ordinary (supersingular) in the usual sense. If $\underline{\omega}_R$ is a free R -module, we have that $\widehat{\mathfrak{A}}^1$ is a formal group law, and there is a coordinate x on $\widehat{\mathfrak{A}}^1$ such that the multiplication by p has the form

$$[p](x) = px + ax^p + \sum_{j=2}^{\infty} c_j x^{j(p-1)+1},$$

where a, c_j are in R and $c_j \in pR$ unless $j \equiv 1 \pmod{p}$. If we assume that $p = 0$ in R , the various a 's glue together to define H , a modular form of level $K(N)$ and weight $p - 1$, defined over \mathbb{F}_p , that is called the *Hasse invariant*. If $W = \text{Spec}(R)$ is an open affine of $\mathcal{M}(N)$ and we denote with ω the differential dual to the coordinate x defined above, we have $H|_W = a\omega^{\otimes p-1}$. We have that H vanishes precisely at the supersingular points of $\mathcal{M}(N)_{\overline{\mathbb{F}}_p}$ and has simple zeros here (so the number of supersingular points is $(p-1)(g-1)$). Moreover, supersingular points actually exist (hence $g \geq 2$). The modular form H can be lifted, non-canonically, to a modular form E_{p-1} defined over \mathbb{Z}_p . We fix once and for all such a lift, everything will not depend on this choice. Over $\text{Spec}(R)$ we can write $E_{p-1}|_{\text{Spec}(R)} = E\omega^{p-1}$ for some $E \in R$. We have $a \equiv E \pmod{p}$.

Let $0 \leq w < 1$ be a rational number, and let us assume that V contains an element of valuation w , denoted p^w . We define

$$\mathcal{M}(N)(w) := \text{Spec}_{\mathcal{M}(N)}(\text{Sym}(\underline{\omega}^{\otimes p-1})/(E_{p-1} - p^w)).$$

Note that $\mathcal{M}(N)(w)$ has a natural moduli interpretation. It classifies couples $((\mathcal{A}, i, \alpha), Y)$, where $(\mathcal{A}/R, i, \alpha)$ is false elliptic curve of level $K(N)$ and Y is a global section of $\underline{\omega}_{\mathcal{A}/R}^{\otimes 1-p}$ such that $YE_{p-1} = p^w$.

Taking the rigidification of the map $\mathfrak{M}(N)(w) \rightarrow \mathfrak{M}(N)$ we get the immersion $\mathfrak{M}(N)^{\text{rig}}(w) \hookrightarrow \mathfrak{M}(N)^{\text{rig}}$, where $\mathfrak{M}(N)^{\text{rig}}(w)$ is the affinoid subdomain of $\mathfrak{M}(N)^{\text{rig}}$ defined by Coleman in [Col97], relative to E_{p-1} . We call $\mathfrak{M}(N)(0)^{\text{rig}}$ the *ordinary locus*, it is an affinoid subdomain of $\mathfrak{M}(N)^{\text{rig}}$: its complement is a finite union of open discs of radius 1, called the *supersingular discs*.

Let $r \geq 1$ be an integer and assume that $0 \leq w < \frac{1}{p^{r-2}(p+1)}$. Let \mathcal{A} be an object of the moduli problem of $\mathcal{M}(N)(w)$, defined over R . In this case $\mathcal{A}[p^r]^1$ has a canonical finite and flat subgroup \mathcal{C}_r of order p^r . This gives a morphism $\mathfrak{M}(N)(w) \rightarrow \mathfrak{M}(N, p^r)$. Its rigidification is a section, defined over $\mathfrak{M}(N)(w)^{\text{rig}}$, of the morphism $\mathfrak{M}(N, p^r)^{\text{rig}} \rightarrow \mathfrak{M}(N)^{\text{rig}}$. We define $\mathfrak{M}(Np^r)(w)^{\text{rig}}$ as the inverse image of $\mathfrak{M}(N)(w)^{\text{rig}}$ with respect to the map $\mathfrak{M}(N, p^r)^{\text{rig}} \rightarrow \mathfrak{M}(N)^{\text{rig}}$. It is an affinoid subdomain of $\mathfrak{M}(Np^r)^{\text{rig}}$ with a finite and étale map to $\mathfrak{M}(N)(w)^{\text{rig}}$. Taking

the normalization of $\mathfrak{M}(N)(w)$ in $\mathfrak{M}(Np^r)(w)^{\text{rig}}$, that is a finite extension of its generic fiber, we obtain $\mathfrak{M}(Np^r)(w)$, a formal model of $\mathfrak{M}(Np^r)(w)^{\text{rig}}$.

If we work ‘at level $K(Np^r)$ ’, we will tacitly assume that $w < \frac{1}{p^{r-2}(p+1)}$, so we have the canonical subgroup of any object of the moduli problem of $\mathcal{M}(Np^r)(w)$. In particular, all our results will hold if w is small enough.

Notation. Let R be a p -adically complete and flat V -algebra. We will say that R is *normal* if R is integrally closed in R_K . We extend this notion to formal schemes in the obvious way.

Proposition 2.1. *We have that $\mathfrak{M}(Np^r)(w)$ is a flat and normal formal scheme over $\text{Spf}(V)$.*

Proof. In the case $r = 0$ this is [Kas04, Proposition 9.5]. If $r \geq 1$ it is a trivial consequence of the definition. \square

Proposition 2.2. *Let R be a normal, flat, and p -adically complete V -algebra. There is a natural bijection between $\mathfrak{M}(Np^r)(w)(R)$ and the set of isomorphism classes of couples $((\mathcal{A}/R, i, \alpha, P), Y)$, where:*

- $(\mathcal{A}/R, i, \alpha, P)$ is an object of the moduli problem, with \mathcal{A} defined over R , of $\mathcal{M}(Np^r)$ and P generates the canonical subgroup of $\mathcal{A}[p^r]^1$;
- Y is a section of $\underline{\omega}_{\mathcal{A}/R}^{\otimes 1-p}$ that satisfies $YE_{p-1} = p^w$.

Proof. This is [Bra12, Proposition 3.2]. \square

The above proposition motivates the following

Definition 2.3. Let $r \geq 0$ be an integer. A $(p$ -adic) test object of level $K(Np^r)$ and growth condition w is a couple $((\mathcal{A}/R, i, \alpha, P, \omega), Y)$, where:

- R is a p -adically complete, flat, and *normal* V -algebra;
- $((\mathcal{A}/R, i, \alpha, P), \omega)$ is a classical test object of level $K(Np^r)$;
- Y is a global section of $\underline{\omega}_{\mathcal{A}/R}^{\otimes 1-p}$ such that $YE_{p-1} = p^w$.

Definition 2.4. Let k be an integer. A p -adic modular form with respect to D , level $K(Np^r)$ and weight k , with coefficients in K and growth condition w , is a rule that assigns to every test object $T = ((\mathcal{A}/R, i, \alpha, P, \omega), Y)$, where T is of level $K(Np^r)$ and growth condition w , an element $f(T) \in R_K$ such that

- $f(T)$ depends only on the isomorphism class of T ;
- if $\varphi: R \rightarrow R'$ is a morphism of p -adically complete, flat, and normal V -algebras and we denote with T' the base change of T to R' , we have $f(T') = \varphi(f(T))$;
- for any $\lambda \in R^*$, we have

$$f((\mathcal{A}/R, i, \alpha, P, \lambda\omega), Y) = \lambda^{-k} f(T).$$

We write $S^D(K, w, K(Np^r), k)$ for the space of such modular forms.

If $h \leq r$ are integers, the natural injective morphism $S^D(K, w, K(Np^h), k) \rightarrow S^D(K, w, K(Np^r), k)$ allows us to consider $S^D(K, w, K(Np^h), k)$ as a subset of $S^D(K, w, K(Np^r), k)$.

Proposition 2.5. *There is an invertible sheaf $\Omega^k(Np^r)(w)$ on $\mathfrak{M}(Np^r)(w)^{\text{rig}}$ such that*

$$S^D(K, w, K(Np^r), k) = H^0(\mathfrak{M}(Np^r)(w)^{\text{rig}}, \Omega^k(Np^r)(w)).$$

Proof. We take for $\Omega^k(Np^r)(w)$ the restriction to $\mathfrak{M}(Np^r)(w)^{\text{rig}}$ of $\Omega^k(Np^r)$. \square

This proposition shows that the natural morphism $S^D(K, w', K(Np^r), k) \rightarrow S^D(K, w, K(Np^r), k)$, where $w' \geq w$ is a rational number that satisfies the same assumptions as w does, is injective.

Remark 2.6. As in Remark 1.5, we can state a stronger result than in the above proposition, namely, that we have a formal model of $\Omega^k(Np^r)(w)$. But, since we do not have an integral model of $\mathfrak{M}(Np^r)(w)$ that satisfies a reasonable moduli property, we do not have an integral model of $\Omega^k(Np^r)(w)$. Later on, we will not even have the formal model of the sheaves we are interested in.

By definition, elements of $S^D(K, K(Np^r), k)$ correspond to sections of $\Omega^k(Np^r)$ over $\mathfrak{M}(Np^r)^{\text{rig}}$, while elements of $S^D(K, w, K(Np^r), k)$ correspond to sections over $\mathfrak{M}(Np^r)(w)^{\text{rig}}$. Elements of $S^D(K, 0, K(Np^r), k)$ are called *convergent* modular forms, while the elements of $S^D(K, w, K(Np^r), k)$, for $w > 0$, are called *overconvergent* modular forms.

3. MORE ON THE CANONICAL SUBGROUP AND THE MAP $d \log$

There is a primitive p -th root of unity ζ_p in V if and only if V contains ξ , a non-trivial root of $x^p - px$, that is, a $(p-1)$ -th root of $-p$. This is an exercise, but it also follows from Oort-Tate theory: by [OT70, pages 8-10], there is an isomorphism of group schemes $\text{Spec}(V[x]/(x^p - 1)) \cong \text{Spec}(V[x]/(x^p - pw_{p-1}x))$, where $w_{p-1} \in \mathbb{Z}_p^*$ is a universal constant. In particular, ζ_p gives a $(p-1)$ -th root of $w_{p-1}p$. We have that the reduction modulo p of w_{p-1} is congruent to $(p-1)!$, so $-w_{p-1}$ admits a $(p-1)$ -th root in \mathbb{Z}_p and we get the desired $(p-1)$ -th root of $-p$. The point of this discussion is that, to simplify the calculations, we prefer to take Oort-Tate theory as *starting point* rather than using it to obtain technical results. In this way our calculations will be completely trivial. From this point of view, $\mu_{p,V}$ is $\text{Spec}(V[x]/(x^p + px))$. From now on, when we work at level $K(Np)$, we will tacitly assume that V contains a fixed $(p-1)$ -th root of $-p$, denoted $(-p)^{1/(p-1)}$ (in particular, V contains also ζ_p). Later on, we will explain a similar assumption for higher levels, equivalent to the fact that V contain a primitive p^r -root of unity.

At level $K(Np)$, the canonical subgroup is given, locally on $\mathcal{M}(N)(w)$, by $\mathcal{C} = \mathcal{C}_1 = \text{Spec}(R[x]/(x^p + \frac{p}{E}x))$, where $E_{p-1}|_{\text{Spec}(R)} = E\omega^{p-1}$ as in Section 2 (the fact that the canonical subgroup arise naturally in this form is the main reason why it is so convenient to use Oort-Tate theory from the very beginning).

Proposition 3.1. *There is $E_1 \in S^D(K, w, K(Np), 1)$ such that $E_1^{q-1} = E_{q-1}$.*

Proof. Let $\text{Spec}(S)$ be the base change of $\text{Spec}(R)$ to $\mathcal{M}(Np)(w)$. By the moduli property of $\mathcal{M}(Np)(w)$, we have a canonical S -point of \mathcal{C} , that gives (using $(-p)^{1/(p-1)}$) a $(p-1)$ -th root of E in S . These roots glue together to define the required E_1 . \square

Let $r \geq 1$ be an integer. Fix an open affine $\text{Spf}(R)$ of $\mathfrak{M}(N)(w)$ such that ω is free, generated by ω as above. Let $\text{Spf}(S_r)$ be the pullback of $\text{Spf}(R)$ to $\mathfrak{M}(Np^r)(w)$ under $\mathfrak{M}(Np^r)(w) \rightarrow \mathfrak{M}(N)(w)$. Let \mathcal{A} be any object of the moduli problem of $\mathfrak{M}(N)(w)$. We know that $\mathcal{A}[p^r]^1$ admits a canonical subgroup \mathcal{C}_r and that the canonical point of $\mathcal{A}_{S_r}[p^r]^1$ generates \mathcal{C}_r . We want to construct a canonical S_r -point of $(\mathcal{C}_r)^\vee$ that is a generator of $(\mathcal{C}_r)^\vee(S_r)$ as $\mathbb{Z}/p^r\mathbb{Z}$ -module. All our schemes

are normal, so it is enough to define a morphism $\mathcal{C}_{r,S_r,K} \rightarrow \mu_{p^r,S_r,K}$. But $\mathcal{C}_{r,S_r,K}$ is a constant group scheme, with $\mathbb{Z}/p^r\mathbb{Z}$ as associated abstract group, and we have P , a canonical generator of $\mathcal{C}_{r,S_r,K}$. So it is enough to define the image of P , or, that is the same, to choose a point of $\mu_{p^r,S_r,K}$. In particular it is natural to assume that V contains ζ_{p^r} , a fixed primitive p^r -th root of unity, and to take it as the image of P . But recall that we want to use Oort-Tate theory as building block, so for us $\mu_{p,V}$ is $\text{Spec}(V[x]/(x^p + px))$ rather than $\text{Spec}(V[x]/(x^p - 1))$. For this reason, we take the following slightly different approach.

Let \mathcal{LT} be the *basic Lubin-Tate group associated to p* , as in [Lan78, Chapter 8], over V . It is a formal group law on $V[[x]]$ that has an action of \mathbb{Z}_p and the multiplication by p is given by $[p](x) = x^p + px$. If $z \in \mathbb{F}_p$, the action of $[z]$ sends x to $[z]x$. All Lubin-Tate groups associated to p are strictly isomorphic, so $\mathcal{LT} \cong \widehat{\mathbb{G}}_{m,V}$. In particular, if G is a finite and flat group scheme over S_r , giving a morphism $G \rightarrow \mathcal{LT}_{S_r}$ is the same as giving a morphism $G \rightarrow \widehat{\mathbb{G}}_{m,S_r}$. Repeating the above discussion, we see that to define a point of $(\mathcal{C}_r)^\vee(S_r)$ it is enough to fix a point of \mathcal{LT} of order exactly p^r . For this reason we fix a sequence $\{\xi_r\}_{r \geq 1}$ of \mathbb{C}_p -points of \mathcal{LT} such that ξ_r has order exactly p^r and $p\xi_{r+1} = \xi_r$ (recall that \mathcal{LT} is denoted additively). By definition, the points of order p of \mathcal{LT} are the points of $\text{Spec}(V[x]/(x^p + px))$, so it is natural to require that ξ_1 is our fixed $(-p)^{1/(p-1)}$. When we will work at level $K(Np^r)$, we will tacitly assume that $\xi_r \in V$, so we have γ_r , a canonical S_r -point of $(\mathcal{C}_r)^\vee$ of order p^r . For various r 's, these morphisms are compatible, and if we let $r \rightarrow \infty$ (and we take $w = 0$) we have a morphism, defined over \mathbb{C}_p

$$\mathcal{C}_\infty \rightarrow \mathcal{LT}[p^\infty],$$

where \mathcal{C}_∞ is the p -divisible group defined by the \mathcal{C}_r 's.

We can study more closely the case $r = 1$. Recall that the canonical subgroup is $\mathcal{C} = \text{Spec}(R[x]/(x^p + \frac{p}{E}))$, where $E \in R$ satisfies $E_{p-1}|_{\text{Spf}(R)} = E\omega^{\otimes p-1}$. Let $S := S_1$. Giving $\gamma = \gamma_1$, our canonical S -point of \mathcal{C}_S^\vee , is the same as giving the morphism $S[x]/(x^p + px) \rightarrow S[x]/(x^p + \frac{p}{E})$, that amounts to choose a $(p-1)$ -th root of E in S . But by definition we have a $(p-1)$ -th root of $-\frac{p}{E}$ in S , and using our fixed ξ_1 we get $E^{1/(p-1)} \in S$. At the end we see that our morphism $\gamma \in \mathcal{C}^\vee(S)$ corresponds to

$$\begin{aligned} \mathcal{C}_S &= \text{Spec}(S[x]/(x^p + \frac{p}{E})) \rightarrow \mathcal{LT}_S = \text{Spf}(S[[x]]) \\ E^{1/(p-1)}x &\leftarrow x \end{aligned}$$

In this way the explicit description of γ is very simple, and the relation between γ and E_1 is completely transparent.

It follows by the explicit description of the canonical subgroup that the module of invariant differential of \mathcal{C} is generated by $d(x)$, with the unique relation $\frac{p}{E}d(x) = 0$, so we have $\underline{\omega}_{\mathcal{C}} \cong R/\frac{p}{E}Rd(x)$. We can be even more precise: let $h: \mathcal{C} \rightarrow \widehat{\mathcal{A}}[p]^1$ be the natural map. We have fixed above a coordinate x of $\widehat{\mathcal{A}}^1$, and we have denoted with ω the differential dual to x . Then, in $\Omega_{\mathcal{C}/R}^1$, we have $h^*(\omega) = \frac{d(x)}{1-E}$. In particular, under the isomorphism $\underline{\omega}_{\mathcal{C}} \cong R/\frac{p}{E}Rd(x)$, we have $h^*(\omega) = d(x)$. Using Oort-Tate theory, we can give an explicit formula for the comultiplication of \mathcal{C} . It is given by

$$c(x) = x \otimes 1 + 1 \otimes x + \frac{1}{p-1} \sum_{i=1}^{p-1} E \frac{w_{p-1}}{w_i w_{p-1}} x^i \otimes x^{p-i},$$

where $w_i \in \mathbb{Z}_p$ are the universal constants of [OT70]. Moreover, the counit of \mathcal{C} is given by $x \mapsto 0$. Having fixed ζ_p , a primitive p -th root of unity in V , one can also write down a morphism $\mathcal{C}_S \rightarrow \text{Spec}(S[x]/(x^p - 1))$ explicitly (clearly, to obtain the morphism corresponding to γ , we need to take as ζ_p the root of unity corresponding to ξ_1 under the isomorphism $R[x]/(x^p - 1) \cong R[x]/(x^p + p)$). This is the approach taken in [AIS11], and we now show that our point of view makes the calculations much easier. Let $\alpha \in S$ be $(p-1)$ -th root of $-\frac{p}{E}$ given by the moduli problem (so we have $\xi_1 = \alpha E^{1/(p-1)}$). Then the map

$$S[x]/(x^p - 1) \rightarrow S[x]/(x^p + \frac{p}{E}x)$$

$$x \mapsto 1 + \frac{1}{1-p} \sum_{i=1}^{p-1} \frac{(-1)^i x^i}{w_i \alpha^i} \left(\sum_{z \in \mathbb{F}_p^*} [x]^{-1} \zeta_p^z \right)^i$$

gives the required morphism $\mathcal{C}_S \rightarrow \text{Spec}(S[x]/(x^p - 1))$. We see that here the formula is much more complicated than the one for γ , in particular, Proposition 3.2 below, that is easy with our approach, requires non trivial calculations in [AIS11].

We continue to use the above notations. Let $\eta = \text{Spec}(\mathbb{K})$ be a generic geometric point of $\text{Spec}(R)$, we write \mathcal{G} for $\pi_1(\text{Spec}(R_K), \eta_K)$. We denote with \overline{R} the direct limit of all R -algebras $T \subseteq \mathbb{K}$ which are normal and such that T_K is finite and étale over R_K . Let G be a finite and flat group scheme over R , killed by p^n . Taking the pullback of the invariant differential $d(x)$ of \mathcal{LT} , we get the map

$$d \log_{G,T}: G^\vee(T) \rightarrow \underline{\omega}_G \otimes_R T/p^n T,$$

where T is as above. We are interested in the map $d \log_G: G^\vee(\overline{R}) \rightarrow \underline{\omega}_G \otimes_R \overline{R}/p^n \overline{R}$ obtained by direct limit. Taking the inverse limit over n , we have the map

$$d \log_{\mathcal{A}}: \text{T}_p((A[p^\infty]^1)^\vee) \otimes_{\mathbb{Z}_p} \widehat{R} \rightarrow \underline{\omega}_{\mathcal{A}/R} \otimes_R \widehat{R},$$

where $\text{T}_p((A[p^\infty]^1)^\vee) = ((A[p^\infty]^1)^\vee)(\overline{R})$ is the Tate module of $\mathcal{A}[p^\infty]^1$.

Proposition 3.2. *We have*

$$d \log_{\mathcal{A}[p]^1, S}(\gamma) \equiv E_1 \pmod{p^{1-w}}.$$

Proof. Clearly we have $\gamma^*(d(x)) = E^{1/(p-1)} d(x)$. We can write $\underline{\omega}_{\mathcal{C}} = \omega R/\frac{p}{E}R$ and $\underline{\omega}_{\mathcal{A}} = \omega R$. But by definition of $\mathcal{M}(N)(w)$, we have $\frac{p}{E} \in p^{1-w}R$, so the natural map $\underline{\omega}_{\mathcal{A}} \rightarrow \underline{\omega}_{\mathcal{C}}$ is an isomorphism modulo p^{1-w} . By definition we have $E_{1|\text{Spf}(R)} = E^{1/(p-1)} \omega \in H^0(\mathfrak{M}(Np)(w), \underline{\omega})$ so we see that $\gamma^*(d(x)) \equiv E_{1|\text{Spf}(S)} \pmod{p^{1-w}}$. \square

From now on we will omit (\overline{R}) in the notation, it should be clear from the context whether we are talking about the group scheme or about the group of points. We also write \overline{R}_z for $\overline{R}/p^z \overline{R}$ (and similarly for other objects).

4. THE HODGE-TATE SEQUENCE

We recall in this section the main technical results of [Bra12] (see also [AIS11]). The proof are exactly the same, so we will omit them. Let $\text{Spf}(R)$ be an open affine of $\mathfrak{M}(N)(w)$, and let \mathcal{A} be the corresponding false elliptic curve defined over R . We assume that $w < \frac{p}{p+1}$.

Definition 4.1. The *Hodge-Tate sequence* of \mathcal{A} is the following sequence of \widehat{R} -modules with semilinear action of \mathcal{G} :

$$0 \rightarrow \underline{\omega}_{\mathcal{A}^\vee/R}^* \otimes_R \widehat{R}(1) \xrightarrow{a_{\mathcal{A}}} \mathrm{T}_p((\mathcal{A}[\varpi^\infty]^1)^\vee) \otimes_{\mathbb{Z}_p} \widehat{R} \xrightarrow{d \log_{\mathcal{A}}} \underline{\omega}_{\mathcal{A}/R} \otimes_R \widehat{R} \rightarrow 0.$$

where $*$ means ‘dual module’, $a_{\mathcal{A}}$ is defined using the $d \log$ -maps of $(\mathcal{A}[p^\infty]^1)^\vee$, and (1) is the Tate-twist.

Theorem 4.2. *The Hodge-Tate sequence is a complex and the map $a_{\mathcal{A}}$ is injective. Moreover, the homology of the sequence is killed by p^v , where $v := \frac{w}{p-1}$. The Hodge-Tate sequence is exact if and only if \mathcal{A} is ordinary. Furthermore, $\ker(d \log_{\mathcal{A}})$ and $\mathrm{Im}(d \log_{\mathcal{A}})$ are free \widehat{R} -modules of rank 1. The latter is generated by $d \log_{\mathcal{A}}((\gamma)_r)$. Finally, taking $\mathrm{Gal}(\overline{R}_K/S_K)$ -invariants, everything remains true also over S (i.e. at level $K(Np)$), but we cannot descend to R (i.e. at level $K(N)$).*

Corollary 4.3. *We have that $\underline{\omega}_{K(Np^r)}^{\mathrm{rig}}$ is globally free if and only if $r \geq 1$. In this case, we have $\underline{\omega}_{K(Np^r)}^{\mathrm{rig}} = \mathcal{O}_{\mathfrak{M}(Np^r)(w)^{\mathrm{rig}}} E_1$.*

This corollary is the reason why the theory is quite different if $r \geq 1$ or not.

Let $\mathcal{D}_r \subseteq (\mathcal{A}[p^r]^1)^\vee$ be the orthogonal of \mathcal{C}_r . Let $w < \frac{1}{p^{r-2}(p+1)}$ as usual (we need to assume $w \leq 1/p$ if $r = 1$). Then the kernel of the map $d \log_{\mathcal{A}[p^r]^1, \overline{R}}: (\mathcal{A}[p^r]^1)^\vee \rightarrow \underline{\omega}_{\mathcal{A}/R} \otimes_R \overline{R}_r$ is \mathcal{D}_r .

Proposition 4.4. *We have a commutative diagram of \mathcal{G} -modules, with exact rows and vertical isomorphisms,*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \ker(d \log_{\mathcal{A}})_{r-v} & \longrightarrow & \mathrm{T}_p((\mathcal{A}[p^\infty]^1)^\vee) \otimes \overline{R}_{r-v} & \longrightarrow & \mathrm{Im}(d \log_{\mathcal{A}})_{r-v} \longrightarrow 0 \\ & & \downarrow \wr & & \parallel & & \downarrow \wr \\ 0 & \longrightarrow & \mathcal{D}_r \otimes \overline{R}_{r-v} & \longrightarrow & (\mathcal{A}[p^r]^1)^\vee \otimes \overline{R}_{r-v} & \longrightarrow & \mathcal{C}_r^\vee \otimes \overline{R}_{r-v} \longrightarrow 0 \end{array}$$

5. MODULAR FORMS OF NON-INTEGRAL WEIGHT

There is a rigid analytic (group) variety \mathcal{W} over \mathbb{Q}_p such that, for any affinoid algebra A , we have $\mathcal{W}(A) = \mathrm{Hom}_{\mathrm{cont}}(\mathbb{Z}_p^*, A^*)$ (functorially). We embed \mathbb{Z} in \mathcal{W} sending $k \in \mathbb{Z}$ to the morphism $t \mapsto t^k$. We have an isomorphism of topological groups $\mathbb{Z}_p \cong \mu_{p-1} \times (1+p)\mathbb{Z}_p$, so, if \mathcal{W}° is the connected component of \mathcal{W} containing the trivial character, we have $\mathcal{W} = \coprod_{\mu_{p-1}^\vee} \mathcal{W}^\circ$, where $\mu_{p-1}^\vee := \mathrm{Hom}_{\mathrm{groups}}(\mu_{p-1}, \mathbb{Q}_p^*)$. Moreover, the map $\chi \mapsto \chi(1+p) - 1$ gives an isomorphism $\mathcal{W}^\circ \cong \mathcal{B}(1^-)$, where $\mathcal{B}(a^-)$ is the open disk of radius a and center the origin. It is well known that any continuous $\chi: \mathbb{Z}_p^* \rightarrow A^*$ is locally analytic. We say that χ is *r-admissible* if the map $\mathbb{Z}_p \rightarrow A^*$ given by $x \mapsto \chi((1+p^{r+1}x))$ is analytic. This is equivalent to

$$|\chi(1+p) - 1| < p^{-\frac{1}{p^{r-1}(p+1)}}.$$

Let $\chi: \mathbb{Z}_p^* \rightarrow K^*$ be r -admissible. It follows that there is $s \in K$ such that $\chi(t) = [t]^i \langle t \rangle^s := [t]^i \exp(s \log(\langle t \rangle))$ for all t with $v(\langle t \rangle - 1) \geq r$, where

- $i \in \mathbb{Z}/(p-1)\mathbb{Z}$;
- $[t]$ means the Teichmüller character applied to the reduction modulo p of t ;
- $\langle t \rangle t := t/[t]$ and $s \in K$ is such that $|s| < p^{-\frac{1}{p-1}+r}$.

Let \mathcal{W}_{r-}° be the subset of \mathcal{W}° corresponding to $\mathcal{B}_{r-} := \mathcal{B}(p^{-\frac{1}{p^{r-1}(p+1)}})$ via $\mathcal{W}^\circ \cong \mathcal{B}(1)$. We write \mathcal{W}_{r-} for $\prod_{\mu_{p-1}} \mathcal{W}_{r-}^\circ$. It follows that χ is r -admissible if and only if $\chi \in \mathcal{W}_{r-}(K)$. Taking *closed* discs \mathcal{C}_r such that $\mathcal{B}_{(r-1)-} \subseteq \mathcal{C}_r \subseteq \mathcal{B}_{r-}$ (where $\mathcal{B}_{0-} := \{0\}$), we obtain an admissible open $\mathcal{W}_r \subseteq \mathcal{W}$. Furthermore $\{\mathcal{W}_r\}_{r \geq 1}$ is an admissible covering of \mathcal{W} and any $\chi \in \mathcal{W}_r(K)$ is r -admissible (but of course the converse is not true). We fix one such $\{\mathcal{W}_r\}_{r \geq 1}$.

Let $r \geq 1$ be an integer and let $\chi: \mathbb{Z}_p^* \rightarrow K^*$ be an r -accessible character and write $\chi(t) = [t]^i \langle t \rangle^s := [t]^i \exp(s \log(\langle t \rangle))$, for t sufficiently close to 1. From now on, when we work at level $K(Np^r)$ we will always assume that

$$w < \min \left((p-1) \left(v(s) - r - \frac{1}{p-1} \right), \frac{1}{p^{r-2}(p+1)} \right).$$

We define the sheaf (of groups) on $\mathfrak{M}(Np^r)(w)$

$$\mathcal{S}_{r,w} := \mathbb{Z}_p^*(1 + p^{r-v} \mathcal{O}_{\mathfrak{M}(Np^r)(w)}).$$

The reason to consider $\mathcal{S}_{r,w}$ is the following: if $x \in \mathcal{S}_{r,w}(\mathrm{Spf}(S_r))$ (where S_r is as in Section 3) we can write $x = ub$, where $u \in \mathbb{Z}_p^*$ and $b \in 1 + p^{r-v} S_r$. It follows from our assumption on w that $x^\chi := \chi(u)b^s$ makes sense, and it is another section of $\mathcal{S}_{r,w}$. Thus, $\mathcal{S}_{r,w}$ is a suitable subsheaf of $\mathcal{O}_{\mathfrak{M}(Np^r)(w)}^*$ where ‘raising to the χ -th power’ makes sense. To define modular forms of weight χ , we need to find an analogue of the $\mathcal{O}_{\mathfrak{M}(Np^r)(w)}^*$ -torsor given by the generators of $\underline{\omega}$, but relative to $\mathcal{S}_{r,w}$, so we can raise to the χ -th power. We proceed in two steps: first of all we find a suitable trivial subsheaf of $\underline{\omega}$ (note that, since we working with the formal models, $\underline{\omega}$ is not trivial even if $r \geq 1$), then, working (via the trivialization) with $\mathcal{O}_{\mathfrak{M}(Np^r)(w)}$, we define the required torsor.

Let $\mathcal{F}_{r,w}$ be the sheaf on $\mathfrak{M}(Np^r)(w)$ defined as follows. Let $\mathrm{Spf}(R) \subseteq \mathfrak{M}(N)(w)$ be an open affine, and write $\mathrm{Spf}(S_r)$ for its pullback to $\mathfrak{M}(Np^r)(w)$. Let $\mathcal{G}_r := \mathrm{Gal}(\overline{R}_K, S_{r,K})$. Let \mathcal{A} be the false elliptic curve corresponding to $\mathrm{Spf}(R)$, we define

$$\mathcal{F}_{r,w}(\mathrm{Spf}(S_r)) := \mathrm{Im}(\mathrm{d} \log_{\mathcal{A}})^{\mathcal{G}_r} \subseteq \underline{\omega}_{\mathcal{A}_{S_r}/S_r}.$$

This the desired trivial subsheaf of $\underline{\omega}$: by the results of Section 4 and Proposition 3.2, we have that $\mathcal{F}_{r,w} = \mathcal{O}_{\mathfrak{M}(Np^r)(w)} E_1$. We can now define the required torsor. Writing \mathfrak{C}_r^\vee for the (constant) sheaf defined by the dual of the canonical subgroup, we have, by Proposition 4.4,

$$(1) \quad \mathcal{F}_{r,w}/p^{r-v} \mathcal{F}_{r,w} \cong \mathfrak{C}_r^\vee \otimes \mathcal{O}_{\mathfrak{M}(Np^r)(w)}/p^{r-v} \mathcal{O}_{\mathfrak{M}(Np^r)(w)}.$$

Having this isomorphism, we define the sheaf (of sets) $\mathcal{F}'_{r,w}$ on $\mathfrak{M}(Np^r)(w)$ as the inverse image of the constant sheaf given by $(\mathfrak{C}_r^\vee)^*$, the subset of \mathfrak{C}_r^\vee of points of order exactly p^r .

Proposition 5.1. *We have that $\mathcal{F}'_{r,w}$ is the trivial $\mathcal{S}_{r,w}$ -torsor generated by E_1 .*

Proof. This follows immediately from Section 4. \square

Definition 5.2. A test object of level $K(Np^r)$ (with $r \geq 1$) and growth condition w is a triple $((\mathcal{A}/S_r, i, \alpha, P), Y, \eta)$, where:

- S_r is a p -adically complete, flat, and normal V -algebra;
- $(\mathcal{A}/S_r, i, \alpha, P)$ is an object of the moduli problem of level $K(Np^r)$;
- Y is a global section of $\underline{\omega}_{\mathcal{A}/S_r}^{\otimes 1-p}$ such that $Y E_{p-1} = p^w$.

- η is a global section of the pullback of $\mathcal{F}'_{r,w}$ to $\mathrm{Spf}(S_r)$.

Remark 5.3. The above definition is different, and not equivalent to Definition 2.3, but the definition of modular form given below, in the case $\chi(t) = t^k$, will be the same as above. Since we are interested in modular forms rather than in test objects, this is not a problem.

Definition 5.4. Let $r \geq 1$ and let χ be an r -accessible character. A p -adic modular form with respect to D , level $K(Np^r)$ and weight χ , with coefficients in K and growth condition w , is a rule that assigns to every test object $T = ((\mathcal{A}/S_r, i, \alpha, P), Y, \eta)$, where T is of level $K(Np^r)$ and growth condition w , an element $f(T) \in S_{r,K}$ such that

- $f(T)$ depends only on the isomorphism class of T ;
- if $\varphi: S_r \rightarrow S'_r$ is a morphism of p -adically complete, flat, and normal V -algebras and we denote with T' the base change of T to S'_r , we have $f(T') = \varphi(f(T))$;
- for any $x \in S_{r,w}(\mathrm{Spf}(S_r)) = \mathbb{Z}_p^*(1 + p^{r-v}S_r)$, we have

$$f(((\mathcal{A}/S_r, i, \alpha, P), Y, x\eta)) = x^{\chi^{-1}} f(T).$$

We write $S^D(K, w, K(Np^r), \chi)$ for the space of such modular forms.

Proposition 5.5. Let $r \geq 1$. On $\mathfrak{M}(Np^r)(w)^{\mathrm{rig}}$ there is an invertible sheaf $\Omega^{\chi}(Np^r)(w)$ such that

$$S^D(K, w, K(Np^r), \chi) = H^0(\mathfrak{M}(Np^r)(w)^{\mathrm{rig}}, \Omega^{\chi}(Np^r)(w)).$$

If $\chi(t) = t^k$ for some integer k , we have a natural isomorphism $\Omega^{\chi}(Np^r)(w) \cong \Omega^k(Np^r)(w) \cong \underline{\omega}^{\otimes k}$, so $S^D(K, w, K(Np^r), \chi) = S^D(K, w, K(Np^r), k)$.

Proof. Let $\mathcal{O}_{\mathfrak{M}(Np^r)(w)}^{(\chi)}$ be $\mathcal{O}_{\mathfrak{M}(Np^r)(w)}$ with the action of $S_{r,w}$ twisted by χ . We take for $\Omega^{\chi}(Np^r)(w)$ the rigidification of

$$\mathcal{H}om_{S_{r,w}}(\mathcal{F}'_{r,w}, \mathcal{O}_{\mathfrak{M}(Np^r)(w)}^{(\chi^{-1})}),$$

where $\mathcal{H}om_{S_{r,w}}(\cdot, \cdot)$ means homomorphisms of sheaves with an action of $S_{r,w}$. It is an invertible sheaf by Proposition 5.1. The second part of the proposition follows by Corollary 4.3. \square

Remark 5.6. As in Remark 2.6, we have a formal model of $\Omega^{\chi}(Np^r)(w)$. We will see that this will be false in the case $r = 0$. In any case, since E_1 is a generator of $\underline{\omega}$ over K , we need to work after having inverted p to have the isomorphism $\Omega^{\chi}(Np^r)(w) \cong \Omega^k(Np^r)(w)$. This is the reason why we consider modular forms with coefficients in K rather than in V .

Let $r \geq h \geq 1$ be integers.. We write $\vartheta_{r,h}: \mathfrak{M}(Np^r)(w) \rightarrow \mathfrak{M}(Np^h)(w)$ for the natural morphism. Its rigidification is finite, étale, and Galois with Galois group $G_{r,h} \subseteq (\mathbb{Z}/p^r\mathbb{Z})^*$ given by the image of $1 + p^h\mathbb{Z}$ under the projection $\mathbb{Z} \rightarrow \mathbb{Z}/p^r\mathbb{Z}$. We let moreover $\vartheta_r: \mathfrak{M}(Np^r)(w) \rightarrow \mathfrak{M}(N)(w)$ be the natural morphism. We have that $\vartheta_r^{\mathrm{rig}}$ is finite and étale with $G_r = (\mathbb{Z}/p^r\mathbb{Z})^*$ as Galois group. Let χ be an r -accessible character and let $f \in S^D(K, w, K(Np^r), \chi)$. For any $a \in G_r$, we define another modular form $f_{|_{\langle a \rangle}}$ via the formula

$$f_{|_{\langle a \rangle}}(\mathcal{A}/S_r, i, \alpha, P, Y, \eta) = f(\mathcal{A}/S_r, i, \alpha, aP, Y, a^{-1}\eta),$$

where the action of G_r on the global sections of $\mathcal{F}'_{r,w}$ is defined by $aE_1 := a^*E_1$, viewing any $a \in G_r$ as an automorphism of $\mathfrak{M}(Np^r)(w)$. Note that this action is compatible, via the isomorphism (1), with the natural action of G_r on $(\mathcal{C}_r^\vee)^*$. The operator $f \mapsto f_{|\langle a \rangle}$ is called a *diamond operator*. It is easy to show that it comes from an action of G_r on $\Omega^\chi(Np^r)(w)$.

Proposition 5.7. *Let $f \in S^D(K, w, K(Np^r), \chi)$. Then f descends to level $K(Np^h)$ if and only if $f_{|\langle a \rangle} = f$ for all $a \in G_{r,h}$.*

Proof. This follows immediately from the definition of modular form using test objects. \square

Example 5.8. We have $E_{1|\langle a \rangle} = [a^{-1}]E_1$ for any $a \in \mathbb{F}_p^*$, so, as we already know, E_1 does not descend to level $K(N)$. On the other hand $E_{p-1|\langle a \rangle} = [a^{-1}]^{p-1}E_1^{\otimes p-1} = E_{p-1}$ as required.

We now study the relations between the sheaf $\Omega^\chi(Np^r)(w)$ for various r . Let $r \geq h \geq 1$ be integers and let χ be an h -accessible character. It follows that χ is also r -accessible.

Proposition 5.9. *We have a natural isomorphism*

$$\sigma_{r,h}: (\vartheta_{r,h,*}\Omega^\chi(Np^r)(w))^{G_{r,h}} \cong \Omega^\chi(Np^h)(w).$$

Furthermore $\sigma_{r,r} = \text{id}$ and, if $n \geq r$ is an integer, we have $\sigma_{n,h} = \sigma_{r,h} \circ \sigma_{n,r}$.

Proof. This is [Bra12, Proposition 6.34]. \square

If χ is r -accessible, we have defined modular forms only of level $K(Np^r)$. We are now ready to remove this restriction, and define also modular forms of level $K(Np^h)$.

Definition 5.10. Let $\chi: \mathbb{Z}_p \rightarrow K^*$ be any continuous character and let $h \geq 0$ be an integer. Choose $r \geq h$ such that χ is r -accessible. A *p -adic modular form with respect to D , level $K(Np^h)$ and weight χ , with coefficients in K and growth condition w* , is a rule that assigns to every test object $T = ((\mathcal{A}/S_r, i, \alpha, P), Y, \eta)$, where T is of level $K(Np^r)$ and growth condition w , an element $f(T) \in S_{r,K}$ such that the conditions of Definition 5.4 are satisfied and moreover

$$f_{|\langle a \rangle} = f$$

for all $a \in G_{r,h}$. We write $S^D(K, w, K(Np^h), \chi)$ for the space of such modular forms.

Remark 5.11. It follows from Proposition 5.9, that the above definition does not depend on the choice of r . Furthermore, it is obvious that Propositions 5.7 and 5.9 stay true in general.

Proposition 5.12. *Let χ be any continuous character and let h be an integer. There is an invertible sheaf $\Omega^\chi(Np^h)(w)$ on $\mathfrak{M}(Np^h)(w)^{\text{rig}}$ such that*

$$S^D(K, w, K(Np^h), \chi) = H^0(\mathfrak{M}(Np^h)(w)^{\text{rig}}, \Omega^\chi(Np^h)(w)).$$

If $\chi(t) = t^k$ for some integer k , we have a natural isomorphism $\Omega^\chi(Np^h)(w) \cong \Omega^k(Np^h)(w) \cong \underline{\omega}^{\otimes k}$, so $S^D(K, w, K(Np^h), \chi) = S^D(K, w, K(Np^h), k)$.

Proof. Choose any $r \geq h$ such that χ is r -accessible. We define

$$\Omega^X(Np^h)(w) := (\vartheta_{r,h,*}\Omega^X(Np^r)(w))^{G_{r,h}}.$$

Since $\mathfrak{M}(Np^r)(w)^{\text{rig}} \rightarrow \mathfrak{M}(Np^h)(w)^{\text{rig}}$ is finite étale with $G_{r,h}$ as Galois group, we have that $\Omega^X(Np^h)(w)$ is an invertible sheaf (see also [Bra12, Propositions 6.24 and 6.27]). \square

Remark 5.13. We finally see that we do not have a formal model of $\Omega^X(N)(w)$. Indeed, the morphism $\mathfrak{M}(Np^r)(w) \rightarrow \mathfrak{M}(N)(w)$ is only generically étale.

We now study what happens for different w 's. Let $w' \geq w$ be a rational number that satisfies the same condition of w . Let $i_{w',w}: \mathfrak{M}(Np^r)(w)^{\text{rig}} \hookrightarrow \mathfrak{M}(Np^r)(w')^{\text{rig}}$ (for any r) be the natural inclusion.

Proposition 5.14. *For any continuous character χ , we have a natural isomorphism*

$$\rho_{w',w}: i_{w',w}^*(\Omega^X(Np^r)(w')) \cong \Omega^X(Np^r)(w).$$

Furthermore $\rho_{w,w} = \text{id}$ and, if $w'' \geq w'$ is a rational number that satisfies the same condition of w , we have $\rho_{w'',w'} = \rho_{w',w} \circ \rho_{w'',w'}$.

Proof. This is [Bra12, Lemma 6.18]. \square

Let χ and χ' be continuous character and let h be an integer. We say that χ and χ' are h -close if the restriction of χ and χ' to $1 + p^h \mathbb{Z}_p$ coincide. We now study the relations between modular forms of weight χ and modular forms of weight χ' .

Proposition 5.15. *Let $0 \leq h \leq r$ be integers and let χ be a continuous character. If χ' is a continuous character that is r -close to χ , then we have*

$$(\Omega^X(Np^r)(w)[\psi^{-1}])^{G_{r,h}} = \Omega^{\chi'}(Np^h)(w),$$

where $\psi := \chi/\chi'$ is considered as a character $G_{r,h} \rightarrow K^*$ and by $\Omega^X(Np^r)(w)[\psi^{-1}]$ we mean that the action of $G_{r,h}$ is twisted by ψ^{-1} .

Proof. We may assume that χ and χ' are r -accessible. We have an isomorphism $S^D(K, w, K(Np^r), \chi) \cong S^D(K, w, K(Np^r), \chi')$ that sends a modular form f to the rule \tilde{f} defined by

$$(\mathcal{A}/S_r, i, \alpha, P, Y, \eta) \mapsto \psi(u)f(\mathcal{A}/S_r, i, \alpha, P, Y, \eta),$$

where $\eta = u(1 + p^{r-v}x)E_1$ (recall that $\mathcal{F}'_{r,w}$ is the trivial $\mathcal{S}_{r,w}$ -torsor generated by E_1). This isomorphism does not respect the action of the diamond operators but induces a G_r -equivariant isomorphism

$$S^D(K, w, K(Np^r), \chi) \cong S^D(K, w, K(Np^r), \chi')[\psi].$$

This means that if $f|_{\langle a \rangle} = \psi(a)^{-1}$ for all $a \in G_{r,h}$, then \tilde{f} is fixed under the action of $G_{r,h}$ and hence descends to level $K(Np^h)$. \square

Corollary 5.16. *Let $0 \leq h \leq r$ be integers and let χ be a continuous character. We have a $G_{r,h}$ -equivariant isomorphism*

$$S^D(K, w, K(Np^r), \chi) \cong \bigoplus_{\chi' \in I} S^D(K, w, K(Np^h), \chi'),$$

where I is a set of representative of the set of continuous characters that are r -close to χ modulo the equivalence relation ‘being h -close’.

Proof. Let us consider the $\mathcal{O}_{\mathfrak{M}(Np^h)(w)^{\text{rig}}}$ -module $\vartheta_{r,h,*}^{\text{rig}} \Omega^\chi(Np^r)(w)$. It has an action of $G_{r,h}$ and it decompose as a direct sum of the eigenspaces of $G_{r,h}$. Hence we have a decomposition

$$\vartheta_{r,h,*}^{\text{rig}} \Omega^\chi(Np^r)(w) = \bigoplus_{\psi: G_{r,h} \rightarrow K^*} (\Omega^\chi(Np^r)(w)[\psi^{-1}])^{G_{r,h}}.$$

Given any $\chi' \in I$, we set $\psi := \chi/\chi'$. By definition, ψ factors through a morphism $G_r \rightarrow K^*$ and hence it define a character $\psi: G_{r,h} \subseteq G_r \rightarrow K^*$. If $\chi'' \in I$ is different from χ' , then the morphism $\varphi: G_{r,h} \rightarrow K^*$ defined by χ'' must be different from ψ , otherwise χ' and χ'' would be h -close. On the other hand, any $\psi: G_{r,h} \rightarrow K^*$ is obtained by some $\chi' \in I$, so the corollary follows by Proposition 5.15. \square

We now explain a functorial property of the sheaves $\Omega^\chi(Np^r)(w)$ that generalizes the fact that we can pullback invariant differentials under an isogeny (see [Bra12, Remark 6.20] for details). Let $r \geq 1$ be an integer and let χ be an r -accessible character. Let $i_{\mathcal{A}}, i_{\mathcal{B}}: \text{Spf}(S_r) \rightarrow \mathfrak{M}(Np^r)(w)$ be two morphisms, corresponding the false elliptic curves \mathcal{A} and \mathcal{B} . Suppose we are given an isogeny $f: \mathcal{B} \rightarrow \mathcal{A}$ whose kernel intersects trivially the canonical subgroup of \mathcal{B} . We obtain an isomorphism

$$f^\chi: i_{\mathcal{B}}^{\text{rig},*} \Omega^\chi(Np^r)(w) \rightarrow i_{\mathcal{A}}^{\text{rig},*} \Omega^\chi(Np^r)(w)$$

that is the natural pullback of invariant differential if χ is an integer. If $0 \leq h \leq r$ is another integer, taking $G_{r,h}$ -invariants we get a similar morphism for $\Omega^\chi(Np^h)(w)$.

Everything we have done so far can be put in analytic families over the weight space, so we see that our sheaf $\Omega^\chi(Np^r)(w)$ really ‘interpolate’ the sheaves $\omega^{\otimes k}$.

Proposition 5.17. *For any integer $r \geq 1$ there is an invertible sheaf $\Omega(Np^h)(w)$, where $h \geq 0$, on $\mathcal{W}_r \times \mathfrak{M}(Np^h)(w)^{\text{rig}}$ such that if $\chi \in \mathcal{W}_r(K)$ is an r -accessible character, then we have a natural isomorphism*

$$(\chi, \text{id})^* \Omega(Np^h)(w) \cong \Omega^\chi(Np^h)(w).$$

If r_1 and r_2 are integers and w_1 and w_2 satisfy the usual condition with respect to r_1 and r_2 , then the restrictions of $\Omega(Np^h)(w_1)$ and $\Omega(Np^h)(w_2)$ to $\mathcal{W}_{r_1} \cap \mathcal{W}_{r_2} \times \mathfrak{M}(Np^h)(w_1)^{\text{rig}} \cap \mathfrak{M}(Np^h)(w_2)^{\text{rig}}$ coincide. Moreover, if $0 \leq h_1 \leq h_2$, then we have a natural action of G_{h_2, h_1} on $(\text{id} \times \vartheta_{h_2, h_1}^{\text{rig}})_ \Omega(Np^{h_2})(w)$ such that*

$$\left((\text{id} \times \vartheta_{h_2, h_1}^{\text{rig}})_* \Omega(Np^{h_2})(w) \right)^{G_{h_2, h_1}} = \Omega(Np^{h_1})(w).$$

Proof. We start assuming $h \geq r$. Let π_i be the natural projection from $\mathcal{W}_r \times \mathfrak{M}(Np^h)(w)^{\text{rig}}$ to the i -th factor. We write $\mathcal{S}_{r,v}$ for $\pi_2^{-1} \mathcal{S}_{r,w}$, and the same for $\mathcal{F}'_{r,w}$. Let $x = ub$ be a local section of $\mathcal{S}_{r,w}$. If $A \otimes B$ is a local section of $\mathcal{O}_{\mathcal{W}_r \times \mathfrak{M}(Np^h)(w)^{\text{rig}}}$ we define $x(A \otimes B)$ as the local section of $\mathcal{O}_{\mathcal{W}_r \times \mathfrak{M}(Np^h)(w)^{\text{rig}}}$ that corresponds to the function

$$(\chi, z) \mapsto \chi(u)A(\chi)b^x B(z),$$

for $\chi \in \mathcal{W}_r(T)$ and $z \in \mathfrak{M}(Np^h)(w)^{\text{rig}}(T)$, where T is any \mathbb{Q}_p -affinoid algebra. We define $\Omega(Np^h)(w)$ as

$$\Omega(Np^h)(w) := \mathcal{H}om_{\mathcal{S}_{r,w}}(\mathcal{F}'_{r,w}, \mathcal{O}_{\mathcal{W}_r \times \mathfrak{M}(Np^h)(w)^{\text{rig}}}).$$

We can prove all the proposition as in the case of a single character, and we can descend to arbitrary h taking $G_{r,h}$ -invariants. \square

6. THE EIGENCURVE

In this section will be mainly interested in level $K(N)$. We fix an r -accessible character χ , where $r \geq 1$ is an integer. We have that $S^D(K, w, K(Np^r), \chi)$ is by definition the global sections of a sheaf on $\mathfrak{M}(Np^r)(w)^{\text{rig}}$, that is an affinoid. Furthermore, we have a formal model of $\Omega^\times(Np^r)(w)$ that is invertible sheaf on $\mathfrak{M}(Np^r)(w)$. It follows (see [Bra12, Proposition 7.1]) that $S^D(K, w, K(Np^r), \chi)$ is a potentially orthonormizable K -Banach module with respect to the sup norm. For modular forms of level $K(N)$ we take G_r -invariants, so we have that $S^D(K, w, K(N), \chi)$ satisfies property (Pr) of [Buz07], that we now recall.

Definition 6.1. Let M be a Banach A -module, where A is an affinoid K -algebra. Following [Buz07, Part I, Section 2], we say that M satisfies the *property (Pr)*, if there is a Banach A -module N such that $M \oplus N$ is potentially orthonormizable.

Let $f \in S^D(K, w, K(Np^r), \chi)$ and let $T = (\mathcal{A}/S_r, i, \alpha, P, Y, \eta)$ be a test object of level $K(Np^r)$ and growth condition w . There is a flat (over V), normal, and p -adically complete S_r -algebra S'_r such that $S_{r,K} \rightarrow S'_{r,K}$ is finite étale and all finite and flat subgroup schemes of $\mathcal{A}[p^r]_K^1$ are defined over $S'_{r,K}$. By normality, all such subgroup schemes extend to subgroups of $\mathcal{A}[p^r]_{S'_r}^1$. We can base change T to S'_r , obtaining the test object T' , and by definition of modular form we have $f(T) = f(T')$. In particular, in the following, we can assume that all finite and flat subgroup schemes of $\mathcal{A}[p^r]^1$ are defined over S_r . Let \mathcal{D} be a finite and flat subgroup scheme of $\mathcal{A}[p^r]^1$ of order p that *intersect trivially the canonical subgroup of $\mathcal{A}[p^r]^1$* . By Remark 1.1, we have that \mathcal{D} defines a subgroup of $\mathcal{A}[p^r]$, and we denote with \mathcal{A}/\mathcal{D} the quotient of \mathcal{A} by this subgroup, that is a false elliptic curve. By our assumption on \mathcal{D} , the level structure on \mathcal{A} defines a level structure on \mathcal{A}/\mathcal{D} , denoted again (i, α, P) . Since \mathcal{D} intersects trivially the canonical subgroup of $\mathcal{A}[p^r]^1$, we have an isomorphism $i_1^* \mathcal{F}'_{r,w} \cong i_2^* \mathcal{F}'_{r,w}$, where $i_1, i_2: \text{Spf}(S_r) \rightarrow \mathfrak{M}(Np^r)(w)$ correspond to \mathcal{A} and \mathcal{A}/\mathcal{D} , respectively. Let η' be the image of η under this isomorphism. By [Bra12, Lemma 7.5], we have that the S_r -point of $\mathfrak{M}(Np^r)(w)$ corresponding to $(\mathcal{A}/\mathcal{D}, i, \alpha, P)$ factors through $\mathfrak{M}(Np^r)(w/p)$. Finally, we obtain a test object $T/\mathcal{D} := ((\mathcal{A}/\mathcal{D})/S_r, i, \alpha, Y, \eta')$ of level $K(Np^r)$ and growth condition w/p .

Definition 6.2. We define the U-operator on $S^D(K, w, K(Np^r), \chi)$ by

$$U f(T) = \frac{1}{p} \sum_{\mathcal{D}} f(T/\mathcal{D}),$$

where the sum is over all \mathcal{D} as above. Taking G_r -invariants, we obtain a similar operator on $S^D(K, w, K(N), \chi)$. By the above discussion, we have that the U-operator factors through $S^D(K, w/p, K(N), \chi)$.

Note that also the U-operator can be put in families.

Proposition 6.3. *If $w > 0$, the U operator on $S^D(K, w, K(N), \chi)$ is completely continuous.*

Proof. We can work at level $K(Np^r)$. The U-operator can be defined via a correspondence on $\mathfrak{M}(Np^r)(w)$ (see [Bra12, Section 7]), so it is a bounded operator. If $w > 0$, the natural map $S^D(K, w/p, K(Np^r), \chi) \rightarrow S^D(K, w, K(Np^r), \chi)$ is completely continuous, so the same is true for U. \square

We have the following result of classicality for our modular forms.

Proposition 6.4 ([Kas09, Theorem 5.1]). *Let $f \in S^D(K, w, K(N), k)$ and suppose that $Uf = af$ for some $a \in K$. If a satisfies $v(a) < k - 1$, then f is classical.*

Using the above notations, we now define the Hecke operators T_l , for $l \neq p$ a prime that does not divide δN . Let $f \in S^D(K, w, K(Np^r), \chi)$. We may assume that all the finite and flat subgroup schemes of order l of \mathcal{A} are defined over S_r . Since in this case it is automatic that any such subgroup scheme \mathcal{D} intersect trivially the canonical subgroup of $\mathcal{A}[p^r]^1$, we can define $T_l f$ as the rule

$$T \mapsto \frac{1}{l+1} \sum_{\mathcal{D}} f(T/\mathcal{D}).$$

Also the operators T_l can be defined using correspondences and can be put in families. In particular, they are bounded operators, but are not completely continuous.

Let $r \geq 1$ be an integer, and assume that $w > 0$ is a rational number sufficiently small. Let \mathcal{Z}_r be the spectral variety associated to the U -operator acting on $H^0(\mathcal{W}_r \times \mathfrak{M}(N)(w)^{\text{rig}}, \Omega(N)(w))$. We have proved that all assumptions needed to use the machine developed by Buzzard in [Buz07] are satisfied, so we have the following

Theorem 6.5. *We have a rigid space $\mathcal{C}_r \subseteq \mathcal{W}_r \times \mathbb{A}_K^{1, \text{rig}}$ equipped with a finite morphism $\mathcal{C}_r \rightarrow \mathcal{Z}_r$. If L is a finite extension of K , then the points of $\mathcal{C}_r(L)$ correspond to systems of eigenvalues of modular forms with growth condition w , finite slope with respect to the U -operator, and coefficients in L . If $x \in \mathcal{C}_r(L)$, let $\mathcal{M}(w)_x$ be the set of modular forms corresponding to x . Then all the elements of $\mathcal{M}(w)_x$ have weight $\pi_1(x) \in \mathcal{W}(L)$ and the U -operator acts on $\mathcal{M}(w)_x$ with eigenvalue $\pi_2(x)^{-1}$. We have that π_1 is, locally on \mathcal{W}_r and on \mathcal{C}_r , finite and surjective. For various r and w , these construction are compatible. Letting $r \rightarrow \infty$ we have $w \rightarrow 0$ and we obtain the global eigencurve $\mathcal{C} \subseteq \mathcal{W} \times \mathbb{A}_K^{1, \text{rig}}$.*

REFERENCES

- [AIS11] Fabrizio Andreatta, Adrian Iovita, and Glenn Stevens. On overconvergent modular forms. preprint. Available at <http://www.mathstat.concordia.ca/faculty/iovita>, 2011.
- [Bra12] Riccardo Brasca. p -adic modular forms of non-integral weight over Shimura curves. 2012. preprint, arXiv:1106.2712v4. To appear in *Compositio Mathematica*.
- [Buz07] Kevin Buzzard. Eigenvarieties. In *L-functions and Galois representations*, volume 320 of *London Math. Soc. Lecture Note Ser.*, pages 59–120. 2007.
- [Col97] Robert F. Coleman. Classical and overconvergent modular forms of higher level. *J. Théor. Nombres Bordeaux*, 9(2):395–403, 1997.
- [DT94] Fred Diamond and Richard Taylor. Nonoptimal levels of mod l modular representations. *Invent. Math.*, 115(3):435–462, 1994.
- [Kas99] Payman L. Kassaei. p -adic modular forms over Shimura curves over \mathbb{Q} . Phd-Thesis. Available at <http://www.mth.kcl.ac.uk/~kassaei/>, 1999.
- [Kas04] Payman L. Kassaei. \mathcal{P} -adic modular forms over Shimura curves over totally real fields. *Compos. Math.*, 140(2):359–395, 2004.
- [Kas09] Payman L. Kassaei. Overconvergence and classicality: the case of curves. *J. Reine Angew. Math.*, 631:109–139, 2009.
- [Lan78] Serge Lang. *Cyclotomic fields*. Springer-Verlag, New York, 1978. Graduate Texts in Mathematics, Vol. 59.
- [OT70] Frans Oort and John Tate. Group schemes of prime order. *Ann. Sci. École Norm. Sup. (4)*, 3:1–21, 1970.

E-mail address: riccardo.brasca@gmail.com

Current address: Max Planck Institute for Mathematics, Bonn, Germany
URL: <http://http://guests.mpim-bonn.mpg.de/ricky/>