

VECTOR BUNDLES ON NON-KAEHLER ELLIPTIC PRINCIPAL BUNDLES

VASILE BRÎNZĂNESCU, ANDREI D. HALANAY, AND GÜNTHER TRAUTMANN

ABSTRACT. We study relatively semi-stable vector bundles and their moduli on non-Kähler principal elliptic bundles over compact complex manifolds of arbitrary dimension. The main technical tools used are the twisted Fourier-Mukai transform and a spectral cover construction. For the important example of such principal bundles, the numerical invariants of a 3-dimensional non-Kähler elliptic principal bundle over a primary Kodaira surface are computed.

1. INTRODUCTION

The study of vector bundles over elliptic fibrations has been a very active area of research in both mathematics and physics over the past twenty years; in fact, there is by now a well understood theory for projective elliptic fibrations (see for example [Don97], [DP08], [Fri89], [FMW99], [BM02],[RMP02], [Bri98], etc.). However, not very much is known about the non-Kähler case; the study of rank 2 vector bundles on non-Kähler elliptic surfaces is done in [BM05],[BM06]. In this article we study relatively semi-stable vector bundles on non-Kähler principal elliptic bundles over complex manifolds of arbitrary dimension with the invariant $\delta \neq 0$. One of the motivations for the study of vector bundles on non-Kähler elliptic n-folds comes from recent developments in superstring theory, where six-dimensional non-Kähler manifolds occur in the context of $\mathcal{N} = 1$ supersymmetric heterotic and type II string compactifications with non-vanishing background H - field; in particular most of the non-Kähler examples appearing in the physics literature so far are non-Kähler principal elliptic fibrations (see [BBDG03], [CCD⁺03], [GP04]). There are also two classes of non-Kähler Calabi-Yau type threefolds appearing in the mathematical and physical literature: one is due to M. Gross (privately communicated to us by A. Căldăraru). Other examples appear in [Add09a], [Add09b],[Kuz08] and [CDH⁺10]. The main technical tools used are the twisted Fourier-Mukai transform, introduced by A.Căldăraru (see [Cal00]) and the spectral cover construction, see [FMW99], [Don95], [BBRMP98], [RMP02].

The paper is organized as follows. In the second section we determine the structure of the relative Jacobian of a principal elliptic bundle as a moduli space and find out that it is the product of the fiber with the basis. In the third and fourth sections, using the relative Jacobian, we adapt the construction of Căldăraru, [Că02], to our case, obtaining a twisted Fourier-Mukai transform. Similar results were obtained in different settings by O.Ben-Bassat [BB09] and I.Burban and B.Kreussler

1991 *Mathematics Subject Classification.* 14J60, 32L05, 14D22, 14F05,32J17,32Q25.

Key words and phrases. non-Kähler principal elliptic bundles, Calabi-Yau type threefolds, holomorphic vector bundles, moduli spaces.

V.Brînzănescu was partially supported by CNCSIS contract 1189/2009-2011.

[BK06]. In the fifth section using this transform and the associated spectral cover we prove that the moduli space of rank n , relatively semi-stable vector bundles is corepresented by the relative Douady space of length n and relative dimension 0 subspaces of the relative Jacobian, see theorem 5.

After reviewing some background results on torus bundles from [Höf93] in the first appendix, in the second appendix we compute the numerical invariants (Hodge and Betti numbers) of a principal elliptic bundle over a primary Kodaira surface and use them to distinguish the non-trivial elliptic bundles. These invariants are also of interest for physicists working on heterotic string-theory models with non-Kähler Calabi-Yau type threefolds as backgrounds.

Acknowledgements

Part of this paper was prepared during the stay at the Kaiserslautern Technical University of A.D. Halanay and V. Brînzănescu with scholarships offered by the Alexander von Humboldt Foundation in the framework of the Stability Pact. The first author expresses his gratitude to the Max-Planck-Institute für Mathematik in Bonn; part of this paper was prepared during his stay there.

2. LINE BUNDLES ON ELLIPTIC PRINCIPAL BUNDLES

In this section we shall be concerned with the study of the (coarse) moduli space of line bundles over a principal elliptic bundle $\pi : X \rightarrow S$, where S is a compact complex manifold, with fiber $E := E_\tau := \mathbb{C}/\Lambda$ ($\Lambda = \mathbb{Z} \oplus \tau\mathbb{Z}$). Among the invariants of such bundles is the homomorphism $\delta : H^1(E, \mathbb{C}) \rightarrow H^2(S, \mathbb{C})$ which is the d_2 -differential $E_2^{0,1} \rightarrow E_2^{2,0}$ of the Leray spectral sequence of the sheaf \mathbb{C}_X with terms $E_2^{p,q} = H^p(S, R^q\pi_*\mathbb{C}_X) \cong H^p(S, \mathbb{C}) \otimes H^q(E, \mathbb{C})$, see also the first appendix for more invariants.

We make the assumption that $\delta \neq 0$. In particular, $X \rightarrow S$ does not have the topology of a product. We should note here that if S is Kähler, then X is non-Kähler if and only if $\delta \neq 0$, see [Höf93].

We shall need in the sequel the following result of Deligne, [Del68], in the formulation of [Höf93, Prop.5.2].

Theorem 1. *Let $X \rightarrow S$ be a principal elliptic bundle. Then the following statements are equivalent:*

- a) *The Leray spectral sequence for \mathbb{C}_X degenerates at the E_2 -level;*
- b) *$\delta : H^1(E, \mathbb{C}) \rightarrow H^2(X, \mathbb{C})$ is the zero map;*
- c) *The restriction map $H^2(X, \mathbb{C}) \rightarrow H^2(E, \mathbb{C})$ to a fibre takes a non-zero value in $H_E^{1,1}$.*

In our case the preceding theorem has a very important consequence

Corollary 1. *Let $X \rightarrow S$ be a principal elliptic bundle with S a compact complex manifold and $\delta \neq 0$. Then for any vector bundle \mathcal{F} over X and any $s \in S$ the bundle $\mathcal{F}|_{X_s}$ has degree 0.*

Proof. Indeed let $r : H^2(X, \mathbb{C}) \rightarrow H^2(E, \mathbb{C})$ be the restriction map. We have that $c_1(\mathcal{L}|_{X_s}) = r(c_1(\mathcal{L})) = 0$ by the theorem. \square

Let us recall now the definition of the Jacobian variety J of a smooth curve C , see for instance [Har77, IV.4]. Let $\text{Pic}^0(C/T)$ for any analytic space T denote the

group

$$\{\mathcal{L} \in \text{Pic}(T \times C) \mid \deg(\mathcal{L}|_{\{t\} \times C}) = 0 \text{ for any } t \in T\} / p^* \text{Pic}(T),$$

where $p : T \times C \rightarrow T$ is the second projection. The Jacobian variety of C will be a variety J , together with an element $\mathcal{P} \in \text{Pic}^0(C/J)$ such that for any analytic space T and any $\mathcal{M} \in \text{Pic}^0(C/T)$ there is a unique morphism $f : T \rightarrow J$ such that $(f \times \text{id}_C)^* \mathcal{P} \sim \mathcal{M}$ in $\text{Pic}^0(T \times C)$, i.e. J represents the functor $T \mapsto \text{Pic}^0(C/T)$. It is well known that J exists for any smooth curve C . When C is an elliptic curve E then $J = E^*$, the dual torus, and \mathcal{P} is called a Poincaré bundle. In this case \mathcal{P} is a line bundle over $E^* \times E$ such that $\mathcal{P}|_{\{[\mathcal{L}]\} \times E} \simeq \mathcal{L}$, and $E^* \simeq \text{Pic}^0(E)$. We pass now to the relative case for elliptic principal bundles.

Definition 1. Let $X \xrightarrow{\pi} S$ be an elliptic principal bundle with typical fibre an elliptic curve E_τ and base S a smooth manifold. Let $F : (\text{An}/S)^{\text{op}} \rightarrow (\text{Sets})$ be the functor from the category of analytic spaces over S to the category of sets, given, for any commutative diagram

$$(2.1) \quad \begin{array}{ccc} X_T & \longrightarrow & X \\ \downarrow \pi & & \downarrow \pi \\ T & \longrightarrow & S, \end{array}$$

where $X_T := X \times_S T$, by

$$F(T) := \{\mathcal{L} \text{ invertible on } X_T \mid \deg(\mathcal{L}|_{X_{T,t}}) = 0, \text{ for all } t \in T\} / \sim,$$

where $\mathcal{L}_1 \sim \mathcal{L}_2$ if there is a line bundle L on T such that $\mathcal{L}_1 \simeq \mathcal{L}_2 \otimes \pi^* L$.

A variety J over S will be called the relative Jacobian of X if

(i) it corepresents the functor F , see [HL97, Def. 2.2.1], i.e. there is a natural transformation $F \xrightarrow{\sigma} \text{Hom}_S(-, J)$ and for any other variety N/S with a natural transformation $F \xrightarrow{\sigma'} \text{Hom}_S(-, N)$ there is a unique S -morphism $J \xrightarrow{\nu} N$ such that $\nu_* \circ \sigma = \sigma'$.

(ii) for any point $s \in S$ the map $F(\{s\}) \rightarrow \text{Hom}_S(\{s\}, J) \simeq J_s$ is bijective. Then each fibre J_s is the Jacobian of the fibre $X_s \simeq E$.

If X is projective, the existence of the relative Jacobian is well known, because it can be identified with the coarse relative moduli space of stable locally free sheaves of rank 1 and degree 0 on the fibres of X , see [HL97], [Cal00]. The relative Jacobian exists also in our non-Kähler case. It is just the product $S \times E^*$ and has the following special properties.

Theorem 2. (i) The functor F is corepresented by $J := S \times E^*$.

(ii) For any point $s \in S$ the map $F(\{s\}) \rightarrow \text{Hom}_S(\{s\}, J) \simeq J_s \simeq E^*$ is bijective.

(iii) The map $\sigma(T)$ is injective for any complex space T .

(iv) The functor F is locally representable by $J = S \times E^*$, i.e. if $U \subset S$ is a trivializing open subset, $\sigma(U)$ is bijective.

For the proof we use the following Seesaw lemma and its Corollary.

Lemma 1. Let $Y \xrightarrow{q} T$ be a principal elliptic bundle over a complex analytic space T with fibre E and let \mathcal{M} be an invertible sheaf on Y such that $\mathcal{M}|_{Y_t} = \mathcal{M}|_{Y_t}$ is trivial on any fibre of q . Then $L = q_* \mathcal{M}$ is locally free of rank 1 and $\mathcal{M} = q^* L$.

Proof. Because the statements are local over T and Y is locally trivial, we may assume that $Y = T \times E$ and that q is the first projection p_1 . We first note that the canonical homomorphism $p_1^*p_{1*}\mathcal{M} \rightarrow \mathcal{M}$ is an isomorphism because each \mathcal{M}_t is trivial so that it is an isomorphism when restricted to a fibre of p_1 . In order to show that $p_{1*}\mathcal{M}$ is invertible we consider the sheaf $\mathcal{M}(a) := \mathcal{M} \otimes p_2^*\mathcal{O}_E(a)$ for some $a \in E$. Then $h^0(\mathcal{M}_t(a)) = 1$ and $h^1(\mathcal{M}_t(a)) = 0$, whereas $h^1(\mathcal{M}_t) = 1$. Let $\phi^i(t) : (R^i p_{1*}\mathcal{M}(a))(t) \rightarrow H^i(X_t, \mathcal{M}_t(a))$ denote the base change homomorphism. Because $\phi^1(t) = 0$ it follows that $R^1 p_{1*}\mathcal{M}(a) = 0$, and by that that $L := p_{1*}\mathcal{M}(a)$ is invertible, see e.g. [Har77, Th.12.11].

Let $\mathbb{C}(a)$ be the sky-scraper sheaf with stalk \mathbb{C} at $a \in E$. Then $\mathcal{M} \otimes p_2^*\mathbb{C}(a) = \mathcal{M}|_{T \times \{a\}}$ and we have the exact sequence

$$0 \rightarrow \mathcal{M} \rightarrow \mathcal{M}(a) \rightarrow \mathcal{M} \otimes p_2^*\mathbb{C}(a) \rightarrow 0,$$

giving rise to the long exact sequence

$$0 \rightarrow p_{1*}\mathcal{M} \rightarrow p_{1*}\mathcal{M}(a) \xrightarrow{\alpha} p_{1*}(\mathcal{M}|_{S \times \{a\}}) \rightarrow R^1 p_{1*}(\mathcal{M}) \rightarrow 0.$$

Denoting by \mathcal{A} the image of α and pulling back to y , we obtain the exact sequence

$$p_1^*p_{1*}\mathcal{M} \xrightarrow{\gamma} p_1^*p_{1*}\mathcal{M}(a) \rightarrow p_1^*\mathcal{A} \rightarrow 0.$$

The restriction of γ to a fibre becomes the canonical map

$$H^0(X_t, \mathcal{M}_t) \otimes \mathcal{O}_{Y_t} \rightarrow H^0(Y_t, \mathcal{M}_t(a)) \otimes \mathcal{O}_{Y_t},$$

which is an isomorphism. Hence $(p_1^*\mathcal{A})_t = 0$ for any $t \in T$, and then also $p_1^*\mathcal{A} = 0$ and finally $\mathcal{A} = 0$. This proves that $p_{1*}\mathcal{M} \cong p_{1*}\mathcal{M}(a) = L$, and we have $\mathcal{M} \cong p_1^*p_{1*}\mathcal{M} \cong p_1^*L$.

One should note here that $p_1^*p_{1*}\mathcal{M}(a) \rightarrow \mathcal{M}(a)$ is not an isomorphism, having $\mathcal{M}|_{T \times \{a\}}$ as its cokernel, because $H^0(X_t, \mathcal{M}_t(a)) \otimes \mathcal{O}_{Y_t} \rightarrow \mathcal{M}_t(a)$ is not an isomorphism. \square

Corollary 2. *Let T be a complex space and $T \times E_\tau \xrightarrow{\theta} T \times E_\tau$ an isomorphism of the form $\theta(t, \alpha) = (t, \alpha + \lambda(t))$ with $\lambda : T \rightarrow E$ a holomorphic map. Then for any invertible sheaf \mathcal{L} on $T \times E_\tau$ with $\deg(\mathcal{L}_t) = 0$ for any $t \in T$, there is an invertible sheaf L on T such that $\theta^*\mathcal{L} \simeq \mathcal{L} \otimes p_1^*L$.*

Proof. Because \mathcal{L}_s has degree 0, $(\theta_s^*\mathcal{L}_s) \otimes \mathcal{L}_s^{-1}$ is trivial on any fiber (recall that a line bundle of degree 0 is isomorphic with its pull-back via a translation, see [BL04], [Mum70]). Now apply the Lemma to $\theta^*\mathcal{L} \otimes \mathcal{L}^{-1}$ \square

Proof of Theorem 2. Let $f : T \rightarrow S$ be a morphism of analytic spaces and let \mathcal{L} be a line bundle on X_T . Let $\{S_i\}$ be an open cover of S that gives a trivialization of the bundle X (that is $X_i := X|_{S_i} \simeq S_i \times E_\tau$). Taking the inverse image of this cover we get an open cover $\{T_i = f^{-1}S_i\}$ of T with the same property. Let us denote $X_{T,i} := X_T|_{T_i}$. There are trivializing maps

$$(2.2) \quad \begin{array}{ccc} X_{T,i} & \xrightarrow[\sim]{\theta_i} & T_i \times E_\tau \\ & \searrow & \swarrow \\ & T_i & \end{array}$$

Let \mathcal{L}_i be the sheaf on $T_i \times E$ defined by $\theta_i^* \mathcal{L}_i = \mathcal{L}|_{X_{T,i}}$ such that $\mathcal{L}_j \simeq (\theta_i \circ \theta_j^{-1})^* \mathcal{L}_i$ over T_{ij} .

Let now \mathcal{P} be a Poincare bundle on $E_\tau^* \times E_\tau$. Then for any i we'll have a **unique** morphism $\phi_i : T_i \rightarrow E_\tau^*$ such that $\mathcal{L}_i \sim (\phi_i \times \text{id})^*(\mathcal{P})$. Taking into account that $\theta_i \circ \theta_j^{-1} = \text{id} \times \theta_{ij}$ and θ_{ij} acts by translations, the preceding corollary implies that $\mathcal{L}_i \sim \mathcal{L}_j$ on T_{ij} . Therefore $\phi_i = \phi_j$ on T_{ij} . So we are given a global morphism $\phi : T \rightarrow E_\tau^*$.

Let now

$$(2.3) \quad \begin{array}{ccc} T & \xrightarrow{\tilde{\phi}} & S \times E_\tau^* = J \\ & \searrow & \swarrow \\ & S & \end{array}$$

be the corresponding map $\tilde{\phi} := (f, \phi)$. This provides us with a map $F(T) \xrightarrow{\sigma(T)} \text{Hom}_S(T, J)$. It is straightforward to check that $\sigma : F \rightarrow \text{Hom}_S(-, J)$ is a morphism of functors. The minimality of J will be proved after the proof of (iv).

ii) Property (ii) follows directly from the definition of the maps $\sigma(\{s\})$.

iii) We show next that each map $\sigma(T) : F(T) \rightarrow \text{Hom}_S(T, J)$ is injective. For that let \mathcal{L} and \mathcal{L}' be two line bundles such that their respective maps ϕ and ϕ' are equal. We need to show that $\pi_*(\mathcal{L}' \otimes \mathcal{L}^{-1}) =: L$ is locally-free.

We have that $\phi_i = \phi'_i$ for any i . This implies that over $X_{T,i}$ we have

$$(2.4) \quad \mathcal{L}_i \sim \mathcal{L}'_i.$$

Because of the relation (2.4) $\mathcal{L}'_t \otimes \mathcal{L}_t^{-1}$ is trivial for any $t \in T$. By the above corollary we obtain that $\pi_*(\mathcal{L}' \otimes \mathcal{L}^{-1})$ is locally free.

iv) Let $U \subset S$ be trivializing for the bundle X such that $X_U \simeq U \times E$. By the universal property of the dual torus E^* , the map $\sigma(U)$ as well as all the maps $\sigma(T)$ for $T \rightarrow U$ are bijective.

Let now $F \xrightarrow{\sigma'} \text{Hom}_S(-, N)$ be any natural transformation. For any $s \in S$ there is the map $\nu_s := \sigma'(\{s\}) \circ \sigma(\{s\})^{-1} : J_s \rightarrow N_s$, thus defining a map $\nu : J \rightarrow N$ over S . In order to show that ν is a morphism, we just remark that ν_s is the restriction of the map $\sigma'(U \times E^*) \circ \sigma(U \times E^*)^{-1}(\text{id}) : J_U \rightarrow N_U$ for a trivializing open subset $U \subset S$ and any $s \in U$.

Finally $\nu_* \circ \sigma = \sigma'$ follows from (iv) of the theorem and the fact that the functor $\text{Hom}_S(-, N)$ is a sheaf, using a trivializing covering for X of S . This completes the proof of (i). \square

Remark 1. *There is a very convenient description of $\sigma(S)$ as follows. Let \mathcal{L} be an arbitrary line bundle over X . We know that $[\mathcal{L}] \in F(S)$ by Corollary 1. By the above proof $\phi := \sigma(S)([\mathcal{L}]) : S \rightarrow S \times E^*$ is given by $\phi(s) = (s, x)$ with $x = [\mathcal{L}|_{X_s}]$.*

It will follow from theorem 3 that the relative Jacobian $J = S \times E^*$ is only a coarse moduli space under our assumption on X . However, by property (iv) of the theorem one can find a system of local universal sheaves which will form a twisted sheaf in the next section as in [Cal00], chapter 4.

3. THE TWISTED UNIVERSAL SHEAF

In the following we replace the relative Jacobian J by $S \times E$ via the canonical isomorphism between E and E^* . Then the local trivialisations $X_i \xrightarrow{\theta_i} S_i \times E$ are at the same time isomorphisms between X_i and $J_i := S_i \times E$. The local universal sheaves \mathcal{U}_i on $X_{iJ} = J \times_S X_i = J_i \times_{S_i} X_i$ are then given as pull backs of the universal sheaf $\mathcal{O}_{E \times E}(\Delta) \otimes p_2^* \mathcal{O}_E(-p_0)$ for the classical Jacobian of the elliptic curve E , after fixing an origin $p_0 \in E$ and where Δ is the diagonal.

Denoting by ρ_i the composition of maps

$$X_{iJ} \xrightarrow{id \times \theta_i} J \times_S (S_i \times E) \simeq S_i \times E \times E \rightarrow E \times E,$$

and by p_X the projection from X_{iJ} to X_i , the local universal sheaf becomes

$$\mathcal{U}_i = \rho_i^*(\mathcal{O}_{E \times E}(\Delta) \otimes p_2^* \mathcal{O}_E(-p_0)) \simeq \mathcal{O}_{X_{iJ}}(\Gamma_i) \otimes p_X^* \mathcal{O}_{X_i}(-s_i),$$

where Γ_i is the inverse of the diagonal (or the graph of the map θ_i) and s_i is the section of X_i corresponding to the reference point p_0 under the isomorphism θ_i , see [Cal00], prop. 4.2.3.

To measure the failure of these bundles to glue to a global universal one let us consider the line bundles $\mathcal{M}_{ij} := \mathcal{U}_i \otimes \mathcal{U}_j^{-1}$ over $J \times_S X_{ij}$. Then the restriction of \mathcal{M}_{ij} to a fibre X_s of the projection $J \times_S X_i \xrightarrow{q_i} J$ is trivial because both \mathcal{U}_j and \mathcal{U}_i restrict to isomorphic sheaves. By Lemma 1 there are invertible sheaves \mathcal{F}_{ij} on $J_{ij} = S_{ij} \times E$ such that $\mathcal{M}_{ij} = q_i^* \mathcal{F}_{ij}$.

This collection of line bundles satisfies the following properties:

1. $\mathcal{F}_{ii} = \mathcal{O}_{J_i}$;
2. $\mathcal{F}_{ji} = \mathcal{F}_{ij}^{-1}$;
3. $\mathcal{F}_{ij} \otimes \mathcal{F}_{jk} \otimes \mathcal{F}_{ki} =: \mathcal{F}_{ijk}$ is trivial, with trivialization induced by the canonical one of $\mathcal{M}_{ij} \otimes \mathcal{M}_{jk} \otimes \mathcal{M}_{ki}$;
4. $\mathcal{F}_{ijk} \otimes \mathcal{F}_{jkl}^{-1} \otimes \mathcal{F}_{kli} \otimes \mathcal{F}_{lij}^{-1}$ is canonically trivial.

These conditions tell us that the collection $\{\mathcal{F}_{ij}\}$ represents a gerbe (see [DP08]) and gives rise to an element $\alpha \in H^2(J, \mathcal{O}_J^*)$. More explicitly, α is defined as follows. We may assume that the sheaves \mathcal{F}_{ij} are already trivial with trivializations $a_{ij} : \mathcal{O}_J \simeq \mathcal{F}_{ij}$ over J_{ij} .

If $c_{ijk} : \mathcal{O}_J \simeq \mathcal{F}_{ijk}$ is the isomorphism which is induced by the canonical trivialization of $\mathcal{M}_{ij} \otimes \mathcal{M}_{jk} \otimes \mathcal{M}_{ki}$, then

$$(3.1) \quad a_{ij} \otimes a_{jk} \otimes a_{ki} = \alpha_{ijk} c_{ijk}$$

with scalar functions α_{ijk} which then define a cocycle for the sheaf \mathcal{O}_J^* , thus defining the class $\alpha \in H^2(J, \mathcal{O}_J^*)$, see [Cal00], sect. 4.3. It is straightforward to prove:

Lemma 2. *The sheaves \mathcal{U}_i can be glued to a global universal sheaf if and only if the class $\alpha = 0$.*

The element α is related to the element $\xi \in H^1(S, \mathcal{O}_S(E))$ which is defined by the cocycle of the elliptic bundle $X \rightarrow S$, using the Ogg-Shafarevich group $\text{III}_S(J)$ of J , see [Cal00], section 4.4. There is an exact sequence $0 \rightarrow \text{Br}(S) \rightarrow \text{Br}(J) \xrightarrow{\pi} \text{III}_S(J) \rightarrow 0$, where $\text{Br}(S) \simeq H^2(J, \mathcal{O}_J^*)$ is the analytic Brauer group of S and $\text{III}_S(J)$ is isomorphic to $H^1(S, \mathcal{O}_S(E))$ in our setting. We have the

Theorem 3. ([Cal00, Th 4.4.1]) $\xi = \pi(\alpha)$.

Because $\xi \neq 0$ in our case, $\alpha \neq 0$, and thus the local universal sheaves cannot be glued to a global universal sheaf by preserving the bundle structure on the elliptic fibres.

4. THE TWISTED FOURIER-MUKAI TRANSFORM

The collection of local universal sheaves above can be considered as an α -twisted sheaf with which one can define a Fourier-Mukai transform. Recall the definition of an α -twisted sheaf on a complex space or on an appropriate scheme X .

Definition 2. *Let $\alpha \in C^2(\mathfrak{U}, \mathcal{O}_X^*)$ be a Čech 2-cocycle, given by an open cover $\mathfrak{U} = \{U_i\}_{i \in I}$ and sections $\alpha_{ijk} \in \Gamma(U_i \cap U_j \cap U_k, \mathcal{O}_X^*)$. An α -twisted sheaf on X will be a pair of families $(\{\mathcal{F}_i\}_{i \in I}, \{\varphi_{ij}\}_{i,j \in I})$ with \mathcal{F}_i a sheaf of \mathcal{O}_X -modules on U_i and $\varphi_{ij} : \mathcal{F}_j|_{U_i \cap U_j} \rightarrow \mathcal{F}_i|_{U_i \cap U_j}$ isomorphisms such that*

- φ_{ii} is the identity for all $i \in I$.
- $\varphi_{ij} = \varphi_{ji}^{-1}$, for all $i, j \in I$.
- $\varphi_{ij} \circ \varphi_{jk} \circ \varphi_{kl}$ is multiplication by α_{ijk} on $\mathcal{F}_i|_{U_i \cap U_j \cap U_k}$ for all $i, j, k \in I$.

It is easy to see that the coherent α -twisted sheaves on X make up an abelian category and thus give rise to a derived category $\mathcal{D}^b(X, \alpha)$. For further properties of α -twisted sheaves, see [Cal00].

With the notation above, the family (\mathcal{U}_i) becomes a twisted sheaf \mathcal{U} w.r.t. the cocycle $p_J^* \alpha$ of the sheaf $\mathcal{O}_{J \times_S X}$ as follows. The trivializations a_{ij} of the \mathcal{F}_{ij} induce isomorphisms $\phi_{ij} : \mathcal{U}_j \simeq \mathcal{U}_i$ which satisfy the definition of a twisted sheaf because of identity 3.1. We also need the dual \mathcal{V} of \mathcal{U} on $J \times_S X$ which locally over S_i is given by

$$\mathcal{V}_i = \rho_i^*(\mathcal{O}_{E \times E}(-\Delta) \otimes p_2^* \mathcal{O}_E(p_0)) \simeq \mathcal{O}_{X_i J}(-\Gamma_i) \otimes p_X^* \mathcal{O}_{X_i}(s_i).$$

It follows that \mathcal{V}_i is α^{-1} -twisted. We let \mathcal{V}^0 and \mathcal{U}^0 denote the extensions of \mathcal{V} and \mathcal{U} to $J \times X$ by zero.

The following theorem supplies us with the main tool for the treatment of the moduli spaces $M_X(n, 0)$ of relatively semistable vector bundles on X of rank n and degree 0 on the fibres X_s in section 5. It is an analog of theorem [Cal00, Th.6.5.4](and also [Că102]):

Theorem 4. *Let $X \xrightarrow{\pi} S$ be an elliptic principal fiber bundle. Let $\alpha \in Br(J)$ be the obstruction to the existence of the universal sheaf on $J \times_S X$ and let \mathcal{U} be the associated $p_J^*(\alpha)$ -twisted universal sheaf on $J \times_S X$ with its dual \mathcal{V} as above.*

Then the twisted Fourier-Mukai transform $\Psi : \mathcal{D}^b(J, \alpha) \rightarrow \mathcal{D}^b(X)$, given by $\Psi(\mathcal{F}) := Rp_{X}(\mathcal{V}^0 \otimes^L Lp_J^* \mathcal{F})$ is an equivalence of categories, where p_J and p_X are the product projections*

$$(4.1) \quad J \xleftarrow{p_J} J \times X \xrightarrow{p_X} X$$

Note here that $\mathcal{V}^0 \otimes^L Lp_J^* \mathcal{F}$ is an object in the category of untwisted sheaves.

Proof. The theorem follows from the Căldăraru's version of the Bridgeland (Orlov, Mukai, etc.)-criterion ([Orl03], [Bri99], [Cal00, Th.3.2.1]), applied to our case. Due to a private communication this criterion works also in the case when α is not torsion¹. It follows that the functor Ψ is fully faithful if and only if for each point $y \in J$ and its skyscraper sheaf $\mathbb{C}(y)$, $\text{Hom}(\Psi(\mathbb{C}(y)), \Psi(\mathbb{C}(y))) = \mathbb{C}$ and for any

¹The authors are indebted to Andrei Căldăraru for this information.

$y_1, y_2 \in J$, $\text{Ext}^i(\Psi(\mathbb{C}(y_1)), \Psi(\mathbb{C}(y_2))) = 0$, unless $y_1 = y_2$ and $0 \leq i \leq \dim(J)$. Moreover Ψ is an equivalence of categories if and only if for any $y \in J$ we have $\Psi(\mathbb{C}(y)) \otimes \omega_X \simeq \Psi(\mathbb{C}(y))$. Note that in our case the canonical bundle is trivial so that the last condition is automatically satisfied.

In order to compute $\Psi(\mathbb{C}(y))$ for a point $y \in J$, let $s \in S_i$ be the image in S and consider $\mathcal{V}^0 \otimes p_J^* \mathbb{C}(y)$. Its support is $(J \times_S X) \cap (\{y\} \times X) = \{y\} \times X_s$. We may therefore identify \mathcal{V} with \mathcal{V}_i and obtain $\mathcal{V}^0|_{\{y\} \times X} \simeq \mathcal{O}_{X_s}(-x + s_i(s))$, where $\theta_i(x) = y$ and s_i denotes the local section of X corresponding to p_0 . Because $p_J^* \mathbb{C}(y) = \mathcal{O}_{y \times X_s}$, we obtain $\Psi(\mathbb{C}(y)) \simeq p_{X*} \mathcal{O}_{X_s}(-x + s_i(s)) \simeq i_{s*} \mathcal{O}_{X_s}(-x + s_i(s))$ by the base change isomorphism for the inclusion i_s which holds in this case because X_s is smooth as is the projection p_X , see [BO95, Lemma 1.3] Using this, we conclude that $\text{Hom}(\Psi(\mathbb{C}(y)), \Psi(\mathbb{C}(y))) = \mathbb{C}$ and we proceed in the same way for Ext . \square

Remark 2. *On can see this result in connection with section 6 of [KO03], since the element $\alpha \in \text{Br}(J)$ is not torsion.*

In the sequel we shall work with the adjoint transform

$$\Phi(-) = R p_{J*}(\mathcal{U}^0 \otimes^L L p_X^*(-))$$

of Ψ , with kernel \mathcal{U}^0 . It is the reverse equivalence, see [BM02, 8.4], [Huy06], [BBHR09] for the untwisted situation.

We need the following special cases of base change properties.

Proposition 1. *For any $s \in S$ let $i_s : X_s \rightarrow X$ and $j_s : \{s\} \times E \rightarrow J$ be the natural inclusions. Then the canonical morphism of functors*

$$(4.2) \quad \mathbf{L}j_s^* \circ \Phi \simeq \Phi_s \circ \mathbf{L}i_s^*,$$

is an isomorphism, where Φ_s is the classical Fourier-Mukai transform associated with the Poincarè bundle over $E \times E$.

Proof. Let \tilde{j}_s denote the inclusion of $J_s \times X_s$ into $J \times_S X$ and let p_{J_s} be the first projection of $J_s \times X_s$. By [BO95, Lemma 1.3] $Lj_s^* p_{J*} = R p_{J_s*} \tilde{j}_s^*$. Then

$$Lj_s^* p_{J*}(\mathcal{V}^0 \otimes p_X^* \mathcal{F}) \simeq R p_{J_s*} \tilde{j}_s^* \mathcal{V}^0 \otimes p_X^* \mathcal{F} \simeq (\mathcal{V}|_{J_s \times X_s}) \otimes p_{X_s}^* i_s^* \mathcal{F},$$

which implies the formula. \square

The following definition is very useful for dealing with the spectral covers in the next section.

Definition 3. ([Muk81]) *We denote by $\Phi^i(\mathcal{F})$ the i -th term of the complex $\Phi(\mathcal{F})$. We say that the sheaf \mathcal{F} is Φ -WIT $_i$ (the weak index theorem holds) if $\Phi^i(\mathcal{F}) \neq 0$ and $\Phi^j(\mathcal{F}) = 0$ for any $j \neq i$. Moreover if \mathcal{F} is WIT $_i$ and $\Phi^i(\mathcal{F})$ is locally free we say that \mathcal{F} is IT $_i$.*

Consider now a rank n vector-bundle \mathcal{F} over the principal elliptic bundle X and denote its restriction to a fibre X_s by \mathcal{F}_s . From Proposition 1 it follows that if \mathcal{F}_s is Φ_s -WIT $_i$ for any s then \mathcal{F} is Φ -WIT $_i$.

5. A SPECTRAL COVER AND VECTOR BUNDLES ON X

In this section we shall apply the twisted Fourier-Mukai transform to the moduli problem for rank- n relatively semi-stable vector bundles on the principal elliptic bundle X . By Deligne's theorem (Theorem 1), the degree of the restriction \mathcal{F}_s

of any vector bundle \mathcal{F} on X is 0 for any $s \in S$. Therefore we consider the set $MS_X(n, 0)$ of rank- n vector bundles on X which are fibrewise semistable and of degree zero, together with its quotient

$$M_X(n, 0) := MS_X(n, 0) / \sim$$

of equivalence classes, where two bundles are defined to be equivalent if they are fibrewise S-equivalent.

Let us recall that a vector bundle \mathcal{E} on a smooth projective curve is called semistable if for any proper subbundle \mathcal{E}' ,

$$\deg(\mathcal{E}') / \text{rank}(\mathcal{E}') \leq \deg(\mathcal{E}) / \text{rank}(\mathcal{E}).$$

For such bundles there is the standard notion of S-equivalence, see e.g. [HL97].

It is well-known that the semistable vector bundles of degree zero on the elliptic curve E are direct sums

$$(5.1) \quad \mathcal{E} = \bigoplus_i \mathcal{A}_{n_i} \otimes \mathcal{O}_E(x_i - p_0),$$

where the \mathcal{A}_n denote the indecomposable Atiyah bundles of degree zero which are inductively defined by nontrivial extensions $0 \rightarrow \mathcal{O}_E \rightarrow \mathcal{A}_n \rightarrow \mathcal{A}_{n-1} \rightarrow 0$ with $\mathcal{A}_1 = \mathcal{O}_E$, see [Ati57] [FMW99, Def. 1.12] or [Tu93]. It follows that each such \mathcal{E} is S-equivalent to a direct sum $\text{gr}(\mathcal{E}) = \bigoplus_j \mathcal{O}_E(y_j - p_0)^{\oplus m_j}$ with pairwise distinct points y_j .

Proposition 2. ([RMP02], [BBRMP98]) *Let \mathcal{F} be a member of $MS_X(n, 0)$. Then (i) \mathcal{F} is Φ -WIT₁.*

(ii) for any $s \in S$ with \mathcal{F}_s as in 5.1, the sheaf $\Phi^1(\mathcal{F}_s)$ is a skyscraper sheaf $\bigoplus_j \mathcal{C}_j$ with $\text{Supp}(\mathcal{C}_j) = \{-y_j\}$, (the point of the dual bundle $\mathcal{O}_E(-y_j + p_0) \simeq \mathcal{O}_E([-y_j] - p_0)$, $[-y_j]$ denoting the divisor of $-y_j \in E$) and $\text{length}(\mathcal{C}_j) = m_j$.

Proof. The first part follows from [BBRMP98] Prop. 2.7 and Cor. 2.12. The second part follows by direct computation and the base change property Proposition 1. \square

Remark 3. *The proof shows that the sheaves $\Phi^1(\mathcal{F}_s)$ and $\Phi^1(\text{gr}(\mathcal{F}_s))$ are the same.*

Remark 4. *The condition for \mathcal{F}_s to be Φ_s -WIT₁ is even equivalent for \mathcal{F}_s to be of degree 0 and semi-stable, see [RMP02].*

For a Φ -WIT₁-sheaf \mathcal{F} on X , we define the **spectral cover** of \mathcal{F} as follows.

Definition 4. *Let \mathcal{F} be a WIT₁ sheaf on X . The spectral cover $C(\mathcal{F})$ of \mathcal{F} is the 0-th Fitting subscheme of J given by the Fitting ideal sheaf $\text{Fitt}_0(\Phi^1(\mathcal{F}))$ of $\Phi^1(\mathcal{F})$.*

Because we work over a non-algebraic manifold and because the image of Φ is not in the derived category of coherent sheaves, but in that of *twisted* sheaves, we need to prove that the Fitting scheme is well-defined in our case. But this follows from the fact that the Fitting ideals are independent of the finite presentation of the local sheaves \mathcal{F}_i of an α -sheaf, see [Eis95, 20.4]. Thus we have well-defined sheaves of ideals $\text{Fitt}_l(\mathcal{F})$ given locally by the ideal sheaves $I_{p_i-l}(\mathcal{F}_i)$ of minors of size $p_i - l$ of the matrix F_i of a local presentation $\mathcal{O}_j^{q_i} \xrightarrow{F_i} \mathcal{O}_j^{p_i} \rightarrow \mathcal{F}_i \rightarrow 0$ over the open set U_i . This sheaf gives us an analytic subspace $V(\text{Fitt}_l(\mathcal{F}))$ called the l -th Fitting scheme by abuse of notation in the analytic category.

By Proposition 2 (ii), for a single fiber X_s , we are given a map $M_{X_s}(n, 0) \rightarrow \text{Sym}^n J_s$ from the moduli space of semistable vector bundles of rank n and degree 0 to the n -th symmetric power of the torus $J_s = \{s\} \times E$, defined by

$$\mathcal{F}_s \mapsto \Sigma_j m_j(s, -y_j).$$

In this way we obtain a map from $M_X(n, 0)$ to $S \times \text{Sym}^n E$, where $\text{Sym}^n E := E^n / \mathfrak{S}_n$ is the n -th symmetric power of E as the quotient of E^n by the symmetric group \mathfrak{S}_n . Then $S \times \text{Sym}^n E$ is a complex manifold of dimension $n + 2$ and can be thought of as the relative space of cycles of degree n in E . We will show that this map is part of a transformation of functors with target $\text{Hom}_S(-, S \times \text{Sym}^n E)$ and that $S \times \text{Sym}^n E$ corepresents the moduli functor $\mathcal{M}_X(n, 0)$ for $M_X(n, 0)$ defined as follows.

For any complex space T over S let the set $\mathcal{M}_X(n, 0)(T)$ be defined by

$$\mathcal{M}_X(n, 0)(T) := \mathcal{MS}_X(n, 0)(T) / \sim,$$

where $\mathcal{MS}_X(n, 0)(T)$ is the set of vector bundles on X_T of rank n and fibre degree 0, and where the equivalence relation $\mathcal{F} \sim \mathcal{G}$ is defined by S -equivalence of the restricted sheaves \mathcal{F}_t and \mathcal{G}_t on the fibres X_{Tt} . The functor property is then defined via pull backs.

We are going to describe the spectral cover as a functor below. For that let $T \rightarrow S$ be a complex space over S and let Φ_T be the Fourier-Mukai transform for the product $J_T \times X_T$ with the pull back \mathcal{U}_T of \mathcal{U} as kernel. By [BBRMP98], Prop. 2.7 and Cor. 2.12, any bundle \mathcal{F}_T in $\mathcal{MS}_X(n, 0)(T)$ is also Φ_T -WIT₁ and admits a spectral cover $C(\mathcal{F}_T) \subset T \times E$ defined by the Fitting ideal $\text{Fitt}_0 \Phi_T^1(\mathcal{F}_T)$.

Lemma 3. *If T is reduced, then $C(\mathcal{F}_T)$ is flat over T .*

Proof. The fibres of $C(\mathcal{F}_T)$ are finite of constant length n as in the case of S above. Because the projection to T is surjective, flatness follows from Douady's criterion in [Dou68]. \square

Lemma 4. *The spectral cover is compatible with base change: For any morphism $h : T' \rightarrow T$ over S and any bundle \mathcal{F}_T in $\mathcal{MS}_X(n, 0)(T)$,*

$$h^* C(\mathcal{F}_T) \simeq C(h^* \mathcal{F}_T)$$

Proof. Because the fibres of the morphisms $p_J : J_T \times X_T \rightarrow J_T$ are 1-dimensional and the sheaves \mathcal{F}_T are locally free, base change holds for $R^1 p_{J*}$, see [BBRMP98] Prop.2.6. When the induced map $J_{T'} \rightarrow J_T$ is denoted by h_J , then

$$h_J^* \Phi_T^1(\mathcal{F}_T) \simeq \Phi_{T'}^1(h^* \mathcal{F}_T)$$

for any $\mathcal{F}_T \in \mathcal{MS}_X(n, 0)(T)$. Since the Fitting ideals are also compatible with base change, the claim follows. \square

The spectral covers $C(\mathcal{F}_T)$ lead us to consider the relative Douady functors

$$\mathcal{D}^n : (An/S)^{op} \rightarrow (\text{Sets}),$$

where (An/S) denotes the category of complex analytic spaces over S and where a set $\mathcal{D}^n(T)$ for a morphism $T \rightarrow S$ is defined as the set of analytic subspaces $Z \subset T \times E$ which are flat over T and have 0-dimensional fibres of constant length n . The Douady functor \mathcal{D}^n is represented by a complex space $D^n(S \times E/S)$ over S , see [Pou69]. For a point $s \in S$, $\mathcal{D}^n(\{s\})$ is the set of 0-dimensional subspaces of length n and can be identified with the symmetric product $\text{Sym}^n(E)$ because

it is well known that the Hilbert-Chow morphism, in our case the Douady-Barlet morphism, $\mathcal{D}^n(\{s\}) \rightarrow \{s\} \times \text{Sym}^n(E)$ is an isomorphism for the smooth curve E , see [Bar75] Ch.V. It is then easy to show that also the relative Douady-Barlet morphism $\mathcal{D}^n(S \times E/S) \rightarrow S \times \text{Sym}^n(E)$ is an isomorphism. This implies that for any complex space T over S there is bijection

$$(5.2) \quad \mathcal{D}^n(T) \xrightarrow{\sim} \text{Hom}_S(T, S \times \text{Sym}^n(E)).$$

One should note here that the behavior of families of cycles is more difficult to describe than of those for the Douady space.

Let now \mathcal{D}_r^n resp. $\mathcal{M}_X(n, 0)_r$ be the restriction of the functors \mathcal{D}^n and $\mathcal{M}_X(n, 0)$ to the category (Anr/S) of reduced complex analytic spaces. By the Lemmas 3 and 4 the spectral covers give rise to a transformation of functors

$$(5.3) \quad \mathcal{M}_X(n, 0)_r \xrightarrow{\gamma} \mathcal{D}_r^n \simeq \text{Hom}_S(-, S \times \text{Sym}^n(E)),$$

where for a reduced space T over S and for a class $[\mathcal{F}_T]$ in $\mathcal{M}_X(n, 0)(T)$ we have $\gamma(T)([\mathcal{F}_T]) = C(\mathcal{F}_T)$. Note that by flatness $C(\mathcal{F}_T)$ depends only on the equivalence class of \mathcal{F}_T . We are now able to present the main theorem which generalises theorem 2.

Theorem 5. *Let $X \rightarrow S$ be an elliptic principal bundle over a compact complex manifold S of arbitrary dimension with invariant $\delta \neq 0$. Then the spectral cover induces a transformation of functors*

$\gamma : \mathcal{M}_X(n, 0)_r \rightarrow \text{Hom}_S(-, S \times \text{Sym}^n(E))$ *with the following properties.*

(i) *The functor $\mathcal{M}_X(n, 0)_r$ is corepresented by $S \times \text{Sym}^n(E)$ via the transformation γ ,*

(ii) *For any point $s \in S$ the induced map $M_{X_s}(n, 0) \rightarrow \text{Sym}^n(E)$ is bijective.*

(iii) *The map $\gamma(T)$ is injective for any reduced complex space T over S .*

(iv) *$\mathcal{M}_X(n, 0)_r$ is locally representable by $S \times \text{Sym}^n(E)$, i.e. if $U \subset S$ is a trivializing open subset for X and T is a complex space over U , then $\gamma(T)$ is bijective.*

Proof. Property (ii) is clear by the construction of the functors.

We begin proving the injectivity in (iii). Let $[\mathcal{F}_1], [\mathcal{F}_2] \in \mathcal{M}_X(n, 0)(T)$ such that $\gamma(T)([\mathcal{F}_1]) = \gamma(T)([\mathcal{F}_2])$. This implies that for every $t \in T$ the spectral covers of $(\mathcal{F}_1)_t$ and $(\mathcal{F}_2)_t$ are the same. If $(\mathcal{F}_1)_t$ is S-equivalent to $\bigoplus_j \mathcal{O}_E(y_j - p_0)^{\oplus m_j}$, then $C((\mathcal{F}_1)_t) = \sum_j m_j(-y_j)$ and visa versa by Atiyah's classification. Hence $(\mathcal{F}_1)_t$ and $(\mathcal{F}_2)_t$ are S-equivalent. But this is precisely the equivalence relation for the classes $[\mathcal{F}_1]$ and $[\mathcal{F}_2]$.

To prove (iv), let $U \subset S$ be an open subset over which X is trivial and let $T \rightarrow U$ be a reduced complex space over U . Then we can assume that $X_T = T \times E$. First we define a map

$$\text{Hom}_S(T, S \times E^n) \xrightarrow{b(T)} \mathcal{M}_X(n, 0)(T)$$

as follows. Given a morphism $(p, f) : T \rightarrow U \times E^n$ over S , let $f_\nu : T \rightarrow E$ be the ν -th component of f . Let then

$$\mathcal{L}_\nu := (f_\nu \times id)^* \mathcal{O}_{E \times E}(-\Delta) \otimes p_2^* \mathcal{O}_E(p_0)$$

on $T \times E$ be the pull back of the dual Poincare bundle. Then the spectral cover of $\mathcal{L}_{\nu, t}$ for any point $t \in T$ consists of the point $f_\nu(t) \in E$. The map $b(T)$ can now be defined by $(p, f) \mapsto [\mathcal{L}_1 \oplus \dots \oplus \mathcal{L}_n]$. This map is obviously \mathfrak{S}_n -equivariant and

thus can be factorized through $\text{Hom}_S(T, S \times \text{Sym}^n(E))$, giving a map

$$\text{Hom}_S(T, S \times \text{Sym}^n(E)) \xrightarrow{\beta(T)} \mathcal{M}_X(n, 0)(T).$$

By construction, $\beta(T)$ is an inverse of $\gamma(T)$.

The proof of (i) is now analogous to that of (i) for Theorem 2, using (iv). \square

APPENDIX A. INVARIANTS OF TORUS BUNDLES

Let M be an n -dimensional compact complex manifold, $T = V/\Lambda$ an m -dimensional complex torus and $X \xrightarrow{\pi} M$ a principal bundle with fiber T . The theory of principal torus bundles is developed in great detail in [Höf93]; see also [BU96]. It is well known that such bundles are described by elements of $H^1(M, \mathcal{O}_M(T))$, where $\mathcal{O}_M(T)$ denotes the sheaf of local holomorphic maps from M to T . Considering the exact sequence of groups

$$0 \rightarrow \Lambda \rightarrow V \rightarrow T \rightarrow 0$$

and taking local sections we obtain the following exact sequence

$$0 \rightarrow \Lambda \rightarrow \mathcal{O}_M \otimes V \rightarrow \mathcal{O}_M(T) \rightarrow 0.$$

Passing to the cohomology we have the long exact sequence

$$\begin{aligned} \cdots \rightarrow H^1(M, \Lambda) \rightarrow H_M^{0,1} \otimes V \rightarrow H^1(M, \mathcal{O}_M(T)) \xrightarrow{c^Z} \\ \xrightarrow{c^Z} H^2(M, \Lambda) \rightarrow H_M^{0,2} \otimes V \rightarrow \cdots \end{aligned}$$

By taking the image of the co-cycle defining the bundle via the map c^Z we obtain a characteristic class $c^Z(X) \in H^2(M, \Lambda) = H^2(M, \mathbb{Z}) \otimes \Lambda$ and also a characteristic class $c(X) \in H^2(M, \mathbb{C}) \otimes V$.

Concerning some important sheaves on X we have (see [Höf93]):

$$(A.1) \quad \mathcal{K}_X = \pi^* \mathcal{K}_M, \quad R^i \pi_* \mathcal{O}_X = \mathcal{O}_M \otimes_{\mathbb{C}} H^{0,i}(T)$$

and the exact sequence

$$(A.2) \quad 0 \rightarrow \Omega_M^1 \rightarrow \pi_* \Omega_X^1 \rightarrow \mathcal{O}_M \otimes_{\mathbb{C}} H^{1,0}(T) \rightarrow 0.$$

All the informations concerning the topology of the bundle $X \rightarrow M$ are given by the following invariants

- a) The exact sequence (A.2) gives rise to an element $\gamma \in \text{Ext}^1(\mathcal{O}_M \otimes H^{1,0}(T), \Omega_M^1) = H^1(\Omega_M^1) \otimes H^{1,0}(T)^*$. Thus γ is a map $H^{1,0}(T) \rightarrow H^{1,1}(M)$.
- b) The first non-trivial d_2 -differential in the Leray spectral sequence ($d_2 : E_2^{0,1} \rightarrow E_2^{2,0}$) of the sheaf \mathbb{C}_X . We obtain in this way a map $\delta : H^1(T, \mathbb{C}) \rightarrow H^2(M, \mathbb{C})$. In the same way we may define the maps $\delta^{\mathbb{Z}} : H^1(T, \mathbb{Z}) \rightarrow H^2(M, \mathbb{Z})$.
- c) The first non-trivial d_2 -differential in the Leray spectral sequence of \mathcal{O}_X , where $d_2 : H^0(R^1 \pi_* \mathcal{O}_X) \rightarrow H^2(\pi_* \mathcal{O}_X)$. Via the identifications (A.1) we get a map $\epsilon : H^{0,1}(T) \rightarrow H^{0,2}(M)$.
- d) The characteristic classes $c^Z(X)$ and $c(X)$ defined above.

These invariants are related by the following theorem of Höfer:

Theorem 6. *Let $X \xrightarrow{\pi} M$ be a holomorphic principal T -bundle. Then:*

- (1) *The Borel spectral sequence ([Hir66, Appendix Two by A.Borel]) ${}^{p,q}E_2^{s,t} = \sum H^{i,s-i}(M) \otimes H^{p-i,t-p+i}(T)$ degenerates on E_3 - level and the d_2 -differential is given by ϵ and γ .*
- (2) *The Leray spectral sequence $E_2^{s,t} = H^s(M, \mathbb{C}) \otimes H^t(T, \mathbb{C})$ degenerates on E_3 - level and the d_2 - differential is given by δ .*
- (3) *Via the identification $H^1(T, \mathbb{Z}) = \text{Hom}(\Lambda, \mathbb{Z})$ the characteristic class $c^{\mathbb{Z}}$ and the map $\delta^{\mathbb{Z}}$ coincide.*
- (4) *δ is determined by $\delta^{\mathbb{Z}}$ via scalar extension.*
- (5) *If $H^2(M)$ has Hodge decomposition then δ determines ϵ and γ and conversely.*

In order to compute the Dolbeault cohomology of X we need to use the Borel spectral sequence because the direct-image sheaves $R^j \pi_* \Omega_X^p$ are non-trivial for $p > 0$ and the Leray spectral sequence is more difficult to use.

APPENDIX B. INVARIANTS OF ELLIPTIC PRINCIPAL BUNDLES OVER SURFACES

In what follows we shall consider fiber bundles with basis M a smooth complex surface and fiber T an elliptic curve. In this case a more detailed description is possible. Let $(1, \tau)$ be a basis of Λ , let $(dt, d\bar{t})$ be a basis of $H^1(T, \mathbb{C})$ given by the decomposition $H^1(T, \mathbb{C}) = H^{1,0}(T) \oplus H^{0,1}(T)$ (another basis on $H^1(T, \mathbb{C})$ is given by the canonical coordinates (dx_1, dx_2)). Assume that $c^{\mathbb{Z}} = a \otimes 1 + b \otimes \tau$, then $c = (a + b \cdot \tau) \otimes 1 = \eta \otimes 1$, with $\eta^{0,2} = 0$. Then we have

$$\begin{aligned} \delta : dt &\mapsto a + \tau \cdot b = \eta \\ d\bar{t} &\mapsto a + \bar{\tau} \cdot b = \bar{\eta} \\ \epsilon : d\bar{t} &\mapsto (a + \bar{\tau} \cdot b)^{0,2} = \bar{\eta}^{0,2} \\ \gamma : dt &\mapsto (a + \tau \cdot b)^{1,1} = \eta^{1,1}. \end{aligned}$$

The only non-zero terms in the Borel spectral sequence are

$$(B.1) \quad \begin{array}{ccc} & H^{p-1,q-2}(M) \otimes H^{1,1}(T) & \\ & \swarrow \gamma \quad \searrow -\epsilon & \\ H^{p,q-1}(M) \otimes H^{0,1}(T) & \oplus & H^{p-1,q}(M) \otimes H^{1,0}(T) \\ & \swarrow \epsilon \quad \searrow \gamma & \\ & H^{p,q+1}(M) \otimes H^{0,0}(T) & \end{array}$$

From now on we shall be concerned with the case when M has trivial canonical bundle. By (A.1) this implies that X also has trivial canonical bundle. The case when M is Kähler was considered by Höfer. This leaves us with the case when M is a primary Kodaira surface. We shall use the preceding diagram to compute the Hodge numbers for X in this case. Recall that the Hodge diamond for a primary Kodaira together with Betti numbers is, see [BPVdV84, V.5].

$$\begin{array}{cccc} & & 1 & & 1 \\ & & 2 & 1 & 3 \\ 1 & & 2 & 1 & 4 \\ & & 1 & 2 & 3 \\ & & 1 & & 1 \end{array}$$

where $h := \text{Rank}(H^{0,1}(M) \otimes H^{1,0}(T) \xrightarrow{\gamma^{(dt)\wedge}} H^{1,2}(M)$ and
 $f := \text{Rank}(H^{1,0}(M) \otimes H^{0,1}(T) \xrightarrow{\epsilon^{(dt)\wedge}} H^{1,2}$. When M is a $K3$ surface we have

$$(B.5) \quad \begin{array}{cccccc} & & & 1 & & 1 \\ & & & 1-g & & 1-e & & 0 \\ & & 1 & & 20-g & & 1-e & 20 \\ & 1 & & 20 & & 20 & & 1 & 42 \\ & & 1-e & & 20-g & & 1 & & 20 \\ & & & 1-e & & 1-g & & & 0 \\ & & & & 1 & & & & 1. \end{array}$$

REFERENCES

- [Add09a] N. Addington. Spinor Sheaves on Singular Quadrics. *ArXiv e-prints*, 2009.
- [Add09b] N. Addington. The Derived Category of the Intersection of Four Quadrics. *ArXiv e-prints*, 2009.
- [Ati57] M. F. Atiyah. Vector bundles over an elliptic curve. *Proc. London Math. Soc.* (3), 7:414–452, 1957. ISSN 0024-6115.
- [Bar75] Daniel Barlet. Espace analytique réduit des cycles analytiques complexes compacts d’un espace analytique complexe de dimension finie. In *Fonctions de plusieurs variables complexes, II (Sém. François Norguet, 1974–1975)*, pages 1–158. Lecture Notes in Math., Vol. 482. Springer, Berlin, 1975.
- [BB09] Oren Ben-Bassat. Twisting derived equivalences. *Trans. Amer. Math. Soc.*, 361(10):5469–5504, 2009. CODEN TAMTAM. ISSN 0002-9947. URL <http://dx.doi.org/10.1090/S0002-9947-09-04832-6>.
- [BBDG03] Katrin Becker, Melanie Becker, Keshav Dasgupta, and Paul S. Green. Compactifications of heterotic theory on non-Kähler complex manifolds. I. *J. High Energy Phys.*, (4):007, 60 pp. (electronic), 2003. ISSN 1126-6708. URL <http://dx.doi.org/10.1088/1126-6708/2003/04/007>.
- [BBHR09] Claudio Bartocci, Ugo Bruzzo, and Daniel Hernández Ruipérez. *Fourier-Mukai and Nahm transforms in geometry and mathematical physics*, volume 276 of *Progress in Mathematics*. Birkhäuser Boston Inc., Boston, MA, 2009. ISBN 978-0-8176-3246-5. xvi+423 pp.
- [BBRMP98] Claudio Bartocci, Ugo Bruzzo, Daniel Hernández Ruipérez, and José M. Muñoz Porras. Mirror symmetry on $K3$ surfaces via Fourier-Mukai transform. *Comm. Math. Phys.*, 195(1):79–93, 1998. CODEN CMPHAY. ISSN 0010-3616. URL <http://dx.doi.org/10.1007/s002200050380>.
- [BK06] Igor Burban and Bernd Kreuzler. On a relative Fourier-Mukai transform on genus one fibrations. *Manuscripta Math.*, 120(3):283–306, 2006. CODEN MSMHB2. ISSN 0025-2611. URL <http://dx.doi.org/10.1007/s00229-006-0013-y>.
- [BL04] Christina Birkenhake and Herbert Lange. *Complex abelian varieties*, volume 302 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2004. ISBN 3-540-20488-1. xii+635 pp.
- [BM02] Tom Bridgeland and Antony Maciocia. Fourier-Mukai transforms for $K3$ and elliptic fibrations. *J. Algebraic Geom.*, 11(4):629–657, 2002. ISSN 1056-3911. URL <http://dx.doi.org/10.1090/S1056-3911-02-00317-X>.
- [BM05] Vasile Brînzănescu and Ruxandra Moraru. Stable bundles on non-Kähler elliptic surfaces. *Comm. Math. Phys.*, 254(3):565–580, 2005. CODEN CMPHAY. ISSN 0010-3616. URL <http://dx.doi.org/10.1007/s00220-004-1269-2>.
- [BM06] Vasile Brînzănescu and Ruxandra Moraru. Twisted Fourier-Mukai transforms and bundles on non-Kähler elliptic surfaces. *Math. Res. Lett.*, 13(4):501–514, 2006. ISSN 1073-2780.
- [BO95] A. Bondal and D. Orlov. Semiorthogonal decomposition for algebraic varieties. In *eprint arXiv:alg-geom/9506012*, 1995.

- [BPVdV84] W. Barth, C. Peters, and A. Van de Ven. *Compact complex surfaces*, volume 4 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1984. ISBN 3-540-12172-2. x+304 pp.
- [Bri98] Tom Bridgeland. Fourier-Mukai transforms for elliptic surfaces. *J. Reine Angew. Math.*, 498:115–133, 1998. CODEN JRMAA8. ISSN 0075-4102. URL <http://dx.doi.org/10.1515/crll.1998.046>.
- [Bri99] Tom Bridgeland. Equivalences of triangulated categories and Fourier-Mukai transforms. *Bull. London Math. Soc.*, 31(1):25–34, 1999. CODEN LMSBBT. ISSN 0024-6093. URL <http://dx.doi.org/10.1112/S0024609398004998>.
- [BU96] Vasile Brînzănescu and Kenji Ueno. Néron-Severi group for torus quasi bundles over curves. In *Moduli of vector bundles (Sanda, 1994; Kyoto, 1994)*, volume 179 of *Lecture Notes in Pure and Appl. Math.*, pages 11–32. Dekker, New York, 1996.
- [Cal00] Andrei Horia Caldararu. *Derived categories of twisted sheaves on Calabi-Yau manifolds*. ProQuest LLC, Ann Arbor, MI, 2000. ISBN 978-0599-72206-4. 196 pp. URL http://gateway.proquest.com/openurl?url_ver=Z39.88-2004&rft_val_fmt=info:ofi/fmt:kev:mtx:dissertation&res_dat=xri:pqdiss&rft_dat=xri:pqdiss:9967459. Thesis (Ph.D.)—Cornell University.
- [Căl02] Andrei Căldăraru. Derived categories of twisted sheaves on elliptic threefolds. *J. Reine Angew. Math.*, 544:161–179, 2002. CODEN JRMAA8. ISSN 0075-4102. URL <http://dx.doi.org/10.1515/crll.2002.022>.
- [CCD⁺03] G. L. Cardoso, G. Curio, G. Dall’Agata, D. Lüst, P. Manousselis, and G. Zoupanos. Non-Kähler string backgrounds and their five torsion classes. *Nuclear Phys. B*, 652(1-3):5–34, 2003. CODEN NUPBBO. ISSN 0550-3213. URL [http://dx.doi.org/10.1016/S0550-3213\(03\)00049-X](http://dx.doi.org/10.1016/S0550-3213(03)00049-X).
- [CDH⁺10] Andrei Căldăraru, Jacques Distler, Simeon Hellerman, Tony Pantev, and Eric Sharpe. Non-birational twisted derived equivalences in abelian GLSMs. *Comm. Math. Phys.*, 294(3):605–645, 2010. CODEN CMPHAY. ISSN 0010-3616. URL <http://dx.doi.org/10.1007/s00220-009-0974-2>.
- [Del68] P. Deligne. Théorème de Lefschetz et critères de dégénérescence de suites spectrales. *Inst. Hautes Études Sci. Publ. Math.*, (35):259–278, 1968. ISSN 0073-8301.
- [Don95] Ron Donagi. Spectral covers. In *Current topics in complex algebraic geometry (Berkeley, CA, 1992/93)*, volume 28 of *Math. Sci. Res. Inst. Publ.*, pages 65–86. Cambridge Univ. Press, Cambridge, 1995.
- [Don97] Ron Y. Donagi. Principal bundles on elliptic fibrations. *Asian J. Math.*, 1(2):214–223, 1997. ISSN 1093-6106.
- [Dou68] A. Douady. Flatness and privilege. *Enseignement Math. (2)*, 14:47–74, 1968. ISSN 0013-8584.
- [DP08] Ron Donagi and Tony Pantev. Torus fibrations, gerbes, and duality. *Mem. Amer. Math. Soc.*, 193(901):vi+90, 2008. CODEN MAMCAU. ISSN 0065-9266. With an appendix by Dmitry Arinkin.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. ISBN 0-387-94268-8; 0-387-94269-6. xvi+785 pp. With a view toward algebraic geometry.
- [FMW99] Robert Friedman, John W. Morgan, and Edward Witten. Vector bundles over elliptic fibrations. *J. Algebraic Geom.*, 8(2):279–401, 1999. ISSN 1056-3911.
- [Fri89] Robert Friedman. Rank two vector bundles over regular elliptic surfaces. *Invent. Math.*, 96(2):283–332, 1989. CODEN INVMBH. ISSN 0020-9910. URL <http://dx.doi.org/10.1007/BF01393965>.
- [GP04] Edward Goldstein and Sergey Prokushkin. Geometric model for complex non-Kähler manifolds with SU(3) structure. *Comm. Math. Phys.*, 251(1):65–78, 2004. CODEN CMPHAY. ISSN 0010-3616. URL <http://dx.doi.org/10.1007/s00220-004-1167-7>.
- [Har77] Robin Hartshorne. *Algebraic geometry*. Springer-Verlag, New York, 1977. ISBN 0-387-90244-9. xvi+496 pp. Graduate Texts in Mathematics, No. 52.
- [Hir66] F. Hirzebruch. *Topological methods in algebraic geometry*. Third enlarged edition. New appendix and translation from the second German edition by R. L. E. Schwarzenberger, with an additional section by A. Borel. Die Grundlehren der Mathematischen Wissenschaften, Band 131. Springer-Verlag New York, Inc., New York, 1966. x+232 pp.

- [HL97] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Aspects of Mathematics, E31. Friedr. Vieweg & Sohn, Braunschweig, 1997. ISBN 3-528-06907-4. xiv+269 pp.
- [Höf93] Thomas Höfer. Remarks on torus principal bundles. *J. Math. Kyoto Univ.*, 33(1): 227–259, 1993. CODEN JMKYAZ. ISSN 0023-608X.
- [Huy06] D. Huybrechts. *Fourier-Mukai transforms in algebraic geometry*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, Oxford, 2006. ISBN 978-0-19-929686-6; 0-19-929686-3. viii+307 pp. URL <http://dx.doi.org/10.1093/acprof:oso/9780199296866.001.0001>.
- [KO03] Anton Kapustin and Dmitri Orlov. Vertex algebras, mirror symmetry, and D-branes: the case of complex tori. *Comm. Math. Phys.*, 233(1):79–136, 2003. CODEN CM-PHAY. ISSN 0010-3616. URL <http://dx.doi.org/10.1007/s00220-002-0755-7>.
- [Kuz08] Alexander Kuznetsov. Derived categories of quadric fibrations and intersections of quadrics. *Adv. Math.*, 218(5):1340–1369, 2008. CODEN ADMTA4. ISSN 0001-8708. URL <http://dx.doi.org/10.1016/j.aim.2008.03.007>.
- [Muk81] Shigeru Mukai. Duality between $D(X)$ and $D(\hat{X})$ with its application to Picard sheaves. *Nagoya Math. J.*, 81:153–175, 1981. CODEN NGMJJA2. ISSN 0027-7630. URL <http://projecteuclid.org/getRecord?id=euclid.nmj/1118786312>.
- [Mum70] David Mumford. *Abelian varieties*. Tata Institute of Fundamental Research Studies in Mathematics, No. 5. Published for the Tata Institute of Fundamental Research, Bombay, 1970. viii+242 pp.
- [Ori03] D. O. Orlov. Derived categories of coherent sheaves and equivalences between them. *Uspekhi Mat. Nauk*, 58(3(351)):89–172, 2003. ISSN 0042-1316. URL <http://dx.doi.org/10.1070/RM2003v058n03ABEH000629>.
- [Pou69] Geneviève Pourcin. Théorème de Douady au-dessus de S . *Ann. Scuola Norm. Sup. Pisa (3)*, 23:451–459, 1969.
- [RMP02] D. Hernández Ruipérez and J. M. Muñoz Porras. Stable sheaves on elliptic fibrations. *J. Geom. Phys.*, 43(2-3):163–183, 2002. CODEN JGPHE5. ISSN 0393-0440. URL [http://dx.doi.org/10.1016/S0393-0440\(02\)00019-0](http://dx.doi.org/10.1016/S0393-0440(02)00019-0).
- [Tu93] Loring W. Tu. Semistable bundles over an elliptic curve. *Adv. Math.*, 98(1):1–26, 1993. CODEN ADMTA4. ISSN 0001-8708. URL <http://dx.doi.org/10.1006/aima.1993.1011>.

”SIMION STOILOW” INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O.BOX 1-764,
014700 BUCHAREST, ROMANIA

E-mail address: Vasile.Brinzanescu@imar.ro

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE,, UNIVERSITY OF BUCHAREST
STR. ACADEMIEI 14
010014 BUCHAREST,ROMANIA

E-mail address: halanay@gt.math.unibuc.ro

UNIVERSITÄT KAISERSLAUTERN, FACHBEREICH MATHEMATIK,
ERWIN-SCHRÖDINGER-STRASSE
D-67663 KAISERSLAUTERN

E-mail address: trm@mathematik.uni-kl.de