

The geometry of variations in Batalin–Vilkovisky formalism

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2013 J. Phys.: Conf. Ser. 474 012024

(<http://iopscience.iop.org/1742-6596/474/1/012024>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 192.68.254.205

This content was downloaded on 28/04/2014 at 10:00

Please note that [terms and conditions apply](#).

The geometry of variations in Batalin–Vilkovisky formalism

Arthemy V. Kiselev

Johann Bernoulli Institute for Mathematics and Computer Science, University of Groningen,
P.O. Box 407, 9700 AK Groningen, The Netherlands

E-mail: A.V.Kiselev@rug.nl

Abstract. We explain why no sources of divergence are built into the Batalin–Vilkovisky (BV) Laplacian, whence there is no need to postulate any *ad hoc* conventions such as “ $\delta(0) = 0$ ” and “ $\log \delta(0) = 0$ ” within BV-approach to quantisation of gauge systems. Remarkably, the geometry of iterated variations does not refer at all to the construction of Dirac’s δ -function as a limit of smooth kernels. We illustrate the reasoning by re-deriving –but not just ‘formally postulating’– the standard properties of BV-Laplacian and Schouten bracket and by verifying their basic inter-relations (e.g., cohomology preservation by gauge symmetries of the quantum master-equation).

Introduction

This is a paper about geometry of variations. We formulate definitions of the objects and structures which are cornerstones of Batalin–Vilkovisky formalism [5, 7, 20, 22, 55]. To confirm the intrinsic self-regularisation of BV-Laplacian, we explain why there are no divergencies in it (such excessive elements are traditionally encoded by using derivatives of Dirac’s δ -distribution). Namely, we specify the geometry in which the following canonical inter-relations between the variational Schouten bracket $[[,]]$ and BV-Laplacian Δ are rigorously proven for any BV-functionals F, G, H :

$$[[F, G \cdot H]] = [[F, G]] \cdot H + (-)^{(|F|-1) \cdot |G|} G \cdot [[F, H]], \quad (1a)$$

$$\Delta(F \cdot G) = \Delta F \cdot G + (-)^{|F|} [[F, G]] + (-)^{|F|} F \cdot \Delta G, \quad (1b)$$

$$\Delta([[F, G]]) = [[\Delta F, G]] + (-)^{|F|-1} [[F, \Delta G]], \quad (1c)$$

$$\Delta^2 = 0 \quad \iff \quad \text{Jacobi}([[,]]) = 0. \quad (1d)$$

There is an immense literature on this subject’s intrinsic difficulties and attempts of regularisation of apparent divergencies in it (e.g., see [12, 13, 25, 50, 51] vs [21]). While the BV-quantisation technique has advanced far from its sources [7, 8], it is still admitted that it lacks sound mathematical consistency ([22, §15] or [3, §3]). The calculus in this field is thus reduced to formal operation with expressions which are expected to render the theory’s main objects and structures. Several *ad hoc* techniques for cancellation of divergencies, allowing one to strike through calculations and obtain meaningful results, are adopted by repetition; we briefly review the plurality of such tricks in what follows.



Our reasoning is independent from such conventional schemes for cancellation of infinities or from other practised roundabouts for regularisation of terms which are believed to be infinite (e.g., by erasing ‘infinite constants’ [11]). In particular, we do not pronounce the traditional password

$$\delta(0) := 0 \tag{2}$$

which lets one enter the existing paradigm and use its quantum alchemy for operation with what remains from Dirac’s δ -distribution.¹ Our message is this: we do not propose to replace ‘bad slogans’ with ‘good slogans,’ which would mean that a choice of conventions is still left to the one who attempts regularisation in the BV-setup. Such deficiency would symptomise that the theory remains a formal procedure. We now focus on the true sources of known difficulties. By analysing the geometry of variations of functionals at a very basic level, we prove the absence of apparently divergent essences. The intrinsically regularised definitions of the BV-Laplacian Δ and Schouten bracket $[\cdot, \cdot]$ are the main result of this paper.

The new understanding leaves intact but substantiates the bulk of results which have been obtained by using various *ad hoc* techniques (that is, explicitly or tacitly referring to the surreal equalities $\delta(0) = 0$ and $\log \delta(0) = 0$); we refer to a detailed review [3] for an account of early developments in BV-formalism. We do not aim at a reformulation or reproduction of any old or recent achievements, accomplishing here a different task.

In fact, we invent nothing new. It is the coupling of dual vector spaces which ensures the intrinsic self-regularisation of BV-Laplacian and validity of equalities (1), with (1c) in particular. Therefore, it would be redundant to start developing any brand-new formalism (cf. [51]); on the other hand, we *prove* properties (1) and not just *postulate* these assertions (cf. [21]).

We employ standard notions, constructions, and techniques from the geometry of jet spaces [28, 40, 45]. Because the geometry of BV-objects is essentially variational, it would be methodologically incomplete to handle them as if the space-time, that is, the base manifold in the bundles of physical fields, were just a point ([27, 48] or [39]). The language of jet spaces is extensively used in the study of BV-models, see [3, 6, 21, 43]: the bundles of jets of sections usually appear in such traditional contexts as calculation of symmetries or conservation laws. In this paper we apply these geometric techniques at a much more profound level and give rigorous definitions for BV-objects. Let us emphasize that we do not aim at extending one’s ability to write more formulas according to a regularly emended system of accepted algorithms; we explicate the genuine nature of objects and their canonical matchings, not taking any formulas for quasi-definitions.

This paper is structured as follows. Containing a brief overview of traditional approaches to regularisation of the BV-formalism, this introduction concludes with a parable; the line of our reasoning is reminiscent to that of *Lettres persanes* by Montesquieu.

In section 1 we describe the true geometry of variations; we first reveal the correspondence between action functionals and infinitesimal shifts of classical trajectories or physical fields. An understanding of nontrivial mechanism of such matching achieved for *one* variation, the picture of *many* variations becomes clear. This approach resolves the obstructions for regularisation of iterated variations in BV-formalism; we remark that Dirac’s δ -function does not appear in section 2 at all.²

In section 2.1 we recall in proper detail the standard construction of Batalin–Vilkovisky (BV) vector bundles with canonically conjugate pairs of ghost parity-even and odd variables.

¹ Another convention is $\log \delta(0) = 0$; we show that natural counterparts of the true geometry of variations lead to this intuitive convention and simultaneously to (2) — none of the two being actually required.

² We refer to [19] for the theory of distributions. Let us specify that singular linear integral operators which emerge in the course of our reasoning will **not** be approached via parametric families of regular linear integral functionals with piecewise continuous or smooth kernels (in which context the notation “ $\delta(0)$ ” for Dirac’s *function* is used in the literature).

In this specific setup we analyse the construction of two distinct couplings of the BV-fibres' ghost parity-homogeneous vector subspaces with their respective duals. In particular, in section 2.2 we focus on the rule of signs which determines the anti-commutation of differential one-forms in the geometry at hand. Applying the geometric concept of iterated variations in section 2.3, we represent the left- and right variations of functionals in terms of left- or right-directed singular linear integral operators; this framework ensures the intrinsic regularisation of iterated variations. We then formulate in section 2.4 the definitions of BV-Laplacian Δ and variational Schouten bracket $[[,]]$ (or *antibracket*). We show that these definitions are operational, amounting to natural, well-defined reconfigurations of the geometry (but not to any hand-made algorithms for cancellation of divergent terms; for those do not appear at all). Our main result, which is contained in section 2.5, is an explicit proof – that is, starting from basic principles – of relations (1). In other words, we neither postulate a validity of these properties nor elaborate a cunning syllogism the aim of which would be to convince why such assertions should hold provided that one knows when various (derivatives of) Dirac's δ -functions must be erased in the course of so arguable a reasoning.

For consistency, we first apply the above theory to a standard derivation of the quantum master-equation from the Schwinger–Dyson condition that essentially eliminates a dependence on the unphysical, ghost parity-odd dimensions (see section 3.1); we also recall here the construction of quantum BV-differential. The point is that neither divergencies nor *ad hoc* cancellations occur in the entire argument. On the same grounds we address in section 3.2 the quantum BV-cohomology preservation by infinitesimal gauge symmetries of the quantum master-equation. (We refer to [7, 8, 20, 22] and also [1, 37, 51] in this context; several methodological comments, which highlight our concept, are placed in section 3 along the lines of a well-known reasoning.)

The paper concludes with a statement that an intrinsic regularisation in the geometry of iterated variations relies on the principle of locality (which manifests also through causality). We argue that a logical complexity of geometric objects grows while they accumulate the (iterated) variations; a conversion of such composite-structure objects into maps which take physical field configurations to numbers entails a decrease of the complexity via a loss of information. Having motivated this claim in section 2, we prove that the logic of analytic reasonings may not be interrupted; for example, the right-hand side of (1c) is not assembled from the would-be constituent blocks ΔF and ΔG for which it is known in advance how they take field configurations to numbers whenever the functionals F and G are given.

The paper explicitly answers the question what variations are — in particular, what iterated variations are. Moreover, we tacitly describe a geometric mechanism which is responsible for the anti-commutation of differential one-forms; such mechanism ensures that the results of calculations match empiric data even if the exterior algebras of forms are introduced by hand. The roots of this principle are none other than the ordering of dual vector spaces which stem in the course of variations in models of nonlinear phenomena (this picture is addressed in section 2.2).

We illustrate our approach with elementary starting section 1 in which we inspect the matching of geometries –one for an action functional, the other for a field's test shift– in the course of derivation of Euler–Lagrange equation of motion in field theory. The second example on pp. 34–36 clarifies the idea specifically in the BV-setup of (anti)fields and (anti)ghosts. We thus provide a pattern for all types of calculations which involve the Schouten bracket and BV-Laplacian in any model.

Historical context: an overview

There is a class of significant papers in which the BV-formalism is developed under assumption that the space-time is a point. Indeed, such hypothesis is equivalent to an agreement that the only admissible sections of bundles over space-time are constant; this implies that even if their

derivatives are nominally present in some formulas, they are always equal to zero. The calculus of variations then reduces to usual differential geometry on the bundles' fibres. It must be noted that publications containing the above assumption did contribute to the subject and in many cases guided its further development (we recall the respective comment in [51] and refer to [12, 22, 27, 39, 48, 53]). Moreover, the no-derivatives reduction sometimes allows one to jump at conclusions which are correct; an integration by parts over the base manifold M^n is restored –whenever possible– at the end of the day. Still this oversimplification is potentially dangerous because variational calculus of integral functionals conceptually exceeds any classical differential geometry on the fibres (see [33] for discussion and [28, 34]). In the variational setup, the objects and their properties become geometrically different from their analogues on usual manifolds even if the terminology is kept unchanged. Here we recall for example that variational multivectors do not split to wedge products of variational one-vectors and likewise, several Leibniz rules are irreparably lost but this can not be noticed when all derivatives equal zero. In fact, it is the abyss between classical geometry of manifolds and geometry of variations for jet spaces of maps of manifolds which motivated our earlier study [34]. Yet the misconception is still present in active research, e.g., see [4, 27, 39, 44].

The fact of incompleteness of such heuristic analogies from usual geometry of manifolds is signalled in [51]. Paradoxically, it is simultaneously not true that a solution of the regularisation problem for BV-Laplacian has no analogues in the case of ODE dynamics on manifolds. From section 1 below it is readily seen that good old techniques persist in the finite-dimensional ODE geometry at the level of standard linear algebra of dual vector spaces.³

The article [51] is a considerable step towards a solution of the regularisation problem in BV-formalism. A weighted, critical overview of various inconsistencies, *ad hoc* practices, and roundabouts is summed up there. The object of [51] was to formulate a self-contained analytic concept which would make the variational calculus of functionals free from divergencies and infinities. Still it remained unclear from [51] what the generality of underlying geometry is and why such self-consistent formalism should actually exist at the level of objects, i.e., beyond a mere ability to write formulas. In particular, it remained unnoticed that the main motivating example –namely, the canonical BV-setup– itself is the only class of geometries in which the technique is grounded.⁴ A correctness but incompleteness of the approach in [51] means the following in practice. Whenever a theorist refers to the formalism of *loc. cit.*, Nature immediately creates a new, principally inobservable essence –a metric field which is denoted by $E(x_1, \dots, x_n; \Gamma)$ in [51]– on top of the electromagnetic and weak gauge connections, as well as the fields for strong force, gravity, or any other gauge fields Γ . It is perhaps this methodological difficulty which hints us why the approach of [51] is considered “formal” by many experts; that conceptual paper remains scarcely known to a wider community.⁵

³ On the other hand, the variational setup highlights the fundamental concept of a physical field as a system with degrees of freedom attached at every point of the space-time M^n ; we focus on this aspect in what follows.

⁴ The integration of closed algebra of gauge symmetries for the quantum master-equation to a group of transformations of the master-action S^h remains a separate problem, which is also addressed in [51]. Suppose that the standard cohomological obstructions to such integration vanish (see section 3.2 below), whence (i) all infinitesimal transformations of the functional S^h are exact, i.e., they are generated by odd ghost-parity elements F , and also (ii) such transformations can be extended from the master-action S^h to evolution of the observables \mathcal{O} . We remark that, unlike it is claimed in [51], neither of the two groups of functionals' transformations is induced by any well-defined change of BV-coordinates; of course, evolutionary vector fields are well-defined objects in that geometry and one could study them regardless of these functionals' transformations. We shall recall in section 3.2 the standard construction of automorphisms for quantum BV-cohomology groups; it illustrates our concept because the notion of quantum gauge symmetries explicitly refers to all basic properties of the BV-Laplacian and Schouten bracket, see (1) on p. 1.

⁵ An attempt to interpret the formalism of [51] in terms of the language of PDE geometry (particularly, in the context of [41], see also [28, 40, 45]) was performed in [23] and published in abridged form in [24]. The construction of Schouten bracket in [23] relies on the notion of variational cotangent bundle [41] and on classical approach to

To demystify the notion of a “metric field $E(x_1, \dots, x_n; \Gamma)$,” we describe in this paper an elementary geometric mechanism for the long-expected but still intuitively paradoxical analytic behaviour of variations. This mechanism implies that Nature is not obliged to respond to the needs of a theorist and create such multi-entry distributions upon request.

Another line of reasoning, which led to much progress in a revision of BV-structures and regularisation of divergences, was pursued in [12, 13]. We recall that the language of *loc. cit.* is functional analytic so that the theory’s objects are viewed as (Dirac’s) distributions (and heat kernels are implemented). According to [12, §1.8], the BV-Laplacian Δ which is used in physical theories is ill-defined because for a given action S over space-time M^n of positive dimension n the object ΔS involves a multiplication of singular distributions (and thus –a quotation from [12] continues– ΔS has the same kind of singularities as appear in one-loop Feynman diagrams). The regularisation technique proposed in [12, 13] stems from analysis of the distributions’ limit behaviour as one approaches the “physical” structures by using regular ones.

The resolution to apparent difficulties is that there are several distinct geometric constructions which yield the same singular linear operators with support on the diagonal (in what follows we study in detail on which space such operators are defined).

We now discuss a peculiar, well-established domain, the very form of existence of which could be hardly believed in. In that theory, there is a serious lack of rigorous definitions for the most elementary objects; at the same time, there is a rapidly growing number of monumental reviews. Whereas the theory’s difficulties are clearly inherited from a deficit of boring rigour at the initial stage, such hardships are proclaimed the theory’s immanent components. At expert level it is mandatory to have a firm knowledge of the built-in difficulties and readily classify the descriptive objects which those apparent obstructions bring into the mathematical apparatus. (There is no firm guarantee that the (un)necessary objects really exist beyond written formulas.) The way of handling inconveniences largely amounts not to resolving them by a thorough study of their origins but to some *ad hoc* methods for hiding their presence. Doing research is thus substituted by practising a ritual.

However, the community of experts who mature in operation with formulas (a part of which are believed to express something objectively existing) maintains a considerable pluralism about a proper way to mask the symptoms of troubles:

- The radicals declare that undefined objects which seem to make trouble must be set equal to zero.
- The revisionist approach prescribes a postfactum erasing of not the entire objects (which are still undefined) but of undesirable elements in those objects’ description.
- A diplomatic viewpoint is that there might be sources of trouble but their contribution to final results is suppressed as soon as the objects’ desired properties are postulated (regardless of the actual presence or absence of such sources and one’s ability to substantiate those properties).

For an external observer, this state-of-the-art could seem atypical for a consistent theory. Indeed, the reliability of its main pillar is a matter of irrational belief.

1. The geometry of variations

Let us first analyse the basic geometry of variations of functionals; by comprehending the full setup of a one-time variation, we shall understand the geometry of many. Specifically, in this

the theory of variations. On one hand, this ensures the validity of Jacobi identity for the bracket (see the *second* half of Eq. (1d) but *not* the first one). But on the other hand, we have showed by a counterexample in [35, §3] that the old approach fails to relate by (1c) the Schouten bracket to BV-Laplacian. In other words, the BV-Laplacian did not entirely generate the variational Schouten bracket, making only Eq. (1b) but not (1c) possible in that geometry (cf. [39]).

section we reveal the interrelation of bundles in the course of integration by parts; we also explain a rigorous construction of iterated variations.

The core of traditional difficulties in this domain is that a use of only fibre bundles π of physical fields, which are subjected to test shifts, is insufficient. We argue that the tangent bundles $T\pi$ to the bundles π may not be discarded (see Fig. 1). For identities (1) to hold one




Figure 1. The fibre bundle π of fields s and vector bundle $T\pi$ of their variations δs .

must substantiate why higher-order variational derivatives are (graded-)permutable whenever one inspects the response of a given functional to shifts of its argument along several directions. To resolve the difficulties, we properly enlarge the space of functionals and adjust a description of the geometry for the functionals' variations: in fact, each variation brings its own copy of the base M^n into the picture (see Fig. 3 on p. 14).

1.1. Notation

We now fix some notations, in most cases matching that from [28] (for a more detailed exposition of these matters, see for example [28, 40, 45]).

Let $\pi: E \rightarrow M$ be a smooth fibre bundle⁶ with m -dimensional fibres $\pi^{-1}(x)$ over points x of a smooth real oriented manifold M of dimension n ; we assume that all mappings, including those which determine the smoothness class of manifolds, are infinitely smooth.

We let x^i denote local coordinates in a chart $U_\alpha \subseteq M^n$ and u_j be the fibre coordinates. We denote by $[\mathbf{u}]$ a differential dependence of the fibre variables (specifically in the BV-setup, a differential dependence $[\mathbf{q}]$ on physical fields and other ghost parity-even variables, and we denote by $[\mathbf{q}^\dagger]$ that of ghost parity-odd BV-variables).

Remark 1.1. We suppose that the initially given bundle π of physical fields is not graded. In what follows, starting with π , we shall construct new bundles whose fibres are endowed with the \mathbb{Z}_2 -valued ghost parity $\text{gh}(\cdot)$. However, our reasoning remains valid for superbundles $\pi^{(0|1)}$ over supermanifolds $M^{(n_0|n_1)}$ ([10, 52]) and to a noncommutative setup of cyclic-invariant words (see [29, 32] and references therein), cf. Fig. 8 below.

We take the infinite jet space $\pi_\infty: J^\infty(\pi) \rightarrow M$ associated with this bundle [15, 45]; a point from the jet space is then $\theta = (x^i, \mathbf{u}, u_{x^i}^k, u_{x^i x^j}^k, \dots, \mathbf{u}_\sigma, \dots) \in J^\infty(\pi)$, where σ is a multi-index and we put $\mathbf{u}_\emptyset \equiv \mathbf{u}$. If $s \in \Gamma(\pi)$ is a section of π , or a *field*, we denote by $j^\infty(s)$ its infinite jet, which is a section $j^\infty(s) \in \Gamma(\pi_\infty)$. Its value at $x \in M$ is $j_x^\infty(s) = (x^i, s^\alpha(x), \frac{\partial s^\alpha}{\partial x^i}(x), \dots, \frac{\partial^{|\sigma|} s^\alpha}{\partial x^\sigma}(x), \dots) \in J^\infty(\pi)$.

We denote by $\mathcal{F}(\pi)$ the properly understood algebra of finite differential order smooth functions on the infinite jet space $J^\infty(\pi)$, see [28, 40] for details. The space of top-degree

⁶ Vector bundles are primary examples but we do not actually use the linear vector space structure of their fibres so that π could be any smooth fibre bundle.

horizontal forms on $J^\infty(\pi)$ is denoted by $\overline{\Lambda}^n(\pi)$; let us also assume that at every $\mathbf{x} \in M$ a volume element $\text{dvol}(\mathbf{x})$ is specified so that its pull-back under π_∞^* is an n -th degree form in $\overline{\Lambda}^n(\pi)$, cf. Remark 1.5 on p. 9.

The highest horizontal cohomology, i.e., the space of equivalence classes of n -forms from $\overline{\Lambda}^n(\pi)$ modulo the image of the horizontal exterior differential \overline{d} on $J^\infty(\pi)$, is denoted by $\overline{H}^n(\pi)$; the equivalence class of $\omega \in \overline{\Lambda}^n(\pi)$ is denoted by $\int \omega \in \overline{H}^n(\pi)$. We assume that sections $s \in \Gamma(\pi)$ are such that integration of functionals $\Gamma(\pi) \rightarrow \mathbb{k}$ by parts is allowed and does not result in any boundary terms (for example, the base manifold is closed, or the sections all have compact support, or decay sufficiently fast towards infinity, or are periodic).

1.2. Euler–Lagrange equations

A derivation of Euler–Lagrange equations \mathcal{E}_{EL} for a given action functional $S = \int \mathcal{L}(\mathbf{x}, [\mathbf{u}]) \text{dvol}(\mathbf{x})$ is a model example which illustrates the correlation of two geometries:⁷ one for “trajectories” $s \in \Gamma(\pi)$ and the other for shifts δs . It is well known that the functional’s response to a test shift δs of its argument $s \in \Gamma(\pi)$ is described by the formula [2, §12]

$$\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} S(s + \varepsilon \cdot \overleftarrow{\delta s}) = \int_M \text{dvol}(\mathbf{x}) \delta s(\mathbf{x}) \cdot \left. \frac{\overleftarrow{\delta \mathcal{L}}(\mathbf{x}, [\mathbf{u}])}{\delta \mathbf{u}} \right|_{j_\mathbf{x}^\infty(s)}. \quad (3)$$

We now claim that this one-step procedure is a correct consequence of definitions but itself not a definition of the functional’s variation. The above formula conceals a longer, nontrivial reasoning of which the right-hand side in (3) is an implication — provided that the functional S will not be varied by using any other test shifts, i.e., if the correspondence $S \mapsto \mathcal{E}_{\text{EL}}$ yields the object \mathcal{E}_{EL} of further study (cf. [2, §13]). Indeed, we notice that the left-hand side of (3) refers to *three* bundles (namely, the fibre bundle π for a section $s \in \Gamma(\pi)$ whose infinite jet is $j^\infty(s) \in \Gamma(\pi_\infty)$, the bundle π_∞ for the integral functional $S \in \overline{H}^n(\pi)$, and the tangent vector bundle $T\pi$ such that $\delta s \in \Gamma(T\pi)$ at the graph of s in π , see Fig. 1. (In what follows, a reference to attachment points $s(\mathbf{x}) \in \pi^{-1}(\mathbf{x})$ will always be implicit in the notation for δs : for a given section $s \in \Gamma(\pi)$, the base manifold M^n is the domain of definition for a test shift $\delta s(\mathbf{x}, s(\mathbf{x})) = \delta s(\mathbf{x})$ that takes values in $T_{s(\mathbf{x})}\pi^{-1}(\mathbf{x})$.) Let us figure out how the domains of definition for the sections s and δs merge to one copy of the manifold M^n over which an integration is performed in the right-hand side of (3). Strictly speaking, from (3) it is unclear whether the variational derivative,

$$\frac{\overleftarrow{\delta \mathcal{L}}(\mathbf{x}, [\mathbf{u}])}{\delta \mathbf{u}} = \sum_{|\sigma| \geq 0} \left(-\frac{\overleftarrow{d}}{d\mathbf{x}} \right)^\sigma \frac{\overrightarrow{\delta \mathcal{L}}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma},$$

stems from one (which would be false) or both (true!) copies of the base M .

To have a clear vision of the variations’ geometry and by this avoid an appearance of phantoms in description, we now vary the action functional S at $s \in \Gamma(\pi)$ along $\delta s \in \Gamma(T\pi)$, commenting on each step we make. In fact, it suffices to figure out where the objects and structures at hand belong to — in particular, we should explain the nature of binary operation \cdot in the right-hand side of conventional formula (3). The key idea is to understand what we are actually doing but not what we have got used to think we do in order to obtain an understandable result [2, §13]. The discovery is that this “multiplication of functions” is a shorthand notation for the

⁷ An arrow over a variational derivative indicates the direction along which the shift δs is transported left- or rightmost. While the objects are non-graded commutative, this indication is not important. It becomes mandatory in the \mathbb{Z}_2 -graded commutative setup (see section 2): likewise, the arrows are also mandatory and fix the direction of rotation for non-commutative cyclic words [29, 32, 36]; note that our formalism is extended verbatim to the variational calculus of such necklaces and their brackets.

canonically defined coupling between vectors and covectors from (co)tangent spaces $W_{s(\mathbf{x})}$ and $W_{s(\mathbf{x})}^\dagger$, respectively, at the points $s(\mathbf{x})$ of fibres $\pi^{-1}(\mathbf{x})$ in the bundle π .

To encode this linear-algebraic setup, let i, j run from 1 to $m = \dim(\pi^{-1}(\mathbf{x})) = \text{rank}(T\pi)$ and take a local basis $\vec{e}_i(\mathbf{y})$ in the tangent spaces $W_{s(\mathbf{y})} = T_{s(\mathbf{y})}(\pi^{-1}(\mathbf{y}))$ at $s(\mathbf{y})$ over base points $\mathbf{y} \in M$. Introduce the dual basis $\vec{e}^{\dagger j}(\mathbf{x})$ in $W_{s(\mathbf{x})}^\dagger$ attached at $s(\mathbf{x})$ over $\mathbf{x} \in M$. By construction, this means that the value

$$\langle \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \rangle \quad (4)$$

is equal to the Kronecker symbol δ_i^j if and only if $\mathbf{x} = \mathbf{y}$ and the vector $\vec{e}_i(\mathbf{y}) \in W_{p_1}$ and covector $\vec{e}^{\dagger j}(\mathbf{x}) \in W_{p_2}^\dagger$ are attached at the same point $p_1 = p_2$ of the fibre $\pi^{-1}(\mathbf{x})$ over $\mathbf{x} = \mathbf{y} \in M$.

The *locality* of this coupling is an absolute geometric postulate: the coupling is not defined whenever $\mathbf{x} \neq \mathbf{y}$ or the values $p_1 = s_1(\mathbf{y})$ and $p_2 = s_2(\mathbf{x})$ of two local sections $s_1, s_2 \in \Gamma(\pi)$ are not equal at $\mathbf{x} = \mathbf{y}$. Physically speaking, the coupling is then not defined because there is no channel of information which would communicate the value $\delta s^i(\mathbf{y}) \cdot \vec{e}_i(\mathbf{y})$ of excitation of the physical field $s \in \Gamma(\pi)$ at a point $\mathbf{y} \in M$ to another point $\mathbf{x} \neq \mathbf{y}$ of the space-time M .

Remark 1.2. Let us remember that the definition of coupling between sections of (co)tangent bundles — i. e., (co)tangent to either a given manifold or a given bundle π which is the case here for Euler–Lagrange equations — forces the congruence $\{\mathbf{x} = \mathbf{y}, s_1(\mathbf{y}) = s_2(\mathbf{x})\}$ of the (co)vectors’ attachment points. We notice further that such congruence mechanism does not refer to any limiting procedure for smooth distributed kernels and regular linear operators on the space of (co)vector fields. Indeed, vectors couple with their duals at a given point regardless of any phantom limiting procedure which would grasp the (co)vector’s values at any other points of the manifold.⁸

Remark 1.3. The coupling is a matching between test-shift vector fields which are tangent to the fibres of π and, on the other hand, with the elements of $\Gamma(T^*\pi)$ which are determined by the Lagrangian \mathcal{L} . This binary operation yields the singular integral operator $\int_M d\mathbf{y} \langle \delta s^i(\mathbf{y}) \vec{e}_i(\mathbf{y}) |$ with support on the diagonal. Independently, the same operator can reappear as the limit in a parametric family of regular integral operators with smooth, distributed kernels. This shows that the same object is constructed by using several algorithms. Yet the analytic behaviour of the limit is determined not only by the limit itself but also by an algorithm how it is attained. Consequently, the object’s analytic properties in the course of derivations could be (and actually, indeed they are) drastically different for different scenarios. This is the key point in a regularisation of the formalism; to achieve this goal, we properly identify the objects which are de facto handled.

Remark 1.4. Referring to a concept of locality of events, this definition of coupling \langle , \rangle ensures a very interesting analytic behaviour of the value $\langle \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger i}(\mathbf{x}) \rangle$ of pairing for dual objects $\vec{e}_i(\mathbf{y})$ and $\vec{e}^{\dagger i}(\mathbf{x})$ at fixed i . Namely, this value is a constant scalar field which equals unit $1 \in \mathbb{K}$ at all points of the manifold M ; the scalar field’s partial derivatives with respect to x^j or y^k , $1 \leq j, k \leq n$, vanish identically. We shall use this property in what follows (see Remark 1.7 on p. 11). We also note that the logarithm of this coupling’s unit value vanishes as well: $\log \langle \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger i}(\mathbf{x}) \rangle = 0$ whenever the coupling is well defined and $1 \leq i \leq m$.

Now let us return to the initial setup in context of Euler–Lagrange equation \mathcal{E}_{EL} and one-step correspondence $S \mapsto \mathcal{E}_{\text{EL}}$, see Fig. 1. We have that $S = \int \mathcal{L}(\mathbf{x}, [\mathbf{u}]) \text{dvol}(\mathbf{x})$ is an integral functional; we let $s \in \Gamma(\pi)$ be a background section (e. g., a sought-for solution of the Euler–Lagrange stationary point equation $\delta S|_s = 0$) and $\delta s \in \Gamma(T\pi)$ be a test shift of s . The linear

⁸ We recall that a similar, purely local geometric principle, not referring to the objects’ values at non-coinciding points, works in the definition of Hirota’s bilinear derivative.

term in a response of $S: \Gamma(\pi) \rightarrow \mathbb{k}$ to a shift of its argument s along δs is (cf.(14) on p. 20)

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} S(s + \varepsilon \overleftarrow{\delta s}) &= \\ &= \sum_{i,j} \sum_{|\sigma| \geq 0} \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle (\delta s^i) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}} \right)^\sigma (\mathbf{y}) \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_\sigma^j} \Big|_{j_\mathbf{x}^\infty(s)} \right\rangle. \end{aligned} \quad (5)$$

Remark 1.5. The rôles of two integral signs in (5) are different. Namely, the volume form $d\text{vol}(\mathbf{x})$ at $\mathbf{x} \in M^n$ comes from the integral functional $S \in \overline{H}^n(\pi)$; should a formal choice of the volume form be different, the Euler–Lagrange equations would also change.⁹ At the same time, the other integral sign $\int d\mathbf{y}$ denotes the singular linear operator $\Gamma(T^*\pi) \rightarrow \mathbb{k}$ with support on the diagonal [19]; in fact, this notation means that a point \mathbf{y} runs through the entire integration domain M .

1.3. Integration by parts

The most interesting things start to happen when one integrates by parts over the domain M^n of test shifts δs . (By default, we let the supports of local perturbations δs be such that no boundary terms appear in the course of integration by parts over M .)

For the sake of transparency let us first consider a model situation when there is just one derivative falling on δs at \mathbf{y} ; all higher-order cases are processed recursively. By the definition of a (partial) derivative $\partial/\partial y^i$, we have that¹⁰

$$\begin{aligned} \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle (\delta s) \frac{\overleftarrow{\partial}}{\partial \mathbf{y}} (\mathbf{y}) \vec{e}(\mathbf{y}), \vec{e}^\dagger(\mathbf{x}) \frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_\sigma} \Big|_{j_\mathbf{x}^\infty(s)} \right\rangle &= \\ &= - \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \delta s(\mathbf{y}) \frac{\overrightarrow{\partial}}{\partial \mathbf{y}} \left\{ \langle \vec{e}(\mathbf{y}), \vec{e}^\dagger(\mathbf{x}) \rangle \frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_\sigma} \Big|_{j_\mathbf{x}^\infty(s)} \right\}. \end{aligned}$$

By using a definition of the partial derivative which falls on the comultiple of δs , we obtain the

⁹ There are natural classes of geometries in which the Lagrangian $\mathcal{L}(\mathbf{x}, [\mathbf{u}])$ in the action S is a well-defined top-degree differential form, e.g., if the unknowns \mathbf{u} are differential one-forms (we recall the Yang–Mills or Chern–Simons gauge theories in this context). Let us remember also that a construction of \mathcal{L} could refer to a choice of volume form $d\text{vol}(\mathbf{x})$ on M^n . For instance, such is the case when the Hodge structure $*$ is involved (the Yang–Mills Lagrangian yields an example: $\mathcal{L} \sim F_{\mu\nu} * F^{\mu\nu}$ in standard notation for the stress tensor). To avoid excessive case-study, we use a uniform notation thus writing $d\text{vol}(\mathbf{x})$ explicitly.

We recall further that the integration measure $d\text{vol}(\mathbf{x}, s(\mathbf{x})) = \sqrt{|\det(g_{\mu\nu}(\mathbf{x}, s))|} d\mathbf{x}$ is field-dependent by virtue of Einstein’s general relativity equations which –in their right-hand sides– absorb the energy-momentum tensor of physical fields $s \in \Gamma(\pi)$. The volume element will be denoted by $d\text{vol}(\mathbf{x})$ in order to emphasize that the space-time M^n is unique: Namely, field-dependent objects interact at its points only if the local geometry of underlying space-time is the same near $\mathbf{x} \in M^n$ for all objects (see Theorem 3 and Remark 2.11 on p. 29 for a realisation of this principle for the smooth manifold M^n endowed with metric tensor $g_{\mu\nu}$).

¹⁰ In the definition of derivative, the calculation of length $|\Delta \mathbf{y}|$ in denominators refers to the standard Euclidean metric in the linear vector spaces which determine coordinate neighbourhoods near points of the manifold M at hand.

difference¹¹

$$\stackrel{\text{def}}{=} - \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \delta s(\mathbf{y}) \cdot \lim_{|\Delta\mathbf{y}| \rightarrow 0} \frac{1}{|\Delta\mathbf{y}|} \left\{ \langle \vec{e}(\mathbf{y} + \Delta\mathbf{y}), \vec{e}(\mathbf{x}) \rangle \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x}}^\infty(s)} - \langle \vec{e}(\mathbf{y}), \vec{e}(\mathbf{x}) \rangle \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x}}^\infty(s)} \right\}.$$

The locality postulate for coupling between (co)vectors \vec{e} and \vec{e}^\dagger forces the equality $\mathbf{y} + \Delta\mathbf{y} = \mathbf{x}$ in the minuend, which yields the two different points at which the restriction of Lagrangian \mathcal{L} to the jet $j^\infty(s)$ of section $s \in \Gamma(\pi)$ is evaluated:

$$= - \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \delta s(\mathbf{y}) \cdot \langle \vec{e}(\mathbf{y}), \vec{e}^\dagger(\mathbf{x}) \rangle \cdot \lim_{|\Delta\mathbf{y}| \rightarrow 0} \frac{1}{|\Delta\mathbf{y}|} \left\{ \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x} + \Delta\mathbf{y}}^\infty(s)} - \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x}}^\infty(s)} \right\}.$$

(Here we use the fact that the scalar product $\langle \cdot, \cdot \rangle$, whenever defined, is the Kronecker symbol.) We continue the equality,

$$\stackrel{\text{def}}{=} - \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle \delta s(\mathbf{y}) \vec{e}(\mathbf{y}), \vec{e}^\dagger(\mathbf{x}) \frac{\vec{\partial}}{\partial \mathbf{x}} \left(\frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x}}^\infty(s)} \right) \right\rangle.$$

We finally recall that the total derivative $d/d\mathbf{x}$ is defined¹² via an application of $\partial/\partial\mathbf{x}$ to restriction to infinite jets $j^\infty(s)$ of sections s at base points \mathbf{x} . Therefore, the above expression is equal to

$$\stackrel{\text{def}}{=} \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle \delta s(\mathbf{y}) \vec{e}(\mathbf{y}), \vec{e}^\dagger(\mathbf{x}) \left(\left(-\frac{\vec{d}}{d\mathbf{x}} \right) \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma} \Big|_{j_{\mathbf{x}}^\infty(s)} \right) \right\rangle.$$

This shows that an integration by parts over the base M in the geometry of test shift δs reappears as integration by parts in the bundle where lives the background section $s \in \Gamma(\pi)$.

Repeating the integration by parts $|\sigma| \geq 0$ times in each term of the sum in (5), we obtain the expression

$$\sum_{i,j} \sum_{|\sigma| \geq 0} \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle \delta s^i(\mathbf{y}) \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \left(\left(-\frac{\vec{d}}{d\mathbf{x}} \right)^\sigma \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma^j} \Big|_{j_{\mathbf{x}}^\infty(s)} \right) \right\rangle.$$

Let us recall once more that the coupling's support is the diagonal in $M \times M$, at points of which the value $\langle \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \rangle$ is the Kronecker symbol δ_i^j . Consequently, we arrive at

$$= \sum_{i,j} \sum_{|\sigma| \geq 0} \int_M d\text{vol}(\mathbf{x}) \delta s^i(\mathbf{x}) \cdot \left(\left(-\frac{\vec{d}}{d\mathbf{x}} \right)^\sigma \frac{\vec{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial \mathbf{u}_\sigma^i} \Big|_{j_{\mathbf{x}}^\infty(s)} \right).$$

¹¹ Here and in the equalities below we suppress the indexes i running through $1, \dots, m$ at $\delta s^i(\mathbf{y})$ and $\vec{e}_i(\mathbf{y})$ or $\vec{e}^{\dagger i}(\mathbf{x})$, or at u_σ^i in the derivative which acts on \mathcal{L} ; we thus avoid an agglomeration of formulas.

¹² By definition, $(\vec{d}/d\mathbf{x}^i)|_{j_{\mathbf{x}}^\infty(s)}(\mathbf{x}) = (\vec{\partial}/\partial x^i(f|_{j_{\mathbf{x}}^\infty(s)}))(\mathbf{x})$ for differential functions f , see [28, 40, 45].

This is formula (3); it is familiar from any textbook on variational principles of classical mechanics (e.g., see [2, §12–13]).

A standard reasoning shows that, whenever a response of the functional’s value $S(s)$ to a test shift of s along any direction δs vanishes, the Euler–Lagrange equation holds:

$$\left. \frac{\overleftarrow{\delta} S}{\delta \mathbf{u}} \right|_{j^\infty(s)} = 0. \tag{6}$$

Its left-hand side belongs to the space $\Gamma(T^*\pi)$ of sections of the cotangent bundle to π .

Remark 1.6. This conclusion tells us that traditional attempts of a brute-force labelling of equations in a given system (6) by using the unknowns \mathbf{u} is not geometric. Indeed, the equations’ left-hand sides are sections of a *vector* bundle, thus forming linear \mathbb{k} -vector spaces so that addition is well defined for the equations within a system. On the other hand, the fibres in the bundle π can be smooth manifolds (i.e., not necessarily being vector spaces) so that one may not add points of those fibres; for such operation is in general not defined at all. Even if π is a vector bundle, the fibres of which are endowed with linear vector space structure, the two structures are not related.

Remark 1.7. The integration by parts transforms a derivative $\partial/\partial \mathbf{y}$ along one copy of the base M to the minus derivative $-\partial/\partial \mathbf{x}$ along the other copy. This produces no visible effect on the mechanism which ensures a restriction onto the diagonal in $M \times M$, i.e., there appears no would-be third term in the Leibniz rule for the product which is defined only on the diagonal. A desperate prescription (2) was introduced in the literature in order to mimick this paradoxical analytic behaviour of the coupling between elements of dual bases.

1.4. Why are variations permutable?

Having outlined the matching of geometries in the course of one sequence of integrations by parts for *one* fixed pair $M \times M \ni (\mathbf{y}, \mathbf{x})$ of copies of the base manifold, we emphasize that such integrations must be performed last, i.e., only when the objects at hand are finally viewed as maps $\Gamma(\pi) \rightarrow \mathbb{k}$.

Should one haste in absence of clear understanding of what is actually being done and for which purpose, further calculation of higher-order variations could predictably but uncontrollably lead to meaningless, manifestly erroneous conclusions (e.g., compare left- and right-hand sides in (7) below).

Namely, there exist integral functionals which determine equal maps $\Gamma(\pi) \rightarrow \mathbb{k}$ but, belonging to different spaces, behave differently in the course of variations, should one attempt any. We say that such functionals are *synonyms*; for instance, see Example 2.4 in the next section for a nontrivial synonym ΔG of the zero functional (cf. Fig. 2). Informally speaking, the composite structure objects with repeated integrals over products $M \times M \times \dots \times M$ of the base retain a kind of memory of the way how they were obtained from primary objects such as the action S . Let us illustrate these claims.

Example 1.1. Let $\delta s_1 \in \Gamma(T\pi)$ be a test shift at $s \in \Gamma(\pi)$ for an integral functional $S = \int \mathcal{L}(\mathbf{x}, [\mathbf{u}]) \, \text{dvol}(\mathbf{x})$ with density \mathcal{L} of positive differential order. (That is, we suppose that some positive-order derivatives are always present in densities of all representatives of the cohomology class $S \in \overline{H}^n(\pi)$; this assumption is not to any extent restrictive but it allows us to not take into account \bar{d} -exact terms whose orders may not be bounded.) By using S , let us construct two new integral functionals. First, we set

$$F = \sum_i \sum_{|\sigma| \geq 0} \int \text{dvol}(\mathbf{x}) \delta s_1^i(\mathbf{x}) \cdot \left(-\frac{\bar{d}}{\text{d}\mathbf{x}} \right)^\sigma \left(\frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_\sigma^i} \right) \in \overline{H}^n(\pi),$$




Figure 2. The synonyms ΔG of zero functional yield constant maps $0: \Gamma(\pi_{BV}) \rightarrow \mathbb{k}$ yet they can nontrivially contribute to larger structures such as $\llbracket F, \Delta G \rrbracket$, see Example 2.4 on p. 34.

so that the mapping $F: \Gamma(\pi) \rightarrow \mathbb{k}$ is defined at $s \in \Gamma(\pi)$ by restriction of the integrand to the jet $j^\infty(s)$ and then by actual integration over M .

Let the other functional $G \in \overline{H}^{2n}(\pi, T\pi)$ be such that its value at the same section $s \in \Gamma(\pi)$ is

$$G(s) = \sum_{i,j} \sum_{|\sigma| \geq 0} \int_M d\mathbf{y} \int_M d\text{vol}(\mathbf{x}) \left\langle (\delta s_1^i) \left(\overleftarrow{\frac{\partial}{\partial \mathbf{y}}} \right)^\sigma (\mathbf{y}) \vec{e}(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_\sigma^j} \Big|_{j_x^\infty(s)} \right\rangle.$$

From the previous section it is clear that F and G are indistinguishable as mappings to \mathbb{k} for every $s \in \Gamma(\pi)$. Yet their variations, i. e., the responses to an extra shift $\delta s_2 \in \Gamma(T\pi)$, are different. Indeed, they are equal to, first,

$$\left(\frac{d}{d\varepsilon_2} \Big|_{\varepsilon_2=0} F \right) (s + \varepsilon_2 \overleftarrow{\delta s_2}) = \sum_{i_1, i_2} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M d\text{vol}(\mathbf{x}) \delta s_2^{i_2}(\mathbf{x}) \delta s_1^{i_1}(\mathbf{x}) \cdot \left\{ \left(-\overrightarrow{\frac{d}{d\mathbf{x}}} \right)^{\sigma_2} \frac{\overrightarrow{\partial}}{\partial u_{\sigma_2}^{i_2}} \left(\left(-\overrightarrow{\frac{d}{d\mathbf{x}}} \right)^{\sigma_1} \frac{\overrightarrow{\partial} \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_{\sigma_1}^{i_1}} \right) \right\} \Big|_{j_x^\infty(s)}.$$

The above formula corresponds to a step-by-step calculation within a naïve approach to the geometry of variations. However, the genuine value of second variation of the integral functional S along δs_1 and then δs_2 at a section s is

$$\left(\frac{d}{d\varepsilon_2} \Big|_{\varepsilon_2=0} G \right) (s + \varepsilon_2 \overleftarrow{\delta s_2}) = \sum_{\substack{i_1, i_2 \\ j_1, j_2}} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M d\mathbf{y}_2 \int_M d\mathbf{y}_1 \int_M d\text{vol}(\mathbf{x}) \left\{ (\delta s_2^{i_2}) \left(\overleftarrow{\frac{\partial}{\partial \mathbf{y}_2}} \right)^{\sigma_2} (\mathbf{y}_2) \langle \vec{e}_{i_2}(\mathbf{y}_2), \vec{e}^{\dagger j_2}(\mathbf{x}) \rangle \cdot (\delta s_1^{i_1}) \left(\overleftarrow{\frac{\partial}{\partial \mathbf{y}_1}} \right)^{\sigma_1} (\mathbf{y}_1) \langle \vec{e}_{i_1}(\mathbf{y}_1), \vec{e}^{\dagger j_1}(\mathbf{x}) \rangle \right\} \cdot \frac{\overrightarrow{\partial}^2 \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_{\sigma_2}^{j_2} \partial u_{\sigma_1}^{j_1}} \Big|_{j_x^\infty(s)}.$$

The analytic distinction between the operators

$$\underbrace{\left(-\frac{\overrightarrow{d}}{d\mathbf{x}}\right)^{\sigma_2} \circ \frac{\overrightarrow{\partial}}{\partial u_{\sigma_2}^{i_2}} \circ \left(-\frac{\overrightarrow{d}}{d\mathbf{x}}\right)^{\sigma_1} \circ \frac{\overrightarrow{\partial}}{\partial u_{\sigma_1}^{i_1}}}_{\text{naïve approach}} \quad \text{and} \quad \underbrace{\left(-\frac{\overrightarrow{d}}{d\mathbf{x}}\right)^{\sigma_1 \cup \sigma_2} \circ \frac{\overrightarrow{\partial}^2}{\partial u_{\sigma_2}^{i_2} \partial u_{\sigma_1}^{i_1}}}_{\text{geometric theory}} \quad (7)$$

reveals why in positive-order Lagrangian models it is forbidden to haste, which would imply that the derivatives along distinct copies of M for variations $\delta s_1, \dots, \delta s_k$ are too early transformed to derivatives along the functional’s own base. Such a conceptual error would repercuss with inexplicable, redundant terms in variations to-follow.

On the other hand, as soon as the product-bundle geometry of iterated variations is properly realized — so that all restrictions to the diagonals are postponed as late as possible, — the variations become (graded-)permutable.¹³ Namely, denote by $|u^i|$, $1 \leq i \leq m$, the overall \mathbb{Z}_2 -valued parities of the fibre coordinates u^i ; the ghost parity $\text{gh}(u^i)$ or individual \mathbb{Z} - or \mathbb{Z}_2 -valued gradings in the bundle π contribute additively to $|u^i|$ and then a residue modulo 2 is taken. Suppose that $\delta s_1 = (\delta s_1^{i_1})$ and $\delta s_2 = (\delta s_2^{i_2})$ are test shifts and $S = \int \mathcal{L}(\mathbf{x}, [\mathbf{u}]) \text{dvol}(\mathbf{x})$ is an integral functional which maps a section $s \in \Gamma(\pi)$ to \mathbb{k} . Then, *after* the integrations by parts in the product-bundle geometry $\pi \times T\pi \times T\pi$ which is described above, there remains

$$\begin{aligned} & \sum_{i_1, i_2} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M \text{dvol}(\mathbf{x}) \delta s_2^{i_2}(\mathbf{x}) \delta s_1^{i_1}(\mathbf{x}) \left(\left(-\frac{\overrightarrow{d}}{d\mathbf{x}}\right)^{\sigma_1 \cup \sigma_2} \frac{\overrightarrow{\partial}^2 \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_{\sigma_2}^{i_2} \partial u_{\sigma_1}^{i_1}} \right) \Big|_{j_{\mathbf{x}}^\infty(s)} \\ &= \sum_{i_1, i_2} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} (-)^{|u^{i_1}| \cdot |u^{i_2}|} \int_M \text{dvol}(\mathbf{x}) \delta s_1^{i_1}(\mathbf{x}) \delta s_2^{i_2}(\mathbf{x}) \left(\left(-\frac{\overrightarrow{d}}{d\mathbf{x}}\right)^{\sigma_1 \cup \sigma_2} \frac{\overrightarrow{\partial}^2 \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_{\sigma_1}^{i_1} \partial u_{\sigma_2}^{i_2}} \right) \Big|_{j_{\mathbf{x}}^\infty(s)}. \end{aligned}$$

Likewise, higher-order iterated variations with $k \geq 2$ test shifts $\delta s_1, \dots, \delta s_k$ are (graded-)permutable with the same rule of signs for permutations of order in which the (graded) partial derivatives $\overrightarrow{\partial} / \partial u_{\sigma_1}^{i_1}, \dots, \overrightarrow{\partial} / \partial u_{\sigma_k}^{i_k}$ fall from the left on the density \mathcal{L} of the functional S . (A case of \mathbb{Z}_2 -graded base manifold $M^{(n_0|n_1)}$ would bring more signs which are also captured in a standard way.)

Let there be $k \geq 2$ variations $\delta s_1, \dots, \delta s_k \in \Gamma(T\pi)$. We finally have that

$$\begin{aligned} & \frac{d}{d\varepsilon_k} \Big|_{\varepsilon_k=0} \circ \dots \circ \frac{d}{d\varepsilon_1} \Big|_{\varepsilon_1=0} S(s + \varepsilon_1 \overleftarrow{\delta s_1} + \dots + \varepsilon_k \overleftarrow{\delta s_k}) = \\ &= \sum_{\substack{i_1, \dots, i_k \\ j_1, \dots, j_k}} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_k| \geq 0}} \int_M d\mathbf{y}_k \dots \int_M d\mathbf{y}_1 \int_M \text{dvol}(\mathbf{x}) \cdot \\ & \left\{ (\delta s_k^{i_k}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_k} \right)^{\sigma_k} (\mathbf{y}_k) \langle \vec{e}_{i_k}(\mathbf{y}_k), \vec{e}^{\dagger j_k}(\mathbf{x}) \rangle \dots (\delta s_1^{i_1}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} (\mathbf{y}_1) \langle \vec{e}_{i_1}(\mathbf{y}_1), \vec{e}^{\dagger j_1}(\mathbf{x}) \rangle \right\} \\ & \cdot \frac{\overrightarrow{\partial}^k \mathcal{L}(\mathbf{x}, [\mathbf{u}])}{\partial u_{\sigma_k}^{j_k} \dots \partial u_{\sigma_1}^{j_1}} \Big|_{j_{\mathbf{x}}^\infty(s)}. \quad (8) \end{aligned}$$

¹³ An idea that iterated variations must be taken at nominally different points \mathbf{x} and \mathbf{y} has been in the air for a long time (let us refer to [38, §1] which contains due credits to E. Witten). A somewhat less obvious fact is that those different points belong to different copies of the manifold M in the product bundle $\pi \times T\pi \times \dots \times T\pi$ over $M \times M \times \dots \times M$.

Whenever any $k - 1$ variation(s) are fixed in the above formula, the co-multiple $|\rangle$ of the remaining, ℓ th variation $\delta s_\ell = \langle \delta s_\ell^{i_\ell}(\mathbf{y}_k) \vec{e}_{i_\ell}(\mathbf{y}_\ell) |$ is an element of the cotangent vector space $T_{s(\mathbf{x})}^* \pi^{-1}(\mathbf{x}) = V_{\mathbf{x}}^\dagger$ at the point $s(\mathbf{x})$ in the fibre $\pi^{-1}(\mathbf{x})$ over a base point $\mathbf{x} \in M^n$.

Remark 1.8. The composite object in the left-hand side of equality (8) is an integral functional in the bundle $\pi \times T\pi \times \dots \times T\pi$ which properly contains the geometry of k variations from $\Gamma(T\pi)$, see Fig. 3. This construction lives not on a Whitney sum $\pi \times_M T\pi \times_M \dots \times_M T\pi$ over




Figure 3. Each variation $\delta s_1, \dots, \delta s_k$ brings its own copy of the base $M^n \ni \mathbf{y}_\ell$ into the product bundle $\pi \times T\pi \times \dots \times T\pi$ over $M \times M \times \dots \times M$.

the base manifold M ; that would force an untimely restriction to the diagonal in the product $M \times M \times \dots \times M$ of bases and hence reproduce the old difficulties of the theory.

1.5. *The spaces of functionals*

The integral functionals $S \in \overline{H}^n(\pi)$, which we have been dealing with until now, are building blocks in a wider class of mappings $\Gamma(\pi) \rightarrow \mathbb{k}$. By viewing elements of $\Gamma(\pi)$ as “points” and functionals from $\overline{H}^n(\pi)$ as “elementary functions” (see [40] and references therein), we consider pointwise-defined (formal sums of) products of such maps, e. g., we let

$$(S_1 \cdot S_2)(s) \stackrel{\text{def}}{=} S_1(s) \cdot S_2(s)$$

for any two already defined functionals S_1 and S_2 ; the binary operation \cdot for their values at $s \in \Gamma(\pi)$ is the usual multiplication of \mathbb{k} -numbers ($\mathbb{k} = \mathbb{R}$ or \mathbb{C}). By definition, we put

$$\overline{\mathfrak{N}}^n(\pi, T\pi) = \bigoplus_{\ell=1}^{+\infty} \bigotimes_{i=1}^{\ell} \bigoplus_{k=0}^{+\infty} \overline{H}^{n(1+k)}(\underbrace{\pi \times T\pi \times \dots \times T\pi}_{k \text{ variations}}).$$

This space contains the linear subspace of *local functionals*,

$$\overline{\mathfrak{M}}^n(\pi) = \bigoplus_{\ell=1}^{+\infty} \bigotimes_{i=1}^{\ell} \overline{H}^n(\pi),$$

for instance, such as the standard weight factor $\exp(\frac{i}{\hbar} S^{\hbar})$ in BV-models with quantum BV-action S^{\hbar} (see section 3.2 below, cf. [9]). The larger space $\overline{\mathfrak{N}}^n(\pi, T\pi) \supseteq \overline{\mathfrak{M}}^n(\pi)$ harbours local functionals *and* their variations of arbitrarily high order. The (products of) integral functionals in $\overline{\mathfrak{M}}^n(\pi) \supset \overline{H}^n(\pi)$ could be viewed as primary objects. In the course of variations, their descendants in $\overline{\mathfrak{N}}^n(\pi, T\pi)$ absorb new test shifts and retain the information about initial building blocks from $\overline{H}^n(\pi)$. This memory governs the analytic behaviour of descendants in operations such as calculation of the BV-Laplacian or taking the Schouten bracket; we also refer to sections 1.4 above and 3.1 in what follows. The composite structure of the bundle

$\pi \times T\pi \times \dots \times T\pi$ is crucial whenever one wants to not only describe initial setup such as a given BV-model but to perform rigorous calculations in it, handling higher-order variations of objects (e. g., third-order variations occur in (1c) on p. 1, see also Example 2.4 on p. 34 below, — and the order is equal to four in property (1d) for the BV-Laplacian Δ to be a differential). The geometric approach to (graded-)permutable variations of functionals makes such calculations well-defined and proofs free from any *ad hoc* regularisation recipes.

2. The geometry of Batalin–Vilkovisky formalism

The geometry of variations which we analysed in the previous section was not specific to a bundle π of unknowns. In this section we first recall a construction of the BV-superbundle whose fibres are endowed with \mathbb{Z}_2 -valued ghost parity. By definition, the BV-bundle $\pi_{\text{BV}}^{(0|1)} = \pi_\infty^*(\zeta_\infty^{(0|1)})$ is induced from the Whitney sum $\zeta^{(0|1)} = \zeta_0 \times_M \zeta_1 \times_M \dots \times_M \zeta_\lambda \times_M \Pi\widehat{\zeta}_0 \times_M \Pi\widehat{\zeta}_1 \times_M \dots \times_M \Pi\widehat{\zeta}_\lambda$ of some \mathbb{Z}_2 -graded vector bundles over M (in what follows we sum up the construction of $\zeta_0, \dots, \zeta_\lambda$ and their parity-reversed duals $\Pi\widehat{\zeta}_0, \dots, \Pi\widehat{\zeta}_\lambda$) by the infinite jet bundle $\pi_\infty: J^\infty(\pi) \rightarrow M$ associated with the smooth fibre bundle π of physical fields.¹⁴

2.1. The BV-zoo

Let a fibre bundle π of physical fields over the base manifold M^n be given and denote by ϕ the fibre coordinates in it. Suppose that

$$S_0 = \int \mathcal{L}_0(\mathbf{x}, [\phi]) \, \text{dvol}(\mathbf{x}) \in \overline{H}^n(\pi)$$

is the action of a field model under study. By using the theory and techniques from section 1 we know how one derives, via the stationary point condition $\overleftarrow{\delta} S|_s = 0$ at $s \in \Gamma(\pi)$ the Euler–Lagrange equations of motion $\mathcal{E}_{\text{EL}} = \{\overleftarrow{\delta} S_0 / \delta \phi = 0\}$ whose left-hand sides belong to the $C^\infty(J^\infty(\pi))$ -module of sections $P_0 = \Gamma(\pi_\infty^*(\zeta_0))$ for the cotangent bundle ζ_0 to π such that $\overleftarrow{\delta} S_0 / \delta \phi|_{j^\infty(s)} \cdot \text{dvol}(\cdot) \in \Gamma(T^*\pi) \otimes_{C^\infty(M)} \Lambda^n(M)$ for any field configuration $s \in \Gamma(\pi)$.

We recall from Remark 1.6 that by following a misfortunate but long-established tradition it is the unknowns ϕ in π but not the global coordinates \mathbf{F} in the fibre of cotangent bundle $T^*\pi$ to π which are used to parametrise the equations within Euler–Lagrange system \mathcal{E}_{EL} at points of the graph of a section $\phi \in \Gamma(\pi)$.

If the model at hand is gauge-invariant, then it admits an off-shell differential dependence $\Phi(\mathbf{x}, [\phi]; [\mathbf{F}]) \equiv 0 \in \Gamma((\pi_\infty \times_M \zeta_{0,\infty})^*(\zeta_1))$ between the left-hand sides \mathbf{F} of equations \mathcal{E}_{EL} . We recall further that the dependence of Noether’s identities Φ on (the derivatives of) \mathbf{F} is *linear* for Euler–Lagrange systems \mathcal{E}_{EL} ; the generators $\mathbf{p}(\mathbf{x}, [\phi]) \in \widehat{P}_1 = \Gamma(\pi_\infty^*(\widehat{\zeta}_1))$ of Noether’s gauge symmetries for S_0 are sections of the bundle $\widehat{\zeta}_1$ which is induced from the dual to ζ_1 with respect to the top-degree horizontal form-valued coupling $\langle \cdot, \cdot \rangle$. Indeed, if

$$0 \equiv \langle \mathbf{p}, \Phi(\mathbf{x}, [\phi]; [\mathbf{F}]) \rangle$$

and Φ is linear in \mathbf{F} or its finite-order derivatives,

$$\Phi(\mathbf{x}, [\phi]; [\mathbf{F}]) = \ell_{\Phi}^{(\mathbf{F})}(\mathbf{F}) \equiv 0,$$

¹⁴ A subtle point, which we reconsider in section 2.1 (see also Remark 1.6), is that the *fibre* bundle π is often identified with the *vector* bundle component ζ_0 in $\zeta^{(0|1)}$. Nevertheless, it is the construction of induced bundle $\pi_\infty^*(\zeta_0 \times_M \dots)$ by using which the physical fields and their derivatives are remembered by the Euler–Lagrange equations (referred to ζ_0), Noether’s identities (in ζ_1), and higher generations of syzygies from $\zeta_2, \dots, \zeta_\lambda$ (if any).

then an integration by parts yields that

$$0 \cong \left\langle (\ell_{\mathbf{\Phi}}^{(\mathbf{F})})^\dagger(\mathbf{p}), \delta S_0 / \delta \phi \right\rangle \cong \vec{\partial}_{(\ell_{\mathbf{\Phi}}^{(\mathbf{F})})^\dagger(\mathbf{p})}^{(\phi)}(S_0).$$

This shows that the evolutionary vector field $\vec{\partial}_{A(\mathbf{p})}^{(\phi)}$ with $A = (\ell_{\mathbf{\Phi}}^{(\mathbf{F})})^\dagger$ and $\mathbf{p} = \mathbf{p}(\mathbf{x}, [\phi])$ is a Noether symmetry of the action S_0 . By reading the above equalities backwards, one obtains the linear Noether relations $\mathbf{\Phi} = A^\dagger(\mathbf{F})$ between the Euler–Lagrange equations of motion.

Likewise, there could in principle appear higher generations of linear identities $\Psi_2(\mathbf{x}, [\phi], [\mathbf{F}]; [\mathbf{\Phi}]) \equiv 0, \dots, \Psi_\lambda(\mathbf{x}, [\phi], [\mathbf{F}], [\mathbf{\Phi}], \dots, [\Psi_{\lambda-2}]; [\Psi_{\lambda-1}]) \equiv 0$ which hold for all ϕ , sections \mathbf{F} in ζ_0 , and so on up to the coordinates $\Psi_{\lambda-2}$. Each i th generation of such identities arises with the respective vector bundle ζ_i with fibre dimension m_i ; the total number of generations is bounded from above by a constant $\lambda \in \mathbb{N} \cup \{0\}$ due to Hilbert’s theorem on syzygies [16]: $0 \leq i \leq \lambda \leq n$, where n is the dimension of base manifold M^n . For example, we have that $\lambda = 1$ for Yang–Mills theory, and $\lambda = 2$ for gravity over a fourfold M^4 .

We denote by \mathbf{F} (alas! at once identifying this global m -tuple in ζ_0 for the equations with the local field variables ϕ), and by $\gamma^\dagger, \mathbf{c}^\dagger, \dots, \mathbf{c}_\lambda^\dagger$ the global fibre coordinates in ζ_1 for Noether’s identities, and so on up to ζ_λ , respectively (see Fig. 4).




Figure 4. The fibre bundle π of physical fields ϕ , the bundle $\zeta^{(0|1)}$ of BV-variables $(\mathbf{q}, \mathbf{q}^\dagger)$, and the vector bundle $T\zeta^{(0|1)}$ of their variations $\delta \mathbf{s} = (\delta s; \delta s^\dagger)$.

In turn, each vector bundle $\zeta_0, \dots, \zeta_\lambda$ brings its \langle, \rangle_i -dual $\widehat{\zeta}_i$ into the picture. (Note that the equations $\overleftarrow{\delta S_0}|_s = 0$ upon $s \in \Gamma(\pi)$ for $S_0 = \int \mathcal{L}(\mathbf{x}, [\phi]) \cdot \text{dvol}(\mathbf{x})$ and all equations’ linear-differential descendants retain the volume form $\text{dvol}(\mathbf{x})$ from the model’s action S_0 at all points $\mathbf{x} \in M^n$.)

We now reverse the parity of linear vector space fibres in $\widehat{\zeta}_0, \dots, \widehat{\zeta}_\lambda$ by introducing the \mathbb{Z}_2 -valued ghost parity $\text{gh}(\cdot)$ and considering the odd neighbours $\Pi \widehat{\zeta}_0, \dots, \Pi \widehat{\zeta}_\lambda$ of the dual vector bundles (see [34, 52] and also Appendix A in [33] for discussion). Let us denote by $\phi^\dagger, \gamma, \mathbf{c}, \dots, \mathbf{c}_\lambda$ the ghost parity-odd global coordinates along linear vector space fibres in $\Pi \widehat{\zeta}_0, \dots, \Pi \widehat{\zeta}_\lambda$, respectively. These variables’ proper names are easily recognized from the standard notation: ϕ replacing \mathbf{F} are the fields and ϕ^\dagger are odd-parity *antifields*, γ are the odd *ghosts* and γ^\dagger are the parity-even *antighosts*, whereas the canonically conjugate variables $\mathbf{c} \leftrightarrow \mathbf{c}^\dagger, \dots, \mathbf{c}_\lambda \leftrightarrow \mathbf{c}_\lambda^\dagger$ are higher ghost-antighost pairs of opposite ghost parities (resp., odd and even). We denote by \mathbf{q} the agglomeration of ghost parity-even variables and by \mathbf{q}^\dagger their respective canonically conjugate parity-odd neighbours.¹⁵

¹⁵ Consider Feynman’s path integral $\int_{\Gamma(\zeta_0)} [D\mathbf{q}] \mathcal{O}([\mathbf{q}], [\mathbf{q}^\dagger])$ of an observable \mathcal{O} over the space of ghost parity-even sections. The BV-Laplacian Δ is the tool which ensures the integral’s effective independence from the unphysical ghost parity-odd variables \mathbf{q}^\dagger , see section 3.1.

Remark 2.1. Let us emphasize that by using the word “parity” we always refer to the ghost parity $\text{gh}(\cdot)$ of objects.¹⁶ In this paper we aim at understanding the geometry of variations so that the graded arithmetic and algebra of derivations play auxiliary rôles. However, as soon as the interaction of geometries is properly fixed, their extension to a \mathbb{Z}_2 -graded setup of superbundle $\pi: E^{(m_0+n_0|m_1+n_1)} \rightarrow M^{(n_0|n_1)}$ of physical fields (possibly, over a base supermanifold $M^{(n_0|n_1)}$) makes no conceptual difficulty ([10], see also [22] and references therein). The theory then becomes bi-graded: it involves (i) the \mathbb{Z}_2 -grading $|\cdot|$ in the ring of field coordinates, which echoes in the \mathbb{Z}_2 -grading of Euler–Lagrange equations of motion, Noether identities, etc., (the model’s action functional S_0 has even grading by default), and (ii) the ghost parity $\text{gh}(\cdot)$, see [52].

The \mathbb{Z}_2 -grading $|\cdot|$ and the ghost parity $\text{gh}(\cdot)$ are independent from each other. We denote by $\mathbf{q} = \mathbf{q}^{(0|1)}$ the ghost parity-even BV-fibre variables, which are then grouped in even- and odd-grading components. Likewise, the ghost parity-odd BV-variables $\mathbf{q}^\dagger = (\mathbf{q}^\dagger)^{(0|1)}$ are arranged in exactly the same way. By construction, the values of \mathbb{Z}_2 -gradings for canonically conjugate variables $(\mathbf{q}, \mathbf{q}^\dagger)$ coincide: we have that $|\mathbf{q}| = |\mathbf{q}^\dagger|$ and $\text{gh}(\mathbf{q}^\dagger) \equiv \text{gh}(\mathbf{q}) + 1 \pmod 2$.

Next, we take the Whitney sum

$$\zeta^{(0|1)} \stackrel{\text{def}}{=} \zeta_0 \times_M \zeta_1 \times_M \dots \times_M \zeta_\lambda \times_M \widehat{\Pi\zeta_0} \times_M \widehat{\Pi\zeta_1} \times_M \dots \times_M \widehat{\Pi\zeta_\lambda}$$

of the double set of dual bundles with opposite ghost parities of fibre coordinates. Finally, let us lift the Whitney sum of infinite jets of those bundles, putting it over the bundle of physical fields by using a pull-back under π_∞ . We denote the resulting bundle over the total space $J^\infty(\pi) \rightarrow M$ by

$$\pi_{\text{BV}}^{(0|1)} = \pi_\infty^*(\zeta_\infty^{(0|1)}).$$

The fibre $W_{\mathbf{x}} = V_{\mathbf{x}} \widehat{\oplus} \Pi V_{\mathbf{x}}^\dagger$ of $\zeta^{(0|1)}$ admits the canonical decomposition in two dual halves of opposite parities;¹⁷ this is shown in Fig. 5.




Figure 5. The BV-fibre is a direct sum of dual vector spaces; one is parity-even and the other is proclaimed ghost parity-odd.

Bearing in mind that the fields ϕ are artificially incorporated into the newly built fibre by ζ_0 , we shall omit an ever-present reference to points $(\mathbf{x}, \phi(\mathbf{x}))$ of jets of sections of the initial bundle π when dealing with variations $\delta \mathbf{s} = (\delta s; \delta s^\dagger)$ for sections s of $\zeta^{(0|1)}$ at $\phi(\mathbf{x})$, see Fig. 4.

¹⁶ By construction, the ghost parities of canonically conjugate BV-variables are complementary modulo 2, that is, to each even-parity variable q there corresponds its odd-parity dual neighbour q^\dagger . Of course, there remains much freedom in a choice of the integer ghost numbers followed by the group homomorphism $(-)^{\text{gh}(\cdot)}: \mathbb{Z} \rightarrow \mathbb{Z}_2$. For example, let (q, q^\dagger) be a pair of conjugate BV-variables; then one balances $\text{gh}(q) = \text{gh}(q^\dagger) \pm 1$ or $\text{gh}(q) = -\text{gh}(q^\dagger) \pm 1$, or by using any other integers such that one is even and the other is odd. Obviously any shift by an even integer (e. g., $\text{gh}(q) \mapsto -\text{gh}(q) = \text{gh}(q) - 2 \cdot \text{gh}(q)$) does not alter any values in the parity group \mathbb{Z}_2 ; this is no more than another way to describe the same theory.

¹⁷ To highlight this duality between ghost parity-even vector space $V_{\mathbf{x}}$ and ghost parity-odd subspace $\Pi V_{\mathbf{x}}^\dagger$ in $W_{\mathbf{x}}$, we use the notation $\widehat{\oplus}$ for their direct sum; whenever a coordinate in $V_{\mathbf{x}}$ is rescaled by const times, the respective conjugate variable in $\Pi V_{\mathbf{x}}^\dagger$ is transformed inverse-proportionally by const^{-1} times, see Remark 2.5 below.

2.2. The signs convention in Nature

The construction of canonically conjugate pairs of global coordinates $(\mathbf{q}, \mathbf{q}^\dagger)$ in the fibres $W_{\mathbf{x}} = V_{\mathbf{x}} \widehat{\oplus} \Pi V_{\mathbf{x}}^\dagger$ refers to a choice of the smooth field of dual bases in the two subspaces of even and odd ghost parity. Suppose that $\vec{e}_i(\mathbf{x})$ is a frame in $V_{\mathbf{x}}$ and $\vec{e}^{\dagger i}(\mathbf{x})$ is its dual in $\Pi V_{\mathbf{x}}^\dagger$, where the index i runs from 1 to the total dimension of even- and odd-parity component in the fibre of $\zeta^{(0|1)}$; we denote by $N = m + m_1 + \dots + m_\lambda$ each of the two dimensions so that the fibre of the Whitney sum $\zeta^{(0|1)}$ has superdimension $(N|N)$.

Let us recall that it is the parity of coordinates \mathbf{q}^\dagger but not of the vectors $\vec{e}^{\dagger i}$ in a basis which is reversed by the operation Π . The odd-parity component in the vector bundle $\zeta^{(0|1)}$ is topologically indistinguishable from $\widehat{\zeta}_0 \times_M \dots \times_M \widehat{\zeta}_\lambda$ but the rules become new for arithmetic in the algebra of coordinate functions on the total space. Therefore, we let the notation $\vec{e}^{\dagger i}(\mathbf{x})$ be identical for the same bases in $V_{\mathbf{x}}^\dagger$ and $\Pi V_{\mathbf{x}}^\dagger$.

Remark 2.2. The presence of *two* dual vector spaces, $V_{\mathbf{x}}$ and $(\Pi)V_{\mathbf{x}}^\dagger$, standardly implies that there are *two* couplings,

$$\langle \cdot, \cdot \rangle: V_{\mathbf{x}} \times (\Pi)V_{\mathbf{x}}^\dagger \rightarrow \mathbb{k} \quad \text{and} \quad \langle \cdot, \cdot \rangle: (\Pi)V_{\mathbf{x}}^\dagger \times V_{\mathbf{x}} \rightarrow \mathbb{k}; \quad (9)$$

we denote both operations in the same way because the order of arguments uniquely determines the choice. Let us remember also that it is not the linear vector space fibres of the superbundle $\zeta^{(0|1)}$ over the bundle π of physical fields but it is the tangent spaces

$$T_{(\mathbf{x}, \phi(\mathbf{x}), s(\mathbf{x}))} \left(V_{\mathbf{x}} \widehat{\oplus} \Pi V_{\mathbf{x}}^\dagger \right) \cong V_{\mathbf{x}} \widehat{\oplus} \Pi V_{\mathbf{x}}^\dagger$$

to those fibres which harbour the variations $\delta s = (\delta s; \delta s^\dagger)$ of sections s of the BV-bundle.

A reason to study the geometry of variations in tangent spaces to the fibres is clear from section 1. In fact, although we have substantiated in section 2.1 that Euler–Lagrange equations and their descendants do form linear vector spaces, this structure is incidental for the BV-formalism while Feynman path integration is not yet begun. The guiding geometric principle is that linear vector spaces appear only in the course of inspection of functionals' responses to infinitesimal test shifts of their arguments.

Couplings (9) are defined only if the linear vector spaces $V_{\mathbf{x}} \ni \delta s(\mathbf{x})$ and $\Pi V_{\mathbf{x}}^\dagger \ni \delta s^\dagger(\mathbf{x})$ are located over the same point $\mathbf{x} \in M^n$ of the base manifold, and over it they are attached as the two components of tangent space $T_{s(\mathbf{x})}(\zeta^{(0|1)})^{-1}(\mathbf{x}, \phi(\mathbf{x}))$, at the same point $s(\mathbf{x}) = s(\mathbf{x}, \phi(\mathbf{x}))$ of fibre in the superbundle $\zeta^{(0|1)}$ over a point $(\mathbf{x}, \phi(\mathbf{x}))$ of the total space for the bundle π of physical fields (see Fig. 4).

A distinction between the vector space $V_{\mathbf{x}}$ and its parity-reversed dual nontrivially determines the couplings' values whenever they are defined. Namely, each of the two finite-dimensional vector spaces is reflexive,

$$\left((V_{\mathbf{x}})^\dagger \right)^\dagger \cong V_{\mathbf{x}} \quad \text{and} \quad \left((\Pi V_{\mathbf{x}}^\dagger)^\dagger \right)^\dagger \cong \Pi V_{\mathbf{x}}^\dagger, \quad (10)$$

but these isomorphisms are not always identity mappings. We have that

$$\langle \vec{e}_i(\mathbf{x}), \vec{e}^{\dagger j}(\mathbf{x}) \rangle = \delta_i^j \quad \text{yet} \quad \langle \vec{e}^{\dagger j}(\mathbf{x}), \vec{e}_i(\mathbf{x}) \rangle = -\delta_i^j, \quad (11)$$

where δ_i^j is the Kronecker symbol whose value is the unit iff $i = j$ and which is set equal to zero otherwise.

Remark 2.3. We claim that this mechanism is responsible, in particular, for the skew-symmetry of various Poisson brackets (e.g., of the parity-odd Schouten bracket). Let us emphasize that this is a principle of order between geometric objects; the concept is not restricted to the BV-setup which we study here. Actually, Eq. (11) is the fundamental reason for differential 1-forms to anticommute¹⁸ (in the class of geometries for which a coupling is defined between the linear vector spaces of co-multiples under the wedge product \wedge ; for instance, such is the case of the Helmholtz criterion $\psi = \delta S / \delta \mathbf{q} \Leftrightarrow \bar{\ell}_\psi^{(\mathbf{q})} = (\bar{\ell}_\psi^{(\mathbf{q})})^\dagger$ for images of the variational derivative [28, 45]). Physically speaking, the binary count by “a vector space,” “not the former, hence its dual,” and “not the dual, but the initial space’s image under central symmetry” builds on the notion of order and realizes the law of the excluded middle.

2.3. Left- and right-variations via operators

Suppose that

$$S = \int \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \, \text{dvol}(\mathbf{x})$$

is an integral functional $\Gamma(\pi_{\text{BV}}) \rightarrow \mathbb{k}$. Let us focus on the correspondence between test shifts $\delta \mathbf{s} = (\delta s; \delta s^\dagger) = \delta s^i \cdot \bar{e}_i + \delta s_i^\dagger \cdot \bar{e}^{\dagger i}$ of BV-fields $s \in \Gamma(\pi_{\text{BV}})$ and, on the other hand, left- or right-acting linear singular integral operators $\overleftarrow{\delta} \mathbf{s}$ and $\overrightarrow{\delta} \mathbf{s}$ which yield the functional’s responses to shifts of its argument \mathbf{s} . By definition, we put

$$\begin{aligned} \overrightarrow{\delta} \mathbf{s} = \int_M \text{d}\mathbf{y} \left\{ (\delta s^i) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}} \right)^\sigma (\mathbf{y}) \cdot \left\langle (\bar{e}^{\dagger i})^\dagger(\mathbf{y}), \bar{e}^{\dagger j}(\cdot) \right\rangle \frac{\overrightarrow{\partial}}{\partial q_\sigma^j} + \right. \\ \left. + (\delta s_i^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}} \right)^\sigma (\mathbf{y}) \cdot \left\langle (\bar{e}_i)^\dagger(\mathbf{y}), \bar{e}_j(\cdot) \right\rangle \frac{\overrightarrow{\partial}}{\partial q_{j,\sigma}^\dagger} \right\} \quad (12a) \end{aligned}$$

and

$$\begin{aligned} \overleftarrow{\delta} \mathbf{s} = \int_M \text{d}\mathbf{y} \left\{ \frac{\overleftarrow{\partial}}{\partial q_\sigma^j} \left\langle \bar{e}^{\dagger j}(\cdot), \dagger(\bar{e}^{\dagger i})(\mathbf{y}) \right\rangle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}} \right)^\sigma (\delta s^i)(\mathbf{y}) + \right. \\ \left. + \frac{\overleftarrow{\partial}}{\partial q_{j,\sigma}^\dagger} \left\langle \bar{e}_j(\cdot), \dagger(\bar{e}_i)(\mathbf{y}) \right\rangle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}} \right)^\sigma (\delta s_i^\dagger)(\mathbf{y}) \right\}. \quad (12b) \end{aligned}$$

The above formulas for directed operators $\overrightarrow{\delta} \mathbf{s}$ and $\overleftarrow{\delta} \mathbf{s}$ contain new notation $(\bar{e}_i)^\dagger$, $(\bar{e}^{\dagger i})^\dagger$ and $\dagger(\bar{e}_i)$, $\dagger(\bar{e}^{\dagger i})$, also referring to an important sign convention which fully determines those adjoint objects. Namely, let us agree that over every $\mathbf{x} \in M^n$ the covectors

$$\bar{e}^{\dagger i}(\mathbf{x}) \left(\frac{\overrightarrow{\partial}}{\partial q_\sigma^i} \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_\mathbf{x}^\infty(\mathbf{s})} + \bar{e}_i(\mathbf{x}) \left(\frac{\overrightarrow{\partial}}{\partial q_{i,\sigma}^\dagger} \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_\mathbf{x}^\infty(\mathbf{s})}$$

and

$$\left(\mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_\sigma^i} \right) \Big|_{j_\mathbf{x}^\infty(\mathbf{s})} \bar{e}^{\dagger i}(\mathbf{x}) + \left(\mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{i,\sigma}^\dagger} \right) \Big|_{j_\mathbf{x}^\infty(\mathbf{s})} \bar{e}_i(\mathbf{x})$$

¹⁸ That is, this argument reveals why a mathematical axiom that differential forms do anticommute in the course of calculations leads to verifiable and relevant theoretic predictions which match experimental data.

are expanded in the cotangent space $T_{\mathbf{s}(\mathbf{x})}^* W_{\mathbf{x}} \cong V_{\mathbf{x}}^{\dagger} \hat{\oplus} (TV_{\mathbf{x}}^{\dagger})^{\dagger}$ with respect to the original basis $(+\vec{e}^{\dagger i}, +\vec{e}_i)$; note the signs (any other convention here would nohow alter the theory's content but it would (in)appropriately modify the signs in (13) below). The normalization of left- and right-adjoint objects $(\vec{e}_i)^{\dagger}$, $(\vec{e}^{\dagger i})^{\dagger}$ and $\dagger(\vec{e}_i)$, $\dagger(\vec{e}^{\dagger i})$ is immediate under assumption that the couplings' equations yield (5) and then (3) after integration by parts — no extra sign factors appear in those formulas. This requirement determines the table

$$\begin{aligned} (\vec{e}^{\dagger i})^{\dagger} &= \vec{e}_i, & \dagger(\vec{e}^{\dagger i}) &= -\vec{e}_i, \\ (\vec{e}_i)^{\dagger} &= -\vec{e}^{\dagger i}, & \dagger(\vec{e}_i) &= \vec{e}^{\dagger i}, \end{aligned} \tag{13}$$

so that the following defining relations hold:

$$\langle (\vec{e}_i)^{\dagger}, \vec{e}_i \rangle = \langle \vec{e}_i, \dagger(\vec{e}_i) \rangle = \langle (\vec{e}^{\dagger i})^{\dagger}, \vec{e}^{\dagger i} \rangle = \langle \vec{e}^{\dagger i}, \dagger(\vec{e}^{\dagger i}) \rangle = +1.$$

Let us notice that the left- and right-acting operation \dagger provides the analogue of left and right \langle, \rangle -dual in this ordered world; the first column in (13) determines a clockwise rotation in the oriented plane spanned by $\vec{e}_i \prec \vec{e}^{\dagger i}$, whereas taking the adjoints $\dagger(\cdot): \vec{e}_i \mapsto \vec{e}^{\dagger i}$ and $\vec{e}^{\dagger i} \mapsto -\vec{e}_i$ induces the counterclockwise rotation in that plane as shown in Fig. 6.




Figure 6. The orientation $\vec{e}_i \prec \vec{e}^{\dagger i}$ and configuration of the left- and right- duals with respect to the couplings \langle, \rangle .

Example 2.1. Identities (13) show up in the directed variations $\overleftarrow{\delta S} \Big|_s^{\delta \mathbf{s}} = \overrightarrow{\delta \mathbf{s}}(S)(s)$ and $\overrightarrow{\delta S} \Big|_s^{\delta \mathbf{s}} = (S) \overleftarrow{\delta \mathbf{s}}(s)$ of an integral functional $S = \int \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \cdot \text{dvol}(\mathbf{x})$. Namely, we have that

$$\begin{aligned} \overleftarrow{\delta S} \Big|_s^{(\delta s, \delta s^{\dagger})} &= \\ &= \int_M \text{d}\mathbf{y} \int_M \text{dvol}(\mathbf{x}) \left\{ (\delta s^i) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}} \right)^{\sigma} (\mathbf{y}) \langle \vec{e}_i(\mathbf{y}), \vec{e}^{\dagger j}(\mathbf{x}) \rangle \left(\frac{\overrightarrow{\partial}}{\partial q_{\sigma}^j} \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} + \right. \\ &\quad \left. + (\delta s_i^{\dagger}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}} \right)^{\sigma} (\mathbf{y}) \langle -\vec{e}^{\dagger i}(\mathbf{y}), \vec{e}_j(\mathbf{x}) \rangle \left(\frac{\overrightarrow{\partial}}{\partial q_{j, \sigma}^{\dagger}} \mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} \right\} \end{aligned} \tag{14a}$$

and

$$\begin{aligned} \overrightarrow{\delta S} \Big|_s^{(\delta s, \delta s^{\dagger})} &= \\ &= \int_M \text{d}\mathbf{y} \int_M \text{dvol}(\mathbf{x}) \left\{ \left(\mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \frac{\overleftarrow{\partial}}{\partial q_{\sigma}^j} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} \langle \vec{e}^{\dagger j}(\mathbf{x}), -\vec{e}_i(\mathbf{y}) \rangle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}} \right)^{\sigma} (\delta s^i)(\mathbf{y}) + \right. \\ &\quad \left. + \left(\mathcal{L}(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \frac{\overleftarrow{\partial}}{\partial q_{j, \sigma}^{\dagger}} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} \langle \vec{e}_j(\mathbf{x}), \vec{e}^{\dagger i}(\mathbf{y}) \rangle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}} \right)^{\sigma} (\delta s_i^{\dagger})(\mathbf{y}) \right\}. \end{aligned} \tag{14b}$$

The operators $\overrightarrow{\delta\mathbf{s}}$ and $\overleftarrow{\delta\mathbf{s}}$ act via ghost-parity graded Leibniz' rule on formal products of integral functionals (and on their images under other infinitesimal variation operators as well), so that the two operators are defined on the entire space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$, see section 2.4.2 below.

Remark 2.4. A reversion $\overleftarrow{\delta\mathbf{s}} \rightleftharpoons \overrightarrow{\delta\mathbf{s}}$ of the direction along $\overleftarrow{\delta\mathbf{s}}$ which such an operator acts means that the initially given operator (for definition, let it be $\overleftarrow{\delta\mathbf{s}}$ which acts to the left) is *destroyed* and in its place the other, opposite-direction operator is created (here it would be $\overrightarrow{\delta\mathbf{s}}$). Note that the variation $\delta\mathbf{s} \in \Gamma(T\pi)$ itself stays unchanged; it is the two realizations of this object via $\overleftarrow{\delta\mathbf{s}}$ and then via $\overrightarrow{\delta\mathbf{s}}$ which differ. (This concept of test shifts as primary geometric objects which contain information about the operators will be essential in Definition 2 of the variational Schouten bracket.)

Remark 2.5. The postulate of duality between $\vec{e}_i(\mathbf{x})$ and $\vec{e}^{\dagger i}(\mathbf{x})$ correlates their transformation laws under dilations: a rescaling $\vec{e}_i \mapsto \text{const} \cdot \vec{e}_i$ with $\text{const} \in \mathbb{k} \setminus \{0\}$ determines the inverse-proportional mapping $\vec{e}^{\dagger i} \mapsto \text{const}^{-1} \cdot \vec{e}^{\dagger i}$ of respective dual vectors. (Likewise, the coordinates in $V_{\mathbf{x}}$ and $\text{PIV}_{\mathbf{x}}^{\dagger}$ are then rescaled by $q^i \mapsto \text{const}^{-1} \cdot q^i$ and $q_i^{\dagger} \mapsto \text{const} \cdot q_i^{\dagger}$, respectively.)

Consider a variation $\delta\mathbf{s} = (\delta s; \delta s^{\dagger}) \in \Gamma(T\zeta^{(0|1)})$ of a BV-section $s \in \Gamma(\zeta^{(0|1)})$ over a given field configuration $\phi \in \Gamma(\pi)$ in the BV-bundle $\pi_{\text{BV}}^{(0|1)}$. The infinitesimal variation vectors $\delta\mathbf{s} = (\delta s; \delta s^{\dagger})$ can be naturally split to ghost parity-homogeneous components:

$$\delta\mathbf{s} = (\delta s; 0) + (0; \delta s^{\dagger}). \quad (15)$$

Here we explicitly use the linear vector space structure in fibres of the tangent bundle $T\zeta^{(0|1)}$. Let us recall that the two homogeneous variations

$$\delta s(\mathbf{x}) = \delta s^i(\mathbf{x}) \cdot \vec{e}_i(\mathbf{x}) \quad \text{and} \quad \delta s^{\dagger}(\mathbf{x}) = \delta s_i^{\dagger}(\mathbf{x}) \cdot \vec{e}^{\dagger i}(\mathbf{x})$$

in the right-hand side of (15) are the canonically dual to each other.

Moreover, by Remark 2.5 it is then possible to have δs and δs^{\dagger} normalized, for every i running from 1 to the dimension N , by the equalities

$$\delta s^i(\mathbf{x}) \cdot \delta s_i^{\dagger}(\mathbf{x}) \equiv +1 \quad (16)$$

at every $\mathbf{x} \in M^n$ where the smooth fields of dual bases \vec{e}_i and $\vec{e}^{\dagger i}$ are defined for the section s . From now on, let us deal only with such *normalized* variations. This implies that the coupling of these geometric objects are “invisible” but still the order in which the co-multiples δs and δs^{\dagger} occur in (11) does determine the signs in various formulas (e. g., in the definition of Schouten bracket, see p. 28 below).

2.4. Definitions of the BV-Laplacian and Schouten bracket

We now combine the geometry of graded-permutable iterated variations, which we explored in section 1 and which absorbs a new copy of the underlying base manifold M^n for each new infinitesimal test shift $\delta\mathbf{s}(\mathbf{x}) \in T_{s(\mathbf{x})}W_{\mathbf{x}}$ of the functionals' arguments at $\mathbf{x} \in M^n$, with the algebra of two couplings (9) between ghost parity-homogeneous halves of infinitesimal variations in the BV-setup $T_{s(\mathbf{x})}W_{\mathbf{x}} \cong V_{\mathbf{x}} \hat{\oplus} \text{PIV}_{\mathbf{x}}^{\dagger}$; the absolute locality of such coupling events is a fundamental principle.

To avoid an agglomeration of formulas and to match the notation with that in section 1, we omit an explicit reference to field configuration $\{\phi(\mathbf{x}), \mathbf{x} \in M^n\}$, indicating only the base points $\mathbf{x} \in M^n$. We also denote by π_{BV} the composite-structure superbundle over M^n (see Fig. 4) so that the notation for the vector bundle of BV-sections' infinitesimal variations is $T\pi_{\text{BV}}$. However,

let us remember that only the linear BV-fibre variables $(\mathbf{q}, \mathbf{q}^\dagger)$ but not the physical fields ϕ are subjected to variations at points $s(\mathbf{x}) \in (\zeta^{(0|1)})^{-1}(\mathbf{x}, s(\mathbf{x}))$ over $(\mathbf{x}, \phi(\mathbf{x})) \in \pi^{-1}(\mathbf{x})$. A brute force labelling of Euler–Lagrange equations by the respective unknowns is an act of will by the one who writes formulas but it is not a prescription from the model’s geometry.

This section contains rigorous, self-regularizing definitions of the BV-Laplacian and Schouten bracket for integral functionals from $\overline{H}^n(\pi_{\text{BV}}) \subsetneq \overline{\mathfrak{M}}^n(\pi_{\text{BV}}) \subsetneq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$. We shall extend the definition to the space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ of products of integral functionals, possibly with earlier-absorbed variations, in the subsequent sections of this paper. We then establish the main properties of these structures and prove relations between them. We note that the definitions which we give here are operational: each of them is a surgery for the couplings and their reconfiguration algorithm. (The locality postulate ensures the restrictions onto diagonals in the product $M \times \dots \times M$ so that those recombinations make sense at every point of M .)

2.4.1. The BV-Laplacian Δ Let us first introduce some shorthand notation. Let $F = \int f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \cdot \text{dvol}(\mathbf{x})$ be an integral functional and $\delta s = (\delta s; 0) + (0; \delta s^\dagger)$ be a variation’s splitting in two ghost parity-homogeneous variations. From section 1 we know that each of the two is referred to its own copy of the base: let it be $\delta s(\mathbf{y}_1)$ and $\delta s^\dagger(\mathbf{y}_2)$ so that formula (5) defines the response of F to an infinitesimal shift of its argument along each of the two directions.

Definition 1. Let $\delta s \in \Gamma(T\pi_{\text{BV}})$ be a test shift normalized by (16) and then split to the sum $(\delta s; 0) + (0; \delta s^\dagger)$ of ghost parity-homogeneous, \langle, \rangle -dual halves. The BV-Laplacian is the linear operator $\Delta: \overline{H}^n(\pi_{\text{BV}}) \rightarrow \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$; for a ghost parity-homogeneous integral functional $F \in \overline{H}^n(\pi)$ and its argument s , the operator Δ is an algorithm for reconfiguration of couplings in the second variation

$$\frac{\text{d}}{\text{d}\varepsilon} \Big|_{\varepsilon=0} \frac{\text{d}}{\text{d}\varepsilon^\dagger} \Big|_{\varepsilon^\dagger=0} F(s + \varepsilon \cdot \overleftarrow{\delta s} + \varepsilon^\dagger \cdot \overleftarrow{\delta s}^\dagger) = \sum_{\substack{i_1, i_2 \\ j_1, j_2}} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M \text{d}\mathbf{y}_1 \int_M \text{d}\mathbf{y}_2 \int_M \text{dvol}(\mathbf{x}) \left\{ \begin{array}{l} (\delta s^{i_1}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} (\mathbf{y}_1) \langle \overleftarrow{e}_{i_1}(\mathbf{y}_1), \overleftarrow{e}^{\dagger j_1}(\mathbf{x}) \rangle \leftarrow \\ \leftarrow (\delta s^\dagger_{i_2}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} (\mathbf{y}_2) \langle -\overleftarrow{e}^{\dagger i_2}(\mathbf{y}_2), \overleftarrow{e}_{j_2}(\mathbf{x}) \rangle \end{array} \right\} \cdot \frac{\overrightarrow{\partial}^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^{j_1} \partial q_{j_2, \sigma_2}^\dagger} \Big|_{j_x^\infty(s)}.$$

This second variation’s integrand contains the couplings \langle, \rangle :

$$T_{s(\mathbf{y}_1)} V_{\mathbf{y}_1} \times T_{s(\mathbf{x})}^* (\Pi) V_{\mathbf{x}} \rightarrow \mathbb{k} \quad \text{and} \quad T_{s(\mathbf{y}_2)} \Pi V_{\mathbf{y}_2}^\dagger \times T_{s(\mathbf{x})}^* (\Pi) V_{\mathbf{x}}^\dagger \rightarrow \mathbb{k}$$

which are defined only if the attachment points coincide for these (co)vectors; an optional presence of the parity reversion operator indicates a possibility of having ghost parity-odd functional F .

At the moment when the object ΔF under construction – or a larger object of which ΔF is an element, see (1c) – is evaluated at a section $s \in \Gamma(\pi_{\text{BV}})$, the integrations by parts carry the derivatives away from the variations’ components: $\overleftarrow{\partial} / \partial \mathbf{y}_i \mapsto \overrightarrow{\partial} / \partial \mathbf{y}_i$ as explained in section 1.3. The third step in definition of Δ acting on F is a surgery algorithm for an on-the-diagonal reattachment of the couplings, see Figure 7. In other words, *after* the integration by parts the

$$\begin{array}{ccc} \langle 1 | & | 3 \rangle & \\ & \langle 2 | & | 4 \rangle \end{array} \quad \longmapsto \quad \begin{array}{ccc} \langle 1 | & & \langle 3 | \\ & | 2 \rangle & | 4 \rangle \end{array}$$

Figure 7. The on-the-diagonal coupling of variations versus taking the trace of bi-linear form.

surgery yields the following:

$$\begin{aligned}
 (\Delta F)\Big|_s^{\delta s} &= \sum_{\substack{i_1, i_2 \\ j_1, j_2}} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M d\mathbf{y}_1 \int_M d\mathbf{y}_2 \int_M \text{dvol}(\mathbf{x}) \cdot \\
 &\cdot \left\{ \delta s^{i_1}(\mathbf{y}_1) \left(-\frac{\vec{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} \underbrace{\langle \vec{e}_{i_1}(\mathbf{y}_1), -\vec{e}^{\dagger i_2}(\mathbf{y}_2) \rangle}_{-1} \cdot \delta s_{i_2}^{\dagger}(\mathbf{y}_2) \left(-\frac{\vec{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} \right\} \\
 &\cdot \left\{ \underbrace{\langle \vec{e}^{\dagger j_1}(\mathbf{x}), \vec{e}_{j_2}(\mathbf{x}) \rangle}_{-1} \cdot \frac{\partial^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^{j_1} \partial q_{j_2, \sigma_2}^{\dagger}} \Big|_{j_{\mathbf{x}}^{\infty}(s)} \right\}. \quad (17)
 \end{aligned}$$

Note that the left-to-right order in $\langle \vec{e}_{i_1}(\mathbf{y}_1), \vec{e}^{\dagger j_1}(\mathbf{x}) \rangle \cdot \langle -\vec{e}^{\dagger i_2}(\mathbf{y}_2), \vec{e}_{j_2}(\mathbf{x}) \rangle$ is preserved by the respective couplings' arguments in $\langle \vec{e}_{i_1}(\mathbf{y}_1), -\vec{e}^{\dagger i_2}(\mathbf{y}_2) \rangle \cdot \langle \vec{e}^{\dagger j_1}(\mathbf{x}), \vec{e}_{j_2}(\mathbf{x}) \rangle$, cf. Fig. 7.

Remark 2.6. Until the moment when the integrations by parts are performed in ΔF , the derivatives $\partial/\partial \mathbf{y}_1$ and $\partial/\partial \mathbf{y}_2$ refer to different copies of the manifold M^n in the base $M^n \times M^n \times M^n$ of the product bundle $\pi_{\text{BV}} \times T\pi_{\text{BV}} \times T\pi_{\text{BV}}$. This implies that the two variations of F in the definition of Δ are graded-permutable between each other and with all other variations falling on $f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])$ whenever ΔF is a constituent element of a larger object (e. g., see (1c–1d) on p. 1).

Remark 2.7. To keep track of multiple copies of the base M^n for functionals and variations (here $\mathbf{x} \in M^n$, $\mathbf{y}_1 \in M^n$, $\mathbf{y}_2 \in M^n$) in the course of integration by parts (see section 1.3), we indicate the respective variations' bases by explicitly writing $\mathbf{q}(\mathbf{y}_1)$ and $\mathbf{q}^\dagger(\mathbf{y}_2)$ in the denominators and we denote by $\partial/\partial \mathbf{y}_1$ and $\partial/\partial \mathbf{y}_2$ the derivatives which now fall on the functional's density $f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])$ — for instance, we do so in Example 2.4 on p. 34 below. Namely, we put

$$\begin{aligned}
 \frac{\overleftarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta q^\alpha(\mathbf{y}_1)} \Big|_{j_{\mathbf{x}}^{\infty}(s)} &= \sum_{|\sigma_1| \geq 0} \left(-\frac{\vec{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} \left(\frac{\vec{\partial} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^\alpha} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} = \quad (18a) \\
 &= \sum_{|\sigma_1| \geq 0} \left(\left(-\frac{\vec{d}}{d\mathbf{y}_1} \right)^{\sigma_1} \frac{\vec{\partial} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^\alpha} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)}
 \end{aligned}$$

and

$$\begin{aligned}
 \frac{\overleftarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta q_{\beta}^{\dagger}(\mathbf{y}_2)} \Big|_{j_{\mathbf{x}}^{\infty}(s)} &= \sum_{|\sigma_2| \geq 0} \left(-\frac{\vec{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} \left(\frac{\vec{\partial} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\beta, \sigma_2}^{\dagger}} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)} = \quad (18b) \\
 &= \sum_{|\sigma_2| \geq 0} \left(\left(-\frac{\vec{d}}{d\mathbf{y}_2} \right)^{\sigma_2} \frac{\vec{\partial} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\beta, \sigma_2}^{\dagger}} \right) \Big|_{j_{\mathbf{x}}^{\infty}(s)}
 \end{aligned}$$

for the ghost parity-homogeneous components of variational derivative. At every point $(\mathbf{x}, \phi(\mathbf{x}), s(\mathbf{x}))$ of the total space for the bundle π_{BV} , and for a given functional F which is assumed ghost parity-homogeneous, we have that

$$\frac{\overleftarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta q^\alpha(\mathbf{y}_1)} \Big|_{j_{\mathbf{x}}^{\infty}(s)} \in T_{s(\mathbf{x})}^*(\Pi)V_{\mathbf{x}} \quad \text{and} \quad \frac{\overleftarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta q_{\beta}^{\dagger}(\mathbf{y}_2)} \Big|_{j_{\mathbf{x}}^{\infty}(s)} \in T_{s(\mathbf{x})}^*(\Pi)V_{\mathbf{x}}^{\dagger}.$$

Let us remember that an attribution of denominators to \mathbf{y}_1 or \mathbf{y}_2 is a matter of notation in (18); whenever happening, everything happens at $\mathbf{x} \in M^n$.

Lemma 1. The BV-Laplacian Δ is independent of a choice of the variation δs normalized by (16).

Indeed, whenever the integrations by parts are performed, products (16) of the dual components are always the same at all points of the intersection of their domains of definition.¹⁹ We illustrate the definition of BV-Laplacian Δ by using Fig. 8; let us notice that it properly




Figure 8. A variational update of the cyclic wor(1)d from [36]: the (anti)words δs and δs^\dagger are pasted into a necklace F according to the graded Leibniz rule. Then they annihilate in such a way that the respective loose ends of the string join, the cyclic order of gems preserved; this yields ΔF .

renders the assertion of Lemma 1 in a wider, noncommutative setup of [36] and [29, 32] (see Remark 1.1 on p. 6).

Corollary 2. In particular, we obtain the equality for immediate numeric value of ΔF at s . Namely, we have that

$$(\Delta F)(s) = \sum_{i_1, i_2} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M \text{dvol}(\mathbf{x}) \left\{ \delta s^{i_1}(\mathbf{y}_1) \cdot \delta s^{i_2}(\mathbf{y}_2) \cdot \left(\left(-\frac{\vec{d}}{d\mathbf{x}} \right)^{\sigma_1 \cup \sigma_2} \frac{\vec{\partial}^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^{i_1} \partial q_{i_2, \sigma_2}^\dagger} \right) \Big|_{j_{\mathbf{x}^\infty}(s)} \right\} \Big|_{\substack{\mathbf{y}_1 = \mathbf{x} \\ \mathbf{y}_2 = \mathbf{x}}} \in \mathbb{k}.$$

By taking one sum containing Kronecker's δ -symbol, one arrives at a conventional formula with a summation over the diagonal:

$$(\Delta F)(s) = \sum_{i=1}^N \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \int_M \text{dvol}(\mathbf{x}) \left(\left(-\frac{\vec{d}}{d\mathbf{x}} \right)^{\sigma_1 \cup \sigma_2} \frac{\vec{\partial}^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\partial q_{\sigma_1}^i \partial q_{i, \sigma_2}^\dagger} \right) \Big|_{j_{\mathbf{x}^\infty}(s)} \stackrel{\text{def}}{=} \sum_{i=1}^N \int_M \text{dvol}(\mathbf{x}) \frac{\overleftarrow{\delta}^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta q^i \delta q_i^\dagger}. \quad (19)$$

We refer to footnote 13 on p. 13 in this context.

¹⁹ The assertion of Lemma 1 extends to the variational Schouten bracket, which is a derivative structure with respect to the BV-Laplacian (see Definition 2 on p. 28). Moreover, the independence of a specific choice of variations implies that their coefficients $(\delta s_1, \delta s_1^\dagger)$ and $(\delta s_2, \delta s_2^\dagger)$, which are built into Δ and $\llbracket \cdot, \cdot \rrbracket$, can be swapped, not altering an object that contains these test shifts δs_1 and δs_2 (see the proof of Lemma 5 on p. 31).

Remark 2.8. The conventional formula

$$\frac{\overleftarrow{\delta}^2 f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta \mathbf{q}(\mathbf{y}_1) \delta \mathbf{q}^\dagger(\mathbf{y}_2)} \Bigg|_{\substack{\mathbf{y}_1 = \mathbf{x} \\ \mathbf{y}_2 = \mathbf{x}}}$$

itself is not the definition of a density of the BV-Laplacian ΔF for an integral functional $F = \int f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \cdot \text{dvol}(\mathbf{x})$. Not containing any built-in sources of divergence, the geometric definition and its implication (19) yield identical results only when one calculates the numeric value $(\Delta F)(s) \in \mathbb{k}$ — but not earlier: structurally different objects (17) and (19) belong to non-isomorphic spaces (so that the former contains more information than the latter), and their analytic behaviour is also different, see Example 2.4 on p. 34.

The following two examples are quoted from [35]; they show that the structure Δ defined above coincides — but only in the simplest situation — with the one which is intuitively known from the literature. We refer to the main Example 2.4 on p. 34 which illustrates the multiple-base geometry in a logically more complex situation of (1c).

Example 2.2. Take a compact, semisimple Lie group G with Lie algebra \mathfrak{g} and consider the corresponding Yang–Mills theory. Write A_i^a for the (coordinate expression of) the gauge potential A — a lower index i because A is a one-form on the base manifold (i. e., a covector), and an upper index a because A is a vector in the Lie algebra \mathfrak{g} of the Lie group G . Defining the field strength \mathcal{F} by $\mathcal{F}_{ij}^a = \partial_i A_j^a - \partial_j A_i^a + f_{bc}^a A_i^b A_j^c$ where f_{bc}^a are the structure constants of the Lie algebra \mathfrak{g} , the Yang–Mills action is²⁰

$$S_{\text{YM}} = \frac{1}{4} \int \mathcal{F}_{ij}^a \mathcal{F}^{a,ij} \, \text{d}^4x,$$

and the full BV-action S_{BV} is²¹

$$S_{\text{BV}} = S_{\text{YM}} + \int A_a^{i\dagger} \left(\frac{\text{d}}{\text{d}x^i} \gamma^a + f_{bc}^a A_i^b \gamma^c \right) \text{d}^4x - \frac{1}{2} \int f_{ab}^c \gamma^a \gamma^b \gamma_c^\dagger \text{d}^4x.$$

Let us calculate the BV-Laplacian of this functional. By Corollary 2, the only terms which survive in $\Delta(S_{\text{BV}})$ are those which contain both A and A^\dagger , or both γ and γ^\dagger . Therefore,

$$\begin{aligned} \Delta(S_{\text{BV}}) &= \int \left(\frac{\overleftarrow{\delta}}{\delta A_j^d} \frac{\overleftarrow{\delta}}{\delta A_d^{j\dagger}} (f_{bc}^a A_a^{i\dagger} A_i^b \gamma^c) - \frac{1}{2} \frac{\overleftarrow{\delta}}{\delta \gamma_d^\dagger} \frac{\overleftarrow{\delta}}{\delta \gamma^d} (f_{ab}^c \gamma^a \gamma^b \gamma_c^\dagger) \right) \text{d}^4x \\ &= \int \left(\frac{\overleftarrow{\delta}}{\delta A_j^d} (f_{bc}^d A_j^b \gamma^c) - \frac{1}{2} \frac{\overleftarrow{\delta}}{\delta \gamma_d^\dagger} (f_{db}^c \gamma^b \gamma_c^\dagger - f_{ad}^c \gamma^a \gamma_c^\dagger) \right) \text{d}^4x \\ &= \int \left(f_{dc}^d \gamma^c - \frac{1}{2} (f_{db}^d \gamma^b - f_{ad}^d \gamma^a) \right) \text{d}^4x = 0. \end{aligned}$$

Let us note also that, since the BV-action S_{BV} is by construction such that the horizontal cohomology class of $[[S_{\text{BV}}, S_{\text{BV}}]]$ is zero, as one easily checks by using Definition 2 below, the functional S_{BV} satisfies quantum master-equation (40) tautologically: both sides are, by independent calculations, equal to zero — should one inspect those values at any section s of the BV-bundle.

²⁰ The action functional S_{YM} is referred to Minkowski flat coordinates such that $\text{dvol}(\mathbf{x}) = \sqrt{|-1|} \text{d}^4x$ in the weak gauge field limit.

²¹ We denote by $A_a^{i\dagger}$ the parity-odd *antifields*, by γ^a the odd *ghosts*, and by γ_a^\dagger the parity-even *antighosts*.

Example 2.3. Consider the nonlinear Poisson sigma model introduced in [11]. Since its fields are not all purely even, we have to generalize all of our reasoning so far to a \mathbb{Z}_2 -graded setup — which is, as noted in Remark 2.1, tedious but straightforward. A verification that $\Delta(S_{\text{CF}})(s) = 0$ for the BV-action S_{CF} of this model and a section s of the respective BV-bundle would, up to minor differences in conventions and notations, proceed just as it does in that paper itself, in section 3.2 thereof — except that no infinite constants or Dirac’s δ -function appear.

Remark 2.9. The BV-Laplacian Δ is extended by using Leibniz’ rule from the space $\overline{H}^n(\pi_{\text{BV}})$ of building blocks in $\overline{\mathfrak{M}}^n(\pi_{\text{BV}})$ to the space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$, see Theorem 3 on p. 29. The couplings’ (re)attachment algorithm then results in formula (1b) on p. 1, which is taken as a *definition* of the variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$, see [39]. In turn, that structure’s extension from $\overline{H}^n(\pi_{\text{BV}}) \times \overline{H}^n(\pi_{\text{BV}})$ to $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \times \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ is immediate (see Theorem 4 below).

The correspondence between Δ and $\llbracket \cdot, \cdot \rrbracket$ is furthered to an equivalence between the property $\Delta^2 = 0$ of BV-Laplacian to be a differential and, on the other hand, Jacobi’s identity for the variational Schouten bracket. We emphasize that the latter *can be* verified within the old approach [41] to geometry of variations. (We refer to [32] for a proof; its crucial idea is that with evolutionary vector fields it does not matter under “whose” total derivatives, $d/d\mathbf{x}$ or $d/d\mathbf{y}_i$, such fields dive.) Nevertheless, the traditional paradigm fails to reveal that the operator Δ is a differential because of a necessity to have the variations graded-permutable and for that, to distinguish between the functionals’ and variations’ domains of definition. Our geometric approach resolves that obstruction and ensures the validity of identities (1c) and (1d) (see Theorems 6 and 8, respectively).

2.4.2. The variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$ The parity-odd Laplacian Δ is the parent object²² which induces the variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$. Namely, the bracket appears in the course of that operator’s extension from the space $\overline{H}^n(\pi_{\text{BV}}) \ni F$ to the space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \supseteq \overline{\mathfrak{M}}^n(\pi_{\text{BV}})$ of local functionals $F_1 \cdots F_\ell$ (it is possible that F_i ’s already contain some normalized variations).

A distinction between *left* and *right* in the directed operators $\overleftarrow{\delta}\mathbf{s}$ and $\overrightarrow{\delta}\mathbf{s}$, the orientation $\vec{e}_i \prec \vec{e}^{\dagger i}$ in the composite BV-fibres $W_{\mathbf{x}} \cong V_{\mathbf{x}} \hat{\oplus} \Pi V_{\mathbf{x}}^{\dagger}$ equipped with two couplings (9), and the ordering of variations $\delta\mathbf{s}_1, \dots, \delta\mathbf{s}_k$ specify the logic of operational Definition 2, which is given in this section.

Remark 2.10. For the sake of brevity, we extend the BV-Laplacian Δ from the space $\overline{H}^n(\pi_{\text{BV}})$ of integral functionals F_1, \dots, F_ℓ to the space $\overline{\mathfrak{M}}^n(\pi_{\text{BV}})$ of local functionals such as $F_1 \cdots F_\ell$, the factors of which do not explicitly contain any built-in variations. To further this extension verbatim onto the full space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \supseteq \overline{\mathfrak{M}}^n(\pi_{\text{BV}})$, one must remember that it is forbidden to break the order in which the directed variation operators $\overrightarrow{\delta}\mathbf{s}_k$ and $\overleftarrow{\delta}\mathbf{s}_k$ appear in the (ordered collection of) objects at hand. (Such concept is illustrated by the third term in (20) below.)

Likewise, we extend Δ to products of just two factors; in the case of arbitrary number $\ell \geq 2$ of building blocks F_1, \dots, F_ℓ one proceeds inductively by using the ghost parity-graded Leibniz rule, then extending Δ onto the vector space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ by linearity.

Let $F = \int f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \text{dvol}(\mathbf{x}_1)$ and $G = \int g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^{\dagger}]) \text{dvol}(\mathbf{x}_2)$ be integral functionals $\Gamma(\pi_{\text{BV}}) \rightarrow \mathbb{k}$ and let $\delta\mathbf{s} = (\delta s; \delta s^{\dagger})$ be a normalized test shift of their product’s argument $s \in \Gamma(\pi_{\text{BV}})$. We now define the operator Δ acting on the element $F \cdot G$ at \mathbf{s} by variations first along $(0; \delta s^{\dagger})$ and then along $(\delta s; 0)$.

²² In particular, the definition of BV-Laplacian logically precedes the construction of Schouten bracket in BV-formalism (although such parity-odd variational Poisson bracket is often introduced through postulated formula (25) in the context of Hamiltonian dynamics and infinite-dimensional completely integrable systems [14, 18, 26, 41, 42]). Indeed, the entire Schouten-bracket machinery of (quantum) BV-cohomology groups and their automorphisms, which we consider in section 3.2, stems from quantum master-equation (40), see p. 41.

According to (14), the object to start with is

$$\int_M d\mathbf{y}_1 \int_M d\mathbf{y}_2 \sum_{\substack{i_1, i_2 \\ j_1, j_2}} \sum_{\substack{|\sigma_1| \geq 0 \\ |\sigma_2| \geq 0}} \left\{ (\delta s^{i_1}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} (\mathbf{y}_1) \langle \vec{e}_{i_1}(\mathbf{y}_1), \vec{e}^{\dagger j_1}(\cdot) \rangle \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^{j_1}} \circ \right. \\ \left. \circ (\delta s_{i_2}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} (\mathbf{y}_2) \langle -\vec{e}^{\dagger i_2}(\mathbf{y}_2), \vec{e}_{j_2}(\cdot) \rangle \frac{\overrightarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger} \right\} \\ \left(\int f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \text{dvol}(\mathbf{x}_1) \cdot \int g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \text{dvol}(\mathbf{x}_2) \right) (s).$$

Their order preserved, the directed operators $\overrightarrow{\delta s}$ and $\overrightarrow{\delta s}^\dagger$ spread over the two factors F and G by the binomial formula because of the Leibniz rule for graded derivations $\overrightarrow{\partial}/\partial q_{\sigma_1}^{j_1}$ and $\overrightarrow{\partial}/\partial q_{j_2, \sigma_2}^\dagger$. Note that whenever the ghost parity-odd object $\overrightarrow{\partial}/\partial q_{j_2, \sigma_2}^\dagger$ overtakes the density f of ghost parity $\text{gh}(F)$, there appears an overall sign factor $(-)^{\text{gh}(F)}$. We thus obtain

$$(\overrightarrow{\delta s} \circ \overrightarrow{\delta s}^\dagger)(F) + (-)^{\text{gh}(F)} \overrightarrow{\delta s}(F) \overrightarrow{\delta s}^\dagger(G) + \overrightarrow{\delta s} \xrightarrow{\cdot \overrightarrow{\delta s}^\dagger(F)} G \\ + (-1)^{\text{gh}(F)} F \cdot (\overrightarrow{\delta s} \circ \overrightarrow{\delta s}^\dagger)(G). \quad (20)$$

The next step is to push right through F its single variations in the middle two terms of the above expression. This yields the equality

$$= (\overrightarrow{\delta s} \circ \overrightarrow{\delta s}^\dagger)(F) \cdot G + (-)^{\text{gh}(F)} \left\{ (F) \overleftarrow{\delta s} \cdot \overrightarrow{\delta s}^\dagger(G) + \overrightarrow{\delta s} \xrightarrow{\cdot (F) \overleftarrow{\delta s}^\dagger} G \right\} \\ + (-)^{\text{gh}(F)} F \cdot (\overrightarrow{\delta s} \circ \overrightarrow{\delta s}^\dagger)(G). \quad (21)$$

We emphasize that the operators $\overleftarrow{\delta s}$ and $\overleftarrow{\delta s}^\dagger$ in the variations $(F) \overleftarrow{\delta s} = \overrightarrow{\delta_{\mathbf{q}} F}$ and $(F) \overleftarrow{\delta s}^\dagger = \overrightarrow{\delta_{\mathbf{q}^\dagger} F}$ are temporarily redirected to the left so that the middle terms in (21) are $(-)^{\text{gh}(F)}$ times

$$\overrightarrow{\delta_{\mathbf{q}} F} \cdot \overleftarrow{\delta_{\mathbf{q}^\dagger} F} + \overleftarrow{\delta_{\mathbf{q}} F} \xrightarrow{\cdot \overrightarrow{\delta_{\mathbf{q}^\dagger} F}} \overrightarrow{G}; \quad (22)$$

this is the input datum for a traditional definition of the variational Schouten bracket (e. g., see [11] vs [39]). Let us remember that the BV-fibres orientation $\mathbf{q} \prec \mathbf{q}^\dagger$ expressed by (9) is built into the last term of (22) even if it is written as follows,

$$(F) \overleftarrow{\delta_{\mathbf{q}}} \cdot \overrightarrow{\delta_{\mathbf{q}^\dagger}}(G) + (F) \overleftarrow{\delta_{\mathbf{q}^\dagger}} \cdot \overrightarrow{\delta_{\mathbf{q}}}(G).$$

Should this be the notation for input, one then usually proclaims that “differential 1-forms anticommute” so that $\langle \delta \mathbf{q}^\dagger \wedge \delta \mathbf{q} \rangle = -\langle \delta \mathbf{q} \wedge \delta \mathbf{q}^\dagger \rangle = -1$ in $\llbracket F, G \rrbracket = \langle \overrightarrow{\delta F} \wedge \overleftarrow{\delta G} \rangle$.

We now are almost in a position to (re)configure the couplings in the four terms of (21). The first term will of course become $\Delta F \cdot G$, and the last will provide $(-)^{\text{gh}(F)} F \cdot \Delta G$; one is here allowed to integrate by parts (as explained in section 1.3) in order to shake the derivatives off δs^{i_1} and $\delta s_{i_2}^\dagger$ prior to evaluation of couplings in the resulting object’s numeric value at its argument s . Yet there remains one more logical step to be done with (22): let us reverse back $\overleftarrow{\delta s} \mapsto \overrightarrow{\delta s}$ and $\overleftarrow{\delta s}^\dagger \mapsto \overrightarrow{\delta s}^\dagger$ so that on one hand, the vertical differentials fall on F but on the other hand, the normalization of the basis which stands near $\delta s^i(\mathbf{y}_1)$ and $\delta s_i^\dagger(\mathbf{y}_2)$ is the *first* not second column

in (13). This yields the following integrand of $(-)^{\text{gh}(F)} \int_M d\mathbf{y}_1 \int_M d\mathbf{y}_2 \int_M d\text{vol}(\mathbf{x}_1) \int_M d\text{vol}(\mathbf{x}_2)$, with a summation over i_1, i_2, j_1, j_2 , and $|\sigma_1| \geq 0, |\sigma_2| \geq 0$,

$$\begin{aligned}
 & \left(f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\sigma_1}^{j_1}} \right) \Big|_{j_{\mathbf{x}_1}^\infty(s)} \langle \overleftarrow{e}^{\dagger j_1}(\mathbf{x}_1), +\overleftarrow{e}_{i_1}(\mathbf{y}_1) \rangle \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} (\delta s^{i_1})(\mathbf{y}_1) \cdot \\
 & \cdot (\delta s_{i_2}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} (\mathbf{y}_2) \langle -\overleftarrow{e}^{\dagger i_2}(\mathbf{y}_2), \overleftarrow{e}_{j_2}(\mathbf{x}_2) \rangle \left(\frac{\overleftarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_{\mathbf{x}_2}^\infty(s)} + \\
 & + \left(f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{j_1, \sigma_1}^\dagger} \right) \Big|_{j_{\mathbf{x}_1}^\infty(s)} \cdot \\
 & \left\{ \begin{array}{l} (\delta s^{i_1}) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_2} (\mathbf{y}_1) \langle \overleftarrow{e}_{i_1}(\mathbf{y}_1) | \\ \langle \overleftarrow{e}_{j_1}(\mathbf{x}_1) | \end{array} \right. \left. \begin{array}{l} | \overleftarrow{e}^{\dagger j_2}(\mathbf{x}_2) \rangle \rangle \\ | -\overleftarrow{e}^{\dagger i_2}(\mathbf{y}_2) \rangle \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_1} (\delta s_{i_2}^\dagger)(\mathbf{y}_2) \rangle \end{array} \right\} \\
 & \left(\frac{\overleftarrow{\partial}}{\partial q_{\sigma_2}^{j_2}} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_{\mathbf{x}_2}^\infty(s)} \cdot
 \end{aligned} \tag{23}$$

The integrations by parts are performed and couplings are reconfigured at the end of the day in exactly same manner as it has been done in Definition 1; let us recall that we now define the BV-Laplacian on a larger space. Namely, the variations couple with the dual variations whereas the differentials of functionals' densities attach to each other.

Definition 2. The *variational Schouten bracket* of two integral functionals

$$F = \int f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \cdot d\text{vol}(\mathbf{x}_1) \quad \text{and} \quad G = \int g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \cdot d\text{vol}(\mathbf{x}_2)$$

is the on-the-diagonal couplings surgery which, by using a normalized test shift $\delta s = (\delta s; 0) + (0; \delta s^\dagger) \in \Gamma(T\pi_{\text{BV}})$, yields the functional from $\mathfrak{N}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ whose construction at a BV-section $s \in \Gamma(\pi_{\text{BV}})$ is²³

$$\begin{aligned}
 & \int_M d\mathbf{y}_1 \int_M d\mathbf{y}_2 \int_M d\mathbf{x}_1 \int_M d\text{vol}(\mathbf{x}_2) \left[\left(f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\sigma_1}^{j_1}} \right) \Big|_{j_{\mathbf{x}_1}^\infty(s)} \right. \\
 & \left. \left\{ \underbrace{\left(-\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} \delta s^{i_1}(\mathbf{y}_1) \langle \overleftarrow{e}_{i_1}(\mathbf{y}_1), -\overleftarrow{e}^{\dagger i_2}(\mathbf{y}_2) \rangle \delta s_{i_2}^\dagger(\mathbf{y}_2) \left(-\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2}}_{-1}, \right. \right. \\
 & \left. \left. \underbrace{\langle \overleftarrow{e}^{\dagger j_1}(\mathbf{x}_1) |}_{-1}, \left. \right. \left. \left(\frac{\overleftarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_{\mathbf{x}_2}^\infty(s)} \right\} \right. \\
 & \left. \left. \underbrace{|\overleftarrow{e}_{j_2}(\mathbf{x}_2) \rangle}_{-1} \right\} \right.
 \end{aligned}$$

²³ Note that the directions of $\partial/\partial \mathbf{y}_i$ are reversed so that the minus signs appear. We emphasize that, prior to the evaluation of reconfigured couplings, the (co)vectors at \mathbf{x}_j channel the partial derivatives to f or g according to the couplings' old arrangement.

$$\begin{aligned}
 & + \left(f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{j_1, \sigma_1}^\dagger} \right) \Big|_{j_{\mathbf{x}_1}^\infty(s)} \\
 & \underbrace{\left\{ \frac{\delta s^{i_1}(\mathbf{y}_1)}{\langle \vec{e}_{j_1}(\mathbf{x}_1) |} \left(-\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_2} \langle \vec{e}_{i_1}(\mathbf{y}_1), (-\vec{e}^{\dagger i_2})(\mathbf{y}_2) \rangle \cdot \left(-\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_1} \delta s_{i_2}^\dagger(\mathbf{y}_2) \right\}}_{+1} \underbrace{\left. \frac{\delta s_{i_2}^\dagger(\mathbf{y}_2)}{|\vec{e}^{\dagger j_2}(\mathbf{x}_2)\rangle} \right\}}_{-1} \\
 & \left(\frac{\overrightarrow{\partial}}{\partial q_{\sigma_2}^{j_2}} g(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \Big|_{j_{\mathbf{x}_2}^\infty(s)}.
 \end{aligned}$$

Note that the inner couplings between variations provide a restriction to the diagonal $\mathbf{y}_1 = \mathbf{y}_2$ and yield the singular integral operators which then act to the right via multiplication by -1 only if $\mathbf{y}_2 = \mathbf{x}_2$ and $\mathbf{y}_1 = \mathbf{x}_1$, respectively. The outer coupling then furnishes the main diagonal $\mathbf{x}_1 = \mathbf{y}_1 = \mathbf{y}_2 = \mathbf{x}_2$, restricting the objects further to the same BV-fibre point in the total space of the BV-bundle. This reveals why over each point of the base M^n the (derivatives of the) densities f and g are restricted to the infinite jet of the same section s ; this also means that, since the moment when the couplings are reconfigured, the volume element $\text{dvol}(\mathbf{x}_1)$ is discarded because appears a new singular linear integral operator with a standard sign $\int d\mathbf{x}_1$.

We conclude the reasoning and sum up the definitions and notations in the following theorem.

Theorem 3. *The BV-Laplacian Δ is the linear operator*

$$\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \times \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \rightarrow \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$$

which acts on products of functionals F and $G \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ by the rule

$$\Delta(F \cdot G) = \Delta(F) \cdot G + (-)^{\text{gh}(F)} \llbracket F, G \rrbracket + (-)^{\text{gh}(F)} F \cdot \Delta G. \tag{24}$$

The variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$ measures the deviation for the BV-Laplacian Δ from being a derivation.

- After integration by parts, Definition 2 implies the renowned coordinate formula

$$\begin{aligned}
 \llbracket F, G \rrbracket = \int \text{dvol}(\mathbf{x}) \left(\frac{\overrightarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta \mathbf{q}} \cdot \frac{\overleftarrow{\delta} g(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta \mathbf{q}^\dagger} - \right. \\
 \left. - \frac{\overrightarrow{\delta} f(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta \mathbf{q}^\dagger} \cdot \frac{\overleftarrow{\delta} g(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger])}{\delta \mathbf{q}} \right). \tag{25}
 \end{aligned}$$

Remark 2.11. Let us recall from Remark 1.5 that the building blocks of local functionals are encoded by equivalence classes of their densities, whereas the underlying integration manifold M^n is endowed with the field-dependent volume element $\text{dvol}(\mathbf{x}, \phi)$. The variational Schouten bracket transforms two given integral functionals F and G into $\llbracket F, G \rrbracket$. For every configuration of physical fields $\phi \in \Gamma(\pi)$, the integration measure is the same in F , G , and $\llbracket F, G \rrbracket$. This is because the couplings are local over points $(\mathbf{x}, \phi(\mathbf{x}))$ in the total space of the bundle π of physical fields, see Remark 2.2 on p. 18; the equality of local sections ϕ at which all (derivatives of) functionals' densities are evaluated ensures the equality of metric tensor elements $g_{\mu\nu}$ in all functionals by virtue of Einstein's general relativity equations.

The operational definition of the *antibracket* $\llbracket \cdot, \cdot \rrbracket$ determines the way how this structure acts on the square $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \times \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ of entire space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ containing formal products of functionals.

Theorem 4. Let $F, G,$ and $H \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be ghost parity-homogeneous functionals. The variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket: \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \times \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \rightarrow \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ has the following properties:

(i) The value of $\llbracket \cdot, \cdot \rrbracket$ at two arguments F and $G \cdot H$ is

$$\llbracket F, G \cdot H \rrbracket = \llbracket F, G \rrbracket \cdot H + (-)^{\text{gh}(F)-1} \text{gh}(G) G \cdot \llbracket F, H \rrbracket. \quad (26)$$

This formula recursively extends to products of arbitrary finite number of factors in the second argument.

(ii) The bracket $\llbracket \cdot, \cdot \rrbracket$ is shifted-graded skew-symmetric:

$$\llbracket F, G \rrbracket = -(-)^{\text{gh}(F)-1 \cdot \text{gh}(G)-1} \llbracket G, F \rrbracket, \quad (27)$$

which extends $\llbracket \cdot, \cdot \rrbracket$ to products of arbitrary finite number of factors taken as its first argument in (26).

(iii) The bracket $\llbracket \cdot, \cdot \rrbracket$ satisfies the shifted-graded Jacobi identity

$$\begin{aligned} (-)^{\text{gh}(F)-1} \text{gh}(H)-1 \llbracket F, \llbracket G, H \rrbracket \rrbracket + (-)^{\text{gh}(F)-1} \text{gh}(G)-1 \llbracket G, \llbracket H, F \rrbracket \rrbracket + \\ + (-)^{\text{gh}(G)-1} \text{gh}(H)-1 \llbracket H, \llbracket F, G \rrbracket \rrbracket = 0, \end{aligned} \quad (28)$$

which stems from graded Leibniz rule (36) for evolutionary vector fields \mathbf{Q}^F defined by the rule $\mathbf{Q}^F(\cdot) \cong \llbracket F, \cdot \rrbracket$ (here the equivalence up to integration by parts is denoted by \cong).

Finally, the variational Schouten bracket extends by linearity to formal sums of elements from $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$.

Proof. The bilinearity of $\llbracket \cdot, \cdot \rrbracket$ is obvious. It is also clear that the terms in $\llbracket F, G \cdot H \rrbracket$ are grouped in two parts: those in which the ghost-parity graded derivations $\overrightarrow{\partial} / \partial \mathbf{q}^\dagger$ act on G and those for H ; the former do not contribute with any extra sign factors whereas the latter do — in a way which depends on the parity $\text{gh}(G)$. This means that $\llbracket F, G \cdot H \rrbracket = \llbracket F, G \rrbracket \cdot H + \dots$; to grasp the sign in front of the term which has been omitted, let us swap the graded multiples G and H . We have that $G \cdot H = (-)^{\text{gh}(G) \text{gh}(H)} H \cdot G$, whence $\llbracket F, G \cdot H \rrbracket = (-)^{\text{gh}(G) \text{gh}(H)} \llbracket F, H \rrbracket \cdot G + \dots$. By recalling that $\text{gh}(\llbracket F, H \rrbracket) = \text{gh}(F) + \text{gh}(H) - 1$, we conclude that

$$\llbracket F, G \cdot H \rrbracket = \llbracket F, G \rrbracket \cdot H + (-)^{\text{gh}(G) \text{gh}(H)} (-)^{\text{gh}(F) + \text{gh}(H) - 1} \text{gh}(G) G \cdot \llbracket F, H \rrbracket,$$

which yields formula (26).

Proving (27) amounts to a count of signs whenever the bracket $\llbracket F, G \rrbracket$ of an ordered pair of ghost parity-graded objects is virtually transformed into $\llbracket G, F \rrbracket$. By using the rule of signs for odd-parity coordinates, $q_{\alpha, \sigma}^\dagger \cdot q_{\beta, \tau}^\dagger = -q_{\beta, \tau}^\dagger \cdot q_{\alpha, \sigma}^\dagger$, we first note that

$$\frac{\overrightarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) = (-)^{\text{gh}(G)-1} \left(g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right) \frac{\overleftarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger},$$

with a similar formula for the left- and right-acting graded derivative of f . By swapping the (variational) derivatives of the densities f and g , we gain the signs $(-)^{\text{gh}(F) \cdot \text{gh}(G)-1}$ and $(-)^{\text{gh}(F)-1 \cdot \text{gh}(G)}$ for the respective terms in (23) on p. 28. Combined together, the two steps accumulate equal factors $(-)^{\text{gh}(F)+1 \cdot \text{gh}(G)-1} = (-)^{\text{gh}(F)-1 \cdot \text{gh}(G)+1} = (-)^{\text{gh}(F)-1 \cdot \text{gh}(G)-1}$. Thirdly, by comparing $(-)^{\text{gh}(F)-1 \cdot \text{gh}(G)-1} \llbracket F, G \rrbracket$ – in which the derivatives of f and g are interchanged and the derivations' directions are reversed – with $\llbracket G, F \rrbracket$, we conclude that the

reconfiguration of couplings in the second term in (23) for the former expression yields *minus* the first term in $\llbracket G, F \rrbracket$. Likewise, the couplings reattachment in the first term of such (23) produces minus the second term in $\llbracket G, F \rrbracket$. This is because the (co)vectors in the differentials of densities remain unswapped, now going in the ‘wrong’ order.

We now refer to [32, Proposition 3] for a proof of property (iii) in a wider, non-commutative setup of cyclic words (cf. [29, 36, 46]). It is remarkable that the reasoning persists within a naïve theory of variations, not referring to our main idea that each test shift brings its own copy of the base M^n into the picture. A key point in the proof is that the rule $\mathbf{Q}^F(\cdot) \cong \llbracket F, \cdot \rrbracket$ naturally associates with functionals F the evolutionary fields \mathbf{Q}^F on the infinite jet superbundles at hand, and with *evolutionary* vector fields it does not matter under ‘whose’ total derivatives such fields dive, obeying their defining property $[\mathbf{Q}^F, \overrightarrow{d}/d\mathbf{x}] = 0$ (i.e., any integrations by parts, which transform the derivatives $\overrightarrow{\partial}/\partial\mathbf{y}_i$ falling on test shifts into total derivatives $\overrightarrow{d}/d\mathbf{x}$ falling on the functionals’ densities, do not mar the outcome even if one attempts to perform such integrations ahead of time). \square

2.5. *Main result: the proof of properties (1c–1d)*

We are ready to *prove* the main interrelations between the BV-Laplacian Δ and variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$. Let us recall that either a validity of these properties was postulated (see [21]) or an *ad hoc* regularization technique was formally employed in the literature in order to mask the seemingly present divergencies (which are actually not there), cf. [22, §15].

Let us fix the terms. In what follows we refer to building blocks from $\overline{H}^n(\pi_{\text{BV}})$ and their descendants – containing reconfigured variations – from $\overline{H}^{n(1+k)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$ as *integral functionals*. Such objects will be used for bases of inductive proofs of Lemmas 5 and 7. We then extend the properties (1c) and $\Delta^2 = 0$ to the space $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}) \supseteq \overline{\mathfrak{M}}^n(\pi_{\text{BV}})$ of *local functionals*, that is, of formal sums of products of (varied descendants of) building blocks.

Lemma 5. Let $F \in \overline{H}^{n(1+k)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$ and $G \in \overline{H}^{n(1+\ell)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$ be two integral functionals; here $k, \ell \geq 0$. Then

$$\Delta(\llbracket F, G \rrbracket) = \llbracket \Delta F, G \rrbracket + (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket. \tag{29}$$

Proof. The key idea is that the structures Δ and $\llbracket \cdot, \cdot \rrbracket$ yield equivalence classes of integral functionals which, after an integration by parts at the end of the day, are *independent* of a choice of the built-in test shifts normalized by (16). Consequently, the composite structure $\Delta(\llbracket \cdot, \cdot \rrbracket)$ does not change under swapping $\delta s_1^\alpha \rightleftharpoons \delta s_2^\beta$, $\delta s_{1,\alpha}^\dagger \rightleftharpoons \delta s_{2,\beta}^\dagger$ of the respective variations $\delta\mathbf{s}_1$ and $\delta\mathbf{s}_2$ in Δ and $\llbracket \cdot, \cdot \rrbracket$. Hence the terms which are skew-symmetric under such exchange necessarily vanish.

For the sake of clarity, let us assume that $F = \int f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \text{dvol}(\mathbf{x}_1)$ and $G = \int g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \text{dvol}(\mathbf{x}_2)$ are just building blocks from the cohomology group $\overline{H}^n(\pi_{\text{BV}})$; this simplification is legitimate because new variations which come from Δ and $\llbracket \cdot, \cdot \rrbracket$ do not interfere with any other test shifts if those are already absorbed by the densities f and g . Suppose that $\delta\mathbf{s}_1$ and $\delta\mathbf{s}_2$ are two normalized variations of a section $s \in \Gamma(\pi_{\text{BV}})$. By definition, we have that²⁴

$$\Delta(\llbracket F, G \rrbracket)(s) = \int_M \text{dz}_1 \int_M \text{dz}_2 \int_M \text{dy}_1 \int_M \text{dy}_2 \int_M \text{dx}_1 \int_M \text{dvol}(\mathbf{x}_2) \cdot \left\{ (\delta s_1^\alpha) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1}(\mathbf{z}_1) \left\langle \vec{e}_\alpha(\mathbf{z}_1), -\vec{e}^{\dagger\alpha}(\mathbf{z}_2) \right\rangle (\delta s_{1,\alpha}^\dagger) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2}(\mathbf{z}_2) \left\langle \vec{e}^{\dagger\alpha}(\cdot), \vec{e}_\alpha(\cdot) \right\rangle \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \right.$$

²⁴ To keep track of their origin, we let the directed derivatives $\partial/\partial\mathbf{y}_i$ or $\partial/\partial\mathbf{z}_j$ remain falling on the respective coefficients in $\delta\mathbf{s}_1$ and $\delta\mathbf{s}_2$; the integration by parts is performed in a standard way prior to the reconfigurations which are shown in the formula.

$$\left[\begin{aligned} & f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\tau_1}^\beta} \langle \overleftarrow{e}^{\dagger\beta}(\mathbf{x}_1) | \left\langle \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\delta s_2^\beta)(\mathbf{y}_1) \overleftarrow{e}_\beta(\mathbf{y}_1), \right. \\ & \quad \left. - \overleftarrow{e}^{\dagger,\beta}(\mathbf{y}_2) (\delta s_{2,\beta}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\mathbf{y}_2) \right\rangle \underline{|\overleftarrow{e}_\beta(\mathbf{x}_2)\rangle} \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) + \\ & + f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \langle \overleftarrow{e}_\beta(\mathbf{x}_1) | \left\langle (\delta s_2^\beta) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\mathbf{y}_1) \overleftarrow{e}_\beta(\mathbf{y}_1), \right. \\ & \quad \left. - \overleftarrow{e}^{\dagger,\beta}(\mathbf{y}_2) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\delta s_{2,\beta}^\dagger)(\mathbf{y}_2) \right\rangle \underline{|\overleftarrow{e}^{\dagger\beta}(\mathbf{x}_2)\rangle} \frac{\overrightarrow{\partial}}{\partial q_{\tau_1}^\beta} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right] \Bigg|_{\substack{j^\infty(s) \\ \mathbf{x}_i = \mathbf{y}_j = \mathbf{z}_k}} . \end{aligned} \right.$$

The partial derivatives $\overrightarrow{\partial}/\partial q_{\sigma_1}^\alpha \circ \overrightarrow{\partial}/\partial q_{\alpha,\sigma_2}^\dagger$ are distributed between the arguments f and g by the graded Leibniz rule. Whenever *none* of the two operators overtakes the density of F , the reconfiguration yields $\llbracket \Delta F, G \rrbracket(\mathbf{s})$. Likewise, if *both* derivatives indexed by α overtake F and an old derivative that fell on g , then we obtain $(-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket(\mathbf{s})$, which is the second term in the right-hand side of (29). We claim that the remaining four terms cancel out by virtue of independence of Δ and $\llbracket \cdot, \cdot \rrbracket$ from a choice of normalized variations. To prove this claim, we consecutively inspect the behaviour of those four terms under a swap $\delta \mathbf{s}_1 \rightleftharpoons \delta \mathbf{s}_2$ of coefficients in the normalized test shifts.

The first and second terms sum up to the difference

$$\begin{aligned} & \left\langle (\delta s_1^\alpha) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\mathbf{z}_1) \overbrace{\overleftarrow{e}_\alpha(\mathbf{z}_1), (-\overleftarrow{e}^{\dagger\alpha})(\mathbf{z}_2)}^{-1} (\delta s_{1,\alpha}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2} (\mathbf{z}_2) \right\rangle \underbrace{\langle \overleftarrow{e}^{\dagger\alpha}(\mathbf{x}_2), \overleftarrow{e}_\alpha(\mathbf{x}_1) \rangle}_{-1} \\ & \cdot \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\tau_1}^\beta} \langle \overleftarrow{e}^{\dagger\beta}(\mathbf{x}_1) | \left\langle \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\delta s_2^\beta)(\mathbf{y}_1) \overbrace{\overleftarrow{e}_\beta(\mathbf{y}_1), (-\overleftarrow{e}^{\dagger\beta})(\mathbf{y}_2)}^{-1} \right. \right. \\ & \quad \left. \left. (\delta s_{2,\beta}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\mathbf{y}_2) \right\rangle \underline{|\overleftarrow{e}_\beta(\mathbf{x}_2)\rangle} \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) + \right. \\ & + \left\langle (\delta s_1^\alpha) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\mathbf{z}_1) \overbrace{\overleftarrow{e}_\alpha(\mathbf{z}_1), (-\overleftarrow{e}^{\dagger\alpha})(\mathbf{z}_2)}^{-1} (\delta s_{1,\alpha}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2} (\mathbf{z}_2) \right\rangle \underbrace{\langle \overleftarrow{e}^{\dagger\alpha}(\mathbf{x}_1), \overleftarrow{e}_\alpha(\mathbf{x}_2) \rangle}_{-1} \\ & \cdot (-)^{\text{gh}(F)-1} \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \langle \overleftarrow{e}_\beta(\mathbf{x}_1) | \left\langle (\delta s_2^\beta) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\mathbf{y}_1) \overbrace{\overleftarrow{e}_\beta(\mathbf{y}_1), (-\overleftarrow{e}^{\dagger\beta})(\mathbf{y}_2)}^{-1} \right. \right. \\ & \quad \left. \left. \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\delta s_{2,\beta}^\dagger)(\mathbf{y}_2) \right\rangle \underline{|\overleftarrow{e}^{\dagger\beta}(\mathbf{x}_2)\rangle} \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \frac{\overrightarrow{\partial}}{\partial q_{\tau_1}^\beta} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]) \right). \quad (30) \end{aligned}$$

Recalling that

$$f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} = (-)^{\text{gh}(F)-1} \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]),$$

let us swap the derivations which fall on f from the left and right; this eliminates the sign $(-)^{\text{gh}(F)-1}$. We proceed likewise for g and then transport the variations $\delta \mathbf{s}_1$ and $\delta \mathbf{s}_2$, exchanging their places (and their rôles with respect to Δ and $\llbracket \cdot, \cdot \rrbracket$). The second term in formula (30)

becomes

$$\begin{aligned} & \left\langle (\delta s_2^\beta) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\mathbf{y}_1) \overbrace{\left(\vec{e}_\beta(\mathbf{y}_1), (-\vec{e}^{\dagger\beta})(\mathbf{y}_2) \right)}^{-1} \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\delta s_{2,\beta}^\dagger)(\mathbf{y}_2) \right\rangle \underbrace{\langle \vec{e}_\beta(\mathbf{x}_1), \vec{e}^{\dagger\beta}(\mathbf{x}_2) \rangle}_{+1} \\ & \cdot \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\sigma_1}^\dagger} \langle \vec{e}^{\dagger\alpha}(\mathbf{x}_1) | \left\langle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\delta s_1^\alpha)(\mathbf{z}_1) \overbrace{\left(\vec{e}_\alpha(\mathbf{z}_1), (-\vec{e}^{\dagger\alpha})(\mathbf{z}_2) \right)}^{-1} (\delta s_{1,\alpha}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_2} \right) (\mathbf{z}_2) \right\rangle \right. \\ & \left. \underbrace{|\vec{e}_\alpha(\mathbf{x}_2)\rangle}_{-1} \frac{\overrightarrow{\partial}}{\partial q_{\tau_1}^\beta} \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]). \end{aligned}$$

It is now readily seen that the first term in (30) and this equivalent expression of its second term are opposite to each other. Indeed, relabel the summation indexes $\alpha \rightleftharpoons \beta$, $\sigma \rightleftharpoons \tau$ so that $\delta s_1^\alpha \rightleftharpoons \delta s_2^\beta$, $\delta s_{1,\alpha}^\dagger \rightleftharpoons \delta s_{2,\beta}^\dagger$, and swap the copies of base manifold M^n by $\mathbf{y} \rightleftharpoons \mathbf{z}$. Due to the second factors in the products $(-1) \cdot (-1) \cdot (-1) \cdot (-1) = +1$ versus $(-1) \cdot (+1) \cdot (-1) \cdot (-1) = -1$, the two terms in (30) cancel out after the integration by parts and evaluation of the couplings in view of (16).

Next, the integrand of $\Delta(\llbracket F, G \rrbracket)(s)$ contains a restriction to the infinite jet $j^\infty(s)$ of the third term, which is

$$\begin{aligned} & \left\langle (\delta s_1^\alpha) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\mathbf{z}_1) \overbrace{\left(\vec{e}_\alpha(\mathbf{z}_1), (-\vec{e}^{\dagger\alpha})(\mathbf{z}_2) \right)}^{-1} (\delta s_{1,\alpha}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2} (\mathbf{z}_2) \right\rangle \underbrace{\langle \vec{e}^{\dagger\alpha}(\mathbf{x}_2), \vec{e}_\alpha(\mathbf{x}_1) \rangle}_{-1} \\ & \cdot \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \left(f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \right) \\ & \underbrace{\langle \vec{e}_\beta(\mathbf{x}_1) | \left\langle (\delta s_2^\beta) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\mathbf{y}_1) \overbrace{\left(\vec{e}_\beta(\mathbf{y}_1), (-\vec{e}^{\dagger\beta})(\mathbf{y}_2) \right)}^{-1} \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\delta s_{2,\beta}^\dagger)(\mathbf{y}_2) \right\rangle}_{+1} \underbrace{|\vec{e}^{\dagger\beta}(\mathbf{x}_2)\rangle}_{+1} \\ & \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} \frac{\overrightarrow{\partial}}{\partial q_{\tau_1}^\beta} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]). \end{aligned}$$

Let the summation indexes be relabelled as above: $\alpha \rightleftharpoons \beta$, $\sigma \rightleftharpoons \tau$, and $\delta s_1^\alpha \rightleftharpoons \delta s_2^\beta$, $\delta s_{1,\alpha}^\dagger \rightleftharpoons \delta s_{2,\beta}^\dagger$ on top of $\mathbf{y} \rightleftharpoons \mathbf{z}$. The transformation of graded derivations falling from the left and right on f is then

$$\begin{aligned} \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \left(f \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \right) & \mapsto \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \left(f \frac{\overleftarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \right) = \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \left((-)^{\text{gh}(F)-1} \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} f \right) = \\ & = (-)^{\text{gh}(F)-2} \cdot (-)^{\text{gh}(F)-1} \left(\frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} f \right) \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} = - \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \left(f \frac{\overleftarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \right). \end{aligned}$$

This minus sign shows that the third term as it was written initially, and the newly produced one in which the variations δs_1 and δs_2 are interchanged have opposite signs. At the same time, these integral functionals must be equal to each other due to independence of Δ and $\llbracket \cdot, \cdot \rrbracket$ of a choice of the test shifts. Therefore, each of those expressions vanishes.

The fourth term is processed analogously; its integrand is

$$\begin{aligned}
 & (-)^{\text{gh}(F)} \left\langle \left(\delta s_1^\alpha \right) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\mathbf{z}_1) \overbrace{\left(\vec{e}_\alpha(\mathbf{z}_1), (-\vec{e}^{\dagger\alpha})(\mathbf{z}_2) \right)}^{-1} \left(\delta s_{1,\alpha}^\dagger \right) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2} (\mathbf{z}_2) \right\rangle \cdot \underbrace{\langle \vec{e}^{\dagger\alpha}(\mathbf{x}_1), \vec{e}_\alpha(\mathbf{x}_2) \rangle}_{-1} \\
 & \cdot \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} f(\mathbf{x}_1, [\mathbf{q}], [\mathbf{q}^\dagger]) \frac{\overleftarrow{\partial}}{\partial q_{\tau_1}^\beta} \\
 & \left\langle \vec{e}^{\dagger\beta}(\mathbf{x}_1) \right| \left\langle \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\delta s_2^\beta)(\mathbf{y}_1) \overbrace{\left(\vec{e}_\beta(\mathbf{y}_1), (-\vec{e}^{\dagger\beta})(\mathbf{y}_2) \right)}^{-1} \left(\delta s_{2,\beta}^\dagger \right) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\mathbf{y}_2) \right\rangle \underbrace{|\vec{e}_\beta(\mathbf{x}_2)\rangle}_{-1} \\
 & \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} g(\mathbf{x}_2, [\mathbf{q}], [\mathbf{q}^\dagger]).
 \end{aligned}$$

The very same procedure of two variations interchange and relabelling restores an almost identical expression in which, however, the parity-odd derivations go in the inverse order $\overrightarrow{\partial}/\partial q_{\beta,\tau_2}^\dagger \circ \overrightarrow{\partial}/\partial q_{\alpha,\sigma_2}^\dagger$. Equal to minus itself, the fourth term vanishes. This concludes the proof. \square

The following example illustrates the assertion of Lemma 5 (but not a technique of its proof which itself accompanies Lemma 1). We use the convention from Remark 2.7, denoting by $d/d\mathbf{y}_i$ or $d/d\mathbf{z}_j$ the total derivatives which act on the functionals' densities at points \mathbf{x}_k ; this keeps track of those derivatives origin and lets us indicate the couplings' values as they appear after the integrations by parts, contributing only with sign factors ± 1 . For the sake of brevity we do not write the (co)vectors \vec{e}_i and $\vec{e}^{\dagger i}$ in the formulas below, referring to the proofs in preceding sections. Likewise, we do not indicate the base point congruences that occur due to the absolute locality of couplings.

An overall comment to Example 2.4 below is that, fully aware of the goal which is to calculate $\Delta(\llbracket F, G \rrbracket)$ or, respectively, $\llbracket \Delta F, G \rrbracket$ and $\llbracket F, \Delta G \rrbracket$, we do not interrupt the logic of our reasoning by attempting to view the intermediate objects $\llbracket F, G \rrbracket$ or ΔF and ΔG as mappings $\Gamma(\pi_{\text{BV}}) \rightarrow \mathbb{k}$, cf. Corollary 2 on p. 24. Such mappings would not be elements of the structures which stand in the left- and right-hand sides of the identity under examination. The slogan is that a step-by-step evaluation is illegal; derivations of the end-product from input data must not be interrupted at half-way.

We also emphasize that the example below is a prototype reasoning which is equally well applicable to any other arguments F and G in (29); a choice of the functionals is here not specific to any model. The point is that equality (29) holds and does not require any manual regularization.

Example 2.4. Consider the integral functionals

$$F = \int q^\dagger q q_{x_1 x_1} dx_1 \quad \text{and} \quad G = \int q_{x_2 x_2}^\dagger \cos q dx_2.$$

Let us show that equality (29) is satisfied for F and G , that is,

$$\Delta(\llbracket F, G \rrbracket) = \llbracket \Delta F, G \rrbracket + \llbracket F, \Delta G \rrbracket, \quad \text{gh}(F) = 1, \quad (31)$$

in the frames of product-bundle geometry of variations and operational definitions of the BV-Laplacian Δ and variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$.

We have

$$\begin{aligned} \llbracket F, G \rrbracket = & \iiint\!\!\!\int dx_1 dx_2 dy_1 dy_2 \left\langle \underbrace{\left(q^\dagger q_{xx} + \frac{d^2}{dy_1^2} (q^\dagger q) \right)}_{x_1} \cdot \underbrace{\langle \delta s_2(y_1), \delta s_2^\dagger(y_2) \rangle}_{+1} \cdot \underbrace{\frac{d^2}{dy_2^2} (\cos q)}_{x_2} \right\rangle \\ & + \iiint\!\!\!\int dx_1 dx_2 dy_1 dy_2 \left\langle \underbrace{(qq_{xx})}_{x_1} \cdot \underbrace{\langle \delta s_2^\dagger(y_2), \delta s_2(y_1) \rangle}_{-1} \cdot \underbrace{(-q_{xx}^\dagger \sin q)}_{x_2} \right\rangle. \end{aligned}$$

Therefore, one side of the expected equality is

$$\begin{aligned} \Delta(\llbracket F, G \rrbracket) = & \int dz_1 \int dz_2 \int dx_1 \int dx_2 \int dy_1 \int dy_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle \delta s_2(y_1), \delta s_2^\dagger(y_2) \rangle}_{+1} \\ & \cdot \left\langle \underbrace{\frac{d^2}{dz_1^2} (1)}_{x_1} \cdot \underbrace{\frac{d^2}{dy_2^2} (\cos q)}_{x_2} + \underbrace{q_{xx}}_{x_1} \cdot \underbrace{\frac{d^2}{dy_2^2} (-\sin q)}_{x_2} + \underbrace{\frac{d^2}{dy_1^2} (1)}_{x_1} \cdot \underbrace{\frac{d^2}{dy_2^2} (\cos q)}_{x_2} + \underbrace{\frac{d^2}{dy_1^2} (q)}_{x_1} \cdot \underbrace{\frac{d^2}{dy_2^2} (-\sin q)}_{x_2} \right\rangle \\ & + \int dz_1 \int dz_2 \int dx_1 \int dx_2 \int dy_1 \int dy_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle \delta s_2^\dagger(y_2), \delta s_2(y_1) \rangle}_{-1} \\ & \cdot \left\langle \underbrace{q_{xx}}_{x_1} \cdot \underbrace{\frac{d^2}{dz_2^2} (-\sin q)}_{x_2} + \underbrace{\frac{d^2}{dz_1^2} (q)}_{x_1} \cdot \underbrace{\frac{d^2}{dz_2^2} (-\sin q)}_{x_2} + \underbrace{(qq_{xx})}_{x_1} \cdot \underbrace{\frac{d^2}{dz_2^2} (-\cos q)}_{x_2} \right\rangle. \end{aligned}$$

The respective pairs of underlined terms cancel out and there remains only

$$\begin{aligned} = & \int \dots \int dz_1 dz_2 dx_1 dx_2 dy_1 dy_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle (qq_{xx})}_{x_1} \cdot \underbrace{\frac{d^2}{dz_2^2} (-\cos q)}_{x_2}}_{-1} \\ & \cdot \underbrace{\langle \delta s_2^\dagger(y_2), \delta s_2(y_1) \rangle}_{-1}. \quad (32) \end{aligned}$$

On the other hand, we obtain that

$$\Delta F = \iiint\!\!\!\int dz_1 dz_2 dx_1 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle q_{xx} + \frac{d^2}{dz_1^2} (q) \rangle}_{x_1},$$

which yields

$$\begin{aligned} \llbracket \Delta F, G \rrbracket = & \int dz_1 \int dz_2 \int dx_1 \int dx_2 \int dy_1 \int dy_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \\ & \cdot \left\langle \underbrace{\left(\frac{d^2}{dy_1^2} (1) + \frac{d^2}{dz_1^2} (1) \right)}_{x_1} \cdot \underbrace{\frac{d^2}{dy_2^2} (\cos q)}_{x_2} \right\rangle \cdot \underbrace{\langle \delta s_2(y_1), \delta s_2^\dagger(y_2) \rangle}_{+1} = 0. \end{aligned}$$

From the fact that the other BV-Laplacian,

$$\Delta G = \iiint\!\!\!\int dz_1 dz_2 dx_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle \frac{d^2}{dz_2^2} (-\sin q) \rangle}_{x_2},$$

does not contain q^\dagger so that the first half of the Schouten bracket $\llbracket F, \Delta G \rrbracket$ drops out, we deduce that

$$\llbracket F, \Delta G \rrbracket = \int \cdots \int dz_1 dz_2 dx_1 dx_2 dy_1 dy_2 \underbrace{\langle \delta s_1(z_1), \delta s_1^\dagger(z_2) \rangle}_{+1} \cdot \underbrace{\langle (qq_{xx})_{x_1} \cdot \frac{d^2}{dz_2^2} (-\cos q) \rangle_{x_2}}_{-1} \cdot \underbrace{\langle \delta s_2^\dagger(y_2), \delta s_2(y_1) \rangle}_{-1}. \quad (33)$$

Consequently, the two sides of (31), namely, $\Delta(\llbracket F, G \rrbracket)$ expressed by (32) and $\llbracket \Delta F, G \rrbracket + \llbracket F, \Delta G \rrbracket$ accumulated in (33), match perfectly for the functionals F and G at hand.

Theorem 6. *Let $F, G \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be two functionals. The Batalin–Vilkovisky Laplacian Δ satisfies the relation*

$$\Delta(\llbracket F, G \rrbracket) = \llbracket \Delta F, G \rrbracket + (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket. \quad (29)$$

In other words, the operator Δ is a graded derivation of the variational Schouten bracket $\llbracket \cdot, \cdot \rrbracket$.

Proof. We prove this by induction over the number of building blocks in each argument of the Schouten bracket in the left hand side of (29). If F and G both belong to $\overline{H}^*(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \cdots \times T\pi_{\text{BV}})$, then Lemma 5 states the assertion, which is the base of induction. To make an inductive step, without loss of generality let us assume that the second argument of $\llbracket \cdot, \cdot \rrbracket$ in (29) is a product of two elements from $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$, each of them containing less multiples from $\overline{H}^*(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \cdots \times T\pi_{\text{BV}})$ than the product. Denote such factors by G and H and recall that by Theorem 4,

$$\llbracket F, G \cdot H \rrbracket = \llbracket F, G \rrbracket \cdot H + (-)^{(\text{gh}(F)-1) \cdot \text{gh}(G)} G \cdot \llbracket F, H \rrbracket.$$

Therefore, using Theorem 3 we have that

$$\begin{aligned} \Delta(\llbracket F, G \cdot H \rrbracket) &= \Delta(\llbracket F, G \rrbracket) \cdot H + (-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket \llbracket F, G \rrbracket, H \rrbracket + (-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket F, G \rrbracket \cdot \Delta H \\ &\quad + (-)^{(\text{gh}(F)-1) \text{gh}(G)} \left(\Delta G \cdot \llbracket F, H \rrbracket + (-)^{\text{gh}(G)} \llbracket G, \llbracket F, H \rrbracket \rrbracket + (-)^{\text{gh}(G)} G \cdot \Delta(\llbracket F, H \rrbracket) \right). \end{aligned}$$

Using the inductive hypothesis in the first and last terms of the right-hand side in the above formula, we continue the equality and obtain

$$\begin{aligned} &= \llbracket \Delta F, G \rrbracket \cdot H + (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket \cdot H + (-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket \llbracket F, G \rrbracket, H \rrbracket \\ &\quad + (-)^{\text{gh}(F) \text{gh}(G)} \llbracket G, \llbracket F, H \rrbracket \rrbracket + (-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket F, G \rrbracket \cdot \Delta H \\ &\quad + (-)^{(\text{gh}(F)-1) \text{gh}(G)} \Delta G \cdot \llbracket F, H \rrbracket + (-)^{\text{gh}(F) \text{gh}(G)} G \cdot \llbracket \Delta F, H \rrbracket \\ &\quad + (-)^{\text{gh}(F) \text{gh}(G)+\text{gh}(F)-1} G \cdot \llbracket F, \Delta H \rrbracket. \end{aligned} \quad (34)$$

On the other hand, let us expand the formula

$$\llbracket \Delta F, G \cdot H \rrbracket + (-)^{\text{gh}(F)-1} \llbracket F, \Delta(G \cdot H) \rrbracket,$$

which is the right hand side of (29) in the inductive claim. We obtain

$$\begin{aligned}
 &= \llbracket \Delta F, G \rrbracket \cdot H + (-)^{\text{gh}(\Delta F)-1} \text{gh}(G) G \cdot \llbracket \Delta F, H \rrbracket \\
 &\quad + (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \cdot H + (-)^{\text{gh}(G)} \llbracket G, H \rrbracket + (-)^{\text{gh}(G)} G \cdot \Delta H \rrbracket \\
 &= \llbracket \Delta F, G \rrbracket \cdot H + (-)^{\text{gh}(F)} \text{gh}(G) G \cdot \llbracket \Delta F, H \rrbracket + (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket \cdot H \\
 &\quad + (-)^{\text{gh}(F)-1} (-)^{\text{gh}(F)-1} \text{gh}(G)-1 \Delta G \cdot \llbracket F, H \rrbracket + (-)^{\text{gh}(F)-1} (-)^{\text{gh}(G)} \llbracket F, \llbracket G, H \rrbracket \rrbracket \\
 &\quad + (-)^{\text{gh}(F)-1} (-)^{\text{gh}(G)} \llbracket F, G \rrbracket \cdot \Delta H + (-)^{\text{gh}(F)-1} (-)^{\text{gh}(G)} (-)^{\text{gh}(F)-1} \text{gh}(G) G \cdot \llbracket F, \Delta H \rrbracket.
 \end{aligned} \tag{35}$$

Comparing (35) with (34), which was derived from the inductive hypothesis, we see that all terms match except for

$$(-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket \llbracket F, G \rrbracket, H \rrbracket + (-)^{\text{gh}(F)} \text{gh}(G) \llbracket G, \llbracket F, H \rrbracket \rrbracket$$

from (34) versus

$$(-)^{\text{gh}(F)+\text{gh}(G)-1} \llbracket F, \llbracket G, H \rrbracket \rrbracket$$

from (35). However, these three terms constitute Jacobi's identity (28) for the variational Schouten bracket. Namely, we have that (cf. [32])

$$\llbracket F, \llbracket G, H \rrbracket \rrbracket = \llbracket \llbracket F, G \rrbracket, H \rrbracket + (-)^{\text{gh}(F)-1} \text{gh}(G)-1 \llbracket G, \llbracket F, H \rrbracket \rrbracket, \tag{36}$$

so that by multiplying both sides of the identity by $(-)^{\text{gh}(F)+\text{gh}(G)-1}$, we fully balance (34) and (35). This completes the inductive step and concludes the proof. \square

Lemma 7. The linear operator

$$\Delta: \overline{H}^{n(1+k)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}}) \longrightarrow \overline{H}^{n(2+k)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$$

is a differential for every $k \geq 0$.

The proof of Lemma 7 is conceptually close to the second and third steps in the proof of Lemma 5. Namely, two normalized variations are swapped in an integral functional within the image of Δ^2 , which yields an indistinguishable result of opposite sign.

Proof. Let δs_1 and δs_2 be normalized test shifts of a section $s \in \Gamma(\pi_{\text{BV}})$, and let $H = \int h(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \cdot \text{dvol}(\mathbf{x})$ be an integral functional. (It suffices to consider a simplified picture $H \in \overline{H}^n(\pi_{\text{BV}})$, not taking into account any built-in variations in the construction of H .) By definition, we have that

$$\begin{aligned}
 \Delta(\Delta H)(s) &= \int_M \text{d}z_1 \int_M \text{d}z_2 \int_M \text{d}y_1 \int_M \text{d}y_2 \int_M \text{dvol}(\mathbf{x}) \cdot \\
 &\quad \cdot \left\{ \left\langle (\delta s_1^\alpha) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_1} \right)^{\sigma_1} (\mathbf{z}_1) \overbrace{\vec{e}_\alpha(\mathbf{z}_1), (-\vec{e}^{\dagger\alpha})(\mathbf{z}_2)}^{-1} (\delta s_{1,\alpha}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{z}_2} \right)^{\sigma_2} (\mathbf{z}_2) \right\rangle \underbrace{\left\langle \vec{e}^{\dagger\alpha}(\mathbf{x}), \vec{e}_\alpha(\mathbf{x}) \right\rangle}_{-1} \right. \\
 &\quad \left. \left\langle (\delta s_2^\beta) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\tau_1} (\mathbf{y}_1) \overbrace{\vec{e}_\beta(\mathbf{y}_1), (-\vec{e}^{\dagger\beta})(\mathbf{y}_2)}^{-1} (\delta s_{2,\beta}^\dagger) \left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\tau_2} (\mathbf{y}_2) \right\rangle \underbrace{\left\langle \vec{e}^{\dagger\beta}(\mathbf{x}), \vec{e}_\beta(\mathbf{x}) \right\rangle}_{-1} \right. \\
 &\quad \left. \left. \frac{\overrightarrow{\partial}}{\partial q_{\sigma_1}^\alpha} \frac{\overrightarrow{\partial}}{\partial q_{\alpha, \sigma_2}^\dagger} \frac{\overrightarrow{\partial}}{\partial q_{\tau_1}^\beta} \frac{\overrightarrow{\partial}}{\partial q_{\beta, \tau_2}^\dagger} h(\mathbf{x}, [\mathbf{q}], [\mathbf{q}^\dagger]) \right\} \Bigg|_{j_{\mathbf{x}}^\infty(s)}.
 \end{aligned}$$

By exchanging the integrand's upper two lines and then relabelling $\alpha \rightleftharpoons \beta$, $\sigma \rightleftharpoons \tau$ so that $\delta s_1^\alpha \rightleftharpoons \delta s_2^\beta$ and $\delta s_{1,\alpha}^\dagger \rightleftharpoons \delta s_{2,\beta}^\dagger$, and by swapping the reference $\mathbf{y} \rightleftharpoons \mathbf{z}$ to copies of the base manifold M^n , we almost recover the initial expression (which should be the case), yet the order in which the parity-odd partial derivatives follow is inverse,

$$\frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \circ \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \mapsto \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger} \circ \frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} = -\frac{\overrightarrow{\partial}}{\partial q_{\alpha,\sigma_2}^\dagger} \circ \frac{\overrightarrow{\partial}}{\partial q_{\beta,\tau_2}^\dagger}.$$

Therefore the integrand of functional $\Delta^2 H$ vanishes, which proves the assertion. \square

Theorem 8. *The Batalin–Vilkovisky Laplacian Δ is a differential: for all $H \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ we have*

$$\Delta^2(H) = 0.$$

Proof. We prove Theorem 8 by induction over the number of building blocks from $\overline{H}^*(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$ in the argument $H \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ of Δ^2 . If $H \in \overline{H}^*(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$ itself is an integral functional, then by Lemma 7 there remains nothing to prove. Suppose now that $H = F \cdot G$ for some $F, G \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$. Then Theorem 3 yields that

$$\Delta^2(F \cdot G) = \Delta \left(\Delta F \cdot G + (-)^{\text{gh}(F)} \llbracket F, G \rrbracket + (-)^{\text{gh}(F)} F \cdot \Delta G \right).$$

Using Theorem 3 again and also Theorem 6, we continue the equality:

$$\begin{aligned} &= \Delta^2 F \cdot G + (-)^{\text{gh}(\Delta F)} \llbracket \Delta F, G \rrbracket + (-)^{\text{gh}(\Delta F)} \Delta F \cdot \Delta G \\ &\quad + (-)^{\text{gh}(F)} \llbracket \Delta F, G \rrbracket + (-)^{\text{gh}(F)} (-)^{\text{gh}(F)-1} \llbracket F, \Delta G \rrbracket \\ &\quad + (-)^{\text{gh}(F)} \Delta F \cdot \Delta G + (-)^{\text{gh}(F)} (-)^{\text{gh}(F)} \llbracket F, \Delta G \rrbracket + (-)^{\text{gh}(F)} (-)^{\text{gh}(F)} F \cdot \Delta^2 G. \end{aligned}$$

By the inductive hypothesis, the first and last terms in the above formula vanish; taking into account that $\text{gh}(\Delta F) = \text{gh}(F) - 1$ in \mathbb{Z}_2 , the terms with $\Delta F \cdot \Delta G$ cancel against each other, as do the terms containing $\llbracket \Delta F, G \rrbracket$ and $\llbracket F, \Delta G \rrbracket$. The proof is complete. \square

3. The quantum master-equation

3.1. The Laplace equation

In this section we inspect the conditions upon functionals $F \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ under which the Feynman path integrals $\int_{\Gamma(\zeta^0)} [Ds] F([s], [s^\dagger])$ are (infinitesimally) independent of the unphysical anti-objects $s^\dagger \in \Gamma(\zeta^1)$. The derivation of such a condition (see equation (39) below) relies on an extra assumption of the translation invariance of a measure in the path integral. It must be noted, however, that we do not define Feynman's integral here and do not introduce that measure which essentially depends on the agreement about the classes of 'admissible' sections $\Gamma(\pi)$ or $\Gamma(\zeta^{(0,1)})$. Consequently, our reasoning is to some extent heuristic.

The basics of path integration, which we recall here for consistency, are standard: they illustrate how the geometry of the BV-Laplacian works in practice. We draw the experts' attention only to the fact that in our notation Ψ is not the gauge fixing fermion Ψ such that the odd-component's section $s^\dagger \in \Gamma(\zeta^1)$ is the restriction of $\delta\Psi/\delta q$ to the jet of a section for ζ^0 ; instead, we let Ψ determine the infinitesimal shift $\dot{q}^\dagger = \delta\Psi/\delta q$ of coordinates along the fibre's parity-odd half. We also note that the preservation of parity is not mandatory here and thus an even-parity $\Psi \in \overline{H}^n(\zeta^0) \hookrightarrow \overline{H}^n(\pi_{\text{BV}})$ is a legitimate choice.

Let $F = \int f(\mathbf{x}, \mathbf{q}, \mathbf{q}^\dagger) \text{dvol}(\mathbf{x}) \in \overline{H}^n(\pi_{\text{BV}})$ be a functional; here and in what follows we proceed over the building blocks of elements from $\overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ by the graded Leibniz

rule. Let $\Psi = \int \psi(\mathbf{y}, \mathbf{q}) \text{dvol}(\mathbf{y}) \in \overline{H}^n(\zeta^0) \hookrightarrow \overline{H}^n(\pi_{\text{BV}})$ be an integral functional which, by assumption, is constant along ghost parity-odd variables: $\Psi(s^\alpha, s^\dagger_\beta) = \Psi(s^\alpha, t^\dagger_\beta)$ for any sections $\{s^\alpha\} \in \Gamma(\zeta^0)$ and $\{s^\dagger_\beta\}, \{t^\dagger_\beta\} \in \Gamma(\zeta^1)$. We investigate under which conditions the path integral $\int_{\Gamma(\zeta^0)} [Ds^\alpha] F(s^\alpha, s^\dagger_\beta): \Gamma(\zeta^1) \rightarrow \mathbb{k}$ is infinitesimally independent of a choice of the anti-objects:

$$\frac{d}{d\varepsilon^\dagger} \Big|_{\varepsilon^\dagger=0} \int_{\Gamma(\zeta^0)} [Ds^\alpha] F\left(s^\alpha, s^\dagger_\beta + \varepsilon^\dagger \frac{\overrightarrow{\delta}\psi}{\delta q^\beta} \Big|_{s^\alpha}\right) = 0 \quad \text{for all } s^\dagger \in \Gamma(\zeta^1). \quad (37)$$

Note that this formula makes sense because the bundles ζ^0 and ζ^1 are dual so that a variational covector in the geometry of ζ^0 acts as a shift vector in the geometry of ζ^1 . The left-hand side of (37) equals

$$\int_{\Gamma(\zeta^0)} [Ds^\alpha] \int_M \text{dvol}(\mathbf{x}) \frac{\overrightarrow{\delta}\psi}{\delta q^\beta}(\mathbf{x}, \mathbf{q}) \Big|_{j_\mathbf{x}^\infty(s^\alpha)} \cdot \frac{\overleftarrow{\delta}f}{\delta q^\dagger_\beta}(\mathbf{x}, \mathbf{q}, \mathbf{q}^\dagger) \Big|_{j_\mathbf{x}^\infty(s^\alpha, s^\dagger_\beta)}, \quad s^\dagger \in \Gamma(\zeta^1).$$

Take any auxiliary section $\delta s = (\delta s^\alpha, \delta s^\dagger_\beta) \in \Gamma(T\zeta^{(0|1)})$ normalized by $\delta s^\alpha(x) \cdot \delta s^\dagger_\alpha(x) \equiv 1$ at every $\mathbf{x} \in M^n$ for each $\alpha = 1, \dots, m + m_1 + \dots + m_\lambda = N$ and blow up the scalar integrand to a pointwise contraction of dual object taking their values in the fibres $T_{(\mathbf{x}, \phi(\mathbf{x}), s(\mathbf{x}))} V_\mathbf{x}$ and $T_{(\mathbf{x}, \phi(\mathbf{x}), s^\dagger(\mathbf{x}))} \Pi V_\mathbf{x}^\dagger$ of $T(\pi_{\text{BV}})$ over $\phi(x)$: for $s = (s^\alpha, s^\dagger_\beta)$ we have

$$\begin{aligned} & \int_M \text{dvol}(\mathbf{x}) \left(\frac{\overrightarrow{\delta}\psi}{\delta q^\alpha} \cdot \frac{\overleftarrow{\delta}f}{\delta q^\dagger_\alpha} \right) \Big|_{j_\mathbf{x}^\infty(s)} \\ &= \int_M \text{dvol}(\mathbf{x}_1) \int_M \text{dvol}(\mathbf{x}_2) \int_M \text{d}\mathbf{y}_1 \int_M \text{d}\mathbf{y}_2 \left(\psi(\mathbf{x}_1, \mathbf{q}) \frac{\overleftarrow{\partial}}{\partial q_{\sigma_1}^{j_1}} \right) \Big|_{j_{\mathbf{x}_1}^\infty(s)} \\ & \cdot \langle \overleftarrow{e}^{\dagger j_1}(\mathbf{x}_1) | \left(\left(\frac{\overleftarrow{\partial}}{\partial \mathbf{y}_1} \right)^{\sigma_1} \delta s^{i_1}(\mathbf{y}_1) \vec{e}_{i_1}(\mathbf{y}_1), -\overleftarrow{e}^{\dagger i_2}(\mathbf{y}_2) \delta s^\dagger_{i_2}(\mathbf{y}_2) \left(\frac{\overrightarrow{\partial}}{\partial \mathbf{y}_2} \right)^{\sigma_2} \right) | \overleftarrow{e}_{j_2}(\mathbf{x}_2) \rangle \\ & \cdot \left(\frac{\overrightarrow{\partial}}{\partial q_{j_2, \sigma_2}^\dagger} f(\mathbf{x}_2, \mathbf{q}, \mathbf{q}^\dagger) \right) \Big|_{j_{\mathbf{x}_2}^\infty(s)}. \end{aligned}$$

In fact, the integrand refers to a definition of the evolutionary vector field \mathbf{Q}^Ψ such that $\mathbf{Q}^\Psi(F) \cong \llbracket \Psi, F \rrbracket$ modulo integration by parts in the building blocks of F , cf. [32]. Due to a special choice of the dependence of Ψ on s only, this is indeed the Schouten bracket $\llbracket \Psi, F \rrbracket$.

To rephrase the indifference of the path integral to a choice of Ψ in terms of an equation upon the functional F alone, we perform integration by parts in Feynman's integral. For this we employ the translation invariance $[Ds] = [D(s - \mu \cdot \delta s)]$ of the functional measure.

Lemma 9. Let $H = \int h(\mathbf{x}, \mathbf{q}, \mathbf{q}^\dagger) \text{dvol}(\mathbf{x}) \in \overline{H}^n(\pi_{\text{BV}}) \subset \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be an integral functional and $\delta s \in \Gamma(T\zeta^0) \hookrightarrow \Gamma(T\zeta^{(0|1)})$ be a shift. Then we have that

$$\int_{\Gamma(\zeta^0)} [Ds^\alpha] \int_M \text{dvol}(\mathbf{x}) \delta s^\nu(\mathbf{x}) \cdot \frac{\overleftarrow{\delta}h}{\delta q^\nu} \Big|_{j_\mathbf{x}^\infty(s^\alpha, s^\dagger_\beta)} = 0,$$

where the section $s^\dagger \in \Gamma(\zeta^1)$ is a parameter.

Proof. Indeed,

$$0 = \frac{d}{d\mu} \Big|_{\mu=0} \int_{\Gamma(\zeta^0)} [Ds^\alpha] H(s^\alpha, s^\dagger_\beta),$$

because the integral contains no parameter $\mu \in \mathbb{k}$. We continue the equality:

$$\begin{aligned} &= \frac{d}{d\mu} \Big|_{\mu=0} \int_{\Gamma(\zeta^0)} [D(s^\alpha - \mu \delta s^\alpha)] H(s^\alpha, s^\dagger_\beta) \\ &= \frac{d}{d\mu} \Big|_{\mu=0} \int_{\Gamma(\zeta^0)} [Ds^\alpha] H(s^\alpha + \mu \delta s^\alpha, s^\dagger_\beta) = \int_{\Gamma(\zeta^0)} [Ds^\alpha] \frac{d}{d\mu} \Big|_{\mu=0} H(s^\alpha + \mu \delta s^\alpha, s^\dagger_\beta), \end{aligned}$$

which yields the helpful formula in the lemma's assertion. \square

Returning to the functionals Ψ and F and denoting $G(s) := \frac{d}{d\ell} \Big|_{\ell=0} F(s + \ell \cdot \overleftarrow{\delta s^\dagger})$, we use the Leibniz rule for the derivative of $H = \Psi \cdot G$:

$$\frac{d}{d\mu} \Big|_{\mu=0} (\Psi \cdot G)(s + \mu \cdot \overleftarrow{\delta s}) = \frac{d}{d\mu} \Big|_{\mu=0} (\Psi)(s + \mu \cdot \overleftarrow{\delta s}) \cdot G(s) + \Psi(s) \cdot \frac{d}{d\mu} \Big|_{\mu=0} (G)(s + \mu \cdot \overleftarrow{\delta s}).$$

Because the path integral over $[Ds^\alpha]$ of the entire expression vanishes by Lemma 9 in which we were ready to proceed by the Leibniz rule over building blocks, we infer that the path integrals of the two terms are opposite. Now take the traces over indexes in both variations. The integral of the first term equals the initial expression for the path integral containing F , i. e., the left-hand side of equation (37). Consequently, if

$$\int_{\Gamma(\zeta^0)} [Ds^\alpha] \Psi(s^\alpha) \cdot \Delta F(s^\alpha, s^\dagger_\beta) = 0 \tag{38}$$

for $\{s^\dagger_\beta\} \in \Gamma(\zeta^1)$ and for all $\Psi \in \overline{H}^n(\zeta^0) \hookrightarrow \overline{H}^n(\pi_{\text{BV}})$, then the path integral over F is infinitesimally independent of a section $\{s^\dagger_\beta\} \in \Gamma(\zeta^1)$.

The condition

$$\Delta F = 0 \tag{39}$$

is sufficient for equation (38), and therefore equation (37), to hold. By specifying a class $\Gamma(\pi_{\text{BV}})$ of admissible sections of the BV-bundle for a concrete field model, and endowing that space of sections with a suitable metric, one could reinstate a path integral analogue of the main lemma in the calculus of variations and then argue that the condition $\Delta F = 0$ is also necessary.

Summarizing, whenever equation (39) holds, one can assign arbitrary admissible values to the odd-parity coordinates; for example, one can let $s^\dagger_\beta(\mathbf{x}) = \delta\psi/\delta q^\beta \Big|_{j_\infty(s^\alpha)}$ for a gauge-fixing integral $\Psi = \int \psi(\mathbf{x}, \mathbf{q}) \text{dvol}(\mathbf{x}) \in \overline{H}^n(\zeta^0)$. This choice is reminiscent of the substitution principle, see [32] and [45].

Laplace's equation (39) ensures the infinitesimal independence from non-physical anti-objects for path integrals of functionals over physical fields – not only in the classical BV-geometry of the bundle π_{BV} , but also in the quantum setup, whenever all objects are tensored with formal power series $\mathbb{k}[[\hbar, \hbar^{-1}]]$ in the Planck constant \hbar . It is accepted that each quantum field s^\hbar contributes to the expectation value of a functional \mathcal{O}^\hbar with the factor $\exp(iS^\hbar(s^\hbar)/\hbar)$, where S^\hbar is the quantum BV-action of the model. Solutions \mathcal{O}^\hbar of the equation $\Delta(\mathcal{O}^\hbar \cdot \exp(iS^\hbar/\hbar)) = 0$ are the *observables*. In particular, the postulate that the unit $1: s^\hbar \mapsto 1 \in \mathbb{k}$ is averaged to unit by the Feynman integral of $1 \cdot \exp(iS^\hbar(s^\hbar)/\hbar)$ over the space of quantum fields s^\hbar normalizes the integration measure and constrains the quantum BV-action by the quantum master-equation (see, e.g., [7, 8, 20, 22, 54]).

Proposition 10. Let S^{\hbar} be the even quantum BV-action (i. e., let it have a density that has an even number of ghost parity-odd coordinates in each of its terms). If the identity

$$\Delta \left(\exp\left(\frac{i}{\hbar} S^{\hbar}\right) \right) = 0$$

holds, then S^{\hbar} satisfies the quantum master-equation:

$$i\hbar \Delta S^{\hbar} = \frac{1}{2} \llbracket S^{\hbar}, S^{\hbar} \rrbracket. \quad (40)$$

Proposition 11. If an even functional \mathcal{O} and the quantum BV-action S^{\hbar} are such that $\Delta(\mathcal{O} \exp(iS^{\hbar}/\hbar)) = 0$ and $\Delta(\exp(iS^{\hbar}/\hbar)) = 0$ hold, respectively, then \mathcal{O} satisfies

$$\Omega^{\hbar}(\mathcal{O}) := -i\hbar \Delta \mathcal{O} + \llbracket S^{\hbar}, \mathcal{O} \rrbracket = 0. \quad (41)$$

We quote the standard proofs of Propositions 10 and 11 from [35] in Appendix Appendix A — yet now we gain a deeper insight on a construction of the quantum BV-differential Ω^{\hbar} .

Remark 3.1. A practical way to fix the signs which arise in the BV-Laplacian and Schouten bracket from the ghost parity and a grading in the case of a superbundle $\pi: E^{(m_0+n_0|m_1+n_1)} \rightarrow M^{(n_0|n_1)}$ of superfields is by a re-derivation of the Laplace equation $\Delta(\mathcal{O} \exp(\frac{i}{\hbar} S^{\hbar})) = 0$ upon an observable \mathcal{O} starting from the Schwinger–Dyson condition,

$$\vec{\partial}_{\vec{\delta}\Psi/\delta\mathbf{q}}^{(\mathbf{q}^\dagger)} \left(\int [D\mathbf{q}] \mathcal{O}([\mathbf{q}], [\mathbf{q}^\dagger]) \exp\left(\frac{i}{\hbar} S^{\hbar}([\mathbf{q}], [\mathbf{q}^\dagger])\right) \right) = 0, \quad (42)$$

which postulates the Feynman path integral’s independence of the non-physical BV-coordinates \mathbf{q}^\dagger with odd ghost parity. Note that the measure in the path integral involves only ghost parity-even objects (whatever be their \mathbb{Z}_2 -grading).

Theorem 12. Let $\mathcal{O} \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be a functional and let the even functional $S^{\hbar} \in \overline{H}^n(\pi_{\text{BV}})$ satisfy quantum master-equation (40). Then the operator Ω^{\hbar} , defined in (41), squares to zero:

$$(\Omega^{\hbar})^2(\mathcal{O}) = 0.$$

Proof. We calculate, using Theorem 6,

$$\begin{aligned} (\Omega^{\hbar})^2(\mathcal{O}) &= \llbracket S^{\hbar}, \llbracket S^{\hbar}, \mathcal{O} \rrbracket - i\hbar \Delta \mathcal{O} \rrbracket - i\hbar \Delta(\llbracket S^{\hbar}, \mathcal{O} \rrbracket - i\hbar \Delta \mathcal{O}) \\ &= \llbracket S^{\hbar}, \llbracket S^{\hbar}, \mathcal{O} \rrbracket \rrbracket - i\hbar \llbracket S^{\hbar}, \Delta \mathcal{O} \rrbracket - i\hbar \llbracket \Delta S^{\hbar}, \mathcal{O} \rrbracket + i\hbar \llbracket S^{\hbar}, \Delta \mathcal{O} \rrbracket + (i\hbar)^2 \Delta^2 \mathcal{O}. \end{aligned}$$

The last term vanishes identically by Theorem 8, while the second term cancels against the fourth term. Using Jacobi’s identity (28) for the Schouten bracket on the first term, we obtain:

$$(\Omega^{\hbar})^2(\mathcal{O}) = -i\hbar \llbracket \Delta S^{\hbar}, \mathcal{O} \rrbracket + \frac{1}{2} \llbracket \llbracket S^{\hbar}, S^{\hbar} \rrbracket, \mathcal{O} \rrbracket = \llbracket -i\hbar \Delta S^{\hbar} + \frac{1}{2} \llbracket S^{\hbar}, S^{\hbar} \rrbracket, \mathcal{O} \rrbracket.$$

Now is the crucial moment in the entire proof. By the logic of our reasoning’s objective, the theorem’s claim is that the operator $(\Omega^{\hbar})^2$ yields zero whenever acting on a functional \mathcal{O} . We accordingly transform the variational Schouten bracket of two terms to the operator realization,

$$\cong \vec{Q}^{-i\hbar \Delta S^{\hbar} + \frac{1}{2} \llbracket S^{\hbar}, S^{\hbar} \rrbracket}(\mathcal{O}),$$

with the evolutionary derivation now acting on the argument. Let us emphasize that a transition from the variational Schouten bracket – which increases the number of bases $M \times \dots \times M$ by construction – to the evolutionary vector field chops a multiplication of geometries by uniquely fixing the field’s generating section.²⁵ But by our initial assumption, this generating section is zero by virtue of (40). Therefore the image of \mathcal{O} under such map vanishes, which proves the assertion. \square

²⁵ It might happen otherwise that a co-multiple of \mathcal{O} under \llbracket, \rrbracket looks like zero as a map of the space $\Gamma(\pi_{\text{BV}})$ yet the bracket with it could still be nonzero, see, e. g., ΔG on p. 36 in Example 2.4.

3.2. Gauge automorphisms of quantum BV-cohomology groups

By using the quantum BV-differential Ω^{\hbar} , let us construct a closed algebra of infinitesimal gauge symmetries for the quantum master-equation (40).

Proposition 13. Let $F \in \overline{\mathfrak{N}^n}(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be an arbitrary odd-parity functional and S^{\hbar} the quantum master-action satisfying (40). Then the infinitesimal shift of the functional S^{\hbar} ,

$$\dot{S}^{\hbar} = \Omega^{\hbar}(F) \iff S^{\hbar} \mapsto S^{\hbar}(\varepsilon) = S^{\hbar} + \varepsilon \cdot \Omega^{\hbar}(F) + \bar{o}(\varepsilon), \quad \varepsilon \in \mathbb{R}, \quad (43)$$

is a symmetry of (40) so that $\Delta(\exp(\frac{i}{\hbar}S^{\hbar}(\varepsilon))) = \bar{o}(\varepsilon)$ in Peano's notation.

- The algebra of infinitesimal gauge symmetries (43) of the quantum master-equation is closed,

$$\left(\frac{d}{d\varepsilon_1} \circ \frac{d}{d\varepsilon_2} - \frac{d}{d\varepsilon_2} \circ \frac{d}{d\varepsilon_1} \right) \Big|_{\varepsilon_i=0} (S^{\hbar}) = \Omega^{\hbar}(\llbracket F_1, F_2 \rrbracket), \quad (44)$$

i.e., the commutator of two even-parity symmetries with respective generators F_1 and F_2 is the infinitesimal gauge symmetry whose generator is the odd Poisson bracket of F_1 and F_2 .

Remark 3.2. The odd-parity generators $F_i \in \overline{\mathfrak{N}^n}(\pi_{\text{BV}}, T\pi_{\text{BV}})$ never evolve in the course of a transformation which is induced by any generator F_j on the quantum BV-action functional S^{\hbar} .

Proof. Assuming a smooth dependence of $S^{\hbar}(\varepsilon)$ on ε , we obtain that²⁶

$$\frac{d}{d\varepsilon} \Delta \left(\exp \left(\frac{i}{\hbar} S^{\hbar}(\varepsilon) \right) \right) = \frac{i}{\hbar} \dot{S}^{\hbar} \cdot \Delta \left(\exp \left(\frac{i}{\hbar} S^{\hbar}(\varepsilon) \right) \right) + \Omega^{\hbar}(\dot{S}^{\hbar}) \cdot \exp \left(\frac{i}{\hbar} S^{\hbar}(\varepsilon) \right).$$

Because $(\Omega^{\hbar})^2 = 0$ by Theorem 12, for \dot{S}^{\hbar} to be an infinitesimal symmetry of the equation $\Delta(\exp(\frac{i}{\hbar}S^{\hbar})) = 0$ it is sufficient that $\dot{S}^{\hbar} = \Omega^{\hbar}(F)$ for some odd-parity functional F .

Second, let

$$\frac{d}{d\varepsilon_i} (S^{\hbar}) = -i\hbar \Delta F_i + \llbracket S^{\hbar}, F_i \rrbracket \quad \text{for } i = 1, 2, \quad \varepsilon_i \in \mathbb{R},$$

and postulate that $\frac{d}{d\varepsilon_i}(F_j) \equiv 0$ for all i and j . Then commutator (44) of even-parity infinitesimal transformations (43) generated by the functionals F_1 and F_2 is

$$\begin{aligned} & \llbracket -i\hbar \Delta F_1 + \llbracket S^{\hbar}, F_1 \rrbracket, F_2 \rrbracket - \llbracket -i\hbar \Delta F_2 + \llbracket S^{\hbar}, F_2 \rrbracket, F_1 \rrbracket \\ & = -i\hbar (\llbracket \Delta F_1, F_2 \rrbracket - \llbracket \Delta F_2, F_1 \rrbracket) + \left(\llbracket \llbracket S^{\hbar}, F_1 \rrbracket, F_2 \rrbracket - \llbracket \llbracket S^{\hbar}, F_2 \rrbracket, F_1 \rrbracket \right). \end{aligned}$$

Because F_1 has odd parity, we swap the factors in $-\llbracket \Delta F_2, F_1 \rrbracket = \llbracket F_1, \Delta F_2 \rrbracket$; likewise, $+\llbracket F_1, \llbracket S^{\hbar}, F_2 \rrbracket \rrbracket$ is the last term in the above expression. From our main Theorem 6 and by Jacobi identity (28) we conclude that the commutator is equal to

$$-i\hbar \Delta(\llbracket F_1, F_2 \rrbracket) + \llbracket S^{\hbar}, \llbracket F_1, F_2 \rrbracket \rrbracket = \Omega^{\hbar}(\llbracket F_1, F_2 \rrbracket),$$

that is, the Schouten bracket of F_1 and F_2 is the new gauge symmetry generator. □

²⁶ This proof is standard: it originates from the cohomological deformation theory for solutions of the Maurer-Cartan equation (e. g., of (40)), see [37].

Remark 3.3. (cf. [51, §5]). The transformation $\exp\left(\frac{i}{\hbar}S^{\hbar}\right) \mapsto \exp\left(\frac{i}{\hbar}S^{\hbar}(\varepsilon)\right)$ for a finite $\varepsilon \in \mathbb{R}$ is determined by the operator $\exp(\varepsilon[\Delta, F])$, where $[\cdot, \cdot]$ is the anticommutator of two odd-parity objects. Indeed, by Theorem 3 we have that

$$\begin{aligned} &\Delta(F \cdot \exp\left(\frac{i}{\hbar}S^{\hbar}\right)) + F \cdot \Delta\left(\exp\left(\frac{i}{\hbar}S^{\hbar}\right)\right) \\ &= \Delta F \cdot \exp\left(\frac{i}{\hbar}S^{\hbar}\right) - \llbracket F, \exp\left(\frac{i}{\hbar}S^{\hbar}\right) \rrbracket - F \cdot \Delta\left(\exp\left(\frac{i}{\hbar}S^{\hbar}\right)\right) + F \cdot \Delta\left(\exp\left(\frac{i}{\hbar}S^{\hbar}\right)\right) \\ &= \frac{i}{\hbar}(-i\hbar \Delta F + \llbracket S^{\hbar}, F \rrbracket) \cdot \exp\left(\frac{i}{\hbar}S^{\hbar}\right) = \frac{i}{\hbar}\dot{S}^{\hbar} \cdot \exp\left(\frac{i}{\hbar}S^{\hbar}\right) = \left.\frac{d}{d\varepsilon}\right|_{\varepsilon=0} \left(\exp\left(\frac{i}{\hbar}S^{\hbar}\right)\right). \end{aligned}$$

Note that the Schouten bracket acts on $\exp\left(\frac{i}{\hbar}S^{\hbar}\right)$ by the Leibniz rule (see Theorem 4) and we then use the equality $-\llbracket F, \frac{i}{\hbar}S^{\hbar} \rrbracket = \frac{i}{\hbar}\llbracket S^{\hbar}, F \rrbracket$ which holds by Theorem 4 again.

Let us now regard the full quantum BV-action as the generating functional for ghost parity-even observables \mathcal{O} , see [54].

Lemma 14. There are no observables \mathcal{O} , other than the identically zero functional, which would be ghost parity-odd.

Proof. Indeed, Eq. (42) implies that the path integral

$$I = \int_{\Gamma(\zeta^0)} [D\mathbf{q}] \mathcal{O}([\mathbf{q}], [\mathbf{q}^\dagger]) \exp\left(\frac{i}{\hbar}S^{\hbar}([\mathbf{q}], [\mathbf{q}^\dagger])\right)$$

over the space of ghost parity-even BV-section components is effectively independent of the ghost parity-odd BV-variables \mathbf{q}^\dagger . Notice further that the ghost parity $\text{gh}(I)$ of this constant function $I([\mathbf{q}^\dagger])$ is equal to that of \mathcal{O} ; the quantum master-action S^{\hbar} is parity-even. Under a (speculative) assumption that an observable \mathcal{O} could be ghost parity-odd, we obtain an odd parity constant. Unless a possibility of their existence is postulated by brute force, this odd-parity constant must be equal to zero, whence the ghost parity-odd functional $\mathcal{O} \in \overline{H}^n(\pi_{\text{BV}}) \subseteq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ itself is zero. \square

In what follows we accept for transparency that there is no grading in the initial geometry of physical fields, i.e., for sections of the bundle $\pi: E^{n+m} \rightarrow M^n$. Let us focus on the standard cohomological approach to quantum BV-models and to their gauge symmetries (cf. [37]).

Lemma 15. Suppose that an infinitesimal shift $S^{\hbar} \mapsto S^{\hbar} + \lambda \cdot \mathcal{O} + \bar{o}(\lambda)$ of the quantum BV-action by using an even-parity functional $\mathcal{O} \in \overline{H}^n(\pi_{\text{BV}}) \subseteq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ does not destroy the quantum master-equation,

$$\left.\frac{d}{d\lambda}\right|_{\lambda=0} \Delta\left(\exp\left(\frac{i}{\hbar}(S^{\hbar} + \lambda \cdot \mathcal{O})\right)\right) = 0.$$

Then the observable \mathcal{O} is Ω^{\hbar} -closed: $-i\hbar \Delta\mathcal{O} + \llbracket S^{\hbar}, \mathcal{O} \rrbracket = 0$.

Proof. The proof literally repeats that of Proposition 13. \square

For a given odd-parity functional $F \in \overline{H}^n(\pi_{\text{BV}})$, we organize the infinitesimal shift (43) of the master-functional S^{\hbar} as follows:

$$\begin{aligned} \dot{S}^{\hbar} &= -i\hbar \Delta(F) + \llbracket S^{\hbar}, F \rrbracket, \\ \dot{\mathcal{O}} &= \llbracket \mathcal{O}, F \rrbracket. \end{aligned}$$

Note that, unless one has that $\Delta F = 0$ incidentally, the transformation of the integral *functional* S^{\hbar} is not induced by any infinitesimal transformation of the BV-variables, that is, by an evolutionary vector field on the horizontal infinite jet space at hand. No earlier than the transformation law $S^{\hbar} \mapsto S^{\hbar}(\varepsilon)$ is postulated, it becomes an act of will to think that the functional F is the generator of parity-preserving evolutionary vector field $\overleftarrow{Q}^F = \overrightarrow{Q}^F$ acting on the BV-variables so that $\dot{\mathcal{O}} \cong \overrightarrow{Q}^F(\mathcal{O})$ for all observables \mathcal{O} .

Furthermore, let us extend the deformation $\mathcal{O} \mapsto \mathcal{O}(\varepsilon)$ of even-parity cocycles $\mathcal{O} \in \ker \Omega^{\hbar}$ to the space of odd-parity functionals $\xi \in \overline{H}^n(\pi_{\text{BV}}) \subseteq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ which produce the coboundaries $\Omega^{\hbar}(\xi)$. Namely, we postulate that

$$\dot{\xi} = \llbracket \xi, F \rrbracket$$

for all such functionals ξ ; here we denote by the dot over ξ its velocity in the course of the transformation generated by a given F . Let us remember however that the law for evolution of the odd-parity functionals ξ which produce the Ω^{\hbar} -coboundaries is different from our earlier postulate (see Proposition 13) that the odd-parity generators F_i of gauge symmetries do not evolve: $dF_i/d\varepsilon_j \equiv 0$ or, in shorthand notation,

$$\dot{F} \equiv 0. \tag{45}$$

We claim that under these hypotheses, the structure of quantum BV-cohomology group remains intact in the course of gauge symmetry transformations of the quantum master-action, $S^{\hbar} \mapsto S^{\hbar}(\varepsilon)$, even though the quantum BV-differential is modified, $\Omega^{\hbar} \mapsto \Omega^{\hbar}(\varepsilon)$, and the cocycles and coboundaries are also deformed.

Theorem 16. *An infinitesimal shift of the quantum BV-cohomology classes induced by (43), (45), and*

$$\begin{aligned} \dot{\mathcal{O}} &= \llbracket \mathcal{O}, F \rrbracket, & \mathcal{O} &\in \ker \Omega^{\hbar}, \\ \dot{\xi} &= \llbracket \xi, F \rrbracket, & \xi &\in \overline{H}^n(\pi_{\text{BV}}) \subseteq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}), \xi \text{ odd}, \end{aligned}$$

yields an isomorphism of the Ω^{\hbar} -cohomology group: under such mapping, every Ω^{\hbar} -closed, even-parity Ω^{\hbar} -cocycle \mathcal{O} becomes $\Omega^{\hbar}(\varepsilon)$ -closed, whereas the transformation of an even-parity coboundary $\Omega^{\hbar}(\xi)$ produces an $\Omega^{\hbar}(\varepsilon)$ -coboundary: $(\Omega^{\hbar}(\xi))(\varepsilon) = \Omega^{\hbar}(\varepsilon)(\xi(\varepsilon))$.

Proof. Let $\mathcal{O} \in \ker \Omega^{\hbar}$ be an even-parity observable and F an odd-parity generator of gauge transformation. Consider the equation $\Omega^{\hbar}(\varepsilon)(\mathcal{O}(\varepsilon)) = 0$ which states that the transformed functional $\mathcal{O}(\varepsilon)$ remains a coboundary. The term which is proportional to ε in this equation's left-hand side is equal to

$$\left. \frac{d}{d\varepsilon} \right|_{\varepsilon=0} \left(-i\hbar \Delta \mathcal{O}(\varepsilon) + \llbracket S^{\hbar}(\varepsilon), \mathcal{O}(\varepsilon) \rrbracket \right) = \Omega^{\hbar}(\dot{\mathcal{O}}) + \llbracket \dot{S}^{\hbar}, \mathcal{O} \rrbracket = \Omega^{\hbar}(\llbracket \mathcal{O}, F \rrbracket) + \llbracket \Omega^{\hbar}(F), \mathcal{O} \rrbracket;$$

recalling once again that $\Omega^{\hbar} = -i\hbar \Delta + \llbracket S^{\hbar}, \cdot \rrbracket$, we continue the equality

$$= -i\hbar \Delta(\llbracket \mathcal{O}, F \rrbracket) + \llbracket S^{\hbar}, \llbracket \mathcal{O}, F \rrbracket \rrbracket + \llbracket -i\hbar \Delta F + \llbracket S^{\hbar}, F \rrbracket, \mathcal{O} \rrbracket.$$

Now by Theorem 6 we obtain that, the observable \mathcal{O} being parity-even,

$$\begin{aligned} &= -i\hbar \llbracket \Delta \mathcal{O}, F \rrbracket + i\hbar \llbracket \mathcal{O}, \Delta F \rrbracket + \llbracket S^{\hbar}, \llbracket \mathcal{O}, F \rrbracket \rrbracket - i\hbar \llbracket \Delta F, \mathcal{O} \rrbracket + \llbracket \mathcal{O}, \llbracket S^{\hbar}, F \rrbracket \rrbracket = \\ &= \llbracket \Omega^{\hbar}(\mathcal{O}), F \rrbracket \cong -\overrightarrow{Q}^F(\Omega^{\hbar}(\mathcal{O})) = 0, \end{aligned}$$

because $[[S^{\hbar}, [\mathcal{O}, F]]] = [[[S^{\hbar}, \mathcal{O}], F] - [\mathcal{O}, [S^{\hbar}, F]]]$ by Jacobi identity (28), because we are inspecting the ε -linear term in the operator $\Omega^{\hbar}(\varepsilon) \circ (\varepsilon = 0 \mapsto \varepsilon \neq 0)$ applied to \mathcal{O} , and \mathcal{O} is an Ω^{\hbar} -cocycle. Therefore, the zero initial condition $\Omega^{\hbar}(\mathcal{O}) = 0$ evolves at zero velocity to the $\Omega^{\hbar}(\varepsilon)$ -cocycle equation $\Omega^{\hbar}(\varepsilon)(\mathcal{O}(\varepsilon)) = 0$ upon $\mathcal{O}(\varepsilon)$.

Likewise, let $\Omega^{\hbar}(\xi)$ be a coboundary for some odd-parity functional ξ which evolves by $\dot{\xi} = [[\xi, F]]$. Then the even-parity observable $\Omega^{\hbar}(\xi) \in \ker \Omega^{\hbar}$ evolves as fast as $[[\Omega^{\hbar}(\xi), F]]$ but simultaneously we have that the mapping Ω^{\hbar} and its argument ξ change. We claim that the two evolutions match so that $(\Omega^{\hbar}(\xi))(\varepsilon)$ is $\Omega^{\hbar}(\varepsilon)$ -exact. Indeed, we have that

$$\begin{aligned} \frac{d}{d\varepsilon} \Big|_{\varepsilon=0} (\Omega^{\hbar}(\varepsilon)(\xi(\varepsilon))) &= \Omega^{\hbar}([[\xi, F]]) + [[\Omega^{\hbar}(F), \xi] \\ &= \underline{-i\hbar} [[\Delta \xi, F] - \underline{i\hbar} [[\xi, \Delta F] + [[S^{\hbar}, [\xi, F]] + \underline{[-i\hbar \Delta F + [S^{\hbar}, F], \xi]}; \end{aligned}$$

by cancelling out the underlined Schouten brackets and then using the Jacobi identity we obtain

$$= \underline{[-i\hbar \Delta \xi, F]} + [[[S^{\hbar}, \xi], F] + [[\xi, [S^{\hbar}, F]] - [[\xi, [S^{\hbar}, F]] = [[\Omega^{\hbar}(\xi), F],$$

which proves our claim.

Summarizing, we see that gauge symmetries of the quantum master-equation induce automorphisms of the Ω^{\hbar} -cohomology group. \square

We conclude that it would be conceptually incorrect to say that the infinitesimal gauge transformations of all functionals in a quantum BV-model are induced by a canonical transformation, determined by the evolutionary vector field \vec{Q}^F acting on the BV-variables. Let us remember that the even-parity quantum master-action $S^{\hbar} \in \overline{H}^n(\pi_{\text{BV}})$ and its descendants, the observables \mathcal{O} evolve by

$$\dot{S}^{\hbar} = -i\hbar \Delta F + [[S^{\hbar}, F] = \Omega^{\hbar}(F), \quad F \in \overline{H}^n(\pi_{\text{BV}}) \subseteq \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}}), \quad F \text{ odd,}$$

and

$$\dot{\mathcal{O}} = [[\mathcal{O}, F].$$

We note that the evolution of the generating functional S^{\hbar}_{BV} is **not** determined by a vector field on the space of BV-variables. Likewise, we recall that the odd-parity arguments ξ of Ω^{\hbar} for the coboundaries $\Omega^{\hbar}(\xi) \sim 0$ do evolve,

$$\dot{\xi} = [[\xi, F],$$

whereas the generators F of gauge symmetries for (40) never change: symbolically,

$$\dot{F} = 0$$

(see Eq. (45) above). In fact, one may think that each F determines a parity-preserving evolutionary vector field \vec{Q}^F on the space of BV-variables, but it is not the objects \vec{Q}^F but the full systems of four distinct evolution equations which encode the deformation of respective functionals. Neither the functionals' attribution to the space of building blocks $\overline{H}^n(\pi_{\text{BV}}) \ni S^{\hbar}$, \mathcal{O} , F nor a functional's parity, $\text{gh}(S^{\hbar}) = \text{gh}(\mathcal{O})$ and $\text{gh}(F) = \text{gh}(\xi)$, completely determines their individual transformation laws.

Remark 3.4. The supports of test shifts δs can be arbitrarily small²⁷ and they can be chosen in such a way that all boundary terms vanish in the course of integration by parts within equivalence classes from the horizontal cohomology groups $\overline{H}^{n(1+k)}(\pi_{\text{BV}} \times T\pi_{\text{BV}} \times \dots \times T\pi_{\text{BV}})$. Let us note also that these integrations by parts (see section 1.3) transport the derivatives from one copy of the base manifold M^n to another copy; this reasoning stays local with respect to base points \mathbf{x} and local volume elements $\text{dvol}(\mathbf{x})$ because the geometric mechanism of locality yields the diagonal in powers of the base manifold. However, an integration by parts in functionals from $\overline{H}^n(\pi_{\text{BV}})$ is a different issue. In fact, it refers to the topology of M^n or to a choice of the class $\Gamma(\pi_{\text{BV}})$ of admissible sections (so that there appear no boundary terms as well). Let us recall that the only place where such global, de Rham cohomology aspect is explicitly used is the proof of Jacobi’s identity for the variational Schouten bracket (see [32]). In turn, Theorems 8 and 12 relate these properties of the bracket \llbracket, \rrbracket to cohomology generators $\Delta^2 = 0$ and $(\Omega^h)^2 = 0$. (The converse is also true: Jacobi’s identity for \llbracket, \rrbracket stems from $\Delta^2 = 0$.) This motivates why the de Rham and quantum BV-cohomologies are interrelated (cf. [5]).

Conclusion

Mathematical models are designed for description of phenomena of Nature; a construction of the models’ objects is not the same as their evaluation at given configurations of the models, which would associate \mathbb{k} -numbers to physical fields $\phi \in \Gamma(\pi)$ in terms of such objects. Namely, consider an Euler–Lagrange model whose primary element is the action functional $S: \Gamma(\pi) \rightarrow \mathbb{k}$. By definition, derivative objects are obtained from S by using natural operations such as $\overleftarrow{\delta}$ or \llbracket, \rrbracket and Δ . The derivative objects’ geometric complexity is greater than that of S because they absorb the domains of definition for test shifts $\delta s_1, \dots, \delta s_k$ of field configurations. We emphasize that such composite structure objects do not yet become maps $\Gamma(\pi) \rightarrow \mathbb{k}$ which would suit well for their evaluation at sections $s \in \Gamma(\pi)$ yielding \mathbb{k} -numbers. The intermediate objects can rather be used as arguments of \llbracket, \rrbracket or Δ in a construction of larger, logically and geometrically more complex objects; we illustrate by Fig. 9 the expansion of analytic structures and their shrinking




Figure 9. The action S as a generator of observables, building blocks of derivative objects as horizontal cohomology classes in products of bundles over $M \times M \times \dots \times M$, and resulting mappings as the objects’ contractions over Whitney’s sum of bundles.

in the course of integration by parts and multiplication of normalized test shifts in reconfigured couplings. Indeed, the derivative objects become multi-linear maps with respect to k -tuples of the variations $\delta s_1, \dots, \delta s_k \in \Gamma(T\pi)$ only when the integrations by parts carry all derivatives away from the test shifts, channelling the derivations to densities of the object’s constituent blocks such as the Lagrangian in the action functional. A surgery of couplings then contracts

²⁷ We recall that the smoothness class of variations δs is determined by smoothness of the frame fields $\vec{e}_i(\mathbf{x})$, $\vec{e}^{\dagger i}(\mathbf{x})$ and coefficient functions $\delta s^i(\mathbf{x})$, $\delta s_i^{\dagger}(\mathbf{x})$.

the values of normalized test shifts by virtue of (16) at every point of the base manifold. This is how maps $\Gamma(\pi) \rightarrow \mathbb{k}$ are obtained.

We conclude that a calculation of composite-structure object may not be interrupted ahead of time. Otherwise speaking, every calculation stretches from its input data to the end value at \mathbf{s} ; independently existing values at \mathbf{s} for the resulting object's constituent elements not always contribute to the sought-for value of the large structure (e.g., consider (1c) on p. 1 and Example 2.4 on p. 34 and try to calculate consecutively the objects ΔF , ΔG , and their Schouten brackets with G and F , respectively, for that example's functionals F and G). Summarizing, it is illegal to construct composite objects step by step, redundantly inspecting the elements' values at field configurations. One must not deviate from a way towards the appointed end of logical reasoning.

In fact, it is us but not Nature who calculates (e.g., the left-hand sides of equations of motion): Nature neither calculates nor evaluates; for there is no built-in mechanism for doing that.²⁸ This implies that there is no ever-growing logical complexity in a description of the Universe; the flow of local, observer-dependent time does not require any perpetual increase of the number $k \geq 0$ of factors in the product-bundle location of objects over $k + 1$ copies of the space-time. Conversely, there always remains a unique copy of the space-time for all local functionals.

The space-time geometry of information transfer is very restrictive: its pointwise locality of events of couplings between dual objects is an absolute principle; by weakening this hypothesis one could create a source of difficulties through causality violation. Consequently, a count of space-time points where the couplings with a given (co)vector occur makes the formalism of singular linear integral operators truly adequate in mathematical models of physical phenomena.²⁹

We finally remark that the product-base approach of bundles $\pi \times T\pi \times \dots \times T\pi$ over $M \times M \times \dots \times M$ to the geometry of variations highlights the concept of physical field as infinite-dimensional system with degrees of freedom which are attached at every point of space-time. The locality principle for (co)vector interaction is the mechanism which distinguishes between space-time points with respect to its (non)Hausdorff topology.

Discussion

Let us finally address two logical aspects of the geometry of variations.

Linear vector space structures Nature is essentially nonlinear; for there is no mechanism which would realize – under a uniform time bound – an arbitrarily large number of replications of an object. This is tautological for those physical fields ϕ which take values in spaces without any linear structure. Moreover, even if there is a brute force labelling of Euler–Lagrange equations by using the fields ϕ , a linear vector space pattern of the equations of motion is not utilized (the same is true for the equations' descendants such as the antifields ϕ^\dagger or (anti)ghosts). Indeed, it

²⁸ The probabilistic approach to evolution of Nature suggests that maxima of transition (and correlation) functions concentrate near the zero loci of such deterministic equations' left-hand sides. At the same time, Noether symmetries of the action S are abundant in the models. Not referring to any actual transformation of a system's components, such symmetries reflect the model's geometry. The analytic machinery of self-regularizing structures yields the invariants – e.g., cohomology classes as in section 3.2 – which constrain the probabilistic laws of evolution.

²⁹ We recall from Remark 1.5 on p. 9 that the volume elements $\text{dvol}(\mathbf{x}, \phi(\mathbf{x})) = \sqrt{|\det(g_{\mu\nu})|} d\mathbf{x}$ are present in the building blocks of composite-structure objects. Let us note further that an association of the weight factors $\text{dvol}(\mathbf{x})$ with point $\mathbf{x} \in M^n$ is intrinsically related to the structure of space-time M^n as topological manifold (cf. [31]). It is readily seen that a discrete tiling of space-time converts the integrations over a measure on it to weighted sums over the points which mark the quantum domains. This links the concept with loop quantum gravity (see e.g. [17, 47, 49]).

is only their the *tangent* spaces whose linear structure is used, in particular, in order to split the variations in ghost parity-homogeneous components. Objects are linearized only in the course of variations under infinitesimal test shifts. For example, this determines the distinction between finite offsets $\Delta\mathbf{x}$ so that $(\mathbf{x}, \mathbf{x} + \Delta\mathbf{x}) \in M \times M$ and infinitesimal test shifts $\delta\mathbf{x}|_{\mathbf{x}} \in T_{\mathbf{x}}M$ which are mapped to the number field \mathbb{k} by covectors $d\mathbf{x}|_{\mathbf{x}} \in T_{\mathbf{x}}^*M$.

Annual reproduction rate for interspecimen breeding of cats and whales An immediate comment on the title of this paragraph is as follows. One could proclaim that the annual reproduction rate for interspecimen breeding of – without loss of generality – cats and whales is equal to zero for a given year. Alternatively, one should understand that such events never happen (not that a given year brought no brood).

This grotesque illustration works equally well for the (co)tangent spaces to fibres of the BV-zoo or, in broad terms, for a definition of Kronecker’s symbol δ_i^j by zero whenever the indices $i \neq j$ do not match so that the couplings in (11) do not eventuate. We argue that, on top of the absolute pointwise locality for couplings (9), a superficial definition of $\langle \cdot, \cdot \rangle$ by zero for mismatching elements \vec{e}_i and $\vec{e}^{\dagger j}$ of dual bases is a mere act of will; in reality those evaluations do not occur. Consequently, the geometry dictates that

$$\log \langle \vec{e}_i(\mathbf{x}), \dagger(\vec{e}_j)(\mathbf{x}) \rangle = \log 1 = 0 \quad \text{and} \quad \log \langle \vec{e}^{\dagger j}(\mathbf{x}), \dagger(\vec{e}^{\dagger i})(\mathbf{x}) \rangle = \log 1 = 0.$$

Combined with the geometric locality principle (4) realized by singular linear integral operators (12), this argument finally resolves the paradoxical, *ad hoc* conventions $\delta(0) = 0$ and $\log \delta(0) = 0$ for Dirac’s distribution.

Acknowledgments

The author thanks the Organizing committee of XXI International conference ‘Integrable systems & quantum symmetries’ (June 11–16, 2013; CVUT Prague, Czech Republic) for cooperation and warm atmosphere during the meeting. These notes follow the lecture course which was read by the author in October 2013 at the Taras Shevchenko National University and Bogolyubov Institute for Theoretical Physics in Kiev, Ukraine; the author is grateful to BITP for hospitality. The author thanks M. A. Vasiliev and A. G. Nikitin for helpful discussions and constructive criticisms.

This research was supported in part by JBI RUG project 103511 (Groningen). A part of this research was done while the author was visiting at the IHÉS (Bures-sur-Yvette); the financial support and hospitality of this institution are gratefully acknowledged.

Appendix A. Proof of Propositions 10 and 11

We need the following two lemmas.

Lemma 17. Let $F \in \overline{H}^n(\pi_{\text{BV}})$ be an even integral functional, let $G \in \overline{\mathfrak{N}}^n(\pi_{\text{BV}}, T\pi_{\text{BV}})$ be another functional, and let $n \in \mathbb{N}_{\geq 1}$. Then

$$\llbracket G, F^n \rrbracket = n \llbracket G, F \rrbracket F^{n-1}.$$

Proof. We use induction on Theorem 4. Note that all signs vanish since F is even, meaning that whenever F is multiplied with any other integral functional, the factors may be freely swapped without this resulting in minus signs. For $n = 1$ the statement is trivial. Suppose the formula holds for some $n \in \mathbb{N}_{>1}$, then $\llbracket G, F^{n+1} \rrbracket =$

$$\llbracket G, F \cdot F^n \rrbracket = \llbracket G, F \rrbracket \cdot F^n + F \cdot \llbracket G, F^n \rrbracket = \llbracket G, F \rrbracket \cdot F^n + nF \cdot \llbracket G, F \rrbracket F^{n-1} = (n + 1) \llbracket G, F \rrbracket \cdot F^n,$$

so that the statement also holds for $n + 1$. □

Lemma 18. Let $F \in \overline{H}^n(\pi_{\text{BV}})$ be an even integral functional, and let $n \in \mathbb{N}_{\geq 2}$. Then

$$\Delta(F^n) = n(\Delta F) \cdot F^{n-1} + \frac{1}{2}n(n-1)[[F, F]] \cdot F^{n-2}.$$

Proof. We use induction and the previous lemma. For $n = 2$ the formula clearly holds by Theorem 3. Suppose that it holds for some $n \in \mathbb{N}_{>2}$, then

$$\begin{aligned} \Delta(F^{n+1}) &= \Delta(F \cdot F^n) = (\Delta F) \cdot F^n + [[F, F^n]] + F \cdot \Delta(F^n) \\ &= (\Delta F) \cdot F^n + n[[F, F]] \cdot F^{n-1} + F \cdot n(\Delta F)F^{n-1} + \frac{1}{2}n(n-1)F \cdot [[F, F]]F^{n-2} \\ &= (n+1)(\Delta F) \cdot F^n + \frac{1}{2}(n+1)n[[F, F]] \cdot F^{n-1}, \end{aligned}$$

so that the statement also holds for $n + 1$. □

Proof of Proposition 10. For convenience, we denote $F = \frac{i}{\hbar}S^{\hbar}$. Then

$$\begin{aligned} 0 &= \Delta(\exp F) = \Delta\left(\sum_{n=0}^{\infty} \frac{1}{n!}F^n\right) = \sum_{n=0}^{\infty} \frac{1}{n!}\Delta(F^n) \\ &= \sum_{n=1}^{\infty} \frac{n}{n!}(\Delta F) \cdot F^{n-1} + \sum_{n=2}^{\infty} \frac{1}{2n!}n(n-1)[[F, F]] \cdot F^{n-2} \\ &= (\Delta F) \cdot \sum_{n=1}^{\infty} \frac{1}{(n-1)!}F^{n-1} + \frac{1}{2}[[F, F]] \cdot \sum_{n=2}^{\infty} \frac{1}{(n-2)!}F^{n-2} \\ &= (\Delta F + \frac{1}{2}[[F, F]]) \cdot \exp F = \left(\frac{i}{\hbar}\Delta S^{\hbar} - \frac{1}{2\hbar^2}[[S^{\hbar}, S^{\hbar}]]\right) \cdot \exp\left(\frac{i}{\hbar}S^{\hbar}\right), \end{aligned}$$

from which the result follows. □

Proof of Proposition 11 (cf. Proposition 13 on p. 42). Again, let us set $F = \frac{i}{\hbar}S^{\hbar}$. We first calculate, using Lemma 17,

$$[[\mathcal{O}, \exp F]] = \sum_{n=0}^{\infty} \frac{1}{n!}[[\mathcal{O}, F^n]] = \sum_{n=1}^{\infty} \frac{n}{n!}[[\mathcal{O}, F]]F^{n-1} = [[\mathcal{O}, F]] \exp F.$$

Then

$$\begin{aligned} 0 &= \Delta(\mathcal{O} \exp F) = (\Delta \mathcal{O}) \exp F + [[\mathcal{O}, \exp F]] + \mathcal{O} \cdot \Delta(\exp F) \\ &= (\Delta \mathcal{O} + [[\mathcal{O}, F]]) \exp F = \left(\Delta \mathcal{O} + \frac{i}{\hbar}[[\mathcal{O}, S^{\hbar}]]\right) \exp\left(\frac{i}{\hbar}S^{\hbar}\right), \end{aligned}$$

from which the assertion follows. □

References

- [1] Alexandrov M, Schwarz A, Zaboronsky O and Kontsevich M 1997 *Int. J. Modern Phys. A* vol **12** 1405–29
- [2] Arnol'd V I 1996 *Mathematical methods of classical mechanics* (Grad. Texts in Math. vol **60**) (New York: Springer-Verlag)
- [3] Barnich G 2000 *Classical and quantum aspects of the extended antifield formalism* (These d'agregation ULB Proc. spring school 'QFT and Hamiltonian systems' (Calimanesti, Romania) pp 1–94 (Preprint hep-th/0011120))
- [4] Barnich G 2010 (AIP Conf. Proc. vol **1307**) *XXIX Workshop on Geometric Methods in Physics* (Białowieża, Poland) 7–18 (Preprint math-ph/1010.0899)

- [5] Barnich G, Brandt F and Henneaux M 1995 *Commun. Math. Phys.* vol **174** 57–91, 93–116 (*Preprint hep-th/9405109*)
- [6] Barnich G, Brandt F and Henneaux M 2000 *Phys. Rep.* **338** 439–569 (*Preprint hep-th/0002245*)
- [7] Batalin I and Vilkovisky G 1981 *Phys. Lett. B* vol **102** 27–31
Batalin I A and Vilkovisky G A 1983 *Phys. Rev. D* vol **29** 2567–82
- [8] Becchi C, Rouet A and Stora R 1976 *Ann. Phys.* **98** 287–321
Tyutin I V 1975 Gauge invariance in field theory and statistical mechanics *Preprint Lebedev FIAN* no 39
- [9] Beilinson A and Drinfeld V 2004 *Chiral algebras* (AMS Colloq. Publ. vol **51**) (Providence, RI: AMS)
- [10] Berezin F A 1987 *Introduction to superanalysis* (Math. Phys. & Appl. Math. vol **9**) (Dordrecht: D. Reidel Publ. Co.)
- [11] Cattaneo A S and Felder G 2000 *Commun. Math. Phys.* vol **212** 591–611
- [12] Costello K J 2007 Renormalisation and the Batalin–Vilkovisky formalism (*Preprint* 0706.1533)
- [13] Costello K 2011 *Renormalization and effective field theory* (Math. Surveys and Monographs vol **170**) (Providence, RI: AMS)
- [14] Dorfman I Ya 1993 *Dirac structures* (J. Wiley & Sons)
- [15] Ehresmann C 1953 *Geometrie Differentielle, Colloq. Inter. du Centre Nat. de la Recherche Scientifique (Strasbourg)* 97–110
- [16] Eisenbud D 2005 *The geometry of syzygies. A second course in commutative algebra and algebraic geometry* (Grad. Texts in Math. vol **229**) (New York: Springer–Verlag)
- [17] Gambini R and Pullin J 1996 *Loops, knots, gauge theories and quantum gravity* (Cambridge Monographs on Math. Phys.) (Cambridge: Cambridge Univ. Press)
- [18] Gel’fand I M and Dorfman I Ja 1981 *Functional Anal. Appl.* vol **14** 223–6
- [19] Gel’fand I M and Shilov G E 1964 *Generalized functions* vol **1** Properties and operations (NY–London: Academic Press)
Gel’fand I M and Shilov G E 1968 *Generalized functions* vol **2** Spaces of fundamental and generalized functions (NY–London: Academic Press)
- [20] Gitman D M and Tyutin I V 1990 *Quantization of fields with constraints* (Springer Ser. Nucl. Part. Phys.) (Berlin: Springer-Verlag)
- [21] Gomis J, París J and Samuel S 1995 *Phys. Rep.* vol **259** 1–145 (*Preprint hep-th/9412228*)
- [22] Henneaux M and Teitelboim C 1992 *Quantization of gauge systems* (Princeton, NJ: Princeton Univ. Press)
- [23] Igonin S, Verbovetsky A and Vitolo R 2002 On the formalism of local variational differential operators *Memorandum* 1641 (Faculty of Math. Sci., Univ. Twente, Enschede, The Netherlands)
- [24] Igonin S, Verbovetsky A and Vitolo R 2004 *Symmetry in nonlinear mathematical physics (Kiev, 2003) Pr. Inst. Mat. Nats. Akad. Nauk Ukr. Mat. Zastos.* vol **50** 1335–42
- [25] de Jonghe F, Siebelink R and Troost W 1993 Hiding anomalies *Phys. Lett. B* vol **306** 295–300
- [26] Kersten P, Krasil’shchik I and Verbovetsky A 2004 *J. Geom. Phys.* vol **50** 273–302
- [27] Khudaverdian H M and Voronov Th Th 2013 *J. Geom. Phys.* vol **64** 31–53
- [28] Kiselev A V 2012 The twelve lectures in the (non)commutative geometry of differential equations *Preprint IHÉS/M/12/13*
- [29] Kiselev A V 2012 *Physics of Particles and Nuclei* vol **43** 663–5 (*Preprint* 1112.5784 [math-ph])
- [30] Kiselev A V 2012 *J. Phys.: Conf. Series* **343** 012058, 1–20 (*Preprint* 1111.3272)
- [31] Kiselev A V 2013 *Proc. 6th Int. Workshop ‘Group analysis of differential equations and integrable systems’ (Protaras, Cyprus)* 111–26 (*Preprint* 1304.0234)
- [32] Kiselev A V 2012 The calculus of multivectors on noncommutative jet spaces *Preprint* 1210.0726 [math.DG]
- [33] Kiselev A V and Krutov A O 2013 Non-Abelian Lie algebroids over jet spaces *Preprint* 1305.4598 [math.DG]
- [34] Kiselev A V and van de Leur J W 2011 *Theoret. and Mathem. Phys.* vol **167** 772–84 (*Preprint* 1006.4227 [math.DG])
- [35] Kiselev A V and Ringers S 2013 On the geometry of the Batalin–Vilkovisky Laplacian *Preprint* 1302.4388 [math.DG]
- [36] Kontsevich M 1993 *The Gel’fand Mathematical Seminars, 1990–1992* eds L Corwin, I Gelfand and J Lepowsky (Boston MA: Birkhäuser) 173–87
- [37] Kontsevich M and Soibelman Y 2009 *Homological mirror symmetry. New developments and perspectives* (Lect. Notes in Phys. vol **757**) eds A Kapustin, M Kreuzer and K-G Schlesinger (Berlin: Springer) 153–219
- [38] Kontsevich M and Vishik S 1994 Determinants of elliptic pseudo-differential operators *Preprint hep-th/9404046*
- [39] Kosmann-Schwarzbach Y 2008 *SIGMA* vol **4**, 5 pp 1–30 (*Preprint* 0710.3098)
- [40] Krasil’shchik I S and Vinogradov A M eds 1999 *Symmetries and conservation laws for differential equations of mathematical physics* (Providence, RI: AMS)

- [41] Kupershmidt B A 1980 *Geometric methods in mathematical physics* (Lecture Notes in Math. vol **775**) *Proc. NSF-CBMS Conf. (Univ. Lowell, Mass.)* (Berlin: Springer) 162–218
 - [42] Magri F 1978 *J. Math. Phys.* vol **19** 1156–62
 - [43] McCloud P 1994 *Class. Quant. Grav.* vol **11** 567–87
 - [44] Merkulov S and Willwacher Th 2010 Grothendieck–Teichmüller and Batalin–Vilkovisky *Preprint* 1012.2467 [math.QA]
 - [45] Olver P J 1993 *Applications of Lie groups to differential equations* (Grad. Texts in Math. vol **107**) 2nd ed (New York: Springer-Verlag)
 - [46] Olver P J and Sokolov V V 1998 *Comm. Math. Phys.* vol **193** 245–68
 - [47] Rovelli C 2004 *Quantum gravity* (Cambridge Monographs on Math. Phys.) (Cambridge: Cambridge Univ. Press)
 - [48] Schwarz A 1993 *Commun. Math. Phys.* vol **155** 249–60 (*Preprint* hep-th/9205088)
 - [49] Thiemann Th 2007 *Modern canonical quantum general relativity* (Cambridge Monographs on Math. Phys.) (Cambridge: Cambridge Univ. Press)
 - [50] Troost W, van Nieuwenhuizen P and van Proeyen A 1990 *Nucl. Phys. B* vol **333** 727–70
 - [51] Voronov B L, Tyutin I V and Shakhverdiev Sh S 1999 *Theoret. and Mathem. Phys.* vol **120** 1026–44 (*Preprint* hep-th/9904215)
 - [52] Voronov T 2002 *Quantization, Poisson brackets, and beyond* (Contemp. Math. vol **315**) ed T Voronov (Providence, RI: AMS) pp 131–68 (*Preprint* math.DG/0105237)
 - [53] Witten E 1990 *Modern Phys. Lett. A* **5** 487–94
 - [54] Zinn-Justin J 1993 *Quantum field theory and critical phenomena* (Int. Ser. of Monographs on Phys. vol **85**) 2nd ed (Oxford: Oxford Univ. Press; New York: The Clarendon Press)
 - [55] Zinn-Justin J 1975 *Trends in Elementary Particle Theory* (Lect. Notes in Phys. vol **37**) eds H Rollnick and K. Dietz (Berlin: Springer) pp 2–39
- Zinn-Justin J 1976 *Méthodes en théorie des champs / Methods in field theory* (École d’Été de Physique Théorique, Session XXVIII, tenue à Les Houches, 28 Juillet–6 Septembre 1975) eds R Balian and J Zinn-Justin (Amsterdam: North-Holland Publ. Co.)