

## CĂLDĂRARU'S CONJECTURE AND TSYGAN'S FORMALITY

DAMIEN CALAQUE, CARLO A. ROSSI, AND MICHEL VAN DEN BERGH

ABSTRACT. In this paper we complete the proof of Căldăraru's conjecture on the compatibility between the module structures on differential forms over poly-vector fields and on Hochschild homology over Hochschild cohomology. In fact we show that twisting with the square root of the Todd class gives an isomorphism of precaluli between these pairs of objects.

Our methods use formal geometry to globalize the local formality quasi-isomorphisms introduced by Kontsevich and Shoikhet (the existence of the latter was conjectured by Tsygan). We also rely on the fact - recently proved by the first two authors - that Shoikhet's quasi-isomorphism is compatible with cap product after twisting with a Maurer-Cartan element.

## CONTENTS

1. Introduction and statement of the main results	2
1.1. The Atiyah and Todd class of a Lie algebroid	2
1.2. Gerstenhaber algebras and precaluli	3
1.3. Poly-vector fields, poly-differential operators, differential forms and Hochschild chains in the Lie algebroid framework	3
1.4. Main results	4
1.5. Căldăraru's conjecture	6
2. Notation and conventions	7
3. Some recollections on Lie algebroids and related topics	7
3.1. Generalities on Lie algebroids	7
3.2. Relative poly-vector fields, poly-differential operators	11
3.3. The sheaf of $L$ -poly-differential operators	11
3.4. The Hochschild complex of $L$ -chains over $R$	12
3.5. The Hochschild–Kostant–Rosenberg Theorem in the Lie algebroid framework	14
4. Fedosov resolutions in the Lie algebroid framework	15
4.1. Introduction	15
4.2. Setup	15
4.3. The (affine) coordinate space of a Lie algebroid	16

---

1991 *Mathematics Subject Classification*. Primary 14F99, 14D99.

*Key words and phrases*. Căldăraru's conjecture, Tsygan formality.

This paper was partly written when the first author was a visitor at the Department of Mathematics of ETH (on leave of absence from Université Lyon 1) thanks to the support of the European Union through a Marie Curie Intra-European Fellowship (contract number MEIF-CT-2007-042212).

The second author has been partially supported by the Fundação para a Ciência e a Tecnologia (FCT / Portugal) while writing the present paper.

The third author is a director of research at the FWO.

4.4.	Fedosov resolutions of $L$ -poly-vector fields and $L$ -poly-differential operators on $R$	18
4.5.	The Fedosov resolution of $L$ -forms on $R$	18
4.6.	The Fedosov resolution of $L$ -chains on $R$	18
5.	Globalization of Tsygan’s formality in the Lie algebroid framework	20
5.1.	Descent for $L_\infty$ -algebras and $L_\infty$ -modules.	20
5.2.	Twisting of $L_\infty$ -algebras and $L_\infty$ -modules	21
5.3.	Compatibility of twisting and descent	22
5.4.	Kontsevich’s and Shoikhet’s formality theorems	23
5.5.	Formality theorem in the ring case	24
5.6.	Functoriality property of Theorem 5.3	28
5.7.	Proof of Theorem 1.3	32
6.	The relationship between Atiyah classes and jet bundles	33
7.	Proof of Theorem 1.1	34
7.1.	A <i>memento</i> of compatibility between cup and cap products	34
7.2.	The proof of Theorem 1.1 in the ring case	38
7.3.	Functoriality properties of the commutative diagram (7.5)	39
7.4.	Proof of Theorem 1.1 in the global case	41
Appendix A.	Explicit formulæ for the $B_\infty$ -structure on poly-differential operators	42
Appendix B.	The precalculus structure on $L$ -chains	44
References		47

## 1. INTRODUCTION AND STATEMENT OF THE MAIN RESULTS

Throughout  $k$  is a ground field of characteristic 0. In this introduction  $(X, \mathcal{O})$  is a ringed site<sup>1</sup> such that  $\mathcal{O}$  is a sheaf of commutative  $k$ -algebras. We fix in addition a Lie algebroid  $\mathcal{L}$  over  $(X, \mathcal{O})$ .

Roughly speaking a Lie algebroid is a sheaf of  $\mathcal{O}$ -modules which is also a sheaf of Lie algebras which acts on  $\mathcal{O}$  by derivations. See §3.1. Standard examples of Lie algebroids are the tangent bundle on a smooth manifold and the holomorphic tangent bundle on a complex manifold. Readers not familiar with Lie algebroids are advised to think of  $\mathcal{L}$  as a tangent bundle (holomorphic or not) for the rest of this introduction. Concepts like “connection” take their familiar meaning in this context. In fact: our main reason for working in the setting of Lie algebroids is that these allow us to treat the algebraic, holomorphic and  $C^\infty$ -cases in a uniform way.

**1.1. The Atiyah and Todd class of a Lie algebroid.** *From now on we make the additional assumption that the Lie algebroid  $\mathcal{L}$  is locally free of rank  $d$  as an  $\mathcal{O}$ -module.*

The Atiyah class  $A(\mathcal{L}) \in \text{Ext}^1(\mathcal{L}, \mathcal{L}^* \otimes \mathcal{L}) = \text{H}^1(X, \mathcal{L}^* \otimes \text{End}_{\mathcal{O}}(\mathcal{L}))$  of  $\mathcal{L}$  may for example be defined as the obstruction against the existence of a global  $\mathcal{L}$ -connection on  $\mathcal{L}$ . See §6 for more details.

---

<sup>1</sup>We work over sites instead of spaces to cover some additional cases which are important for algebraic geometry (like algebraic spaces and Deligne–Mumford stacks). Readers not interested in such generality may assume that  $(X, \mathcal{O})$  is just a ringed space.

The  $i$ -th scalar Atiyah class  $a_i(\mathcal{L})$  of  $\mathcal{L}$  is defined as

$$a_i(\mathcal{L}) = \text{tr}\left(\bigwedge^i A(\mathcal{L})\right) \in H^i(X, \bigwedge^i \mathcal{L}^*),$$

where  $\bigwedge^i$  is the map

$$\bigwedge^i : (\mathcal{L}^* \otimes \text{End}(\mathcal{L}))^{\otimes i} \rightarrow \bigwedge^i \mathcal{L}^* \otimes \text{End}(\mathcal{L})$$

given by composition on  $\text{End}(\mathcal{L})^{\otimes i}$  and the exterior product on  $(\mathcal{L}^*)^{\otimes i}$  and where  $\text{tr}$  is the usual trace on  $\text{End}(\mathcal{L})$ , extended linearly to a map  $\bigwedge^i \mathcal{L}^* \otimes \text{End}(\mathcal{L}) \rightarrow \bigwedge^i \mathcal{L}^*$ .

The Todd class  $\text{td}(\mathcal{L})$  of  $\mathcal{L}$  is derived from the Atiyah class  $A(\mathcal{L})$  by the following familiar formula:

$$(1.1) \quad \text{td}(\mathcal{L}) = \det\left(\frac{A(\mathcal{L})}{1 - \exp(-A(\mathcal{L}))}\right) \in \bigoplus_{i \geq 0} H^i(X, \bigwedge^i \mathcal{L}^*),$$

where the function

$$(1.2) \quad q(x) = \frac{x}{1 - \exp(-x)}$$

is extended to  $\bigwedge \mathcal{L}^* \otimes_{\mathcal{O}} \text{End}(\mathcal{L})$  via its formal Taylor expansion. In this way the Todd class  $\text{td}(\mathcal{L})$  of  $\mathcal{L}$  can be expressed in terms of the scalar Atiyah classes of  $\mathcal{L}$ .

**1.2. Gerstenhaber algebras and precalculi.** By definition a Gerstenhaber algebra is a graded vector space equipped with a Lie bracket  $[-, -]$  of degree zero and a commutative, associative cup product  $\cup$  of degree one<sup>2</sup> such that the Leibniz rule is satisfied

$$[a, b \cup c] = [a, b] \cup c + (-1)^{|a|(|b|+1)} b \cup [a, c]$$

If  $A$  is a Gerstenhaber algebra then a precalculus [12] over  $A$  is a quadruple  $(A, M, \iota, L)$  where  $M$  is a graded vector space and  $\iota : A \otimes M \rightarrow M$  and  $L : A \otimes M \rightarrow M$  are linear maps of degree 1 and 0 respectively such that  $\iota$  makes  $M$  into an  $(A[-1], \cup)$ -module and  $L$  makes  $M$  into an  $(A, [-, -])$ -Lie module and such that the following compatibilities hold for  $a, b \in A$

$$(1.3) \quad \iota_a L_b - (-1)^{(|a|+1)|b|} L_b \iota_a = \iota_{[a, b]}$$

$$(1.4) \quad L_a \iota_b + (-1)^{|a|+1} \iota_a L_b = L_{a \cup b}$$

A precalculus is *not* the same as a Gerstenhaber module. The second equation in the previous display is not correct for a Gerstenhaber module.

Below  $\iota$  will be referred to as “contraction” and  $L$  as the “Lie derivative”. Furthermore we will often write  $a \cap m$  for  $\iota_a(m)$  and as such refer to it as the “cap product”.

**1.3. Poly-vector fields, poly-differential operators, differential forms and Hochschild chains in the Lie algebroid framework.** For a Lie algebroid  $\mathcal{L}$  the sheaves of  $\mathcal{L}$ -poly-vector fields and  $\mathcal{L}$ -differential forms are defined as

$$T_{\text{poly}}^{\mathcal{L}}(X) = \bigoplus_{n \geq -1} \bigwedge^{n+1} \mathcal{L}, \quad \Omega^{\mathcal{L}}(X) = \bigoplus_{n \leq 0} \bigwedge^{-n} \mathcal{L}^*$$

where the wedge products are taken over  $\mathcal{O}_X$ .

<sup>2</sup>Note that our grading conventions are shifted with respect to the usual ones.

The sheaf  $T_{\text{poly}}^{\mathcal{L}}(X)$  becomes a sheaf of Gerstenhaber algebras when endowed with the trivial differential, the Lie algebroid version of the Schouten–Nijenhuis Lie bracket and the exterior product. Our grading convention is such that the Lie bracket and wedge product are of degree 0 and 1 respectively.

We equip  $\Omega^{\mathcal{L}}(X)$  with the trivial(!) differential<sup>3</sup>, and also with the contraction operator and Lie derivative with respect to  $\mathcal{L}$ -poly-vector fields. In this way the pair  $(T_{\text{poly}}^{\mathcal{L}}(X), \Omega^{\mathcal{L}}(X))$  becomes a sheaf of precalculi. In our conventions the contraction operator and Lie derivative have degrees 1 and 0 respectively.

The Lie algebroid generalization of the sheaf of  $\mathcal{L}$ -poly-differential operators is denoted by  $D_{\text{poly}}^{\mathcal{L}}(X)$  [2, 28]. It is the tensor algebra over  $\mathcal{O}$  of the universal enveloping algebra of  $\mathcal{L}$  (see §3.3 below).

The sheaf  $D_{\text{poly}}^{\mathcal{L}}(X)$  has similar properties as the standard sheaf of poly-differential operators on  $X$  (see e.g. [17]). In particular it is a differential graded Lie algebra (shortly, from now on, a DG-Lie algebra) and also a Gerstenhaber algebra *up to homotopy*. For the definition of the differential, the Lie bracket (of degree 0) and the cup product (of degree 1) see §3.3.

The sheaf of  $\mathcal{L}$ -Hochschild chains  $C_{\text{poly}}^{\mathcal{L}}(X)$  may be defined as the  $\mathcal{O}$ -dual of  $D_{\text{poly}}^{\mathcal{L}}(X)$  (although we use a slightly different approach). Furthermore there is a differential  $b_H$  as well as actions  $\cap, \mathbb{L}$  of  $D_{\text{poly}}^{\mathcal{L}}(X)$  on  $C_{\text{poly}}^{\mathcal{L}}(X)$  which make the pair  $(D_{\text{poly}}^{\mathcal{L}}(X), C_{\text{poly}}^{\mathcal{L}}(X))$  into a precalculus up to homotopy. We refer to §3.4.

Finally, we recall that there is a Hochschild–Kostant–Rosenberg (HKR for short) quasi-isomorphism from  $T_{\text{poly}}^{\mathcal{L}}(X)$  to  $D_{\text{poly}}^{\mathcal{L}}(X)$ ; dually, there is a HKR quasi-isomorphism from  $C_{\text{poly}}^{\mathcal{L}}(X)$  to  $\Omega^{\mathcal{L}}(X)$ . As in the classical case where  $\mathcal{L}$  is the tangent bundle neither of these HKR quasi-isomorphisms is compatible with the Gerstenhaber and precalculus structures up to homotopy.

**1.4. Main results.** Now we consider the derived category  $D(X)$  of sheaves of  $k$ -vector spaces over  $X$ . When equipped with the derived tensor product this becomes a symmetric monoidal category. Furthermore, viewed as objects in  $D(X)$ , both  $T_{\text{poly}}^{\mathcal{L}}(X)$  and  $D_{\text{poly}}^{\mathcal{L}}(X)$  are honest Gerstenhaber algebras and their combination with  $\Omega^{\mathcal{L}}(X)$  and  $C_{\text{poly}}^{\mathcal{L}}(X)$  yields precalculi.

Our first main result relates the Todd class of a Lie algebroid (as discussed in §1.1) to the failure of the HKR isomorphisms to preserve these precalculus structures.

**Theorem 1.1.** *Let  $\mathcal{L}$  be a locally free Lie algebroid of rank  $d$  over the ringed site  $(X, \mathcal{O}_X)$ . Then we have the following commutative diagram of precalculi in the category  $D(X)$ :*

$$(1.5) \quad \begin{array}{ccc} T_{\text{poly}}^{\mathcal{L}}(X) & \xrightarrow{\text{HKR} \circ \iota_{\sqrt{\text{td}(\mathcal{L})}}} & D_{\text{poly}}^{\mathcal{L}}(X) \\ \Downarrow & & \Downarrow \\ \Omega^{\mathcal{L}}(X) & \xleftarrow{(\sqrt{\text{td}(\mathcal{L})} \wedge -) \circ \text{HKR}} & C_{\text{poly}}^{\mathcal{L}}(X), \end{array}$$

<sup>3</sup>The De Rham differential  $d_L$  on  $\Omega^{\mathcal{L}}(X)$  is not part of the precalculus structure. In the operadic setting of [12],  $d_L$  appears as a unary operation and not as a differential.

where the vertical arrows indicate actions and the horizontal arrows are isomorphisms. Here  $\wedge$  denotes the left multiplication in  $\Omega^\mathcal{L}(X)$  and  $\iota$  denotes the contraction action of  $\Omega^\mathcal{L}(X)$  on  $T^\mathcal{L}(X)$ .<sup>4</sup>

The convention that wavy arrows indicate actions will be used throughout below.

The following corollary will be applied to Căldăraru's conjecture below.

**Corollary 1.2.** *There is a commutative diagram of precalculi:*

$$(1.6) \quad \begin{array}{ccc} \bigoplus_{m,n \geq 0} H^m(X, \wedge^n \mathcal{L}) & \xrightarrow{\text{HKR} \circ \iota \circ \sqrt{\text{td}(\mathcal{L})}} & \mathbb{H}^\bullet(X, D_{\text{poly}}^\mathcal{L}(X)) \\ \downarrow \wr & & \downarrow \wr \\ \bigoplus_{m,n \geq 0} H^m(X, \wedge^n \mathcal{L}^*) & \xleftarrow{(\sqrt{\text{td}(\mathcal{L}) \wedge -} \circ \text{HKR})} & \mathbb{H}^\bullet(X, C_{\text{poly}}^\mathcal{L}(X)). \end{array}$$

with  $\mathbb{H}^\bullet(X, -)$  denoting the hypercohomology functor.

*Proof.* This follows by applying the functor  $\mathbb{H}^\bullet(X, -)$  to the commutative diagram (1.5).  $\square$

If we consider only the Lie brackets and the Lie algebra actions then the horizontal isomorphisms in the commutative diagram (1.5) are obtained from the horizontal arrows in diagram (1.7) below, which is part of our second main result:

**Theorem 1.3.** *Assume that  $\mathbb{R} \subset k$ . Let  $\mathcal{L}$  be a locally free Lie algebroid of rank  $d$  over the ringed site  $(X, \mathcal{O})$ . There exist sheaves of differential graded Lie algebras  $(\mathfrak{g}_i^\mathcal{L}, d_i, [\ , \ ]_i)$  and sheaves of DG-Lie modules  $(\mathfrak{m}_i^\mathcal{L}, b_i, L_i)$  over them as well as  $L_\infty$ -quasi-isomorphisms  $\mathfrak{U}_\mathcal{L}$  from  $\mathfrak{g}_1^\mathcal{L}$  to  $\mathfrak{g}_2^\mathcal{L}$  and  $\mathfrak{S}_\mathcal{L}$  from  $\mathfrak{m}_2^\mathcal{L}$  to  $\mathfrak{m}_1^\mathcal{L}$ , which fit into the following commutative diagram:*

$$(1.7) \quad \begin{array}{ccccccc} T_{\text{poly}}^\mathcal{L}(X) & \hookrightarrow & \mathfrak{g}_1^\mathcal{L} & \xrightarrow{\mathfrak{U}_\mathcal{L}} & \mathfrak{g}_2^\mathcal{L} & \longleftarrow & D_{\text{poly}}^\mathcal{L}(X) \\ \downarrow \wr & & \downarrow \wr & & \downarrow \wr & & \downarrow \wr \\ \Omega^\mathcal{L}(X) & \hookrightarrow & \mathfrak{m}_1^\mathcal{L} & \xleftarrow{\mathfrak{S}_\mathcal{L}} & \mathfrak{m}_2^\mathcal{L} & \longleftarrow & C_{\text{poly}}^\mathcal{L}(X) \end{array}$$

where the hooked arrows are strict (i.e. DG-Lie) quasi-isomorphisms.

1.4.1. *Comments on the results and the proofs.* The proof of Theorems 1.1 and 1.3 depends on the simultaneous globalization of a number of local formality results due to Kontsevich [17] (see also [18]), Tsygan [26], Shoikhet [21] and the first two authors [4–6]. This globalization is performed by a functorial version of formal geometry [7] (see also [29]).

The proof of Theorem 1.1 roughly speaking involves the construction of a morphism up to homotopy between the precalculus structures up to homotopy on  $(T_{\text{poly}}^\mathcal{L}(X), \Omega^\mathcal{L}(X))$  and  $(D_{\text{poly}}^\mathcal{L}(X), C_{\text{poly}}^\mathcal{L}(X))$ . In this paper we do not construct a full “precalculus $_\infty$ ”-quasi-isomorphism between these structures (in the case that  $\mathcal{L}$  is a tangent bundle this has been done in [12] using operadic methods; actually, in *loc. cit.*, the authors work in the “calculus $_\infty$ ” setting, encoding also the De Rham

<sup>4</sup>Note that normally we view  $\Omega^\mathcal{L}(X)$  as a module over  $T^\mathcal{L}(X)$ . In the definition of the horizontal arrows in the diagram (1.5) the opposite actions appear for reasons that are mysterious to the authors.

differential, which is not part of the precalculus structure as observed before). On the other hand, in contrast to *loc. cit.*, the results we prove are explicit and this fact is essential to recover Căldăraru's conjecture as formulated in [9] (see Theorem 1.4 below).

We are able to obtain such explicit results by starting with the local quasi-isomorphisms of Kontsevich and Shoikhet which are given by explicit formulæ (in contrast to, say, Tamarkin's local  $G_\infty$ -quasi-isomorphism [24]). While these are *a priori* only  $L_\infty$ -quasi-isomorphisms they are nonetheless compatible with products up to homotopy [6, 17] in a strong explicit sense and this turns out to be enough for our purposes.

As the local quasi-isomorphisms of Kontsevich and Shoikhet are defined over  $\mathbb{R}$  (see §5.4) we have to assume  $\mathbb{R} \subset k$  in the statement of Theorem 1.3. However enough coefficients are rational (and computable), which in turn allows us to prove Theorem 1.1 over an arbitrary field of characteristic zero. This idea was already used in [7]. See §7.3.1. For Theorem 1.3 we could likely have started with a Tamarkin-style local quasi-isomorphism [24] defined over  $\mathbb{Q}$ , but since the coefficients of such a local quasi-isomorphism are not explicit, the result would not be immediately applicable to Theorem 1.1.

The existence of the upper horizontal isomorphism in (1.5) has been proved independently in [7, 11], while its explicit form has been computed in [7]. The existence of the lower horizontal isomorphism has been shown in [12]. As observed above, our approach *via* Kontsevich's and Shoikhet's local formality formulæ allows us to compute it explicitly.

**1.5. Căldăraru's conjecture.** Assume now that  $X$  is a smooth algebraic or complex variety. Căldăraru's conjecture (stated originally in the algebraic case) asserts the existence of various compatibilities between the Hochschild (co)homology and tangent (co)homology of  $X$  (see below). For the full statement we refer to [9]. The results in this paper complete the proof of Căldăraru's conjecture.

We now explain this in more detail. The Hochschild (co)homology [23] of  $X$  is defined as

$$\begin{aligned} \mathrm{HH}^n(X) &= \mathrm{Ext}_{\mathcal{O}_{X \times X}}^n(\mathcal{O}_\Delta, \mathcal{O}_\Delta) & (n \geq 0) \\ \mathrm{HH}_n(X) &= \mathrm{Tor}_{-n}^{\mathcal{O}_{X \times X}}(\mathcal{O}_\Delta, \mathcal{O}_\Delta) & (n \leq 0) \end{aligned}$$

where  $\Delta \subset X \times X$  is the diagonal. From these definitions it is clear that  $\mathrm{HH}^\bullet(X)$  has a canonical algebra structure (by the Yoneda product) and  $\mathrm{HH}_\bullet(X)$  is a module over it.

Furthermore if we put  $\mathcal{L} = T_X$  then it is proved in [8] (and partially in [30]) that there are isomorphisms of algebras and modules

$$\begin{array}{ccc} \mathrm{HH}^\bullet(X) & \longrightarrow & \mathbb{H}^\bullet(X, D_{\mathrm{poly}}^\mathcal{L}(X)) \\ \downarrow \wr & & \downarrow \wr \\ \mathrm{HH}_\bullet(X) & \longrightarrow & \mathbb{H}^\bullet(X, C_{\mathrm{poly}}^\mathcal{L}(X)) \end{array}$$

where on the right-hand side we consider only the part of the precalculus given by the cup and cap product.

We define the tangent (co)homology of  $X$  by

$$\mathrm{HT}^\bullet(X) = \bigoplus \mathrm{H}^\bullet(X, \bigwedge^\bullet T_X), \quad \mathrm{H}\Omega_\bullet(X) = \bigoplus \mathrm{H}^\bullet(X, \Omega_X^{-\bullet}).$$

where now  $\Omega_X^\bullet$  denotes the graded sheaf of differential forms on  $X$ .

The commutative diagram (1.6) then yields the following

**Theorem 1.4** (“Căldăraru’s conjecture”). *For a smooth algebraic or complex variety  $X$  over  $k$  there is a commutative diagram of  $k$ -algebras and modules*

$$(1.8) \quad \begin{array}{ccc} \mathrm{HT}^\bullet(X) & \xrightarrow{\mathrm{HKR} \circ \iota_{\sqrt{\mathrm{td}(X)}}} & \mathrm{HH}^\bullet(X) \\ \Downarrow & & \Downarrow \\ \mathrm{H}\Omega_\bullet(X) & \xleftarrow{(\sqrt{\mathrm{td}(X)} \wedge -) \circ \mathrm{HKR}} & \mathrm{HH}_\bullet(X), \end{array}$$

where  $\mathrm{td}(X)$  is the Todd class for  $\mathcal{L} = T_X$ .

Theorem 1.4 completes the proof of the parts of Căldăraru’s conjecture [9] which do not depend on  $X$  being proper. The cohomological part (the upper row in the above diagram) had already been proved in [7] and is also an unpublished result of Kontsevich.

In the proper case there is an additional assertion in Căldăraru’s conjecture which involves the natural bilinear form on  $\mathrm{HH}_\bullet(X)$ . We do not consider this assertion in the present paper as it has already been proved by Markarian [19] and Ramadoss [20]. If we combine Theorem 1.8 with the results of Markarian and Ramadoss we obtain a full proof of Căldăraru’s conjecture. Let us also mention that in the compact Calabi-Yau case Căldăraru’s conjecture has been proved in [15].

*Acknowledgment.* We express our gratitude to the anonymous referees for the careful reading of the manuscript. Their comments have helped us improve the paper.

## 2. NOTATION AND CONVENTIONS

As stated already we always work over a ground field  $k$  of characteristic 0; unadorned tensor products are over  $k$ .

Most objects we consider are equipped with a topology which will be explicitly specified when needed. However if an object is introduced without a specific topology, or if the topology is not clear from the context, then it is assumed to be equipped with the discrete topology.

Many objects we will encounter are  $\mathbb{Z}$ -graded. Koszul’s sign rule is always assumed in this context. For a double or higher complex we apply the sign rule with respect to total degree.

## 3. SOME RECOLLECTIONS ON LIE ALGEBROIDS AND RELATED TOPICS

**3.1. Generalities on Lie algebroids.** In this section  $R$  is a commutative  $k$ -algebra.

**Definition 3.1.** A **Lie algebroid**  $L$  over  $R$  is a Lie algebra over  $k$  which is in addition an  $R$ -module and is endowed with an anchor map  $\rho : L \rightarrow \mathrm{Der}_k(R)$  satisfying the compatibility

$$(3.1) \quad [l_1, r l_2] = \rho(l_1)(r) l_2 + r [l_1, l_2], \quad r \in R, \quad l_i \in L, \quad i = 1, 2.$$

The basic example of a Lie algebroid over  $R$  is  $L = \text{Der}_k(R)$  with the identity anchor map and the commutator Lie bracket.

If  $L$  is a Lie algebroid then  $R \oplus L$  is a Lie algebra with Lie bracket  $[(r, l), (r', l')] = (\rho(l)(r') - \rho(l')(r), [l, l'])$ . We define the **universal enveloping algebra**  $U_R(L)$  of  $L$  to be the quotient of the enveloping algebra associated to the Lie algebra  $R \oplus L$  by the relation  $r \otimes l = rl$  ( $r \in R, l \in R \oplus L$ ).

For the sake of simplicity, below we will usually omit the anchor map  $\rho$  from the notation, unless it is necessary for the sake of clarity.

The universal enveloping algebra of a Lie algebroid satisfies a universal property similar to that of an ordinary enveloping algebra. This implies for example that the anchor map  $\rho$  uniquely extends to an algebra morphism from  $U_R(L)$  to  $\text{End}_k(R)$ , or equivalently: it yields a left  $U_R(L)$ -module structure on  $R$ . *For reasons which will become clear later we assume that our Lie algebroids are free of rank  $d$  over  $R$ .*

**3.1.1.  $L$ -poly-vector fields and  $L$ -differential forms over  $R$ .** To a Lie algebroid  $L$  over  $R$  we associate

$$(3.2) \quad T_{\text{poly}}^L(R) = \bigoplus_{n \geq -1} \wedge_R^{n+1} L,$$

$$(3.3) \quad \Omega^L(R) = \bigoplus_{n \leq 0} \wedge_R^{-n} L^*, \quad L^* = \text{Hom}_R(L, R).$$

We refer to (3.2) and (3.3) as the spaces of  **$L$ -poly-vector fields** and  **$L$ -forms** on  $R$ .

As an exterior algebra  $T_{\text{poly}}^L(R)$  has a wedge product which we denote by  $\cup$  (“the cup product”). The extension of the Lie bracket on  $L$  to a bi-derivation on  $T_{\text{poly}}^L(R)$  defines a Lie bracket which is called the Schouten–Nijenhuis bracket and is denoted by  $[-, -]$ . Note that with our grading conventions the cup product has degree one and the Lie bracket has degree zero. The cup product and the Lie bracket make  $T_{\text{poly}}^L(R)$  into a (shifted) Gerstenhaber algebra with trivial differential.

On the other hand,  $\Omega^L(R)$  is obviously a graded algebra with respect to the wedge product. In addition there is an analogue  $d_L$  of the De Rham differential on  $\Omega^L(R)$ , which is given on generators by

$$\begin{aligned} d_L(r)(l) &= l(r) \\ d_L(l^*)(l_1, l_2) &= l_1(l^*(l_2)) - l_2(l^*(l_1)) - l^*([l_1, l_2]), \end{aligned}$$

for  $r \in R, l, l_i \in L, i = 1, 2, l^* \in L^*$ , and is extended uniquely by Leibniz’s rule.

The natural contraction operation of  $L$ -forms on  $R$  with respect to  $L$ -poly-vector fields is denoted by  $\cap$  (the “cap product”). The Lie derivative  $L$  of  $L$ -forms on  $R$  with respect to  $L$ -poly-vector fields is specified in the usual way *via* Cartan’s homotopy formula as the commutator of  $d_L$  and the contraction. The pair

$$((T_{\text{poly}}^L(R), [-, -], \cup), (\Omega^L(R), \cap, L))$$

forms a precalculus (see §1.2).

**3.1.2.  $L$ -connections.** As usual  $L$  is a Lie algebroid over  $R$ .

**Definition 3.2.** Let  $M$  be an  $R$ -module  $M$ . An  **$L$ -connection** on  $M$  is a  $k$ -linear map  $\nabla$  from  $M$  to  $L^* \otimes_R M$ , which satisfies Leibniz’s rule

$$(3.4) \quad \nabla(rm) = d_L(r) \otimes m + r\nabla m, \quad r \in R, \quad m \in M.$$



The  $L$ -connection  $\nabla$  is said to be **flat**, if  $\nabla^2 = 0$ . Equivalently, the assignment  $l \mapsto \nabla_l$ , where  $\nabla_l$  denotes the action of  $\nabla$  followed by contraction with respect to  $l$ , defines a Lie algebra morphism from  $L$  to  $\text{End}_k(M)$ .

If we let  $l \in L$  act as  $\nabla_l$  then a flat  $L$ -connection on  $M$  extends to a left  $U_R(L)$ -module structure on  $M$ .

Furthermore a flat  $L$ -connection  $\nabla$  on  $M$  can be extended to a differential (denoted by the same symbol) on the graded  $R$ -module  $\Omega^L(R) \otimes_R M$  via Leibniz's rule

$$\nabla(\omega \otimes_R m) = d_L \omega \otimes_R m + (-1)^{|\omega|} \omega \wedge \nabla m, \quad \omega \in \Omega^L(R), \quad m \in M.$$

**3.1.3.  $L$ -differential operators over  $R$ .** In this section we define the algebra of poly-differential operators of a Lie algebroid and we list some of its properties. We give some explicit formulæ along the lines of [6].

As in the case of ordinary Lie algebras  $U_R(L)$  (see §3.1) may be naturally filtered by giving  $R$  filtered degree 0 and  $L$  filtered degree 1. In particular

$$F_0 U_R(L) = R, \quad F_1 U_R(L) = R \oplus L,$$

We view  $U_R(L)$  as an  $R$ -central bimodule via the natural embedding of  $R$  into  $U_R(L)$ . Explicitly, if we denote this embedding by  $i$  then<sup>5</sup>

$$(3.5) \quad rD \stackrel{!}{=} Dr \stackrel{\text{def}}{=} i(r)D, \quad r \in R, D \in U_R(L)$$

Moreover  $U_R(L)$  is an  $R$ -coalgebra [28], i.e.  $U_R(L)$  possesses an  $R$ -linear coproduct  $\Delta : U_R(L) \rightarrow U_R(L) \otimes_R U_R(L)$  and an  $R$ -linear counit, satisfying the usual axioms. The comultiplication actually takes values in

$$\begin{aligned} & (U_R(L) \otimes_R U_R(L))' \\ & = \left\{ \sum_j D_j \otimes E_j \in U_R(L) \otimes_R U_R(L) \mid \forall r \in R : \sum_j D_j i(r) \otimes E_j = \sum_j D_j \otimes E_j i(r) \right\} \end{aligned}$$

which is an  $R$ -algebra even though  $U_R(L) \otimes_R U_R(L)$  is not.

The comultiplication  $\Delta$  and counit  $\epsilon$  are given by similar formulæ as in the Lie algebra case

$$(3.6) \quad \begin{aligned} \Delta(r) &= r \otimes_R 1 = 1 \otimes_R r & r \in R \\ \Delta(l) &= l \otimes_R 1 + 1 \otimes_R l & l \in L \\ \Delta(DE) &= D_{(1)} E_{(1)} \otimes D_{(2)} E_{(2)} & D, E \in U_R(L) \\ \epsilon(D) &= D(1) \end{aligned}$$

In the third formula we have used Sweedler's convention. The expression on the right-hand side is well defined because it is the product inside the algebra  $(U_R(L) \otimes_R U_R(L))'$ . In the fourth formula we have used the natural action of  $U_R(L)$  on  $R$  (see §3.1).

The algebra (better: in the terminology of [2, 28] “the Hopf algebroid”)  $U_R(L)$  may be thought of as an algebra of  $L$ -differential operators on  $R$ : in the case  $L = \text{Der}_k(R)$  and  $R$  smooth over  $k$  then  $U_R(L)$  coincides with the algebra of differential operators on  $R$ .

---

<sup>5</sup>Note that there is an at first sight more natural right  $R$ -module structure on  $U_R(L)$  given by the formula  $Dr = Di(r)$ . This alternative right module structure will not be used in this paper.

3.1.4. *L-jets*. Let  $(U_R(L))_{\leq n}$  be the elements of degree  $\leq n$  with respect to the canonical filtration on  $U_R(L)$  introduced in §3.1.3. The  $L$ - $n$ -jets are defined as

$$J^n L = \text{Hom}_R(U_R(L)_{\leq n}, R)$$

(this is unambiguous, as the left and right  $R$ -modules structures on  $U_R(L)$  are the same, see (3.5)). We also put

$$(3.7) \quad \mathcal{J}L = \text{Hom}_R(U_R(L), R) = \text{proj} \lim_n J^n L \quad (\text{as } U_R(L) = \text{inj} \lim_n (U_R(L))_{\leq n}).$$

$\mathcal{J}L$  has a natural commutative algebra structure obtained from the comultiplication on  $U_R(L)$ . Thus for  $\phi_1, \phi_2 \in \mathcal{J}L$ ,  $D \in U_R(L)$  we have

$$(\phi_1 \phi_2)(D) = \phi_1(D_{(1)}) \phi_2(D_{(2)}),$$

and the unit in  $\mathcal{J}L$  is given by the counit on  $U_R(L)$ .

In addition  $\mathcal{J}L$  has two commuting left  $U_R(L)$ -module structures which we now elucidate. First of all there are two distinct monomorphisms of  $k$ -algebras

$$\begin{aligned} \alpha_1 : R &\rightarrow \mathcal{J}L : r \mapsto (D \mapsto r\epsilon(D)), \\ \alpha_2 : R &\rightarrow \mathcal{J}L : r \mapsto (D \mapsto D(r)). \end{aligned}$$

It will be convenient to write  $R_i = \alpha_i(R)$  and to view  $\mathcal{J}L$  as an  $R_1 - R_2$ -bimodule.

There are also two distinct commuting actions by derivations of  $L$  on  $\mathcal{J}L$ . Let  $l \in L$ ,  $\phi \in \mathcal{J}L$ ,  $D \in U_R(L)$ .

$$\begin{aligned} {}^1\nabla_l(\phi)(D) &= l(\phi(D)) - \phi(lD) \\ {}^2\nabla_l(\phi)(D) &= \phi(Dl) \end{aligned}$$

Again it will be convenient to write  $L_i$  for  $L$  acting by  ${}^i\nabla$ . Then  ${}^i\nabla$  defines a flat  $L_i$ -connection on  $\mathcal{J}L$ , considered as an  $R_i$ -module. The connection  ${}^1\nabla$  is the well-known **Grothendieck connection**. It follows that  $\mathcal{J}L$  is a  $U_R(L)_1 - U_R(L)_2$ -bimodule (with both  $U_R(L)_1$  and  $U_R(L)_2$  acting on the left).

The  $U_R(L)_2$  action on  $\mathcal{J}L$  takes the very simple form

$$(D \cdot \phi)(E) = \phi(ED)$$

(for  $D, E \in U_R(L)_2$ ,  $\phi \in \mathcal{J}L$ ).

Define  $\epsilon : \mathcal{J}L \rightarrow R$  by  $\epsilon(\phi) = \phi(1)$  and put  $J^c L = \ker \epsilon$ . Then  $\mathcal{J}L$  is complete for the  $J^c L$ -adic topology and the filtration on  $\mathcal{J}L$  induced by (3.7) coincides with the  $J^c L$ -adic filtration. If we filter  $\mathcal{J}L$  with the  $J^c L$ -adic filtration then we obtain

$$(3.8) \quad \text{gr } \mathcal{J}L = S_R L^*$$

and the  $R_1$  and  $R_2$ -action on the r.h.s. of this equation coincide (here and below the letter  $S$  stands for ‘‘symmetric algebra’’).

The induced actions on  $\text{gr } \mathcal{J}L = S_R L^*$  of  $l \in L$ , considered as an element of  $L_1$  and  $L_2$ , are given by the contractions  $i_{-l}$  and  $i_l$ , respectively.

In case  $R$  is the coordinate ring of a smooth affine algebraic variety and  $L = \text{Der}_k(R)$  then we may identify  $\mathcal{J}L$  with the completion  $R\widehat{\otimes}R$  of  $R \otimes R$  at the kernel of the multiplication map  $R \otimes R \rightarrow R$ . The two actions of  $R$  on  $\mathcal{J}L$  are respectively  $R\widehat{\otimes}1$  and  $1\widehat{\otimes}R$ .

Similarly a derivation on  $R$  can be extended to  $R\widehat{\otimes}R$  in two ways by letting it act respectively on the first and second factor. Since derivations are continuous they act on adic completions and hence in particular on  $\mathcal{J}L$ . This provides the two actions of  $L$  on  $\mathcal{J}L$ .

In the sequel we will view the action labelled by “1” as the default action. I.e. we will usually not write the 1 explicitly.

**3.2. Relative poly-vector fields, poly-differential operators.** We need relative poly-differential operators and poly-vector fields. So assume that  $A \rightarrow B$  is a morphism of commutative  $k$ -algebras. Then

$$\begin{aligned} T_{\text{poly},A}(B) &= \bigoplus_{n \geq -1} T_{\text{poly},A}^n(B) \\ D_{\text{poly},A}(B) &= \bigoplus_{n \geq -1} D_{\text{poly},A}^n(B) \end{aligned}$$

where  $T_{\text{poly},A}^n(B) = \bigwedge_B^{n+1} \text{Der}_A(B)$ . Similarly,  $D_{\text{poly},A}^n(B) \subseteq \text{Hom}_A(B^{\otimes_A(n+1)}, B)$  consists of those  $A$ -linear maps from  $B^{\otimes_A(n+1)}$  to  $B$ , which are  $A$ -linear differential operators on  $B$  in each argument.

It is easy to see that  $T_{\text{poly},A}(B)$  is a Gerstenhaber algebra when equipped with the Schouten bracket and the exterior product. Similarly  $D_{\text{poly},A}(B)$  is a graded subspace of the relative Hochschild complex  $C_A^\bullet(B)$  and since differential operators are closed under composition one easily sees that it is in fact a sub- $B_\infty$ -algebra, see Appendix A for more details on  $B_\infty$ -algebras.

If  $A$  and  $B$  are DG-algebras then we equip  $T_{\text{poly},A}(B)$ ,  $D_{\text{poly},A}(B)$  with the total differentials  $[d_B, -]$  and  $[d_B, -] + d_H$  where  $d_H$  denotes the Hochschild differential. Similar results now apply.

### 3.3. The sheaf of $L$ -poly-differential operators.

**Definition 3.3.** For a Lie algebroid  $L$  over  $R$  we define the graded vector space  $D_{\text{poly}}^L(R)$  of  **$L$ -poly-differential operators on  $R$**  as the tensor algebra over  $R$  of  $U_R(L)$  with shifted degree, i.e.

$$D_{\text{poly}}^L(R) = \bigoplus_{n \geq -1} U_R(L)^{\otimes_R(n+1)}.$$

The action of  $U_R(L)$  on  $R$  extends to a map

$$(3.9) \quad D_{\text{poly}}^{L,n}(R) \rightarrow \text{Hom}_k(R^{\otimes n+1}, R)$$

defined by

$$(D_1 \otimes \cdots \otimes D_{n+1})(r_1 \otimes \cdots \otimes r_{n+1}) \mapsto D_1(r_1) \cdots D_{n+1}(r_{n+1})$$

whose image lies in the space  $D_{\text{poly}}(R)$  of poly-differential operators on  $R$ .

$D_{\text{poly}}^L(R)$  is a  $B_\infty$ -algebra. In particular it is a DG-Lie algebra and furthermore it is a Gerstenhaber algebra up to homotopy. In Appendix A we give the formulæ for the full  $B_\infty$ -structure. Here we content ourselves by reminding the reader of the basic operations.

The **Gerstenhaber bracket** on  $D_{\text{poly}}^L(R)$  is defined by

$$(3.10) \quad [D_1, D_2] = D_1\{D_2\} - (-1)^{|D_1||D_2|} D_2\{D_1\}, \quad D_i \in D_{\text{poly}}^L(R), \quad i = 1, 2.$$

where

$$D_1\{D_2\} = \sum_{i=0}^{|D_1|} (-1)^{i|D_2|} (\text{id}^{\otimes i} \otimes \Delta^{|D_2|} \otimes \text{id}^{\otimes |D_1|-i})(D_1) \cdot (1^{\otimes i} \otimes D_2 \otimes 1^{\otimes |D_1|-i}).$$

It is a Lie bracket of degree 0. The special element  $\mu = 1 \otimes_R 1 \in D_{\text{poly}}^{L,1}(R) = U_R(L) \otimes_R U_R(L)$  satisfies  $[\mu, \mu] = 0$ . The **Hochschild differential** is defined as the operator  $d_H = [\mu, -]$ .

The **cup product** on  $D_{\text{poly}}^L(R)$  is defined by

$$(3.11) \quad D_1 \cup D_2 = (-1)^{(|D_1|-1)(|D_2|-1)} D_1 \otimes_R D_2$$

(See also Appendix A for an explicit derivation of the previous Formula).

One may now show that these operations make the 4-tuple  $(D_{\text{poly}}^L(R), d_H, [\ , \ ], \cup)$  into a Gerstenhaber algebra up to homotopy (see Lemma A.1). Indeed if  $R$  is smooth over  $k$  and  $L = \text{Der}_k(R)$  is the tangent bundle then the operations we have defined are the same as those one obtains from the identification  $D_{\text{poly}}^L(R) = D_{\text{poly}}(R)$  where we view the right-hand side as a sub- $B_\infty$ -algebra of the Hochschild complex  $C^\bullet(R)$  of  $R$  (cfr. §3.2).

It is in fact, as we explain now, not necessary to verify that we have defined a homotopy Gerstenhaber structure on  $D_{\text{poly}}^L(R)$ . Indeed the results can be obtained directly from the known results for the Hochschild complex (see [13, 14]). Similarly it is not necessary to write explicit formulæ for  $[-, -]$  and  $\cup$  (or for the whole  $B_\infty$ -structure for that matter). This point of view will be useful when we consider Hochschild chains as in that case the formulæ become more complicated.

The  $L_2$ -action on  $JL$  commutes with the  $R_1$ -action (see §3.1.4) so we obtain a ring homomorphism

$$U_{R_2}(L_2) \rightarrow D_{R_1}(JL) : D \mapsto (\theta \mapsto D(\theta)).$$

and hence a map

$$(3.12) \quad D_{\text{poly}}^{L_2}(R_2) \rightarrow D_{\text{poly}, R_1}(JL).$$

of Gerstenhaber algebras up to homotopy. The right-hand side has an  $R_1$ -connection given by  $[\ ^1\nabla, -]$  and it follows from [7, Prop. 4.2.4, Lemma 4.3.4] that the left-hand side of (3.12) is given by the horizontal sections for this connection.

Now as discussed in §3.2, we know that  $D_{\text{poly}, R_1}(JL)$  is a  $B_\infty$ -algebra and it is an easy verification that the braces and the differential, which make up the  $B_\infty$ -structure, are horizontal for  $[\ ^1\nabla, -]$ . Hence the  $B_\infty$ -structure on  $D_{\text{poly}, R_1}(JL)$  descends to  $D_{\text{poly}}^L(R)$  and one verifies that its basic operations are indeed given by the formulæ we gave earlier.

**3.4. The Hochschild complex of  $L$ -chains over  $R$ .** We start with the following definition.

**Definition 3.4.** For a Lie algebroid  $L$  over  $R$ , the graded  $R$ -module

$$(3.13) \quad C_{\text{poly}, p}^L(R) = \begin{cases} JL^{\widehat{\otimes}_{R-p}}, & p < 0 \\ R, & p = 0, \end{cases}$$

is called the space of **Hochschild  $L$ -chains over  $R$** .

Our aim in this section will be to show that the pair

$$(D_{\text{poly}}^L(R), C_{\text{poly}}^L(R))$$

is a precalculus up to homotopy. We will do this without relying on explicit formulæ (as they are quite complicated). Instead we will reduce to a relative version

of [6] which discusses Hochschild (co)homology. Explicit formulæ are given in Appendix B.

Let us first remind the reader that if  $A$  is a  $k$ -algebra then the pair  $(C^\bullet(A), C_\bullet(A))$  consisting of the spaces of Hochschild cochains and chains is a precalculus up to homotopy. For  $C^\bullet(A)$  this is just the (shifted) homotopy Gerstenhaber structure which we have already mentioned in §3.3 and which was introduced in [13, 14].

The full precalculus structure up to homotopy on  $(C^\bullet(A), C_\bullet(A))$  is a more intricate object. A complete treatment in a very general setting has been given in [6]. It is shown that the precalculus structure can be obtained from two interacting  $B_\infty$ -module structures on  $C_\bullet(A)$ . These  $B_\infty$ -module structures are obtained from brace-type operations. For more operadic approaches see [12].

Although we do not really use them, for the benefit of the reader we state the well-known formulæ for the contraction, the Lie derivative and the differential. If  $P \in C^{m-1}(A) = \text{Hom}(A^{\otimes m}, A)$  and  $(a_0 | \cdots | a_t) \in C_{-t}(A) = A^{\otimes t+1}$  then we have

$$\begin{aligned} \iota_P(a_0 | \cdots | a_t) &= (a_0 P(a_1, \dots, a_m) | a_{m+1} | \cdots | a_t) \\ L_P(a_0 | \cdots | a_t) &= \sum_{i=0}^{t-m+1} (-1)^{(m-1)i} (a_0 | \cdots | a_{i-1} | P(a_i, \dots, a_{i+m-1}) | a_{i+m} | \cdots | a_t) \\ &\quad + \sum_{l=t-m+2}^{t+1} (-1)^{lt} (P(a_l, \dots, a_t, a_0, \dots, a_{m-t+l-2}) | a_{m-t+l-1} | \cdots | a_{l-1}) \end{aligned}$$

The differential  $b_H$  is defined as  $L_\mu$  where  $\mu$  is the multiplication, considered as an element of  $\text{Hom}(A^{\otimes 2}, A)$ .

To construct the precalculus structure up to homotopy on  $(D_{\text{poly}}^L(R), C_{\text{poly}}^L(R))$  we proceed as in §3.3. We first define an object that is larger than  $C_{\text{poly}}^L(R)$ .

**Definition 3.5.** The space of  $L$ -poly-jets over  $R$  is the *completed* space of relative Hochschild chains  $\widehat{C}_{R_1, \bullet}(JL)$ . Explicitly

$$(3.14) \quad \widehat{C}_{R_1, \bullet}(JL) = \bigoplus_{p \leq 0} JL^{\widehat{\otimes}_{R_1} -p-1}$$

The Grothendieck connection  ${}^1\nabla$  on  $JL$  (see §3.1.4) yields a connection on  $\widehat{C}_{R_1, \bullet}(JL)$  by Leibniz's rule to which we also refer to as the Grothendieck connection. The following result was proved in [3].

**Proposition 3.6.** *For a Lie algebroid  $L$  over a commutative ring  $R$  as above, there is an isomorphism of graded vector spaces*

$$(3.15) \quad \widehat{C}_{R_1, \bullet}(JL)^{{}^1\nabla} \rightarrow C_{\text{poly}}^L(R)$$

which sends

$$\phi_1 \otimes \phi_2 \otimes \cdots \otimes \phi_p \in \widehat{C}_{R_1, -p}(JL)^{{}^1\nabla}$$

to

$$\epsilon(\phi_1)\phi_2 \otimes \cdots \otimes \phi_p \in C_{\text{poly}, 1-p}^L(R)$$

*Proof.* The arguments of the proof of [3, Prop. 1.11], can be repeated almost *verbatim*.  $\square$

The formulæ from [6] for the Hochschild complexes now yield that

$$(D_{\text{poly}, R_1}(JL), \widehat{C}_{R_1, \bullet}(JL)) \subset (C_{R_1}^{\text{cont}, \bullet}(JL), \widehat{C}_{R_1, \bullet}(JL))$$

is a precalculus up to homotopy. Furthermore one verifies that the formulæ in [6] are compatible with the Grothendieck connection  ${}^1\nabla$ . Hence the precalculus descends to one on

$$(3.16) \quad (D_{\text{poly}, R_1}(JL)^{{}^1\nabla}, \widehat{C}_{R_1, \bullet}(JL)^{{}^1\nabla}) = (D_{\text{poly}}^L(R), C_{\text{poly}}^L(R))$$

where we use (3.12) as well as Proposition 3.6.

It remains to check that this construction coincides with the standard one for a smooth commutative algebra. Namely, if  $R/k$  is smooth and  $L$  is the tangent bundle then we have

$$D_{\text{poly}}^L(R) = D_{\text{poly}}(R)$$

We also have  $JL = R \widehat{\otimes} R$  (see §3.1.4) and in this way we obtain an isomorphism

$$(3.17) \quad C_{\text{poly}, -p}^L(R) = (R \widehat{\otimes} R)^{\widehat{\otimes}_{R_1} p} \rightarrow R^{\widehat{\otimes} p+1} : (r_1 \widehat{\otimes} s_1) \widehat{\otimes} \cdots \widehat{\otimes} (r_p \widehat{\otimes} s_p) \mapsto (r_1 \cdots r_p) \widehat{\otimes} s_1 \widehat{\otimes} \cdots \widehat{\otimes} s_p$$

which yields an isomorphism of graded vector spaces

$$C_{\text{poly}}^L(R) = \widehat{C}_{\bullet}(R)$$

Thus we have an isomorphism of pairs of graded vector spaces

$$(3.18) \quad (D_{\text{poly}}^L(R), C_{\text{poly}}^L(R)) = (D_{\text{poly}}(R), \widehat{C}_{\bullet}(R))$$

The right-hand side is a precalculus up to homotopy (as it is basically a pair of spaces of Hochschild chains/cochains).

**Lemma 3.7.** *The precalculus up to homotopy on the right-hand side of (3.18) is the same one as the one we have constructed on the left-hand side.*

*Proof.* Note that going from the pair  $(k, R)$  to  $(R, JL)$  is a base extension by  $R$  (since  $JL = R \widehat{\otimes} R$ ). Since the formulæ in [6] are clearly compatible with base extension we have that the precalculus structure on

$$(3.19) \quad (D_{\text{poly}, R}(JL), \widehat{C}_{R, \bullet}(JL)) = (R \widehat{\otimes} D_{\text{poly}}(R), R \widehat{\otimes} \widehat{C}_{\bullet}(R))$$

is obtained by base extension from the one on

$$(D_{\text{poly}}(R), \widehat{C}_{\bullet}(R))$$

Furthermore one checks that the Grothendieck connections on  $D_{\text{poly}, R}(JL)$  and  $\widehat{C}_{R, \bullet}(JL)$  under the isomorphism (3.19) act by the standard Grothendieck connection on the copy of  $R$  appearing on the left of  $\widehat{\otimes}$  and trivially on  $D_{\text{poly}}(R)$ ,  $\widehat{C}_{\bullet}(R)$ . Hence its invariants are precisely  $D_{\text{poly}}(R)$ ,  $\widehat{C}_{\bullet}(R)$ . This finishes the proof.  $\square$

**3.5. The Hochschild–Kostant–Rosenberg Theorem in the Lie algebroid framework.** We recall the Lie algebroid version of the famous cohomological Hochschild–Kostant–Rosenberg (shortly, HKR) quasi-isomorphism; for a proof, we refer to [2].

**Theorem 3.8.** *We consider a Lie algebroid  $L$  over  $R$  in the sense of Definition 3.1, which is assumed to be free of rank  $d$  over  $R$ .*

*Then, the map*

$$(3.20) \quad \text{HKR}(l_1 \wedge \cdots \wedge l_p) = (-1)^{\frac{p(p-1)}{2}} \frac{1}{p!} \sum_{\sigma \in \mathfrak{S}_p} (-1)^{\sigma} l_{\sigma(1)} \otimes_R \cdots \otimes_R l_{\sigma(p)}$$

defines a quasi-isomorphism of complexes from  $(T_{\text{poly}}^L(R), 0)$  to  $(D_{\text{poly}}^L(R), d_{\text{H}})$ .

There is a dual version of Theorem 3.8, which will also be needed.

**Theorem 3.9.** *The quasi-isomorphism (3.20) induces the quasi-isomorphism*

$$\text{HKR}(a) = a \circ \text{HKR}$$

of complexes from  $(C_{\text{poly}}^L(R), b_{\text{H}})$  to  $(\Omega^L(R), 0)$ .

#### 4. FEDOSOV RESOLUTIONS IN THE LIE ALGEBROID FRAMEWORK

**4.1. Introduction.** The aim of this section is to discuss Fedosov resolutions [10] in the Lie algebroid framework. These are needed to formulate and prove the globalization result, which in turn leads to the main results.

To help the reader understand our algebraic setup (which was inspired by [29]) we give some motivation for the definitions in the subsequent sections. For the sake of exposition we assume in this introduction that  $X$  is some kind of  $d$ -dimensional smooth space and  $\mathcal{L}$  is an appropriate version of the tangent bundle of  $X$ .

One of the applications of formal geometry is the globalization of local *coordinate dependent* constructions. For example using the Darboux Lemma it is trivial to quantize a symplectic manifold locally but such local quantizations are coordinate dependent and they do not globalize easily. The same is true for the local formality morphisms (see §5.4 below for more details) which we use in this paper.

The idea is then to replace  $X$  by a much larger infinite dimensional space  $X^{\text{coord}} \rightarrow X$  that parametrizes formal local coordinate systems on  $X$ . For example if  $X$  is an algebraic variety then the fiber at  $x \in X$  in  $X^{\text{coord}}$  is given by the  $k$ -algebra isomorphisms  $\hat{\mathcal{O}}_{X,x} \rightarrow k[[t_1, \dots, t_d]]$ . An equivalent way of saying this is that  $X^{\text{coord}}$  universally trivializes the jet bundle  $(\hat{\mathcal{O}}_{X,x})_{x \in X}$  over  $X$ .

Local constructions can be tautologically globalized to  $X^{\text{coord}}$  and this should be followed by some type of descent for  $X^{\text{coord}}/X$ . A general procedure to do this is to resolve  $\mathcal{O}_X$  by a De Rham-type complex over  $\mathcal{O}_{X^{\text{coord}}}$  but this does not really work as the fibers of  $X^{\text{coord}} \rightarrow X$  are not contractible.

However in the aforementioned examples the local constructions are all compatible with *linear* coordinate changes. So if we define  $X^{\text{aff}} = X^{\text{coord}}/\text{Gl}_d$  then the constructions descend to  $X^{\text{aff}}$  and as the fibers of  $X^{\text{aff}}/X$  are contractible we can descend further to  $X$ .

In this paper we work over a general locally free Lie algebroid  $\mathcal{L}$  rather than  $T_X$ . In this setting we define the analogue of  $X^{\text{coord}}$  as the space which universally trivializes the space of jet bundles for  $\mathcal{L}$  (see §3.1.4).

**4.2. Setup.** As a general principle we work on the presheaf level in this paper, performing sheafification only as the very last step of the constructions. This means that we may throughout replace all spaces by rings and locally free sheaves may be treated as free modules.

As before we consider a Lie algebroid  $L$  over a ring  $R$  in the sense of Definition 3.1, *i.e.*  $L$  is free of rank  $d$  over  $R$ .

First we discuss Fedosov resolutions of  $L$ -poly-vector fields and  $L$ -poly-differential operators as Gerstenhaber algebras up to homotopy, referring to [7] for details. Finally, we discuss Fedosov resolutions of  $\Omega^L(R)$  (see (3.3)) and  $C_{\text{poly}}^L(R)$  (see (3.13)) which are compatible with the precalculus structure up to homotopy.

**4.3. The (affine) coordinate space of a Lie algebroid.** For a Lie algebroid  $L$  over  $R$  as above, its **coordinate space**  $R^{\text{coord},L}$  has been introduced and discussed in details in [7, 27], to which we refer for a more extensive treatment.

As explained in §4.1, the main property of  $R^{\text{coord},L}$  is the existence of an isomorphism of  $R^{\text{coord},L}$ -algebras

$$(4.1) \quad t : R^{\text{coord},L} \widehat{\otimes}_{R_1} \mathcal{J}L \rightarrow R^{\text{coord},L} \llbracket x_1, \dots, x_d \rrbracket = R^{\text{coord},L} \widehat{\otimes} F, \quad F = k \llbracket x_1, \dots, x_d \rrbracket,$$

and  $R^{\text{coord},L}$  is universal with respect to this property, *i.e.* if there is an  $R$ -algebra  $W$ , such that there is a  $W$ -linear isomorphism  $W \widehat{\otimes}_{R_1} \mathcal{J}L \rightarrow W \llbracket x_1, \dots, x_d \rrbracket$ , then there exists a unique morphism  $R^{\text{coord},L} \rightarrow W$ .

We note in particular that in contrast to  $\mathcal{J}L$  the ring  $R^{\text{coord},L}$  is *not* an adic topological ring: it is equipped with the discrete topology (like  $R$ ).

**Example 4.1.** Assume  $R = k[x_1, \dots, x_d]$  and  $L = \text{Der}_k(R)$ . As explained in [27, §6.1.5][29] we have

$$R^{\text{coord},L} = R[y_{i,\alpha} : i = 1, \dots, d, \alpha \in \mathbb{N}^d \setminus \{0\}]_{\det(y_{i,e_j})},$$

where  $e_j$  is the  $j$ -th standard basis vector in  $\mathbb{Z}^d$ , and the subscript  $\det(y_{i,e_j})$  refers to the localization at the indicated element. As in this case  $X = \text{Spec } R$  has global coordinates  $x_1, \dots, x_n$  the coordinate ring of the jet bundle  $\mathcal{J}L$  is equal to  $R \llbracket y_1, \dots, y_d \rrbracket$  where  $y_i$  is a local version of the global coordinate  $x_i$ . The morphism  $t$  is the ‘‘universal Taylor expansion’’ morphism

$$t(y_i) = \sum_{\alpha} y_{i,\alpha} t^{\alpha}.$$

As a consequence of the universal property of  $R^{\text{coord},L}$ ,  $R^{\text{coord},L}$  admits an action of  $\text{GL}_d(k)$ , such that the following identity holds true on  $R^{\text{coord},L} \widehat{\otimes} F$  for  $A \in \text{GL}_d(k)$

$$(A^{-1} \widehat{\otimes} A)|_{\mathcal{J}L} = \text{Id}_{\mathcal{J}L}$$

where  $\mathcal{J}L$  is considered as a subalgebra of  $R^{\text{coord},L} \widehat{\otimes} F$  through (4.1).

By means of  $R^{\text{coord},L}$ , we consider the graded algebra  $C^{\text{coord},L} = \Omega_{R^{\text{coord},L}} \otimes_{\Omega_{R_1}} \Omega^{L_1}(R_1)$ . It has the structure of a DG-algebra with differential  $d_{C^{\text{coord},L}} = d_{\Omega_{R^{\text{coord},L}}} \otimes_{\Omega_{R_1}} 1 + 1 \otimes_{\Omega_{R_1}} d_{L_1}$ , and inherits from  $R^{\text{coord},L}$  a rational  $\text{GL}_d(k)$ -action.

The universal isomorphism (4.1) extends to an isomorphism

$$(4.2) \quad t : C^{\text{coord},L} \widehat{\otimes}_{R_1} \mathcal{J}L \rightarrow C^{\text{coord},L} \llbracket x_1, \dots, x_d \rrbracket,$$

where we used the respective obvious identifications

$$\begin{aligned} C^{\text{coord},L} \widehat{\otimes}_{R^{\text{coord},L}} (R^{\text{coord},L} \widehat{\otimes}_{R_1} \mathcal{J}L) &\cong \Omega_{R^{\text{coord},L}} \widehat{\otimes}_{\Omega_{R_1}} (\Omega^{L_1}(R_1) \otimes_{R_1} \mathcal{J}L) \cong C^{\text{coord},L} \widehat{\otimes}_{R_1} \mathcal{J}L, \\ C^{\text{coord},L} \widehat{\otimes}_{R^{\text{coord},L}} (R^{\text{coord},L} \widehat{\otimes} F) &\cong C^{\text{coord},L} \llbracket x_1, \dots, x_d \rrbracket. \end{aligned}$$

We endow the graded algebra on the left-hand, resp. right-hand, side of (4.2) with the following natural differential

$$(4.3) \quad {}^1\nabla^{\text{coord}} = d_{\Omega_{R^{\text{coord},L}}} \widehat{\otimes}_{\Omega_{R_1}} 1 + 1 \widehat{\otimes}_{\Omega_{R_1}} {}^1\nabla, \quad \text{resp.}$$

$$(4.4) \quad d = d_{C^{\text{coord},L}} \widehat{\otimes} 1,$$

where  ${}^1\nabla$  has been introduced in Subsection 3.4. Both (4.3) and (4.4) are, by construction, flat  $C^{\text{coord},L}$ -connections on the respective spaces, and the obvious inclusions from  $C^{\text{coord},L}$  into  $C^{\text{coord},L} \widehat{\otimes}_{R_1} \mathcal{J}L$  and  $C^{\text{coord},L} \llbracket x_1, \dots, x_d \rrbracket$  are morphisms of DG-algebras.



The main property of the connections (4.3) and (4.4) lies in the existence of a **canonical Maurer–Cartan element in  $C^{\text{coord},L}$** : namely, according to [27, Subsection 1.6] and [7, Subsection 5.2], there exists a unique element  $\omega$  of  $C^{\text{coord},L} \widehat{\otimes} \text{Der}(F)$  of degree 1, satisfying

$$t \circ {}^1\nabla^{\text{coord}} \circ t^{-1} - \text{d} = \omega,$$

where the expression on the left-hand side is naturally viewed as a  $C^{\text{coord},L}$ -linear derivation of  $F$ . Furthermore,  $\omega$  satisfies the Maurer–Cartan equation in the DG-Lie algebra  $C^{\text{coord},L} \widehat{\otimes} \text{Der}(F)$ , i.e.

$$\text{d}\omega + \frac{1}{2}[\omega, \omega] = 0,$$

(which implies that  $\text{d} + [\omega, \bullet]$  is a flat connection on  $C^{\text{coord},L}[[x_1, \dots, x_d]]$ ) and the verticality condition

$$(4.5) \quad \iota_v \omega = 1 \otimes v, \quad v \in \mathfrak{gl}_d(k)$$

(here,  $\iota_v$  on the left-hand side denotes the contraction operation on  $C^{\text{coord},L}$  with respect to  $v$ , coming from the infinitesimal action of  $\mathfrak{gl}_d(k)$  on  $R^{\text{coord},L}$ ;  $v$  on the right-hand side denotes the linear vector field associated to  $v$ , acting on  $F$ ).

Finally, we consider the **affine coordinate space  $R^{\text{aff},L}$**  of a Lie algebroid  $L$  over  $R$ : it is simply the  $\text{GL}_d(k)$ -invariant ring

$$R^{\text{aff},L} = (R^{\text{coord},L})^{\text{GL}_d(k)}$$

It is an  $R$ -algebra in an obvious way, and enjoys a universal property similar to the one satisfied by  $R^{\text{coord},L}$ , for which we refer to [7, Subsection 5.4].

**Example 4.2.** Continuing Example 4.1, assume  $R = k[x_1, \dots, x_d]$  and  $L = \text{Der}_k(R)$ . We now have

$$R^{\text{aff},L} = R[y_{i,\alpha} : i = 1, \dots, d, |\alpha| \geq 2],$$

where  $|\bullet|$  denotes the norm of a multiindex in  $\mathbb{N}^d$ . We observe that  $R^{\text{aff},L}$  is an (infinite) polynomial ring, while  $R^{\text{coord},L}$  is not, due to the localization.

Similarly, we have the DG-algebra  $C^{\text{aff},L} = \Omega_{R^{\text{aff},L}} \otimes_{\Omega_{R_1}} \Omega^{L_1}(R_1)$ , with differential  $\text{d}_{C^{\text{aff},L}} = \text{d}_{\Omega_{R^{\text{aff},L}}} \otimes_{\Omega_{R_1}} 1 + 1 \otimes_{\Omega_{R_1}} \text{d}_{L_1}$ . We may further consider the graded algebra

$$C^{\text{aff},L} \widehat{\otimes}_{R^{\text{aff},L}} (R^{\text{aff},L} \widehat{\otimes}_{R_1} \mathcal{J}L) \cong \Omega_{R^{\text{aff},L}} \widehat{\otimes}_{\Omega_{R_1}} (\Omega^{L_1}(R_1) \otimes_{R_1} \mathcal{J}L) \cong C^{\text{aff},L} \widehat{\otimes}_{R_1} \mathcal{J}L,$$

endowed with the natural differential

$${}^1\nabla^{\text{aff}} = \text{d}_{\Omega_{R^{\text{aff},L}}} \widehat{\otimes}_{\Omega_{R_1}} 1 + 1 \widehat{\otimes}_{\Omega_{R_1}} {}^1\nabla,$$

making the natural inclusion  $C^{\text{aff},L} \hookrightarrow C^{\text{aff},L} \widehat{\otimes}_{R_1} \mathcal{J}L$  into a morphism of DG-algebras. Obviously,  ${}^1\nabla^{\text{coord}}$  descends by its very construction to  $C^{\text{aff},L} \widehat{\otimes}_{R_1} \mathcal{J}L$  and identifies with  ${}^1\nabla^{\text{aff}}$ .

**Lemma 4.3.**  *$R^{\text{aff},L}$  is of the form  $S \otimes R$  where  $S$  is an (infinitely generated) polynomial ring.*

*Proof.* See [7, §5.3]. □

Note that this depends on our standing assumption that  $L$  is free and furthermore the decomposition  $R^{\text{aff},L} = S \otimes R$  is not canonical.

**4.4. Fedosov resolutions of  $L$ -poly-vector fields and  $L$ -poly-differential operators on  $R$ .** In this section, we recall briefly the main results of [7, §4.3], to which we refer for more details. We consider relative poly-differential operators and poly-vector fields (see §3.2) in the following situation:  $(A, d_A) = (C^{\text{aff}, L}, d_{C^{\text{aff}, L}})$  and  $(B, d_B) = (C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL, {}^1\nabla^{\text{aff}})$ .

**Theorem 4.4.** *For a Lie algebroid  $L$  over  $R$  as above, there exist quasi-isomorphisms of Gerstenhaber algebras up to homotopy*

(4.6)

$$(T_{\text{poly}}^L(R), 0, [ , ], \cup) = (T_{\text{poly}}^{L_2}(R_2), 0, [ , ], \cup) \hookrightarrow (T_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}, [ , ], \cup),$$

(4.7)

$$(D_{\text{poly}}^L(R), d_{\text{H}}, [ , ], \cup) = (D_{\text{poly}}^{L_2}(R_2), d_{\text{H}}, [ , ], \cup) \hookrightarrow (D_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + d_{\text{H}}, [ , ], \cup).$$

*Proof.* We refer to [7] for details. For example the map (4.7) is derived by suitable base extension from (3.12). For the fact that the maps are quasi-isomorphisms we refer to [7, Prop. 7.3.1].  $\square$

**4.5. The Fedosov resolution of  $L$ -forms on  $R$ .** We consider the precalculus  $(\Omega^L(R), 0, L, \cap)$  of  $L$ -forms over the Gerstenhaber algebra  $(T_{\text{poly}}^L(R), 0, [ , ], \cup)$ , described in §3.1: we describe now a well-suited resolution of  $(\Omega^L(R), 0, L, \cap)$  which is compatible with the Fedosov resolution  $(T_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}, [ , ], \cup)$  from Theorem 4.4.

**Theorem 4.5.** *For a Lie algebroid  $L$  over  $R$  as above, there exists a quasi-isomorphism of precalculi as in the following commutative diagram <sup>6</sup>:*

$$\begin{array}{ccc} (T_{\text{poly}}^L(R), 0, [ , ], \cup) = (T_{\text{poly}}^{L_2}(R_2), 0, [ , ], \cup) & \hookrightarrow & (T_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}, [ , ], \cup) , \\ \downarrow \wr & & \downarrow \wr \\ (\Omega^L(R), 0, L, \cap) = (\Omega^{L_2}(R_2), 0, L, \cap) & \hookrightarrow & (\Omega_{C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL / C^{\text{aff}, L}}, {}^1\nabla^{\text{aff}}, L, \cap) \end{array}$$

*the vertical arrows denoting the contraction and Lie derivative.*

*Proof.* We refer to [7, §4.3.3]: we observe that the construction of the quasi-isomorphism uses a dualization of the construction of the quasi-isomorphism (4.6), and that contraction operations and differentials are preserved by the above quasi-isomorphism, whence all algebraic structures are preserved.  $\square$

**4.6. The Fedosov resolution of  $L$ -chains on  $R$ .** We consider the DG-algebra  $(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL, {}^1\nabla^{\text{aff}})$ , and to it we associate the  $C^{\text{aff}, L}$ -relative Hochschild chain complex, i.e.

$$\begin{aligned} \widehat{C}_{C^{\text{aff}, L}, \bullet}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL) &= \bigoplus_{p \leq 0} (C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL)^{\widehat{\otimes}_{C^{\text{aff}, L}}(-p+1)} \\ &\cong \bigoplus_{p \leq 0} (C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL)^{\widehat{\otimes}_{R_1}(-p+1)} = \\ &= C^{\text{aff}, L} \widehat{\otimes}_{R_1} \widehat{C}_{R_1, \bullet}(JL), \end{aligned}$$

<sup>6</sup> $\Omega_A$ , for a topological  $k$ -algebra  $A$ , denotes the continuous De Rham complex. A similar convention holds for an extension of topological algebras  $B/A$ .

Further, we have the identification

$$C^{\text{aff},L} \widehat{\otimes}_{R_1} \widehat{C}_{R_1,\bullet}(JL) \cong \Omega_{R^{\text{aff},L}} \widehat{\otimes}_{\Omega_{R_1}} \left( \Omega^{L_1}(R_1) \widehat{\otimes}_{R_1} \widehat{C}_{R_1,\bullet}(JL) \right).$$

and one checks that the differentials coming from the Grothendieck connection on each side are the same. I.e.

$${}^1\nabla^{\text{aff}} = d_{\Omega_{R^{\text{aff},L}}} \widehat{\otimes} 1 + 1 \widehat{\otimes} {}^1\nabla,$$

**Proposition 4.6.** *For a Lie algebroid  $L$  over  $R$  as above, the cohomology of  $(\widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}})$  is concentrated in degree 0, where*

$$H^0(\widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}) \cong C_{\text{poly}}^L(R).$$

*Proof.* Taking the inverse of (3.15) we obtain a morphism

$$C_{\text{poly}}^L(R) \cong \widehat{C}_{R_1,\bullet}(JL)^{1\nabla} \hookrightarrow \widehat{C}_{R_1,\bullet}(JL)$$

which extends to a morphism

$$(4.8) \quad (C_{\text{poly}}^L(R), 0) \rightarrow (\Omega_{R^{\text{aff},L}} \widehat{\otimes}_{\Omega_{R_1}} (\Omega^{L_1}(R_1) \widehat{\otimes}_{R_1} \widehat{C}_{R_1,\bullet}(JL)), d_{\Omega_{R^{\text{aff},L}}} \otimes 1 + 1 \otimes {}^1\nabla)$$

We will show that it is a quasi-isomorphism. To this end we make use of the identification  $R^{\text{aff},L} = S \otimes R$  given in Lemma 4.3. The right-hand side of the extended morphism becomes

$$(\Omega_S \widehat{\otimes} (\Omega^{L_1}(R_1) \widehat{\otimes}_{R_1} \widehat{C}_{R_1,\bullet}(JL)), d_S \otimes 1 + 1 \otimes {}^1\nabla)$$

Using a filtration argument together with a suitable version of Poincaré's Lemma for  $S$ , the previous complex is quasi-isomorphic to

$$(\Omega^{L_1}(R_1) \otimes_{R_1} \widehat{C}_{R_1,\bullet}(JL), {}^1\nabla)$$

It remains to show that for each  $p \leq 0$

$$(\Omega^{L_1}(R_1) \otimes_{R_1} \widehat{C}_{R_1,p}(JL), {}^1\nabla)$$

has cohomology in degree 0. Filtering this complex with respect to the  $J$ -adic filtration and taking the associated graded complex one verifies that one obtains

$$(\Omega^L(R) \otimes_{R_1} S(L^*)^{\otimes -p-1}, d)$$

where the differential  $d$  is obtained from the action of  $L$  on  $S(L^*)^{\otimes p+1}$  by contraction. Using again a suitable version of Poincaré's Lemma one finds that the resulting complex is indeed exact in degrees  $< 0$ .  $\square$

**Theorem 4.7.** *For a Lie algebroid  $L$  over  $R$  as above, there is a quasi-isomorphism of precalculi up to homotopy as in the following commutative diagram:*

$$\begin{array}{ccc} (D_{\text{poly}}^L(R), d_H, [\ , \ ], \cup) = (D_{\text{poly}}^{L_2}(R_2), d_H, [\ , \ ], \cup) & \xrightarrow{\quad} & (D_{\text{poly},C^{\text{aff},L}}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + d_H, [\ , \ ], \cup) , \\ \Downarrow & & \Downarrow \\ (C_{\text{poly}}^L(R), b_H, L, \cap) & \xrightarrow{\quad} & (\widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + b_H, L, \cap) \end{array}$$

*the vertical arrows denoting the contraction and Lie derivative.*

## 5. GLOBALIZATION OF TSYGAN'S FORMALITY IN THE LIE ALGEBROID FRAMEWORK

The present section is devoted to the proof of Theorem 1.3. We first briefly review some basic facts on  $L_\infty$ -algebras,  $L_\infty$ -modules and related morphisms. This is discussed in [7, §6] for  $L_\infty$ -morphisms. Here we add a discussion on the descent procedure for  $L_\infty$ -modules over  $L_\infty$ -algebras and related morphisms.

Then, we add a short *excursus* on Kontsevich's and Shoikhet's formality theorems: we focus on the main properties of both formality morphisms, without delving into the technical details of their respective constructions.

Finally, we give the main lines, along which the globalization of Tsygan's formality can be proved: the proof is a combination of the properties of Kontsevich's and Shoikhet's  $L_\infty$ -morphisms with the Fedosov resolutions from §4.

**5.1. Descent for  $L_\infty$ -algebras and  $L_\infty$ -modules.** We discuss a series of descent scenarios for  $L_\infty$ -algebras,  $L_\infty$ -modules and related morphisms, which are modelled after the formalism for descent of differential forms in differential geometry. The verification of the results in this section are along the same lines as [27, §7.6, §7.7]. To clearly separate all the various cases we have numbered them.

(1) To start it is convenient to work over an arbitrary DG operad  $\mathcal{O}$  with underlying graded operad  $\tilde{\mathcal{O}}$  (thus we forget the differential on  $\mathcal{O}$ ). Assume that  $\mathfrak{g}$  is an algebra over  $\mathcal{O}$  and consider a set of  $\tilde{\mathcal{O}}$ -derivations  $(\iota_v)_{v \in \mathfrak{s}}$  of degree  $-1$  on  $\mathfrak{g}$  ( $\mathfrak{s}$  is an index set, without any additional structure). Put  $L_v = d_{\mathfrak{g}}\iota_v + \iota_v d_{\mathfrak{g}}$ . This is a derivation of  $\mathfrak{g}$  of degree zero which commutes with  $d_{\mathfrak{g}}$ . Put

$$(5.1) \quad \mathfrak{g}^{\mathfrak{s}} = \{w \in \mathfrak{g} \mid \forall v \in \mathfrak{s} : \iota_v w = L_v w = 0\}$$

It is easy to see that  $\mathfrak{g}^{\mathfrak{s}}$  is an algebra over  $\mathcal{O}$  as well. Informally we will call such a set of derivations  $(\iota_v)_{v \in \mathfrak{s}}$  an  $\mathfrak{s}$ -action.

(2) Assume that  $M$  is a  $\mathfrak{g}$ -module and assume that  $\mathfrak{s}$  also acts on  $M$ , in a way compatible with the action of  $\mathfrak{s}$  on  $\mathfrak{g}$ , i.e. a general element  $v$  of  $\mathfrak{s}$  determines an operator  $\iota_v$  on  $M$ , such that Leibniz's rule holds true for the operations  $\tilde{\mathcal{O}}(n) \otimes (\mathfrak{g}^{\otimes n-1} \otimes M) \rightarrow M$ . Again, we set  $L_v = d_M \iota_v + \iota_v d_M$ , which is a derivation of degree 0 on  $M$  compatible with the derivations  $L_v$  on  $\mathfrak{g}$ ,  $d_M$  being the differential on  $M$ .

(3) The above constructions apply in particular if  $\mathfrak{g}$  is an  $L_\infty$ -algebra. Assume that it has Taylor coefficients  $Q_n$ ,  $n \geq 1$ . Then  $L_v$  is defined by means of  $d_{\mathfrak{g}} = Q_1$ , and the derivation property of  $\iota_v$  reads as

$$(5.2) \quad \iota_v(Q_n(x_1, \dots, x_n)) = \sum_{i=1}^n (-1)^{\sum_{j=1}^{i-1} |x_j| + i} Q_n(x_1, \dots, \iota_v x_i, \dots, x_n), \quad x_j \in \mathfrak{g}, \quad j = 1, \dots, n.$$

Under these conditions the  $L_\infty$ -structure descends to  $\mathfrak{g}^{\mathfrak{s}}$ .

(4) Similarly if  $M$  is an  $L_\infty$ -module over  $\mathfrak{g}$  defined by Taylor coefficients  $R_n$  then the compatibility condition is

$$(5.3) \quad \begin{aligned} \iota_v(R_n(x_1, \dots, x_n; m)) &= \sum_{i=1}^n (-1)^{\sum_{j=1}^{i-1} |x_j| + i} R_n(x_1, \dots, \iota_v x_i, \dots, x_n; m) + \\ &+ (-1)^{\sum_{i=1}^n |x_i| + n - 1} R_n(x_1, \dots, x_n; \iota_v m), \quad m \in M, \quad x_j \in \mathfrak{g}, \quad j = 1, \dots, n. \end{aligned}$$

If this holds true then  $M^{\mathfrak{s}}$  becomes an  $L_\infty$ -module over  $\mathfrak{g}^{\mathfrak{s}}$ .

(5) We also need descent for  $L_\infty$ -morphisms. This does not immediately fall under the operadic framework given in (1), (2) but it is easy enough to give explicit formulæ like (5.2), (5.3). Thus assume  $\psi : \mathfrak{g} \rightarrow \mathfrak{h}$  is an  $L_\infty$ -morphism between  $L_\infty$ -algebras with  $\mathfrak{s}$ -action. Under the following compatibility condition

$$(5.4) \quad \iota_v(\psi_n(x_1, \dots, x_n)) = \sum_{i=1}^n (-1)^{\sum_{j=1}^{i-1} |x_j| + (i-1)} \psi_n(x_1, \dots, \iota_v x_i, \dots, x_n)$$

$x_j \in \mathfrak{g}$ ,  $j = 1, \dots, n$ ,  $n \geq 1$ ,  $\psi$  descends to an  $L_\infty$ -morphism  $\psi^{\mathfrak{s}} : \mathfrak{g}^{\mathfrak{s}} \rightarrow \mathfrak{h}^{\mathfrak{s}}$ .

(6) Let  $\psi : \mathfrak{g} \rightarrow \mathfrak{h}$  be a morphism between  $L_\infty$ -algebras with  $\mathfrak{s}$ -action such that the descent condition (5.4) holds, and let  $N$  be an  $L_\infty$ -module over  $\mathfrak{h}$  equipped with a compatible  $\mathfrak{s}$ -action. Let  $N_\psi$  be the pullback of  $N$  along  $\psi$ . Then the  $\mathfrak{s}$ -action on  $N_\psi$  is compatible with the  $\mathfrak{s}$ -action on  $\mathfrak{g}$ .

(7) Now assume that  $\mathfrak{g}$  is an  $L_\infty$ -algebra and  $M, N$  are  $L_\infty$ -modules over  $\mathfrak{g}$ . Assume that all objects are equipped with an  $\mathfrak{s}$ -action and that the descent conditions are satisfied.

Assume that  $\varphi : M \rightarrow N$  is an  $L_\infty$ -module morphism. Then the condition for  $\varphi$  to descend to an  $L_\infty$ -morphism  $M^{\mathfrak{s}} \rightarrow N^{\mathfrak{s}}$  is

$$(5.5) \quad \begin{aligned} \iota_v(\varphi_n(x_1, \dots, x_n; m)) &= \sum_{i=1}^n (-1)^{\sum_{j=1}^{i-1} |x_j| + (i-1)} \varphi_n(x_1, \dots, \iota_v x_i, \dots, x_n; m) + \\ &+ (-1)^{\sum_{i=1}^n |x_i| + n} \varphi_n(x_1, \dots, x_n; \iota_v m), \end{aligned}$$

for  $m \in M$ ,  $x_j \in \mathfrak{g}$ ,  $j = 1, \dots, n$ ,  $n \geq 1$ .

**5.2. Twisting of  $L_\infty$ -algebras and  $L_\infty$ -modules.** We refer to [10, §2], for a very detailed exposition of  $L_\infty$ -algebras,  $L_\infty$ -modules and the associated twisting procedures. See also [31].

**Convention.** We will work with infinite sums. We assume throughout that the occurring sums are convergent and that standard series manipulations are allowed. This will be the case in our applications.

If  $(\mathfrak{g}, Q)$  is an  $L_\infty$ -algebra then the Maurer-Cartan equation is defined as

$$(5.6) \quad \sum_{j=1}^{\infty} \frac{1}{j!} Q_n(\underbrace{\omega, \dots, \omega}_j) = 0,$$

and a solution  $\omega \in \mathfrak{g}_1$  is called a Maurer-Cartan element (MC element for short). Below we will only use DG-Lie algebras and in this case (5.6) reduces to the finite sum

$$d\omega + \frac{1}{2}[\omega, \omega] = 0.$$

A MC element defines a new “twisted” DG-Lie structure on  $\mathfrak{g}$  (denoted by  $\mathfrak{g}_\omega$ ) with Taylor coefficients

$$Q_{\omega,n}(x_1, \dots, x_n) = \sum_j \frac{1}{j!} Q_{n+j}(\underbrace{\omega, \dots, \omega}_j, x_1, \dots, x_n), \quad n \geq 1$$

If  $\mathfrak{g}$  is actually a DG-Lie algebra then twisting keeps the bracket but changes the differential to

$$d_\omega = d_{\mathfrak{g}} + [\omega, -].$$

If  $\mathfrak{h}$  is another  $L_\infty$  algebra,  $\psi$  is an  $L_\infty$ -morphism from  $\mathfrak{g}$  to  $\mathfrak{h}$  and  $\omega$  is a MC element in  $\mathfrak{g}$  then

$$(5.7) \quad \psi(\omega) = \sum_{n \geq 1} \frac{1}{n!} \psi_n(\underbrace{\omega, \dots, \omega}_n)$$

is a MC element in  $\mathfrak{h}$ .

We may also twist  $\psi$  with respect to  $\omega$ , so as to get an  $L_\infty$ -morphism  $\psi_\omega$  from  $\mathfrak{g}_\omega$  to  $\mathfrak{g}_{\psi(\omega)}$ , where

$$\psi_{\omega,n}(x_1, \dots, x_n) = \sum_{j \geq 0} \frac{1}{j!} \psi_{n+j}(\underbrace{\omega, \dots, \omega}_j, x_1, \dots, x_n), \quad n \geq 1.$$

If  $M$  is an  $L_\infty$ -module over a DG-Lie algebra with Taylor coefficients  $R_n$  and  $\omega \in \mathfrak{g}_1$  is a MC element then we may define a twisted  $L_\infty$  structure on  $M_\omega$  over  $\mathfrak{g}_\omega$  by the formula

$$R_{\omega,n}(x_1, \dots, x_n; m) = \sum_{j \geq 0} \frac{1}{j!} R_{n+j}(\underbrace{\omega, \dots, \omega}_j, x_1, \dots, x_n; m), \quad n \geq 0$$

If  $\mathfrak{g}$  is a DG-Lie algebra and  $M$  is a DG-Lie module over  $\mathfrak{g}$  then twisting keeps the  $\mathfrak{g}$ -action on  $M$  but changes the differential on  $M$  to

$$d_\omega = d + \omega \bullet.$$

Twisting of modules is compatible with pullback. More precisely if  $\psi : \mathfrak{g} \rightarrow \mathfrak{h}$  is an  $L_\infty$ -morphism,  $N$  is an  $L_\infty$ -module over  $\mathfrak{h}$  and  $\omega \in \mathfrak{g}_1$  is a MC element then we have

$$(5.8) \quad (N_{\psi(\omega)})_{\psi_\omega} = (N_\psi)_\omega$$

If  $\varphi : M \rightarrow N$  is an  $L_\infty$ -morphism of DG-Lie modules over the DG-Lie algebra  $\mathfrak{g}$  and  $\omega$  is a MC element in  $\mathfrak{g}_1$  then we obtain a twisted  $L_\infty$ -morphism  $\varphi_\omega : M_\omega \rightarrow N_\omega$  which is defined by

$$(5.9) \quad \varphi_{\omega,n}(x_1, \dots, x_n; m) = \sum_{j \geq 0} \frac{1}{j!} \varphi_{n+j}(\underbrace{\omega, \dots, \omega}_j, x_1, \dots, x_n; m), \quad n \geq 1.$$

**5.3. Compatibility of twisting and descent.** Assume now that  $\mathfrak{g}$  is a DG-Lie algebra equipped with an  $\mathfrak{s}$ -action and that  $\omega \in \mathfrak{g}_1$  is a MC element. Then  $\mathfrak{s}$  still acts on  $\mathfrak{g}_\omega$ , where we forget here about the differential: in fact, the concept of an  $\mathfrak{s}$ -action only refers to the underlying Lie algebra structure on  $\mathfrak{g}$ . However  $\mathfrak{g}^\mathfrak{s}$  and  $\mathfrak{g}_\omega^\mathfrak{s}$  will be different (as the Lie derivative  $L_v$  for  $v \in \mathfrak{s}$  will be different).

If  $(M, R)$  is an  $L_\infty$ -module over  $\mathfrak{g}$  which is also equipped with a compatible  $\mathfrak{s}$ -action then the  $\mathfrak{s}$ -actions on  $\mathfrak{g}_\omega$  and  $M_\omega$  are compatible provided the following condition holds

$$(5.10) \quad R_n(\iota_v \omega, x_2, \dots, x_n; m) = 0, x_i \in \mathfrak{g}, i = 2, \dots, n, n \geq 2, m \in M.$$

This condition is automatic if  $M$  is a DG-Lie module.

If  $\psi : \mathfrak{g} \rightarrow \mathfrak{h}$  is an  $L_\infty$ -morphism of DG-Lie algebras equipped with an  $\mathfrak{s}$ -action and the descent condition (5.4) is satisfied for  $\psi$  then an easy computation (see e.g. [27, §7.7]) shows that the same descent condition will be satisfied for  $\psi_\omega$  if the following condition holds

$$(5.11) \quad \psi_n(\iota_v \omega, x_2, \dots, x_n) = 0, x_i \in \mathfrak{g}, i = 2, \dots, n, s \in \mathfrak{s}, n \geq 2,$$

Furthermore if in this setting  $N$  is an  $L_\infty$ -module over  $\mathfrak{h}$  with compatible  $\mathfrak{s}$ -action such that the compatibility condition (5.10) holds then the corresponding condition will hold for  $N_\psi$ .

Similarly if we have an  $L_\infty$ -morphism  $\varphi : M \rightarrow N$  between DG-Lie modules over a DG-Lie algebra  $\mathfrak{g}$  such that  $\mathfrak{g}, M, N$  are equipped with compatible  $\mathfrak{s}$ -actions in such a way that the descent condition (5.5) holds for  $\varphi$  then the same descent condition will be satisfied for  $\varphi_\omega$  if the following condition holds

$$\varphi_n(\iota_v \omega, x_2, \dots, x_n; m) = 0, m \in M, x_i \in \mathfrak{g}, i = 2, \dots, n, n \geq 2, s \in \mathfrak{s}.$$

**5.4. Kontsevich's and Shoikhet's formality theorems.** In this brief section, we quote (without proofs) Kontsevich's and Shoikhet's formality theorems, along with the relevant properties, which we will need later in the proof of globalization results.

We consider the algebra  $F = k[[x_1, \dots, x_d]]$  of formal power series in  $d$  variables over a field  $k$  containing  $\mathbb{R}$ .

To  $F$ , we associate the DG-Lie algebras  $(T_{\text{poly}}(F), 0, [\ , \ ])$ , resp.  $(D_{\text{poly}}(F), d_H, [\ , \ ])$ , of formal poly-vector fields, resp. formal poly-differential operators, on  $F$ ; further, we consider the DG-Lie modules  $(\Omega_F, 0, L)$ , resp.  $(\widehat{C}_\bullet(F), b_H, L)$ , over  $(T_{\text{poly}}(F), 0, [\ , \ ])$ , resp.  $(D_{\text{poly}}(F), d_H, [\ , \ ])$ , where  $\Omega_F$  denotes the continuous De Rham complex of  $F$  with De Rham differential  $d$ , and  $\widehat{C}_\bullet(F)$  is the continuous Hochschild chain complex of  $F$ .

The following is Kontsevich's celebrated "formality" result.

**Theorem 5.1** ([17]). *There is an  $L_\infty$ -quasi-isomorphism*

$$\mathcal{U} : (T_{\text{poly}}(F), 0, [\ , \ ]) \rightarrow (D_{\text{poly}}(F), d_H, [\ , \ ]),$$

enjoying the following properties:

- i) *the first Taylor coefficient of  $\mathcal{U}$  coincides with the Hochschild–Kostant–Rosenberg quasi-isomorphism*

$$\text{HKR}(\partial_{i_1} \wedge \dots \wedge \partial_{i_p}) = (-1)^{\frac{p(p-1)}{2}} \frac{1}{p!} \sum_{\sigma \in \mathfrak{S}_p} (-1)^\sigma \partial_{i_{\sigma(1)}} \otimes \dots \otimes \partial_{i_{\sigma(p)}}$$

- from the DG-vector space  $(T_{\text{poly}}(F), 0)$  to the DG-vector space  $(D_{\text{poly}}(F), d_H)$ .  
ii) *If  $n \geq 2$ , and  $\gamma_i, i = 1, \dots, n$ , are elements of  $T_{\text{poly}}^0(F)$ , then*

$$\mathcal{U}_n(\gamma_1, \dots, \gamma_n) = 0.$$

iii) If  $n \geq 2$ ,  $\gamma_1$  is a linear vector field on  $F$  (i.e. an element of  $\mathfrak{gl}_d$ ),  $\gamma_i$ ,  $i = 2, \dots, n$  are general elements of  $T_{\text{poly}}(F)$ , then

$$\mathcal{U}_n(\gamma_1, \gamma_2, \dots, \gamma_n) = 0.$$

By composing the action  $L$  of  $D_{\text{poly}}(F)$  on  $\widehat{C}_\bullet(F)$  with the  $L_\infty$ -quasi-isomorphism  $\mathcal{U}$  from Theorem 5.1,  $\widehat{C}_\bullet(F)$  inherits an  $L_\infty$ -module structure over the DG-Lie algebra  $(T_{\text{poly}}(F), 0, [ , ])$ .

The first part of the following theorem was a conjecture by Tsygan [26] which has been proved by Shoikhet in [21]. The second part has been proved in [10].

**Theorem 5.2.** *There is an  $L_\infty$ -quasi-isomorphism*

$$\mathcal{S} : (\widehat{C}_\bullet(F), \mathfrak{b}_H, L \circ \mathcal{U}) \rightarrow (\Omega_F, 0, L)$$

of  $L_\infty$ -modules over the DG-Lie algebra  $(T_{\text{poly}}(F), 0, [ , ])$ , enjoying the following properties:

i) the 0-th Taylor coefficient of  $\mathcal{S}$  coincides with the Hochschild–Kostant–Rosenberg quasi-isomorphism

$$\text{HKR}((a_0 | \dots | a_p)) = \frac{1}{p!} a_0 da_1 \cdots da_p$$

from the DG-vector space  $(\widehat{C}_\bullet(F), \mathfrak{b}_H)$  to the DG-vector space  $(\Omega_F, 0)$ .

ii) If  $n \geq 1$ ,  $\gamma_1$  is a linear vector field on  $F$ ,  $\gamma_i$ ,  $i = 2, \dots, n$  are general elements of  $T_{\text{poly}}(F)$ , and  $c$  is a general element of  $\widehat{C}_\bullet(F)$ , then

$$\mathcal{S}_n(\gamma_1, \dots, \gamma_n; c) = 0.$$

**5.5. Formality theorem in the ring case.** This section is devoted to the proof of a Tsygan-like formality theorem in the case of a Lie algebroid  $L$  over a  $k$ -algebra  $R$ , such that  $L$  is free over  $R$  of rank  $d$ : the proof combines Shoikhet’s formality theorem 5.2 with the Fedosov resolutions from §4.

**Theorem 5.3.** *Assume  $\mathbb{R} \subset k$ . For a Lie algebroid  $L$  over  $R$  as above, there exist DG-Lie algebras  $(\mathfrak{g}_i^L, d_i, [ , ]_i)$ , DG-Lie modules  $(\mathfrak{m}_i^L, \mathfrak{b}_i, L_i)$  over  $\mathfrak{g}_i$ ,  $i = 1, 2$ , and  $L_\infty$ -quasi-isomorphisms  $\mathfrak{U}_L$  from  $\mathfrak{g}_1^L$  to  $\mathfrak{g}_2^L$  and  $\mathfrak{S}_L$  from  $\mathfrak{m}_2^L$  to  $\mathfrak{m}_1^L$ , which fit into the following commutative diagram:*

$$(5.12) \quad \begin{array}{ccccccc} T_{\text{poly}}^L(R) & \hookrightarrow & \mathfrak{g}_1^L & \xrightarrow{\mathfrak{U}_L} & \mathfrak{g}_2^L & \longleftarrow & D_{\text{poly}}^L(R) \\ \downarrow \mathfrak{L} & & \downarrow \mathfrak{L}_1 & & \downarrow \mathfrak{L}_2 & & \downarrow \mathfrak{L} \\ \Omega^L(R) & \hookrightarrow & \mathfrak{m}_1^L & \xleftarrow{\mathfrak{S}_L} & \mathfrak{m}_2^L & \longleftarrow & C_{\text{poly}}^L(R) \end{array}$$

such that the induced maps

$$T_{\text{poly}}^L(R) \rightarrow \mathbf{H}^\bullet(D_{\text{poly}}^L(R), d_H), \quad \mathbf{H}^\bullet(C_{\text{poly}}^L(R), \mathfrak{b}_H) \rightarrow \Omega^L(R)$$

on (co)homology coincide with the respective HKR-quasi-isomorphisms. The morphisms indicated by hooked arrows are actual quasi-isomorphisms of DG-Lie algebras and DG-Lie modules respectively.



*Proof.* The first step in the proof of Theorem 5.3 may be borrowed from [7, §7.3]. Namely, we consider the following graded vector spaces:

$$\begin{aligned} C^{\text{coord},L} \widehat{\otimes} T_{\text{poly}}(F) &\cong T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), \\ C^{\text{coord},L} \widehat{\otimes} D_{\text{poly}}(F) &\cong D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), \\ C^{\text{coord},L} \widehat{\otimes} \Omega_F &\cong \Omega_{C^{\text{coord},L} \widehat{\otimes} F / C^{\text{coord},L}}, \\ C^{\text{coord},L} \widehat{\otimes} \widehat{C}_{\bullet}(F) &\cong \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L} \widehat{\otimes} F), \end{aligned}$$

where the DG-algebra  $C^{\text{coord},L}$  has been introduced in §4.3, and where  $\widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L} \widehat{\otimes} F)$  denotes the  $C^{\text{coord},L}$ -relative Hochschild chain complex of the DG-algebra  $C^{\text{coord},L} \widehat{\otimes} F$ .

The Maurer–Cartan form on  $C^{\text{coord},L} \widehat{\otimes} F$  introduced in §4.3 defines a twisted differential  $d_{\omega} = d + \omega$  on the listed graded vector spaces and as explained in §5.2  $d_{\omega}$  is compatible with the respective DG-Lie algebra and DG-Lie module structures.

Thus, formal geometry provides us with the following DG-Lie algebras and respective DG-Lie modules:

$$\begin{array}{ccc} (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d_{\omega}, [\ , \ ]), & & (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d_{\omega} + d_{\text{H}}, [\ , \ ]), \\ \Downarrow & & \Downarrow \\ (\Omega_{C^{\text{coord},L} \widehat{\otimes} F / C^{\text{coord},L}}, d_{\omega}, L) & & (\widehat{C}_{C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d_{\omega} + b_{\text{H}}, L) \end{array}$$

We repeat that, viewing all DG-Lie algebra and DG-Lie module structures above as  $L_{\infty}$ -structures, the differential  $d_{\omega}$  is the twist of the standard structures with respect to the MC element  $\omega$  of  $C^{\text{coord},L} \widehat{\otimes} \text{Der}(F) = T_{\text{poly},C^{\text{coord},L}}^0(C^{\text{coord},L} \widehat{\otimes} F)$ .

The  $L_{\infty}$ -quasi-isomorphism  $\mathcal{U}$  of Theorem 5.1 extends  $C^{\text{coord},L}$ -linearly to an  $L_{\infty}$ -quasi-isomorphism

$$\mathcal{U}_L : (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d, [\ , \ ]) \rightarrow (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d + d_{\text{H}}, [\ , \ ]).$$

The composition of the DG-Lie action  $L$  of  $D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F)$  on  $\widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L} \widehat{\otimes} F)$  with the  $L_{\infty}$ -quasi-isomorphism  $\mathcal{U}_L$  endows the latter graded vector space with a structure of  $L_{\infty}$ -module over the DG-Lie algebra  $T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F)$ , which is obtained by  $C^{\text{coord},L}$ -base extension of the corresponding  $L_{\infty}$ -module structure of  $\widehat{C}_{\bullet}(F)$  over  $T_{\text{poly}}(F)$ .

Accordingly, the  $L_{\infty}$ -quasi-isomorphism  $\mathcal{S}$  of Theorem 5.2 extends to an  $L_{\infty}$ -quasi-isomorphism of  $L_{\infty}$ -modules

$$\mathcal{S}_L : (\widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L} \widehat{\otimes} F), d + b_{\text{H}}, L \circ \mathcal{U}_L) \rightarrow (\Omega_{C^{\text{coord},L} \widehat{\otimes} F / C^{\text{coord},L}}, d, L)$$

both viewed as  $L_{\infty}$ -modules over  $T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F)$ .

As outlined in §5.2 we may apply the twisting procedures for  $L_{\infty}$ -algebras,  $L_{\infty}$ -modules and  $L_{\infty}$ -morphisms to the present case, where the MC element is the Maurer–Cartan form  $\omega$ : thus, we get an  $L_{\infty}$ -morphism  $\mathcal{U}_{L,\omega}$

$$\mathcal{U}_{L,\omega} : (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d_{\omega}, [\ , \ ]) \rightarrow (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F), d_{\omega} + d_{\text{H}}, [\ , \ ]),$$

where here and below we used Property *ii*) of Theorem 5.1, which yields that the MC element  $\mathcal{U}(\omega)$  equals  $\omega$ .

The  $L_{\infty}$ -morphism  $\mathcal{U}_{L,\omega}$  yields an  $L_{\infty}$ -module structure on  $\widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L} \widehat{\otimes} F)$  over the  $\omega$ -twisted DG-Lie algebra  $T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L} \widehat{\otimes} F)$ .

Translating (5.8) to the present case we have

$$\begin{aligned} \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}F), d_\omega + b_H, L \circ \mathcal{U}_{L,\omega} \right) \\ = \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}F), d + b_H, L \circ \mathcal{U}_L \right)_\omega \end{aligned}$$

from which we get an  $L_\infty$ -quasi-isomorphism

$$\mathcal{S}_{L,\omega} : \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}F), d_\omega + b_H, L \circ \mathcal{U}_{L,\omega} \right) \rightarrow \left( \Omega_{C^{\text{coord},L}\widehat{\otimes}F/C^{\text{coord},L}}, d_\omega, L \right)$$

of  $L_\infty$ -modules.

Using the isomorphism (4.2) we obtain isomorphisms of DG-Lie algebras and respective DG-Lie modules

$$\begin{aligned} (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}F), d_\omega, [ , ]) &\cong (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}}, [ , ]), \\ (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}F), d_\omega + d_H, [ , ]) &\cong (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + d_H, [ , ]), \\ \left( \Omega_{C^{\text{coord},L}\widehat{\otimes}F/C^{\text{coord},L}}, d_\omega, L \right) &\cong \left( \Omega_{C^{\text{coord},L}\widehat{\otimes}_{R_1}JL/C^{\text{coord},L}}, {}^1\nabla^{\text{coord}}, L \right), \\ \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}F), d_\omega + b_H, L \right) &\cong \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + b_H, L \right), \end{aligned}$$

an  $L_\infty$ -morphism

$$\mathcal{U}_L^{\text{coord}} : (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}}, [ , ]) \rightarrow (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + d_H, [ , ]),$$

which yields an  $L_\infty$ -module structure on  $\widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL)$  over  $T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL)$ , and finally an  $L_\infty$ -morphism

$$\mathcal{S}_L^{\text{coord}} : \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + b_H, L \circ \mathcal{U}_L^{\text{coord}} \right) \rightarrow \left( \Omega_{C^{\text{coord},L}\widehat{\otimes}F/C^{\text{coord},L}}, {}^1\nabla^{\text{coord}}, L \right).$$

We recall from §4.3 that there is a rational action of  $\text{GL}_d(k)$  on  $C^{\text{coord},L}$ , which extends in a natural way to a (topological) rational action on all DG-Lie algebras and DG-Lie modules above. The previous actions determine infinitesimally actions of  $\mathfrak{gl}_d(k)$  on all DG-Lie algebras and DG-Lie modules considered so far in the sense of §5.2.

The  $L_\infty$ -morphism  $\mathcal{U}_{L,\omega}$  descends with respect to the action of the set  $\mathfrak{s} = \mathfrak{gl}_d(k)$  (using the notation of §5.2), because the descent condition (5.11) is satisfied as a consequence of Property *iii*) of Theorem 5.1 and of the verticality property (4.5) of  $\omega$ .

Similarly, Property *ii*) of Theorem 5.2, together with the verticality property of  $\omega$  implies that  $\mathcal{S}_{L,\omega}$  descends with respect to the action of  $\mathfrak{gl}_d(k)$  (see §5.3). Summarizing all arguments so far, and because of the compatibility of the  $\text{GL}_d(k)$ -action with the isomorphism (4.2), we get  $L_\infty$ -morphisms

$$\begin{aligned} (U_L^{\text{coord}})^{\mathfrak{gl}_d(k)} : (T_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}}, [ , ])^{\mathfrak{gl}_d(k)} \\ \rightarrow (D_{\text{poly},C^{\text{coord},L}}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + d_H, [ , ])^{\mathfrak{gl}_d(k)} \end{aligned}$$

and

$$\begin{aligned} (S_L^{\text{coord}})^{\mathfrak{gl}_d(k)} : \left( \widehat{C}_{C^{\text{coord},L},\bullet}(C^{\text{coord},L}\widehat{\otimes}_{R_1}JL), {}^1\nabla^{\text{coord}} + b_H, L \circ \mathcal{U}_L^{\text{coord}} \right)^{\mathfrak{gl}_d(k)} \\ \rightarrow \left( \Omega_{C^{\text{coord},L}\widehat{\otimes}F/C^{\text{coord},L}}, {}^1\nabla^{\text{coord}}, L \right)^{\mathfrak{gl}_d(k)}, \end{aligned}$$

Repeating almost *verbatim* the arguments at the end of [7, §7.3.3], there are obvious isomorphisms of DG-Lie algebras and DG-Lie modules

$$\begin{aligned} (T_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}, [\ , \ ] ) &\cong (T_{\text{poly}, C^{\text{coord}, L}}(C^{\text{coord}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{coord}}, [\ , \ ] )^{\mathfrak{gl}_d(k)}, \\ (D_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + d_{\text{H}}, [\ , \ ] ) &\cong (D_{\text{poly}, C^{\text{coord}, L}}(C^{\text{coord}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{coord}} + d_{\text{H}}, [\ , \ ] )^{\mathfrak{gl}_d(k)}, \\ (\Omega_{C^{\text{aff}, L} \widehat{\otimes}_F / C^{\text{aff}, L}}, {}^1\nabla^{\text{aff}}, L) &\cong (\Omega_{C^{\text{coord}, L} \widehat{\otimes}_F / C^{\text{coord}, L}}, {}^1\nabla^{\text{coord}}, L)^{\mathfrak{gl}_d(k)}, \\ (\widehat{C}_{C^{\text{aff}, L}, \bullet}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + b_{\text{H}}, L) &\cong (\widehat{C}_{C^{\text{coord}, L}, \bullet}(C^{\text{coord}, L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{coord}} + b_{\text{H}}, L)^{\mathfrak{gl}_d(k)}. \end{aligned}$$

We now set

$$\begin{aligned} \mathfrak{g}_1^L &= T_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), & \mathfrak{g}_2^L &= D_{\text{poly}, C^{\text{aff}, L}}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), \\ \mathfrak{m}_1^L &= \Omega_{C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL / C^{\text{aff}, L}}, & \mathfrak{m}_2^L &= \widehat{C}_{C^{\text{aff}, L}, \bullet}(C^{\text{aff}, L} \widehat{\otimes}_{R_1} JL), \end{aligned}$$

and  $\mathfrak{U}_L = \mathfrak{U}_L^{\text{aff}}$ ,  $\mathfrak{S}_L = \mathfrak{S}_L^{\text{aff}}$ : combining all the results so far, we get the commutative diagram (5.12), and to prove the claim, it remains to show that  $\mathfrak{U}_L$  and  $\mathfrak{S}_L$  are  $L_\infty$ -quasi-isomorphisms.

The proof of the fact that  $\mathfrak{U}_L$  is a quasi-isomorphism can be found in [7, §7.3.4]; the proof of the fact that  $\mathfrak{S}_L$  is a quasi-isomorphism is dual. We will now sketch it.

The  $L_\infty$ -morphism  $\mathfrak{S}_L$  is obtained from  $\mathcal{S}_{L, \omega}$  using the isomorphism (4.2) and by (5.9) the Taylor components of  $\mathcal{S}_{L, \omega}$  are given by

$$\begin{aligned} \mathcal{S}_{L, \omega, n}(\gamma_1, \dots, \gamma_n; c) &= \sum_{m \geq 0} \frac{1}{m!} \mathcal{S}_{L, n+m}(\underbrace{\omega, \dots, \omega}_m, \gamma_1, \dots, \gamma_n; c), \\ \gamma_i &\in T_{\text{poly}, C^{\text{coord}, L}}(C^{\text{coord}, L} \widehat{\otimes} F), \quad c \in \widehat{C}_{C^{\text{coord}, L}, \bullet}(C^{\text{coord}, L} \widehat{\otimes} F). \end{aligned}$$

$T_{\text{poly}, C^{\text{coord}, L}}(C^{\text{coord}, L} \widehat{\otimes} F)$ ,  $D_{\text{poly}, C^{\text{coord}, L}}(C^{\text{coord}, L} \widehat{\otimes} F)$ ,  $\Omega_{C^{\text{coord}, L} \widehat{\otimes}_F / C^{\text{coord}, L}}$  and  $\widehat{C}_{C^{\text{coord}, L}, \bullet}(C^{\text{coord}, L} \widehat{\otimes} F)$  are bi-graded complexes: the first degree is the natural degree coming from  $C^{\text{coord}, L}$ , while the second degree is associated to poly-vector degree, (shifted) Hochschild degree, (negative) form degree and (negative) Hochschild degree respectively.

The component  $\mathcal{S}_{L, \omega, 0}$  can be written into a sum

$$\mathcal{S}_{L, \omega, 0}(c) = \sum_{n \geq 0} \frac{1}{n!} \mathcal{S}_n(\underbrace{\omega, \dots, \omega}_n; c);$$

the grading property of the  $L_\infty$ -quasi-isomorphism  $\mathcal{S}$  of Theorem 5.2 implies that the component  $\mathcal{S}_{L, \omega, 0}^n$  of  $\mathcal{S}_{L, \omega, 0}$  indexed by  $n$  has bi-degree  $(n, -n)$ .

Dualizing [7, Lemma 7.3.2], and using Property *i*) of Theorem 5.2, we get the following commutative diagram of graded vector spaces:

$$(5.13) \quad \begin{array}{ccc} C_{\text{poly}}^L(R) & \hookrightarrow & C^{\text{coord}, L} \widehat{\otimes} \widehat{C}_{\bullet}(F), \\ \text{HKR} \downarrow & & \downarrow \mathcal{S}_{L, \omega, 0}^0 \\ \Omega^L(R) & \hookrightarrow & C^{\text{coord}, L} \widehat{\otimes} \Omega_F \end{array}$$

where the morphism HKR on the left vertical arrow has been defined in Theorem 3.9.

The twisting procedure and the descent procedure by the isomorphism (4.2) produce the commutative diagram

$$\begin{array}{ccc}
C_{\text{poly}}^L(R) & \hookrightarrow & \widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L}\widehat{\otimes}_{R_1} JL) \\
\text{HKR} \downarrow & & \downarrow \mathfrak{S}_{L,0}^0 \\
\Omega^L(R) & \hookrightarrow & \Omega_{C^{\text{aff},L}\widehat{\otimes}_{R_1} JL/C^{\text{aff},L}}
\end{array}$$

out of the commutative diagram (5.13); the above bi-gradings naturally translate into bi-gradings on  $\Omega_{C^{\text{aff},L}\widehat{\otimes}F/C^{\text{aff},L}}$  and  $\widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L}\widehat{\otimes}F)$ . The component  $\mathfrak{S}_{L,0}$  is a sum of terms  $\mathfrak{S}_{L,0}^n$ ,  $n \geq 0$ , of bi-degree  $(n, -n)$ .

We now prove that the morphisms  $\mathfrak{S}_{L,0}$  and  $\mathfrak{S}_{L,0}^0$  coincide at the level of cohomology. For this, we consider on the double complexes  $\Omega_{C^{\text{aff},L}\widehat{\otimes}F/C^{\text{aff},L}}$  and  $\widehat{C}_{C^{\text{aff},L},\bullet}(C^{\text{aff},L}\widehat{\otimes}F)$  the filtration with respect to the second degree: then, the corresponding spectral sequences degenerate at their first terms, because of the results of §4.5, §4.6, and the resulting complexes consist of single columns  $(\Omega^L(R), 0)$  and  $(C_{\text{poly}}^L(R), \mathfrak{b}_H)$ . Thus, the respective second terms of the spectral sequences coincide with  $\Omega^L(R)$  and with  $H^\bullet(C_{\text{poly}}^L(R), \mathfrak{b}_H)$ . Since both spectral sequences degenerate at their first term (i.e. the cohomology with respect to the first degree is concentrated in degree 0),  $\mathfrak{S}_{L,0}$  and  $\mathfrak{S}_{L,0}^0$  obviously coincide at the level of cohomology, and this ends the proof.  $\square$

**5.6. Functoriality property of Theorem 5.3.** We consider two Lie algebroids  $(L, R)$ ,  $(M, S)$  as above.

**Definition 5.4.** An **algebraic morphism** from  $(L, R)$  to  $(M, S)$  consists of a pair  $(\ell, \lambda)$ , where *i*)  $\lambda$  is a  $k$ -algebra morphism from  $R$  to  $S$ , and *ii*)  $\ell$  is a Lie algebra morphism from  $L$  to  $M$ , enjoying the following compatibility properties with respect to the corresponding anchor maps:

$$\lambda(\ell(r)) = \ell(\lambda(r)), \quad \ell(rl) = \lambda(r)\ell(l), \quad r \in R, \quad l \in L.$$

The universal property of the universal enveloping algebra of a Lie algebroid yields, for any algebraic morphism  $\varphi = (\ell, \lambda)$  from  $(L, R)$  to  $(M, S)$ , a Hopf algebroid morphism  $\varphi_D : U_R(L) \rightarrow U_S(M)$ . Thus,  $(\ell, \lambda)$  defines a morphism  $\varphi_D$  of  $B_\infty$ -algebras from  $D_{\text{poly}}^L(R)$  to  $D_{\text{poly}}^M(S)$ : in particular, it restricts to a morphism of Gerstenhaber algebras up to homotopy.

Further, the algebraic morphism  $\varphi$  defines a morphism  $\varphi_T : T_{\text{poly}}^L(R) \rightarrow T_{\text{poly}}^M(S)$  by extending (via the  $S$ -linear wedge product) the assignment

$$\varphi_T : S \otimes_R L \rightarrow M : s \otimes_R l \mapsto s\ell(l)$$

Since  $(\ell, \lambda)$  preserves the anchor map and Lie bracket, we have a morphism of Gerstenhaber algebras from  $T_{\text{poly}}^L(R)$  to  $T_{\text{poly}}^M(S)$ .

**Proposition 5.5.** *We assume  $(L, R)$ ,  $(M, S)$  to be Lie algebroids over  $R$  and  $S$  respectively, and  $\varphi = (\ell, \lambda)$  to be an algebraic morphism between them as in Definition 5.4; we further assume that the morphism*

$$\varphi_T : S \otimes_R L \rightarrow M : s \otimes_R l \mapsto s\ell(l)$$

*is an isomorphism of  $S$ -modules.*

The morphism  $(\ell, \lambda)$  determines a morphism of DG-algebras

$$(\Omega^L(R), d_L) \xrightarrow{\varphi_\Omega} (\Omega^M(S), d_M),$$

which satisfies

$$(5.14) \quad \varphi_\Omega(\gamma \cap \omega) = \varphi_T(\gamma) \cap \varphi_\Omega(\omega), \quad \gamma \in T_{\text{poly}}^L(R), \quad \omega \in \Omega^L(R),$$

and a morphism of DG-algebras

$$\varphi_J : \mathcal{J}L \rightarrow \mathcal{J}M,$$

which satisfies

$$(5.15) \quad \lambda(\alpha(E)) = \varphi_J(\alpha)(\varphi_D(E)), \quad \alpha \in \mathcal{J}L, \quad E \in \mathcal{U}_R(L),$$

$$(5.16) \quad \varphi_J({}^1\nabla_l \alpha) = {}^1\nabla_{\ell(l)} \varphi_J(\alpha), \quad \alpha \in \mathcal{J}L, \quad l \in L,$$

$$(5.17) \quad \varphi_J({}^2\nabla_l \alpha) = {}^2\nabla_{\ell(l)} \varphi_J(\alpha), \quad \alpha \in \mathcal{J}L, \quad l \in L,$$

and which commutes with the algebra monomorphisms  $\alpha_i$ ,  $i = 1, 2$  (see §3.1.4).

*Proof.* Since  $\varphi_T$  is an isomorphism of  $S$ -modules, we define  $\varphi_\Omega$  on  $L$ -differential forms on  $R$  via

$$\varphi_\Omega(r) = \lambda(r), \quad \varphi_\Omega(l^*)(sl(l)) = s\lambda(l^*(l)), \quad r \in R, \quad s \in S, \quad l \in L, \quad l^* \in L^*,$$

and we extend it to  $\Omega^R(L)$  by  $R$ -linearity and by multiplicativity with respect to the wedge product.

To prove that  $\varphi_\Omega$  intertwines  $d_L$  and  $d_M$ , it suffices to verify the claim on  $R$  and  $L^*$ . In the first case, we have

$$\varphi_\Omega(d_L(r))(sl(l)) = s\lambda(d_L(r)(l)) = s\lambda(l(r)) = sl(l)(\lambda(r)) = sl(l)(\varphi_\Omega(r)) = d_M(\varphi_\Omega(r))(sl(l)),$$

for a general element  $r$  of  $R$ ,  $s$  of  $S$  and  $l$  of  $L$ , while in the second case we have

$$\begin{aligned} \varphi_\Omega(d_L l^*)(s_1 \ell(l_1), s_2 \ell(l_2)) &= s_1 s_2 \lambda(d_L l^*(l_1, l_2)) \\ &= s_1 s_2 \lambda(l_1(l^*(l_2))) - s_1 s_2 \lambda(l_2(l^*(l_1))) - s_1 s_2 \lambda(l^*([l_1, l_2])) \\ &= s_1 \ell(l_1)(s_2 \lambda(l^*(l_2))) + s_1 s_2 \ell(l_1)(\lambda(l^*(l_2))) - \\ &\quad - s_2 \ell(l_2)(s_1 \lambda(l^*(l_1))) - s_1 s_2 \ell(l_2)(\lambda(l^*(l_1))) - \\ &\quad - s_1 \ell(l_1)(s_2 \lambda(l^*(l_2))) + s_2 \ell(l_2)(s_1 \lambda(l^*(l_1))) - s_1 s_2 \lambda(l^*([l_1, l_2])) \\ &= (s_1 \ell(l_1))(\varphi_\Omega(l^*)(s_2 \ell(l_2))) - (s_2 \ell(l_2))(\varphi_\Omega(l^*)(s_1 \ell(l_1))) - \\ &\quad - \varphi_\Omega(l^*)(s_1 \ell(l_1)(s_2 \ell(l_2)) - s_2 \ell(l_2)(s_1 \ell(l_1)) + s_1 s_2 \ell([l_1, l_2])) \\ &= d_M(\varphi_\Omega(l^*))(s_1 \ell(l_1), s_2 \ell(l_2)). \end{aligned}$$

By compatibility with wedge products, it suffices to prove (5.14) for  $\gamma$  in  $R$  or in  $L$ , and for a general  $\omega$ : we check exemplarily the claim for  $\gamma$  in  $L$ , i.e.

$$\begin{aligned} \varphi_\Omega(l \cap \omega)(s_1 \ell(l_1), \dots, s_p \ell(l_p)) &= s_1 \cdots s_p \lambda((l \cap \omega)(l_1, \dots, l_p)) \\ &= s_1 \cdots s_p \lambda(\omega(l, l_1, \dots, l_p)) \\ &= \varphi_\Omega(\omega)(\ell(l), s_1 \ell(l_1), \dots, s_p \ell(l_p)) \\ &= (\varphi_T(l) \cap \varphi_\Omega(\omega))(s_1 \ell(l_1), \dots, s_p \ell(l_p)). \end{aligned}$$

We now define the morphism  $\varphi_J$  on  $\mathcal{J}L$ : for a general element  $\alpha$  of  $\mathcal{J}L$ , we set

$$\varphi_J(\alpha)(s) = s\lambda(\alpha(1)), \quad \varphi_J(\alpha)(s\ell(l_1) \cdots \ell(l_p)) = s\lambda(\alpha(l_1 \cdots l_p)), \quad s \in S, \quad l_i \in L.$$

It is sufficient to define  $\varphi_J$  on such elements of  $U_S(M)$ , since, being  $\varphi_T$  an isomorphism of  $S$ -modules, a general element of  $U_S(M)$  is a sum of elements of the form

$$\begin{aligned} (s_1 \ell(l_1)) \cdots (s_p \ell(l_p)) &= s_1 \ell(l_1) s_2 \ell(l_2) \cdots s_p \ell(l_p) \\ &= s_1 (\ell(l_1)(s_2)) \ell(l_2) \cdots s_p \ell(l_p) + s_1 s_2 \ell(l_1) \ell(l_2) \cdots s_p \ell(l_p) = \cdots, \end{aligned}$$

where the product has to be understood in  $U_S(M)$ .

Since  $\varphi_D$  is defined by extending  $\lambda$  and  $\ell$  in a way compatible with the Lie algebroid structure of  $U_R(L)$ , (5.15) follows immediately.

As for (5.16), it suffices to check the identity on  $R$  and on elements of  $U_S(M)$  of the form  $s \ell(l_1) \cdots \ell(l_p)$ . In the first case, we have for  $s \in S$ ,  $l \in L$

$$\begin{aligned} ({}^1\nabla_{\ell(l)} \varphi_J(\alpha))(s) &= \ell(l)(\varphi_J(\alpha)(s)) - \varphi_J(\alpha)(\ell(l)s) \\ &= \ell(l)(s\lambda(\alpha(1))) - \varphi_J(\alpha)(\ell(l)s) \\ &= \ell(l)(s)\lambda(\alpha(1)) + s\ell(l)\lambda(\alpha(1)) - \varphi_J(\alpha)(\ell(l)(s)) - \varphi_J(\alpha)(s\ell(l)) \\ &= s\ell(l)\lambda(\alpha(1)) - s\lambda(\alpha(l)) \\ &= s\lambda({}^1\nabla_l \alpha)(1) \\ &= \varphi_J({}^1\nabla_l \alpha)(s). \end{aligned}$$

As for the second case, we have for  $\alpha \in \mathcal{J}L$ ,  $l, l_i \in L$ ,  $i = 1, \dots, p$ ,  $s \in S$

$$\begin{aligned} ({}^1\nabla_{\ell(l)} \varphi_J(\alpha))(s \ell(l_1) \cdots \ell(l_p)) &= \ell(l)(\varphi_J(\alpha)(s \ell(l_1) \cdots \ell(l_p))) - \varphi_J(\alpha)(\ell(l)s \ell(l_1) \cdots \ell(l_p)) \\ &= \ell(l)(s\lambda(\alpha(l_1 \cdots l_p))) - \varphi_J(\alpha)(\ell(l)s \ell(l_1) \cdots \ell(l_p)) \\ &= \ell(l)(s)\lambda(\alpha(l_1 \cdots l_p)) + s\ell(l)(\lambda(\alpha(l_1 \cdots l_p))) \\ &\quad - \varphi_J(\alpha)(\ell(l)(s)\ell(l_1) \cdots \ell(l_p)) - \varphi_J(\alpha)(s\ell(l)\ell(l_1) \cdots \ell(l_p)) \\ &= s\lambda(\ell(\alpha(l_1 \cdots l_p))) - s\lambda(\alpha)(\ell l_1 \cdots l_p) \\ &= \varphi_J({}^1\nabla_l \alpha)(s \ell(l_1) \cdots \ell(l_p)). \end{aligned}$$

The identity (5.17) as well as the compatibility with  $\alpha_i$ ,  $i = 1, 2$  are verified by similar computations.  $\square$

Assume now that  $\varphi = (\ell, \lambda) : (L, R) \rightarrow (M, S)$  is as in the previous lemma and that  $\varphi_T : S \otimes_R L \rightarrow M$  is an isomorphism. As always we assume that  $L$  (and hence  $M$ ) is free of rank  $d$ . Looking at associated graded objects we see that the extended map

$$(5.18) \quad S_1 \otimes_{R_1} \mathcal{J}L \rightarrow \mathcal{J}M : s \otimes \alpha \mapsto s\varphi_J(\alpha)$$

is an isomorphism. Hence any  $R_1$ -linear differential operator on  $\mathcal{J}L$  can be extended to an  $S_1$ -linear differential operator on  $\mathcal{J}M$ . We use this to define a map

$$\varphi_D : D_{R_1}(\mathcal{J}L) \rightarrow D_{S_1}(\mathcal{J}M)$$

and a corresponding map of  $B_\infty$ -algebras

$$(5.19) \quad \varphi_D : D_{\text{poly}, R_1}(\mathcal{J}L) \rightarrow D_{\text{poly}, S_1}(\mathcal{J}M)$$

such that the following diagram is commutative

$$(5.20) \quad \begin{array}{ccc} D_{\text{poly}}^L(R) & \xrightarrow{\varphi_D} & D_{\text{poly}}^M(S) \\ \downarrow & & \downarrow \\ D_{\text{poly},R_1}(JL) & \xrightarrow{\varphi_D} & D_{\text{poly},S_1}(JM) \end{array}$$

where the vertical monomorphisms have been defined in (3.12).

An easy computation shows that  $\varphi_D$  in (3.12) commutes with the action of the Grothendieck connection  $[{}^1\nabla, -]$ . It follows by the discussion in §3.3 that if we take the invariants for  $[{}^1\nabla, -]$  of the lower line in (5.20) we obtain the upper line.

We extend  $\varphi_J$  to a map of graded vector spaces

$$\varphi_C : \widehat{C}_{R,\bullet}(JL) \rightarrow \widehat{C}_{S,\bullet}(JM) : \alpha_1 \otimes \cdots \otimes \alpha_n \mapsto \varphi_J(\alpha_1) \otimes \cdots \otimes \varphi_J(\alpha_n)$$

which is again essentially just base extension over  $S/R$ . This map obviously commutes with the Grothendieck connection  ${}^1\nabla$ . We obtain a map of pairs of graded vector spaces

$$(\varphi_D, \varphi_C) : (D_{\text{poly},R}(JL), \widehat{C}_{R,\bullet}(JL)) \rightarrow (D_{\text{poly},S}(JM), \widehat{C}_{S,\bullet}(JM))$$

and as this map is just base extension over  $S/R$  it is compatible with all structures defined in [6], hence in particular with the DG-Lie algebra and DG-Lie module structures and also with the precalculi up to homotopy.

Taking invariants for  ${}^1\nabla$  and using (3.16) we obtain a commutative diagram of precalculus structure up to homotopy

$$\begin{array}{ccc} (D_{\text{poly}}^L(R), d_H, [ , ], \cup) & \xrightarrow{\varphi_D} & (D_{\text{poly}}^M(S), d_H, [ , ], \cup) \\ \Downarrow & & \Downarrow \\ (C_{\text{poly}}^L(R), b_H, L, \cap) & \xrightarrow{\varphi_C} & (C_{\text{poly}}^M(S), b_H, L, \cap) \end{array}$$

One also obtains from Proposition 5.5 a commutative diagram of precalculi.

$$\begin{array}{ccc} (T_{\text{poly}}^L(R), 0, [ , ], \cup) & \xrightarrow{\varphi_T} & (T_{\text{poly}}^M(S), 0, [ , ], \cup) \\ \Downarrow & & \Downarrow \\ (\Omega^L(R), 0, L, \cap) & \xrightarrow{\varphi_\Omega} & (\Omega^M(S), 0, L, \cap) \end{array}$$

Furthermore from (5.18) and the universal property of coordinate spaces (see (4.1)) we obtain an  $R$ -algebra morphism from  $R^{\text{coord},L}$  to  $S^{\text{coord},M}$ . It extends further to a morphism of DG-algebras from  $C^{\text{coord},L}$  to  $C^{\text{coord},M}$  thanks to (5.16) and the fact that  $\varphi_\Omega$  is a morphism of DG-algebras from  $\Omega^L(R)$  to  $\Omega^M(S)$ .

Finally, the algebraic morphism  $(\ell, \lambda)$  induces precalculi morphisms (up to homotopy) between all corresponding Fedosov resolutions, since the monomorphism  $\alpha_2$  and the connection  ${}^2\nabla$ , which are needed in the construction of the Fedosov resolutions of §4 (we refer to [7] for more details thereabout), have been proved to be preserved by  $(\ell, \lambda)$ .

As a consequence of these arguments, we deduce the following theorem, which expresses the functoriality properties of the commutative diagram (5.12) of Theorem 5.3.

**Theorem 5.6.** *For a general algebraic morphism  $\varphi = (\ell, \lambda)$  from  $(L, R)$  to  $(M, S)$  as in Definition 5.4, which induces an isomorphism  $S \otimes_R L \cong M$  of  $S$ -modules, and such that  $L$  is free of rank  $d$ , the  $L_\infty$ -quasi-isomorphisms  $\mathfrak{U}_L, \mathfrak{U}_M, \mathfrak{S}_L$  and  $\mathfrak{S}_M$  of DG-Lie algebras and DG-Lie modules fit into the commutative diagram*

$$(5.21) \quad \begin{array}{ccccccc} T_{\text{poly}}^L(R) & \hookrightarrow & \mathfrak{g}_1^L & \xrightarrow{\mathfrak{U}_L} & \mathfrak{g}_2^L & \longleftarrow & D_{\text{poly}}^L(R) \\ \varphi_T \downarrow & & \varphi_T \downarrow & & \varphi_D \downarrow & & \varphi_D \downarrow \\ T_{\text{poly}}^M(S) & \hookrightarrow & \mathfrak{g}_1^M & \xrightarrow{\mathfrak{U}_M} & \mathfrak{g}_2^M & \longleftarrow & D_{\text{poly}}^M(S) \\ \mathcal{L} \downarrow & & \mathcal{L}_1 \downarrow & & \mathcal{L}_2 \downarrow & & \mathcal{L} \downarrow \\ \Omega^M(S) & \hookrightarrow & \mathfrak{m}_1^M & \xleftarrow{\mathfrak{S}_M} & \mathfrak{m}_2^M & \longleftarrow & C_{\text{poly}}^M(S) \\ \varphi_\Omega \uparrow & & \varphi_\Omega \uparrow & & \varphi_J \uparrow & & \varphi_J \uparrow \\ \Omega^L(R) & \hookrightarrow & \mathfrak{m}_1^L & \xleftarrow{\mathfrak{S}_L} & \mathfrak{m}_2^L & \longleftarrow & C_{\text{poly}}^L(R) \end{array}$$

where we have borrowed notation from Proposition 5.5; all such morphisms are compatible with respect to the composition of algebraic morphisms between Lie algebroids.

Note that Theorem 5.6 makes no reference to the (homotopy) precalculus structures which we discussed above; we will need these below.

**5.7. Proof of Theorem 1.3.** We now collect the results of §5.5 and §5.6 to give the proof of Theorem 1.3, *via* a well-suited gluing procedure.

We consider a ringed site  $(X, \mathcal{O})$ , and a sheaf of Lie algebroids  $\mathcal{L}$ , such that  $\mathcal{L}$  is locally free of rank  $d$  over  $\mathcal{O}$ . We replace  $X$  by its full subcategory of objects  $U$  such that  $\mathcal{L}(U)$  is free over  $\mathcal{O}(U)$ . This does not change the category of sheaves.

All sheaves of DG-Lie algebras and DG-Lie modules in the commutative diagram (1.7) are obtained by sheafifying the corresponding presheaves of DG-Lie algebras and DG-Lie modules, i.e.

$$\begin{aligned} U &\rightarrow T_{\text{poly}}^{\mathcal{L}(U)}(\mathcal{O}(U)), & U &\rightarrow D_{\text{poly}}^{\mathcal{L}(U)}(\mathcal{O}(U)), \\ U &\rightarrow \Omega^{\mathcal{L}(U)}(\mathcal{O}(U)), & U &\rightarrow C_{\text{poly}}^{\mathcal{L}(U)}(\mathcal{O}(U)). \end{aligned}$$

Since  $\mathcal{L}$  is locally free of order  $d$  over  $\mathcal{O}$ , for a morphism  $V \rightarrow U$  in  $X$ , the corresponding restriction morphism  $(\mathcal{O}(U), \mathcal{L}(U)) \rightarrow (\mathcal{O}(V), \mathcal{L}(V))$ , yields an isomorphism

$$\mathcal{O}(V) \otimes_{\mathcal{O}(U)} \mathcal{L}(U) \cong \mathcal{L}(V).$$

Thus, any restriction morphism as above may be viewed as an algebraic morphism between Lie algebroids, satisfying the isomorphism property of Theorem 5.6.

If we then consider the DG-Lie algebras and DG-Lie modules

$$U \rightarrow \mathfrak{g}_i^{\mathcal{L}(U)}, \quad U \rightarrow \mathfrak{m}_i^{\mathcal{L}(U)}, \quad i = 1, 2,$$

Theorem 5.3 produces, for any  $U$  in  $X$ ,  $L_\infty$ -quasi-isomorphisms  $\mathfrak{U}_{\mathcal{L}(U)}$  and  $\mathfrak{S}_{\mathcal{L}(U)}$  which fit into a commutative diagram (5.12). By Theorem 5.6 these are actually morphisms of presheaves.

Sheafifying all presheaves and morphisms between presheaves concludes the proof.



## 6. THE RELATIONSHIP BETWEEN ATIYAH CLASSES AND JET BUNDLES

In the present section we review some technical results from [7, §8], to which we refer for more details. We need only the main notation and conventions for use in §7.

For a field  $k$  of characteristic 0, we consider a sheaf  $\mathcal{L}$  of Lie algebroids over a ringed site  $(X, \mathcal{O})$ , which is locally free of rank  $d$  over  $\mathcal{O}$ .

We have a short exact sequence of  $\mathcal{O}_1$ - $\mathcal{O}_2$ -bimodules

$$0 \rightarrow \mathcal{L}^* \rightarrow J^1\mathcal{L} \rightarrow \mathcal{O} \rightarrow 0$$

where  $\mathcal{O}_i$ ,  $i = 1, 2$ , denotes a copy of  $\mathcal{O}$  embedded in  $J\mathcal{L}$  via the monomorphism  $\alpha_i$  and where  $J^1\mathcal{L}$  was introduced in §3.1.4.

For a general  $\mathcal{O}$ -module  $\mathcal{E}$ , tensoring over  $\mathcal{O}_2$  yields a short exact sequence

$$0 \longrightarrow \mathcal{L}^* \otimes_{\mathcal{O}} \mathcal{E} \longrightarrow J^1\mathcal{L} \otimes_{\mathcal{O}_2} \mathcal{E} \longrightarrow \mathcal{E} \longrightarrow 0$$

which we will call the  $\mathcal{L}$ -Atiyah sequence. The  $\mathcal{L}$ -Atiyah class  $A_{\mathcal{L}}(\mathcal{E})$  of  $\mathcal{E}$  over  $\mathcal{L}$  is the extension class of this sequence in  $\text{Ext}_{\mathcal{O}}^1(\mathcal{E}, \mathcal{L}^* \otimes_{\mathcal{O}} \mathcal{E})$ . As explained in §1.1, if  $\mathcal{E}$  is a vector bundle, the  $i$ -th scalar Atiyah class  $a_{\mathcal{L},i}(\mathcal{E})$  of  $\mathcal{E}$  is defined as

$$(6.1) \quad a_{\mathcal{L},i}(\mathcal{E}) = \text{tr}\left(\bigwedge^i A_{\mathcal{L}}(\mathcal{E})\right) \in \text{H}^i(X, \bigwedge^i \mathcal{L}^*),$$

Below we will only consider the case  $\mathcal{E} = \mathcal{L}$ . In that case we simplify the notation to

$$A(\mathcal{L}) = A_{\mathcal{L}}(\mathcal{L}), \quad a_i(\mathcal{L}) = a_{i,\mathcal{L}}(\mathcal{L})$$

Observe that the  $a_i(\mathcal{L})$  are cohomology classes. We now outline how we may realize them as explicit cocycles.

By the very construction of  $C^{\text{coord},\mathcal{L}}$  and  $C^{\text{aff},\mathcal{L}}$ , there are natural morphisms of DG-algebras

$$(6.2) \quad \Omega^{\mathcal{L}_2}(X) \xrightarrow{\theta} C^{\text{aff},\mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1} \hookrightarrow C^{\text{coord},\mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1} \cong C^{\text{coord},\mathcal{L}} \widehat{\otimes} \Omega_F.$$

The differentials on the first three DG-algebras are the natural ones (see §4.3). The differential on the fourth DG-algebra is  $d + L_{\omega}$  for a certain MC element  $\omega \in C^{\text{coord},\mathcal{L}} \widehat{\otimes} \text{Der}(F)$  and  $d$  the natural differential. See again §4.3.

The MC element  $\omega$  can be expressed as

$$\omega = \eta_{\alpha} \omega_{\alpha,i} \partial_{x_i}, \quad i = 1, \dots, d,$$

where  $\eta_{\alpha}$  is in  $C^{\text{coord},\mathcal{L}}$  and has degree 1,  $\omega_{\alpha,i}$  belongs to  $F$  and  $\partial_{x_i} = \partial/\partial x_i$ .

If we define  $\Xi$  to be the matrix with entries

$$(6.3) \quad \Xi_{ij} = \eta_{\alpha} d_F(\partial_{x_j} \omega_{\alpha,i}) \in C^{\text{coord},\mathcal{L}} \widehat{\otimes} \Omega_F,$$

where  $d_F$  is the De Rham differential on  $\Omega_F$ , then on the nose we have

$$\text{Tr}(\Xi^n) \in C^{\text{coord},\mathcal{L}} \widehat{\otimes} \Omega_F$$

Furthermore it is true that

$$(d + L_{\omega})(\text{Tr}(\Xi^n)) = 0$$

It is shown in [7, §8] that  $\text{Tr}(\Xi^n)$  is actually the image of a (necessarily unique) element in  $C^{\text{aff},\mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1}$ . Abusing notation somewhat we will still write this

element as  $\text{Tr}(\Xi^n)$ . It is still a cocycle and in this way represents an element of

$$\overline{\text{Tr}(\Xi^n)} \in \Gamma(X, \mathbb{H}^{2n}(C^{\text{aff}, \mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1}))$$

which maps naturally to the hypercohomology

$$\mathbb{H}^{2n}(X, C^{\text{aff}, \mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1})$$

Further, we observe that the injection  $\Omega^{\mathcal{L}}(X) \xrightarrow{\theta} C^{\text{aff}, \mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1}$  of DG-algebras is a quasi-isomorphism, as discussed in §4.5. Thus  $\theta$  induces an isomorphism

$$\bigoplus_{m,n} \mathbb{H}^m(X, \wedge^n \mathcal{L}^*) = \mathbb{H}^\bullet(X, \Omega^{\mathcal{L}}(X)) \xrightarrow{\mathbb{H}(\theta)} \mathbb{H}^\bullet(X, C^{\text{aff}, \mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1})$$

The following identity is [7, eq. (8.8)]

$$(6.4) \quad a_n(\mathcal{L}) = \mathbb{H}(\theta)^{-1} \left( \overline{\text{Tr}(\Xi^n)} \right), \quad n \geq 1,$$

which indeed expresses  $a_n(\mathcal{L})$  in terms of the explicit cocycle  $\text{Tr}(\Xi^n)$ .

## 7. PROOF OF THEOREM 1.1

The aim of this Section is to prove Theorem 1.1 which implies Căldăraru's conjecture (Theorem 1.4) as has been outlined in the introduction.

For this purpose, we first remind the reader of the main result of [6] about compatibility between cap products. We then prove a ring-theoretical globalized version of this result (compare to the proof of Theorem 5.3). By functoriality (see §5.6), we obtain the sheaf-theoretical globalization. Finally, using results of [6], we compute explicitly the isomorphism appearing in the compatibility between cap products, which we identify with the action of the homological HKR-quasi-isomorphism followed by left multiplication by the square root of the (modified) Todd class.

**7.1. A *memento* of compatibility between cup and cap products.** In this section, we present a *memento* of the main results of [6, 7] concerning compatibility between cup and cap products respectively.

First of all as before  $F$  is the algebra of formal power series in  $d$  variables over the field  $k$  which we assume to contain  $\mathbb{R}$  for now. We recall the existence of (homotopy) Gerstenhaber algebra structures on  $T_{\text{poly}}(F)$  and  $D_{\text{poly}}(F)$ , which together with  $\Omega_F$  and  $\widehat{C}_\bullet(F)$  yield (homotopy) precalculi [6].

We recall also the  $L_\infty$ -quasi-isomorphisms  $\mathcal{U}$  introduced in Theorem 5.1 and  $\mathcal{S}$  introduced in Theorem 5.2. We denote by  $\mathcal{U}_n$ ,  $n \geq 1$ , resp.  $\mathcal{S}_n$ ,  $n \geq 0$ , the  $n$ -th Taylor component of  $\mathcal{U}$ , resp.  $\mathcal{S}$ .

We further consider a commutative DG-algebra  $(\mathfrak{m}, d_{\mathfrak{m}})$ . The precalculus structures on  $(T_{\text{poly}}(F), \Omega_F)$  and  $(D_{\text{poly}}(F), \widehat{C}_\bullet(F))$ , can be extended by  $\mathfrak{m}$ -linearity to precalculi  $(T_{\text{poly}}^{\mathfrak{m}}(F), \Omega_F^{\mathfrak{m}}) = (T_{\text{poly}}(F) \widehat{\otimes} \mathfrak{m}, \Omega_F \widehat{\otimes} \mathfrak{m})$  and  $(D_{\text{poly}}^{\mathfrak{m}}(F), \widehat{C}_\bullet^{\mathfrak{m}}(F)) = (D_{\text{poly}, \mathfrak{m}}(F \widehat{\otimes} \mathfrak{m}), \widehat{C}_{\bullet, \mathfrak{m}}(F \widehat{\otimes} \mathfrak{m}))$ .

**Convention.** Below we will work with potentially infinite series with coefficients in  $\mathfrak{m}$ . We make the standard assumption that we are in a setting where all these series converge and standard series manipulations are allowed. In our actual application all series will be finite for degree reasons.

A MC element  $\gamma$  of  $T_{\text{poly}}^{\mathfrak{m}}(F)$  can be written as a sum

$$\gamma = \gamma_{-1} + \gamma_0 + \gamma_1 + \gamma_2 + \cdots,$$

where  $\gamma_i$  is an element of  $T_{\text{poly}}^{\mathfrak{m}}(F)$  of poly-vector degree  $i$ ,  $i \geq -1$ , which satisfies the Maurer–Cartan equation

$$d_{\mathfrak{m}}\gamma + \frac{1}{2}[\gamma, \gamma] = 0.$$

We denote by  $\mathcal{U}(\gamma)$  the image of a MC element  $\gamma$  as above with respect to  $\mathcal{U}$  (see (5.7)). This is again a MC element. Further, we set

$$\begin{aligned} \mathcal{U}_{\gamma,1}(\gamma_1) &= \sum_{n \geq 0} \frac{1}{n!} \mathcal{U}_{n+1}(\underbrace{\gamma, \dots, \gamma}_n, \gamma_1), \quad \gamma_1 \in T_{\text{poly}}^{\mathfrak{m}}(F), \\ \mathcal{S}_{\gamma,0}(c) &= \sum_{n \geq 0} \frac{1}{n!} \mathcal{S}_n(\underbrace{\gamma, \dots, \gamma}_n; c), \quad c \in \widehat{C}_{\bullet}^{\mathfrak{m}}(F). \end{aligned}$$

Since  $\mathcal{U}$  and  $\mathcal{S}_{\gamma}$  are  $L_{\infty}$ -quasi-isomorphisms,  $\mathcal{U}_{\gamma,1}$  and  $\mathcal{S}_{\gamma,0}$  are both quasi-isomorphisms of DG-vector spaces.

**Theorem 7.1.** *For a general commutative DG-algebra  $(\mathfrak{m}, d_{\mathfrak{m}})$  as above, and for a general MC element  $\gamma$  of  $T_{\text{poly}}^{\mathfrak{m}}(F)$ ,  $\mathcal{U}_{\gamma,1}$  and  $\mathcal{S}_{\gamma,0}$  descend to quasi-isomorphisms of (homotopy) precalculi, fitting into the commutative diagram*

$$\begin{array}{ccc} \left( T_{\text{poly}}^{\mathfrak{m}}(F), d_{\mathfrak{m}} + [\gamma, \bullet], [ \ , \ ], \cup \right) & \xrightarrow{\mathcal{U}_{\gamma,1}} & \left( D_{\text{poly}}^{\mathfrak{m}}(F), d_{\mathfrak{m}} + d_{\text{H}} + [\mathcal{U}(\gamma), \bullet], [ \ , \ ], \cup \right), \\ \Downarrow & & \Downarrow \\ \left( \Omega_F^{\mathfrak{m}}, d_{\mathfrak{m}} + L_{\gamma}, L, \cap \right) & \xleftarrow{\mathcal{S}_{\gamma,0}} & \left( \widehat{C}_{\bullet}^{\mathfrak{m}}(F), d_{\mathfrak{m}} + b_{\text{H}} + L_{\mathcal{U}(\gamma)}, L, \cap \right) \end{array}$$

in the sense that  $\mathcal{U}_{\gamma,1}$  and  $\mathcal{S}_{\gamma,0}$  preserve Lie brackets, Lie actions, cup and cap products up to homotopy.

Kontsevich [17] has first stated and proved that  $\mathcal{U}_{\gamma,1}$  defines a quasi-isomorphism of Gerstenhaber algebras up to homotopy from  $T_{\text{poly}}^{\mathfrak{m}}(F)$  to  $D_{\text{poly}}^{\mathfrak{m}}(F)$  in the sense specified above. We observe that the identity  $\mathcal{U}_{\gamma,1}([\gamma_1, \gamma_2]) = [\mathcal{U}_{\gamma,1}(\gamma_1), \mathcal{U}_{\gamma,1}(\gamma_2)]$  at the level of cohomology, for  $\gamma_i$  in  $T_{\text{poly}}^{\mathfrak{m}}(F)$ ,  $i = 1, 2$ , holds true, because  $\mathcal{U}$  is an  $L_{\infty}$ -morphism: in particular, there is a homotopy operator describing the compatibility with Lie brackets, expressible in terms of the Taylor components of  $\mathcal{U}$  twisted by the MC element  $\gamma$ . On the other hand, the identity  $\mathcal{U}_{\gamma,1}(\gamma_1 \cup \gamma_2) = \mathcal{U}_{\gamma,1}(\gamma_1) \cup \mathcal{U}_{\gamma,1}(\gamma_2)$  at the level of cohomology comes from a more complicated identity up to homotopy: in this situation, the homotopy operator is not expressible in terms of the Taylor components of  $\mathcal{U}$ . For an explicit description of the homotopy operator, we refer to [6, 7, 18].

The actual formulation of Theorem 7.1 has been first proposed as a conjecture in the particular case, where  $\gamma$  is a (formal) Poisson structure, by Shoikhet [21]: this conjecture has been first proved in [22] only in degree 0 and later in [5] for all degrees. A more general result has been stated and proved in [6], to which we refer for more details. The identity  $\mathcal{S}_{\gamma,0}(L_{\mathcal{U}_{\gamma,1}}(\gamma_1)(c)) = L_{\gamma_1}(\mathcal{S}_{\gamma,0}(c))$  at the level of cohomology, for  $\gamma_1$  in  $T_{\text{poly}}^{\mathfrak{m}}(F)$ ,  $c$  in  $\widehat{C}_{\bullet}^{\mathfrak{m}}(F)$ , is a consequence of the fact that  $\mathcal{S}_{L,\gamma}$  is an  $L_{\infty}$ -morphism of  $L_{\infty}$ -modules (in particular, there is a homotopy

formula involving the Taylor components of  $\mathcal{U}$  and  $\mathcal{S}$ , twisted by  $\gamma$ ). The identity  $\mathcal{S}_{\gamma,0}(\mathcal{U}_{\gamma,1}(\gamma_1) \cap c) = \gamma_1 \cap \mathcal{S}_{\gamma,0}(c)$  at the level of cohomology holds true in virtue of a homotopy formula, but the corresponding homotopy operator does not involve the Taylor components of  $\mathcal{U}$  and  $\mathcal{S}$ : such an operator has been explicitly described in [6].

We briefly review in section §7.1.1 below the construction of the homotopy operator for the compatibility between cap products.

*7.1.1. The homotopy formula for the compatibility between cap products.* For later computations, we write down the explicit homotopy operator for the compatibility between the  $\cap$ -actions: namely, for a MC element  $\gamma$  as in Theorem 7.1, for  $\gamma_1$  a general element of  $T_{\text{poly}}^m(F)$  and  $c$  a general element of  $\widehat{C}_\bullet^m(F)$ , we have the homotopy relation

$$(7.1) \quad \begin{aligned} \mathcal{S}_{\gamma,0}(\mathcal{U}_{\gamma,1}(\gamma_1) \cap c) - \gamma_1 \cap \mathcal{S}_{\gamma,0}(c) &= (d_m + L_\gamma)\mathcal{H}_\gamma^{\mathcal{S}}(\gamma_1, c) + \mathcal{H}_\gamma^{\mathcal{S}}(d_m \gamma_1 + [\gamma, \gamma_1], c) + \\ &+ (-1)^{|\gamma_1|} \mathcal{H}_\gamma^{\mathcal{S}}(\gamma_1, d_m c + b_H c + L_{\mathcal{U}(\gamma)} c), \end{aligned}$$

where

$$(7.2) \quad \mathcal{H}_\gamma^{\mathcal{S}}(\gamma_1, c) = \sum_{n \geq 0} \frac{1}{n!} \sum_{\Gamma \in \mathcal{G}_{n+1, m+1}^{\mathcal{S}}} \overset{\circ}{W}_{D, \Gamma} \mathcal{S}_\Gamma(\gamma_1, \underbrace{\gamma, \dots, \gamma}_n, c),$$

$c$  being of Hochschild degree  $-m$ .

In (7.2), the second sum is over “ $\mathcal{S}$ -admissible graphs” of type  $(n+1, m+1)$ : these are directed graphs with  $n+2$  vertices of the first type and  $m+1$  cyclically ordered vertices of the second type and with an orientation of the outgoing edges from vertices of the first type, and with a special vertex of the first type, labelled by 0. The vertices of the second type can be only endpoints of edges, and  $\mathcal{S}$ -admissible graphs do not contain edges starting and ending at the same vertex; finally, the vertex 0 has only incoming edges.

To the vertex 1 of the first type of an  $\mathcal{S}$ -admissible graph  $\Gamma$  is assigned the poly-vector field  $\gamma_1$ : the number of outgoing edges from 1 equals the poly-vector degree of  $\gamma_1$  plus 1. To any other vertex of the first type, except 0, is assigned a copy of the MC element  $\gamma$ . To the  $i$ -th vertex of the second type is assigned the  $i+1$ -th component of the Hochschild chain  $c$ . Pictorially, here is an  $\mathcal{S}$ -admissible graph of type  $(4, 5)$ , with corresponding coloring by poly-vector fields and Hochschild chains:




Figure 1 - An  $\mathcal{S}$ -admissible graph of type  $(4, 5)$

The differential form  $S_\Gamma(\gamma_1, \underbrace{\gamma_2, \dots, \gamma_n}_n, c)$  is defined explicitly in [5, 6, 21].

More important for our purposes is the integral weight  $\overset{\circ}{W}_{D,\Gamma}$ , for a general  $S$ -admissible graph of type  $(n + 1, m + 1)$ ,

$$(7.3) \quad \overset{\circ}{W}_{D,\Gamma} = \int_{\overset{\circ}{\mathcal{Y}}_{n+1,m+1}^+} \omega_{D,\Gamma}.$$

First of all,  $\overset{\circ}{\mathcal{Y}}_{n+1,m+1}^+$  denotes the codimension-1-submanifold (with corners) of the compactified configuration space  $\mathcal{D}_{n+1,m+1}^+$  of  $n + 1$  points in the punctured unit disk  $D^\times$  and  $m + 1$  cyclically oriented points in  $S^1$ , consisting of configurations of points, where the point labelled by 1 moves on a smooth curve from the origin to the first point  $\bar{1}$  (with respect to the cyclic order) in  $S^1$ . Graphically,




Figure 2 - A general configuration of points in  $\overset{\circ}{\mathcal{Y}}_{n+1,m+1}^+$

In Figure 1, the dashed line represents the curve, along which the point 1 (labelled as “o”) moves. The differential form  $\omega_{D,\Gamma}$  associated to a graph in  $\mathcal{G}_{n+1,m+1}^S$  is a product of smooth 1-forms on  $\mathcal{D}_{n+1,m+1}^+$ : the basic ingredient is a slight modification of the exterior derivative of Kontsevich’s angle function, see [6, 17] for more details.

For the globalization procedure of the compatibility between cap products, we need the following technical Lemma, which corresponds, in the present framework, to Theorem 5.2, *ii*).

**Lemma 7.2.** *If  $\Gamma$  is an  $S$ -admissible graph in  $\mathcal{G}_{n+1,m+1}^S$ ,  $n \geq 1$ , and at least one of the poly-vector fields  $\gamma_i$ ,  $i \neq 1$ , is linear on  $F$ , then*

$$\overset{\circ}{W}_{D,\Gamma} S_\Gamma(\gamma_1, \gamma_2, \dots, \gamma_{n+1}, c) = 0.$$

*Proof.* The first point of the first type in  $\overset{\circ}{\mathcal{Y}}_{n+1,m+1}^+$ , by the very construction of  $\overset{\circ}{\mathcal{Y}}_{n+1,m+1}^+$ , moves from the origin 0 to the first point in  $S^1$  with respect to the cyclic order: to the former point is associated the poly-vector field  $\gamma_1$ . Any other point associated to a vertex of the first type moves freely in the punctured unit disk  $D^\times$ .

Without loss of generality we assume  $\gamma_2$  to be an  $\mathbf{m}$ -valued linear vector field: the valence (i.e. the number of outgoing edges) of the corresponding vertex of the first type is 1, while the linearity of  $\gamma_2$  implies that there can be at most one incoming

edge to the vertex corresponding to  $\gamma_2$ . This follows from the construction of the differential form  $\mathcal{S}_\Gamma(\gamma_1, \gamma_2, \dots, \gamma_{n+1}, c)$ .

Thus, we may safely restrict to  $S$ -admissible graphs  $\Gamma$ , such that the vertex 2 has valence exactly 1 and with at most one incoming edge.

If the vertex labelled by 2 does not have incoming edges, the corresponding integral weight  $\overset{\circ}{W}_{D,\Gamma}$  vanishes by dimensional reasons: in fact, we integrate a 1-form (corresponding to the only outgoing edge from 2) over a 2-dimensional submanifold (with corners) of  $D^\times$ .

If the vertex labelled by 2 has exactly one incoming and one outgoing edge, we may apply [6, Lemma 6.1], to yield the vanishing of the corresponding weight  $\overset{\circ}{W}_{D,\Gamma}$ .  $\square$

**7.2. The proof of Theorem 1.1 in the ring case.** We will first assume that the ground field contains  $\mathbb{R}$ . At the end of the section we will show how to get rid of this restriction.

We consider a Lie algebroid  $L$  over  $R$  as in Definition 3.1 free of rank  $d$  over  $R$ . Then, we set  $(\mathfrak{m}, d_{\mathfrak{m}}) = (C^{\text{coord},L}, d)$ , where  $d = d_{\Omega_{R^{\text{coord}},L}} \widehat{\otimes}_{\Omega_{R_1}} 1 + 1 \widehat{\otimes}_{\Omega_{R_1}} d_{L_1}$  (see §4.3 for more details), and the Maurer–Cartan form is an  $\mathfrak{m}$ -valued vector field on  $F$  obeying

$$d\omega + \frac{1}{2}[\omega, \omega] = 0.$$

By Theorem 5.1, *ii*) we have  $\mathcal{U}(\omega) = \omega$ . Furthermore one checks that by degree reasons  $\mathcal{U}_\omega$  and  $\mathcal{S}_\omega$  yield finite sums when evaluated on specific elements. The same goes for the associated homotopies. So the results of §7.1 apply.

Combining the arguments of the proof of Theorem 5.3 with Theorem 7.1 we get the following commutative diagram of precalculus structures up to homotopy (7.4)

$$\begin{array}{ccc} (T_{\text{poly}, C^{\text{aff},L}}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}}, [\ , \ ], \cup) & \xrightarrow{\mathfrak{U}_{L,1}} & (D_{\text{poly}, C^{\text{aff},L}}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + d_{\text{H}}, [\ , \ ], \cup) \\ \Downarrow & & \Downarrow \\ (\Omega_{C^{\text{aff},L} \widehat{\otimes}_{R_1} JL / C^{\text{aff},L}}, {}^1\nabla^{\text{aff}}, L, \cap) & \xleftarrow{\mathfrak{S}_{L,0}} & (\widehat{C}_{C^{\text{aff},L}, \bullet}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL), {}^1\nabla^{\text{aff}} + b_{\text{H}}, L, \cap) \end{array}$$

The fact that  $\mathfrak{U}_{L,1}$  preserves the respective Lie brackets up to homotopy is a consequence of the fact that  $\mathfrak{U}_L$  is an  $L_\infty$ -morphism; similarly, the fact that  $\mathfrak{S}_{L,0}$  preserves the Lie module structure up to homotopy is a consequence of the fact that  $\mathfrak{S}_L$  is an  $L_\infty$ -morphism of  $L_\infty$ -modules.

On the other hand,  $\mathfrak{U}_{L,1}$  is compatible with respect to the products labelled by  $\cup$  up to homotopy by the results of [7, §10.1].

As for the compatibility between the actions labelled by  $\cap$  up to homotopy, we first observe that the homotopy formula (7.1) is well-defined in the case  $(\mathfrak{m}, d_{\mathfrak{m}}) = (C^{\text{coord},L}, d)$  and  $\gamma = \omega$ , with the same notation as above: by the same arguments as in the proof of Theorem 5.3 it remains to prove that the homotopy operator (7.2) descends to a homotopy operator

$$\mathfrak{H}_L^{\mathcal{S}} : T_{\text{poly}, C^{\text{aff},L}}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL) \otimes \widehat{C}_{C^{\text{aff},L}, \bullet}(C^{\text{aff},L} \widehat{\otimes}_{R_1} JL) \rightarrow \Omega_{C^{\text{aff},L} \widehat{\otimes}_{R_1} JL / C^{\text{aff},L}}.$$

This holds true as a consequence of Lemma 7.2 together with the verticality property of the Maurer–Cartan form  $\omega$ , see §4.3.

If we now couple the commutative diagram (7.4) with the results of §4.4, §4.5 and §4.6, and using the same notation introduced at the end of the proof of Theorem 5.3 we get the following commutative diagram of precalculi up to homotopy

$$(7.5) \quad \begin{array}{ccccccc} T_{\text{poly}}^L(R) & \hookrightarrow & \mathfrak{g}_1^L & \xrightarrow{\mathfrak{U}_{L,1}} & \mathfrak{g}_2^L & \longleftarrow & D_{\text{poly}}^L(R) \\ \downarrow & & \downarrow & & \downarrow & & \downarrow \\ \Omega^L(R) & \hookrightarrow & \mathfrak{m}_1^L & \xleftarrow{\mathfrak{S}_{L,0}} & \mathfrak{m}_2^L & \longleftarrow & C_{\text{poly}}^L(R) \end{array} .$$

The quasi-isomorphisms  $\mathfrak{U}_{L,1}$  and  $\mathfrak{S}_{L,0}$  are obtained from  $\mathcal{U}_{L,\omega,1}$  and  $\mathcal{S}_{L,\omega,0}$  respectively by means of the descent procedure: since  $\omega$  is an  $\mathfrak{m}$ -valued vector field in  $T_{\text{poly}}^{\mathfrak{m}}(F) = \mathfrak{g}_1^L$ , for  $\mathfrak{m} = C^{\text{coord},L}$ , we can use the results of [7, §10.1], and [6, §6], to evaluate explicitly  $\mathcal{U}_{L,\omega,1}$  and  $\mathcal{S}_{L,\omega,0}$ , namely

$$(7.6) \quad \mathcal{U}_{L,\omega,1} = \text{HKR} \circ \iota_{j(\omega)}, \quad \mathcal{S}_{L,\omega,0} = j(\omega) \wedge \text{HKR},$$

where

$$(7.7) \quad j(\omega) = \det \sqrt{\frac{\Xi}{\exp(\frac{\Xi}{2}) - \exp(-\frac{\Xi}{2})}},$$

with  $\Xi$  as defined in (6.3). To interpret (7.7) one should expand the right-hand side formally in terms of  $\text{Tr}(\Xi^n)$  and then substitute the expression for  $\Xi$  given in (6.3). This yields an element of  $C^{\text{coord},L} \widehat{\otimes} \Omega_F$  of degree  $2n$ . Thus  $j(\omega)$  is a sum of elements in  $C^{\text{coord},L} \widehat{\otimes} \Omega_F$  of even total degree.

By the discussion in §6 the element  $\text{Tr}(\Xi^n) \in C^{\text{coord},L} \widehat{\otimes} \Omega_F$  may be interpreted as an element in  $C^{\text{aff},\mathcal{L}} \widehat{\otimes}_{\mathcal{O}_1} \Omega_{J\mathcal{L}/\mathcal{O}_1}$  via the inclusions (6.2). Hence the same holds for  $j(\omega)$ . We keep the same notation for this reinterpreted version of  $j(\omega)$ .

We thus get the following formulæ:

$$(7.8) \quad \mathfrak{U}_{L,1} = \text{HKR} \circ \iota_{j(\omega)}, \quad \mathfrak{S}_{L,0} = j(\omega) \wedge \text{HKR}.$$

**7.3. Functoriality properties of the commutative diagram (7.5).** The computations in the proof of Proposition 5.5 imply the following theorem, expressing the functoriality properties of the commutative diagram (7.5).

**Theorem 7.3.** *For a general algebraic morphism  $(\ell, \lambda)$  from  $(L, R)$  to  $(M, S)$  as in Definition 5.4, which induces an isomorphism  $S \otimes_R L \cong M$  of  $S$ -modules, and such that  $L$  is free of rank  $d$  over  $R$  there exist quasi-isomorphisms  $\mathfrak{U}_{L,1}$ ,  $\mathfrak{U}_{M,1}$ ,*

$\mathfrak{S}_{L,0}$  and  $\mathfrak{S}_{M,0}$ , fitting into the commutative diagram of precalculi up to homotopy

$$(7.9) \quad \begin{array}{ccccccc} T_{\text{poly}}^L(R) & \hookrightarrow & \mathfrak{g}_1^L & \xrightarrow{\mathfrak{U}_{L,1}=\text{HKR} \circ \iota_{j(\omega_L)}} & \mathfrak{g}_2^L & \hookleftarrow & D_{\text{poly}}^L(R) \\ \varphi_T \downarrow & & \varphi_T \downarrow & & \varphi_D \downarrow & & \varphi_D \downarrow \\ T_{\text{poly}}^M(S) & \hookrightarrow & \mathfrak{g}_1^M & \xrightarrow{\mathfrak{U}_{M,1}=\text{HKR} \circ \iota_{j(\omega_M)}} & \mathfrak{g}_2^M & \hookleftarrow & D_{\text{poly}}^M(S) \\ \mathcal{L} \downarrow & & \mathcal{L}_1 \downarrow & & \mathcal{L}_2 \downarrow & & \mathcal{L} \downarrow \\ \Omega^M(S) & \hookrightarrow & \mathfrak{m}_1^M & \xleftarrow{\mathfrak{S}_{M,0}=j(\omega_M) \wedge \text{HKR}} & \mathfrak{m}_2^M & \hookleftarrow & C_{\text{poly}}^M(S) \\ \varphi_\Omega \uparrow & & \varphi_\Omega \uparrow & & \varphi_J \uparrow & & \varphi_J \uparrow \\ \Omega^L(R) & \hookrightarrow & \mathfrak{m}_1^L & \xleftarrow{\mathfrak{S}_{L,0}=j(\omega_L) \wedge \text{HKR}} & \mathfrak{m}_2^L & \hookleftarrow & C_{\text{poly}}^L(R) \end{array}$$

where we borrow notation from Proposition 5.5, and where  $\omega_L$  and  $\omega_M$ , denote the Maurer–Cartan form on  $C^{\text{coord},L}$  and  $C^{\text{coord},M}$  respectively. The precalculus structures up to homotopy on  $(\mathfrak{g}_i^*, \mathfrak{m}_i^*)$ ,  $*$  =  $L, M$ ,  $i$  =  $1, 2$ , are defined as in §5.5. Moreover the implied homotopies are in a similar way functorial for algebraic morphisms  $(\ell, \lambda)$  from  $(L, R)$  to  $(M, S)$  satisfying  $S \otimes_R L \cong M$ .

Almost all important objects appearing in Theorem 7.3 have already appeared in Theorem 5.6, hence the functoriality properties extend to the present situation. The commutativity of the upper and lower squares involving  $j(\omega)$  follows from the compatibility of the inclusions (6.2) with the base extension  $S/R$ . The functoriality properties of the implied homotopies are verified in the same way. See [7, Lemma 10.1.1] for results on  $\mathfrak{U}_{*,1}$  and related homotopies; in virtue of Lemma 7.2, the homotopy expressing the compatibility of  $\mathfrak{S}_{*,0}$  with cap products descends correctly on  $C^{\text{aff},*}$ , and the functoriality properties of such a homotopy follow along the same lines of the functoriality properties in Theorem 5.6, as the homotopy under consideration is expressed in terms of scalar combinations of poly-differential operators associated to certain graphs, as the  $L_\infty$ -quasi-isomorphisms of Kontsevich and Shoikhet.

**7.3.1. Arbitrary base fields.** We now briefly indicate how we may replace  $k$  by a general field of characteristic zero. Our arguments depend on the existence of a number of explicit homotopies. These homotopies are constructed as scalar linear combinations of poly-differential operators indexed by certain graphs, where the scalars depend only on the corresponding graphs. For the arguments to work the coefficients need to satisfy certain linear equations. These equations have a solution over  $\mathbb{R}$  (given that over this field we have homotopies that work). Thus they have a solution over any field of characteristic zero.

We will now be more specific. We refer to [7, §10.4] for what concerns Lie brackets and cup products; here we concentrate on the compatibility between cap products. We embed  $k$  in a field  $K$  containing  $\mathbb{R}$ . By virtue of [6, §6],  $\mathcal{U}_{\gamma,1}$  and  $\mathcal{S}_{\gamma,0}$  are defined over  $\mathbb{Q}$  and thus  $k$  (while they are *a priori* defined over  $\mathbb{R} \subset K$ ). Then observe that equation (7.1) is linear in the coefficients  $\overset{\circ}{W}_{D,\Gamma}$  of  $\mathcal{H}_\gamma^S$ . Since we already have a solution of these equations in  $\mathbb{R} \subset K$ , we get one in  $k$  by applying any projection  $K \rightarrow k$ .



**7.4. Proof of Theorem 1.1 in the global case.** Let  $(X, \mathcal{O})$  be a ringed site and  $\mathcal{L}$  be a locally free sheaf of Lie algebroids over  $\mathcal{O}$  of rank  $d$ . We denote by  $D(X)$  the derived category of sheaves of  $k$ -vector spaces over  $X$ . According to the results of Section 3, transported to the framework of sheaves of  $k$ -vector spaces,  $(T_{\text{poly}}^{\mathcal{L}}(X), \Omega^{\mathcal{L}}(X))$  and  $(D_{\text{poly}}^{\mathcal{L}}(X), C_{\text{poly}}^{\mathcal{L}}(X))$  are precalculi up to homotopy. Therefore, viewed as objects of  $D(X)$  they are genuine precalculi.

Additionally, the sheafification procedure can be applied to the commutative diagram (7.9), in virtue of the results of §7.3 (using the fact that the homotopies are functorial as well) : if we further consider the resulting commutative diagram of sheaves of  $k$ -vector spaces in the derived category  $D(X)$ , then using (6.4) we get the commutative diagram of precalculi

$$\begin{array}{ccccc}
 T_{\text{poly}}^{\mathcal{L}}(X) & \xrightarrow{\quad} & \mathfrak{g}_1^{\mathcal{L}} & & \\
 \downarrow \text{HKR} \circ \iota_{\tilde{\text{td}}(\mathcal{L})^{1/2}} & \searrow & \downarrow \text{HKR} \circ \iota_{j(\omega)} & \searrow & \\
 & \Omega^{\mathcal{L}}(X) & \xrightarrow{\quad} & \mathfrak{m}_1^{\mathcal{L}} & \\
 D_{\text{poly}}^{\mathcal{L}}(X) & \xrightarrow{\quad} & \mathfrak{g}_2^{\mathcal{L}} & \xrightarrow{\quad} & \mathfrak{m}_2^{\mathcal{L}} \\
 \downarrow & \swarrow & \downarrow & \swarrow & \downarrow \\
 & C_{\text{poly}}^{\mathcal{L}}(X) & \xrightarrow{\quad} & \mathfrak{m}_2^{\mathcal{L}} & \\
 & \uparrow \text{td}(\mathcal{L})^{1/2} \wedge \text{HKR} & & \uparrow j(\omega) \wedge \text{HKR} & \\
 & & & & 
 \end{array}$$

where all horizontal and vertical arrows represent isomorphisms in the derived category  $D(X)$ . Here  $\tilde{\text{td}}(\mathcal{L})$  is the modified Todd class of  $\mathcal{L}$  which is obtained by replacing the function  $q(x)$  in the definition of the Todd class (see (1.2)) by

$$\tilde{q}(x) = \frac{x}{e^{x/2} - e^{-x/2}}.$$

Hence at this point we have proved Theorem 1.1 provided that we replace the Todd class by the modified one. To obtain the result for the ordinary Todd class we follow the method of [7, §10.3]. We have

$$\begin{aligned}
 \tilde{\text{td}}(\mathcal{L}) &= \text{td}(\mathcal{L}) \det(e^{-A(\mathcal{L})/2}) \\
 &= \text{td}(\mathcal{L}) e^{-\text{Tr}(A(\mathcal{L}))/2} \\
 &= \text{td}(\mathcal{L}) e^{-a_1(\mathcal{L})/2}
 \end{aligned}$$

In other words it is sufficient to prove that  $(\iota_{e^{-a_1(\mathcal{L})/4}}, e^{a_1(\mathcal{L})/4} \wedge -)$  defines an automorphism of the precalculus  $(T_{\text{poly}}^{\mathcal{L}}(X), \Omega^{\mathcal{L}}(X))$ .

Via the inclusions (6.2) together with (6.4) we may as well prove that  $(\iota_{e^{-\text{Tr}(\Xi)/4}}, e^{\text{Tr}(\Xi)/4} \wedge -)$  defines an automorphism of the precalculus  $(C^{\text{coord}} \widehat{\otimes} T_{\text{poly}}(F), C^{\text{coord}} \widehat{\otimes} \Omega_F)$  or equivalently that  $(\iota_{\text{Tr}(\Xi)}, -\text{Tr}(\Xi) \wedge -)$  act as derivations. The fact that  $\iota_{\text{Tr}(\Xi)}$  is a derivation with respect to the cup product and Lie bracket has been checked in [7, §10.3]. So it remains to show compatibility with the cap product and Lie derivative.

As  $\text{Tr}(\Xi) = \sum_{i,\alpha} \eta_\alpha d_F(\partial_i \omega_\alpha^i)$  we first derive some identities for  $\iota_{d_F b}$  and  $d_F b \wedge -$  with  $b$  in  $F$ .

First we claim

$$(7.10) \quad d_F b \wedge (D \cap \sigma) = -\iota_{d_F b}(D) \cap \sigma + (-1)^{|D|+1} D \cap (d_F b \wedge \sigma)$$

for  $b \in F$ ,  $D \in T_{\text{poly}}(F)$ ,  $\sigma \in \Omega_F$ . If  $D = D_1 \cup D_2$  and (7.10) holds for  $D_1$ ,  $D_2$  then it holds for  $D$  as well. To see this note

$$\begin{aligned} d_F b \wedge ((D_1 \cup D_2) \cap \sigma) &= d_F b \wedge (D_1 \cap (D_2 \cap \sigma)) \\ &= -\iota_{d_F b}(D_1) \cap (D_2 \cap \sigma) + (-1)^{|D_1|+1} D_1 \cap (d_F b \wedge (D_2 \cap \sigma)) \\ &= -\iota_{d_F b}(D_1) \cap (D_2 \cap \sigma) - (-1)^{|D_1|+1} D_1 \cap \iota_{d_F b}(D_2) \cap \sigma \\ &\quad + (-1)^{|D_1|+|D_2|} D_1 \cap D_2 \cap (d_F b \wedge \sigma) \\ &= -\iota_{d_F b}(D_1 \cup D_2) \cap \sigma + (-1)^{|D_1 \cup D_2|+1} (D_1 \cup D_2) \cap (d_F b \wedge \sigma) \end{aligned}$$

So we only have to consider the case where  $D$  is a function or a vector field. The case that  $D$  is a function is trivial so assume that  $D$  is a vector field. In that case we find for the right-hand side of (7.10)

$$\begin{aligned} -\iota_{d_F b}(D) \cap \sigma + (-1)^{|D|+1} D \cap (d_F b \wedge \sigma) &= D b \cap \sigma - D b \wedge \sigma + d_F b \wedge (D \cap \sigma) \\ &= d_F b \wedge (D \cap \sigma) \end{aligned}$$

which is equal to the left-hand side of (7.10).

For the Lie derivative we use  $L_D = [d_F, D \cap -]$ . It is clear that  $d_F$  and  $d_F b \wedge -$  commute. We then compute using (7.10)

$$\begin{aligned} d_F b \wedge L_D \sigma &= d_F b \wedge (d_F(D \cap \sigma) - (-1)^{|D|+1} D \cap d_F \sigma) \\ &= -d_F(d_F b \wedge (D \cap \sigma)) + (-1)^{|D|} d_F b \wedge (D \cap d_F \sigma) \\ &= d_F(\iota_{d_F b}(D) \cap \sigma) + (-1)^{|D|} d_F(D \cap (d_F b \wedge \sigma)) \\ &\quad + (-1)^{|D|+1} \iota_{d_F b}(D) \cap d\sigma - D \cap (d_F b \wedge d_F \sigma) \\ &= L_{\iota_{d_F b} D}(\sigma) + (-1)^{|D|} L_D(d_F b \wedge \sigma) \end{aligned}$$

If  $\eta$  is an odd element in  $C^{\text{coord}}$  then  $\iota_{\eta d_F b} D = \eta \iota_{d_F b} D$  and  $L_{\iota_{\eta d_F b} D}(\sigma) = L_{\eta \iota_{d_F b} D}(\sigma) = -\eta L_{\iota_{d_F b} D}(\sigma)$ . Using this we find

$$\text{Tr}(\Xi) \wedge (D \cap \sigma) = -\iota_{\text{Tr}(\Xi)}(D) \cap \sigma + D \cap (\text{Tr}(\Xi) \wedge \sigma)$$

and

$$\text{Tr}(\Xi) \wedge L_D \sigma = -L_{\iota_{\text{Tr}(\Xi)} D}(\sigma) + L_D(\text{Tr}(\Xi) \wedge \sigma)$$

We conclude that  $(\iota_{\text{Tr}(\Xi)}, -\text{Tr}(\Xi) \wedge -)$  does indeed define a derivation of precalculus.

#### APPENDIX A. EXPLICIT FORMULÆ FOR THE $B_\infty$ -STRUCTURE ON POLY-DIFFERENTIAL OPERATORS

In this appendix and the next one we develop the precalculus structure on  $L$ -chains over  $L$ -cochains up to homotopy. The results in these appendices are provided for background and are not essential for the results in the body of the paper.

The graded vector space  $V = D_{\text{poly}}^L(R)$  is naturally a  $B_\infty$ -algebra. This means that the cofree coassociative coalgebra (with counit)  $T(V)$  is canonically equipped with the structure of a DG bialgebra. The notion of  $B_\infty$ -algebra has been introduced in [1]; though, we make use here mainly of the  $B_\infty$ -algebra structure given by braces [13, 14], to which we refer for more details, see also [6, Sections 1,2].

The corresponding associative product  $m$  on  $T(V)$  is uniquely determined by its Taylor components  $m_{p,q} : T^p(V) \otimes T^q(V) \rightarrow V$ . We have  $m_{p,q} = 0$  if  $p \neq 1$  and

$$\begin{aligned}
& \stackrel{(A.1)}{m_{1,q}}(D \otimes (D_1 \otimes \cdots \otimes D_q)) = D\{D_1, \dots, D_q\} = \\
& = \sum_{1 \leq i_1 \leq \cdots \leq i_q \leq |D| + \sum_{b=1}^{q-1} |D_b| + 1} (-1)^{\sum_{k=1}^q |D_k|(i_k - 1)} \\
& \left( 1^{\otimes(i_1-1)} \otimes \Delta^{|D_1|} \otimes 1^{\otimes(i_2-i_1-|D_1|-1)} \otimes \Delta^{|D_2|} \otimes \cdots \otimes 1^{\otimes(i_q-i_{q-1}-|D_{q-1}|-1)} \otimes \Delta^{|D_q|} \otimes 1^{\otimes(|D|+\sum_{b=1}^{q-1}|D_b|-i_q)} \right) (D) \\
& \left( 1^{\otimes(i_1-1)} \otimes D_1 \otimes 1^{\otimes(i_2-i_1-|D_1|-1)} \otimes D_2 \otimes \cdots \otimes 1^{\otimes(i_q-i_{q-1}-|D_{q-1}|-1)} \otimes D_q \otimes 1^{\otimes(|D|+\sum_{b=1}^{q-1}|D_b|-i_q)} \right),
\end{aligned}$$

for elements  $D, D_i, i = 1, \dots, q$ , of  $D_{\text{poly}}^L(R)$ , where  $|-|$  denotes the (shifted) degree of elements of  $D_{\text{poly}}^L(R)$ : accordingly, we have  $|D\{D_1, \dots, D_q\}| = |D| + \sum_{a=1}^q |D_a|$  and thus all brace operations are of degree zero. In the sum (A.1), we have  $1 \leq i_1, i_k + |D_k| + 1 \leq i_{k+1}, k = 1, \dots, q-1, i_q + |D_q| \leq |D| + \sum_{a=1}^q |D_a| + 1$ . The sign conventions are taken from [6]. The brace operations (A.1) satisfy an infinite family of quadratic identities (see e.g. [6]), which are equivalent to the associativity of the product  $m$ .

We define the cup product by means of the brace operations, see also [6, 13], *via* the assignment

$$(A.2) \quad D_1 \cup D_2 = (-1)^{|D_1|+1} \mu\{D_1, D_2\}, \quad D_i \in D_{\text{poly}}^L(R), \quad i = 1, 2.$$

It is obvious that the cup product has (shifted) degree 1. An easy verification using Formula (A.1) shows that the previous definition of cup product coincides with the one given in Formula (3.11).

We now have the following compatibilities

**Lemma A.1.** *The degree 0 operation (3.10) and the degree 1 operation (A.2) satisfy the following properties:*

$$(A.3)$$

$$[D_1, D_2] = -(-1)^{|D_1||D_2|} [D_2, D_1],$$

$$(A.4)$$

$$[D_1, [D_2, D_3]] = [[D_1, D_2], D_3] + (-1)^{|D_1||D_2|} [D_2, [D_1, D_3]],$$

$$(A.5)$$

$$D_1 \cup D_2 = (-1)^{(|D_1|-1)(|D_2|-1)} D_2 \cup D_1 \pm \left( d_{\text{H}}(D_1\{D_2\}) - (d_{\text{H}}D_1)\{D_2\} - (-1)^{|D_1|} D_1\{d_{\text{H}}D_2\} \right),$$

$$(A.6)$$

$$D_1 \cup (D_2 \cup D_3) = (D_1 \cup D_2) \cup D_3,$$

and

$$(A.7)$$

$$\begin{aligned}
[D_1, D_2 \cup D_3] &= [D_1, D_2] \cup D_3 + (-1)^{|D_1|(|D_2|-1)} D_2 \cup [D_1, D_3] + (-1)^{|D_1|} \left( d_{\text{H}}(D_1\{D_2, D_3\}) - \right. \\
&\quad \left. (d_{\text{H}}D_1)\{D_2, D_3\} - (-1)^{|D_1|} D_1\{d_{\text{H}}D_2, D_3\} - (-1)^{|D_1|+|D_2|} D_1\{D_2, d_{\text{H}}D_3\} \right),
\end{aligned}$$

for general elements  $D_i$  of  $D_{\text{poly}}^L(R)$ ,  $i = 1, 2, 3$ , and where  $d_{\text{H}} = [\mu, \bullet]$ ,  $\mu = 1 \otimes_R 1$ .

APPENDIX B. THE PRECALCULUS STRUCTURE ON  $L$ -CHAINS

We need results from [6, 25] about algebraic structures on Hochschild (co)chains, which have to be adapted to the Lie algebroid framework.

According to [6, 25], there are two distinct, non-compatible, left  $B_\infty$ -module structures on the Hochschild chain complex of  $A$ , viewed as a  $B_\infty$ -algebra with respect to the brace operations (A.1). Equivalently, we view the two left  $B_\infty$ -module structures on the Hochschild chain complex as the data of two left actions  $m_{L,i}$ ,  $i = 1, 2$ , on the left comodule cofreely cogenerated by the Hochschild chain complex of  $A$  over the coalgebra cofreely cogenerated by the Hochschild cochain complex of  $A$ .

These results can be applied to the present situation with due changes:  $\widehat{C}_{R,\bullet}(JL)$  has two left  $B_\infty$ -module structures over the  $B_\infty$ -algebra  $D_{\text{poly}}^L(R)$ .

We borrow the main notation and sign conventions from [6]. We denote by  $m_{L,i}$ ,  $i = 1, 2$  the two left  $B_\infty$ -module structures on  $\widehat{C}_{R,\bullet}(JL)$ : they are uniquely determined by their Taylor components

$$\begin{aligned}
\text{(B.1)} \quad & \left( m_{L,1}^{1;q,r}(P \otimes (Q_1 \otimes \cdots \otimes Q_q) \otimes a \otimes (R_1 \otimes \cdots \otimes R_r)) \right) (D) = \\
& = \sum_{\substack{l = -|a| - \sum_{b=1}^q |Q_b| - q + 1 \pmod{-|a|+1} \\ l \leq j_1 \leq \cdots \leq j_q \leq -|a| \\ 1 \leq k_1 \leq \cdots \leq k_r \leq |a| + |P| + l}} (-1)^{l(-|a|-l+1) + \sum_{b=1}^q |Q_b|(j_b-l) + \sum_{c=1}^r |R_c|(k_c-l-1)} \\
& \sigma^{-|a|-l+1}(a) \left( (\Delta^{(|P| + \sum_{b=1}^q |Q_b| + \sum_{c=1}^r |R_c|)} \otimes 1^{\otimes(-|a|-|P| - \sum_{b=1}^q |Q_b| - \sum_{c=1}^r |R_c|)})(D) \right. \\
& \left( 1^{(j_1-l)} \otimes \Delta^{|Q_1|} \otimes \cdots \otimes 1^{\otimes(j_q-j_{q-1}-|Q_{q-1}|-1)} \otimes \Delta^{|Q_q|} \otimes 1^{\otimes(-|a|-j_q-|Q_q|+k_1)} \otimes \Delta^{|R_1|} \otimes \cdots \otimes \right. \\
& \left. \otimes 1^{\otimes(k_r-k_{r-1}-|R_{r-1}|-1)} \otimes \Delta^{|R_r|} \otimes 1^{\otimes(|a|+|P| + \sum_{b=1}^q |Q_b| + \sum_{c=1}^{r-1} |R_c| + l - k_r - 1)} \otimes 1^{\otimes(-|a|-|P| - \sum_{b=1}^q |Q_b| - \sum_{c=1}^r |R_c|)} \right) (P) \\
& \left. \left( 1^{(j_1-l)} \otimes Q_1 \otimes \cdots \otimes 1^{\otimes(j_q-j_{q-1}-|Q_{q-1}|-1)} \otimes Q_q \otimes 1^{\otimes(-|a|-j_q-|Q_q|+k_1)} \otimes R_1 \otimes \cdots \otimes \right. \right. \\
& \left. \left. \otimes 1^{\otimes(k_r-k_{r-1}-|R_{r-1}|-1)} \otimes R_r \otimes 1^{\otimes(|a|+|P| + \sum_{b=1}^q |Q_b| + \sum_{c=1}^{r-1} |R_c| + l - k_r - 1)} \otimes 1^{\otimes(-|a|-|P| - \sum_{b=1}^q |Q_b| - \sum_{c=1}^r |R_c|)} \right) \right),
\end{aligned}$$

where  $\sigma$  is the operator on  $\widehat{C}_{R,\bullet}(JL)$  defined via

$$\sigma(a)(D_0 \otimes \cdots \otimes D_{-|a|}) = a(D_1 \otimes \cdots \otimes D_{-|a|} \otimes D_0), \quad D_i \in U_R(L), \quad i = 0, \dots, -|a|,$$

which obviously satisfies  $\sigma^{-|a|+1} = \text{id}$ , and the indices in the summation satisfy  $l \leq j_1$ ,  $j_i + |Q_i| + 1 \leq j_{i+1}$ ,  $i = 1, \dots, q-1$ ,  $j_q + |Q_q| \leq -|a|$ ,  $k_i + |R_i| + 1 \leq k_{i+1}$ ,  $i = 1, \dots, r-1$ ,  $k_r + |R_r| \leq |a| + |P| + \sum_{b=1}^q |Q_b| + \sum_{j=1}^r |R_j| + l - 1$ , and

$$\begin{aligned}
\text{(B.2)} \quad & \left( m_{L,2}^{0;0,r}(a \otimes (R_1 \otimes \cdots \otimes R_r)) \right) (D) = \\
& = \sum_{1 \leq i_1 \leq \cdots \leq i_p \leq -|a|} (-1)^{\sum_{c=1}^r |R_c|(i_c-1)} \\
& a \left( \left( 1^{\otimes i_1} \otimes \Delta^{|R_1|} \otimes 1^{\otimes(i_2-i_1-|R_1|-1)} \otimes \Delta^{|R_2|} \otimes \cdots \otimes 1^{\otimes(i_r-i_{r-1}-|R_{r-1}|-1)} \otimes \Delta^{|R_r|} \otimes 1^{\otimes(|D| + \sum_{c=1}^{r-1} |R_c| - i_r)} \right) (D) \right. \\
& \left. \left( 1^{\otimes i_1} \otimes R_1 \otimes 1^{\otimes(i_2-i_1-|R_1|-1)} \otimes R_2 \otimes \cdots \otimes 1^{\otimes(i_r-i_{r-1}-|R_{r-1}|-1)} \otimes R_r \otimes 1^{\otimes(|D| + \sum_{c=1}^{r-1} |R_c| - i_r)} \right) \right),
\end{aligned}$$

where the summation is over indices  $i_1, \dots, i_r$ , such that  $1 \leq i_1$ ,  $i_k + |D_k| + 1 \leq i_{k+1}$ ,  $k = 1, \dots, p-1$ ,  $i_p + |D_p| \leq -|a|$ . We observe that the components of  $m_L$ , resp.  $m_{L,2}$ , are non-trivial only if  $p \leq 1$ , with no restrictions on  $q, r$ , resp. only if  $q = r = 0$ , with no restrictions on  $p$ .

It is not difficult but quite tedious to verify that both (B.1) and (B.2) have degree 0 and satisfy an infinite family of quadratic relations involving braces.

The Taylor components of  $m_{L,i}$ ,  $i = 1, 2$ , permit to define a pairing of degree 0 between  $D_{\text{poly}}^L(R)$  and  $\widehat{C}_{R,\bullet}(JL)$  via

$$(B.3) \quad L_D a = m_{L,1}^{1,0,0}(D \otimes a) + (-1)^{|D|} m_{L,2}^{0,0,1}(a \otimes D), \quad D \in D_{\text{poly}}^L(R), \quad a \in \widehat{C}_{R,\bullet}(JL).$$

Similarly, we may consider two distinct pairings between  $D_{\text{poly}}^L(R)$  and  $\widehat{C}_{R,\bullet}(JL)$ : for  $\mu$  as above,

$$(B.4) \quad D \cap a = (-1)^{|D|} m_{L,1}^{1,1,0}(\mu \otimes D \otimes a),$$

$$(B.5) \quad a \cap D = (-1)^{|a|} m_{L,1}^{1,0,1}(\mu \otimes a \otimes D), \quad D \in D_{\text{poly}}^L(R), \quad a \in \widehat{C}_{R,\bullet}(JL).$$

It follows from their very definition that both (B.4) and (B.5) have degree 1.

**Lemma B.1.** *The pairing (B.3) of degree 0 and the pairings (B.4) and (B.5) of degree 1 satisfy the following properties:*

$$(B.6)$$

$$L_{[D_1, D_2]} a = L_{D_1}(L_{D_2} a) - (-1)^{|D_1||D_2|} L_{D_2}(L_{D_1} a),$$

$$(B.7)$$

$$D \cap a = (-1)^{(|D|-1)(|a|-1)} a \cap D \pm \left( b_{\mathbb{H}}(m_{L,1}^{1,0,0}(D \otimes a) - m_{L,1}^{1,0,0}(d_{\mathbb{H}} D \otimes a) - (-1)^{|D|} m_{L,1}^{1,0,0}(D \otimes b_{\mathbb{H}} a)) \right),$$

$$(B.8)$$

$$D_1 \cap (D_2 \cap a) = (D_1 \cup D_2) \cap a,$$

$$(B.9)$$

$$(a \cap D_1) \cap D_2 = a \cap (D_1 \cup D_2),$$

$$(B.10)$$

$$\begin{aligned} L_{D_1}(D_2 \cap a) &= [D_1, D_2] \cap a + (-1)^{|D_1|(|D_2|-1)} D_2 \cap L_{D_1} a + (-1)^{|D_1|} \left( b_{\mathbb{H}}(m_{L,1}^{1,1,0}(D_1 \otimes D_2 \otimes a)) \right. \\ &\quad \left. - m_{L,1}^{1,1,0}(d_{\mathbb{H}} D_1 \otimes D_2 \otimes a) - (-1)^{|D_1|} m_{L,1}^{1,1,0}(D_1 \otimes d_{\mathbb{H}} D_2 \otimes a) \right. \\ &\quad \left. - (-1)^{|D_1|+|D_2|} m_{L,1}^{1,1,0}(D_1 \otimes D_2 \otimes b_{\mathbb{H}} a) \right), \end{aligned}$$

$$(B.11)$$

$$\begin{aligned} L_{D_1}(a \cap D_2) &= L_{D_1} a \cap D_2 + (-1)^{|D_1|(|a|-1)} a \cap [D_1, D_2] + (-1)^{|D_1|} \left( b_{\mathbb{H}}(m_{L,1}^{1,0,1}(D_1 \otimes a \otimes D_2)) \right. \\ &\quad \left. - m_{L,1}^{1,0,1}(d_{\mathbb{H}} D_1 \otimes a \otimes D_2) - (-1)^{|D_1|} m_{L,1}^{1,0,1}(D_1 \otimes b_{\mathbb{H}} a \otimes D_2) \right. \\ &\quad \left. - (-1)^{|D_1|+|a|} m_{L,1}^{1,0,1}(D_1 \otimes a \otimes d_{\mathbb{H}} D_2) \right), \end{aligned}$$

and finally

(B.12)

$$\begin{aligned}
& \mathbb{L}_{D_1 \cup D_2} a + (-1)^{(|D_1|-1)(|D_2|-1)} \mathbb{L}_{D_2 \cup D_1} a \\
&= \left( D_1 \cap \mathbb{L}_{D_2} a + (-1)^{(|D_1|-1)(|D_2|+|a|-1)} \mathbb{L}_{D_2} a \cap D_1 \right) \\
&\quad + (-1)^{|a|(|D_2|-1)} \left( \mathbb{L}_{D_1} a \cap D_2 + (-1)^{(|D_1|+|a|-1)(|D_2|-1)} D_2 \cap \mathbb{L}_{D_1} a \right) \\
&\quad + (-1)^{(|D_2|-1)} \left( [D_1, D_2] \cap a + (-1)^{(|a|-1)(|D_1|+|D_2|-1)} a \cap [D_1, D_2] \right) \\
&\quad + (-1)^{|D_2|} \mathfrak{b}_H(\mathfrak{m}_{L,2}^{0,0,2}(a \otimes R_1 \otimes R_2)) - (-1)^{|D_1|} \mathfrak{m}_{L,2}^{0,0,2}(\mathfrak{b}_H a \otimes D_1 \otimes D_2) \\
&\quad + (-1)^{|D_2|} \mathfrak{m}_{L,2}^{0,0,2}(a \otimes \mathfrak{d}_H D_1 \otimes D_2) + (-1)^{|D_1|+|D_2|} \mathfrak{m}_{L,2}^{0,0,2}(a \otimes D_1 \otimes \mathfrak{d}_H D_2) \\
&\quad + (-1)^{|D_1|} \mathfrak{b}_H(\mathfrak{m}_{L,2}^{0,0,2}(a \otimes D_2 \otimes D_1)) - (-1)^{|D_2|} \mathfrak{m}_{L,2}^{0,0,2}(\mathfrak{b}_H a \otimes D_2 \otimes D_1) \\
&\quad + (-1)^{|D_1|} \mathfrak{m}_{L,2}^{0,0,2}(a \otimes \mathfrak{d}_H D_2 \otimes D_1) + (-1)^{|D_1|+|D_2|} \mathfrak{m}_{L,2}^{0,0,2}(a \otimes D_2 \otimes \mathfrak{d}_H D_1),
\end{aligned}$$

for a general element  $a$  of  $\widehat{C}_{R,\bullet}(JL)$  and general elements  $D, D_i, i = 1, 2$ , of  $D_{\text{poly}}^L(R)$ , and where  $\mathfrak{b}_H = \mathbb{L}_\mu$ , for  $\mu$  as before.

As for Lemma A.1, the proof essentially makes use of the brace identities, of the fact that  $\mathfrak{m}_{L,i}, i = 1, 2$ , is a left action with respect to the brace operations, and of the fact that  $\mathfrak{m}_{L,1}$  and  $\mathfrak{m}_{L,2}$  satisfy a weak compatibility, as explained in more details in [6].

Both actions  $\mathfrak{m}_{L,1}$  and  $\mathfrak{m}_{L,2}$  are compatible with the Grothendieck connection, i.e.

$$\begin{aligned}
{}^1\nabla_l \left( \mathfrak{m}_{L,1}^{1,q,r}(D \otimes Q_1 \otimes \cdots \otimes a \otimes R_1 \otimes \cdots) \right) &= \mathfrak{m}_{L,1}^{1,q,r}(D \otimes Q_1 \otimes \cdots \otimes {}^1\nabla_l a \otimes R_1 \otimes \cdots), \quad q, r \geq 0, \\
{}^1\nabla_l \left( \mathfrak{m}_{L,2}^{0,0,r}(D_1 \otimes \cdots \otimes a) \right) &= \mathfrak{m}_{L,2}^{p,0,0}(D_1 \otimes \cdots \otimes {}^1\nabla_l a), \quad p \geq 0,
\end{aligned}$$

for  $D, D_i (i = 1, \dots, p), Q_j (j = 1, \dots, q), R_k (k = 1, \dots, r)$  elements of  $D_{\text{poly}}^L(R)$ , and  $a$  of  $\widehat{C}_{R,\bullet}(JL)$ . Both identities follow from the fact that  ${}^1\nabla$  commutes with the operator  $\sigma$  and from the fact that  $U_R(L)$  is a Hopf algebroid, in particular, the comultiplication is an algebra morphism.

Then, in virtue of Lemma B.1, the pairings (B.3), (B.4) and (B.5) are compatible with the Grothendieck connection, implying in particular that the Hochschild differential is also compatible therewith. By the very same arguments, Formulæ (B.7), (B.8), (B.9), (B.10), (B.11) and (B.12) are compatible with the Grothendieck connection, whence  $(\text{Ker}({}^1\nabla) \cap \widehat{C}_{R,\bullet}(JL), \mathfrak{b}_H, \mathbb{L}, \cap)$ , where  $\cap$  denotes here both (B.4) and (B.5), inherits a structure of precalculus up to homotopy over the Gerstenhaber algebra  $(D_{\text{poly}}^L(R), \mathfrak{d}_H, [\ , \ ], \cup)$  up to homotopy.

For the sake of completeness, we write down explicit formulæ for the Hochschild differential  $\mathfrak{b}_H$  on the complex of Hochschild  $L$ -chains on  $R$  and for the pairing (B.5) between  $D_{\text{poly}}^L(R)$  and  $C_{\text{poly}}^L(R)$ ; in [8], we will deduce the same formulæ in the framework of homological algebra and derived functors. Explicitly,

$$\begin{aligned}
\mathfrak{b}_H(a) &= a \circ \mathfrak{d}_H, \\
a \cap D &= (-1)^{|a|} a(D \otimes_R \bullet), \quad a \in C_{\text{poly}}^L(R), \quad D \in D_{\text{poly}}^L(R).
\end{aligned}$$

We observe that (B.6) implies that  $\mathfrak{b}_H$ , the Hochschild differential on  $L$ -chains, is compatible with respect to (B.3), and that (B.10) and B.11, in the special case

$D_1 = \mu$ , imply that  $b_H$  satisfies Leibniz's rule with respect to (B.4) and (B.5) respectively.

Thus, combining these arguments with Proposition 3.6, we have the following important

**Theorem B.2.** *For a Lie algebroid  $L$  over the ring  $R$  as above, the twist of (B.3), (B.4), (B.5) and of the Hochschild differential  $b_H$  with respect to the isomorphism (3.15) endow  $C_{\text{poly}}^L(R)$  with a structure of precalculus up to homotopy over the Gerstenhaber algebra  $(D_{\text{poly}}^L(R), d_H, [ , ], \cup)$  up to homotopy.*

## REFERENCES

- [1] H. J. Baues, *The double bar and cobar constructions*, Compositio Math. **43** (1981), no. 3, 331–341. MR632433 (83f:55006)
- [2] Damien Calaque, *Formality for Lie algebroids*, Comm. Math. Phys. **257** (2005), no. 3, 563–578. MR2164943 (2007d:53151)
- [3] Damien Calaque, Vasily Dolgushev, and Gilles Halbout, *Formality theorems for Hochschild chains in the Lie algebroid setting*, J. Reine Angew. Math. **612** (2007), 81–127. MR2364075 (2008j:53153)
- [4] Damien Calaque and Carlo A. Rossi, *Lectures on Duflo isomorphisms in Lie algebra and complex geometry*, EMS Series of Lectures in Mathematics, European Mathematical Society (EMS), Zürich, 2011. MR2816610
- [5] ———, *Shoikhet's conjecture and Duflo isomorphism on (co)invariants*, SIGMA Symmetry Integrability Geom. Methods Appl. **4** (2008), Paper 060, 17. MR2434938 (2009g:16016)
- [6] ———, *Compatibility with cap-products in Tsygan's formality and homological Duflo isomorphism*, Lett. Math. Phys. **95** (2011), no. 2, 135–209. MR2764337
- [7] Damien Calaque and Michel Van den Bergh, *Hochschild cohomology and Atiyah classes*, Adv. Math. **224** (2010), no. 5, 1839–1889. MR2646112 (2011i:14037)
- [8] Damien Calaque, Carlo A. Rossi, and Michel Van den Bergh, *Hochschild (co)homology for Lie algebroids*, IMRN **2010** (2010), no. 21, 4098–4136. MR2738352 (2011m:14026)
- [9] Andrei Căldăraru, *The Mukai pairing. II. The Hochschild-Kostant-Rosenberg isomorphism*, Adv. Math. **194** (2005), no. 1, 34–66. MR2141853 (2006a:14029)
- [10] Vasily Dolgushev, *A formality theorem for Hochschild chains*, Adv. Math. **200** (2006), no. 1, 51–101. MR2199629 (2006m:16010)
- [11] Vasily Dolgushev, Dmitry Tamarkin, and Boris Tsygan, *The homotopy Gerstenhaber algebra of Hochschild cochains of a regular algebra is formal*, J. Noncommut. Geom. **1** (2007), no. 1, 1–25. MR2294189 (2008c:18007)
- [12] ———, *Formality of the homotopy calculus algebra of Hochschild (co)chains* (2008), available at [arXiv:0807.5117](https://arxiv.org/abs/0807.5117).
- [13] Murray Gerstenhaber and Alexander A. Voronov, *Higher-order operations on the Hochschild complex*, Funktsional. Anal. i Prilozhen. **29** (1995), no. 1, 1–6, 96 (Russian, with Russian summary); English transl., Funct. Anal. Appl. **29** (1995), no. 1, 1–5. MR1328534 (96g:18006)
- [14] Ezra Getzler and John Jones, *Operads, homotopy algebra and iterated integrals for double loop spaces*, available at [arXiv:hep-th/9403055](https://arxiv.org/abs/hep-th/9403055).
- [15] Daniel Huybrechts and Marc Nieper-Wißkirchen, *Remarks on derived equivalences of Ricci-flat manifolds*, Math. Zeit. **267** (2011), no. 3–4, 939–963. MR2776067
- [16] M. Kapranov, *Rozansky-Witten invariants via Atiyah classes*, Compositio Math. **115** (1999), no. 1, 71–113. MR1671737 (2000h:57056)
- [17] Maxim Kontsevich, *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. **66** (2003), no. 3, 157–216. MR2062626 (2005i:53122)
- [18] Dominique Manchon and Charles Torossian, *Cohomologie tangente et cup-produit pour la quantification de Kontsevich*, Ann. Math. Blaise Pascal **10** (2003), no. 1, 75–106 (French, with English summary). MR1990011 (2004f:53106)
- [19] Nikita Markarian, *The Atiyah class, Hochschild cohomology and the Riemann-Roch theorem*, J. Lond. Math. Soc. (2) **79** (2009), no. 1, 129–143. MR2472137 (2010d:14020)
- [20] Ajay C. Ramadoss, *The Mukai pairing and integral transforms in Hochschild homology*, Mosc. Math. J. **10** (2010), no. 3, 629–645. MR2732577

- [21] Boris Shoikhet, *A proof of the Tsygan formality conjecture for chains*, Adv. Math. **179** (2003), no. 1, 7–37. MR2004726 (2004m:53161)
- [22] ———, *Tsygan formality and Duflo formula*, Math. Res. Lett. **10** (2003), no. 5-6, 763–775. MR2024732 (2004m:53162)
- [23] Richard G. Swan, *Hochschild cohomology of quasiprojective schemes*, J. Pure Appl. Algebra **110** (1996), no. 1, 57–80. MR1390671 (97j:19003)
- [24] Dmitry Tamarkin, *Another proof of M. Kontsevich formality theorem for  $\mathbb{R}^n$*  (1998), available at [arXiv:math.QA/980302](https://arxiv.org/abs/math/980302).
- [25] Dmitry Tamarkin and Boris Tsygan, *Cyclic formality and index theorems*, Lett. Math. Phys. **56** (2001), no. 2, 85–97. EuroConférence Moshé Flato 2000, Part II (Dijon). MR1854129 (2003e:19008)
- [26] Boris Tsygan, *Formality conjectures for chains*, Differential topology, infinite-dimensional Lie algebras, and applications, Amer. Math. Soc. Transl. Ser. 2, vol. 194, Amer. Math. Soc., Providence, RI, 1999, pp. 261–274. MR1729368 (2001g:53161)
- [27] Michel Van den Bergh, *On global deformation quantization in the algebraic case*, J. Algebra **315** (2007), no. 1, 326–395. MR2344349 (2008i:14018)
- [28] Ping Xu, *Quantum groupoids*, Comm. Math. Phys. **216** (2001), no. 3, 539–581. MR1815717 (2002f:17033)
- [29] Amnon Yekutieli, *Deformation quantization in algebraic geometry*, Adv. Math. **198** (2005), no. 1, 383–432. MR2183259 (2006j:53131)
- [30] ———, *Private communication*.
- [31] ———, *Continuous and twisted  $L_\infty$  morphisms*, J. Pure Appl. Algebra **207** (2006), no. 3, 575–606. MR2265541 (2008i:53131)

(Damien Calaque) DEPARTMENT OF MATHEMATICS, ETH ZÜRICH, 8092 ZÜRICH, SWITZERLAND,  
 ON LEAVE FROM INSTITUT CAMILLE JORDAN, CNRS & UNIVERSITÉ LYON 1, FRANCE  
*E-mail address:* [damien.calaque@math.ethz.ch](mailto:damien.calaque@math.ethz.ch), [calaque@math.univ-lyon1.fr](mailto:calaque@math.univ-lyon1.fr)

(Carlo A. Rossi) MPIM BONN, VIVATSGASSE 7, 53111 BONN, GERMANY  
*E-mail address:* [rossi@mpim-bonn.mpg.de](mailto:rossi@mpim-bonn.mpg.de)

(Michel Van den Bergh) DEPARTEMENT WNI, HASSELT UNIVERSITY, AGORALAAN, 3590 DIEPEN-  
 BEEK, BELGIUM  
*E-mail address:* [michel.vandenbergh@uhasselt.be](mailto:michel.vandenbergh@uhasselt.be)