

UPPER BOUNDS FOR THE NUMBER OF SOLUTIONS TO QUARTIC THUE EQUATIONS

SHABNAM AKHTARI

ABSTRACT. We will give upper bounds for the number of integral solutions to quartic Thue equations. Our main tool here is a logarithmic curve $\phi(x, y)$ that allows us to use the theory of linear forms in logarithms. This manuscript improves the results of author's earlier work with Okazaki [2] by giving special treatments to forms with respect to their signature.

1. INTRODUCTION

In this paper, we will study binary quartic forms with integer coefficients; i.e. polynomials of the shape

$$F(x, y) = a_0x^4 + a_1x^3y + a_2x^2y^2 + a_3xy^3 + a_4y^4,$$

with $a_i \in \mathbb{Z}$, $i \in \{0, 1, 2, 3, 4\}$. We aim to give upper bounds for the number of solutions to the equation

$$(1) \quad |F(x, y)| = 1.$$

Here we will count (x, y) and $(-x, -y)$ as one solution. Let $M(F)$ be the Mahler measure of $F(x, y)$. In [2] we used some ideas of Stewart [16] to bound the number of solutions with $|y| < M(F)^6$. We will slightly modify those ideas and use them to give an upper bound for the solutions of (1) with $|y| < M(F)^{3.5}$. Then we will improve the main result in [2] by giving better upper bounds for the number of solutions (x, y) with large $|y| \geq M(F)^{3.5}$ to equation (1). The following is the main result of this manuscript.

Theorem 1.1. *Let $F(x, y)$ be an irreducible quartic binary form with integer coefficients. The Diophantine equation (1) has at most U_F (see the table below) solutions in integers x and y , provided that the discriminant of F is greater than D_0 , where D_0 is an explicitly computable constant.*

Signature of F	U_F
(0, 2)	6
(2, 1)	14
(4, 0)	26

The reason for having different upper bounds for forms with different signature in Theorem 1.1, relies upon the fact that the number fields generated over \mathbb{Q} by a root of the equation $F(x, 1) = 0$ have a rings integers with different numbers of fundamental units.

2010 *Mathematics Subject Classification.* 11J86, 11D45.

Key words and phrases. Thue Equations, Linear Forms in Logarithms.

One can use the method of this manuscript to deal with particular quartic Thue equations, where more information about the coefficients of the quartic form are available. Therefore, in applications, the strong condition on the size of discriminant may be removed.

The equation

$$x^4 - 4x^3y - x^2y^2 + 4xy^3 + y^4 = 1$$

has exactly 8 solutions $(x, y) = (0, 1), (1, 0), (1, 1), (-1, 1), (4, 1), (-1, 4), (8, 7), (-7, 8)$ (see [11] for a proof). The author is not aware of any binary quartic forms $F(x, y)$ for which the equation $F(x, y) = 1$ has more than 8 solutions. Magma [7] did not find any solution to

$$x^4 - 4x^3y - x^2y^2 + 4xy^3 + y^4 = -1.$$

We will always assume that $F(x, y)$ is irreducible. In fact, when the form $F(x, y)$ is not irreducible over $\mathbb{Z}[x, y]$, we are in a much simpler situation. In general, equation (1) may have infinitely many integral solutions; $F(x, y)$ could, for instance, be a power of a linear or indefinite binary quadratic form that represents unity. If $F(x, y)$ is not an irreducible form, however, we may very easily derive a stronger version of our main theorem under the assumption that $F(x, y)$ is neither a 4th power of a linear form nor a second power of a quadratic form. Suppose that $F(x, y)$ is reducible and can be factored over $\mathbb{Z}[x, y]$ as follows

$$F(x, y) = F_1(x, y)F_2(x, y),$$

with $\deg(F_1) \leq \deg(F_2)$ and F_1 irreducible over $\mathbb{Z}[x, y]$. Therefore, the following equations must be satisfied:

$$(2) \quad F_1(x, y) = \pm 1$$

and

$$(3) \quad F_2(x, y) = \pm 1.$$

This means the number of solutions to (1) is no more than the minimum of number of solutions to (2) and (3). First suppose that F_1 is a linear form. Then the equation (3) can be written as a polynomial of degree 3 in x and therefore there are no more than 12 complex solutions to above equations. Now let us suppose that F_1 is a quadratic form. Using Bézout's theorem from classical algebraic geometry concerning the number of common points of two plane algebraic curves, we conclude that (1) has at most 16 integral solutions.

In this manuscript, we give new and sharper bounds for the number of solutions to equation (1). The bound given in [2] is improved here mostly due to some adjustment in the definition of the logarithmic curve $\phi(x, y)$ in Section 6. We also study the geometry of binary forms with respect to their signature to get amore precise understanding of the distribution of solutions to (1). We also appeal to a result of Voutier (Proposition 2.2) to estimate the height of algebraic numbers in the number field generated over \mathbb{Q} by a root of the equation $F(x, 1) = 0$. These allow us to extend our method introduced in [2] for quartic forms that split in \mathbb{R} to all quartic forms.

2. PRELIMINARIES

Let $f(x) = a_n x^n + \dots + a_1 x + a_0$ be the minimal polynomial of an algebraic number $\alpha \neq 0$. Suppose that over \mathbb{C} ,

$$f(x) = a_n(x - \alpha_1) \dots (x - \alpha_n).$$

We put

$$M(\alpha) = M(f) = |a_n| \prod_{i=1}^n \max(1, |\alpha_i|).$$

$M(\alpha)$ is known as the Mahler measure of α .

The Mahler measure of a binary form $G(x, y)$ is defined to be equal to the Mahler measure of the polynomial $G(x, 1)$. In [8], Mahler showed, for polynomial G of degree n and discriminant D_G , that

$$(4) \quad M(G) \geq \left(\frac{D_G}{n^n} \right)^{\frac{1}{2n-2}}.$$

Let $F(x, y)$ be a quartic form that factors over \mathbb{C} as follows

$$F(x, y) = a_0(x - \alpha_1 y)(x - \alpha_2 y)(x - \alpha_3 y)(x - \alpha_4 y).$$

The discriminant D of F is given by

$$D = D_F = a_0^6(\alpha_1 - \alpha_2)^2(\alpha_1 - \alpha_3)^2(\alpha_1 - \alpha_4)^2(\alpha_2 - \alpha_3)^2(\alpha_2 - \alpha_4)^2(\alpha_3 - \alpha_4)^2.$$

We call forms F and G equivalent if they are equivalent under $GL_2(\mathbb{Z})$ -action; i.e. if there exist integers a_1, a_2, a_3 and a_4 such that

$$F(a_1 x + a_2 y, a_3 x + a_4 y) = G(x, y)$$

for all x, y , where $a_1 a_4 - a_2 a_3 = \pm 1$. We denote by N_F the number of solutions in integers x and y of the Diophantine equation (1). If F and G are equivalent then $N_F = N_G$ and $D_F = D_G$.

Suppose there is a solution (x_0, y_0) to equation (1). Since

$$\gcd(x_0, y_0) = 1,$$

there exist integers $x_1, y_1 \in \mathbb{Z}$ with

$$x_0 y_1 - x_1 y_0 = 1.$$

Then

$$F^*(1, 0) = 1,$$

where

$$F^*(x, y) = F(x_0 x + x_1 y, y_0 x + y_1 y).$$

Therefore, F^* is a monic form equivalent to F . From now on we will assume F is monic.

Let $\mathbb{Q}(\alpha_1)^\sigma$ be the embeddings of the real number field $\mathbb{Q}(\alpha_1)$ in \mathbb{R} , $1 \leq \sigma \leq n$, where $\{\alpha_1, \alpha_2, \dots, \alpha_n\}$ are roots of $F(x, 1) = 0$. We respectively have n Archimedean valuations of $\mathbb{Q}(\alpha_1)$:

$$|\rho|_\sigma = \left| \rho^{(\sigma)} \right|, \quad 1 \leq \sigma \leq n.$$

We enumerate simple ideals of $\mathbb{Q}(\alpha)$ by indices $\sigma > n$ and define non-Archimedean valuation of $\mathbb{Q}(\alpha)$ by the formulas

$$|\rho|_\sigma = (\text{Norm } \mathfrak{p})^{-k},$$

where

$$k = \text{ord}_{\mathfrak{p}}(\alpha), \quad \mathfrak{p} = \mathfrak{p}_{\sigma}, \quad \sigma > n,$$

for any $\rho \in \mathbb{Q}^*(\alpha)$. Then we have the *product formula* :

$$\prod_1^{\infty} |\rho|_{\sigma} = 1, \quad \rho \in \mathbb{Q}(\alpha).$$

Note that $|\rho|_{\sigma} \neq 1$ for only finitely many ρ . We should also remark that if $\sigma_2 = \bar{\sigma}_1$, i.e.,

$$\sigma_2(x) = \bar{\sigma}_1(x) \quad \text{for} \quad x \in \mathbb{Q}(\alpha),$$

then the valuations $|\cdot|_{\sigma_1}$ and $|\cdot|_{\sigma_2}$ are equal. We define the *absolute logarithmic height* of ρ as

$$h(\rho) = \frac{1}{2n} \sum_{\sigma=1}^{\infty} |\log |\rho|_{\sigma}|.$$

Proposition 2.1. *For every non-zero algebraic number α , we have $h(\alpha^{-1}) = h(\alpha)$. For algebraic numbers $\alpha_1, \dots, \alpha_n$, we have*

$$h(\alpha_1 \dots \alpha_n) \leq h(\alpha_1) + \dots + h(\alpha_n)$$

and

$$h(\alpha_1 + \dots + \alpha_n) \leq \log n + h(\alpha_1) + \dots + h(\alpha_n).$$

Proof. See [4] for proof. □

Proposition 2.2. (Voutier [17]) *Suppose α is a non-zero algebraic number of degree n which is not a root of unity. If $n \geq 2$ then*

$$h(\alpha) = \frac{1}{n} \log M(\alpha) > \frac{1}{4} \left(\frac{\log \log n}{\log n} \right)^3.$$

Let α and β be two algebraic numbers. Then the following inequalities hold (see [4]):

$$(5) \quad h(\alpha + \beta) \leq \log 2 + h(\alpha) + h(\beta)$$

and

$$(6) \quad h(\alpha\beta) \leq h(\alpha) + h(\beta).$$

Lemma 2.3. (Mahler [8]) *If a and b are distinct zeros of polynomial $P(x)$ with degree n , then we have*

$$|a - b| \geq \sqrt{3}(n)^{-(n+2)/2} M(P)^{-n+1},$$

where $M(P)$ is the Mahler measure of P .

Lemma 2.4. *Let $f(x) = a_n x^n + \dots + a_1 x + a_0$ be an irreducible polynomial of degree n and α_m be one of its roots. For $f'(x)$ the derivative of f , we have*

$$2^{-(n-1)^2} \frac{|D_f|}{M(f)^{2n-2}} \leq |f'(\alpha_m)| \leq \frac{n(n+1)}{2} H(f) (\max(1, |\alpha_m|))^{n-1},$$

where D_f is the discriminant, $M(f)$ is the Mahler measure and $H(f)$ is the naive height of f .

Proof. The right hand side inequality is trivial by noticing that

$$f'(x) = na_n x^{n-1} + \dots + a_1 x.$$

To see the left hand side inequality, observe that for α_i, α_j , two distinct roots of $f(x)$, we have

$$|\alpha_i - \alpha_j| \leq 2 \max(1, |\alpha_i|) \max(1, |\alpha_j|).$$

Then

$$\begin{aligned} |f'(\alpha_m)| &= \prod_{i=1, i \neq m}^n |\alpha_i - \alpha_m| \geq \prod_{i=1, i \neq m}^n \frac{|\alpha_i - \alpha_m|}{\max(1, |\alpha_i|) \max(1, |\alpha_m|)} \\ &\geq 2^{n-1-n(n-1)} \prod_{j=1}^n \prod_{i=1, i \neq j}^n \frac{|\alpha_i - \alpha_j|}{\max(1, |\alpha_i|) \max(1, |\alpha_j|)} \\ &= 2^{-(n-1)^2} \frac{|D_F|}{M(F)^{2n-2}}. \end{aligned}$$

□

Suppose that \mathbb{K} is an algebraic number field of degree d over \mathbb{Q} embedded in \mathbb{C} . If $\mathbb{K} \subset \mathbb{R}$, we put $\chi = 1$, and otherwise $\chi = 2$. We are given numbers $\alpha_1, \dots, \alpha_n \in \mathbb{K}^*$ with absolute logarithm heights $h(\alpha_j)$, $1 \leq j \leq n$. Let $\log \alpha_1, \dots, \log \alpha_n$ be arbitrary fixed non-zero values of the logarithms. Suppose that

$$A_j \geq \max\{dh(\alpha_j), |\log \alpha_j|\}, \quad 1 \leq j \leq n.$$

Now consider the linear form

$$L = b_1 \log \alpha_1 + \dots + b_n \log \alpha_n,$$

with $b_1, \dots, b_n \in \mathbb{Z}$ and with the parameter

$$B = \max\{1, \max\{b_j A_j / A_n : 1 \leq j \leq n\}\}.$$

For brevity we put

$$\Omega = A_1 \dots A_n,$$

$$C(n) = C(n, \chi) = \frac{16}{n! \chi} e^n (2n+1+2\chi)(n+2)(4n+4)^{n+1} \left(\frac{1}{2} en\right)^\chi,$$

$$C_0 = \log(e^{4.4n+7} n^{5.5} d^2 \log(en)),$$

$$W_0 = \log(1.5eBd \log(ed)).$$

The following is the main result of [10].

Proposition 2.5 (Matveev). *If $\log \alpha_1, \dots, \log \alpha_n$ are linearly independent over \mathbb{Z} and $b_n \neq 0$, then*

$$\log |L| > -C(n)C_0W_0d^2\Omega.$$

3. SUMMARY OF THE PROOF

Suppose that (x, y) is an integral solution to equation (1). Then we have

$$(x - \alpha_1 y)(x - \alpha_2 y)(x - \alpha_3 y)(x - \alpha_4 y) = \pm 1.$$

Therefore, for some $1 \leq i \leq 4$,

$$|x - \alpha_i y| \leq 1.$$

Definition. We say the pair of solution (x, y) is related to α_i if

$$|x - \alpha_i y| = \min_{1 \leq j \leq 4} |x - \alpha_j y|.$$

Proposition 3.1. *Let $F(x, y)$ be an irreducible monic binary quartic form with integer coefficients and Mahler measure $M(F)$. The Diophantine equation (1) has at most N_1 (see the table below) solutions in integers x and y with $0 < y < M(F)^{3.5}$, provided that the discriminant of F is greater than D_0 , where D_0 is an explicitly computable constant.*

Signature of F	N_1
(0, 2)	5
(2, 1)	9
(4, 0)	12

Since $F(x, y)$ is monic, $(1, 0)$ is a trivial solution to $F(x, y) = 1$. We will need to define a subset of solutions to (1), called \mathfrak{A} (see Section 8 for details). This set contains the trivial solution $(1, 0)$ and 5 other pairs of solution only when $F(x, 1) = 0$ has 4 real roots.

Signature of F	$ \mathfrak{A} $
(0, 2)	1
(2, 1)	1
(4, 0)	6

Proposition 3.2. *Let $F(x, y)$ be an irreducible binary quartic form with integer coefficients and Mahler measure $M(F)$. The Diophantine equation (1) has at most N_2 (see the table below) solutions in integers x and y with $y \geq M(F)^{3.5}$, provided that the discriminant of F is greater than D_0 , where D_0 is an explicitly computable constant.*

Signature of F	N_2
(0, 2)	0
(2, 1)	4
(4, 0)	8

4. SOLUTIONS WITH SMALL y ; THE PROOF OF PROPOSITION 3.1

We may suppose that $F(x, y)$ is a monic form and has the smallest Mahler measure among all monic forms that are equivalent to F . Assume that $F(x, 1) = 0$ has r real roots and $2s$ non-real roots ($r + 2s = 4$).

Let Y_0 be a positive real number. Following Stewart [16] and Bombieri and Schmidt [3], we will estimate the solutions (x, y) to (1) for which $0 < y \leq Y_0$. For binary form

$$F(x, y) = (x - \alpha_1 y) \dots (x - \alpha_n y)$$

put

$$L_i(x, y) = x - \alpha_i y$$

for $i = 1, \dots, n$. Then

Lemma 4.1. *Suppose F is a monic binary form. Then for every solutions (x, y) of (1) we have*

$$\frac{1}{L_i(x, y)} - \frac{1}{L_j(x, y)} = (\beta_j - \beta_i)y,$$

where β_1, \dots, β_n are such that the form

$$J(u, w) = (u - \beta_1 w) \dots (u - \beta_n w)$$

is equivalent to F .

Proof. This is Lemma 4 of [16] and Lemma 3 of [3], by taking (x_0, y_0) equal to $(1, 0)$. \square

For every solution $(x, y) \neq (1, 0)$ of (1), fix $j = j(x, y)$ with

$$|L_j(x, y)| \geq 1.$$

Then, by Lemma 4.1,

$$(7) \quad \frac{1}{|L_i(x, y)|} \geq |\beta_j - \beta_i||y| - 1.$$

For complex conjugate $\bar{\beta}_j$ of β_j , where $j = j(x, y)$, we also have

$$\frac{1}{|L_i(x, y)|} \geq |\bar{\beta}_j - \beta_i||y| - 1.$$

Hence

$$\frac{1}{|L_i(x, y)|} \geq |\operatorname{Re}(\beta_j) - \beta_i||y| - 1,$$

where $\operatorname{Re}(\beta_j)$ is the real part of β_j . We now choose an integer $m = m(x, y)$ with $|\operatorname{Re}(\beta_j) - \beta_j| \leq 1/2$, and we obtain

$$(8) \quad \frac{1}{|L_i(x, y)|} \geq \left(|m - \beta_i| - \frac{1}{2} \right) |y| - 1,$$

for $i = 1, \dots, n$.

Definition. For $1 \leq i \leq n$, Let \mathfrak{X}_i be the set of solutions to (1) with $1 \leq y \leq Y_0$ and $|L_i(x, y)| \leq \frac{1}{2y}$.

Remark When α_k and α_l are complex conjugates, $\mathfrak{X}_l = \mathfrak{X}_k$ and therefore we only need to consider $r + s$ different sets \mathfrak{X}_i .

Lemma 4.2. *Suppose (x_1, y_1) and (x_2, y_2) are two distinct solutions in \mathfrak{X}_i with $y_1 \leq y_2$. Then*

$$\frac{y_2}{y_1} \geq \frac{2}{7} \max(1, |\beta_i(x_1, y_1) - m(x_1, y_1)|).$$

Proof. This is Lemma 5 of [16] and Lemma 4 of [3]. \square

Lemma 4.3. *Suppose (x, y) is a solution to (1) with $y > 0$ and $|L_i(x, y)| > \frac{1}{2y}$. Then*

$$|m(x, y) - \beta_i(x, y)| \leq \frac{7}{2}.$$

Proof. This is Lemma 6 of [16]. \square

By Lemma 4.1 the form

$$J(u, w) = (u - \beta_1 w) \dots (u - \beta_n w)$$

is equivalent to $F(x, y)$ and therefore the form

$$\hat{J}(u, w) = (u - (\beta_1 - m)w) \dots (u - (\beta_n - m)w)$$

is also equivalent to $F(x, y)$. Therefore, since we assumed that F has the smallest Mahler measure among its equivalent forms, we get

$$(9) \quad \prod_{i=1}^n \max(1, |\beta_i(x, y) - m(x, y)|) \geq M(F).$$

For each set \mathfrak{X}_i that is not empty, let $(x^{(i)}, y^{(i)})$ be the element with the largest value of y . Let \mathfrak{X} be the set of solutions of (1) with $1 \leq y \leq Y_0$ minus the elements $(x^{(1)}, y^{(1)}), \dots, (x^{(r+s)}, y^{(r+s)})$. Suppose that, for integer i , the set \mathfrak{X}_i is non-empty. Index the elements of \mathfrak{X}_i as

$$(x_1^{(i)}, y_1^{(i)}), \dots, (x_v^{(i)}, y_v^{(i)}),$$

so that $y_1^{(i)} \leq \dots \leq y_v^{(i)}$ (note that $(x_v^{(i)}, y_v^{(i)}) = (x^{(i)}, y^{(i)})$). By Lemma 4.2

$$\frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq \frac{y_{k+1}^{(i)}}{y_k^{(i)}}$$

for $k = 1, \dots, v-1$. Hence

$$\prod_{(x, y) \in \mathfrak{X} \cap \mathfrak{X}_i} \frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq Y_0.$$

For (x, y) in \mathfrak{X} but not in \mathfrak{X}_i we have

$$\frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq 1.$$

By Lemma 4.3. Thus

$$\prod_{(x, y) \in \mathfrak{X}} \frac{2}{7} \max\left(1, \left|\beta_i(x_k^{(i)}, y_k^{(i)})\right|\right) \leq Y_0.$$

Comparing this with (9) and since we have at most $r+s$ different sets \mathfrak{X}_i , we obtain

$$(10) \quad \left(\left(\frac{2}{7}\right)^4 M(F)\right)^{|\mathfrak{X}|} \leq Y_0^{r+s}.$$

If D_F satisfies the following numerical inequality

$$D_F \geq 4^4 \left(\frac{7}{2}\right)^{4 \times 6 \times 65}$$

then we have

$$\left(\frac{D_F}{4^4}\right)^{\frac{1}{6 \times 65}} \geq \left(\frac{7}{2}\right)^4.$$

From here and the fact that $M(F) \geq \left(\frac{D_F}{4^4}\right)^{\frac{1}{6}}$ (see (4)), we conclude that if the discriminant is large enough then $M(F)$ will be large enough to satisfy

$$\left(\frac{2}{7}\right)^4 M(F) \geq M(F)^{64/65}.$$

By (10),

$$(11) \quad |\mathfrak{X}| < (r+s) \frac{65 \log Y_0}{64 \log M(F)}.$$

When $F(x, y)$ has signature $(4, 0)$, choose $\theta_1 > 0$ such that

$$\frac{65}{16} \left(\frac{11}{6} + \theta_1\right) < 8.$$

From (10), we conclude that in this case $|\mathfrak{X}|$ is at most 7 and therefore (1) has at most 11 solutions with $1 \leq y < M(f)^{\frac{11}{6} + \theta_1}$.

When $F(x, y)$ has signature $(2, 1)$ choose $\theta_2 > 0$ such that

$$3 \times \frac{65}{64} \left(\frac{11}{6} + \theta_2\right) < 6.$$

From (10), we conclude that in this case $|\mathfrak{X}|$ is at most 5 and therefore (1) has at most 8 solutions with $1 \leq y < M(f)^{\frac{11}{6} + \theta_2}$.

We can repeat the similar argument for forms with signature $(0, 2)$ and choose $\theta_3 > 0$ such that

$$2 \times \frac{65}{64} \left(\frac{11}{6} + \theta_3\right) < 4.$$

This will give us at most 5 solutions with $1 \leq y < M(f)^{\frac{11}{6} + \theta_3}$. But for this case, we have more to say in the next section.

Lemma 4.4. *Let F be a binary form of degree $n \geq 3$ with integer coefficients and nonzero discriminant D . For every pair of integers (x, y) with $y \neq 0$*

$$\min_{\alpha} \left| \alpha - \frac{x}{y} \right| \leq \frac{2^{n-1} n^{n-1/2} (M(F))^{n-2} |F(x, y)|}{|D(F)|^{1/2} |y|^n},$$

where the minimum is taken over the zeros α of $F(z, 1)$.

Proof. This is Lemma 3 of [16]. □

Lemma 4.5. *Let $F(x, y)$ be a quartic binary form with discriminant D , where $|D| \geq D_0$. Suppose that α_i is a root of $F(z, 1) = 0$. Suppose that $\theta > 0$. Then related to α_i , there is at most 1 solution for equation (1) in integers x and y with $\left(\frac{11}{6} + \theta\right) < y < M(F)^{3.5}$.*

Proof. Assume that (x_1, y_1) and (x_2, y_2) are two distinct solutions to (1), both related to α_i with $y_2 > y_1 > M(F)^2$. By Lemma 4.4, we have

$$\begin{aligned} \left| \frac{x_2}{y_2} - \frac{x_1}{y_1} \right| &\leq \left| \alpha_i - \frac{x_1}{y_1} \right| + \left| \alpha_i - \frac{x_2}{y_2} \right| \\ &\leq \frac{2^3 4^{7/2} (M(F))^2}{|D(F)|^{1/2} |y_1|^4} + \frac{2^3 4^{7/2} (M(F))^2}{|D(F)|^{1/2} |y_2|^4} \\ &\leq \frac{2^4 4^{7/2} (M(F))^2}{|D(F)|^{1/2} |y_1|^4}. \end{aligned}$$

Since (x_1, y_1) , (x_2, y_2) are distinct, we have $|x_2 y_1 - x_1 y_2| \geq 1$. Therefore,

$$\left| \frac{1}{y_1 y_2} \right| \leq \left| \frac{x_2}{y_2} - \frac{x_1}{y_1} \right| \leq \frac{M(F)^2}{|y_1|^4}.$$

This is because we assumed that D_F is large. Thus,

$$(12) \quad \frac{y_1^3}{M(F)^2} \leq y_2.$$

Following Stewart [16], we define δ_j , for $j = 1, 2$, by

$$y_j = M(F)^{1+\delta_j}.$$

By (4) the Mahler measure of F is large and (12) implies that

$$3\delta_1 \leq \delta_2.$$

From here, we conclude that

$$y_2 > M(F)^{3.5}.$$

In other words, related to each root α_i , there exists at most 1 solution in x and y with $M(F)^{\frac{11}{6}+\theta_1} < y < M(F)^{3.5}$. \square

5. PROOF OF THE MAIN THEOREM FOR FORMS WITH SIGNATURE (0, 2)

We will first show that if a pair of integer (x, y) satisfies $F(x, y) = \pm 1$ and is related to a non-real root α of $F(x, 1) = 0$ then

$$|y| < M(F)^{9/4}.$$

Lemma 5.1. *For quartic binary form $F(x, y)$, let α be a non-real root of $F(x, 1) = 0$. If a pair of integer (x, y) satisfies $F(x, y) = \pm 1$ and is related to α then*

$$(13) \quad |y| \leq \frac{2^{\frac{19}{4}}}{(\sqrt{3}|D_F|)^{1/4}} M(F)^{9/4}.$$

Proof. Let $\alpha = \tau + it$, with $t \neq 0$, be a non-real root of $F(x, 1) = 0$. If a solution (x, y) of (1) is related to α then $\bar{\alpha}$, the complex conjugate of α , is also a root of $F(x, 1) = 0$ and we have

$$\left| \frac{x}{y} - \alpha \right| = \frac{\left| \frac{x}{y} - \alpha \right| + \left| \frac{x}{y} - \bar{\alpha} \right|}{2} \geq \frac{|\alpha - \bar{\alpha}|}{2}.$$

Moreover, if $\beta \neq \alpha$ is a root of $F(x, 1) = 0$ then

$$\left| \frac{x}{y} - \beta \right| \geq \frac{\left| \frac{x}{y} - \alpha \right| + \left| \frac{x}{y} - \beta \right|}{2} \geq \frac{|\beta - \alpha|}{2}.$$

Thus

$$\begin{aligned} \frac{1}{|y|^4} &= \left| \frac{x}{y} - \alpha \right| \prod_{\alpha_i \neq \alpha} \left| \frac{x}{y} - \alpha_i \right| \\ &\geq \frac{|\alpha - \bar{\alpha}|}{2} \prod_{\alpha_i \neq \alpha} \frac{|\alpha - \alpha_i|}{2} \\ &= |\alpha - \bar{\alpha}| |f'(\alpha)| 2^{-4}. \end{aligned}$$

By Lemma 2.3,

$$|\alpha - \bar{\alpha}| \geq \sqrt{3}(4)^{-3} M(F)^{-3}.$$

This, together with Lemma 2.4, shows that

$$\frac{1}{|y|^4} \geq \sqrt{3} 2^{-19} \frac{|D_F|}{M(F)^9}.$$

This completes our proof. \square

Suppose that (x_1, y_1) and (x_2, y_2) are two solutions related to a fixed root α , with $M(f)^{\frac{17}{12} + \theta_3} < y_1 < y_2 < M(F)^{9/4}$. Similar to the proof of Lemma 4.5, we define δ_j , for $j = 1, 2$, by

$$y_j = M(F)^{1 + \delta_j}.$$

Inequality (12) implies that

$$3\delta_1 \leq \delta_2.$$

From here, we conclude that if $y_1 \geq M(F)^{\frac{17}{12} + \theta_3}$ then

$$y_2 \geq M(F)^{9/4}.$$

In other words, related to each root α_i , there exists at most 1 solution in x and y with $M(F)^{\frac{17}{12} + \theta_3} < y < M(F)^{9/4}$.

When $F(x, y)$ has signature $(0, 2)$ Choose $\theta_3 > 0$ such that

$$2 \times \frac{65}{64} \left(\frac{17}{12} + \theta_3 \right) < 3.$$

From (10), we conclude that in this case $|\mathfrak{X}|$ is at most 2 and therefore (1) has at most 4 solutions with $1 \leq y < M(f)^{\frac{17}{12} + \theta_3}$.

Since F is monic, $(1, 0)$ is a trivial solution to equation (1). Therefore, when F has signature $(0, 2)$ the number of solutions to (1) does not exceed 6.

6. TRANSCENDENTAL CURVE $\phi(x, y)$

Fix a positive integer k . Define

$$(14) \quad \phi_m(x, y) = \log \left| \frac{D^{\frac{1}{4k}}(x - y\alpha_m)}{|f'(\alpha_m)|^{\frac{1}{k}}} \right|$$

and

$$(15) \quad \phi(x, y) = (\phi_1(x, y), \phi_2(x, y), \phi_3(x, y), \phi_4(x, y)).$$

Let $\|\phi(x, y)\|$ be the L_2 norm of the vector $\phi(x, y)$.

Remark. In [2], a logarithmic curve $\phi(x, y)$ is defined by taking $k = 3$. The new general definition of $\phi(x, y)$ in this paper gives us the freedom of choosing k

large enough to make our approximations sharper. In order to have our estimations correct it is sufficient to take $k = 90$.

Lemma 6.1. *Suppose that (x, y) is a solution to the equation $F(x, y) = 1$, where F is the binary form in Theorem 1.1. If*

$$|x - \alpha_i y| = \min_{1 \leq j \leq 4} |x - \alpha_j y|$$

then

$$\|\phi(x, y)\| \leq 6 \log \frac{1}{|x - \alpha_i y|} + \|\phi(1, 0)\|.$$

Proof. Let us assume that

$$|x - \alpha_{s_j} y| < 1, \quad \text{for } 1 \leq j \leq p$$

and

$$|x - \alpha_{b_k} y| \geq 1, \quad \text{for } 1 \leq k \leq 4 - p,$$

where $1 \leq p, s_j, b_k \leq 4$. We have

$$\prod_k |x - \alpha_{b_k} y| = \frac{1}{\prod_j |x - \alpha_{s_j} y|}.$$

Therefore, for any $1 \leq k \leq 4 - p$, we have

$$\log |x - \alpha_{b_k} y| \leq p \log \frac{1}{|x - \alpha_i y|}.$$

Since

$$|x - \alpha_i y| = \min_{1 \leq j \leq 4} |x - \alpha_j y|,$$

we also have

$$|\log |x - \alpha_{s_j} y|| \leq |\log |x - \alpha_i y||.$$

From here, we conclude that

$$\begin{aligned} \|\phi(x, y)\| &\leq \|\phi(1, 0)\| + (4 - p)p |\log |x - \alpha_i y|| + p |\log |x - \alpha_i y|| \\ &= \|\phi(1, 0)\| + (5p - p^2) |\log |x - \alpha_i y||. \end{aligned}$$

The function $A(p) = 5p - p^2$ obtains its maximum value 6 over $p \in \{1, 2, 3, 4\}$. \square

In the following lemma we approximate the size of $f'(\alpha)$ in terms of the discriminant and heights of f , where f' is the derivative of the polynomial f and α is a root of $f = 0$.

Lemma 6.2. *Suppose that F is a monic quartic binary form. Then $(1, 0)$ is a solution to the equation $|F(x, y)| = 1$ and*

$$\|\phi(1, 0)\| \leq 4 \log \left(2^{9/k} |D|^{\frac{-3}{4k}} M(F)^{\frac{6}{k}} \right),$$

Proof. By the definition of $\phi(x, y)$,

$$\|\phi(1, 0)\| \leq \sum_{m=1}^4 \log \left| \frac{D^{\frac{1}{4k}}}{|f'(\alpha_m)|^{\frac{1}{k}}} \right|$$

Substituting the lower bound for the $|f'(\alpha_m)|^{\frac{1}{k}}$ from Lemma 2.4 completes the proof. \square

7. EXPONENTIAL GAP PRINCIPLE

Here, our goal is to show

Proposition 7.1. *Suppose that (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are three pairs of non-trivial solutions to (1) with*

$$|x_j - \alpha_4 y_j| < 1,$$

and $|y_j| > M(F)^{3.5}$, for $j \in \{1, 2, 3\}$. If $\|\phi(x_1, y_1)\| \leq \|\phi(x_2, y_2)\| \leq \|\phi(x_3, y_3)\|$ then

$$\|\phi(x_3, y_3)\| > 0.00014 \exp\left(\frac{\|\phi(x_1, y_1)\|}{6}\right).$$

In [2], we showed that when $F(x, y)$ splits in \mathbb{R} , i.e. when the signature is $(4, 0)$, assuming that $y_3 > y_1 > M(F)^6$, one can get the following inequality

$$\|\phi(x_3, y_3)\| > \exp\left(\frac{\|\phi(x_1, y_1)\|}{6}\right) 2\sqrt{3} \log^4 \frac{1 + \sqrt{5}}{2},$$

that is sharper than the inequality in Proposition 7.1. This makes the value of D_0 in our main theorem smaller and does not decrease the upper bound for the number of solutions.

Observe that for three pairs of solutions in Proposition 7.1, the three points $\phi(x_1, y_1)$, $\phi(x_2, y_2)$ and $\phi(x_3, y_3)$ form a triangle Δ . To establish Proposition 7.1, we will find a lower bound and an upper bound for the area of Δ . Then comparing these bounds, Proposition 7.1 will be proven. The length of each side of Δ is less than $2\|\phi(x_3, y_3)\|$. Lemma 7.2 gives an upper bound for the height of Δ . Let $(x, y) \neq (1, 0)$ be a solution to (1) and let $t = \frac{x}{y}$. We have

$$\phi(x, y) = \phi(t) = \sum_{i=1}^4 \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{k}}} \mathbf{b}_i,$$

where,

$$\begin{aligned} \mathbf{b}_1 &= \frac{1}{4}(3, -1, -1, -1), & \mathbf{b}_2 &= \frac{1}{4}(-1, 3, -1, -1), \\ \mathbf{b}_3 &= \frac{1}{4}(-1, -1, 3, -1), & \mathbf{b}_4 &= \frac{1}{4}(-1, -1, -1, 3), \end{aligned}$$

Without loss of generality, we will assume that α_4 is a real root and for the pair of solution (x, y) we have

$$|x - \alpha_4 y| \leq 1.$$

We may write (see the definition of $\phi(x, y)$ in (15))

$$(16) \quad \phi(x, y) = \phi(t) = \sum_{i=1}^3 \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{k}}} \mathbf{c}_i + E_4 \mathbf{b}_4,$$

where, for $1 \leq i \leq 3$,

$$\mathbf{c}_i = \mathbf{b}_i + \frac{1}{3} \mathbf{b}_4, \quad E_4 = \log \frac{|t - \alpha_4|}{|f'(\alpha_4)|^{\frac{1}{k}}} - \frac{1}{3} \sum_{i=1}^3 \log \frac{|t - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{k}}}$$

One can easily observe that

$$\mathbf{c}_i \perp \mathbf{b}_4, \text{ for } 1 \leq i \leq 4.$$

Lemma 7.2. *Let*

$$\mathbf{L}_4 = \sum_{i=1}^3 \log \frac{|\alpha_4 - \alpha_i|}{|f'(\alpha_i)|^{\frac{1}{2}}} \mathbf{c}_i + z \mathbf{b}_4, \quad z \in \mathbb{R}.$$

Suppose that $(x, y) \neq (1, 0)$ is a pair of solution to (1) with

$$|x - \alpha_4 y| = \min_{1 \leq j \leq 4} |x - \alpha_j y|$$

and $y \geq M(F)^{3.5}$. Then the distance between $\phi(x, y)$ and the line \mathbf{L}_4 is less than

$$\exp\left(\frac{-\|\phi(x, y)\|}{6}\right).$$

Proof. The proof goes exactly as the proof of Lemma 6.3 in [2]. \square

Lemma 7.2 shows that the height of Δ is at most

$$2 \exp\left(\frac{-\|\phi(x_1, y_1)\|}{6}\right).$$

Therefore, the area of Δ is less than

$$(17) \quad 2 \|\phi(x_3, y_3)\| \exp\left(\frac{-\|\phi(x_1, y_1)\|}{6}\right).$$

Let us now estimate the area of Δ from below. Since

$$F(x, y) = (x - \alpha_1 y)(x - \alpha_2 y)(x - \alpha_3 y)(x - \alpha_4 y) = \pm 1,$$

we conclude that $x - \alpha_i y$ is a unit in $\mathbb{Q}(\alpha_i)$ when (x, y) is a pair of solution to (1). Suppose that (x_1, y_1) and (x_2, y_2) are two pairs of non-trivial solutions to (1). Then

$$\phi(x_1, y_1) - \phi(x_2, y_2) = \left(\log \left| \frac{x_1 - \alpha_1 y_1}{x_2 - \alpha_1 y_2} \right|, \dots, \log \left| \frac{x_1 - \alpha_4 y_1}{x_2 - \alpha_4 y_2} \right| \right) = \vec{e}.$$

Since $\frac{x_1 - \alpha_i y_1}{x_2 - \alpha_i y_2}$ is a unit in $\mathbb{Q}(\alpha_i)$, by Proposition 2.2, we have

$$\|\vec{e}\| = 8h \left(\frac{x_1 - \alpha_i y_1}{x_2 - \alpha_i y_2} \right) > 2 \left(\frac{\log \log 4}{\log 4} \right)^3 > 0.026.$$

Now we can estimate each side of Δ from below to conclude that the area of the triangle Δ is greater than

$$\sqrt{3} \left(\frac{\log \log 4}{\log 4} \right)^6 > 0.00029.$$

Comparing this with (17) we conclude that

$$2 \|\phi(x_3, y_3)\| \exp\left(\frac{-\|\phi(x_1, y_1)\|}{6}\right) > 0.00029.$$

Proposition 7.1 is immediate from here.

Note that when $\mathbb{Q}(\alpha_i)$ is a totally real field, we have a better upper bound for the size of \vec{e} ;

$$\|\vec{e}\| \geq 4 \log^2 \frac{1 + \sqrt{5}}{2}$$

(see exercise 2 on page 367 of [14]). Now we can estimate each side of Δ from below by

$$4\sqrt{3} \log^4 \frac{1 + \sqrt{5}}{2}.$$

In order to study the curve $\phi(t)$, we will consider some well-known geometric properties of the unit group U of $K = \mathbb{Q}(\alpha)$, where α is a root of $F(x, 1) = 0$. Let r be the number of real conjugate fields of K and $2s$ the number of complex conjugate fields of K . Then by Dirichlet's unit theorem, the ring of integers O_K contains $r + s - 1$ fundamental units and there are three possibilities:
 If $F(x, 1) = 0$ has no real roots, we call F a form of signature $(0, 2)$.
 If $F(x, 1) = 0$ has 4 real roots, we call F a form of signature $(4, 0)$.
 If $F(x, 1) = 0$ has 2 real roots, we call F a form of signature $(2, 1)$.

Here we are working with quartic forms ($r + 2s = 4$) we have $r + s - 1 = \frac{r}{2} + 1$ fundamental units in O_K . Let τ be the obvious restriction of the embedding of $\mathbb{Q}(\alpha)$ in \mathbb{C}^4 ; i.e. $\tau(u) = (u_1, u_2, u_3, u_4)$, where u_i are algebraic conjugates of u . By Dirichlet's unit theorem, we have a sequence of mappings

$$(18) \quad \tau : U \mapsto V \subset \mathbb{C}^4$$

and

$$(19) \quad \log : V \mapsto \Lambda,$$

where V is the image of the map τ , Λ is a $r + s - 1$ -dimensional lattice, and the mapping \log is defined as follows:

For $(x_1, \dots, x_{r+s-1}) \in V$,

$$\log(x_1, \dots, x_{r+s-1}) = (\log |x_1|, \dots, \log |x_{r+s-1}|),$$

where r and s are defined in Dirichlet's unit Theorem. We have $r + s - 1 \leq 3$. If (x, y) is a pair of solutions to (1) then

$$(x - \alpha_j y)$$

is a unit in $\mathbb{Q}(\alpha_i)$. Suppose that $\mathbb{Q}(\alpha_i)$ is a real number field and

$$\lambda_2, \dots, \lambda_{r+s}$$

are fundamental units of $\mathbb{Q}(\alpha_i)$ and are chosen so that

$$\log(\tau(\lambda_2)), \dots, \log(\tau(\lambda_{r+s}))$$

are respectively first to $r + s - 1$ successive minimas of the lattice Λ . Let us assume that

$$\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\|,$$

form a *reduced* basis for the lattice Λ , so that

$$(20) \quad \phi(x, y) = \phi(1, 0) + \sum_{k=2}^{r+s} m_k \log(\tau(\lambda_k)), \quad m_k \in \mathbb{Z},$$

with

$$(21) \quad \|m_k \log(\tau(\lambda_k))\| \leq \|\phi(x, y) - \phi(1, 0)\|.$$

8. GEOMETRY OF $\phi(x, y)$

Lemma 8.1. *For every fixed integer m , there are at most $2r + 2s - 2$ solutions (x, y) to (1) for which in (20), $m_{r+s} = m$.*

Proof. Let S be the $(r + s - 1)$ -dimensional affine space of all vectors

$$\phi(1, 0) + \sum_{i=2}^{r+s} \mu_i \log(\tau(\lambda_i)) \quad (\mu_i \in \mathbb{R}).$$

Let $\mu_{r+s} = m$. Then the points

$$\phi(1, 0) + \sum_{i=2}^{r+s-1} \mu_i \log(\tau(\lambda_i)) + m \log(\tau(\lambda_{r+s}))$$

form an $(r + s - 2)$ -dimensional hyperplane S_1 of S . Put $f(t) = F(t, 1)$. For $t \in \mathbb{R}$, define $y(t)$ and $x(t)$ as follows:

$$\begin{aligned} y(t) &:= |f(t)|^{-1/4}, \\ x(t) &:= ty(t). \end{aligned}$$

Similar to $\phi(x, y)$, we define the curve $\phi(t)$ on \mathbb{R} :

$$\phi(t) = (\phi_1(t), \phi_2(t), \phi_3(t), \phi_4(t)),$$

where, for $1 \leq m \leq 4$

$$\phi_m(t) = \log \left| \frac{D^{\frac{1}{4k}}(x(t) - \alpha_m y(t))}{|f'(\alpha_m)|^{\frac{1}{k}}} \right|.$$

Observe that for an integral solution (x, y) to (1) and $\phi(x, y)$ defined in (15), we have

$$\phi(x, y) = \phi\left(\frac{x}{y}\right).$$

Let $\vec{N} = (N_1, N_2, N_3, N_4) \in S$ be the normal vector of S_1 . Then the number of times that the curve $\phi(t)$ intersects S_1 equals the number of solutions in t to

$$(22) \quad \vec{N} \cdot \phi(t) = 0.$$

We have

$$\lim_{t \rightarrow \alpha_i^+} \log |t - \alpha_i| = -\infty$$

and

$$\lim_{t \rightarrow \alpha_i^-} \log |t - \alpha_i| = -\infty.$$

Note that if α_i is a non-real root of $F(x, 1)$ then $\bar{\alpha}_i$, the complex conjugate of α_i is also a root and we have

$$\log |t - \alpha_i| = \log |t - \bar{\alpha}_i|.$$

If $\alpha_1, \dots, \alpha_r$ are the reals roots and $\alpha_{r+1}, \dots, \alpha_{r+s}, \alpha_{r+s+1}, \dots, \alpha_{r+2s}$ are non-real roots with $\alpha_{r+s+k} = \bar{\alpha}_{r+k}$, then the derivative $\frac{d}{dt} (\vec{N} \cdot \phi(t))$ can be written as $\frac{P(t)}{Q(t)}$, where $Q(t) = (t - \alpha_1) \dots (t - \alpha_r)(t - \alpha_{r+1}) \dots (t - \alpha_{r+s})$ and $P(t)$ is a polynomial of degree $r + s - 1$. Therefore, the derivative has at most $r + s - 1$ zeros and consequently, the equation (22) can not have more than $2r + 2s - 2$ solutions. \square

Lemma 8.2. *Let F be an irreducible monic quartic form. Suppose that (x, y) is a solution to the Thue equation $F(x, y) = \pm 1$ with $y \geq M(F)^{3.5}$. Then*

$$\|\phi(1, 0)\| < \|\phi(x, y)\|.$$

Proof. Let $\alpha_1, \dots, \alpha_n$ be the roots of $F(z, 1) = 0$. Then

$$\left(\frac{x}{y} - \alpha_1\right)\left(\frac{x}{y} - \alpha_2\right)\left(\frac{x}{y} - \alpha_3\right)\left(\frac{x}{y} - \alpha_4\right) = \frac{\pm 1}{y^4}.$$

There must exist a root α_j so that $\left|\frac{x}{y} - \alpha_j\right| \geq \frac{1}{y}$. By Lemma 2.4 and since $y \geq M(F)^{3.5}$, the absolute value of the term $\phi_j(x, y)$ alone exceeds $\|\phi(1, 0)\|$ (see Lemma 6.2). \square

Recall that we assumed $F(x, y)$ is a monic form. Therefore $(1, 0)$ is a trivial solution to the equation $F(x, y) = \pm 1$.

Definition of the set \mathfrak{A} . Assume that equation (1) has more than $2r + 2s - 2$ solutions. Then we can list $2r + 2s - 3$ solutions $(x_i, y_i) \neq (1, 0)$ ($1 \leq i \leq 2r + 2s - 3$), so that $r_i = \|\phi(x_i, y_i)\|$ are the smallest among all $\|\phi(x, y)\|$, where (x, y) varies over all non-trivial pairs of solutions. We denote the set of all these $2r + 2s - 3$ solutions and the trivial solution $(1, 0)$ by \mathfrak{A} .

Corollary 8.3. *Let $(x, y) \notin \mathfrak{A}$ be a solution to (1) with $y > M(F)^{3.5}$. Then*

$$\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\| \leq 2\|\phi(x, y)\|.$$

Proof. Since we have assumed that $\|\log(\tau(\lambda_2))\| \leq \dots \leq \|\log(\tau(\lambda_{r+s}))\|$, it is enough to show that $\|\log(\tau(\lambda_{r+s}))\| \leq 2\|\phi(x, y)\|$. By Lemma 8.1, there is at least one solution $(x_0, y_0) \in \mathfrak{A}$ so that

$$\phi(x, y) - \phi(x_0, y_0) = \sum_{i=2}^{r+s} k_i \log(\tau(\lambda_i)),$$

with $k_n \neq 0$. Since $\{\log(\tau(\lambda_i))\}$ is a reduced basis for the lattice Λ in (19), we conclude that

$$\|\log(\tau(\lambda_{r+s}))\| \leq \|\phi(x, y) - \phi(x_0, y_0)\| \leq 2\|\phi(x, y)\|.$$

\square

Lemma 8.4. *Suppose that (x, y) is a solution to (1) with $y \geq M(F)^{3.5}$. We have*

$$\|\phi(x, y)\| \geq \frac{1}{2} \log\left(\frac{|D|^{1/12}}{2}\right).$$

Proof. Let α_i and α_j be two distinct roots of quartic polynomial $F(x, 1)$. We have

$$\begin{aligned} \left| e^{\phi_i(1,0) - \phi_i(x,y)} - e^{\phi_j(1,0) - \phi_j(x,y)} \right| &= \left| \frac{1}{x - y\alpha_i} - \frac{1}{x - y\alpha_j} \right| \\ &= \frac{|\alpha_i - \alpha_j| |y|}{|x - y\alpha_i| |x - y\alpha_j|} \\ &\geq \frac{|\alpha_i - \alpha_j|}{|x - y\alpha_i| |x - y\alpha_j|}. \end{aligned}$$

Since $|\phi_i| < \|\phi\|$ and $\|\phi(1, 0)\| \leq \|\phi(x, y)\|$, we may conclude

$$\left(2e^{2\|\phi(x, y)\|}\right)^6 \geq \prod_{1 \leq i < j \leq 4} \left| \frac{1}{x - y\alpha_i} - \frac{1}{x - y\alpha_j} \right| \geq \sqrt{D}.$$

□

9. PROOF OF PROPOSITION 3.2 FOR FORMS OF SIGNATURE (4, 0)

Let $(x, y) \neq (1, 0)$ be a pair of solution to (1) and

$$\phi(x, y) = \phi(1, 0) + \sum_{k=2}^{r+s} m_k \log(\tau(\lambda_k)), \quad m_k \in \mathbb{Z}.$$

Let us set $t = \frac{x}{y}$ and define

$$T_{i,j}(t) := \log \left| \frac{(t - \alpha_i)(\alpha_4 - \alpha_j)}{(t - \alpha_j)(\alpha_4 - \alpha_i)} \right|.$$

Notice that

$$\begin{aligned} T_{i,j}(x, y) = T_{i,j}(t) &= \log \left| \frac{\alpha_4 - \alpha_i}{\alpha_4 - \alpha_j} \right| + \log \left| \frac{t - \alpha_j}{t - \alpha_i} \right| \\ &= \log \left| \frac{\alpha_4 - \alpha_i}{\alpha_4 - \alpha_j} \right| + \log \left| \frac{x - \alpha_j y}{x - \alpha_i y} \right| \\ (23) \qquad \qquad \qquad &= \log |\lambda_{i,j}| + \sum_{k=2}^4 m_i \log \frac{|\lambda_k|}{|\lambda'_k|}, \end{aligned}$$

where $\lambda_{i,j} = \log \left| \frac{\alpha_4 - \alpha_i}{\alpha_4 - \alpha_j} \right|$ and λ_k and λ'_k are fundamental units in $\mathbb{Q}(\alpha_j)$ and $\mathbb{Q}(\alpha_i)$, respectively.

Lemma 9.1. *Let (x, y) be a pair of solution to (1) with $|y| \geq M(F)^{3.5}$. Then there exists a pair (i, j) for which*

$$|T_{i,j}(x, y)| < \exp \left(-\frac{\|\phi(t)\|}{6} \right).$$

Proof. This is a consequence of Lemma 7.2 and the proof goes exactly the same as the proof of Lemma 6.8 in [2]. □

Let index σ be the isomorphism from $\mathbb{Q}(\alpha_i)$ to $\mathbb{Q}(\alpha_j)$ such that $\sigma(\alpha_i) = \alpha_j$. We may assume that $\sigma(\lambda_i) = \lambda'_i$ for $i = 2, 3, 4$. Let (x_1, y_1) , (x_2, y_2) , (x_3, y_3) be three distinct solutions to (1) with

$$y_k \geq M(F)^{3.5}$$

and

$$|x_k - \alpha_4 y_k| = \min_{1 \leq i \leq 4} |x_k - \alpha_i y_k| \quad k \in \{1, 2, 3\}.$$

This assumption will lead us to a contradiction at the end of this section, implying that related to each real root of $F(x, 1) = 0$, there are at most 2 solutions with $y \geq M(F)^{3.5}$. Recall that related to a non-real root, there exists no such solution.

Put $r_k = \|\phi(x_k, y_k)\|$ and assume that $r_1 \leq r_2 \leq r_3$. We will apply Matveev's lower bound to

$$T_{i,j}(x_3, y_3) = \log |\lambda_{i,j}| + \sum_{k=2}^{r+s} m_k \log \frac{|\lambda_k|}{|\lambda'_k|},$$

where (i, j) is chosen so that Lemma 9.1 is satisfied and $m_k \in \mathbb{Z}$. Forms of signature $(0, 2)$ have one fundamental unit and forms of signature $(2, 1)$ have two fundamental units. Moreover, if $\lambda_{i,j}$ is a unit then we can write $T_{i,j}(x, y)$ as a linear form in fewer number of logarithms. We remark here that when dealing with linear forms in 2 logarithms, one can use sharper lower bounds (see for example [6]).

Suppose that λ is a unit in the number field and λ' is its algebraic conjugate. We have

$$h(\lambda') = h(\lambda) = \frac{1}{8} |\log(\tau(\lambda))|_1,$$

where h is the logarithmic height and $|\cdot|_1$ is the L_1 norm on \mathbb{R}^4 and the mappings τ and \log are defined in (18) and (19). So we have

$$h(\lambda) = \frac{1}{8} |\log(\tau(\lambda))|_1 \leq \frac{\sqrt{4}}{8} \|\log(\tau(\lambda))\|,$$

where $\|\cdot\|$ is the L_2 norm on \mathbb{R}^{r+s-1} . Since α_4, α_i and α_j have degree 4 over \mathbb{Q} , the number field $\mathbb{Q}(\alpha_4, \alpha_i, \alpha_j)$ has degree $d \leq 24$ over \mathbb{Q} . So when λ is a unit

$$(24) \quad \max\left\{dh\left(\frac{\lambda}{\lambda'}\right), \left|\log\left(\left|\frac{\lambda}{\lambda'}\right|\right)\right|\right\} \leq \max\left\{24h\left(\frac{\lambda}{\lambda'}\right), \left|\log\left(\left|\frac{\lambda}{\lambda'}\right|\right)\right|\right\} \leq 12 \|\log(\tau(\lambda))\|.$$

In order to apply Theorem 2.5 to $T_{i,j}(x, y)$, we will take, for $k > 1$,

$$A_k = 12 \|\log(\tau(\lambda))\|.$$

By (21) and Lemma 8.2, we have

$$A_k \leq 24 \|\phi(x, y)\|, \quad \text{for } k > 1.$$

Now we need to estimate A_1 .

Lemma 9.2. *Let F be an irreducible binary quartic form with integral coefficients. Assume that (x, y) is a solution to (1) with $y \geq M(F)^{3.5}$. Then, we have*

$$h\left(\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j}\right) \leq 2 \log 2 + 2 \|\phi(x, y)\|.$$

Proof. Let $\beta_i = x - y\alpha_i$. We have

$$\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j} = \frac{\beta_k - \beta_i}{\beta_k - \beta_j}.$$

Thus by (5) and (6),

$$h\left(\frac{\alpha_k - \alpha_i}{\alpha_k - \alpha_j}\right) \leq 2 \log 2 + 4h(\beta_k).$$

To complete the proof, we will show that

$$h(\beta_k) \leq \frac{1}{2} \|\phi(x, y)\|.$$

Set

$$v_i = \log |\beta_i| = \phi_i(x, y) - \phi_i(1, 0) \quad \text{for } i \in \{1, 2, 3, 4\}$$

and

$$\vec{v} = (v_1, v_2, v_3, v_4).$$

Since β_k is a unit, we have

$$h(\beta_k) = \frac{1}{8} \sum_{i=1}^4 |v_i| = \frac{1}{8} (s_1, s_2, s_3, s_4) \cdot \vec{v},$$

where $s_1, s_2, s_3, s_4 \in \{+1, -1\}$. Since $\|(s_1, s_2, s_3, s_4)\| = 2$, we obtain

$$h(\beta_k) \leq \frac{1}{4} \|\vec{v}\|.$$

On the other hand, we have

$$\|\vec{v}\| \leq \|\phi(x, y)\| + \|\phi(1, 0)\| \leq 2\|\phi(x, y)\|.$$

This completes our proof. \square

Set, for $k \in \{1, 2, 3\}$,

$$r_k = \|\phi(x_k, y_k)\|.$$

We may take

$$A_1 = 48 \log 2 + 48r_1$$

(recall that $\alpha_1, \alpha_i, \alpha_j$ are algebraic conjugates). Since $m_1 = 1$, we will put

$$B = \max\{1, \max\{m_j A_j / A_1 : 1 \leq j \leq 4\}\}.$$

Since we assumed that $\tau(\lambda_i), 2 \leq i \leq r+s-1$ are successive minimas for the lattice Λ , we have

$$|m_j A_j| \leq r_3 + \|\phi(1, 0)\| < 2r_3.$$

Thus we may take $B = \frac{r_3}{12}$ (see (21)).

Proposition 2.5 implies that for a constant number K ,

$$\log T_{i,j}(x_3, y_3) > -K r_1^4 \log r_3.$$

Comparing this with Lemma 9.1, we have

$$\left(\frac{-r_3}{6}\right) > -K r_1^4 \log r_3,$$

or

$$\frac{r_3}{\log r_3} < 6K r_1^4.$$

Thus there is a computable constant number K_1 , so that

$$(25) \quad r_3 < K_1 r_1^4,$$

This is because r_3 is large enough by Lemma 8.4. But by Lemma 7.1 we have

$$r_3 > 0.00014 \exp\left(\frac{r_1}{6}\right).$$

This is a contradiction, for by Lemma 8.4,

$$r_1 \geq \frac{1}{2} \log \left(\frac{|D|^{\frac{1}{12}}}{2} \right)$$

and D is large. Thus, there are at most 2 solutions (x, y) with $y \geq M(F)^{3.5}$ related to each real root α_i . The proof of Proposition 3.2 for forms with signature $(4, 0)$ is complete now. This argument can be used for forms with signature $(2, 1)$, as well and will give us an bound of 17 upon the number of solutions to (1) for this case. In the next section, we will see that for forms of signature $(2, 1)$ we do not need to consider the set \mathfrak{A} , as the lattice generated by the fundamental units of the corresponding number field is contained in a plane.

10. PROOF OF PROPOSITION 3.2 FOR FORMS OF SIGNATURE (2, 1)

Let $F(x, y)$ be a quartic form of signature (2, 1). Suppose that (x_1, y_1) , (x_2, y_2) and (x_3, y_3) are three pairs of non-trivial solution to (1) with

$$|x_j - \alpha_4 y_j| < 1,$$

and $|y_j| > M(F)^{3.5}$, for $j \in \{1, 2, 3\}$. We will assume α_4 is real (recall that related to a non-real root, there exists no solution (x, y) with $|y| > M(F)^{3.5}$). The real number field $\mathbb{Q}(\alpha_4)$ has two fundamental units λ_2, λ_3 chosen so that $\log(\tau(\lambda_2)), \log(\tau(\lambda_3))$ are successive minimas of the lattice Λ , with

$$(26) \quad \|\log(\tau(\lambda_2))\| \|\log(\tau(\lambda_3))\| \geq \frac{2}{\sqrt{3}} \text{Vol}(\Lambda),$$

where $\text{Vol}(\Lambda)$ is the volume of fundamental parallelepiped of lattice Λ . If (x, y) is a solution to $|F(x, y)| = 1$, then

$$\phi(x, y) \in \phi(1, 0) + \Lambda = \Lambda_1.$$

Note that

$$\text{Vol}(\Lambda) = \text{Vol}(\Lambda_1).$$

For distinct pairs of solution (x_1, y_1) , (x_2, y_2) and (x_3, y_3) , three vectors $\phi(x_1, y_1)$, $\phi(x_2, y_2)$ and $\phi(x_3, y_3)$ generate a sub-lattice of Λ_1 with the volume of fundamental parallelepiped equal to $2A$. Therefore,

$$(27) \quad 2A \geq \text{Vol}(\Lambda_1) = \text{Vol}(\Lambda).$$

On the other hand, by (17), A , the area of Δ is less than

$$2 \|\phi(x_3, y_3)\| \exp\left(\frac{-\|\phi(x_1, y_1)\|}{6}\right).$$

This, together with (27) gives

$$(28) \quad \|\phi(x_3, y_3)\| > \frac{\text{Vol}(\Lambda)}{4} \exp\left(\frac{\|\phi(x_1, y_1)\|}{6}\right).$$

Let us replace Proposition 7.1 by the above inequality when $F(x, y)$ has signature (2, 1).

Put $r_k = \|\phi(x_k, y_k)\|$ and assume that $r_1 \leq r_2 \leq r_3$. We will apply Matveev's lower bound to

$$T_{i,j}(x_3, y_3) = \log|\lambda_{i,j}| + \sum_{k=2}^{r+s} m_k \log \frac{|\lambda_k|}{|\lambda'_k|},$$

where (i, j) is chosen so that Lemma 9.1 is satisfied and $m_k \in \mathbb{Z}$. Similar to Section 9, let index σ be the isomorphism from $\mathbb{Q}(\alpha_i)$ to $\mathbb{Q}(\alpha_j)$ such that $\sigma(\alpha_i) = \alpha_j$. We may assume that $\sigma(\lambda_i) = \lambda'_i$ for $i = 2, 3, 4$. Recall that related to a non-real root, there exists no such solution.

We use our estimation from Section 9 for A_1 and B :

$$A_1 = 48 \log 2 + 48r_1$$

and

$$B = \frac{r_3}{12}.$$

For fundamental units λ_k , with $k \in \{2, 3\}$, using (24), we may put

$$A_k = 12 \|\log(\tau(\lambda))\|.$$

Theorem 2.5 and (26) imply that for a constant number K ,

$$\log T_{i,j}(x_3, y_3) > -K r_1 \log r_3 \text{Vol}(\Lambda).$$

Comparing this with Lemma 9.1, we have

$$\left(\frac{-r_3}{6}\right) > -K r_1 \log r_3 \text{Vol}(\Lambda),$$

or

$$\frac{r_3}{\log r_3} < 6K r_1 \text{Vol}(\Lambda).$$

Thus, since r_3 is large enough by Lemma 8.4, there is a computable constant number K_2 , so that

$$r_3 < K_2 r_1 \text{Vol}(\Lambda)$$

(compare this with (25)). But by (28) we have

$$r_3 > \frac{\text{Vol}(\Lambda)}{4} \exp\left(\frac{r_1}{6}\right).$$

Consequently, for some positive constant K_3 ,

$$r_1 > K_3 \exp\left(\frac{r_1}{6}\right).$$

This is a contradiction by Lemma 8.4 and since the discriminant is large. Thus, there are at most 2 solutions (x, y) with $y \geq M(F)^{3.5}$ related to each real root α_i . The proof of Proposition 3.2 and Theorem 1.1 are complete now.

11. ACKNOWLEDGEMENTS

I would like to thank Professor Ryotaro Okazaki for his helpful comments and suggestions. Most of this work has been done during my visit to mathematisches forschungsinstitut oberwolfach and Max-Planck institute for mathematics in Bonn. I am very grateful to both institutes for support and hospitality.

REFERENCES

- [1] S. Akhtari, The method of Thue-Siegel for binary quartic forms, *Acta Arith.* 141 (2010), no. 1, 1–31.
- [2] S. Akhtari, R. Okazaki, The quartic Thue equations, *J. Number Theory* 130 (2010), no. 1, 40–60.
- [3] E. Bombieri, W. M. Schmidt, On Thue's equation, *Invent. Math.* 88 (1987), 69–81.
- [4] Y. Bugeaud, Bornes effectives pour les solutions des équations en S-unités et des équations de Thue-Mahler, *J. Number Theory* 71 (1998), 227–244.
- [5] J.H. Evertse and K. Györy, Effective finiteness results for binary forms with given discriminant, *Compositio Math.* 79 (1991), 169–204.
- [6] N. Gouillon, Explicit lower bounds for linear forms in two logarithms. *J. Théor. Nombres Bordeaux* 18 (2006), no. 1, 125–146.
- [7] Magma, <http://magma.maths.usyd.edu.au/magma/>.
- [8] K. Mahler, An inequality for the discriminant of a polynomial, *Michigan Math. J.* 11 (1964), 257–262.
- [9] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers, *Izv. Ross. Akad. Nauk Ser. Mat.* 62 (1998), 81–136, translation in *Izv. Math.* 62 (1998), 723–772.
- [10] E. M. Matveev, An explicit lower bound for a homogeneous rational linear form in logarithms of algebraic numbers II, *Izv. Ross. Akad. Nauk Ser. Mat.* 64 (2000), 125–180, translation in *Izv. Math.* 64 (2000), 1217–1269.
- [11] M. Mignotte, A. Pethö and R. Roth, Complete solutions of a family of quartic Thue and index form equations, *Math. Comp.* 65 (1996), 341–354.

- [12] L.J. Mordell, Diophantine Equations, Academic Press, London and New York (1969).
- [13] R. Okazaki, Geometry of a cubic Thue equation, Publ. Math. Debrecen. 61 (2002), 267-314.
- [14] M. Pohst and H. Zassenhause, Algorithmic Algebraic Number Theory, Cambridge University Press (1989).
- [15] W.M. Schmidt, Diophantine approximations and Diophantine equations, Lecture Notes in Mathematics, Springer Verlag (2000).
- [16] C. L. Stewart, On the number of solutions of polynomial congruences and Thue equations, Journal of American Math. Soc. 4 (1991), 793-838.
- [17] P. Voutier, An effective lower bound for the height of algebraic numbers, Acta Arith. 74 (1996), 81-95.
- [18] P.G. Walsh, On the number of large integer points on elliptic curves, Acta Arith. 138 (2009), no. 4, 317-327.

CRM, UNIVERSITY OF MONTREAL, P.O. BOX 6128, CENTRE-VILLE STATION, MONTREAL H3C 3J7

E-mail address: akhtari@crm.umontreal.ca