

A Jordan-Hölder Theorem for Differential Algebraic Groups

Phyllis J. Cassidy* and Michael F. Singer†

Abstract

We show that a differential algebraic group can be filtered by a finite subnormal series of differential algebraic groups such that successive quotients are almost simple, that is have no normal subgroups of the same type. We give a uniqueness result, prove several properties of almost simple groups and, in the ordinary differential case, classify almost simple linear differential algebraic groups.

1 Introduction

Let (k, δ) be a differential field of characteristic zero with derivation δ and let $k[\delta]$ be the ring of linear differential operators with coefficients in k . It has been known for a long time (see [40] for a brief history and [2],[34],[45] for more recent algorithmic results) that one has a form of unique factorization for this ring:

Given $L \in k[\delta]$, $L \notin k$, there exist irreducible $L_i \in k[\delta]$, $i = 1, \dots, r$ such that $L = L_1 \cdots L_r$. Furthermore, if $L = \tilde{L}_1 \cdots \tilde{L}_s$, $\tilde{L}_i \in k[\delta]$, \tilde{L}_i irreducible, then $r = s$ and, for some permutation σ of the subscripts, there exist nonzero $R_i, S_i \in k[\delta]$ with $\deg(R_i) < \deg(\tilde{L}_i)$ such that $L_{\sigma(i)}R_i = S_i\tilde{L}_i$ for $i = 1, \dots, r$.

A direct generalization of this to partial differential operators fails as the following example shows. In [6], Blumberg considered the following third order linear partial differential

*Department of Mathematics, Smith College, MA 01063; Department of Mathematics, The City College of New York, New York, NY 10038, USA, email: pcassidy@smith.edu

†Department of Mathematics, North Carolina State University, Box 8205, Raleigh, NC 27695-8205, USA, email: singer@math.ncsu.edu. The second author was partially supported by NSF Grant CCR-0634123 and he would also like to thank the Max Planck Institut für Mathematik and the Courant Institute of Mathematical Sciences for their hospitality. This work was begun during a visit to the first institute in the Spring of 2006 and continued during a stay as a Visiting Member at the second institute in the Fall of 2008.

operator

$$L = \delta_x^3 + x\delta_x^2\delta_y + 2\delta_x^2 + 2(x+1)\delta_x\delta_y + \delta_x + (x+2)\delta_y$$

over the differential field $\mathbb{C}(x, y)$ where $\delta_x = \frac{\partial}{\partial x}$ and $\delta_y = \frac{\partial}{\partial y}$. He showed that this operator has two factorizations

$$\begin{aligned} L &= (\delta_x + 1)(\delta_x + 1)(\delta_x + x\delta_y) \\ &= (\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y)(\delta_x + 1) . \end{aligned}$$

and that the operator $\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y$ could not be further factored in any differential extension of $\mathbb{C}(x, y)$ (this example is also discussed in [18]).

To confront this situation one must recast the factorization problem in other terms. For ordinary differential operators, the factorization result is best restated in terms of $k[\delta]$ -modules. If one considers the $k[\delta]$ -module $M_L = k[\delta]/k[\delta]L$, then the above factorization result can be stated as a Jordan-Hölder type theorem: M_L has a composition series $M_L = M_0 \supset M_1 \supset \dots \supset M_r = \{0\}$ where successive quotients M_{i-1}/M_i are simple and any two such series have the same length and, after a possible renumbering, isomorphic quotients. Casting questions in terms of ideals and modules in the partial differential case yields many interesting results (cf., [18, 19, 20, 17], [16], [32]). Other approaches to factoring linear operators are contained in [4],[39, 38], [43]. In [44], Tsarev alludes to an approach to factorization through the theory of certain abelian categories that has points of contact with the approach presented here.

We will take an alternate approach - via differential algebraic groups. This stems from the seemingly trivial observation that the solutions of a linear differential equation form a group under addition. In fact, this is an example of a linear differential algebraic group, that is a group of matrices whose entries lie in a differential field and satisfy some fixed set differential equations (here we identify $\{y \in k \mid L(y) = 0\}$ with the group $\left\{ \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} \mid L(y) = 0 \right\}$.) We shall prove a Jordan-Hölder type theorem for differential algebraic groups (of which linear differential algebraic groups are a special case.) This will encompass the factorization result for linear ordinary differential equations, allow us to give a “factorization” result for linear partial differential equations and allow us to reduce the study of general differential algebraic groups to the study of *almost simple* groups (to be defined below) and their extensions.

The rest of the paper is organized as follows. In Section 2, we shall review some definitions and facts from differential algebra and the theory of linear differential groups. We will state and prove a Jordan-Hölder Theorem for linear differential groups and show how it applies to the example of Blumberg above. In Section 3, we will discuss the structure of almost simple groups in more detail.

The authors would like to acknowledge the influence of Ellis Kolchin. The first author discussed her notions of solid and the core (what we call strongly connected and the strong

identity component) with him in the 1980's and he suggested a version of the Jordan-Hölder Theorem below.

2 Jordan-Hölder Theorem

Let (k, Δ) be a differential field of characteristic zero with commuting set of derivations $\Delta = \{\delta_1, \dots, \delta_m\}$ and (\mathbb{U}, Δ) a k -universal field containing k (see [24] for definitions of this and other concepts from differential algebra). In this section, we will consider differential algebraic varieties defined over k (for example, subsets of \mathbb{U}^n of elements satisfying polynomial differential equations with coefficients in k) and refer to these as Δ - k -sets and we will similarly use the terminology Δ - k -morphisms and Δ - k -group (see [25] for the formal definition of these concepts or, equivalently one could work with definable sets and definable groups in the theory of differential fields, (cf., [31])). If no k is mentioned, we assume that objects are defined over \mathbb{U} . Although we will state and prove our results in this general context, readers who wish to restrict themselves to *linear* differential algebraic groups or just solutions of systems of homogeneous linear differential equations, will find almost all that is needed (with one notable exception which we will discuss later) in the papers [11, 12].

2.1 Differential Type and Typical Differential Transcendence Degree

The key concepts used in this paper are *differential type* and *typical differential transcendence degree*. We briefly review their definitions. Let $\eta = (\eta_1, \dots, \eta_n) \in \mathbb{U}^n$ and let $\Theta(s)$ be the set of products of elements from Δ of total degree at most s . In Section II.12 of [24], Kolchin shows that there is a numerical polynomial (i.e., a polynomial taking on integer values on the integers), the *differential dimension polynomial*, $\omega_{\eta/k}(s)$ of degree at most m such that for sufficiently large values of s , we have $\omega_{\eta/k}(s) = \text{tr.deg.}_k k(\theta(\eta_i))_{\theta \in \Theta(s), 1 \leq i \leq n}$ (see Example 2.29 for further discussions concerning calculating $\omega_{\eta/k}(s)$ and also [26] for further information, algorithms and references concerning this polynomial). Writing $\omega_{\eta/k}(s) = \sum_{0 \leq i \leq m} a_i \binom{s+i}{i}$, one defines the *differential type* τ of $\eta = (\eta_1, \dots, \eta_n)$ over k to be the degree of $\omega_{\eta/k}(s)$ and the *typical differential transcendence degree* to be the coefficient a_τ . If X is a Δ - k -set, we define its differential type $\tau(X)$ to be the maximum of the types of its elements with entries in \mathbb{U} and its typical differential transcendence degree $a_\tau(X)$ to be the maximum of the typical differential transcendence degree of its elements with entries in \mathbb{U} and of differential type $\tau(X)$. Informally, we can say that the elements of X depend on $\tau(X)$ independent variables and $a_\tau(X)$ arbitrary functions (this is made precise by Theorem 7, p. 119 of [24]). We shall need the following facts concerning the behavior of the type under morphisms.

Lemma 2.1 *Let X and Y be irreducible Δ - k -sets and $f : X \rightarrow Y$ be an injective dominant Δ - k -morphism. Then $\tau(X) = \tau(Y)$.*

Proof. Let $x = (x_1, \dots, x_n)$ be a generic point of X over k . By assumption $y = f(x)$ is a generic point of Y over k and, since f is differentially rational, $k \langle y \rangle \subset k \langle x \rangle$. Note that \mathbb{U} is universal over $k \langle y \rangle$ as well. We will show that $k \langle y \rangle = k \langle x \rangle$. Let σ be a differential isomorphism of $k \langle x \rangle$ over $k \langle y \rangle$ into \mathbb{U} . Since $\sigma(y) = y$ and $\sigma(y) = \sigma(f(x)) = f(\sigma(x))$ we have that $f(\sigma(x)) = f(x)$. Since f is injective, we have that $\sigma(x) = x$. If some coordinate x_i of x were not in $k \langle y \rangle$, there would be an isomorphism of $k \langle x \rangle$ over $k \langle y \rangle$ into \mathbb{U} which would move x_i . Therefore we can conclude that $k \langle x \rangle = k \langle y \rangle$. Proposition 15(b), p. 117 of [24] implies the conclusion. ■

Corollary 2.2 *Let X and Y be irreducible Δ - k -sets and $f : X \rightarrow Y$ be an injective Δ - k -morphism. Then $\tau(X) \leq \tau(Y)$.*

Proof. Let V be the closure of $f(Y)$. The map $f : X \rightarrow V$ is injective and dominant so $\tau(X) = \tau(V)$. Proposition 15(b), p. 117 of [24] implies that $\tau(V) \leq \tau(Y)$. ■

Let G be a Δ - k -group and H a Δ - k -subgroup of G . Kolchin showed ([25] Ch. IV.4) that the coset space G/H has the structure of a Δ - k -set (in fact, a Δ - k -homogeneous space for G) and showed that

$$\tau(G) = \max(\tau(H), \tau(G/H)). \quad (1)$$

Furthermore, he showed that if $\tau = \tau(G)$, then

$$a_\tau(G) = a_\tau(H) + a_\tau(G/H) \quad (2)$$

where $a_\tau(H) = 0$ if $\tau(G) > \tau(H)$ with a similar convention for $a_\tau(G/H)$.

Remarks 2.3 1. The fact that G/H has the structure of a k -differential algebraic set also follows from the model-theoretic concept of *elimination of imaginaries* which holds for differentially closed fields ([29], p. 57).

2. The simplest (to define, not necessarily to understand) examples of differential algebraic groups are zero sets of systems of linear homogeneous differential equations in one indeterminate. Cassidy [11] has shown that these are the only Δ -subgroups of $\mathbb{G}_a(\mathbb{U}) = (\mathbb{U}, +)$. In this case, quotients are easy to construct. If $H \subset G \subset \mathbb{G}_a(\mathbb{U})$ and H is defined by the vanishing of L_1, \dots, L_t , then the image in \mathbb{U}^t of G under the map $y \mapsto (L_1(y), \dots, L_t(y))$ is a Δ -subgroup of \mathbb{U}^n isomorphic to G/H . Grigoriev and Schwarz [18, 19, 20, 17] have made extensive use of the differential type and typical transcendence degree in their studies of systems of linear partial differential equations and refer to the pair (τ, a_τ) as the *gauge* of the system.

3. As we mentioned earlier, if one restricts one's attention to *linear* differential algebraic groups (as defined by Cassidy [11]), one will find all needed facts in the works of Cassidy and [24] except for the equations (1) and (2), which one can just assume when reading the following results and proofs.

2.2 Strongly Connected Groups, Almost Simple Groups and Isogeny

2.2.1 Strongly Connected Groups

We now define the *strong identity component* of a Δ - k -group G . Let \mathcal{S} be the set of all Δ - \mathbb{U} -subgroups $H \subset G$ such that $\tau(H) = \tau(G)$ and $a_\tau(H) = a_\tau(G)$. Note that from (1) and (2), this is the same as the set of Δ - \mathbb{U} -subgroups $H \subset G$ such that $\tau(G/H) < \tau(G)$. We claim that if $H_1, H_2 \in \mathcal{S}$, then $H_1 \cap H_2 \in \mathcal{S}$. To see this, note that we have an injection of $H_1/H_1 \cap H_2$ into G/H_2 , so by (1) we have $\tau(H_1/H_1 \cap H_2) < \tau(G)$. Again by (1), we have that $\tau(G/H_1 \cap H_2) = \max(\tau(G/H_1), \tau(H_1/H_1 \cap H_2)) < \tau(G)$. Therefore $H_1 \cap H_2 \in \mathcal{S}$.

Since the differential algebraic topology is noetherian, the set \mathcal{S} has minimal elements. The above argument shows that there is a unique minimal element and justifies the following definition.

Definition 2.4 *Let G be a Δ - k -group. The strong identity component G_0 of G is defined to be the smallest Δ - \mathbb{U} -subgroup H of G such that $\tau(G/H) < \tau(G)$. We say that G is strongly connected if $G_0 = G$.*

Note that G_0 is contained in the identity component G^0 . Therefore, every strongly connected Δ -group is connected.

Remarks 2.5 1. In the usual theory of algebraic groups, one can define the identity component of a group G to be the smallest subgroup G^0 such that G/G^0 is finite and define a group to be connected if $G = G^0$. The notions of strong identity component and strongly connected are meant to be refinements in the context of Δ -groups of these former concepts.

2. If $\sigma : \mathbb{U} \rightarrow \mathbb{U}$ is a Δ - k -automorphism of \mathbb{U} , then $\sigma(G_0)$ is again a minimal element of \mathcal{S} and so must coincide with G_0 . Therefore, Corollary 2, p.77 of [25] implies that G_0 is defined over k . Note that if $\phi : G \rightarrow G$ is a Δ -automorphism of G , then $\phi(G_0)$ is again a minimal element of \mathcal{S} and so must equal G_0 . Therefore G_0 is a characteristic subgroup of G . In particular G_0 is a *normal* subgroup of G and if H is a normal subgroup of G , the strong identity component of H is again normal in G .

3. The analogue of the fact that if G is an algebraic group, every connected algebraic subgroup is contained in the identity component of G is the following: If G is a Δ -group of type τ , then every strongly connected Δ -subgroup H of type τ is contained in the strong

identity component of G . To see this, first note that the usual isomorphism theorems hold for Δ -groups ([25], Chapter IV). Since G_0 is normal in G , HG_0 is a Δ -subgroup of G and HG_0 is isomorphic to $H/G \cap G_0$ as Δ -groups. Since the canonical inclusion $H/H \cap G_0 \hookrightarrow G/G_0$ is injective, Corollary 2.2 implies that $\tau(H/H \cap G_0) < \tau(G) = \tau(H)$. Since H is strongly connected, $H \cap G_0 = H$.

4. The analogue of the fact that if G, G' are algebraic groups, and $\phi : G \rightarrow G'$ is a homomorphism, then, the image of the identity component is contained in the identity component of G' is the following: If G, G' are Δ -groups of the same type, and $\phi : G \rightarrow G'$ is a homomorphism, then, the image of the strong identity component of G is contained in the strong identity component of G' . This follows immediately from the preceding remark.

5. Let G be a strongly connected Δ -group. Equation (1) implies that the image of G under a homomorphism of Δ -groups is also strongly connected, and is either trivial or has the same type.

The following Lemma is useful in showing a group is strongly connected.

Lemma 2.6 *Let G be a Δ -group and let H and N be strongly connected subgroups, with $N \triangleleft G$ and $\tau(H) = \tau(N)$. Then, the Δ -subgroup HN of G is strongly connected and $\tau(HN) = \tau(H) = \tau(N)$. Furthermore, $a_\tau(HN) = a_\tau(H) + a_\tau(N) - a_\tau(H \cap N)$.*

Proof. Without loss of generality, we may assume that $H \not\subseteq N$. We have $\tau(HN) = \max\{\tau(HN/N), \tau(N)\} = \max\{\tau(H/H \cap N), \tau(N)\}$. Since H is strongly connected and $H \cap N \neq H$, we have $\tau(H/H \cap N) = \tau(H)$. Therefore $\tau(HN) = \tau(H) = \tau(N)$. Remark 2.5.3 implies that H and N are contained in $(HN)_0$. This implies that $HN \subset (HN)_0$ so HN is strongly connected. The final statement concerning the typical dimension (as well as the statement concerning differential type) is contained in ([41] Cor. 4.3, p. 485; see also [25], Cor. 4, p. 110). ■

Examples 2.7 Let k be a differentially closed ordinary differential field with derivation δ (cf., [29, 28] or [23], where these are referred to as constrainedly closed).

1. Let G be the additive group $\mathbb{G}_a(k)$. G has type 1 and typical differential transcendence degree 1. Cassidy [11] showed that any proper Δ -subgroup H is of the form $H = \{a \in k \mid L(a) = 0\}$ for some linear operator L . The type of such a group is 0 and its typical differential transcendence degree is d where d is the order of L . Therefore the type of G/H for any proper H is equal to the type of G and so G is strongly connected.

2. Let $G_1 = G$ as above and $G_2 = \mathbb{G}_a(C)$ where $C = \{c \in k \mid \delta(c) = 0\}$, that is, the subgroup of constants. The type of G_2 is 0 and the typical differential transcendence degree is 1. Using equations (1) and (2), we see that $\tau(G_1 \times G_2) = \max(\tau(G_1), \tau(G_2)) = 1$ and $d(G_1 \times G_2) = d(G_1) = 1$. The group G_1 is minimal with respect to having the same type and typical transcendence degree as $G_1 \times G_2$ and so is the strong identity component of this group. Note that $G_1 \times G_2$ is irreducible as a $\{\delta\}$ - k -variety.

To state the analogue of the Jordan-Hölder Theorem, we need two more definitions: the first is the appropriate notion of “simple” and the second is the relevant notion of “equivalent”. The next two subsections deal with these notions.

2.2.2 Almost Simple Groups

Definition 2.8 A Δ - k -group G is almost simple if for any normal proper Δ - k -subgroup H of G we have $\tau(H) < \tau(G)$.

Note that an almost simple group is strongly connected. We shall use the term *simple* to denote a group (resp., algebraic group, Δ -group) whose only proper normal subgroup (resp., normal algebraic subgroup, normal Δ -subgroup) is the trivial group and the term *quasisimple* to denote a group (resp., algebraic group, Δ -group) G such that $G/Z(G)$ is simple and $Z(G)$ is finite, where $Z(G)$ is the center of G .

Examples 2.9 Let k be as in Example 2.7 and C its field of constants.

1. The group $\mathbb{G}_a(C)$ has type 0 with only the trivial group as a proper subgroup so it is simple (and so almost simple).
2. The group $\mathbb{G}_m(C)$ has type 0 with only finite subgroups as proper algebraic (or δ -) subgroups so it is quasisimple.
3. The group $\mathbb{G}_a(k)$ has type 1 and any proper subgroup has type 0 or -1 (if it is $\{0\}$). Therefore $\mathbb{G}_a(k)$ is almost simple.
4. If G is a quasisimple linear algebraic group defined over C , then, as we shall show in Section 3 that $G(k)$ and $G(C)$ are quasisimple δ -groups.

In Section 3, we shall show that for ordinary differential fields, these are essentially the only almost simple *linear* δ -groups.

Example 2.10 Let $\Delta = \{\delta_1, \delta_2\}$ and \mathbb{U} be a universal Δ -field. Let $G \subset \mathbb{G}_a(\mathbb{U})$ be the additive group of solutions of $\delta_1 y = f(y)$ where $f \in \mathbb{U}[\delta_2]$, the ring of linear operators in the derivation δ_2 . Following Proposition 2.45 of [42], we show that G is almost simple. If $H \subset G$ is a proper Δ -subgroup of G , then there would exist a linear operator $L \notin [\delta_1 y - f(y)]$ vanishing on H . Since $\delta_1 u = f(u)$ for all $u \in H$, we can assume that this operator is a polynomial in δ_2 alone. If this operator would have order d , then for any $v \in H$, we have $k(v, \delta_x v, \delta_x^2 v, \dots) = k(v, \delta_x v, \dots, \delta_x^{d-1} v)$. Since $\delta_t v \in k(v, \delta_x v, \dots, \delta_x^{d-1} v)$, this latter field equals $k \langle v \rangle$ which must therefore be of finite transcendence degree over k . Therefore H has type 0 and G is almost simple.

It is not hard to see that a finite normal subgroup of a connected algebraic group must be central. The next proposition is an analogous result for strongly connected Δ -groups.

Proposition 2.11 *Let G be a strongly connected Δ -group. Then every normal Δ -subgroup of smaller type is central.*

We immediately have the following corollaries.

Corollary 2.12 *Let G be an almost simple Δ -group. Any proper normal Δ -subgroup is central.*

Proof. For any Δ -group G , the strong identity component, G_0 , is a normal subgroup of the same type so $G = G_0$, that is G is strongly connected. By definition any proper normal Δ -subgroup is of smaller type. ■

Corollary 2.13 *Let G be an almost simple Δ -group. Then $G/Z(G)$ is a simple Δ -group.*

Proof of Proposition 2.11: Let N be a normal Δ -subgroup of G . We define a Δ -map

$$\alpha : G \times N \rightarrow N$$

by the formula

$$\alpha(g, a) = gag^{-1}, \quad g \in G, a \in N.$$

One checks that $\alpha(gh, a) = \alpha(g, \alpha(h, a))$ and $\alpha(1, a) = a$ and so α defines an action of G on N . For fixed $a \in N$, the map $\alpha_a(g) = gag^{-1}$ is a Δ -map from G to N that is constant on the left cosets of $Z_G(a) = \{g \in G \mid gag^{-1} = a\}$. Note that $Z_G(a)$ is a Δ -subgroup of G ([25], Section IV.4, Cor. 2(b)).

Theorem 3, p.105 of [25] implies that there is a Δ -map $\beta : G/Z_G(a) \rightarrow N$ such that $\pi \circ \beta = \alpha_a$, where π is the canonical quotient map. The image of $G/Z_G(a)$ under β is the same as the image of G under α_a , namely Ga . Let V be the Δ -closure of Ga .

Note that $\alpha_a(g) = \alpha_a(h)$ implies that $gag^{-1} = hah^{-1}$ so $h^{-1}g \in Z_G(z)$. Therefore β is an injective Δ -map. Corollary 2.2 implies that $\tau(G/Z_G(a)) \leq \tau(H) < \tau(G)$. Since G is strongly connected, we have $Z_G(a) = G$. ■

2.2.3 Isogeny

Definition 2.14 *1) Let G, H be Δ - k -groups. A Δ - k -morphism $\phi : G \rightarrow H$ is an isogeny if it is surjective and $\tau(\ker \phi) < \tau(G)$.*

2) Two Δ - k -groups H_1, H_2 are Δ - k -isogenous if there exists a Δ - k -group G and isogenies $\phi_i : G \rightarrow H_i$ for $i = 1, 2$.

The notion of isogenous in the present context generalizes the notion of isogenous in the theory of algebraic groups (cf., [37]) where two algebraic groups H_1, H_2 are isogenous if there is an algebraic group G and morphisms $\phi_i : G \rightarrow H_i$ for $i = 1, 2$ with finite kernel.

Example 2.15 Let $G = \mathrm{SL}_6(\mathbb{C})$. The center of G is isomorphic to the cyclic group of order 6. Let C_2 and C_3 be cyclic subgroups of this center of orders 2 and 3 respectively. Let $H_1 = \mathrm{SL}_6(\mathbb{C})/C_2$ and $H_2 = \mathrm{SL}_6(\mathbb{C})/C_3$ and let $\phi_i : G \rightarrow H_i$ be the obvious projections. These show H_1 and H_2 are isogenous in both the category of algebraic groups and the category of Δ -groups. Note that they are not isomorphic since their centers are finite groups of different orders.

Remarks 2.16 1. If G_1 and G_2 are strongly connected Δ -groups and there is an isogeny $\phi : G_1 \rightarrow G_2$, then G_1 and G_2 are isogenous. However, the converse is false, as the above example shows.

2. If $\phi : G_1 \rightarrow G_2$ is an isogeny then equations (1) and (2) imply that $\tau(G_1) = \tau(G_2)$ and $a_\tau(G_1) = a_\tau(G_2)$.

3. The composite of isogenies of strongly connected Δ -groups is an isogeny.

4. If G_1 and G_2 are strongly connected of the same type, then it is easy to see that $G_1 \times G_2$ is strongly connected. Suppose G'_1 and G'_2 are strongly connected and $\phi_i : G_i \rightarrow G'_i$ is an isogeny, $i = 1, 2$. Then, $\phi_1 \times \phi_2 : G_1 \times G_2 \rightarrow G'_1 \times G'_2$, with kernel $\ker \phi_1 \times \ker \phi_2$ is an isogeny.

Proposition 2.17 *Let G_1 and G_2 be Δ - k -groups. The following are equivalent:*

1. *There exists a strongly connected Δ - k -group H and isogenies $\phi_1 : H \rightarrow G_1$ and $\phi_2 : H \rightarrow G_2$:*

$$\begin{array}{ccc} & H & \\ \phi_1 \swarrow & & \searrow \phi_2 \\ G_1 & & G_2 \end{array} .$$

2. *There exists a strongly connected Δ - k -group K and isogenies $\psi_1 : G_1 \rightarrow K$ and $\psi_2 : G_2 \rightarrow K$:*

$$\begin{array}{ccc} G_1 & & G_2 \\ \psi_1 \searrow & & \swarrow \psi_2 \\ & K & \end{array} .$$

Proof. Assume the first statement. Put $H_1 = \phi_1(\ker \phi_2)$, and $H_2 = \phi_2(\ker \phi_1)$. Clearly, $H_1 = \phi_1(\ker \phi_1 \ker \phi_2)$, and $H_2 = \phi_2(\ker \phi_1 \ker \phi_2)$. Therefore,

$$\begin{aligned} G_1/H_1 &= \phi_1(H) / \phi_1(\ker \phi_2) = \phi_1(H) / \phi_1(\ker \phi_1 \ker \phi_2) = H / (\ker \phi_1 \ker \phi_2) \\ G_2/H_2 &= \phi_2(H) / \phi_2(\ker \phi_1) = \phi_2(H) / \phi_2(\ker \phi_1 \ker \phi_2) = H / (\ker \phi_1 \ker \phi_2). \end{aligned}$$

Set $K = H / (\ker \phi_1 \ker \phi_2)$. Since K is the image of the strongly connected group H , it is strongly connected. Since $\tau(\ker \phi_1 \ker \phi_2) < \tau(H)$, the projections $\psi_i : G_i \rightarrow G_i/H_i \simeq K$ are isogenies.

Now assume the second statement. Let $G = \{(g_1, g_2) \in G_1 \times G_2 \mid \psi_1(g_1) = \psi_2(g_2)\}$, that is, the pull-back or fiber product. One sees that the natural projections $\phi_1 : G \rightarrow G_1$ and $\phi_2 : G \rightarrow G_2$ are surjective. This implies that $\tau(G) \geq \tau(G_1)$. The kernel of the ϕ_i 's is contained in $\ker \psi_1 \times \ker \psi_2$ and so $\tau(\ker \phi_1) \leq \tau(\ker \psi_1 \times \ker \psi_2) < \tau(G_1)$. Since $G_1 \simeq G / \ker(\phi_1)$ we have $\tau(G) = \tau(G_1)$ and ϕ_1 is an isogeny. Similar reasoning shows that ϕ_2 is an isogeny. \blacksquare

The following result is a reworking of Theorem 6, [37] in the context of Δ - k -groups.

Proposition 2.18 *Isogeny is an equivalence relation on the set of strongly connected Δ -groups. If G_1 and G_2 are isogenous strongly connected Δ -groups, with $\tau = \tau(G_1) = \tau(G_2)$, then there is a bijective correspondence between the strongly connected Δ -subgroups of G_1 with type τ and those of G_2 with type τ . Let J_1 and K_1 be Δ -subgroups of G_1 corresponding to the Δ -subgroups J_2 and K_2 of G_2 . Then, $J_1 \subseteq K_1$ if and only if $J_2 \subseteq K_2$, and $J_1 \trianglelefteq K_1$ if and only if $J_2 \trianglelefteq K_2$. In this case, K_1/J_1 and K_2/J_2 are isogenous. If G_1, G_2 , and G'_1, G'_2 are pairs of isogenous groups, then $G_1 \times G'_1$ and $G_2 \times G'_2$ are isogenous.*

Proof. Since reflexivity and symmetry follow immediately from the definition, we need only show that isogeny is transitive. So, let $G_1, G_2, H, G_2, G_3, H'$ be strongly connected Δ -groups, and let $\phi_i : H \rightarrow G_i$, $i = 1, 2$, $\phi'_i : H' \rightarrow G_i$, $i = 2, 3$, be isogenies. Then, $H \times H'$ is strongly connected, and $\phi_2 \times \phi'_2 : H \times H' \rightarrow G_2 \times G_2$ is surjective. Let $K = ((\phi_2 \times \phi'_2)^{-1}(\text{Diag } G_2 \times G_2))_0$, where $(\dots)_0$ denotes the strong identity component. Let $\pi_H, \pi_{H'}$, be the projections from $H \times H'$ to H , and to H' . We have the following diagram

$$\begin{array}{ccccc}
& & & & K \\
& & & \nearrow \pi_H & \searrow \pi_{H'} \\
& & H & & H' \\
\phi_1 \swarrow & & \phi_2 \searrow & & \phi'_2 \swarrow & \phi'_3 \searrow \\
G_1 & & G_2 & & G_3
\end{array}$$

We claim that the restrictions of π_H and $\pi_{H'}$ to K are isogenies. Since the composition of isogenies is an isogenies, our claim implies that G_1 and G_3 are isogenous and so the isogeny relation is an equivalence relation.

To prove our claim, first note that (1) implies that $\tau(H \times H') = \tau(H) = \tau(H') = \tau(G_2 \times G_2)$, and $\tau(\ker(\phi_2 \times \phi'_2)) = \tau(\ker \phi_2 \times \ker \phi'_2) < \tau(H \times H')$. If $(h, h') \in K \cap \ker \pi_H$ then $h = 1$ and, since $\phi_2(h) = \phi'_2(h')$, we have $h' \in \ker \phi'_2$. Therefore $\ker \pi_H = 1 \times \ker \phi'_2$ and so $\tau(\ker \pi_H) < \tau(H)$. Similarly, $\tau(\ker \pi_{H'}) < \tau(H')$. We know that $\tau(K) = \tau(G_2) = \tau(H)$. Since $\tau(\ker \pi_H) < \tau(H)$, it follows that $a_\tau(\pi_H(K)) = a_\tau(H)$. Since H is strongly connected, $\pi_H(K) = H$, and, therefore, π_H is an isogeny. Similarly, $\pi_{H'}$ is an isogeny and this shows that isogeny relation is an equivalence relation.

Suppose we have the following isogeny diagram:

$$\begin{array}{ccc} & H & \\ \phi_1 \swarrow & & \searrow \phi_2 \\ G_1 & & G_2 \end{array} .$$

We define the following correspondence between strongly connected Δ -subgroups of G_1 of type $\tau = \tau(G_1)$ and strongly connected Δ -subgroups of G_2 of type $\tau = \tau(G_2)$: Subgroups J_1 of G_1 and J_2 of G_2 correspond if there exists a strongly connected Δ -subgroup P of H such that $\phi_i|_P$ maps P to J_i and is an isogeny, $i = 1, 2$. We claim that given J_1 , there is a unique choice for P , namely, the strong identity component of $\phi_1^{-1}(J_1)$. First note that for any such P , Remark 2.16.2 implies that $\tau(P) = \tau$ and $a_\tau(P) = a_\tau(J_1) = a_\tau(J_2)$. Furthermore any such P must lie in $\phi_1^{-1}(J_1)$ and, since it is assumed to be strongly connected, it must be contained in the strong identity component $(\phi_1^{-1}(J_1))_0$ of $\phi_1^{-1}(J_1)$. Since $\tau(\ker \phi_1) < \tau$ we have that $\tau(\ker \phi_1 \cap \phi_1^{-1}(J_1)) < \tau$ and so ϕ restricted to $(\phi_1^{-1}(J_1))_0$ is an isogeny. In particular $\tau((\phi_1^{-1}(J_1))_0) = \tau$ and $a_\tau((\phi_1^{-1}(J_1))_0) = a_\tau(G_1)$. Since $P \subset (\phi_1^{-1}(J_1))_0$ and both of these groups are strongly connected we have $P = (\phi_1^{-1}(J_1))_0$. A similar argument shows that $P = (\phi_2^{-1}(J_2))_0$. It follows easily that the correspondence is unique.

Now, we shall show that if $J_i, K_i, i = 1, 2$, are corresponding strongly connected Δ -subgroups of G_i of type τ , then

$$J_1 \subseteq K_1 \iff J_2 \subseteq K_2.$$

Let $P, Q, i = 1, 2$, be the unique strongly connected Δ -subgroups of H of type τ such that $\phi_i(P) = J_i$ and $\phi_i(Q) = K_i$. P is the strong identity component of $\phi_i^{-1}(J_i)$ and Q is the strong identity component of $\phi_i^{-1}(K_i)$. $J_i \subseteq K_i$ implies that $\phi_i^{-1}(J_i) \subseteq \phi_i^{-1}(K_i)$. $\tau(P) = \tau(Q) = \tau$. Since P is a strongly connected Δ -subgroup of $\phi_i^{-1}(K_i)$ of the same type as the latter, it follows, as above, that P is contained in its strong identity component Q . The reverse implication is proved the same way. Now suppose that J_1 is normal in K_1 . Then, $\phi_1^{-1}(J_1)$ is normal in $\phi_1^{-1}(K_1)$ and has the same type. By Remark 2.5.4, P is normal in $\phi_1^{-1}(K_1)$. Therefore, P is normal in Q . Since ϕ_2 is surjective, J_2 is normal in K_2 . To finish the proof, we need only show that K_1/J_1 and K_2/J_2 are isogenous. We now define a map $\phi : Q/P \rightarrow K_1/J_1$ by $\phi(q \bmod P) = \phi_1(q) \bmod J_1$ for any $q \in Q$. To see that this is well defined, let $q_1, q'_1 \in Q$, and suppose $q_1 \bmod P = q'_1 \bmod P$. Then, $\phi_1(q_1 - q'_1) \in J_1$. Therefore, $\phi_1(q_1) \bmod J_1 = \phi_1(q'_1) \bmod J_1$. So, the map $\phi : Q/P \rightarrow K_1/J_1$ is well defined. It is clearly a homomorphism. We claim that it is surjective. Let $k_1 \in K_1$. There exists $q \in Q$ such that $\phi_1(q) = k_1$ and so $\phi(q \bmod P) = k_1 \bmod J_1$. A calculation shows that the kernel of ϕ is $[Q \cap \phi_1^{-1}(J_1)]/P$. The strong identity component P of $\phi_1^{-1}(J_1)$ is contained in $Q \cap \phi_1^{-1}(J_1)$. Thus, $\tau([Q \cap \phi_1^{-1}(J_1)]/P) < \tau(\phi_1^{-1}(J_1)) = \tau(Q) = \tau(Q/P)$ since Q is strongly connected. So, $\phi : Q/P \rightarrow K_1/J_1$ is an isogeny. Similar reasoning shows that one can define a map $\psi : Q/P \rightarrow K_2/J_2$ and that it too is an isogeny. \blacksquare

An immediate corollary is

Corollary 2.19 *Let G_1 and G_2 be isogenous strongly connected Δ -groups. Either both are almost simple or neither is.*

Another corollary is

Corollary 2.20 *Let G_1 and G_2 be isogenous strongly connected Δ -groups. Either both are commutative or neither is.*

The proof of Corollary 2.20 depends on the following concept and lemma.

Definition 2.21 *Let G be a Δ - k -group. The differential commutator $D_\Delta(G)$ is the smallest Δ - k -subgroup of G containing the commutator subgroup of G .*

An example of Cassidy (*cf.*, p.111 of [25]) shows that $D_\Delta(G)$ may be strictly larger than the commutator subgroup of G .

Lemma 2.22 *Let G be a Δ - k -group. If G is strongly connected and not commutative, then $\tau(D_\Delta(G)) = \tau(G)$.*

Proof. Let a be in G . Let $c_a : G \rightarrow G$ be the Δ -map that sends x to $axa^{-1}x^{-1}$. If $D_\Delta(G)$ has smaller type than G , then Proposition 2.11 implies that $D_\Delta(G)$ is contained in the center $Z(G)$. In this case we have $axy a^{-1}y^{-1}x^{-1} = axa^{-1}(aya^{-1}y^{-1})x^{-1} = (axa^{-1}x^{-1})(aya^{-1}y^{-1})$ and therefore c_a is a Δ -homomorphism of G into $D_\Delta(G)$. Since G is strongly connected, $c_a(G) = \{1\}$. Thus, $a \in Z(G)$. It follows that G is commutative. ■

Proof of Corollary 2.20: Let K be a Δ -group such that for G_1, G_2 , and K we have the commutative diagram:

$$\begin{array}{ccc} G_1 & & G_2 \\ & \psi_1 \searrow & \swarrow \psi_2 \\ & & K \end{array}$$

with isogenies ψ_i . Assume G_1 is commutative. Then, K is commutative. Suppose G_2 is not commutative. Lemma 2.22 implies that $\tau(D_\Delta(G_2)) = \tau(G_2) = \tau(K)$. Since ψ_2 is an isogeny, $\tau(\psi_2(D_\Delta(G_2))) = \tau(K)$ as well. Since $\psi_2(D_\Delta(G_2)) \subset D_\Delta(K)$ we have that $D_\Delta(K) \neq \{1\}$, a contradiction. ■

We give two examples of infinite families of isogenous, pairwise nonisomorphic Δ -groups.

Example 2.23 Let $\Delta = \{\delta\}$ and \mathbb{U} be a Δ -universal field with Δ -constants C . We shall exhibit an infinite number of almost simple Δ -groups that are isogenous to $\mathbb{G}_a(\mathbb{U})$ but are pairwise non-isomorphic. For each $n \in \mathbb{N}$, let

$$G_n = \left\{ g(a, b) = \begin{pmatrix} a & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a \in \mathbb{U}^*, b \in \mathbb{U}, (\delta a)a^{-1} = \delta^n(b) \right\}.$$

The map $\alpha : G_n \rightarrow \mathbb{G}_a(\mathbb{U})$ given by $\alpha(g(a, b)) = (\delta a)a^{-1}$ is a surjective homomorphism with kernel

$$K_n = \{g(a, b) \mid \delta a = 0, \delta^n(b) = 0\}.$$

Note that G_n has differential type 1 and that K_n has differential type 0 so α is an isogeny and therefore G_n is isogenous to $\mathbb{G}_a(\mathbb{U})$.

Since $\mathbb{G}_a(\mathbb{U})$ is almost simple, Corollary 2.19 implies that G_n is almost simple.

We now claim that for $n \neq m, n, m \in \mathbb{N}$, G_n and G_m are not Δ -isomorphic. To see this, note first that the set of unipotent matrices in any G_n is precisely $\{g(1, b) \mid \delta^n(b) = 0\}$. This is a Δ -group of differential type 0 and typical differential transcendence degree equal to n . The property of being unipotent is preserved under Δ -isomorphism ([11], Proposition 35) so our claim is proved.

Example 2.24 Let \mathbb{U} be a $\Delta = \{\delta_x, \delta_t\}$ differentially closed field. Let C be the field of Δ -constants of \mathbb{U} and $x, t \in \mathbb{U}$ satisfy $\delta_x(x) = \delta_t(t) = 1, \delta_x(t) = \delta_t(x) = 0$. Let

$$H = \{y \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t y - \delta_x^2 y = 0\}$$

be the subgroup of $\mathbb{G}_a(\mathbb{U})$ defined by the Heat Equation. We shall exhibit an infinite number of almost simple Δ -groups that are isogenous to H but not isomorphic to H . In particular, we shall show that if $\lambda \in \mathbb{U}$ is any nonzero solution of $\delta_t \lambda + \delta_x^2 \lambda = 0$, then H is isogenous to

$$H_\lambda = \{z \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t z - \delta_x^2 z + 2 \frac{\delta_x \lambda}{\lambda} \delta_x z = 0\}$$

and that there are an infinite number of λ whose corresponding groups are pairwise non-isomorphic.

The groups H and H_λ have differential type 1 and, from Example 2.10, we know that they are almost simple. An elementary calculation shows that the map $\phi(z) = (1/\lambda)\delta_x z$ is a homomorphism from H_λ onto H . The kernel of ϕ is C . Therefore these groups are isogenous.

For any nonzero $c \in C$,

$$\lambda_c = e^{c^2 t} \cos cx$$

is a solution of $\delta_t \lambda + \delta_x^2 \lambda = 0$. Abusing notation, we define

$$H_c = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u = \delta_{xx} u + 2(c \tan cx) \delta_x u\}$$

Each of these groups is isogenous to \mathbb{G}_a so they are all pairwise isogenous. We shall show that for nonzero $c, d \in C$ with $c/d \notin \mathbb{Q}$, the groups H_c and H_d are not isomorphic. The proof of this claim follows from the special nature of the differential coordinate rings $\mathbb{U}\{H_c\}_\Delta$ and $\mathbb{U}\{H_d\}_\Delta$ of the groups. Note that as rings we may write

$$\mathbb{U}\{H_c\}_\Delta = \mathbb{U}[y, \delta_x y, \dots, \delta_x^n y, \dots] \text{ and } \mathbb{U}\{H_d\}_\Delta = \mathbb{U}[z, \delta_x z, \dots, \delta_x^n z, \dots]$$

where y and its δ_x -derivatives are algebraically independent over \mathbb{U} and a similar statement is true for z . As δ_x -rings these two rings are δ_x -isomorphic to a δ_x -differential polynomial ring $\mathbb{U}\{Y\}_{\delta_x}$ and this will be a key fact in the proof of our claim. What distinguishes these rings is their δ_t -structure, which is given by the defining equations of the groups.

Let us assume (with the aim of arriving at a contradiction) that H_c and H_d are isomorphic as Δ -groups and let $\psi : H_c \rightarrow H_d$ be a differential isomorphism. This isomorphism induces a Δ -isomorphism $\psi^* : \mathbb{U}\{H_d\}_{\Delta} \rightarrow \mathbb{U}\{H_c\}_{\Delta}$. Identifying both $\mathbb{U}\{H_c\}$ and $\mathbb{U}\{H_d\}_{\Delta}$ with the δ_x -ring $\mathbb{U}\{Y\}_{\delta_x}$, the map ψ^* induces a δ_x -automorphism of $\mathbb{U}\{Y\}_{\delta_x}$ and *a fortiori* of the quotient field $\mathbb{U}\langle Y \rangle_{\delta_x}$. The automorphisms of this latter field have been classified by Ritt (as part of his differential Lüroth theorem) in his book ([36], p. 56). He shows that any δ_x -automorphism σ of this latter field must be of the form

$$\sigma(Y) = \frac{aY + b}{cY + d}$$

where $a, b \in \mathbb{U}$. It is not hard to see that if σ furthermore maps $\mathbb{U}\{Y\}_{\delta_x}$ to itself, we must have that $\sigma(Y) = aY + b$ for some $a, b \in \mathbb{U}$. In particular, ψ^* must be of the form $\psi^*(z) = ay + b$ for some $a, b \in \mathbb{U}$ and, since ψ is assumed to be a homomorphism as well, we must have $b = 0$ and so $\psi^*(z) = ay$. The following calculation will show that this leads to a contradiction.

We have that

$$\begin{aligned}\delta_t z &= \delta_{xx}z + 2(c \tan cx)\delta_x z \\ \delta_t y &= \delta_{xx}y + 2(d \tan dx)\delta_x y\end{aligned}$$

Calculating $\delta_t(\psi^*(z))$ and $\psi^*(\delta_t z)$ we have

$$\begin{aligned}\delta_t(\psi^*(z)) &= a\delta_{xx}y + (2ad(\tan dx))\delta_x y + \delta_t ay \\ \psi^*(\delta_t z) &= a\delta_{xx}y + (2\delta_x a + 2ac(\tan cx))\delta_x y + (\delta_{xx}a + 2c(\tan cx)\delta_x a)y\end{aligned}$$

Since $\delta_t(\psi^*(z)) = \psi^*(\delta_t z)$ and $y, \delta_x y, \delta_{xx}y$ are algebraically independent, we must have

$$ad(\tan dx) = \delta_x a + ac(\tan cx) \quad (3)$$

$$\delta_t a = \delta_{xx}a + 2c(\tan cx)\delta_x a \quad (4)$$

Equation (3) implies that

$$a = \alpha \frac{\cos cx}{\cos dx}$$

where $\delta_x \alpha = 0$. Inserting this expression into equation (4) we have

$$\frac{\delta_t \alpha}{\alpha} = c^2 + \frac{2d^2}{\cos^2 dx} - d^2 - \frac{2c^2}{\cos^2 cx}. \quad (5)$$

Since $\delta_x(\frac{\delta_t \alpha}{\alpha}) = 0$, we must have that the right hand side of equation (5) is some constant γ . Evaluating the right hand side at $x = 0$ and $x = \frac{2\pi}{c}$, we have

$$d^2 - c^2 = \gamma \quad (6)$$

$$\left(\frac{2}{\cos^2(2d\pi/c)} - 1\right)d^2 - c^2 = \gamma \quad (7)$$

Note that $\cos(2d\pi/c) \neq 0$ since we are assuming $c/d \notin \mathbb{Q}$. From these equations, we can conclude that $\cos^2(2d\pi/c) = 1$ which implies that $d/c \in \mathbb{Q}$, a contradiction. Therefore, for any set of constants S , whose members are pairwise independent over the integers, this yields a family $\{H_c\}_{c \in S}$ of isogenous Δ -subgroups of \mathbb{G}_a that are pairwise nonisomorphic.

We now turn to another choice of λ : $\lambda = x$. This situation was already considered by Cartan ([9], p.145-146). Letting

$$H_x = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u = \delta_{xx} u - \frac{2}{x} \delta_x u\},$$

Cartan states (without proof) that H and H_x are not ‘‘isomorphes holoédriques’’. We shall show that H and H_x are not isomorphic.

Let us assume that H and H_x are isomorphic as Δ -groups and let $\psi : H \rightarrow H_x$ be a differential isomorphism. Let

$$\mathbb{U}\{H\}_\Delta = \mathbb{U}[y, \delta_x y, \dots, \delta_x^n y, \dots] \text{ and } \mathbb{U}\{H_x\}_\Delta = \mathbb{U}[z, \delta_x z, \dots, \delta_x^n z, \dots]$$

where y and its δ_x -derivatives are algebraically independent over \mathbb{U} and a similar statement is true for z . As above, we have a differential isomorphism $\psi^* : \mathbb{U}\{H_x\}_\Delta \rightarrow \mathbb{U}\{H\}_\Delta$ and we may assume that $\psi^*(z) = ay$. The following calculation will show that this leads to a contradiction. Since ψ^* is also a Δ -isomorphism, we must have that $\psi^*(\delta_t z) = \delta_t(\psi^*(z))$. We have

$$\psi^*(\delta_t z) = a\delta_x^2 y + (2\delta_x a - \frac{2}{x}a)\delta_x y + (\delta_x^2 a - \frac{2}{x}\delta_x a)y \quad (8)$$

$$\delta_t(\psi^*(z)) = a\delta_x^2 y + (\delta_t a)y. \quad (9)$$

Since $\delta_x^2 y, \delta_x y$ and y are algebraically independent over \mathbb{U} , comparing coefficients of like terms in (8) and (9), we have

$$x\delta_x a - a = 0 \quad (10)$$

$$x\delta_x^2 a - 2\delta_x a - x\delta_t a = 0. \quad (11)$$

Equation (10) implies that $a = cx$ where $\delta_x c = 0$. Therefore $x\delta_x^2 a = 0$. This, combined with equation (11), implies that $0 = 2\delta_x a + x\delta_t a = 2c + (\delta_t c)x^2$. Therefore $0 = \delta_x(2c + (\delta_t c)x^2) = 2\delta_t cx$, so $\delta_t c = c = 0$. We therefore have that $a = 0$. This contradicts the fact that ψ^* is

an isomorphism and completes the proof that H and H_x are not Δ -isomorphic.

The Heat Equation $\delta_t y = \delta_x^2 y$ and the related equations $\delta_t z = \delta_x^2 z - \frac{\delta_t \lambda}{\lambda} \delta_x z$ arise from the study of conservation laws and generalized symmetries (cf., [5], [33]), which we now briefly describe. Let L be a homogeneous linear Δ -polynomial. A vector $\mathbf{F} = (T(y), X(y))$ of differential polynomials is called a *conserved vector* if $\nabla \mathbf{F}(u) = \delta_t T(u) + \delta_x X(u) = 0$ for all solutions of $L(y) = 0$. The equation $\delta_t T(u) + \delta_x X(u) = 0$ is called a *conservation law* for $L(y) = 0$. If L is the Heat Equation, then it follows from [33] that the conservation laws for $L(y) = 0$ are generated (as a C -vector space) by conservation laws corresponding to conserved vectors of the form

$$\mathbf{F} = (\lambda y, -\lambda \delta_x y + (\delta_x \lambda) u) \quad (12)$$

where λ satisfies $\delta_y \lambda + \delta_x^2 \lambda = 0$. In [5], Bluman and Kumei introduce the notion of a *potential symmetry* corresponding to each conservation law. The potential symmetry corresponding to the conserved vector (12) is given by the system

$$\begin{aligned} \delta_x z &= \lambda y \\ \delta_t z &= \lambda \delta_x y - (\delta_x \lambda) y. \end{aligned}$$

This latter system defines a Δ -group $G \subset \mathbb{G}_a \times \mathbb{G}_a$ that has been extensively studied for $\lambda = 1$ and $\lambda = x$. Letting π_1 and π_2 be the projections of $\mathbb{G}_a \times \mathbb{G}_a$ onto the first and second coordinates of (y, z) , one can show that $\pi_1(G) = H$, $\pi_2(G) = H_\lambda$ and the π_i are isogenies. This gives another proof that the groups H and H_λ are isogenous.

Further studies of the algebraic and model theoretic structure of the solutions of the Heat Equation and related differential groups can be found in the thesis [42] of Sonat Suer.

2.3 Jordan-Hölder Theorem

We can now state and prove the following analogue of the Jordan-Hölder Theorem (cf., Theorem 6, [37]).

Theorem 2.25 *Let G be a strongly connected Δ - k -group with $\tau(G) = \tau \neq -1$ (i.e., $G \neq \{1\}$). There exists a normal sequence $G = G_0 \triangleright G_1 \triangleright \dots \triangleright G_r = \{1\}$ of Δ - k -subgroups such that for each $i = 0, \dots, r-1$*

1. G_i is a strongly connected group with $\tau(G_i) = \tau$,
2. $a_\tau(G_0) > a_\tau(G_1) > \dots > a_\tau(G_{r-1})$, and
3. G_i/G_{i+1} is almost simple.

If $G = H_0 \triangleright H_1 \triangleright \dots \triangleright H_s = \{1\}$ is another normal sequence as above, then $r = s$ and there exists a permutation π of $\{0, \dots, r\}$ such that $G_{\pi(i)}/G_{\pi(i)+1}$ is k -isogenous to H_i/H_{i+1} .

Proof. If G is not almost simple, there exists a normal proper Δ - k -subgroup H with $\tau(H) = \tau(G) = \tau$. Since G is strongly connected, we have that $a_\tau(H) < a_\tau(G)$. Among such subgroups let H have largest $a_\tau(H)$ and let G_1 be the strong identity component of H . Note that, by definition, $\tau(G_1) = \tau(H)$. Since G_1 is a characteristic subgroup of H , it is normal in G . We have that $a_\tau(G) = a_\tau(G/G_1) + a_\tau(G_1)$ so $a_\tau(G/G_1) \neq 0$. Therefore $\tau(G/G_1) = \tau$. For any normal Δ - k -subgroup K of G with $K \supset G_1$, maximality of $a_\tau(H) = a_\tau(G_1)$ implies that $a_\tau(K) = a_\tau(G_1)$ so $\tau(K/G_1) < \tau$. Therefore G/G_1 is almost simple. Proceeding by induction on $a_\tau(G_i)$ yields the required sequence of Δ - k -subgroups.

To prove the last sentence in the Theorem we proceed as in the proof of the Jordan-Hölder Theorem as given in ([27], Ch. 1,§3). For each $i = 0, \dots, r-1$, $j = 0, \dots, s-1$ let

$$G_{i,j} = G_{i+1}(H_j \cap G_i) .$$

The G_{ij} give a refinement of the sequence of G_i 's:

$$G = G_{0,0} \triangleright G_{0,1} \triangleright \dots \triangleright G_{0,s-1} \triangleright G_1 = G_{1,0} \triangleright G_{1,2} \triangleright \dots \triangleright \{0\}$$

Similarly, define for each $j = 0, \dots, s-1$, $i = 0, \dots, r$,

$$H_{j,i} = H_{j+1}(G_i \cap H_j) .$$

This yields a refinement of the H_j . By Lemma 3.3 of [27] we have that

$$G_{i,j}/G_{i,j+1} \text{ is isomorphic to } H_{j,i}/H_{j,i+1}$$

(the isomorphism obviously being an isomorphism of Δ - k -groups.) We now claim that for any $i = 0, \dots, r-1$ there is at exactly one $j, 0 \leq j \leq s-1$ such that $\tau(G_{i,j}/G_{i,j+1}) = \tau$ and that for this value of j we have that $G_{i,j}/G_{i,j+1}$ is isogenous to G_i/G_{i+1} (and a similar statement will hold for the sequence of $H_{j,i}$). This and the previous sentence then imply that $r = s$ and the final statement of the Theorem is true.

For each $i = 0, \dots, r-1$, have that

$$a_\tau(G_i) = a_\tau(G_{i,0}/G_{i,1}) + \dots + a_\tau(G_{i,s-1}/G_{i+1,0}) + a_\tau(G_{i+1})$$

(recall that $G_{i+1,0} = G_{i+1}$). Since $a_\tau(G_i) > a_\tau(G_{i+1})$ we have that some $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$ and so $\tau(G_{i,j}/G_{i,j+1}) = \tau$. Equation (1) furthermore implies that $\tau(G_{i,j}/G_{i+1}) = \tau$. Let j be the smallest integer such that $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$.

We will show that $G_i = G_{i,0} = G_{i,1} = \dots = G_{i,j}$. Note that $\tau = \tau(G_{i,j}/G_{i,j+1}) \leq \tau(G_{i,j}/G_{i+1}) \leq \tau(G_i/G_{i+1}) = \tau$. Therefore $\tau(G_{i,j}/G_{i+1}) = \tau$. Furthermore, $\tau = \tau(G_i/G_{i+1}) \geq \tau(G_{i,t}/G_{i+1}) \geq \tau(G_{i,j}/G_{i+1}) = \tau$ so $\tau(G_{i,t}/G_{i+1}) = \tau$ for any $t = 0, 1, \dots, j-1$. Since $a_\tau(G_{i,t}/G_{i,t+1}) = 0$ for $t = 0, 1, \dots, j-1$, we have $\tau(G_{i,t}/G_{i,t+1}) < \tau$. Since G_i/G_{i+1} is strongly connected, $G_{i,1}/G_{i+1} \triangleleft G_i/G_{i+1}$ and $\tau((G_{i,0}/G_{i+1})/(G_{i,1}/G_{i+1})) = \tau(G_{i,0}/G_{i,1}) < \tau$, we have $G_{i,0} = G_{i,1}$. Continuing in a similar fashion, we see $G_{i,0} = \dots = G_{i,j}$.

From this discussion we can conclude that $G_{i,j} = G_i$ and $G_{i,j+1} \triangleleft G_i$. Let $\pi : G_i/G_{i+1} \longrightarrow G_{i,j}/G_{i,j+1}$ be the canonical projection. The kernel is a proper normal subgroup so, again by almost simplicity, it must have smaller type. Therefore G_i/G_{i+1} is isogenous to $G_{i,j}/G_{i,j+1}$.

We now turn to the uniqueness claim. Assume that $\tau(G_{i,t}/G_{i,t+1}) = \tau$ for some $t > j$. We would then have $\tau(G_{i,t}/G_{i+1}) = \tau$ and so $\tau(G_{i,j+1}/G_{i+1}) = \tau$. Since G_i/G_{i+1} is almost simple and $G_{i,j+1} \triangleleft G_{i,j} = G_i$, we must have $G_{i,j+1} = G_{i,j}$. contradicting $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$. ■

Remark 2.26 As is clear from the above proof the analogue of the results of Zassenhaus and Schreier (cf., [27], Chapter I,§3, Lemma 3.3 and Theorem3.4) stating that two normal sequences have isomorphic refinements is true in the context of Δ -groups. This follows from the fact that the isomorphism theorems are true in this context.

Corollary 2.27 *Let G and H be isogenous strongly connected Δ -groups with normal sequences $G = G_0 \triangleright G_1 \triangleright \dots \triangleright G_r = \{1\}$ and $H = H_0 \triangleright H_1 \triangleright \dots \triangleright H_s = \{1\}$ as in Theorem 2.25. Then $r = s$ and there exists a permutation π of $\{0, \dots, r\}$ such that $G_{\pi(i)}/G_{\pi(i)+1}$ is k -isogenous to H_i/H_{i+1} .*

Proof. Immediate from Theorem 2.25 and Proposition 2.18. ■

Remarks 2.28 1. One can consider groups G that are not strongly connected and apply Theorem 2.25 first to the strong identity component G_0 , then to the strong identity component of G/G_0 , etc. This will yield a more complete Jordan-Hölder Theorem.

2. In Example 2.30 below, we show that the uniqueness up to isogeny in Theorem 2.25 cannot be strengthened to give uniqueness up to isomorphism.

3. In [3], Baudisch presents a Jordan-Hölder Theorem for superstable groups G . He shows that such a group has a sequence of definable subgroups $(1) \triangleleft H_0 \triangleleft H_1 \triangleleft \dots \triangleleft H_r \triangleleft G$ such that H_{i+1}/H_i is infinite and either abelian or simple modulo a finite center and H_r is of finite index in G . Although this is of a similar nature as our result, his result does not address the structure of abelian groups nor does he address the question of uniqueness. A key step in understanding the relationship between Baudisch's result and ours would be to understand the relationship between U-rank and the type and typical differential transcendence degree (cf., [42]).

Example 2.29 We now return to the example of Blumberg mentioned in the Introduction:

$$L = \delta_x^3 + x\delta_x^2\delta_y + 2\delta_x^2 + 2(x+1)\delta_x\delta_y + \delta_x + (x+2)\delta_y \quad (13)$$

$$= (\delta_x + 1)(\delta_x + 1)(\delta_x + x\delta_y) \quad (14)$$

$$= (\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y)(\delta_x + 1) . \quad (15)$$

We shall show how this example fits into the above discussion.

Let $\Delta = \{\delta_x, \delta_y\}$ and k a differentially closed Δ -field. Recall from Remark 2.3.2 that the gauge of a Δ - k -set X is the pair $(\tau(X), a_\tau(X))$ and order these lexicographically. Let R be any nonzero homogeneous linear differential polynomial in δ_x and δ_y of order d and let $G_R \subset K$ be the solutions of $R = 0$. In [24], Kolchin gives a general method for computing differential dimension polynomials $\omega_{\eta/k}(t)$ in terms of a characteristic set for the defining ideal I of η over k . When this ideal is generated by linear homogeneous differential polynomials $\{L_1(Y), \dots, L_r(Y)\}$ in one variable, one does not need this full machinery. In this restricted case, if $\tilde{L}_1, \dots, \tilde{L}_s$ is a reduced Gröbner basis (with respect to a graded monomial order) of the left ideal $\langle L_1, \dots, L_r \rangle$ in $k[\delta_1, \dots, \delta_m]$, then $\tilde{L}_1(Y), \dots, \tilde{L}_s(Y)$ is a characteristic set of I^1 . The differential dimension polynomial can be computed from the leading terms of this set (with respect to the same monomial order, which is orderly; see Lemma 16, Ch. 0.17 and Theorem 6, Ch. II.12 of [24]). For $\Delta = \{\delta_x, \delta_y\}$ and k a differentially closed Δ -field, this can be made explicit. Identifying the monomial $\delta_x^i \delta_y^j$, we use the order $(i, j) > (i', j')$ if $i + j > i' + j'$ or $i + j = i' + j'$ and $i > i'$. If $E = \{(i_1, j_1), \dots, (i_t, j_t)\}$ represent the leading terms of a reduced Groebner basis of a left ideal of $k[\delta_x, \delta_y]$, we can assume that $i_1 < i_2 < \dots < i_t$ and $j_1 > j_2 > \dots > j_t$. Let

$$W = \{(i, j) \in \mathbb{N}^2 \mid i \geq i_1, j \geq j_t, (i, j) \notin \cup_{(i', j') \in E} ((i', j') + \mathbb{N}^2)\}$$

and let $d = i_1 + j_t$. Geometrically, W is the set of points in the quarter-plane $(i_1, j_t) + \mathbb{N}^2$ below the “stairs” $\cup_{(i', j') \in E} ((i', j') + \mathbb{N}^2)$. Lemma 16, Ch. 0.17 and Theorem 6, Ch. II.12 of [24], imply that

$$\omega(s) = \binom{s+2}{2} - \binom{s-d+2}{2} + |W| = ds + \frac{3d-d^2}{2} + |W|.$$

For example, if the leading monomial of an operator R is $\delta_x^i \delta_y^j$ with $i + j = d$, then the differential dimension polynomial of the differential ideal $\langle R(Y) \rangle$ is $\omega(s) = \binom{s+2}{2} - \binom{s-d+2}{2} = ds + \frac{3d-d^2}{2}$. Therefore the gauge of G_R is $(1, d)$. If $H \subset G_R$ is a proper Δ -subgroup, then the defining differential ideal J contains a linear differential polynomial such that the leading term of its associated operator is not divisible by the leading term of R . The stairs corresponding to leading terms of a characteristic set of the ideal of J strictly contain $(1, d) + \mathbb{Z}^2$. One sees from this that the associated differential dimension polynomial either has degree 0 or leading coefficient less than d . This implies that either the type of H is smaller than the type of G_R or the typical transcendence degree of H is smaller than d . In particular, this implies that G_R is strongly connected.

The factorization in (14) yields a sequence of subgroups of G_L as follows. Let G_1 be the solutions of $(\delta_x + 1)(\delta_x + x\delta_y)(Y) = 0$ and let G_2 be the solutions of $(\delta_x + x\delta_y)(Y) = 0$. As noted above, each of these groups is strongly connected. The quotients G/G_1 and G_1/G_2

¹[8], p.84-85; since the coefficients of the $L_i(Y)$ lie in a field, we can assume that the coefficient of the highest term is 1. This implies that the leader and separant are 1 and one sees that the algorithm for producing a reduced Groebner basis in $k[\delta_1, \dots, \delta_n]$ and the algorithm for producing a characteristic set both lead to the same result.

are both isomorphic to the group defined by $(\delta_x + 1)(Y) = 0$. Therefore each quotient in the sequences $G \supset G_1 \supset G_2 \supset \{0\}$ is strongly connected and has gauge $(1, 1)$. Let H be one of these quotients and K a proper subgroup of H . If the gauge of H is $(1, 1)$, then $\tau(K/H) = 0$, contradicting the fact that H is strongly connected. Therefore $a_1(K) = 0$ and so $\tau(K) = 0$. Therefore these quotients are almost simple and so the sequence satisfies the conclusion of the Theorem.

The factorization in (15) yields a sequence of groups as follows. Let H_2 be the group defined by $(\delta_x + 1)(Y) = 0$. We then have the sequence $G \supset H_2 \supset \{0\}$. The Theorem implies that the group G/H_2 cannot be almost simple. In fact the proof of the Theorem tells us how to refine this sequence. Let $H_1 = H_2 + G_2$ (note that this sum is direct since the groups have a trivial intersection). Lemma 2.6 implies this group is strongly connected. The ideal defining this group is the intersection of the ideals defining each of the summands, i.e.

$$\langle (\delta_x + 1)(Y) \rangle \cap \langle (\delta_x + x\delta_y)(Y) \rangle = \langle L_1(Y), L_2(Y) \rangle$$

where

$$\begin{aligned} L_1 &= x\delta_x^2\delta_y + x^2\delta_x\delta_y^2 - \delta_x^2 - \delta_x\delta_y + x^2\delta_y^2 - \delta_x - \delta_y - x\delta_y \\ L_2 &= \delta_x^3 - x^2\delta_x\delta_y^2 + 3\delta_x^2 + 2x\delta_x\delta_y + 3\delta_x\delta_y - x^2\delta_y^2 + 2\delta_x + 2x\delta_y + 3\delta_y \end{aligned}$$

form a Gröbner basis (with respect to an orderly ranking with $\delta_x > \delta_y$) of this left ideal (cf., Example 5 of [18]. The above polynomials were calculated using the `Ore_algebra` and `Groebner` packages in Maple and differ slightly from those presented in [18]). One can use this representation to see directly that H_1 is strongly connected. Looking at leading terms and using Lemma 16 of Ch. 0.17 of [24], we see that the gauge is $(1, 2)$. Furthermore, we see that if this ideal is contained in a larger ideal J of the same gauge, then J must be generated by a single operator of order 2. Using the techniques of [18] one can show that L_1 and L_2 do not have a common right factor of order 2 so this is impossible. Therefore H_1 is strongly connected of gauge $(1, 2)$. This yields the sequence $G \supset H_1 \supset H_2 \supset \{0\}$ where successive quotients are almost simple. We will now compare successive quotients of the two sequences:

H_2 : We will construct an isogeny between G/G_1 and H_2 . Consider the operator $T = (\delta_x + 1)(\delta_x + x\delta_y)$. This operator maps G onto G/G_1 , which is naturally isomorphic to the solution space of $(\delta_x + 1)(Y) = 0$, an almost simple (and therefore strongly connected) group. The group $T(H_2)$ must therefore be either all of G/G_1 or of rank less than 1 (and therefore $\{0\}$ since H_2 is strongly connected). If $T(H_2) = 0$, then H_2 would be annihilated by both $(\delta_x + 1)(\delta_x + x\delta_y)$ and $(\delta_x + 1)$, but this is impossible since the ideal $\langle (\delta_x + 1)(\delta_x + x\delta_y), (\delta_x + 1) \rangle$ contains δ_y and so is of rank 0. Therefore $T(H_2) = G/G_1$ and T defines an isogeny. In fact, since $H_2 = \{fe^{-x} \mid f_x = 0\}$ one can show that T is an isomorphism of H_2 onto G/G_1 .

H_1/H_2 : This group is $(H_2 + G_2)/H_2$ which is isomorphic to $G_2/(H_2 \cap G_2)$. The group $\overline{H_2 \cap G_2}$ is precisely the common zeroes of $(\delta_x + 1)(Y) = 0$ and $(\delta_x + x\delta_y)(Y) = 0$, that is,

just the element 0. Therefore G_2 is isogenous (in fact, isomorphic) to H_1/H_2 .

G/H_1 : We will show that G/H_1 is isogenous to G_1/G_2 . To see this, consider the map $\overline{G_1} \hookrightarrow G \rightarrow G/H_1$. The kernel of this map is $G_1 \cap H_1$, that is those solutions of $(\delta_x + 1)(\delta_x + x\delta_y)(Y) = 0$ that are the sum of a solution of $(\delta_x + 1)(Y) = 0$ and a solution of $(\delta_x + x\delta_y)(Y) = 0$. This clearly contains G_2 but is not all of G_1 , so we have a homomorphism $G_1/G_2 \rightarrow G/H_1$, that is not the zero map. Since G_1/G_2 is almost simple, the image of this map has gauge $(1, 1)$ and the kernel has rank at most 0. Since G/H_1 is almost simple, the image must be all of G/H_1 . Therefore we have an isogeny of G_1/G_2 onto G/H_1 . This isogeny has a nontrivial kernel; it is the solutions of $(\delta_x + 1)(Y) = 0, \delta_y(Y) = 0$. Although *this isogeny* has a nontrivial kernel, we claim that G_1/G_2 is nonetheless isomorphic to G/H_1 . To show this it is enough to show the following. Let K_1 be the solutions of $(\delta_x + 1)(Y) = 0$ and K_2 be the solutions of $(\delta_x + 1)(Y) = 0, \delta_y(Y) = 0$. We have that G_1/G_2 is isomorphic to K_1 and G/H_1 is isomorphic (by the above) to K_1/K_2 . The homomorphism $\delta_y : \mathbb{G}_a \rightarrow \mathbb{G}_a$ maps K_1 surjectively to K_1 and its kernel in K_1 is K_2 . Therefore K_1/K_2 is isomorphic to K_1 and our claim is proved.

Example 2.30 We shall use the groups defined in Example 2.24 to construct an example of a Δ -group having two normal sequences with isogenous but non-isomorphic quotients.

Let $\Delta = \{\delta_x, \delta_t\}$ and $k = \mathbb{U}$ and C be the Δ -constants of \mathbb{U} . Let K_1 be the solutions of $(\delta_x^2 - \delta_t)(Y) = 0$ and K_2 be the solutions of $(\delta_x - x\delta_t)(Y) = 0$. Let $G = K_1 + K_2$. Lemma 2.6 implies that this group is strongly connected. In fact its annihilating ideal is the intersection of $\langle \delta_x^2 - \delta_t \rangle$ and $\langle \delta_x - x\delta_t \rangle$ in $k\{\delta_x, \delta_t\}$ which is $\langle L \rangle$ where

$$\begin{aligned} L &= x\delta_x^3 - x^2\delta_x^2\delta_t - 2\delta_x^2 - x\delta_x\delta_t + x^2\delta_t^2 + 2\delta_t \\ &= (x\delta_x - x^2\delta_t - 2)(\delta_x^2 - \delta_t) \\ &= (x\delta_x^2 - x\delta_t - 2\delta_x)(\delta_x - x\delta_t). \end{aligned} \tag{16}$$

One sees directly that G is strongly connected since a subgroup defined by the vanishing of a single operator must be strongly connected. Consider the two sequences of subgroups: $G \supset K_1 \supset \{0\}$ and $G \supset K_2 \supset \{0\}$. We claim that each quotient in either of these sequences is almost simple and, while K_1 is isogenous to G/K_2 , it is not isomorphic to either G/K_2 or K_2 .

The almost simplicity of K_1 and K_2 follows from the observation in Example 2.10. In particular any quotient of K_1 or K_2 by a Δ -subgroup of smaller type will also be almost simple. Therefore $G/K_1 \simeq K_2/(K_1 \cap K_2)$ and $G/K_2 \simeq K_1/(K_1 \cap K_2)$ are almost simple. Clearly K_1 is isogenous to G/K_2 and K_2 is isogenous to G/K_1 .

We now claim that K_1 is not isomorphic to either G/K_2 or K_2 . Note that K_2 has typical differential transcendence degree 2 while K_1 has typical differential transcendence degree 1, so these two groups cannot be isogenous. From (16), one sees that G/K_2 is isomorphic to the group of solutions of $(x\delta_x^2 - x\delta_t - 2\delta_x)Y = 0$. This is the group H_x of Example 2.24

and K_1 is the group H of the Heat Equation in this latter example. As we have shown in Example 2.24 these groups are not isogenous.

3 Almost Simple Groups

In this section we will develop some facts concerning almost simple Δ -groups. We will use the terms Δ -group, Δ -subgroup, etc to mean that the objects are all defined over \mathbb{U} . Similarly almost simple, irreducible, etc. refer to these properties with respect to \mathbb{U} , a differentially closed Δ -field.

3.1 Quasisimple Linear Algebraic Groups are Almost Simple Δ -groups

We begin with the following lemma.

Lemma 3.1 *Let (\mathbb{U}, Δ) be a differentially closed differential field, $\Delta' \subset \Delta$ and G a non-commutative connected algebraic subgroup of $\mathrm{GL}_n(\mathbb{U})$ with finite center, that is defined over the field C' of Δ' -constants of \mathbb{U} . Then the Δ' -subgroup $G(C')$ of G equals its own normalizer in G .*

Proof. First note that since C' is algebraically closed, $G(C')$ is Zariski dense in G and is non-commutative and connected. Let N be the normalizer of $G(C')$ in G . For $g \in N$ and $c \in G(C')$, we have that $g^{-1}cg = c_1 \in G(C')$. For any $\delta' \in \Delta'$, differentiating this latter equation yields $-g^{-1}\delta'g \cdot g^{-1}cg + g^{-1}c\delta'g = 0$ and therefore

$$c^{-1}\ell\delta'g c = \ell\delta'g$$

where $\ell\delta'g = \delta'g \cdot g^{-1}$ is called the logarithmic derivative of g . It is well known that $\ell\delta'g \in \mathfrak{g}$, the lie algebra of G (see [24], Ch. V.22 or [30], Sec.5, for an exposition of properties of the logarithmic derivative in the linear case). Since $G(C')$ is Zariski dense in G , this latter equation implies that every element of the \mathbb{U} -span W of $\ell\delta'g$ is left invariant under the adjoint action of G on \mathfrak{g} . Corollary 3.2 of [21] implies that W is annihilated by the adjoint action of \mathfrak{g} on \mathfrak{g} , that is W lies in the center of \mathfrak{g} . Theorems 3.2 and 4.2 in Chapter 4 of [21] imply that the center of \mathfrak{g} is the lie algebra of the center of G . Since the center of G is finite, we must have that $\ell\delta'g = 0$, that is $\delta'g = 0$ and so $g \in G(C')$. ■

Proposition 3.2 *Let (\mathbb{U}, Δ) be a differentially closed differential field with Δ -constants C and let $G \subset \mathrm{GL}_n(\mathbb{U})$ be a non-commutative quasisimple linear algebraic group defined over C . If N is a proper normal Δ -subgroup of G , then N is finite. In particular, if G is quasisimple (resp., simple) as an algebraic group, then it is quasisimple (resp., simple) as a Δ -group and therefore in either case is almost simple.*

Proof. Let N be a proper normal Δ -subgroup of G and N^0 its identity component (in the differential algebraic topology). N^0 is again normal in G . Let $H \subset G$ be the Zariski closure of N^0 . One sees that H is connected in the Zariski topology and is again normal in G . We wish to show that H is finite. If not, then, since G is quasisimple as an algebraic group, we must have that $H = G$. This implies that N^0 is a Zariski dense Δ -subgroup of G , connected in the differential algebraic topology. Theorem 19 of [14] states that there exists a finite set of derivations Δ' that are \mathbb{U} -linear combinations of the elements of Δ such that N^0 is conjugate in $\mathrm{GL}_n(\mathbb{U})$ to $G(C')$ where $C' = \{c \in \mathbb{U} \mid \delta'(c) = 0, \forall \delta' \in \Delta'\}$. Expanding Δ if necessary we may assume $\Delta' \subset \Delta$ (this does not change the hypotheses or conclusions). Lemma 3.1 implies that N^0 is its own normalizer so $N^0 = G$, contradicting the fact that N is a proper subgroup of G . ■

3.2 Almost Simple Linear Δ -Groups

We now restrict ourselves to *linear* differential algebraic groups. In this situation, we are able to derive additional properties of almost simple groups and, in the case of ordinary differential fields, give a strong classification of almost simple linear Δ -groups.

Proposition 3.3 *Let G be a non-trivial commutative simple linear Δ -group. Then either G is finite of prime order or is isomorphic to $\mathbb{G}_a(C)$.*

Proof. Assume $G \subset \mathrm{GL}_n(\mathbb{U})$ for some positive integer n . If G is finite then it clearly must be of prime order. Let us now assume that G is infinite. Since the identity component G^0 is a normal Δ -subgroup of finite index, we must have that $G = G^0$ so G is connected. Therefore \overline{G} , the Zariski closure of G in $\mathrm{GL}_n(\mathbb{U})$, is also connected and commutative. Since \mathbb{U} is algebraically closed, we may assume that $\overline{G} = \mathbb{G}_m^p(\mathbb{U}) \times \mathbb{G}_a^q(\mathbb{U})$ for some non-negative integers p, q and that G is Zariski dense in $\mathbb{G}_m^p(\mathbb{U}) \times \mathbb{G}_a^q(\mathbb{U})$. Assume that $p > 0$ and let π be the projection of \overline{G} onto some factor of $\mathbb{G}_m^p(\mathbb{U})$. We have that $\pi(G(\mathbb{U}))$ is Zariski dense in $\mathbb{G}_m(\mathbb{U})$ and so by Proposition 31 of [11], $\pi(G(\mathbb{U}))$ contains $\mathbb{G}_m(C)$. This would imply that G would have a proper nontrivial normal Δ -subgroup. Therefore $p = 0$ and $\overline{G} = \mathbb{G}_a^q(\mathbb{U})$ for some q . Let $0 \neq a = (a_1, \dots, a_q) \in G$. The subgroup Ca is Δ -closed and Δ -isomorphic to $\mathbb{G}_a(C)$. Therefore G must be isomorphic to $\mathbb{G}_a(C)$. ■

In the theory of linear algebraic groups, one knows that the commutator subgroup of linear algebraic group is closed in the Zariski topology and therefore that a quasisimple linear algebraic group is perfect, that is, it equals its commutator subgroup (in fact, every element of a semisimple algebraic group is a commutator [35]). As we have already mentioned following Definition 2.21, in contrast to the algebraic case, the commutator subgroup of a differential algebraic group need not be closed in the Kolchin topology (Ch. IV.5, [25]). Nonetheless, we have the following proposition

Proposition 3.4 *Let G be an infinite almost simple linear Δ -group. Then either G is commutative or $D_\Delta(G) = G$.*

Proof. Assume G is not commutative and suppose that $D_\Delta(G) \neq G$. Since $D_\Delta(G)$ is a normal subgroup of G , Corollary 2.12 implies that $D_\Delta(G)$ lies in the center $Z(G)$ of G . Therefore $G/Z(G)$ is commutative. Since every proper normal subgroup of G is central, the group $G/Z(G)$ is simple. Proposition 3.3 implies that $G/Z(G)$ is isomorphic to $\mathbb{G}_a(C)$. In particular, $0 = \tau(G/Z(G)) = \tau(G)$. This implies that $\tau(Z(G)) = -1$, that is, $Z(G)$ is finite. Since $D_\Delta(G)$ is connected we must have that $D_\Delta(G)$ is trivial, so G is commutative, a contradiction. \blacksquare

We do not have an example of a non-commutative almost simple linear differential algebraic group whose commutator subgroup is not closed in the Kolchin topology. In fact, as we show below in Proposition 3.8, non-commutative almost simple linear Δ -groups of *type at most 1* are equal to their commutator subgroups.

We derive one more property of the center. Let $G \subset \mathrm{GL}_n$ be a Δ -group and \overline{G} its Zariski closure. Let \overline{G}_u and \overline{G}_r be the unipotent and (solvable) radical of \overline{G} . One can easily show that $\overline{G}_u \cap G$ is the unique maximal normal unipotent differential algebraic subgroup of G and $(\overline{G}_r \cap G)^0$ is the unique maximal connected solvable Δ -subgroup of G . We shall denote $\overline{G}_u \cap G$ by G_u and $(\overline{G}_r \cap G)^0$ by G_r and refer to these as the unipotent radical and radical respectively.

Proposition 3.5 *Let G be an infinite almost simple linear Δ -group.*

1. *If G is non-commutative, then the unipotent radical G_u of G equals the identity component $(Z(G))^0$ of the center of G .*
2. *If G is commutative then either G is isomorphic to $G_m(C)$ or G is isogenous to an almost simple subgroup of $G_a(\mathbb{U})$.*

Proof. Let $G \subset \mathrm{GL}_n$ and let \overline{G} be the Zariski closure of G .

1. Assume that G is a non-commutative group. The commutator subgroup $(\overline{G}, \overline{G})$ of \overline{G} is a Zariski closed subgroup of \overline{G} and, *a fortiori*, is Kolchin closed. Therefore it contains $D_\Delta(G) = G$. Since $(\overline{G}, \overline{G})$ is Zariski closed, it must also contain \overline{G} . Therefore $\overline{G} = (\overline{G}, \overline{G})$ is perfect.

The group \overline{G} has a Levi decomposition

$$\overline{G} = \overline{G}_u \rtimes P$$

where P is a reductive algebraic group. Since \overline{G} is perfect, P is perfect. The solvable radical $R(P)$ of P has finite intersection with $[P, P]$ (Lemma, p. 125 [22]) so $R(h)$ is finite.

Therefore P must be semisimple. This implies that the radical \overline{G}_r of \overline{G} equals \overline{G}_u . Therefore, $G_r = G_u$. This implies that $Z(G)^0 \subset G_u$. Since $D_\Delta(G) = G$, we have that G_u is a proper normal Δ -subgroup of G and so is central. Therefore $Z(G)^0 = G_u$.

2. Assume G is commutative so \overline{G} is also commutative. Therefore $\overline{G} \simeq \mathbb{G}_m(\mathbb{U})^s \times (\mathbb{G}_a(\mathbb{U}))^t$ for some integers s, t . The map $l\Delta : \mathbb{G}_m(\mathbb{U}) \rightarrow (\mathbb{G}_a(\mathbb{U}))^m$ given by $l\Delta(u) = (\delta_1(u)u^{-1}, \dots, \delta_m(u)u^{-1})$ is a Δ -homomorphism so we have a Δ -homomorphism

$$\phi = (l\Delta^s, id^t) : G \rightarrow (\mathbb{G}_a(\mathbb{U}))^{ms} \times (\mathbb{G}_a(\mathbb{U}))^t.$$

The kernel of this map is $G \cap (\mathbb{G}_m(C))^s \subset (\mathbb{G}_m(\mathbb{U}))^s \subset (\mathbb{G}_m(\mathbb{U}))^s \times (\mathbb{G}_a(\mathbb{U}))^t$. Let $\pi : (\mathbb{G}_a(\mathbb{U}))^{ms} \times (\mathbb{G}_a(\mathbb{U}))^t \rightarrow \mathbb{G}_a(\mathbb{U})$ be a projection onto one of the $\mathbb{G}_a(\mathbb{U})$ factors. Since G is almost simple, the group $\pi \circ \phi(G) \subset \mathbb{G}_a(\mathbb{U})$ is a nontrivial almost simple Δ -group or is the trivial group. If it is a nontrivial almost simple group, then the map $\pi \circ \phi$ is an isogeny since its kernel in G is a proper subgroup. If this group is trivial for all projections π , then G must be a subgroup of $\mathbb{G}_m(C)^s$. This implies that G is a connected algebraic subgroup of $\mathbb{G}_m(C)^s$. Such groups are isomorphic to $(\mathbb{G}_m(C))^\ell$ for some ℓ . Since G is almost simple, we have that $\ell = 1$. ■

Remarks 3.6 1. We cannot strengthen the conclusion of part 2. above. Let (k, δ) be an ordinary differential field and let $G \subset \mathbb{G}_m \times \mathbb{G}_a$ be the graph of the homomorphism $l\delta : \mathbb{G}_m \rightarrow \mathbb{G}_a$. G contains the group $\mathbb{G}_m(C) \times 0$ and so cannot be unipotent. Therefore G is isogenous to $\mathbb{G}_a(\mathbb{U})$ (via the projection onto the second factor) but is not isomorphic to any subgroup of $\mathbb{G}_a(\mathbb{U})$.

2. For information concerning commutative unipotent differential algebraic groups, see [13]. Nonetheless, we have no general classification of almost simple proper Δ -subgroups of $\mathbb{G}_a(\mathbb{U})$ (except in the ordinary differential case; see below).

The above propositions yield the following Theorem. We shall denote by $\mathbb{U} \cdot \Delta$ the \mathbb{U} span of Δ . The elements of $\mathbb{U} \cdot \Delta$ are derivations on \mathbb{U} .

Theorem 3.7 *Let G be an almost simple linear Δ -group of type τ .*

1. *If G is non-commutative, then*

(a) $G = D_\Delta(G)$,

(b) $G_u = (Z(G))^0$, and

(c) *There exists a finite commuting subset $\Delta^* \subset \mathbb{U} \cdot \Delta$, $|\Delta^*| = m - \tau$, such that $G/Z(G) \simeq H(C_{\Delta^*})$ where H is a simple algebraic group defined over \mathbb{Q} and $C_{\Delta^*} \subset \mathbb{U}$ is the field of Δ^* -constants.*

2. *If G is commutative, then G is either isogenous to a Δ -subgroup of $\mathbb{G}_a(\mathbb{U})$ or is isomorphic to $\mathbb{G}_m(C)$.*

Proof. Statement 1(a) follows from Proposition 3.4. Statement 1(b) follows from Proposition 3.5.1. Statement 2. follows from Proposition 3.5.2. To verify statement 1(c), note that Corollary 2.13 states that $G/Z(G)$ is a simple Δ -group. Its Zariski closure must be a simple linear algebraic group. The main result of [14] gives the conclusion of 1(c). ■

3.3 Almost Simple Linear Δ -Groups of Differential Type ≤ 1

We note that this includes *ordinary* linear Δ -groups. Restricting ourselves to Δ -groups of differential type at most 1 allows us to sharpen Proposition 3.4.

Proposition 3.8 *Let G be an infinite almost simple linear Δ -group with $\tau(G) \leq 1$. Then either G is commutative or perfect.*

Proof. Assume that G is not commutative. Corollary 2.13 states that $G/Z(G)$ is a simple linear differential algebraic group. Theorem 17 of [14] implies that $G/Z(G)$ is differentially isomorphic to a group $G'(F)$ G' is a simple non-commutative algebraic group and F is a field of constants with respect to a set of \mathbb{U} -linear combinations of elements of Δ . For an arbitrary group H , let $C_m(H) = \{x_1 y_1 x_1^{-1} y_1^{-1} \cdots x_m y_m x_m^{-1} y_m^{-1} \mid x_i, y_i \in H\}$. Since G' is simple, we know that $G' = C_1(G')$ (cf., [35]). We will show that $G = C_s(G)$ for some s .

Let X_m be the Kolchin closure of $C_m(G)$ in G . Let $\pi : G \rightarrow G/Z(G)$ be the canonical projection. Since $G' = C_1(G')$, we have that $\pi(C_1(G)) = \pi(G)$. Therefore $\pi(X_1) = G/Z(G)$. Since the group $Z(G)$ has smaller type than G , we have that $\tau(G) = \tau(G/Z(G))$ and $a_\tau(G) = a_\tau(G/Z(G))$. Therefore $\tau(X_1) = \tau(G)$ and furthermore $a_\tau(X_1) = a_\tau(G)$.

For each $i = 1, 2, \dots$ let $\omega_i(s) = a_i \binom{s+1}{1} + b_i$ be the differential dimension polynomial of a generic point of X_i and $\omega(s) = a \binom{s+1}{1} + b$ be the differential dimension polynomial of G (we include the possibility that $a = 0$). Since $X_1 \subset X_2 \subset \dots \subset G$, we have that $\omega_1(s) \leq \omega_2(s) \leq \dots \leq \omega(s)$. As we have just shown, $a_1 = a$ so we must have $a_1 = a_2 = \dots = a$. Therefore, $b_1 \leq b_2 \leq \dots \leq b$. Since these are all integers, we must have that $b_r = b_{r+1} = \dots$ for some r . This implies that $\omega_r(s) = \omega_{r+1}(s) = \dots$ and so $X_r = X_{r+1} = \dots$ (cf., Proposition 2, Ch.III.5 [24]).

We will now show (following a similar proof for algebraic sets) that $G = C_{2r}$. For any $c \in C_r(G)$ the map $d \mapsto dc$ sends $C_r(G)$ to $C_{2r}(G)$ and therefore sends X_r to X_r . A similar argument shows that for any $c \in X_r$, we have that multiplication by c on the left sends X_r to X_r . Therefore X_r is a Δ -subgroup of G . Since $C_r(G)$ is invariant under conjugation by elements of G , we have that X_r is a normal subgroup of G of the same type and so must equal G . Since C_r is constructible, it contains an open subset of its closure, that is an open subset of G . For any $g \in G$, we have that gC_r^{-1} intersects C_r . Therefore $g \in C_r \cdot C_r = C_{2r}$. ■

We can also strengthen Theorem 3.7.

Theorem 3.9 *Let G be a nontrivial almost simple linear Δ -group with $\tau(G) \leq 1$. Let $|\Delta| = m$.*

1. *If G is non-commutative, then there exists a quasisimple algebraic group H defined over \mathbb{Q} and a commuting linearly independent set of $m-1$ derivations $\Delta' \subset \mathbb{U}\Delta$ such that G is Δ -isomorphic to either $H(C)$ or $H(C')$ where C are the Δ -constants and C' are the Δ' constants of \mathbb{U} .*
2. *If G is commutative, then G is either isogenous to an almost simple Δ -subgroup $G' \subset \mathbb{G}_a(\mathbb{U})$ with $\tau(G') = 1$ or is Δ -isomorphic to $\mathbb{G}_a(C)$ or $\mathbb{G}_m(C)$.*

Proof. 1. Let G be a non-commutative almost simple Δ -group with $\tau(G) \leq 1$. We shall first show that $Z(G)$ is finite.

If $\tau(G) = 0$, then $\tau(Z) = -1$ and so $Z(G)$ is finite. Now assume $\tau(G) = 1$. We shall rely heavily on the results of [1]. Theorem 3.7.1(c) implies that $G/Z(G) \simeq H(F)$, where H is a simple algebraic group defined over \mathbb{Q} and F is an ordinary differentially closed field with respect some derivation δ that is a \mathbb{U} linear combination of elements of Δ . The field of constants of F is the field C of Δ -constants of \mathbb{U} . Therefore there is an exact sequence

$$1 \longrightarrow Z(G) \xrightarrow{\text{incl.}} G \xrightarrow{\alpha} H \longrightarrow 1.$$

Let $\tilde{G} = \alpha^{-1}(H(C))$. We then have the exact sequence

$$1 \longrightarrow Z(G) \xrightarrow{\text{incl.}} \tilde{G} \xrightarrow{\alpha} H(C) \longrightarrow 1.$$

Since the type of $Z(G)$ is at most 0 and the type of $H(C)$ is 0, we must have that the type of \tilde{G} is 0, that is, \tilde{G} is a group of finite Morley rank. The group $H(C)$ is a simple algebraic group of constant matrices and so is a simple δ -group as well. Therefore every normal subgroup of \tilde{G} is in $Z(G)$. Proposition 3.8 implies that \tilde{G} is perfect. Therefore the results of [1] imply that $Z(G)$ is finite.

Since $Z(G)$ is finite and $G/Z(G)$ is a simple group, we have that G is a quasisimple group. Theorem 3.11 in the Appendix states that there exists a quasisimple algebraic group H defined over \mathbb{Q} and a commuting linearly independent set of derivations $\Delta' \subset \mathbb{U}\Delta$ such that G is Δ -isomorphic $H(C')$ where C' are the Δ' constants of \mathbb{U} . Since the type of G is at most 1, we have that either Δ' has m or $m-1$ elements. If Δ' has m elements, then the Δ' constants and the Δ -constants coincide so G is Δ -isomorphic to $H(C)$, where C is the field of Δ -constants.

2. From Theorem 3.7, we know that a commutative almost simple Δ -group G is either isogenous to a subgroup G' of $\mathbb{G}_a(\mathbb{U})$ or isomorphic to $\mathbb{G}_m(C)$. If $\tau(G) = 0$, let u be a nonzero element of G' . The group $H = C \cdot u$ is a Δ -subgroup of G' with $\tau(H) = 0$. Since G' is also almost simple, we must have $G' = H$ and so G' is isomorphic to $\mathbb{G}_a(C)$. Therefore

G is isogenous to $\mathbb{G}_a(C)$.

We will now show that any almost simple group G that is isogenous to $\mathbb{G}_a(C)$ is isomorphic to $\mathbb{G}_a(C)$. Since G is isogenous to $\mathbb{G}_a(C)$, there exists a Δ -group G' and Δ -surjective homomorphisms $\alpha_1 : G' \rightarrow G$, $\alpha_2 : G' \rightarrow \mathbb{G}_a(C)$ where α_1 and α_2 have finite kernels. If we can show that α_2 has trivial kernel, then G' would be isomorphic to $\mathbb{G}_a(C)$ and so α_1 would have trivial kernel as well. This would further imply that α_1 is an isomorphism.

Therefore we must show the following: If there is a surjective Δ -homomorphism $\alpha : G \rightarrow \mathbb{G}_a(C)$ with finite kernel H then H is trivial. We first show that G is commutative. To see this, note that the map $\beta : G \times G \rightarrow G$ given by $\beta(g, h) = ghg^{-1}h^{-1}$ has image contained in H . Since G is connected, the image must be $\{1\}$. Therefore G is commutative.

Let $|H| = n$. Since \mathbb{G}_a is torsion free, we have that the torsion subgroup of G is H . Therefore, H is the kernel of the homomorphism $\gamma : G \rightarrow G$, $\gamma(g) = g^n$. The differential type of $\gamma(G)$ must therefore be the same as the type of G . Since G is strongly connected, we have $\gamma(G) = G$. This implies that if $h \in H$, $h \neq 1$, there exists a $g \in G \setminus H$ such that $g^n = h$. This element g would then be a torsion element not in H , a contradiction. Therefore H is trivial. ■

Corollary 3.10 *Assume that $|\Delta| = 1$, that is, \mathbb{U} is an ordinary differential field. Let G be an almost simple Δ -group.*

1. *If G is non-commutative, then there exists a quasisimple algebraic group H defined over \mathbb{Q} such that G is Δ -isomorphic to $H(\mathbb{U})$ or $H(C)$.*
2. *If G is commutative, then either G is isogenous to $\mathbb{G}_a(\mathbb{U})$ or Δ -isomorphic to $\mathbb{G}_a(C)$ or $\mathbb{G}_m(C)$.*

Proof. Corollary 3.10.1 follows immediately from Theorem 3.9.1. Corollary 3.10.2 follows from Theorem 3.9.2 once one notes that any proper Δ -subgroup of $\mathbb{G}_a(\mathbb{U})$ has type 0. ■

Example 2.23 shows that even in the case of ordinary differential fields, there are many non-isomorphic almost simple Δ -groups that are isogenous to $\mathbb{G}_a(\mathbb{U})$.

Appendix: Quasisimple Linear Δ -Groups

In this section we show how the results of [14]² give the following result that was needed in the proof of Theorem 3.9.

²See also [10] for a useful discussion of the results of [14].

Theorem 3.11 *Let G be a quasisimple linear Δ -group. There exists a quasisimple algebraic group H defined over \mathbb{Q} , a commuting basis $\tilde{\Delta}$ of $\mathbb{U}\Delta$ such that G is Δ -isomorphic to $H(C')$ where C' are the Δ' constants of \mathbb{U} for some $\Delta' \subset \tilde{\Delta}$.*

This result will follow from the following three propositions. The first two appear in [14]. Note that in this latter paper, the author uses the term *simple* to mean quasisimple (cf, [14], p. 222). We use the same notation as in Theorem 3.11

Proposition 3.12 ([14], Corollary 1, p. 228) *Let G be a connected Zariski dense Δ -subgroup of a semisimple algebraic subgroup H of $\mathrm{GL}_n(\mathbb{U})$. Then, G is quasisimple if and only if H is quasisimple.*

Proposition 3.13 ([14], Theorem 19, p. 232) *Let G be a connected Zariski dense Δ -subgroup of a quasisimple algebraic group H where H is defined over the constants of Δ . Then, G is conjugate to $H(C')$ where C' are the Δ' constants of some commuting linearly independent set of derivations of $\mathbb{U}\Delta$.*

Proposition 3.14 *Let G be a quasisimple Δ -subgroup of $\mathrm{GL}_n(\mathbb{U})$. Then, there exists a Δ -rational isomorphism ϕ from G onto a Zariski dense Δ -subgroup of a quasisimple algebraic group H .*

Proof. From Theorem 4.3, Chapter VIII of [22] we have that the Zariski closure \overline{G} of G can be written as a semidirect product

$$R_u(\overline{G}) \rtimes H,$$

where $R_u(\overline{G})$ is the unipotent radical of \overline{G} and H is a reductive algebraic subgroup of \overline{G} and is H is connected.

We claim that H is semisimple. First note that \overline{G} is perfect, that is, $\overline{G} = [\overline{G}, \overline{G}]$. To see this, Theorem 3.7 implies that $G = D_\Delta(G)$. The group $[\overline{G}, \overline{G}]$ is Zariski closed and so is Kolchin closed. Since $[\overline{G}, \overline{G}] \subset \overline{G}$ contains $[G, G]$, it contains $D_\Delta(G) = G$ and therefore must equal \overline{G} . Let $\phi : \overline{G} \rightarrow H$ be the projection with kernel $R_u(\overline{G})$. We have $H = \phi(\overline{G}) = \phi([\overline{G}, \overline{G}]) = [H, H]$, that is, H is perfect. The solvable radical $R(H)$ has finite intersection with $[H, H]$ (Lemma, p.125 [22]), so $R(H)$ is finite. Therefore H is semisimple.

We now claim that ϕ restricted to G is an isomorphism of G onto a Zariski dense subgroup of H . The group $\phi(\overline{G})$ is Zariski closed, from which it follows that $H = \phi(\overline{G})$ is the Zariski closure of $\phi(G)$. To see that ϕ is injective on G , note that, since G is quasisimple, $\ker \phi|_G$ is either finite or all of G . If $G = \ker \phi|_G$, we would have $G \subset R_u(\overline{G})$, and so G would not be quasisimple. Therefore $\ker \phi|_G \subset R_u(\overline{G})$ is a finite unipotent group and so must be trivial.

Since $\phi(G)$ is a connected quasisimple Zariski dense Δ -subgroup of the semisimple group H , Proposition 3.12 implies that H is quasisimple. ■

Proof of Theorem 3.11. From Proposition 3.14, we may assume that G is a connected Zariski dense Δ -subgroup of a quasisimple algebraic group H defined over \mathbb{U} . A fundamental theorem of Chevalley ([15]³), states that H is \mathbb{U} -isomorphic to an algebraic group defined over \mathbb{Q} . Therefore, we may further assume that H itself is a quasisimple group defined over $\mathbb{Q} \subset C$, the constants of Δ . Proposition 3.13 implies that G is Δ -isomorphic to $H(C')$ where C' are the Δ' constants of some commuting linearly independent set of derivations of $\mathbb{U}\Delta$. Proposition 7, Chapter 0 of [25] states that we may extend Δ' to a basis $\tilde{\Delta}$ of $\mathbb{U}\Delta$. ■

References

- [1] Tuna Altinel and Gregory Cherlin. On central extensions of algebraic groups. *J. Symbolic Logic*, 64(1):68–74, 1999.
- [2] Moulay A. Barkatou, Thomas Cluzeau, and Jacques-Arthur Weil. Factoring partial differential systems in positive characteristic. In *Differential equations with symbolic computation*, Trends Math., pages 213–238. Birkhäuser, Basel, 2005. With an appendix by Marius van der Put.
- [3] Andreas Baudisch. On superstable groups. *J. London Math. Soc. (2)*, 42(3):452–464, 1990.
- [4] Richard Beals and Elena A. Kartashova. Constructive factorization of linear partial differential operators. *Teoret. Mat. Fiz.*, 145(2):165–180, 2005.
- [5] George W. Bluman and Sukeyuki Kumei. *Symmetries and differential equations*, volume 81 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1989.
- [6] Henry Blumberg. *Über algebraische Eigenschaften von linearin homogene Differentialausdrücken*. PhD thesis, Georg-August-Universität, 1912.
- [7] Armand Borel. Properties and linear representations of Chevalley groups. In *Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, N.J., 1968/69)*, Lecture Notes in Mathematics, Vol. 131, pages 1–55. Springer, Berlin, 1970.
- [8] Giuseppa Carrà Ferro. A survey on differential Gröbner bases. In *Gröbner bases in symbolic analysis*, volume 2 of *Radon Ser. Comput. Appl. Math.*, pages 77–108. Walter de Gruyter, Berlin, 2007.
- [9] Elie Cartan. Les groupes de transformations continues, infinis, simple. *Annales scientifiques de l'École Normale Supérieure, Sér. 3*, 26:93–161, 1909.

³See [7] (especially 3.3(6) and 3.3(1)) for a survey of these notes and [10] for a discussion of the literature reproving and extending this result.

- [10] Phyllis Joan Cassidy. Quasisimple differential algebraic groups. Preprint; available at www.math.ncsu.edu/~singer/Quasisimple.pdf, 2009.
- [11] Phyllis Joan Cassidy. Differential algebraic groups. *Amer. J. Math.*, 94:891–954, 1972.
- [12] Phyllis Joan Cassidy. The differential rational representation algebra on a linear differential algebraic group. *J. Algebra*, 37(2):223–238, 1975.
- [13] Phyllis Joan Cassidy. Unipotent differential algebraic groups. In *Contributions to algebra (collection of papers dedicated to Ellis Kolchin)*, pages 83–115. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1977.
- [14] Phyllis Joan Cassidy. The classification of the semisimple differential algebraic groups and the linear semisimple differential algebraic Lie algebras. *J. Algebra*, 121(1):169–238, 1989.
- [15] Claude Chevalley. *Classification des groupes algébriques semi-simples*. Springer-Verlag, Berlin, 2005. Collected works. Vol. 3, Edited and with a preface by P. Cartier, With the collaboration of Cartier, A. Grothendieck and M. Lazard.
- [16] Thomas Cluzeau and Alban Quadrat. Factoring and decomposing a class of linear functional systems. *Linear Algebra Appl.*, 428(1):324–381, 2008.
- [17] Dima Grigoriev. Weak Bézout inequality for D-modules. *J. Complexity*, 21(4):532–542, 2005.
- [18] Dima Grigoriev and Fritz Schwarz. Factoring and solving linear partial differential equations. *Computing*, 73(2):179–197, 2004.
- [19] Dima Grigoriev and Fritz Schwarz. Loewy and primary decompositions of D -modules. *Adv. in Appl. Math.*, 38(4):526–541, 2007.
- [20] Dima Grigoriev and Fritz Schwarz. Non-holonomic overideals of differential ideals in two variables and absolute factoring. [arXiv:math.AP/0811.1368v1](https://arxiv.org/abs/math/0811.1368v1), 2008.
- [21] Gerhard Hochschild. *Basic Theory of Algebraic Groups and Lie Algebras*, volume 75 of *Graduate Texts in Mathematics*. Springer Verlag, Berlin, 1976.
- [22] James E. Humphreys. *Linear Algebraic Groups*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1975.
- [23] E. R. Kolchin. Constrained extensions of differential fields. *Advances in Math.*, 12:141–170, 1974.
- [24] Ellis R. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, New York, 1976.

- [25] Ellis R. Kolchin. *Differential algebraic groups*, volume 114 of *Pure and Applied Mathematics*. Academic Press Inc., Orlando, FL, 1985.
- [26] M. V. Kondratieva, A. B. Levin, A. V. Mikhalev, and E. V. Pankratiev. *Differential and difference dimension polynomials*, volume 461 of *Mathematics and its Applications*. Kluwer Academic Publishers, Dordrecht, 1999.
- [27] Serge Lang. *Algebra*. Addison Wesley, New York, 3rd edition, 1993.
- [28] David Marker. *Model theory*, volume 217 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002. An introduction.
- [29] David Marker, Margit Messmer, and Anand Pillay. *Model theory of fields*, volume 5 of *Lecture Notes in Logic*. Springer-Verlag, Berlin, 1996.
- [30] Claude Mitschi and Michael F. Singer. Solvable-by-finite groups as differential Galois groups. *Ann. Fac. Sci. Toulouse*, XI(3):403 – 423, 2002.
- [31] Anand Pillay. Some foundational questions concerning differential algebraic groups. *Pacific J. Math.*, 179(1):179–200, 1997.
- [32] Jean-Francois Pommaret. *Partial differential control theory*, volume 530 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 2001. Vol. I. Mathematical tools; Vol. II. Control systems, With a foreword by Jan C. Willems.
- [33] Roman O. Popovych, Michael Kunzinger, and Nataliya M. Ivanova. Conservation laws and potential symmetries of linear parabolic equations. *Acta Appl. Math.*, 100(2):113–185, 2008.
- [34] Marius van der Put and Michael F. Singer. *Galois Theory of Linear Differential Equations*, volume 328 of *Grundlehren der mathematischen Wissenschaften*. Springer, Heidelberg, 2003.
- [35] Rimhak Ree. Commutators in semi-simple algebraic groups. *Proc. Amer. Math. Soc.*, 15:457–460, 1964.
- [36] Joseph F. Ritt. *Differential Algebra*, volume 33 of *American Mathematical Society Colloquium Publication*. American Mathematical Society, New York, 1950.
- [37] Maxwell Rosenlicht. Some basic theorems on algebraic groups. *Amer. J. Math.*, 78:401–443, 1956.
- [38] Ekaterina S. Shemyakova. Parametric factorizations of second-, third- and fourth-order linear partial differential operators with a completely factorable symbol on the plane. *Math. Comput. Sci.*, 1(2):225–237, 2007.

- [39] Ekaterina S. Shemyakova and Franz Winkler. Obstacles to the factorization of partial differential operators into several factors. *Programmirovaniye*, 2:17–25, 2007. translation in *Program. Comput. Software* 33 (2007) no. 2, 67–73.
- [40] Michael F. Singer. Testing reducibility of linear differential operators: a group theoretic perspective. *Applicable Algebra in Engineering, Communication and Computing*, 7:77–104, 1996.
- [41] William Yu Sit. Typical differential dimension of the intersection of linear differential algebraic groups. *J. Algebra*, 32(3):476–487, 1974.
- [42] Sonat Suer. *Model Theory of Differentially Closed Fields with Several Commuting Derivations*. PhD thesis, University of Illinois at Urbana-Champaign, 2007.
- [43] Sergey P. Tsarev. Factorization of linear partial differential operators and the Darboux method for integrating nonlinear partial differential equations. *Teoret. Mat. Fiz.*, 122(1):144–160, 2000.
- [44] Sergey P. Tsarev. Factorization of linear differential operators and systems. In *Algebraic theory of differential equations*, volume 357 of *London Math. Soc. Lecture Note Ser.*, pages 111–131. Cambridge Univ. Press, Cambridge, 2009.
- [45] Min Wu. On the factorization of differential modules. In *Differential equations with symbolic computation*, Trends Math., pages 239–254. Birkhäuser, Basel, 2005.

A Jordan-Hölder Theorem for Differential Algebraic Groups

Phyllis J. Cassidy* and Michael F. Singer†

Abstract

We show that a differential algebraic group can be filtered by a finite subnormal series of differential algebraic groups such that successive quotients are almost simple, that is have no normal subgroups of the same type. We give a uniqueness result, prove several properties of almost simple groups and, in the ordinary differential case, classify almost simple linear differential algebraic groups.

1 Introduction

Let (k, δ) be a differential field of characteristic zero with derivation δ and let $k[\delta]$ be the ring of linear differential operators with coefficients in k . It has been known for a long time (see [40] for a brief history and [2],[34],[46] for more recent algorithmic results) that one has a form of unique factorization for this ring:

Given $L \in k[\delta]$, $L \notin k$, there exist irreducible $L_i \in k[\delta]$, $i = 1, \dots, r$ such that $L = L_1 \cdots L_r$. Furthermore, if $L = \tilde{L}_1 \cdots \tilde{L}_s$, $\tilde{L}_i \in k[\delta]$, \tilde{L}_i irreducible, then $r = s$ and, for some permutation σ of the subscripts, there exist nonzero $R_i, S_i \in k[\delta]$ with $\deg(R_i) < \deg(\tilde{L}_i)$ such that $L_{\sigma(i)}R_i = S_i\tilde{L}_i$ for $i = 1, \dots, r$.

A direct generalization of this to partial differential operators fails as the following example shows. In [6], Blumberg considered the following third order linear partial differential

*Department of Mathematics, Smith College, MA 01063; Department of Mathematics, The City College of New York, New York, NY 10038, USA, email: pcassidy@smith.edu

†Department of Mathematics, North Carolina State University, Box 8205, Raleigh, NC 27695-8205, USA, email: singer@math.ncsu.edu. The second author was partially supported by NSF Grant CCR-0634123 and he would also like to thank the Max Planck Institut für Mathematik and the Courant Institute of Mathematical Sciences for their hospitality. This work was begun during a visit to the first institute in the Spring of 2006 and continued during a stay as a Visiting Member at the second institute in the Fall of 2008.

operator (which he attributes to E. Landau)

$$L = \delta_x^3 + x\delta_x^2\delta_y + 2\delta_x^2 + 2(x+1)\delta_x\delta_y + \delta_x + (x+2)\delta_y$$

over the differential field $\mathbb{C}(x, y)$ where $\delta_x = \frac{\partial}{\partial x}$ and $\delta_y = \frac{\partial}{\partial y}$. He showed that this operator has two factorizations

$$\begin{aligned} L &= (\delta_x + 1)(\delta_x + 1)(\delta_x + x\delta_y) \\ &= (\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y)(\delta_x + 1). \end{aligned}$$

and that the operator $\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y$ could not be further factored in any differential extension of $\mathbb{C}(x, y)$ (this example is also discussed in [17]).

To confront this situation one must recast the factorization problem in other terms. For ordinary differential operators, the factorization result is best restated in terms of $k[\delta]$ -modules. If one considers the $k[\delta]$ -module $M_L = k[\delta]/k[\delta]L$, then the above factorization result can be stated as a Jordan-Hölder type theorem: M_L has a composition series $M_L = M_0 \supset M_1 \supset \dots \supset M_r = \{0\}$ where successive quotients M_{i-1}/M_i are simple and any two such series have the same length and, after a possible renumbering, isomorphic quotients. Casting questions in terms of ideals and modules in the partial differential case yields many interesting results (cf., [17, 18, 19, 16], [15], [32]). Other approaches to factoring linear operators are contained in [4],[39, 38], [44]. In [45], Tsarev alludes to an approach to factorization through the theory of certain abelian categories that has points of contact with the approach presented here.

We will take an alternate approach - via differential algebraic groups. This stems from the seemingly trivial observation that the solutions of a linear differential equation form a group under addition. In fact, this is an example of a linear differential algebraic group, that is, a group of matrices whose entries lie in a differential field and satisfy some fixed set of differential equations (here we identify $\{y \in k \mid L(y) = 0\}$ with the group $\left\{ \begin{pmatrix} 1 & y \\ 0 & 1 \end{pmatrix} \mid L(y) = 0 \right\}$.) We shall prove a Jordan-Hölder type theorem for differential algebraic groups (of which linear differential algebraic groups are a special case.) This will encompass the factorization result for linear ordinary differential equations, allow us to give a “factorization” result for linear partial differential equations and allow us to reduce the study of general differential algebraic groups to the study of *almost simple* groups (to be defined below) and their extensions.

The rest of the paper is organized as follows. In the remainder of this section we review some basic definitions and facts from differential algebra. In Section 2, we shall review some definitions and facts from differential algebra and the theory of linear differential algebraic groups. We will state and prove a Jordan-Hölder Theorem for linear differential algebraic groups and show how it applies to the example of Landau above. In Section 3, we will discuss the structure of almost simple groups in more detail.

The authors would like to acknowledge the influence of Ellis Kolchin. The first author discussed her notions of solid and the core (what we call strongly connected and the strong

identity component) with him in the 1980's and he suggested a version of the Jordan-Hölder Theorem below. In addition, we would like to thank William Sit for explaining the connection between Gröbner bases of left ideals in the ring of differential operators and characteristic sets of linear differential ideals (see the discussion in Example 2.31).

1.1 Differential Algebra Preliminaries

Throughout, we let \mathbb{Z} = the ring of integers, \mathbb{Q} = the field of rational numbers, \mathbb{C} = the field of complex numbers.

We refer to [9, 10, 11, 12, 13, 24, 25] for the basic concepts of differential algebraic geometry and differential algebraic groups. All rings have characteristic zero. We fix a set $\Delta = \{\delta_1, \dots, \delta_m\}$ of commuting derivation operators. We often use the prefix Δ - to indicate the Δ -differential structure on a ring, field, group, *etc.* If k is a Δ -field, and R and S are Δ - k -algebras, a k -homomorphism $\varphi : R \rightarrow S$ is a Δ - k -homomorphism if $\varphi \circ \delta = \delta \circ \varphi$, $\delta \in \Delta$. Let k be a Δ -field. The k -algebra of differential operators generated by Δ is denoted by $k[\Delta]$, and the set of monomials $\theta = \delta_1^{e_1} \dots \delta_m^{e_m}$ in $k[\Delta]$ by Θ . The *order* of θ is the degree of θ .

If $\eta = (\eta_1, \dots, \eta_n)$ is a family of elements of a Δ - k -algebra, the Δ - k -algebra $k\{\eta\}$ is the k -algebra $k[\Theta\eta]$, where $\Theta\eta$ is the family of derivatives $(\theta\eta_j)_{\theta \in \Theta, 1 \leq j \leq n}$ of the coordinates of η . Thus, if z_1, \dots, z_n are elements of a Δ - k -algebra such that the family $(\theta z_j)_{\theta \in \Theta, 1 \leq j \leq n}$ is algebraically independent over k , the Δ - k -algebra $k\{z\} = k\{z_1, \dots, z_n\}$ is called the Δ -polynomial algebra. Suppose $k\{\eta\}$ is an integral domain. Set $k\langle\eta\rangle$ equal to its quotient field. We say that $k\{\eta\}$ (*resp.* $k\langle\eta\rangle$) is a *finitely generated Δ - k -algebra* (*resp.* Δ - k -field). The quotient field $k\langle z\rangle$ of the Δ -polynomial ring is called the field of Δ -rational functions. Let $(P_i)_{i \in I}$ be a family of Δ -polynomials in $k\{z_1, \dots, z_n\}$. The Δ -ideal $[(P_i)_{i \in I}]$ equals the ideal $((\Theta P_i)_{i \in I})$.

Fix a Δ -field k . A Δ - k -field K is *differentially closed* if given a prime Δ -ideal I in $K\{z_1, \dots, z_n\}$ and differential polynomial $Q \notin I$, there exists $\eta = (\eta_1, \dots, \eta_n) \in K^n$, with

$$Q(\eta) \neq 0, \quad \text{and } P(\eta) = 0 \quad \text{for all } P \in I.$$

Equivalently, K is differentially closed if every system of differential polynomial equations and inequations that has a solution with coordinates in a Δ - K -field has a solution with coordinates in K (see [23], [27], [28], [31]). We say that a Δ - k -field \mathbb{U} is *k -universal* if, in addition to being differentially closed, \mathbb{U} satisfies the useful property that if I is a prime Δ -ideal in $k\{z_1, \dots, z_n\}$, there is in \mathbb{U}^n a *generic zero* of I . A generic zero η is a zero of I that has the property that if P is in $k\{z_1, \dots, z_n\}$ and $P(\eta) = 0$, then P is in I . We fix k and \mathbb{U} .

We define on *affine n -space* \mathbb{U}^n both the Zariski and Kolchin topologies. The *Kolchin topology*, which is finer than the Zariski topology, is also Noetherian. A subset X of \mathbb{U}^n is a Δ -variety if it is *Kolchin closed*, *i.e.*, X is the set of zeros of a set of Δ -polynomials

with coefficients in \mathbb{U} . By the Ritt Basis Theorem, there is a finite set P_1, \dots, P_r of Δ -polynomials with coefficients in \mathbb{U} such that X is the set of zeros of P_1, \dots, P_r . Call P_1, \dots, P_r *defining Δ -polynomials* of X . The radical Δ -ideal $I = \sqrt{[P_1, \dots, P_r]}$ in $\mathbb{U}\{y\}$ generated by P_1, \dots, P_r , is called the *defining Δ -ideal* of X . If P_1, \dots, P_r have coefficients in k , we say that X is *defined over k* , and call it a Δ - k -variety. Let Y be a Δ -subvariety of a Δ - k -variety X . We do not assume that Y is defined over k . However, there is a finitely generated delta-field extension K of k such that Y is a Δ - K -variety. X is *irreducible* if and only if I is prime. The Ritt Basis Theorem implies that X is a finite union of maximal irreducible Δ -subvarieties. This implies that the Kolchin topology is Noetherian. In this paper, we will usually assume that X is irreducible, with defining Δ -ideal I . The *residue class ring* $\mathbb{U}\{z_1, \dots, z_n\}/I$ is a Δ - \mathbb{U} -algebra and an integral domain, which we denote by $\mathbb{U}\{X\}$. As in algebraic geometry, we regard the residue classes of the Δ -indeterminates as coordinates on X , and call the elements of $\mathbb{U}\{X\}$ *Δ -polynomial functions* on X . If X is a Δ - k -variety, the Δ - k -subalgebra $k\{X\}$, consisting of the elements of $\mathbb{U}\{X\}$ that have coefficients in k , is Δ - k -isomorphic to $k\{z_1, \dots, z_n\}/(I \cap k\{z_1, \dots, z_n\})$. If X is irreducible, the elements of the quotient field $\mathbb{U}\langle X \rangle$ of $\mathbb{U}\{X\}$ (*resp.* $k\langle X \rangle$ of $k\{X\}$) are called Δ -rational functions (*resp.* Δ - k -rational functions) on X . A Δ -rational function f in $\mathbb{U}\langle X \rangle$ is *everywhere defined* if for every $\eta \in X$, there exist $p, q \in \mathbb{U}\{X\}$ such that $q(\eta) \neq 0$, and $f(\eta) = \frac{p(\eta)}{q(\eta)}$. In contrast with rational functions on algebraic varieties, an everywhere defined Δ -rational function need not be in the ring of Δ -polynomial functions on X . Indeed, it may not have a global denominator. Let $\eta = (\eta_1, \dots, \eta_n)$ be in \mathbb{U}^n , and let σ be a Δ - k -automorphism of \mathbb{U} . We define $\sigma\eta = (\sigma\eta_1, \dots, \sigma\eta_n)$. $\sigma(X)$ is a Δ -variety. A necessary and sufficient condition that X be defined over k is that $\sigma(X) \subseteq X$ (equivalently, $\sigma(X) = X$) for every Δ - k -automorphism of \mathbb{U} .

If X is an irreducible Δ - k -variety, an element $\eta = (\eta_1, \dots, \eta_n)$ is generic over k for X if η is a generic zero of $I \cap k\{z_1, \dots, z_n\}$, where I is the defining Δ -ideal of X . The Δ -field $k\langle \eta \rangle$ is Δ - k -isomorphic to $k\langle X \rangle$. It represents in \mathbb{U} the field of rational functions on X .

If $X \subseteq \mathbb{U}^n$ and $Y \subseteq \mathbb{U}^p$ are irreducible Δ - k -varieties, a Δ -*morphism* $f : X \rightarrow Y$ is a map whose p coordinate functions are everywhere defined Δ -rational functions. If they are Δ - k -rational, we call f a Δ - k -morphism. We should note that if f is a Δ -morphism, $f(X)$ and its Kolchin closure are irreducible, and are defined over k if f is a Δ - k -morphism. A Δ -*morphism* $f : X \rightarrow Y$ is *dominant* if the Kolchin closure of $f(X)$ is Y . In this case, $f(X)$ contains a Kolchin dense open subset of Y ($f(X)$ is *constructible*). If X is an arbitrary Δ - k -variety with components X_1, \dots, X_s , and Y is a Δ - k -variety with components Y_1, \dots, Y_t , we can define a Δ -morphism to be an s -tuple of Δ -morphisms $f_i : X_i \rightarrow Y_{j_i}$, $i = 1, \dots, s$.

If $X \subseteq \mathbb{U}^n$ and $Y \subseteq \mathbb{U}^p$ are Δ -varieties, then the Cartesian product $X \times Y$ is a Δ -subvariety of \mathbb{U}^{n+p} , and the projection maps are Δ -morphisms.

An *affine Δ -group* is a group G whose underlying set is a Δ -subvariety of \mathbb{U}^n for some n , and whose group laws are Δ -morphisms. A Δ -homomorphism of affine Δ -groups is a homomorphism that is also a Δ -morphism of Δ -varieties. As in algebraic group theory, G has a finite number of connected components. We call G a Δ - k -group if its group laws and

components are defined over k and its identity element has coordinates in k . Note that a Δ -subgroup of G is not necessarily a Δ - k -subgroup, but is defined over a finitely generated Δ - k -field in \mathbb{U} . Not all affine Δ -groups are linear (that is, isomorphic to a Δ -subgroup of $\mathrm{GL}_n(\mathbb{U})$ for some n). In this paper, we will assume that a linear Δ -group is embedded in $\mathrm{GL}_n(\mathbb{U})$, for some n .

In [25], Kolchin develops axiomatically a theory of Δ - k -groups and Δ - k -varieties, extending the affine theory. In Chapter V.3 of [25], Kolchin proves that a Δ - k -variety X has a k -affine open dense subset U . In the corollary, p. 140, he shows that by extending k perhaps to a finitely generated Δ -extension field, we may assume that there is a covering of X by open dense subsets, each of which is defined over k and k -affine. Although we will state and prove our results in the general context of [25], readers who wish to restrict themselves to *linear* Δ -groups, or just to the solution sets of linear homogeneous differential equations, will find almost all that is needed (with one notable exception which we will discuss later) in the papers [10, 11]. Another source is Anand Pillay's foundational paper [31], which treats Δ -groups as *definable groups* in differentially closed fields. The theory of definable groups is particularly well-suited to interpret Kolchin's axiomatic approach.

2 Jordan-Hölder Theorem

2.1 Differential Type and Typical Differential Transcendence Degree

The key concepts used in this paper are *differential type* and *typical differential transcendence degree* (typical differential dimension). We briefly review their definitions. Let k be a Δ -field finitely generated over \mathbb{Q} .

Let $\eta = (\eta_1, \dots, \eta_n)$ be in \mathbb{U}^n and let $\Theta(s)$ be the set of monomial operators in Θ of order $\leq s$. In Section II.12 of [24], Kolchin shows that there is a numerical polynomial $\omega_{\eta/k}(s)$ (*i.e.*, a polynomial taking on integer values on \mathbb{Z}), called the *differential dimension polynomial*, of degree $\leq m$, such that for large values of s , $\omega_{\eta/k}(s)$ is the transcendence degree over k of $k((\theta\eta_i))_{\theta \in \Theta(s), 1 \leq i \leq n}$ for all sufficiently large values of s . See Example 2.31 for further discussion about the computation of the differential dimension polynomial, and also [24] for further information, and algorithms, as well as references. We may write

$$\omega_{\eta/k}(s) = \sum_{i=0}^m a_i \binom{s+i}{i}, \quad a_i \in \mathbb{Z}.$$

If I is a prime Δ -ideal in the differential polynomial ring $\mathbb{U}\{z_1, \dots, z_n\}$, and k is a differential field of definition of I , we may define the differential dimension polynomial of I to be $\omega_{\eta/k}(s)$, where η is a generic zero of $I \cap k\{z_1, \dots, z_n\}$. $\omega_{\eta/k}(s)$ is independent of the choice of k and η . If X is the Kolchin closed subset of \mathbb{U}^n defined by I , and η is generic for X

over a field k of definition, we define the dimension polynomial $\omega(X)$ to be $\omega_{\eta/k}(s)$.

The dimension polynomial $\omega(X)$ is not, however, a differential birational invariant. This means that it makes no sense to speak of the dimension polynomial of an arbitrary irreducible Δ - k -variety. There are, however, two important differential birational invariants singled out by Kolchin: the *degree* τ and the *leading coefficient* a_τ of $\omega(X)$. If X is any irreducible Δ - k -subvariety of \mathbb{U}^n , we call the degree of $\omega(X)$ the *differential type* (or Δ -type) $\tau(X)$ of X , and the leading coefficient the *typical differential dimension* (or Δ -dimension) $a_\tau(X)$ of X . Let X be a Δ - k -variety. As we mentioned in Section 1.1, we may assume that k is such that there is a finite set of Kolchin open k -affine subsets U of X covering X . There exists a positive integer n such that each open set U in the covering is Δ - k -rationally isomorphic to a Δ - k -subvariety Y of \mathbb{U}^n . If X is irreducible, so are U and Y . In this case, we define $\tau(U)$ to be $\tau(Y)$, and $a_\tau(U)$ to be $a_\tau(Y)$. Since Δ -type and typical Δ -dimension are differential birational invariants, all the open sets in the covering have the same Δ -type τ and typical Δ -dimension a_τ . So, we can define $\tau(X) = \tau$ and $a_\tau(X) = a_\tau$. We should note that a point of X is generic for X over k if and only if whenever U is a k -affine k -open subset of X containing the point, and Y is a Δ - k -subvariety of \mathbb{U}^n that is Δ - k -rationally isomorphic to U , the coordinate n -tuple $\eta = (\eta_1, \dots, \eta_n)$ is generic over k for Y . By abuse of language, we will call the generic point η . Thus, $\tau(X)$ is the degree and $a_\tau(X)$ is the leading coefficient of $\omega_{\eta/k}(s)$. If X is not irreducible, we define $\tau = \tau(X)$ to be the maximum of the differential type of its irreducible components, and $a_\tau(X)$ to be the maximum of their typical differential dimensions. The connected components of a Δ - k -group G all have the same Δ -type and typical Δ -dimension. We should also mention that if X is an irreducible Δ -subvariety of a Δ - k -group or Δ - k -homogeneous space and σ is a Δ -automorphism of \mathbb{U} over k , then $\tau(\sigma(X)) = \tau(X)$ and $a_\tau(\sigma(X)) = a_\tau(X)$.

Grigoriev and Schwarz, [16, 17, 18, 19], have made extensive use of differential type τ and typical differential dimension a_τ in their studies of systems of linear homogeneous partial differential equations. Following Grigoriev and Schwarz, we will call the pair $(\tau(X), a_\tau(X))$ the *gauge* of X .

Let C_k be the field of Δ -constants of k . A *transformation* of Δ is a set of derivation operators on k , denoted by the symbol Δ^c , where $c = (c_{ij})_{1 \leq i, j \leq m} \in GL_m(C_k)$ and Δ^c is the set

$$\delta'_i = \sum_{j=1}^m c_{ij} \delta_j, \quad i = 1, \dots, m$$

Every Δ -field extension of k is also a Δ^c -extension of k . Suppose $L = k\langle \eta \rangle$, with $\eta = (\eta_1, \dots, \eta_m)$ a finite family of elements of L . In Theorem 7 of [24], Kolchin established the significance of Δ -type and typical Δ -transcendence degree by showing that if $\tau = \tau_{\eta/k}$ is the Δ -type of η/k , then there is a transformation Δ^c of Δ and a subset Δ' of cardinality τ such that L is a finitely generated Δ' -field extension of k of Δ' -differential transcendence degree a_τ . In fact, there is a Zariski open dense subset of $GL_m(C_k)$ with this property. Since every Δ -extension field is a Δ^c -field extension and conversely, and Δ -dimension polynomials of

Δ - k -varieties are invariant under such transformations, we may assume that $\Delta^c = \Delta$, and, thus, $\Delta' \subset \Delta$. Let K be a Δ -subfield of L containing k . Then, K is a Δ' -finitely generated Δ' -extension field of k (Proposition 14, p. 112 of [24]).

Since the Δ -type of η/k and the typical Δ -transcendence degree of η/k are differential birational invariants, we can refer to them as the Δ -type of L over k and the typical Δ -transcendence degree of L over k .

From the classical interpretation of Δ -type and typical Δ -dimension, one expects the truth of the following lemma.

Lemma 2.1 *Let L be a Δ -finitely generated Δ -extension field of k , and let K be a Δ -subfield of L containing k . Then, the Δ -type of K/k is less than or equal to the Δ -type of L/k , and the typical Δ -transcendence degree of K/k is less than or equal to the typical Δ -transcendence degree of L/k .*

Proof. Let $\Delta' \subset \Delta$ be such that L/k is a Δ' -finitely generated extension field. Then, since K/k is also Δ' -finitely generated, if Δ'' is any subset of Δ of larger cardinality than Δ' , K is Δ'' -algebraic over k ([24], Proposition 12, p. 109. See also [42], Proposition 2.4, p. 478.) It follows that the Δ -type of K/k is less than or equal to the Δ -type of L/k . If the inequality of Δ -types is strict, so is the inequality of Δ -transcendence degrees. So, suppose Δ -type (K/k) = Δ -type (L/k). Then, the typical Δ -transcendence degree of K/k is the cardinality of a Δ' -transcendence basis of K/k . Since it can be extended to a Δ' -transcendence basis of L/k (Theorem 4(d), p. 105 of [24]), it follows that the Δ -transcendence degree of K/k must be less than or equal to that of L/k . ■

Corollary 2.2 *Let X and Y be Δ - k -varieties, and let $f : X \rightarrow Y$ be a dominant Δ - k -morphism. Then, $\tau(X) \geq \tau(Y)$ and $a_\tau(X) \geq a_\tau(Y)$.*

Proof. Let η be generic for X over k . Then, $\zeta = f(\eta)$ is generic for Y over k . Therefore, $k \langle \zeta \rangle$ is a Δ -subfield of $k \langle \eta \rangle$. The conclusion then follows from the preceding lemma. ■

Lemma 2.3 *Let X and Y be irreducible Δ - k -varieties and $f : X \rightarrow Y$ be an injective dominant Δ - k -morphism. Then $\tau(X) = \tau(Y)$.*

Proof. Let $\eta = (\eta_1, \dots, \eta_n)$ be a generic point of X over k . By assumption $\zeta = f(\eta)$ is a generic point of Y over k and, since f is differentially rational, $k \langle \zeta \rangle \subset k \langle \eta \rangle$. Note that \mathbb{U} is universal over $k \langle \zeta \rangle$ as well. We will show that $k \langle \zeta \rangle = k \langle \eta \rangle$. Let σ be a differential isomorphism of $k \langle \eta \rangle$ over $k \langle \zeta \rangle$ into \mathbb{U} . Since $\sigma(\zeta) = \zeta$ and $\sigma(\zeta) = \sigma(f(\eta)) = f(\sigma(\eta))$ we have that $f(\sigma(\eta)) = f(\eta)$. Since f is injective, we have that $\sigma(\eta) = \eta$. If some coordinate η_i of η were not in $k \langle \zeta \rangle$, there would be an isomorphism of $k \langle \eta \rangle$ over $k \langle \zeta \rangle$ into \mathbb{U} which would move η_i . Therefore we can conclude that $k \langle \eta \rangle = k \langle \zeta \rangle$. Proposition 15(b), p. 117 of [24] implies the conclusion. ■

Corollary 2.4 *Let X and Y be irreducible Δ - k -varieties and $f : X \rightarrow Y$ be an injective Δ - k -morphism. Then $\tau(X) \leq \tau(Y)$.*

Proof. Let V be the closure of $f(Y)$. The map $f : X \rightarrow V$ is injective and dominant so $\tau(X) = \tau(V)$. Proposition 15(b), p. 117 of [24] implies that $\tau(V) \leq \tau(Y)$. ■

Let G be a Δ - k -group and H a Δ - k -subgroup of G . Kolchin showed ([25] Ch. IV.4) that the coset space G/H has the structure of a Δ - k -variety (in fact, a Δ - k -homogeneous space for G) and showed that

$$\tau(G) = \max(\tau(H), \tau(G/H)). \quad (1)$$

Furthermore, he showed that if $\tau = \tau(G)$, then

$$a_\tau(G) = a_\tau(H) + a_\tau(G/H) \quad (2)$$

where $a_\tau(H) = 0$ if $\tau(G) > \tau(H)$ with a similar convention for $a_\tau(G/H)$.

Remarks 2.5 1. The fact that G/H has the structure of a Δ - k -variety also follows from the model-theoretic concept of *elimination of imaginaries* which holds for differentially closed fields ([28], p. 57).

2. The simplest (to define, not necessarily to understand) examples of differential algebraic groups are zero sets of systems of linear homogeneous differential equations in one indeterminate. Cassidy [10] has shown that these are the only Δ -subgroups of $\mathbb{G}_a(\mathbb{U}) = (\mathbb{U}, +)$. In this case, quotients are easy to construct. If $H \subset G \subset \mathbb{G}_a(\mathbb{U})$ and H is defined by the vanishing of L_1, \dots, L_t , then the image in \mathbb{U}^t of G under the map $y \mapsto (L_1(y), \dots, L_t(y))$ is a Δ -subgroup of \mathbb{U}^n isomorphic to G/H .

3. As we mentioned earlier, if one restricts one's attention to *linear* differential algebraic groups (as defined by Cassidy [10]), one will find all needed facts in the works of Cassidy and [24] except for the equations (1) and (2), which one can just assume when reading the following results and proofs.

2.2 Strongly Connected Groups, Almost Simple Groups and Isogeny

2.2.1 Strongly Connected Groups

We now define the *strong identity component* of a Δ - k -group G . Let \mathcal{S} be the set of all Δ -subgroups $H \subset G$ such that $\tau(H) = \tau(G)$ and $a_\tau(H) = a_\tau(G)$. Note that from (1) and (2), this is the same as the set of Δ -subgroups $H \subset G$ such that $\tau(G/H) < \tau(G)$. We claim that if $H_1, H_2 \in \mathcal{S}$, then $H_1 \cap H_2 \in \mathcal{S}$. To see this, note that we have an injection of $H_1/H_1 \cap H_2$ into G/H_2 , so by (1) we have $\tau(H_1/H_1 \cap H_2) < \tau(G)$. Again by (1), we have

that $\tau(G/H_1 \cap H_2) = \max(\tau(G/H_1), \tau(H_1/H_1 \cap H_2)) < \tau(G)$. Therefore $H_1 \cap H_2 \in \mathcal{S}$.

Since the Kolchin topology is Noetherian, the set \mathcal{S} has minimal elements. The above argument shows that there is a unique minimal element and justifies the following definition.

Definition 2.6 *Let G be a Δ - k -group. The strong identity component G_0 of G is defined to be the smallest Δ - \mathbb{U} -subgroup H of G such that $\tau(G/H) < \tau(G)$. We say that G is strongly connected if $G_0 = G$.*

Note that G_0 is contained in the identity component G^0 . Therefore, every strongly connected Δ -group is connected.

Remarks 2.7 1. In the usual theory of algebraic groups, one can define the identity component of a group G to be the smallest subgroup G^0 such that G/G^0 is finite and define a group to be connected if $G = G^0$. The notions of strong identity component and strongly connected are meant to be refinements in the context of Δ -groups of these former concepts.

2. If σ is a Δ - k -automorphism of \mathbb{U} , then $\sigma(G_0)$ is again a minimal element of \mathcal{S} and so must coincide with G_0 . Therefore, Corollary 2, p.77 of [25] implies that G_0 is defined over k . Note that if $\phi : G \rightarrow G$ is a Δ -automorphism of G , then $\phi(G_0)$ is again a minimal element of \mathcal{S} and so must equal G_0 . Therefore G_0 is a characteristic subgroup of G . In particular G_0 is a *normal* subgroup of G and if H is a normal Δ -subgroup of G , the strong identity component of H is again normal in G .

3. The analogue of the fact that if G is an algebraic group, every connected algebraic subgroup is contained in the identity component of G is the following: If G is a Δ -group of type τ , then every strongly connected Δ -subgroup H of type τ is contained in the strong identity component of G . To see this, first note that the usual isomorphism theorems hold for Δ -groups ([25], Chapter IV). Since G_0 is normal in G , HG_0 is a Δ -subgroup of G and HG_0/G_0 is isomorphic to $H/G \cap G_0$ as Δ -groups. Since the canonical inclusion $H/H \cap G_0 \hookrightarrow G/G_0$ is injective, Corollary 2.4 implies that $\tau(H/H \cap G_0) < \tau(G) = \tau(H)$. Since H is strongly connected, $H \cap G_0 = H$.

4. The analogue of the fact that if G, G' are algebraic groups, and $\phi : G \rightarrow G'$ is a homomorphism, then, the image of the identity component is contained in the identity component of G' is the following: If G, G' are Δ -groups of the same type, and $\phi : G \rightarrow G'$ is a homomorphism, then, the image of the strong identity component of G is contained in the strong identity component of G' . This follows immediately from the preceding remark.

5. Let G be a strongly connected Δ -group. Let G' be a nontrivial Δ -group, and let $\varphi : G \rightarrow G'$ be a surjective Δ -homomorphism. Then, G' is strongly connected and the type $\tau(G')$ equals the type τ of G . For, Corollary 2.2 says that $\tau(G') \leq \tau(G) = \tau$. Therefore, the strong connectivity of G implies that $\tau(G') = \tau$. Suppose G' is not strongly connected. Then, there exists a nontrivial Δ -group G'' such that $\tau(G'')$ is less than τ , and a

surjective Δ -homomorphism $\varphi' : G' \longrightarrow G''$. Then, $\varphi' \circ \varphi$ is a surjective Δ -homomorphism from G onto G'' , contradicting the strong connectivity of G .

The following lemma is useful in showing a group is strongly connected.

Lemma 2.8 *Let G be a Δ -group and let H and N be strongly connected subgroups, with $N \triangleleft G$ and $\tau(H) = \tau(N)$. Then, the Δ -subgroup HN of G is strongly connected and $\tau(HN) = \tau(H) = \tau(N)$. Furthermore, $a_\tau(HN) = a_\tau(H) + a_\tau(N) - a_\tau(H \cap N)$.*

Proof. Without loss of generality, we may assume that $H \not\subset N$. We have $\tau(HN) = \max\{\tau(HN/N), \tau(N)\} = \max\{\tau(H/H \cap N), \tau(N)\}$. Since H is strongly connected and $H \cap N \neq H$, we have $\tau(H/H \cap N) = \tau(H)$. Therefore $\tau(HN) = \tau(H) = \tau(N)$. Remark 2.7.3 implies that H and N are contained in $(HN)_0$. This implies that $HN \subset (HN)_0$ so HN is strongly connected. The final statement concerning the typical dimension (as well as the statement concerning differential type) is contained in ([42] Cor. 4.3, p. 485; see also [25], Cor. 4, p. 109). ■

Examples 2.9 Let \mathbb{U} be an ordinary universal differential field with derivation δ .

1. Let G be the additive group $\mathbb{G}_a(\mathbb{U})$. G has type 1 and typical differential dimension 1. Cassidy [10] showed that any proper Δ -subgroup H is of the form $H = \{a \in k \mid L(a) = 0\}$ for some linear differential operator L . The type of such a group is 0 and its typical differential dimension is d where d is the order of L . Therefore the type of G/H for any proper H is equal to the type of G and so G is strongly connected.

2. Let $G_1 = G$ as above and $G_2 = \mathbb{G}_a(C)$ where $C = \{c \in k \mid \delta(c) = 0\}$. The type of G_2 is 0 and the typical differential dimension is 1. Using equations (1) and (2), we see that $\tau(G_1 \times G_2) = \max(\tau(G_1), \tau(G_2)) = 1$ and $a_\tau(G_1 \times G_2) = a_\tau(G_1) = 1$. The group G_1 is minimal with respect to having the same type and typical differential dimension as $G_1 \times G_2$ and so is the strong identity component of this group. Note that $G_1 \times G_2$ is connected.

To state the analogue of the Jordan-Hölder Theorem, we need two more definitions: the first is the appropriate notion of “simple” and the second is the relevant notion of “equivalent”. The next two subsections deal with these notions.

2.2.2 Almost Simple Groups

Definition 2.10 *An infinite Δ -group G is almost simple if for any normal proper Δ -subgroup H of G we have $\tau(H) < \tau(G)$. A Δ - k -group is almost k -simple if for any normal Δ - k -subgroup H of G we have $\tau(H) < \tau(G)$.*

If G is almost simple, then G is almost k -simple. Note that an almost simple group is strongly connected. We shall use the term *simple* to denote a group (resp., algebraic

group, Δ -group) whose only proper normal subgroup (resp., normal algebraic subgroup, normal Δ -subgroup) is the trivial group, and the term *quasisimple* to denote a group (resp., algebraic group, Δ -group) G such that $G/Z(G)$ is simple and $Z(G)$ is finite, where $Z(G)$ is the center of G . If G is an algebraic k -group (resp. Δ - k -group), we will call G k -simple (k -quasisimple) if every proper normal algebraic k -subgroup (resp. Δ - k -subgroup) is trivial (resp. finite).

Examples 2.11 Let \mathbb{U} be as in Example 2.9 and $C_{\mathbb{U}}$ its field of constants.

1. The group $\mathbb{G}_a(C_{\mathbb{U}})$ has type 0 with only the trivial group as a proper subgroup so it is simple (and so almost simple).
2. The group $\mathbb{G}_m(C_{\mathbb{U}})$ has type 0 with only finite subgroups as proper algebraic (or δ -) subgroups so it is quasisimple.
3. The group $\mathbb{G}_a(\mathbb{U})$ has type 1 and any proper subgroup has type 0 or -1 (if it is $\{0\}$). The proper δ -subgroups of $G = \mathbb{G}_a(\mathbb{U})$ are the kernels of nonzero linear differential operators $L \in \mathbb{U}[\delta]$. It follows, as E. Cartan noted in [8], that every nontrivial δ -homomorphic image of G is isomorphic to G . Therefore $\mathbb{G}_a(\mathbb{U})$ is almost simple.
4. If G is a quasisimple linear algebraic group defined over $C_{\mathbb{U}}$, then, as we shall show in Section 3 that $G(\mathbb{U})$ and $G(C_{\mathbb{U}})$ are quasisimple δ -groups.

In Section 3, we shall show that for ordinary differential fields, these are essentially the only almost simple *linear* δ -groups.

Example 2.12 Let $\Delta = \{\delta_t, \delta_x\}, \mathbb{U}$ be a universal Δ -field and z a differential indeterminate. Let $G \subset \mathbb{G}_a(\mathbb{U})$ be the additive group of solutions of $\delta_t y = f(y)$, $f \in \mathbb{U}[\delta_x]$. Following Proposition 2.45 of [43], we show that G is almost simple. The type of G is 1 (see also Example 2.31). If $H \subset G$ is a proper Δ -subgroup of G , then there would exist a linear operator $g \in \mathbb{U}[\Delta]$ such that $g(z)$ is not in the differential ideal $[\delta_t z - f(z)]$ generated by $\delta_t z - f(z)$ in $\mathbb{U}\{z\}$ and such that $H \subset \{y \in G \mid g(y) = 0\}$. We may assume that $g \in \mathbb{U}[\delta_x]$. If g has order d , then for any $y \in H$, we have $k(y, \delta_x y, \delta_x^2 y, \dots) = k(y, \delta_x y, \dots, \delta_x^{d-1} y)$. Since $\delta_t y \in k(y, \delta_x y, \dots, \delta_x^{d-1} y)$, this field equals $k \langle y \rangle$ which must therefore be of finite transcendence degree over k . Therefore H has type 0 and G is almost simple.

It is not hard to see that a finite normal subgroup of a connected algebraic group must be central. The next proposition is an analogous result for strongly connected Δ -groups.

Proposition 2.13 *Let G be a strongly connected Δ -group. Then every normal Δ -subgroup of smaller type is central.*

We immediately have the following corollaries.

Corollary 2.14 *Let G be an almost simple Δ -group. Any proper normal Δ -subgroup is central.*

Proof. For any Δ -group G , the strong identity component, G_0 , is a normal Δ -subgroup of the same type, and so $G = G_0$, that is, G is strongly connected. By definition any proper normal Δ -subgroup is of smaller type. ■

If G is an almost k -simple Δ - k -group, the strong identity component is a normal Δ - k -subgroup of the same type, and so, $G = G_0$. By definition, any proper normal Δ - k -subgroup has smaller type, and hence is central.

Corollary 2.15 *Let G be an almost simple Δ -group. Then $G/Z(G)$ is a simple Δ -group.*

Proof of Proposition 2.13: Let N be a normal Δ -subgroup of G . We define a Δ -rational map

$$\alpha : G \times N \rightarrow N$$

by the formula

$$\alpha(g, a) = gag^{-1}, \quad g \in G, a \in N.$$

One checks that $\alpha(gh, a) = \alpha(g, \alpha(h, a))$ and $\alpha(1, a) = a$ and so α defines an action of the Δ -group G on N . For fixed $a \in N$, the map $\alpha_a(g) = gag^{-1}$ is a Δ -rational map from G to N that is constant on the left cosets of $Z_G(a) = \{g \in G \mid gag^{-1} = a\}$. Note that $Z_G(a)$ is a Δ -subgroup of G ([25], Section IV.4, Cor. 2(b)).

Theorem 3, p.105 of [25] implies that there is a Δ -morphism $\beta : G/Z_G(a) \rightarrow N$ such that $\pi \circ \beta = \alpha_a$, where π is the canonical quotient map. The image of $G/Z_G(a)$ under β is the same as the image of G under α_a , namely Ga . Let V be the Δ -closure of Ga .

Note that $\alpha_a(g) = \alpha_a(h)$ implies that $gag^{-1} = hah^{-1}$ so $h^{-1}g \in Z_G(z)$. Therefore β is an injective Δ -map. Corollary 2.4 implies that $\tau(G/Z_G(a)) \leq \tau(N) < \tau(G)$. Since G is strongly connected, we have $Z_G(a) = G$. ■

2.2.3 Isogeny

Definition 2.16 1) *Let G, H be strongly connected Δ - k -groups. A Δ - k -morphism $\phi : G \rightarrow H$ is an isogeny if it is surjective and $\tau(\ker \phi) < \tau(G)$.*

2) *Two strongly connected Δ - k -groups H_1, H_2 are Δ - k -isogenous if there exists a strongly connected Δ - k -group G and isogenies $\phi_i : G \rightarrow H_i$ for $i = 1, 2$.*

The notion of isogenous in the present context generalizes the notion of isogenous in the theory of algebraic groups (cf., [37]) where two connected algebraic groups H_1, H_2 are isogenous if there is an algebraic group G and morphisms $\phi_i : G \rightarrow H_i$ for $i = 1, 2$ with finite kernel.

Example 2.17 Let $G = \mathrm{SL}_6(\mathbb{C})$. The center of G is isomorphic to the cyclic group of order 6. Let C_2 and C_3 be cyclic subgroups of this center of orders 2 and 3 respectively. Let $H_1 = \mathrm{SL}_6(\mathbb{C})/C_2$ and $H_2 = \mathrm{SL}_6(\mathbb{C})/C_3$ and let $\phi_i : G \rightarrow H_i$ be the obvious projections. These show H_1 and H_2 are isogenous in both the category of algebraic groups and the category of Δ -groups. Note that they are not isomorphic since their centers are finite groups of different orders.

Remarks 2.18 1. If G_1 and G_2 are strongly connected Δ -groups and there is an isogeny $\phi : G_1 \rightarrow G_2$, then G_1 and G_2 are isogenous. However, the converse is false, as the above example shows.

2. If $\phi : G_1 \rightarrow G_2$ is an isogeny then equations (1) and (2) imply that $\tau(G_1) = \tau(G_2)$ and $a_\tau(G_1) = a_\tau(G_2)$.

3. The composite of isogenies of strongly connected Δ -groups is an isogeny.

4. If G_1 and G_2 are strongly connected of the same type, then it is easy to see that $G_1 \times G_2$ is strongly connected. Suppose G'_1 and G'_2 are strongly connected and $\phi_i : G_i \rightarrow G'_i$ is an isogeny, $i = 1, 2$. Then, $\phi_1 \times \phi_2 : G_1 \times G_2 \rightarrow G'_1 \times G'_2$, with kernel $\ker \phi_1 \times \ker \phi_2$ is an isogeny.

Proposition 2.19 *Let G_1 and G_2 be strongly connected Δ - k -groups. The following are equivalent:*

1. *There exists a strongly connected Δ - k -group H and isogenies $\phi_1 : H \rightarrow G_1$ and $\phi_2 : H \rightarrow G_2$:*

$$\begin{array}{ccc} & H & \\ \phi_1 \swarrow & & \searrow \phi_2 \\ G_1 & & G_2 \end{array} .$$

2. *There exists a strongly connected Δ - k -group K and isogenies $\psi_1 : G_1 \rightarrow K$ and $\psi_2 : G_2 \rightarrow K$:*

$$\begin{array}{ccc} G_1 & & G_2 \\ \psi_1 \searrow & & \swarrow \psi_2 \\ & K & \end{array} .$$

Proof. Note that all the Δ -groups and Δ -homomorphisms are defined over k . Assume the first statement. Put $H_1 = \phi_1(\ker \phi_2)$, and $H_2 = \phi_2(\ker \phi_1)$. Clearly, $H_1 = \phi_1(\ker \phi_1 \ker \phi_2)$, and $H_2 = \phi_2(\ker \phi_1 \ker \phi_2)$. Therefore,

$$\begin{aligned} G_1/H_1 &= \phi_1(H) / \phi_1(\ker \phi_2) = \phi_1(H) / \phi_1(\ker \phi_1 \ker \phi_2) = H / (\ker \phi_1 \ker \phi_2) \\ G_2/H_2 &= \phi_2(H) / \phi_2(\ker \phi_2) = \phi_2(H) / \phi_2(\ker \phi_1 \ker \phi_2) = H / (\ker \phi_1 \ker \phi_2). \end{aligned}$$

Set $K = H / (\ker \phi_1 \ker \phi_2)$. Since K is the image of the strongly connected group H , it is strongly connected. Since $\tau(\ker \phi_1 \ker \phi_2) < \tau(H)$, the projections $\psi_i : G_i \rightarrow G_i/H_i \simeq K$

are isogenies.

Now assume the second statement. Let $G = \{(g_1, g_2) \in G_1 \times G_2 \mid \psi_1(g_1) = \psi_2(g_2)\}$, that is, the pull-back or fiber product. One sees that the natural projections $\phi_1 : G \rightarrow G_1$ and $\phi_2 : G \rightarrow G_2$ are surjective. This implies that $\tau(G) \geq \tau(G_1)$. The kernel of the ϕ_i 's is contained in $\ker \psi_1 \times \ker \psi_2$ and so $\tau(\ker \phi_1) \leq \tau(\ker \psi_1 \times \ker \psi_2) < \tau(G_1)$. Since $G_1 \simeq G / \ker(\phi_1)$ we have $\tau(G) = \tau(G_1)$ and ϕ_1 is an isogeny. Similar reasoning shows that ϕ_2 is an isogeny. \blacksquare

The following result is a reworking of Theorem 6, [37] in the context of Δ - k -groups.

Proposition 2.20 *Isogeny is an equivalence relation on the set of strongly connected Δ - k -groups. If G_1 and G_2 are isogenous strongly connected Δ - k -groups, with $\tau = \tau(G_1) = \tau(G_2)$, then there is a bijective correspondence between the strongly connected Δ - k -subgroups of G_1 with type τ and those of G_2 with type τ . Let J_1 and K_1 be Δ - k -subgroups of G_1 corresponding to the Δ - k -subgroups J_2 and K_2 of G_2 . Then, $J_1 \subseteq K_1$ if and only if $J_2 \subseteq K_2$, and $J_1 \trianglelefteq K_1$ if and only if $J_2 \trianglelefteq K_2$. In this case, the strongly connected Δ - k -groups K_1/J_1 and K_2/J_2 are isogenous and have type τ . If G_1, G_2 , and G'_1, G'_2 are pairs of isogenous Δ - k -groups, then the Δ - k -groups $G_1 \times G'_1$ and $G_2 \times G'_2$ are isogenous and have type τ .*

Proof. Since reflexivity and symmetry follow immediately from the definition, we need only show that isogeny is transitive. So, let $G_1, G_2, H, G_2, G_3, H'$ be strongly connected Δ - k -groups with type τ , and let $\phi_i : H \rightarrow G_i, i = 1, 2, \phi'_i : H' \rightarrow G_i, i = 2, 3$, be isogenies. Then, the Δ - k -group $H \times H'$ is strongly connected, and $\phi_2 \times \phi'_2 : H \times H' \rightarrow G_2 \times G_2$ is surjective. Let $K = ((\phi_2 \times \phi'_2)^{-1}(\text{Diag } G_2 \times G_2))_0$, where $(\dots)_0$ denotes the strong identity component. Then, K is a strongly connected Δ - k -group. Let $\pi_H, \pi_{H'}$, be the projections from $H \times H'$ to H , and to H' . We have the following diagram

$$\begin{array}{ccccc}
 & & K & & \\
 & & \swarrow \pi_H & & \searrow \pi_{H'} \\
 & H & & & H' \\
 \phi_1 \swarrow & & \phi_2 \searrow & & \phi'_2 \swarrow & \phi'_3 \searrow \\
 G_1 & & G_2 & & G_3
 \end{array}$$

We claim that the restrictions of the Δ - k homomorphisms π_H and $\pi_{H'}$ to K are isogenies. Since the composition of isogenies is an isogenies, our claim implies that G_1 and G_3 are isogenous and so the isogeny relation is an equivalence relation.

To prove our claim, first note that (1) implies that $\tau(H \times H') = \tau(H) = \tau(H') = \tau(G_2 \times G_2)$, and $\tau(\ker(\phi_2 \times \phi'_2)) = \tau(\ker \phi_2 \times \ker \phi'_2) < \tau(H \times H')$. If $(h, h') \in K \cap \ker \pi_H$ then $h = 1$ and, since $\phi_2(h) = \phi'_2(h')$, we have $h' \in \ker \phi'_2$. Therefore $\ker \pi_H = 1 \times \ker \phi'_2$ and so $\tau(\ker \pi_H) < \tau(H)$. Similarly, $\tau(\ker \pi_{H'}) < \tau(H')$. We know that

$\tau(K) = \tau(G_2) = \tau(H)$. Since $\tau(\ker \pi_H) < \tau(H)$, it follows that $a_\tau(\pi_H(K)) = a_\tau(H)$. Since H is strongly connected, $\pi_H(K) = H$, and, therefore, π_H is an isogeny. Similarly, $\pi_{H'}$ is an isogeny and this shows that the isogeny relation is an equivalence relation on the set of strongly connected Δ - k -groups.

Suppose we have the following isogeny diagram of strongly connected Δ - k -groups:

$$\begin{array}{ccc} & H & \\ \phi_1 \swarrow & & \searrow \phi_2 \\ G_1 & & G_2 \end{array}.$$

We define the following correspondence between strongly connected Δ - k -subgroups of G_1 of type $\tau = \tau(G_1)$ and strongly connected Δ - k -subgroups of G_2 of type $\tau = \tau(G_2)$: Subgroups J_1 of G_1 and J_2 of G_2 correspond if there exists a strongly connected Δ - k -subgroup P of H such that $\phi_i|_P$ maps P to J_i and is an isogeny, $i = 1, 2$. We claim that given J_1 , there is a unique choice for P , namely, the strong identity component of $\phi_1^{-1}(J_1)$, which is, of course, a strongly connected Δ - k -subgroup of H . First note that for any such P , Remark 2.18.2 implies that $\tau(P) = \tau$ and $a_\tau(P) = a_\tau(J_1) = a_\tau(J_2)$. Furthermore any such P must lie in $\phi_1^{-1}(J_1)$ and, since it is assumed to be strongly connected, it must be contained in the strong identity component $(\phi_1^{-1}(J_1))_0$ of $\phi_1^{-1}(J_1)$. Since $\tau(\ker \phi_1) < \tau$ we have that $\tau(\ker \phi_1 \cap \phi^{-1}(J_1)) < \tau$ and so ϕ restricted to $(\phi_1^{-1}(J_1))_0$ is an isogeny. In particular $\tau((\phi_1^{-1}(J_1))_0) = \tau$ and $a_\tau((\phi_1^{-1}(J_1))_0) = a_\tau(G_1)$. Since $P \subset (\phi_1^{-1}(J_1))_0$ and both of these groups are strongly connected we have $P = (\phi_1^{-1}(J_1))_0$. A similar argument shows that $P = (\phi_2^{-1}(J_2))_0$. It follows easily that the correspondence is unique.

Now, we shall show that if $J_i, K_i, i = 1, 2$, are corresponding strongly connected Δ - k -subgroups of G_i of type τ , then

$$J_1 \subseteq K_1 \iff J_2 \subseteq K_2.$$

Let $P, Q, i = 1, 2$, be the unique strongly connected Δ - k -subgroups of H of type τ such that $\phi_i(P) = J_i$ and $\phi_i(Q) = K_i$. P is the strong identity component of $\phi_i^{-1}(J_i)$ and Q is the strong identity component of $\phi_i^{-1}(K_i)$. $J_i \subseteq K_i$ implies that $\phi_i^{-1}(J_i) \subseteq \phi_i^{-1}(K_i)$. $\tau(P) = \tau(Q) = \tau$. Since P is a strongly connected Δ - k -subgroup of $\phi_i^{-1}(K_i)$ of the same type as the latter, it follows, as above, that P is contained in its strong identity component Q . The reverse implication is proved the same way. Now suppose that J_1 is normal in K_1 . Then, $\phi_1^{-1}(J_1)$ is normal in $\phi_1^{-1}(K_1)$ and has the same type. By Remark 2.7.2, P is normal in $\phi_1^{-1}(K_1)$. Therefore, P is normal in Q . Since ϕ_2 is surjective, J_2 is normal in K_2 . To finish the proof, we need only show that the Δ - k -groups K_1/J_1 and K_2/J_2 are isogenous and have the same type τ . First, note that Remark 2.7.5 says that K_i/J_i are strongly connected and have type τ . We now define a map $\phi : Q/P \rightarrow K_1/J_1$ by $\phi(q \bmod P) = \phi_1(q) \bmod J_1$ for any $q \in Q$. To see that this is well defined, let $q_1, q'_1 \in Q$, and suppose $q_1 \bmod P = q'_1 \bmod P$. Then, $\phi_1(q_1 - q'_1) \in J_1$. Therefore, $\phi_1(q_1) \bmod J_1 = \phi_1(q'_1) \bmod J_1$. So, the map $\phi : Q/P \rightarrow K_1/J_1$ is well defined. It is clearly a homomorphism. We claim that it is surjective. Let $k_1 \in K_1$. There exists $q \in Q$ such

that $\phi_1(q) = k_1$ and so $\phi(q \bmod P) = k_1 \bmod J_1$. A calculation shows that the kernel of ϕ is $[Q \cap \phi_1^{-1}(J_1)]/P$. The strong identity component P of $\phi_1^{-1}(J_1)$ is contained in $Q \cap \phi_1^{-1}(J_1)$. Thus, $\tau([Q \cap \phi_1^{-1}(J_1)]/P) < \tau(\phi_1^{-1}(J_1)) = \tau(Q) = \tau(Q/P)$ since Q is strongly connected. So, $\phi : Q/P \rightarrow K_1/J_1$ is an isogeny. Similar reasoning shows that one can define a map $\psi : Q/P \rightarrow K_2/J_2$ and that it too is an isogeny. The above proof also shows that J_1 is defined over k if and only if J_2 is defined over k . ■

An immediate corollary is:

Corollary 2.21 *Let G_1 and G_2 be isogenous strongly connected Δ - k -groups. Either both are almost simple or neither is.*

Proof. Suppose G_1 is almost simple. Let N be a normal Δ -subgroup of G_2 . There exists a finitely generated Δ - k -subfield k' of \mathbb{U} such that N is defined over k' . Since k' is finitely Δ -generated over k , \mathbb{U} is universal over k' . Moreover, as we remarked in Section 2.1, the type of G_2 is invariant under the choice of field of definition. If the type of N is $\tau = \tau(G_2) = \tau(G_1)$, Proposition 2.20 gives us a proper normal Δ - k' -subgroup of G_1 of type τ , contradicting the almost simplicity of G_1 . ■

Another corollary is

Corollary 2.22 *Let G_1 and G_2 be isogenous strongly connected Δ -groups. Either both are commutative or neither is.*

The proof of Corollary 2.22 depends on the following concept and lemma.

Definition 2.23 *Let G be a Δ - k -group. The differential commutator group $D_\Delta(G)$ is the smallest Δ -subgroup of G containing the commutator subgroup of G . It is defined over k .*

An example of Cassidy (*cf.*, p.111 of [25]) shows that $D_\Delta(G)$ may be strictly larger than the commutator subgroup of G .

Lemma 2.24 *Let G be a Δ - k -group. If G is strongly connected and not commutative, then $\tau(D_\Delta(G)) = \tau(G)$.*

Proof. Let a be in G . Let $c_a : G \rightarrow G$ be the Δ -map that sends x to $axa^{-1}x^{-1}$. If $D_\Delta(G)$ has smaller type than G , then Proposition 2.13 implies that $D_\Delta(G)$ is contained in the center $Z(G)$. In this case we have $axya^{-1}y^{-1}x^{-1} = axa^{-1}(aya^{-1}y^{-1})x^{-1} = (axa^{-1}x^{-1})(aya^{-1}y^{-1})$ and therefore c_a is a Δ -homomorphism of G into $D_\Delta(G)$. Since G is strongly connected, $c_a(G) = \{1\}$. Thus, $a \in Z(G)$. It follows that G is commutative. ■

Proof of Corollary 2.22: Let K be a Δ - k -group such that for G_1, G_2 , and K we have the commutative diagram:

$$\begin{array}{ccc} G_1 & & G_2 \\ & \psi_1 \searrow & \swarrow \psi_2 \\ & & K \end{array}$$

with isogenies ψ_i . Assume G_1 is commutative. Then, K is commutative. Suppose G_2 is not commutative. Lemma 2.24 implies that $\tau(D(G_2)) = \tau(G_2) = \tau(K)$. Since ψ_2 is an isogeny, $\tau(\psi_2(D_\Delta(G_2))) = \tau(K)$ as well. Since $\psi_2(D_\Delta(G_2)) \subset D_\Delta(K)$ we have that $D_\Delta(K) \neq \{1\}$, a contradiction. \blacksquare

We give two examples of infinite families of isogenous, pairwise nonisomorphic Δ -groups.

Example 2.25 Let $\Delta = \{\delta\}$ and \mathbb{U} be a Δ -universal field with Δ -constants C . We shall exhibit an infinite number of almost simple Δ -groups that are isogenous to $\mathbb{G}_a(\mathbb{U})$ but are pairwise nonisomorphic. For each $n \in \mathbb{N}$, let

$$G_n = \left\{ g(a, b) = \begin{pmatrix} a & 0 & 0 \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix} \mid a \in \mathbb{G}_m(\mathbb{U}), b \in \mathbb{G}_a(\mathbb{U}), (\delta a)a^{-1} = \delta^n(b) \right\}.$$

The map $\alpha : G_n \rightarrow \mathbb{G}_a(\mathbb{U})$ given by $\alpha(g(a, b)) = (\delta a)a^{-1}$ is a surjective homomorphism with kernel

$$K_n = \{g(a, b) \mid \delta a = 0, \delta^n(b) = 0\}.$$

Note that G_n has differential type 1 and that K_n has differential type 0; so α is an isogeny and therefore G_n is isogenous to $\mathbb{G}_a(\mathbb{U})$.

We shall now show that G_n is strongly connected. Let $(G_n)_0$ be the strong identity component of G_n and let $\pi_1(g(a, b)) = a$ and $\pi_2(g(a, b)) = b$ be the projections of G_n onto $\mathbb{G}_m(\mathbb{U})$ and $\mathbb{G}_a(\mathbb{U})$ respectively. Since these latter groups are strongly connected, these projections are surjective when restricted to $(G_n)_0$ as well. Using π_1 we see that for any $g(a, b) \in G_n$ there exists $b_1 \in \mathbb{G}_a(\mathbb{U})$ such that $g(a, b_1) \in (G_n)_0$ and therefore $g(a, b) = g(1, b - b_1)g(a, b_1)$. Note that $\delta^n(b - b_1) = 0$. This implies that the group $H_n = \{b \in \mathbb{G}_a(\mathbb{U}) \mid \delta^n(b) = 0\}$ is mapped surjectively onto $G_n/(G_n)_0$ via the map $b \mapsto g(1, b) \bmod (G_n)_0$. Using π_2 , a similar argument shows that the map $c \mapsto g(c, 0) \bmod (G_n)_0$ maps $\mathbb{G}_m(C) = \{c \in \mathbb{U} \mid c \neq 0, \delta(c) = 0\}$ surjectively onto $G_n/(G_n)_0$. Therefore $G_n/(G_n)_0$ is isomorphic to a quotient of H_n and a quotient of $\mathbb{G}_m(C)$. Taking a fundamental system of solutions of the differential equation $\delta^n = 0$, we see that H_n is isomorphic as a Δ -group to $(C^n, +)$. Therefore, any quotient Δ -group is torsion-free. This implies that $G_n/(G_n)_0$ is torsion-free. But, the Δ -group $G_n/(G_n)_0$ is also a quotient of $\mathbb{G}_m(C)$, whose torsion group is Kolchin dense. Therefore, either the torsion group of $G_n/(G_n)_0$ is Kolchin dense, or $G_n/(G_n)_0$ is trivial. Since the latter must hold, G_n is strongly connected.

Since $\mathbb{G}_a(\mathbb{U})$ is almost simple, Corollary 2.21 implies that G_n is almost simple.

We now claim that for $n \neq m, n, m \in \mathbb{N}$, G_n and G_m are not Δ -isomorphic. To see this, note first that the set of unipotent matrices in any G_n is precisely $\{g(1, b) \mid \delta^n(b) = 0\}$. This is a Δ -group of differential type 0 and typical differential dimension equal to n . The property of being unipotent is preserved under Δ -isomorphism ([10], Proposition 35) so our claim is proved. Note also that there does not exist an isogeny from $\mathbb{G}_a(\mathbb{U})$ onto G_n , and, therefore, this example does not contradict Example 2.11.3.

Example 2.26 Let \mathbb{C} be the field of complex numbers, and let k be the subfield $\mathbb{C} \left(x, t, \{e^{c^2 t}, e^{icx}\}_{c \in \mathbb{C}} \right)$ of the field \mathbb{M} of functions meromorphic on \mathbb{C}^2 . Let $\Delta = \{\delta_x, \delta_t\}$, where δ_x acts on \mathbb{M} as $\frac{\partial}{\partial x}$ and δ_t acts as $\frac{\partial}{\partial t}$. Then, \mathbb{M} is a Δ -field, and k is a Δ -subfield of \mathbb{M} . Let \mathbb{U} be a Δ -field universal over k , and with field C of constants, as usual. We do not assume that \mathbb{M} is a subfield of \mathbb{U} . Let

$$H = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u - \delta_x^2 u = 0\}$$

be the subgroup of $\mathbb{G}_a(\mathbb{U})$ defined by the Heat Equation. We shall exhibit an infinite family of almost simple Δ -groups that are isogenous to H but not Δ -isomorphic to H . In particular, we shall show that if $\lambda \in \mathbb{U}$ is any nonzero solution of $\delta_t \lambda + \delta_x^2 \lambda = 0$, then H is isogenous to

$$H_\lambda = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u - \delta_x^2 u + 2 \frac{\delta_x \lambda}{\lambda} \delta_x u = 0\}$$

and that there are an infinite number of $\lambda \in k$ whose corresponding groups are pairwise not Δ -isomorphic.

The groups H and H_λ have differential type 1 and, from Example 2.12, we know that they are almost simple. An elementary calculation shows that the map $\phi(u) = (1/\lambda)\delta_x u$ is a homomorphism from H_λ onto H . The kernel of ϕ is $\mathbb{G}_a(C)$. Therefore these groups are isogenous.

For any nonzero $c \in \mathbb{C}$,

$$\lambda_c = e^{c^2 t} \cos cx$$

is a solution in k of $\delta_t \lambda + \delta_x^2 \lambda = 0$. We define

$$H_c = H_{\lambda_c} = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u = \delta_x^2 u + 2(c \tan cx) \delta_x u\}.$$

Each of these groups is isogenous to H so they are all pairwise isogenous. We shall show that for nonzero $c, d \in C$ with $c/d \notin \mathbb{Q}$, the groups H_c and H_d are not isomorphic. The proof of this claim follows from the special nature of the differential coordinate rings $\mathbb{U}\{H_c\}_\Delta$ and $\mathbb{U}\{H_d\}_\Delta$ of the groups.

$$\begin{aligned} \mathbb{U}\{H_c\}_\Delta &= \mathbb{U}\{z\}_\Delta, \text{ where } \delta_t z = \delta_x^2 z + 2(c \tan cx) \delta_x z, \\ \mathbb{U}\{H_d\}_\Delta &= \mathbb{U}\{y\}_\Delta, \text{ where } \delta_t y = \delta_x^2 y + 2(d \tan dx) \delta_x y. \end{aligned}$$

Note that both groups are defined over k . The form of the defining differential equations imply that

$$\mathbb{U}\{H_c\}_\Delta = \mathbb{U}\{H_c\}_{\{\delta_x\}} = \mathbb{U}[z, \delta_x z, \dots, \delta_x^n z, \dots] \text{ and } \mathbb{U}\{H_d\}_\Delta = \mathbb{U}\{H_d\}_{\{\delta_x\}} = \mathbb{U}[y, \delta_x y, \dots, \delta_x^n y, \dots].$$

where y and its δ_x -derivatives are algebraically independent over \mathbb{U} and a similar statement is true for z . As δ_x -rings these two rings are δ_x -isomorphic to a δ_x -differential polynomial ring $\mathbb{U}\{Y\}_{\delta_x}$ and this will be a key fact in the proof of our claim. What distinguishes these rings is their δ_t -structure, which is given by the defining equations of the groups. The derivation operator δ_t acts as an "evolutionary derivation" on the δ_x -differential ring.

Let us assume (with the aim of arriving at a contradiction) that H_c and H_d are isomorphic as Δ -groups and let $\psi : H_d \rightarrow H_c$ be a differential isomorphism. This isomorphism induces a Δ -isomorphism $\psi^* : \mathbb{U}\{H_c\}_\Delta \rightarrow \mathbb{U}\{H_d\}_\Delta$, defined by the formula $\psi^*(f) = f$ composed with ψ . Note that ψ^* restricted to U is the identity automorphism. Identifying both $\mathbb{U}\{H_c\}$ and $\mathbb{U}\{H_d\}_\Delta$ with the δ_x -ring $\mathbb{U}\{Y\}_{\{\delta_x\}}$, the map ψ^* induces a δ_x -automorphism of $\mathbb{U}\{Y\}_{\{\delta_x\}}$ and *a fortiori* of the quotient field $\mathbb{U} \langle Y \rangle_{\{\delta_x\}}$. The δ_x - \mathbb{U} -automorphisms of $\mathbb{U} \langle Y \rangle_{\delta_x}$ have been classified by Ritt (as part of his differential Lüroth theorem) in his book ([36], p. 56). He shows that any δ_x -automorphism σ of this latter field must be of the form

$$\sigma(Y) = \frac{aY + b}{cY + d}$$

where $a, b \in \mathbb{U}$. It is not hard to see that if σ furthermore maps $\mathbb{U}\{Y\}_{\{\delta_x\}}$ to itself, we must have that $\sigma(Y) = aY + b$ for some $a, b \in \mathbb{U}$. In particular, ψ^* must be of the form $\psi^*(z) = ay + b$ for some $a, b \in \mathbb{U}$ and, since ψ is assumed to be a homomorphism of additive groups as well, we must have $b = 0$ and so $\psi^*(z) = ay$. The following calculation will show that this leads to a contradiction.

We have that

$$\begin{aligned} \delta_t z &= \delta_x^2 z + 2(c \tan cx) \delta_x z \\ \delta_t y &= \delta_x^2 y + 2(d \tan dx) \delta_x y \end{aligned}$$

Note that c goes with the domain of the comorphism and d with its range. Calculating $\delta_t(\psi^*(z))$ and $\psi^*(\delta_t z)$ we have

$$\begin{aligned} \delta_t(\psi^*(z)) &= a\delta_x^2 y + (2ad(\tan dx))\delta_x y + \delta_t ay \\ \psi^*(\delta_t z) &= a\delta_x^2 y + (2\delta_x a + 2ac(\tan cx))\delta_x y + (\delta_x^2 a + 2c(\tan cx)\delta_x a)y \end{aligned}$$

Since $\delta_t(\psi^*(z)) = \psi^*(\delta_t z)$ and $y, \delta_x y, \delta_x^2 y$ are algebraically independent, we must have

$$ad(\tan dx) = \delta_x a + ac(\tan cx) \quad (3)$$

$$\delta_t a = \delta_x^2 a + 2c(\tan cx)\delta_x a \quad (4)$$

Equation (3) implies that

$$a = \alpha \frac{\cos cx}{\cos dx}$$

where α is in the field of constants of δ_x in \mathbb{U} . Inserting this expression into equation (4) we have

$$\frac{\delta_t \alpha}{\alpha} = c^2 + \frac{2d^2}{\cos^2 dx} - d^2 - \frac{2c^2}{\cos^2 cx}. \quad (5)$$

Since $\delta_x(\frac{\delta_t \alpha}{\alpha}) = 0$, we must have that the right-hand side of equation (5) is some constant $\gamma \in C$. A linear disjointness argument implies that γ is in the field of constants of k . In particular, it is a complex number. So, the terms of equations (6) and (7) are complex functions. Evaluating the right-hand side at $x = 0$ and $x = \frac{2\pi}{c}$, we have

$$d^2 - c^2 = \gamma \quad (6)$$

$$\left(\frac{2}{\cos^2(2d\pi/c)} - 1\right)d^2 - c^2 = \gamma \quad (7)$$

Note that $\cos(2d\pi/c) \neq 0$ since we are assuming that d/c is neither an integer nor half an integer. From these equations, we can conclude that $\cos^2(2d\pi/c) = 1$. Therefore, $\cos(2d\pi/c) = 1$ or -1 . So, properties of the complex cosine function tell us that $2\pi d/c$ equals $2n\pi$ or $(2n+1)\pi$. So, d/c is either an integer or d/c is half an integer, a contradiction. Therefore, for any set of constants S , whose members are pairwise independent over the integers, this yields a family $\{H_c\}_{c \in S}$ of isogenous Δ -subgroups of \mathbb{G}_a that are pairwise nonisomorphic.

We now turn to another choice of λ : $\lambda = x$. This situation was already considered by Cartan ([8], p.145-146). Letting

$$H_x = \{u \in \mathbb{G}_a(\mathbb{U}) \mid \delta_t u = \delta_x^2 u - \frac{2}{x} \delta_x u\},$$

Cartan states (without proof) that H and H_x are not “isomorphes holoédriques”. We shall show that H and H_x are not Δ -isomorphic.

Let us assume that H and H_x are isomorphic as Δ -groups and let $\psi : H \rightarrow H_x$ be a differential isomorphism. Let

$$\mathbb{U}\{H\}_\Delta = \mathbb{U}[y, \delta_x y, \dots, \delta_x^n y, \dots] \text{ and } \mathbb{U}\{H_x\}_\Delta = \mathbb{U}[z, \delta_x z, \dots, \delta_x^n z, \dots]$$

where y and its δ_x -derivatives are algebraically independent over \mathbb{U} and a similar statement is true for z . As above, we have a differential isomorphism $\psi^* : \mathbb{U}\{H_x\}_\Delta \rightarrow \mathbb{U}\{H\}_\Delta$ and we may assume that $\psi^*(z) = ay$. The following calculation will show that this leads to a contradiction. Since ψ^* is also a Δ - \mathbb{U} -isomorphism, we must have that $\psi^*(\delta_t z) = \delta_t(\psi^*(z))$. We have

$$\psi^*(\delta_t z) = a\delta_x^2 y + (2\delta_x a - \frac{2}{x}a)\delta_x y + (\delta_x^2 a - \frac{2}{x}\delta_x a)y \quad (8)$$

$$\delta_t(\psi^*(z)) = a\delta_x^2 y + (\delta_t a)y. \quad (9)$$

Since $\delta_x^2 y, \delta_x y$ and y are algebraically independent over \mathbb{U} , comparing coefficients of like terms in (8) and (9), we have

$$x\delta_x a - a = 0 \quad (10)$$

$$x\delta_x^2 a - 2\delta_x a - x\delta_t a = 0. \quad (11)$$

Equation (10) implies that $a = cx$ where c is a δ_x -constant in \mathbb{U} . $\delta_x c = 0$. Therefore $x\delta_x^2 a = 0$. This, combined with equation (11), implies that $0 = 2\delta_x a + x\delta_t a = 2c + (\delta_t c)x^2$. Therefore $0 = \delta_x(2c + (\delta_t c)x^2) = 2\delta_t cx$, so $\delta_t c = c = 0$. We therefore have that $a = 0$. This contradicts the fact that ψ^* is an isomorphism and completes the proof that H and H_x are not Δ -isomorphic.

The Heat Equation $\delta_t y = \delta_x^2 y$ and the related equations $\delta_t z = \delta_x^2 z - \frac{\delta_t \lambda}{\lambda} \delta_x z$ arise from the study of conservation laws and generalized symmetries (cf., [5], [33]), which we now briefly describe. Let L be a homogeneous linear Δ -polynomial. A vector $\mathbf{F} = (T(y), X(y))$ of differential polynomials is called a *conserved vector* if $\nabla \mathbf{F}(u) = \delta_t T(u) + \delta_x X(u) = 0$ for all solutions of $L(y) = 0$. The equation $\delta_t T(u) + \delta_x X(u) = 0$ is called a *conservation law* for $L(y) = 0$. If L is the Heat Equation, then it follows from [33] that the conservation laws for $L(y) = 0$ are generated (as a C -vector space) by conservation laws corresponding to conserved vectors of the form

$$\mathbf{F} = (\lambda y, -\lambda \delta_x y + (\delta_x \lambda) y) \quad (12)$$

where λ satisfies $\delta_y \lambda + \delta_x^2 \lambda = 0$. In [5], Bluman and Kumei introduce the notion of a *potential symmetry* corresponding to each conservation law. The potential symmetry corresponding to the conserved vector (12) is given by the system

$$\begin{aligned} \delta_x z &= \lambda y \\ \delta_t z &= \lambda \delta_x y - (\delta_x \lambda) y. \end{aligned}$$

This latter system defines a Δ -group $G \subset \mathbb{G}_a \times \mathbb{G}_a$ that has been extensively studied for $\lambda = 1$ and $\lambda = x$. Letting π_1 and π_2 be the projections of $\mathbb{G}_a \times \mathbb{G}_a$ onto the first and second coordinates of (y, z) , one can show that $\pi_1(G) = H$, $\pi_2(G) = H_\lambda$ and the π_i are isogenies. This gives another proof that the groups H and H_λ are isogenous.

Further studies of the algebraic and model-theoretic structure of the solutions of the Heat Equation and related differential algebraic groups can be found in the thesis [43] of Sonat Suer.

2.3 Jordan-Hölder Theorem

We can now state and prove the following analogue of the Jordan-Hölder Theorem (cf., Theorem 6, [37]).

Theorem 2.27 *Let G be a strongly connected Δ -group with $\tau(G) = \tau \neq -1$ (i.e., $G \neq \{1\}$). There exists a normal sequence $G = G_0 \triangleright G_1 \triangleright \dots \triangleright G_r = \{1\}$ of Δ -subgroups such that for each $i = 0, \dots, r-1$*

1. G_i is a strongly connected group with $\tau(G_i) = \tau$,

2. $a_\tau(G_0) > a_\tau(G_1) > \dots > a_\tau(G_{r-1})$, and

3. G_i/G_{i+1} is almost simple of type τ and typical differential dimension $a_\tau(G_i) - a_\tau(G_{i+1})$.

If $G = H_0 \triangleright H_1 \triangleright \dots \triangleright H_s = \{1\}$ is another normal sequence as above, then $r = s$ and there exists a permutation π of $\{0, \dots, r\}$ such that $G_{\pi(i)}/G_{\pi(i)+1}$ is k -isogenous to H_i/H_{i+1} .

Proof. If G is not almost simple, there exists a normal proper Δ -subgroup H with $\tau(H) = \tau(G) = \tau$. Since G is strongly connected, we have that $a_\tau(H) < a_\tau(G)$. Among such subgroups let H have largest $a_\tau(H)$ and let G_1 be the strong identity component of H . Note that, by definition, $\tau(G_1) = \tau(H)$. Since G_1 is a characteristic subgroup of H , it is normal in G . We have that $a_\tau(G) = a_\tau(G/G_1) + a_\tau(G_1)$ so $a_\tau(G/G_1) \neq 0$. Therefore $\tau(G/G_1) = \tau$. For any normal Δ -subgroup K of G with $K \supset G_1$, maximality of $a_\tau(H) = a_\tau(G_1)$ implies that $a_\tau(K) = a_\tau(G_1)$ so $\tau(K/G_1) < \tau$. Therefore G/G_1 is almost simple. The type of G/G_1 is τ and its typical dimension is $a_\tau(G) - a_\tau(G_1)$. Proceeding by induction on $a_\tau(G_i)$ yields the required sequence of Δ -subgroups.

To prove the last sentence in the theorem we proceed as in the proof of the Jordan-Hölder Theorem as given in ([26], Ch. 1, §3). For each $i = 0, \dots, r-1$, $j = 0, \dots, s-1$ let

$$G_{i,j} = G_{i+1}(H_j \cap G_i).$$

The G_{ij} give a refinement of the sequence of G_i 's:

$$G = G_{0,0} \triangleright G_{0,1} \triangleright \dots \triangleright G_{0,s-1} \triangleright G_1 = G_{1,0} \triangleright G_{1,1} \triangleright \dots \triangleright \{0\}$$

Similarly, define for each $j = 0, \dots, s-1$, $i = 0, \dots, r$,

$$H_{j,i} = H_{j+1}(G_i \cap H_j).$$

This yields a refinement of the H_j . By Lemma 3.3 of [26] we have that

$$G_{i,j}/G_{i,j+1} \text{ is isomorphic to } H_{j,i}/H_{j,i+1}$$

(the isomorphism obviously being an isomorphism of Δ - k -groups.) We now claim that for any $i = 0, \dots, r-1$ there is at exactly one j , $0 \leq j \leq s-1$ such that $\tau(G_{i,j}/G_{i,j+1}) = \tau$ and that for this value of j we have that $G_{i,j}/G_{i,j+1}$ is isogenous to G_i/G_{i+1} (and a similar statement will hold for the sequence of $H_{j,i}$). This and the previous sentence then imply that $r = s$ and the final statement of the theorem is true.

For each $i = 0, \dots, r-1$, have that

$$a_\tau(G_i) = a_\tau(G_{i,0}/G_{i,1}) + \dots + a_\tau(G_{i,s-1}/G_{i+1,0}) + a_\tau(G_{i+1})$$

(recall that $G_{i+1,0} = G_{i+1}$). Since $a_\tau(G_i) > a_\tau(G_{i+1})$ we have that some $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$ and so $\tau(G_{i,j}/G_{i,j+1}) = \tau$. Equation (1) furthermore implies that

$\tau(G_{i,j}/G_{i+1}) = \tau$. Let j be the smallest integer such that $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$.

We will show that $G_i = G_{i,0} = G_{i,1} = \dots = G_{i,j}$. Note that $\tau = \tau(G_{i,j}/G_{i,j+1}) \leq \tau(G_{i,j}/G_{i+1}) \leq \tau(G_i/G_{i+1}) = \tau$. Therefore $\tau(G_{i,j}/G_{i+1}) = \tau$. Furthermore, $\tau = \tau(G_i/G_{i+1}) \geq \tau(G_{i,t}/G_{i+1}) \geq \tau(G_{i,j}/G_{i+1}) = \tau$ so $\tau(G_{i,t}/G_{i+1}) = \tau$ for any $t = 0, 1, \dots, j-1$. Since $a_\tau(G_{i,t}/G_{i,t+1}) = 0$ for $t = 0, 1, \dots, j-1$, we have $\tau(G_{i,t}/G_{i,t+1}) < \tau$. Since G_i/G_{i+1} is strongly connected, $G_{i,1}/G_{i+1} \triangleleft G_i/G_{i+1}$ and $\tau((G_{i,0}/G_{i+1})/(G_{i,1}/G_{i+1})) = \tau(G_{i,0}/G_{i,1}) < \tau$, we have $G_{i,0} = G_{i,1}$. Continuing in a similar fashion, we see $G_{i,0} = \dots = G_{i,j}$.

From this discussion we can conclude that $G_{i,j} = G_i$ and $G_{i,j+1} \triangleleft G_i$. Let $\pi : G_i/G_{i+1} \rightarrow G_{i,j}/G_{i,j+1}$ be the canonical projection. The kernel is a proper normal subgroup so, again by almost simplicity, it must have smaller type. Therefore G_i/G_{i+1} is isogenous to $G_{i,j}/G_{i,j+1}$.

We now turn to the uniqueness claim. Assume that $\tau(G_{i,t}/G_{i,t+1}) = \tau$ for some $t > j$. We would then have $\tau(G_{i,t}/G_{i+1}) = \tau$ and so $\tau(G_{i,j+1}/G_{i+1}) = \tau$. Since G_i/G_{i+1} is almost simple and $G_{i,j+1} \triangleleft G_{i,j} = G_i$, we must have $G_{i,j+1} = G_{i,j}$, contradicting $a_\tau(G_{i,j}/G_{i,j+1}) \neq 0$. ■

Remarks 2.28 1. As is clear from the above proof the analogue of the results of Zassenhaus and Schreier (cf., [26], Chapter I, §3, Lemma 3.3 and Theorem 3.4) stating that two normal sequences have isomorphic refinements is true in the context of Δ -groups. This follows from the fact that the isomorphism theorems are true in this context.

2. A parallel theorem, in which the groups in the subnormal series and the quotients are Δ - k -groups, and the quotients are almost k -simple, holds for strongly connected Δ - k -groups.

Corollary 2.29 *Let G and H be isogenous strongly connected Δ -groups with normal sequences $G = G_0 \triangleright G_1 \triangleright \dots \triangleright G_r = \{1\}$ and $H = H_0 \triangleright H_1 \triangleright \dots \triangleright H_s = \{1\}$ as in Theorem 2.27. Then $r = s$ and there exists a permutation π of $\{0, \dots, r\}$ such that $G_{\pi(i)}/G_{\pi(i)+1}$ is k -isogenous to H_i/H_{i+1} .*

Proof. Immediate from Theorem 2.27 and Proposition 2.20. ■

Remarks 2.30 1. One can consider groups G that are not strongly connected and apply Theorem 2.27 first to the strong identity component G_0 , then to the strong identity component of G/G_0 , etc. This will yield a more complete Jordan-Hölder Theorem.

2. In Example 2.32 below, we show that the uniqueness up to isogeny in Theorem 2.27 cannot be strengthened to give uniqueness up to isomorphism.

3. In [3], Baudisch presents a Jordan-Hölder Theorem for superstable groups G . He shows that such a group has a sequence of definable subgroups $(1) \triangleleft H_0 \triangleleft H_1 \triangleleft \dots \triangleleft H_r \triangleleft G$ such that H_{i+1}/H_i is infinite and either abelian or simple modulo a finite center and H_r is of finite index in G . Although this is of a similar nature as our result, his result does not address the structure of abelian groups nor does he address the question of uniqueness. A key step in understanding the relationship between Baudisch's result and ours would

be to understand the relationship between U-rank and the type and typical differential transcendence degree (cf., [43]).

Example 2.31 We now return to the example of Landau mentioned in the Introduction:

$$L = \delta_x^3 + x\delta_x^2\delta_y + 2\delta_x^2 + 2(x+1)\delta_x\delta_y + \delta_x + (x+2)\delta_y \quad (13)$$

$$= (\delta_x + 1)(\delta_x + 1)(\delta_x + x\delta_y) \quad (14)$$

$$= (\delta_x^2 + x\delta_x\delta_y + \delta_x + (x+2)\delta_y)(\delta_x + 1) . \quad (15)$$

We shall show how this example fits into the above discussion.

Let $\Delta = \{\delta_x, \delta_y\}$ and k a differentially closed Δ -field. Recall from Section 2.1 that the gauge of a Δ - k -variety X is the pair $(\tau(X), a_\tau(X))$ and order these lexicographically. Let R be any nonzero homogeneous linear differential polynomial in δ_x and δ_y of order d and let $G_R \subset k$ be the solutions of $R = 0$. In [24], Kolchin gives a general method for computing differential dimension polynomials $\omega_{\eta/k}(s)$ in terms of a characteristic set for the defining ideal I of η over k . When this ideal is generated by linear homogeneous differential polynomials $\{L_1(z), \dots, L_r(z)\}$ in one variable, one does not need this full machinery. In this restricted case, if G_1, \dots, G_s is a reduced Gröbner basis (with respect to a graded monomial order) of the left ideal $\langle L_1, \dots, L_r \rangle$ in $k[\delta_1, \dots, \delta_m]$, then $G_1(z), \dots, G_s(z)$ is a characteristic set of I^1 . The differential dimension polynomial can be computed from the leading terms of this set (with respect to the same monomial order, which is orderly; see Lemma 16, Ch. 0.17 and Theorem 6, Ch. II.12 of [24]). For $\Delta = \{\delta_x, \delta_y\}$ and k a differentially closed Δ -field, this can be made explicit. Identifying the monomial $\delta_x^i\delta_y^j$, we use the order $(i, j) > (i', j')$ if $i + j > i' + j'$ or $i + j = i' + j'$ and $i > i'$. If $E = \{(i_1, j_1), \dots, (i_t, j_t)\}$ represent the leading terms of a reduced Groebner basis of a left ideal of $k[\delta_x, \delta_y]$, we can assume that $i_1 < i_2 < \dots < i_t$ and $j_1 > j_2 > \dots > j_t$. Let

$$W = \{(i, j) \in \mathbb{N}^2 \mid i \geq i_1, j \geq j_t, (i, j) \notin \cup_{(i', j') \in E} ((i', j') + \mathbb{N}^2)\}$$

and let $d = i_1 + j_t$. Geometrically, W is the set of points in the quarter-plane $(i_1, j_t) + \mathbb{N}^2$ below the “stairs” $\cup_{(i', j') \in E} ((i', j') + \mathbb{N}^2)$. Lemma 16, Ch. 0.17 and Theorem 6, Ch. II.12 of [24], imply that

$$\omega(s) = \binom{s+2}{2} - \binom{s-d+2}{2} + |W| = ds + \frac{3d-d^2}{2} + |W|.$$

¹In [22, 30], the authors define and prove the existence of Gröbner bases for a wide class of rings. For left ideals J in $k[\delta_1, \dots, \delta_m]$ their definition reduces to a set $\{G_1, \dots, G_m\} \subset J$ such that the leading term of each element of J be divisible by the leading term of some G_i . This guarantees that $\{G_1(z), \dots, G_m(z)\}$ is coherent. If $\{G_1, \dots, G_m\}$ is reduced in the usual sense of Gröbner bases, then $\{G_1(z), \dots, G_m(z)\}$ is autoreduced. Using the criterion of ([24], Lemma 2, p.167 or [41], Ex. 8.17) and the fact that the condition of [22, 30] guarantees that $\{G_1(z), \dots, G_m(z)\}$ generates $I \cap k\{z\}_1$ where I is the differential ideal generated by the $L_i(z)$ and $k\{z\}_1$ is the set of homogeneous linear differential polynomials, one can see that $\{G_1(z), \dots, G_m(z)\}$ is a characteristic set of I . We thank William Sit for these references and explanations.

For example, if the leading monomial of an operator R is $\delta_x^i \delta_y^j$ with $i + j = d$, then the differential dimension polynomial of the differential ideal $[R(z)]$ generated by $R(z)$ is $\omega(s) = \binom{s+2}{2} - \binom{s-d+2}{2} = ds + \frac{3d-d^2}{2}$. Therefore the gauge of G_R is $(1, d)$. If $H \subset G_R$ is a proper Δ -subgroup, then the defining differential ideal J contains a linear differential polynomial such that the leading term of its associated operator is not divisible by the leading term of R . The stairs corresponding to leading terms of a characteristic set of the ideal J strictly contain $(1, d) + \mathbb{Z}^2$. One sees from this that the associated differential dimension polynomial either has degree 0 or leading coefficient less than d . This implies that either the type of H is smaller than the type of G_R or the typical transcendence degree of H is smaller than d . In particular, this implies that G_R is strongly connected.

The factorization in (14) yields a sequence of subgroups of G_L as follows. Let G_1 be the solutions of $(\delta_x + 1)(\delta_x + x\delta_y)(z) = 0$ and let G_2 be the solutions of $(\delta_x + x\delta_y)(z) = 0$. As noted above, each of these groups is strongly connected. The quotients G/G_1 and G_1/G_2 are both isomorphic to the group defined by $(\delta_x + 1)(z) = 0$. Therefore each quotient in the sequences $G \supset G_1 \supset G_2 \supset \{0\}$ is strongly connected and has gauge $(1, 1)$. Let H be one of these quotients and K a proper subgroup of H . If the gauge of H is $(1, 1)$, then $\tau(K/H) = 0$, contradicting the fact that H is strongly connected. Therefore $a_1(K) = 0$ and so $\tau(K) = 0$. Therefore these quotients are almost simple and so the sequence satisfies the conclusion of .

The factorization in (15) yields a sequence of groups as follows. Let H_2 be the group defined by $(\delta_x + 1)(z) = 0$. We then have the sequence $G \supset H_2 \supset \{0\}$. The Theorem implies that the group G/H_2 cannot be almost simple. In fact the proof of the theorem tells us how to refine this sequence. Let $H_1 = H_2 + G_2$ (note that this sum is direct since the groups have a trivial intersection). Lemma 2.8 implies this group is strongly connected. The ideal defining this group is the intersection of the ideals defining each of the summands, i.e.

$$\langle (\delta_x + 1)(z) \rangle \cap \langle (\delta_x + x\delta_y)(z) \rangle = \langle L_1(z), L_2(z) \rangle$$

where

$$\begin{aligned} L_1 &= x\delta_x^2\delta_y + x^2\delta_x\delta_y^2 - \delta_x^2 - \delta_x\delta_y + x^2\delta_y^2 - \delta_x - \delta_y - x\delta_y \\ L_2 &= \delta_x^3 - x^2\delta_x\delta_y^2 + 3\delta_x^2 + 2x\delta_x\delta_y + 3\delta_x\delta_y - x^2\delta_y^2 + 2\delta_x + 2x\delta_y + 3\delta_y \end{aligned}$$

form a Gröbner basis (with respect to an orderly ranking with $\delta_x > \delta_y$) of this left ideal (cf., Example 5 of [17]. The above polynomials were calculated using the `Ore_algebra` and `Groebner` packages in Maple and differ slightly from those presented in [17]). One can use this representation to see directly that H_1 is strongly connected. Looking at leading terms and using Lemma 16 of Ch. 0.17 of [24], we see that the gauge is $(1, 2)$. Furthermore, we see that if this ideal is contained in a larger ideal J of the same gauge, then J must be generated by a single operator of order 2. Using the techniques of [17] one can show that L_1 and L_2 do not have a common right factor of order 2 so this is impossible. Therefore H_1 is strongly connected of gauge $(1, 2)$. This yields the sequence $G \supset H_1 \supset H_2 \supset \{0\}$

where successive quotients are almost simple. We will now compare successive quotients of the two sequences:

H_2 : We will construct an isogeny between G/G_1 and H_2 . Consider the operator $T = (\delta_x + 1)(\delta_x + x\delta_y)$. This operator maps G onto G/G_1 , which is naturally isomorphic to the solution space of $(\delta_x + 1)(z) = 0$, an almost simple (and therefore strongly connected) group. The group $T(H_2)$ must therefore be either all of G/G_1 or of rank less than 1 (and therefore $\{0\}$ since H_2 is strongly connected). If $T(H_2) = 0$, then H_2 would be annihilated by both $(\delta_x + 1)(\delta_x + x\delta_y)$ and $(\delta_x + 1)$, but this is impossible since the ideal $\langle (\delta_x + 1)(\delta_x + x\delta_y), (\delta_x + 1) \rangle$ contains δ_y and so is of rank 0. Therefore $T(H_2) = G/G_1$ and T defines an isogeny. In fact, since $H_2 = \{fe^{-x} \mid f_x = 0\}$ one can show that T is an isomorphism of H_2 onto G/G_1 .

H_1/H_2 : This group is $(H_2 + G_2)/H_2$ which is isomorphic to $G_2/(H_2 \cap G_2)$. The group $H_2 \cap G_2$ is precisely the common zeroes of $(\delta_x + 1)(z) = 0$ and $(\delta_x + x\delta_y)(z) = 0$, that is, just the element 0. Therefore G_2 is isogenous (in fact, isomorphic) to H_1/H_2 .

G/H_1 : We will show that G/H_1 is isogenous to G_1/G_2 . To see this, consider the map $\overline{G_1} \hookrightarrow G \rightarrow G/H_1$. The kernel of this map is $G_1 \cap H_1$, that is those solutions of $(\delta_x + 1)(\delta_x + x\delta_y)(z) = 0$ that are the sum of a solution of $(\delta_x + 1)(z) = 0$ and a solution of $(\delta_x + x\delta_y)(z) = 0$. This clearly contains G_2 but is not all of G_1 , so we have a homomorphism $G_1/G_2 \rightarrow G/H_1$, that is not the zero map. Since G_1/G_2 is almost simple, the image of this map has gauge $(1, 1)$ and the kernel has rank at most 0. Since G/H_1 is almost simple, the image must be all of G/H_1 . Therefore we have an isogeny of G_1/G_2 onto G/H_1 . This isogeny has a nontrivial kernel; it is the solutions of $(\delta_x + 1)(z) = 0, \delta_y(z) = 0$. Although *this isogeny* has a nontrivial kernel, we claim that G_1/G_2 is nonetheless isomorphic to G/H_1 . To show this it is enough to show the following. Let K_1 be the solutions of $(\delta_x + 1)(z) = 0$ and K_2 be the solutions of $(\delta_x + 1)(z) = 0, \delta_y(z) = 0$. We have that G_1/G_2 is isomorphic to K_1 and G/H_1 is isomorphic (by the above) to K_1/K_2 . The homomorphism $\delta_y : \mathbb{G}_a \rightarrow \mathbb{G}_a$ maps K_1 surjectively to K_1 and its kernel in K_1 is K_2 . Therefore K_1/K_2 is isomorphic to K_1 and our claim is proved.

Example 2.32 We shall use the groups defined in Example 2.26 to construct an example of a Δ -group having two normal sequences with isogenous but nonisomorphic quotients.

Let $\Delta = \{\delta_x, \delta_t\}$ and $k = \mathbb{U}$ and C be the Δ -constants of \mathbb{U} . Let K_1 be the solutions of $(\delta_x^2 - \delta_t)(z) = 0$ and K_2 be the solutions of $(\delta_x - x\delta_t)(z) = 0$. Let $G = K_1 + K_2$. Lemma 2.8 implies that this group is strongly connected. In fact its annihilating ideal is the intersection of $\langle \delta_x^2 - \delta_t \rangle$ and $\langle \delta_x - x\delta_t \rangle$ in $k\{\delta_x, \delta_t\}$ which is $\langle L \rangle$ where

$$\begin{aligned} L &= x\delta_x^3 - x^2\delta_x^2\delta_t - 2\delta_x^2 - x\delta_x\delta_t + x^2\delta_t^2 + 2\delta_t \\ &= (x\delta_x - x^2\delta_t - 2)(\delta_x^2 - \delta_t) \\ &= (x\delta_x^2 - x\delta_t - 2\delta_x)(\delta_x - x\delta_t). \end{aligned} \tag{16}$$

One sees directly that G is strongly connected since a subgroup defined by the vanishing of a single operator must be strongly connected. Consider the two sequences of subgroups:

$G \supset K_1 \supset \{0\}$ and $G \supset K_2 \supset \{0\}$. We claim that each quotient in either of these sequences is almost simple and, while K_1 is isogenous to G/K_2 , it is not isomorphic to either G/K_2 or K_2 .

The almost simplicity of K_1 and K_2 follows from the observation in Example 2.12. In particular any quotient of K_1 or K_2 by a Δ -subgroup of smaller type will also be almost simple. Therefore $G/K_1 \simeq K_2/(K_1 \cap K_2)$ and $G/K_2 \simeq K_1/(K_1 \cap K_2)$ are almost simple. Clearly K_1 is isogenous to G/K_2 and K_2 is isogenous to G/K_1 .

We now claim that K_1 is not isomorphic to either G/K_2 or K_2 . Note that K_2 has typical differential transcendence degree 2 while K_1 has typical differential transcendence degree 1, so these two groups cannot be isogenous. From (16), one sees that G/K_2 is isomorphic to the group of solutions of $(x\delta_x^2 - x\delta_t - 2\delta_x)z = 0$. This is the group H_x of Example 2.26 and K_1 is the group H of the Heat Equation in this latter example. As we have shown in Example 2.26 these groups are not isomorphic.

3 Almost Simple Δ -Groups

In this section we will develop some facts concerning almost simple Δ -groups. We will use the terms Δ -group, Δ -subgroup, etc. to mean that the objects are all defined over \mathbb{U} . Similarly almost simple, irreducible, etc. refer to these properties with respect to \mathbb{U} , a differentially closed Δ -field.

3.1 Quasisimple Linear Algebraic Groups are Almost Simple Δ -groups

We begin with the following lemma.

Lemma 3.1 *Let (\mathbb{U}, Δ) be a differentially closed differential field, $\Delta' \subset \Delta$ and G a non-commutative connected algebraic subgroup of $\mathrm{GL}_n(\mathbb{U})$ with finite center, that is defined over the field C' of Δ' -constants of \mathbb{U} . Then the Δ' -subgroup $G(C')$ of G equals its own normalizer in G .*

Proof. First note that since C' is algebraically closed, $G(C')$ is Zariski dense in G and is non-commutative and connected. Let N be the normalizer of $G(C')$ in G . For $g \in N$ and $c \in G(C')$, we have that $g^{-1}cg = c_1 \in G(C')$. For any $\delta' \in \Delta'$, differentiating this latter equation yields $-g^{-1}\delta'g \cdot g^{-1}cg + g^{-1}c\delta'g = 0$ and therefore

$$c^{-1}\ell\delta'g \ c = \ell\delta'g$$

where $\ell\delta'g = \delta'g \cdot g^{-1}$ is called the logarithmic derivative of g . It is well known that $\ell\delta'g \in \mathfrak{g}$, the lie algebra of G (see [24], Ch. V.22 or [29], Sec.5, for an exposition of properties of the

logarithmic derivative in the linear case). Since $G(C')$ is Zariski dense in G , this latter equation implies that every element of the \mathbb{U} -span W of $\ell\delta'g$ is invariant under the adjoint action of G on \mathfrak{g} . Corollary 3.2 of [20] implies that W is annihilated by the adjoint action of \mathfrak{g} on \mathfrak{g} , that is W lies in the center of \mathfrak{g} . Theorems 3.2 and 4.2 in Chapter 4 of [20] imply that the center of \mathfrak{g} is the lie algebra of the center of G . Since the center of G is finite, we must have that $\ell\delta'g = 0$, that is $\delta'g = 0$ and so $g \in G(C')$. ■

Proposition 3.2 *Let (\mathbb{U}, Δ) be a differentially closed differential field with Δ -constants C and let $G \subset \mathrm{GL}_n(\mathbb{U})$ be a non-commutative quasisimple linear algebraic group defined over C . If N is a proper normal Δ -subgroup of G , then N is finite. In particular, if G is quasisimple (resp., simple) as an algebraic group, then it is quasisimple (resp., simple) as a Δ -group and therefore in either case is almost simple.*

Proof. Let N be a proper normal Δ -subgroup of G and N^0 its identity component (in the Kolchin topology). N^0 is again normal in G . Let $H \subset G$ be the Zariski closure of N^0 . One sees that H is connected in the Zariski topology and is again normal in G . We wish to show that H is finite. If not, then, since G is quasisimple as an algebraic group, we must have that $H = G$. This implies that N^0 is a Zariski dense Δ -subgroup of G , connected in the Kolchin topology. Theorem 19 of [13] states that there exists a finite set of derivations Δ' that are \mathbb{U} -linear combinations of the elements of Δ such that N^0 is conjugate in $\mathrm{GL}_n(\mathbb{U})$ to $G(C')$ where $C' = \{c \in \mathbb{U} \mid \delta'(c) = 0, \forall \delta' \in \Delta'\}$. Expanding Δ if necessary we may assume $\Delta' \subset \Delta$ (this does not change the hypotheses or conclusions). Lemma 3.1 implies that N^0 is its own normalizer so $N^0 = G$, contradicting the fact that N is a proper subgroup of G . ■

3.2 Almost Simple Linear Δ -Groups

We now restrict ourselves to *linear* differential algebraic groups. In this situation, we are able to derive additional properties of almost simple groups and, in the case of ordinary differential fields, give a strong classification of almost simple linear Δ -groups.

Proposition 3.3 *Let G be a nontrivial commutative simple linear Δ -group. Then either G is finite of prime order or is isomorphic to $\mathbb{G}_a(C)$.*

Proof. Assume $G \subset \mathrm{GL}_n(\mathbb{U})$ for some positive integer n . If G is finite then it clearly must be of prime order. Let us now assume that G is infinite. Since the identity component G^0 is a normal Δ -subgroup of finite index, we must have that $G = G^0$ so G is connected. Therefore \overline{G} , the Zariski closure of G in $\mathrm{GL}_n(\mathbb{U})$, is also connected and commutative. Since \mathbb{U} is algebraically closed, we may assume that $\overline{G} = \mathbb{G}_m^p(\mathbb{U}) \times \mathbb{G}_a^q(\mathbb{U})$ for some non-negative integers p, q . Assume that $p > 0$ and let π be the projection of \overline{G} onto some factor of $\mathbb{G}_m^p(\mathbb{U})$. We have that $\pi(G(\mathbb{U}))$ is Zariski dense in $\mathbb{G}_m(\mathbb{U})$ and so by Proposition 31 of [10],

$\pi(G(\mathbb{U}))$ contains $\mathbb{G}_m(C)$. This would imply that G would have a proper nontrivial normal Δ -subgroup. Therefore $p = 0$ and $\overline{G} = \mathbb{G}_a^q(\mathbb{U})$ for some p . Let $0 \neq a = (a_1, \dots, a_q) \in G$. The subgroup Ca is Δ -closed and Δ -isomorphic to $\mathbb{G}_a(C)$. Therefore G must be isomorphic to $\mathbb{G}_a(C)$. ■

If we assume that k is algebraically closed and G is a commutative k -simple Δ - k -group, then, we can only say that G is either finite or isomorphic to a proper Δ - k -subgroup of $\mathbb{G}_a(\mathbb{U})$. For, if G is Zariski dense in $\mathbb{G}_a^q(\mathbb{U})$, let π be the projection of $\mathbb{G}_a^q(\mathbb{U})$ onto $\mathbb{G}_a(\mathbb{U})$. Then, G is delta-isomorphic to $\pi(G)$, but, we do not know that the latter group contains a point rational over k .

In the theory of linear algebraic groups, one knows that the commutator subgroup of a linear algebraic group is closed in the Zariski topology and therefore that a quasisimple linear algebraic group is perfect, that is, it equals its commutator subgroup (in fact, every element of a semisimple algebraic group is a commutator [35]). As we have already mentioned following Definition 2.23, in contrast to the algebraic case, the commutator subgroup of a differential algebraic group need not be closed in the Kolchin topology (Ch. IV.5, [25]). Nonetheless, we have the following proposition

Proposition 3.4 *Let G be an infinite almost simple linear Δ -group. Then either G is commutative or $D_\Delta(G) = G$.*

Proof. Assume G is not commutative and suppose that $D_\Delta(G) \neq G$. Since $D_\Delta(G)$ is a normal Δ -subgroup of G , Corollary 2.14 implies that $D_\Delta(G)$ lies in the center $Z(G)$ of G . Therefore $G/Z(G)$ is commutative. Since every proper normal Δ -subgroup of G is central, the group $G/Z(G)$ is simple. Proposition 3.3 implies that $G/Z(G)$ is isomorphic to $\mathbb{G}_a(C)$. In particular, $0 = \tau(G/Z(G)) = \tau(G)$. This implies that $\tau(Z(G)) = -1$, that is, $Z(G)$ is finite. Since $D_\Delta(G)$ is connected we must have that $D_\Delta(G)$ is trivial, so G is commutative, a contradiction. ■

Suppose G is an infinite almost simple linear Δ -group. An argument based on the proof of Proposition 3.4 breaks down at the last step. We do not know a priori that a simple Δ -subgroup of $\mathbb{G}_a(\mathbb{U})$ has type 0 unless \mathbb{U} is an ordinary differential field. We also do not have an example of a non-commutative almost simple linear differential algebraic group whose commutator subgroup is not closed in the Kolchin topology. In fact, as we show below in Proposition 3.8, non-commutative almost simple linear Δ -groups of *type at most 1* are perfect.

We derive one more property of the center. Let $G \subset \mathrm{GL}_n$ be a Δ -group and \overline{G} its Zariski closure. Let \overline{G}_u be the unipotent radical and \overline{G}_r be the solvable radical of \overline{G} . One can easily show that $\overline{G}_u \cap G$ is the unique maximal normal unipotent differential algebraic subgroup of G and $(\overline{G}_r \cap G)^0$ is the unique maximal connected solvable Δ -subgroup of G . We shall

denote $\overline{G}_u \cap G$ by G_u and $(\overline{G}_r \cap G)^0$ by G_r and refer to these as the unipotent radical and radical respectively. If G is defined over k , so are G_u and G_r .

Proposition 3.5 *Let G be an infinite almost simple linear Δ -group.*

1. *If G is non-commutative, then the unipotent radical G_u of G equals the identity component $(Z(G))^0$ of the center of G .*
2. *If G is commutative then either G is isomorphic to $G_m(C)$ or G is isogenous to an almost simple subgroup of $G_a(\mathbb{U})$.*

Proof. Let $G \subset \mathrm{GL}_n$ and let \overline{G} be the Zariski closure of G .

1. Assume that G is a non-commutative group. The commutator subgroup $(\overline{G}, \overline{G})$ of \overline{G} is a Zariski closed subgroup of \overline{G} and, *a fortiori*, is Kolchin closed. Therefore it contains $D_\Delta(G) = G$. Since $(\overline{G}, \overline{G})$ is Zariski closed, it must also contain \overline{G} . Therefore $\overline{G} = (\overline{G}, \overline{G})$ is perfect.

The group \overline{G} has a Levi decomposition

$$\overline{G} = \overline{G}_u \rtimes P$$

where P is a reductive algebraic group. Since \overline{G} is perfect, P is perfect. The solvable radical P_r of P has finite intersection with (P, P) (Lemma, p. 125 [21]) so P_r is finite. Therefore P must be semisimple. This implies that the radical \overline{G}_r of \overline{G} equals \overline{G}_u . Therefore, $G_r = G_u$. This implies that $Z(G)^0 \subset G_u$. Since $D_\Delta(G) = G$, we have that G_u is a proper normal Δ -subgroup of G and so is central. Therefore $Z(G)^0 = G_u$.

2. Assume G is commutative so \overline{G} is also commutative. Therefore $\overline{G} \simeq \mathbb{G}_m(\mathbb{U})^s \times (\mathbb{G}_a(\mathbb{U}))^t$ for some integers s, t . The map $l\Delta : \mathbb{G}_m(\mathbb{U}) \rightarrow (\mathbb{G}_a(\mathbb{U}))^m$ given by $l\Delta(u) = (\delta_1(u)u^{-1}, \dots, \delta_m(u)u^{-1})$ is a Δ -homomorphism so we have a Δ -homomorphism

$$\phi = (l\Delta^s, id^t) : G \rightarrow (\mathbb{G}_a(\mathbb{U}))^{ms} \times (\mathbb{G}_a(\mathbb{U}))^t.$$

The kernel of this map is $G \cap (\mathbb{G}_m(C))^s \subset (\mathbb{G}_m(\mathbb{U}))^s \subset (\mathbb{G}_m(\mathbb{U}))^s \times (\mathbb{G}_a(\mathbb{U}))^t$. Let $\pi : (\mathbb{G}_a(\mathbb{U}))^{ms} \times (\mathbb{G}_a(\mathbb{U}))^t \rightarrow \mathbb{G}_a(\mathbb{U})$ be a projection onto one of the $\mathbb{G}_a(\mathbb{U})$ factors. Since G is almost simple, the group $\pi \circ \phi(G) \subset \mathbb{G}_a(\mathbb{U})$ is a nontrivial almost simple Δ -group or is the trivial group. If it is a nontrivial almost simple group, then the map $\pi \circ \phi$ is an isogeny since its kernel in G is a proper subgroup. If this group is trivial for all projections π , then G must be a subgroup of $\mathbb{G}_m(C)^s$. This implies that G is a connected algebraic subgroup of $\mathbb{G}_m(C)^s$. Such groups are isomorphic to $(\mathbb{G}_m(C))^\ell$ for some ℓ . Since G is almost simple, we have that $\ell = 1$. ■

Remarks 3.6 1. We cannot strengthen the conclusion of part 2. above. Let (\mathbb{U}, δ) be an ordinary differential field and let $G \subset \mathbb{G}_m(\mathbb{U}) \times \mathbb{G}_a(\mathbb{U})$ be the graph of the homomorphism

$l\delta : \mathbb{G}_m(\mathbb{U}) \rightarrow \mathbb{G}_a(\mathbb{U})$. G contains the group $\mathbb{G}_m(C) \times 0$ and so cannot be unipotent. Therefore G is isogenous to $\mathbb{G}_a(\mathbb{U})$ (via the projection onto the second factor) but is not isomorphic to any subgroup of $\mathbb{G}_a(\mathbb{U})$.

2. For information concerning commutative unipotent differential algebraic groups, see [12]. Nonetheless, we have no general classification of almost simple proper Δ -subgroups of $\mathbb{G}_a(\mathbb{U})$ (except in the ordinary differential case; see below).

The above propositions yield the following theorem. We shall denote by $\mathbb{U} \cdot \Delta$ the \mathbb{U} span of Δ . The elements of $\mathbb{U} \cdot \Delta$ are derivations on \mathbb{U} .

Theorem 3.7 *Let G be an almost simple linear Δ -group of type τ .*

1. *If G is non-commutative, then:*

(a) $G = D_\Delta(G)$,

(b) $G_u = (Z(G))^0$, and

(c) *There exists a finite commuting subset $\Delta' \subset \mathbb{U} \cdot \Delta$, $|\Delta'| = m - \tau$, such that $G/Z(G) \simeq H(C_{\Delta'})$ where H is a simple algebraic group defined over \mathbb{Q} and $C_{\Delta'} \subset \mathbb{U}$ is the field of Δ' -constants.*

2. *If G is commutative, then G is either isogenous to a Δ -subgroup of $\mathbb{G}_a(\mathbb{U})$ or is isomorphic to $\mathbb{G}_m(C)$.*

Proof. Statement 1(a) follows from Proposition 3.4. Statement 1(b) follows from Proposition 3.5.1. Statement 2. follows from Proposition 3.5.2. To verify statement 1(c), note that Corollary 2.15 states that $G/Z(G)$ is a simple Δ -group. Its Zariski closure must be a simple linear algebraic group. The main result of [13] gives the conclusion of 1(c). ■

3.3 Almost Simple Linear Δ -Groups of Differential Type ≤ 1

We note that this includes *ordinary* linear Δ -groups. Restricting ourselves to Δ -groups of differential type at most 1 allows us to sharpen Proposition 3.4.

Proposition 3.8 *Let G be an infinite almost simple linear Δ -group with $\tau(G) \leq 1$. Then either G is commutative or perfect.*

Proof. Assume that G is not commutative. Corollary 2.15 states that $G/Z(G)$ is a simple linear differential algebraic group. Theorem 17 of [13] implies that $G/Z(G)$ is differentially isomorphic to a group $G'(F)$ where G' is a simple non-commutative algebraic group and F is a field of constants with respect to a set of \mathbb{U} -linear combinations of elements of Δ . For an arbitrary group H , let $C_m(H) = \{x_1 y_1 x_1^{-1} y_1^{-1} \cdots x_m y_m x_m^{-1} y_m^{-1} \mid x_i, y_i \in H\}$. Since G' is

simple, we know that $G' = C_1(G')$ (cf., [35]). We will show that $G = C_s(G)$ for some s .

Let X_m be the Kolchin closure of $C_m(G)$ in G . Let $\pi : G \rightarrow G/Z(G)$ be the canonical projection. Since $G' = C_1(G')$, we have that $\pi(C_1(G)) = \pi(G)$. Therefore $\pi(X_1) = G/Z(G)$. Since the group $Z(G)$ has smaller type than G , we have that $\tau(G) = \tau(G/Z(G))$ and $a_\tau(G) = a_\tau(G/Z(G))$. Therefore $\tau(X_1) = \tau(G)$ and furthermore $a_\tau(X_1) = a_\tau(G)$.

For each $i = 1, 2, \dots$ let $\omega_i(s) = a_i \binom{s+1}{1} + b_i$ be the differential dimension polynomial of a generic point of X_i and $\omega(s) = a \binom{s+1}{1} + b$ be the differential dimension polynomial of G (we include the possibility that $a = 0$). Since $X_1 \subset X_2 \subset \dots \subset G$, we have that $\omega_1(s) \leq \omega_2(s) \leq \dots \leq \omega(s)$. As we have just shown, $a_1 = a$ so we must have $a_1 = a_2 = \dots = a$. Therefore, $b_1 \leq b_2 \leq \dots \leq b$. Since these are all integers, we must have that $b_r = b_{r+1} = \dots$ for some r . This implies that $\omega_r(s) = \omega_{r+1}(s) = \dots$ and so $X_r = X_{r+1} = \dots$ (cf., Proposition 2, Ch.III.5 [24]).

We will now show (following a similar proof for algebraic sets) that $G = C_{2r}$. For any $c \in C_r(G)$ the map $d \mapsto dc$ sends $C_r(G)$ to $C_{2r}(G)$ and therefore sends X_r to X_r . A similar argument shows that for any $c \in X_r$, we have that multiplication by c on the left sends X_r to X_r . Therefore X_r is a Δ -subgroup of G . Since $C_r(G)$ is invariant under conjugation by elements of G , we have that X_r is a normal subgroup of G of the same type and so must equal G . Since C_r is constructible, it contains an open subset of its closure, that is an open subset of G . For any $g \in G$, we have that gC_r^{-1} intersects C_r . Therefore $g \in C_r \cdot C_r = C_{2r}$.

■

We can also strengthen Theorem 3.7.

Theorem 3.9 *Let G be an infinite almost simple linear Δ -group with $\tau(G) \leq 1$. Let $|\Delta| = m$.*

1. *If G is non-commutative, then there exists a quasisimple algebraic group H defined over \mathbb{Q} and a commuting linearly independent set of $m-1$ derivations $\Delta' \subset \mathbb{U}\Delta$ such that G is Δ -isomorphic to either $H(C)$ or $H(C')$ where C are the Δ -constants and C' are the Δ' constants of \mathbb{U} .*
2. *If G is commutative, then G is either isogenous to an almost simple Δ -subgroup $G' \subset \mathbb{G}_a(\mathbb{U})$ with $\tau(G') = 1$ or is Δ -isomorphic to $\mathbb{G}_a(C)$ or $\mathbb{G}_m(C)$.*

Proof. 1. Let G be a non-commutative almost simple Δ -group with $\tau(G) \leq 1$. We shall first show that $Z(G)$ is finite.

If $\tau(G) = 0$, then $\tau(Z) = -1$ and so $Z(G)$ is finite. Now assume $\tau(G) = 1$. We shall rely heavily on the results of [1]. Theorem 3.7.1(c) implies that $G/Z(G) \simeq H(F)$, where H is a simple algebraic group defined over \mathbb{Q} and F is an ordinary differentially closed field with

respect to some derivation δ that is a \mathbb{U} -linear combination of elements of Δ . The field of constants of F is the field C of Δ -constants of \mathbb{U} . Therefore there is an exact sequence

$$1 \longrightarrow Z(G) \xrightarrow{\text{incl.}} G \xrightarrow{\alpha} H \longrightarrow 1,$$

where α is a Δ -homomorphism. Let $\tilde{G} = \alpha^{-1}(H(C))$. We then have the exact sequence

$$1 \longrightarrow Z(G) \xrightarrow{\text{incl.}} \tilde{G} \xrightarrow{\alpha} H(C) \longrightarrow 1.$$

Since the type of $Z(G)$ is at most 0 and the type of $H(C)$ is 0, we must have that the type of \tilde{G} is 0, that is, \tilde{G} is a group of finite Morley rank. The group $H(C)$ is a simple algebraic group of constant matrices and so is a simple δ -group as well. Therefore every normal subgroup of \tilde{G} is in $Z(G)$. Proposition 3.8 implies that \tilde{G} is perfect. Therefore the results of [1] imply that $Z(G)$ is finite.

Since $Z(G)$ is finite and $G/Z(G)$ is a simple group, we have that G is a quasisimple Δ -group. Theorem 3.11 in the Appendix states that there exists a quasisimple algebraic group H defined over \mathbb{Q} and a commuting linearly independent set of derivations $\Delta' \subset \mathbb{U}\Delta$ such that G is Δ -isomorphic $H(C')$ where C' are the Δ' constants of \mathbb{U} . Since the type of G is at most 1, we have that either Δ' has m or $m - 1$ elements. If Δ' has m elements, then the Δ' constants and the Δ -constants coincide so G is Δ -isomorphic to $H(C)$, where C is the field of Δ -constants.

2. From Theorem 3.7, we know that a commutative almost simple Δ -group G is either isogenous to a subgroup G' of $\mathbb{G}_a(\mathbb{U})$ or isomorphic to $\mathbb{G}_m(C)$. If $\tau(G) = 0$, let u be a nonzero element of G' . The group $H = C \cdot u$ is a Δ -subgroup of G' with $\tau(H) = 0$. Since G' is also almost simple, we must have $G' = H$ and so G' is isomorphic to $\mathbb{G}_a(C)$. Therefore G is isogenous to $\mathbb{G}_a(C)$.

We will now show that any almost simple group G that is isogenous to $\mathbb{G}_a(C)$ is isomorphic to $\mathbb{G}_a(C)$. Since G is isogenous to $\mathbb{G}_a(C)$, there exists a strongly connected Δ -group G' and surjective Δ -homomorphisms $\alpha_1 : G' \rightarrow G$, $\alpha_2 : G' \rightarrow \mathbb{G}_a(C)$ where α_1 and α_2 have finite kernels. If we can show that α_2 has trivial kernel, then G' would be isomorphic to $\mathbb{G}_a(C)$ and so α_1 would have trivial kernel as well. This would further imply that α_1 is an isomorphism.

Therefore we must show the following: If there is a surjective Δ -homomorphism $\alpha : G \rightarrow \mathbb{G}_a(C)$ with finite kernel H then H is trivial. Since G is also isogenous to $\mathbb{G}_a(C)$, G is commutative (Corollary 2.22).

Let $|H| = n$. Since \mathbb{G}_a is torsion free, we have that the torsion subgroup of G is H . Therefore, H is the kernel of the homomorphism $\gamma : G \rightarrow G$, $\gamma(g) = g^n$. The differential type of $\gamma(G)$ must therefore be the same as the type of G . Since G is strongly connected, we have $\gamma(G) = G$. This implies that if $h \in H$, $h \neq 1$, there exists a $g \in G \setminus H$ such that $g^n = h$. This element g would then be a torsion element not in H , a contradiction. Therefore H is trivial. ■

Corollary 3.10 *Assume that $|\Delta| = 1$, that is, \mathbb{U} is an ordinary differential field. Let G be an almost simple Δ -group.*

1. *If G is non-commutative, then there exists a quasisimple algebraic group H defined over \mathbb{Q} such that G is Δ -isomorphic to $H(\mathbb{U})$ or $H(C)$.*
2. *If G is commutative, then either G is isogenous to $\mathbb{G}_a(\mathbb{U})$ or Δ -isomorphic to $\mathbb{G}_a(C)$ or $\mathbb{G}_m(C)$.*

Proof. Corollary 3.10.1 follows immediately from Theorem 3.9.1. Corollary 3.10.2 follows from Theorem 3.9.2 once one notes that any proper Δ -subgroup of $\mathbb{G}_a(\mathbb{U})$ has type 0. Note that 2 is no longer true for almost k -simple Δ -groups. ■

Example 2.25 shows that even in the case of ordinary differential fields, there are many nonisomorphic almost simple Δ -groups that are isogenous to $\mathbb{G}_a(\mathbb{U})$.

Appendix: Quasisimple Linear Δ -Groups

In this appendix we show how the results of [13]² give the following result that was needed in the proof of Theorem 3.9.

Theorem 3.11 *Let G be a quasisimple linear Δ -group. There exists a quasisimple algebraic group H defined over \mathbb{Q} , a commuting basis $\tilde{\Delta}$ of $\mathbb{U}\Delta$ such that G is Δ -isomorphic to $H(C')$ where C' are the Δ' constants of \mathbb{U} for some $\Delta' \subset \tilde{\Delta}$.*

This result will follow from the following three propositions. The first two appear in [13]. Note that in this latter paper, the author uses the term *simple* to mean quasisimple (cf, [13], p. 222). We use the same notation as in Theorem 3.11

Proposition 3.12 ([13], Corollary 1, p. 228) *Let G be a connected Zariski dense Δ -subgroup of a semisimple algebraic subgroup H of $\mathrm{GL}_n(\mathbb{U})$. Then, G is quasisimple if and only if H is quasisimple.*

Proposition 3.13 ([13], Theorem 19, p. 232) *Let G be a connected Zariski dense Δ -subgroup of a quasisimple algebraic group H where H is defined over the constants of Δ . Then, G is conjugate to $H(C')$ where C' are the Δ' constants of some commuting linearly independent set of derivations of $\mathbb{U}\Delta$.*

²See also [9] for a useful discussion of the results of [13].

Proposition 3.14 *Let G be a quasisimple Δ -subgroup of $\mathrm{GL}_n(\mathbb{U})$. Then, there exists a Δ -rational isomorphism ϕ from G onto a Zariski dense Δ -subgroup of a quasisimple algebraic group H .*

Proof. From Theorem 4.3, Chapter VIII of [21] we have that the Zariski closure \overline{G} of G can be written as a semidirect product

$$\overline{G}_u \rtimes H,$$

where \overline{G}_u is the unipotent radical of \overline{G} and H is a reductive algebraic subgroup of \overline{G} and is H is connected.

We claim that H is semisimple. First note that \overline{G} is perfect, that is, $\overline{G} = (\overline{G}, \overline{G})$. To see this, Theorem 3.7 implies that $G = D_\Delta(G)$. The group $(\overline{G}, \overline{G})$ is Zariski closed and so is Kolchin closed. Since $(\overline{G}, \overline{G}) \subset \overline{G}$ contains (G, G) , it contains $D_\Delta(G) = G$ and therefore must equal \overline{G} . Let $\phi : \overline{G} \rightarrow H$ be the projection with kernel \overline{G}_u . We have $H = \phi(\overline{G}) = \phi((\overline{G}, \overline{G})) = (H, H)$, that is, H is perfect. The solvable radical H_r has finite intersection with (H, H) (Lemma, p.125 [21]), so H_r is finite. Therefore H is semisimple.

We now claim that ϕ restricted to G is an isomorphism of G onto a Zariski dense subgroup of H . The group $\phi(\overline{G})$ is Zariski closed, from which it follows that $H = \phi(\overline{G})$ is the Zariski closure of $\phi(G)$. To see that ϕ is injective on G , note that, since G is quasisimple, $\ker \phi|_G$ is either finite or all of G . If $G = \ker \phi|_G$, we would have $G \subset \overline{G}_u$, and so G would not be quasisimple. Therefore $\ker \phi|_G \subset \overline{G}_u$ is a finite unipotent group and so must be trivial.

Since $\phi(G)$ is a connected quasisimple Zariski dense Δ -subgroup of the semisimple group H , Proposition 3.12 implies that H is quasisimple. ■

Proof of Theorem 3.11. From Proposition 3.14, we may assume that G is a connected Zariski dense Δ -subgroup of a quasisimple algebraic group H defined over \mathbb{U} . A fundamental theorem of Chevalley ([14]³), states that H is isomorphic to an algebraic group defined over \mathbb{Q} . Therefore, we may further assume that H itself is a quasisimple group defined over $\mathbb{Q} \subset C$, the constants of Δ . Proposition 3.13 implies that G is Δ -isomorphic to $H(C')$ where C' are the Δ' constants of some commuting linearly independent set of derivations of $\mathbb{U}\Delta$. Proposition 7, Chapter 0 of [25] states that we may extend Δ' to a basis $\tilde{\Delta}$ of $\mathbb{U}\Delta$. ■

References

- [1] Tuna Altinel and Gregory Cherlin. On central extensions of algebraic groups. *J. Symbolic Logic*, 64(1):68–74, 1999.

³See [7] (especially 3.3(6) and 3.3(1)) for a survey of these notes and [9] for a discussion of the literature reproving and extending this result.

- [2] Moulay A. Barkatou, Thomas Cluzeau, and Jacques-Arthur Weil. Factoring partial differential systems in positive characteristic. In *Differential equations with symbolic computation*, Trends Math., pages 213–238. Birkhäuser, Basel, 2005. With an appendix by Marius van der Put.
- [3] Andreas Baudisch. On superstable groups. *J. London Math. Soc. (2)*, 42(3):452–464, 1990.
- [4] Richard Beals and Elena A. Kartashova. Constructive factorization of linear partial differential operators. *Teoret. Mat. Fiz.*, 145(2):165–180, 2005.
- [5] George W. Bluman and Sukeyuki Kumei. *Symmetries and differential equations*, volume 81 of *Applied Mathematical Sciences*. Springer-Verlag, New York, 1989.
- [6] Henry Blumberg. *Über algebraische Eigenschaften von linearin homogene Differentialausdrücken*. PhD thesis, Georg-August-Universität, 1912.
- [7] Armand Borel. Properties and linear representations of Chevalley groups. In *Seminar on Algebraic Groups and Related Finite Groups (The Institute for Advanced Study, Princeton, N.J., 1968/69)*, Lecture Notes in Mathematics, Vol. 131, pages 1–55. Springer, Berlin, 1970.
- [8] Elie Cartan. Les groupes de transformations continues, infinis, simple. *Annales scientifiques de l'École Normale Supérieure, Sér. 3*, 26:93–161, 1909.
- [9] Phyllis Joan Cassidy. Quasisimple differential algebraic groups. Preprint; available at www.math.ncsu.edu/~singer/Quasisimple.pdf, 2009.
- [10] Phyllis Joan Cassidy. Differential algebraic groups. *Amer. J. Math.*, 94:891–954, 1972.
- [11] Phyllis Joan Cassidy. The differential rational representation algebra on a linear differential algebraic group. *J. Algebra*, 37(2):223–238, 1975.
- [12] Phyllis Joan Cassidy. Unipotent differential algebraic groups. In *Contributions to algebra (collection of papers dedicated to Ellis Kolchin)*, pages 83–115. Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1977.
- [13] Phyllis Joan Cassidy. The classification of the semisimple differential algebraic groups and the linear semisimple differential algebraic Lie algebras. *J. Algebra*, 121(1):169–238, 1989.
- [14] Claude Chevalley. *Classification des groupes algébriques semi-simples*. Springer-Verlag, Berlin, 2005. Collected works. Vol. 3, Edited and with a preface by P. Cartier, With the collaboration of Cartier, A. Grothendieck and M. Lazard.
- [15] Thomas Cluzeau and Alban Quadrat. Factoring and decomposing a class of linear functional systems. *Linear Algebra Appl.*, 428(1):324–381, 2008.

- [16] Dima Grigoriev. Weak Bézout inequality for D-modules. *J. Complexity*, 21(4):532–542, 2005.
- [17] Dima Grigoriev and Fritz Schwarz. Factoring and solving linear partial differential equations. *Computing*, 73(2):179–197, 2004.
- [18] Dima Grigoriev and Fritz Schwarz. Loewy and primary decompositions of D -modules. *Adv. in Appl. Math.*, 38(4):526–541, 2007.
- [19] Dima Grigoriev and Fritz Schwarz. Non-holonomic overideals of differential ideals in two variables and absolute factoring. [arXiv:math.AP/0811.1368v1](https://arxiv.org/abs/math/0811.1368v1), 2008.
- [20] Gerhard Hochschild. *Basic Theory of Algebraic Groups and Lie Algebras*, volume 75 of *Graduate Texts in Mathematics*. Springer Verlag, Berlin, 1976.
- [21] James E. Humphreys. *Linear Algebraic Groups*. Graduate Texts in Mathematics. Springer-Verlag, New York, 1975.
- [22] Mariano Insa and Franz Pauer. Gröbner bases in rings of differential operators. In *Gröbner bases and applications (Linz, 1998)*, volume 251 of *London Math. Soc. Lecture Note Ser.*, pages 367–380. Cambridge Univ. Press, Cambridge, 1998.
- [23] Ellis R. Kolchin. Constrained extensions of differential fields. *Advances in Math.*, 12:141–170, 1974.
- [24] Ellis R. Kolchin. *Differential Algebra and Algebraic Groups*. Academic Press, New York, 1976.
- [25] Ellis R. Kolchin. *Differential algebraic groups*, volume 114 of *Pure and Applied Mathematics*. Academic Press Inc., Orlando, FL, 1985.
- [26] Serge Lang. *Algebra*. Addison Wesley, New York, 3rd edition, 1993.
- [27] David Marker. *Model theory*, volume 217 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2002. An introduction.
- [28] David Marker, Margit Messmer, and Anand Pillay. *Model theory of fields*, volume 5 of *Lecture Notes in Logic*. Springer-Verlag, Berlin, 1996.
- [29] Claude Mitschi and Michael F. Singer. Solvable-by-finite groups as differential Galois groups. *Ann. Fac. Sci. Toulouse*, XI(3):403 – 423, 2002.
- [30] Franz Pauer. Gröbner bases with coefficients in rings. *J. Symbolic Comput.*, 42(11-12):1003–1011, 2007.
- [31] Anand Pillay. Some foundational questions concerning differential algebraic groups. *Pacific J. Math.*, 179(1):179–200, 1997.

- [32] Jean-Francois Pommaret. *Partial differential control theory*, volume 530 of *Mathematics and its Applications*. Kluwer Academic Publishers Group, Dordrecht, 2001. Vol. I. Mathematical tools; Vol. II. Control systems, With a foreword by Jan C. Willems.
- [33] Roman O. Popovych, Michael Kunzinger, and Nataliya M. Ivanova. Conservation laws and potential symmetries of linear parabolic equations. *Acta Appl. Math.*, 100(2):113–185, 2008.
- [34] Marius van der Put and Michael F. Singer. *Galois Theory of Linear Differential Equations*, volume 328 of *Grundlehren der mathematischen Wissenschaften*. Springer, Heidelberg, 2003.
- [35] Rimhak Ree. Commutators in semi-simple algebraic groups. *Proc. Amer. Math. Soc.*, 15:457–460, 1964.
- [36] Joseph F. Ritt. *Differential Algebra*, volume 33 of *American Mathematical Society Colloquium Publication*. American Mathematical Society, New York, 1950.
- [37] Maxwell Rosenlicht. Some basic theorems on algebraic groups. *Amer. J. Math.*, 78:401–443, 1956.
- [38] Ekaterina S. Shemyakova. Parametric factorizations of second-, third- and fourth-order linear partial differential operators with a completely factorable symbol on the plane. *Math. Comput. Sci.*, 1(2):225–237, 2007.
- [39] Ekaterina S. Shemyakova and Franz Winkler. Obstacles to the factorization of partial differential operators into several factors. *Programmirovaniye*, 2:17–25, 2007. translation in *Program. Comput. Software* 33 (2007) no. 2, 67–73.
- [40] Michael F. Singer. Testing reducibility of linear differential operators: a group theoretic perspective. *Applicable Algebra in Engineering, Communication and Computing*, 7:77–104, 1996.
- [41] William Y. Sit. The Ritt-Kolchin theory for differential polynomials. In *Differential algebra and related topics (Newark, NJ, 2000)*, pages 1–70. World Sci. Publ., River Edge, NJ, 2002.
- [42] William Yu Sit. Typical differential dimension of the intersection of linear differential algebraic groups. *J. Algebra*, 32(3):476–487, 1974.
- [43] Sonat Suer. *Model Theory of Differentially Closed Fields with Several Commuting Derivations*. PhD thesis, University of Illinois at Urbana-Champaign, 2007.
- [44] Sergey P. Tsarev. Factorization of linear partial differential operators and the Darboux method for integrating nonlinear partial differential equations. *Teoret. Mat. Fiz.*, 122(1):144–160, 2000.

- [45] Sergey P. Tsarev. Factorization of linear differential operators and systems. In *Algebraic theory of differential equations*, volume 357 of *London Math. Soc. Lecture Note Ser.*, pages 111–131. Cambridge Univ. Press, Cambridge, 2009.
- [46] Min Wu. On the factorization of differential modules. In *Differential equations with symbolic computation*, Trends Math., pages 239–254. Birkhäuser, Basel, 2005.