



RESEARCH PAPER

# An ERF2-like transcription factor regulates production of the defense sesquiterpene capsidiol upon *Alternaria alternata* infection

Na Song<sup>1,2,\*</sup>, Lan Ma<sup>1,\*</sup>, Weiguang Wang<sup>3,\*</sup>, Huanhuan Sun<sup>1</sup>, Lei Wang<sup>1</sup>, Ian T. Baldwin<sup>4, ID</sup> and Jinsong Wu<sup>1,†, ID</sup>

<sup>1</sup> Yunnan Key Laboratory for Wild Plant Resources, Kunming Institute of Botany, Chinese Academy of Sciences, Lanhei Road 132, 650201, Kunming, China

<sup>2</sup> University of Chinese Academy of Science, Beijing 10049, China

<sup>3</sup> Key Laboratory of Chemistry in Ethnic Medicinal Resources of State Ethnic Affairs Commission and Ministry of Education, Yunnan Minzu University, 650031, Kunming, China

<sup>4</sup> Department of Molecular Ecology, Max Planck Institute for Chemical Ecology, 07745 Jena, Germany

\* These authors contributed equally to this work.

† Correspondence: [jinsongwu@mail.kib.ac.cn](mailto:jinsongwu@mail.kib.ac.cn)

Received 11 March 2019; Editorial decision 4 July 2019; Accepted 5 July 2019

Editor: Steven Spoel, University of Edinburgh, UK

## Abstract

Capsidiol is a sesquiterpenoid phytoalexin produced in *Nicotiana* and *Capsicum* species in response to pathogen attack. Whether capsidiol plays a defensive role and how its biosynthesis is regulated in the wild tobacco *Nicotiana attenuata* when the plant is attacked by *Alternaria alternata* (tobacco pathotype), a notorious necrotrophic fungus causing brown spot disease, are unknown. Transcriptome analysis indicated that a metabolic switch to sesquiterpene biosynthesis occurred in young leaves of *N. attenuata* after *A. alternata* inoculation: many genes leading to sesquiterpene production were strongly up-regulated, including the capsidiol biosynthetic genes *5-epi-aristolochene synthase (EAS)* and *5-epi-aristolochene hydroxylase (EAH)*. Consistently, the level of capsidiol was increased dramatically in young leaves after fungal inoculation, from not detectable in mock control to  $50.68 \pm 3.10 \mu\text{g g}^{-1}$  fresh leaf at 3 d post-inoculation. Capsidiol-reduced or capsidiol-depleted plants, which were generated by silencing *EAHs* or *EASs* by virus-induced gene silencing, were more susceptible to the fungus. In addition, this sesquiterpene when purified from infected plants exhibited strong anti-fungal activities against *A. alternata in vitro*. Furthermore, an ERF2-like transcription factor was found to positively regulate capsidiol production and plant resistance through the direct transactivation of a capsidiol biosynthetic gene, *EAS12*. Taken together, our results demonstrate that capsidiol, a phytoalexin highly accumulated in *N. attenuata* plants in response to *A. alternata* infection, plays an important role in pathogen resistance independent of jasmonate and ethylene signaling pathways, and its biosynthesis is transcriptionally regulated by an ERF2-like transcription factor.

**Keywords:** 5-epi-Aristolochene hydroxylase (EAH), 5-epi-aristolochene synthase (EAS), ethylene responsive factor, *Nicotiana attenuata*, phytoalexin, sesquiterpene, virus induced gene silencing (VIGS).

## Introduction

Plants are constantly attacked by a wide variety of microbial pathogens. In response, they activate a large number of intricate defense mechanisms, including the formation of reactive oxygen species, physical reinforcement of cell walls, and production of phytohormones, antimicrobial proteins, and metabolites (Glazebrook, 2005; Ahuja *et al.*, 2012; Mengiste, 2012). The class of small molecules known as phytoalexins is produced by plants *de novo* in response to pathogen attack, and is an important part of the plant defense repertoire (Ahuja *et al.*, 2012). Arabidopsis mutants impaired in the production of the phytoalexin camalexin are more susceptible to infection by necrotrophic fungi, such as *Alternaria brassicicola* (Nafisi *et al.*, 2007), *Botrytis cinerea* (Kliebenstein *et al.*, 2005), and *Plectosphaerella cucumerina* (Sanchez-Vallet *et al.*, 2010). In *Nicotiana* species, two phytoalexins have recently received attention: scopoletin (El Oirdi *et al.*, 2010; Sun *et al.*, 2014b) and capsidiol (Mialoundama *et al.*, 2009; Shibata *et al.*, 2010, 2016; Grosskinsky *et al.*, 2011).

Capsidiol has been proposed to be an important ‘chemical weapon’ against pathogens in *Nicotiana* species. This bicyclic sesquiterpene is produced via cyclization of farnesyl pyrophosphate (FPP) to 5-*epi*-aristolochene by 5-*epi*-aristolochene synthase (EAS), followed by two hydroxylation reactions catalysed by 5-*epi*-aristolochene dihydroxylase (EAH) (Facchini and Chappell, 1992; Ralston *et al.*, 2001). Capsidiol exhibits toxicity towards many pathogens, including *Phytophthora capsici* and *B. cinerea* (Stoessl *et al.*, 1972; Ward *et al.*, 1974). Recently, molecular evidence also supports its role in non-host resistance against *P. infestans* in *N. benthamiana*, as *NbEAS*- or *NbEAH*-silenced plants were highly susceptible (Shibata *et al.*, 2010).

Ethylene response factors (ERFs) are transcription factors that play crucial roles in plant immunity (Huang *et al.*, 2016). In Arabidopsis, several ERFs have been identified as important regulators in *Botrytis* resistance, such as ORA59, ERF1, and RAP2.2 (Solano *et al.*, 1998; Berrocal-Lobo *et al.*, 2002; Pré *et al.*, 2008; Zhao *et al.*, 2012). Most ERFs are able to bind specifically to DNA sequences containing a GCC (GCC box) and/or a dehydration-responsive element/C-repeat (DRE/CRT box, A/GCCGAC) (Hao *et al.*, 1998, 2002). However, the ERF RAP2.2 binds the consensus sequence ATCTA in the promoter region of *phytoene synthase* and *phytoene desaturase* to regulate carotenoid biosynthesis (Welsch *et al.*, 2007).

The necrotrophic fungal pathogen *Alternaria alternata* (tobacco pathotype) causes brown spot disease in *Nicotiana tabacum* (LaMondia, 2001) and many other *Nicotiana* species, including the wild tobacco *Nicotiana attenuata* (Schuck *et al.*, 2014). Inoculation with *A. alternata* elicits the activation of both jasmonate (JA) and ethylene signaling pathways in *N. attenuata* plants, which subsequently lead to the accumulation of the phytoalexins scopoletin and its glycoside form, scopolin (Sun *et al.*, 2014b; Li and Wu, 2016; Sun *et al.*, 2017). Currently it is not known if fungus-inoculated *N. attenuata* plants produce other phytoalexins, such as capsidiol. If they do, how their biosynthesis is transcriptionally regulated is a significant question.

In this study, capsidiol was identified and confirmed to be an important phytoalexin produced in *N. attenuata* when

challenged by *A. alternata*, and its regulation by an ERF2-like transcription factor was investigated in detail.

## Materials and methods

### Plant and fungal materials

Seeds of the 31st generation of an inbred line of *Nicotiana attenuata* were used as the wild-type (WT) genotype. Ethylene-deficient and -insensitive (*irACO* and *Ov-etr1*, respectively), and JA-deficient (*irAOC*) *N. attenuata* plants were generated previously (von Dahl *et al.*, 2007; Kallenbach *et al.*, 2012). Seed germination and plant growth were conducted as described by Krügel *et al.* (2002).

*Alternaria alternata* was grown and inoculated into leaves as described by Sun *et al.* (2014a).

### RNA-seq data processing and analysis

Source-sink transition leaves of rosette-staged WT plants (35 d old) were detached and inoculated with *A. alternata* for 1 d, when only a few of fungal hyphae had penetrated into leaf tissues (Sun *et al.*, 2014a). Total RNA of three biological replicates of mock (WT\_0L\_M, with sample names S716, S717, and S719) or inoculated leaf samples (WT\_0L\_Inf, with sample names S20, S722, and S726) were isolated with TRIzol reagent (Invitrogen). RNA sequencing was conducted by Shanghai OE-Biotech (<http://www.oebiotech.com/>) with the Illumina HiSeq 2000.

Sequencing was performed at 8 G depth, and mapped to the *N. attenuata* reference genome sequence. The relative abundance of the transcripts was measured with the FPKM (RPKM) method, which measures the transcript abundance as RPKM (reads per kilobase of exon model per million mapped reads). The differential expression between mock and 1 d post-inoculation (dpi) samples with a cutoff of 2-fold change and its significance were calculated.

The raw sequence data reported in this paper have been deposited in the Genome Sequence Archive in BIG Data Center, Beijing Institute of Genomics (BIG), Chinese Academy of Sciences, under accession numbers CRA001483 (publicly accessible at <http://bigd.big.ac.cn/gsa>).

### Generation of *NaEAS*- and *NaEAH*-silenced virus-induced gene silencing plants

A 579 bp *NaEAS* cDNA fragment amplified with primers (Z003\_F and Z004\_R; Supplementary Table S1) and a 424 bp fragment of *NaEAH* cloned with primers (Z047\_F and Z048\_R; Supplementary Table S1) were individually inserted into pTV00 (Ratcliff *et al.*, 2001) in reverse orientations. *Agrobacterium tumefaciens* (strain GV3101) harboring these constructs was mixed with the strain with pBINTRA and inoculated into *N. attenuata* leaves generating *NaEAS*- and *NaEAH*-silenced plants (virus-induced gene silencing (VIGS) *NaEAS*s and VIGS *NaEAH*s). The *A. tumefaciens*-mediated transformation procedure was performed as described previously (Saedler and Baldwin, 2004). To monitor the progress of VIGS, *phytoene desaturase* (*PDS*) was also silenced. Silencing *PDS* results in the visible bleaching of green tissues (Saedler and Baldwin, 2004; Wu *et al.*, 2008) about 2–3 weeks after the inoculation. When the leaves of *PDS*-silenced plants began to bleach, the young leaves of VIGS plants and empty vector-inoculated plants (EV plants) were selected for further experiments. Around 20 plants were inoculated with each construct, and usually 10 biological replicates per construct exhibiting efficient silencing were used for each experiment, and all VIGS experiments were repeated twice.

### Purification and quantification of capsidiol in *N. attenuata* after infection

Around 500 g leaves that had been inoculated with *A. alternata* for 3 d were collected for capsidiol extraction. Leaves were twice extracted with 70% acetone (2 liters) at room temperature. The solvent was evaporated

and suspended in water, and then extracted with ethyl acetate. The ethyl acetate-soluble fraction (10 g) was decolorized on an MCI gel ([www.gls.co.jp](http://www.gls.co.jp)) with methanol: H<sub>2</sub>O (90:10) to obtain a yellow gum (7 g), which was subsequently purified with a silica gel column with a chloroform: acetone gradient system (from 10:0, 9:1, 8:2, 7:3, 6:4, to 1:1) to yield six main fractions (A–F). Fraction B (chloroform: acetone, 9:1; 2 g) was subjected to repeated chromatography over silica gel (petroleum ether: acetone, from 30:1 to 1:1) to yield fractions B1–B4. Fraction B3 (petroleum ether: acetone, 10:1) was separated further with an RP-18 column (acetonitrile: H<sub>2</sub>O, 30:70). The obtained crude capsidiol was further purified by semi-preparative HPLC (3 ml min<sup>-1</sup>, UV detection at  $\lambda_{\text{max}}=202$  nm, acetonitrile: H<sub>2</sub>O, 40:60; ZORBAX SB-C18 column (5  $\mu\text{m}$ , 9.4×250 mm, Agilent 1200, USA) to yield capsidiol (30 mg, >99.5% purity). The purified capsidiol showed the same characteristic NMR data and HPLC retention times as the authentic standard provided by Prof. J. Chappell (University of Kentucky, USA).

*Alternaria alternata*-elicited capsidiol levels were determined by HPLC by reference to the authentic capsidiol standard. Each leaf was inoculated with four agar plugs with fungal mycelium, and a 1.5×1.5 cm<sup>2</sup> area of the leaf lamina was harvested at 1 or 3 dpi (around 200 mg fresh mass). Samples were ground in liquid nitrogen and twice extracted with 1 ml dichloromethane; 2 ml extracts were dried in a SpeedVac concentrator (Eppendorf), and finally dissolved in 1 ml methanol for HPLC. At a flow rate of 1 ml min<sup>-1</sup>, 10  $\mu\text{l}$  of each sample was injected onto a ZORBAX SB-C18 column (5  $\mu\text{m}$ , 4.6×250 mm) (Agilent 1260). The mobile phase was composed of solvent A (water), and solvent B (acetonitrile) was used in a isocratic elution (40% of B). Capsidiol was detected at 202 nm, with a retention time of 10.38 min. The standard capsidiol was dissolved in methanol at six concentrations (6.25, 12.5, 25, 50, 100, and 200  $\mu\text{g ml}^{-1}$ ) to create an external standard curve that was used to calculate fungus-induced capsidiol levels.

#### Bioassays for the inhibition of *A. alternata* growth by capsidiol *in vitro*

The inhibition of *A. alternata* mycelium growth by capsidiol *in vitro* was tested in Petri dishes by sub-culturing a 3 mm-diameter mycelium plug on potato dextrose agar (PDA) medium containing various concentrations of capsidiol for 6 d in the dark at 25 °C. Twenty milligrams of capsidiol was dissolved in 10 ml methanol, and added to the PDA medium at final concentrations of 0, 50, 100, and 200  $\mu\text{g ml}^{-1}$ . PDA plates with 1% methanol served as controls. Photos were taken every 2 d, and the area of mycelium growth was calculated with ImageJ (<http://imagej.nih.gov/ij/>).

#### Quantification of scopoletin and scopolin

Around 0.2 g leaf samples at 6 dpi of EV, VIGS NaEAS and VIGS NaEAH plants were harvested and ground to fine powder in liquid nitrogen. The levels of scopoletin and scopolin were determined by HPLC-MS/MS as described in Sun *et al.* (2014b).

#### Real-time PCR

Total RNA was extracted from a 1.5×1.5 cm<sup>2</sup> area of leaf lamina that encompassed the inoculation site with TRI reagent (Invitrogen). For each treatment, four to five replicate biological samples were collected. cDNA was synthesized from 500 ng total RNA with reverse transcriptases (Thermo Fisher Scientific). Real-time PCR was performed on a CFX Connect qPCR System (Bio-Rad) with iTaq Universal SYBR Green Supermix (Bio-Rad) and gene-specific primers as described (Wu *et al.*, 2013). For each analysis, a linear standard curve (obtained from threshold cycle number versus log DNA quantity) was constructed by using a dilution series of a specific cDNA sample, and the transcript levels of unknown samples were calculated according to the standard curve. Finally, the relative transcript levels of target genes were obtained by dividing the extrapolated transcript levels of the target genes by the levels of a house-keeping gene, *NaActin 2*, as an internal standard from the same sample. The transcript abundance of *NaActin 2* was not altered in leaves inoculated with *A. alternata* at 1 and 3 dpi (Xu *et al.*, 2018). All primers are listed in Supplementary Table S1 at JXB online.

#### Transient expression assays

To determine the subcellular localization of NaERF2-like, a 35S::ERF2-eGFP construct was used for transient expression in *N. attenuata* protoplasts. The full-length coding sequence of *NaERF2-like* was amplified by primers Z141\_F and Z142\_R (Supplementary Table S1) and inserted into vector pM999 via *SacI* and *XhoI*. The method of protoplast isolation and transient transformation was adopted from (Yoo *et al.*, 2007) with some modifications. In brief, mesophyll protoplasts were isolated from source-sink transition leaves, and 20  $\mu\text{g}$  of plasmids was transfected into 2×10<sup>5</sup> protoplasts with polyethylene glycol (PEG) solution (0.4 g ml<sup>-1</sup> PEG 4000, 0.2 M mannitol, 0.1 M CaCl<sub>2</sub>). Transformed cells were cultured in solution (4 mM MES, 0.5 M mannitol, 20 mM KCl) for 18 h in the dark and images were obtained with a fluorescence microscope (Leica DM5500 B) with excitation at 488 nm for the green fluorescent protein (GFP) signal.

For a transient transactivation assay, the promoter region of *NaEAS12* (–1671 to –1; numbered from the first ATG) and firefly luciferase (*LUC*) were amplified and cloned into pCAMBIA1301. Next, the PCR fragment including the promoter region of *NaEAS12*, *LUC*, and Nos terminator was subcloned into pMD18 via *HindIII* and *SacI* to reduce the size of the vector and increase the transformation efficiency. Similarly, 35S::NaERF2-like-2HA-eGFP with Nos terminator was also subcloned into pMD18 vector. *Nicotiana attenuata* protoplasts were prepared as describe above, and transformed with both NaEAS12<sub>Promoter</sub>::LUC and 35S::NaERF2-like-2HA-eGFP, or with NaEAS12<sub>Promoter</sub>::LUC and 35S::2HA-eGFP as control. After transformation, the protoplasts were subjected to RNA extraction with a PrimeScript™ RT reagent kit with gDNA Eraser followed by a gene expression assay. All the experiments were repeated twice with similar results.

#### Yeast-one-hybrid assay

The Matchmaker yeast one-hybrid system (Clontech) was used to test the binding of NaERF2-like and the NaEAS12 promoter *in vitro* according to the user manual. The promoter region of *NaEAS12* (pEAS12-b; –926 to –699; numbered from the first ATG), which contained the candidate binding motif of EM13 (5′-tagattATCTaattctact-3′), was inserted into pAbAi vector with *HindIII* and *XhoI*. The bait construct was linearized and integrated into the genome of yeast strain Y1HGGold. The full-length coding sequence of *NaERF2-like* was introduced into pGADT7 AD vector via *Clal* and *XhoI*, and then the construct was transformed into the yeast cells containing the bait. The positive clones were analysed on SD/–His/–Leu medium supplied with 200 ng ml<sup>-1</sup> (final concentration) 3-amino-1,2,4-triazole (3-AT). Y1HGGold [pGADT7/pEAS12-b-AbAi] was used as a negative control and Y1HGGold [pGADT7 Rec-p53/p53-AbAi] was used as positive control.

#### Electromobility shift assays

The full-length coding sequence of *NaERF2-like* was cloned in-frame into the *EcoRI*–*XhoI* sites of the pET28a (+), and His–NaERF2-like was expressed and purified with Ni-NTA agarose (Qiagen). A biotin-labeled probe, EM13 (5′-tagattATCTaattctact-3′), and a mutant probe (5′-tagattAATTaattctact-3′) were synthesized by Sangon Biotech (Shanghai). The detection of the binding of the recombinant protein and the probes (300 ng of recombinant protein and 30 ng labeled probe) was carried out with a chemiluminescence electrophoretic mobility shift assay (EMSA) kit (Beyotime Biotechnology) according to the protocol suggested by the manufacturer.

#### Generation of *Ov-NaERF2-like* plants and chromatin immune-precipitation assay

The full-length cDNA of NaERF2-like with two hemagglutinin (HA: YPYDVPDYA) epitopes at its C-terminus was cloned into pCAMBIA1301 vector after the 35S promoter with the in-fusion technique (Clontech). *Nicotiana attenuata* plants (31st generation of an inbred line) were transformed by *Agrobacterium tumefaciens* with this construct according to Krügel *et al.* (2002). T1 seeds were screened for single T-DNA inserts (1:3 segregation of hygromycin resistance), and two lines



of T2 (Ov-NaERF2-like line 1 and 2) with HA signals (see Results, Supplementary Fig. S2) were selected and used in this study.

Chromatin immunoprecipitation was performed with EpiQuik Plant ChIP Kit (EPIGENTEK) according to the user manual. The source-sink transition leaves of Ov-NaERF2-like line 2 at 1 dpi (1 g) were used for ChIP assays. A ChIP grade anti-HA antibody (Abcam) was used to immunoprecipitate the protein-DNA complex, and the precipitated DNA was further purified for real time PCR by using primer sets from *NaEAS12* promoter (5'-CACTTTAACCCCGGTAAGT-3' and 5'-CACTTCTCAGATTCTCCAGTTTGG-3') and *NaActin 2* gene (5'-GGTCGTACCACCGGTATTGTG-3' and 5'-GTCAAGACGGAGAATGGCATG-3') for negative control. The ChIP experiments were performed twice with similar results. Chromatins that were precipitated without antibody served as the negative controls.

## Results

### *Transcriptome analysis reveals the strong regulation of sesquiterpene biosynthetic genes in N. attenuata after A. alternata inoculation*

Previously, we have demonstrated that, in response to *A. alternata* inoculation, *N. attenuata* plants activate both JA and ethylene signaling pathways to regulate the biosynthesis of the phytoalexins scopoletin and its  $\beta$ -glycoside form, scopolin (Sun et al., 2014b, 2017; Li and Wu, 2016). As *feruloyl-CoA 6'-hydroxylase 1* (*NaF6'H1*), the key enzyme gene for scopoletin biosynthesis, was highly elicited at 1 d post-inoculation (dpi) when the infection was at an early stage and only a few fungal hyphae had been observed to penetrate into leaf tissues via stomata (Sun et al., 2014a), transcriptome analysis was performed in *A. alternata*-inoculated *N. attenuata* leaves at 1 dpi to identify secondary metabolites that could potentially act similarly to scopoletin. Notably, a set of genes involved in terpene synthesis were strongly up-regulated, including *thiolase*, *HMG-CoA synthase*, *HMG-CoA reductase*, *mevalonate kinase*, *phosphomevalonate kinase*, *diphosphomevalonate decarboxylase* (*MVPP decarboxylase*), *isopentenyl-diphosphate delta-isomerase*, *FPP synthase*, the capsidiol biosynthetic genes *5-epi-aristolochene synthases* (*EASs*) and *5-epi-aristolochene 1,3-dihydroxylases* (*EAHs*), and the solavetivone biosynthetic genes *premnaspirodiene oxygenase* and *premnaspirodiene synthase* (Fig. 1A; Supplementary Table S2). The *squalene synthase* involved in triterpene biosynthesis was down-regulated (Fig. 1A; Supplementary Table S2). These results strongly indicate that the sesquiterpene biosynthetic pathway is activated during the inoculation of *A. alternata*.

To confirm the regulation of terpene biosynthesis genes in response to *A. alternata*, we performed quantitative real time PCR on *N. attenuata* samples collected at 1 dpi. Compared with mock infection controls, the transcripts of *MVPP decarboxylase*, *HMG-CoA reductase*, and *FPP synthase* at 1 dpi were increased 322-, 53-, and 14-fold, respectively (Fig. 1B–D), while the transcripts of *squalene synthase* were reduced by 75% (Fig. 1E). More importantly, transcripts of the capsidiol biosynthetic genes *NaEASs* and *NaEAHs*, which are encoded by a multi-gene family (Supplementary Table S2) and thus were detected by primers designed at the consensus region, increased 368- and 40-fold compared with control (Fig. 1F, G). These data fully confirmed the transcriptomics result of strong activation of the sesquiterpene biosynthesis pathway in *N. attenuata* after inoculation.

### *Accumulation of capsidiol in N. attenuata in response to A. alternata inoculation*


Since *NaEAS* and *NaEAH* are key genes in the capsidiol biosynthesis pathway, we expected to see an increase in capsidiol production after *A. alternata* inoculation. The levels of capsidiol at 1 and 3 dpi in the young source-sink transition leaves (0 leaves) and mature leaves (+3 leaves) were determined by HPLC. The +3 leaves are three phyllotaxic positions older than 0 leaves, and are more susceptible to *A. alternata* (Sun et al., 2014a). Our results indicated that capsidiol was not detectable in the 0 leaves of mock control, but its level was increased to  $4.45 \pm 0.72 \mu\text{g g}^{-1}$  fresh leaf in 0 leaves at 1 dpi, and to  $50.68 \pm 3.10 \mu\text{g g}^{-1}$  at 3 dpi (Fig. 2). Interestingly, this compound was also detected in the susceptible +3 leaves, but its level was only about one-third of that of 0 leaves, with  $1.66 \pm 1.00 \mu\text{g g}^{-1}$  fresh leaf at 1 dpi, and  $14.54 \pm 4.39 \mu\text{g g}^{-1}$  at 3 dpi (Fig. 2). These results indicate that (i) capsidiol is highly elicited in *N. attenuata* leaves after inoculation, and (ii) the lower accumulation of capsidiol in mature leaves may account for their susceptibility to the fungus.

### *Capsidiol accumulation is essential for A. alternata resistance in N. attenuata*

To further evaluate the role of capsidiol in *N. attenuata* resistance against *A. alternata*, we silenced *NaEASs* and *NaEAHs* separately with their conserved sequences, as both are members of a multi-gene family. Compared with mock controls, transcripts of *NaEAHs* were dramatically induced in young leaves of *N. attenuata* plants transformed with empty vector (EV) at 2 dpi; however, plants transformed with the *NaEAHs*-silencing construct (VIGS *NaEAHs*) showed a 92% reduction in *NaEAHs* transcripts compared with EV plants with the same treatments, indicating effective silencing of the *NaEAHs* (Fig. 3A). We also investigated the capsidiol level in EV and VIGS *NaEAHs* plants at 3 dpi. A capsidiol level of  $57.72 \pm 5.88 \mu\text{g g}^{-1}$  fresh leaf was detected in EV plants, while only  $8.99 \pm 1.56 \mu\text{g g}^{-1}$  fresh leaf was found in VIGS *NaEAHs* plants (Fig. 3B).

Because a small amount of capsidiol was still present in VIGS *NaEAHs* plants, we attempted to generate additional capsidiol-depleted *N. attenuata* plants by silencing *NaEASs*. Expression of *NaEASs* was successfully silenced, as only 7% of the transcripts of *NaEASs* were detected at 2 dpi in VIGS *NaEASs* plants compared with EV plants (Fig. 3A). More importantly, the *A. alternata*-elicited capsidiol level at 3 dpi was abolished in VIGS *NaEASs* plants, with only 0.2% of the level detected in EV plants, which was comparable to the levels quantified in the mock controls of EV plants (Fig. 3B). From these data, we infer that *NaEASs* genes are crucial for *A. alternata*-elicited capsidiol production.

To test whether capsidiol-reduced or -depleted plants are more susceptible to *A. alternata*, young leaves of EV, VIGS *NaEAHs*, and VIGS *NaEASs* plants were inoculated with the fungus. Two independent VIGS experiments showed significantly increased lesion diameters in VIGS *NaEAHs* (121% of EV plants) and VIGS *NaEASs* plants (131% of EV plants) (Fig. 3C). We did not observe changes in scopoletin and scopolin, two important phytoalexins involved in *A. alternata* resistance (Fig. 3B). These results strongly indicate that capsidiol plays an important role in defending against *A. alternata*.




**Fig. 1.** The regulation of terpene biosynthetic genes in *N. attenuata* in response to *A. alternata* inoculation at 1 d post-inoculation (dpi). Genes involved in terpene synthesis were strongly regulated during transcriptome analysis in three biological replicate samples of mock and 1 dpi. (A) Transcripts of all enzymes marked in red font were up-regulated at 1 dpi, while *squalene synthase* (SS), shown with green font, was down-regulated. (B–G) To validate this regulation, relative mean transcripts ( $\pm$ SE) of *HMG-CoA reductase* (B), *MVAPP decarboxylase* (C), *FPPS* (D), *squalene synthase* (E), *EASs* (F), and *EAHs* (G) were measured by real-time PCR in four biological replicate 0 leaves at 1 dpi. Leaves without inoculation were collected as controls (Mock 1 d). Both *EASs* and *EAHs* were detected by primers conserved in gene family members. Asterisks indicate level of significant difference between Mock 1 d and 1 dpi samples (Student's *t*-test: \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.005$ ). DMAPP, 3'-dimethylallyl diphosphate; EA, 5-*epi*-aristolochene; EAH, 5-*epi*-aristolochene hydroxylase; EAS, 5-*epi*-aristolochene synthase; FPP, farnesyl diphosphate; FPPS, farnesyl pyrophosphate synthase; GGPP, geranylgeranyl diphosphate; GPP, geranyl diphosphate; HMG-CoA, 3-hydroxy-3-methyl-glutaryl-CoA; IPP, isopentenyl diphosphate; MVA, mevalonic acid; MVAP, mevalonic acid 5-phosphate; MVAPP, mevalonic acid 5-diphosphate; PO, premnaspirodiene oxidase; PS, premnaspirodiene synthase.

#### *Capsidiol* exhibits anti-fungal activity against *A. alternata* in vitro

To test whether capsidiol has a direct impact on fungal growth or not, we purified 30 mg capsidiol from 500 g of

*A. alternata*-inoculated leaves, and applied this compound in various concentrations to the growth medium to evaluate its inhibitory activity on fungal growth *in vitro*. The fungi were grown on PDA plates containing 50 or 100  $\mu\text{g ml}^{-1}$  capsidiol, and we observed that fungal growth was reduced to 56.6% or



**Fig. 2.** Accumulation of capsidiol in 0 (young) and +3 (mature) leaves after *A. alternata* inoculation. Mean ( $\pm$ SE) capsidiol levels were determined by HPLC in five biological replicate 0 and +3 leaves at 1 and 3 dpi. Asterisks indicate level of significant difference between mock and infected samples (Student's *t*-test: \* $P$ <0.05, \*\* $P$ <0.01, \*\*\* $P$ <0.005). ND, not detectable.

43.8%, respectively, of that of controls. Application of  $200 \mu\text{g ml}^{-1}$  capsidiol resulted in further reduction in growth to 37.1% of control (Fig. 4A, B). These results suggest that capsidiol at a concentration observed *in planta* has a direct impact on the fungal growth *in vitro*.

#### *Alternaria alternata*-induced capsidiol accumulation is not dependent on JA and ethylene signaling

JA and ethylene signaling pathways are crucial for biosynthesis of the phytoalexin scopoletin. To investigate the roles of these two signaling pathways in capsidiol biosynthesis, we measured the levels of capsidiol and transcripts of *NaEASs* and *NaEAHs* after *A. alternata* inoculation in WT, JA-deficient (*irAOC*), ethylene-deficient (*irACO*), and ethylene-insensitive (*Ov-etr1*) plants generated previously (von Dahl et al., 2007; Kallenbach et al., 2012). We found that the induction levels of *NaEASs* and *NaEAHs* by *A. alternata* were similar in WT, *irAOC*, *irACO*, and *Ov-etr1* plants at both 1 and 3 dpi (Fig. 5A, B). In addition, no significant differences of capsidiol production were observed in WT, *irAOC*, *irACO*, and *Ov-etr1* plants at 3 dpi (Fig. 5C). Thus, our data indicated that *A. alternata*-induced capsidiol accumulation is not dependent on JA and ethylene signaling.

#### *NaERF2-like*, a transcription factor highly induced in young leaves, is required to mount a capsidiol-based defense

Since we observed increases in both capsidiol production and *NaEASs* and *NaEAHs* transcripts in *N. attenuata* leaves after fungal inoculation, we hypothesized that the transcription factors regulating expression of *NaEASs* and *NaEAHs* were also increased. Thus, we silenced the expression of the six fungus-elicited transcription factor genes with the most abundant transcripts after fungal inoculation (Supplementary Table S3), namely *ethylene-responsive transcription factor ABR1-like* (*NaERF ABR1-like*; gene accession no.: XM\_019374371), *zinc finger protein ZAT12-like* (*NaZAT12*; gene accession no.: XM\_019368773.1), *probable WRKY transcription factor 40* (*NaWRKY40*; gene accession no.: XM\_019402562.1), *probable*

*WRKY transcription factor 43* (*NaWRKY43*; gene accession no.: XM\_019375046.1), *probable WRKY transcription factor 61* (*NaWRKY61*; gene accession no.: XM\_019371308.1), and *ethylene-responsive transcription factor 2-like* (*NaERF2-like*; gene accession no.: XM\_019399671.1), to identify the regulator(s) responsible for capsidiol biosynthesis. *Alternaria alternata*-elicited *NaEAS12* was not altered in plants silenced with *NaERF ABR1-like*, *NaZAT12*, *NaWRKY40*, *NaWRKY43*, or *NaWRKY61* (Supplementary Fig. S1).

*NaERF2-like* (gene accession no.: XM\_019399671.1), one of the top six transcription factors strongly up-regulated in response to fungal inoculation in our transcriptome analysis, was highly induced in *N. attenuata* 0 leaves at both 1 and 3 dpi (Fig. 6A; Supplementary Table S3). The *NaERF2-like* protein exhibited nuclear localization in the protoplast of *N. attenuata* when its eGFP fusion protein was driven by a constitutive 35S promoter (Fig. 6B).


To investigate the role of the *NaERF2-like* transcription factor in capsidiol biosynthesis in detail, we silenced the gene by VIGS, and then measured the levels of capsidiol and transcripts of *NaEASs* and *NaEAHs* after fungal inoculation. *NaERF2-like* transcripts were highly elicited at 2 dpi in EV plants; in contrast, VIGS *NaERF2-like* plants showed an 87% reduction in *NaERF2-like* transcripts after the same treatment (Fig. 7A). Compared with EV plants at 2 dpi, the transcripts of *NaEASs* and *NaEAHs* in VIGS *NaERF2-like* plants were reduced by 75% and 62%, respectively (Fig. 7B, C). As expected, *A. alternata*-induced capsidiol level in VIGS *NaERF2-like* plants at 3 dpi was reduced by 68% when compared with EV plants (Fig. 7E). However, the fungus-elicited transcriptional levels of *NaF6'H1* were not affected in VIGS *NaERF2-like* plants (Fig. 7D). These results indicate that silencing *NaERF2-like* substantially decreases *A. alternata*-elicited transcription of *NaEASs* and *NaEAHs*, and consequently, capsidiol level, without affecting scopoletin-based defense.

In addition, silencing *NaERF2-like* led to plants being more susceptible to *A. alternata*, as significantly larger lesions were observed in VIGS *NaERF2-like* plants at 6 dpi in two independent VIGS experiments (Fig. 7F).

#### *NaERF2-like* directly regulates the capsidiol biosynthetic gene *NaEAS12*

Next, we explored the mechanism by which *NaERF2-like* regulates capsidiol-based resistance. We hypothesized that *NaERF2-like* might directly regulate genes in the capsidiol biosynthetic pathway. Since *NaEAS12* expression was greatly reduced in *NaERF2-like*-silenced plants (Fig. 8A), we selected this gene to test whether its promoter could be directly activated by *NaERF2-like*.

Several lines of evidence were consistent with the idea that *NaERF2-like* directly regulates the capsidiol biosynthetic gene *NaEAS12*. When the *NaERF2-like* gene was overexpressed in the protoplasts of *N. attenuata*, the *luciferase* gene (*LUC*) driven by the *NaEAS12* promoter showed a 5.5-fold increase in expression (Fig. 8B), indicating that the overexpression of *NaERF2-like* enhanced the transcriptional activity of *NaEAS12* promoter. From the promoter region of *NaEAS12*, an ATCTA motif, previously shown to be the binding site of RAP2.2 in




**Fig. 3.** Silencing expression of *NaEAHs* or *NaEASs* dramatically reduces *A. alternata*-induced capsidiol levels and plant resistance without affecting scopoletin. (A) Mean ( $\pm$ SE) relative *A. alternata*-induced transcripts of *NaEAHs* and *NaEASs* as measured by real-time PCR in five replicate young leaves of EV, VIGS *NaEAHs*, and VIGS *NaEASs* plants at 2 dpi. (B) Mean ( $\pm$ SE) capsidiol and scopoletin (including scopolin) levels were determined by HPLC in five replicate young leaves of EV, VIGS *NaEAHs*, and VIGS *NaEASs* plants at 3 dpi. (C) Mean ( $\pm$ SE) diameter of necrotic lesions in eight replicate young leaves of EV, VIGS *NaEAHs*, and VIGS *NaEASs* plants infected with *A. alternata* for 6 d. As both *NaEAH* and *NaEAS* are encoded by large gene families, conserved cDNA regions were used for silencing and real time PCR. Asterisks indicate level of significant difference between EV and VIGS leaves (Student's *t*-test: \* $P$ <0.05, \*\*\* $P$ <0.005). ND, not detectable.

Arabidopsis (Welsch *et al.*, 2007), was identified and confirmed as the binding site for the ERF2-like by yeast one-hybrid, electrophoretic mobility shift assay (EMSA) and chromatin immunoprecipitation-based qPCR (ChIP-qPCR). Yeast one-hybrid experiments revealed that the ERF2-like protein could bind to the *NaEAS12* promoter fragment EAS12-b (located from -699 to -926 bp upstream of the starting codon), which contained the ATCTA motif (Fig. 9). Further EMSA experiments indicated that NaERF2-like could directly bind to the

biotin labeled probe EM13 (5'-tagattATCTAattctact-3'), but not to the mutated one (5'-tagattAATTAattctact-3') (Fig. 9). To further confirm these results *in vivo*, we generated a transgenic line ectopically expressing the NaERF2-like protein fused with two HA tags at the C-terminus (35S::NaERF2-like-2HA) for use in ChIP-qPCR. The fusion protein could be detected by a commercial HA antibody (Supplemental Fig. S2), suggesting that the protein was successfully overexpressed in *N. attenuata* plants. Transgenic plants were inoculated with






**Fig. 4.** Capsidiol exhibits anti-fungal activity *in vitro*. (A) Growth of *A. alternata* mycelium at day 6 in PDA with capsidiol at final concentration of 0, 50, 100, and 200  $\mu\text{g ml}^{-1}$ . PDA plates with 1% methanol served as controls. (B) The area of *A. alternata* mycelium growth in PDA with different concentrations of capsidiol was determined by ImageJ. Data were collected every 2 d. Asterisks indicate level of significant difference between mock and infected samples (Student's *t*-test: \*\*\* $P < 0.005$ ).

*A. alternata* and sampled at 1 dpi. We found that the HA-tagged NaERF2-like protein bound the *NaEAS12* promoter at a site that encompassed the ACTCA motif (Fig. 9).

#### Overexpression of NaERF2-like does not alter plant resistance, but increases *A. alternata*-induced NaEAS12 gene expression and capsidiol levels

To further understand the role of *NaERF2-like* in pathogen defense, we investigated *NaEAS12* gene expression, capsidiol level, and plant resistance in WT and two *NaERF2-like* overexpression (Ov-*NaERF2-like*) lines. The ectopic overexpression of *NaERF2-like* fused with two HA tags substantially increased the basal and induced transcriptional levels of *NaERF2-like* (Fig. 10A) without affecting the plants' morphology and size. As expected, Ov-*NaERF2-like* line 1 and Ov-*NaERF2-like* line 2 showed 178% and 219% of the *NaEAS12* expression of WT when 0 leaves were inoculated at 1 d (Fig. 10A). Consistently, capsidiol levels attained values 149% and 175% of that of WT in Ov-*NaERF2-like* lines 1 and 2 at 3 dpi, respectively (Fig. 10B). However, we found no difference in lesion diameters in both overexpression lines compared with WT (Supplementary Fig. S2), indicating that overexpression of *NaERF2-like* increased the gene expression of *NaEAS12*, and subsequently capsidiol biosynthesis, but had only a minor effect on the plants' resistance.



**Fig. 5.** JA and ethylene pathways play a minor role in *A. alternata*-induced transcription of *NaEASs* and *NaEAHs*, and capsidiol biosynthesis. (A, B) Mean ( $\pm$ SE) relative *A. alternata*-induced *NaEASs* (A) and *NaEAHs* (B) transcripts as measured by real-time PCR in five biological replicates of young leaves (0 leaves) in WT, irAOC (JA deficient), irACO (ethylene deficient), and Ov-etr1 (ethylene insensitive) plants at 1 and 3 dpi. (C) Mean ( $\pm$ SE) capsidiol levels determined by HPLC in five replicate 0 leaves of WT, irAOC, irACO, and Ov-etr1 plants at 3 dpi. ND, not detectable.


## Discussion

Phytoalexins are important 'chemical weapons' employed by plants in defending against pathogens. In addition to scopoletin and scopolin, two phytoalexins regulated by JA and ethylene signaling pathways in response to *A. alternata* infection (Sun et al., 2014b; Li and Wu, 2016), we demonstrate in this study that capsidiol is another important phytoalexin produced by *N. attenuata*, and its biosynthesis is not dependent on JA and ethylene signaling pathways, but is transcriptionally regulated by a transcription factor, NaERF2-like.

### Capsidiol is an important phytoalexin produced in *N. attenuata* in response to *A. alternata* infection

Capsidiol was initially isolated from pepper fruit after treatments with various pathogens, including *Phytophthora capsici*, *Botrytis cinerea*, and *Fusarium oxysporum* (Stoessl et al., 1972). Later, this compound was also identified from infected *Nicotiana*





**Fig. 6.** NaERF2-like is highly elicited after *A. alternata* inoculation and is targeted to the nucleus. (A) Mean ( $\pm$ SE) relative *A. alternata*-induced *NaERF2-like* transcripts as measured by real-time PCR in four replicate 0 leaves at 1 and 3 dpi. Asterisks indicate level of significant difference between mock and infected samples (Student's *t*-test: \*\*\* $P$ <0.005). (B) Two fusion proteins, 35S::eGFP and 35S::NaERF2-like-eGFP, were expressed transiently in *N. attenuata* protoplasts. NaERF2-like was found to be targeted to the nucleus. Images were taken in bright field (left), UV (middle) and GFP channels (right).

species (Bailey *et al.*, 1975; Guedes *et al.*, 1982; Mialoundama *et al.*, 2009; Shibata *et al.*, 2010; Grosskinsky *et al.*, 2011; Shibata *et al.*, 2016). In our *N. attenuata*–*A. alternata* pathosystem, we found that a large number of genes were strongly regulated in response to *A. alternata* inoculation at 1 dpi during transcriptome analysis. Many of these genes were involved in the biosynthesis of sesquiterpenes, capsidiol, and solavetivone (Fig. 1; Supplementary Table S2). Indeed, the levels of capsidiol were dramatically induced to  $50.68 \pm 3.10 \mu\text{g g}^{-1}$  fresh leaf at 3 dpi in young 0 leaves (Fig. 2). These findings suggest that capsidiol is involved in the resistance of *N. attenuata* to *A. alternata* infection.


Ideally, the benefits of a putative resistant trait should be determined in plants differing only in a single gene that controls the defense trait and are otherwise identical (Bergelson *et al.*, 1996). In this study, virus-induced gene silencing of *NaEAHs* or *NaEASs* was used to manipulate the production of capsidiol, and the results revealed that capsidiol-reduced or -depleted plants were more susceptible to *A. alternata* (Fig. 3). In response to *A. alternata* inoculation, capsidiol level was increased to  $50.68 \pm 3.10 \mu\text{g g}^{-1}$  in young 0 leaves at 3 dpi (Fig. 2). Considering that this capsidiol level was averaged from leaf laminae of  $1.5 \times 1.5 \text{ cm}^2$  after inoculation, we speculated that the concentration of capsidiol *in planta* at the interface of fungal

infection would be higher than  $50 \mu\text{g g}^{-1}$  fresh leaf. Therefore, we tested the inhibition rate of fungal growth *in vitro* by supplement of capsidiol at concentrations of 50, 100, and  $200 \mu\text{g ml}^{-1}$ . The fungal growth was reduced by 43.4% in PDA plates after 6 d with  $50 \mu\text{g ml}^{-1}$  of capsidiol (Fig. 4), indicating that capsidiol had strong anti-fungal activities against *A. alternata* when it was applied *in vitro* at a concentration similar to those *in planta*.

Thus, our results demonstrate that capsidiol is an important phytoalexin involved in the defense mechanism of *N. attenuata* against *A. alternata*, and the high level of capsidiol accumulated in the young 0 leaves accounts for their high resistance to *A. alternata* infection. Whether solavetivone plays a role in resistance is unknown. Transcriptome data, especially the strong up-regulation of *premnaspirodiene oxygenases* and *premnaspirodiene synthases* at 1 dpi (Fig. 1 and Supplementary Table S2), are consistent with a defensive role of solavetivone. However, this question needs further investigation.

#### Regulation of capsidiol biosynthesis

Several reports indicate that the phytoalexin production is influenced by endogenous plant hormones. Increasing cytokinin (CK) levels in *N. tabacum* plants by exogenous CK




**Fig. 7.** Silencing *NaERF2-like* impairs *A. alternata*-induced transcripts of *NaEASs* and *NaEAHs*, capsidiol levels and plant resistance without affecting *NaF6'H1* transcript accumulation. (A–D) Mean ( $\pm$ SE) relative *A. alternata*-induced *NaERF2-like* (A), *NaEASs* (B), *NaEAHs* (C), and *NaF6'H1* (D) transcript abundance as measured by real-time PCR in five replicate young leaves of EV and VIGS *NaERF2-like* plants at 2 dpi. (E) Capsidiol levels determined by HPLC in five replicate young leaves of EV and VIGS *NaERF2-like* plants at 3 dpi. (F) Mean ( $\pm$ SE) diameter of necrotic lesions recorded in eight replicate young leaves of EV and VIGS *NaERF2-like* plants infected with *A. alternata* for 6 d. Two independent VIGS experiments returned similar results. Asterisks indicate level of significant difference between EV and VIGS leaves (Student's *t*-test: \* $P < 0.05$ , \*\*\* $P < 0.005$ ). ND, not detectable.

application or overexpression of the bacterial *isopentenyl transferase* gene enhanced their resistance to the hemibiotrophic bacterium *Pseudomonas syringae* by increasing levels of capsidiol and scopoletin (Grosskinsky et al., 2011). This CK-mediated resistance is independent of the salicylic acid, JA, and ethylene signaling pathways (Grosskinsky et al., 2011). Abscisic acid (ABA) negatively regulates elicitor-induced biosynthesis of capsidiol in *N. plumbaginifolia*; a 2-fold increase in capsidiol synthesis was observed in ABA-deficient mutants compared with WT plants when exposed to cellulose or *B. cinerea* (Mialoundama et al., 2009). In addition, when *N. benthamiana*

was inoculated with *Phytophthora infestans*, pathogen-induced capsidiol and *NbEAS* and *NbEAH* expression were abolished in plants with *ethylene insensitive 2* (*NbEIN2*) silenced, suggesting that the ethylene signaling pathway is essential for capsidiol production (Shibata et al., 2010; Ohtsu et al., 2014).

In contrast to the observation in *N. benthamiana* inoculated with *P. infestans*, our experiments did not support the role of ethylene signaling in capsidiol elicitation in the *N. attenuata*–*A. alternata* pathosystem. Compared with WT plants, both capsidiol production and transcripts of *NaEAHs* and *NaEASs* after *A. alternata* challenge were induced to the




**Fig. 8.** *Alternaria alternata*-elicited *NaEAS12* transcripts are largely dependent on *NaERF2-like*, and transient overexpression of *NaERF2-like* leads to transactivation of *NaEAS12* promoter. (A) Mean ( $\pm$ SE) relative *A. alternata*-induced *NaEAS12* transcripts as measured by real-time PCR in five replicate young leaves of EV and VIGS *NaERF2-like* plants at 2 dpi. Asterisks indicate level of significant difference between EV and VIGS leaves at 2 dpi (Student's *t*-test: \*\*\* $P$ <0.005). (B) Transcript abundance of *NaERF2-like-2HA-eGFP* and *LUC* in *N. attenuata* protoplasts co-expressing *NaEAS12* promoter::LUC and 35S::2HA-eGFP or 35S::*NaERF2-like-2HA-eGFP* measured in three biological samples. Experiments were repeated twice with similar results. Asterisks indicate level of significant difference between control and *NaERF2-like* overexpressing protoplasts (Student's *t*-test: \* $P$ <0.05; \*\*\* $P$ <0.005).

same high levels in *irACO* (ethylene-deficient) and *Ov-etr1* (ethylene-insensitive) plants (Fig. 5), two transgenic plants generated previously by von Dahl *et al* (2007). To further support this idea, we also silenced *NaEIN2* by VIGS. *Alternaria alternata*-elicited *NaF6'H1* was dramatically reduced in VIGS *NaEIN2* plants, which was consistent with our previous finding that ethylene signaling was essential for scopoletin biosynthesis (Sun *et al.*, 2017), but *NaEAHs* and *NaEASs* transcripts were induced to levels similar to those of plants transformed with empty vector (Supplementary Fig. S3). This result also indicated that blocking the ethylene signaling pathway had little effect on the expression of these two key capsidiol biosynthesis

genes. Thus, we concluded that the ethylene signaling pathway did not play a critical role in capsidiol production and gene expression of *NaEAHs* and *NaEASs* in *N. attenuata* after *A. alternata* challenge. Whether or not ethylene is involved in capsidiol biosynthesis is likely dependent on the pathosystem. Additionally, our experiments did not indicate a role for JA signaling in the regulation of capsidiol biosynthesis, as transcripts of *NaEAHs* and *NaEASs* were equivalent in WT and JA-deficient *irAOC* plants at 1 and 3 dpi (Fig. 5).

Transcription factors integrate developmental and environmental signals by interacting with the promoter regions of enzyme genes, thus enhancing or suppressing their expression to

### NaEAS12 promoter



**Fig. 9.** The binding of NaERF2-like and *NaEAS12* promoter as demonstrated by yeast one-hybrid, EMSA and chromatin immunoprecipitation. The *NaEAS12* promoter structure indicated the pEAS12-b (–669 to –926 numbered from the ATG) for yeast one-hybrid, the probe EM13 (5′-tagattATCTaattctact-3′) for EMSA, and the pEAS12-13 (–669 to –781 numbered from the ATG) for ChIP assays. The red letters ‘AT’ indicate the mutated positions. Yeast one-hybrid analysis revealed that NaERF2-like could bind to EAS12-b as the yeast cells could grow on the SD/–His/–Leu medium supplied with 200 ng ml<sup>–1</sup> (final concentration) 3-AT; Y1HGold [pGADT7/pEAS12-b-AbAi] was used as a negative control, and Y1HGold [pGADT7 Rec-p53/p53-AbAi] was used as a positive control. EMSA demonstrated that His tagged NaERF2-like could bind to probe EM13 but not to the mutated one. The mutant probe (5′-tagattAATaattctact-3′) served as a negative control in EMSA. ChIP-real time PCR data indicated NaERF2-like bound to the promoter of *NaEAS12*. Negative controls were without antibody (no Ab) and with HA antibody but using primers detecting *NaActin 2*. Asterisks indicate level of significant difference between no Ab and with Ab in pEAS12-13 (Student’s *t*-test: \*\**P*<0.01).


regulate the biosynthesis of secondary metabolites (Yang *et al.*, 2012). For example, the sesquiterpene phytoalexin gossypol in cotton is regulated by GaWRKY1 (Xu *et al.*, 2004), and EREB58 mediates the JA-induced production of sesquiterpene volatiles in maize (Li *et al.*, 2015); artemisinin, a sesquiterpene lactone found exclusively in *Artemisia annua*, is controlled by bZIP1 and glandular trichome-specific WRKY1 (Zhang *et al.*, 2015; Chen *et al.*, 2017).

Despite the great role of capsidiol in resistance, the transcriptional regulation of its biosynthesis is still not clear. Due to the high level of gene expression seen in the capsidiol biosynthetic pathway, we hypothesized that the transcription factors regulating these genes must also be highly expressed during the initial infection period of the fungus. We performed a screen of the six most up-regulated transcription factors. Indeed, NaERF2-like was identified as being a positive regulator of plant resistance and capsidiol production. When compared with EV plants, those with a silenced *NaERF2-like* gene accumulated fewer transcripts of *NaEASs* and *NaEAHs*, as well as lower levels of capsidiol (Fig. 7). Consistently, *NaERF2-like*-silenced plants were susceptible to the fungus (Fig. 7).

Both *NaEAS* and *NaEAH* are members of a multi-gene family. Since *NaEAS12* expression is greatly reduced in *NaERF2-like*-silenced plants (Fig. 8), we selected this gene to test whether its promoter could be directly activated by NaERF2-like. Several lines of evidence support that NaERF2-like directly regulates the capsidiol biosynthetic gene *NaEAS12*, including (i) the binding of NaERF2-like protein to the promoter region of *NaEAS12*, which was supported by yeast one-hybrid, EMSA and ChIP-qPCR experiments (Fig. 9), and (ii) the *NaEAS12* promoter was activated in response to transient *NaERF2-like* overexpression (Fig. 8); this result is further confirmed by stable transgenic lines of *Ov-NaERF2-like*, which exhibit increased *NaEAS12* transcripts and higher levels of capsidiol accumulation (Fig. 10). Currently, it is not known how other *NaEASs* and *NaEAHs* are regulated by NaERF2-like. Further experiments are needed to test whether or not they are regulated in a way similar to *NaEAS12*.

The ectopic overexpression of *NaERF2-like* increased the gene expression of *NaEAS12* and capsidiol production after fungal inoculation, but did not result a higher resistance in *Ov-NaERF2-like* line 1 and *Ov-NaERF2-like* line 2 (Fig. 10).





**Fig. 10.** Overexpression of *NaERF2-like* enhances *NaEAS12* expression and capsidiol production in stable transformation plants. (A) Mean ( $\pm$ SE) relative *A. alternata*-induced *NaERF2-like* and *NaEAS12* transcripts as measured by real-time PCR in five replicate 0 leaves of WT, Ov-*NaERF2-like* line 1, and Ov-*NaERF2-like* line 2 plants at 1 dpi. (B) Mean ( $\pm$ SE) capsidiol levels were determined by HPLC in five replicate 0 leaves of WT, Ov-*NaERF2-like* line 1, and Ov-*NaERF2-like* line 2 plants at 3 dpi. Asterisks indicate level of significant difference between WT and Ov-*NaERF2-like* lines after inoculation (Student's *t*-test: \* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.005$ ). ND, not detectable.

This was likely due to the reduced inhibition efficiency of fungal growth when capsidiol levels exceeded  $50 \mu\text{g g}^{-1}$  fresh leaf. The fungal growth was reduced by 43.4% in PDA plates after 6 d with  $50 \mu\text{g ml}^{-1}$  of capsidiol (Fig. 4). When 2-fold capsidiol ( $100 \mu\text{g ml}^{-1}$ ) was added into the PDA plates, the inhibition effect was only increased from 43.4 to 56.2% (Fig. 4). These results indicated that the inhibition rate of fungal growth by capsidiol was not linear with concentration, but rather decreased when capsidiol concentration exceeded  $50 \mu\text{g ml}^{-1}$ , and this could be the reason why *NaERF2-like* overexpression plants had higher levels of capsidiol but not higher resistance.

#### Accession numbers

Sequence data from this article can be found in the GenBank data library under accession numbers: XM\_019375732.1 (*HMG-CoA synthase*), XM\_019375278.1 (*MVAPP decarboxylase*), XM\_019403732.1 (*FPP synthase*), XM\_019409657.1

(*squalene synthase*), XM\_019408556.1 (*EAS12*), and XM\_019399671.1 (*NaERF2-like*).

#### Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. *Alternaria alternata*-elicited *NaEAS12* transcripts in EV and plants individually silenced with the top five up-regulated transcription factors.

Fig. S2. Overexpression of *NaERF2-like* does not affect plant resistance.

Fig. S3. Silencing *NaEIN2* has a great impact on *A. alternata*-induced transcripts of *NaF6'H1* but does not affect transcripts of *NaEAHs* and *NaEASs*.

Table S1. Primers used in this study.

Table S2. Transcriptome analysis revealed regulation of genes involved in sesquiterpene biosynthesis in three biological replicate *N. attenuata* leaves after *A. alternata* inoculation at 1 d.

Table S3. Transcriptome analysis revealed top six highly elicited transcriptional factor genes in three biological replicate *A. alternata*-inoculated *N. attenuata* leaves at 1 d.

#### Acknowledgements

We thank Prof. Joe Chappell (University of Kentucky, USA) for the capsidiol standard, the Service Center for Experimental Biotechnology of Kunming Institute of Botany, the Chinese Academy of Sciences (CAS), for plant growth support. This project was supported by the National Science Foundation of China (NSFC Grant No. 31670262) and the Key Project of Applied Basic Research Program of Yunnan (2014FA040) to JW, NSFC grant (No. 31700231), the Applied Basic Research Program of Yunnan (2017FB048) and CAS "Light of West China" Program to LM.

#### References

- Ahuja I, Kissen R, Bones AM. 2012. Phytoalexins in defense against pathogens. *Trends in Plant Science* **17**, 73–90.
- Bailey JA, Burden RS, Vincent GG. 1975. Capsidiol: antifungal compound produced in *Nicotiana tabacum* and *Nicotiana clevelandii* following infection with tobacco necrosis virus. *Phytochemistry* **14**, 597–597.
- Bergelson J, Purrington CB, Palm CJ, Lopez-Gutierrez JC. 1996. Costs of resistance: a test using transgenic *Arabidopsis thaliana*. *Proceedings of the Royal Society B: Biological Sciences* **263**, 1659–1663.
- Berrolcal-Lobo M, Molina A, Solano R. 2002. Constitutive expression of ETHYLENE-RESPONSE-FACTOR1 in *Arabidopsis* confers resistance to several necrotrophic fungi. *The Plant Journal* **29**, 23–32.
- Chen M, Yan T, Shen Q, et al. 2017. GLANDULAR TRICHOME-SPECIFIC WRKY 1 promotes artemisinin biosynthesis in *Artemisia annua*. *New Phytologist* **214**, 304–316.
- El Oirdi M, Trapani A, Bouarab K. 2010. The nature of tobacco resistance against *Botrytis cinerea* depends on the infection structures of the pathogen. *Environmental Microbiology* **12**, 239–253.
- Facchini PJ, Chappell J. 1992. Gene family for an elicitor-induced sesquiterpene cyclase in tobacco. *Proceedings of the National Academy of Sciences, USA* **89**, 11088–11092.
- Glazebrook J. 2005. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual Review of Phytopathology* **43**, 205–227.
- Grosskinsky DK, Naseem M, Abdelmohsen UR, et al. 2011. Cytokinins mediate resistance against *Pseudomonas syringae* in tobacco through increased antimicrobial phytoalexin synthesis independent of salicylic acid signaling. *Plant Physiology* **157**, 815–830.

- Guedes MEM, Kuc J, Hammerschmidt R, Bostock R.** 1982. Accumulation of 6 sesquiterpenoid phytoalexins in tobacco leaves infiltrated with *Pseudomonas lachrymans*. *Phytochemistry* **21**, 2987–2988.
- Hao D, Ohme-Takagi M, Sarai A.** 1998. Unique mode of GCC box recognition by the DNA-binding domain of ethylene-responsive element-binding factor (ERF domain) in plant. *The Journal of Biological Chemistry* **273**, 26857–26861.
- Hao D, Yamasaki K, Sarai A, Ohme-Takagi M.** 2002. Determinants in the sequence specific binding of two plant transcription factors, CBF1 and NtERF2, to the DRE and GCC motifs. *Biochemistry* **41**, 4202–4208.
- Huang PY, Catinot J, Zimmerli L.** 2016. Ethylene response factors in Arabidopsis immunity. *Journal of Experimental Botany* **67**, 1231–1241.
- Kallenbach M, Bonaventure G, Gilardoni PA, Wissgott A, Baldwin IT.** 2012. *Empoasca* leafhoppers attack wild tobacco plants in a jasmonate-dependent manner and identify jasmonate mutants in natural populations. *Proceedings of the National Academy of Sciences, USA* **109**, E1548–57.
- Kliebenstein DJ, Rowe HC, Denby KJ.** 2005. Secondary metabolites influence *Arabidopsis/Botrytis* interactions: variation in host production and pathogen sensitivity. *The Plant Journal* **44**, 25–36.
- Krügel T, Lim M, Gase K, Halitschke R, Baldwin IT.** 2002. *Agrobacterium*-mediated transformation of *Nicotiana attenuata*, a model ecological expression system. *Chemoecology* **12**, 177–183.
- LaMondia JA.** 2001. Outbreak of brown spot of tobacco caused by *Alternaria alternata* in connecticut and massachusetts. *Plant Disease* **85**, 230.
- Li J, Wu J.** 2016. Scopolin, a glycoside form of the phytoalexin scopoletin, is likely involved in the resistance of *Nicotiana attenuata* against *Alternaria alternata*. *Journal of Plant Pathology* **98**, 641–644.
- Li S, Wang H, Li F, Chen Z, Li X, Zhu L, Wang G, Yu J, Huang D, Lang Z.** 2015. The maize transcription factor EREB58 mediates the jasmonate-induced production of sesquiterpene volatiles. *The Plant Journal* **84**, 296–308.
- Mengiste T.** 2012. Plant immunity to necrotrophs. *Annual Review of Phytopathology* **50**, 267–294.
- Mialoundama AS, Heintz D, Debayle D, Rahier A, Camara B, Bouvier F.** 2009. Abscisic acid negatively regulates elicitor-induced synthesis of capsidiol in wild tobacco. *Plant Physiology* **150**, 1556–1566.
- Nafisi M, Goregaoker S, Botanga CJ, Glawischnig E, Olsen CE, Halkier BA, Glazebrook J.** 2007. *Arabidopsis* cytochrome P450 monooxygenase 71A13 catalyzes the conversion of indole-3-acetaldoxime in camalexin synthesis. *The Plant Cell* **19**, 2039–2052.
- Ohtsu M, Shibata Y, Ojika M, Tamura K, Hara-Nishimura I, Mori H, Kawakita K, Takemoto D.** 2014. Nucleoporin 75 is involved in the ethylene-mediated production of phytoalexin for the resistance of *Nicotiana benthamiana* to *Phytophthora infestans*. *Molecular Plant-Microbe Interactions* **27**, 1318–1330.
- Pré M, Atallah M, Champion A, De Vos M, Pieterse CM, Memelink J.** 2008. The AP2/ERF domain transcription factor ORA59 integrates jasmonic acid and ethylene signals in plant defense. *Plant Physiology* **147**, 1347–1357.
- Ralston L, Kwon ST, Schoenbeck M, Ralston J, Schenk DJ, Coates RM, Chappell J.** 2001. Cloning, heterologous expression, and functional characterization of 5-*epi*-aristolochene-1,3-dihydroxylase from tobacco (*Nicotiana tabacum*). *Archives of Biochemistry and Biophysics* **393**, 222–235.
- Ratcliff F, Martin-Hernandez AM, Baulcombe DC.** 2001. Technical Advance. Tobacco rattle virus as a vector for analysis of gene function by silencing. *The Plant Journal* **25**, 237–245.
- Saedler R, Baldwin IT.** 2004. Virus-induced gene silencing of jasmonate-induced direct defences, nicotine and trypsin proteinase-inhibitors in *Nicotiana attenuata*. *Journal of Experimental Botany* **55**, 151–157.
- Sanchez-Vallet A, Ramos B, Bednarek P, Lopez G, Pislewska-Bednarek M, Schulze-Lefert P, Molina A.** 2010. Tryptophan-derived secondary metabolites in *Arabidopsis thaliana* confer non-host resistance to necrotrophic *Plectosphaerella cucumerina* fungi. *The Plant Journal* **63**, 115–127.
- Schuck S, Weinhold A, Luu VT, Baldwin IT.** 2014. Isolating fungal pathogens from a dynamic disease outbreak in a native plant population to establish plant-pathogen bioassays for the ecological model plant *Nicotiana attenuata*. *PLoS One* **9**, e102915.
- Shibata Y, Kawakita K, Takemoto D.** 2010. Age-related resistance of *Nicotiana benthamiana* against hemibiotrophic pathogen *Phytophthora infestans* requires both ethylene- and salicylic acid-mediated signaling pathways. *Molecular Plant-Microbe Interactions* **23**, 1130–1142.
- Shibata Y, Ojika M, Sugiyama A, Yazaki K, Jones DA, Kawakita K, Takemoto D.** 2016. The full-size ABCG transporters Nb-ABCG1 and Nb-ABCG2 function in pre- and postinvasion defense against *Phytophthora infestans* in *Nicotiana benthamiana*. *The Plant Cell* **28**, 1163–1181.
- Solano R, Stepanova A, Chao Q, Ecker JR.** 1998. Nuclear events in ethylene signaling: a transcriptional cascade mediated by ETHYLENE-INSENSITIVE3 and ETHYLENE-RESPONSE-FACTOR1. *Genes & Development* **12**, 3703–3714.
- Stoessel A, Unwin CH, Ward EWB.** 1972. Postinfectious inhibitors from plants I. Capsidiol, an antifungal compound from *Capsicum frutescens*. *Phytopathology* **74**, 141–152.
- Sun H, Hu X, Ma J, Hettenhausen C, Wang L, Sun G, Wu J, Wu J.** 2014a. Requirement of ABA signalling-mediated stomatal closure for resistance of wild tobacco to *Alternaria alternata*. *Plant Pathology* **63**, 1070–1077.
- Sun H, Song N, Ma L, Li J, Ma L, Wu J, Wu J.** 2017. Ethylene signalling is essential for the resistance of *Nicotiana attenuata* against *Alternaria alternata* and phytoalexin scopoletin biosynthesis. *Plant Pathology* **66**, 277–284.
- Sun H, Wang L, Zhang B, Ma J, Hettenhausen C, Cao G, Sun G, Wu J, Wu J.** 2014b. Scopoletin is a phytoalexin against *Alternaria alternata* in wild tobacco dependent on jasmonate signalling. *Journal of Experimental Botany* **65**, 4305–4315.
- von Dahl CC, Winz RA, Halitschke R, Kuhnemann F, Gase K, Baldwin IT.** 2007. Tuning the herbivore-induced ethylene burst: the role of transcript accumulation and ethylene perception in *Nicotiana attenuata*. *The Plant Journal* **51**, 293–307.
- Ward EWB, Unwin CH, Stoessel A.** 1974. Postinfectious inhibitors from plants 13. Fungitoxicity of phytoalexin, capsidiol, and related sesquiterpenes. *Canadian Journal of Botany – Revue Canadienne De Botanique* **52**, 2481–2488.
- Welsch R, Maass D, Voegel T, Dellapenna D, Beyer P.** 2007. Transcription factor RAP2.2 and its interacting partner SINAT2: stable elements in the carotenogenesis of Arabidopsis leaves. *Plant Physiology* **145**, 1073–1085.
- Wu J, Wang L, Baldwin IT.** 2008. Methyl jasmonate-elicited herbivore resistance: does MeJA function as a signal without being hydrolyzed to JA? *Planta* **227**, 1161–1168.
- Wu J, Wang L, Wünsche H, Baldwin IT.** 2013. Narboh D, a respiratory burst oxidase homolog in *Nicotiana attenuata*, is required for late defense responses after herbivore attack. *Journal of Integrative Plant Biology* **55**, 187–198.
- Xu Y, Wang J, Wang S, Wang J, Chen X.** 2004. Characterization of GaWRKY1, a cotton transcription factor that regulates the sesquiterpene synthase gene (+)- $\delta$ -cadinene synthase-A. *Plant Physiology* **135**, 507–515.
- Xu Z, Song N, Ma L, Fang D, Wu J.** 2018. NaPDR1 and NaPDR1-like are essential for the resistance of *Nicotiana attenuata* against fungal pathogen *Alternaria alternata*. *Plant Diversity* **40**, 68–73.
- Yang C, Fang X, Wu X, Mao Y, Wang L, Chen X.** 2012. Transcriptional regulation of plant secondary metabolism. *Journal of Integrative Plant Biology* **54**, 703–712.
- Yoo SD, Cho YH, Sheen J.** 2007. *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nature Protocols* **2**, 1565–1572.
- Zhang F, Fu X, Lv Z, et al.** 2015. A basic leucine zipper transcription factor, AabZIP1, connects abscisic acid signaling with artemisinin biosynthesis in *Artemisia annua*. *Molecular Plant* **8**, 163–175.
- Zhao Y, Wei T, Yin KQ, Chen Z, Gu H, Qu LJ, Qin G.** 2012. *Arabidopsis* RAP2.2 plays an important role in plant resistance to *Botrytis cinerea* and ethylene responses. *New Phytologist* **195**, 450–460.