

## Ecology of *Thioploca* spp.: Nitrate and Sulfur Storage in Relation to Chemical Microgradients and Influence of *Thioploca* spp. on the Sedimentary Nitrogen Cycle

JAKOB ZOPFI,<sup>1\*</sup> THOMAS KJÆR,<sup>2</sup> LARS P. NIELSEN,<sup>2</sup> AND BO BARKER JØRGENSEN<sup>1</sup>

Max Planck Institute for Marine Microbiology, D-28359 Bremen, Germany,<sup>1</sup> and Institute of Biological Sciences, University of Aarhus, DK-8000 Aarhus C, Denmark<sup>2</sup>

Received 27 November 2000/Accepted 25 September 2001

Microsensors, including a recently developed  $\text{NO}_3^-$  biosensor, were applied to measure  $\text{O}_2$  and  $\text{NO}_3^-$  profiles in marine sediments from the upwelling area off central Chile and to investigate the influence of *Thioploca* spp. on the sedimentary nitrogen metabolism. The studies were performed in undisturbed sediment cores incubated in a small laboratory flume to simulate the environmental conditions of low  $\text{O}_2$ , high  $\text{NO}_3^-$ , and bottom water current. On addition of  $\text{NO}_3^-$  and  $\text{NO}_2^-$ , *Thioploca* spp. exhibited positive chemotaxis and stretched out of the sediment into the flume water. In a core densely populated with *Thioploca*, the penetration depth of  $\text{NO}_3^-$  was only 0.5 mm and a sharp maximum of  $\text{NO}_3^-$  uptake was observed 0.5 mm above the sediment surface. In sediments with only few *Thioploca* spp.,  $\text{NO}_3^-$  was detectable down to a depth of 2 mm and the maximum consumption rates were observed within the sediment. No chemotaxis toward nitrous oxide ( $\text{N}_2\text{O}$ ) was observed, which is consistent with the observation that *Thioploca* does not denitrify but reduces intracellular  $\text{NO}_3^-$  to  $\text{NH}_4^+$ . Measurements of the intracellular  $\text{NO}_3^-$  and  $\text{S}^0$  pools in *Thioploca* filaments from various depths in the sediment gave insights into possible differences in the migration behavior between the different species. Living filaments containing significant amounts of intracellular  $\text{NO}_3^-$  were found to a depth of at least 13 cm, providing final proof for the vertical shuttling of *Thioploca* spp. and nitrate transport into the sediment.

Although the ability of microorganisms to oxidize reduced sulfur compounds with nitrate as the electron acceptor has been known for about one hundred years (1) and several pure cultures have been obtained and studied (see, e.g., references 39 and 42), only a little is known about the ecological significance of this type of metabolism. For instance, the first report showing a clear coupling between the sulfur and nitrogen cycles in the marine environment was sulfide-driven denitrification at the oxic-anoxic interface in the water column of the Central Baltic Sea (3).

It was recently discovered that sulfide-oxidizing bacteria of the genus *Thioploca* possess large nitrate-filled vacuoles (10). *Thioploca* is highly abundant in the shelf sediments along Peru and Chile (11, 12, 14, 29), and it was therefore suspected that these organisms play a major role in coupling the biogeochemical cycles of nitrogen and sulfur in upwelling areas (10). The observation of benthic *Thioploca* filaments in the upwelling area of the Arabian Sea (20) and the finding of both *Thioploca* and the spherical, nitrate-storing bacterium *Thiomargarita* off Namibia (34) support this conclusion. Nitrate-storing sulfide-oxidizing bacteria have also been observed at hydrothermal vents and cold seeps and in organic-rich sediments (23, 24, 44).

Currently, none of the nitrate-storing sulfur bacteria is in pure culture, and alternative methods have to be applied to study their physiology and ecology. Enzyme preparations from

partially purified *Beggiatoa* samples showed high nitrate reductase and ribulose-1,5-bisphosphate carboxylase-oxygenase activity and provide the first biochemical evidence for the use of nitrate as a electron acceptor for sulfide oxidation and chemoautotrophic growth (23). Incubation experiments with partially purified *Thioploca* filaments revealed that sulfide ( $\text{H}_2\text{S}$ ) was first rapidly oxidized to  $[\text{S}^0]$ , which was then further oxidized to sulfate ( $\text{SO}_4^{2-}$ ) in a second independent step. Intravacuolar  $[\text{NO}_3^-]$  served as the electron acceptor and was reduced to ammonium ( $\text{NH}_4^+$ ) (25). Radiolabeled bicarbonate ( $\text{H}^{14}\text{CO}_3^-$ ) and  $[2\text{-}^{14}\text{C}]\text{acetate}$  were assimilated, indicating that *Thioploca* is a facultative chemolithoautotroph capable of mixotrophic growth (22, 25). Whole-core incubations in small laboratory flumes have helped to unveil the chemotactic behavior of *Thioploca* under changing environmental conditions. *Thioploca* showed positive chemotaxis toward nitrate and low sulfide concentrations ( $<100\ \mu\text{M}$ ) but a phobic reaction toward oxygen and high sulfide concentrations (17). These observations and the finding of mostly vertically oriented living filaments several centimeters deep in the sediment led to the suggestion that *Thioploca* shuttles up and down between  $\text{NO}_3^-$ -rich bottom water and  $\text{H}_2\text{S}$ -containing sediment (10, 17).

The samples for this study were collected at stations within and off the Bay of Concepción in central Chile, where the species composition and annual dynamics of the *Thioploca* population are known from a previous study (36). On the shelf, the community was composed mainly of *T. araucae*, *T. chileae*, and a yet undescribed form of *Thioploca* with much shorter cells. This so-called short-cell morphotype (SCM) (35, 36) is usually found in deeper sediment layers than the two other

\* Corresponding Author. Present address: Institute of Biology and Danish Center for Earth System Science, University of Southern Denmark, Odense, Campusvej 55, DK-5230 Odense M, Denmark, Phone: 45 6550 2745. Fax: 45 6593 0457. E-mail: jzopfi@biology.sdu.dk.

species and is characterized by rounded cells and a cell length/diameter ratio of  $\leq 0.48$ . The SCM is closely related but not identical to the known *Thioploca* species as revealed by partial 16S rDNA analysis (35). Within the Bay of Concepción, large vacuolated filaments cover the sediment surface during part of the year. They are, apart from the lack of sheaths, phenotypically and phylogenetically almost identical to *T. araucae* (36, 40).

The aim of this study was to gain information about the ecology of *Thioploca* spp. and the influence of these organisms on the nitrogen and sulfur cycles in the habitat. By the use of microsensors, we show that the nitrate uptake of the sediment is strongly influenced by *Thioploca*. Chemotaxis toward inorganic nitrogen compounds was studied, and measurements of  $[S^0]$  and  $[NO_3^-]$  concentrations were used to test the concept of a vertical shuttling between the nitrate-rich bottom water and the deeper sediment layers. We also demonstrate that nitrate is transported by *Thioploca* down to a sediment depth of at least 13 cm.

#### MATERIALS AND METHODS

**Abbreviations.** Nitrate and elemental sulfur are generally abbreviated by  $NO_3^-$  and  $S^0$ , respectively. For the intracellular pools  $[NO_3^-]$  and  $[S^0]$  are used.

**Study area.** The continental shelf region off the Concepción Bay (central Chile) is characterized by intense seasonal upwelling. Between austral late spring and early fall, southern and southwestern winds prevail and the northward-flowing Sub-Antarctic Surface water is forced off the coast, leading to upwelling of Equatorial Subsurface water from the Poleward Undercurrent at 100 to 400 m (37). The Equatorial Subsurface water is characterized by high salinity (34.4 to 34.8‰), low temperature (8.5 to 10.5°C), low oxygen concentrations ( $< 20 \mu M$ ), but high nitrate (about  $25 \mu M$ ) and nutrient concentrations (37). Upwelling off central Chile is intermittent and usually lasts between 2 and 7 days (16). Primary and secondary productivity greatly increase when the nutrient-rich water is transported up into the euphotic zone. For the coastal upwelling area off central Chile, a primary production of  $9.6 \text{ g of C m}^{-2} \text{ day}^{-1}$  has been reported (10), which is one of the highest observed in marine environments. Due to the lack of oxygen and sufficient amounts of alternative electron acceptors, sedimentary organic matter is almost exclusively degraded by sulfate-reducing bacteria (41). The sulfate reduction rates reported for this area ( $170$  to  $4,670 \text{ nmol cm}^{-3} \text{ day}^{-1}$ ) [(8)] are among the highest observed in coastal margins, but free sulfide concentrations in these sediments are surprisingly low, indicating an efficient reoxidation of sulfide (8; J. Zopfi, M. E. Böttcher, and B. B. Jørgensen, submitted for publication).

**Sampling and site description.** During January and February 1997, we repeatedly sampled sediment from three stations within and off the Bay of Concepción (Fig. 1). The sediment was collected from the research vessel *Kay Kay* of the University of Concepción (Concepción, Chile) by means of a small gravity corer. The cores were stored onboard at 4°C in a refrigerator and were transported on the day of sampling to the Marine Biological Station of the University of Concepción in Dichato, where all experiments were performed.

Station 4 ( $36^\circ 38' 08'' \text{ S}$ ;  $073^\circ 02' 03'' \text{ W}$ ) was 24 m deep and located within the bay (Fig. 1). The sediment at this station was highly sulfidic (up to  $1,200 \mu M$  at 7 cm deep [Zopfi et al., submitted]) and was uniformly black below the brownish uppermost 3 to 4 mm. The top 4.5 to 5 cm of the sediment was a flocculent ooze, with mass accumulation of *Beggiatoa* spp. The sediment of Station 7 ( $36^\circ 36' 05'' \text{ S}$ ;  $073^\circ 00' 06'' \text{ W}$ ; 32 m deep), at the mouth of the bay, was covered with a brown, spongy layer 1.5 cm thick, which was densely populated by *Thioploca* spp. Below this layer was a 0.4-cm-thick band of black iron sulfide, followed by gray-brownish sediment. Burrows of sediment-dwelling organisms were observed. The sediment of Station 18 ( $35^\circ 30' 08'' \text{ S}$ ;  $073^\circ 07' 06'' \text{ W}$ ; 88 m deep) had a similar structure, with a spongy layer 0.5 to 1 cm thick, a thin black layer (0.2 to 0.3 cm thick), and then gray sediment below. *Thioploca*, however, was much less abundant, and no animal burrows were observed. The concentration of free sulfide was  $< 6 \mu M$  down to a depth of 20 cm at Stations 7 and 18. During the sampling period, the bottom-water concentration of  $O_2$  was  $< 2$  to  $7 \mu M$  at all stations. The  $NO_3^-$  concentrations were about  $6 \mu M$  at Station 4, and 7 to  $24 \mu M$  at Stations 7 and 18 (36).




FIG. 1. Sampling sites within the Bay of Concepción and on the adjacent continental shelf off central Chile.

**Microsensor measurements and flux calculations.** A Clark-type  $O_2$  microsensor with a guard cathode was used to measure oxygen microprofiles (28). Nitrate was measured with a microbiosensor consisting of an electrochemical  $N_2O$  microsensor surrounded by an outer casing (19). A 100 to 200- $\mu m$ -long reaction chamber was formed between the tip of the internal  $N_2O$  sensor and the impermeable membrane at the tip of the outer casing. An  $N_2O$  reductase-deficient culture of *Agrobacterium* immobilized in the reaction chamber transformed  $NO_3^-$  and  $NO_2^-$  to  $N_2O$ , which was detected by the  $N_2O$  microsensor. Due to this design, the microsensor measured the combined  $NO_3^-$ ,  $NO_2^-$ , and  $N_2O$  concentrations in a sample. The  $N_2O$  microsensor was constructed like the  $O_2$  microsensor (28) but the cathode was plated with silver and the electrolyte consisted of 0.5 M NaOH and 0.5 M KCl. The  $N_2O$  microsensor was polarized at  $-1.2 \text{ V}$  against an Ag/AgCl anode immersed in the electrolyte. The current from the microsensor was measured with a custom-made pA meter and recorded on a strip chart recorder. The active silver surface of the  $N_2O$  sensing cathode is also sensitive toward  $H_2S$ . However, the concentrations of free sulfide in the top centimeters of the Station 7 and 18 sediment was usually  $< 1 \mu M$ , and hence no interference with  $H_2S$  was anticipated. Calibrations and measurements were done at the same temperature. The detection limit of the sensor was about  $3 \mu M NO_3^-$ , and the linear range was 3 to  $70 \mu M NO_3^-$ . The sensitivity for  $N_2O$  was about  $1 \mu M$ , and the signal was linear up to least  $1,000 \mu M$ . The response time for 90% signal intensity was about 50 s. A resting time of 1 min was used for each depth step in a profile. The position of the sediment surface was determined for each profile using a dissection microscope (magnification,  $\times 10$  to  $\times 50$ ).

Fluxes across the sediment-water interface were calculated from the upper linear part of the microsensor profiles by Fick's first law of diffusion,  $J = -D \times dC/dx$ , where  $J$  is the flux (in micromoles per square centimeter per second),  $D$  is the diffusion coefficient (in square centimeters per second), and  $dC/dx$  is the concentration gradient (in micromoles per cubic centimeter per centimeter). The activity profiles were calculated as described in detail elsewhere (T. Kjær, L.-H. Larsen, and N. P. Revsbech, unpublished data). Tabulated diffusion coefficients were recalculated for the *in situ* temperature and salinity (4, 21). Since the sediment was highly porous in the top 1 cm, the same diffusion coefficients of  $1.8 \times 10^{-5}$  and  $1.41 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1}$  for  $O_2$  and  $NO_3^-$  respectively, were used for the sediment and water phase.

**Experimental setup.** To simulate the environmental conditions prevailing off the coast, measurements were done in a small flow chamber where deaerated surface water from Station 7 was circulated. A small aquarium pump was used to create flows of approximately 4 to  $6 \text{ cm s}^{-1}$  about 2 cm above the sediment surface. The flow chamber consisted of two horizontal Plexiglas plates that were separated with a spacer. Both Plexiglas plates contained a central hole ( $50 \text{ cm}^2$ ). A sediment core was brought into the flow through the hole in the bottom plate, and the microsensors were introduced through the hole in the top plate. A dissolved- $O_2$  concentration of about  $5 \mu M$  was maintained in the circulating water by adjusting the area of air-exposed seawater at the top hole, and the water was kept at the *in situ* temperature of  $12^\circ \text{C}$  by a thermostatted circulating cooler.

For the chemotaxis experiments,  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ , or  $\text{N}_2\text{O}$  was added to nitrate-depleted flume water and the behavior of *Thioploca* was observed from above through a dissection microscope. The number of filaments emerging from their sheaths was determined by setting the focus plane at about 2 mm above the sediment surface and by counting the filaments penetrating the plane. The length of a filament was determined with a measuring eyepiece and by focusing down from the filament tip to the sediment surface.

**Extraction and analysis of  $[\text{NO}_3^-]$  and  $[\text{S}^0]$  in *Thioploca*.** Bundles of *Thioploca* filaments from different sediment depths of Station 7 were picked out and aligned in a film of seawater on a microscope slide. Forceps and needles were used to rip the sheath apart so that intact single filaments could be isolated.

The length ( $l$ ) and diameter ( $d$ ) of each filament were determined, and the biovolume ( $V$ ) was calculated ( $V = \pi l d^2/4$ ). According to the cell diameter and the cell-length-to-diameter ratio, the organism was identified as *T. araucae*, *T. chilense*, or SCM (36). With the same formula, the volume ratio between the cytoplasm and vacuole was determined for each species by using the following cell dimensions (length, diameter): *T. araucae*, 14.4  $\mu\text{m}$ , 15.4  $\mu\text{m}$ ; *T. chilense*, 35.5  $\mu\text{m}$ , 24.2  $\mu\text{m}$ ; SCM, 35.5  $\mu\text{m}$ , 9.4  $\mu\text{m}$ . For all three species, a mean cytoplasm thickness of 1  $\mu\text{m}$  was used (36; H. Schulz, personal communication).

A single filament was then picked up on the tip of a purpose-made glass needle and left to dry in air for a few minutes, so that the filaments died and the cells cracked. Nitrate was extracted from the filament by dipping the glass needle for 5 s into a droplet of 20  $\mu\text{l}$  of demineralized water.

Nitrate was analyzed by the cadmium reduction method (13), with some adaptations to small amounts. The cadmium column typically consisted of a 3-cm-long glass tube (inner diameter, 1.1 mm) with a slightly coiled, 1-mm-wide cadmium rod inside. A bent glass capillary with a pointed tip was mounted on the upper end of the column, and a straight capillary was mounted on the bottom. The total system could contain about 10  $\mu\text{l}$  of liquid, and once activated the column was always kept filled with buffer solution. The column was held almost vertical, and when the upper tip was dipped in liquid, gravity created a water flow of about 30  $\mu\text{l min}^{-1}$  through the column. Capillary forces in the pointed tip prevented intrusion of air when the tip was out of water. The 20- $\mu\text{l}$  sample with the extracted nitrate was sucked up, immediately followed by 40  $\mu\text{l}$  of buffer solution. Below the reduction column the sample and buffer solution was collected in a 300- $\mu\text{l}$  well of a microplate. Aliquots (20  $\mu\text{l}$ ) of reagents were added by the standard procedure, and 200  $\mu\text{l}$  of demineralized water was mixed in as well. The color intensity was read at 670 nm, and corrections for turbidity, if any, were made by reading at 405 nm.  $\text{NO}_3^-$  standard solutions (20  $\mu\text{l}$  each) were processed parallel with the filament extracts and used for calibration. Linearity was observed up to 250  $\mu\text{M}$ . The reduction efficiency was checked by using  $\text{NO}_2^-$  standards, and the cadmium column was reactivated or renewed whenever required. The detection limit of the  $\text{NO}_3^-$  assay was about 20 pmol. We analyzed filaments with biovolumes from 0.0006 to 0.0154  $\text{mm}^3$ ; the detection limit was therefore equivalent to an  $[\text{NO}_3^-]$  concentration of 33 mM for the smallest filament and 2 mM for the largest filament. No  $\text{NO}_3^-$  was detected when empty sheaths were analyzed or when a filament was dried and extracted a second time, thus confirming that no significant contamination or loss of nitrate occurred in the procedure.

After extraction of nitrate, the filament was air dried again and immersed in 50  $\mu\text{l}$  of methanol for extraction of  $[\text{S}^0]$ . The complete dissolution of sulfur globules was verified by extraction time series and light microscopy. Elemental sulfur in the extract was quantified as cyclo-octasulfur ( $\text{S}_8$ ) by high-performance liquid chromatography. Separation was done on a Zorbax ODS column (125 by 4 mm, 5  $\mu\text{m}$ ; Knauer, Germany) with methanol (100%; high-performance liquid chromatography grade) as the eluent at a flow rate of 1  $\text{ml min}^{-1}$ .  $\text{S}_8$  eluted after 3.5 min and was detected at 265 nm. The detection limit was 43 pmol and was equivalent to an  $\text{S}^0$  concentration of 72 mM for the smallest filament and 3 mM for the largest filament. Repeated measurements of  $[\text{S}^0]$  in filaments of the same species inhabiting a common sheath showed a variability of <10% relative standard deviation. No loss of  $\text{S}^0$  was observed during the preceding  $[\text{NO}_3^-]$  extraction.

Elemental sulfur in the bulk sediment was extracted from Zn-preserved samples with pure methanol for 16 h on a rotary shaker; the sediment-to-extractant ratio was about 1:20 (wet wt/vol).  $\text{S}^0$  in the filtered (0.45- $\mu\text{m}$ -pore-size filter) extracts was quantitated as described above. The variability within triplicate  $\text{S}^0$  extractions was <14%.

**Statistical treatment.** The correlation between  $[\text{NO}_3^-]$  and  $[\text{S}^0]$  was determined and tested for significance by the method of Spearman (30). The  $[\text{NO}_3^-]/[\text{S}^0]$  ratios of filaments from different depth intervals were tested for similarity with the H test of Kruskal and Wallis (30).




FIG. 2. Laboratory flume measurements of  $\text{O}_2$  and  $\text{NO}_3^-$  concentrations in sediments with a high (Station 7) and very low (Station 18) density of *Thioploca* spp. The broken line indicates the sediment-water interface.

## RESULTS AND DISCUSSION

### Influence of *Thioploca* on $\text{NO}_3^-$ profiles and uptake rates.

Nitrate has been measured in biofilms and lake sediments by liquid ion exchanger (LIX)-based microsensors (7, 32, 38), but similar measurements in marine environments were not possible due to the interference of  $\text{Cl}^-$  ions. In this study we used a recently developed  $\text{NO}_3^-$  biosensor that allowed us to measure microprofiles in sediments from the upwelling system off central Chile and to study the influence of *Thioploca* on the  $\text{NO}_3^-$  uptake. Huettel et al. (17) suggested that filament protrusion may be a strategy to overcome the diffusion limitation to  $\text{NO}_3^-$  uptake imposed by the boundary layer and that *Thioploca* may thereby outcompete  $\text{NO}_3^-$ -consuming bacteria in the sediment. To test this hypothesis, we incubated sediment from two different stations in the flume under similar conditions to those described above and measured the oxygen and nitrate microprofiles.

At the time of sampling, the sediment of Station 7 was densely populated by *Thioploca* spp., with a total wet biomass of 44  $\text{g m}^{-2}$  (36) and the filaments stretched out of the sediment when nitrate was present in the flume water. *Thioploca* was much less abundant in the second core from Station 18, where the wet biomass was only 9  $\text{g m}^{-2}$  (36). Protruding filaments were only sporadically observed in the core studied. This difference in *Thioploca* spp. abundance was clearly reflected in the  $\text{O}_2$  and  $\text{NO}_3^-$  profiles (Fig. 2). The profiles from Station 7 measured in the vicinity of *Thioploca* filaments were oddly shaped, and  $\text{O}_2$  and  $\text{NO}_3^-$  hardly penetrated to the sediment surface. Since the filaments protruded into the flume water, maximum  $\text{NO}_3^-$  uptake rates occurred above the sediment surface (Fig. 3) and the  $\text{NO}_3^-$  penetration was only 0.5 mm. Station 18 showed an  $\text{O}_2$  microprofile normal for marine




FIG. 3. Vertical distribution of  $\text{NO}_3^-$  uptake rates in sediment with a high (Station 7) and low (Station 18) density of *Thioploca* filaments. The broken line indicates the sediment-water interface.

sediments, with a diffusive boundary layer thickness of about 1 mm and a maximal  $\text{O}_2$  penetration depth of 1 mm. Nitrate penetrated about 2 mm into the sediment, and the linear range of the  $\text{NO}_3^-$  gradient (Fig. 2) and maximal uptake rates were both within the sediment (Fig. 3). The profile structure and penetration depth were very similar to those found in organic-rich lake sediments (19, 38).

Interestingly, the diffusive boundary layer at Station 7 was  $>1.5$  mm, considerably thicker than at Station 18 (Fig. 2). A likely explanation for this could be that *Thioploca* filaments protruding from the sediment impede the water flow, which leads to a thickening of the boundary layer and thus to a lower diffusive exchange across the sediment-water interface. Thus, the oxygen uptake at Station 7 was only  $0.26 \pm 0.11 \text{ mmol m}^{-2} \text{ day}^{-1}$  ( $n = 6$ ) compared to  $0.74 \pm 0.13 \text{ mmol m}^{-2} \text{ day}^{-1}$  ( $n = 6$ ) at Station 18. The calculated diffusive  $\text{NO}_3^-$  uptake at Station 7 was  $5.45 \pm 1.17 \text{ mmol m}^{-2} \text{ day}^{-1}$  ( $n = 4$ ), about 55% higher than at Station 18 ( $3.53 \pm 0.76 \text{ mmol m}^{-2} \text{ day}^{-1}$ ;  $n = 5$ ). In summary, *Thioploca* spp. have a major influence on the sedimentary  $\text{NO}_3^-$  metabolism. They penetrate up through the diffusive boundary layer, cause a reduced  $\text{NO}_3^-$  penetration depth, move the maximum  $\text{NO}_3^-$  uptake upward, and increase the areal  $\text{NO}_3^-$  uptake rate.

An interesting effect of the *Thioploca* community might be that nitrate-reducing bacteria in the sediment are outcompeted for  $\text{NO}_3^-$  and that denitrification consequently plays a minor role in sediments densely inhabited by *Thioploca* spp. The situation, however, is probably more complex, as indicated by denitrification measurements. Despite the presence of *Thioploca* spp., denitrification rates of 4.5 and 9  $\text{mmol m}^{-2} \text{ day}^{-1}$  were determined for Station 7 by adding  $100 \mu\text{M}$   $^{15}\text{NO}_3^-$  to the flume water (L. P. Nielsen, unpublished data). These values are slightly higher than usually observed in normal coastal sediments at similar  $\text{NO}_3^-$  concentrations (references 2 and 15 and references therein) and suggest only an incomplete sup-




FIG. 4. Response of *Thioploca* spp. to addition of  $10 \mu\text{M}$   $\text{NO}_2^-$  to the flume water as indicated by the number of filaments protruding  $>2$  mm out of the sediment and by the total length of all filaments exposed to flume water.

pression of denitrification. Nitrate for denitrification may be supplied by leakage from *Thioploca* filaments or via advective transport of bottom water into the sediment. Advective transport becomes progressively more important with increasing flow velocities (9) and may indeed have been an important process, because the top 1 to 2 cm of the sediment had a spongy consistency and was very porous due to the *Thioploca* mat (26). Furthermore, the in situ flow velocity of the bottom water near the sediment may well exceed the  $5 \text{ cm s}^{-1}$  that we have used in our experiments.

**Response of *Thioploca* filaments to  $\text{NO}_2^-$  and  $\text{N}_2\text{O}$ .** The chemotactic behavior of *Thioploca* spp. toward  $\text{O}_2$ ,  $\text{NO}_3^-$ , and  $\text{H}_2\text{S}$  was studied by Huettel et al. (17). However,  $\text{NO}_2^-$  and  $\text{N}_2\text{O}$  are also intermediates and by-products of nitrification and nitrate reduction processes (15, 18) and can be found in the oxygen minimum zone of upwelling areas (5, 6). We studied whether *Thioploca* spp. show chemotaxis toward  $\text{NO}_2^-$  and  $\text{N}_2\text{O}$  and whether they can utilize them as electron acceptors by using cores from Station 7 and adding  $\text{NO}_2^-$  or  $\text{N}_2\text{O}$  to nitrate-free flume water. On addition of  $10 \mu\text{M}$   $\text{NO}_2^-$ , the number and length of protruding filaments rapidly increased and reached a maximum after 1.2 h (Fig. 4). Single bundles were observed through the dissection microscope, and before the  $\text{NO}_2^-$  addition the filament tips were moving in and out of the sheath with frequent reversals just at the sediment-water interface. About 30 s after the  $\text{NO}_2^-$  addition, the reversals stopped and all the filaments moved upward, protruding from the sheath. When the  $\text{NO}_2^-$  concentration dropped below  $2.6 \mu\text{M}$ , the filaments began to retreat until the initial positions were reestablished (Fig. 4). A similar sequence was observed when  $\text{NO}_3^-$  was added (data not shown). It cannot completely be ruled out that the real tactic trigger was  $\text{NO}_3^-$  produced by nitrifying bacteria. However, the fast reaction of *Thioploca* and the low  $\text{O}_2$  concentrations limiting nitrification in the setup do support a direct response to  $\text{NO}_2^-$ .

In contrast to  $\text{NO}_3^-$  and  $\text{NO}_2^-$ , *Thioploca* filaments did not show chemotaxis toward  $\text{N}_2\text{O}$ , suggesting that nitrous oxide may not be used as an electron acceptor for sulfide oxidation. Nitrous oxide is the obligate precursor for dinitrogen formation during denitrification and can be used by most, although not all, denitri-




FIG. 5. Microprofile of N<sub>2</sub>O in sediment from Station 7. Nitrous oxide was added to the flume water after the depletion of nitrate.

ying bacteria as an electron acceptor (45). Reduction of N<sub>2</sub>O to ammonium has never been reported. The absence of a response to N<sub>2</sub>O therefore supports the finding of Otte et al. (25) that ammonium rather than dinitrogen, as previously assumed (10), is the terminal product of nitrate reduction.

Microprofiles of N<sub>2</sub>O were measured during the chemotaxis experiment, and an average profile ( $n = 3$ ) is depicted in Fig. 5. Since *Thioploca* did not stretch out from the sediment, the N<sub>2</sub>O profile exhibited a regular diffusive boundary layer of about 0.5 mm. However, even without the contribution of *Thioploca* spp., N<sub>2</sub>O was rapidly ( $15.1 \pm 1.5 \text{ mmol m}^{-2} \text{ day}^{-1}$ ) consumed within the first 3.5 mm of the sediment, demonstrating the potential for sedimentary dinitrogen formation.

**Internal [S<sup>0</sup>] and [NO<sub>3</sub><sup>-</sup>] concentrations.** A concentration range of 150 to 500 nmol mm<sup>-3</sup> has been reported for the [NO<sub>3</sub><sup>-</sup>] content of *Thioploca* cells (10), but to date nothing is known about the variability within the different species or about the [NO<sub>3</sub><sup>-</sup>]/[S<sup>0</sup>] ratio within individual filaments and whether it changes with depth. Additionally, *Thioploca* filaments have been found down to 26 cm deep (36), but it was not clear whether they were still alive and contained [NO<sub>3</sub><sup>-</sup>]. Therefore, *Thioploca* filaments were collected from different depths of a core from Station 7 and the concentrations of [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] were determined. The statistical analysis of the results from the depth intervals 0 to 1 cm, 1 to 2 cm, 2 to 3 cm, 3 to 4 cm, 4 to 7 cm, and 7 to 10 cm did not indicate significant differences in the [NO<sub>3</sub><sup>-</sup>]/[S<sup>0</sup>] ratios between the different sections ( $P > 0.1$ ). By the same statistical method, it was shown that the two lowest intervals from 10 to 13 cm and 13 to 16 cm, although not different from each other, were different from the first six intervals, with a high probability ( $P < 0.001$ ). Based on these results, the measurements of [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] from 0 to 10 cm and from 10 to 16 cm deep, respectively, are grouped together (Fig. 6A and B).

In the upper part (0 to 10 cm) of the core, *T. araucae* (57%), and *T. chileae* (30%) were the dominant mat-forming species

(Fig. 6A), with the rest being SCM thioplocas. The SCM were more abundant (81%) in the deeper section. Similarly, Schulz et al. (36) found the maximum *T. araucae* and *T. chileae* biomass close to the sediment-water interface whereas the SCM were most abundant below 7 cm deep. The difference in the species composition was also reflected in the [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] concentrations. Whereas [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] varied over similar concentration ranges (up to ~500 nmol mm<sup>-3</sup>) in the upper section (Fig. 6A), the values were significantly shifted toward lower [NO<sub>3</sub><sup>-</sup>] and higher [S<sup>0</sup>] concentrations in the 10- to 16-cm-deep section. Both the SCM and *T. chileae* stored [S<sup>0</sup>] up to ~800 nmol mm<sup>-3</sup>. However, the maximum [S<sup>0</sup>] concentration determined for *T. araucae* was only 355 nmol mm<sup>-3</sup> (Fig. 6B), which nicely corresponded to a lower volume ratio between cytoplasm and vacuole. Whereas SCM and *T. chileae* typically have a *C/V* ratio of 0.47, it is only 0.23 in *T. araucae*.

Based on results from chemotaxis experiments, it was concluded that *Thioploca* spp. fill their vacuoles with nitrate at the sediment surface. Then they migrate into deeper sediment layers, where they oxidize sulfide to [S<sup>0</sup>] and SO<sub>4</sub><sup>2-</sup> until low [NO<sub>3</sub><sup>-</sup>] concentrations are reached and the upward movement is induced again (10, 17). Such a behavior would imply that (i) there exists a negative relationship between the two storage compounds in a filament and (ii) filaments with a high [NO<sub>3</sub><sup>-</sup>]/[S<sup>0</sup>] ratio are found predominantly at the sediment surface whereas filaments rich in [S<sup>0</sup>] but depleted of [NO<sub>3</sub><sup>-</sup>] are found in deeper sections. This, however, seems to be valid only for SCM thioplocas, where a negative correlation between [S<sup>0</sup>] and [NO<sub>3</sub><sup>-</sup>] was found ( $r = -0.44$ ;  $n = 24$ ;  $P < 0.05$ ; Spearman test) and a decreasing [NO<sub>3</sub><sup>-</sup>]/[S<sup>0</sup>] ratio correlated with the sediment depth ( $r^2 = 0.71$ ). No significant correlations ( $P > 0.05$ ) were found in *T. araucae* or in *T. chileae*, which suggests that they migrate in a different manner. A reason for the lack of correlation was found when the average [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] concentrations of all individuals of a species in a given depth interval were calculated (Fig. 7). Despite the variability in the [NO<sub>3</sub><sup>-</sup>]/[S<sup>0</sup>] ratios, the average concentra-




FIG. 6. (A and B) Concentrations of  $[\text{NO}_3^-]$  and  $[\text{S}^0]$  in single *Thioploca* filaments collected at different sediment depths from Station 7. (C) Storage concentration in sheathless sulfur bacteria filaments from the sediment surface (0 to 2 cm) at Station 4. Open circles, *T. araucae*; solid circles, *T. chileae*; open triangles, SCM; dotted symbols, filaments from the deepest sediment section (13 to 16 cm); open diamonds, sheathless filaments.

tions were surprisingly constant down several centimeters deep. This is probably best explained by a continuous and rapid shuttling of *Thioploca* filaments relative to the metabolic rate. A further consequence of the continuous shuttling is that the filaments frequently reach the sediment surface, where they can recharge their  $[\text{NO}_3^-]$  storage. Ample supply of nitrate, on the other hand, could allow *Thioploca* to oxidize  $\text{H}_2\text{S}$  directly to  $\text{SO}_4^{2-}$  instead of forming  $[\text{S}^0]$  first. If this is true, it also explains why no clear correlation between  $[\text{S}^0]$  and  $[\text{NO}_3^-]$  was detected in *T. araucae* and *T. chileae*, respectively. However, final proof for a vertical shuttling of *Thioploca* spp. and nitrate transport into the sediment comes from the finding of living filaments with substantial  $[\text{NO}_3^-]$  concentrations ( $31 \pm 59 \text{ nmol mm}^{-3}$ ;  $n = 9$  [Fig. 6B]) in the 13- to 16-cm-deep sediment section. Calculations based on the turnover time of  $[\text{NO}_3^-]$  (8 to 10

days [25]) and a migration velocity of  $5 \text{ mm h}^{-1}$  (17) showed that the internal reservoir of electron acceptor in *Thioploca* cells is sufficient to reach such depths.

For comparison,  $[\text{NO}_3^-]$  and  $[\text{S}^0]$  concentrations were also measured in the free filaments living at the sediment surface of Station 4 (Fig. 6C). Their  $[\text{NO}_3^-]$  content was  $42 \pm 27 \text{ nmol mm}^{-3}$ , i.e., much lower and less variable than in the 0- to 10-cm-deep section of Station 7 (Fig. 6A). It was also considerably lower than reported for *Beggiatoa* spp. from Monterey Canyon and Guaymas Basin (23). In general,  $[\text{NO}_3^-]$  and  $[\text{S}^0]$  concentrations of the Station 4 filaments were similar to those of *Thioploca* spp. found below 10 cm at Station 7. This was probably due to the high  $\text{H}_2\text{S}$  content (up to  $1,200 \mu\text{M}$ ) in this sediment. Based on the finding of a phobic response to sulfide concentrations of  $>500 \mu\text{M}$  (17) and the high morphological and phylogenetic similarity to *T. araucae*, it was even suggested that these filaments might actually be thioplocas that had moved out of their sheaths or did not produce them under the prevailing environmental conditions (40).

**Intracellular versus extracellular pools of  $\text{S}^0$  and  $\text{NO}_3^-$ .** In sediments of a Danish fjord and in the Santa Barbara basin, it was observed that elemental sulfur was associated primarily with *Beggiatoa* filaments (31, 43). To quantify the contribution of  $[\text{S}^0]$  to the total pool of  $\text{S}^0$  in the sediment, we manually collected all *Thioploca* filaments present in a core from Station 7 and determined the  $[\text{S}^0]$  and  $\text{S}^0$  concentrations in the remaining bulk sediment (Fig. 8). In contrast to the two other studies, we found that  $[\text{S}^0]$  made up maximally 27% of the total  $\text{S}^0$  pool. Furthermore, the two sulfur pools exhibited different distribution patterns. Whereas the  $[\text{S}^0]$  corresponded to the distribution of *Thioploca* spp. (33) and decreased gradually with depth, the  $\text{S}^0$  was maximal at a depth of 5 cm. Because the turnover times of the two sulfur pools may be different, one cannot draw quantitative conclusions about the relative significance of *Thioploca*-associated and chemical sulfide oxidation, but it clearly demonstrates that other processes contribute to sulfide oxidation and sedimentary  $\text{S}^0$  formation. The distribution of chemical species ( $\text{H}_2\text{S}$ ,  $\text{Fe}^{2+}$ , and  $\text{S}_2\text{O}_3^{2-}$ ) in the pore water suggests that reducible iron oxides may also be an important oxidant for pore water  $\text{H}_2\text{S}$  (41; Zopfi et al., submitted).

From the known amount of  $[\text{S}^0]$  and the average  $[\text{NO}_3^-]/[\text{S}^0]$ -ratio in a given depth interval, one can calculate the amount of nitrate being accumulated by *Thioploca* cells and transported into the sediment. If the  $[\text{NO}_3^-]$  were completely released from the cells, it would lead to pore water concentrations of about 1 mM (Fig. 8). There is evidence in the published literature that some measurements of pore water  $\text{NO}_3^-$  were influenced by  $[\text{NO}_3^-]$  released from vacuolated sulfur bacteria during pore water sampling. For example Henrichs and Farrington (14) noted that the  $\text{NO}_3^-$  pore water concentrations in *Thioploca*-containing sediment were higher than in the overlying seawater. In sediments from the Peru upwelling area,  $\text{NO}_3^-$  was found in significant concentrations down to an unusual depth of 10 cm and the maximum concentrations (up to  $102 \mu\text{M}$ ) were conspicuously high (11). In later publications where unusually high nitrate concentrations were observed, it was already suspected that they were affected by  $\text{NO}_3^-$  released from disrupted cells (27, 41).

Over recent years, a variety of approaches have been applied to estimate the contribution of *Thioploca* spp. to sulfide oxi-






FIG. 7. Vertical distribution of the mean [NO<sub>3</sub><sup>-</sup>] and [S<sup>0</sup>] concentrations in the different *Thioploca* spp. from Station 7.

ation. Although the reported values vary from 3 to 91% (8, 10, 25, 41), most estimates fall in the range of 20 to 30%. However, even if *Thioploca* spp. were not the dominant player in sulfide oxidation in these sediments, they are most significant for the sedimentary nitrogen cycling. Organic matter in the sediments off Concepción Bay is almost exclusively degraded via sulfate reduction (equation 1) (41).



If only 25% of the formed sulfide is oxidized by *Thioploca* spp. according to equation 2, this would lead to an 83% increase of the sedimentary ammonium production. Since ammonium is not lost from the environment, in contrast to N<sub>2</sub>, and since nitrogen tends to be the limiting factor for phytoplankton in the marine environment, this form of nutrient regeneration




FIG. 8. Intracellular (solid symbols) and extracellular (open symbols) pools of S<sup>0</sup> and NO<sub>3</sub><sup>-</sup> in sediment of Station 7. The broken line indicates the sediment surface.

could have considerable consequences for the primary productivity in the area.

#### ACKNOWLEDGMENTS

The staff of Dichato, the crew of R/V *Kay Kay*, and all members of the *Thioploca* '97 expedition, V. A. Gallardo, J. G. Kuenen, S. Otte, H. Schulz, B. Strotmann, and A. Teske are thanked for their help and cooperation. A. Rusch is acknowledged for help with the statistical analysis, and T. Ferdelman and two anonymous reviewers are thanked for valuable comments on the manuscript.

This work was supported by the German Ministry for Education, Science, Research and Technology (to J.Z.) and the Max-Planck Society.

#### REFERENCES

1. Beijerinck, M. W. 1904. Phénomènes de réduction produits par les microbes. Arch. Nederl. Sci. Exactes Nat. **9**:131–157.
2. Binnerup, S. J., K. Jensen, N. P. Revsbech, M. H. Jensen, and J. Sørensen. 1992. Denitrification, dissimilatory reduction of nitrate to ammonium, and nitrification in a bioturbated estuarine sediment as measured with <sup>15</sup>N and microsensor techniques. Appl. Environ. Microbiol. **58**:303–313.
3. Brettar, I., and G. Rheinheimer. 1991. Denitrification in the central Baltic: evidence for H<sub>2</sub>S-oxidation as a motor of denitrification at the oxic anoxic interface. Mar. Ecol. Prog. Ser. **77**:157–169.
4. Broecker, W. S., and T. H. Peng. 1974. Gas exchange rates between air and sea. Tellus **26**:21–35.
5. Codispoti, L. A., G. E. Friedrich, T. T. Packard, H. E. Glover, P. J. Kelly, R. W. Spinrad, R. T. Barber, J. W. Elkins, B. B. Ward, F. Lipschultz, and N. Lostaunau. 1986. High nitrite levels off Northern Peru: a signal of instability in the marine denitrification rate. Science **233**:1200–1202.
6. Copin-Montégut, C., and P. Raimbault. 1994. The peruvian upwelling near 15°S in August 1986. Results of continuous measurements of physical and chemical properties between 0 and 220 m depth. Deep-Sea Res. **41**:439–467.
7. de Beer, D., and J.-P. R. A. Sweerts. 1989. Measurement of nitrate gradients with an ion-selective microelectrode. Anal. Chim. Acta. **219**:351–356.
8. Ferdelman, T. G., C. Lee, S. Pantoja, J. Harder, B. Bebout, and H. Fossing. 1997. Sulfate reduction and methanogenesis in a *Thioploca*-dominated sediment off the coast of Chile. Geochim. Cosmochim. Acta. **61**:3065–3079.
9. Forster, S., M. Huettel, and W. Ziebis. 1996. Impact of boundary layer flow velocity on oxygen utilization in coastal sediments. Mar. Ecol. Prog. Ser. **143**:173–185.
10. Fossing, H., V. A. Gallardo, B. B. Jørgensen, M. Huettel, L. P. Nielsen, H. Schulz, D. Canfield, S. Foster, R. N. Glud, J. K. Gundersen, J. Kuever, N. B. Ramsing, A. Teske, B. Thamdrup, and O. Ulloa. 1995. Concentration and transport of nitrate by the mat-forming sulphur bacterium *Thioploca*. Nature **374**:713–715.
11. Froelich, P. N., M. A. Arthur, W. C. Burnett, M. Deakin, V. Hensley, R. Jahnke, L. Kaul, K.-H. Kim, K. Roe, A. Soutar, and C. Vathakanon. 1988. Early diagenesis of organic matter in Peru continental margin sediments: phosphorite precipitation. Mar. Geol. **80**:309–343.
12. Gallardo, V. A. 1977. Large benthic microbial communities in sulphide biota under Peru-Chile subsurface countercurrent. Nature **268**:331–332.
13. Grasshoff, K., M. Erhardt, and K. Kremling. 1983. Methods of sea water analysis. Verlag Chemie, Weinheim, Germany.
14. Henrichs, S. M., and J. W. Farrington. 1984. Peru upwelling region sediments near 15°S. I. Remineralization and accumulation of organic matter. Limnol. Oceanogr. **29**:1–19.
15. Herbert, R. A. 1999. Nitrogen cycling in coastal marine ecosystems. FEMS Microbiol. Rev. **23**:563–590.
16. Hill, A. E., B. M. Hickey, F. A. Shillington, P. T. Strub, K. H. Brink, E. D. Barton, and A. C. Thomas. 1998. Eastern ocean boundaries coastal segment (E), p. 29–67. In A. R. Robinson and K. H. Brink (ed.), The sea, vol. 11. John Wiley & Sons, Inc., New York, N.Y.
17. Huettel, M., S. Forster, S. Klöser, and H. Fossing. 1996. Vertical migration in the sediment dwelling sulfur bacteria *Thioploca* spp. in overcoming diffusion limitations. Appl. Environ. Microbiol. **62**:1863–1872.
18. Knowles, R. 1982. Denitrification. Microbiol. Rev. **46**:43–70.
19. Larsen, L. H., T. Kjær, and N. P. Revsbech. 1997. A microscale NO<sub>3</sub><sup>-</sup> biosensor for environmental applications. Anal. Chem. **69**:3527–3531.
20. Levin, L., J. Gage, P. Lamont, L. Cammidge, C. Martin, A. Patience, and J. Crooks. 1997. Infaunal community structure in a low-oxygen, organic rich habitat on the Oman continental slope, NW Arabian Sea, p. 223–230. In Proceedings of the 30th Marine Biology Symposium.
21. Li, Y. H., and S. Gregory. 1974. Diffusion of ions in sea water and in deep-sea sediments. Geochim. Cosmochim. Acta **38**:703–714.
22. Maier, S., and V. A. Gallardo. 1984. Nutritional characteristics of two marine thioplocas determined by autoradiography. Arch. Microbiol. **139**:218–220.
23. McHatton, S. C., J. P. Barry, H. W. Jannasch, and D. C. Nelson. 1996. High nitrate concentrations in vacuolate, autotrophic marine *Beggiatoa* spp. Appl. Environ. Microbiol. **62**:954–958.
24. Namasaraev, B. B., L. E. Dulov, G. A. Dubinina, T. I. Zemskaya, L. Z. Granina, and E. V. Karabanov. 1994. Bacterial synthesis and destruction of organic matter in microbial mats of Lake Baikal. Microbiology **63**:193–197.
25. Otte, S., J. G. Kuenen, L. P. Nielsen, H. W. Pearl, J. Zopfi, H. N. Schulz, A. Teske, B. Strotmann, V. A. Gallardo, and B. B. Jørgensen. 1999. Nitrogen, carbon and sulfur metabolism in natural *Thioploca* samples. Appl. Environ. Microbiol. **65**:3148–3157.
26. Reimers, C. E. 1982. Organic matter in anoxic sediments off Central Peru: relations of porosity, microbial decomposition and deformation properties. Mar. Geol. **46**:175–197.
27. Reimers, C. E., K. C. Ruttenberg, D. E. Canfield, M. B. Christiansen, and J. B. Martin. 1996. Porewater pH and authigenic phases formed in the uppermost sediments of Santa Barbara Basin. Geochim. Cosmochim. Acta **60**:4037–4057.
28. Revsbech, N. P. 1989. An oxygen microelectrode with a guard cathode. Limnol. Oceanogr. **34**:472–476.
29. Rosenberg, R., W. E. Arntz, E. C. de Flores, L. A. Flores, G. Carabajal, I. Finger, and J. Tarazona. 1983. Benthos biomass and oxygen deficiency in the upwelling system off Peru. J. Mar. Res. **41**:263–279.
30. Sachs, L. 1997. Angewandte Statistik, 8th ed. Springer-Verlag, Berlin, Germany.
31. Schimmelmann, A., and M. Kastner. 1993. Evolutionary changes over the last 1000 years of reduced sulfur phases and organic carbon in varved sediments of the Santa Barbara Basin, California. Geochim. Cosmochim. Acta **57**:67–78.
32. Schramm, A., L. H. Larsen, N. P. Revsbech, N. B. Ramsing, R. Amann, K.-H. Schleifer. 1996. Structure and function of a nitrifying biofilm as determined by in situ hybridization and the use of microelectrodes. Appl. Environ. Microbiol. **62**:4641–4647.
33. Schulz, H. N., B. B. Jørgensen, H. A. Fossing, and N. B. Ramsing. 1996. Community structure of filamentous, sheath-building sulfur bacteria. *Thioploca* spp. off the coast of Chile. Appl. Environ. Microbiol. **62**:1855–1862.
34. Schulz, H. N., T. Brinkhoff, T. G. Ferdelman, M. Hernández Mariné, A. Teske, and B. B. Jørgensen. 1999. Dense populations of a giant sulfur bacterium in Namibian shelf sediments. Science **284**:389–544.
35. Schulz, H. N. 1999. Nitrate-storing sulfur bacteria in sediments of coastal upwelling. Ph.D. thesis. University of Bremen, Bremen, Germany.
36. Schulz, H. N., B. Strotmann, V. A. Gallardo, and B. B. Jørgensen. 2000. Population study of filamentous sulfur bacteria *Thioploca* spp. off the Bay of Concepción, Chile. Mar. Ecol. Prog. Ser. **200**:117–126.
37. Strub, P. T., J. M. Mesias, V. Montecino, J. Rutlland, and J. Salinas. 1998. Coastal ocean circulation off western south America, p. 273–313. In A. R. Robinson and K. H. Brink (ed.), The sea, vol. 11. John Wiley & Sons, Inc., New York, N.Y.
38. Sweerts, J. P. R. A., and D. de Beer. Microelectrode measurements of nitrate gradients in the littoral and profundal sediments of a meso-eutrophic lake (Lake Veichten, The Netherlands). Appl. Environ. Microbiol. **55**:754–757.
39. Taylor, B. F., D. S. Hoare, and S. L. Hoare. 1971. *Thiobacillus denitrificans* as an obligate chemolithotroph. Isolation and growth studies. Arch. Microbiol. **78**:193–204.
40. Teske, A., M. L. Sogin, L. P. Nielsen, and H. W. Jannasch. 1999. Phylogenetic relationships of large marine *Beggiatoa*. Syst. Appl. Microbiol. **22**:39–44.
41. Thamdrup, B., and D. E. Canfield. 1996. Pathways of carbon oxidation in continental margin sediments off central Chile. Limnol. Oceanogr. **41**:1629–1650.
42. Timmer ten Hoor, A. 1975. A new type of thiosulphate oxidizing, nitrate reducing microorganism: *Thiomicrospira denitrificans* sp. nov. Neth. J. Sea Res. **9**:343–351.
43. Troelsen, H., and B. B. Jørgensen. 1982. Seasonal dynamics of elemental sulfur in two coastal sediments. Estuarine Coastal Shelf Sci. **15**:255–266.
44. Zemskaya, T. I., B. B. Namasaraev, N. M. Dultseva, T. A. Khanaeva, L. P. Golobokova, G. A. Dubinina, L. E. Dulov, and E. Wada. 2001. Ecophysiological characteristics of the mat forming bacterium *Thioploca* in bottom sediments of the Frolikha Bay, northern Baikal. Microbiology **70**:335–341.
45. Zumft, W. G., and P. M. H. Kroneck. 1990. Metabolism of nitrous oxide, p. 37–55. In N. P. Revsbech and J. Sørensen (ed.), Denitrification in soil and sediments. Plenum Press, New York, N.Y.