

Quantifying the efficiency and biases of forest *Saccharomyces* sampling strategies

Primrose J. Boynton¹  | Vienna Kowalik² | Doreen Landermann¹ | Eva H. Stukenbrock^{1,3}

¹ Environmental Genomics Research Group, Max-Planck Institute for Evolutionary Biology, Plön, Germany

² Ecology and Evolution Unit, Okinawa Institute of Science and Technology, Okinawa, Japan

³ Botanisches Institut, Christian-Albrechts Universität, Botanisches Institut, Kiel, Germany

Correspondence

Primrose J. Boynton, Environmental Genomics Research Group, Max-Planck Institute for Evolutionary Biology, August-Thienemann-Str. 2, Plön 24306, Germany.
Email: pboynton@evolbio.mpg.de

Funding information

Max Planck Fellowship

Abstract

Saccharomyces yeasts are emerging as model organisms for ecology and evolution, and researchers need environmental *Saccharomyces* isolates to test ecological and evolutionary hypotheses. However, methods for isolating *Saccharomyces* from nature have not been standardized, and isolation methods may influence the genotypes and phenotypes of studied strains. We compared the effectiveness and potential biases of an established enrichment culturing method against a newly developed direct plating method for isolating forest floor *Saccharomyces* spp. In a European forest, enrichment culturing was both less successful at isolating *Saccharomyces paradoxus* per sample collected and less labour intensive per isolated *S. paradoxus* colony than direct isolation. The two methods sampled similar *S. paradoxus* diversity: The number of unique genotypes sampled (i.e., genotypic diversity) per *S. paradoxus* isolate and average growth rates of *S. paradoxus* isolates did not differ between the two methods, and growth rate variances (i.e., phenotypic diversity) only differed in one of three tested environments. However, enrichment culturing did detect rare *Saccharomyces cerevisiae* in the forest habitat and also found two *S. paradoxus* isolates with outlier phenotypes. Our results validate the historically common method of using enrichment culturing to isolate representative collections of environmental *Saccharomyces*. We recommend that researchers choose a *Saccharomyces* sampling method based on resources available for sampling and isolate screening. Researchers interested in discovering new *Saccharomyces* phenotypes or rare *Saccharomyces* species from natural environments may also have more success using enrichment culturing. We include step-by-step sampling protocols in the supplemental materials.

KEYWORDS

enrichment culture, environmental isolates, forest, *Saccharomyces cerevisiae*, *Saccharomyces paradoxus*, soil

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2019 The Authors. *Yeast* published by John Wiley & Sons Ltd


1 | INTRODUCTION

Naturally occurring *Saccharomyces* populations are models for ecology and evolution (Boynton & Greig, 2014). Use of these models has led to exciting discoveries about microbial ecology and evolution; for example, adaptation to climate can lead to speciation (Leducq et al., 2014), domesticated *Saccharomyces cerevisiae* is more phenotypically diverse than wild *Saccharomyces paradoxus* (Warringer et al., 2011), and interspecific hybrids can have high fitnesses in stressful environments (Bernardes, Stelkens, & Greig, 2017; Stelkens, Brockhurst, Hurst, Miller, & Greig, 2014). These studies made inferences based on the phenotypes and genotypes of isolates collected from wild and domesticated substrates. And *Saccharomyces* substrates are diverse: Wild substrates include tree bark, insect guts, fresh leaves, leaf litter, soil, fruits, and parasitic *Cyttaria* galls, (Glushakova, Ivannikova, Naumova, Chernov, & Naumov, 2007; Kowallik & Greig, 2016; Libkind et al., 2011; Mortimer & Polsinelli, 1999; Sampaio & Goncalves, 2008; Stefanini et al., 2012), and domesticated substrates include wine, beer, bread, kimchi, kombucha, palm wine, and pulque, among many other substrates (Boynton & Greig, 2016; Carbonetto, Ramsayer, Nidelet, Legrand, & Sicard, 2018; Estrada-Godina et al., 2001; Ezeronye & Okerentugba, 2001; Gallone et al., 2016; Greenwalt, Steinkraus, & Ledford, 2000; Jeong, Jung, Lee, Jin, & Jeon, 2013). *Saccharomyces* yeasts are also a single clade in the diverse polyphyletic group of yeasts (single-celled fungi that reproduce by budding or fission; Kurtzman, Fell, & Boekhout, 2011). These diverse yeasts inhabit floral nectar, extreme environments, soils, and insect bodies, among many other habitats (Buzzini, Turchetti, & Yurkov, 2018; Chappell & Fukami, 2018; Stefanini, 2018; Yurkov, 2018). One challenge of environmental yeast sampling is to minimize sampling biases so researchers can assure that observed diversity patterns are not artefacts of their chosen sampling strategy.

Enrichment culturing is a reliable and frequently used method for isolating difficult-to-culture bacteria, archaea, and eukaryotic microbes, including *Saccharomyces*, from natural environments (Korzhenkov et al., 2019; Li, Podar, & Morgan-Kiss, 2016; Schlegel & Jannasch, 1967; Sniegowski, Dombrowski, & Fingerhahn, 2002; Figure 1a). Microbiologists have been relying on enrichment cultures for over a century (Beijerinck, 1961) and have used them to isolate many of the model *Saccharomyces* strains commonly used in laboratory studies (Johnson et al., 2004; Liti et al., 2009; Sniegowski et al., 2002). To isolate a microbe using enrichment culturing, a researcher adds a small amount of natural material to a growth medium designed to be hospitable to the target microbe and inhospitable to other microbes (Liti, Warringer, & Blomberg, 2017; Schlegel & Jannasch, 1967). If the enrichment medium is well designed, the target microbe is expected to grow in abundance, and after some incubation time, this enrichment culture can be streaked to a solid medium and colonies of the target microbe can be easily isolated. An alternative to enrichment culturing is to spread a microbial substrate directly onto a selective solid medium, with or without dilution, and to pick colonies that morphologically resemble the target microbe (Glushakova et al., 2007; Stefanini et al., 2012; Figure 1b).

Because it can be difficult to isolate *Saccharomyces* from natural substrates, many investigations of wild *Saccharomyces* rely on enrichment culturing, usually in high-sugar, acidic media (Charron, Leducq, Bertin, Dube, & Landry, 2014; Robinson, Pinharanda, & Bensasson, 2016; Sniegowski et al., 2002; Sweeney, Kuehne, & Sniegowski, 2004). Comparative studies of *Saccharomyces* genomes have been carried out using collections of *Saccharomyces* strains isolated using various strategies, including both enrichment and direct culturing (Liti et al., 2009; Peter et al., 2018). However, isolation strategy can influence the genotypes and phenotypes of isolated microbes: Previous studies have documented higher genotypic diversity among bacteria isolated using direct plating compared with enrichment culturing, and

(a) Enrichment cultures



(b) Direct plating




FIGURE 1 Schematic illustration of sampling strategies used to isolate *Saccharomyces* for this project. (a) Enrichment culturing and (b) direct plating. Photo: Doreen Landermann [Colour figure can be viewed at wileyonlinelibrary.com]

resulting dirty liquid was pipetted on each of two plates containing the solid modified selective medium PIM1 (3-g yeast extract, 5-g peptone, 10-g sucrose, 3-g malt extract, 1-mg chloramphenicol, 80-ml ethanol, 5.2-ml 1 M HCl, and 20-g agar per litre; Kowallik & Greig, 2016; Sniegowski et al., 2002). Liquid was spread on plates using sterile glass beads, and plates were left open in a laminar flow hood until dry. Plates were incubated for 3 days at 30°C before colonies were picked.

For enrichments (Figure 1b), material was mixed with 10 ml of the liquid selective medium PIM1 (composition as for solid PIM1 but without agar) in a 15-ml sterile tube, mixtures were inverted, and tubes were incubated, slightly open and without shaking, at 30°C. After 10 days, a sterile wooden stick was inserted into each enrichment tube and a small amount of the liquid (approximately 50 µl) was streaked onto a single plate containing the solid selective medium PIM2 (20-g methyl-(alpha)-D-glucopyranoside, 1-ml 5% Antifoam Y-30 emulsion, 6.7-g yeast nitrogen base without amino acids, 4-ml 1 M HCl, and 20-g agar per litre; Kowallik & Greig, 2016; Sniegowski et al., 2002), and plates were incubated 4 days at 30°C before colonies were picked.

We include these procedures as step-by-step protocols for the convenience of future researchers in Data S1.

2.2 | Yeast identification

After incubation, we streaked colonies with yeast-like morphology to fresh YPD medium (10-g yeast extract, 20-g peptone, 20-g dextrose, and 25-g agar per litre). For each method, up to six (March and April sampling days) or 12 (June and July sampling days) colonies per sample were selected. After 1 day of growth on YPD at 30°C, cultures were frozen at -80°C in 20% glycerol, and a small amount of each culture was transferred to sporulation medium (20-g potassium acetate, 2.2-g yeast extract, 0.5-g dextrose, 870-mg complete amino acid mixture, and 25-g agar per litre). Any cultures with bacteria-like morphology on YPD medium (slimy culture and/or cells smaller than 1 micron across) were not frozen and were discarded. Sporulation cultures were incubated for at least 3 days at room temperature before being screened under a compound microscope for *Saccharomyces*-like asci (tetrads).

All cultures producing tetrads were identified using sequencing of the internal transcribed sequence (ITS), a region neighbouring rRNA-coding DNA (Schoch et al., 2012). We sequenced every strain using the ITS1/ITS4 primer pair (White, Bruns, Lee, & Taylor, 1990). PCR mixes were 7–15 µl in volume and contained one yeast colony, 0.5-µM each primer, and either 50% Phusion® High-Fidelity PCR master mix with HF buffer or 1× HF-buffer, 100-µM dNTP mix, 3% DMSO, and 1 U/50 µl Phusion DNA polymerase. PCR reactions were cycled at 98°C for 30 s and then 35 cycles of 98°C for 5 s, 62°C for 20 s, and 72°C for 30 s, plus a 10-min terminal extension at 72°C. PCR products were cleaned using illustra™ ExoProStar™ according to the manufacturer's instructions and sequenced on an ABI 3130xl sequencer with BigDye™ Terminator v3.1 chemistry.

ITS sequences were compared with sequences from the type or neotype strains of *S. paradoxus*, *S. cerevisiae*, *Saccharomyces*

kudriavzevii, and *Saccharomyces mikatae* (Genbank accession numbers NR_138272.1, NR_111007.1, KY105195.1, and KY105198.1). If a sequence did not align with *Saccharomyces* sequences, we compared the sequence with all sequences in the NCBI database from type strains using BLAST (Zhang, Schwartz, Wagner, & Miller, 2000). If the sequence aligned with *Saccharomyces* sequences but had more than one base pair different from its closest match, we supplemented ITS sequences with sequences from the gene for translation elongation factor 1 using primers EF1-983F and EF1-2212R (Rehner & Buckley, 2005) using the protocols above, but with a PCR annealing temperature of 57°C. In some cases, cultures originating from apparent single colonies were in fact mixtures of two yeast species. We counted these colonies as *Saccharomyces* if sequences from one of the species was *Saccharomyces*.

2.3 | Growth rates

We compared the distributions of maximum growth rates between the two groups of *S. paradoxus* strains (strains collected using enrichment culturing and strains collected using direct plating) in three liquid media. The media were liquid PIM1, a minimal yeast medium (1.7-g yeast nitrogen base without amino acids and ammonium, 5-g ammonium sulphate, and 2.5-g dextrose per litre), and liquid YPD (composition as for solid YPD, but without agar). To avoid confounding effects of environmental source (i.e., combination of substrate, date collected, and tree), we compared growth rates for pairs of *S. paradoxus* strains originating from the same environmental source. In other words, we collected a data set of *S. paradoxus* growth rates from two groups of strains with equal representations of combinations of substrate, date collected, and tree, and differing only in the method used to isolate the strains. To ensure that all isolates were pure *S. paradoxus* cultures (some cultures that came from what appeared to be single colonies during isolation were found to be mixtures of multiple species after ITS sequencing), we streaked all isolates used for growth rate measurements to single-colony cultures a second time. We confirmed that these single-colony cultures were *S. paradoxus* by mating them with an *S. paradoxus* tester strain (NCYC 3708, α , *ura3::KANMX*, *ho::HYGMX*). In total, 110 isolates (55 from each sampling method) were measured.

Growth rates were measured using an Epoch 2 microplate reader (Biotek Instrument, Inc., Winooski, VT, USA) and calculated using the included Gen5 software version 3.03.14 (Biotek Instrument, Inc.). We first inoculated strains in 0.2 ml of each liquid medium in a 96-well microplate and incubated cultures without shaking or measurement in the microplate reader at 30°C for 24 hr to condition strains to microplate reader conditions. We then transferred 2 µl from each culture to 198-µl fresh medium in a new microplate and incubated the new microplate under the same conditions for 20–24 hr, except for PIM1, in which we grew strains up to 60 hours. OD_{660} was measured during the second incubation every 10 min, and maximum growth rate (MOD_{660}/min) was calculated from the maximum slope of each growth curve over four points (30 min total) using Gen5 software. Reported growth rates for each isolate are means of three replicates.

2.4 | Genotyping

Nine microsatellite loci were identified by searching for common *S. cerevisiae* repeats in the reference genome of *S. paradoxus* strain CBS432 (Liti et al., 2009; Young, Sloan, & Van Riper, 2000) and by adapting previously published *S. cerevisiae* microsatellite loci for *S. paradoxus* (González Techera, Jubany, Carrau, & Gaggero, 2001; Legras, Ruh, Merdinoglu, & Karst, 2005). Primers were designed using Primer3 2.3.4 in Geneious 8.1.8 (Untergasser et al., 2012, <https://www.geneious.com>). Seven microsatellite loci were three-nucleotide repeats; one locus was two-nucleotide repeats; and one locus was four-nucleotide repeats. All loci are described in Table 2. Some loci were complex, including repeats with different sequences; when analysing data, we assumed that alleles of these loci with the same length had the same sequence.

All *S. paradoxus* strains for which growth rates were measured (see above) were genotyped. We amplified microsatellite regions as previously described (Babiker & Tautz, 2015; Hardouin et al., 2015), with slight modifications. Reactions were carried out in 5- μ l PCR mixes containing one colony of each *S. paradoxus* isolate, 2.5- μ l 2 \times Qiagen Multiplex PCR master mix, and 0.2- μ M each primer. Forward primers were labelled with either FAM, HEX, or NED at the five-prime end, and we multiplexed 4–5 primer pairs in each reaction. PCR cycling, dilution, and denaturation were carried out as previously described (Babiker & Tautz, 2015; Hardouin et al., 2015); fragments were run on an ABI 3730 DNA analyser and were analysed using Geneious 8.1.8 with microsatellite plugin version 1.4.4. Genescan ROX-500 (ThermoFisher Scientific) was used as a size standard. All nine microsatellite loci showed variation in the collection of *S. paradoxus* isolates: The lowest number of length polymorphisms detected for any locus was two and the maximum was 10.

2.5 | Statistical analyses

We compared sampling success across substrates (leaf litter or soil) and methods (direct plating or enrichment) using a generalized linear mixed-effects model with probability of isolating *Saccharomyces* (including both *S. paradoxus* and *S. cerevisiae*) as the response variable, substrate and method as fixed effects, and tree and date as random effects. We selected the best model using a top-down strategy, comparing Akaike's information criteria after removing predictors from a full model one by one. Models were calculated using the lme4 package in R version 3.6.0 (Bates, Machler, Bolker, & Walker, 2015; R Development Core Team, 2019).

We compared growth rate distributions by first comparing variances using Levene's test for homogeneity of variance (Levene, 1960) and then comparing medians using paired Wilcoxon signed rank tests. We visualized relationships among genotypes using a neighbour-joining tree of Edwards' genetic distance (Edwards, 1971). Genotypes detected per sample (excluding samples in which only a single isolate was measured) were compared between methods using a paired Wilcoxon signed rank test, and total genotypes isolated were

TABLE 2 Information about microsatellite loci used to genotype *Saccharomyces paradoxus* isolates

Name	Chromosome	Repeat sequence	Fragment length (CBS 432)	Forward primer	Reverse primer	Reference
Chrom16.CTT	16	CTT	223	FAM-CCTCATGGGTTTCGTCGTCT	CGGCTTTGGAAATCCTGGACT	This study
Chrom11.TTG	11	TTG	274	FAM-ATCCTGCTTTGGTCGAAGA	GCTAACTCCGCTACTCACCC	This study
Chrom4.TAAA	4	TAAA	212	FAM-CGGGGTTTTCAATCTTTTGAAAAC	CGGCACCTACCTATTTACCAAGTAAT	Modified from IV-TAAA (Legras et al., 2005)
Chrom4.CAA	4	CAA, CAG	261	HEX-GATCCACCATGGGACCACAA	AGCCATTGACTCTGCTTGTCT	This study
Chrom15.CAA	15	CAA, CAG	295	HEX-TTGGGAATGGCGCTACTTT	CCCGTGAACAGCACCATAT	Modified from XV-CAAB (Legras et al., 2005)
Chrom2.ATT	2	TAT	223	FAM-GGTTAACCTGCCTGTTGAGGA	GCATCTCCGTCCTCCAAACA	Modified from SCAAT2 (González Techera et al., 2001; Legras et al., 2005)
Chrom6.CA	6	AC	184	HEX-GCGGAGGGCTTATCATCTT	CCTCGTATATCCGTCCTCGC	This study
Chrom11.CAA	11	GCA, CAA	284	HEX-GGTGCCTGAAGTGGAAAGGT	GCTCGCTGATGTTGTTCCTG	This study
Chrom2.TAG	2	CTA, CAA, CAG	249	NED-GCCAGGCCAGATAATCAGCA	ACCAGCCTGGATATGAGGGT	This study

compared between methods using the bootstrap method described in Chao et al. (2014) with 50 replications. Statistics were calculated using R version 3.6.0 (R Development Core Team, 2019) and the poppr, ape, car, and iNEXT packages (Fox & Weisberg, 2019; Hsieh, Ma, & Chao, 2019; Kamvar, Tabima, & Grünwald, 2014; Paradis & Schliep, 2018). Graphics were produced using the ggplot2 package and FigTree v.1.4.3 (Wickham, 2016, <http://tree.bio.ed.ac.uk/software/figtree/>).

3 | RESULTS

3.1 | Influence of sampling method on *Saccharomyces* isolation success

Direct plating was more successful than enrichment culturing for isolating *Saccharomyces* spp. from natural substrates ($z = 6.1$, $p < .001$; Tables 3, 4 and Figure 2). We found *Saccharomyces* isolates in 45% of direct plating samples and 19% of enrichment culturing samples. However, enrichment culturing produced the only *S. cerevisiae* found in this study: We found six *S. cerevisiae* isolates from a single enrichment culture from Tree 3 in March of 2017. All other *Saccharomyces* isolates found in this study were *S. paradoxus*. Other detected yeast species included *Saccharomycodes ludwigii*, *Torulaspora delbrueckii*, *Pichia membranifaciens*, and *Hanseniaspora osmophila*, all of which have previously been found alongside *Saccharomyces* spp. in beverage fermentations (Domizio et al., 2011; Gschaedler, 2017).

Although the direct plating method was more successful than the enrichment method, it was also more labour intensive (Table 5). We found more colonies with *S. paradoxus*-like morphology, including colonies that belonged to non-*Saccharomyces* genera, using the direct plating method (969) than using the enrichment method (284), and we screened all of these colonies for tetrad formation. As a result, we screened more than three times as many colonies for tetrads when using the direct plating method than we did using the enrichment method. After screening for tetrads and ITS sequencing, only 32% of the total isolated direct plating colonies were *S. paradoxus*, compared with 74% of enrichment colonies.

Both methods isolated *Saccharomyces* colonies from both substrates, most trees, and all timepoints (Figure 2). We had significantly more sampling success on soil than leaf litter substrates ($z = 5.7$, $p < .001$, Table 4), but other relationships among sampling success, sampling method, and sampling environments were idiosyncratic. For

TABLE 4 Model summary table (Model 4)

Variable	Estimate	Std. error	z	p
Intercept	-2.5324	0.4599	-5.506	<.001
Method (plating)	1.5494	0.2559	6.054	<.001
Substrate (soil)	1.4445	0.2542	5.683	<.001

example, direct plating did not produce any *Saccharomyces* isolates from Tree 6, whereas three enrichment samples from this tree isolated *S. paradoxus*, and enrichments produced more *Saccharomyces* isolates in March than direct plating did (Figure 2). Because our sampling effort was not the same for all trees at all months, we did not model tree habitat or sampling month as fixed effects; instead, we modelled these parameters as random effects and found that models including tree and month fit the data better than models without tree and month (Table 3).

3.2 | Phenotype diversity of sampled *Saccharomyces paradoxus*

Growth rate distributions did not differ between the two methods in PIM1 and the minimal medium and differed slightly in variance in the YPD medium (Figure 3 and Tables 6, 7). Median growth rates did not differ significantly between the two methods in any of the three tested media (Table 6), and variances in growth rate only differed significantly in YPD (Levene's test $F(1, 108) = 5.42$, $p = .022$, Table 7), with enrichment cultures isolating a wider variance of *S. paradoxus* growth rates in YPD than direct plating (Figure 3c). When two outlier strains were removed (Figure 3c), this difference disappeared, $F(1, 106) = 3.59$, $p = .06$.

3.3 | Genotype diversity of sampled *Saccharomyces paradoxus*

The two isolation methods sampled equivalent genotype diversities, both across and within samples. In total, we found 21 unique clonal genotypes (Figure 4). The minimum number of clones per genotype was one and the maximum was 55 (Figure S1). The enrichment method discovered 17 genotypes (95% confidence interval [12.1, 21.9]), and the direct plating method discovered 12 genotypes (95% confidence interval [8.8, 15.2]), but this difference was not significant

TABLE 3 Model selection (mixed-effects generalized linear model)

Model	Fixed effects	Random effects	AIC	Compared with	Better model
1	Method + substrate + method:substrate	1 tree + 1 month	447.20		
2	Method + substrate + method:substrate	1 month	470.94	Model 1	1
3	Method + substrate + method:substrate	1 tree	451.99	Model 1	1
4	Method + substrate	1 tree + 1 month	446.76	Model 1	4
5	Method	1 tree + 1 month	480.34	Model 4	4
6	Substrate	1 tree + 1 month	485.71	Model 4	4

Note. The bolded row indicates the best-fitting model.




FIGURE 2 Percentages of samples in which *Saccharomyces* could be detected using (a) enrichment cultures or (b) direct plating. Bars represent all samples for each category of sampling, and shading represents *Saccharomyces* species. *Saccharomyces paradoxus* and *Saccharomyces cerevisiae* were the only detected *Saccharomyces* species

TABLE 5 Colonies processed and sampling success for each method

Method	Total colonies picked	Sequenced colonies with <i>Saccharomyces</i> -like ascus morphology (%)	<i>Saccharomyces</i> isolates (%)
Enrichment	284	246 (87)	211 (74)
Plating	969	344 (35)	307 (32)

because the two confidence intervals overlap (Figure 4). Enrichment cultures sampled more genotypes per sample (mean = 1.71, samples with only one isolate excluded) than direct plating cultures (mean = 1.57), but this difference was also not significant (Wilcoxon signed rank test paired $V = 27$, $p = .61$). All genotypes except one were homozygous at all loci. The single heterozygous genotype was present in two isolates sampled from leaf litter beneath Tree 7 in June of 2017; the enrichment sampling and direct plating methods each isolated one of the two heterozygous isolates.

4 | DISCUSSION

4.1 | Direct plating detects *Saccharomyces paradoxus* more frequently than enrichment culturing

Enrichment culturing did not increase *Saccharomyces* sampling success per collected forest leaf litter and soil sample compared with direct plating, in spite of researchers' long history of using enrichment

culturing to isolate *Saccharomyces* from forest environments (Kowallik & Greig, 2016; Naumov, Naumova, & Sniegowski, 1998; Sniegowski et al., 2002). We expect reliable *Saccharomyces* isolation from this forest using direct plating to be a result of high *S. paradoxus* abundance on forest floor substrates. Indeed, a previous study determined that hundreds to tens of thousands of *S. paradoxus* cells can occupy a gram of leaf litter near the bases of oak trees in this forest (Kowallik & Greig, 2016). These quantitative observations were made by serially diluting enrichment cultures and estimating the number of *S. paradoxus* cells per gram of leaf litter based on the highest dilution in which *S. paradoxus* could be found. We expect direct plating to be less successful in environments in which *Saccharomyces* are rarer, and note that enrichment culturing is frequently used to isolate *Saccharomyces* from tree bark, which may be a habitat with lower *Saccharomyces* density than the forest floor habitats we sampled (Kowallik et al., 2015; Sniegowski et al., 2002). *S. paradoxus* abundance can also vary over time, with spikes after environmental changes such as rain events (Anderson et al., 2018; Glushakova et al., 2007). It is possible that environmental conditions at other locations, or characteristics of non-European *S. paradoxus* populations, would result in different sampling successes using these two methods from that reported here.

It was not possible to completely standardize quantities of sampled natural material when comparing direct and enrichment-based sampling methods. We collected a larger volume of material for direct cultures (~5 ml) than for enrichment cultures (~2 ml), but the proportion of the original enrichment sample ultimately screened for *Saccharomyces* colonies depends on processes occurring during enrichment. For direct plating, we screened 400 μ l of the 25-ml total suspension of soil




FIGURE 3 Histograms representing distributions of growth rates for *Saccharomyces paradoxus* clones isolated using enrichment culturing and direct plating in (a) PIM1, (b) minimal, and (c) YPD media [Colour figure can be viewed at wileyonlinelibrary.com]

TABLE 6 Median growth rate comparisons between sampling methods

Tested medium	Wilcoxon signed rank test paired V	p
PIM1	818.5	.69
Minimal	911.5	.24
YPD	733	.76

TABLE 7 Variance in growth rate comparisons between sampling methods

Tested medium	F	df	p
PIM1	0.99	1, 108	.32
Minimal	3.17	1, 108	.078
YPD	5.42	1, 108	.022




FIGURE 4 Genotype rarefaction curves of genotypes detected. Thick lines represent average genotypes observed as a function of isolates sampled and shaded areas represent 95% standard errors [Colour figure can be viewed at wileyonlinelibrary.com]

or leaf litter material in water for *Saccharomyces* colonies. For enrichment culturing, we potentially screened all 2 ml of collected material (if *S. paradoxus* cells present in low cell numbers at the start of enrichment culturing grew to high cell numbers during the enrichment incubation) or none of the collected material (if other microbes in enrichments inhibited *S. paradoxus* growth). *S. paradoxus* sampling success in enrichment cultures depends on the composition of the cosampled microbial community, and as a result, we chose to standardize by maximum number of colonies screened (six or 12) instead of by volume of material collected. Researchers adapting our methods could adjust the amount of material collected, the volume of liquid plated, or the number of colonies screened to optimize the methods to their own systems.

4.2 | Both isolation methods sampled similar *Saccharomyces paradoxus* diversities from forest substrates

Overall, enrichment culturing and direct plating collected similar phenotypic and genotypic diversity (Figures 3 and 4). We found no evidence that enrichment culturing selected for fitter individuals in the enrichment medium than direct plating (Figure 3a). Although we genotyped many representatives of the same clonal genotypes (Figure S1), clonal reproduction inside of enrichment cultures did not decrease sampled diversity. High clonality in a local area is common for wild *S. paradoxus* populations and is most likely a result of extensive asexual reproduction in natural habitats (Tsai, Bensasson, Burt, & Koufopanou, 2008; Xia et al., 2017). We also found no evidence for sexual outcrossing in the enrichment cultures

themselves. If outcrossing had occurred during enrichment, we would expect to have seen heterozygous F1 offspring among the genotypes isolated using enrichment. Instead, the only heterozygous genotype in our collection was isolated from a single environment using both enrichment and plating methods and was unlikely to have arisen during enrichment.

The enrichment method did isolate some outlier *S. paradoxus* phenotypes and *S. cerevisiae* that the direct plating method did not (Figures 2 and 3c). We did not find many of these outliers, but we speculate that diverse interactions with microbes in enrichments may have led to isolation of outlier phenotypes and *Saccharomyces* spp. For example, the isolated outlier *Saccharomyces* may have come from enrichments containing bacteria that promoted outlier *S. paradoxus* or *S. cerevisiae* growth at the expense of other *S. paradoxus* genotypes. Microbial diversity across enrichment cultures may similarly explain our idiosyncratic sampling success across months and trees (Figure 2). For example, it is possible that a bacterium that inhibits *S. paradoxus* growth in the enrichment medium was more common in summer than spring months, resulting in lower enrichment sampling success in summer.

4.3 | Recommendations for future yeast sampling

Our results identified a trade-off between resources spent on sampling and resources spent on sequencing: Enrichment culturing was less successful than direct plating at finding *Saccharomyces* per sample collected, but more successful per ITS region sequenced (Figure 2 and Table 5). Researchers with a few precious samples are therefore better off isolating *Saccharomyces* using direct plating, especially if *Saccharomyces* is common on their substrates. Conversely, if samples are easy to get but funds available for sequencing are limited, researchers may prefer to use enrichment culturing or to use direct plating with more phenotypic screening tests than we used. For example, researchers using the direct plating protocol might replica-plate colonies to a second selective medium such as PIM2, which did not increase sampling biases in the enrichment cultures in our sampling, to reduce the number of non-*Saccharomyces* colonies that must be sequenced. Researchers with limited time or freezer space who would like assurances that most picked colonies are *Saccharomyces* may also prefer enrichment culturing or direct plating with additional selective steps.

Although, on average, both methods sampled similar phenotypic and genotypic diversity, our isolation of outlier isolates using enrichments suggests that researchers targeting outliers may also prefer enrichment culturing. For example, researchers sampling environments to find unusual *Saccharomyces* phenotypes for applied biotechnology (e.g., food microbiology and drug discovery) may uncover more diversity using enrichment culturing. Researchers interested in detecting rare *Saccharomyces* species in an environment (e.g., *S. cerevisiae* from our study forest and *S. mikatae* and *Saccharomyces eubayanus* from European forests; Alsammar et al., 2018) may also have more success using enrichment culturing.

5 | CONCLUSIONS

Our results validated use of enrichment culturing for isolating diverse and representative collections of *S. paradoxus* from natural material. We found no evidence that processes during enrichment culturing decrease the diversity of sampled *Saccharomyces* spp. and weak evidence that these processes may in fact increase sampled diversity. Although it is generally a good idea to standardize sampling methods within a study as much as possible, conclusions from studies comparing *Saccharomyces* genotype and phenotype diversity from a variety of sources, including culture collections, are likely to be reliable (Strope et al., 2015; Warringer et al., 2011) and the diversity found in culture collections is likely to be representative of natural *Saccharomyces* diversity in sampled environments. In addition to validating the frequently used enrichment method for isolating *Saccharomyces* spp., this study provides a reliable direct method for isolating *Saccharomyces* spp. and describes a set of microsatellite markers that can be used to conveniently identify *S. paradoxus* genotype diversity. The utility of *Saccharomyces* as an ecology and evolutionary model relies on our understanding of its natural history, and we hope that these and other improvements in field sampling methods will empower researchers to explore the environmental contexts of these exciting microbial model organisms.

ACKNOWLEDGEMENTS

We would like to thank Danielle Stevens and Tjorben Nawroth for help in the field, Jenna Gallie and the Gallie lab for help with growth curves, Michael Habig for advice on statistics, and Amine Hassani for helpful conversations on microbial diversity in enrichment cultures. Thank you to Christoph Freiherr von Fürstenberg-Plessen for permission to work in the Nehnten forest. This work was supported through a Max Planck Fellowship to Eva H. Stukenbrock.

DATA ACCESSIBILITY

All data for this project, including a list of identified yeasts, sampling success data for each soil and leaf litter sample, isolate growth rates, and isolate genotype data, have been deposited in the Edmond repository (doi:10.17617/3.2i).

ORCID

Primrose J. Boynton  <https://orcid.org/0000-0003-0720-8966>

REFERENCES

- Alsammar, H. F., Naseeb, S., Brancia, L. B., Gilman, R. T., Wang, P., & Delneri, D. (2018). Targeted metagenomics approach to capture the biodiversity of *Saccharomyces* genus in wild environments. *Environmental Microbiology Reports.*, 11(2), 206–214. <https://doi.org/10.1111/1758-2229.12724>
- Anderson, J. B., Kasimer, D., Xia, W., Schröder, N. C. H., Cichowicz, P., Lionello, S., ... Kohn, L. M. (2018). Persistence of resident and transplanted genotypes of the undomesticated yeast *Saccharomyces paradoxus* in forest soil. *mSphere*, 3(3), e00211–e00218. <https://doi.org/10.1128/mSphere.00211-18>
- Babiker, H., & Tautz, D. (2015). Molecular and phenotypic distinction of the very recently evolved insular subspecies *Mus musculus hegolandicus* ZIMMERMANN, 1953. *BMC Evolutionary Biology*, 15, 160. <https://doi.org/10.1186/s12862-015-0439-5>
- Bates, D., Machler, M., Bolker, B. M., & Walker, S. C. (2015). Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1), 1–48.
- Beijernick, M. W. (1961). Enrichment culture studies with urea bacteria (T. D. Brock, Trans.). In T. D. Brock (Ed.), *Milestones in microbiology*. London, United Kingdom: Prentice-Hall International.
- Bernardes, J. P., Stelkens, R. B., & Greig, D. (2017). Heterosis in hybrids within and between yeast species. *Journal of Evolutionary Biology*, 30(3), 538–548. <https://doi.org/10.1111/jeb.13023>
- Boynton, P. J., & Greig, D. (2014). The ecology and evolution of non-domesticated *Saccharomyces* species. *Yeast*, 31(12), 449–462. <https://doi.org/10.1002/yea.3040>
- Boynton, P. J., & Greig, D. (2016). Species richness influences wine ecosystem function through a dominant species. *Fungal Ecology*, 22, 61–72. <https://doi.org/10.1016/j.funeco.2016.04.008>
- Buzzini, P., Turchetti, B., & Yurkov, A. (2018). Extremophilic yeasts: The toughest yeasts around? *Yeast*, 35(8), 487–497. <https://doi.org/10.1002/yea.3314>
- Carbonetto, B., Ramsayer, J., Nidelet, T., Legrand, J., & Sicard, D. (2018). Bakery yeasts, a new model for studies in ecology and evolution. *Yeast*, 35(11), 591–603. <https://doi.org/10.1002/yea.3350>
- Chao, A., Gotelli, N. J., Hsieh, T. C., Sander, E. L., Ma, K. H., Colwell, R. K., & Ellison, A. M. (2014). Rarefaction and extrapolation with Hill numbers: A framework for sampling and estimation in species diversity studies. *Ecological Monographs*, 84(1), 45–67. <https://doi.org/10.1890/13-0133.1>
- Chappell, C. R., & Fukami, T. (2018). Nectar yeasts: A natural microcosm for ecology. *Yeast*, 35(6), 417–423. <https://doi.org/10.1002/yea.3311>
- Charron, G., Leducq, J. B., Bertin, C., Dube, A. K., & Landry, C. R. (2014). Exploring the northern limit of the distribution of *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* in North America. *FEMS Yeast Research*, 14(2), 281–288. <https://doi.org/10.1111/1567-1364.12100>
- Domizio, P., Romani, C., Comitini, F., Gobbi, M., Lencioni, L., Mannazzu, I., & Ciani, M. (2011). Potential spoilage non-Saccharomyces yeasts in mixed cultures with *Saccharomyces cerevisiae*. *Annals of Microbiology*, 61(1), 137–144. <https://doi.org/10.1007/s13213-010-0125-1>
- Dunbar, J., White, S., & Forney, L. (1997). Genetic diversity through the looking glass: Effect of enrichment bias. *Applied and Environmental Microbiology*, 63(4), 1326–1331.
- Edwards, A. W. F. (1971). Distance between populations on the basis of gene frequencies. *Biometrics*, 27(4), 873–881. <https://doi.org/10.2307/2528824>
- Estrada-Godina, A. R., Cruz-Guerrero, A. E., Lappe, P., Ulloa, M., Garcia-Garibay, M., & Gomez-Ruiz, L. (2001). Isolation and identification of killer yeasts from Agave sap (aguamiel) and pulque. *World Journal of Microbiology & Biotechnology*, 17(6), 557–560. <https://doi.org/10.1023/a:1012210106203>
- Ezeronye, O. U., & Okerentugba, P. O. (2001). Genetic and physiological variants of yeast selected from palm wine. *Mycopathologia*, 152(2), 85–89. <https://doi.org/10.1023/a:1012323721012>
- Fox, J., & Weisberg, S. (2019). *An {R} companion to applied regression (third edition)*. Thousand Oaks, CA: Sage.
- Gallone, B., Steensels, J., Prahl, T., Soriaga, L., Saels, V., Herrera-Malaver, B., ... Verstrepen, K. J. (2016). Domestication and divergence of *Saccharomyces cerevisiae* beer yeasts. *Cell*, 166(6), 1397–1410. <https://doi.org/10.1016/j.cell.2016.08.020>

- Glushakova, A. M., Ivannikova, Y. V., Naumova, E. S., Chernov, I. Y., & Naumov, E. I. (2007). Massive isolation and identification of *Saccharomyces paradoxus* yeasts from plant phyllosphere. *Microbiology*, 76(2), 205–210. <https://doi.org/10.1134/S0026261707020129>
- González Techera, A., Jubany, S., Carrau, F. M., & Gaggero, C. (2001). Differentiation of industrial wine yeast strains using microsatellite markers. *Letters in Applied Microbiology*, 33(1), 71–75. <https://doi.org/10.1046/j.1472-765X.2001.00946.x>
- Greenwalt, C. J., Steinkraus, K. H., & Ledford, R. A. (2000). Kombucha, the fermented tea: Microbiology, composition, and claimed health effects. *Journal of Food Protection*, 63(7), 976–981. <https://doi.org/10.4315/0362-028x-63.7.976>
- Gschaedler, A. (2017). Contribution of non-conventional yeasts in alcoholic beverages. *Current Opinion in Food Science*, 13, 73–77. <https://doi.org/10.1016/j.cofs.2017.02.004>
- Hardouin, E. A., Orth, A., Teschke, M., Darvish, J., Tautz, D., & Bonhomme, F. (2015). Eurasian house mouse (*Mus musculus* L.) differentiation at microsatellite loci identifies the Iranian plateau as a phylogenetic hotspot. *BMC Evolutionary Biology*, 15, 26. <https://doi.org/10.1186/s12862-015-0306-4>
- Hsieh, T. C., Ma, K. H., & Chao, A. (2019). iNEXT: iNterpolation and EXTrapolation for species diversity. R package version 2.0.19. <http://chao.stat.nthu.edu.tw/blog/software-download/>.
- Jeong, S. H., Jung, J. Y., Lee, S. H., Jin, H. M., & Jeon, C. O. (2013). Microbial succession and metabolite changes during fermentation of dongchimi, traditional Korean watery kimchi. *International Journal of Food Microbiology*, 164(1), 46–53. <https://doi.org/10.1016/j.ijfoodmicro.2013.03.016>
- Johnson, L. J., Koufopanou, V., Goddard, M. R., Hetherington, R., Schafer, S. M., & Burt, A. (2004). Population genetics of the wild yeast *Saccharomyces paradoxus*. *Genetics*, 166(1), 43–52. <https://doi.org/10.1534/genetics.166.1.43>
- Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: An R package for genetic analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ*, 2, e281. <https://doi.org/10.7717/peerj.281>
- Korzhenkov, A. A., Toshchakov, S. V., Bargiela, R., Gibbard, H., Ferrer, M., Teplyuk, A. V., ... Golyshina, O. V. (2019). Archaea dominate the microbial community in an ecosystem with low-to-moderate temperature and extreme acidity. *Microbiome*, 7(11), 11. <https://doi.org/10.1186/s40168-019-0623-8>
- Kowallik, V. (2015). The natural ecology of *Saccharomyces* yeasts. (Dr. rer. nat.), Christian Albrechts University of Kiel.
- Kowallik, V., & Greig, D. (2016). A systematic forest survey showing an association of *Saccharomyces paradoxus* with oak leaf litter. *Environmental Microbiology Reports*, 8(5), 833–841. <https://doi.org/10.1111/1758-2229.12446>
- Kowallik, V., Miller, E., & Greig, D. (2015). The interaction of *Saccharomyces paradoxus* with its natural competitors on oak bark. *Molecular Ecology*, 24(7), 1596–1610. <https://doi.org/10.1111/mec.13120>
- Kurtzman, C. P., Fell, J. W., & Boekhout, T. (2011). Definition, classification and nomenclature of the yeasts. In C. P. Kurtzman, J. W. Fell, & T. Boekhout (Eds.), *The yeasts, a taxonomic study* (Vol. 1) (pp. 87–110). London: Elsevier.
- Leducq, J. B., Charron, G., Samani, P., Dube, A. K., Sylvester, K., James, B., ... Landry, C. R. (2014). Local climatic adaptation in a widespread microorganism. *Proceedings of the Royal Society B-Biological Sciences*, 281(1777), 20132472. <https://doi.org/10.1098/rspb.2013.2472>
- Legras, J., Ruh, O., Merdinoglu, D., & Karst, F. (2005). Selection of hyper-variable microsatellite loci for the characterization of *Saccharomyces cerevisiae* strains. *International Journal of Food Microbiology*, 102(1), 73–83. <https://doi.org/10.1016/j.ijfoodmicro.2004.12.007>
- Levene, H. (1960). Robust tests for equality of variances. In I. Olkin, & H. Hotelling (Eds.), *Contributions to probability and statistics: Essays in honor of Harold Hotelling* (pp. 278–292). Palo Alto: Stanford University Press.
- Li, W., Podar, M., & Morgan-Kiss, R. M. (2016). Ultrastructural and single-cell-level characterization reveals metabolic diversity in a microbial eukaryote community from an ice-covered Antarctic lake. *Applied and Environmental Microbiology*, 82(12), 3659–3670. <https://doi.org/10.1128/AEM.00478-16>
- Libkind, D., Hittinger, C. T., Valerio, E., Goncalves, C., Dover, J., Johnston, M., ... Sampaio, J. P. (2011). Microbe domestication and the identification of the wild genetic stock of lager-brewing yeast. *Proceedings of the National Academy of Sciences of the United States of America*, 108(35), 14539–14544. <https://doi.org/10.1073/pnas.1105430108>
- Liti, G., Carter, D. M., Moses, A. M., Warringer, J., Parts, L., James, S. A., ... Louis, E. J. (2009). Population genomics of domestic and wild yeasts. *Nature*, 458(7236), 337–341. <https://doi.org/10.1038/nature07743>
- Liti, G., Warringer, J., & Blomberg, A. (2017). Isolation and laboratory domestication of natural yeast strains. *Cold Spring Harb Protoc*, 2017(8), pdb.prot089052. <https://doi.org/10.1101/pdb.prot089052>
- Mortimer, R., & Polsinelli, M. (1999). On the origins of wine yeast. *Research in Microbiology*, 150(3), 199–204. [https://doi.org/10.1016/s0923-2508\(99\)80036-9](https://doi.org/10.1016/s0923-2508(99)80036-9)
- Naumov, G. I., Naumova, E. S., & Sniegowski, P. D. (1998). *Saccharomyces paradoxus* and *Saccharomyces cerevisiae* are associated with exudates of North American oaks. *Canadian Journal of Microbiology*, 44(11), 1045–1050. <https://doi.org/10.1139/cjm-44-11-1045>
- Oda, Y., Larimer, F. W., Chain, P. S. G., Malfatti, S., Shin, M. V., Vergez, L. M., ... Harwood, C. S. (2008). Multiple genome sequences reveal adaptations of a phototrophic bacterium to sediment microenvironments. *Proceedings of the National Academy of Sciences of the United States of America*, 105(47), 18543–18548. <https://doi.org/10.1073/pnas.0809160105>
- Paradis, E., & Schliep, K. (2018). ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, 35(3), 526–528.
- Peter, J., De Chiara, M., Friedrich, A., Yue, J. X., Pflieger, D., Bergstrom, A., ... Schacherer, J. (2018). Genome evolution across 1,011 *Saccharomyces cerevisiae* isolates. *Nature*, 556(7701), 339–344. <https://doi.org/10.1038/s41586-018-0030-5>
- R Development Core Team (2019). *R: A language and environment for statistical computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Rehner, S. A., & Buckley, E. (2005). A *Beauveria* phylogeny inferred from nuclear ITS and EF1-alpha sequences: Evidence for cryptic diversification and links to *Cordyceps teleomorphs*. *Mycologia*, 97(1), 84–98. <https://doi.org/10.3852/mycologia.97.1.84>
- Robinson, H. A., Pinharanda, A., & Bensasson, D. (2016). Summer temperature can predict the distribution of wild yeast populations. *Ecology and Evolution*, 6(4), 1236–1250. <https://doi.org/10.1002/ece3.1919>
- Sampaio, J. P., & Goncalves, P. (2008). Natural populations of *Saccharomyces kudriavzevii* in Portugal are associated with oak bark and are sympatric with *S. cerevisiae* and *S. paradoxus*. *Applied and Environmental Microbiology*, 74(7), 2144–2152. <https://doi.org/10.1128/aem.02396-07>
- Schlegel, H. G., & Jannasch, H. W. (1967). Enrichment cultures. *Annual Reviews Microbiology*, 21, 49–70. <https://doi.org/10.1146/annurev.mi.21.100167.000405>
- Schoch, C. L., Seifert, K. A., Huhndorf, S., Robert, V., Spouge, J. L., Levesque, C. A., ... Fungal Barcoding, C. (2012). Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode marker for Fungi. *Proceedings of the National Academy of Sciences of*

- the United States of America, 109(16), 6241–6246. <https://doi.org/10.1073/pnas.1117018109>
- Sniegowski, P. D., Dombrowski, P. G., & Fingerhahn, E. (2002). *Saccharomyces cerevisiae* and *Saccharomyces paradoxus* coexist in a natural woodland site in North America and display different levels of reproductive isolation from European conspecifics. *FEMS Yeast Research*, 1(4), 299–306. <https://doi.org/10.1111/j.1567-1364.2002.tb00048.x>
- Stefanini, I. (2018). Yeast-insect associations: It takes guts. *Yeast*, 35(4), 315–330. <https://doi.org/10.1002/yea.3309>
- Stefanini, I., Dapporto, L., Legras, J. L., Calabretta, A., Di Paola, M., De Filippo, C., ... Cavalieri, D. (2012). Role of social wasps in *Saccharomyces cerevisiae* ecology and evolution. *Proceedings of the National Academy of Sciences of the United States of America*, 109(33), 13398–13403. <https://doi.org/10.1073/pnas.1208362109>
- Stelkens, R. B., Brockhurst, M. A., Hurst, G. D. D., Miller, E. L., & Greig, D. (2014). The effect of hybrid transgression on environmental tolerance in experimental yeast crosses. *Journal of Evolutionary Biology*, 27(11), 2507–2519. <https://doi.org/10.1111/jeb.12494>
- Strope, P. K., Skelly, D. A., Kozmin, S. G., Mahadevan, G., Stone, E. A., Magwene, P. M., ... McCusker, J. H. (2015). The 100-genomes strains, an *S. cerevisiae* resource that illuminates its natural phenotypic and genotypic variation and emergence as an opportunistic pathogen. *Genome Research*, 25(5), 762–774. <https://doi.org/10.1101/gr.185538.114>
- Sweeney, J. Y., Kuehne, H. A., & Sniegowski, P. D. (2004). Sympatric natural *Saccharomyces cerevisiae* and *S. paradoxus* populations have different thermal growth profiles. *FEMS Yeast Research*, 4(4–5), 521–525. [https://doi.org/10.1016/s1567-1356\(03\)00171-5](https://doi.org/10.1016/s1567-1356(03)00171-5)
- Tsai, I. J., Bensasson, D., Burt, A., & Koufopanou, V. (2008). Population genomics of the wild yeast *Saccharomyces paradoxus*: Quantifying the life cycle. *Proceedings of the National Academy of Sciences of the United States of America*, 105(12), 4957–4962. <https://doi.org/10.1073/pnas.0707314105>
- Untergasser, A., Cutcutache, I., Koressaar, T., Ye, J., Faircloth, B. C., Remm, M., & Rozen, S. G. (2012). Primer3-new capabilities and interfaces. *Nucleic Acids Research*, 40(15), e115. <https://doi.org/10.1093/nar/gks596>
- Warringer, J., Zorgo, E., Cubillos, F. A., Zia, A., Gjuvsland, A., Simpson, J. T., ... Blomberg, A. (2011). Trait variation in yeast is defined by population history. *PLoS Genetics*, 7(6), 30. <https://doi.org/10.1371/journal.pgen.1002111>
- White, T. J., Bruns, T., Lee, S., & Taylor, J. (1990). Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In M. A. Innis, D. H. Gelfand, J. J. Sninsky, & T. J. White (Eds.), *PCR protocols: A guide to methods and applications* (pp. 315–322). San Diego: Academic Press.
- Wickham, H. (2016). *ggplot2: Elegant graphics for data analysis*. New York: Springer.
- Xia, W., Nielly-Thibault, L., Charron, G., Landry, C. R., Kasimer, D., Anderson, J. B., & Kohn, L. M. (2017). Population genomics reveals the structure at the individual, host-tree scale and persistence of genotypic variants of the undomesticated yeast *Saccharomyces paradoxus* in a natural woodland. *Molecular Ecology*, 26(4), 995–1007. <https://doi.org/10.1111/mec.13954>
- Young, E. T., Sloan, J. S., & Van Riper, K. (2000). Trinucleotide repeats are clustered in regulatory genes in *Saccharomyces cerevisiae*. *Genetics*, 154(3), 1053–1068.
- Yurkov, A. M. (2018). Yeasts of the soil—Obscure but precious. *Yeast*, 35(5), 369–378. <https://doi.org/10.1002/yea.3310>
- Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning DNA sequences. *Journal of Computational Biology*, 7(1–2), 203–214. <https://doi.org/10.1089/10665270050081478>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Boynton PJ, Kowallik V, Landermann D, Stukenbrock EH. Quantifying the efficiency and biases of forest *Saccharomyces* sampling strategies. *Yeast*. 2019;1–12. <https://doi.org/10.1002/yea.3435>