

GENERALIZED ADJOINT ACTIONS

ARKADY BERENSTEIN AND VLADIMIR RETAKH

ABSTRACT. The aim of this paper is to generalize the classical formula $e^x y e^{-x} = \sum_{k \geq 0} \frac{1}{k!} (ad\ x)^k(y)$ by replacing e^x with any formal power series $f(x) = 1 + \sum_{k \geq 1} a_k x^k$. We also obtain combinatorial applications to q -exponentials, q -binomials, and Hall-Littlewood polynomials.

1. NOTATION AND MAIN RESULTS

One of the most fundamental tools in Lie theory, the adjoint action of Lie groups on their Lie algebras, is based on the following formula:

$$(1.1) \quad e^x y e^{-x} = e^{ad\ x}(y) = \sum_{k \geq 0} \frac{1}{k!} (ad\ x)^k(y),$$

where $(ad\ x)^k(y) = [x, [x, \dots, [x, y], \dots]]$ and $[a, b] = ab - ba$.

The aim of this paper is to generalize (1.1) by replacing e^t with any formal power series

$$(1.2) \quad f = f(t) = 1 + \sum_{k \geq 1} a_k t^k$$

over a field \mathbb{k} .

For any formal power series (1.2) over \mathbb{k} define polynomials

$$P_k(t) = P_{f,k}(t) = (-1)^k \det \begin{pmatrix} 1 & a_1 t & a_2 t^2 & \dots & a_k t^k \\ 1 & a_1 & a_2 & \dots & a_k \\ 0 & 1 & a_1 & \dots & a_{k-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & a_1 \end{pmatrix}$$

for $k = 0, 1, 2, \dots$ (with the convention that $P_0(t) = 1$). Clearly, $P_k(1) = 0$ for $k \geq 1$. Using Cramer's rule with respect to the last column, one obtains a recursion $P_k(t) = a_k t^k - \sum_{i=1}^k a_i P_{k-i}(t)$.

The following result is, probably, well-known (for readers' convenience, we prove it in Section 2).

Theorem 1.1. *For any power series $f(t)$ as in (1.2), one has*

$$(1.3) \quad \frac{f(tx)}{f(x)} = \sum_{k \geq 0} P_{f,k}(t) \cdot x^k$$

and

$$(1.4) \quad P_{f,k}(st) = \sum_{i=0}^k P_{f,i}(s) P_{f,k-i}(t) t^i$$

for all $k \geq 0$.

This work was partially supported by the NSF grant DMS-1403527 (A. B.) and by the NSA grant H98230-14-1-0148 (V. R.).

Furthermore, for any algebra \mathcal{A} over \mathbb{k} , a subset $\mathbf{q} = \{q_1, \dots, q_k\} \subset \mathbb{k}$, $x, y \in \mathcal{A}$, and $k \geq 1$ define

$$(ad\ x)^{\mathbf{q}}(y) = [x, [x, \dots, [x, y]_{q_1}, \dots]_{q_{k-1}}]_{q_k}$$

where $[a, b]_q := ab - qba$. It is easy to see that

$$(1.5) \quad (ad\ x)^{\mathbf{q}}(y) = \sum_{j=0}^k (-1)^j e_j(q_1, \dots, q_k) \cdot x^{k-j} y x^j ,$$

where $e_j(q_1, \dots, q_k)$ is the j -th elementary symmetric function.

Theorem 1.2. *Let \mathcal{A} be \mathbb{k} -algebra and suppose that f is any power series (1.2) with $a_k \neq 0$ for $k \geq 1$. Then*

$$(1.6) \quad f(x) y f(x)^{-1} = y + \sum_{k \geq 1} a_k (ad\ x)^{\mathbf{q}_k}(y) ,$$

for any $x, y \in \mathcal{A}$, where $\mathbf{q}_k = \{q_{1k}, \dots, q_{kk}\}$ is the set of roots of $P_{f,k}(t)$.

Remark 1.3. A formula for $f(x) y f(x)^{-1}$ without assumption that all $a_k \neq 0$ is given in Proposition 2.3.

Remark 1.4. Strictly speaking, the formula (1.6), similarly to (1.1) requires a completion of \mathcal{A} . One can bypass this by replacing x with $\tau \cdot x$ where τ is a purely transcendental element of \mathbb{k} so that the right hand side of (1.6) becomes a power series in τ (and, maybe extending \mathbb{k} if it lack such an element).

Remark 1.5. The subsets \mathbf{q}_k may belong to an extension of \mathbb{k} , however, the operators $(ad\ x)^{\mathbf{q}_k}$ are defined over \mathbb{k} due to (1.5) because all symmetric functions in \mathbf{q}_k belong to \mathbb{k} .

It is easy to see that if $a_k = \frac{1}{k!}$ for all k , then $P_k(t) = \frac{(t-1)^k}{k!}$ which immediately recovers (1.1). Suppose now that $a_k = \frac{1}{[k]_q!}$ for all k , where $k_q! = [1]_q \cdots [k]_q$ is the q -factorial and $[\ell]_q = 1 + q + \cdots + q^{\ell-1}$. We will show (Proposition 2.5) that $P_{f,k}(t) = \frac{(t-1)(t-q) \cdots (t-q^{k-1})}{[k]_q!}$ for $f(t) = e_q^t = \sum_{k \geq 0} \frac{t^k}{[k]_q!}$, therefore, recover the following famous result (see e.g., [3]).

Theorem 1.6. *Let $e_q^x = \sum_{k \geq 0} \frac{x^k}{[k]_q!}$ be the q -exponential. Then $e_q^x \cdot y \cdot (e_q^x)^{-1} = \sum_{k \geq 0} \frac{1}{[k]_q!} (ad\ x)^{\{1, q, \dots, q^{k-1}\}}(y)$.*

On the other hand, combining Theorem 1.1 and Proposition 2.5, we recover the following well-known properties of q -exponentials and q -binomials:

$$e_q^{q^n x} = e_q^x \left(1 + \sum_{k=1}^n \frac{(q^n - 1)(q^n - q) \cdots (q^n - q^{k-1})}{[k]_q!} x^k \right)$$

for $n \geq 0$, in particular,

$$e_q^{q^n x} = e_q^x \cdot (1 + (q-1)x)$$

and

$$1 + \sum_{k=1}^n \frac{(q^n - 1)(q^n - q) \cdots (q^n - q^{k-1})}{[k]_q!} x^k = \prod_{i=1}^n (1 + (q-1)q^{i-1}x) .$$

We conclude with a curious observation that the polynomials $P_{f,k}(t)$ are related to the Hall-Littlewood symmetric polynomials.

Proposition 1.7. *Suppose that $f(t) = \prod_{k \geq 1} (1 - x_k t)$. Then*

$$P_{f,k}(t) = Q_{(k)}(\mathbf{x}; t)$$

for all $k \geq 0$, where $\mathbf{x} = \{x_k, k \geq 0\}$ is viewed as an infinite set of variables, $Q_\lambda(\mathbf{x}; t)$ is Hall-Littlewood polynomial ([2, Section 3.2]), and (k) is a one-row Young diagram with k cells. In particular,

$$Q_{(k)}(\mathbf{x}; t) = (-1)^k \det \begin{pmatrix} 1 & -e_1 t & e_2 t^2 & \dots & (-1)^k e_k t^k \\ 1 & -e_1 & e_2 & \dots & (-1)^k e_k \\ 0 & 1 & -e_1 & \dots & (-1)^{k-1} e_{k-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & -e_1 \end{pmatrix}$$

for all $k \geq 0$, where $e_k = e_k(\mathbf{x})$ is the k -th elementary symmetric function.

Acknowledgments. We gratefully acknowledge the support of Centre de Recerca Matemàtica, Barcelona, where this work was accomplished. We are grateful to Eric Rains for pointing out the connection with Hall-Littlewood polynomials.

2. PROOFS

Proof of Theorem 1.1. We need the following well-known fact (attributed to Wronski, see e.g., [1]).

Lemma 2.1. *Let f be any formal power series (1.2). Then $\frac{1}{f(t)} = 1 + \sum_{k \geq 1} D_k(f) t^k$, where*

$$D_k(f) = (-1)^k \det \begin{pmatrix} a_1 & a_2 & a_3 & \dots & a_k \\ 1 & a_1 & a_2 & \dots & a_{k-1} \\ 0 & 1 & a_1 & \dots & a_{k-2} \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & a_1 \end{pmatrix}$$

(with the convention $D_0(f) = 1$).

The following generalization of Lemma 2.1 is, apparently, well-known (for readers' convenience we prove it here).

Lemma 2.2. *Let $f(t) = 1 + \sum_{k \geq 1} a_k t^k$, $g(t) = 1 + \sum_{k \geq 1} b_k t^k$ be formal power series. Then*

$$\frac{g(t)}{f(t)} = \sum_{k \geq 0} D_k(g, f) t^k,$$

where $D_k(g, f) = (-1)^k \det \begin{pmatrix} 1 & b_1 & \dots & b_{k-1} & b_k \\ 1 & a_1 & \dots & a_{k-1} & a_k \\ 0 & 1 & \dots & a_{k-2} & a_{k-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & a_1 \end{pmatrix}$ (with the convention $D_0(g, f) = 1$).

Proof. Indeed, using Lemma 2.1, we obtain (with the convention $b_0 = 1$):

$$\frac{g(t)}{f(t)} = g(t) \cdot \frac{1}{f(t)} = \left(\sum_{i \geq 0} b_i t^i \right) \left(\sum_{j \geq 0} D_j(f) t^j \right) = \sum_{k \geq 0} d_k t^k$$

where

$$d_k = \sum_{i=0}^k b_i D_{k-i}(f) = (-1)^k \det \begin{pmatrix} b_0 & b_1 & \dots & b_{k-1} & b_k \\ 1 & a_1 & \dots & a_{k-1} & a_k \\ 0 & 1 & \dots & a_{k-2} & a_{k-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & a_1 \end{pmatrix}$$

by Cramer rule because $b_i D_{k-i}(f) = (-1)^k \det \begin{pmatrix} 0 & 0 & \dots & b_i & \dots & 0 \\ 1 & a_1 & \dots & a_i & \dots & a_k \\ 0 & 1 & \dots & a_{i+1} & \dots & a_{k-1} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots \\ 0 & 0 & \dots & 1 & a_1 & a_2 \\ 0 & 0 & \dots & \dots & 1 & a_1 \end{pmatrix}$.

The lemma is proved. \square

Then taking $b_k = a_k t^k$ for $k \geq 1$ in Lemma 2.2, we obtain (1.3).

To prove (1.4), compute $\frac{f(stx)}{f(x)}$ in two ways, using the first assertion:

$$\frac{f(stx)}{f(x)} = \sum_{k \geq 0} P_{f,k}(st) \cdot x^k$$

and $\frac{f(stx)}{f(x)} = \frac{f(stx)}{f(tx)} \cdot \frac{f(tx)}{f(x)} = \left(\sum_{i \geq 0} P_{f,i}(s) \cdot (tx)^i \right) \left(\sum_{j \geq 0} P_{f,j}(t) \cdot x^j \right)$. Comparing the coefficients of x^k in both series, we obtain (1.4).

Theorem 1.1 is proved. \square

Proof of Theorem 1.2. We need the following result.

Proposition 2.3. *For any power series f as in (1.2) one has:*

$$(a) P_{f,k}(t) = \sum_{j=0}^k a_{k-j} D_j(f) \cdot t^j \text{ for all } k \geq 0.$$

$$(b) f(x)yf(x)^{-1} = y + z_1 + z_2 + \dots, \text{ where } z_k = \sum_{i=0}^k a_i D_{k-i}(f) \cdot x^i y x^{k-i} \text{ for all } k \geq 1.$$

Proof. Prove (a). Indeed, using Lemma 2.1, we obtain:

$$\frac{f(tx)}{f(x)} = \sum_{i,j \geq 0} (a_i t^i x^i) (D_j(f) x^j) = \sum_{k \geq 0} \left(\sum_{i=0}^k a_i D_{k-i}(f) \cdot t^i \right).$$

This together with Theorem 1.1 proves (a).

Prove (b) now. Indeed,

$$f(x)yf(x)^{-1} = \sum_{i,j \geq 0} (a_i x^i) y (D_j(f) x^j) = \sum_{k \geq 0} \left(\sum_{j=0}^k a_i D_{k-i}(f) \cdot x^i y x^{k-i} \right)$$

This proves (b). \square

Now we can finish the proof of Theorem 1.2. Indeed, suppose that $P_{f,k}(t)$ is factored as

$$P_{f,k}(t) = a_k (t - q_{1k}) \cdots (t - q_{kk}).$$

Then, by Proposition 2.3(a), $a_i D_{k-i} = a_k (-1)^{k-i} e_{k-i}(q_{1k}, \dots, q_{kk})$ for $i = 0, \dots, k$. Therefore, in the notation of Proposition 2.3(b),

$$z_k = \sum_{i=0}^k a_k (-1)^{k-i} e_{k-i}(q_{1k}, \dots, q_{kk}) \cdot x^i y x^{k-i} = a_k (ad x)^{\mathfrak{q}_k}(y)$$

for all $k \geq 1$, which together with Proposition 2.3(b) verifies (1.6).

Theorem 1.2 is proved. \square

Proposition 2.4. $P_{e_q^t, k}(t) = \frac{(t-1)(t-q) \cdots (t-q^{k-1})}{[k]_q!}$ for all $k \geq 1$.

Proof. It suffices to show that $P_{e_q^t, k}(q^a) = 0$ for all $0 \leq a < k$. We proceed by induction in such pairs (a, k) . If $a = 0$, then we have nothing to prove since $P_{f, k}(1) = 0$ for all f .

Using Theorem 1.1, we obtain:

$$P_{f, k}(q^a) = \sum_{i=0}^k P_{f, i}(q^b) P_{f, k-i}(q^{a-b}) q^{(a-b)i} .$$

Taking $f = e_q^t$, $1 \leq b \leq a < k$, and using the inductive hypothesis, this gives $P_{f, k}(q^a) = 0$ for any $1 \leq a < k$.

The proposition is proved. \square

Corollary 2.5. For all $k \geq 1$ one has: $\det \begin{pmatrix} 1 & \frac{t}{[1]_q} & \frac{t^2}{[2]_q} & \cdots & \frac{t^k}{[k]_q} \\ 1 & \frac{1}{[1]_q} & \frac{1}{[2]_q} & \cdots & \frac{1}{[k]_q} \\ 0 & 1 & \frac{1}{[1]_q} & \cdots & \frac{1}{[k-1]_q} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ 0 & 0 & \cdots & 1 & \frac{1}{[1]_q} \end{pmatrix} = \frac{(1-t)(q-t) \cdots (q^{k-1}-t)}{[k]_q!} .$

Proof of Proposition 1.7. Indeed, if $f(t)$ is as in Proposition 1.7, then

$$\frac{f(tu)}{f(u)} = \prod_{k \geq 1} \frac{1 - x_k t u}{1 - x_k u} = \sum_{k \geq 0} Q_{(k)}(\mathbf{x}; t) u^k$$

by [2, Equations (2.10) and (2.13)]. This and Theorem 1.1 imply that $P_{f, k} = Q_{(k)}(\mathbf{x}; t)$ for all $k \geq 0$, which proves the first assertion of Proposition 1.7.

To prove the second assertion, note that $a_k = (-1)^k e_k(\mathbf{x})$ for all $k \geq 0$ because of the well-known formula (see e.g., [2, Section 1.2]):

$$\prod_{k \geq 1} (1 - x_k t) = \sum_{k \geq 0} (-1)^k e_k(\mathbf{x}) t^k .$$

This and the first assertion of Proposition 1.7 imply the second assertion of the proposition. \square

REFERENCES

- [1] A. Inselberg, On determinants of Toeplitz-Hessenberg matrices arising in power series, *J. Math. Anal. Appl.* **63** (1978), no. 2, 347–353.
- [2] I. Macdonald, *Symmetric functions and Hall polynomials* Second edition. With contributions by A. Zelevinsky. Oxford Mathematical Monographs. Oxford Science Publications. The Clarendon Press, Oxford University Press, New York, 1995.
- [3] A. Volkov, Beyond the “pentagon identity,” *Lett. Math. Phys.* **39** (1997), no. 4, 393–397.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403, USA
E-mail address: arkadiy@math.uoregon.edu

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, PISCATAWAY, NJ 08854, USA
E-mail address: vretakh@math.rutgers.edu