IRREGULAR PRIMES WITH RESPECT TO GENOCCHI NUMBERS AND ARTIN'S PRIMITIVE ROOT CONJECTURE

SU HU, MIN-SOO KIM, PIETER MOREE, AND MIN SHA

Dedicated to the memory of Prof. Christopher Hooley (1928-2018)

ABSTRACT. We introduce and study a variant of Kummer's notion of (ir)regularity of primes which we call G-(ir)regularity and is based on Genocchi rather than Bernoulli numbers. We say that an odd prime p is G-irregular if it divides at least one of the Genocchi numbers $G_2, G_4, \dots, G_{p-3}$, and G-regular otherwise. We show that, as in Kummer's case, G-irregularity is related to the divisibility of some class number. Furthermore, we obtain some results on the distribution of G-irregular primes. In particular, we show that each primitive residue class contains infinitely many G-irregular primes and establish non-trivial lower bounds for their number up to a given bound x as x tends to infinity. As a byproduct, we obtain some results on the distribution of primes in arithmetic progressions with a prescribed near-primitive root.

1. INTRODUCTION

1.1. The classical case. The n -th Bernoulli polynomial $B_n(x)$ is implicitly defined as the coefficient of t^n in the generating function

$$\frac{te^{xt}}{e^t - 1} = \sum_{n=0}^{\infty} B_n(x) \frac{t^n}{n!},$$

where e is the base of the natural logarithm. For $n \geq 0$ the Bernoulli numbers B_n are defined by $B_n = B_n(0)$. It is well-known that $B_0 = 1$, and $B_n = 0$ for every odd integer $n > 1$.

In this paper, p always denotes a prime. An odd prime p is said to be *B-irregular* if p divides the numerator of at least one of the Bernoulli numbers $B_2, B_4, \dots, B_{p-3}$, and *B-regular* otherwise. The first twenty B-irregular primes are

$$37, 59, 67, 101, 103, 131, 149, 157, 233, 257, 263, \\ 271, 283, 293, 307, 311, 347, 353, 379, 389.$$

The notion of B-irregularity has an important application in algebraic number theory. Let $\mathbb{Q}(\zeta_p)$ be the p -th cyclotomic field, and h_p its class number. Denote by h_p^+ the class number of $\mathbb{Q}(\zeta_p + \zeta_p^{-1})$ and put $h_p^- = h_p/h_p^+$. Kummer proved that h_p^- is an integer (now called the *relative class number* of $\mathbb{Q}(\zeta_p)$) and gave the following characterization (see [34, Theorem 5.16]). (Here and in the sequel, for any integer $n \geq 1$, ζ_n denotes an n -th primitive root of unity.)

2010 *Mathematics Subject Classification.* 11B68, 11A07, 11R29.

Key words and phrases. Irregular prime, Bernoulli number and polynomial, Euler number and polynomial, Genocchi number, refined class number, primitive root, Artin's primitive root conjecture.

Theorem 1.1 (Kummer). *An odd prime p is B-irregular if and only if $p \mid h_p^-$.*

Kummer showed that if $p \nmid h_p^-$, then the Diophantine equation $x^p + y^p = z^p$ does not have an integer solution x, y, z with p coprime to xyz , cf. [34, Chapter 1].

We remark that for any odd prime p , $p \mid h_p^-$ if and only if $p \mid h_p$; see [34, Theorem 5.34]. The class number h_p^+ is rather more elusive than h_p^- . Here Vandiver made the conjecture that $p \nmid h_p^+$ for any odd prime p .

Jensen [16] was the first to prove that there are infinitely many B-irregular primes. More precisely, he showed that there are infinitely many B-irregular primes not of the form $4n + 1$. This was generalized by Montgomery [22], who showed that 4 can be replaced by any integer greater than 2. To the best of our knowledge, the following result due to Metsänkylä [21] has not been improved upon.

Theorem 1.2 (Metsänkylä [21]). *Given an integer $m > 2$, let $\mathbb{Z}_m^*$ be the multiplicative group of the residue classes modulo m , and let H be a proper subgroup of $\mathbb{Z}_m^*$. Then, there exist infinitely many B-irregular primes not lying in the residue classes in H .*

Let $\mathcal{P}_B$ be the set of B-irregular primes. Carlitz [3] gave a simple proof of the infinitude of this set, and recently Luca, Pizarro-Madariaga and Pomerance [20, Theorem 1] made this more quantitative by showing that

$$(1.1) \quad \mathcal{P}_B(x) \geq (1 + o(1)) \frac{\log \log x}{\log \log \log x}$$

as $x \rightarrow \infty$. (Here and in the sequel, if S is a set of natural numbers, then $S(x)$ denotes the number of elements in S not exceeding x .) Heuristical arguments suggest a much stronger result (consistent with numerical data; see, for instance, [2, 11]).

Conjecture 1.3 (Siegel [29]). *Asymptotically we have*

$$\mathcal{P}_B(x) \sim \left(1 - \frac{1}{\sqrt{e}}\right) \pi(x),$$

where $\pi(x)$ denotes the prime counting function.

The reasoning behind this conjecture is as follows. We assume that the numerator of B_{2k} is not divisible by p with probability $1 - 1/p$. Therefore, assuming the independence of divisibility by distinct primes, we expect that p is B-regular with probability

$$\left(1 - \frac{1}{p}\right)^{\frac{p-3}{2}},$$

which with increasing p tends to $e^{-1/2}$.

For any positive integers a, d with $\gcd(a, d) = 1$, let $\mathcal{P}_B(d, a)$ be the set of B-irregular primes congruent to a modulo d . We pose the following conjecture, which suggests that B-irregular primes are uniformly distributed in arithmetic progressions. It is consistent with numerical data; see Table 2.

Conjecture 1.4. *For any positive integers a, d with $\gcd(a, d) = 1$, asymptotically we have*

$$\mathcal{P}_B(d, a)(x) \sim \frac{1}{\varphi(d)} \left(1 - \frac{1}{\sqrt{e}}\right) \pi(x),$$

where φ denotes Euler's totient function.

Although we know that there are infinitely many B-irregular primes, it is still not known whether there are infinitely many B-regular primes!

1.2. Some generalizations. In [3] Carlitz studied (ir)regularity with respect to Euler numbers.

The Euler numbers E_n are a sequence of integers defined by the relation

$$\frac{2}{e^t + e^{-t}} = \sum_{n=0}^{\infty} E_n \frac{t^n}{n!}.$$

It is easy to see that $E_0 = 1$, and $E_n = 0$ for any odd $n \geq 1$. Euler numbers, just as Bernoulli numbers, can be defined via special polynomial values. The Euler polynomials $E_n(x)$ are implicitly defined as the coefficient of t^n in the generating function

$$\frac{2e^{xt}}{e^t + 1} = \sum_{n=0}^{\infty} E_n(x) \frac{t^n}{n!}.$$

Then, $E_n = 2^n E_n(1/2)$, $n = 0, 1, 2, \dots$.

An odd prime p is said to be *E-irregular* if it divides at least one of the integers $E_2, E_4, \dots, E_{p-3}$, and *E-regular* otherwise. The first twenty E-irregular primes are

$$19, 31, 43, 47, 61, 67, 71, 79, 101, 137, 139, \\ 149, 193, 223, 241, 251, 263, 277, 307, 311.$$

Vandiver [31] proved that if p is E-regular, then the Diophantine equation $x^p + y^p = z^p$ does not have an integer solution x, y, z with p coprime to xyz . This criterion makes it possible to discard many of the exponents Kummer could not handle:

$$37, 59, 103, 131, 157, 233, 257, 271, 283, 293, \dots$$

Carlitz [3] showed that there are infinitely many E-irregular primes. Luca, Pizarro-Madariaga and Pomerance in [20, Theorem 2] showed that the number of E-irregular primes up to x satisfies the same lower bound as in (1.1). Regarding their distribution, we currently only know that there are infinitely many E-irregular primes not lying in the residue classes $\pm 1 \pmod{8}$, which was proven by Ernvall [5].

Like the B-(ir)regularity of primes, their E-(ir)regularity also relates to the divisibility of class numbers of cyclotomic fields; see [6, 7].

Let $\mathcal{P}_E$ be the set of E-irregular primes, and let $\mathcal{P}_E(d, a)$ be the set of E-irregular primes congruent to a modulo d . It was conjectured and tested in [8, Section 2] that:

Conjecture 1.5. *Asymptotically we have*

$$\mathcal{P}_E(x) \sim \left(1 - \frac{1}{\sqrt{e}}\right) \pi(x).$$

As in the case of B-irregular primes, we pose the following conjecture.

Conjecture 1.6. *For any positive integers a, d with $\gcd(a, d) = 1$, asymptotically we have*

$$\mathcal{P}_E(d, a)(x) \sim \frac{1}{\varphi(d)} \left(1 - \frac{1}{\sqrt{e}}\right) \pi(x).$$

This conjecture is consistent with computer calculations; see Table 3.

Later on, Ernvall [6, 7] introduced χ -irregular primes and proved the infinitude of such irregular primes for any Dirichlet character χ , including B-irregular primes and E-irregular primes as special cases. In addition, Hao and Parry [10] defined m -regular primes for any square-free integer m , and Holden [13] defined irregular primes by using the values of zeta functions of totally real number fields.

In this paper, we introduce a new kind of irregular prime based on Genocchi numbers and study their distribution in detail.

1.3. Regularity with respect to Genocchi numbers. The Genocchi numbers G_n are defined by the relation

$$\frac{2t}{e^t + 1} = \sum_{n=1}^{\infty} G_n \frac{t^n}{n!}.$$

It is well-known that $G_1 = 1$, $G_{2n+1} = 0$ for $n \geq 1$, and $(-1)^n G_{2n}$ is an odd positive integer. The Genocchi numbers G_n are related to Bernoulli numbers B_n by the formula

$$(1.2) \quad G_n = 2(1 - 2^n)B_n.$$

In view of the definitions of G_n and $E_n(x)$, we directly obtain

$$(1.3) \quad G_n = nE_{n-1}(0), \quad n \geq 1.$$

In analogy with Kummer and Carlitz, we here define an odd prime p to be *G-irregular* if it divides at least one of the integers $G_2, G_4, \dots, G_{p-3}$, and *G-regular* if it does not. The first twenty G-irregular primes are

$$17, 31, 37, 41, 43, 59, 67, 73, 89, 97, 101, 103, \\ 109, 113, 127, 131, 137, 149, 151, 157.$$

Clearly, if an odd prime p is B-irregular, then it is also G-irregular.

Recall that a *Wieferich prime* is an odd prime p such that $2^{p-1} \equiv 1 \pmod{p^2}$, which arose in the study of Fermat's Last Theorem. So, if an odd prime p is a Wieferich prime, then p divides G_{p-1} , and otherwise it does not divide G_{p-1} . Currently there are only two Wieferich prime known, namely 1093 and 3511. If there are further ones, they are larger than 6.7×10^{15} ; see [4]. Both 1093 and 3511 are G-irregular primes. However, 1093 is B-regular, and 3511 is B-irregular.

As in the classical case, the G-irregularity of primes can be linked to the divisibility of some class numbers of cyclotomic fields. Let S be the set of infinite places of $\mathbb{Q}(\zeta_p)$ and T the set of places above the prime 2. Denote by $h_{p,2}$ the (S, T) -refined class number of $\mathbb{Q}(\zeta_p)$. Similarly, let $h_{p,2}^+$ be the refined class number of $\mathbb{Q}(\zeta_p + \zeta_p^{-1})$ with respect to its infinite places and places above the prime 2 (for the definition of the refined class number of global fields, we refer to Gross [9, Section 1] or Hu and Kim [15, Section 2]). Define

$$h_{p,2}^- = h_{p,2}/h_{p,2}^+.$$

It turns out that $h_{p,2}^-$ is an integer (see [15, Proof of Proposition 3.4]).

Theorem 1.7. *Let p be an odd prime. Then, if p is G -irregular, we have $p \mid h_{p,2}^-$. If furthermore p is not a Wieferich prime, the converse is also true.*

1.3.1. *Global distribution of G -irregular primes.* Let g be a non-zero integer. For an odd prime $p \nmid g$, let $\text{ord}_p(g)$ be the multiplicative order of g modulo p , that is, the smallest positive integer k such that $g^k \equiv 1 \pmod{p}$.

Theorem 1.8. *An odd prime p is G -regular if and only if it is B -regular and satisfies $\text{ord}_p(4) = (p-1)/2$.*

Note that $\text{ord}_p(4) \mid (p-1)/2$. Using quadratic reciprocity it is not difficult to show (see Proposition 2.2) that if $p \equiv 1 \pmod{8}$, then $\text{ord}_p(4) \neq (p-1)/2$. Thus Theorem 1.8 has the following corollary.

Corollary 1.9. *Primes p satisfying $p \equiv 1 \pmod{8}$ are G -irregular.*

Although the B -irregular primes are very mysterious, the set of primes p such that $\text{ord}_p(4) = (p-1)/2$ is far less so. Its distributional properties are analyzed in detail in Proposition 1.12. The special case $a = d = 1$ in combination with Theorem 1.8, yields the following estimate.

Theorem 1.10. *Let $\mathcal{P}_G$ be the set of G -irregular primes. Let $\epsilon > 0$ be arbitrary and fixed. Then we have, for every x sufficiently large,*

$$\mathcal{P}_G(x) > \left(1 - \frac{3}{2}A - \epsilon\right) \frac{x}{\log x},$$

with A the Artin constant

$$(1.4) \quad A = \prod_{\text{prime } p} \left(1 - \frac{1}{p(p-1)}\right) = 0.3739558136192022880547280543464 \dots$$

Note that $1 - 3A/2 = 0.4390662795 \dots$

Using Siegel's heuristic, one arrives at the following conjecture.

Conjecture 1.11. *Asymptotically we have*

$$\mathcal{P}_G(x) \sim \left(1 - \frac{3A}{2\sqrt{e}}\right) \pi(x) \ (\approx 0.6597765 \cdot \pi(x)).$$

Some numerical evidence for Conjecture 1.11 is given in Section 6. The heuristic behind this conjecture is straightforward. Under the Generalized Riemann Hypothesis (GRH) it can be shown that the set of primes p such that $\text{ord}_p(4) = (p-1)/2$ has density $3A/2$ (Proposition 1.12 with $a = d = 1$). By Siegel's heuristic one expects a fraction $3A/(2\sqrt{e})$ of these to be B -regular. The conjecture follows on invoking Theorem 1.8.

1.3.2. *G -irregular primes in prescribed arithmetic progressions.* Using Proposition 1.12 we can give a non-trivial lower bound for the number of G -irregular primes in a prescribed arithmetic progression. Recall that if $S \subseteq T$ are sets of natural numbers, then the relative density of S in T is defined as

$$\lim_{x \rightarrow \infty} \frac{S(x)}{T(x)},$$

if this limit exists. We use the notations $\gcd$ and lcm for greatest common divisor, respectively least common multiple, but often will write (a, b) , rather than $\gcd(a, b)$. We also use the “big O” notation O , and we write O_ρ to emphasize the dependence of the implied constant on some parameter (or a list of parameters) ρ .

Proposition 1.12. *Given two coprime positive integers a and d , we put*

$$(1.5) \quad \mathcal{Q}(d, a) = \{p > 2 : p \equiv a \pmod{d}, \text{ord}_p(4) = (p-1)/2\}.$$

Let ϵ be arbitrary and fixed. Then, for every x sufficiently large we have

$$(1.6) \quad \mathcal{Q}(d, a)(x) < \frac{(\delta(d, a) + \epsilon)}{\varphi(d)} \frac{x}{\log x}$$

with

$$\delta(d, a) = c(d, a)R(d, a)A,$$

where

$$R(d, a) = 2 \prod_{p|(a-1, d)} \left(1 - \frac{1}{p}\right) \prod_{p|d} \left(1 + \frac{1}{p^2 - p - 1}\right),$$

and

$$c(d, a) = \begin{cases} 3/4 & \text{if } 4 \nmid d; \\ 1/2 & \text{if } 4 \mid d, 8 \nmid d, a \equiv 1 \pmod{4}; \\ 1 & \text{if } 4 \mid d, 8 \nmid d, a \equiv 3 \pmod{4}; \\ 1 & \text{if } 8 \mid d, a \not\equiv 1 \pmod{8}; \\ 0 & \text{if } 8 \mid d, a \equiv 1 \pmod{8}, \end{cases}$$

is the relative density of the primes $p \not\equiv 1 \pmod{8}$ in the set of primes $p \equiv a \pmod{d}$.

Under GRH, we have

$$(1.7) \quad \mathcal{Q}(d, a)(x) = \frac{\delta(d, a)}{\varphi(d)} \frac{x}{\log x} + O_d\left(\frac{x \log \log x}{\log^2 x}\right).$$

We discuss the connection of this result and Artin’s primitive root conjecture in Section 3. A numerical demonstration of the estimate (1.7) is given in Section 6 for some choices of a and d . By Proposition 2.2, in case $8 \mid d$ and $a \equiv 1 \pmod{8}$, we in fact have $\mathcal{Q}(d, a) = \emptyset$ and so $\mathcal{Q}(d, a)(x) = 0$.

Combination of Theorem 1.8 and Proposition 1.12 yields directly the following result.

Theorem 1.13. *Given two coprime positive integers a and d , we put*

$$\mathcal{P}_G(d, a) = \{p : p \equiv a \pmod{d} \text{ and } p \text{ is } G\text{-irregular}\}.$$

Let ϵ be arbitrary and fixed. For every x sufficiently large, we have

$$(1.8) \quad \mathcal{P}_G(d, a)(x) > \frac{(1 - \delta(d, a) - \epsilon)}{\varphi(d)} \frac{x}{\log x},$$

where $\delta(d, a)$ is defined in Proposition 1.12.

Under GRH we have

$$(1.9) \quad \mathcal{P}_G(d, a)(x) \geq \frac{(1 - \delta(d, a))}{\varphi(d)} \frac{x}{\log x} + O_d\left(\frac{x \log \log x}{\log^2 x}\right).$$

Note that the inequality (1.8) (with $a = d = 1$) yields Theorem 1.10 as a special case.

An easy analysis (see (2.6) in Section 2.2) shows that $1 - \delta(d, a) > 0$, and so we obtain the following corollary, which can be compared with Theorem 1.2.

Corollary 1.14. *Each primitive residue class contains a subset of G -irregular primes having positive density.*

Moreover, by Corollary 1.9 in case $a \equiv 1 \pmod{8}$ and $8|d$ we have $1 - \delta(d, a) = 1$ and in fact $\mathcal{P}_G(d, a)(x) = \pi(x; d, a)$, where

$$\pi(x; d, a) = \#\{p \leq x : p \equiv a \pmod{d}\}.$$

In the remaining cases we have $\delta(d, a) > 0$, and so $1 - \delta(d, a) < 1$. However, $1 - \delta(d, a)$ can be arbitrarily close to 1 (see Proposition 2.3), and the same holds for the relative density of $\mathcal{P}_G(d, a)$ by Theorem 1.13.

Corollary 1.14 taken by itself is not a deep result and can be easily proved directly, see Moree and Sha [26, Proposition 1.6].

The reasoning that leads us to Conjecture 1.11 in addition with the assumption that B-irregular primes are equidistributed over residue classes with a fixed modulus, suggests that the following conjecture might be true.

Conjecture 1.15. *Given two coprime positive integers a and d , asymptotically we have*

$$\mathcal{P}_G(d, a)(x) \sim \left(1 - \frac{\delta(d, a)}{\sqrt{e}}\right) \pi(x; d, a),$$

where $\delta(d, a)$ is defined in Proposition 1.12.

Numerical evidence for Conjecture 1.15 is presented in Section 6. Note that this conjecture implies Conjecture 1.11 (choosing $a = d = 1$). On observing that $\delta(a, d) = 0$ if and only if $8 \mid d$ and $a \equiv 1 \pmod{8}$, it also implies the following conjecture.

Conjecture 1.16. *Consider the subset of G -regular primes in the primitive residue class $a \pmod{d}$. It has a positive density, provided we are not in the case $8 \mid d$ and $a \equiv 1 \pmod{8}$.*

2. PRELIMINARIES

In this section, we gather some results which are used later on.

2.1. Elementary results. For a primitive Dirichlet character χ with an odd conductor f , the generalized Euler numbers $E_{n, \chi}$ are defined by (see [17, Section 5.1])

$$2 \sum_{a=1}^f \frac{(-1)^a \chi(a) e^{at}}{e^{ft} + 1} = \sum_{n=0}^{\infty} E_{n, \chi} \frac{t^n}{n!}.$$

For any odd prime p , let ω_p be the Teichmüller character of $\mathbb{Z}/p\mathbb{Z}$. Any multiplicative character of $\mathbb{Z}/p\mathbb{Z}$ is of the form ω_p^k for some $1 \leq k \leq p-1$. In particular, the odd characters are ω_p^k with $k = 1, 3, \dots, p-2$.

The following lemma, formulated and proved in the notation of [17], is an analogue of a well-known result for the generalized Bernoulli numbers; see [34, Corollary 5.15].

Lemma 2.1. *Suppose that p is an odd prime and k, n are non-negative integers. Then $E_{k, \omega_p^{n-k}} \equiv E_n(0) \pmod{p}$.*

Proof. By [17, Proposition 5.4], for arbitrary integers $k, n \geq 0$, we have

$$(2.1) \quad E_{k, \omega_p^{n-k}} = \int_{\mathbb{Z}_p} \omega_p^{n-k}(a) a^k d\mu_{-1}(a).$$

By [17, Proposition 2.1 (1)], $E_n(0)$ also can be expressed as a p -adic integral, namely

$$(2.2) \quad E_n(0) = \int_{\mathbb{Z}_p} a^n d\mu_{-1}(a).$$

Since $\omega_p(a) \equiv a \pmod{p}$, we have

$$(2.3) \quad \omega_p^{n-k}(a) \equiv a^{n-k} \pmod{p} \quad \text{and} \quad \omega_p^{n-k}(a) a^k \equiv a^n \pmod{p}.$$

From (2.1), (2.2) and (2.3), we deduce that

$$E_{k, \omega_p^{n-k}} - E_n(0) = \int_{\mathbb{Z}_p} (\omega_p^{n-k}(a) a^k - a^n) d\mu_{-1}(a) \equiv 0 \pmod{p}. \quad \square$$

Recall that $\mathcal{Q}(d, a)$ is defined in (1.5). For ease of notation we put

$$(2.4) \quad \mathcal{Q} = \mathcal{Q}(1, 1) = \{p > 2 : \text{ord}_p(4) = (p-1)/2\}.$$

For the understanding of the distribution of the primes in $\mathcal{Q}$, it turns out to be very useful to consider their residues modulo 8.

Proposition 2.2. *For $j = 1, 3, 5, 7$ we put $\mathcal{Q}_j = \mathcal{Q}(8, j)$. We have*

$$\mathcal{Q} = \mathcal{Q}_1 \cup \mathcal{Q}_3 \cup \mathcal{Q}_5 \cup \mathcal{Q}_7,$$

with $\mathcal{Q}_1 = \emptyset$, and, for $j = 3, 5$,

$$\mathcal{Q}_j = \{p : p \equiv j \pmod{8}, \text{ord}_p(2) = p-1\},$$

and, furthermore,

$$\mathcal{Q}_7 = \{p : p \equiv 7 \pmod{8}, \text{ord}_p(2) = (p-1)/2\}.$$

Proof. If $p \equiv 1 \pmod{8}$, then by quadratic reciprocity $2^{(p-1)/2} \equiv 1 \pmod{p}$, and we conclude that $\text{ord}_p(4) \mid (p-1)/4$ and hence $\mathcal{Q}_1 = \emptyset$.

Note that

$$\text{ord}_p(4) = \begin{cases} \text{ord}_p(2) & \text{if } \text{ord}_p(2) \text{ is odd;} \\ \text{ord}_p(2)/2 & \text{otherwise.} \end{cases}$$

In case $p \equiv \pm 3 \pmod{8}$, we have $2^{(p-1)/2} \equiv -1 \pmod{p}$, and so $\text{ord}_p(2)$ must be even. The assumption that p is in $\mathcal{Q}$ now implies that $\text{ord}_p(2) = 2 \cdot \text{ord}_p(4) = p-1$. In case $p \equiv 7 \pmod{8}$, we have $2^{(p-1)/2} \equiv 1 \pmod{p}$, and so $\text{ord}_p(2)$ must be odd. The assumption that p is in $\mathcal{Q}$ now implies that $\text{ord}_p(2) = \text{ord}_p(4) = (p-1)/2$. $\square$

2.2. The size of $\delta(d, a)$. In this section we study the extremal behaviour of the quantity $\delta(d, a)$ defined in Proposition 1.12. We put

$$G(d) = \prod_{p|d} \left(1 + \frac{1}{p^2 - p - 1}\right), \quad F(d) = \frac{\varphi(d)}{d} G(d).$$

An easy calculation gives that

$$G(d) = \frac{1}{A} \prod_{p \nmid d} \left(1 - \frac{1}{p(p-1)}\right) = \frac{1}{A} \left(1 + O\left(\frac{1}{q}\right)\right),$$

where q is the smallest prime not dividing d . Trivially, $G(d) < 1/A$, and $G(d) < 1/(2A)$ when d is odd.

It is a classical result (see, for instance, [1, Theorem 13.14]), that

$$\liminf_{d \rightarrow \infty} \frac{\varphi(d)}{d} \log \log d = e^{-\gamma},$$

where γ is the Euler-Mascheroni constant ($\gamma = 0.577215664901532 \dots$). The proof is in essence an application of Mertens' theorem (see [1, Theorem 13.13])

$$(2.5) \quad \prod_{p \leq x} \left(1 - \frac{1}{p}\right) \sim \frac{e^{-\gamma}}{\log x}.$$

An easy variation of the latter proof yields

$$\liminf_{d \rightarrow \infty} AF(d) \log \log d = e^{-\gamma}.$$

Recall that

$$R(d, a) = 2G(d) \prod_{p|b} \left(1 - \frac{1}{p}\right) = 2G(d) \frac{\varphi(b)}{b}, \text{ with } b = (a-1, d).$$

Note that

$$2F(d) \leq R(d, a) \leq 2G(d)/(2, d) < 1/A,$$

and hence $\delta(d, a) = 0$ or

$$(2.6) \quad 0 < AF(d) \leq \delta(d, a) \leq 2AG(d)/(2, d) < 1.$$

Proposition 2.3. *We have*

$$\liminf_{d \rightarrow \infty} \min_{\substack{1 \leq a < d \\ (a, d) = 1 \\ \delta(d, a) > 0}} \delta(d, a) \log \log d = e^{-\gamma} \quad \text{and} \quad \limsup_{d \rightarrow \infty} \max_{\substack{1 \leq a < d \\ (a, d) = 1}} \delta(d, a) = 1.$$

Proof. From the above remarks it follows that the limit inferior and superior are $\geq e^{-\gamma}$, respectively ≤ 1 . We consider two infinite families of pairs (a, d) to show that these bounds are actually sharp.

Let $n \geq 3$ be arbitrary. Put $d_n = \prod_{3 \leq p \leq n} p$. We have $c(4d_n, 1) = 1/2$ and

$$\delta(4d_n, 1) = (1 + o(1)) \prod_{2 \leq p \leq n} (1 - 1/p), \quad (n \rightarrow \infty)$$

TABLE 1. Some values of $\delta(4d_n, 1)$ and $\delta(8d_n, a_n)$

n	10^3	10^4	10^5	10^6	10^7
$\delta(4d_n, 1) \approx$	0.080954	0.060884	0.048752	0.040638	0.034833
$\delta(4d_n, 1)e^\gamma \log \log(4d_n) \approx$	0.989659	0.997633	0.999422	0.999851	0.999960
$\delta(8d_n, a_n) \approx$	0.999872	0.999990	0.999999	0.9999999	0.99999999

by Proposition 1.12. Using Mertens' theorem (2.5) and the prime number theorem, we deduce that

$$\delta(4d_n, 1) \sim \frac{e^{-\gamma}}{\log n} \sim \frac{e^{-\gamma}}{\log \log(4d_n)}, \quad (n \rightarrow \infty),$$

and so the limit inferior actually equals $e^{-\gamma}$.

Put

$$a_n = \begin{cases} 2 + 3d_n & \text{if } d_n \equiv 7 \pmod{8}; \\ 2 + d_n & \text{otherwise.} \end{cases}$$

We have $a_n \not\equiv 1 \pmod{8}$, $1 \leq a_n < 8d_n$, $(a_n, 8d_n) = 1$, and $(a_n - 1, 8d_n)$ is a power of two. We infer that $R(8d_n, a_n) = 1/A + O(1/n)$ and $c(8d_n, a_n) = 1$, and so $\delta(8d_n, a_n) = 1 + O(1/n)$, showing that the limit superior equals 1. $\square$

The two constructions in the above proof are put to the test in Table 1. The table also gives an idea of how fast the lower bound $1 - \delta(4d_n, 1)$ for the relative density of the set $\mathcal{P}_G(4d_n, 1)$ established in Theorem 1.13, tends to 1.

3. SOME RESULTS RELATED TO ARTIN'S PRIMITIVE ROOT CONJECTURE

It is natural to wonder whether the set $\mathcal{Q}$, see (2.4), is an infinite set or not. This is closely related to Artin's primitive root conjecture stating that if $g \neq -1$ or a square, then infinitely often $\text{ord}_p(g) = p - 1$ (which is maximal by Fermat's little theorem). In case g is a square, the maximal order is $(p - 1)/2$ and one can wonder whether this happens infinitely often. If this is so for $g = 4$, then our set $\mathcal{Q}$ is infinite. We now go into a bit more technical detail.

We say that a set of primes $\mathcal{P}$ has density $\delta(\mathcal{P})$ and satisfies a Hooley type estimate, if

$$(3.1) \quad \mathcal{P}(x) = \delta(\mathcal{P}) \frac{x}{\log x} + O\left(\frac{x \log \log x}{\log^2 x}\right),$$

where the implied constant may depend on $\mathcal{P}$.

Let $g \notin \{-1, 0, 1\}$ be an integer. Put

$$\mathcal{P}_g = \{p : \text{ord}_p(g) = p - 1\}.$$

Artin in 1927 conjectured that this set, when g is not a square, is infinite and also conjectured a density for it. To this day, this conjecture is open; see [25] for a survey. Hooley [14] proved in 1967 that if the Riemann Hypothesis holds for the number fields

$\mathbb{Q}(\zeta_n, g^{1/n})$ with all square-free n (this is a weaker form of the GRH), then the estimate (3.1) holds for the set $\mathcal{P}_g$ with

$$\delta(g) := \delta(\mathcal{P}_g) = \sum_{n=1}^{\infty} \frac{\mu(n)}{[\mathbb{Q}(\zeta_n, g^{1/n}) : \mathbb{Q}]},$$

where μ is the Möbius function; and he also showed that $\delta(g)/A$ is rational and explicitly determined its value, with A the Artin constant (see (1.4)). For example, in case $g = 2$ we have $\delta(2) = A$.

By the Chebotarev density theorem, the density of primes $p \equiv 1 \pmod{n}$ such that $\text{ord}_p(g) \mid (p-1)/n$ is equal to $1/[\mathbb{Q}(\zeta_n, g^{1/n}) : \mathbb{Q}]$. Note that in order to ensure that $\text{ord}_p(g) = p-1$, it is enough to show that there is no prime q such that $\text{ord}_p(g) \mid (p-1)/q$. By inclusion and exclusion we are then led to expect that the set $\mathcal{P}_g$ has natural density $\delta(g)$. The problem with establishing this rigorously is that the Chebotarev density theorem only allows one to take finitely many splitting conditions into account. Let us now consider which result we can obtain on restricting to the primes $q \leq y$. Put

$$(3.2) \quad \delta_y(g) = \sum_{P(n) \leq y} \frac{\mu(n)}{[\mathbb{Q}(\zeta_n, g^{1/n}) : \mathbb{Q}]},$$

where $P(n)$ denotes the largest prime factor of n . Now we may apply the Chebotarev density theorem and we obtain that

$$(3.3) \quad \mathcal{P}_g(x) \leq (\delta_y(g) + \epsilon) \frac{x}{\log x},$$

where $\epsilon > 0$ is arbitrary and x, y are sufficiently large (where sufficiently large may depend on the choice of ϵ).

Completing the sum in (3.2) and using that $[\mathbb{Q}(\zeta_n, g^{1/n}) : \mathbb{Q}] \gg_g n\varphi(n)$ (see [33, Proposition 4.1]) and the classical estimate $\varphi(n)^{-1} = O((\log \log n)/n)$, we obtain that

$$\delta_y(g) = \delta(g) + O_g \left(\sum_{n \geq y} \frac{1}{n\varphi(n)} \right) = \delta(g) + O_g \left(\frac{\log \log y}{y} \right).$$

On combining this with (3.3) we obtain the estimate

$$(3.4) \quad \mathcal{P}_g(x) \leq (\delta(g) + \epsilon) \frac{x}{\log x},$$

where $\epsilon > 0$ is arbitrary and x, y are sufficiently large (where sufficiently large may depend on the choices of ϵ and g).

For any integer $g \notin \{-1, 0, 1\}$ and any integer $t \geq 1$, put

$$\mathcal{P}(g, t) = \{p : p \equiv 1 \pmod{t}, \text{ord}_p(g) = (p-1)/t\}.$$

Now, if the Riemann Hypothesis holds for the number fields $\mathbb{Q}(\zeta_{nt}, g^{1/nt})$ with all square-free n , then Hooley's proof can be easily extended, resulting in the estimate (3.1) for the set $\mathcal{P}(g, t)$ with density

$$(3.5) \quad \delta(g, t) = \sum_{n=1}^{\infty} \frac{\mu(n)}{[\mathbb{Q}(\zeta_{nt}, g^{1/nt}) : \mathbb{Q}]},$$

and with $\delta(g, t)/A$ a rational number; see [18]. This number was first computed explicitly by Wagstaff [33, Theorem 2.2], which can be done much more compactly and elegantly these days using the character sum method of Lenstra et al. [19].

By Wagstaff's work [33] we have $\delta(\mathcal{Q}) = \delta(4, 2) = 3A/2$. Alternatively it is an easy and instructive calculation to determine $\delta(4, 2)$ oneself. Since $\sqrt{2} \in \mathbb{Q}(\zeta_n)$ if and only if $8 \mid n$, we see that if $4 \nmid n$, then $[\mathbb{Q}(\zeta_{2n}, 2^{1/2n}) : \mathbb{Q}] = \varphi(2n)n$ and so by (3.5),

$$\delta(4, 2) = \sum_{n=1}^{\infty} \frac{\mu(n)}{\varphi(2n)n} = \sum_{2 \nmid n} \frac{\mu(n)}{\varphi(n)n} + \sum_{2 \mid n} \frac{\mu(n)}{2\varphi(n)n} = \frac{3}{4} \sum_{2 \nmid n} \frac{\mu(n)}{\varphi(n)n} = \frac{3}{2}A,$$

where we use the fact that

$$\sum_{\substack{n=1 \\ (m,n)=1}}^{\infty} \mu(n)f(n) = \prod_{p \nmid m} (1 - f(p))$$

holds certainly true if the sum is absolutely convergent and $f(n)$ is a multiplicative function defined on the square free integers (cf. Moree and Zumalacárregui [27, Appendix A.1], where a similar problem with $g = 9$ instead of $g = 4$ is considered).

The following result generalizes the above to the case where we require the primes in $\mathcal{P}(g, t)$ to also be in some prescribed arithmetic progression. It follows from Lenstra's work [18], who introduced Galois theory into the subject.

Theorem 3.1. *Let $1 \leq a \leq d$ be coprime integers. Let $t \geq 1$ be an integer. Put*

$$\mathcal{P}(g, t, d, a) = \{p : p \equiv 1 \pmod{t}, p \equiv a \pmod{d}, \text{ord}_p(g) = (p-1)/t\}.$$

Let σ_a be the automorphism of $\mathbb{Q}(\zeta_d)$ determined by $\sigma_a(\zeta_d) = \zeta_d^a$. Let $c_a(m)$ be 1 if the restriction of σ_a to the field $\mathbb{Q}(\zeta_d) \cap \mathbb{Q}(\zeta_m, g^{1/m})$ is the identity and $c_a(m) = 0$ otherwise. Put

$$\delta(g, t, d, a) = \sum_{n=1}^{\infty} \frac{\mu(n)c_a(nt)}{[\mathbb{Q}(\zeta_d, \zeta_{nt}, g^{1/nt}) : \mathbb{Q}]}.$$

Then, assuming RH for all number fields $\mathbb{Q}(\zeta_d, \zeta_{nt}, g^{1/nt})$ with n square-free, we have

$$(3.6) \quad \mathcal{P}(g, t, d, a)(x) = \delta(g, t, d, a) \frac{x}{\log x} + O_{g,t,d} \left(\frac{x \log \log x}{\log^2 x} \right),$$

Unconditionally we have the weaker statement that

$$(3.7) \quad \mathcal{P}(g, t, d, a)(x) \leq (\delta(g, t, d, a) + \epsilon) \frac{x}{\log x},$$

where $\epsilon > 0$ is arbitrary and x is sufficiently large (where sufficiently large may depend on the choice of ϵ, g, t, d and a).

It seems that this result has not been formulated in the literature. It is a simple combination of two cases each of which have been intensively studied, namely the primes having a near-primitive root ($d = 1, t > 1$), and the primes in arithmetic progression having a prescribed primitive root ($t = 1$).

As before $\delta(g, t, d, a)/A$ is a rational number that can be explicitly computed. The case $g = t = 2, d = 8$ and $a = 7$ is one of the most simple cases. This is a lucky

coincidence, as in our proof of Proposition 1.12 we will apply Theorem 3.1 to determine $\delta(\mathcal{Q}_7) = \delta(2, 2, 8, 7)$.

Note that $\mathcal{P}(g, t, d, a) \subseteq \{p : p \equiv 1 \pmod{t}, p \equiv a \pmod{d}, \text{ord}_p(g) \mid (p-1)/t\}$. It is shown in [26, Theorem 1.3] that if the latter set is not empty, then it contains a positive density subset of primes that are not in $\mathcal{P}(g, t, d, a)$.

4. PROOFS OF THE MAIN RESULTS

It suffices to prove Theorems 1.7 and 1.8 and Proposition 1.12.

4.1. Proof of Theorem 1.7. By [15, Proposition 3.4], we obtain

$$h_{p,2}^- = (-1)^{\frac{p-1}{2}} 2^{2-p} E_{0,\omega_p} E_{0,\omega_p^3} \cdots E_{0,\omega_p^{p-2}}.$$

Using Lemma 2.1 and (1.3), we then infer that

$$\begin{aligned} h_{p,2}^- &\equiv (-1)^{\frac{p-1}{2}} 2^{2-p} E_1(0) E_3(0) \cdots E_{p-2}(0) \\ &\equiv \frac{(-1)^{\frac{p-1}{2}} 2^{2-p}}{(p-1)!!} G_2 G_4 \cdots G_{p-3} G_{p-1} \pmod{p}. \end{aligned}$$

So, if p is G-irregular, then $p \mid h_{p,2}^-$. Conversely, if $p \mid h_{p,2}^-$ and p is not a Wieferich prime, then by (1.2) we first have $p \nmid G_{p-1}$, and thus p is G-irregular. $\square$

4.2. Proof of Theorem 1.8. We first recall a fact about Bernoulli numbers that any odd prime p does not divide the denominators of the Bernoulli numbers $B_2, B_4, \dots, B_{p-3}$ (this follows from the von Staudt-Clausen theorem). Now, given an odd prime p , if it is G-regular, then there is no $1 \leq k \leq (p-3)/2$ such that p divides the integer G_{2k} , that is, $2(1-2^{2k})B_{2k}$ by (1.2); and so p is B-regular and $\text{ord}_p(4) = (p-1)/2$.

Conversely, if p is B-regular and $\text{ord}_p(4) = (p-1)/2$, then p does not divide the denominators of the Bernoulli numbers $B_2, B_4, \dots, B_{p-3}$ and $p \nmid 2^{2k} - 1$ for $1 \leq k \leq (p-3)/2$. Consequently p does not divide any integer $G_{2k} = 2(1-2^{2k})B_{2k}$ with $1 \leq k \leq (p-3)/2$, which implies that p is G-regular. $\square$

4.3. Proof of Proposition 1.12. The proof relies on Theorem 3.1. We only establish the assertion under GRH, as the proof of the unconditional result is very similar. Namely, it uses the unconditional estimate (3.7) instead of (3.6).

It is enough to prove the result in case $8 \mid d$. In fact, in case $8 \nmid d$ we lift the congruence class $a \pmod{d}$ to congruence classes with modulus $\text{lcm}(8, d)$. The ones among those that are $\not\equiv 1 \pmod{8}$ have relative density $AR(\text{lcm}(8, d), a) = AR(d, a)$ (as $R(d, a)$ only depends on the odd prime factors of d). The one that is $\equiv 1 \pmod{8}$ (if it exists at all) has relative density zero. It follows that the relative density of the unlifted congruence equals $c(d, a)R(d, a)A$ with $c(d, a)$ the relative density of the primes $p \not\equiv 1 \pmod{8}$ in the congruence class $a \pmod{d}$. The easy determination of $c(d, a)$ is left to the interested reader.

From now on we assume that $8 \mid d$. We can write $a \equiv j \pmod{8}$ for some $j \in \{1, 3, 5, 7\}$ and distinguish three cases.

Case I: $j = 1$. By Proposition 2.2 the set $\mathcal{Q}(d, a)$ is empty and the result holds trivially true.

Case II: $j \in \{3, 5\}$. By Proposition 2.2,

$$\mathcal{Q}(d, a) = \{p : p \equiv a \pmod{d}, \text{ord}_p(2) = p - 1\}.$$

By Theorem 3.1, under GRH, this set has density $\delta(2, 1, d, a)$. For arbitrary g, d, a the third author determined the rational number $\delta(g, 1, d, a)/A$, see [23, Theorem 1] or [24, Theorem 1.2]. On applying his result, the proof of this subcase is then completed.

Case III: $j = 7$. By Proposition 2.2,

$$\mathcal{Q}(d, a) = \{p : p \equiv a \pmod{d}, \text{ord}_p(2) = (p - 1)/2\}.$$

For simplicity we write $\delta = \delta(\mathcal{Q}(d, a))$. By Theorem 3.1 we have

$$(4.1) \quad \delta = \delta(2, 2, d, a) = \sum_{n=1}^{\infty} \frac{\mu(n)c_a(2n)}{[\mathbb{Q}(\zeta_d, \zeta_{2n}, 2^{1/2n}) : \mathbb{Q}]}.$$

In case n is even, then trivially $\mathbb{Q}(\sqrt{-1}) \subseteq \mathbb{Q}(\zeta_d) \cap \mathbb{Q}(\zeta_{2n}, 2^{1/2n})$. As σ_a acts by conjugation on $\mathbb{Q}(\sqrt{-1})$, cf. [24, Lemma 2.2], and not as the identity, it follows that $c_a(2n) = 0$.

Next assume that n is odd and square-free. Then by [24, Lemma 2.4] we infer that

$$\mathbb{Q}(\zeta_d) \cap \mathbb{Q}(\zeta_{2n}, 2^{1/2n}) = \mathbb{Q}(\zeta_{(d,n)}, \sqrt{2}).$$

Since

$$\sigma_a|_{\mathbb{Q}(\sqrt{2})} = \text{id.} \quad \text{and} \quad \sigma_a|_{\mathbb{Q}(\zeta_{(d,2n)})} \begin{cases} = \text{id.} & \text{if } a \equiv 1 \pmod{(d, 2n)}; \\ \neq \text{id.} & \text{otherwise,} \end{cases}$$

we conclude that

$$c_a(2n) = \begin{cases} 1 & \text{if } a \equiv 1 \pmod{(d, 2n)}; \\ 0 & \text{otherwise,} \end{cases}$$

with ‘id.’ a shorthand for identity. Note that the assumptions on a, d and n imply that $a \equiv 1 \pmod{(d, 2n)}$ iff $a \equiv 1 \pmod{2(d, n)}$ iff $a \equiv 1 \pmod{(d, n)}$. It follows that (4.1) simplifies to

$$\delta = \sum_{\substack{2 \nmid n \\ a \equiv 1 \pmod{(d, n)}}} \frac{\mu(n)}{[\mathbb{Q}(\zeta_d, \zeta_{2n}, 2^{1/2n}) : \mathbb{Q}]}.$$

When n is odd and square-free, using [24, Lemma 2.3] we obtain

$$[\mathbb{Q}(\zeta_d, \zeta_{2n}, 2^{1/2n}) : \mathbb{Q}] = [\mathbb{Q}(\zeta_{\text{lcm}(d, 2n)}, 2^{1/2n}) : \mathbb{Q}] = n\varphi(\text{lcm}(d, 2n)) = n\varphi(\text{lcm}(d, n)).$$

We thus get

$$\varphi(d)\delta = \sum_{\substack{2 \nmid n \\ a \equiv 1 \pmod{(d, n)}}} \frac{\mu(n)\varphi(d)}{n\varphi(\text{lcm}(d, n))}.$$

Put

$$w(n) = \frac{n\varphi(\text{lcm}(d, n))}{\varphi(d)}.$$

In this notation we obtain

$$\delta = \frac{1}{\varphi(d)} \sum_{\substack{2 \nmid n \\ a \equiv 1 \pmod{(d,n)}}} \frac{\mu(n)}{w(n)},$$

where the argument in the sum is multiplicative in n . Using [24, Lemma 3.1] and the notation used there and in [24, Theorem 1.2], we find

$$\begin{aligned} \varphi(d)\delta &= S(1) - S_2(1) = 2S(1) = 2A(a, d, 1) \\ &= 2A \prod_{p|(a-1, d)} \left(1 - \frac{1}{p}\right) \prod_{p|d} \left(1 + \frac{1}{p^2 - p - 1}\right) = \delta(d, a), \end{aligned}$$

as was to be proved. $\square$

5. OUTLOOK

A small improvement of the upper bound (1.6) (and consequently the lower bound (1.8)) would be possible if instead of the estimate (3.7) a Vinogradov type estimate for $\mathcal{P}(g, t, d, a)(x)$ could be established, say

$$(5.1) \quad \mathcal{P}(g, t, d, a)(x) \leq \delta(g, t, d, a) \frac{x}{\log x} + O_{g,t,d} \left(\frac{x(\log \log x)^2}{\log^{5/4} x} \right).$$

Vinogradov [32] established the above result in case $a = d = t = 1$. Establishing (5.1) seems technically quite involved. Recent work by Pierce et al. [28] offers perhaps some hope that one can even improve on the error term in (5.1).

6. SOME NUMERICAL EXPERIMENTS

In this section, using the Bernoulli numbers modulo p function developed by David Harvey in Sage [30] (see [2, 11, 12] for more details and improvements) and the `euler_number` function in Sage, we provide numerical evidence for the truth of Conjectures 1.4, 1.6, 1.11 and 1.15 and also for (1.7) in Proposition 1.12.

The Bernoulli numbers modulo p function returns the values of $B_0, B_2, \dots, B_{p-3}$ modulo p , and so by checking whether there is a zero value we can determine whether p is B-irregular. For checking the E-irregularity, we use the `euler_number` function in Sage to compute and store Euler numbers and then use the definition of E-irregularity. It would be a separate project to test large E-irregular primes, cf. [2, 11, 12].

In the tables, we only record the first six digits of the decimal parts.

In Tables 2 and 3 the ratio $\mathcal{P}_B(d, a)(x)/\pi(x; d, a)$, respectively $\mathcal{P}_E(d, a)(x)/\pi(x; d, a)$ is recorded for $x = 10^5$ in the column ‘experimental’ for various choices of d and a , and in the column ‘theoretical’ the limit value predicted by Conjecture 1.4 is given.

Table 4 gives the ratio $\mathcal{P}_G(x)/\pi(x)$ for various values of x , and the value in the column ‘theoretical’ is the limit value $1 - 3A/(2\sqrt{e})$ predicted by Conjecture 1.11.

Table 5 gives the ratio $\mathcal{P}_G(d, a)(x)/\pi(x; d, a)$ for $x = 5 \cdot 10^6$ in the column ‘experimental’ for various choices of d and a , and the corresponding limit values $1 - c(d, a)R(d, a)A/\sqrt{e}$ predicted by Conjecture 1.15 are in the column ‘theoretical’.

TABLE 2. The ratio $\mathcal{P}_B(d, a)(x)/\pi(x; d, a)$ for $x = 10^5$

d	a	experimental	theoretical
3	2	0.394424	0.393469
4	1	0.388877	
5	4	0.397071	
7	4	0.391005	
9	8	0.387742	
12	5	0.390203	
15	13	0.389858	
20	13	0.385191	

TABLE 3. The ratio $\mathcal{P}_E(d, a)(x)/\pi(x; d, a)$ for $x = 10^5$

d	a	experimental	theoretical
3	2	0.395672	0.393469
4	1	0.388040	
5	4	0.397071	
7	4	0.393504	
9	8	0.391494	
12	5	0.388127	
15	13	0.399002	
20	13	0.385191	

TABLE 4. The ratio $\mathcal{P}_G(x)/\pi(x)$

x	experimental	theoretical
10^5	0.661592	0.659776
10^6	0.659558	
$2 \cdot 10^6$	0.660860	
$3 \cdot 10^6$	0.661413	
$4 \cdot 10^6$	0.660683	
$5 \cdot 10^6$	0.660864	

TABLE 5. The ratio $\mathcal{P}_G(d, a)(x)/\pi(x; d, a)$ for $x = 5 \cdot 10^6$

d	a	experimental	theoretical
3	1	0.728296	0.727821
5	2	0.643010	0.641870
4	1	0.771512	0.773184
20	9	0.757311	0.761246
12	11	0.460584	0.455642
20	19	0.528567	0.522493
8	7	0.550086	0.546368
24	13	0.634191	0.637094

TABLE 6. The ratio $\mathcal{Q}(d, a)(x)/\pi(x; d, a)$ for $x = 5 \cdot 10^6$

d	a	experimental	theoretical
3	1	0.449049	0.448746
5	2	0.589614	0.590456
4	1	0.374664	0.373955
20	9	0.395498	0.393637
12	11	0.898284	0.897493
20	19	0.789316	0.787275
8	7	0.747300	0.747911
24	13	0.598815	0.598329

Finally, Table 6 gives the ratio $\mathcal{Q}(d, a)(x)/\pi(x; d, a)$ for $x = 5 \cdot 10^6$ in the column ‘experimental’ for various choices of d and a . In the column ‘theoretical’, there is the corresponding relative density $\delta(d, a)$ predicted in (1.7) and known to be true under GRH.

In view of the definition of the constant $c(d, a)$ in Proposition 1.12, there are four cases excluding the case $8|d$ and $a \equiv 1 \pmod{8}$ (which gives $c(d, a) = 0$). For each of these four cases there are two instances in Tables 5 and 6.

ACKNOWLEDGEMENT

The authors would like to thank the referee for careful reading and valuable comments. This work was supported by the National Natural Science Foundation of China, Grant No. 11501212. The research of Min-Soo Kim and Min Sha was also supported by the Kyungnam University Foundation Grant, 2017, respectively a Macquarie University Research Fellowship. The authors thank Bernd Kellner for pointing out a link with the Genocchi numbers and suggesting the references [6, 10, 13], and Peter Stevenhagen for very helpful feedback. They also thank Alexandru Ciolan for proofreading earlier versions.

REFERENCES

- [1] T.M. Apostol, *Introduction to analytic number theory*, Undergraduate Texts in Mathematics, Springer-Verlag, New York-Heidelberg, 1976.
- [2] J.P. Buhler and D. Harvey, *Irregular primes to 163 million*, Math. Comp. **80** (2011), 2435–2444.
- [3] L. Carlitz, *Note on irregular primes*, Proc. Amer. Math. Soc. **5** (1954), 329–331.
- [4] F.G. Dorais and D. Klyve, *A Wieferich prime search up to 6.7×10^{15}* , J. Integer Sequences **14** (2011), Article 11.9.2.
- [5] R. Ernvall, *On the distribution mod 8 of the E-irregular primes*, Ann. Acad. Sci. Fenn. Ser. A I, **1** (1975), 195–198.
- [6] R. Ernvall, *Generalized Bernoulli numbers, generalized irregular primes, and class number*, Ann. Univ. Turku., Ser. AI **178** (1979), 72 pp.
- [7] R. Ernvall, *Generalized irregular primes*, Mathematika **30** (1983), 67–73.
- [8] R. Ernvall and T. Metsänkylä, *Cyclotomic invariants and E-irregular primes*, Math. Comp. **32** (1978), 617–629.
- [9] B. Gross, *On the values of abelian L-functions at $s = 0$* , J. Fac. Sci. Univ. Tokyo **35** (1988), 177–197.

- [10] F.H. Hao and C.J. Parry, *Generalized Bernoulli numbers and m -regular primes*, Math. Comp. **43** (1984), 273–288.
- [11] W. Hart, D. Harvey and W. Ong, *Irregular primes to two billion*, Math. Comp. **86** (2017), 3031–3049.
- [12] D. Harvey, *A multimodular algorithm for computing Bernoulli numbers*, Math. Comp. **79** (2010), 2361–2370.
- [13] J. Holden, *Irregularity of prime numbers over real quadratic fields*, Algorithmic number theory (J.P. Buhler, ed.), Springer, Lect. Notes Comput. Sci. **1423** (1998), 454–462.
- [14] C. Hooley, *On Artin’s conjecture*, J. Reine Angew. Math. **225** (1967), 209–220.
- [15] S. Hu and M.-S. Kim, *The $(S, \{2\})$ -Iwasawa theory*, J. Number Theory **158** (2016), 73–89.
- [16] K.L. Jensen, *Om talteoretiske Egenskaber ved de Bernoulliske Tal*, Nyt Tidsskr. Mat. **26**, Afd. B (1915), 73–83.
- [17] M.-S. Kim and S. Hu, *On p -adic Hurwitz-type Euler zeta functions*, J. Number Theory **132** (2012), 2977–3015.
- [18] H.W. Lenstra, Jr., *On Artin’s conjecture and Euclid’s algorithm in global fields*, Invent. Math. **42** (1977), 202–224.
- [19] H.W. Lenstra, Jr., P. Moree and P. Stevenhagen, *Character sums for primitive root densities*, Math. Proc. Cambridge Philos. Soc. **157** (2014), 489–511.
- [20] F. Luca, A. Pizarro-Madariaga and C. Pomerance, *On the counting function of irregular primes*, Indag. Math. **26** (2015), 147–161.
- [21] T. Metsänkylä, *Distribution of irregular prime numbers*, J. Reine Angew. Math. **282** (1976), 126–130.
- [22] H.L. Montgomery, *Distribution of irregular primes*, Illinois J. Math. **9** (1965), 553–558.
- [23] P. Moree, *On primes in arithmetic progression having a prescribed primitive root*, J. Number Theory **78** (1999), 85–98.
- [24] P. Moree, *On primes in arithmetic progression having a prescribed primitive root. II*, Funct. Approx. Comment. Math. **39** (2008), 133–144.
- [25] P. Moree, *Artin’s primitive root conjecture – a survey*, Integers **12A** (2012), A13.
- [26] P. Moree and M. Sha, *Primes in arithmetic progressions and nonprimitive roots*, B. Aust. Math. Soc., to appear, <https://arxiv.org/abs/1901.02650>.
- [27] P. Moree and A. Zumalacárregui, *Salajan’s conjecture on discriminating terms in an exponential sequence*, J. Number Theory **160** (2016), 646–665.
- [28] L.B. Pierce, C.L. Turnage-Butterbaugh and M.M. Wood, *An effective Chebotarev density theorem for families of number fields, with an application to ℓ -torsion in class groups*, arXiv:1709.09637, preprint, 2017.
- [29] C.L. Siegel, *Zu zwei Bemerkungen Kummers*, Nachr. Akad. d. Wiss. Goettingen, Math. Phys. Kl. **II** (1964), 51–62.
- [30] The Sage Developers, *SageMath, the Sage Mathematics Software System (Version 8.2)*, 2018, <http://www.sagemath.org>.
- [31] H.S. Vandiver, *Note on Euler number criteria for the first case of Fermat’s last theorem*, Amer. J. Math. **62** (1940), 79–82.
- [32] A.I. Vinogradov, *Artin’s L -series and his conjectures*, (English. Russian original), Proc. Steklov Inst. Math. **112**, 124–142 (1971); translation from Trudy Mat. Inst. Steklov **112**, 123–140 (1971).
- [33] S.S. Wagstaff, Jr., *Pseudoprimes and a generalization of Artin’s conjecture*, Acta Arith. **41** (1982), 141–150.
- [34] L.C. Washington, *Introduction to Cyclotomic Fields*, 2nd ed., Springer-Verlag, New York, 1997.

DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY, GUANGZHOU 510640, CHINA

E-mail address: mahusu@scut.edu.cn

DIVISION OF MATHEMATICS, SCIENCE, AND COMPUTERS, KYUNGNAM UNIVERSITY, 7(WORYEONG-DONG) KYUNGNAMDAEHAK-RO, MASANHAPPO-GU, CHANGWON-SI, GYEONGSANGNAM-DO 51767, REPUBLIC OF KOREA

E-mail address: mskim@kyungnam.ac.kr

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, VIVATSGASSE 7, D-53111 BONN, GERMANY

E-mail address: moree@mpim-bonn.mpg.de

DEPARTMENT OF COMPUTING, MACQUARIE UNIVERSITY, SYDNEY, NSW 2109, AUSTRALIA

E-mail address: shamin2010@gmail.com