

RESEARCH



# A meromorphic extension of the 3D index

Stavros Garoufalidis<sup>1,2\*</sup> and Rinat Kashaev<sup>3</sup>

\*Correspondence:  
stavros@math.gatech.edu  
http://www.math.gatech.edu/~stavros  
<sup>1</sup>School of Mathematics, Georgia  
Institute of Technology, Atlanta,  
GA 30332-0160, USA  
Full list of author information is  
available at the end of the article  
To Don Zagier, on the occasion  
of his 65th birthday.

## Abstract

Using the locally compact abelian group  $\mathbb{T} \times \mathbb{Z}$ , we assign a meromorphic function to each ideal triangulation of a 3-manifold with torus boundary components. The function is invariant under all 2–3 Pachner moves, and thus is a topological invariant of the underlying manifold. If the ideal triangulation has a strict angle structure, our meromorphic function can be expanded into a Laurent power series whose coefficients are formal power series in  $q$  with integer coefficients that coincide with the 3D index of (Dimofte et al. in *Adv Theor Math Phys* 17(5):975–1076, 2013). Our meromorphic function can be computed explicitly from the matrix of the gluing equations of a triangulation, and we illustrate this with several examples.

**Keywords:** Ideal triangulations, 3-manifolds, 3D index, Locally compact abelian groups, Quantum dilogarithm, State integrals, Meromorphic functions

## Contents

1	Introduction	2
1.1	The 3D index of Dimofte–Gaiotto–Gukov	2
1.2	Our results	2
1.3	Discussion	3
2	Building blocks	4
2.1	The tetrahedral weight	4
2.2	The quantum dilogarithm	6
2.3	The pentagon identity and the Pachner 2–3 move	9
3	The state integral	11
3.1	Definition of the state integral	11
3.2	Identification with the 3D index of Dimofte–Gaiotto–Gukov	16
3.3	Invariance under 2–3 Pachner moves	18
3.4	The singularities of $I_{\mathcal{T}}(q)$	19
4	Examples and computations	20
4.1	A non-1-efficient triangulation with two tetrahedra	20
4.2	The $4_1$ knot	23
4.3	The sister of the $4_1$ knot	24
4.4	The unknot	25
4.5	The trefoil	26
4.6	The $5_2$ knot	26
4.7	The $6_1$ knot	27
	Appendix A: A quantum dilogarithm over the LCA group $\mathbb{T} \times \mathbb{Z}$	28
	Appendix B: The quantum dilogarithm and the Beta pentagon relation	31
	References	34

© The Author(s) 2018. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## 1 Introduction

### 1.1 The 3D index of Dimofte–Gaiotto–Gukov

In a recent breakthrough in mathematical physics, the physicists Dimofte, Gaiotto and Gukov [8, 9] introduced the *3D index*, a powerful new invariant of an ideal triangulation  $\mathcal{T}$  of a compact orientable 3-manifold  $M$  with non-empty boundary consisting of tori. The 3D index was motivated by the study of the low energy limit of a famous 6-dimensional  $(2, 0)$  superconformal field theory, and seems to contain a great deal of information about the geometry and topology of the ambient manifold. For suitable ideal triangulations, the 3D index is a collection of formal Laurent power series in a variable  $q$ , parametrized by a choice of peripheral homology class, i.e., an element of  $H_1(\partial M, \mathbb{Z})$ .

Physics predicts that the 3D index is independent of the triangulation  $\mathcal{T}$  and that it is a topological invariant of the ambient manifold. However, there is a subtlety: The 3D index itself (which is a sum over some  $q$ -series over a lattice) is only defined for suitable triangulations, and it is invariant under 2–3 moves of such triangulations. It is not known whether suitable triangulations are connected under 2–3 moves, and it is known that some 3-manifolds (for instance, the unknot) have no suitable triangulation. It was shown in [10, 12] that a triangulation is suitable if and only if it is 1-efficient, i.e., has no normal surfaces which are topologically 2-spheres or tori. Thus, the connected sum of two nontrivial knots, or the Whitehead double of a nontrivial knot has no 1-efficient triangulations. With some additional work, one can extract from the 3D index a topological invariant of hyperbolic 3-manifolds [12].

This partial success in constructing a topological invariant suggests the existence of an invariant of ideal triangulations unchanged under all 2–3 Pachner moves. The construction of such an invariant is the goal of our paper. Indeed, to any ideal triangulation, we associate an invariant which is a meromorphic function of the peripheral variables, and for triangulations with strict angle structures, the coefficients of its expansion into Laurent series coincides with the 3D index of [8]. Our meromorphic function is an example of a topological invariant associated with the self-dual locally compact abelian group (abbreviated LCA group)  $\mathbb{T} \times \mathbb{Z}$ . A more detailed formulation of our results follows.

In a sense, our paper does the opposite from that of [13]. In the latter paper, we expressed state integral invariants (which are analytic functions in a cut plane) in terms of  $q$ -series, whereas in the present paper we assemble  $q$ -series into meromorphic state-integral invariants. Our work illustrates the principle that some state integrals can be formulated in terms of  $q$ -series and vice versa.

### 1.2 Our results

Fix an ideal triangulation  $\mathcal{T}$  of an oriented 3-manifold  $M$  whose boundary consists of  $r$  tori, and choose peripheral curves that form a basis of  $H_1(\partial M, \mathbb{Z})$ . To simplify notation, we will present our results only in the case when  $M$  has a single torus boundary though our statements and proofs remain valid in the general case.

After a choice of a meridian and longitude, we can identify the complex torus  $\mathbb{T}_M = H^1(\partial M, \mathbb{C}^*)$  with  $(\mathbb{C}^*)^2$  where the latter is given by the coordinates  $(e_\mu, e_\lambda)$ . Throughout the paper,  $q$  will denote a complex number inside the unit disk:  $|q| < 1$ . When  $q = -e^h$  with  $\operatorname{Re}(h) < 0$  and  $z \in \mathbb{C}$ , we define  $(-q)^z = e^{zh}$ . For  $r, s \in \mathbb{Q}$ , we define the associated  $q$ -rays of the complex torus by

$$\Sigma_{r,s} = \{(e_\mu, e_\lambda) \in (\mathbb{C}^*)^2 \mid e_\mu^r e_\lambda^s \in q^{\mathbb{N}}\}, \tag{1}$$

where  $\mathbb{N} = \mathbb{Z}_{\geq 0}$  is the set of nonnegative integers. A shifted  $q$ -ray is a subset of the complex torus of the form  $\varepsilon q^t \Sigma_{r,s}$  for some  $t \in \mathbb{Q}$  and  $\varepsilon = \pm 1$ .

**Theorem 1.1** *With the above assumptions, there exists a meromorphic function*

$$I_{\mathcal{T}}(q): (\mathbb{C}^*)^2 \ni (e_\mu, e_\lambda) \mapsto I_{\mathcal{T},e_\mu,e_\lambda}(q) \in \mathbb{C} \cup \{\infty\}$$

with the following properties:

- (a)  $I_{\mathcal{T}}(q)$  is invariant under 2–3 Pachner moves.
- (b)  $I_{\mathcal{T}}(q)$  is given by a balanced state integral depending only on the Neumann–Zagier matrices of the gluing equations of  $\mathcal{T}$ .
- (c) The singularities of  $I_{\mathcal{T}}(q)$  are contained in the union of finitely many shifted  $q$ -rays.

**Theorem 1.2** *When  $\mathcal{T}$  has a strict angle structure, we have a Laurent series expansion*

$$I_{\mathcal{T},e_\mu,e_\lambda}(q) = \sum_{(m,e) \in \mathbb{Z}^2} e_\mu^m e_\lambda^e I_{\mathcal{T}}(m, e)(q^2) \tag{2}$$

convergent on the unit torus  $|e_\mu| = |e_\lambda| = 1$ , where  $I_{\mathcal{T}}(m, e)$  is the 3D index of [9].

Fix a 3-manifold  $M$  as above and consider the set  $\mathcal{S}_M$  of all ideal triangulations of  $M$  that admit a strict angle structure.

**Corollary 1.3** *Although it is not known yet if  $\mathcal{S}_M$  is connected or not by 2–3 Pachner moves, the 3D index of [9] is constant on  $\mathcal{S}_M$ .*

### 1.3 Discussion

In a series of papers [2–4, 15], topological invariants of (ideally triangulated) 3-manifolds have been constructed from certain self-dual LCA groups equipped with quantum dilogarithm functions. The main idea of those constructions is the following. Fixing a self-dual LCA group with a Gaussian exponential and a quantum dilogarithm function, one assigns a state-integral invariant to an ideal triangulation decorated by a pre-angle structure (in the cited papers this is called *shape structure*) that is a choice of a strict angle structure within each ideal tetrahedron, but the angles do not have to add up to  $2\pi$  around the edges of the ideal triangulation. The resulting state integral is often the germ of a meromorphic function on the (affine vector) space of real-valued pre-angle structures. This affine space has an (affine vector) subspace of complex-valued angle structures (the pre-angle structures that add up to  $2\pi$  around each geometric edge of the triangulation). The above meromorphic function is either infinity or restricts to a meromorphic function on the space of complex-valued angle structures. When the latter happens, the state integral depends only on the peripheral angle monodromy. This way, we obtain an invariant of ideal triangulations which depends on the peripheral angle monodromy.

The above construction is general and, in particular, it applies to the invariants constructed in [2–4] and [17]. Our goal is to give a self-contained presentation in the case of the self-dual LCA group  $\mathbb{T} \times \mathbb{Z}$  with a quantum dilogarithm first found by Woronowicz in [21] and to relate this invariant to the 3D index of Dimofte–Gaiotto–Gukov [8].

## 2 Building blocks

### 2.1 The tetrahedral weight

In this section, we define the tetrahedral weight which is the building block of our state integral. We give a self-contained treatment of the symmetries and identities that it satisfies.

Below, we will often consider expansions of meromorphic functions defined on open annuli or punctured disks, examples of which are given in Eqs. (20), (23), (24). These Laurent expansions (not to be confused with the formal Laurent series which involve only finitely many negative powers and arbitrarily many positive powers) are well known in complex analysis and their existence, convergence and manipulation follows from Cauchy’s theorem. A detailed discussion of this can be found, for example, in [1].

As a warm up, recall the Pochhammer symbol

$$(x; q)_m := \prod_{i=0}^{m-1} (1 - q^i x), \quad m \in \mathbb{N} \cup \{\infty\}, \tag{3}$$

where  $\mathbb{N} := \mathbb{Z}_{\geq 0}$  and we always assume that  $|q| < 1$ . The next lemma summarizes the well-known properties of the Pochhammer symbol  $(x; q)_\infty$ .

**Lemma 2.1** *The Pochhammer symbol  $(x; q)_\infty$  has the following properties.*

- (a) *It is an entire function of  $x$  with simple zeros  $x \in q^{-\mathbb{N}}$ .*
- (b) *It satisfies the  $q$ -difference equation*

$$(x; q)_\infty = (1 - x)(qx; q)_\infty. \tag{4}$$

- (c) *It has convergent power series expansions*

$$\frac{1}{(x; q)_\infty} = \sum_{n=0}^{\infty} \frac{x^n}{(q; q)_n}, \quad |x| < 1, \tag{5a}$$

$$(x; q)_\infty = \sum_{n=0}^{\infty} (-1)^n \frac{q^{\frac{1}{2}n(n-1)} x^n}{(q; q)_n}, \quad \forall x \in \mathbb{C}. \tag{5b}$$

For the proof of part (c), see for instance [22, Prop. 2].

Consider the function

$$G_q(z) = \frac{(-qz^{-1}; q)_\infty}{(z; q)_\infty} = \frac{\left(1 + \frac{q}{z}\right) \left(1 + \frac{q^2}{z}\right) \left(1 + \frac{q^3}{z}\right) \dots}{(1 - z)(1 - qz)(1 - q^2z) \dots}. \tag{6}$$

The next lemma summarizes its properties.

**Lemma 2.2** *The function  $G_q(z)$  defined in (6) has the following properties.*

- (a) *It is a meromorphic function of  $z \in \mathbb{C}^* := \mathbb{C} \setminus \{0\}$  with simple zeros and poles in  $-q^{1+\mathbb{N}}$  and  $q^{-\mathbb{N}}$ , respectively, and with essential singularities at  $z = 0$  and  $z = \infty$ .*
- (b) *It satisfies the  $q$ -difference equation*

$$G_q(qz) = (1 - z)(1 + z^{-1})G_q(z) \tag{7}$$

and the involution equation

$$G_q(-qz) = \frac{1}{G_q(z^{-1})}. \tag{8}$$

(c) It has a convergent Laurent series expansion in the punctured unit disk  $0 < |z| < 1$ :

$$G_q(z) = \sum_{n \in \mathbb{Z}} J(n)(q)z^n, \tag{9}$$

where

$$J(n)(q) := \sum_{k=(-n)_+}^{\infty} \frac{q^{\frac{k(k+1)}{2}}}{(q)_k(q)_{n+k}}, \quad (n)_+ := \max\{n, 0\}, \tag{10}$$

is a well-defined element of  $\mathbb{Z}[[q]]$ , analytic in the disk  $|q| < 1$ .

Parts (a) and (b) follow easily from the product expansion of the Pochhammer symbol, and part (c) follows from (5a)–(5b).

The tetrahedral weight is a function  $\psi^0(z, w)$  defined by:

$$\psi^0(z, w) := c(q) G_q(-qz)G_q(w^{-1})G_q(wz^{-1}), \tag{11}$$

where

$$c(q) := \frac{(q; q)_{\infty}^2}{(q^2; q^2)_{\infty}}. \tag{12}$$

The properties of this function are summarized in the following lemma.

**Lemma 2.3** *The function  $\psi^0(z, w)$  defined in (11) has the following properties.*

(a) It is a meromorphic function of  $(z, w) \in (\mathbb{C}^*)^2$  with zeros in

$$z \in q^{\mathbb{N}}, \quad \text{or} \quad w^{-1} \in -q^{1+\mathbb{N}}, \quad \text{or} \quad z^{-1}w \in -q^{1+\mathbb{N}} \tag{13}$$

and poles in

$$z \in -q^{-1-\mathbb{N}}, \quad \text{or} \quad w \in q^{\mathbb{N}}, \quad \text{or} \quad zw^{-1} \in q^{\mathbb{N}}. \tag{14}$$

(b) It satisfies the  $q$ -difference equations

$$w^{-1}\psi^0(qz, w) - \psi^0(z, w) - q^{-1}z^{-1}\psi^0(z, w) = 0, \tag{15a}$$

$$w\psi^0(qz, w) - \psi^0(z, w) - qz\psi^0(z, w) = 0. \tag{15b}$$

(c) It satisfies the  $\mathbb{Z}/2$  and  $\mathbb{Z}/3$ -invariance equations

$$\psi^0(z, w) = \psi^0(-q^{-1}w^{-1}, -q^{-1}z^{-1}), \tag{16a}$$

$$\psi^0(z, w) = \psi^0(-q^{-1}z^{-1}w, -q^{-1}z^{-1}) = \psi^0(-q^{-1}w^{-1}, zw^{-1}). \tag{16b}$$

(d) *In the domain*

$$1 < |w| < |z| < |q|^{-1}, \tag{17}$$

*we have the absolutely convergent expansion*

$$\psi^0(z, w) = c(q) \sum_{e, m \in \mathbb{Z}} z^e w^m \sum_{\substack{k_1, k_2, k_3 \in \mathbb{Z} \\ k_1 - k_3 = e; k_3 - k_2 = m}} (-q)^{k_1} J(k_1)(q) J(k_2)(q) J(k_3)(q), \tag{18}$$

*where the interior sum is a well-defined element of  $\mathbb{Z}[[q]]$ , analytic in the disk  $|q| < 1$ .*

These properties follow from the definition of  $\psi^0$  and the properties of  $G_q$  listed in Lemma 2.2.

### 2.2 The quantum dilogarithm

In this subsection, we identify the tetrahedral weight function  $\psi^0(z, w)$  with (the reciprocal of) the quantum dilogarithm function  $\psi(z, m)$  on the self-dual LCA group  $\mathbb{T} \times \mathbb{Z}$ , given by [15, Eqn.97]

$$\psi(z, m) = \frac{(-q^{1-m}/z; q^2)_\infty}{(-q^{1-m}z; q^2)_\infty}. \tag{19}$$

In the context of quantum groups, this function first appeared in [21]. In Appendix A, we explain how formula (19) fits the general definition of a quantum dilogarithm over the LCA group  $\mathbb{T} \times \mathbb{Z}$ .

Lemma 2.1 and the above definition imply the following properties of the function  $\psi(z, w)$ .

**Lemma 2.4** *The function  $\psi(z, m)$  defined in (19) has the following properties.*

- (a) *It is a meromorphic function of  $z$  with simple poles and zeros at  $z \in -q^{-1-|m|-2\mathbb{N}}$  and  $z \in -q^{1+|m|+2\mathbb{N}}$ , respectively.*
- (b) *It is analytic in the annulus  $0 < |z| < |q|^{-1-|m|}$  (which always includes the unit circle  $|z| = 1$ ) where it has an absolutely convergent Laurent series expansion*

$$\psi(z, m) = \sum_{e \in \mathbb{Z}} I^\Delta(m, e)(q) z^e, \tag{20}$$

*where*

$$I^\Delta(m, e)(q) = (-q)^e I_\Delta(m, e)(q^2) \tag{21}$$

*is related to the tetrahedron index  $I_\Delta$  of [8] given by*

$$I_\Delta(m, e)(q) = \sum_{n=(-e)_+}^{\infty} (-1)^n q^{\frac{1}{2}n(n+1) - (n+\frac{1}{2}e)m} \frac{1}{(q)_n (q)_{n+e}} \in \mathbb{Z}[[q^{\frac{1}{2}}]] \tag{22}$$

*and  $(e)_+ := \max\{0, e\}$  and  $(q)_n := (q; q)_n = \prod_{i=1}^n (1 - q^i)$ .*

The next theorem connects the tetrahedral weight function  $\psi^0(z, w)$  with the above function  $\psi(z, m)$ .

**Theorem 2.5** (a) *In the domain (17), we have the identity*

$$\psi^0(z, w) = \sum_{m \in \mathbb{Z}} \psi(z, m)w^m, \tag{23}$$

where the sum is absolutely convergent.

(b) *In the domain (17), we have an absolutely convergent double Laurent series expansion*

$$\psi^0(z, w) = \sum_{e, m \in \mathbb{Z}} I^\Delta(m, e)(q)z^e w^m. \tag{24}$$

*Proof* We let RHS denote the sum in the right-hand side of Eq. (23). RHS is absolutely convergent in the domain (17) and it can be explicitly calculated by using Ramanujan’s  ${}_1\psi_1$ -summation formula. The detailed computation appears in [15, Eqn. (98)], and the result reads

$$\text{RHS} = \frac{(q^2; q^2)_\infty (z^{-2}; q^2)_\infty}{(w^{-2}; q^2)_\infty (w^2 z^{-2}; q^2)_\infty} \left( \frac{\theta_q(zw^{-2})}{\theta_q(z)} + w \frac{\theta_q(qw^2/z)}{\theta_q(q/z)} \right), \tag{25}$$

where

$$\theta_q(x) := \sum_{k \in \mathbb{Z}} q^{k^2} x^k = (q^2; q^2)_\infty (-qx; q^2)_\infty (-q/x; q^2)_\infty \tag{26}$$

is the Jacobi theta function, and the second equality in (26) is the Jacobi triple product identity. By using Lemma 2.6 (see below) and the Jacobi triple product identity, we can further simplify the right-hand side of (25), thus getting

$$\begin{aligned} \text{RHS} &= \frac{(q; q)_\infty^2 (z^{-1}; q)_\infty (-qw; q)_\infty (-qz/w; q)_\infty}{(q^2; q^2)_\infty (-qz; q)_\infty (w^{-1}; q)_\infty (w/z; q)_\infty} \\ &= c(q) G_q(-qz) G_q(w^{-1}) G_q(wz^{-1}) \\ &= \psi^0(z, w). \end{aligned}$$

This concludes the proof of the part (a). Part (b) follows from (23) combined with (20).  $\square$

**Lemma 2.6** *For any choice of the square root  $p := \sqrt{q}$ , we have the identity*

$$\frac{\theta_q(zw^{-2})}{\theta_q(z)} + w \frac{\theta_q(qw^2/z)}{\theta_q(q/z)} = \frac{(q; q)_\infty}{(q^2; q^2)_\infty} \frac{\theta_p(pw/z) \theta_p(w/p)}{\theta_p(z/p)}. \tag{27}$$

*Proof* Denoting the left-hand side of (27) as  $f(z^{-1}, w)$ , we have

$$f(u, w) = h(u, w)/g(u),$$

where

$$g(u) := \theta_q(u)\theta_q(qu), \quad h(u, w) := \theta_q(qu)\theta_q(uw^2) + w\theta_q(u)\theta_q(quw^2).$$

First, we rewrite  $g(u)$  as follows:

$$\begin{aligned} g(u) &= (q^2; q^2)_\infty^2 (-qu; q^2)_\infty (-q^2u; q^2)_\infty (-q/u; q^2)_\infty (-1/u; q^2)_\infty \\ &= (q^2; q^2)_\infty^2 (-qu; q)_\infty (-1/u; q)_\infty = \frac{(q^2; q^2)_\infty^2}{(q; q)_\infty} \theta_p(pu). \end{aligned} \tag{28}$$

Then, we transform  $h(u/w, w)$  as follows:

$$\begin{aligned} h\left(\frac{u}{w}, w\right) &= \sum_{k, l \in \mathbb{Z}} q^{k^2+l^2} u^{k+l} w^{l-k} (q^k + q^l w) \\ &= \sum_{k, l \in \mathbb{Z}} q^{k^2-2kl+2l^2} u^k w^{2l-k} (q^{k-l} + q^l w), \end{aligned} \tag{29}$$

where in the second equality we have shifted the summation variable  $k \mapsto k - l$ . Next, we write out separately the sum over even and odd  $k$ :

$$\begin{aligned} h\left(\frac{u}{w}, w\right) &= \sum_{k, l \in \mathbb{Z}} q^{2k^2+2(l-k)^2} u^{2k} w^{2l-2k} \\ &\quad \times \left( q^{2k-l} + q^l w + q^{4k+1-2l} u w^{-1} (q^{2k+1-l} + q^l w) \right) \\ &= \sum_{k, l \in \mathbb{Z}} q^{2k^2+2l^2+k} u^{2k} w^{2l} \left( q^{-l} + q^l w + q^{2k+1-2l} u w^{-1} (q^{1-l} + q^l w) \right), \end{aligned} \tag{30}$$

where, this time, in the second equality we have shifted the summation variable  $l \mapsto l + k$ . Now, we can absorb both summations by using the definition of the  $\theta$ -function:

$$\begin{aligned} h\left(\frac{u}{w}, w\right) &= \theta_{q^2}(qu^2) (\theta_{q^2}(q^{-1}w^2) + w\theta_{q^2}(qw^2)) \\ &\quad + qu\theta_{q^2}(q^3u^2) (qw^{-1}\theta_{q^2}(q^{-3}w^2) + \theta_{q^2}(q^{-1}w^2)). \end{aligned} \tag{31}$$

Finally, by using the functional equation

$$\theta_q(q^{2l}x) = q^{-l^2} x^{-l} \theta_q(x), \quad \text{for all } l \in \mathbb{Z}, \tag{32}$$

with  $q \mapsto q^2, x = w^2$  and  $l = -1$  in the second term, we arrive at the following factorized formula

$$h\left(\frac{u}{w}, w\right) = (\theta_{q^2}(qu^2) + qu\theta_{q^2}(q^3u^2)) (\theta_{q^2}(q^{-1}w^2) + w\theta_{q^2}(qw^2)) = s(qu)s(w), \tag{33}$$

where

$$\begin{aligned} s(w) &:= \theta_{q^2}(q^{-1}w^2) + w\theta_{q^2}(qw^2) = \sum_{k \in \mathbb{Z}} q^{2k^2} w^{2k} (q^{-k} + q^k w) \\ &= \sum_{k \in \mathbb{Z}} p^{4k^2} w^{2k} (p^{-2k} + p^{2k} w) = \sum_{k \in \mathbb{Z}} (p^{(2k)^2-2k} w^{2k} + p^{(2k+1)^2-2k-1} w^{2k+1}) \\ &= \sum_{k \in \mathbb{Z}} p^{k^2-k} w^k = \theta_p(w/p). \end{aligned} \tag{34}$$

Thus, we have obtained the equality

$$h(u, w) = \theta_p(puw)\theta_p(w/p), \tag{35}$$

and formula (27) follows straightforwardly. □



In summary, the tetrahedral weight  $\psi^0(z, w)$  is given by two sum formulas and a product formula, and the last of which implies the meromorphicity of the function and the location of its zeros and poles:

$$\psi^0(z, w) = \sum_{m \in \mathbb{Z}} \psi(z, m)w^m \tag{36}$$

$$= \sum_{e, m \in \mathbb{Z}} I^\Delta(m, e)(q)z^e w^m \tag{37}$$

$$= c(q) G_q(-qz)G_q(w^{-1})G_q(wz^{-1}). \tag{38}$$

For completeness, the next lemma summarizes the  $q$ -difference equations and the symmetries of  $\psi(z, w)$ .

**Lemma 2.7** *The function  $\psi(z, m)$  defined in (19) satisfies the following  $q$ -difference equations*

$$\psi(qz, m + 1) - \psi(z, m) - q^{-m-1}z^{-1}\psi(z, m) = 0, \tag{39a}$$

$$\psi(qz, m - 1) - \psi(z, m) - q^{-m+1}z\psi(z, m) = 0. \tag{39b}$$

*The above equations characterize the meromorphic function  $\psi(z, m)$  up to multiplication by a function of  $q$ , analytic in the unit disk  $|q| < 1$ .*

These follow from the definition of  $\psi(z, w)$  and Eqs. (4), (23) and (15a)–(15b). Alternatively, they can be derived from Eq. (20) and the symmetries of the tetrahedron index  $I_\Delta$  given in [10, Thm.3.2].

For completeness, the next lemma summarizes the symmetries of  $I^\Delta(m, e)$ .

**Lemma 2.8** *For all integers  $m$  and  $e$ , we have the  $\mathbb{Z}/2$  and  $\mathbb{Z}/3$ -invariance equations:*

$$I^\Delta(m, e)(q) = (-q)^{e+m}I^\Delta(-e, -m)(q), \tag{40a}$$

$$I^\Delta(m, e)(q) = (-q)^e I^\Delta(-e - m, m)(q) = (-q)^{e+m}I^\Delta(e, -e - m)(q). \tag{40b}$$

*As a consequence, we have another  $\mathbb{Z}/2$ -invariance equation*

$$I^\Delta(m, e) = I^\Delta(-m, m + e). \tag{41}$$

These follow from Eqs. (24) and (16a)–(16b). Additionally, they follow from Eqs. (21) and the symmetries of the tetrahedron index  $I_\Delta(m, e)$  given in [10, Thm.3.2].

### 2.3 The pentagon identity and the Pachner 2–3 move

Let us recall the pentagon identity for the tetrahedron index  $I_\Delta$  from [10, Thm.3.7].

$$I_\Delta(m_1 - e_2, e_1)I_\Delta(m_2 - e_1, e_2) = \sum_{e_3 \in \mathbb{Z}} q^{e_3} I_\Delta(m_1, e_1 + e_3)I_\Delta(m_2, e_2 + e_3)I_\Delta(m_1 + m_2, e_3) \tag{42}$$

for all integers  $m_1, m_2, e_1, e_2$ . Replacing  $q$  by  $q^2$  in (42) and using Eq. (21), it follows that the tetrahedron index  $I^\Delta$  satisfies the equation

$$\begin{aligned}
 & I^\Delta(m_1 - e_2, e_1)I^\Delta(m_2 - e_1, e_2) \\
 &= \sum_{e_3 \in \mathbb{Z}} (-q)^{-e_3} I^\Delta(m_1, e_1 + e_3)I^\Delta(m_2, e_2 + e_3)I^\Delta(m_1 + m_2, e_3). \tag{43}
 \end{aligned}$$

To remove the factor  $(-q)^{e_3}$  in the above equation, we apply Eq. (40b) to the two terms of the left-hand side and to the last term in the right-hand side, and obtain

$$\begin{aligned}
 & I^\Delta(e_1, -e_1 + e_2 - m_1)I^\Delta(e_2, e_1 - e_2 - m_2) \\
 &= \sum_{e_3 \in \mathbb{Z}} I^\Delta(m_1, e_1 + e_3)I^\Delta(m_2, e_2 + e_3)I^\Delta(e_3, -e_3 - m_1 - m_2).
 \end{aligned}$$

Setting

$$x = e_1, \quad y = -e_1 + e_2 - m_1, \quad u = e_2, \quad v = e_1 - e_2 - m_2, \quad e_3 = -z,$$

we rewrite the latter equality as

$$I^\Delta(x, y)I^\Delta(u, v) = \sum_{z \in \mathbb{Z}} I^\Delta(u - x - y, x - z)I^\Delta(-z, z + v + y)I^\Delta(-u - v + x, u - z).$$

Applying Eq. (41) to both functions on the left-hand side of the above equation, and after a linear change in variables we obtain

$$\begin{aligned}
 I^\Delta(x, y)I^\Delta(u, v) &= \sum_{z \in \mathbb{Z}} I^\Delta(-u - y, -x - z) \\
 &\quad \times I^\Delta(-z, z + x + y + u + v)I^\Delta(-x - v, -u - z).
 \end{aligned}$$

Applying Eq. (41) to the first and third terms of the right-hand side of the above equation, we obtain

$$\begin{aligned}
 I^\Delta(x, y)I^\Delta(u, v) &= \sum_{z \in \mathbb{Z}} I^\Delta(u + y, -x - y - u - z)I^\Delta(-z, z + x + y + u + v) \\
 &\quad \times I^\Delta(x + v, -x - u - v - z).
 \end{aligned}$$

Finally, changing the summation variable  $z \mapsto z - x - y - u - v$ , we obtain equation

$$I^\Delta(x, y)I^\Delta(u, v) = \sum_{z \in \mathbb{Z}} I^\Delta(u + y, v - z)I^\Delta(x + y + u + v - z, z)I^\Delta(x + v, y - z), \tag{44}$$

which coincides with a special (constant) form of the beta pentagon equation [16, Eqn.(2)] for the LCA group  $\mathbb{Z}$ . Using the fact that the beta pentagon relation is stable under Fourier transformation, see [16, Sec.2], we also conclude that the function


$$\phi(x, y) := \sum_{m, n \in \mathbb{Z}} x^m y^{-n} I^\Delta(m, n) = \psi^0(x, 1/y) \tag{45}$$

satisfies the constant beta pentagon equation for the circle LCA group  $\mathbb{T} = \{z \in \mathbb{C} \mid |z| = 1\}$ :

$$\phi(x, y)\phi(u, v) = \int_{\mathbb{T}} \phi(uy, v/z)\phi(xyuv/z, z)\phi(xv, y/z) \frac{dz}{2\pi iz}. \tag{46}$$

Equivalently, the function  $\psi^0(z, w)$ , thought of as a distribution on  $\mathbb{T}^2$ , satisfies the integral identity

$$\psi^0(x, y)\psi^0(u, v) = \int_{\mathbb{T}} \psi^0\left(\frac{u}{y}, \frac{v}{z}\right)\psi^0\left(\frac{xuz}{yv}, z\right)\psi^0\left(\frac{x}{v}, \frac{y}{z}\right) \frac{dz}{2\pi iz}, \tag{47}$$



**Fig. 1** The angles of an ideal tetrahedron with ordered vertices

where all variables belong to the unit circle  $\mathbb{T}$ . The distributional interpretation of  $\psi^0(z, w)$  means that its restriction to  $\mathbb{T}^2$  should be obtained by approaching  $\mathbb{T}^2$  from the domain (17), which is the domain of absolute convergence of the double series (24), i.e.,

$$\psi^0(z, w) = \psi_{+0,+0}^0(z, w) := \lim_{\alpha, \gamma \rightarrow 0} \psi_{\alpha, \gamma}^0(z, w), \quad |z| = |w| = 1, \tag{48}$$

$\alpha > 0, \gamma > 0, \alpha + \gamma < \pi$

where

$$\psi_{\alpha, \gamma}^0(z, w) := \psi^0\left(z(-q)^{-\frac{\alpha+\gamma}{\pi}}, w(-q)^{-\frac{\alpha}{\pi}}\right) \tag{49a}$$

$$= c(q) G_q(z(-q)^{\frac{\beta}{\pi}}) G_q(w^{-1}(-q)^{\frac{\alpha}{\pi}}) G_q(z^{-1}w(-q)^{\frac{\gamma}{\pi}}), \tag{49b}$$

where  $\alpha + \beta + \gamma = \pi$ . The positive parameters  $\alpha, \beta, \gamma$  satisfying the condition  $\alpha + \beta + \gamma = \pi$  (which we call pre-angle structure) can be identified with the dihedral angles of a positively oriented ideal hyperbolic tetrahedron. These angles are placed on the edges of a tetrahedron with ordered vertices according to Fig. 1.

Moreover, the constant distributional beta pentagon identity (47) is a constant limit of the analytically continued non-constant identity

$$\psi_{\alpha_3, \gamma_3}^0(x, y) \psi_{\alpha_1, \gamma_1}^0(u, v) = \int_{\mathbb{T}} \psi_{\alpha_0, \gamma_0}^0\left(\frac{u}{y}, \frac{v}{z}\right) \psi_{\alpha_2, \gamma_2}^0\left(\frac{xuz}{yv}, z\right) \psi_{\alpha_4, \gamma_4}^0\left(\frac{x}{v}, \frac{y}{z}\right) \frac{dz}{2\pi iz}, \tag{50}$$

where  $(\alpha_i, \beta_i, \gamma_i)$  are pre-angle structures on five tetrahedra  $T_i$  for  $i = 0, \dots, 4$  which are compatible, i.e., satisfy the linear relations

$$\alpha_1 = \alpha_0 + \alpha_2, \quad \alpha_3 = \alpha_2 + \alpha_4, \quad \gamma_1 = \gamma_0 + \alpha_4, \quad \gamma_3 = \alpha_0 + \gamma_4, \quad \gamma_2 = \gamma_1 + \gamma_3. \tag{51}$$

Notice that a compatible angle structure satisfies the balancing condition for the interior edge 13:  $\beta_0 + \gamma_2 + \beta_4 = 2\pi$ . Such an identity for Faddeev’s quantum dilogarithm appeared in [3, Prop.1].

### 3 The state integral

#### 3.1 Definition of the state integral

Fix an ideal triangulation  $\mathcal{T}$  of an oriented 3-manifold with  $N$  tetrahedra  $T_i$  for  $i = 1, \dots, N$ . The invariant is defined as follows:

- (a) Assign variables  $x_i$  for  $i = 1, \dots, N$  to  $N$  edges of  $\mathcal{T}$ .
- (b) Choose a strictly positive pre-angle structure  $\theta = (\alpha, \beta, \gamma)$  at each tetrahedron. Here,  $\alpha$  is the angle of the 01 and 23 edges,  $\beta$  is the angle of the 02 and 13 edges, and  $\gamma$  is the angle of the 03 and 12 edges. The angles are normalized so that at each tetrahedron, their sum is  $\pi$ .
- (c) The weight of a tetrahedron  $T$  is given by

$$B(T, x, \theta) = \psi^0 \left( (-q)^{-\frac{\alpha+\gamma}{\pi}} \frac{X_\alpha}{X_\gamma}, (-q)^{-\frac{\alpha}{\pi}} \frac{X_\beta}{X_\gamma} \right) \tag{52a}$$

$$= c(q) G_q \left( (-q)^{\frac{\alpha}{\pi}} \frac{X_\gamma}{X_\beta} \right) G_q \left( (-q)^{\frac{\beta}{\pi}} \frac{X_\alpha}{X_\gamma} \right) G_q \left( (-q)^{\frac{\gamma}{\pi}} \frac{X_\beta}{X_\alpha} \right), \tag{52b}$$

where

$$X_\alpha := x_{01}x_{23}, \quad X_\beta := x_{02}x_{13}, \quad X_\gamma := x_{03}x_{12},$$

and  $x_{ij}$  is the variable at edge  $ij$  of the tetrahedron.

- (d) Define

$$I_{\mathcal{T},\theta}^{\text{pre}}(q) = \int_{\mathbb{T}^N} \prod_{i=1}^N B(T_i, x, \theta) d\mu(x), \tag{53}$$

where  $d\mu(x) = (2\pi i)^{-N} \prod_{i=1}^N dx_i/x_i$  is the normalized Haar measure on  $\mathbb{T}^N$ .

Recall the exponent matrices  $(\overline{A}|\overline{B}|\overline{C})$  of the edge gluing equations of  $\mathcal{T}$  [6, 19, 20]. These are  $N \times N$  matrices with integer entries which determine the gluing equations

$$\prod_{j=1}^N z_j^{\overline{A}_{i,j}} (z'_j)^{\overline{B}_{i,j}} (z''_j)^{\overline{C}_{i,j}} = 1, \quad i = 1, \dots, N, \tag{54}$$

where  $z' = 1/(1 - z)$  and  $z'' = 1 - 1/z$ , and as usual we have  $zz'z'' = -1$ .

For a vector  $x = (x_1, \dots, x_N)$  of nonzero complex numbers, and a vector  $v = (v_1, \dots, v_N)$  of integers, define  $x^v = \prod_{i=1}^N x_i^{v_i}$ . Also, for a matrix  $A$ , let  $A_i$  denote its  $i$ th column.

The next proposition implies that the integral (53) (and even the integrand) depends on only the Neumann–Zagier matrices of the gluing equations of the triangulation  $\mathcal{T}$ .

**Proposition 3.1** *With the above notation and for  $i = 1, \dots, N$ , we have:*

$$B(T_i, x, \theta) = c(q) G_q \left( (-q)^{\frac{\alpha_i}{\pi}} x^{(\overline{C}-\overline{B})_i} \right) G_q \left( (-q)^{\frac{\beta_i}{\pi}} x^{(\overline{A}-\overline{C})_i} \right) G_q \left( (-q)^{\frac{\gamma_i}{\pi}} x^{(\overline{B}-\overline{A})_i} \right). \tag{55}$$

*It follows that  $I_{\mathcal{T},\theta}(q)$  depends on only the matrices  $\overline{A}, \overline{B}, \overline{C}$  and  $\theta$ .*

*Proof* Let  $e_i \in \mathbb{Z}^N$  denote the  $i$ th coordinate vector (1 in position  $i$  and 0 otherwise). It suffices to show that

$$\left( \frac{X_\gamma}{X_\beta} \right)_i = x^{(\overline{C}-\overline{B})e_i}, \quad \left( \frac{X_\alpha}{X_\gamma} \right)_i = x^{(\overline{A}-\overline{C})e_i}, \quad \left( \frac{X_\beta}{X_\alpha} \right)_i = x^{(\overline{B}-\overline{A})e_i}. \tag{56}$$

This follows from the fact that the matrices  $\overline{A}, \overline{B}$  and  $\overline{C}$  indexed by edges  $\times$  tetrahedra record the number of times a shape  $z_j$  (resp.,  $z'_j, z''_j$ ) of the  $j$ th tetrahedron appears around an edge  $e_i$ . □

Our choice of  $\theta$  and Eq. (17) implies that the integrand in (53) is an analytic function of  $x \in \mathbb{T}^N$ , and therefore the integral converges. We are interested in two affine vector subspaces of  $\mathbb{C}^{3N}$ :

- $\mathcal{A}_{\mathcal{T}}$ , the space of complexified pre-angle structures, i.e.,  $\theta \in \mathbb{C}^{3N}$  such that  $\alpha_i + \beta_i + \gamma_i = \pi$  for  $i = 1, \dots, N$ .
- $\mathcal{B}_{\mathcal{T}} \subset \mathcal{A}_{\mathcal{T}}$ , the affine subspace of  $\mathcal{A}_{\mathcal{T}}$  that consists of balanced complexified pre-angle structures, that is the sum of the angles around each edge of  $\mathcal{T}$  is  $2\pi$ . The points of  $\mathcal{B}$  are also known as complex-valued angle structures on  $\mathcal{T}$ .

$\mathcal{A}_{\mathcal{T}}$  and  $\mathcal{B}_{\mathcal{T}}$  are complex affine subspaces of  $\mathbb{C}^{3N}$  of dimension  $2N$  and  $N + 1$ , respectively, [18].

The integral in (53) extends to a meromorphic function of  $\theta \in \mathcal{A}_{\mathcal{T}}$ , regular when  $\text{Re}(\theta) > 0$ . Our task is to show that this extension restricts to a meromorphic function on  $\mathcal{B}_{\mathcal{T}}$ . To do so, it will be convenient to parametrize  $\mathcal{A}_{\mathcal{T}}$ . This breaks the symmetry in the definition of the integral; however, it is a useful gauge to draw conclusions.

Consider a vector  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_N) \in \mathbb{C}^N$  and complex numbers  $\mu, \lambda$  defined by

$$\begin{aligned} \bar{A}\alpha + \bar{B}\beta + \bar{C}\gamma = \pi\mathbf{2} + \varepsilon, & \quad v_{\mu} \cdot \alpha + v'_{\mu} \cdot \beta + v''_{\mu} \cdot \gamma = \mu, \\ v_{\lambda} \cdot \alpha + v'_{\lambda} \cdot \beta + v''_{\lambda} \cdot \gamma = \lambda, & \end{aligned} \tag{57}$$

where  $\mu$  and  $\lambda$  are the sums of the angles along the meridian and the longitude curves,  $\mathbf{2} \in \mathbb{C}^N$  is the vector with all coordinates 2 and  $v_{\mu}, v_{\lambda}$  and their primed versions are the vectors of the meridian and longitude cusp equations. Note that  $\varepsilon_1 + \dots + \varepsilon_N = 0$ . A *quad* is a choice of a pair of opposite edges in a tetrahedron. Choosing a quad allows one to eliminate the angle variable of those edges using the equation  $\alpha_i + \beta_i + \gamma_i = \pi$ . Note that each tetrahedron has 3 quads, and hence  $\mathcal{T}$  has  $3^N$  quads. If  $Q$  is a system of  $N$  quads obtained by choosing one quad from each tetrahedron of  $\mathcal{T}$ , let  $Q(\theta) \in \mathbb{C}^N$  be defined so that its  $i$ th component is the angle associated with the corresponding quad in  $T_i$ , i.e.,  $Q(\theta)_i = \alpha_i$  if, for example, the quad of the  $i$ th tetrahedron is (01), (23). In effect,  $Q(\theta)$  chooses one of the 3 angles for every tetrahedron of  $\mathcal{T}$ . Together with Eqs. (57), we get a linear map

$$\mathcal{A}_{\mathcal{T}} \rightarrow \mathbb{C}^N \times \mathbb{C}^N \times \mathbb{C}^2, \quad \theta \mapsto (Q(\theta), \varepsilon, \mu, \lambda).$$

For  $i, j \in \{1, \dots, N\}$ , let  $\pi_{ij} : \mathbb{C}^N \times \mathbb{C}^N \times \mathbb{C}^2 \rightarrow \mathbb{C}^{N-1} \times \mathbb{C}^{N-1} \times \mathbb{C}^2$  denote the projection that removes the  $i$ th entry of the first copy of  $\mathbb{C}^N$  and the  $j$ th entry of the second copy of  $\mathbb{C}^N$ . The next proposition describes a parametrization of  $\mathcal{A}_{\mathcal{T}}$ .

**Proposition 3.2** *For every  $j$ , there exists a system of quads  $Q$  of  $\mathcal{T}$  and an  $i$  such that the composition  $\mathcal{A}_{\mathcal{T}} \rightarrow \mathbb{C}^N \times \mathbb{C}^N \times \mathbb{C}^2 \xrightarrow{\pi_{ij}} \mathbb{C}^{N-1} \times \mathbb{C}^{N-1} \times \mathbb{C}^2$  is an affine linear isomorphism.*

*Proof* Consider the standard system of quads  $Q$  of  $\mathcal{T}$ , i.e., the one that chooses the (01), (23) edges of each tetrahedron of  $\mathcal{T}$ . It follows that  $Q(\theta) = \alpha$ . Eliminating  $\beta$  from Eq. (57), we obtain that

$$A\alpha + B\gamma = \pi\mathbf{2} + \varepsilon + v, \tag{58}$$

where

$$A = \bar{A} - \bar{B}, \quad B = \bar{C} - \bar{B}, \quad v = -\bar{B}\mathbf{1}\pi.$$

Consider the matrix  $(A'|B')$  obtained from  $(A|B)$  by replacing the  $j$ th row of  $(A|B)$  with the peripheral cusp equation corresponding to the meridian. Neumann–Zagier prove that  $(A'|B')$  is the upper half of a symplectic  $2N \times 2N$  matrix [19]. In fact, we can take the first row of the bottom half of the symplectic matrix to be the peripheral cusp equation corresponding to the longitude. If  $B'$  is invertible, then we can solve for  $\gamma$  from Eq. (58) and deduce that  $\theta$  is determined by  $(\alpha, \pi_j(\varepsilon), \mu)$ . Using the longitude cusp equation, it follows that  $\theta$  is determined by  $(\pi_i(\alpha), \pi_j(\varepsilon), \mu, \lambda)$ .

When  $B'$  is not invertible, using the fact that  $(A'|B')$  is the upper half of a symplectic matrix and [7, Lem.A.3], it follows that we can always find a system of quads  $Q$  for which the corresponding matrix  $B'$  is invertible. The result follows.  $\square$

Without loss of generality, we can assume that Proposition 3.2 holds for the standard system of quads, and that  $i = j = N$ . After a change in variables  $x_i \rightarrow x_i/x_N$  for  $i = 1, \dots, N - 1$ , the integral  $I_{T,\theta}(q)$  reduces to an  $N - 1$ -dimensional integral since the integrand is independent of  $x_N$ .

The next lemma shows that after a change in variables,  $I_{T,\theta}(q)$  is expressed as an integral whose contour (a product of tori, with radi  $|q|$  raised to linear forms of  $\alpha$ ) depends only on  $\alpha$  and whose integrand depends only on  $(\varepsilon, \mu, \lambda)$ .

**Proposition 3.3** *With the above assumptions, there exists an edge column vector  $\eta$  whose coordinates are affine linear forms in  $\alpha$ , a contour  $C_\alpha$ , and affine linear forms  $v_i, v'_i, v''_i$  in the variables  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_{N-1})$  and  $\mu, \lambda$  such that after the change in variables  $x_i = (-q)^{n_i}y_i$ , we have*

$$\begin{aligned} I_{T,\theta}^{\text{pre}}(q) &= c(q)^N \int_{C_\alpha} d\mu(y) \\ &\times \prod_{i=1}^N G_q \left( (-q)^{v_i(\varepsilon,\mu,\lambda)} y^{(\bar{C}-\bar{B})_i} \right) \\ &\times G_q \left( (-q)^{v'_i(\varepsilon,\mu,\lambda)} y^{(\bar{A}-\bar{C})_i} \right) \\ &\times G_q \left( (-q)^{v''_i(\varepsilon,\mu,\lambda)} y^{(\bar{B}-\bar{A})_i} \right). \end{aligned} \tag{59}$$

*Proof* Recall Eqs. (56) and (58). If  $x = (-q)^n y$  then

$$x^{(\bar{C}-\bar{B})e_i} = (-q)^{\alpha_i + e_i^T (\bar{C}-\bar{B})^T \eta} y^{(\bar{C}-\bar{B})e_i} = (-q)^{\alpha_i + e_i^T B^T \eta} y^{Be_i}$$

and likewise for the cyclic permutations. Using Eq. (55), and the above equalities, we are looking for an edge vector  $\eta$  such that  $\alpha + B^T \eta$ ,  $\beta + (A - B)^T \eta$  and  $\gamma - A^T \eta$  depend on only  $\varepsilon, \mu, \lambda$ . Since the sum of these three vectors is constant, it suffices to find  $\eta$  such that  $\alpha - B^T \eta$  and  $\gamma - A^T \eta$  depend on only  $\varepsilon, \mu, \lambda$ . Using the fact that  $B'$  is invertible, we can take  $\eta = (B'^T)^{-1} \alpha$ . Solving for  $\gamma$  from Eq. (58), we obtain that

$$\gamma - A'^T \eta = B'^{-1}(\pi \mathbf{2} + \varepsilon + v) - (B')^{-1} A' \alpha - (A')^T ((B')^T)^{-1} \alpha = B'^{-1}(\pi \mathbf{2} + \varepsilon + v).$$

The last equality follows from the fact that  $(A'|B')$  is the upper half of a symplectic matrix, and hence  $(B')^{-1} A'$  is symmetric. Finally observe that  $|x_i| = 1$ ; hence,  $|y_i| = |q|^{n_i}$  where  $n_i$  are linear forms in  $\alpha$ . Hence, the  $y$ -contour  $C_\alpha$  is a product of tori whose radii depend linearly on  $\alpha$ . This completes the proof.  $\square$

*Remark 3.4* With the assumptions of Proposition 3.3, there exists an edge column vector  $\eta'$  whose coordinates are affine linear forms in  $\alpha, \mu, \lambda$ , a contour  $C$ , and affine linear forms  $\xi_i, \xi'_i, \xi''_i$  in the variables  $\varepsilon = (\varepsilon_1, \dots, \varepsilon_{N-1}), \mu$  and  $\lambda$  such that after the change in variables  $y_i = (-q)^{\eta'_i} z_i$ , we have

$$I_{T,\theta}^{\text{pre}}(q) = c(q)^N \int_C d\mu(z) \prod_{i=1}^N G_q \left( (-q)^{\xi_i(\varepsilon,\mu,\lambda)} z^{(\overline{C}-\overline{B})_i} \right) \times G_q \left( (-q)^{\xi'_i(\varepsilon,\mu,\lambda)} z^{(\overline{A}-\overline{C})_i} \right) G_q \left( (-q)^{\xi''_i(\varepsilon,\mu,\lambda)} z^{(\overline{B}-\overline{A})_i} \right), \tag{60}$$

where  $\xi_i(0, \mu, \lambda), \xi'_i(0, \mu, \lambda)$  and  $\xi''_i(0, \mu, \lambda)$  are  $\mathbb{Z}$ -linear combinations of  $1, \mu/pi, \lambda/(2\pi)$ . This follows from the symplectic properties of the Neumann–Zagier matrices [19] (compare also with the matrices  $(A'|B')$  in Eq. (67) below).

The contours in Eqs. (59) and (60) depend on a positive pre-angle structure, but the integrals are independent of the choice of the pre-angle structure. When we balance, i.e., set  $\varepsilon = 0$ , there are two possibilities: Either we can move the contour by a small isotopy in order to avoid the singularities of the integrand, or we cannot do so. In the former case, the new contour is canonically defined from the old contour. In the latter case, we apply the residue theorem to change the integration contour, and the residue contribution is either finite or infinite. In the latter case, by definition, our meromorphic function is infinity.

*Remark 3.5* An example of an integral where the latter case occurs is the following integral

$$I_\varepsilon(C) := \int_C \frac{dz}{z} G_q(z) G_q((-q)^\varepsilon z^{-1}), \tag{61}$$

where  $C$  is the contour  $|z| = 1 - \delta$  for  $\delta > 0$  small. The singularities of the integrand are  $z \in q^{-\mathbb{N}}$  and  $z \in (-q)^\varepsilon q^{\mathbb{N}}$ . When  $\varepsilon$  approaches zero, the contour is pinched from two sides by the singularities at  $z = (-q)^\varepsilon$  and  $z = 1$ . To avoid this pinching, by applying the residue theorem, we move the contour of integration  $C$  to the other side of 1, i.e., the contour  $C'$  given by  $|z| = 1 + \delta$ :

$$I_\varepsilon(C) = I_\varepsilon(C') - 2\pi i \operatorname{Res}_{z=1} \left( \frac{1}{z} G_q(z) G_q((-q)^\varepsilon z^{-1}) \right) = I_\varepsilon(C') + 2\pi i \frac{(-q; q)_\infty}{(q; q)_\infty} G_q((-q)^\varepsilon). \tag{62}$$

Even though  $I_\varepsilon(C')$  is regular at  $\varepsilon = 0$ , the contribution from the residue is singular because of the simple pole of  $G_q((-q)^\varepsilon)$  at  $\varepsilon = 0$ . Thus, we conclude that  $I_0(C) = \infty$ .

The next proposition defines the meromorphic function that appears in Theorem 1.1.

**Proposition 3.6** *Setting  $\varepsilon = 0$  (i.e., balancing the edges), and assuming that we find a contour  $C$ , we obtain a meromorphic function of  $(e_\mu, e_\lambda) := ((-q)^{\mu/\pi}, (-q)^{\lambda/(2\pi)}) \in (\mathbb{C}^*)^2$*

$$I_{T,e_\mu,e_\lambda}(q) = c(q)^N \int_C d\mu(z) \prod_{i=1}^N G_q \left( (-q)^{\xi_i(0,\mu,\lambda)} z^{(\overline{C}-\overline{B})_i} \right) G_q \left( (-q)^{\xi'_i(0,\mu,\lambda)} z^{(\overline{A}-\overline{C})_i} \right) \times G_q \left( (-q)^{\xi''_i(0,\mu,\lambda)} z^{(\overline{B}-\overline{A})_i} \right).$$

*The above integral is absolutely convergent.*

Let us phrase our previous proposition in more invariant language. Recall the complex torus  $\mathbb{T}_M = H^1(\partial M, \mathbb{C}^*)$ . Define the map

$$\varpi_{\mathcal{T}} : \mathcal{B}_{\mathcal{T}} \rightarrow \mathbb{T}_M, \quad \delta \mapsto (-q)^{\text{hol}(\delta)/\pi} \tag{63}$$

for a simple closed curve  $\delta$  of  $\partial M$ , where  $\text{hol}(\delta)$  denotes the angle holonomy along  $\delta$ . Proposition 3.6 states that the restriction  $I_{\mathcal{T}}^{\text{bal}}$  of the meromorphic function  $I_{\mathcal{T}}^{\text{pre}}$  on  $\mathcal{B}$  is the pullback of a meromorphic function  $I_{\mathcal{T}}$  on  $\mathbb{T}_M$ . In other words, the following diagram commutes:

$$\begin{array}{ccc} \mathcal{A}_{\mathcal{T}} & \longleftarrow \mathcal{B}_{\mathcal{T}} & \xrightarrow{\varpi_{\mathcal{T}}} \mathbb{T}_M \\ & \searrow I_{\mathcal{T}}^{\text{pre}} & \downarrow I_{\mathcal{T}} \\ & & \mathbb{C} \end{array} \tag{64}$$

### 3.2 Identification with the 3D index of Dimofte–Gaiotto–Gukov

In this subsection, we discuss the Laurent expansion of the meromorphic function  $I_{\mathcal{T}, e_{\mu}, e_{\lambda}}(q)$  on the real torus  $|e_{\mu}| = |e_{\lambda}| = 1$ , under the assumption that  $\mathcal{T}$  supports a strict angle structure. As we will show, the coefficients of the Laurent series are the 3D index of Dimofte–Gaiotto–Gukov. This will conclude the proof of Theorem 1.2.

The next lemma (for  $h = 0$ ) is based on the idea that the upper half part of a symplectic matrix with integer entries is a pair of coprime matrices.

**Lemma 3.7** *Suppose  $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  is a symplectic matrix with integer entries where  $A, B, C, D$  are  $N \times N$  matrices. Let  $(A'|B')$  denote the upper  $(N + h) \times 2N$  part of  $M$  and consider  $r, s \in \mathbb{Z}^N$  that satisfy the equation*

$$A'r + B's = \begin{pmatrix} 0 \\ v \end{pmatrix}$$

for a vector  $v \in \mathbb{Z}^h$ . Then, there exists  $k \in \mathbb{Z}^{N-h}$  such that

$$r = -B'^T \begin{pmatrix} v \\ k \end{pmatrix}, \quad s = A'^T \begin{pmatrix} v \\ k \end{pmatrix}.$$

*Proof* We apply the symplectic matrix  $M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  to the vector  $\begin{pmatrix} r \\ s \end{pmatrix}$  and define  $k \in \mathbb{Z}^{N-h}$  so that  $k' := \begin{pmatrix} v \\ k \end{pmatrix} \in \mathbb{Z}^N$  satisfies  $M \begin{pmatrix} r \\ s \end{pmatrix} = \begin{pmatrix} 0 \\ k' \end{pmatrix}$ . Then,

$$\begin{pmatrix} r \\ s \end{pmatrix} = M^{-1} \begin{pmatrix} 0 \\ k' \end{pmatrix} = \begin{pmatrix} D^T & -B^T \\ -C^T & A^T \end{pmatrix} \begin{pmatrix} 0 \\ k' \end{pmatrix} = \begin{pmatrix} -B^T k' \\ A^T k' \end{pmatrix}.$$

The result follows. □

*Proof* (of Theorem 1.2) Fix an ideal triangulation  $\mathcal{T}$  of  $M$  with  $N$  tetrahedra and let  $\bar{A}, \bar{B}$  and  $\bar{C}$  denote the Neumann–Zagier matrices describing gluing Eqs. (54). If we eliminate



the shape  $z'_i = -1/(z_i z''_i)$ , we obtain the gluing equations in the form:

$$\prod_{j=1}^N z_j^{A_{i,j}} (z''_j)^{B_{i,j}} = (-1)^{v_i}, \quad i = 1, \dots, N, \tag{65}$$

where

$$A = \bar{A} - \bar{B}, \quad B = \bar{C} - \bar{B}.$$

Notice that  $\bar{A} - \bar{C} = A - B, \bar{B} - \bar{C} = -B$ . Fix a strict angle structure  $\theta$  and use Proposition 3.1 to write the integrand of  $I_{T,\theta}^{\text{pre}}$  as follows:

$$\begin{aligned} \prod_{i=1}^N B(T_i, x, \theta) &= \prod_{i=1}^N \psi^0 \left( (-q)^{-\frac{\alpha_i + \gamma_i}{\pi}} \frac{X_{\alpha_i}}{X_{\gamma_i}}, (-q)^{-\frac{\alpha_i}{\pi}} \frac{X_{\beta_i}}{X_{\gamma_i}} \right) \\ &= \prod_{i=1}^N \psi^0 \left( (-q)^{-\frac{\alpha_i + \gamma_i}{\pi}} x^{(\bar{A} - \bar{C})_i}, (-q)^{-\frac{\alpha_i}{\pi}} x^{(\bar{B} - \bar{C})_i} \right) \\ &= \prod_{i=1}^N \psi^0 \left( (-q)^{-\frac{\alpha_i + \gamma_i}{\pi}} x^{(A - B)_i}, (-q)^{-\frac{\alpha_i}{\pi}} x^{-(B)_i} \right). \end{aligned}$$

Use Eq. (24) to expand the integrand of  $I_{T,\theta}^{\text{pre}}$  as a convergent series on the torus  $|x_i| = 1$  (for  $i = 1, \dots, N$ ):

$$\begin{aligned} \prod_{i=1}^N B(T_i, x, \theta) &= \sum_{r, s \in \mathbb{Z}^N} I^\Delta(r_1, s_1) \dots I^\Delta(r_N, s_N) x^{\sum_i (A - B)_i s_i} x^{-\sum_i (B)_i r_i} (-q)^{-\frac{1}{\pi} \sum_i (\alpha_i + \gamma_i) s_i + \alpha_i r_i} \end{aligned}$$

where  $r = (r_1, \dots, r_N) \in \mathbb{Z}^N$  and  $s = (s_1, \dots, s_N) \in \mathbb{Z}^N$ . Now, we can compute the absolutely convergent integral  $I_{T,\theta}^{\text{pre}}$  by integrating over the torus. After interchanging summation and integration (justified by uniform convergence) and applying the residue theorem, we obtain a sum over  $r, s \in \mathbb{Z}^N$  such that  $(A - B)s - Br = 0$ :


$$\begin{aligned} I_{T,\theta}^{\text{pre}} &= \int_{\mathbb{T}^N} d\mu(x) \prod_{i=1}^N B(T_i, x, \theta) \\ &= \sum_{r, s} I^\Delta(r_1, s_1) \dots I^\Delta(r_N, s_N) (-q)^{-\frac{1}{\pi} \sum_i (\alpha_i + \gamma_i) s_i + \alpha_i r_i} \int_{\mathbb{T}^N} d\mu(x) x^{\sum_i (A - B)_i s_i - \sum_i (B)_i r_i} \\ &= \sum_{r, s: (A - B)s - Br = 0} I^\Delta(r_1, s_1) \dots I^\Delta(r_N, s_N) (-q)^{-\frac{1}{\pi} \sum_i (\alpha_i + \gamma_i) s_i + \alpha_i r_i}. \end{aligned}$$

Collecting further terms whose meridian and (half) longitude holonomy is a fixed integer, we obtain the uniformly convergent sum:

$$I_{T,\theta}^{\text{pre}}(q) = \sum_{m, e \in \mathbb{Z}} (-q)^{\frac{m\mu}{\pi}} (-q)^{\frac{e\lambda}{2\pi}} \sum_{r, s} I^\Delta(r_1, s_1) \dots I^\Delta(r_N, s_N), \tag{66}$$

where  $r, s \in \mathbb{Z}^N$  satisfy the equation

$$A's + B'(-r - s) = \begin{pmatrix} 0 \\ v \end{pmatrix}, \quad v = \begin{pmatrix} m \\ e \end{pmatrix} \tag{67}$$



**Fig. 2** The ordered 2–3 Pachner move

for the matrix  $(A'|B')$  (where  $A', B'$  are  $(N + 1) \times N$  matrices) obtained from  $(A|B)$  after we remove any one row of it and replace it with two rows of the meridian and half-longitude monodromy. Neumann–Zagier [19] prove that  $(A'|B')$  can be completed to a symplectic matrix. Using this and Lemma 3.7, it follows that there exists  $k \in \mathbb{Z}^{N-1}$  such that

$$s = -B'^T \begin{pmatrix} v \\ k \end{pmatrix}, \quad -r - s = A'^T \begin{pmatrix} v \\ k \end{pmatrix}.$$

Let  $a_i$  and  $b_i$  for  $i = 1, \dots, N$  denote the  $i$ th column of  $A'$  and  $B'$ , respectively, and let  $k' = \begin{pmatrix} v \\ k \end{pmatrix}$ . Using the above and Eqs. (40a), (41), we obtain that

$$I^\Delta(r_i, s_i) = I^\Delta(-a_i \cdot k' + b_i \cdot k', -b_i \cdot k') = I^\Delta(-b_i \cdot k', a_i \cdot k')$$

for  $i = 1, \dots, N$ . Combined with Eq. (66), this gives

$$\begin{aligned} I_{\mathcal{T}, \theta}^{\text{pre}}(q) &= \sum_{m, e \in \mathbb{Z}} (-q)^{\frac{m\mu}{\pi}} (-q)^{\frac{e\lambda}{2\pi}} \sum_{k \in \mathbb{Z}^{N-1}} \prod_{i=1}^N I^\Delta(r_i, s_i)(q) \\ &= \sum_{m, e \in \mathbb{Z}} (-q)^{\frac{m\mu}{\pi}} (-q)^{\frac{e\lambda}{2\pi}} \sum_{k \in \mathbb{Z}^{N-1}} \prod_{i=1}^N I^\Delta(-b_i \cdot k', a_i \cdot k')(q) \\ &= \sum_{m, e \in \mathbb{Z}} (-q)^{\frac{m\mu}{\pi}} (-q)^{\frac{e\lambda}{2\pi}} \sum_{k \in \mathbb{Z}^{N-1}} \prod_{i=1}^N (-q)^{v \cdot k'} I_\Delta(-b_i \cdot k', a_i \cdot k')(q^2), \end{aligned}$$

where the last equality follows from Eq. (21) and  $v = (1, \dots, 1, m, e) \in \mathbb{Z}^{N+1}$ . The latter sum coincides with the 3D index of [9]; see also [12, Sec.4.5]. This completes the proof of Theorem 1.2.  $\square$

### 3.3 Invariance under 2–3 Pachner moves

Next, we prove the invariance of the meromorphic function  $I_{\mathcal{T}}$  under 2–3 Pachner moves. Consider two ideal triangulations  $\mathcal{T}$  and  $\tilde{\mathcal{T}}$  with  $N$  and  $N + 1$  tetrahedra that are related by a 2–3 Pachner move as in Fig. 2 and choose  $\theta$  and  $\tilde{\theta}$  compatible positive angle structures on  $\mathcal{T}$  and  $\tilde{\mathcal{T}}$  that satisfy Eq. (51). In particular, this means that the sum of angles around the interior edge of the 3 tetrahedra is  $2\pi$ .

Note that  $\tilde{\theta}$  determines  $\theta$ , but not vice versa. Let us denote the linear map  $\tilde{\theta} \mapsto \theta$  by  $\theta = m_3^2(\tilde{\theta})$ . The commutative diagram (64) gives a commutative diagram

$$\begin{array}{ccccc}
 \mathcal{A}_{\tilde{\mathcal{T}}} & \longleftarrow & \mathcal{B}_{\tilde{\mathcal{T}}} & \xrightarrow{\varpi_{\tilde{\mathcal{T}}}} & \mathbb{T}_M \\
 m_3^2 \downarrow & & m_3^2 \downarrow & & \parallel \\
 \mathcal{A}_{\mathcal{T}} & \longleftarrow & \mathcal{B}_{\mathcal{T}} & \xrightarrow{\varpi_{\mathcal{T}}} & \mathbb{T}_M
 \end{array} \tag{68}$$

Let  $\tilde{\theta}$  and  $\theta$  be positive pre-angle structures. We claim that

$$I_{\mathcal{T},\theta}^{\text{pre}}(q) = I_{\tilde{\mathcal{T}},\tilde{\theta}}^{\text{pre}}(q). \tag{69}$$

This follows by separating the integration variable of the inner edge of the 2–3 move in the integral  $I_{\tilde{\mathcal{T}},\tilde{\theta}}^{\text{pre}}(q)$  and applying the pentagon identity (50) to that variable. Since a meromorphic function is uniquely determined by its values on positive pre-angle structures and the map  $m_3^2 : \mathcal{A}_{\tilde{\mathcal{T}}} \rightarrow \mathcal{A}_{\mathcal{T}}$  is onto, it follows that  $I_{\tilde{\mathcal{T}}}^{\text{pre}} = I_{\mathcal{T}}^{\text{pre}} \circ m_3^2$ . Now restrict to  $\mathcal{B}_{\tilde{\mathcal{T}}}$ , use the above commutative diagram and the fact that  $m_3^2$  is onto. It follows that  $I_{\tilde{\mathcal{T}}}^{\text{bal}} = I_{\mathcal{T}}^{\text{bal}} \circ m_3^2$ . Using once again the commutative diagram and the fact that  $\varpi_{\mathcal{T}}$  and  $\varpi_{\tilde{\mathcal{T}}}$  are onto, it follows that  $I_{\tilde{\mathcal{T}}} = I_{\mathcal{T}}$ .

### 3.4 The singularities of $I_{\mathcal{T}}(q)$

The singularities of the integrand of  $I_{\mathcal{T},e_{\mu},e_{\lambda}}(q)$  are given by Lemma 2.2. To determine the singularities of the meromorphic function  $I_{\mathcal{T}}$ , perform one integral at a time and use the next lemma.

**Lemma 3.8** *Suppose  $f(z)$  is a meromorphic function of  $z$  with singularities on  $q^{-\mathbb{N}}$ . Fix positive integers  $a_1, \dots, a_p > 0$  and  $b_1, \dots, b_n > 0$ , let  $s = (s_1, \dots, s_p)$ ,  $t = (t_1, \dots, t_n)$  and consider the integral:*

$$F(s, t) = \int_{\mathcal{C}} \prod_{i=1}^p f(s_i z^{a_i}) \prod_{j=1}^n f(t_j z^{-b_j}) \frac{dz}{z},$$

where  $\mathcal{C}$  is a contour that separates  $q^{-1-\mathbb{N}}$  from  $q^{\mathbb{N}}$ . Then,  $F(s, t)$  is a meromorphic function of  $(s, t)$  with singularities at a subset of

$$\left\{ (s, t) \mid s_i^{b_j} t_j^{a_i} \in q^{-a_i \mathbb{N} - b_j \mathbb{N}} \text{ for some } 1 \leq i \leq p, 1 \leq j \leq n \right\}. \tag{70}$$

*Proof* The singularities of the integrand is the set  $\Sigma^-(s) \cup \Sigma^+(t)$  where

$$\Sigma^-(s) = \cup_{i=1}^p \{z^{a_i} \in s_i^{-1} q^{-\mathbb{N}}\}, \quad \Sigma^+(t) = \cup_{j=1}^n \{z^{b_j} \in t_j q^{\mathbb{N}}\}.$$

As long as  $\Sigma^-(s) \cup \Sigma^+(t)$  does not touch the contour  $\mathcal{C}$ ,  $F(s, t)$  is regular. It follows that if  $F(s, t)$  is singular when pinching occurs, in other words we must have  $z^{a_i} = s_i^{-1} q^{-k}$  and  $z^{b_j} = t_j q^l$  for some  $i, j$  and some  $k, l \in \mathbb{N}$ . Thus,  $z^{a_i b_j} = (s_i^{-1} q^{-k})^{b_j} = (t_j q^l)^{a_i}$ . The result follows.  $\square$

### 4 Examples and computations

#### 4.1 A non-1-efficient triangulation with two tetrahedra

It is traditional in hyperbolic geometry to illustrate theorems concerning ideally triangulated manifolds with the case of the standard triangulation of the complement of the  $4_1$  knot. In our examples, we will deviate from this principle and begin by giving a detailed computation of the state integral for the case of a non-1-efficient ideal triangulation with two tetrahedra. This illustrates Propositions 3.1, 3.2, 3.3, 3.6 and Theorem 1.1, and also points out the inapplicability of Theorem 1.2.

Ideal triangulations can be efficiently described, constructed and manipulated by SnapPy and Regina [5,6], and we will follow their description below. In particular, ideal triangulations can be uniquely reconstructed by their isometry signature, and the latter is a string of letters and numbers. There are exactly 10 ideal triangulations with two tetrahedra of manifolds with one cusp, given in Table 3 of [11], 9 of them are 1-efficient, and one of them with isometry signature  $cPcbbbdei$  is not. Although we will not use it, this triangulation is not ordered. The underlying 3-manifold  $M$  is the union of a  $T(2, 4)$  torus link with the  $T(1, 3)$  (trefoil) torus knot. In Regina format, the tetrahedron gluings of the triangulation  $\mathcal{T}$  of  $cPcbbbdei$  are given by:

tet	glued to	(012)	(013)	(023)	(123)
0		1(130)	1(023)	1(021)	1(132)
1		0(032)	0(201)	0(013)	0(132)

The edges of  $\mathcal{T}$  are given by:

tet	edge	01	02	03	12	13	23
0		0	0	0	0	1	1
1		0	0	0	1	0	1

and the triangle faces of  $\mathcal{T}$  are given by:

tet	face	012	013	023	123
0		0	1	2	3
1		2	0	1	3

The gluing equations, in SnapPy format and with the Regina ordering of the edges, are given by:

$$\begin{pmatrix} 1 & 1 & 2 & 1 & 2 & 1 \\ 1 & 1 & 0 & 1 & 0 & 1 \\ 0 & -1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 & 0 & 0 \end{pmatrix}.$$

The  $\bar{A}$ ,  $\bar{B}$  and  $\bar{C}$  matrices are given by:

$$\bar{A} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad \bar{B} = \begin{pmatrix} 1 & 2 \\ 1 & 0 \end{pmatrix}, \quad \bar{C} = \begin{pmatrix} 2 & 1 \\ 0 & 1 \end{pmatrix}.$$

The holonomy of the meridian and the longitude is given by:

$$\mu = -\beta_0 + \alpha_1, \quad \lambda = 2\gamma_0.$$

The  $\varepsilon$  variables are given by:

$$\varepsilon_0 = \alpha_0 + \beta_0 + 2\gamma_0 + \alpha_1 + 2\beta_1 + \gamma_1 - 2\pi, \quad \varepsilon_1 = \alpha_0 + \beta_0 + \alpha_1 + \gamma_1 - 2\pi. \quad (71)$$

Using  $\alpha_i + \beta_i + \gamma_i = \pi$  for  $i = 0, 1$ , we can solve the above equations in terms of the variables  $\alpha_0, \varepsilon_0, \mu, \lambda$ :

$$\begin{aligned} \alpha_0 &= \alpha_0, & \alpha_1 &= \pi - \alpha_0 - \lambda/2 + \mu, \\ \beta_0 &= \pi - \alpha_0 - \lambda/2, & \beta_1 &= \varepsilon_0 - \lambda/2, \\ \gamma_0 &= \lambda/2, & \gamma_1 &= \alpha_0 - \varepsilon_0 + \lambda - \mu. \end{aligned} \quad (72)$$

This illustrates Proposition 3.2.

If  $x_i$  are the variables of the  $i$ th edge for  $i = 0, 1$  and  $\theta$  is a pre-angle structure, then

$$\begin{aligned} B(\mathcal{T}, x, \theta)(q) &= c(q)^2 G_q \left( (-q)^{\frac{\alpha_0}{\pi}} \frac{x_0 x_0}{x_0 x_1} \right) G_q \left( (-q)^{\frac{\beta_0}{\pi}} \frac{x_0 x_1}{x_0 x_0} \right) G_q \left( (-q)^{\frac{\gamma_0}{\pi}} \frac{x_0 x_1}{x_0 x_1} \right) \\ &\quad \times G_q \left( (-q)^{\frac{\alpha_1}{\pi}} \frac{x_0 x_1}{x_0 x_0} \right) G_q \left( (-q)^{\frac{\beta_1}{\pi}} \frac{x_0 x_1}{x_0 x_1} \right) G_q \left( (-q)^{\frac{\gamma_1}{\pi}} \frac{x_0 x_0}{x_0 x_1} \right) \\ &= c(q)^2 G_q \left( (-q)^{\frac{\alpha_0}{\pi}} \frac{x_0}{x_1} \right) G_q \left( (-q)^{\frac{\beta_0}{\pi}} \frac{x_1}{x_0} \right) G_q \left( (-q)^{\frac{\gamma_0}{\pi}} \right) \\ &\quad \times G_q \left( (-q)^{\frac{\alpha_1}{\pi}} \frac{x_1}{x_0} \right) G_q \left( (-q)^{\frac{\beta_1}{\pi}} \right) G_q \left( (-q)^{\frac{\gamma_1}{\pi}} \frac{x_0}{x_1} \right). \end{aligned} \quad (73)$$

When  $\theta$  is positive, the absolutely convergent state integral is given by:

$$\begin{aligned} I_{\mathcal{T}, \theta}^{\text{pre}}(q) &= \frac{c(q)^2}{(2\pi i)^2} \int_{\mathbb{T}^2} G_q \left( (-q)^{\frac{\alpha_0}{\pi}} \frac{x_0}{x_1} \right) G_q \left( (-q)^{\frac{\beta_0}{\pi}} \frac{x_1}{x_0} \right) G_q \left( (-q)^{\frac{\gamma_0}{\pi}} \right) \\ &\quad \times G_q \left( (-q)^{\frac{\alpha_1}{\pi}} \frac{x_1}{x_0} \right) G_q \left( (-q)^{\frac{\beta_1}{\pi}} \right) G_q \left( (-q)^{\frac{\gamma_1}{\pi}} \frac{x_0}{x_1} \right) \frac{dx_0 dx_1}{x_0 x_1}. \end{aligned} \quad (74)$$

After rescaling  $x_0 \rightarrow x_0/x_1$ , the integral is free of the  $x_1$ -variable and is given by:

$$\begin{aligned} I_{\mathcal{T}, \theta}^{\text{pre}}(q) &= \frac{c(q)^2}{2\pi i} G_q \left( (-q)^{\frac{\gamma_0}{\pi}} \right) G_q \left( (-q)^{\frac{\beta_1}{\pi}} \right) \\ &\quad \times \int_{\mathbb{T}} G_q \left( (-q)^{\frac{\alpha_0}{\pi}} x_0 \right) G_q \left( (-q)^{\frac{\beta_0}{\pi}} \frac{1}{x_0} \right) G_q \left( (-q)^{\frac{\alpha_1}{\pi}} \frac{1}{x_0} \right) G_q \left( (-q)^{\frac{\gamma_1}{\pi}} x_0 \right) \frac{dx_0}{x_0}. \end{aligned} \quad (75)$$

Using Eq. (73), the above integral becomes:

$$\begin{aligned} I_{\mathcal{T}, \theta}^{\text{pre}}(q) &= \frac{c(q)^2}{2\pi i} G_q \left( (-q)^{\frac{\lambda}{2\pi}} \right) G_q \left( (-q)^{\frac{\varepsilon_0 - \lambda/2}{\pi}} \right) \int_{\mathbb{T}} G_q \left( (-q)^{\frac{\alpha_0}{\pi}} x_0 \right) \\ &\quad \times G_q \left( (-q)^{1 + \frac{-\alpha_0 - \lambda/2}{\pi}} \frac{1}{x_0} \right) G_q \left( (-q)^{1 + \frac{-\alpha_0 - \lambda/2 + \mu}{\pi}} \frac{1}{x_0} \right) G_q \left( (-q)^{\frac{\alpha_0 - \varepsilon_0 + \lambda - \mu}{\pi}} x_0 \right) \frac{dx_0}{x_0}. \end{aligned} \quad (76)$$

Applying the change in variables

$$(-q)^{\frac{\varepsilon_0}{\pi}} x_0 = y_0, \tag{77}$$

the above integral becomes:

$$\begin{aligned} I_{T,\theta}^{\text{pre}}(q) &= \frac{c(q)^2}{2\pi i} G_q \left( (-q)^{\frac{\lambda}{2\pi}} \right) G_q \left( (-q)^{\frac{\varepsilon_0 - \lambda/2}{\pi}} \right) \\ &\quad \times \int_{C_\alpha} G_q(y_0) G_q \left( (-q)^{1 - \frac{\lambda/2}{\pi}} \frac{1}{y_0} \right) G_q \left( (-q)^{1 + \frac{-\lambda/2 + \mu}{\pi}} \frac{1}{y_0} \right) \\ &\quad \times G_q \left( (-q)^{\frac{-\varepsilon_0 + \lambda - \mu}{\pi}} y_0 \right) \frac{dy_0}{y_0}, \end{aligned} \tag{78}$$

where  $C_\alpha$  is the torus  $|y_0| = |q^{\frac{\varepsilon_0}{\pi}}|$  and the integrand depends on  $\varepsilon_0, \mu, \lambda$  and  $y_0$ . Notice that when  $\theta$  is positive, Eq. (73) implies that  $\varepsilon_0 = \gamma_0 + \beta_1 > 0$  and  $\varepsilon_1 = -\gamma_0 - \beta_1 < 0$ . Moreover, we can set  $\varepsilon_0 = 0$  and obtain the uniformly convergent balanced integral

$$\begin{aligned} I_{T,\theta}^{\text{bal}}(q) &= \frac{c(q)^2}{2\pi i} G_q \left( (-q)^{\frac{\lambda}{2\pi}} \right) G_q \left( (-q)^{\frac{-\lambda/2}{\pi}} \right) \\ &\quad \times \int_{C_\alpha} G_q(y_0) G_q \left( (-q)^{1 - \frac{\lambda/2}{\pi}} \frac{1}{y_0} \right) \\ &\quad \times G_q \left( (-q)^{1 + \frac{-\lambda/2 + \mu}{\pi}} \frac{1}{y_0} \right) G_q \left( (-q)^{\frac{\lambda - \mu}{\pi}} y_0 \right) \frac{dy_0}{y_0}, \end{aligned} \tag{79}$$

which only depends on  $(e_\mu, e_\lambda) = ((-q)^{\mu/\pi}, (-q)^{\lambda/(2\pi)})$ . To simplify notation further, let  $(s, t) = (e_\mu, e_\lambda)$ . Then, we get:

$$\begin{aligned} I_{T,s,t}(q) &= \frac{c(q)^2}{2\pi i} G_q(t) G_q(t^{-1}) \\ &\quad \times \int_C G_q(y_0) G_q \left( (-q)t^{-1}y_0^{-1} \right) G_q \left( (-q)st^{-1}y_0^{-1} \right) G_q \left( s^{-1}t^2y_0 \right) \frac{dy_0}{y_0}, \end{aligned} \tag{80}$$

where  $C$  is the torus  $|y_0| = |q|^{\delta_0}$  for small  $\delta_0 > 0$ . This is the meromorphic function of Theorem 1.1. Next, we compute its singularities, starting from the singularities of the integrand. Using part (a) of Lemma 2.3, we see that the singularities of the integrand are given by

$$y_0 \in \Sigma^-(s, t) \cup \Sigma^+(s, t),$$

where

$$\Sigma^-(s, t) = q^{-\mathbb{N}} \cup st^{-2}q^{-\mathbb{N}}, \quad \Sigma^+(s, t) = (-q)t^{-1}q^{\mathbb{N}} \cup (-q)st^{-1}q^{\mathbb{N}}.$$

The contour of integration has to separate  $\Sigma^-(s, t)$  from  $\Sigma^+(s, t)$ . The integral can only be singular when pinching occurs, that is, for  $(s, t)$  such that  $\Sigma^-(s, t)$  intersects  $\Sigma^+(s, t)$ . This happens precisely when

$$t \in -q^{\mathbb{Z} \setminus \{0\}} \quad \text{or} \quad s^{-1}t \in -q^{\mathbb{Z} \setminus \{0\}}. \tag{81}$$

Using the notation of the  $q$ -rays (1), the above set is given by

$$-q\Sigma_{0,1} \cup -q^{-1}\Sigma_{0,-1} \cup -q\Sigma_{-1,1} \cup -q^{-1}\Sigma_{1,-1}.$$

Thus, the above integral is a meromorphic function which is regular on the complement of the set  $\Sigma_{\mathcal{T}}$ . Note that some of the points of the set (81) might be regular points of the integral, this happens for instance when the residue at simple poles vanishes. Note also that the set (81) is disjoint from the real torus  $|s| = |t| = 1$ , so the integral can be expanded into Laurent series convergent on the real torus  $|s| = |t| = 1$ . However, the prefactor  $G_q(t)G_q(t^{-1})$  has singularities on the set  $t \in q^{\mathbb{Z}}$  (and those are actual, i.e., not removable), which prevent the meromorphic function  $I_{\mathcal{T},s,t}(q)$  from being expanded into Laurent series on the torus  $|s| = |t| = 1$ . In conclusion, the meromorphic function  $I_{\mathcal{T},s,t}(q)$  is regular in the complement of the shifted  $q$ -rays

$$\Sigma_{0,1} \cup \Sigma_{0,-1} \cup -q\Sigma_{0,1} \cup -q^{-1}\Sigma_{0,-1} - q\Sigma_{-1,1} \cup -q^{-1}\Sigma_{1,-1}. \tag{82}$$

**4.2 The  $4_1$  knot**

Next, we discuss the example of the  $4_1$  knot, giving the first and last steps of the above computations, and asking the reader to fill in the intermediate steps.

The gluing equations, in `Snappy` format and with the `Regina` ordering of the edges, are given by:

$$\begin{pmatrix} 2 & 1 & 0 & 2 & 1 & 0 \\ 0 & 1 & 2 & 0 & 1 & 2 \\ 1 & 0 & 0 & 0 & 0 & -1 \\ 1 & 1 & 1 & 1 & -1 & -3 \end{pmatrix}.$$

The above matrix determines the following information. The  $\bar{A}$ ,  $\bar{B}$  and  $\bar{C}$  matrices are given by:

$$\bar{A} = \begin{pmatrix} 2 & 2 \\ 0 & 0 \end{pmatrix}, \quad \bar{B} = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad \bar{C} = \begin{pmatrix} 0 & 0 \\ 2 & 2 \end{pmatrix}.$$

The holonomy of the meridian and the longitude is given by:

$$\mu = \alpha_0 - \gamma_1, \quad \lambda = \alpha_0 + \beta_0 + \gamma_0 + \alpha_1 - \beta_1 - 3\gamma_1.$$

The  $\varepsilon$  variables are given by:

$$\varepsilon_0 = 2\alpha_0 + \beta_0 + 2\alpha_1 + \beta_1 - 2\pi, \quad \varepsilon_1 = \beta_0 + 2\gamma_0 + \beta_1 + 2\gamma_1 - 2\pi.$$

Using  $\alpha_i + \beta_i + \gamma_i = \pi$  for  $i = 0, 1$ , we can solve the above equations in terms of the variables  $\alpha_0, \varepsilon_0, \mu, \lambda$ :

$$\begin{aligned} \alpha_0 &= \alpha_0, & \alpha_1 &= \alpha_0 + \lambda/2 - \mu, \\ \beta_0 &= \pi - 2\alpha_0 + \varepsilon_0 - \lambda/2, & \beta_1 &= \pi - 2\alpha_0 - \lambda/2 + 2\mu, \\ \gamma_0 &= \alpha_0 - \varepsilon_0 + \lambda/2, & \gamma_1 &= \alpha_0 - \mu. \end{aligned} \tag{83}$$

Let  $\theta$  be a positive semi-angle structure. Rescaling  $x_0 \rightarrow x_0/x_1$ , applying the change in variables

$$(-q)^{-\frac{\alpha_0}{\pi}} x_0 = y_0, \tag{84}$$

letting  $(s, t) = (e_\mu, e_\lambda)$ , and following the steps of the previous example we obtain that the state integral is given by

$$I_{\mathcal{T}_{4_1, \theta}}^{\text{pre}}(q) = \frac{c(q)^2}{2\pi i} \int_{C_\alpha} \frac{dy_0}{y_0} G_q(y_0^{-1}) G_q(s^{-1}y_0^{-1}) G_q((-q)^{-\varepsilon_0}ty_0^{-1}) G_q(s^{-1}ty_0^{-1}) \\ \times G_q(-q(-q)^{\varepsilon_0}t^{-1}y_0^2) G_q(-qs^2t^{-1}y_0), \tag{85}$$

where the contour of integration  $C_\alpha$  (determined by (84)) is given by  $|y_0| = |q^{-\frac{\alpha_0}{\pi}}|$ . Moreover, we can set  $\varepsilon_0 = 0$ , and obtain the meromorphic function of the  $4_1$  knot:

$$I_{\mathcal{T}_{4_1, s, t}}(q) = \frac{c(q)^2}{2\pi i} \int_C \frac{dy_0}{y_0} G_q(y_0^{-1}) G_q(s^{-1}y_0^{-1}) G_q(ty_0^{-1}) G_q(s^{-1}ty_0^{-1}) \\ \times G_q(-qt^{-1}y_0^2) G_q(-qs^2t^{-1}y_0), \tag{86}$$

where  $C$  is the torus  $|y_0| = 1^+$ .

### 4.3 The sister of the $4_1$ knot

Next, we present the invariant for the sister  $m003$  of the  $4_1$  knot. The gluing equations, in SnapPy format and with the Regina ordering of the edges, are given by:

$$\begin{pmatrix} 2 & 0 & 1 & 2 & 0 & 1 \\ 0 & 2 & 1 & 0 & 2 & 1 \\ 0 & -2 & 0 & 2 & 0 & 0 \\ 0 & -1 & 0 & 2 & -1 & 0 \end{pmatrix}.$$

Using the above matrix, we can compute the  $\bar{A}, \bar{B}, \bar{C}$  matrices, the holonomy of the meridian and longitude, the  $\varepsilon$  variables, and express all variables in terms of the variables  $\alpha_0, \varepsilon_0, \mu, \lambda$ :

$$\begin{aligned} \alpha_0 &= \alpha_0, & \alpha_1 &= \alpha_0 - \varepsilon_0 + \lambda, \\ \beta_0 &= \alpha_0 - \varepsilon_0 + \lambda - \mu/2, & \beta_1 &= \alpha_0 - \varepsilon_0 + \mu/2, \\ \gamma_0 &= \pi - 2\alpha_0 + \varepsilon_0 - \lambda + \mu/2, & \gamma_1 &= \pi - 2\alpha_0 + 2\varepsilon_0 - \lambda - \mu/2. \end{aligned} \tag{87}$$

Let  $\theta$  be a positive semi-angle structure. Rescaling  $x_0 \rightarrow x_0/x_1$ , applying the change in variables

$$(-q)^{\frac{\alpha_0}{\pi}} x_0 = y_0, \tag{88}$$

letting  $(s, t) = (e_\mu, e_\lambda)$ , and following the steps of the previous example we obtain that the state integral is given by

$$I_{\mathcal{T}_{m003, \theta}}^{\text{pre}}(q) = \frac{c(q)^2}{2\pi i} \int_{C_\alpha} \frac{dy_0}{y_0} G_q((-q)^{1+2\varepsilon_0}s^{-1/2}t^{-2}y_0^{-2}) \\ \times G_q((-q)^{1+\varepsilon_0}s^{1/2}t^{-2}y_0^{-2}) G_q(y_0) G_q((-q)^{-\varepsilon_0}s^{1/2}y_0) G_q((-q)^{-\varepsilon_0}t^2y_0) \\ \times G_q((-q)^{-\varepsilon_0}s^{-1/2}t^2y_0), \tag{89}$$



where the contour of integration  $C_\alpha$  (determined by (88)) is given by  $|y_0| = |(-q)^{\frac{\alpha_0}{\pi}}|$ . Moreover, we can set  $\varepsilon_0 = 0$ , and obtain the meromorphic function of the sister of the 4<sub>1</sub> knot:

$$I_{\mathcal{T}_{m003,s,t}}(q) = \frac{c(q)^2}{2\pi i} \int_C \frac{dy_0}{y_0} G_q(-qs^{-1/2}t^{-2}y_0^{-2}) G_q(-qs^{1/2}t^{-2}y_0^{-2}) \times G_q(y_0) G_q(s^{1/2}y_0) G_q(t^2y_0) G_q(s^{-1/2}t^2y_0), \tag{90}$$

where  $C$  is the torus  $|y_0| = 1^+$ .

**4.4 The unknot**

Next, we compute the invariant for the unknot, and find a surprise: We can compute the integral exactly. The unknot has three triangulations with two tetrahedra given in Table 3 of [11]. One of them has isometry signature cMcabbgds. The gluing equations, in SnapPy format and with the Regina ordering of the edges, are given by:

$$\begin{pmatrix} 1 & 2 & 2 & 2 & 2 & 2 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & -2 \\ 1 & 0 & 0 & 0 & -1 & 0 \end{pmatrix}.$$

Following the steps of the previous examples, we have:

$$\varepsilon_0 = \alpha_0 + 2\beta_0 + 2\gamma_0 + 2\alpha_1 + 2\beta_1 + 2\gamma_1 - 2\pi, \quad \varepsilon_1 = \alpha_0 - 2\pi, \tag{91}$$

which simplifies to  $\varepsilon_0 = -\alpha_0 + 2\pi$  and in particular is bigger than  $\pi$  when  $\theta$  is positive. We can express all variables in terms of the variables  $\alpha_0, \varepsilon_0, \mu, \lambda$ :

$$\begin{aligned} \alpha_0 &= 2\pi - \varepsilon_0, & \alpha_1 &= -\pi + \varepsilon_0 + \lambda + \mu/2, \\ \beta_0 &= \beta_0, & \beta_1 &= 2\pi - \varepsilon_0 - \lambda, \\ \gamma_0 &= -\pi - \beta_0 + \varepsilon_0, & \gamma_1 &= -\mu/2. \end{aligned} \tag{92}$$

After rescaling  $x_0 \rightarrow x_0/x_1$ , applying the change in variables

$$(-q)^{-\frac{\beta_0}{\pi}} x_0 = y_0, \tag{93}$$

letting  $(s, t) = (e_\mu, e_\lambda)$ , and following the steps of the previous example we obtain that the state integral is given by

$$I_{\mathcal{T}_{\text{unknot},\theta}}^{\text{pre}}(q) = \frac{c(q)^2}{2\pi i} G_q((-q)^{2-\varepsilon_0}) G_q(s^{-1/2}) G_q((-q)^{2-\varepsilon_0}t^{-2}) G_q((-q)^{-1+\varepsilon_0}s^{1/2}t^2) \times \int_{C_\alpha} \frac{dy_0}{y_0} G_q(-q^{-1}y_0) G_q(y_0^{-1}), \tag{94}$$

where  $C_\alpha$  is the torus  $|y_0| = |(-q)^{-\frac{\beta_0}{\pi}}|$ . On the other hand, Eqs. (7) and (8) give that

$$G_q(-q^{-1}y_0) G_q(y_0^{-1}) = \frac{1}{(1+y_0)(1+q^{-1}y_0)},$$

so that

$$I_{\mathcal{T}_{\text{unknot},\theta}^{\text{bal}}}(q) = \frac{c(q)^2}{2\pi i} G_q((-q)^{2-\varepsilon_0}) G_q(s^{-1/2}) G_q((-q)^{2-\varepsilon_0}t^{-2}) G_q((-q)^{-1+\varepsilon_0}s^{1/2}t^2) \times \int_{C_\alpha} \frac{dy_0}{y_0} \frac{1}{(1+y_0)(1+q^{-1}y_0)}. \tag{95}$$

When  $\theta$  is positive, the contour  $C_\alpha$  encircles both singularities  $-1$  and  $-q$  of the integrand, and a residue calculation reveals that the integral is zero. It follows that

$$I_{\mathcal{T}_{\text{unknot},s,t}}(q) = 0. \tag{96}$$

#### 4.5 The trefoil

Next, we compute the invariant for the trefoil. The gluing equations, in SnapPy format and with the Regina ordering of the edges, are given by:

$$\begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 0 \\ 1 & 2 & 2 & 2 & 1 & 2 \\ 0 & 0 & -1 & 1 & 0 & 0 \\ 1 & 0 & -4 & 4 & -1 & 0 \end{pmatrix}.$$

The state integral is given by

$$I_{\mathcal{T}_{3_1,\theta}^{\text{pre}}}(q) = \frac{c(q)^2}{2\pi i} G_q((-q)^{1+\varepsilon_0/2}s^2t^{-1}) G_q((-q)^{1+\varepsilon_0/2}s^{-2}t) \times \int_{C_\alpha} \frac{dy_0}{y_0} G_q(y_0^{-1}) G_q(s^{-1}y_0^{-1}) G_q((-q)^{-\varepsilon_0/2}s^3t^{-1}y_0) G_q((-q)^{-\varepsilon_0/2}s^{-2}ty_0), \tag{97}$$

where the contour of integration  $C_\alpha$  is given by  $|y_0| = |(-q)^{\frac{-\alpha_1}{\pi}}|$ . Moreover, we can set  $\varepsilon_0 = 0$  and obtain the meromorphic function given by:

$$I_{\mathcal{T}_{3_1,s,t}}(q) = \frac{c(q)^2}{2\pi i} G_q(-qs^2t^{-1}) G_q(-qs^{-2}t) \times \int_C \frac{dy_0}{y_0} G_q(y_0^{-1}) G_q(s^{-1}y_0^{-1}) G_q(s^3t^{-1}y_0) G_q(s^{-2}ty_0), \tag{98}$$

where  $C$  is the torus  $|y_0| = 1^+$ .

#### 4.6 The $5_2$ knot

Next, we present the invariant of the  $5_2$  knot. The gluing equations, in SnapPy format (we use the homological longitude and the Regina ordering of the edges), are given by:

$$\begin{pmatrix} 1 & 1 & 0 & 1 & 0 & 0 & 1 & 1 & 0 \\ 0 & 1 & 1 & 0 & 0 & 2 & 0 & 1 & 1 \\ 1 & 0 & 1 & 1 & 2 & 0 & 1 & 0 & 1 \\ -1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 2 & 0 & -3 & 1 & 0 & -2 & 0 & 0 & 1 \end{pmatrix}.$$

After rescaling  $x_i \rightarrow x_i/x_2$  and a change in variables  $x_0 = (-q)^{\alpha_0}y_0$  and  $x_1 = (-q)^{-\alpha_1/2}s^{1/2}t^{1/2}y_1$ , the state integral is given by

$$\begin{aligned}
 I_{T_{5_2,\theta}}^{\text{pre}}(q) &= \frac{c(q)^3}{(2\pi i)^2} \int_{C_\alpha} \frac{dy_0 dy_1}{y_0 y_1} G_q(y_0^{-1}) G_q(sy_0^{-1}) G_q((-q)^{-\varepsilon_1}s^2y_0^{-1}) \\
 &\quad \times G_q(-qs^{-2}t^{-1}y_0y_1^{-2}) G_q((-q)^{1+\varepsilon_0/4}y_0y_1^{-1}) \\
 &\quad \times G_q((-q)^{1+3\varepsilon_0/4+\varepsilon_1}s^{-3}t^{-1}y_0y_1^{-1}) \\
 &\quad \times G_q((-q)^{-\varepsilon_0/4}y_1) G_q((-q)^{-3\varepsilon_0/4}sty_1) G_q(sty_1^2), \tag{99}
 \end{aligned}$$

where  $C_\alpha$  is the torus given by  $|y_0| = |(-q)^{-\frac{\alpha_0}{\pi}}|$  and  $|y_1s^{1/2}t^{1/2}| = |(-q)^{\frac{\alpha_1}{2\pi}}|$ . Moreover, we can set  $\varepsilon_0 = \varepsilon_1 = 0$  and obtain the meromorphic function given by:

$$\begin{aligned}
 I_{T_{5_2,s,t}}(q) &= \frac{c(q)^3}{(2\pi i)^2} \int_C \frac{dy_0 dy_1}{y_0 y_1} G_q(y_0^{-1}) G_q(sy_0^{-1}) G_q(s^2y_0^{-1}) G_q(-qs^{-2}t^{-1}y_0y_1^{-2}) \\
 &\quad \times G_q(-qy_0y_1^{-1}) G_q(-qs^{-3}t^{-1}y_0y_1^{-1}) G_q(y_1) G_q(sty_1) G_q(sty_1^2), \tag{100}
 \end{aligned}$$

where  $C$  is the torus given by  $|y_0| = 1^+$  and  $|y_1s^{1/2}t^{1/2}| = 1^-$ .

**4.7 The  $6_1$  knot**

Finally, we present the invariant of the  $6_1$  knot. The gluing equations, in SnapPy format (we use the homological longitude and the Regina ordering of the edges), are given by:

$$\begin{pmatrix}
 1 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 1 & 1 & 0 \\
 0 & 1 & 0 & 1 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 1 \\
 0 & 1 & 1 & 0 & 0 & 2 & 0 & 1 & 1 & 1 & 0 & 0 \\
 1 & 0 & 1 & 1 & 2 & 0 & 0 & 0 & 1 & 0 & 1 & 1 \\
 -1 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 \\
 -1 & 1 & 0 & 0 & 1 & 1 & 0 & -1 & 0 & 0 & 0 & -2
 \end{pmatrix}.$$

After rescaling  $x_i \rightarrow x_i/x_3$  and a change in variables

$$x_0 = (-q)^{\alpha_2}y_0, \quad x_1 = (-q)^{\alpha_0}y_1, \quad x_2 = (-q)^{-\alpha_1/2}t^{1/2}y_2 \tag{101}$$

the state integral is given by

$$\begin{aligned}
 I_{T_{6_1,\theta}}^{\text{pre}}(q) &= \frac{c(q)^4}{(2\pi i)^3} \int_{C_\alpha} \frac{dy_0 dy_1 dy_2}{y_0 y_1 y_2} G_q(y_0^{-1}) G_q(y_1^{-1}) G_q(sy_1^{-1}) G_q((-q)^{\varepsilon_2}s^{-2}y_0^{-1}y_1) \\
 &\quad \times G_q(-qs^{-1}t^{-1}y_1y_2^{-2}) G_q((-q)^{1/2-\varepsilon_0/4+\varepsilon_1/4}t^{-1}y_2^{-1}) \\
 &\quad \times G_q((-q)^{3/2+\varepsilon_0/4+3\varepsilon_1/4}y_0y_2^{-1}) \\
 &\quad \times G_q((-q)^{3/2+3\varepsilon_0/4+\varepsilon_1/4}t^{-1}y_0y_1y_2^{-1}) G_q((-q)^{-1/2-3\varepsilon_0/4-\varepsilon_1/4}ty_1^{-1}y_2) \\
 &\quad \times G_q((-q)^{1/2+\varepsilon_0/4-\varepsilon_1/4-\varepsilon_2}s^2ty_0y_1^{-1}y_2) G_q((-q)^{-1/2-\varepsilon_0/4-3\varepsilon_1/4}y_0^{-1}y_1y_2) \\
 &\quad \times G_q(-qty_2^2), \tag{102}
 \end{aligned}$$

where  $C_\alpha$  is the torus given by  $|y_0| = |(-q)^{\frac{-\alpha_2}{\pi}}|$ ,  $|y_1| = |(-q)^{\frac{-\alpha_0}{\pi}}|$  and  $|y_2 t^{1/2}| = |(-q)^{\frac{\alpha_1}{2\pi}}|$ . Moreover, we can set  $\varepsilon_0 = \varepsilon_1 = \varepsilon_2 = 0$  and obtain the meromorphic function given by:

$$\begin{aligned}
 I_{\mathcal{T}_{61},s,t}(q) &= \frac{c(q)^4}{(2\pi i)^3} \int_{C_\alpha} \frac{dy_0 dy_1 dy_2}{y_0 y_1 y_2} G_q(y_0^{-1}) G_q(y_1^{-1}) G_q(sy_1^{-1}) G_q(s^{-2}y_0^{-1}y_1) \\
 &\times G_q(-qs^{-1}t^{-1}y_1 y_2^{-2}) G_q((-q)^{1/2}t^{-1}y_2^{-1}) G_q((-q)^{3/2}y_0 y_2^{-1}) \\
 &\times G_q((-q)^{3/2}t^{-1}y_0 y_1 y_2^{-1}) G_q((-q)^{-1/2}t y_1^{-1} y_2) \\
 &\times G_q((-q)^{1/2} s^2 t y_0 y_1^{-1} y_2) G_q((-q)^{-1/2} y_0^{-1} y_1 y_2) G_q(-qt y_2^2), \tag{103}
 \end{aligned}$$

where  $C$  is the torus given by  $|y_0| = 1^+$ ,  $|y_1| = 1^+$  and  $|y_2 t^{1/2}| = 1^-$ .

**Author details**

<sup>1</sup>School of Mathematics, Georgia Institute of Technology, Atlanta, GA 30332-0160, USA, <sup>2</sup>Max Planck Institute for Mathematics, Vivatsgasse 7, 53111 Bonn, Germany, <sup>3</sup>Section de Mathématiques, Université de Genève, 2-4 rue du Lièvre, Case Postale 64, 1211 Geneva 4, Switzerland.

**Acknowledgements**

Open access funding provided by Max Planck Society. The authors wish to thank R. Siejakowski for a careful reading of the manuscript and for pointing out typographical errors, and A. Mellit for enlightening discussions. S.G. was supported in part by grant DMS-14-06419 of the US National Science Foundation. R.K. was supported in part by the Swiss National Science Foundation and the center of excellence grant Center for quantum geometry of Moduli Spaces of the Danish National Research Foundation. The paper was presented by S.G. in a conference at the Max-Planck Institute in Bonn (Modular Forms are Everywhere) in honor of D. Zagier’s 65th birthday. S.G. wishes to thank the organizers, and particularly K. Bringmann and D. Zagier for their invitation and hospitality.

**Appendix A: A quantum dilogarithm over the LCA group  $\mathbb{T} \times \mathbb{Z}$**

The function (19) with  $q$  real has been introduced and studied in the functional analytic context of Hilbert spaces and quantum  $E(2)$  group by Woronowicz in [21]. In this section, we derive some of its operator properties by using the theory of quantum dilogarithms over Pontryagin self-dual LCA groups developed in [2, 15].

Throughout the section, for a Hilbert space  $H$ , we write  $A: H \rightarrow H$ , if  $A$  is a not necessarily bounded linear operator in  $H$  whose domain is dense in  $H$ . The Hermitian conjugate of  $A$  will be denoted  $A^*$ . Below, we will use freely the standard Dirac’s bra-ket notation.

The group  $\mathbb{T} \times \mathbb{Z}$  is a self-dual LCA group with the *gaussian exponential*

$$\langle \cdot \rangle: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{T}, \quad \langle z, m \rangle = z^m, \quad \text{for all } (z, m) \in \mathbb{T} \times \mathbb{Z}. \tag{104}$$

The *Fourier kernel* is fixed as the co-boundary of the gaussian exponential

$$\langle z, m; w, n \rangle := \frac{\langle zw, m + n \rangle}{\langle z, m \rangle \langle w, n \rangle} = z^n w^m. \tag{105}$$

We define a unitary *Fourier operator*  $F: L^2(\mathbb{T} \times \mathbb{Z}) \rightarrow L^2(\mathbb{T} \times \mathbb{Z})$  by the integral kernel

$$\langle z, m | F | w, n \rangle = \langle z, m; w, n \rangle. \tag{106}$$

For any (measurable) function  $f: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{C}$ , we associate three normal operators as follows. The multiplication operator by  $f$ :

$$f(q): L^2(\mathbb{T} \times \mathbb{Z}) \rightarrow L^2(\mathbb{T} \times \mathbb{Z}), \quad \langle z, m | f(q) = f(z, m) \langle z, m |, \tag{107}$$

its unitary conjugate by the Fourier operator

$$f(p) := Ff(q)F^*, \tag{108}$$

and the unitary conjugate of the latter by the inverse of the (unitary) multiplication operator  $\langle q \rangle$  by the gaussian exponential (104):

$$f(p + q) := \langle q \rangle^* f(p) \langle q \rangle. \tag{109}$$

We remark that all three operators  $f(q)$ ,  $f(p)$  and  $f(p + q)$  have spectrum given by the closure of the image of  $f$ .

**Lemma A.1** *The function*

$$\eta: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{C}_{\neq 0}, \quad (z, m) \mapsto -zq^{1-m}, \tag{110}$$

satisfies the following operator equations:

$$\eta(q)\eta(p) = q^2\eta(p)\eta(q), \quad \eta(q)^*\eta(p) = \eta(p)\eta(q)^*, \quad \eta(p + q) = -\eta(p)\eta(q). \tag{111}$$

*Proof* In the Hilbert space  $L^2(\mathbb{Z})$ , define a self-adjoint operator  $h$  by

$$\langle m|h = m\langle m|, \quad \text{for all } m \in \mathbb{Z}, \tag{112}$$

and a unitary operator  $z$  by

$$\langle m|z = \langle m - 1|, \quad \text{for all } m \in \mathbb{Z}. \tag{113}$$

These operators satisfy the commutation relation

$$[h, z] := hz - zh = z, \tag{114}$$

which is verified as follows:

$$\langle m|[h, z] = \langle m|h z - \langle m|z h = m\langle m - 1| - (m - 1)\langle m - 1| = \langle m|z. \tag{115}$$

Similarly, in the Hilbert space  $L^2(\mathbb{T})$ , we define a self-adjoint operator  $H$  by

$$\langle z|H = z \frac{\partial}{\partial z} \langle z|, \quad \text{for all } z \in \mathbb{T}, \tag{116}$$

and a unitary operator  $Z$  by

$$\langle z|Z = z\langle z|, \quad \text{for all } z \in \mathbb{T}. \tag{117}$$

It is easily verified that

$$[H, Z] := HZ - ZH = Z. \tag{118}$$

Moreover, if  $J: L^2(\mathbb{Z}) \rightarrow L^2(\mathbb{T})$  is the isomorphism defined by the integral kernel

$$\langle z|J|m \rangle = z^m, \quad \text{for all } (z, m) \in \mathbb{T} \times \mathbb{Z}, \tag{119}$$

then we have the equalities

$$Jh = HJ, \quad Jz = ZJ. \tag{120}$$

Identifying  $H, Z, h, z$  with their natural counterparts in  $L^2(\mathbb{T} \times \mathbb{Z})$ , we have

$$Fh = HF, \quad Fz = ZF, \quad FH = -hF, \quad FZ = z^{-1}F, \tag{121}$$

and

$$h(q) = \langle q|h, \quad Z(q) = \langle q|Z, \quad H(q) = \langle q|(H+h), \quad z(q) = \langle q|zZ^{-1} \tag{122}$$

so that

$$f(q) = f(Z, h), \quad f(p) = f(z^{-1}, H), \quad f(p+q) = f(z^{-1}Z, H+h) \tag{123}$$

for any  $f: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{C}$ , where, in the right-hand sides, the functions with operator arguments are understood in the spectral sense. In the case of  $f = \eta$ , we thus have

$$\eta(q) = -Zq^{1-h}, \quad \eta(p) = -z^{-1}q^{1-H}, \quad \eta(p+q) = -z^{-1}Zq^{1-H-h} \tag{124}$$

and the relations (111) are verified straightforwardly. □

Next, observe that the function

$$\mu: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{C}, \quad (z, m) \mapsto (\eta(z, m); q^2)_\infty, \tag{125}$$

nowhere vanishes and satisfies the operator five-term identity

$$\mu(p)\mu(q) = \mu(q)\mu(p+q)\mu(p) \tag{126}$$

as a consequence of Lemma A.1 and the formal power series identity in non-commuting indeterminates

$$(v; q^2)_\infty (u; q^2)_\infty = (u; q^2)_\infty (-vu; q^2)_\infty (v; q^2)_\infty, \quad uv = q^2vu, \tag{127}$$

which is equivalent to the  $q$ -binomial formula

$$\sum_{n=0}^{\infty} \frac{(a; q^2)_n}{(q^2; q^2)_n} z^n = \frac{(az; q^2)_\infty}{(z; q^2)_\infty}, \tag{128}$$

see [14]. Indeed, by substituting  $u$  by  $\eta(q)$  and  $v$  by  $\eta(p)$ , we convert (127) into (126). Lemma A.1 also implies that the function

$$\phi_q: \mathbb{T} \times \mathbb{Z} \rightarrow \mathbb{T}, \quad (z, m) \mapsto \mu(z, m)/\mu(1/z, m) = \mu(z, m)/\overline{\mu(z, m)}, \tag{129}$$

satisfies the unitarized version of identity (126):

$$\phi_q(\mathfrak{p})\phi_q(\mathfrak{q}) = \phi_q(\mathfrak{q})\phi_q(\mathfrak{p} + \mathfrak{q})\phi_q(\mathfrak{p}). \tag{130}$$

Moreover,  $\phi_q$  satisfies an inversion relation, see below (131), which allows us to identify it as an example of a quantum dilogarithm over  $\mathbb{T} \times \mathbb{Z}$ .

**Lemma A.2** *The function (129) satisfies the following inversion relation:*

$$\phi_q(z, m)\phi_q(1/z, -m) = z^m = \langle z, m \rangle, \quad \text{for all } (z, m) \in \mathbb{T} \times \mathbb{Z}. \tag{131}$$

*Proof* We have

$$\begin{aligned} \phi_q(z, m)\phi_q(1/z, -m) &= \frac{(-q^{1-m}z; q^2)_\infty (-q^{1+m}/z; q^2)_\infty}{(-q^{1-m}/z; q^2)_\infty (-q^{1+m}z; q^2)_\infty} \\ &= \frac{\theta_q(zq^{-m})}{\theta_q(zq^m)} = \frac{\theta_q(q^{-2m}zq^m)}{\theta_q(zq^m)} = q^{-m^2} (zq^m)^m = z^m, \end{aligned} \tag{132}$$

where in the second equality we have used (26) and in the fourth equality the functional equation (32). □

**Appendix B: The quantum dilogarithm and the Beta pentagon relation**

In this section, we generalize the result of [2] to include the LCA groups which do not admit division by 2. Note that most of the equations of this section (for instance, (137), (140), (145), (149)) are valid as distributions.

Let  $A$  be a self-dual LCA group with a gaussian exponential  $\langle \cdot \rangle: A \rightarrow \mathbb{T}$  and the Fourier kernel

$$\langle x; y \rangle = \frac{\langle x + y \rangle}{\langle x \rangle \langle y \rangle}, \quad \text{for all } (x, y) \in A^2. \tag{133}$$

For a nonnegative integer  $n \in \mathbb{N}$ , denote

$$[n] := \mathbb{Z}_{\geq 0} \cap \mathbb{Z}_{\leq n} \quad \text{for all } n \in \mathbb{Z}_{\geq 0}. \tag{134}$$

According to [2], a bounded function

$$f: [4] \times A \rightarrow \mathbb{C}, \quad (i, x) \mapsto f_i(x), \tag{135}$$

is called of *Faddeev type* if it satisfies the non-constant version of the operator pentagon relation

$$f_1(\mathfrak{p})f_3(\mathfrak{q}) = f_4(\mathfrak{q})f_2(\mathfrak{p} + \mathfrak{q})f_0(\mathfrak{p}). \tag{136}$$

The latter is equivalent to the functional integral identity

$$\tilde{f}_1(x)\tilde{f}_3(u)\langle x; u \rangle = \int_A \tilde{f}_4(u - z)\tilde{f}_2(z)\tilde{f}_0(x - z)\langle z \rangle dz, \quad \text{for all } (x, u) \in A^2, \tag{137}$$

where

$$\tilde{f}_i(x) := (F^{-1}f_i)(x) = \int_A \langle x; -y \rangle f_i(y) dy. \tag{138}$$

Defining

$$\hat{f}_i(x) := \tilde{f}_i(x)\langle x \rangle, \tag{139}$$

we rewrite (137) in the form

$$\hat{f}_1(x)\hat{f}_3(u) = \int_A \langle x - z; z - u \rangle \hat{f}_4(u - z)\hat{f}_2(z)\hat{f}_0(x - z) dz \quad \text{for all } (x, u) \in A^2, \tag{140}$$

We call a subgroup  $B \subset A$  isotropic if it satisfies the condition

$$\langle b; b' \rangle = 1, \quad \text{for all } (b, b') \in B^2. \tag{141}$$

**Lemma B.1** *Let a bi-character  $\chi : A^2 \rightarrow \mathbb{T}$  be such that*

$$\langle x; y \rangle = \chi(x, y)\chi(y, x), \quad \text{for all } (x, y) \in A^2, \tag{142}$$

and  $B \subset A$  an isotropic subgroup. Then, for any function  $f_i(x)$  of Faddeev type, the function

$$g : [4] \times A^2 \rightarrow \mathbb{C}, \quad (i, x, y) \mapsto g_i(x, y) := \chi(x, y) \int_B \hat{f}_i(x + b)\langle b; y \rangle db \tag{143}$$

is automorphic, i.e., satisfies

$$\begin{aligned} g_i(x + b, y) &= \chi(-y, b)g_i(x, y), & g_i(x, y + b) &= \chi(x, b)g_i(x, y), \\ &\text{for all } (i, b, x, y) \in [4] \times B \times A^2, \end{aligned} \tag{144}$$

and satisfies the integral identity

$$g_1(x, y)g_3(u, v) = \int_{A/B} g_4(u - z, v + z - x)g_2(z, y + v)g_0(x - z, y + z - u) dz \tag{145}$$

*Proof* The automorphicity properties (144) are verified in a straightforward manner, while to derive the integral identity (145), we write

$$\begin{aligned} &\frac{g_1(x, y)g_3(u, v)}{\chi(x, y)\chi(u, v)} \\ &= \int_{A \times B^2} \langle x + b - z; z - u - c \rangle \langle b; y \rangle \langle c; v \rangle \hat{f}_4(u + c - z)\hat{f}_2(z) \\ &\quad \times \hat{f}_0(x + b - z) d(z, b, c) \\ &= \int_{A \times B^2} \langle x - z; z - u \rangle \langle b; y + z - u \rangle \langle c; v + z - x \rangle \hat{f}_4(u + c - z)\hat{f}_2(z) \\ &\quad \times \hat{f}_0(x + b - z) d(z, b, c) \\ &= \int_A \frac{\langle x - z; z - u \rangle g_4(u - z, v + z - x)\hat{f}_2(z)g_0(x - z, y + z - u)}{\chi(u - z, v + z - x)\chi(x - z, y + z - u)} dz \\ &= \int_A \frac{g_4(u - z, v + z - x)\hat{f}_2(z)g_0(x - z, y + z - u)}{\chi(u - z, v)\chi(x - z, y)} dz \end{aligned} \tag{146}$$



so that

$$\begin{aligned}
 g_1(x, y)g_3(u, v) &= \int_A \chi(z, y + v)g_4(u - z, v + z - x)\hat{f}_2(z)g_0(x - z, y + z - u) dz \\
 &= \int_{(A/B) \times B} \chi(z + b, y + v)g_4(u - z - b, v + z + b - x)\hat{f}_2(z + b) \\
 &\quad \times g_0(x - z - b, y + z + b - u) d(z, b) \\
 &= \int_{(A/B) \times B} \chi(z + b, y + v)\chi(v + y, b)g_4(u - z, v + z - x)\hat{f}_2(z + b) \\
 &\quad \times g_0(x - z, y + z - u) d(z, b) \\
 &= \int_{A/B} g_4(u - z, v + z - x)g_2(z, y + v)g_0(x - z, y + z - u) dz. \tag{147}
 \end{aligned}$$

□

Now, we point out that the integral identity (145) is an equivalent form of the automorphic Beta pentagon identity. Namely, if we define

$$\phi_i(x, y) := g_i(-y, x + y) \Leftrightarrow g_i(x, y) = \phi_i(x + y, -x), \tag{148}$$

then

$$\phi_1(x, y)\phi_3(u, v) = \int_{A/B} \phi_4(u + y, v - z)\phi_2(x + y + u + v - z, z)\phi_0(x + v, y - z) dz. \tag{149}$$

In the case of the quantum dilogarithm  $\phi_q$  constructed in Appendix A, we have

$$A = \mathbb{T} \times \mathbb{Z}, \quad \langle z, m \rangle = z^m, \quad \langle z, m; w, n \rangle = z^m w^n \tag{150}$$

so that

$$\begin{aligned}
 \hat{\phi}_q(z, m) &= z^m \int_{\mathbb{T} \times \mathbb{Z}} \phi_q(t, k)\langle z, m; 1/t, -k \rangle d(t, k) = z^m \int_{\mathbb{T} \times \mathbb{Z}} \phi_q(t, k)z^{-k}t^{-m} d(t, k) \\
 &= z^m \int_{\mathbb{T} \times \mathbb{Z}} \psi(1/t, k)z^{-k}t^{-m} d(t, k) = z^m \int_{\mathbb{T}} \psi^0(1/t, 1/z)t^{-m} \frac{dt}{2\pi it}.
 \end{aligned}$$

We choose

$$B = \mathbb{Z} \subset \mathbb{T} \times \mathbb{Z} \tag{151}$$

and

$$\chi(\langle z, m \rangle, \langle w, n \rangle) = w^m, \tag{152}$$

so that the automorphic factors are trivial:

$$\chi(\langle z, m \rangle, \langle 1, n \rangle) = 1, \quad \text{for all } (z, m, n) \in \mathbb{T} \times \mathbb{Z}^2. \tag{153}$$

Thus,

$$\begin{aligned}
 g((z, m), (w, n)) &= \chi((z, m), (w, n)) \sum_{k \in \mathbb{Z}} \hat{\phi}_q(z, m+k) \langle 1, k; w, n \rangle \\
 &= w^m \sum_{k \in \mathbb{Z}} \hat{\phi}_q(z, m+k) w^k = \sum_{k \in \mathbb{Z}} \hat{\phi}_q(z, k) w^k \\
 &= \int_{\mathbb{T}} \sum_{k \in \mathbb{Z}} (zw/t)^k \psi^0(1/t, 1/z) \frac{dt}{2\pi it} \\
 &= \int_{\mathbb{T}} \delta_{\mathbb{T}}(zw/t) \psi^0(1/t, 1/z) \frac{dt}{2\pi it} = \psi^0(1/(zw), 1/z)
 \end{aligned}$$

and

$$\phi((z, m), (w, n)) = g((1/w, -n), (zw, m+n)) = \psi^0(1/z, w). \quad (154)$$

Taking into account the symmetry of the Beta pentagon identity under the negation of all arguments, we conclude that  $\psi^0(1/z, w)$  and  $\psi^0(z, 1/w)$  both satisfy the Beta pentagon identity.

Received: 31 August 2017 Accepted: 19 October 2018 Published online: 3 December 2018

## References

- Ahlfors, L.: Complex analysis. In: An Introduction to the Theory of Analytic Functions of One Complex Variable, International Series in Pure and Applied Mathematics, 3rd ed., McGraw-Hill Book Co., New York (1978)
- Andersen, J.E., Kashaev, R.: Complex Quantum Chern–Simons (2014). Preprint [arXiv:1409.1208](https://arxiv.org/abs/1409.1208)
- Andersen, J.E., Kashaev, R.: A new formulation of the Teichmüller TQFT (2013). Preprint [arXiv:1305.4291](https://arxiv.org/abs/1305.4291)
- Andersen, J.E., Kashaev, R.: A TQFT from quantum Teichmüller theory. *Commun. Math. Phys.* **330**(3), 887–934 (2014)
- Burton, B.: Regina, software for 3-manifold topology and normal surface theory. <http://regina.sourceforge.net>. Accessed 30 Jan 2015
- Culler, M., Dunfield, N., Weeks, J.: SnapPy, a computer program for studying the topology of 3-manifolds. <http://snappy.computop.org>. Accessed 30 Jan 2015
- Dimofte, T., Garoufalidis, S.: The quantum content of the gluing equations. *Geom. Topol.* **17**(3), 1253–1315 (2013)
- Dimofte, T., Gaiotto, D., Gukov, S.: 3-manifolds and 3d indices. *Adv. Theor. Math. Phys.* **17**(5), 975–1076 (2013)
- Dimofte, T., Gaiotto, D., Gukov, S.: Gauge theories labelled by three-manifolds. *Commun. Math. Phys.* **325**(2), 367–419 (2014)
- Garoufalidis, S.: The 3D index of an ideal triangulation and angle structures. *Ramanujan J.* **40**(3), 573–604 (2016)
- Garoufalidis, S., Hodgson, C., Hoffman, N., Rubinstein, H.: The 3D-index and normal surfaces. III. *J. Math.* **60**(1), 289–352 (2016)
- Garoufalidis, S., Hodgson, C., Rubinstein, H., Segerman, H.: 1-efficient triangulations and the index of a cusped hyperbolic 3-manifold. *Geom. Topol.* **19**(5), 2619–2689 (2015)
- Garoufalidis, S., Kashaev, R.: From state integrals to  $q$ -series. *Math. Res. Lett.* **24**(3), 781–801 (2017)
- Kashaev, R.: The  $q$ -Binomial Formula and the Rogers Dilogarithm Identity (2004). Preprint [arXiv:math/0407078](https://arxiv.org/abs/math/0407078)
- Kashaev, R.: The Yang–Baxter relation and gauge invariance (2015). Preprint [arXiv:1510.03043](https://arxiv.org/abs/1510.03043)
- Kashaev, R.: Beta pentagon relations. *Theoret. and Math. Phys.* **181**(1), 1194–1205 (2014) (Russian version appears in *Theoret. Mat. Fiz.* **1**, 81 (2014), **1**, 73–85)
- Kashaev, R., Luo, F., Vartanov, G.: A TQFT of Turaev–Viro type on shaped triangulations. *Ann. Henri Poincaré* **17**(5), 1109–1143 (2016)
- Luo, F., Tillmann, S.: Angle structures and normal surfaces. *Trans. Am. Math. Soc.* **360**(6), 2849–2866 (2008)
- Neumann, W., Zagier, D.: Volumes of hyperbolic three-manifolds. *Topology* **24**(3), 307–332 (1985)
- Thurston, W.: The geometry and topology of 3-manifolds, Universitext. Springer, Berlin (1977). <http://msri.org/publications/books/gt3m>
- Woronowicz, S.: Operator equalities related to the quantum  $E(2)$  group. *Commun. Math. Phys.* **144**(2), 417–428 (1992)
- Zagier, D.: The Dilogarithm Function, *Frontiers in Number Theory, Physics, and Geometry*, II, pp. 3–65. Springer, Berlin (2007)