

REIDEMEISTER CLASSES IN LAMPLIGHTER TYPE GROUPS

EVGENIJ TROITSKY

ABSTRACT. We prove that for any automorphism ϕ of the restricted wreath product $\mathbb{Z}_2 \text{wr} \mathbb{Z}^k$ and $\mathbb{Z}_3 \text{wr} \mathbb{Z}^{2d}$ the Reidemeister number $R(\phi)$ is infinite, i.e. these groups have the property R_∞ .

For $\mathbb{Z}_3 \text{wr} \mathbb{Z}^{2d+1}$ and $\mathbb{Z}_p \text{wr} \mathbb{Z}^k$, where $p > 3$ is prime, we give examples of automorphisms with finite Reidemeister numbers. So these groups do not have the property R_∞ .

For these groups and $\mathbb{Z}_m \text{wr} \mathbb{Z}$, where m is relatively prime to 6, we prove the twisted Burnside-Frobenius theorem (TBFT_f): if $R(\phi) < \infty$, then it is equal to the number of equivalence classes of finite-dimensional irreducible unitary representations fixed by the action $[\rho] \mapsto [\rho \circ \phi]$.

INTRODUCTION

The *Reidemeister number* $R(\phi)$ of an automorphism ϕ of a (countable discrete) group G is the number of its *Reidemeister* or *twisted conjugacy classes*, i.e. the classes of the twisted conjugacy equivalence relation: $g \sim hg\phi(h^{-1})$, $h, g \in G$. Denote by $\{g\}_\phi$ the class of g .

The following two interrelated problems are in the mainstream of the study of Reidemeister numbers.

In [5] A.Fel'shtyn and R.Hill conjectured that $R(\phi)$ is equal to the number of fixed points of the associated homeomorphism $\hat{\phi}$ of the unitary dual \hat{G} (the set of equivalence classes of irreducible unitary representations of G), if one of these numbers is finite. The action of $\hat{\phi}$ on the class of a representation ρ is defined as $[\rho] \mapsto [\rho \circ \phi]$. This conjecture is called TBFT (twisted Burnside-Frobenius theorem). This statement can be considered as a generalization to infinite groups and to the twisted case the classical Burnside-Frobenius theorem: the number of conjugacy classes of a finite group is equal to the number of equivalence classes of its irreducible representations. The TBFT conjecture (more precisely, its modification TBFT_f, taking into account only finite-dimensional representations) was proved for polycyclic-by-finite groups in [9, 13]. Preliminary and related results, examples and counter-examples can be found in [5, 8, 10, 6, 26, 11, 15, 27].

Also A.Fel'shtyn and co-authors formulated the second problem (a historical overview can be found in [7]): the problem of description of the class of groups having the following R_∞ property: $R(\phi) = \infty$ for any automorphism $\phi : G \rightarrow G$. Thus, the second problem is in some sense complementary to the first one: the question about TBFT has no sense for R_∞ groups (formally having a positive answer). The property R_∞ was studied very intensively during the last years and was proved and disproved for many groups (see a bibliography overview in [7] and [15], and very recent papers [3, 24, 12] and the literature therein). For Jiang type spaces the property R_∞ has some direct topological consequences (see e.g.

2000 *Mathematics Subject Classification.* 20C; 20E45; 20E22; 20E26; 20F16; 22D10.

Key words and phrases. Reidemeister number, R_∞ -group, twisted conjugacy class, residually finite group, restricted wreath product, lamplighter group.

This work is supported by the Russian Science Foundation under grant 16-11-10018.

[19]). Relations with group growth are discussed e.g. in [17]. Concerning applications of Reidemeister numbers in Dynamics we refer to [20, 4].

The R_∞ property was studied for the lamplighter group $\mathbb{Z}_2 \text{wr} \mathbb{Z}$ and some its generalizations being restricted wreath products with \mathbb{Z} in [18, 25, 24]. In particular, in [18] it was proved that most part of groups of the form $\mathbb{Z}_q \text{wr} \mathbb{Z}$ are not R_∞ groups. More precisely, it is an R_∞ group if and only if $(q, 6) \neq 0$. In contrast with this result, in [24] it is proved that the generalizations $\Gamma_d(q)$ of the lamplighter group always have the R_∞ property for $d > 2$ (for $d = 2$ one has $\Gamma_d(q) \cong \mathbb{Z}_q \text{wr} \mathbb{Z}$). The groups $\Gamma_d(q)$ admit a Cayley graph isomorphic to a Diestel-Leader graph $DL_d(q)$. The lamplighter group and its generalizations attracted a lot of attention recently, in particular due to its relations with automata groups, self-similar groups, and branch groups (see e.g. [1]).

For groups under consideration in the present paper, even for $k = 2$, the situation is much more complicated, because \mathbb{Z} has only one automorphism with finite Reidemeister number, namely $-\text{Id}$, and its square has infinite Reidemeister number, but for $\mathbb{Z} \oplus \mathbb{Z}$ we have a lot of automorphisms with finite Reidemeister numbers, and many of them have finite Reidemeister numbers for all their iterations (see, e.g. [4]).

In the present paper we prove that for any automorphism ϕ of the restricted wreath product $\mathbb{Z}_2 \text{wr} \mathbb{Z}^k$ (Theorem 2.3) and $\mathbb{Z}_3 \text{wr} \mathbb{Z}^{2d}$ (Theorem 4.1) the Reidemeister number $R(\phi)$ is infinite, i.e. these groups have the property R_∞ .

For $\mathbb{Z}_3 \text{wr} \mathbb{Z}^{2d+1}$ (Theorem 4.1) and $\mathbb{Z}_p \text{wr} \mathbb{Z}^k$, where $p > 3$ is prime, (Theorem 3.5) we give examples of automorphisms with finite Reidemeister numbers. So these groups do not have the property R_∞ .

For these groups and $\mathbb{Z}_m \text{wr} \mathbb{Z}$, where m is relatively prime to 6, we prove in Theorem 5.1 the twisted Burnside-Frobenius theorem (TBFT_f): if $R(\phi) < \infty$, then it is equal to the number of equivalence classes of finite-dimensional irreducible unitary representations fixed by $\hat{\phi} : \rho \mapsto \rho \circ \phi$.

This gives (probably first) examples of finitely generated residually finite but not almost polycyclic groups (having infinitely generated subgroup, [23, p. 4]), for which the TBFT is true.

ACKNOWLEDGEMENT: The author is indebted to A. Fel'shtyn for helpful discussions in the Max-Planck Institute for Mathematics (Bonn) in February, 2017 and the MPIM for supporting this visit.

The author is grateful to L. Alania, R. Jimenez Benitez, and V. Manuilov for valuable advises and suggestions.

This work is supported by the Russian Science Foundation under grant 16-11-10018.

1. PRELIMINARIES

The following easy statement is well known:

Proposition 1.1. *Suppose, H is a ϕ -invariant normal subgroup of G and $\bar{\phi} : G/H \rightarrow G/H$ is the induced automorphism. Then ϕ induces an epimorphism of each Reidemeister class of ϕ onto some Reidemeister class of $\bar{\phi}$. In particular, one has $R(\bar{\phi}) \leq R(\phi)$.*

Denote by $C(\phi)$ the fixed point subgroup. The following much more non-trivial statement can be extracted from [16] (see also [9]):

Lemma 1.2. *In the above situation $R(\phi|_H) \leq R(\phi) \cdot |C(\bar{\phi})|$.*

It is well known (see [8]) the following.

Lemma 1.3. *For an abelian group G the Reidemeister class of the unit element is a subgroup, and the other classes are corresponding cosets.*

The following statement is very useful in the field.

Lemma 1.4. *A right shift by $g \in G$ maps Reidemeister classes of ϕ onto Reidemeister classes of $\tau_{g^{-1}} \circ \phi$, where τ_g is the inner automorphism: $\tau_g(x) = gxg^{-1}$. In particular, $R(\tau_g \circ \phi) = R(\phi)$.*

Proof. Indeed,

$$xy\phi(x^{-1})g = x(yg)g^{-1}\phi(x^{-1})g = x(yg)(\tau_{g^{-1}} \circ \phi)(x^{-1}).$$

□

Also we need the following statement ([14], [11, Prop. 3.4]):

Lemma 1.5. *Let $\phi : G \rightarrow G$ be an automorphism of a finitely generated residually finite group G with $R(\phi) < \infty$ (in particular, G can be a finitely generated abelian group). Then the subgroup of fixed elements is finite: $|C(\phi)| < \infty$.*

Note, that this is not correct for infinitely generated groups, see [27].

Combining this lemma with some results of [16] one can prove:

Lemma 1.6. *Suppose in the situation of Lemma 1.1 that G/H is a finitely generated residually finite group. Then $R(\phi) < \infty$ if and only if $R(\bar{\phi}) < \infty$ and $R(\tau_g \phi') < \infty$ for any $g \in G$.*

2. THE CASE OF $\mathbb{Z}_2 \text{ wr } \mathbb{Z}^k$

Let $\Gamma := \mathbb{Z}_2 \text{ wr } \mathbb{Z}^k$ be a restricted wreath product. In other words,

$$\Gamma = \bigoplus_{x \in \mathbb{Z}^k} (\mathbb{Z}_2)_{(x)} \rtimes_{\alpha} \mathbb{Z}^k, \quad (\mathbb{Z}_2)_{(x)} \cong \mathbb{Z}_2, \quad \alpha(y)(\delta_x) := \delta_{y+x},$$

where $y \in \mathbb{Z}^k$ and δ_x is a unique non-trivial element of $(\mathbb{Z}_2)_{(x)} \subset \Gamma$. The direct sum supposes only finitely many non-trivial components for each element (in contrast with the direct product corresponding to the (unrestricted) wreath product).

The group Γ is a finitely generated metabelian group, in particular, residually finite (see e.g. [22]).

Let $\phi : \Gamma \rightarrow \Gamma$ be an automorphism. We will prove that $R(\phi) = \infty$. Denote $\Sigma := \bigoplus_{x \in \mathbb{Z}^k} (\mathbb{Z}_2)_{(x)} \subset \Gamma$. Then Σ is a characteristic subgroup as the torsion subgroup. Denote the restriction of ϕ by $\phi' : \Sigma \rightarrow \Sigma$, and the quotient automorphism by $\bar{\phi} : \mathbb{Z}^k \rightarrow \mathbb{Z}^k$.

If $R(\phi) < \infty$, then $R(\bar{\phi}) < \infty$ by Proposition 1.1. Hence, by Lemma 1.5, $\bar{\phi}$ has finitely many fixed elements. Thus, by Lemma 1.2, $R(\phi') < \infty$. Hence, to prove that $R(\phi) = \infty$, it is sufficient to verify that $R(\phi') = \infty$.

Since Σ is abelian, the results of e.g. [2] imply that

$$(1) \quad \phi'(\alpha(g)(h)) = \alpha(\bar{\phi}(g))(\phi'(h)), \quad h \in \Sigma, \quad g \in \mathbb{Z}^k.$$

Any element of Σ is a finite sum of some elements δ_x . Let

$$(2) \quad \phi'(\delta_0) = \delta_{x(1)} + \cdots + \delta_{x(n)}.$$

The following lemma generalizes [18, Prop. 2.1] from the case $k = 1$ to arbitrary k .

Lemma 2.1. *In (2) one has $n = 1$. Moreover, ϕ' is a permutation of δ_x 's.*

Proof. First of all, apply (1) to $h = \delta_0$. We have:

$$(3) \quad \phi'(\delta_g) = \phi'(\alpha(g)(h)) = \alpha(\overline{\phi}(g))(\phi'(\delta_0)).$$

Thus, for any $g \in \mathbb{Z}^k$, the element $\phi'(\delta_g)$ is obtained by the appropriate shift of indexes in the right side expression in (2).

Now suppose that $n \neq 1$, and $\phi'(h) = \delta_0$ for some $h = \delta_{r(1)} + \dots + \delta_{r(t)}$. Then $t \neq 1$, because the statement of the lemma for ϕ' and its inverse are equivalent. Denote by $T \subset \mathbb{Z}^k$ the support of $\phi'(\delta_0)$, i.e.,

$$T = \{x(1), \dots, x(n)\}.$$

Denote by T_1, \dots, T_t the supports of $\phi'(\delta_{r(1)}), \dots, \phi'(\delta_{r(t)})$ respectively. They are appropriate distinct shifts of T . Denote $S := T_1 \cup \dots \cup T_t$ (without cancellations). After cancellations in Σ (i.e. excluding of points in S covered by an even number of T_j 's) we should obtain only one point, namely, 0.

Introduce now the notion of $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)$ -vertex $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[R] \in R$ for any bounded subset $R \subset \mathbb{Z}^k$, where $(\sigma_1, \dots, \sigma_k)$ is a permutation of $(1, \dots, k)$ and $\varepsilon_i = \pm 1$. We define it inductively in the following way: R_{k-1} is the subset of points of R with minimal (if $\varepsilon_1 = -1$) or maximal (if $\varepsilon_1 = +1$) coordinate number σ_1 , R_{k-2} is the subset of points of R_{k-1} with minimal (if $\varepsilon_2 = -1$) or maximal (if $\varepsilon_2 = +1$) coordinate number σ_2 , and so on. Then R_0 is one point. This point we define to be $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[R]$. This point also can be considered as a lexicographic maximum of points of R for the ordering σ and the inverse direction of that coordinates, where $\sigma_j = -1$, i.e. a *lexicographic maximum with respect to* $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)$.

Evidently, $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_j]$ is a \mathbb{Z}^k -shift of $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T]$ and $T_j = T_i$ if and only if

$$(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_j] = (\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_i]$$

for at least one (thus, for any) $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)$. Hence, in our situation, they are distinct.

We claim that for any $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)$ the vertex $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[S]$ coincides with $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_j]$ for one and only one j and is not covered by other points. Indeed, the uniqueness follows from the argument above. If it is covered by some point of T_j other than $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_j]$, then it would be not the lexicographic maximum w.r.t. $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)$, because $(\varepsilon_1\sigma_1, \varepsilon_2\sigma_2, \dots, \varepsilon_k\sigma_k)[T_j] \in S$ would be greater.

Thus no vertex will be canceled. Thus they all coincide with 0 and $S = \{0\}$. Hence, $n = r = 1$.

Together with the argument at the beginning of the proof, this gives the second statement. \square

By this lemma, we can define $x_0 \in \mathbb{Z}^k$ by $\phi'(\delta_0) = \delta_{x_0}$. Equation (3) can be written now as

$$(4) \quad \phi'(\delta_y) = \delta_{y'}, \quad y' := \overline{\phi}(y) + x_0 \in \mathbb{Z}^k.$$

Lemma 2.2. *If δ_{x_1} and δ_{x_2} belong to the same Reidemeister class of ϕ' , then*

$$(5) \quad \overline{\phi}^t(x_1) + \overline{\phi}^{t-1}(x_0) + \dots + \overline{\phi}(x_0) + x_0 = x_2$$

or

$$(6) \quad \overline{\phi}^t(x_2) + \overline{\phi}^{t-1}(x_0) + \dots + \overline{\phi}(x_0) + x_0 = x_1$$

for some integer t .

Proof. By Lemma 1.3, the elements δ_{x_1} and δ_{x_2} belong to the same Reidemeister class of ϕ' if and only if $\delta_{x_1} - \delta_{x_2} = h - \phi'(h)$ for some $h \in \Sigma$. Representing h as $h = \delta_{u(1)} + \cdots + \delta_{u(t)}$ (with distinct summands) and applying (4) one has

$$\delta_{x_1} - \delta_{x_2} = h - \phi'(h) = \sum_{j=1}^t [\delta_{u(j)} - \delta_{u(j)}'].$$

This is the same in Σ as

$$\delta_{x_1} + \delta_{x_2} = \sum_{j=1}^t [\delta_{u(j)} + \delta_{u(j)}'].$$

Since all $\delta_{u(j)}$ are distinct, all $\delta_{u(j)'}$ are distinct too, by Lemma 2.1. So the cancellation on the right can be only when $\delta_{u(j)} = \delta_{u(i)'}$. So one of $\delta_{u(j)}$ should be equal to δ_{x_1} , one of $\delta_{u(i)'}$ should be equal to δ_{x_2} (or vice versa), and the remaining δ 's should annihilate. Thus, after the appropriate renumbering of $1, \dots, t$, we have in the first case:

$$x_1 = u(1), \quad u(1)' = u(2), \quad \dots \quad u(t-1)' = u(t), \quad u(t)' = x_2,$$

or

$$\begin{aligned} \overline{\phi}(x_1) + x_0 &= u(2), \\ \overline{\phi}^2(x_1) + \overline{\phi}(x_0) + x_0 &= u(2)' = u(3), \\ \overline{\phi}^3(x_1) + \overline{\phi}^2(x_0) + \overline{\phi}(x_0) + x_0 &= u(3)' = u(4), \\ &\dots \dots \dots \\ \overline{\phi}^t(x_1) + \overline{\phi}^{t-1}(x_0) + \cdots + \overline{\phi}(x_0) + x_0 &= u(t)' = x_2. \end{aligned}$$

In the second case we need to interchange x_1 and x_2 :

$$\overline{\phi}^t(x_2) + \overline{\phi}^{t-1}(x_0) + \cdots + \overline{\phi}(x_0) + x_0 = x_1.$$

□

Theorem 2.3. *The group $\Gamma = \mathbb{Z}_2 \text{ wr } \mathbb{Z}^k$ has the property R_∞ .*

Proof. One can reduce the proof of $R(\phi) = \infty$ to the case $x_0 = 0$. Indeed, consider the element $w := -x_0 \in \mathbb{Z}^k \subset \Gamma$ and the corresponding inner automorphism $\tau_w : \Gamma \rightarrow \Gamma$. Then by Lemma 1.4, $R(\tau_w \circ \phi) = R(\phi)$. On the other hand, by the definition of a semidirect product,

$$(\tau_w \circ \phi)'(\delta_0) = \alpha(w)(\phi'(\delta_0)) = \alpha(-x_0)(\delta_{x_0}) = \delta_0.$$

So, suppose $x_0 = 0$. Then (5) and (6) take the form $\overline{\phi}^t(x_1) = x_2$ for some integer t . Thus, it is sufficient to prove that $\overline{\phi} : \mathbb{Z}^k \rightarrow \mathbb{Z}^k$ has infinitely many orbits.

For this purpose denote by $A \in GL_k(\mathbb{Z})$ the matrix of $\overline{\phi}$. Let us show that each orbit intersects the first coordinate axis not more than in 2 points. Denote by $(x, 0, \dots, 0) = x \cdot e_1$, $x \neq 0$, one point from the intersection, and let $\overline{\phi}^n(x \cdot e_1) = x \cdot \overline{\phi}^n(e_1) = x \cdot y \cdot e_1$ be another intersection point. Since A^n as an element of $GL_k(\mathbb{Z})$ has the first column (r_1, \dots, r_k) such that $\gcd(r_1, \dots, r_k) = 1$ (because the expansion of the determinant by the first column has the form $\pm 1 = r_1 \cdot R_1 + \cdots + r_k \cdot R_k$). Thus, $y = \pm 1$ and $xy = \pm x$. □

3. THE CASE OF \mathbb{Z}_p wr \mathbb{Z}^k FOR $p > 3$

A part of argument in this section will be close to some argument of [18]. Suppose now that $\Gamma = \mathbb{Z}_p \text{ wr } \mathbb{Z}^k$ for a general prime p . We conserve the notation Σ for the normal subgroup $\oplus \mathbb{Z}_p$.

Now δ_x is a generator of a subgroup A_x isomorphic to \mathbb{Z}_p , and $p \cdot \delta_x = 0$. Suppose

$$(7) \quad \phi'(\delta_0) = m_1 \delta_{x(1)} + \cdots + m_n \delta_{x(n)}.$$

Then as above,

$$(8) \quad \phi'(\delta_x) = m_1 \delta_{\overline{\phi(x)+x(1)}} + \cdots + m_n \delta_{\overline{\phi(x)+x(n)}}.$$

First of all we need an analog of Lemma 2.1.

In the general situation instead of the sets T_j we need $(T_j, s_j \cdot \vec{m})$, which are some shifts with multiplication of (T, \vec{m}) , where $\vec{m} = (m_1, \dots, m_n)$. Fortunately (and that is why we have restricted ourselves to the prime order case) if $m \in \mathbb{Z}_p$, $m \neq 0$, then it generates \mathbb{Z}_p . That is why the sum of several elements with the same $T_{j(1)} = \cdots = T_{j(r)}$ either has the same support $= T_{j(1)}$ and coefficients vector $(m_{j(1)} + \cdots + m_{j(r)})\vec{m}$, or completely annihilates, when $m_{j(1)} + \cdots + m_{j(r)} = 0 \pmod p$.

So, after cancellations we may assume that all supports T_j are distinct and repeat the remaining part of the proof of Lemma 2.1 and obtain

Lemma 3.1. *If p is prime, one has*

$$\phi'(\delta_0) = m \cdot \delta_{x_0}$$

for some $x_0 \in \mathbb{Z}^k$ and $0 \neq m \in \mathbb{Z}_p$.

Let us note, that generally $m \neq 1$ in this situation. For example, in \mathbb{Z}_3 one can take $s = m = 2$ and $sm = 4 = 1 \pmod 3$.

For other elements we have

$$(9) \quad \phi'(\delta_x) = m \delta_{\overline{\phi(x)+x_0}}.$$

To calculate $R(\phi')$ we need to calculate the index of the image of $(1 - \phi')\Sigma$ in Σ .

Suppose first that $x_0 = 0$,

$$\phi'(\delta_x) = m \delta_{\overline{\phi(x)}}.$$

Then for any x we have $(1 - \phi')$ -invariant subgroup

$$(10) \quad \cdots \oplus A_{\overline{\phi^{-1}(x)}} \oplus A_x \oplus A_{\overline{\phi(x)}} \oplus A_{\overline{\phi^2(x)}} \oplus \cdots.$$

In contrast with the case $k = 1$ considered in [18], the corresponding orbit of $\overline{\phi}$ can be infinite or finite, but not necessary of length 2.

Let us note that since $\tau_g \overline{\phi} = \overline{\phi}$, they have the same orbit structure.

Lemma 3.2. *If an orbit is infinite, then the corresponding restriction of $1 - \phi'$ on the subgroup (10) is not an epimorphism.*

Proof. Indeed, under the appropriate description,

$$1 - \phi' : (\dots, 0, 0, a_1, a_2, \dots, a_r, 0, 0, \dots) \mapsto (\dots, 0, -ma_1, a_1 - ma_2, a_2 - ma_3, \dots, a_r, 0, 0, \dots)$$

If $a_1 \neq 0$ and $a_r \neq 0$, then $-ma_1 \neq 0$ and the length of non-trivial part increases. Thus, elements concentrated in one summand, e.g. δ_x , are not in the image. \square

If the orbit is finite of length s , the matrix of $(1 - \phi')$ has the form

$$(11) \quad E - M = \begin{pmatrix} 1 & 0 & \cdots & 0 & -m \\ -m & 1 & 0 & & 0 \\ 0 & -m & 1 & 0 & \\ 0 & 0 & -m & 1 & \ddots & \vdots \\ \vdots & & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & \cdots & 0 & -m & 1 \end{pmatrix}$$

and is an epimorphism if and only if its determinant is not zero (for prime p):

$$(12) \quad \det(E - M) = 1 - m^s \not\equiv 0 \pmod{p}.$$

Lemma 3.3. *For a non-trivial orbit of $\bar{\phi} : \mathbb{Z}^k \rightarrow \mathbb{Z}^k$, there exist infinitely many orbits of the same cardinality (finite or infinite).*

If $R(\bar{\phi}) < \infty$ there is a unique trivial orbit.

Proof. Let $(r_1, \dots, r_k) \neq 0$ be a point of the orbit. Since $\bar{\phi}$ as an element of $GL(\mathbb{Z}, k)$ preserves $\gcd(r_1, \dots, r_k)$, the elements $(i \cdot r_1, \dots, i \cdot r_k)$, $i \in \mathbb{Z}$, $i > 2$, belong to distinct orbits of the same cardinality.

If there is a non-zero fixed point x_* of ϕ' , then there is an infinite series $\{s \cdot x_*\}$, $s \in \mathbb{Z}$, of fixed points. By Lemma 1.5 this contradicts to $R(\bar{\phi}) < \infty$. \square

Lemma 3.4. *If $\bar{\phi}$ has an infinite orbit, $R(\phi') = \infty$ and $R(\phi) = \infty$.*

If $\bar{\phi}$ has only finite orbits, there are two possibilities:

- 1) $R(\phi') = R(\phi) = \infty$. *This occurs if, at least for one orbit, the corresponding restriction of $1 - \phi'$ is not an epimorphism.*
- 2) $R(\phi') = 1$. *This occurs if, for all orbits, the corresponding restriction of $1 - \phi'$ is an epimorphism.*

If we have one of this cases for ϕ' , then the same is true for all $\tau_g \circ \phi'$.

Proof. Let us note that $\tau_g = \tau_{g'} : \Sigma \rightarrow \Sigma$ if $g^{-1}g' \in \Sigma$, so all automorphisms $\tau_g \circ \phi' : \Sigma \rightarrow \Sigma$ are described by $g \in \mathbb{Z}^k$. In this case $\tau_g \circ \phi' = \alpha(g) \circ \phi'$ and the sizes of above invariant groups are the same as for ϕ' .

As it was explained above all possible automorphisms ϕ' corresponding to a given $\bar{\phi} = \tau_g \circ \bar{\phi}$ are completely defined by m and x_0 such that $\phi'(\delta_0) = m \cdot \delta_{x_0}$, and each pair (m, x_0) defines some ϕ' and ϕ . The relation

$$\tau_{y_0} \circ \phi'(\delta_0) = \alpha(y_0) \circ \phi'(\delta_0) = m \cdot \delta_{y_0 + x_0}$$

shows that all automorphisms of Σ with the same m differ from each other by an appropriate τ_{y_0} and vice versa.

Hence, if $\bar{\phi}$ has an infinite orbit and we take for ϕ , some $g \in \mathbb{Z}^k$ such that $\tau_g \circ \phi'(\delta_0) = m \cdot \delta_0$, then there exists an appropriate invariant subgroup of ϕ' (over this orbit) with a non-epimorphic restriction of $1 - \tau_g \circ \phi'$ (Lemma 3.2). Since we have infinitely many such orbits (Lemma 3.3) then $R(\phi') = \infty$ and $R(\phi) = \infty$.

Now we will describe the matrix form of restrictions onto invariant subgroups of an arbitrary ϕ' (i.e. not necessary $x_0 = 0$) for the case of finite orbits of $\bar{\phi}$. Suppose, $\phi'(\delta_0) = m\delta(x_0)$, s is the length of the orbit $x_1, \bar{\phi}(x_1), \dots, \bar{\phi}^{s-1}(x_1)$, t is the length of the

orbit $x_0, \bar{\phi}(x_0), \dots, \bar{\phi}^{t-1}(x_0)$, we have

$$\phi'(\delta_{x_1}) = \alpha(\bar{\phi}(x_1))\phi'(\delta_0) = m \cdot \delta_{\bar{\phi}(x_1)+x_0},$$

$$\phi'(\delta_{\bar{\phi}(x_1)+x_0}) = \alpha(\bar{\phi}(\bar{\phi}(x_1) + x_0))\phi'(\delta_0) = m \cdot \delta_{\bar{\phi}^2(x_1)+\bar{\phi}(x_0)+x_0},$$

$$\phi'(\delta_{\bar{\phi}^2(x_1)+\bar{\phi}(x_0)+x_0}) = \alpha(\bar{\phi}(\bar{\phi}^2(x_1) + \bar{\phi}(x_0) + x_0))\phi'(\delta_0) = m \cdot \delta_{\bar{\phi}^3(x_1)+\bar{\phi}^2(x_0)+\bar{\phi}(x_0)+x_0}, \dots$$

In order to estimate the length r of the underlying orbit

$$(13) \quad x_1 \mapsto \bar{\phi}(x_1) + x_0 \mapsto \bar{\phi}^2(x_1) + \bar{\phi}(x_0) + x_0 \mapsto \bar{\phi}^3(x_1) + \bar{\phi}^2(x_0) + \bar{\phi}(x_0) + x_0 \mapsto \dots$$

remark that

$$(14) \quad y := \bar{\phi}^{t-1}(x_0) + \bar{\phi}^{t-2}(x_0) + \dots + x_0 = 0.$$

Indeed,

$$\bar{\phi}(y) - y = \bar{\phi}^t(x_0) - x_0 = 0.$$

Hence, if $y \neq 0$, we have a non-trivial fixed point for $\bar{\phi}$. This contradicts $R(\bar{\phi}) < \infty$ (as in the proof of Lemma 3.3).

Equality (14) and the definition of s imply

$$(15) \quad x_1 = \bar{\phi}^r(x_1) + \bar{\phi}^{r-1}(x_0) + \dots + \bar{\phi}(x_0) + x_0, \text{ where } r = \text{lcm}(s, t).$$

Suppose now that $\bar{\phi}$ does not have infinite orbits. Then the length of any orbit is bounded by $\text{lcm}(l_1, \dots, l_k)$, where l_i is the length of the orbit of the element e_i of the standard base. More precisely, any orbit length is a divisor of $\text{lcm}(l_1, \dots, l_k)$. In particular, for any s and t , as above, $r = \text{lcm}(t, s)$ is a divisor of $\text{lcm}(l_1, \dots, l_k)$. Hence, by (15) the length r' of (13) is some divisor of r and so of $\text{lcm}(l_1, \dots, l_k)$.

If the underlying orbit (13) does not start from the point $x_1 = 0$, one can obtain infinitely many underlying orbits by multiplying x_1 by different positive integers $j = 1, 2, \dots$. More precisely, some of them can coincide, but for a sufficiently large j , the point jx_1 will not be an element of the orbit (13), etc. So, there is infinitely many distinct orbits. Moreover, we can find a sufficiently large J such that for any $j \geq J$, the distances between jx_1 and $\bar{\phi}(j \cdot x_1) = j \cdot \bar{\phi}(x_1)$ are more than $x_0, \bar{\phi}(x_0) + x_0, \dots, \bar{\phi}^{t-1}(x_0) + \dots + \bar{\phi}(x_0) + x_0$. Evidently, for these orbits the length $r' = r$ (not only a divisor of r). So among the orbits starting in $x_1, 2x_1, \dots$, we have infinitely many orbits of length r and some finite number $< J$ of orbits of some length dividing r . Then the Reidemeister number of the restriction on the subgroup over the union of these orbits is finite if and only if $1 - m^r \not\equiv 0 \pmod{p}$. But then for any divisor r' of r we have, for $r'' := r/r'$,

$$1 - m^r = (1 - m^{r'})(1 + m^{r'} + m^{2r'} + \dots + m^{r'-1}) \not\equiv 0 \pmod{p}.$$

Hence, $1 - m^{r'} \not\equiv 0 \pmod{p}$. Thus, $1 - \phi'$ is an epimorphism over the orbits from the above finite series too. In particular, for the ‘‘initial’’ orbit (13).

It remains to discuss the case of $x_1 = 0$. In this case the length of the orbit is t . Considering some $x_1 \neq 0$ with the length of $\bar{\phi}$ -orbit equal to some $s \neq 0$ we arrive as above to $1 - m^r \not\equiv 0 \pmod{p}$, where $r = \text{lcm}(s, t)$. Since t divides $r = \text{lcm}(s, t)$, we obtain $1 - m^t \not\equiv 0 \pmod{p}$ similarly to the case of r' above. Thus $1 - \phi'$ is an epimorphism over this orbit too. Hence, it is an epimorphism in entire Σ .

As it was explained in the beginning of the proof, to vary ϕ' is the same as to consider various $\tau_g \circ \phi'$ for a fixed ϕ' . This completes the proof. \square

Theorem 3.5. *If $p > 3$, we can find an automorphism with finite Reidemeister number. Thus, $\mathbb{Z}_p \text{ wr } \mathbb{Z}^k$ does not have the property R_∞ if $p > 3$.*

Proof. For $p > 3$ consider $\bar{\phi} = -\text{Id}$. Then $R(\bar{\phi}) = 2^k$ and all non-trivial orbits are of length 2. Define ϕ' by $x_0 = 0$ and some non-zero $m \in \mathbb{Z}_p$, satisfying $1 - m^2 \not\equiv 0 \pmod{p}$. Take e.g. $m = 2$ (cf. [18, p. 879]). Then $1 - m^2 = -3 \not\equiv 0 \pmod{p}$ for any prime $p > 3$. \square

The intermediate case of $p = 3$ will be studied in the next section and the answer will depend on parity of k .

4. THE CASE OF $\mathbb{Z}_3 \text{ wr } \mathbb{Z}^k$

Theorem 4.1. *The group $\Gamma = \mathbb{Z}_3 \text{ wr } \mathbb{Z}^k$ has the property R_∞ for odd k and does not have the property R_∞ for even k .*

Proof. By Lemma 3.2, $R(\phi)$ may be finite only if all orbits of $\bar{\phi}$ are finite.

First, note that in this case m can be equal to 1 or 2.

If $m = 1$, then $1 - m^s \equiv 0 \pmod{3}$ not depending on the length s of the corresponding orbit. Keeping in mind the argument from the previous section, in particular, Lemma 3.3, we obtain $R(\phi) = \infty$.

If $m = 2$, then $1 - m^r \equiv 0 \pmod{3}$ for even r and $1 - m^s \not\equiv 0 \pmod{3}$ for odd r .

Denote by $M \in GL(k, \mathbb{Z})$ the matrix of $\bar{\phi}$. Since all orbits are finite, in particular, the orbits of the elements of the standard base, we have $M^r = E$ for some $r \in \mathbb{Z}$, $r > 1$ (see the previous section for more detail).

If $k = 2d + 1$, then $\det(M - \lambda E) = 0$ has at least one real solution λ_0 . It must be a root of 1 of degree r . Thus, $\lambda_0 = \pm 1$. If $\lambda_0 = 1$, then $\det(E - M) = 0$ and $R(\bar{\phi}) = \infty$. If $\lambda_0 = -1$, then r is even, and $\bar{\phi}$ has an orbit of even length $s = 2m$. Let t be the length of the $\bar{\phi}$ -orbit of x_0 , where $\phi'(\delta_0) = \delta_{x_0}$. Then, as in the proof of Lemma 3.4, we can detect infinitely many orbits of length $r = \text{lcm}(s, t)$. Since r is even, the restriction of $1 - \phi'$ on the subgroup, related the underlying orbit of the form (13) is not an epimorphism. Since there is infinitely many such orbits, $R(\phi') = \infty$. Then $R(\phi) = \infty$, as above.

If $k = 2$, consider M to be the generator $\begin{pmatrix} 0 & 1 \\ -1 & -1 \end{pmatrix}$ of a subgroup in $GL(2, \mathbb{Z})$ isomorphic to \mathbb{Z}_3 (see, e.g. [21, p. 179]). For this M , $R(\bar{\phi}) = \det(E - M) = 3$. It has only orbits of length 3 (except of the trivial one). The same is true for $\underbrace{M \oplus \cdots \oplus M}_d$ in \mathbb{Z}^{2d}

with $R(\bar{\phi}) = 3^d$. Then the lengths of all underlying orbits are some powers of 3 (except maybe of the trivial one) and $1 - \phi'$ is an epimorphism (for ϕ defined by this $\bar{\phi}$, $m = 2$, and arbitrary x_0). As in the previous section, this means that all $\tau_g \circ \phi'$ have $R(\tau_g \circ \phi') = 1$ and $R(\phi) = R(\bar{\phi}) < \infty$. \square

Remark 4.2. If $k = 1$, in particular, is odd, we obtain the R_∞ property for $\mathbb{Z}_3 \text{ wr } \mathbb{Z}$. This is a particular case of [18].

5. TWISTED BURNSIDE-FROBENIUS THEOREM

Theorem 5.1. *Suppose, Γ is $\mathbb{Z}_m \text{ wr } \mathbb{Z}$, where m is relatively prime to 6, or $\mathbb{Z}_p \text{ wr } \mathbb{Z}^k$ for a prime $p > 3$ and an arbitrary k , or $\mathbb{Z}_3 \text{ wr } \mathbb{Z}^{2d}$. Then the twisted Burnside-Frobenius theorem is true for Γ .*

Proof. In all these cases for ϕ with $R(\phi) < \infty$, we have $R(\tau_g \circ \phi') = 1$ (for prime cases this is proved in Lemma 3.4, for \mathbb{Z}_m wr \mathbb{Z} a similar statement can be easily extracted from the proof of Theorem 2.3 in [18]).

Evidently, $\{h\}_{\phi'} \subset \{h\}_{\phi}$, $h \in \Sigma$. Thus, $R(\phi') = 1$ implies that only one Reidemeister class of ϕ is mapped onto the class $\{e\}_{\bar{\phi}}$. By Lemma 1.4 $R(\tau_g \circ \phi') = 1$ implies the same for other classes $\{g\}_{\bar{\phi}}$. Thus, $\pi : \mathbb{Z}_p$ wr $\mathbb{Z}^k \rightarrow \mathbb{Z}^k$ induces a bijection of Reidemeister classes, or, speaking geometrically, the Reidemeister classes of ϕ are cylinders over Reidemeister classes of $\bar{\phi}$. Then $R = R(\phi) = R(\bar{\phi}) = \# \text{Fix}(\widehat{\bar{\phi}})$, because the TBFT is true for an automorphism of a finitely generated Abelian group (see [4]). If ρ_1, \dots, ρ_R are these representations, then $\rho_1 \circ \pi, \dots, \rho_R \circ \pi$ are some pairwise non-equivalent $\widehat{\bar{\phi}}$ -fixed representations of Γ . Then $R(\phi) \leq \# \text{Fix}(\widehat{\bar{\phi}})$. The opposite inequality is always true ([9, The proof of Theorem 5.2]).

An alternative argument is to see that, since the classes are cylinders, the dimension of the space of shifts of indicator functions of Reidemeister classes of ϕ is the same as for $\bar{\phi}$, and in particular, is finite. Then the TBFT_f follows from Lemma 3.8 of [9]. \square

REFERENCES

- [1] LAURENT BARTHOLDI, MARKUS NEUHAUSER, AND WOLFGANG WOESS. Horocyclic products of trees. *J. Eur. Math. Soc. (JEMS)* **10**, No. 3, 771–816, 2008.
- [2] M. J. CURRAN. Automorphisms of semidirect products. *Math. Proc. R. Ir. Acad.* **108**, No. 2, 205–210, 2008.
- [3] KAREL DEKIMPE AND DACIBERG GONÇALVES. The R_∞ property for abelian groups. *Topol. Methods Nonlinear Anal.* **46**, No. 2, 773–784, 2015.
- [4] A. FEL'SHTYN. Dynamical zeta functions, Nielsen theory and Reidemeister torsion. *Mem. Amer. Math. Soc.* **147**, No. 699, xii+146, 2000.
- [5] A. FEL'SHTYN AND R. HILL. The Reidemeister zeta function with applications to Nielsen theory and a connection with Reidemeister torsion. *K-Theory* **8**, No. 4, 367–393, 1994.
- [6] A. FEL'SHTYN, F. INDUKAEV, AND E. TROITSKY. Twisted Burnside theorem for two-step torsion-free nilpotent groups. In *C*-algebras and elliptic theory. II*, Trends in Math., pages 87–101. Birkhäuser, 2008.
- [7] A. FEL'SHTYN, YU. LEONOV, AND E. TROITSKY. Twisted conjugacy classes in saturated weakly branch groups. *Geometriae Dedicata* **134**, 61–73, 2008.
- [8] A. FEL'SHTYN AND E. TROITSKY. A twisted Burnside theorem for countable groups and Reidemeister numbers. In C. Consani and M. Marcolli, editors, *Noncommutative Geometry and Number Theory*, pages 141–154. Vieweg, Braunschweig, 2006.
- [9] A. FEL'SHTYN AND E. TROITSKY. Twisted Burnside-Frobenius theory for discrete groups. *J. Reine Angew. Math.* **613**, 193–210, 2007.
- [10] A. FEL'SHTYN, E. TROITSKY, AND A. VERSHIK. Twisted Burnside theorem for type II_1 groups: an example. *Math. Res. Lett.* **13**, No. 5, 719–728, 2006.
- [11] ALEXANDER FEL'SHTYN, NIKITA LUCHNIKOV, AND EVGENIJ TROITSKY. Twisted inner representations. *Russian Journal of Mathematical Physics* **22**, No. 3, 301–306, 2015.
- [12] ALEXANDER FEL'SHTYN AND TIMUR NASYBULOV. The R_∞ and S_∞ properties for linear algebraic groups. *J. Group Theory* **19**, No. 5, 901–921, 2016.
- [13] ALEXANDER FEL'SHTYN AND EVGENIJ TROITSKY. Geometry of Reidemeister classes and twisted Burnside theorem. *J. K-Theory* **2**, No. 3, 463–506, 2008.
- [14] ALEXANDER FEL'SHTYN AND EVGENIJ TROITSKY. Twisted conjugacy classes in residually finite groups. Arxiv e-print 1204.3175, 2012.
- [15] ALEXANDER FEL'SHTYN AND EVGENIJ TROITSKY. Aspects of the property R_∞ . *J. Group Theory* **18**, No. 6, 1021–1034, 2015.
- [16] D. GONÇALVES. The coincidence Reidemeister classes on nilmanifolds and nilpotent fibrations. *Topology and Its Appl.* **83**, 169–186, 1998.

- [17] D. GONÇALVES AND P. WONG. Twisted conjugacy classes in exponential growth groups. *Bull. London Math. Soc.* **35**, No. 2, 261–268, 2003.
- [18] DACIBERG GONÇALVES AND PETER WONG. Twisted conjugacy classes in wreath products. *Internat. J. Algebra Comput.* **16**, No. 5, 875–886, 2006.
- [19] DACIBERG GONÇALVES AND PETER WONG. Twisted conjugacy classes in nilpotent groups. *J. Reine Angew. Math.* **633**, 11–27, 2009.
- [20] B. JIANG. *Lectures on Nielsen Fixed Point Theory*, volume 14 of *Contemp. Math.* Amer. Math. Soc., Providence, RI, 1983.
- [21] MORRIS NEWMAN. *Integral matrices*. Academic Press, New York-London, 1972. Pure and Applied Mathematics, Vol. 45.
- [22] DEREK J. S. ROBINSON. *A course in the theory of groups*, volume 80 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1996.
- [23] DANIEL SEGAL. *Polycyclic groups*. Number 82 in Cambridge Tracts in Mathematics. Cambridge University Press, Cambridge, 1983.
- [24] MELANIE STEIN, JENNIFER TABACK, AND PETER WONG. Automorphisms of higher rank lamplighter groups. *Internat. J. Algebra Comput.* **25**, No. 8, 1275–1299, 2015.
- [25] JENNIFER TABACK AND PETER WONG. The geometry of twisted conjugacy classes in wreath products. In *Geometry, rigidity, and group actions*, Chicago Lectures in Math., pages 561–587. Univ. Chicago Press, Chicago, IL, 2011. (arXiv:0805.1372).
- [26] E. TROITSKY. Noncommutative Riesz theorem and weak Burnside type theorem on twisted conjugacy. *Funct. Anal. Pril.* **40**, No. 2, 44–54, 2006. In Russian, English translation: *Funct. Anal. Appl.* **40** (2006), No. 2, 117–125.
- [27] EVGENIJ TROITSKY. Two examples related to the twisted Burnside-Frobenius theory for infinitely generated groups. *Fundam. Appl. Math.* **21**, No. 5, 231–239, 2016. (in Russian, English translation in Journal of Mathematical Sciences (New York) to appear, preprint version <http://www.mpim-bonn.mpg.de/preblob/5689>).

DEPT. OF MECH. AND MATH., MOSCOW STATE UNIVERSITY, 119991 GSP-1 MOSCOW, RUSSIA
E-mail address: troitsky@mech.math.msu.su
URL: <http://mech.math.msu.su/~troitsky>