

# AN EXPLICIT CONDUCTOR FORMULA FOR $GL(n) \times GL(1)$

ANDREW CORBETT

ABSTRACT. We prove an explicit formula for the conductor of an irreducible, admissible representation of  $GL(n, F)$  twisted by a character of  $F^\times$  where the field  $F$  is local and non-archimedean. As a consequence, we quantify the number of character twists of such a representation of fixed conductor.

## 1. THE PROBLEM OF THE TWISTED CONDUCTOR

Let  $F$  denote a non-archimedean local field of characteristic zero and let  $n \geq 2$ . For an irreducible, admissible representation  $\pi$  of  $GL(n, F)$  and a quasi-character  $\chi$  of  $F^\times$ , we can form the twist  $\chi\pi = (\chi \circ \det) \otimes \pi$ . Our main result, Theorem 2.7, is an explicit formula for the conductor  $a(\chi\pi)$ , c.f. the Artin conductor, as defined in §3.1. This formula is given by

$$a(\chi\pi) = a(\pi) + \Delta_\chi(\pi) - \delta_\chi(\pi) \quad (1)$$

where  $\Delta_\chi(\pi)$  and  $\delta_\chi(\pi)$  are non-negative integers as defined in Theorem 2.7; they denote a *dominant* and a *non-twist-minimal interference* term, respectively. We give detailed analysis of these terms in §4.2, answering questions such as “for what number of  $\chi$  is there interference present?”

As an example, computing  $a(\chi\pi)$  in the limit  $a(\chi) \rightarrow \infty$  is straightforward: from Proposition 2.2 and Equation (5) we deduce

$$a(\chi\pi) = na(\chi) \quad (2)$$

whenever  $a(\chi) > a(\pi)$ . In this case  $\Delta_\chi(\pi) = na(\chi) - a(\pi)$  and  $\delta_\chi(\pi) = 0$ . Bushnell–Henniart extend (2) by proving the upper bound<sup>1</sup>

$$a(\chi\pi) \leq \max\{a(\pi), a(\chi)\} + (n-1)a(\chi), \quad (3)$$

surrendering to a weaker bound in the region  $0 \leq a(\chi) \leq a(\pi)$ . Nevertheless, this bound is sharp in that it is attained for *some*  $\pi$  and  $\chi$ ; as in (2) for example.

However, in general such examples become sparse, rendering (3) as rather coarse as one averages over  $\chi$  with  $a(\chi) \asymp a(\pi)$ . In such cases, evaluating the integers  $\Delta_\chi(\pi)$  and  $\delta_\chi(\pi)$  exactly is of crucial importance for numerous problems in analytic number theory.

In this paper we consider applications to studying  $a(\chi\pi)$  in a quantitative fashion; for example, counting the number of  $\chi$  for which  $a(\chi\pi)$  is equal to a given integer. See §4. This analysis is applicable in practice should one need to understand  $a(\chi\pi)$  on average.

---

*Date:* 15<sup>th</sup> January 2018.

<sup>1</sup>Inequality (3) is a special case of both [2, Theorem 1] and indeed our own Theorem 2.7. See Corollary 2.10 for a more precise inequality.

Indeed, our formula may be utilised when studying the analytic behaviour of  $L$ -functions attached to automorphic forms. In particular it applies in conjunction with the following two techniques: taking harmonic  $GL(1)$ -averages and applying the functional equation for  $GL(n) \times GL(1)$ - $L$ -functions. For example, conductors of such character twists arise in the work of Nelson–Pitale–Saha [13] who address the quantum unique ergodicity conjecture for holomorphic cusp forms with “powerful” level (see [13, Remarks 1.9 & 3.16]). The current record for upper and lower bounds for the sup-norm of a Maaß-newform on  $GL(2)$  in the level aspect [17–19] also depends crucially on the  $n = 2$  case of Theorem 2.7.

An instance where (1) is applied constructively is carried out in [4], once again when  $n = 2$ . Originally, in [1], Brunault computed the value of ramification indices of modular parameterisation maps of various elliptic curves over  $\mathbb{Q}$ . Whenever the newform attached to  $E$  is “twist minimal”, Brunault could prove that this index was trivial (equal to 1), holding in particular whenever the conductor of  $E$  is square-free. This problem has now been completely solved by Saha and the present author [4]. In our solution, it is the degenerate cases of (1), with non-trivial  $\Delta_\chi(\pi)$  and  $\delta_\chi(\pi)$ , that give rise to the few examples of non-trivial ramification indices.

These results all concern the case  $n = 2$ , where the conductor formula for twists of supercuspidal representations was given by Tunnell [22, Proposition 3.4] in his thesis (see [4, Lemma 2.7] for the general case). Tunnell himself applied his formula to count isomorphism classes of supercuspidal representations of fixed odd conductor [22, Theorem 3.9]. He used this observation in his proof of the local Langlands correspondence for  $GL(2, F)$  in the majority of cases.

Our present result is suggestive of similar applications: a bound for local Whittaker newforms (and a corresponding global sup-norm bound) in the level aspect; bounds for matrix coefficients of local representations, and estimates relating to the Voronoï summation problem for  $GL(n)$ , to name a few.

In §2 we describe how irreducible, admissible representations of  $GL(n, F)$  are classified and then go on to give a full account of our main result. This classification assumes the least amount of necessary information in order to give a completely explicit formula. In §3 we give a uniform proof of our main result for the quasi-square-integrable representations (see Proposition 2.2); these representations are used as building blocks to arrive at the general case. Lastly, in §4, we provide a detailed analysis of the terms  $\Delta_\chi(\pi)$  and  $\delta_\chi(\pi)$  as found in (1).

## 2. AN EXPLICIT FORMULA FOR TWISTED CONDUCTORS

Here we give full details of the formula proposed in (1). We first describe the formula for quasi-square-integrable representations of  $GL(n, F)$ , which is then used to build the result in its full generality.

**2.1. The Langlands classification for  $GL(n, F)$ .** Let  $\mathcal{A}_F(n)$  denote the set of (equivalence classes of) the irreducible, admissible representations of  $GL(n, F)$ . The natural building blocks that describe  $\mathcal{A}_F(n)$  are the *quasi-square-integrable*

representations; these are those  $\pi \in \mathcal{A}_F(n)$  for which there is an  $\alpha \in \mathbb{R}$  such that  $|\cdot|^\alpha \pi$  has square-integrable matrix coefficients on  $\mathrm{GL}(n, F)$  modulo its centre.

The ‘Langlands classification’ (due to Bernstein–Zelevinsky in this case) describes the structure of each representation in the graded ring  $\mathcal{A}_F = \bigoplus_{n \geq 1} \mathcal{A}_F(n)$  in terms of the subset  $\mathcal{S}\mathcal{G}_F$  of quasi-square-integrable representations. By [25, Theorems 9.3 & 9.7], one deduces an addition law  $\boxplus$  on  $\mathcal{S}\mathcal{G}_F$ , by which  $\mathcal{S}\mathcal{G}_F$  generates a free commutative monoid  $\Lambda$ . The classification is then the assertion that there is a bijection between  $\mathcal{A}_F$  and the semi-group of non-identity elements in  $\Lambda$ , thus endowing  $\mathcal{A}_F$  with the addition law  $\boxplus$ . Crucially, the maps  $(\mathcal{A}_F, \boxplus) \rightarrow (\mathbb{C}, \cdot)$ , given by applying  $L$ - or  $\varepsilon$ -factors, are homomorphisms of semi-groups (see [23, §2.5] for their definitions). Both expositions [14, 23] provide excellent background on this topic.

The upshot of this classification is that for any  $\pi \in \mathcal{A}_F(n)$  there exists a unique partition  $n_1 + \cdots + n_r = n$  alongside a collection of quasi-square-integrable representations  $\pi_i \in \mathcal{S}\mathcal{G}_F \cap \mathcal{A}_F(n_i)$  for  $1 \leq i \leq r$  such that

$$\pi = \pi_1 \boxplus \cdots \boxplus \pi_r \quad (4)$$

and for any quasi-character  $\chi$  of  $F^\times$  we have

$$a(\chi\pi) = a(\chi\pi_1) + \cdots + a(\chi\pi_r). \quad (5)$$

Equation (5) follows from the definition of the  $\varepsilon$ -factor and formula (11) in §3.1.

## 2.2. The formula for quasi-square-integrable representations.

**Definition 2.1.** An irreducible, admissible representation  $\pi$  of  $\mathrm{GL}(n, F)$  is called *twist minimal* if  $a(\pi)$  is the smallest of the integers  $a(\chi\pi)$  as  $\chi$  varies over the quasi-characters of  $F^\times$ .

For quasi-square-integrable representations, the notion of twist-minimality is sufficient to give an exact formula for the conductor of their twist.

**Proposition 2.2.** *Let  $\pi$  be an irreducible, admissible, quasi-square-integrable representation of  $\mathrm{GL}(n, F)$  and let  $\chi$  be a quasi-character of  $F^\times$ . Then*

$$a(\chi\pi) \leq \max\{a(\pi), na(\chi)\} \quad (6)$$

with equality in (6) whenever  $\pi$  is twist minimal or  $a(\pi) \neq na(\chi)$ .

*Remark 2.3.*

*Remark 2.4.* In practice, one handles those  $\pi \in \mathcal{S}\mathcal{G}_F \cap \mathcal{A}_F(n)$  which are *not* twist minimal as follows. Tautologically, write  $\pi = \mu\pi^{\min}$  where  $\mu$  is a quasi-character of  $F^\times$  and  $\pi^{\min}$  is twist minimal. Then Proposition 2.2 implies that  $a(\chi\pi) = \max\{a(\pi^{\min}), na(\chi\mu)\}$ . In particular,  $n \mid a(\pi)$  if  $\pi$  is not twist minimal.

Let us briefly mention the conductor formula of Bushnell–Henniart–Kutzko [3, Theorem 6.5] for  $\mathrm{GL}(n) \times \mathrm{GL}(m)$ -pairs of *supercuspidal* representations. There they deploy the full structure theory of supercuspidal representations to prove a detailed identity relating the conductor to the respective inducing data of the given supercuspidal representations. However, this formula is difficult to apply in practice. Indeed, our own Proposition 2.2 may be derived from their work.

Comparing the  $m = 1$  case of [3] to our present result, our formula is simpler and holds uniformly on the larger set  $\mathcal{S}\mathcal{G}_F$  which contain not only the supercuspidals but also, for example, the special representations (c.f. the known formula for the Steinberg representation [16, p. 18]). Accordingly, we give an elementary proof of Proposition 2.2. This promotes our observation that, as far as the conductor is concerned, the set  $\mathcal{S}\mathcal{G}_F$ , or rather its subset of twist minimal elements, contains sufficient and necessary information to explicitly determine the conductor of any twist.

We defer our proof of Proposition 2.2 until §3.4. The arguments made there are also used to prove the following result on the central character:

**Proposition 2.5.** *Let  $\pi$  be an irreducible, admissible, quasi-square-integrable representation of  $GL(n, F)$  with central character  $\pi|_{F^\times} = \omega_\pi$ . Then*

$$a(\omega_\pi) \leq \frac{a(\pi)}{n}. \quad (7)$$

*Remark 2.6.* The central character of a quasi-square-integrable representation has relatively small conductor. In general, highly ramified central characters arise due to the components in a given  $\pi_1 \boxplus \cdots \boxplus \pi_r$  for  $r \geq 2$ . For this reason, such representations should be handled separately, as is distinguished in this work.

**2.3. The general formula.** We arrive at our main result, having defined the necessary and sufficient set of properties for the representations  $\pi \in \mathcal{A}_F$  in order to give a complete and explicit formula for the conductor of its twists.

**Theorem 2.7.** *Let  $\pi$  be an irreducible, admissible representation of  $GL(n, F)$  given in terms of quasi-square-integrable representations  $\pi_i$  of  $GL(n_i, F)$ , as described in (4), where  $n = n_1 + \cdots + n_r$  and  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$ . Let  $\chi$  be a quasi-character of  $F^\times$ . Then*

$$a(\chi\pi) = a(\pi) + \Delta_\chi(\pi) - \delta_\chi(\pi)$$

where  $\Delta_\chi$  and  $\delta_\chi$  are semi-group homomorphisms  $(\mathcal{A}_F, \boxplus) \rightarrow (\mathbb{Z}_{\geq 0}, +)$  defined by their values on the representations  $\pi_i \in \mathcal{S}\mathcal{G}_F$  as follows:

$$\Delta_\chi(\pi_i) = \begin{cases} \max\{n_i a(\chi) - a(\pi_i), 0\} & \text{if } a(\chi) \neq a(\mu_i) \\ 0 & \text{if } a(\chi) = a(\mu_i) \end{cases}$$

and

$$\delta_\chi(\pi_i) = \begin{cases} a(\pi_i) - \max\{a(\pi_i^{\min}), n_i a(\chi\mu_i)\} & \text{if } a(\chi) = a(\mu_i) \\ 0 & \text{if } a(\chi) \neq a(\mu_i) \end{cases}$$

where  $\pi_i^{\min}$  is twist minimal and  $\mu_i$  a quasi-character of  $F^\times$  such that  $\pi_i = \mu_i \pi_i^{\min}$ .

*Remark 2.8.* Both terms  $\Delta_\chi(\pi)$  and  $\delta_\chi(\pi)$  are non-negative for any  $\pi$  and  $\chi$ .

*Proof.* By applying Proposition 2.2 to Equation (5) we have

$$a(\chi\pi) = \sum_{i=1}^r \max\{a(\pi_i^{\min}), n_i a(\chi\mu_i)\}. \quad (8)$$

In particular, if  $a(\chi) \neq a(\mu_i)$  for a given  $1 \leq i \leq r$  then the respective summand in (8) is equal to  $\max\{a(\pi_i^{\min}), n_i a(\chi \mu_i)\} = \max\{a(\pi_i), n_i a(\chi)\}$ . Whenever we have  $a(\chi) \neq a(\mu_i)$  we also have  $a(\pi_i) \geq \max\{a(\pi_i^{\min}), n_i a(\chi \mu_i)\}$ .  $\square$

*Remark 2.9.* In the special case  $n = 2$ , we prove Theorem 2.7 in [4, Lemma 2.7]. In general, one should understand the non-vanishing of  $\delta_\chi(\pi)$  as occurring rarely, whereas  $\Delta_\chi(\pi)$  describes the dominant or ‘‘usual’’ behaviour of  $a(\chi\pi)$ . We make these statements explicit in a quantitative sense in §4.2.

**Corollary 2.10.** *Let  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$  and  $\chi$  be as in Theorem 2.7 with  $\pi_i = \mu_i \pi_i^{\min}$  for twist minimal representations  $\pi_i^{\min}$ . Define the ‘totally minimal’ representation  $\pi^{\min} = \pi_1^{\min} \boxplus \cdots \boxplus \pi_r^{\min}$  and let  $\Omega_\chi(\pi) = \{1 \leq i \leq r : a(\pi_i) > n_i a(\chi)\}$ . Then*

$$a(\pi^{\min}) \leq a(\chi\pi) \leq a(\pi) + a(\chi) \left( n - \sum_{i \in \Omega_\chi(\pi)} n_i \right). \quad (9)$$

*Proof.* The lower bound of (9) follows immediately from (8). Note that for  $i \in \Omega_\chi(\pi)$  we have  $\Delta_\chi(\pi_i) = \delta_\chi(\pi_i) = 0$ . The upper bound is then obtained by bounding  $\Delta_\chi(\pi_i)$  and  $\delta_\chi(\pi_i)$  respectively by  $a(\pi_i) + n_i a(\chi)$  for  $i \notin \Omega_\chi(\pi)$ .  $\square$

*Proof of Inequality (3).* We recover Bushnell–Henniart’s bound (3) using Corollary 2.10. If  $a(\chi) > a(\pi)$  then  $a(\pi) = na(\chi)$ , by (8). Whilst if  $a(\chi) \leq a(\pi)$  then (3) is a special case of (9) since we have  $\Omega_\chi(\pi) = \emptyset$  and each  $n_i \geq 1$ .  $\square$

### 3. CONDUCTORS OF TWISTS VIA DIVISION ALGEBRAS

In this section we provide proofs for Propositions 2.2 and 2.5. These results apply to all quasi-square-integrable representations uniformly, as is reflected in our proof. In particular, our conductor formula bypasses many of the complications occurring in the formula for supercuspidal representations given in [3].

**3.1. Notation and definition of the conductor.** Let  $\pi$  denote an irreducible, admissible representation of  $GL(n, F)$  and  $\chi$  a quasi-character of  $F^\times$ . Let  $\tilde{\pi}$  be the contragredient representation and  $\omega_\pi$  the central character of  $\pi$  respectively.

**3.1.1. The non-archimedean local field.** We denote by  $\mathfrak{o}$  the ring of integers of  $F$ ;  $\mathfrak{p}$  the maximal ideal of  $\mathfrak{o}$ ;  $\varpi$  a choice of uniformising parameter, that is a generator of  $\mathfrak{p}$ , and  $q = \#(\mathfrak{o}/\mathfrak{p})$ . Let  $|\cdot|$  denote the absolute value on  $F$ , normalised so that  $|\varpi| = q^{-1}$  and  $v_F$  the valuation on  $F$  defined via  $|x| = q^{-v_F(x)}$  for  $x \in F$ . We define open neighbourhoods  $U_F(m)$  of 1 in  $U_F(0) = \mathfrak{o}^\times$  by  $U_F(m) = 1 + \varpi^m \mathfrak{o}$  for  $m > 0$ . Define the integer  $a(\chi)$  to be the least  $m \geq 0$  such that  $\chi(U_F(m)) = \{1\}$ . Let  $K = GL(n, \mathfrak{o})$  and for each  $m \geq 0$  let  $K_1(m)$  be the subgroup of  $K$  that stabilises the row vector  $(0, \dots, 0, 1)$ , from the right, modulo  $\mathfrak{p}^m$ .

**3.1.2. The floor and ceiling functions.** For each  $\alpha \in \mathbb{R}$  let  $\lfloor \alpha \rfloor$  denote the *floor* of  $\alpha$ , defined via  $\lfloor \alpha \rfloor = m$  if and only if  $m \in \mathbb{Z}$  and  $m \leq \alpha < m + 1$ . Similarly, let  $\lceil \alpha \rceil$  denote the *ceiling* of  $\alpha$  if and only if  $\lceil \alpha \rceil \in \mathbb{Z}$  such that  $\lceil \alpha \rceil - 1 < \alpha \leq \lceil \alpha \rceil$ .

3.1.3. *Epsilon constants and the conductor.* Here we define the integer  $a(\pi)$ , the conductor of  $\pi$ . This subsumes the definition of  $a(\chi)$  when  $n = 1$ , for which the following notions were originally founded in Tate’s thesis [20].

Let  $\psi$  be an additive character of  $F$  of exponent  $n(\psi) = \min\{m : \psi|_{\mathfrak{p}^m} = 1\}$ . Godement–Jacquet prove the existence of  $\varepsilon$ -factors  $\varepsilon(s, \pi, \psi) \in \mathbb{C}[q^{-s}, q^s]$  in [7, Theorem 3.3, (4)]. Applying the functional equation twice, one obtains

$$\varepsilon(s, \pi, \psi)\varepsilon(1-s, \tilde{\pi}, \psi) = \omega_\pi(-1). \quad (10)$$

Hence  $\varepsilon(s, \pi, \psi)$  is a unit in  $\mathbb{C}[q^{-s}, q^s]$ ; that is, a  $\mathbb{C}^\times$ -constant multiple of an integral power of  $q^{-s}$ . Explicitly, using [7, (3.3.5)] we prove

$$\varepsilon(s, \pi, \psi) = \varepsilon(1/2, \pi, \psi) q^{(a(\pi) - n(\psi)n)(\frac{1}{2} - s)}, \quad (11)$$

in which we *implicitly define the conductor*  $a(\pi)$ . After the proof of the local Langlands corresponds for  $\mathrm{GL}(n, F)$  [8], the conductor  $a(\pi)$  coincides with the Artin conductor of an  $n$ -dimensional Weil–Deligne representation. A fundamental property of  $\varepsilon$ -factors is that  $\varepsilon(s, \chi\pi, \psi) = \prod_{i=1}^r \varepsilon(s, \chi\pi_i, \psi)$  for  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$ , as in (4) (see [7, Theorem 3.4]). This observation proves (5) by applying (11). Moreover, if  $\pi$  is generic, the conductor  $a(\pi)$  may be interpreted in terms of newform theory as we now explain.

3.1.4. *Conductors of generic representations and newform theory.* Legibility permitting, the letter  $\mathcal{G}$  indicates that all representations in  $\mathcal{S}\mathcal{G}_F$  are *generic*. By showing so for the regular representation of  $\mathrm{GL}(n, F)$ , of a given central character, Jacquet shows that all discrete series representations are generic [9, Theorem 2.1, (3)]. By the Langlands classification (denoting as in (4)), any  $\pi \in \mathcal{A}_F(n)$  as is generic (a.k.a. “non-degenerate”) if and only if  $\pi$  is equivalent to the (irreducible) representation induced from the *external* tensor product  $\pi_1 \boxtimes \cdots \boxtimes \pi_r$  on the parabolic subgroup associated to  $n_1 + \cdots + n_r$  (by [25, Theorem 9.7, (a)]). The elements of  $\mathcal{S}\mathcal{G}_F$  correspond to those irreducible representations with  $r = 1$ .

Assume that  $\pi$  is generic. Then the conductor  $a(\pi)$  may be equivalently constructed in a language more familiar to the theory of automorphic forms: let us re-define the conductor  $a(\pi)$  of  $\pi$  to be the least non-negative integer  $m$  such that  $\pi$  contains a non-zero  $K_1(m)$ -fixed vector.

The fundamental theorem of newform theory is that the space of  $K_1(a(\pi))$ -fixed vectors is one-dimensional. This theorem is due to Gelfand–Každan [6] in the present context. The coincidence of the definitions for  $a(\pi)$  given in §3.1.3 and §3.1.4 is proved by Jacquet–Piatetski-Shapiro–Shalika [11, Théorème (5)].

3.2. **Central simple division algebras.** Let  $D$  be a division algebra over  $F$  of dimension  $[D : F] = n^2$ . Let  $\mathrm{Nrd} = \mathrm{Nrd}_D$  denote the reduced norm on  $D$ . (See [12, §4.1] for a pleasant construction.) Any valuation on  $D$  may be obtained via composing the reduced norm with a valuation on  $F$  (see [21, Theorem 1.4]); let us normalise such a choice by  $v_D = v_F \circ \mathrm{Nrd}$ .



**3.2.1. Unit groups.** Define a basis of neighbourhoods of  $1 \in D^\times$  by  $U_D(m) = \{x \in D^\times : v_D(x - 1) \geq m\}$  for  $m > 0$  and let  $U_D(0) = \ker(v_D)$ . Note that if  $n = 1$  (so that  $D = F$ ) we recover  $U_D(m) = U_F(m)$ . It is an important fact that the norm map  $\text{Nrd}: D^\times \rightarrow F^\times$  is surjective (see [24, Prop. 6, Ch. X-2, p. 195] for instance). Upon restriction to the above neighbourhoods, for each  $m \geq 0$  we have  $\text{Nrd}(U_D(m)) = U_D(m) \cap F$ .

**Lemma 3.1.** *For each  $m \geq 0$  we have  $U_D(m) \cap F^\times = U_F(\lceil m/n \rceil)$ .*

*Proof.* For all  $a \in F^\times$  we have  $v_D(a) = v_F(\text{Nrd}(a)) = v_F(a^n) = nv_F(a)$ ; the definition of  $U_F(\lceil m/n \rceil)$  is then equivalent to that of the intersection.  $\square$

**Lemma 3.2.** *For each  $m \geq 0$  we have  $\text{Nrd}(U_D(m)) = U_F(\lceil m/n \rceil)$ .*

*Proof.* This follows by applying Lemma 3.1 to  $\text{Nrd}(U_D(m)) = U_D(m) \cap F$ .  $\square$

**3.2.2. The level of a representation of  $D^\times$ .** If  $\chi$  is a quasi-character of  $F^\times$  and  $\pi'$  an irreducible, admissible representation of  $D^\times$ , analogous to the unramified case we form the twist  $\chi\pi' = (\chi \circ \text{Nrd}) \otimes \pi'$ . Define the *level*  $l(\pi')$  of  $\pi'$  to be the least non-negative integer  $m$  such that  $\pi'(U_D(m)) = \{1\}$ .

**Lemma 3.3.** *Let  $\pi'$  be an irreducible, admissible representation of  $D^\times$ . The conductor  $a(\pi')$  as defined via the  $\varepsilon$ -factor (mutatis mutandis as in §3.1.3) is related to the level  $l(\pi')$  by the formula*

$$a(\pi') = l(\pi') + n - 1.$$

*Proof.* This is proved in [12, §4.3], stated explicitly in [12, (4.3.4)]. To assist (mathematical) translation, we remark on the following: their unit groups  $V_j$  equal our  $U_D(j)$  for  $j \geq 0$ . Fix their character  $\chi \in \text{Hom}(V_j/V_{j+1}, \mathbb{C}^\times)$  by defining  $\chi = \pi'|_{V_j}$  and choosing  $j = l(\pi') - 1$ . Then their  $c \in D$ , “der Kontrolleur von  $\chi$ ”, satisfies  $v_D(c) = -a(\chi) = -a(\pi')$ ; it is constructed in [12, (4.3.1)] from where we have  $v_D(c) = -n - j$ , noting the non-triviality of  $\chi$  on  $V_j$ . All together this implies  $a(\pi') = n + j = n + l(\pi') - 1$ .  $\square$

**Lemma 3.4.** *Let  $\chi$  be a quasi-character of  $F^\times$ . Then*

$$l(\chi \circ \text{Nrd}) = na(\chi) - n + 1.$$

*Proof.* By Lemma 3.2 we consider  $\chi$  restricted to  $U_F(\lceil m/n \rceil)$  for each  $m \geq 0$ . By the minimality of the  $a(\chi)$ , the character  $\chi \circ \text{Nrd}$  is trivial on  $U_D(m)$  whenever

$$n(a(\chi) - 1) \leq m - 1. \tag{12}$$

By the minimality of the level, we have equality in (12) when  $m = l(\chi \circ \text{Nrd})$ .  $\square$

**3.3. The Jacquet–Langlands correspondence for division algebras.** This special case of functoriality stipulates a bijection between the following:

- The set of equivalence classes of irreducible, admissible representations of  $GL(n, F)$ , with unitary central character, which are *square-integrable modulo centre*. These are precisely the square-integrable elements of  $\mathcal{S}\mathcal{G}_F \cap \mathcal{A}_F(n)$ .

- The set of equivalence classes of irreducible, admissible representations of  $D^\times$  with unitary central character where  $D$  is a central-simple  $F$ -algebra of dimension  $n^2$ .

*Remark 3.5.* Discrete series representations of the above two types have unitary central characters. In the above bijection, if  $\pi$  corresponds to  $\pi'$  then their central characters agree:  $\omega_\pi = \omega_{\pi'}$ . Moreover,  $\chi\pi$  corresponds to  $\chi\pi'$  for any quasi-character  $\chi$ . As a consequence of the Peter–Weyl theorem, the representations of  $D^\times$  are finite dimensional (since  $D^\times$  is compact modulo centre).

The correspondence as stated here is due to Rogawski [15, Theorem 5.8], where the original case  $n = 2$  was famously proved by Jacquet–Langlands [10]. The most general statement allows one to replace  $D^\times$  with  $GL(m, D)$  where  $D$  has dimension  $d^2$  and  $m$  must satisfy  $n = md$ . This is established in [5] by Deligne–Kazhdan–Vignéras.

**3.4. The main proofs.** Here we provide a stand alone proof of our main result in the quasi-square-integrable case. Assume the hypotheses and notations of Propositions 2.2 and 2.5; in particular,  $\pi \in \mathcal{S}\mathcal{G}_F$ .

3.4.1. *Proof of Proposition 2.2.* The following lemma reduces the proof to the case where  $\pi$  is just square-integrable.

**Lemma 3.6.** *For all quasi-characters  $\chi$  with  $a(\chi) = 0$  we have  $a(\chi\pi) = a(\pi)$ .*

*Proof.* Let  $m \geq 0$ . As each  $\pi \in \mathcal{S}\mathcal{G}_F$  is generic, the space  $\pi^{K_1(m)}$  of  $K_1(m)$ -fixed vectors in  $\pi$  is non-zero if and only if  $(\chi\pi)^{K_1(m)} \neq \{0\}$ .  $\square$

Henceforth we assume  $\pi$  to be square-integrable. The generalised Jacquet–Langlands correspondence implies  $a(\chi\pi) = a(\chi\pi')$  where  $\pi'$  is the irreducible, admissible, unitary representation of  $D^\times$  associated to  $\pi$  as determined by [15, Theorem 5.8]. The proof of Proposition 2.2 now follows by applying Lemmas 3.3 and 3.4 to the following.

**Lemma 3.7.** *Let  $\pi'$  be an irreducible, admissible, unitary representation of  $D^\times$  and  $\chi$  a quasi-character of  $F^\times$ . Then*

$$l(\chi\pi') \leq \max\{l(\pi'), l(\chi \circ \text{Nrd})\} \quad (13)$$

with equality in (13) whenever  $\pi'$  is twist minimal or  $l(\pi') \neq l(\chi \circ \text{Nrd})$ .

*Proof.* By definition,  $(\chi\pi')(x) = \chi(\text{Nrd}(x))\pi'(x)$  for every  $x \in U_D(m)$  with  $m \geq 0$ . One immediately obtains (13) by minimality. Equality also follows in the given cases, noting that twist minimality in  $a(\pi')$  is equivalent to twist minimality in  $l(\pi')$  since they are linearly related (by Lemma 3.3).  $\square$

3.4.2. *Proof of Proposition 2.5.* Recall that  $\pi'|_{F^\times} = \omega_\pi$ . Taking  $m = l(\pi')$  in Lemma 3.1 we deduce that

$$a(\omega_\pi) \leq \left\lceil \frac{l(\pi')}{n} \right\rceil < \frac{a(\pi) - n + 1}{n} + 1 = \frac{a(\pi) + 1}{n}.$$

Thus we infer that  $na(\omega_\pi) \leq a(\pi)$  as required.



## 4. CHARACTERS PRESERVING THE CONDUCTOR UNDER TWISTING

The goal of this section is twofold: in §4.1 we count the number of characters  $\chi$  such that  $a(\chi\pi)$  is equal to a given integer. Then, in §4.2, we explicitly analyse the behaviour of the dominant and interference terms of Theorem 2.7. These questions are motivated by their applications to analytic number theory.

## 4.1. Sets of twist-fixing characters.

4.1.1. *Characters of a given conductor.* The valuation  $v_F$  defines a split exact sequence  $1 \rightarrow \mathfrak{o}^\times \rightarrow F^\times \xrightarrow{v_F} \mathbb{Z} \rightarrow 1$ . We thus write any quasi-character  $\chi$  on  $F^\times$  as  $\chi(x) = \chi'(x)q^{-v_F(x)m}$  for some  $m \in \mathbb{Z}$  and a character  $\chi'$  of  $F^\times$  such that  $\chi'(\varpi) = 1$ , denoting the space of such  $\chi'$  by  $\mathfrak{X}$  so that the unitary dual of  $\mathfrak{o}^\times$  satisfies  $\hat{\mathfrak{o}}^\times \cong \mathfrak{X}$ . With interest in characters that fix the conductor under twisting, we define the following  $\mathfrak{X}$ -subsets:

$$\mathfrak{X}(k) = \{\chi \in \mathfrak{X} : a(\chi) \leq k\}; \quad \mathfrak{X}'(k) = \{\chi \in \mathfrak{X} : a(\chi) = k\}; \quad (14)$$

and

$$\mathfrak{X}'_\pi(k, j) = \{\chi \in \mathfrak{X} : a(\chi) = k \text{ and } a(\chi\pi) = j\} \quad (15)$$

for some  $k, j \geq 0$ .

Our present point of departure shall be to give sharp upper bounds for the set  $\mathfrak{X}'_\pi(k, j)$  via Corollary 2.10. In contrast, the cardinalities of  $\mathfrak{X}(k)$  and  $\mathfrak{X}'(k)$  are straightforward to compute.

**Lemma 4.1.** *For each  $k \geq 1$ ,  $\#\mathfrak{X}(k) = q^{k-1}(q-1)$ ,  $\#\mathfrak{X}'(1) = q-2$ , and for  $k \geq 2$ ,  $\#\mathfrak{X}'(k) = q^{k-2}(q-1)^2$ .*

*Proof.* Consider the subgroup series  $\{1\} = \mathfrak{X}(0) \leq \mathfrak{X}(1) \leq \dots \leq \mathfrak{X}(k) \leq \mathfrak{X}$ . For  $k \geq l \geq k/2 \geq 1$ , we have  $\mathfrak{X}(k)/\mathfrak{X}(l) \cong U_F(l)/U_F(k) \cong \mathfrak{o}/\mathfrak{p}^{k-l}$ . In particular, taking  $l = k-1$  and noting  $\mathfrak{X}(1) \cong (\mathfrak{o}/\mathfrak{p})^\times$ , one counts the given cardinalities inductively. The number  $\#\mathfrak{X}'$  is obtained by subtraction.  $\square$

We remark that in [4, Lemmas 2.1 & 2.2] we counted the elements  $\chi \in \mathfrak{X}'(k)$  for which  $a(\chi\mu)$  remains fixed for a given  $\mu \in \mathfrak{X}'(k)$ , characterising the existence of such elements as  $q$  becomes small. In the present work we consider a “non-Abelian” variant of this result by characterising the set  $\mathfrak{X}'_\pi(k, j)$ .

4.1.2. *The quasi-square-integrable case.* First consider those  $\pi \in \mathcal{S}\mathcal{G}_F \cap \mathcal{A}_F(n)$  so that Proposition 2.2 applies. For integers  $k, j \geq 0$ , if either  $\pi$  is twist minimal or  $k \neq a(\pi)/n$  then

$$\mathfrak{X}'_\pi(k, j) = \begin{cases} \mathfrak{X}'_\pi(k) & \text{if } j = \max\{a(\pi), nk\} \\ \emptyset & \text{if } j \neq \max\{a(\pi), nk\}. \end{cases} \quad (16)$$

These cases correspond to  $\ell = 0$  as given in the following lemma.

**Lemma 4.2.** *For each  $\pi \in \mathcal{S}\mathcal{G}_F \cap \mathcal{A}_F(n)$  write  $\pi = \mu\pi^{\min}$  for a twist minimal representation  $\pi^{\min}$ . For integers  $j, k \geq 0$  we have  $\mathfrak{X}'_\pi(k, j) = \emptyset$  unless*

$a(\pi^{\min}) \leq j \leq \max\{a(\pi), nk\}$ , in which case there exists an  $\ell \geq 0$  such that  $j = \max\{a(\pi), nk\} - \ell$  and

$$\#\mathfrak{X}'_{\pi}(k, j) \leq \#\mathfrak{X}(k - \lfloor \frac{\ell}{n} \rfloor). \quad (17)$$

*Proof.* If either  $\pi$  is minimal or  $k \neq a(\pi)/n$  then the lemma follows by (16). Hence assume  $a(\pi) = kn$  and  $\pi = \mu\pi^{\min}$  where  $\pi^{\min}$  is twist minimal with  $a(\pi^{\min}) < a(\pi)$  and  $\mu \in \mathfrak{X}'(k)$ . Then  $\mathfrak{X}'_{\pi}(k, j) = \emptyset$  unless  $a(\pi^{\min}) \leq j \leq nk$ . In this case, if there exists a  $\chi \in \mathfrak{X}'(k)$  such that  $\max\{a(\pi^{\min}), na(\chi\mu)\} = j$  then there are  $\#\mathfrak{X}(\lfloor j/n \rfloor)$  of them as we must have  $\chi \in \mu^{-1}\mathfrak{X}(\lfloor j/n \rfloor)$ .  $\square$

4.1.3. *The general case.* Lemma 4.2 may be assembled to describe all of  $\mathcal{A}_F(n)$ .

**Proposition 4.3.** *Let  $\pi$  be an irreducible, admissible representation of  $\mathrm{GL}(n, F)$  and  $\pi^{\mathrm{tot}}$  a totally minimal representation attached to  $\pi$  as in Corollary 2.10. For integers  $j, k \geq 0$  we have  $\mathfrak{X}'_{\pi}(k, j) = \emptyset$  unless  $j$  satisfies inequality (9), in which case*

$$\#\mathfrak{X}'_{\pi}(k, j) \leq \#\mathfrak{X}(\lfloor \frac{j}{n} \rfloor). \quad (18)$$

*Proof.* Write  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$  as in (4). Set  $j' = l - \sum_{i \notin \Phi} a(\chi\pi_i)$  where  $\Phi = \{1 \leq i \leq r : a(\pi_i) = n_i a(\chi) \text{ and } \pi_i = \mu_i \pi_i^{\min} \text{ for } a(\pi_i^{\min}) < a(\pi_i)\}$ . Then (16) applies when  $i \notin \Phi$ . We are left answering the question of how many  $\chi \in \mathfrak{X}'(k)$  satisfy

$$j' = \sum_{i \in \Phi} \max\{a(\pi_i^{\min}), n_i a(\chi\mu_i)\}.$$

For each summand, this is answered by Lemma 4.2. We conclude that  $j'$  satisfies  $\sum_{i \in \Phi} a(\pi_i^{\min}) \leq j' \leq \sum_{i \in \Phi} \max\{a(\pi_i), nk\}$ , as per (9), and for some  $i \in \Phi$  we have

$$\#\mathfrak{X}'_{\pi}(k, j) \leq \#\mathfrak{X}\left(\max\left\{\left\lfloor \frac{a(\pi_i^{\min})}{n_i} \right\rfloor, a(\chi\mu_i)\right\}\right). \quad (19)$$

$\square$

*Remark 4.4.* The upper bound  $q^{j/n}$  for (18) typically occurs elsewhere, working against upper bounds for a typical summation problem averaging over  $a(\chi\pi)$ . Proposition 4.3 acts to counteract such trivially bounded terms. The bound (18) may be improved to (19) with the general notation of  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$ .

**4.2. The leading and interference terms.** Here we detail the asymptotic behaviour of  $\Delta_{\chi}(\pi)$  and  $\delta_{\chi}(\pi)$ . Our first port of call is to describe the rarity with which the interference term satisfies  $\delta_{\chi}(\pi) \neq 0$ . The following lemma follows directly from the definition of  $\delta_{\chi}(\pi)$  in Theorem 2.7.

**Lemma 4.5** (Absence of interference). *Let  $\pi$  be an irreducible, admissible representation of  $\mathrm{GL}(n, F)$  written, as in (4), in terms of irreducible, quasi-square-integrable representations,  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$ . Recall that  $\pi_i \in \mathcal{S}\mathcal{G}_F$  is a representation of  $\mathrm{GL}(n_i, F)$  for  $1 \leq i \leq r$ . Let  $\chi$  be a quasi-character of  $F^{\times}$ .*

- (1) *We have  $\delta_{\chi}(\pi) = 0$  if  $n_i \nmid a(\pi_i)$  for each  $1 \leq i \leq r$ .*

- (2) If there exists an  $1 \leq i \leq r$  such that  $n_i \mid a(\pi)$  then  $\delta_\chi(\pi_i) = 0$  if  $a(\pi_i) \neq n_i a(\chi)$ .
- (3) Suppose there exists an  $1 \leq i \leq r$  such that  $a(\pi_i) = n_i a(\chi)$ , then  $\delta_\chi(\pi_i) = 0$  if and only if  $a(\chi \mu_i) = a(\chi)$  where  $\pi_i = \mu_i \pi_i^{\min}$  is written as the  $\mu_i$ -twist of a twist minimal representation  $\pi_i^{\min}$ .

Our second port of call is to quantify the rarity described in Lemma 4.5.

**Lemma 4.6** (Quantitative interference). *Let  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$  as in Lemma 4.5. Suppose  $\chi \in \mathfrak{X}$  and that for some  $1 \leq i \leq r$  we have  $\delta_\chi(\pi_i) \neq 0$ . Write  $\pi_i = \mu_i \pi_i^{\min}$  as per part (3) of Lemma 4.5. Then, for each  $0 < j \leq a(\pi_i) - a(\pi_i^{\min})$  satisfying  $j \equiv a(\pi_i) \pmod{n_i}$ , there are precisely*

$$\#\mathfrak{X}\left(\frac{a(\pi_i) - j}{n}\right) \quad (20)$$

characters  $\chi \in \mathfrak{X}$  such that  $\delta_\chi(\pi_i) = a(\pi_i) - j$ . The number of  $\chi \in \mathfrak{X}(a(\pi_i)/n)$  satisfying  $\delta_\chi(\pi_i) = a(\pi_i)$  is

$$(q-2) \cdot \#\mathfrak{X}\left(\frac{a(\pi_i)}{n} - 1\right). \quad (21)$$

*Proof.* The number in (20) is determined by the necessity that

$$\chi \in \mu_i^{-1} \mathfrak{X}\left(\frac{a(\pi_i) - j}{n}\right).$$

Similarly, we count up to the number in (21) by observing that  $\chi \in \mathfrak{X}(a(\pi_i)/n_i)$  but  $\chi$  is not an element of  $\mathfrak{X}((a(\pi_i)/n_i) - 1)$  nor  $\mu_i^{-1} \mathfrak{X}((a(\pi_i)/n_i) - 1)$ .  $\square$

**Proposition 4.7** (Dominant behaviour). *In each case of Lemma 4.5 for which  $\chi$  and  $\pi = \pi_1 \boxplus \cdots \boxplus \pi_r$  satisfy  $\delta_\chi(\pi) = 0$ , we have the “dominant” conductor formula*

$$a(\chi\pi) = \sum_{i=1}^r \max\{a(\pi_i), n_i a(\chi)\}. \quad (22)$$

#### ACKNOWLEDGEMENT

The author would like to express gratitude to the Max-Planck-Institut für Mathematik, Bonn for providing welcoming hospitality and an invigorating mathematical environment in which this project was conceived and birthed.

#### REFERENCES

1. F. Brunault, *On the ramification of modular parametrizations at the cusps*, J. Théor. Nombres Bordeaux **28** (2016), no. 3, 773–790.
2. C. Bushnell and G. Henniart, *An upper bound on conductors for pairs*, J. Number Theory **65** (1997), no. 2, 183–196.
3. C. Bushnell, G. Henniart, and P. Kutzko, *Local Rankin-Selberg convolutions for  $GL_n$ : explicit conductor formula*, J. Amer. Math. Soc. **11** (1998), no. 3, 703–730.
4. A. Corbett and A. Saha, *On the order of vanishing of newforms at cusps*, Math. Res. Lett. (2018), To Appear.

5. P. Deligne, D. Kazhdan, and M.-F. Vignéras, *Représentations des algèbres centrales simples  $p$ -adiques*, Representations of reductive groups over a local field, Travaux en Cours, Hermann, Paris, 1984, pp. 33–117.
6. I. Gelfand and D. Každan, *Representations of the group  $GL(n, K)$  where  $K$  is a local field*, Lie groups and their representations (Proc. Summer School, Bolyai János Math. Soc., Budapest, 1971), Halsted, New York, 1975, pp. 95–118.
7. R. Godement and H. Jacquet, *Zeta functions of simple algebras*, Lecture Notes in Mathematics, Vol. 260, Springer-Verlag, Berlin-New York, 1972.
8. M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by V. Berkovich.
9. H. Jacquet, *Generic representations*, Non-commutative harmonic analysis (Actes Colloq., Marseille-Luminy, 1976), Springer, Berlin, 1977, pp. 91–101. Lecture Notes in Math., Vol. 587.
10. H. Jacquet and R. Langlands, *Automorphic forms on  $GL(2)$* , Lecture Notes in Mathematics, Vol. 114, Springer-Verlag, Berlin-New York, 1970.
11. H. Jacquet, I. Piatetski-Shapiro, and J. Shalika, *Conducteur des représentations du groupe linéaire*, Math. Ann. **256** (1981), no. 2, 199–214.
12. H. Koch and E.-W. Zink, *Zur Korrespondenz von Darstellungen der Galoisgruppen und der zentralen Divisionsalgebren über lokalen Körpern (der zahme Fall)*, Math. Nachr. **98** (1980), 83–119.
13. P. Nelson, A. Pitale, and A. Saha, *Bounds for Rankin–Selberg integrals and quantum unique ergodicity for powerful levels*, J. Amer. Math. Soc. **27** (2014), no. 1, 147–191.
14. D. Prasad and A. Raghuram, *Representation theory of  $GL(n)$  over non-Archimedean local fields*, School on Automorphic Forms on  $GL(n)$ , ICTP Lect. Notes, vol. 21, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2008, pp. 159–205.
15. J. Rogawski, *Representations of  $GL(n)$  and division algebras over a  $p$ -adic field*, Duke Math. J. **50** (1983), no. 1, 161–196.
16. D. Rohrlich, *Elliptic curves and the Weil–Deligne group*, Elliptic curves and related topics, CRM Proc. Lecture Notes, vol. 4, Amer. Math. Soc., Providence, RI, 1994, pp. 125–157.
17. A. Saha, *On sup-norms of cusp forms of powerful level*, ArXiv e-prints (2014), Online.
18. ———, *Hybrid sup-norm bounds for maass newforms of powerful level*, ArXiv e-prints (2015), Online.
19. ———, *Large values of newforms on  $GL(2)$  with highly ramified central character*, Int. Math. Res. Not. IMRN (2016), no. 13, 4103–4131.
20. J. Tate, *Fourier analysis in number fields, and Hecke’s zeta-functions*, Algebraic Number Theory (Proc. Instructional Conf., Brighton, 1965), Thompson, Washington, D.C., 1967, pp. 305–347.
21. J.-P. Tignol and A. Wadsworth, *Value functions on simple algebras, and associated graded rings*, Springer Monographs in Mathematics, Springer, Cham, 2015.
22. J. Tunnell, *On the local Langlands conjecture for  $GL(2)$* , Invent. Math. **46** (1978), no. 2, 179–200.
23. T. Wedhorn, *The local Langlands correspondence for  $GL(n)$  over  $p$ -adic fields*, School on Automorphic Forms on  $GL(n)$ , ICTP Lect. Notes, vol. 21, Abdus Salam Int. Cent. Theoret. Phys., Trieste, 2008, pp. 237–320.
24. A. Weil, *Basic number theory*, Classics in Mathematics, Springer-Verlag, Berlin, 1973, Second edition.
25. A. Zelevinsky, *Induced representations of reductive  $p$ -adic groups. II. On irreducible representations of  $GL(n)$* , Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 2, 165–210.