

Determinant of Laplacian on tori of constant positive curvature with one conical point

Victor Kalvin, Alexey Kokotov

July 16, 2018

Abstract. We find an explicit expression for the zeta-regularized determinant of (the Friedrichs extensions) of the Laplacians on a compact Riemann surface of genus one with conformal metric of curvature 1 having a single conical singularity of angle 4π .

1 Introduction

Let X be a compact Riemann surface of genus one and let $P \in X$. According to [1], Cor. 3. 5. 1, there exists *at most* one conformal metric on X of constant curvature 1 with a (single) conical point of angle 4π at P . The following simple construction shows that such a metric, $m(X, P)$, in fact always exists (and due to [1] is unique).

Consider the spherical triangle $T = \{(x_1, x_2, x_3) \in S^2 \subset \mathbb{R}^3 : x_1 \geq 0, x_2 \geq 0, x_3 \geq 0\}$ with all three angles equal to $\pi/2$. Gluing two copies of T along their boundaries, we get the Riemann sphere \mathbb{P} with metric m of curvature 1 and three conical points P_1, P_2, P_3 of conical angle π . Consider the two-fold covering

$$\mu : X(Q) \rightarrow \mathbb{P} \tag{1.1}$$

ramified over P_1, P_2, P_3 and some point $Q \in \mathbb{P} \setminus \{P_1, P_2, P_3\}$. Lifting the metric m from \mathbb{P} to the compact Riemann surface $X(Q)$ of genus one via μ , one gets the metric μ^*m on $X(Q)$ which has curvature 1 and the unique conical point of angle 4π at the preimage $\mu^{-1}(Q)$ of Q . Clearly, any compact surface of genus one is (biholomorphically equivalent to) $X(Q)$ for some $Q \in \mathbb{P} \setminus \{P_1, P_2, P_3\}$. Now let X be an arbitrary compact Riemann surface of genus one and let P be any point of X . Take $Q \in \mathbb{P}$ such that $X = X(Q)$ and consider the automorphism $\alpha : X \rightarrow X$ (the translation) of X sending P to $\mu^{-1}(Q)$. Then

$$m(X, P) = \alpha^*(\mu^*(m)) = (\mu \circ \alpha)^*(m).$$

Now introduce the scalar (Friedrichs) self-adjoint Laplacian $\Delta(X, P) := \Delta^{m(X, P)}$ on X corresponding to the metric $m(X, P)$. For any P and Q from X the operators $\Delta(X, P)$ and $\Delta(X, Q)$ are isospectral and, therefore, the ζ -regularized (modified, i. e. with zero

modes excluded) determinant $\det\Delta(X, P)$ is independent of $P \in X$ and, therefore, is a function on moduli space \mathcal{M}_1 of Riemann surfaces of genus one. The main result of the present work is the following explicit formula for this function:

$$\det\Delta(X, P) = C_1 |\Im\sigma| |\eta(\sigma)|^4 F(t) = C_2 \det\Delta^{(0)}(X) F(t), \quad (1.2)$$

where σ is the b -period of the Riemann surface X , C_1 and C_2 are absolute constants, η is the Dedekind eta-function, $\Delta^{(0)}$ is the Laplacian on X corresponding to the flat conformal metric of unit volume, the surface X is represented as the two-fold covering of the Riemann sphere $\mathbb{C}P^1$ ramified over the points $0, 1, \infty$ and $t \in \mathbb{C} \setminus \{0, 1\}$, and

$$F(t) = \frac{|t|^{\frac{1}{24}} |t-1|^{\frac{1}{24}}}{(|\sqrt{t}-1| + |\sqrt{t}+1|)^{\frac{1}{4}}}. \quad (1.3)$$

As it is well-known, the moduli space \mathcal{M}_1 coincides with the quotient space

$$(\mathbb{C} \setminus \{0, 1\}) / G,$$

where G is a finite group of order 6, generated by transformations $t \rightarrow \frac{1}{t}$ and $t \rightarrow 1-t$. A direct check shows that $F(t) = F(\frac{1}{t})$ and $F(t) = F(1-t)$ and, therefore, the right hand side of (1.2) is in fact a function on \mathcal{M}_1 .

Remark 1. *Using the classical relation (see, e. g. [2] (3.35))*

$$t = - \left(\frac{\Theta_{[0]}^1(0|\sigma)}{\Theta_{[1]}^0(0|\sigma)} \right)^4,$$

one can rewrite the right hand side as a function σ only.

The classical (see [9]) relation $\det\Delta^{(0)} = C |\Im\sigma| |\eta(\sigma)|^4$ used in (1.2), implies that (1.2) can be considered as a version of Polyakov's formula (relating determinants of the Laplacians corresponding to two *smooth* metrics in the same conformal class) for the case of two conformally equivalent metrics on a torus: one of them is smooth and flat, another is of curvature one and has one (very special) singular point.

2 Metrics on the base and on the covering

Here we find an explicit expression for the metric m on the Riemann sphere $\mathbb{P} = \mathbb{C}P^1$ of curvature 1 and with three conical singularities at $P_1 = 0$, $P_2 = 1$ and $P_3 = \infty$.

The stereographic projection (from the south pole) maps the spherical triangle T onto quarter of the unit disk $\{z \in \mathbb{C}; |z| \leq 1, 0 \leq \text{Arg } z \leq \pi/2\}$. The conformal map

$$z \mapsto w = \left(\frac{1+z^2}{1-z^2} \right)^2 \quad (2.1)$$

sends this quarter of the disk to the upper half-plane H ; the corner points $i, 0, 1$ go to the points $0, 1$ and ∞ on the real line. The push forward of the standard round metric

$$\frac{4|dz|^2}{(1+|z|^2)^2}$$

on the sphere by this map gives rise to the metric

$$m = \frac{|dw|^2}{|w||w-1|(|\sqrt{w}+1|+|\sqrt{w}-1|)^2} \quad (2.2)$$

on H ; clearly, the latter metric can be extended (via the same formula) to $\mathbb{C}P^1$. The resulting curvature one metric on $\mathbb{C}P^1$ (also denoted by m) has three conical singularities of angle π : at $w = 0$, $w = 1$ and $w = \infty$.

Consider a two-fold covering of the Riemann sphere by a compact Riemann surface $X(t)$ of genus 1

$$\mu : X(t) \rightarrow \mathbb{C}P^1 \quad (2.3)$$

ramified over four points: $0, 1, \infty$ and $t \in \mathbb{C} \setminus \{0, 1\}$. Clearly, the pull back metric μ^*m on $X(t)$ is a curvature one metric with exactly one conical singularity. The singularity is a conical point of angle 4π located at the point $\mu^{-1}(t)$.

3 Determinant of Laplacian as function of critical value t

The analysis from [3] in particular implies that one can introduce the standard Ray-Singer ζ -regularized determinant of the (Friedrichs) self-adjoint Laplacian Δ^{μ^*m} in $L_2(X(t), \mu^*m)$

$$\det \Delta^{\mu^*m} := \exp\{-\zeta'_{\Delta^{\mu^*m}}(0)\},$$

where $\zeta'_{\Delta^{\mu^*m}}$ is the operator zeta-function. In this section we establish a formula for the variation of $\zeta'_{\Delta^{\mu^*m}}(0)$ with respect to the parameter t (the fourth ramification point of the covering (2.3)). The derivation of this formula coincides almost verbatim with the proof of [3, Proposition 6.1], therefore, we will give only few details.

For the sake of brevity we identify the point t of the base $\mathbb{C}P^1$ with its (unique) preimage $\mu^{-1}(t)$ on $X(t)$.

Let $Y(\lambda; \cdot)$ be the (unique) special solution of the Helmholtz equation (here λ is the spectral parameter) $(\Delta^m - \lambda)Y = 0$ on $X \setminus \{t\}$ with asymptotics $Y(\lambda)(x) = \frac{1}{x} + O(x)$ as $x \rightarrow 0$, where $x(P) = \sqrt{\mu(P) - t}$ is the distinguished holomorphic local parameter in a vicinity of the ramification point $t \in X(t)$ of the covering (2.3). Introduce the complex-valued function $\lambda \mapsto b(\lambda)$ as the coefficient near x in the asymptotic expansion

$$Y(x, \bar{x}; \lambda) = \frac{1}{x} + c(\lambda) + a(\lambda)\bar{x} + b(\lambda)x + O(|x|^{2-\epsilon}) \text{ as } x \rightarrow 0.$$

The following variational formula is proved in [3, Proposition 6.1]:

$$\partial_t(-\zeta'_{\Delta^{\mu^*m}}(0)) = \frac{1}{2}(b(0) - b(-\infty)). \quad (3.1)$$

The value $b(0)$ is found in [3, Lemma 4.2]: one has the relation

$$b(0) = -\frac{1}{6}S_{Sch}(x)\Big|_{x=0}, \quad (3.2)$$

where S_{Sch} is the Schiffer projective connection on the Riemann surface $X(t)$.

Since $\lambda = -\infty$ is a local regime, in order to find $b(-\infty)$ the solution Y can be replaced by a local solution with the same asymptotic as $x \rightarrow 0$. A local solution \hat{Y} with asymptotic

$$\hat{Y}(u, \bar{u}; \lambda) = \frac{1}{u} + \hat{c}(\lambda) + \hat{a}(\lambda)\bar{u} + \hat{b}(\lambda)u + O(|u|^{2-\epsilon}) \text{ as } u \rightarrow 0$$

in the local parameter $u^2 = z - s$ was constructed in [3, Lemma 4.1] by separation of variables; here z and $w = \mu(P)$ (resp. s and t) are related by (2.1) (resp. by (2.1) with $z = s$ and $w = t$) and $\hat{b}(-\infty) = \frac{1}{2} \frac{\bar{s}}{1+|s|^2}$. One can easily find the coefficients $A(t)$ and $B(t)$ of the Taylor series $u = A(t)x + B(t)x^3 + O(x^5)$. As a local solution replacing Y we can take $A(t)\hat{Y}$. This immediately implies $b(-\infty) = A^2(t)\hat{b}(-\infty) - B(t)/A(t)$. A straightforward calculation verifies that

$$b(-\infty) = \partial_t \log \left(|t||t-1|(|\sqrt{t}+1|+|\sqrt{t}-1|)^2 \right)^{1/4}. \quad (3.3)$$

Observe that the right hand side in (3.3) is actually the value of $\partial_w \log \rho(w, \bar{w})^{-1/4}$ at $w = t$, where $\rho(w, \bar{w})$ is the conformal factor of the metric (2.2); this is also a direct consequence of [8, Lemma 4].

Using (3.1) together with (3.2) and (3.3), we are now able to derive an explicit formula for $\det \Delta^{\mu^* m}$.

4 Explicit formula for the determinant

Equations (3.1), (3.2) and (3.3) imply that the determinant of the Laplacian $\det \Delta^{\mu^* m} = \exp\{-\zeta'_{\Delta^{\mu^* m}}(0)\}$ can be represented as a product

$$\det \Delta^{\mu^* m} = C |\Im \sigma| |\tau(t)|^2 \left| \frac{1}{|t||t-1|(|\sqrt{t}+1|+|\sqrt{t}-1|)^2} \right|^{1/8} \quad (4.1)$$

where $\tau(t)$ is the value of the Bergman tau-function (see [4], [5], [6]) on the Hurwitz space $H_{1,2}(2)$ of two-fold genus one coverings of the Riemann sphere, having ∞ as a ramification point at the covering, ramified over $1, 0, \infty$ and t . More specifically, τ is a solution of the equation

$$\frac{\partial \log \tau}{\partial t} = -\frac{1}{12} S_B(x)|_{x=0}$$

where S_B is the Bergman projective connection on the covering Riemann surface $X(t)$ of genus one and x is the distinguished holomorphic parameter in a vicinity of the ramification point t of $X(t)$. We remind the reader that the Bergman and the Schiffer projective connections are related via the equation

$$S_{Sch}(x) = S_B(x) - 6\pi(\Im \sigma)^{-1} v^2(x)$$

where v is the normalized holomorphic differential on $X(t)$ and that the Rauch variational formula (see, e. g., [4]) implies the relation

$$\frac{\partial \log \Im \sigma}{\partial t} = \frac{\pi}{2} (\Im \sigma)^{-1} v^2(x)|_{x=0}.$$

The needed explicit expression for τ can be found e. g. in [6, f-la (18)] (it is a very special case of the explicit formula for the Bergman tau-function on general coverings of arbitrary genus and degree found in [5] as well as of a much earlier formula of Kitaev and Korotkin for hyperelliptic coverings [7]). Namely, [6, f-la (18)] implies that

$$\tau = \eta^2(\sigma) \left[\frac{v(\infty)^3}{v(P_1)v(P_2)v(Q)} \right]^{\frac{1}{12}}, \quad (4.2)$$

where P_1 and P_2 are the points of the $X(t)$ lying over 0 and 1, Q is a point of $X(t)$ lying over t and ∞ denotes the point of the covering curve $X(t)$ lying over the point at infinity of the base $\mathbb{C}P^1$; v is an arbitrary nonzero holomorphic differential on $X(t)$; and, say, $v(P_1)$ is the value of this differential in the distinguished holomorphic parameter at P_1 . (One has to take into account that $\tau = \tau_I^{-2}$, where τ_I is from [6].) Taking

$$v = \frac{dw}{\sqrt{(w(w-1)(w-t))}},$$

and using the following expressions for the distinguished local parameters at P_1, P_2, Q and ∞

$$x = \sqrt{w}; \quad x = \sqrt{w-1}; \quad x = \sqrt{w-t}; \quad x = \frac{1}{\sqrt{w}}$$

one arrives at the relations (where \sim means = up to insignificant constants like ± 2 , etc.)

$$v(P_1) \sim \frac{1}{\sqrt{t}}; \quad v(P_2) \sim \frac{1}{\sqrt{t-1}}; \quad v(Q) \sim \frac{1}{\sqrt{t(t-1)}}; \quad v(\infty) \sim 1.$$

These relations together with (4.2) and (4.1) imply (1.2).

References

- [1] Ching-Li Chai, Chang-Shou Lin, Chin-Lung Wang, Mean field equation, hyperelliptic curves and modular forms: I, Cambridge Journal of Mathematics, Vol. 3, N 1-2, 2015
- [2] C. Clemens, A scrapbook of complex curve theory, Grad. Studies in Math., Vol 55
- [3] V. Kalvin, A. Kokotov, Metrics of constant positive curvature, Hurwitz spaces and $\det\Delta$, IMRN, 2018; in press; arXiv:1612.08660
- [4] A. Kokotov, D. Korotkin, Tau-functions on Hurwitz spaces, Mathematical Physics, Analysis and Geometry, 7 (2004), no. 1, 47–96.
- [5] A. Kokotov, D. Korotkin, Isomonodromic tau-function of Hurwitz Frobenius manifolds, Int. Math. Res. Not. IMRN (2006), pp. 1-34
- [6] A.Kokotov, I. Strachan, On the isomonodromic tau-function for the Hurwitz spaces of branched coverings of genus zero and one, Mathematical Research Letters, 12, 2005, no. 5-6, 857-875.
- [7] V. Kitaev, D. Korotkin, On solutions of the Schlesinger equations in terms of theta-functions, International Mathematics Research Notices, 1998, no. 17,877-905.
- [8] V. Kalvin, On Determinants of Laplacians on Compact Riemann Surfaces Equipped with Pullbacks of Conical Metrics by Meromorphic Functions, in preparation
- [9] J. Polchinski, Evaluation of the one loop string path integral. Comm. Math. Phys. 104 (1986), no. 1, 37–47